



Main Manuscript for

On the Stratospheric Chemistry Of Mid-Latitude Wildfire Smoke

Susan Solomon^{a,*}, Kimberlee Dube^b, Kane Stone^a, Pengfei Yu^c, Doug Kinnison^d, Owen B. Toon^e, Susan E. Strahan^f, Karen H. Rosenlof^g, Robert Portmann^g, Sean Davis^g, William Randel^d, Peter Bernath^{h,i}, Chris Boone^h, Charles G. Bardeen^d, Adam Bourassa^b, Daniel Zawada^b, Doug Degenstein^b

^aDepartment of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139; ^bInstitute for Space and Atmospheric Science, University of Saskatchewan, Saskatoon, Saskatchewan, Canada; ^cInstitute for Environmental and Climate Research, Jinan University, Guangzhou, China; ^dAtmospheric Chemistry Observations and Modeling, National Center for Atmospheric Research, Boulder, CO 80307; ^eDepartment of Atmospheric and Oceanic Sciences and Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303; ^fGESTAR, NASA Goddard Space Flight Center, Greenbelt, MD, 20771; ^gNOAA Chemical Sciences Laboratory, Boulder CO 80305; ^hDepartment of Chemistry and Biochemistry, Old Dominion University, Norfolk, VA 23529; ⁱDepartment of Chemistry, University of Waterloo, Waterloo, Ontario, Canada

*Susan Solomon

Email: solos@mit.edu

Author Contributions: S.S., K.D., K.S., P.Y., O.B.T., and S.E.S conceptualized the work; P.B., C.B., D.Z., and D.D. contributed to the data acquisition; R.P., K.H.R, C.G.B. and P.Y devised the model runs; A. B., K.D., K.S., and P.Y. conducted the data and model analysis; S.S., K. D., and K. S. drafted the manuscript following discussions with author team members from all institutions; and all authors revised the paper.

Competing Interest Statement: The authors declare no conflict of interest.

Classification: Physical Sciences – Earth, Atmospheric, and Planetary Sciences

Keywords: Stratospheric ozone, smoke, wildfire, chemistry

Abstract

Massive Australian wildfires lofted smoke directly into the stratosphere in the austral summer of 2019/20. The smoke led to increases in optical extinction throughout the mid-latitudes of the southern hemisphere that rivalled substantial volcanic perturbations. Previous studies have assumed that the smoke became coated with sulfuric acid and water, and would deplete the ozone layer through heterogeneous chemistry on those surfaces, as is routinely observed following volcanic enhancements of the stratospheric sulfate layer. Here observations of extinction and reactive nitrogen species from multiple independent satellites that sampled the smoke region are compared to one another and to model calculations for the first time. The data display a strong decrease in reactive nitrogen concentrations with increased aerosol extinction in the stratosphere, which is a known fingerprint for key heterogeneous chemistry on sulfate/H₂O particles (specifically the hydrolysis of N₂O₅ to form HNO₃). This chemical shift affects not only reactive nitrogen but also chlorine and reactive hydrogen species, and is expected to cause mid-latitude ozone layer depletion. Comparison of the model ozone to observations suggests that N₂O₅ hydrolysis contributed to reduced ozone, but additional chemical and/or dynamical processes are also important. These findings suggest that if wildfire smoke injection into the stratosphere increases sufficiently in frequency and magnitude as the world warms due to climate change, ozone recovery under the Montreal Protocol could be impeded, at least sporadically. Modelled austral mid-latitude total ozone loss was about 1% in March 2020, which is significant compared to expected ozone recovery of about 1% per decade.

Significance Statement

Large wildfires have been observed to inject smoke into the stratosphere, raising questions about their potential to affect the stratospheric ozone layer that protects life on Earth from biologically-damaging ultraviolet radiation. Multiple observations of aerosol and NO₂ concentrations from three independent satellite instruments are used here together with model calculations to identify decreases in stratospheric NO₂ concentrations following major Australian 2019/20 wildfires. The data confirm that important chemistry did occur on the smoke particle surfaces. The observed behavior in NO₂ with increasing particle concentrations is a marker for surface chemistry that contributes to mid-latitude ozone depletion. The results indicate that increasing wildfire activity in a warming world may slow the recovery of the ozone layer.

Main Text

Introduction

Recent large-scale wildfire events in many parts of the world including British Columbia in 2017 (1) and the Australian “black summer” fires in 2020 (2,3) have injected substantial loadings of smoke particles directly into the stratosphere via the outflow from towering Pyrocumulonimbus (PyroCb) towers. While wildfires have occurred for many thousands of years, evidence suggests that their scale and frequency are increasing with global warming (4). The properties and composition of wildfire smoke particles have been studied in the troposphere and stratosphere using both in-situ and lidar methods (e.g., 5, 6, 7, 8) and stratospheric smoke plume heights have been documented with satellite observations (9,10). Further, repeated transects through fire smoke observed by instruments onboard passenger aircraft have shown that wildfire smoke particles acquire a liquid coating (11), a finding supported by multi-wavelength lidar studies (5, 12). Satellite observations revealed large increases in organic species in the gas phase associated with stratospheric smoke (13), including acetone and CH₃OH (as would be expected due to incomplete combustion in biomass burning). They also provided spectral evidence that the smoke particles themselves contained organic material, (e.g., carboxylic acids as noted in wood smoke in ref. 14). Stratospheric single-particle measurements of smoke have detected

internally mixed particles containing not only organic compounds (i.e., black and organic carbon) but also sulfates (8).

Stratospheric aerosol extinction ratios (relative to a purely molecular atmosphere) associated with the 2020 Australian fires were comparable to those following the eruption of the Calbuco volcano in April 2015 and blanketed the Southern Hemisphere (SH) mid-latitudes (3,15). Even under non-volcanic conditions, the stratosphere contains a layer of liquid sulfuric acid/water particles that can drive significant mid-latitude ozone depletion chemistry (16). Sufficiently explosive volcanic eruptions inject additional sulfur into the stratosphere, which ultimately increases the sulfuric acid abundances and can greatly enhance the particle surface areas. Reactions occurring on such particles affect reactive nitrogen (NO and NO_2 ; the sum of the two is called NO_x here). NO and NO_2 exchange rapidly with one another in the daytime stratosphere depending on ozone abundances, temperatures, and photolysis rates. Therefore, the sum of both species is more robust to variability in temperature, ozone, or solar angle than either alone. NO_x reductions in turn affect ClO and OH radicals, and all of these species participate in catalytic cycles that deplete ozone. Major volcanic eruptions of the past half-century have been shown to enhance mid-latitude stratospheric ozone destruction (17,18,19,20). The 2015 Calbuco event resulted in observable reductions in mid latitude SH ozone concentrations in the lower stratosphere, consistent with calculations of chemical depletion (21). The 2020 Australian fires were associated with similar SH ozone reductions (15). The smoke's radiative properties also locally warmed the lower stratosphere by up to a few degrees (15, 22).

A detailed model study of the 2020 Australian wildfire particles assumed that the particles became coated with sulfuric acid (22) and hence displayed similar mid-latitude chemistry to background and volcanic stratospheric aerosols (see Methods). That work estimated that heterogeneous reactions involving wildfire-enhanced aerosols could reduce SH mid-latitude stratospheric ozone by about 5-10 Dobson Units (DU) from July-August of 2020. However, some studies have argued that wildfire smoke might form glassy surfaces (see reference 6) in the lowermost stratosphere, which would likely display quite different chemical reactivity from liquids. Tropospheric studies have shown that wildfire particles contain differing mixes of soot, primary organics, and secondary organic compounds as well as minerals and salts (23), dependent on such factors as the type of fuel (rainforest, woodland, etc.) and state of the fire (smoldering, flaming, etc.). While some studies have suggested somewhat reduced uptake of N_2O_5 when aerosols are coated with organics (e.g., 24, 25), other work indicates differing behavior depending upon specific composition and such factors as whether the organic coatings are straight chain or branched (e.g., 26). Therefore, observations and modelling studies that can improve the understanding of the impacts of wildfire smoke on stratospheric composition and chemistry are needed, and are the goal of this paper.

We use satellite observations of NO_x species and aerosols together with model simulations (from 22) to examine the role of the 2020 Australian wildfire smoke in mid-latitude stratospheric NO_x chemistry. The abundance of stratospheric NO_x has long been known to be a key marker for mid-latitude heterogeneous chemistry on liquid sulfate aerosols, particularly when aerosols are enhanced (e.g., in major volcanic eruptions, 17, 27) as described in detail below. We demonstrate that the satellite NO_x observations provide strong evidence that stratospheric wildfire smoke drives important chemistry that can be expected to contribute to ozone depletion as long as stratospheric chlorine abundances remain elevated.

Results

We make use of three satellite records to examine the behavior of stratospheric reactive nitrogen after the Australian fires, i.e., NO_2 data from the Optical Spectrograph and InfraRed Imager System (OSIRIS, see reference 28) the Stratospheric Aerosol and Gases Experiment on the International Space Station (SAGEIII/ISS, reference 29), as well as the Atmospheric Chemistry

Experiment (ACE, reference 30). We also present ≈ 750 nm extinction ratio data from both OSIRIS and SAGE III, as well as from the Ozone Mapping and Profiler Suite Limb Profiler (OMPS/LP) from (31, 32). SAGE III/ISS and OSIRIS both employ absorption for NO_2 measurement at visible wavelengths, while ACE uses Fourier transform infrared spectroscopy. All three make use of limb-viewing geometry, either through direct solar occultation (SAGE III/ISS and ACE) or limb scattering (OSIRIS).

Figure 1 presents monthly averaged stratospheric aerosol extinction ratio time series from the three instruments, demonstrating broad consistency between the datasets. The observations display a large perturbation to SH mid-latitude aerosol extinction ratio due to the 2020 Australian fires. As noted in previous studies using OMPS (3), the mid-latitude aerosol extinction ratio perturbation after these fires was comparable to that following the substantial eruption of Mt. Calbuco in 2015; the OSIRIS and SAGE data shown in Figure 1 provide independent support for this conclusion. The eruption of the Ulawun volcano in the tropics in 2019 affected the aerosol loading in the tropics shortly before the 2020 fires in all three datasets. While coverage in OSIRIS and SAGE III/ISS is more limited due to their limb viewing geometries, the three instruments nonetheless suggest similar timing and spread of the 2020 wildfire smoke. Figure S1 presents the extinction ratios at 675 nm as estimated by the model in (22), and shows good general agreement with OMPS despite the small difference in the wavelengths available for each. Smoke particles accumulate water, increasing extinction and providing added surface area to drive faster heterogeneous chemistry, just as added sulfate does following volcanic eruptions, but with different hygroscopicity (see Methods).

Figure 2 displays the corresponding monthly averaged satellite NO_x anomalies obtained from the OSIRIS and SAGE III/ISS NO_2 data at 18.5 km (as described in Methods), along with the NO_x change calculated in the model. OSIRIS data indicate that 2020 NO_x was lower than all previous years since 2002 throughout a broad range of latitude for multiple months, from 30-60°S (see Figure 2), so this region was selected for focused study in this paper. OSIRIS data display larger variability at lower latitudes (particularly below about 19 km), making identification of wildfire impacts challenging equatorward of about 30°S. Further, Ulawun may have perturbed tropical NO_x , but the abrupt NO_x change in early 2020 (at least for latitudes poleward of 30°S) suggests that the fires dominated at these latitudes. SAGE III/ISS NO_x data show very similar timing and spread of the mid-latitude anomaly to OSIRIS, albeit with more limited coverage. ACE NO_2 data also have limited coverage, and are shown in Figure S2; these are not converted to NO_x here both because of coverage limitations and because ACE measures both NO and NO_2 directly but the NO retrieval is still under development. Nonetheless, ACE NO_2 data display consistent features to the other datasets. Observed 2020 anomalies in reactive nitrogen species at 18.5 km from 30-60°S relative to other available years are at least 20% or larger in all three satellite datasets, a large change. Magnitudes of the NO_x perturbations from OSIRIS and SAGE data are different in part because of differences in coverage as well as the number of available sunrise versus sunset data points in each. Figure 2 also shows the spread of the ensemble mean NO_x change at 18.5 km calculated in the model, defined as the difference between the smoke versus no smoke runs, and the model is in good general agreement with the data. Note that the model calculations shown here did not include the Ulawun eruption, and thus reflect purely the calculated NO_x change from smoke particle chemistry. A latitude height plot for March comparing the model and OSIRIS data is presented in Figure S3, again showing broad consistency between the model and the data. The 18.5 km altitude was selected for focus in this paper in order to balance OSIRIS data quality (better at higher rather than lower altitudes) and levels displaying extensive SH NO_x perturbations as shown in Figs. 2 and S3.

While dynamical contributions to the anomalies cannot be ruled out, Figure 3 (top panel) probes this region in more detail and shows that for February and March monthly averages at 18.5 km, the OSIRIS NO_x amounts are lower than observed in any previous year of the available 20-year record, strongly suggestive that the wildfires drove the change. ACE data presented in Figure S4

also display record lows for March and April 2020 in a record spanning 17 years. The chemical mechanism responsible is discussed further below. The OSIRIS data also indicate a large influence of the Calbuco aerosols on NO_x , beginning in the latter half of 2015 and extending into 2016. Indeed, while 2020 OSIRIS data show the lowest NO_x observed in this region in February and March, the second lowest is 2016 after Calbuco. By August of 2020 the wildfire smoke impact on NO_x appears to have diminished in the OSIRIS observations, and concentrations in the latter half of the year are within the range of other years. Figure 3 (bottom panel) presents the calculated NO_x concentrations from 20 realizations in the model for the smoke and no smoke cases. The OSIRIS observations for March indicate about 1×10^9 molec/ cm^3 after the fires versus about 1.4×10^9 in other years, and the model results are very close to these values.

The primary chemical mechanism driving NO_x reductions with increasing stratospheric sulfate aerosols and its implications for mid-latitude ozone losses have long been known (16, 17, 33). Even for background aerosols, these processes decrease mid-latitude ozone column abundances by several percent compared with estimates using only gas-phase chemistry for current levels of stratospheric chlorine loading. As chlorine abundances diminish in the future because of the phaseout of chlorofluorocarbons under the Montreal Protocol, the ozone depletion can be expected to decrease and eventually flip sign to positive values (34), but depletion can be expected through the mid-21st century. Further, these reactions were responsible for enhanced mid-latitude ozone destruction following several past volcanic eruptions (e.g., El Chichon and Pinatubo, see 19, 20).

The principal lower stratospheric photochemical mechanism is well established: NO is converted entirely to NO_2 at night, which goes on to form NO_3 and then to N_2O_5 . The NO_3 intermediate photolyzes rapidly in daytime so the formation of N_2O_5 is only rapid at night. Hence N_2O_5 is an important nighttime reservoir for NO_x . A critical reaction under warm mid-latitude conditions is the heterogeneous hydrolysis of N_2O_5 , which converts reactive nitrogen to HNO_3 – a process that does not occur in the gas phase. N_2O_5 photolyzes fairly rapidly during the day (order of hours), while HNO_3 photolysis is much slower in the lower stratosphere (order of a week or more). Nighttime conversion of N_2O_5 to HNO_3 therefore reduces NO_x , which in turn means that the NO_2 concentration available to form ClONO_2 is reduced, leading to an increase in ozone-destroying ClO. Reductions in NO_x influence HO_x radicals as well (19, 35), which are also important for ozone loss chemistry. N_2O_5 hydrolysis on sulfuric acid/water particles has been extensively studied in the laboratory, and occurs with high efficiency (36) at essentially all atmospheric temperatures. More recent studies have shown that BrONO_2 hydrolysis is also important for heterogeneous HNO_3 formation under these conditions (37), while ClONO_2 hydrolysis contributes at colder conditions (i.e., temperatures below about 195K, see 33). Here we use satellite observations to probe whether similar composition changes occur due to wildfire smoke. Because HNO_3 concentrations are much larger than those of NO_x at the altitude range considered, NO_x is a better indicator of this chemistry than HNO_3 would be. Observations also indicate that some HNO_3 was taken up by these particles (38), perhaps due to their high organic content (39).

A key point first made by (17) is the role of non-linear chemistry that occurs with increasing aerosol loading. While the rate of N_2O_5 hydrolysis increases rapidly at lower aerosol content, the reaction saturates when HNO_3 is formed fast enough to remove essentially all the N_2O_5 formed in a given night, due to slow release by HNO_3 photolysis the following day in the lower stratosphere. Further increases in aerosols then cannot significantly increase the reaction rate, because N_2O_5 is already being destroyed as fast as it can be produced (i.e., formation of NO_3 and hence N_2O_5 through the nighttime $\text{NO}_2 + \text{O}_3$ reaction becomes the rate-limiting step).

This heterogeneous chemistry leads to a characteristic curve of decreasing NO_x abundances versus increasing aerosols (see reference 17, Fig. 2), a diagnostic fingerprint of this chemistry. Figure 4 presents such curves for 40–45°S at 18.5 km using available SAGE III/ISS sunrise and sunset NO_x data, OSIRIS NO_x data, and NO_x calculated in the smoke model. Observations and

model results are de-seasonalized by month using all available years of data for each instrument (see Methods). High extinction values are observed without low NO_x in January, 2020 when the plume had freshly entered the stratosphere, suggesting that the timescale for the chemistry is of order one month. The 2020 observations reveal the expected decay, in agreement with the model. We note that the rate-limiting gas-phase reaction $\text{NO}_2 + \text{O}_3 \rightarrow \text{NO}_3 + \text{O}_2$ is faster by about 7% due to the smoke-induced warming in the March ensemble mean (212.15K versus 210.9K) at 40-45°S, 18.5 km. This temperature change is, however, only a small contribution to the modelled NO_x changes compared to the more than three-fold March surface area change and hence increased N₂O₅ hydrolysis rate due to the smoke. The OSIRIS data suggest that the peak 2020 NO_x reduction driven by the wildfires may have exceeded that from the Calbuco volcano, but it should be recalled that OSIRIS did not sample this region immediately after that eruption and did promptly sample the smoke. OSIRIS data suggest that the N₂O₅ hydrolysis reaction reached its saturation limit on the wildfire particles, a finding also suggested by the SAGEIII/ISS data albeit less clearly because of the limited coverage. The model is less clear regarding saturation but is in broad agreement with the decline. Figure S5 shows that similar behavior is observed in the ACE NO₂ record as well, despite limitations of coverage and differences in the wavelengths of the extinction measurements.

Figure 4 strongly supports the view that the Australian wildfire particles drive hydrolysis of N₂O₅ in a manner that is similar to sulfate particles. Thus, the presence of organic matter along with sulfate (15) apparently did not render the particles sufficiently glassy to inhibit the uptake of water needed to allow N₂O₅ hydrolysis. Figure 4 supports the view taken by Yu et al. (22) that the 2020 wildfire aerosols behaved like sulfate particles insofar as their mid-latitude heterogeneous chemistry is concerned.

We next compare modelled and observed midlatitude ozone changes but do not consider Antarctic ozone hole behavior. Polar stratospheric clouds (PSCs) are responsible for the extreme austral springtime ozone losses found in the Antarctic through heterogeneous chlorine and bromine chemistry (40, 41) and are enhanced after volcanic eruptions (42). They are composed in part of liquid sulfuric acid, water, and nitric acid. It is plausible that the Australian smoke particles may have enhanced PSC reactions, and perhaps influenced mid-latitude ozone indirectly through transport of reduced ozone values from the ozone hole but this chemistry is not examined here. The model used here did not include the wildfire aerosols in their PSC reaction set, allowing us to isolate the mid-latitude chemistry alone (i.e., as distinct from any transport from the ozone hole region at polar latitudes) with high confidence.

Figure 5 compares weekly and zonally averaged observed total ozone anomalies to the changes obtained between the smoke versus no smoke model runs (see Methods). The differences between smoke and no smoke runs explicitly isolates the impact of the chemistry included in the model, while the observational anomalies will reflect not only these chemical effects but any others that may be occurring, as well as any dynamical changes. The total ozone anomalies at southern mid-latitudes from the free-running ensemble mean of the model results display important similarities in morphology with time and latitude, but are considerably smaller than observed. Ozone reductions at low latitudes near -10 to -20S in the model are not observed in the data, perhaps due to dynamical variability or incomplete smoke chemistry. Low ozone anomalies (lowest 25th percentile of the record) occur near -50 to -55S in late March but are larger than that suggested in the model, and variable increases also seen near -40 to -55S in April to May suggest dynamical fluctuations. Notably, OMI observes low total ozone in the -40 to -50S latitude band from late May through August 1, too early in the year for substantial polar depletion (although transport from polar regions may well contribute later in the year). The model also suggests reduced ozone from smoke chemistry throughout this period at those latitudes, but the calculated ozone loss is smaller than observed. Dynamical variability could contribute to the observed changes, and it is plausible that the heating from the smoke plume altered the stratospheric circulation; these factors are not examined here. A combination of dynamical

variability and chemical contributions to the anomalous mid-latitude ozone changes may be occurring, or additional chemical processes on the smoke particles not considered here may be important. Overall, the comparison suggests that the smoke chemistry indicated by the NO_x observations as represented in this model did contribute to ozone reductions that appear to occur in the observations, but also shows that other factors are highly likely to be important.

Discussion

Multiple satellite datasets for stratospheric aerosol extinction and NO_x perturbations following the Australian wildfires of 2020 have been compared to one another and to recently published model calculations in this paper. Record-low NO_x abundances in the SH mid-latitude lower stratosphere were measured by OSIRIS and ACE. SAGE III/ISS observations only extend over about the past three years, but also display exceptionally low NO_x in 2020, comparable to the anomalies found in the other two instruments. The observed seasonal and latitudinal changes in NO_x near 18.5 km are broadly consistent with modelling results. Most important, the satellite data from both OSIRIS and SAGE III/ISS indicate large decreases in NO_x abundances which saturate with increasing aerosol extinction values, in good agreement with the model. This characteristic behavior has been previously studied (17). Its occurrence in three sets of satellite measurements presented here strongly suggests that the wildfire aerosols drove hydrolysis of N₂O₅ on wet particles. Thus, the evidence indicates that this key heterogeneous reaction was indeed enhanced on the Australian fire smoke surfaces, just as it is following major volcanic eruptions including e.g., Calbuco in the SH in 2015. The impact of that volcanic eruption on NO_x was also shown to be discernible in the OSIRIS data, but was smaller than that obtained following the 2020 fires in the monthly averages for February and March.

Decreased NO_x and NO₂ related to N₂O₅ hydrolysis is expected to be accompanied by increases in HO_x and ClO, which drives net decreases in mid-latitude ozone following major volcanic eruptions (19, 20). Decreases in mid-latitude SH ozone of up to 8 DU were also observed after the Australian fires, and these reductions began prior to the formation of the springtime Antarctic ozone hole, indicating a local origin rather than transport of low ozone air from the ozone hole. Model results display chemical ozone decreases from about -40 to -50S that are similar in their evolution from March through August but smaller in magnitude (with peak values of about 15 DU). The discrepancy may be due to dynamics, or to added chemistry not represented in the model. We note that the Australian smoke was unusual in that it came largely from eucalyptus trees (3,6), and whether similar chemistry can occur on other sources of stratospheric smoke more typical of other landscapes is not known. Laboratory work to elucidate the heterogeneous reaction rates that may occur in the stratosphere on mixed organic/sulfate particles is badly needed, as well as field and ongoing satellite observations to better understand their composition and chemistry.

Overall, this work provides strong evidence that the Australian forest fires of 2020 resulted in chemical impacts on mid-latitude stratospheric NO_x in a manner similar to that observed following volcanic eruptions. Modelled austral mid-latitude total ozone loss was about 1% in March 2020, which is significant in magnitude (albeit limited in space and time) as compared to expected ongoing southern hemisphere mid-latitude ozone recovery due to the Montreal Protocol of about 1% p[REDACTED]. The results suggest that this chemistry contributed to but did not fully capture the observed ozone changes following the fires. These findings are important given the uncertainties surrounding the chemistry that may occur on and in smoke particles. Our findings are suggestive that the Australian fire smoke did behave like sulfate aerosols and might therefore also have affected liquid PSCs and the Antarctic ozone loss in 2020 as well. Further work is required to examine that important chemistry and dynamics in detail. Finally, evidence strongly suggests that wildfire frequency and spatial extent has already increased, and will continue to increase in the future due to climate change until ecosystem changes reduce available fuels. Our

findings support the view that heterogeneous chemistry on wildfire smoke particles from pyroCb that reach the stratosphere represents a new and important chemistry-climate coupling mechanism that temporarily decreased SH mid-latitude ozone in 2020. Future fires in a warming world could display larger or more persistent impacts if wildfires become more frequent and/or intense, but there are many uncertainties including the chemical unknowns discussed herein. This paper has highlighted the need for further examination both of the chemistry of wildfire smoke in the stratosphere and the projected recovery of the ozone layer using coupled chemistry-climate-vegetation models.

Materials and Methods

NO₂ data are available since 2002 from the Optical Spectrograph and InfraRed Imager System (OSIRIS) instrument based upon limb-scattered solar radiation, version 7.1. OSIRIS data have previously been compared to other measurements including the solar occultation method for NO₂ employed by the Stratospheric Aerosol and Gases Experiment (SAGE III/ISS, version 5.1, see 29). Broad agreement between the SAGE III/ISS dataset and OSIRIS NO₂ has been demonstrated (43). The OSIRIS data for the morning orbit node are shifted to a common local solar time of 12:00 pm. OSIRIS and SAGE III/ISS data are converted to NO_x using the photochemical box model described in (44). NO₂ observations are also presented in the supplement from the Atmospheric Chemistry Experiment (ACE), which employs Fourier Transform Infrared (FTIR) spectroscopy in solar occultation mode at sunrise and sunset as well (45). Total ozone data are from the Ozone Monitoring Instrument (46), referred to as OMI.

Monthly mean values are calculated for each of the instrument datasets for any month containing at least five measurements. Months with fewer than five measurements in a given latitude-altitude bin are not shown. High latitude winter data are always missing in OSIRIS, SAGEIII/ISS, and ACE due to lack of sunlight for the measurement. SAGE III/ISS and ACE data are generally limited by their orbits and occultation opportunities. Other gaps indicate data dropouts, especially for the aging OSIRIS instrument. The data are de-seasonalized by subtracting the overall mean value for a given month of the year from that month (i.e. The overall mean January is subtracted from each individual January).

Extinction ratio data at approximately 750 nm are available from both OSIRIS and SAGE III, as well as a third instrument, the Ozone Mapping and Profiler Suite Limb Profiler (OMPS/LP). ACE includes two imagers which measure aerosol extinction at 525 nm and 1020 nm. OMPS extinction data shown here use the tomographic retrieval developed at the University of Saskatchewan (32). SAGE III/ISS and OSIRIS both measure limb extinction at visible/near infrared wavelengths, providing a useful comparison to the tomographic inversion approach used with OMPS.

Observations are compared to modelling results for the 2019-2020 Australian fires from the Community Aerosol and Radiation Model for Atmospheres (CARMA) coupled with the Community Earth System Model (CESM-CARMA) presented in reference (22). The model includes 56 vertical layers from the surface of the Earth to about 45 km, and a resolution of about 2° in latitude and longitude. The model was spun up in specified dynamics mode nudged to the Goddard Earth Observing System version 5 analysis (GEOS-5) from mid-summer to the end of 2019, after which 20 perturbed initial condition runs were carried out in free-running mode from December 29, 2019 to the end of 2020. Smoke was input from December 29–31, 2019 and on January 4, 2020, the dates when pyrocumulonimbus were observed in the stratosphere (47).

The 20 smoke ensemble members are paired with a control run (no smoke) with the same initial conditions. Several test cases were conducted to probe the sensitivity of the aerosol abundances and lofting to the amount of injected material and the percentage of black carbon in the initial smoke plume. Results are shown from the case that agreed best with observations, in which it

was assumed that the amount of smoke was 3 x that injected by the Pacific North-west wildfire events and 2.5% black carbon.

Heterogeneous chemistry and extinction are enhanced in hygroscopic particles that pick up water and swell, providing increased surface areas as well as water content. Like sulfate aerosols, organics are assumed to pick up water in the model, but with a lower hygroscopicity. The adopted hygroscopicity of sulfate is 0.8 while the adopted hygroscopicity of organics is 0.5 and that of black carbon is 0.1. Therefore, the smoke particle sizes are not as large as those that would occur for comparable sulfate particles. We determine the swelling of the mixed particles differently from pure sulfuric acid, based on weight percent calculation. Details are in reference 48 (A6.1 for pure sulfuric acid; A6.2 for mixed particles). Available model calculations of extinction ratio used here are at a slightly different wavelength than the observations, 675 nm.

Acknowledgments. SS and KS were funded in part by NSF 1906719 and 1848863. DK was funded in part by NASA grant (80NSSC19K0952). PY was funded by the second Tibetan Plateau Scientific Expedition and Research Program (2019QZKK0604) and by the second Tibetan Plateau Scientific Expedition and Research Program (2019QZKK0604) and National Science Foundation of China (42121004). OBT was funded by the Open Philanthropy Foundation. SES is funded by the NASA Atmospheric Composition Modeling and Analysis Program. This research was enabled by the computational and storage resources of NCAR's Computational and Information Systems Laboratory (CISL), sponsored by the NSF. Cheyenne: HPE/SGI ICE XA System (NCAR Community Computing). Boulder, CO: National Center for Atmospheric Research. <https://doi.org/10.5065/D6RX99HX>. The ORISIS and ACE missions and data products are funded by the Canadian Space Agency. The authors thank the OMPS teams at NASA GSFC for producing high quality OMPS Level 1 data that enable the University of Saskatchewan data product retrievals and the SAGE III/ISS team at NASA Langley for data production and advice.

Data Availability. Model output used is available at https://osf.io/6j8cb/?view_only=72f53447bf464a2bbcc1dfc32d492bab. OSIRIS data are available at <https://research-groups.usask.ca/osiris/data-products.php#OSIRISLevel2DataProducts>. SAGE III/ISS data are available at https://doi.org/10.5067/ISS/SAGEIII/SOLAR_HDF4_L2-V5.1. OMPS aerosol data are available at <https://zenodo.org/record/4029555>. ACE data are available through the following sign-up link: <https://database.scisat.ca/l2signup.php>. OMI ozone data were obtained from <https://acd-ext.gsfc.nasa.gov/anonftp/toms/>.

References

1. O. Torres *et al.*, Stratospheric injection of massive smoke plume From Canadian boreal fires in 2017 as Seen by DSCOVR-EPIC, CALIOP, and OMPS-LP Observations. *J. Geophys Res. Atmos.* **125**, e2020JD032579 doi:10.1029/2020JD032579 (2020).
2. S. Khaykin *et al.*, The 2019/20 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude. *Commun. Earth Environ.* **1**, 22 (2020).
3. D. A. Peterson, *et al.*, Australia's Black Summer pyrocumulonimbus super outbreak reveals potential for increasingly extreme stratospheric smoke events. *npj Clim Atmo Sci.* **4**, 1-16 (2021).
4. S. Seneviratne *et al.*, "Weather and climate extreme events in a changing climate", in IPCC sixth assessment report, chapter 11, WMO/UNEP, 2021.
5. H. Baars, *et al.*, The unprecedented 2017-2018 stratospheric smoke event: decay phase and aerosol properties observed with the EARLINET. *Atmos. Chem. Phys.* **19**, 15183-15198 (2019).

6. A. Ansmann, *et al.*, Tropospheric and stratospheric wildfire smoke profiling with lidar: mass, surface area, CCN, and INP retrieval. *Atmos. Chem. Phys.* **21**, 9779-9807 (2021).
7. M. Haarig *et al.*, Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke. *Atm. Chem. Phys.* **18**, 11847–11861 (2018).
8. D. M. Murphy *et al.*, Radiative and chemical implications of the size and composition of aerosol particles in the existing or modified global stratosphere. *Atmos. Chem. Phys.* **21**, 8915-8932 (2021).
9. M. Fromm, A. Tupper, D. Rosenfeld, R. Servranckx, R. McRae, Violent pyro-convective storm devastates Australia's capital and pollutes the stratosphere. *Geophys. Res. Lett.* **33**, 1-5 (2006).
10. D. A. Peterson *et al.*, Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke. *npj Clim Atmos Sci* **1**, 30 (2018).
11. J. Ditas *et al.*, Strong Impact of wildfires on the abundance and aging of black carbon in the lowermost stratosphere. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E11595-E11603 (2018).
12. R. Engelmann *et al.*, Wildfire smoke, Arctic haze, and aerosol effects on mixed-phase and cirrus clouds over the North Pole region during MOSAiC: an introduction. *Atmos. Chem. Phys.* **21**, 13,397-13,423 (2021).
13. C. D. Boone, P. F. Bernath, M. D. Fromm, Pyrocumulonimbus stratospheric plume injections measured by the ACE-FTS. *Geophys. Res. Lett.*, **47**, e2020GL088442. <https://doi.org/10.1029/2020GL088442> (2020).
14. M. Zhong, M. Jang, Dynamic light absorption of biomass-burning organic carbon photochemically aged under natural sunlight. *Atmos. Chem. Phys.* **14**, 1517-1525 (2014).
15. L. A. Rieger, W. J. Randel, A. E. Bourassa, S. Solomon, Stratospheric temperature and ozone anomalies associated with the 2020 Australian new year fires. *Geophys. Res. Lett.*, in press (2021).
16. J. M Rodriguez, M. K. W. Ko, N. D. Sze, Role of heterogeneous conversion of N₂O₅ on sulphate aerosols in global ozone losses. *Nature*. **352**, 134-137 (1991).
17. M. Prather, Catastrophic loss of stratospheric ozone in dense volcanic clouds. *J. Geophys. Res. Atmos.* **97**, 10187-10191 (1992).
18. D. J. Hofmann, S. Solomon, Ozone depletion through heterogeneous chemistry following the eruption of the El Chichon volcano. *J. Geophys. Res.*, **94**, 5029-5041 (1989).
19. Solomon, S. *et al.*, The role of aerosol variations in anthropogenic ozone depletion at northern mid-latitudes. *J. Geophys. Res.*: **101**, 6713-6727 (1996).
20. E. L. Fleming, C. H. Jackman, D. K. Weisenstein, M. K. W. Ko, The impact of interannual variability on multidecadal total ozone simulations. *J. Geophys. Res.* **112**, D10310 doi:10.1029/2006JD007953 (2007).
21. K. A. Stone *et al.*, Observing the Impact of Calbuco Volcanic Aerosols on South Polar Ozone Depletion in 2015. *J. Geophys. Res.: Atm.* **122**, 11,862-11,879 (2017).
22. P. Yu *et al.*, Persistent stratospheric warming due to 2019-2020 Australian wildfire smoke. *Geophys. Res. Lett.* **48**, e2021GL092609 doi:10.1029/2021GL092609 (2021).
23. A. L. Hodshire, *et al.*, Aging Effect on Biomass Burning Aerosol Mass and Composition: A Critical Review of Field and Laboratory Studies. *Environ. Sci. Technol.* **53**, 10007-10022 (2019).

24. J. A. Thornton, P. D. Abbatt, N₂O₅ reaction on submicron sea salt aerosol: kinetics, products, and the effect of surface active organics, *J. Phys. Chem. A*, **109**, 10004–10012, doi:10.1021/jp054183t, (2005).
25. T. H. Bertram, J. A. Thornton, Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride. *Atmos. Chem. Phys.* **9**, 8351–8363, doi:10.5194/acp-9-8351-2009 (2009).
26. E. N. Escorcia, S.J. Sjostedt, J. P. D. Abbatt, Kinetics of N₂O₅ Hydrolysis on Secondary Organic Aerosol and Mixed Ammonium Bisulfate – Secondary Organic Particles. *J. Phys. Chem.* **114**, 13112–13121 (2010).
27. D. W. Fahey *et al.*, *In situ* measurements constraining the role of sulphate aerosols in mid-latitude ozone depletion. *Nature*. **363**, 509–514 (1993).
28. E J Llewellyn *et al.*, The OSIRIS instrument on the Odin spacecraft. *Can. J. Phys.* **82**(6), 411–422 (2004).
29. M. Cisewski *et al.*, The Stratospheric Aerosol and Gas Experiment (SAGE III) on the International Space Station (ISS) Mission. *Sensors, Systems, and Next-Generation Satellites XVIII*, **9241**, 924107. International Society for Optics and Photonics (2014).
30. P. F. Bernath, Atmospheric Chemistry Experiment (ACE): Mission overview. *Geophys. Res. Lett.* **32**, L15S01, doi:10.1029/2005GL022386 (2005).
31. A.E. Bourassa *et al.*, Odin-OSIRIS stratospheric aerosol data product and SAGE III intercomparison. *Atmos. Chem. Phys.* **12**, 605–614 (2012).
32. D.J. Zawada *et al.*, Tomographic retrievals of ozone with the OMPS Limb Profiler: algorithm description and preliminary results. *Atmos. Meas. Tech.*, **11**, 2375–2393, (2018).
33. Hanson, D. H., A. R. Ravishankara, S. Solomon, Heterogeneous reactions in sulfuric acid aerosols: a framework for model calculations. *J. Geophys. Res.*, **99**, 3615–3629 (1994).
34. X. X. Tie, G. Brasseur, The response of stratospheric ozone to volcanic-eruptions: Sensitivity to atmospheric chlorine loading. *Geophys. Res. Lett.*, **22**, 3035–3038. doi:10.1029/95GL03057 (1994).
35. P. O. Wennberg *et al.*, Removal of Stratospheric O₃ by Radicals: In Situ Measurements of OH, HO₂, NO, NO₂, ClO, and BrO. *Science*. **266**, 398–404 (1994).
36. M. Mozurkewich, J. G. Calvert, Reaction probability of N₂O₅ on aqueous aerosols, *J. Geophys. Res.* **93**, 15889, doi:10.1029/JD093iD12p15889 (1988).
37. D. R. Hanson, A. R. Ravishankara, E. R. Lovejoy, Reaction of BrONO₂ with H₂O on submicron sulfuric acid aerosol and the implications for the lower stratosphere. *J. Geophys. Res. Atmos.* **101**, 9063–9069 (1996).
38. M. J. Schwartz *et al.*, Australian New Year's PyroCb Impact on Stratospheric Composition. *Geophys. Res. Lett.* **47**, e2020GL090831 (2020).
39. S. R. Handley, D. Clifford, D. J. Donaldson, Photochemical Loss of Nitric Acid on Organic Films: a Possible Recycling Mechanism for NO_x. *Environ. Sci. Technol.* **41**, 3898–3903 (2007).
40. S. Solomon, R. R. Garcia, F. S. Rowland, D. J. Wuebbles, On the depletion of Antarctic ozone. *Nature*. **321**, 755–758 (1986).
41. Solomon, S. Stratospheric ozone depletion: A review of concepts and history. *Rev. Geophys.*, **37**, 275–316, (1999).
42. R. W. Portmann *et al.*, Role of aerosol variations in anthropogenic ozone depletion in the polar regions. *J. Geophys Res.* **101**, 22,991–23,006 (1996).

43. K. Dube, *et al.*, Accounting for the photochemical variation in stratospheric NO₂ in the SAGE III/ISS solar occultation retrieval. *Atmos. Meas. Tech.* **14**, 557–566 (2021).
44. C.A. McLinden, S. C. Olsen B. Hannegan, O. Wild, M.J. Prather, J. Sundet. Stratospheric ozone in 3-D models: A simple chemistry and the cross-tropopause flux. *J. Geophys. Res.-Atmos.*, **105**, 14653–14665 (2000).
45. P. E. Sheese *et al.*, Validation of ACE-FTS version 3.5 NO_y species profiles using correlative satellite measurements. *Atmos. Meas. Tech.* **9**, 5781–5810 (2016).
46. P. F. Levelt *et al.*, The Ozone Monitoring Instrument: overview of 14 years in space. *Atmos. Chem. Phys.* **18**, 5699-5745 (2018).
47. G. P. Kablick, D. R. Allen, M. D. Fromm, G. E. Nedoluha, Australian pyroCb smoke generates synoptic-scale stratospheric anticyclones. *Geophys. Res. Lett.*, **47**, e2020GL088101 (2005).
48. P. Yu *et al.*, Evaluations of tropospheric aerosol properties simulated by the community earth system model with a sectional aerosol microphysics scheme. *J. Adv. Model. Earth Syst.* **7**, 865-914 (2015).
49. C. A. Mears, F. J. Wentz, Construction of the Remote Sensing Systems V3.2 Atmospheric Temperature Records From the MSU and AMSU Microwave Sounders, *J. Atm. Ocean Tech.*, **26**, 1040-1056 (2009).
50. P. Braesicke, J. Neu *et al.* Chapter 3, Update on Global Ozone, in *Scientific Assessment of Ozone Depletion: 2018, Global Ozone Research and Monitoring Project – Report No. 58*. 588 (2018).

Figure Captions




Figure 1. Monthly mean lower stratospheric aerosol extinction observations at around 750 nm (defined as the ratio to gas phase molecules only), available since 2015 from OMPS (panel A), OSIRIS (panel B) and SAGE III/ISS (panel C). Data represent an average for the lower stratosphere, weighted by the microwave limb sounder temperature weighting function, which is centered around 100 mb. The year 2015 is selected as the start date to capture the Calbuco eruption, which displayed similar SH extinction levels to those obtained after the 2020 Australian fires. Gray regions indicate missing values (see Methods). SAGE III/ISS and OSIRIS data shown here are the average of sunrise and sunset occultations.


Figure 2. Monthly averaged 18.5 km level anomalies (percent) in de-seasonalized available years of OSIRIS NO_x (panel A) and sunrise (B) and sunset (C) SAGE III/ISS data, along with the difference between the smoke and no smoke model runs for 2020 (D). Gray regions in the data indicate missing values (see Methods). The hatched regions on the OSIRIS panel show where the 2020 anomaly is greater than the maximum, or less than the minimum anomaly over all the data from 2002-2019.

Figure 3. NO_x concentrations (molecules/cm³) by month, averaged from 30-60°S at 18.5 km. Top panel presents twenty years of OSIRIS measurements, with 2015 and 2016 (after the Calbuco eruption) and 2020 and 2021 (after the Australian black summer fires) distinguished from others by the indicated colors. Bottom panel shows twenty realizations of 2020 both with (red) and without (gray) smoke for the same latitudes and altitude as calculated in the model.


Figure 4. Monthly mean deseasonalized NO_x versus aerosol extinction at 18.5 km and from 45°S to 40°S for the model (panel A, OSIRIS (B), and SAGE III/ISS (C and D). 2015 and 2016 (after the Calbuco eruption) and 2020 and 2021 (after the Australian black summer fires) are in colors, while the other years make up the gray points. Outliers greater than 4 median absolute deviations from the median were removed from the OSIRIS NO_x and SAGE III/ISS NO_x data. Model points are averaged results for each month from each of the 20 ensemble members.

Figure 5. Calculated weekly averaged changes in total column ozone for the ensemble mean of the smoke minus no smoke runs in the model (top). These model runs did not allow smoke particles to pick up HNO_3 and form PSCs. Observed anomalies in 2020 total column ozone from OMI observations (bottom). The OMI data represent anomalies calculated after the time series has been linearly detrended over the period from 2005-2020 (see Methods). Large negative anomalies during 2020 are indicate by line hatching if they are lower than the 25th percentile.






OSIRIS, NO_x, 18.5 km, 30–60°S



CESM-CARMA, NO_x, 18.5 km, 30–60°S



● 2015 ● 2016 ● 2020 ● 2021

