

# Ligand and Solvent Effects on CO<sub>2</sub> Insertion into Group 10 Metal Alkyl Bonds

Anthony P. Deziel,<sup>a</sup> Matthew R. Espinosa,<sup>a</sup> Ljiljana Pavlovic,<sup>b</sup> David J. Charboneau,<sup>a</sup> Nilay Hazari,<sup>a,\*</sup> Kathrin H. Hopmann<sup>b,\*</sup> & Brandon Q. Mercado<sup>a</sup>

<sup>a</sup>Department of Chemistry, Yale University, P. O. Box 208107, New Haven, Connecticut, 06520, USA. E-mail: [nilay.hazari@yale.edu](mailto:nilay.hazari@yale.edu).

<sup>b</sup>Department of Chemistry, UiT The Arctic University of Norway, N-9307 Tromsø, Norway. E-mail: [kathrin.hopmann@uit.no](mailto:kathrin.hopmann@uit.no).


---

## Abstract

The insertion of carbon dioxide into metal element  $\sigma$ -bonds is an important elementary step in many catalytic reactions for carbon dioxide valorization. Here, the insertion of carbon dioxide into a family of Group 10 alkyl complexes of the type (<sup>R</sup>PBP)M(CH<sub>3</sub>) (<sup>R</sup>PBP = B(NCH<sub>2</sub>PR<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>; R = Cy or <sup>t</sup>Bu; M = Ni or Pd) to generate  $\kappa^1$ -acetate complexes of the form (<sup>R</sup>PBP)M{OC(O)CH<sub>3</sub>} is investigated. This involved the preparation and characterization of a number of new complexes supported by the unusual <sup>R</sup>PBP ligand, which features a central boryl donor that exerts a strong *trans*-influence, and the identification of a new decomposition pathway that results in C–B bond formation. In contrast to other Group 10 methyl complexes supported by pincer ligands, carbon dioxide insertion into (<sup>R</sup>PBP)M(CH<sub>3</sub>) is facile and occurs at room temperature because of the high *trans*-influence of the boryl donor. Given the mild conditions for carbon dioxide insertion, we perform a rare kinetic study on carbon dioxide insertion into a late-transition metal alkyl species using (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>). These studies demonstrate that the Dimroth-Reichardt parameter for a solvent correlates with the rate of carbon dioxide insertion and that Lewis acids do not promote insertion. DFT calculations indicate that insertion into (<sup>t</sup>BuPBP)M(CH<sub>3</sub>) (M = Ni or Pd) proceeds via an S<sub>E</sub>2 mechanism and we compare the reaction pathway for carbon dioxide insertion into Group 10 methyl complexes with insertion into Group 10 hydrides. Overall, this work provides fundamental insight that will be valuable for the development of improved and new catalysts for carbon dioxide utilization.


## Introduction

Carbon dioxide (CO<sub>2</sub>) is an attractive target as a chemical feedstock due to its low cost, non-toxic nature, and abundance.<sup>1</sup> Although there are several chemicals currently industrially prepared from CO<sub>2</sub>, such as urea and carbonates,<sup>2</sup> only a small fraction of our available CO<sub>2</sub> is used as a source of carbon for either commodity<sup>1h</sup> or fine<sup>3</sup> chemical synthesis. The full exploitation of CO<sub>2</sub> as a feedstock is limited by its kinetic and thermodynamic stability, which poses a scientific challenge for developing synthetic routes to make more valuable chemicals from CO<sub>2</sub>. Utilizing homogeneous transition metal catalysts is one attractive approach for solving kinetic problems associated with the conversion of CO<sub>2</sub> into more valuable chemicals and a variety of systems have been developed for the thermal, electrochemical, and photochemical conversion of CO<sub>2</sub>.<sup>1,3</sup> An elementary step which is proposed in many of these catalytic reactions is CO<sub>2</sub> insertion into an M-E σ-bond (for example E = H, OR, NR<sub>2</sub>, or CR<sub>3</sub>) to form a product with a M-O bond (Eq 1).<sup>4</sup> To date, the majority of catalysts feature late transition metals,<sup>1,3</sup> likely due to the relative weakness of the M-O bonds that are formed,<sup>4</sup> which makes cleavage of the M-O bond and subsequent transformation more facile.



The crucial role of CO<sub>2</sub> insertion into late transition metal element σ-bonds in catalysis means that understanding the pathway for insertion is important for rationally designing improved catalytic systems.<sup>4</sup> In particular, understanding the impact of the ancillary ligand, solvent, and additives that are commonly used in catalysis, such as Lewis acids, is valuable for the development of optimized catalysts and reaction conditions. Although the influence of these factors on CO<sub>2</sub> insertion into metal hydrides has been studied,<sup>5</sup> there is limited information on how these variables affect CO<sub>2</sub> insertion into metal alkyl bonds.<sup>6,7,8,9,10,11,12,13</sup> These reactions are particularly important because they can result in the generation of products containing C-C bonds from CO<sub>2</sub>, which has been identified as a high priority research target by the United States National Academies of Science.<sup>2</sup> For example, Group 10 catalysts have been used for the reductive carboxylation of a variety of alkyl halides and pseudo halides.<sup>3h,3i</sup> In these reactions, C-C bonds are proposed to form between CO<sub>2</sub> and the alkyl electrophile via the insertion of CO<sub>2</sub> into a metal alkyl bond.

A major limitation in studying CO<sub>2</sub> insertion into Group 10 metal alkyl bonds is the paucity of systems that are stable and react under mild conditions.<sup>14</sup> In fact, the majority of well-defined



**Figure 1:** **a)** Previous examples of pincer-supported Group 10 methyl complexes that undergo CO<sub>2</sub> insertion to form  $\kappa^1$ -acetate complexes. **b)** <sup>R</sup>PBP supported palladium complexes studied in this work, which undergo CO<sub>2</sub> insertion under milder conditions.

examples of CO<sub>2</sub> insertion reactions into Group 10 metal alkyls require elevated temperatures, long reaction times, and do not give quantitative yields of products, which makes it difficult to perform kinetic studies (Figure 1a). Due to the need for high temperatures, there has been a focus on complexes supported by pincer ligands, which often result in complexes with high thermal stability. For example, Wendt *et al.* demonstrated that treatment of (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>) (<sup>t</sup>BuPCP = 2,6-C<sub>6</sub>H<sub>3</sub>(CH<sub>2</sub>P<sup>t</sup>Bu<sub>2</sub>)<sub>2</sub>) with 4 atm of CO<sub>2</sub> at 80 °C results in complete conversion to the  $\kappa^1$ -acetate complex (<sup>t</sup>BuPCP)Pd{OC(O)CH<sub>3</sub>}, but the reaction takes two days.<sup>15</sup> Similarly, we reported that the reaction of 1 atm of CO<sub>2</sub> with the related compound (<sup>t</sup>BuPCP)Ni(CH<sub>3</sub>) requires heating to 150 °C to form the acetate complex (<sup>t</sup>BuPCP)Ni{OC(O)CH<sub>3</sub>}, but only approximately 75% of the acetate product is formed.<sup>16</sup> Changing the pincer ligand from <sup>t</sup>BuPCP to a system containing an sp<sup>3</sup>-hybridized carbon donor or replacing one of the phosphine arms of the pincer ligand with a hemilabile nitrogen donor only leads to small changes in the rate of reaction.<sup>17</sup> A notable example of a Group 10 methyl complex that does not contain a pincer ligand and inserts CO<sub>2</sub> at room temperature is an unusual anionic palladium dimethyl complex, with a pendant Lewis acid, prepared by Jordan and co-workers.<sup>7</sup> Although this system inserts CO<sub>2</sub> rapidly it requires careful handling as it is unstable at room temperature. Overall, in order to gain detailed information on the factors that influence the rate of CO<sub>2</sub> insertion there is a need for stable alkyl complexes that cleanly react with CO<sub>2</sub> under mild conditions.


In this work, we describe the insertion of CO<sub>2</sub> into palladium and nickel methyl complexes supported by <sup>R</sup>PBP (<sup>R</sup>PBP = B(NCH<sub>2</sub>PR<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub><sup>-</sup>; R = Cy or <sup>t</sup>Bu) pincer ligands, which contain a strongly donating central boryl donor (Figure 1b). As a result of the strong *trans*-influence of the boryl donor, these complexes insert CO<sub>2</sub> at significantly milder conditions (room temperature) than previously reported pincer systems. This enables us to measure the kinetics of CO<sub>2</sub> insertion and study the effect of modulating the pincer ligand and the solvent on CO<sub>2</sub> insertion. We demonstrate that the Dimroth-Reichardt *E*<sub>T</sub>(30) parameter<sup>18</sup> for a solvent correlates with the rate of CO<sub>2</sub> insertion and show that Lewis acids, which are often present in catalysis, do not promote CO<sub>2</sub> insertion. Our experimental studies are complemented by computational investigations to understand the pathway for CO<sub>2</sub> insertion and identify the structure of key transition states and intermediates. Overall, our work is one of the first quantitative investigations of the impact of the ancillary ligand, solvent, and LAs on CO<sub>2</sub> insertion into metal alkyl bonds and will assist in the development of improved and new catalysts for CO<sub>2</sub> utilization.

## Results and Discussion

### *Synthesis of <sup>R</sup>PBP Supported Nickel and Palladium Methyl Complexes and Reactivity with CO<sub>2</sub>*


Previous studies on CO<sub>2</sub> insertion into pincer supported Group 10 metal hydrides demonstrate that the rate of CO<sub>2</sub> insertion increases as the *trans*-influence of the donor opposite to the hydride ligand increases.<sup>19</sup> As a result, nickel and palladium hydride complexes supported by <sup>Cy</sup>PSiP (<sup>Cy</sup>PSiP = Si(Me)(2-PCy<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>)<sub>2</sub>) ligands, which contain a strongly donating silyl ligand *trans* to the hydride insert CO<sub>2</sub> more rapidly than the corresponding species containing <sup>R</sup>PCP ligands. We hypothesized that a similar trend may also occur for nickel and palladium methyl species. Unfortunately, the previously described complexes (<sup>Cy</sup>PSiP)M(CH<sub>3</sub>) (M = Ni or Pd) undergo an intramolecular ligand rearrangement,<sup>20</sup> as does (<sup>t</sup>BuPSiP)Pd(CH<sub>3</sub>) (<sup>t</sup>BuPSiP = Si(Me)(2-P<sup>t</sup>Bu<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>)<sub>2</sub>, see SI) which complicates studies of CO<sub>2</sub> insertion. Therefore, we prepared nickel and palladium methyl complexes supported by <sup>R</sup>PBP ligands (<sup>R</sup>PBP = B(NCH<sub>2</sub>PR<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub><sup>-</sup>; R = Cy or <sup>t</sup>Bu) that feature a boryl donor, which has previously been shown to exert a large *trans*-influence,<sup>21</sup> opposite the methyl ligand.

(<sup>Cy</sup>PBP)Ni(CH<sub>3</sub>) was synthesized via an analogous route to that previously described for (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>).<sup>21h</sup> Initially, NiCl<sub>2</sub>•DME was treated with <sup>Cy</sup>PB<sup>H</sup>P to generate (<sup>Cy</sup>PBP)NiCl in 68% yield (Scheme 1). The subsequent reaction of (<sup>Cy</sup>PBP)NiCl with MeMgCl generates



**Scheme 1:** Synthesis of  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$ .


$(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  in 26% yield. The major cause of the low yield of  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  is the need to perform sequential recrystallizations to remove magnesium salts. Specifically, considerable care is required to remove  $\text{MgCl}_2$  by-products from the Grignard reagent, which cause the methyl species to convert back to the chloride complex.  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  was characterized in the solid-state using X-ray crystallography (Figure 2) and has an analogous distorted square planar geometry to  $(^{\text{tBu}}\text{PBP})\text{Ni}(\text{CH}_3)$ .<sup>21h</sup> The P(1)-Ni(1)-P(2) bond angle is  $158.67(10)^\circ$ , which is similar to that observed in  $(^{\text{tBu}}\text{PBP})\text{Ni}(\text{CH}_3)$  ( $157.58(2)^\circ$ ). The B(1)-Ni(1)-C(1) bond angle is almost linear ( $179.1(4)^\circ$ ), which presumably results in the boryl ligand exerting the maximum possible *trans*-influence on the methyl ligand. The Ni(1)-B(1) bond length in  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  is  $1.927(9)$  Å, which is comparable to other species with a Ni-boryl bond.<sup>21h,22</sup> The nickel methyl bond length is  $2.019(9)$  Å, which is significantly shorter than the nickel methyl bond length in  $(^{\text{tBu}}\text{PBP})\text{Ni}(\text{CH}_3)$ , which is  $2.059(2)$  Å. The longer bond length in  $(^{\text{tBu}}\text{PBP})\text{Ni}(\text{CH}_3)$  is likely due to steric factors, as the larger steric properties of the *tert*-butyl substituents in  $(^{\text{tBu}}\text{PBP})\text{Ni}(\text{CH}_3)$  may cause the nickel



**Figure 2.** Solid-state structure of  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  with thermal ellipsoids at 30% probability. Hydrogen atoms are omitted for clarity. Selected distances (Å) and angles ( $^\circ$ ): Ni(1)-B(1)  $1.927(9)$ , Ni(1)-C(1)  $2.019(9)$ , Ni(1)-P(1)  $2.148(2)$ , Ni(1)-P(2)  $2.138(2)$ , B(1)-Ni(1)-C(1)  $179.1(4)$ , B(1)-Ni(1)-P(1)  $80.1(3)$ , C(1)-Ni(1)-P(1)  $100.6(3)$ , B(1)-Ni(1)-P(2)  $78.7(3)$ , C(1)-Ni(1)-P(2)  $100.6(3)$ , P(1)-Ni(1)-P(2)  $158.67(10)$ .


methyl bond to elongate. The increased steric demand of the <sup>t</sup>BuPBP ligand was confirmed using buried volume calculations (SambVca 2.1<sup>23</sup>) using the crystal structures of (<sup>R</sup>PBP)Ni(CH<sub>3</sub>), which show that the buried volume of (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>) is 92.1% compared to 89.3% for (<sup>Cy</sup>PBP)Ni(CH<sub>3</sub>) (see SI).

Palladium chloride complexes supported by <sup>R</sup>PBP (R = Cy or <sup>t</sup>Bu) ligands were synthesized through the reaction of the protonated <sup>R</sup>PB<sup>H</sup>P ligand with commercially available [(η<sup>3</sup>-allyl)PdCl]<sub>2</sub> in yields of 69% and 63%, respectively. This is different to a previous route utilized to metallate the <sup>t</sup>BuPBP ligand, which used the more expensive Pd(COD)Cl<sub>2</sub> as the palladium source.<sup>21k</sup> Treatment of (<sup>R</sup>PBP)PdCl with MeLi results in the formation of the (<sup>R</sup>PBP)Pd(CH<sub>3</sub>) in yields of 25% for (<sup>Cy</sup>PBP)Pd(CH<sub>3</sub>) and 60% for (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (Scheme 2). In an analogous fashion to (<sup>Cy</sup>PBP)Ni(CH<sub>3</sub>), it is important to remove alkali halide by-products as they can cause the palladium methyl species to be converted back to the palladium chloride precursor and the multiple recrystallizations required for (<sup>Cy</sup>PBP)Pd(CH<sub>3</sub>) lowers the yield. (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) was characterized using X-ray crystallography (Figure 3). The geometry around palladium is distorted square planar. The P(1)-Pd(1)-P(2) bond angle is 153.74(3)°, which is a greater deviation from linearity than that observed in a related complex with a <sup>t</sup>BuPCP ligand. Specifically in (<sup>t</sup>BuPCP)PdCl P-Pd-P is 165.79(2)°. <sup>24</sup> As a consequence, the fourth coordination site in pincer complexes supported by <sup>t</sup>BuPBP is likely more sterically accessible than in species supported by <sup>t</sup>BuPCP ligands. The Pd(1)-B(1) bond length in (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) is 2.027(4) Å, which is significantly longer than the Pd-B bond length in the only other crystallographically characterized palladium complex with a <sup>t</sup>BuPBP ligand, (<sup>t</sup>BuPBP)PdCl (Pd-B is 1.972(4) Å). This is consistent with the methyl ligand exerting a significantly stronger *trans*-influence than the chloride ligand.




**Scheme 2:** Synthesis of (<sup>R</sup>PBP)Pd(CH<sub>3</sub>) (R = Cy or <sup>t</sup>Bu).

We treated complexes of the type (<sup>R</sup>PBP)M(CH<sub>3</sub>) (M = Ni or Pd, R = Cy or <sup>t</sup>Bu) with 1 atm of CO<sub>2</sub> in C<sub>6</sub>D<sub>6</sub> at room temperature (Scheme 3). In the cases of (<sup>R</sup>PBP)Pd(CH<sub>3</sub>) (R = Cy or <sup>t</sup>Bu) NMR spectroscopy indicated clean and essentially quantitative CO<sub>2</sub> insertion to the corresponding




**Figure 3.** Solid-state structure of  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  with thermal ellipsoids at 30% probability. Hydrogen atoms are omitted for clarity. Selected distances (Å) and angles ( $^\circ$ ): Pd(1)-B(1) 2.027(4), Pd(1)-C(1) 2.185(4), Pd(1)-P(1) 2.3030(8), Pd(1)-P(2) 2.3035(8), B(1)-Pd(1)-C(1) 178.09(15), B(1)-Pd(1)-P(1) 77.36(12), C(1)-Pd(1)-P(1) 103.49(10), B(1)-Pd(1)-P(2) 77.33(12), C(1)-Pd(1)-P(2) 102.02(10), P(1)-Pd(1)-P(2) 153.74(3).

acetate complexes at room temperature. The reactions took 1 and 6 hours to reach completion for  $(^{\text{Cy}}\text{PBP})\text{Pd}(\text{CH}_3)$  and  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$ , respectively. These are the most rapid insertions of  $\text{CO}_2$  into pincer-supported Group 10 metal alkyl complexes observed to date. The acetate products were isolated and  $(^t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  was characterized using X-ray crystallography (Figure 4). The solid-state structure of  $(^t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  has the same distorted square planar geometry as  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  and confirms  $\kappa^1$ -binding of the acetate. As expected, the Pd(1)-B(1) bond length (1.975(2) Å) in  $(^t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  is significantly shorter than the Pd-B bond length in  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  (2.027(4) Å) due to the weaker *trans*-influence of the acetate ligand. Further, the increased Pd-O bond length in  $(^t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  (2.1861(14) Å) in comparison to  $(^t\text{BuPCP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  (2.1279(19) Å)<sup>15</sup> is consistent with the increased *trans*-influence of the  $^t\text{BuPBP}$  ligand in comparison to the related  $^t\text{BuPCP}$  ligand. In contrast, the insertion of  $\text{CO}_2$  into  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  was not clean. Although NMR spectroscopy suggested that




**Scheme 3:** Generation of  $(^{\text{R}}\text{PBP})\text{M}\{\text{OC}(\text{O})\text{CH}_3\}$  ( $\text{M} = \text{Ni}, \text{R} = \text{Cy}; \text{M} = \text{Pd}, \text{R} = \text{Cy}$  or  $^t\text{Bu}$ ) through  $\text{CO}_2$  insertion into  $(^{\text{R}}\text{PBP})\text{M}(\text{CH}_3)$ .



**Figure 4.** Solid-state structure of  $(^t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  with thermal ellipsoids at 30% probability. Hydrogen atoms are omitted for clarity. Selected distances (Å) and angles (°): Pd(1)-B(1) 1.975(2), Pd(1)-O(1) 2.1861(14), Pd(1)-P(1) 2.3256(5), Pd(1)-P(2) 2.3336(5), B(1)-Pd(1)-O(1) 174.99(7), B(1)-Pd(1)-P(1) 77.32(7), O(1)-Pd(1)-P(1) 100.17(4), B(1)-Pd(1)-P(2) 79.00(7), O(1)-Pd(1)-P(2) 103.05(4), P(1)-Pd(1)-P(2) 155.805(18).


$(^{\text{Cy}}\text{PBP})\text{Ni}\{\text{OC}(\text{O})\text{CH}_3\}$ , which was not isolated, was the major product, free ligand and several unidentified decomposition products were present (see SI). Additionally, it took approximately 4 days at room temperature for all of the  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$  to disappear. Overall, from these initial qualitative examples of  $\text{CO}_2$  insertion into  $(^{\text{Cy}}\text{PBP})\text{M}(\text{CH}_3)$  ( $\text{M} = \text{Ni}$  or  $\text{Pd}$ ) and  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  we can make the following conclusions: (i) The insertion of  $\text{CO}_2$  into  $(^{\text{Cy}}\text{PBP})\text{Pd}(\text{CH}_3)$  is faster than the corresponding insertion into  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$ . This is in agreement with results for  $\text{CO}_2$  insertion into  $(^t\text{BuPCP})\text{M}(\text{CH}_3)$  ( $\text{M} = \text{Ni}$  or  $\text{Pd}$ )<sup>15-16</sup> but stands in contrast to computational studies of  $\text{CO}_2$  insertion into  $(^t\text{BuPCP})\text{MH}$  ( $\text{M} = \text{Ni}$  or  $\text{Pd}$ ), where insertion is proposed to be faster for nickel.<sup>19</sup> (ii) The insertion of  $\text{CO}_2$  into  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  is significantly faster than the corresponding insertion into  $(^t\text{BuPCP})\text{Pd}(\text{CH}_3)$ .<sup>15</sup> This suggests that the rate of  $\text{CO}_2$  insertion can be increased by having a stronger *trans*-influence donor opposite the methyl, similar to what has been observed for  $\text{CO}_2$  insertion into Group 10 metal hydrides.<sup>19</sup> (iii) The faster insertion of  $\text{CO}_2$  into  $(^{\text{Cy}}\text{PBP})\text{Pd}(\text{CH}_3)$  compared to  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  suggests that steric factors are important.

In contrast to the quantitative or relatively clean insertion of  $\text{CO}_2$  observed for  $(^{\text{Cy}}\text{PBP})\text{Pd}(\text{CH}_3)$ ,  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$ , and  $(^{\text{Cy}}\text{PBP})\text{Ni}(\text{CH}_3)$ , insertion into  $(^t\text{BuPBP})\text{Ni}(\text{CH}_3)$  was not clean and was significantly slower than for the other methyl compounds. After 10 days at room temperature, approximately 30% of  $(^t\text{BuPBP})\text{Ni}(\text{CH}_3)$  was still observed, along with 50% of a product we propose to be  $(^t\text{BuPBP})\text{Ni}\{\text{OC}(\text{O})\text{CH}_3\}$  on the basis of its similar  $^{31}\text{P}$  NMR chemical shift to  $(^{\text{Cy}}\text{PBP})\text{Ni}\{\text{OC}(\text{O})\text{CH}_3\}$ , and 20% of unidentified decomposition products (Scheme 4). Attempts



**Scheme 4:** CO<sub>2</sub> insertion into (tBuPBP)Ni(CH<sub>3</sub>).

to independently synthesize (tBuPBP)Ni{OC(O)CH<sub>3</sub>} via reactions between silver acetate and (tBuPBP)NiCl or sodium acetate and (tBuPBP)Ni(OTf) were unsuccessful. Although, initially a new resonance is observed in the <sup>31</sup>P NMR spectra of these reactions with the same chemical shift as the proposed insertion product with CO<sub>2</sub> (84.9 ppm), this resonance disappears over time, indicating that (tBuPBP)Ni{OC(O)CH<sub>3</sub>} is likely unstable in solution. This decomposition prohibited isolation and full characterization of (tBuPBP)Ni{OC(O)CH<sub>3</sub>}. Nevertheless, since the rate of CO<sub>2</sub> insertion into (tBuPBP)Ni(CH<sub>3</sub>) is presumably independent of the decomposition of the corresponding insertion product, we can make qualitative conclusions about the rate of insertion. We hypothesize that the slower rate of CO<sub>2</sub> insertion is related to the increased steric bulk of (tBuPBP)Ni(CH<sub>3</sub>), which makes it harder for CO<sub>2</sub> to interact with the complex. When (tBuPBP)Ni(CH<sub>3</sub>) was treated with CO<sub>2</sub> at higher temperatures even larger amounts of decomposition products were observed. Control experiments indicate that (tBuPBP)Ni(CH<sub>3</sub>) is not stable even when heated under an N<sub>2</sub> atmosphere. Although we were unable to identify all of the decomposition products of (tBuPBP)Ni(CH<sub>3</sub>), an unusual dinuclear species (tBuPB<sup>Me</sup>P)<sub>2</sub>Ni<sub>2</sub>(μ-N<sub>2</sub>) (tBuPB<sup>Me</sup>P = MeB(NCH<sub>2</sub>P<sup>t</sup>Bu<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>), containing two nickel(0) centers and a bridging dinitrogen ligand crystallized out of solution (Figure 5). In this complex, the nickel methyl ligand has undergone reductive coupling with the boryl ligand to form a new C-B bond. The boron atom no longer coordinates to the nickel center and the two phosphorus donors of the new bidentate tBuPB<sup>Me</sup>P ligands do not coordinate to the same nickel center but instead coordinate to two different nickel atoms. The geometry around the nickel centers is distorted trigonal planar and the dinitrogen ligand coordinates in an end-on fashion. The N-N bond distance of 1.145(8) Å is slightly elongated compared to that of free N<sub>2</sub> (1.098 Å), consistent with weak activation of N<sub>2</sub> and what has been observed in related nickel dinitrogen complexes.<sup>25</sup> The decomposition of (tBuPBP)Ni(CH<sub>3</sub>) into (tBuPB<sup>Me</sup>P)<sub>2</sub>Ni<sub>2</sub>(μ-N<sub>2</sub>) demonstrates a new reductive pathway for decomposition of complexes supported by <sup>R</sup>PBP ligands. However, this is likely a particular problem for the tBuPBP ligand when




**Figure 5. a)** Decomposition of  $(t\text{BuPBP})\text{Ni}(\text{CH}_3)$  under  $\text{N}_2$  at  $60\text{ }^\circ\text{C}$ . **b)** Solid-state structure of  $(t\text{BuPB}^{\text{Me}}\text{P})_2\text{Ni}_2(\mu\text{-N}_2)$  with thermal ellipsoids at 30% probability. The model is near the crystallographic inversion center; half of the structure is generated by symmetry. Hydrogen atoms are omitted for clarity. Selected distances ( $\text{\AA}$ ) and angles ( $^\circ$ ):  $\text{Ni}(1)\text{-N}(1)$  1.854(4),  $\text{Ni}(1)\text{-P}(1)$  2.293(11),  $\text{Ni}(1)\text{-P}(2)$  2.164(11),  $\text{N}(1)\text{-N}(2)$  1.145(8),  $\text{P}(1)\text{-Ni}(1)\text{-P}(2)$  122.0(5),  $\text{N}(1)\text{-Ni}(1)\text{-P}(1)$  118.9,  $\text{N}(1)\text{-Ni}(1)\text{-P}(2)$  110.8(3),  $\text{Ni}(1)\text{-N}(1)\text{-N}(2)$  178.7(5).

it is coordinated to Ni because of steric constraints associated with the bulky *tert*-butyl substituents and small size of nickel.


#### *Kinetic Studies of $\text{CO}_2$ Insertion into $(t\text{BuPBP})\text{Pd}(\text{CH}_3)$*

It has traditionally been challenging to perform kinetics experiments on  $\text{CO}_2$  insertion into Group 10 alkyl complexes either because of the instability of the complexes<sup>14</sup> or the harsh conditions required for insertion.<sup>16,17</sup> In contrast, the insertion of  $\text{CO}_2$  into  $(t\text{BuPBP})\text{Pd}(\text{CH}_3)$  under mild conditions enables kinetics experiments to be performed. Initially,  $^1\text{H}$  or  $^{31}\text{P}$  NMR spectroscopy was used to monitor the reaction of  $(t\text{BuPBP})\text{Pd}(\text{CH}_3)$  with  $\text{CO}_2$  as a function of time (Figure 6a). It was possible to measure the concentrations of both the  $(t\text{BuPBP})\text{Pd}(\text{CH}_3)$  starting material and the  $(t\text{BuPBP})\text{Pd}\{\text{OC}(\text{O})\text{CH}_3\}$  product, which enabled us to confirm the reaction is clean and



**Figure 6.** Representative traces for the insertion of CO<sub>2</sub> into (tBuPBP)Pd(CH<sub>3</sub>) at 40 °C in C<sub>6</sub>D<sub>6</sub> with 1 atm of CO<sub>2</sub> showing **a)** the concentrations of (tBuPBP)Pd(CH<sub>3</sub>) and (tBuPBP)Pd{OC(O)CH<sub>3</sub>} as a function of time and **b)** the ln of the concentration of (tBuPBP)Pd(CH<sub>3</sub>) as a function of time.

quantitative. An excess of CO<sub>2</sub> was used to ensure that the reactions were performed under pseudo-first order conditions with the concentration of CO<sub>2</sub> measured using quantitative <sup>13</sup>C NMR spectroscopy (see SI).<sup>26</sup> Initial experiments in C<sub>6</sub>D<sub>6</sub> indicate that the reaction is first order in both [(tBuPBP)Pd(CH<sub>3</sub>)] and [CO<sub>2</sub>], demonstrating that the overall rate law is  $k_1[(tBuPBP)Pd(CH_3)][CO_2]$  (Figures 6b and 7). The  $k_{obs}$  values obtained from a plot of  $\ln([(tBuPBP)Pd(CH_3)])$  versus time were divided by the concentration of CO<sub>2</sub> to obtain values of  $k_1$  (see SI). Monitoring the rate of the reaction in C<sub>6</sub>D<sub>6</sub> at various temperatures allowed for the activation parameters to be determined through Eyring analysis (see SI). The enthalpy of activation,  $\Delta H^\ddagger$ ,  $10.9 \pm 1.1$  kcal/mol, for insertion into (tBuPBP)Pd(CH<sub>3</sub>) is significantly lower than that for (tBuPCP)Pd(CH<sub>3</sub>) ( $\Delta H^\ddagger = 17.4 \pm 1.7$  kcal/mol),<sup>26</sup> which is the most closely related system



**Figure 7.** Reaction order in [CO<sub>2</sub>] of the insertion of CO<sub>2</sub> into (tBuPBP)Pd(CH<sub>3</sub>) in C<sub>6</sub>D<sub>6</sub> at 40 °C.


for which an experimental activation energy for CO<sub>2</sub> insertion has been measured. This is consistent with the hypothesis that the strong *trans*-influence of the <sup>t</sup>BuPBP ligand weakens the palladium methyl bond and destabilizes the ground state. The entropies of activation, ΔS<sup>‡</sup>, for (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) and (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>) are within error of each other at -32.2 ± 3.2 and -28.2 ± 4.54 cal/molK, respectively. The similar negative values suggest that the nature of the rate-limiting transition state in both reactions are analogous and is in agreement with two molecules combining to form one molecule in the transition state (*vide infra*). The overall difference in ΔG<sup>‡</sup> at 298 K between (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (20.5 ± 2.1 kcal/mol) and (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (25.8 ± 2.6 kcal/mol) is consistent with the much faster rate of insertion observed for the <sup>t</sup>BuPBP system.

In CO<sub>2</sub> insertion reactions into metal hydrides, it has been demonstrated that the solvent can have a major influence on the rate of insertion.<sup>5</sup> To date there are no studies exploring how the rate of CO<sub>2</sub> insertion into metal alkyls varies as the solvent is changed. We determined the rate of CO<sub>2</sub> insertion (k<sub>1</sub>) into (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) in toluene-*d*<sub>8</sub>, benzene-*d*<sub>6</sub>, 1,4-dioxane, THF, and glyme (Table 1). There was only a small solvent effect with approximately a 1.9-fold difference in k<sub>1</sub> between the slowest (toluene-*d*<sub>8</sub>) and fastest (glyme) solvent (Table 1). This compares with a 30-fold increase in the rate of CO<sub>2</sub> insertion into an iridium hydride in a similar set of solvents.<sup>5a</sup> We propose that the smaller impact of solvent in this case is because there is less build-up of charge in the rate-determining transition state for CO<sub>2</sub> insertion (*vide infra*) compared to the corresponding transition state for metal hydrides. To determine if the variation in the rate of insertion correlated with any solvent parameters, ln(k<sub>1</sub>) was plotted against the dielectric constant, Gutmann Acceptor Number (AN),<sup>27</sup> or Dimroth-Reichardt E<sub>T</sub>(30) parameter<sup>18</sup> of the solvent. No significant correlation was observed between the rate of CO<sub>2</sub> insertion and the dielectric constant

Entry	Solvent	k <sub>1</sub> (M <sup>-1</sup> s <sup>-1</sup> × 10 <sup>-2</sup> ) <sup>a</sup>	Rate Enhancement
1	Toluene- <i>d</i> <sub>8</sub>	1.1	1
2	Benzene- <i>d</i> <sub>6</sub>	1.3	1.2
3	1,4-Dioxane	1.5	1.4
4	Tetrahydrofuran	1.8	1.6
5	Glyme	2.1	1.9

**Table 1.** Effect of Solvent on rate of CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) at 40 °C and 1 atm of CO<sub>2</sub>.

<sup>a</sup>These values are the average of two trials and the errors are ±10%.



**Figure 8.** Plot of  $\ln(k_1)$  for the insertion of  $\text{CO}_2$  into  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  versus the Dimroth-Reichardt  $E_T(30)$  solvent parameter for various solvents at  $40^\circ\text{C}$  and 1 atm of  $\text{CO}_2$ .

or AN of a solvent (see SI). In contrast, an excellent correlation was observed between the  $E_T(30)$  value of a solvent and the rate of  $\text{CO}_2$  insertion (Figure 8), suggesting that  $E_T(30)$  is a good predictor of the rate of  $\text{CO}_2$  insertion. A similar relationship was observed for  $\text{CO}_2$  insertion into metal hydrides and hydroxides,<sup>5b</sup> raising the possibility that this correlation is general across a variety of distinct metal-element  $\sigma$ -bonds. It is likely that  $E_T(30)$  is a better predictor of  $\text{CO}_2$  insertion rates because it explicitly considers specific solute/solvent interactions, which are not accounted for when the solvent is modelled using either dielectric constant or AN.<sup>5b</sup> Unfortunately, we were unable to extend the correlation between the rate of  $\text{CO}_2$  insertion into  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  and  $E_T(30)$  to more polar solvents. This is because these solvents led to the decomposition of  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  or resulted in rates that were too fast to measure using NMR spectroscopy. For example,  $\text{CO}_2$  insertion into  $(^t\text{BuPBP})\text{Pd}(\text{CH}_3)$  is complete in less than 10 minutes at room temperature in pyridine and benzonitrile, which is too fast to obtain reliable kinetics by NMR spectroscopy. Nevertheless, given the slow rates of  $\text{CO}_2$  insertion previously observed into pincer-supported Group 10 alkyls, the completion of reactions in less than 10 minutes at room temperature is unprecedented.

Lewis acids have been proposed to increase the rate of  $\text{CO}_2$  insertion into a variety of different metal-element  $\sigma$ -bonds, but these effects are often dependent on the pathway of insertion and the reaction conditions.<sup>5,7-8,9b,28</sup> As a result, it is difficult to predict when a Lewis acid will increase the rate of a  $\text{CO}_2$  insertion reaction. We evaluated the impact of a variety of different Lewis acids on the rate of  $\text{CO}_2$  insertion into  $(^t\text{BuPCP})\text{Pd}(\text{CH}_3)$  using THF as the solvent (Table 2).<sup>29</sup> Although


Entry	Additive (20 equiv.)	$k_{\text{obs}}$ ( $\text{s}^{-1} \times 10^{-3}$ ) <sup>b</sup>	Rate Enhancement
1	None	1.6	1
2	(NBu <sub>4</sub> )PF <sub>6</sub>	2.3	1.4
3	LiNTf <sub>2</sub> <sup>a</sup>	2.2	1.4
4	NaNTf <sub>2</sub> <sup>a</sup>	2.1	1.3
5	KNTf <sub>2</sub> <sup>a</sup>	2.3	1.4
6	LiPF <sub>6</sub>	2.4	1.5
7	LiBPh <sub>4</sub> •3DME	2.5	1.6
8	LiOTf	2.1	1.3

**Table 2.** Effect of Lewis Acids on rate of CO<sub>2</sub> insertion into (tBuPBP)Pd(CH<sub>3</sub>) in THF at 40 °C and 1 atm of CO<sub>2</sub>. <sup>a</sup>NTf = triflimide. <sup>b</sup>These values are the average of two trials and the errors are  $\pm 10\%$ .

there is an increase in rate when the reaction was performed in the presence of 20 equivalents of a variety of Lewis acids, a control experiment using 20 equivalents of (nBu<sub>4</sub>)PF<sub>6</sub>, which does not contain a Lewis acid, gives the same enhancement. This indicates that there is no increase in rate caused by Lewis acids and instead it is likely the case that increasing the ionic strength of the solution results in a slight enhancement in the rate of CO<sub>2</sub> insertion. Consistent with this hypothesis, all of the different Lewis acids evaluated give approximately the same increase in the rate of CO<sub>2</sub> insertion. At this stage, it is unclear why no Lewis acid effect is observed in the insertion of CO<sub>2</sub> into (tBuPBP)Pd(CH<sub>3</sub>), given that Li<sup>+</sup> has been demonstrated to increase the rate of CO<sub>2</sub> insertion into (CO)<sub>5</sub>W(CH<sub>3</sub>).<sup>8</sup> However, in the next section, DFT calculations are performed to understand the exact nature of the transition states involved in CO<sub>2</sub> insertion into (tBuPBP)Pd(CH<sub>3</sub>).

### Computational Studies of CO<sub>2</sub> Insertion into (<sup>R</sup>PBP)Pd(CH<sub>3</sub>) and Related Species

Previous studies investigating CO<sub>2</sub> insertion into PCP supported palladium and nickel methyl complexes have proposed two plausible pathways for insertion: (i) a 1,2-insertion (*inner sphere*) or (ii) an S<sub>E</sub>2 pathway (*outer sphere*).<sup>16,17c,26</sup> In the 1,2-insertion pathway, the O-bound acetate product is formed in one step and CO<sub>2</sub> interacts with the metal in the key transition state (Figure 9a). This means that both the C–C and M–O bonds are formed at the same time. In contrast, the S<sub>E</sub>2 pathway involves two steps and starts with electrophilic attack of CO<sub>2</sub> on the coordinated methyl, without any direct interaction between CO<sub>2</sub> and the metal (Figure 9b). This results in the formation of the new C–C bond. In the second step, the intermediate containing an acetate-metal



**Figure 9.** Two plausible mechanisms for CO<sub>2</sub> insertion into pincer-supported Group 10 methyls: **a)** 1,2-insertion and **b)** S<sub>E</sub>2.


ion-pair rearranges via TS2 to form the *O*-bound acetate adduct. Here, we used DFT methods to understand the pathway for CO<sub>2</sub> insertion into (t<sup>Bu</sup>PBP)M(CH<sub>3</sub>) (M = Ni or Pd). We also independently calculated CO<sub>2</sub> insertion into (t<sup>Bu</sup>PCP)Pd(CH<sub>3</sub>) in order to directly compare it with the t<sup>Bu</sup>PBP systems.

Our calculations indicate that CO<sub>2</sub> insertion into (t<sup>Bu</sup>PBP)Pd(CH<sub>3</sub>) proceeds via an S<sub>E</sub>2 pathway (Figure 10 & Table 3, Figures S43-S44). The rate-limiting step is the initial attack of CO<sub>2</sub> on the palladium methyl group (TS1\_Out), which has a computed barrier of 19.0 kcal/mol (298 K). This is within error of the activation barrier measured experimentally (*vide supra*). Although there is no interaction between CO<sub>2</sub> and the metal in TS1\_Out, we note that there are two C-H⋯OCO interactions between the *tert*-butyl substituents of the t<sup>Bu</sup>PBP ligand and CO<sub>2</sub> (2.43 and 2.74 Å,

Complex	TS1_Out (kcal/mol)	TS1_Inn (kcal/mol)
(t <sup>Bu</sup> PBP)Pd(CH <sub>3</sub> )	<b>19.0</b> (23.9)	25.1 (30.6)
(MePBP)Pd(CH <sub>3</sub> )	22.5 (23.4)	<b>19.4</b> (21.0)
(t <sup>Bu</sup> PBP)Ni(CH <sub>3</sub> )	<b>24.5</b> (27.8)	31.9 (33.7)
(t <sup>Bu</sup> PCP)Pd(CH <sub>3</sub> )	<b>24.2</b> (29.2)	36.7 (41.1)
(MePCP)Pd(CH <sub>3</sub> )	26.8 (30.0)	<b>25.0</b> (27.8)

**Table 3.** Calculated free energy barriers for inner and outer sphere C-CO<sub>2</sub> bond formation with different pincer-supported palladium and nickel complexes (kcal/mol, B3LYP-D3/def2-TZVPPD[CPCM]/B3LYP-D3/def2-SVP[CPCM], 298 K). For comparison, results for the ωB97XD functional are given in parenthesis, showing the same mechanistic trends.

respectively), which provide charge stabilization. This may explain why the addition of Lewis acids to the system does not result in a rate enhancement (*vide supra*). The second step of the  $S_E2$  pathway has a very small barrier and involves rotation of the acetate-palladium ion pair to form the *O*-bound acetate product (Figure 10). In contrast, the barrier for the alternative 1,2-insertion is significantly higher at 25.1 kcal/mol (**TS1\_Inn**, Figure 10, Figure S45). The high barrier is likely



**Figure 10.** Top: Computed free energy profile (B3LYP-D3/def2-TZVPPD[CPCM]/B3LYP-D3/def2-SVP[CPCM], 298K) for CO<sub>2</sub> insertion into (tBuPBP)Pd(CH<sub>3</sub>), Bottom: Optimized transition state structures for CO<sub>2</sub> insertion into (R<sup>t</sup>PBP)Pd(CH<sub>3</sub>), with R = tBu or Me. All energies are in kcal/mol.

caused by the bulky *tert*-butyl substituents of the <sup>t</sup>BuPBP ligand, which make it difficult for CO<sub>2</sub> to approach palladium without steric repulsion. In order to investigate this hypothesis, we calculated CO<sub>2</sub> insertion into (<sup>Me</sup>PBP)Pd(CH<sub>3</sub>), where the *tert*-butyl substituents on the <sup>t</sup>BuPBP ligand have been replaced with methyl substituents. For CO<sub>2</sub> insertion into (<sup>Me</sup>PBP)Pd(CH<sub>3</sub>) the 1,2-insertion pathway is preferred by 3.1 kcal/mol over the S<sub>E2</sub> mechanism (Figures 10 & S46-S47). Thus, the size of the pincer ligand has a decisive role in promoting either a 1,2-insertion (small phosphine substituents) or an S<sub>E2</sub> pathway (large phosphine substituents). Similar ligand effects were previously observed for NHC-Cu complexes.<sup>30</sup>

The <sup>t</sup>BuPCP supported complex, (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>), also shows a preference for the S<sub>E2</sub> pathway, with a computed barrier for insertion of 24.2 kcal/mol (**TS\_Out**, Figure S48-S49). Consistent with previous results,<sup>26</sup> the 1,2-insertion mechanism is unlikely for (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>) due to the high barrier of 36.7 kcal/mol (**TS\_Inn**, Figure S50).<sup>31</sup> Thus, both (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) and (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>) show a clear preference for an S<sub>E2</sub> mechanism. In the rate limiting transition state (**TS\_Out**), (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) displays a longer C–CO<sub>2</sub> bond (0.09 Å) and a smaller CO<sub>2</sub> angle (1.7°) compared to (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>), which indicates an earlier TS. The lower barrier for (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (by 5.2 kcal/mol) is in agreement with our kinetic studies, showing that CO<sub>2</sub> insertion is much faster for (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>). The decrease in the barrier for CO<sub>2</sub> insertion is also matched by the greater thermodynamic driving force for CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) compared with (<sup>t</sup>BuPCP)Pd(CH<sub>3</sub>) (-23.3 kcal/mol versus -15.8 kcal/mol). These results highlight how a donor atom with strong *trans*-influence opposite to the methyl can lower the barrier for insertion and suggest that the enhancement is due to destabilization of the ground state as opposed to a kinetic effect associated with stabilization of the transition state.

The slower qualitative rate of CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>) compared to (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (*vide supra*) is surprising given that CO<sub>2</sub> insertion into nickel hydrides is typically faster than palladium hydrides.<sup>19</sup> Calculations on CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>) confirm our experimental results as the calculated barrier is 24.5 kcal/mol at 298K, more than 5 kcal/mol higher than for (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) (B3LYP-D3, Table 3). The lowest energy pathway for CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>) also follows an S<sub>E2</sub> mechanism (Figure S51-52). Geometrical analysis of the rate-determining transition states for CO<sub>2</sub> insertion into (<sup>t</sup>BuPBP)Ni(CH<sub>3</sub>) and (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>) shows that the Ni–P bonds are shorter by around 0.15 Å compared with the corresponding Pd–P

bonds due to the smaller atomic radius of nickel. We propose that the increased steric congestion around nickel results in the higher barrier for CO<sub>2</sub> insertion relative to the palladium system.<sup>32</sup> This is in agreement with previous results showing that CO<sub>2</sub> insertion reactions into nickel hydrides are highly sensitive to steric factors.<sup>5a</sup> It is likely, however, that there is no inherent electronic reason for CO<sub>2</sub> insertion into nickel methyl complexes to be slower than palladium methyl complexes, and if a nickel complex could be prepared where steric factors were not relevant,<sup>33</sup> we expect to see comparable rates of insertion.

## Conclusions

In this work, we have demonstrated that nickel and palladium methyl complexes supported by <sup>R</sup>PBP ligands undergo CO<sub>2</sub> insertion under significantly milder conditions than any previously reported pincer-ligated Group 10 alkyl species. This is because of the strong *trans*-influence of the central boryl donor in <sup>R</sup>PBP type ligands, which destabilizes the M–CH<sub>3</sub> bond. Qualitatively, we show that CO<sub>2</sub> insertion into (<sup>R</sup>PBP)Pd(CH<sub>3</sub>) is faster than CO<sub>2</sub> insertion into (<sup>R</sup>PBP)Ni(CH<sub>3</sub>), which is likely due to steric factors. In contrast, we propose that steric factors are less significant in CO<sub>2</sub> insertion into Group 10 hydrides, and as a result nickel hydrides typically undergo faster insertion than palladium hydrides.<sup>19</sup> Given the rapid rate of insertion into <sup>R</sup>PBP-ligated complexes, we were able to complete a rare kinetic study on CO<sub>2</sub> insertion into a metal alkyl complex. Our results demonstrate that the solvent parameter that is best for predicting the rate of CO<sub>2</sub> insertion is the Dimroth-Reichardt *E*<sub>T</sub>(30) parameter. This is analogous to what we have observed for transition metal hydrides,<sup>5b</sup> but the dependence of the insertion rate on the solvent is significantly less pronounced in the case of the metal alkyl complex. We hypothesize that this is because there is less charge build-up in the transition state for carbon dioxide insertion into a metal alkyl complex compared to related metal hydrides. Consistent with there being only a small build-up of charge in the transition state, we did not observe any enhancement from Lewis acids in CO<sub>2</sub> insertion reactions into (<sup>t</sup>BuPBP)Pd(CH<sub>3</sub>). DFT calculations indicate that the proposed mechanism for insertion into (<sup>t</sup>BuPBP)M(CH<sub>3</sub>) (M = Ni or Pd) follows an S<sub>E</sub>2 pathway, but this is dependent on steric factors and compounds with less bulky substituents on the phosphine donors are more likely to undergo insertion via a 1,2-insertion mechanism. Overall, our results indicate that the fastest systems for CO<sub>2</sub> insertion into a Group 10 methyl will have a strong *trans*-influence donor opposite the methyl, be able to undergo insertion in solvents with high *E*<sub>T</sub>(30) values, and not have sterically bulky ancillary ligands that prevent the approach of CO<sub>2</sub>. Future research in our group will aim to

extend these principles to other metal-element  $\sigma$ -bonds and apply them to improve catalytic reactions.

## Acknowledgements

NH acknowledges support from National Science Foundation through Grant CHE-1953708. We thank Dr. Hee Won Suh for preliminary work on this project and Dr. Fabian Menges for assistance obtaining high resolution mass spectrometry data with equipment funded by NSF MRI Grant CHE-1828190. KHH and LjP thank the Research Council of Norway (300769), the Tromsø Research Foundation (Grant No. TFS2016KHH), NordForsk (Grant No. 85378) and Notur - The Norwegian Metacenter for Computational Science (No. nn9330k).

## Additional information

Additional information about selected experiments, NMR spectra, and other details are available via the Internet.

## Competing Financial Interests

The authors declare no competing financial interests.

## References

- (a) Quadrelli, E. A.; Centi, G.; Duplan, J.-L.; Perathoner, S. Carbon Dioxide Recycling: Emerging Large-Scale Technologies with Industrial Potential. *ChemSusChem* **2011**, *4*, 1194-1215; (b) Schneider, J.; Jia, H.; Muckerman, J. T.; Fujita, E. Thermodynamics and Kinetics of CO<sub>2</sub>, CO, and H<sup>+</sup> Binding to the Metal Centre of CO<sub>2</sub> Reduction Catalysts. *Chem. Soc. Rev.* **2012**, *41*, 2036-2051; (c) Appel, A. M.; Bercaw, J. E.; Bocarsly, A. B.; Dobbek, H.; DuBois, D. L.; Dupuis, M.; Ferry, J. G.; Fujita, E.; Hille, R.; Kenis, P. J. A.; Kerfeld, C. A.; Morris, R. H.; Peden, C. H. F.; Portis, A. R.; Ragsdale, S. W.; Rauchfuss, T. B.; Reek, J. N. H.; Seefeldt, L. C.; Thauer, R. K.; Waldrop, G. L. Frontiers, Opportunities, and Challenges in Biochemical and Chemical Catalysis of CO<sub>2</sub> Fixation. *Chem. Rev.* **2013**, *113*, 6621-6658; (d) Wang, W.-H.; Himeda, Y.; Muckerman, J. T.; Manbeck, G. F.; Fujita, E. CO<sub>2</sub> Hydrogenation to Formate and Methanol as an Alternative to Photo- and Electrochemical CO<sub>2</sub> Reduction. *Chem. Rev.* **2015**, *115*, 12936-12973; (e) Limbach, M. 'Chapter Four - Acrylates from Alkenes and CO<sub>2</sub>, the Stuff That Dreams Are Made of.' In *Advances in Organometallic Chemistry*. Ed. Pérez, P. J. Academic Press, 2015. pp 175-202; (f) Bernskoetter, W. H.; Hazari, N. Reversible Hydrogenation of Carbon Dioxide to Formic Acid and Methanol: Lewis Acid Enhancement of Base Metal Catalysts. *Acc. Chem. Res.* **2017**, *50*, 1049-1058; (g) Artz, J.; Müller, T. E.; Thenert, K.; Kleinekorte, J.; Meys, R.; Sternberg, A.; Bardow, A.; Leitner, W. Sustainable Conversion of Carbon Dioxide: An Integrated Review of Catalysis and Life Cycle Assessment. *Chem. Rev.* **2018**, *118*, 434-504; (h) Burkart, M. D.; Hazari, N.; Tway, C. L.; Zeitler, E. L. Opportunities and Challenges for Catalysis in Carbon Dioxide Utilization. *ACS Catal.* **2019**, *9*, 7937-7956; (i) Hepburn, C.; Adlen, E.; Beddington, J.; Carter, E. A.; Fuss, S.; Mac Dowell, N.; Minx, J. C.; Smith, P.; Williams, C. K. The Technological and Economic Prospects for CO<sub>2</sub> Utilization and Removal. *Nature* **2019**, *575*, 87-97; (j) Zhang, Z.; Pan, S.-Y.; Li, H.; Cai, J.; Olabi, A. G.; Anthony, E. J.; Manovic, V. Recent Advances in Carbon Dioxide Utilization. *Renew. Sustain. Energy Rev.* **2020**, *125*, 109799; (k) Gao, W.; Liang, S.; Wang, R.; Jiang, Q.; Zhang, Y.; Zheng, Q.; Xie, B.; Toe, C. Y.; Zhu, X.; Wang, J.; Huang, L.; Gao, Y.; Wang, Z.; Jo, C.; Wang, Q.; Wang, L.; Liu, Y.; Louis, B.; Scott, J.; Roger, A.-C.; Amal, R.; He, H.; Park, S.-E. Industrial Carbon Dioxide Capture and Utilization: State of the Art and Future Challenges. *Chem. Soc. Rev.* **2020**, *49*, 8584-8686.
2. National Academies of Sciences, Engineering, and Medicine. *Gaseous Carbon Waste Streams Utilization: Status and Research Needs*. The National Academies Press, Washington DC, 2019.

3. (a) Huang, K.; Sun, C.-L.; Shi, Z.-J. Transition-Metal Catalyzed C-C Bond Formation Through the Fixation of Carbon Dioxide. *Chem. Soc. Rev.* **2011**, *40*, 2435-2452; (b) Omae, I. Recent Developments in Carbon Dioxide Utilization for the Production of Organic Chemicals. *Coord. Chem. Rev.* **2012**, *256*, 1384-1405; (c) Tsuji, Y.; Fujihara, T. Carbon Dioxide as a Carbon Source in Organic Transformation: Carbon-Carbon Bond Forming Reactions by Transition-Metal Catalysts. *Chem. Commun.* **2012**, *48*, 9956-9964; (d) Yeung, C. S.; Dong, V. M. Making C-C Bonds from Carbon Dioxide via Transition-Metal Catalysis. *Top. Catal.* **2014**, *57*, 1342-1350; (e) Liu, Q.; Wu, L.; Jackstell, R.; Beller, M. Using Carbon Dioxide as a Building Block in Organic Synthesis. *Nat. Commun.* **2015**, *6*, 5933; (f) Yu, D.; Teong, S. P.; Zhang, Y. Transition Metal Complex Catalyzed Carboxylation Reactions with CO<sub>2</sub>. *Coord. Chem. Rev.* **2015**, *293-294*, 279-291; (g) Liu, A.-H.; Yu, B.; He, L.-N. Catalytic Conversion of Carbon Dioxide to Carboxylic Acid Derivatives. *Greenhouse Gases: Sci. Technol.* **2015**, *5*, 17-33; (h) Börjesson, M.; Moragas, T.; Gallego, D.; Martin, R. Metal-Catalyzed Carboxylation of Organic (Pseudo)halides with CO<sub>2</sub>. *ACS Catal.* **2016**, *6*, 6739-6749; (i) Tortajada, A.; Juliá-Hernández, F.; Börjesson, M.; Moragas, T.; Martin, R. Transition-Metal-Catalyzed Carboxylation Reactions with Carbon Dioxide. *Angew. Chem. Int. Ed.* **2018**, *57*, 15948-15982; (j) Obst, M.; Pavlovic, L.; Hopmann, K. H. Carbon-Carbon Bonds with CO<sub>2</sub>: Insights from Computational Studies. *J. Organomet. Chem.* **2018**, *864*, 115-127; (k) Yang, Y.; Lee, J.-W. Toward Ideal Carbon Dioxide Functionalization. *Chem. Sci.* **2019**, *10*, 3905-3926; (l) Hong, J.; Li, M.; Zhang, J.; Sun, B.; Mo, F. C-H Bond Carboxylation with Carbon Dioxide. *ChemSusChem* **2019**, *12*, 6-39; (m) Yeung, C. S. Photoredox Catalysis as a Strategy for CO<sub>2</sub> Incorporation: Direct Access to Carboxylic Acids from a Renewable Feedstock. *Angew. Chem. Int. Ed.* **2019**, *58*, 5492-5502.
4. Hazari, N.; Heimann, J. E. Carbon Dioxide Insertion into Group 9 and 10 Metal-Element  $\sigma$ -Bonds. *Inorg. Chem.* **2017**, *56*, 13655-13678.
5. (a) Heimann, J. E.; Bernskoetter, W. H.; Hazari, N.; Mayer, J. M. Acceleration of CO<sub>2</sub> Insertion into Metal Hydrides: Ligand, Lewis Acid, and Solvent Effects on Reaction Kinetics. *Chem. Sci.* **2018**, *8*, 6629-6638; (b) Heimann, J. E.; Bernskoetter, W. H.; Hazari, N. Understanding the Individual and Combined Effects of Solvent and Lewis Acid on CO<sub>2</sub> Insertion into a Metal Hydride. *J. Am. Chem. Soc.* **2019**, *141*, 10520-10529.
6. For an example of CO<sub>2</sub> insertion into a Sc alkyl bond see: LeBlanc, F. A.; Berkefeld, A.; Piers, W. E.; Parvez, M. Reactivity of Scandium  $\beta$ -Diketiminato Alkyl Complexes with Carbon Dioxide. *Organometallics* **2012**, *31*, 810-818.
7. For an example of CO<sub>2</sub> insertion into a Zr alkyl bond see: Lau, K.-C.; Petro, B. J.; Bontemps, S.; Jordan, R. F. Comparative Reactivity of Zr- and Pd-Alkyl Complexes with Carbon Dioxide. *Organometallics* **2013**, *32*, 6895-6898.
8. For an example of CO<sub>2</sub> insertion into a Cr alkyl bond see: Darensbourg, D. J.; Rokicki, A. Reduction of Carbon Dioxide and Carbonyl Sulfide by Anionic Group VIB Metal Hydrides and Alkyls. Carbon-Hydrogen and Carbon-Carbon Bond Formation Processes and the Structure of [PNP][Cr(CO)<sub>5</sub>SC(O)H]. *J. Am. Chem. Soc.* **1982**, *104*, 349-350.
9. For examples of CO<sub>2</sub> insertion into W-alkyl bonds see: (a) Darensbourg, D. J.; Hanckel, R. K.; Bauch, C. G.; Pala, M.; Simmons, D.; White, J. N. A Kinetic Investigation of Carbon Dioxide Insertion Processes Involving Anionic Tungsten-Alkyl and -Aryl Derivatives: Effects of Carbon Dioxide Pressure, Counterions, and Ancillary Ligands. Comparisons with Migratory Carbon Monoxide Insertion Processes. *J. Am. Chem. Soc.* **1985**, *107*, 7463-7473; (b) Darensbourg, D. J.; Pala, M. Cation-Anion Interaction in the [Na-kryptofix-221][W(CO)<sub>5</sub>O<sub>2</sub>CH] Derivative and its Relevance in Carbon Dioxide Reduction Processes. *J. Am. Chem. Soc.* **1985**, *107*, 5687-5693; (c) Darensbourg, D. J.; Grottsch, G. Stereochemical Studies of the Carbon Dioxide Insertion Reactions into the Tungsten-Alkyl Bond. *J. Am. Chem. Soc.* **1985**, *107*, 7473-7476; (d) Darensbourg, D. J.; Kudasoski, R. Metal-Induced Transformations of Carbon Dioxide. Carbon-Carbon Bond-Forming Processes Involving Anionic Group VIB Metal Derivatives, and the X-ray structure of [PNP][cis-MeW(CO)<sub>4</sub>PMe<sub>3</sub>]. *J. Am. Chem. Soc.* **1984**, *106*, 3672-3673.
10. For examples of CO<sub>2</sub> insertion into Fe-alkyl bonds see: (a) Ittel, S.; Tolman, C.; English, A.; Jesson, J. The Chemistry of 2-Naphthyl bis[bis(dimethylphosphino)ethane] Hydride Complexes of Iron, Ruthenium, and Osmium. 2. Cleavage of sp and sp<sup>3</sup> Carbon-Hydrogen, Carbon-Oxygen, and Carbon-Halogen bonds. Coupling of Carbon Dioxide and Acetonitrile. *J. Am. Chem. Soc.* **1978**, *100*, 7577-7585; (b) Arafa, I. M.; Shin, K.; Goff, H. M. Carbon Monoxide and Carbon Dioxide Carbon-Metal Bond Insertion Chemistry of Alkyliron(III) Porphyrin Complexes. *J. Am. Chem. Soc.* **1988**, *110*, 5228-5229; (c) Allen, O. R.; Dalgarno, S. J.; Field, L. D.; Jensen, P.; Turnbull, A. J.; Willis, A. C. Addition of CO<sub>2</sub> to Alkyl Iron complexes, Fe(PP)<sub>2</sub>Me<sub>2</sub>. *Organometallics* **2008**, *27*, 2092-2098; (d) Lau, K.-C.; Jordan, R. F. Reactivity of (Pyridine-Diimine) Fe Alkyl Complexes with Carbon Dioxide. *Organometallics* **2016**, *35*, 3658-3666.
11. For examples of CO<sub>2</sub> insertion into Ru-alkyl bonds see reference 10a and: (a) Allen, O. R.; Dalgarno, S. J.; Field, L. D.; Jensen, P.; Willis, A. C. Insertion of CO<sub>2</sub> into the Ru-C Bonds of cis- and trans-Ru(dmpe)<sub>2</sub>Me<sub>2</sub> (dmpe = Me<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PMe<sub>2</sub>). *Organometallics* **2009**, *28*, 2385-2390; (b) Hartwig, J. F.; Bergman, R. G.; Andersen, R. A. Insertion Reactions of Carbon Monoxide and Carbon Dioxide with Ruthenium Benzyl, Arylamido, and Aryloxide

complexes: A Comparison of the Reactivity of Ruthenium-Carbon, Ruthenium-Nitrogen, and Ruthenium-Oxygen Bonds. *J. Am. Chem. Soc.* **1991**, *113*, 6499-6508.

12. For examples of CO<sub>2</sub> insertion into Rh-alkyl bonds see: (a) Darensbourg, D. J.; Groetsch, G.; Wiegrefe, P.; Rheingold, A. L. Insertion Reactions of Carbon Dioxide with Square-Planar Rhodium Alkyl and Aryl Complexes. *Inorg. Chem.* **1987**, *26*, 3827-3830; (b) Ostapowicz, T. G.; Hölscher, M.; Leitner, W. CO<sub>2</sub> Insertion into Metal-Carbon Bonds: A Computational Study of RhI Pincer Complexes. *Chem. Eur. J.* **2011**, *17*, 10329-10338; (c) Suga, T.; Saitou, T.; Takaya, J.; Iwasawa, N. Mechanistic Study of the Rhodium-Catalyzed Carboxylation of Simple Aromatic Compounds with Carbon Dioxide. *Chem. Sci.* **2017**, *8*, 1454-1462; (d) Pavlovic, L.; Vaitla, J.; Bayer, A.; Hopmann, K. H. Rhodium-Catalyzed Hydrocarboxylation: Mechanistic Analysis Reveals Unusual Transition State for Carbon-Carbon Bond Formation. *Organometallics* **2018**, *37*, 941-948.

13. For examples of CO<sub>2</sub> insertion into Cu-alkyl bonds see: (a) Ikariya, T.; Yamamoto, A. Preparation and Properties of Ligand-Free Methylcopper and of Copper Alkyls Coordinated with 2,2'-Bipyridyl and Tricyclohexylphosphine. *J. Organomet. Chem.* **1974**, *72*, 145-151; (b) Mankad, N. P.; Gray, T. G.; Laitar, D. S.; Sadighi, J. P. Synthesis, Structure, and CO<sub>2</sub> Reactivity of a Two-Coordinate (Carbene)copper(I) Methyl Complex. *Organometallics* **2004**, *23*, 1191-1193; (c) Sakaki, S.; Ohkubo, K. Ab initio MO Study of Carbon Dioxide Insertion into a Methyl-Copper(I) Bond. Critical Difference from CO<sub>2</sub> Insertion into a Hydrogen-Copper(I) Bond. *Organometallics* **1989**, *8*, 2970-2973.

14. The following examples describe CO<sub>2</sub> insertion into Ni(I) alkyl species. In both cases, the starting materials and products are highly unstable, which precludes detailed kinetic studies. (a) Diccianni, J. B.; Hu, C. T.; Diao, T. Insertion of CO<sub>2</sub> Mediated by a (Xantphos)Ni-alkyl Species. *Angew. Chem. Int. Ed.* **2019**, *58*, 13865-13868; (b) Somerville, R. J.; Odena, C.; Obst, M. F.; Hazari, N.; Hopmann, K. H.; Martin, R. Ni(I)-alkyl Complexes Bearing Phenanthroline Ligands: Experimental Evidence for CO<sub>2</sub> Insertion at Ni(I) Centers. *J. Am. Chem. Soc.* **2020**, *142*, 10936-10941.

15. Johansson, R.; Jarenmark, M.; Wendt, O. F. Insertion of Carbon Dioxide into (PCP)Pd<sup>II</sup>-Me Bonds. *Organometallics* **2005**, *24*, 4500-4502.

16. Schmeier, T. J.; Hazari, N.; Incarvito, C. D.; Raskatov, J. R. Exploring the Reactions of CO<sub>2</sub> with PCP Supported Nickel Complexes. *Chem. Commun.* **2011**, *47*, 1824-1826.

17. (a) Jonasson, K. J.; Wendt, O. F. Synthesis and Characterization of a Family of POCOP Pincer Complexes with Nickel: Reactivity Towards CO<sub>2</sub> and Phenylacetylene. *Chem. Eur. J.* **2014**, *20*, 11894-11902; (b) Mousa, A. H.; Bendix, J.; Wendt, O. F. Synthesis, Characterization, and Reactivity of PCN Pincer Nickel Complexes. *Organometallics* **2018**, *37*, 2581-2593; (c) Mousa, A. H.; Polukeev, A. V.; Hansson, J.; Wendt, O. F. Carboxylation of the Ni-Me Bond in an Electron-Rich Unsymmetrical PCN Pincer Nickel Complex. *Organometallics* **2020**, *39*, 1553-1560.

18. (a) Reichardt, C. Empirical Parameters of the Polarity of Solvents. *Angew. Chem. Int. Ed.* **1965**, *4*, 29-40; (b) Reichardt, C. Solvatochromic Dyes as Solvent Polarity Indicators. *Chem. Rev.* **1994**, *94*, 2319-2358.


19. Suh, H.-W.; Schmeier, T. J.; Hazari, N.; Kemp, R. A.; Takase, M. K. Experimental and Computational Studies of the Reaction of Carbon Dioxide with Pincer-Supported Nickel and Palladium Hydrides. *Organometallics* **2012**, *31*, 8225-8236.

20. Mitton, S. J.; McDonald, R.; Turculet, L. Nickel and Palladium Silyl Pincer Complexes: Unusual Structural Rearrangements that Involve Reversible Si-C(sp<sup>3</sup>) and Si-C(sp<sup>2</sup>) Bond Activation. *Angew. Chem. Int. Ed.* **2009**, *48*, 8568-8571.

21. (a) Segawa, Y.; Yamashita, M.; Nozaki, K. Syntheses of PBP Pincer Iridium Complexes: A Supporting Boryl Ligand. *J. Am. Chem. Soc.* **2009**, *131*, 9201-9203; (b) Segawa, Y.; Yamashita, M.; Nozaki, K. Diphenylphosphino- or Dicyclohexylphosphino-Tethered Boryl Pincer Ligands: Syntheses of PBP Iridium(III) Complexes and Their Conversion to Iridium-Ethylene Complexes. *Organometallics* **2009**, *28*, 6234-6242; (c) Hill, A. F.; Lee, S. B.; Park, J.; Shang, R.; Willis, A. C. Analogies between Metallaboratranes, Triboronates, and Boron Pincer Ligand Complexes. *Organometallics* **2010**, *29*, 5661-5669; (d) Hasegawa, M.; Segawa, Y.; Yamashita, M.; Nozaki, K. Isolation of a PBP-Pincer Rhodium Complex Stabilized by an Intermolecular C-H  $\sigma$  Coordination as the Fourth Ligand. *Angew. Chem. Int. Ed.* **2012**, *51*, 6956-6960; (e) Ogawa, H.; Yamashita, M. Platinum Complexes Bearing a Boron-Based PBP Pincer Ligand: Synthesis, Structure, and Application as a Catalyst for Hydrosilylation of 1-Decene. *Dalton Trans.* **2013**, *42*, 625-629; (f) Miyada, T.; Yamashita, M. Oxygenation of a Ruthenium Complex Bearing a PBP-Pincer Ligand Inducing the Formation of a Boronate Ligand with a Weak Ru-O Bond. *Organometallics* **2013**, *32*, 5281-5284; (g) Lin, T.-P.; Peters, J. C. Boryl-Metal Bonds Facilitate Cobalt/Nickel-Catalyzed Olefin Hydrogenation. *J. Am. Chem. Soc.* **2014**, *136*, 13672-13683; (h) Curado, N.; Maya, C.; López-Serrano, J.; Rodríguez, A. Boryl-Assisted Hydrogenolysis of a Nickel-Methyl Bond. *Chem. Commun.* **2014**, *50*, 15718-15721; (i) Ríos, P.; Curado, N.; López-Serrano, J.; Rodríguez, A. Selective Reduction of Carbon Dioxide to Bis(silyl)acetal Catalyzed by a PBP-Supported Nickel Complex. *Chem. Commun.* **2016**, *52*, 2114-2117; (j) Ríos, P.; Rodríguez, A.; Lopez-Serrano, J. Mechanistic Studies on the Selective Reduction of CO<sub>2</sub> to the Aldehyde Level by a Bis(phosphino)boryl (PBP)-Supported Nickel Complex. *ACS Catal.*

- 2016**, *6*, 5715-5723; (k) Ding, Y.; Ma, Q.-Q.; Kang, J.; Zhang, J.; Li, S.; Chen, X. Palladium(II) Complexes Supported by PBP and POCOP Pincer Ligands: A Comparison of their Structure, Properties and Catalytic Activity. *Dalton Trans.* **2019**, *48*, 17633-17643.
22. Adhikari, D.; Huffman, J. C.; Mindiola, D. J. Structural Elucidation of a Nickel Boryl Complex. A Recyclable Borylation Ni(II) Reagent of Bromobenzene. *Chem. Commun.* **2007**, 4489-4491.
23. (a) Clavier, H.; Nolan, S. P. Percent Buried Volume for Phosphine and N-Heterocyclic Carbene Ligands: Steric Properties in Organometallic Chemistry. *Chem. Commun.* **2010**, *46*, 841-861; (b) Falivene, L.; Credendino, R.; Poater, A.; Petta, A.; Serra, L.; Oliva, R.; Scarano, V.; Cavallo, L. SambVca 2. A Web Tool for Analyzing Catalytic Pockets with Topographic Steric Maps. *Organometallics* **2016**, *35*, 2286-2293.
24. Kimmich, B. F. M.; Marshall, W. J.; Fagan, P. J.; Hauptman, E.; Bullock, R. M. Palladium Complexes with PCP Ligands: Preparation and Structural Studies of Two Forms of  $[(PCP)Pd(OH_2)]^+BF_4^-$ . *Inorg. Chim. Acta* **2002**, *330*, 52-58.
25. (a) Beck, R.; Shoshani, M.; Krasinkiewicz, J.; Hatnean, J. A.; Johnson, S. A. Synthesis and Chemistry of Bis(triisopropylphosphine) Nickel(I) and Nickel(0) Precursors. *Dalton Trans.* **2013**, *42*, 1461-1475; (b) Harman, W. H.; Lin, T.-P.; Peters, J. C. A  $d^{10}$  Ni-(H<sub>2</sub>) Adduct as an Intermediate in H-H Oxidative Addition across a Ni-B Bond. *Angew. Chem. Int. Ed.* **2014**, *53*, 1081-1086; (c) Kim, Y.-E.; Kim, J.; Lee, Y. Formation of a Nickel Carbon Dioxide Adduct and its Transformation Mediated by a Lewis Acid. *Chem. Commun.* **2014**, *50*, 11458-11461; (d) Kim, Y.-E.; Oh, S.; Kim, S.; Kim, O.; Kim, J.; Han, S. W.; Lee, Y. Phosphinite-Ni(0) Mediated Formation of a Phosphide-Ni(II)-OCOOME Species via Uncommon Metal-Ligand Cooperation. *J. Am. Chem. Soc.* **2015**, *137*, 4280-4283; (e) Charboneau, D. J.; Balcells, D.; Hazari, N.; Lant, H. M. C.; Mayer, J. M.; Melvin, P. R.; Mercado, B. Q.; Morris, W. D.; Repisky, M.; Suh, H.-W. Dinitrogen-Facilitated Reversible Formation of a Si-H Bond in a Pincer-Supported Ni Complex. *Organometallics* **2016**, *35*, 3154-3162.
26. Johnson, M. T.; Johansson, R.; Kondrashov, M. V.; Steyl, G.; Ahlquist, M. S. G.; Roodt, A.; Wendt, O. F. Mechanisms of the CO<sub>2</sub> Insertion into (PCP) Palladium Allyl and Methyl  $\sigma$ -Bonds. A Kinetic and Computational Study. *Organometallics* **2010**, *29*, 3521-3529.
27. Mayer, U.; Gutmann, V.; Gerger, W. The Acceptor Number — A Quantitative Empirical Parameter for the Electrophilic Properties of Solvents. *Monatsh. Chem.* **1975**, *106*, 1235-1257.
28. Charboneau, D. J.; Brudvig, G. W.; Hazari, N.; Lant, H. M.; Saydjari, A. K. Development of an Improved System for the Carboxylation of Aryl Halides Through Mechanistic Studies. *ACS Catal.* **2019**, *9*, 3228-3241.
29. THF was selected as the solvent due to the improved solubility of many Lewis acids in THF compared to other ethereal solvents.
30. García-López, D.; Pavlovic, L.; Hopmann, K. H. To Bind or Not to Bind: Mechanistic Insights into C-CO<sub>2</sub> Bond Formation with Late Transition Metals. *Organometallics* **2020**, *39*, 1339-1347.
31. We also calculated CO<sub>2</sub> insertion into  $(^{Me}PCP)Pd(CH_3)$ , where the *tert*-butyl substituents on the <sup>t</sup>BuPCP ligand have been replaced with methyl substituents. The calculations indicate that for  $(^{Me}PCP)Pd(CH_3)$ , the 1,2-insertion pathway is preferred over the S<sub>E</sub>2 mechanism by 1.8 kcal/mol (Table 3, Figures S53-S54). This is consistent with the hypothesis that the steric properties of the pincer ligand play a large role in determining the pathway for CO<sub>2</sub> insertion.
32. We note that other computational protocols ( $\omega$ B97XD, PBE0-D3BJ) provide the same reactivity trends as B3LYP-D3 (Table S3).
33. Our results do not explain Wendt's observation (see references 17b and 17c) that CO<sub>2</sub> insertion into  $(^{tBu}PC^{iPr}N)Ni(CH_3)$  ( $^{tBu}PC^{iPr}N = 1-(3-((di-tert-butylphosphino)methyl)phenyl)-N,N$ -diisopropyl-methanamine)) is faster than insertion into  $(^{tBu}PC^{Me}N)Ni(CH_3)$  ( $^{tBu}PC^{Me}N = 1-(3-((di-tert-butylphosphino)methyl)phenyl)-N,N$ -methyl-methanamine)). In this case, ligand hemilability may play a role and it is possible that the more sterically bulky  $^{tBu}PC^{iPr}N$  is more likely to decoordinate.

## TOC Graphic



- *Rapid insertion due to trans-influence of boryl ligand*
- *Computations show  $\text{CO}_2$  inserts through an  $\text{S}_{\text{E}}2$  pathway*
- *Mild conditions allow for kinetic studies*

Trans Effect	Metal Effect	Steric Effect	Solvent Effect
PBP > PCP	Pd > Ni	Cy > <sup>t</sup> Bu	Follows $\text{E}_{\text{T}}(30)$