

An application of the Gaussian correlation inequality to the small deviations for a Kolmogorov diffusion*

Marco Carfagnini[†]

Abstract

We consider an iterated Kolmogorov diffusion X_t of step n . The small ball problem for X_t is solved by means of the Gaussian correlation inequality. We also prove Chung's laws of iterated logarithm for X_t both at time zero and infinity.

Keywords: Diffusion processes, Kolmogorov diffusion, functional limit laws, small ball problem.

MSC2020 subject classifications: Primary 60F17, Secondary 60F15; 60G51; 60J65.

Submitted to ECP on January 2, 2022, final version accepted on February 22, 2022.

Contents

1	Introduction	1
2	The setting and main results	3
3	Proofs	4
	References	6

1 Introduction

Let $\{X_t\}_{0 \leq t \leq T}$ be an \mathbb{R}^n -valued stochastic process with continuous paths such that $X_0 = 0$ a.s. where $T > 0$ is fixed. Denote by $W_0(\mathbb{R}^n)$ the space of \mathbb{R}^n -valued continuous functions on $[0, T]$ starting at zero. Given a norm $\|\cdot\|$ on $W_0(\mathbb{R}^n)$, the small ball problem for X_t consists in finding the rate of explosion of

$$-\log \mathbb{P} (\|X\| < \varepsilon)$$

as $\varepsilon \rightarrow 0$. More precisely, a process X_t is said to satisfy a *small deviation principle* with rates α and β if there exist a constant $c > 0$ such that

$$\lim_{\varepsilon \rightarrow 0} -\varepsilon^\alpha |\log \varepsilon|^\beta \log \mathbb{P} (\|X\| < \varepsilon) = c. \quad (1.1)$$

The values of α, β and c depend on the process X_t and on the chosen norm on $W_0(\mathbb{R}^n)$. Small deviation principles have many applications including metric entropy estimates

*Research was supported in part by NSF Grants DMS-1712427 and DMS-1954264.

[†]University of Connecticut, United States of America. E-mail: marco.carfagnini@uconn.edu

and Chung's law of the iterated logarithm. We refer to the survey paper [13] for more details.

We say that a process X_t satisfies *Chung's law of the iterated logarithm* (LIL) as $t \rightarrow \infty$ (resp. as $t \rightarrow 0$) with rate $a \in \mathbb{R}_+$ if there exists a constant C such that

$$\liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^a \max_{0 \leq s \leq t} |X_s| = C, \quad a.s. \quad (1.2)$$

(resp. $\liminf_{t \rightarrow 0} \left(\frac{\log |\log t|}{t} \right)^a \max_{0 \leq s \leq t} |X_s| = C$ a.s.). When X_t is a Brownian motion, it was proven in a famous paper by K.-L. Chung in 1948 that (1.2) holds with $a = \frac{1}{2}$ and $C = \frac{\pi}{\sqrt{8}}$. To find the rates α and β such that the limit in (1.1) exists, and then find the constant c is an extremely hard problem in general. Even the estimation of the rate of explosion of (1.1) is usually a difficult problem. Indeed, as can be surmised in [10, 15], the small ball problem for Gaussian processes is equivalent to metric entropy problems in functional analysis. In [11] and [20] a Brownian sheet in Hölder and uniform norm is considered, and the integrated Brownian motion in the uniform norm is the content of [8], and the m -fold integrated Brownian motion in both the uniform and L^2 -norm is considered in [4]. In [17] and [3] a small deviation principle and Chung's LIL are proved for a class of stochastic integrals and for a hypoelliptic Brownian motion on the Heisenberg group. When X_t is a Gaussian process with stationary increments, upper and lower bounds on (1.1) can be found in [19, 16].

In this paper we consider the Kolmogorov diffusion of step n .

Definition 1.1. Let $T > 0$ and b_t be a one-dimensional standard Brownian motion. The stochastic process $\{X_t\}_{0 \leq t \leq T}$ on \mathbb{R}^n defined by

$$X_t := \left(b_t, \int_0^t b_{t_2} dt_2, \int_0^t \int_0^{t_2} b_{t_3} dt_3 dt_2, \dots, \int_0^t \int_0^{t_2} \dots \int_0^{t_{n-1}} b_{t_n} dt_n \dots dt_2 \right)$$

is the Kolmogorov diffusion of step n .

$\{X_t\}_{0 \leq t \leq T}$ is a Markov process with generator given by $L = \frac{1}{2} \frac{\partial^2}{\partial x_1^2} + \sum_{d=2}^n x_{d-1} \frac{\partial}{\partial x_d}$. In particular, when $n = 2$ X_t is the Markov process associated to the differential operator $L = \frac{1}{2} \frac{\partial^2}{\partial x^2} + x \frac{\partial}{\partial y}$ and it was first introduced by A. N. Kolmogorov in [9], where he obtained an explicit expression for its transition density. Later, L. Hörmander in [6] used L as the simplest example of a hypoelliptic second order differential operator. More precisely, the operator L satisfies the weak Hörmander condition. $\{X_t\}_{0 \leq t \leq T}$ is a Gaussian process and its law μ is a Gaussian measure on the Banach space $(W_0(\mathbb{R}^n), \|\cdot\|)$, where

$$\|f\| := \max_{0 \leq t \leq T} |f(t)|, \quad \forall f \in W_0(\mathbb{R}^n).$$

The main result of this paper is Theorem 2.6, where we prove the small deviation principle (1.1) for X_t with rates $\alpha = 2$, $\beta = 0$, and constant $c = \frac{\pi}{\sqrt{8}}$. Our proof relies on the Gaussian correlation inequality (GCI), see e.g. [18, 12], applied to the Gaussian measure μ on $W_0(\mathbb{R}^n)$. A different application of the GCI to estimate small balls probabilities is given in [14]. In Theorem 2.6 we also state Chung's LIL at time zero and infinity for X_t with rates given by $a = \frac{1}{2}$ and $a = \frac{2n-1}{2}$ respectively.

The stochastic processes considered in [8, 17, 3] all satisfy a scaling property, that is, there exists a scaling constant $\delta \in (0, \infty)$ such that $X_{\varepsilon t} \stackrel{(d)}{=} \varepsilon^\delta X_t$. Properties of Gaussian measures on Banach spaces and scaling properties have been used to show the existence of a small deviation principle for some processes such as a Brownian motion with values in a finite dimensional Banach space in [5] and an integrated Brownian motion in [8]. Moreover, in [8, 17, 3] the scaling rate δ coincides with the rate of Chung's LIL at infinity,

and the small deviations' rates are given by $\beta = 0$, $\alpha = \frac{1}{\delta}$. The Kolmogorov diffusion does not satisfy a scaling property with respect to the standard Euclidean norm, and the small deviations rate α is not related to the Chung's LIL rate.

Lastly, large deviations and Chung's LIL at time zero for the limsup of the Kolmogorov diffusion are discussed in Section [1, Section 4.2] and [2, Example 3.5] respectively.

The paper is organized as follows. In Section 2 we collect some examples and state the main result of this paper, namely, small deviation principle and Chung's LIL at time zero and infinity for a step n Kolmogorov diffusion. Section 3 contains the proof of the main result.

2 The setting and main results

Notation 2.1. 4 Let X_t be an \mathbb{R}^n -valued stochastic process with $X_0 = 0$ a.s. Then X_t^* denotes the process defined by

$$X_t^* := \max_{0 \leq s \leq t} |X_s|,$$

where $|\cdot|$ denotes the Euclidean norm.

Notation 2.2. [Dirichlet eigenvalues in \mathbb{R}^n] We denote by $\lambda_1^{(n)}$ the lowest Dirichlet eigenvalue of $-\frac{1}{2}\Delta_{\mathbb{R}^n}$ on the unit ball in \mathbb{R}^n .

Let us collect some examples of Chung's LIL and small deviation principle.

Example 2.3. [Brownian motion] Let X_t be a standard Brownian motion. Then $X_{\varepsilon t} \stackrel{(d)}{=} \varepsilon^{\frac{1}{2}} X_t$, and it satisfies the small deviation principle

$$\lim_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P}(X_T^* < \varepsilon) = \lambda_1^{(1)} T, \quad (2.1)$$

where $\lambda_1^{(1)}$ is defined in Notation 2.2, see e.g. [7, Lemma 8.1]. Moreover, in a famous paper by K.-L. Chung in 1948 it was proven that

$$\liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^{\frac{1}{2}} \max_{0 \leq s \leq t} |X_s| = \sqrt{\lambda_1^{(1)}} \quad a.s. \quad (2.2)$$

Example 2.4. [Integrated Brownian motion]. Let $X_t := \int_0^t b_s ds$, where b_s is a one-dimensional standard Brownian motion. It is easy to see that $X_{\varepsilon t} \stackrel{(d)}{=} \varepsilon^{\frac{3}{2}} X_t$. In [8] it is shown that there exists a finite constant $c_0 > 0$ such that

$$\liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^{\frac{3}{2}} \max_{0 \leq s \leq t} |X_s| = c_0 \quad a.s. \quad (2.3)$$

and (2.3) was used to prove that

$$\lim_{\varepsilon \rightarrow 0} -\varepsilon^{\frac{2}{3}} \log \mathbb{P}(X_1^* < \varepsilon) = c_0^{\frac{2}{3}}.$$

Example 2.5. [Iterated integrated Brownian motion] Let b_t be a one-dimensional Brownian motion starting at zero. Denote by $X_1(t) := b_t$ and

$$X_d(t) := \int_0^t X_{d-1}(s) ds, \quad t \geq 0, d \geq 2,$$

the d -fold integrated Brownian motion for a positive integer d . Note that $X_d(\varepsilon t) \stackrel{(d)}{=} \varepsilon^{\frac{2d-1}{2}} X_d(t)$. In [4] it was shown that for any integer d there exists a constant $\gamma_d > 0$ such

that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} -\varepsilon^{\frac{2}{2d-1}} \log \mathbb{P} \left(\max_{0 \leq t \leq 1} |X_d(t)| < \varepsilon \right) &= \gamma_d^{\frac{2}{2d-1}}, \\ \liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^{\frac{2d-1}{2}} \max_{0 \leq s \leq t} |X_d(s)| &= \gamma_d \quad a.s. \end{aligned} \quad (2.4)$$

Our main object is the Kolmogorov diffusion on \mathbb{R}^n defined by

$$X_t := (X_1(t), \dots, X_n(t)),$$

where

$$X_d(t) := \int_0^t \int_0^{t_2} \cdots \int_0^{t_{d-1}} b_{t_d} dt_d \cdots dt_2, \quad \text{for } d = 3, \dots, n,$$

and $X_2(t) := \int_0^t b_s ds$, $X_1(t) := b_t$, where b_t is a one-dimensional standard Brownian motion. Note that $X_d(\varepsilon t) \stackrel{(d)}{=} \varepsilon^{\frac{2d-1}{2}} X_d(t)$ for all $d = 1, \dots, n$, and hence the process X_t does not have a scaling property with respect to the Euclidean norm $|\cdot|$ in \mathbb{R}^n .

Theorem 2.6. *Let $T > 0$ and X_t be the Kolmogorov diffusion on \mathbb{R}^n . Then*

$$\lim_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P} (X_T^* < \varepsilon) = \lambda_1^{(1)} T, \quad (2.5)$$

$$\liminf_{t \rightarrow 0} \sqrt{\frac{\log |\log t|}{t}} \max_{0 \leq s \leq t} |X_s| = \sqrt{\lambda_1^{(1)}} \quad a.s. \quad (2.6)$$

$$\liminf_{t \rightarrow \infty} \left(\frac{\log \log t}{t} \right)^{\frac{2n-1}{2}} \max_{0 \leq s \leq t} |X_s| = \gamma_n \quad a.s. \quad (2.7)$$

where $\lambda_1^{(1)}$ is defined in Notation 2.2, and γ_n is given by (2.4)

Remark 2.7. By (2.2) and Brownian inversion, it follows that a standard Brownian motion satisfies Chung's LIL at time zero and infinity with rate $a = \frac{1}{2}$, and it satisfies a small deviation principle with rate $\alpha = 2$. By (2.5), the n -step Kolmogorov diffusion X_t satisfies the same small deviation principle as a one-dimensional standard Brownian motion. As far as Chung's LIL for X_t is concerned, the first component dominates when $t \rightarrow 0$ with rate $a = \frac{1}{2}$, and the n -th component dominates as $t \rightarrow \infty$ with rate $a = \frac{2n-1}{2}$.

3 Proofs

Proof of Theorem 2.6. Let us first prove the small deviation principle (2.5). One has that $\mathbb{P} (X_T^* < \varepsilon) \leq \mathbb{P} (b_T^* < \varepsilon)$, and hence by (2.1) it follows that

$$\lambda_1^{(1)} T \leq \liminf_{\varepsilon \rightarrow 0} -\varepsilon^2 \mathbb{P} (X_T^* < \varepsilon).$$

Let us now show that

$$\limsup_{\varepsilon \rightarrow 0} -\varepsilon^2 \mathbb{P} (X_T^* < \varepsilon) \leq \lambda_1^{(1)} T.$$

For any $x_1, \dots, x_n \in (0, 1)$ such that $x_1 + \cdots + x_n = 1$ we have that

$$\begin{aligned} \mathbb{P} (X_T^* < \varepsilon) &\geq \mathbb{P} \left(\max_{0 \leq t \leq T} |X_1(t)| < x_1 \varepsilon, \dots, \max_{0 \leq t \leq T} |X_n(t)| < x_n \varepsilon, \right) \\ &\geq \mathbb{P} \left(\max_{0 \leq t \leq T} |X_1(t)| < x_1 \varepsilon \right) \cdots \mathbb{P} \left(\max_{0 \leq t \leq T} |X_n(t)| < x_n \varepsilon \right), \end{aligned}$$

where in the second line we used the Gaussian correlation inequality for the law of the process $\{X_t\}_{0 \leq t \leq T}$ which is a Gaussian measure on $W_0(\mathbb{R}^n)$. Thus,

$$-\varepsilon^2 \log \mathbb{P}(X_T^* < \varepsilon) \leq -\sum_{d=1}^n \varepsilon^2 \log \mathbb{P}\left(\max_{0 \leq t \leq T} |X_d(t)| < x_d \varepsilon\right). \quad (3.1)$$

Note that, for any $d = 2, \dots, n$

$$\max_{0 \leq t \leq T} |X_d(t)| \leq \int_0^T \int_0^{t_2} \cdots \int_0^{t_{d-2}} \max_{0 \leq t \leq T} |X_2(t)| dt_{d-1} \cdots dt_2 = \frac{T^{d-2}}{(d-2)!} \max_{0 \leq t \leq T} |X_2(t)|,$$

and hence

$$\mathbb{P}\left(\max_{0 \leq t \leq T} |X_2(s)| < \frac{(d-2)!}{T^{d-2}} x_d \varepsilon\right) \leq \mathbb{P}\left(\max_{0 \leq t \leq T} |X_d(s)| < x_d \varepsilon\right), \quad (3.2)$$

and by [8, Theorem 1.1] we have that, for any $d = 2, \dots, n$

$$\begin{aligned} 0 &\leq \limsup_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P}\left(\max_{0 \leq t \leq T} |X_d(t)| < x_d \varepsilon\right) \\ &\leq \lim_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P}\left(\max_{0 \leq t \leq T} |X_2(t)| < \frac{(d-2)!}{T^{d-2}} x_d \varepsilon\right) \\ &= \lim_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P}\left(\max_{0 \leq t \leq T} \left| \int_0^t b_s ds \right| < \frac{(d-2)!}{T^{d-2}} x_d \varepsilon\right) = 0. \end{aligned} \quad (3.3)$$

Thus, by (3.1) and (3.2)

$$-\varepsilon^2 \log \mathbb{P}(X_T^* < \varepsilon) \leq -\varepsilon^2 \log \mathbb{P}(b_T^* < x_1 \varepsilon) - \sum_{d=2}^n \varepsilon^2 \log \mathbb{P}\left(\max_{0 \leq t \leq T} |X_2(t)| < \frac{(d-2)!}{T^{d-2}} x_d \varepsilon\right),$$

and by (3.3) and (2.1) it follows that

$$\limsup_{\varepsilon \rightarrow 0} -\varepsilon^2 \log \mathbb{P}(X_T^* < \varepsilon) \leq \frac{\lambda_1^{(1)}}{x_1^2} T.$$

The result follows by letting x_1 go to one.

Let us now prove (2.6). By (2.2) and Brownian time inversion one has that

$$\liminf_{t \rightarrow 0} \sqrt{\frac{\log |\log t|}{t}} \max_{0 \leq s \leq t} |b_s| = \sqrt{\lambda_1} \quad a.s. \quad (3.4)$$

Note that

$$\begin{aligned} |b_s|^2 &\leq |X_s|^2 = |b_s|^2 + \sum_{d=2}^n |X_d(s)|^2 \\ &\leq |b_s|^2 + \max_{0 \leq u \leq s} |b_u|^2 \sum_{d=2}^n \frac{s^{2d-2}}{(d-1)!^2}, \end{aligned}$$

and hence

$$\begin{aligned} \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |b_s|^2 &\leq \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |X_s|^2 \\ &\leq \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |b_s|^2 \left(1 + \sum_{d=2}^n \frac{t^{2d-2}}{(d-1)!^2}\right) \end{aligned}$$

By (3.4) it follows that, for any $d = 2, \dots, n$

$$\lim_{t \rightarrow 0} t^{2d-2} \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |b_s|^2 = 0 \quad a.s.$$

and thus

$$\liminf_{t \rightarrow 0} \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |X_s|^2 = \liminf_{t \rightarrow 0} \frac{\log |\log t|}{t} \max_{0 \leq s \leq t} |b_s|^2 = \lambda_1 \quad a.s.$$

which completes the proof of (2.6). Let us now prove (2.7). Set

$$\phi(t) := \frac{\log \log t}{t}.$$

By (2.4) we have that, for any $d = 1, \dots, n-1$

$$\liminf_{t \rightarrow \infty} \phi(t)^{\frac{2n-1}{2}} \max_{0 \leq s \leq t} |X_d(s)| = \liminf_{t \rightarrow \infty} \phi(t)^{n-d} \phi(t)^{\frac{2d-1}{2}} \max_{0 \leq s \leq t} |X_d(s)| = 0 \quad a.s. \quad (3.5)$$

since $\phi(t) \rightarrow 0$ as $t \rightarrow \infty$. Note that

$$|X_n(s)|^2 \leq |X_s|^2 = \sum_{d=1}^{n-1} |X_d(s)|^2 + |X_n(s)|^2,$$

and hence

$$\begin{aligned} \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_n(s)|^2 &\leq \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_s|^2 \\ &\leq \sum_{d=1}^{n-1} \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_d(s)|^2 + \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_n(s)|^2. \end{aligned}$$

Thus, by (2.4) and (3.5) it follows that

$$\liminf_{t \rightarrow \infty} \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_s|^2 = \liminf_{t \rightarrow \infty} \phi(t)^{2n-1} \max_{0 \leq s \leq t} |X_n(s)|^2 = \gamma_n^2 \quad a.s.$$

and (2.7) is proven. \square

References

- [1] Gérard Ben Arous and Jing Wang, *Very rare events for diffusion processes in short time*, 2019.
- [2] Marco Carfagnini, Juraj Földes, and David P. Herzog, *A functional law of the iterated logarithm for weakly hypoelliptic diffusions at time zero*, 2021.
- [3] Marco Carfagnini and Maria Gordina, *Small deviations and Chung's law of iterated logarithm for a hypoelliptic Brownian motion on the Heisenberg group*, 2020.
- [4] Xia Chen and Wenbo V. Li, *Quadratic functionals and small ball probabilities for the m -fold integrated Brownian motion*, Ann. Probab. **31** (2003), no. 2, 1052–1077. MR1964958
- [5] Alejandro de Acosta, *Small deviations in the functional central limit theorem with applications to functional laws of the iterated logarithm*, Ann. Probab. **11** (1983), no. 1, 78–101. MR682802
- [6] Lars Hörmander, *Hypoelliptic second order differential equations*, Acta Math. **119** (1967), 147–171. MR0222474
- [7] Nobuyuki Ikeda and Shinzo Watanabe, *Stochastic differential equations and diffusion processes*, second ed., North-Holland Mathematical Library, vol. 24, North-Holland Publishing Co., Amsterdam, 1989. MRMR1011252
- [8] Davar Khoshnevisan and Zhan Shi, *Chung's law for integrated Brownian motion*, Trans. Amer. Math. Soc. **350** (1998), no. 10, 4253–4264. MR1443196

- [9] A. Kolmogoroff, *Über das Gesetz des iterierten Logarithmus*, Ann. of Math. **35** (1934), no. 1, 116–117. MR1503147
- [10] James Kuelbs and Wenbo V. Li, *Metric entropy and the small ball problem for Gaussian measures*, J. Funct. Anal. **116** (1993), no. 1, 133–157. MR1237989
- [11] James Kuelbs and Wenbo V. Li, *Small ball estimates for Brownian motion and the Brownian sheet*, J. Theoret. Probab. **6** (1993), no. 3, 547–577. MR1230346
- [12] Rafał Latała and Dariusz Matlak, *Royen's proof of the Gaussian correlation inequality*, Geometric aspects of functional analysis, Lecture Notes in Math., vol. 2169, Springer, Cham, 2017, pp. 265–275. MR3645127
- [13] W. V. Li and Q.-M. Shao, *Gaussian processes: inequalities, small ball probabilities and applications*, Stochastic processes: theory and methods, Handbook of Statist., vol. 19, North-Holland, Amsterdam, 2001, pp. 533–597. MR1861734
- [14] Wenbo V. Li, *A Gaussian correlation inequality and its applications to small ball probabilities*, Electron. Comm. Probab. **4** (1999), 111–118. MR1741737
- [15] Wenbo V. Li and Werner Linde, *Approximation, metric entropy and small ball estimates for Gaussian measures*, Ann. Probab. **27** (1999), no. 3, 1556–1578. MR1733160
- [16] Ditlev Monrad and Holger Rootzén, *Small values of Gaussian processes and functional laws of the iterated logarithm*, Probab. Theory Related Fields **101** (1995), no. 2, 173–192. MR1318191
- [17] Bruno Rémillard, *On Chung's law of the iterated logarithm for some stochastic integrals*, Ann. Probab. **22** (1994), no. 4, 1794–1802. MR1331204
- [18] Thomas Royen, *A simple proof of the Gaussian correlation conjecture extended to some multivariate gamma distributions*, Far East J. Theor. Stat. **48** (2014), no. 2, 139–145. MR3289621
- [19] Qi Man Shao, *A note on small ball probability of a Gaussian process with stationary increments*, J. Theoret. Probab. **6** (1993), no. 3, 595–602. MR1230348
- [20] Michel Talagrand, *The small ball problem for the Brownian sheet*, Ann. Probab. **22** (1994), no. 3, 1331–1354. MR1303647

Acknowledgments. The author thanks Davar Khoshnevisan and Zhan Shi for suggesting the Gaussian correlation inequality [18, 12].

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴ , ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>