

Atlantic circulation change still uncertain

K. Halimeda Kilbourne^{a*}, Alan D. Wanamaker^b, Paola Moffa-Sanchez^c, David J. Reynolds^d, Daniel E. Amrhein^e, Paul G. Butler^d, Geoffrey Gebbie^f, Marlos Goes^g, Malte F. Jansen^h, Christopher M. Littleⁱ, Madelyn Mette^j, Eduardo Moreno-Chamarro^k, Pablo Ortega^k, Bette L. Otto-Bliesner^e, Thomas Rossby^l, James Scourse^d, and Nina M. Whitney^f

^aUniversity of Maryland Center for Environmental Science, Chesapeake Biological Laboratory, USA,

^bDepartment of Geological and Atmospheric Sciences, Iowa State University, USA,

^cGeography Department, Durham University, DH1 3LE, UK,

^dCentre for Geography and Environmental Sciences, University of Exeter, Penryn, Cornwall, TR10 9EZ, UK,

^eClimate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, Colorado, USA,

^dCentre for Geography and Environmental Sciences, University of Exeter, Penryn, Cornwall, TR10 9EZ, UK,

^fWoods Hole Oceanographic Institution, USA

^gCooperative Institute of Marine and Atmospheric Studies, University of Miami, and Atlantic Oceanographic and Meteorological Laboratory, National Oceanic and Atmospheric Administration, Miami, USA,

^hDepartment of the Geophysical Sciences, The University of Chicago, Chicago, IL, USA,

ⁱOceanography Department, Atmospheric and Environmental Research, Inc., USA,

^jU.S. Geological Survey, St. Petersburg Coastal and Marine Science Center, St. Petersburg, Florida, USA,

^kBarcelona Supercomputing Center, Barcelona, Spain,

^lGraduate School of Oceanography, University of Rhode Island, USA,

* corresponding author

Deep oceanic overturning circulation in the Atlantic (Atlantic Meridional Overturning Circulation, AMOC) is projected to decrease in the future in response to anthropogenic warming. Caesar et al.¹ argue that an AMOC slowdown started in the 19th century and intensified during the mid-20th century. Although the argument and selected evidence proposed have some merits, we find that their conclusions might be different if a more complete array of data available in the North Atlantic region had been considered. We argue that the strength of AMOC over recent centuries is still poorly constrained and the expected slowdown may not have started yet.

Recently, Moffa-Sanchez et al.² compiled a comprehensive set of paleoclimate proxy data from the North Atlantic and Arctic regions using objective criteria for identifying high-quality datasets of ocean conditions spanning the last two millennia (Figure 1). Although no direct (singular) proxy for AMOC exists, the paleoceanographic proxy data compiled by Moffa-Sanchez et al.² highlight the spatial and temporal complexities of ocean state in modern times and the recent past. When all the available proxy records potentially related to AMOC variability and 20th century observational datasets are considered, the time history of the AMOC system becomes less certain. In contrast, selecting only a subset of proxy records that share similar trends, as performed by Caesar et al.¹, provides an incomplete perspective on AMOC changes through time.

Increased data availability in recent decades has enabled a shift in the fields of paleoceanography and paleoclimatology toward more objective and transparent data selection in studies aimed at quantitatively reconstructing past variability. Such screening methods tend to minimize the impact of spurious or less reliable records on analyses and work to enhance the common signal in proxy records. Additionally, analyzing networks of suitable and carefully selected data enables robust uncertainty estimates on the resulting reconstructions, which is essential in providing confidence in the results and the ability to compare information across

60 disciplines. Key to such work is identifying robust criteria and weighting schemes that objectively
61 identify and utilize the most reliable data. Caesar et al.¹ use a variety of proxy records in their
62 analysis, but do not identify the reasoning or criteria for selecting those records over many
63 others that are likely related to aspects of AMOC dynamics (see the recent review²).

64 Objective and inclusive data selection standards are especially important when addressing
65 AMOC, which is a system composed of many different components that can behave differently
66 at different latitudes, depths, and timescales³ and looking at any singular index of AMOC
67 inherently oversimplifies the system. The complex signals in the available AMOC-related proxy
68 variables over recent centuries support this notion², though many of these studies were not
69 considered by Caesar et al.¹

70 In addition to the need for objective standards, we argue that most of the records
71 compiled in the Caesar et al. paper have substantial caveats that were not discussed.
72 Reconstructing the strength of AMOC more than a few decades ago relies upon paleoclimate
73 and paleoceanographic proxies because direct measurements are unavailable. Some proxies
74 are more directly related to components of AMOC variability than others, and some sites are
75 better situated to record specific oceanographic and atmospheric processes than others. The
76 limited scope of data utilized combined with the inherent uncertainties in the proxies and
77 conflicting evidence from other sources, leaves the question open whether the available
78 evidence supports the conclusion that AMOC is currently undergoing an unprecedented
79 shift/weakening.

80 Key information and rationale about the records included are lacking in Caesar et al.¹.
81 For example, the Rahmstorf et al.⁴ AMOC reconstruction used by Caesar et al.¹ is based on the
82 subpolar North Atlantic temperature minus the Northern Hemisphere mean temperature, each
83 constructed from tree ring and ice core records, and a scaling coefficient derived from one
84 climate model. These data are land-based estimates influenced by atmospheric conditions, not

necessarily robust indicators of marine temperatures, and the resulting index is strongly impacted by the global warming signal⁵. Furthermore, subpolar gyre sea surface temperatures (SSTs) are an unreliable indicator of AMOC variability^{5,6} because SST can have multiple drivers and the spatial AMOC/SST fingerprints used for such reconstructions are temporally non-stationary^{2,5}. Variables related to marine biological processes used as evidence by Caesar et al.¹ are potentially problematic as they are not directly responding to the AMOC and their signal may be compromised by other non-physical factors. For instance, the Sherwood et al.⁷ study provides nitrogen isotopic evidence of a shift in nutrient dynamics since the 19th century in the northwestern Atlantic which they attribute to local changes in water masses, and others⁴ have linked to AMOC. The interpretation of this proxy is predicated on stable nitrogen utilization and nitrogen isotope signatures in the system despite massive anthropogenic perturbation of the global N cycle over the study period⁸. Additional evidence used to infer an AMOC slowdown by Caesar et al.¹ come from sortable silt records off Cape Hatteras⁹, which are arguably one of the most direct proxies available for near-bottom water current speed¹⁰. However, this proxy assumes that the position of the bottom current is stationary through time and that these deep flow changes are representative of AMOC strength. Similar methods have been used to examine the other parts of the deep AMOC limb, including the Nordic Overflows with results that are not consistent with conclusions reached by Caesar et al.¹ (for example, see^{11, 12, 13}), yet these records were not considered.

Finally, the proxy data presented by Caesar et al.¹ need to be reconciled with observations of AMOC and AMOC-related variables in the 20th and 21st centuries. Caesar et al.¹ plot a trend derived from Smeed et al.¹⁴ to support their supposition that AMOC has significantly decreased in recent decades. However, the decreasing trend measured in RAPID data between 2004 and 2012 is really more of a stepwise shift¹⁴ and is likely a part of decadal-scale variability with increases in AMOC from 1960 to the early 2000s^{15, 16}. To date, the RAPID array observations are too short to resolve multidecadal and longer-scale variability. Some

indirect or partial AMOC measures over the instrumental era permit investigation into decadal-
to-multidecadal variability and suggest a modest decline in transport¹⁷, others show no trend¹⁸,
¹⁹, and one record²⁰ shows a recent strengthening of the AMOC at subpolar latitudes. While
diverse regional responses are plausible amidst a large-scale AMOC decline, work remains to
understand the origin of such discrepancies.

These apparently contradictory results may be reconciled with more information
regarding the spatial and temporal scales of variability involved in each dataset as well as the
sensitivity and fidelity of the proxies to record aspects of AMOC during a large global climate
perturbation. Real and interesting subtleties and discrepancies in the data still exist, and any
impression that the historical AMOC evolution is confidently known from a subset of the
available data is misleading until the conflicts are resolved. Instead, highlighting apparent
contradictions will help us with the work of reconciling the data and answering the important
question of whether the AMOC and/or its components have indeed slowed down in recent
centuries.




Figure 1. Available well-dated northern North Atlantic paleoceanographic records include proxies for temperature, salinity, sea ice, and ocean circulation. A full list is in Supplementary Information Table 1. Surface (circles) and deep ocean records (squares) screened by Moffa-Sanchez et al.² (white) are compared with the subset of data (red) used by Caesar et al.¹ The red diamonds are only presented in Caesar et al.¹ and include: biological productivity, nutrient records, and intermediate water temperatures. Multiple cores/archives in the same location are offset for visibility. Source locations, original studies, and figure-making software credits are in Supplementary Information Table 1.

Author contributions: K.H.K, A.D.W, P.M-S., and D.J.R. drafted the manuscript. All authors contributed to discussions in the conception of the work, writing, and editing the manuscript.

Acknowledgements: K.H.K. acknowledges funding from NOAA grant NA20OAR4310481. D.E.A and B.L.O-B. acknowledge support from the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under cooperative agreement no. 1852977. NW acknowledges support from the NOAA Climate and Global Change

Postdoctoral Fellowship. MFJ acknowledges support from NSF award OCE-1846821. UMCES contribution 6062.

References

1. Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N. & Rahmstorf, S. Current Atlantic Meridional Overturning Circulation weakest in last millennium. *Nature Geoscience* **14**, 118–120 (2021).
2. Moffa-Sánchez, P. *et al.* Variability in the Northern North Atlantic and Arctic Oceans Across the Last Two Millennia: A Review. *Paleoceanography and Paleoclimatology* **34**, 1399–1436 (2019).
3. Gu, S., Liu, Z. & Wu, L. Time Scale Dependence of the Meridional Coherence of the Atlantic Meridional Overturning Circulation. *Journal of Geophysical Research: Oceans* **125**, e2019JC015838 (2020).
4. Rahmstorf, S. *et al.* Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation. *Nature Clim. Change* **5**, 475–480 (2015).
5. Little, C. M., Zhao, M. & Buckley, M. W. Do Surface Temperature Indices Reflect Centennial-Timescale Trends in Atlantic Meridional Overturning Circulation Strength? *Geophysical Research Letters* **47**, e2020GL090888 (2020).
6. Keil, P. *et al.* Multiple drivers of the North Atlantic warming hole. *Nature Climate Change* **10**, 667–671 (2020).
7. Sherwood, O. A., Lehmann, M. F., Schubert, C. J., Scott, D. B. & McCarthy, M. D. Nutrient regime shift in the western North Atlantic indicated by compound-specific $\delta^{15}\text{N}$ of deep-sea gorgonian corals. *Proc Natl Acad Sci USA* **108**, 1011 (2011).
8. Gruber, N. & Galloway, J. N. An Earth-system perspective of the global nitrogen cycle. *Nature* **451**, 293–296 (2008).
9. Thornalley, D. J. R. *et al.* Anomalously weak Labrador Sea convection and Atlantic overturning during the past 150 years. *Nature* **556**, 227–230 (2018).
10. McCave, I. N., Thornalley, D. J. R. & Hall, I. R. Relation of sortable silt grain-size to deep-sea current speeds: Calibration of the ‘Mud Current Meter’. *Deep Sea Research Part I: Oceanographic Research Papers* **127**, 1–12 (2017).
11. Moffa-Sánchez, P., Hall, I. R., Thornalley, D. J. R., Barker, S. & Stewart, C. Changes in the strength of the Nordic Seas Overflows over the past 3000 years. *Quaternary Science Reviews* **123**, 134–143 (2015).
12. Mjell, T. L., Ninnemann, U. S., Kleiven, H. F. & Hall, I. R. Multidecadal changes in Iceland Scotland Overflow Water vigor over the last 600 years and its relationship to climate. *Geophysical Research Letters* **43**, 2111–2117 (2016).
13. Moffa-Sánchez, P. & Hall, I. R. North Atlantic variability and its links to European climate over the last 3000 years. *Nature Communications* **8**, 1726 (2017).
14. Smeed, D. A. *et al.* The North Atlantic Ocean Is in a State of Reduced Overturning. *Geophysical Research Letters* **45**, 1527–1533 (2018).
15. Karspeck, A. R. *et al.* Comparison of the Atlantic meridional overturning circulation between 1960 and 2007 in six ocean reanalysis products. *Climate Dynamics* 1–26 (2017).
16. Willis, J. K. Can in situ floats and satellite altimeters detect long-term changes in Atlantic Ocean overturning? *Geophysical Research Letters* **37**, (2010).

- 187 17. Piecuch, C. G. Likely weakening of the Florida Current during the past century revealed
188 by sea-level observations. *Nature Communications* **11**, 3973 (2020).
- 189 18. Yashayaev, I. & Loder, J. W. Recurrent replenishment of Labrador Sea Water and
190 associated decadal-scale variability. *Journal of Geophysical Research: Oceans* **121**, 8095–8114
191 (2016).
- 192 19. Rossby, T., Chafik, L. & Houpert, L. What can Hydrography Tell Us About the Strength of
193 the Nordic Seas MOC Over the Last 70 to 100 Years? *Geophysical Research Letters* **47**,
194 e2020GL087456 (2020).
- 195 20. Desbruyères, D. G., Mercier, H., Maze, G. & Daniault, N. Surface predictor of overturning
196 circulation and heat content change in the subpolar North Atlantic. *Ocean Science* **15**, 809–817
197 (2019).
- 198