

Yeasts from temperate forests

Authors:

Simone Mozzachiodi^{1*}, Feng-Yan Bai², Petr Baldrian³, Graham Bell⁴, Kyria Boundy-Mills⁵, Pietro Buzzini⁶, Neža Čadež⁷, Francisco Cubillos Riffo^{8,9}, Sofia Dashko¹⁰, Roumen Dimitrov¹¹, Kaitlin J. Fisher¹², Brian R. Gibson¹³, Dilnora Gouliamova¹¹, Duncan Greig¹⁴, Lina Heistinger¹⁵, Chris Todd Hittinger¹², Marina Jecmenica¹⁶, Vassiliki Koufopanou¹⁷, Christian R. Landry^{18,19,20,21,22}, Tereza Mašínová³, Elena S. Naumova²³, Dana Opulente²⁴, Jacqueline J. Peña²⁵, Uroš Petrović^{26,27}, Isheng Jason Tsai²⁸, Benedetta Turchetti⁶, Pablo Villarreal^{8,9}, Andrey Yurkov²⁹, Gianni Liti^{1*} and Primrose Boynton^{30*}

Affiliations:

¹Université Côte d'Azur, CNRS, INSERM, IRCCyN, Nice, France; ²State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China; ³Laboratory of Environmental Microbiology, Institute of Microbiology of the Czech Academy of Sciences, Videnska 1083, 14220 Praha 4, Czech Republic; ⁴Biology Department and Redpath Museum, McGill University, Québec, Canada; ⁵Department of Food Science and Technology, University of California Davis, Davis, CA 95616 USA; ⁶Department of Agriculture, Food and Environmental Sciences & Industrial Yeasts Collection DBVPG, University of Perugia, Italy; ⁷University of Ljubljana, Biotechnical Faculty, Food Science and Technology Department, Jamnikarjeva 101, 1000 Ljubljana, Slovenia; ⁸Universidad de Santiago de Chile, Facultad de Química y Biología, Departamento de Biología, Santiago, Chile; ⁹Millennium Institute for Integrative Biology (iBio), Santiago, Chile; ¹⁰DSM Food Specialties, Center for Food Innovation, Alexander Fleminglaan 1, 2613 AX, Delft, The Netherlands; ¹¹Institute of Microbiology, Bulgarian Academy of Sciences, G. Bonchev, 26, Sofia, Bulgaria; ¹²Laboratory of Genetics, Wisconsin Energy Institute, DOE Great Lakes Bioenergy Research Center, Center for Genomic Science Innovation, J. F. Crow Institute for the Study of Evolution, University of Wisconsin-Madison, Madison, WI 53726-4084, USA; ¹³Technische Universität Berlin, Institute of Food Technology and Food Chemistry, Chair of Brewing and Beverage Technology, Ackerstr. 76, 13355 Berlin, Germany; ¹⁴Centre for Life's Origins and Evolution, University College London, Gower Street, London WC1N 6BT, UK; ¹⁵ETH Zurich, Department of Biology, Institute of Biochemistry, Switzerland; ¹⁶University of Natural Resources and Life Sciences (BOKU) Vienna; ¹⁷Department of Life Sciences, Imperial College London, Silwood Park, Ascot, Berks, SL5 7PY, UK; ¹⁸Département de Biochimie, de Microbiologie et de Bio-informatique, Faculté des Sciences et de Génie, Université Laval, G1V 0A6, Canada; ¹⁹Institut de Biologie Intégrative et des Systèmes, Université Laval, G1V 0A6, Canada; ²⁰PROTEO, Le regroupement québécois de recherche sur la fonction, l'ingénierie et les applications des protéines, Université Laval, G1V 0A6, Canada; ²¹Centre de Recherche sur les Données Massives, Université Laval, G1V 0A6, Canada; ²²Département de Biologie, Faculté des Sciences et de Génie, Université Laval, G1V 0A6, Canada; ²³State Research Institute of Genetics and Selection of Industrial Microorganisms of National Research Centre "Kurchatov Institute", Moscow, Russia; ²⁴Department of Biology, Villanova University, Villanova, Pennsylvania, USA; ²⁵Department of Plant Biology, University of Georgia, Athens, Georgia 30602; ²⁶University of Ljubljana, Biotechnical Faculty, Department of Biology, Večna pot 111, SI-1000 Ljubljana, Slovenia; ²⁷Jožef Stefan Institute, Department of Molecular and Biomedical Sciences, Jamova cesta 39, SI-1000 Ljubljana, Slovenia; ²⁸Biodiversity Research Center, Academia Sinica, Taiwan; ²⁹Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Inhoffenstrasse 7B, 38124, Braunschweig, Germany; ³⁰Biology department, Wheaton College, 26 E. Main St., Norton, MA, USA

*Correspondence should be addressed to PBo (boynton_primrose@wheatoncollege.edu), GL (gianni.liti@unice.fr), SM (simone.mozzachiodi@gmail.com)

Running title: Temperate forest yeasts

Keywords: isolation, biodiversity, *Saccharomyces*, *Lachancea*, *Komagataella*, *Cryptococcus*

Take-aways:

- Temperate forests are a widespread global biome, and host diverse yeasts.
- Well-studied model yeasts from other fields are in temperate forests.
- Isolation strategies depend on target yeasts and researcher questions.
- Temperate forest yeasts are potentially exciting biotechnological resources.
- Advancing technology is revealing yeast growth and life cycle secrets.

Abstract

Yeast are ubiquitous in temperate forests. While this broad habitat is well-defined, the yeasts inhabiting it and their life cycles, niches, and contributions to ecosystem functioning are less understood. Yeasts are present on nearly all sampled substrates in temperate forests worldwide. They associate with soils, macroorganisms, and other habitats, and no doubt contribute to broader ecosystem-wide processes. Researchers have gathered information leading to hypotheses about yeasts' niches and their life cycles based on physiological observations in the laboratory as well as genomic analyses, but the challenge remains to test these hypotheses in the forests themselves. Here we summarize the habitat and global patterns of yeast diversity, give some information on a handful of well-studied temperate forest yeast genera, discuss the various strategies to isolate forest yeasts, and explain temperate forest yeasts' contributions to biotechnology. We close with a summary of the many future directions and outstanding questions facing researchers in temperate forest yeast ecology. Yeasts present an exciting opportunity to better understand the hidden world of microbial ecology in this threatened and global habitat.

1. Introduction

Yeast, defined here as predominantly unicellular fungi that often reproduce asexually by budding or fission (Kurtzman et al., 2011; Boekhout et al., 2021a), and dimorphic fungi that have filamentous and unicellular phases, are ubiquitous throughout Earth's biomes (Péter et al., 2017; Boekhout et al., 2021a); however, they are often overlooked in favor of sporocarp-forming fungi. This is especially true in temperate forests, where researchers have paid considerable attention to decomposer and mycorrhizal fungi, many of which produce large macroscopic fruiting bodies (Albarracín et al., 2013; Tedersoo et al., 2010; Boddy & Hiscox 2016). Nevertheless, yeasts have colonized a diverse variety of available forest substrates, including soils and the microbiomes of plants, animals, and other fungi (Péter et al., 2017; Yurkov, 2018; Stefanini, 2018; Starmer & Lachance, 2011; Yurkov et al., 2012; Boekhout et al., 2021a). In addition to being ubiquitous, yeasts are involved in multiple ecosystem processes, acting as free-living saprobes, mutualists, parasites, and other symbionts (Starmer & Lachance, 2011). They likely have profound impacts on ecosystem functioning in temperate forests. However, most of mycologists' and ecologists' current knowledge about yeasts in temperate forests comes from *ex-situ* studies of model organisms. In response, research on natural history and ecology throughout this fascinating fungal group in the temperate forest environment is currently expanding.

Because yeasts are so diverse, they have the potential to have diverse impacts on temperate forest ecosystems. Genomic surveys have associated yeasts with higher stress tolerance, simpler carbon utilisation, and higher nitrogen uptake compared to filamentous fungi

(Treseder & Lennon 2015; Romero-Olivares et al., 2021), although these genomic surveys are biased toward model organisms. In temperate forests, yeast communities respond to changes in forest properties including tree age, density, management, and wood deposition, and these changes may feed back on forest ecosystem function (Birkhofer et al., 2012; Yurkov et al., 2012; Yurkov et al., 2012b; Yurkov et al., 2016). While data on yeasts' impacts on temperate forest ecosystem functioning is still in its infancy, temperate forest yeasts may impart ecosystem resilience in the face of climate change. For example, researchers speculate that stress-tolerant yeasts will respond positively to stressful environmental changes relative to filamentous fungi, leading to decreased filamentous fungal decomposition and increased carbon storage in forest environments (Treseder & Lennon 2015).

In this review, we define temperate forests broadly as forests in the temperate zone, found between boreal and tropical forests. Overall, forests cover nearly a third of global land area, and 16% of forested area is occupied by temperate forests (10 million km²) (FAO & UNEP 2020). Transitions between temperate and boreal, or temperate and tropical, are not sharp. Temperate forest vegetation is diverse, including coniferous, deciduous, and mixed forests in Köppen classification climate groups C and D (Beck et al., 2018). This vegetation ranges from nearly boreal (continental, with pronounced cold seasons) to subtropical with mild climates (only rarely with temperatures below 0 °C) and can include climates with a pronounced dry season (Mediterranean climate). In several geographic areas, temperate forests border large grasslands, including the North American great plains and tallgrass prairie, Patagonian grasslands, Eurasian steppes, and Humid Pampas (Myster (ed.) 2012). The majority of temperate forests are located in the northern hemisphere (Beck et al., 2018).

Most forests in the temperate zone are included in two terrestrial biomes (Olson & Dinerstein 2002), the temperate broadleaf and mixed forests biome and the temperate coniferous forest biome. The former includes most broadleaf and mixed forests in both hemispheres, and the latter includes Alpine, coastal Scandinavian, Asian (Himalayan and Tian Shan), Mediterranean, and coastal North American Pacific forests. Dominant tree families include Fagaceae (e.g. *Betula*, *Fagus*, *Nothofagus*, *Quercus*), Pinaceae (e.g. *Abies*, *Picea*, *Pinus*), Cupressaceae (e.g. *Fitzroya*, *Juniperus*, *Sequoia*), and, exclusively in the southern hemisphere, Araucariaceae (e.g. *Agathis*, *Araucaria*), Podocarpaceae (e.g. *Dacrydium*, *Podocarpus*, *Saxeogathea*), and tree ferns (*Cyatheales*) (Reyna et al., 2018; Mao et al., 2012; Beard., 1990; Brock et al., 2016). Temperate forests also include unique and interesting plant communities such as dry Mediterranean-type forests (e.g. Oak and Eucalyptus forests), humid temperate rainforests (e.g. Valdivian forest in South America) and relict laurel forests (e.g. Macaronesian Laurisilva) (Cowling et al., 1996; Claudino-Sales 2019; Noh et al., 2019).

Human activities have had a large impact on forests, mainly through continuous deforestation (e.g. for timber and energy) and forest conversion into agricultural lands (FAO &

UNEP 2020). Additionally, global warming and fires are devastating for forest ecosystems, with fire destroying 1% of temperate forest area in 2015 (FAO & UNEP 2020). Recently recognized biodiversity hotspots in temperate forests highlight the unique character of these ecosystems and need for forest conservation (Myers et al., 2000). We expect high numbers of endemic plants and animals to similarly harbor high numbers of endemic microorganisms, including yeasts.

In this review, we address the current knowledge of yeast diversity in temperate forests, their ecology, and their biotechnological applications. We discuss biases in the literature and how an unbiased picture of yeasts' roles and biodiversity will emerge from future studies.

2. Yeast biodiversity and abundance in temperate forests

2.1 Yeast distributions from environmental sequencing

Based on environmental sequencing data, yeasts are widespread globally in temperate forest soil and leaf litter (Figure 1), but their frequencies tend to be low. However, there are potential biases and errors in environmental sequencing due to incomplete databases, detection of dead or dormant cells, technical biases, and different detection limits between sequence- and culture-based studies; conclusions from environmental sequencing should therefore be made cautiously and confirmed using culture-based studies whenever possible (Lücking et al., 2020). A comprehensive screening of temperate forest soil and leaf litter internal transcribed spacer (ITS) barcode sequences deposited in the GlobalFungi database (Vetrovsky et al., 2020) and analysed for this review revealed 3,783,412 total yeast sequences belonging to 859 yeast species hypotheses (SHs); yeasts here were "yeast" or "facultative yeast" as in Polme et al. (2020), with some manual curation (Table S1); and only samples with more than 5000 fungal sequences were considered. Fungal ITS-targeting primers perform well with all important fungal lineages containing yeasts as well as individual yeasts in a mock community (Větrovský et al., 2016; 2019) and no significant PCR bias is observed when comparing the yeast share in metagenomic data (without PCR) and ITS amplicon data (Tláskal et al., 2021); nor does there seem to be important DNA extraction bias against selected yeast taxa (Větrovský et al., 2016). This screening showed that the median yeast abundance in soil was significantly higher than in litter (Wilcoxon Rank-Sum test, $W = 722187$, $p = 2.9 \times 10^{-15}$). Yeasts were generally not common: only 12% of soil samples and 3% of litter samples included more than 8% yeast ITS sequences (Figure S1). Richness ranged from 1-31 in soil and 1-35 in litter (Table S1). Because of geographical sampling biases (Figure 1) and the overall low frequencies of yeasts in most individual samples, we could not conduct analyses correlating yeast incidence with season, climate, or host plant. However, we

anecdotally noticed a weak amount of seasonality in yeast communities; this seasonality remains to be investigated with targeted studies.

The global temperate forest yeast community emerging from barcode sequencing includes some widespread and dominant taxa. Specifically, the most frequently detected classes containing yeasts were *Tremellomycetes* (87.7% of yeast sequences), *Dothiomycetes* (4.6%) and *Eurotiomycetes* (3.1%). The most prevalent yeast species, with prevalence meaning the percent of temperate forest and soil samples on which a species was found, were *Saitozyma podzolica*, *Solicoccozyma terricola*, *Apotrichum xylopini/porosum*, *Tausonia pullulans*, and *Cutaneotrichosporon moniliforme* (Table S2). These five yeast species are widespread and cosmopolitan, and have broad climatic niches (Větrovský et al., 2019).

2.2 Yeasts in culture collections

While culture-independent methods to profile microbial species and their functions are gaining prominence, the vast majority of the current knowledge about the biodiversity and distributions of yeasts in temperate forests has been garnered through direct isolation and subsequent study of yeast cultures. Currently, thousands of yeast strains isolated from temperate forests are preserved in both public and private microbial culture collections. Researchers have used collections to demonstrate that yeast communities in natural unmanaged forests can be more species-rich than in managed ones (Yurkov et al., 2012; Yurkov et al., 2016); below-ground communities reflect properties of plant cover, such as canopy density and diversity; and seasonality impacts total quantity and species richness of yeasts (Yurkov et al. 2016). However, these conclusions rely on targeted sampling, which is currently limited by locality, season and substrates.

Many culture collections throughout the globe contain lots of yeast isolates from temperate forests. In many cases, researchers can search culture collection catalogs for keywords or substrate, such as “soil”, “forest”, or “bark” to help identify temperate forest yeasts. The Global Catalog of Microorganisms (<https://gcm.wdcm.org>), organized with the World Data Centre for Microorganisms (WDCM), has combined the catalogs of over 140 public microbe collections to streamline the process of locating a specific yeast species or yeasts isolated from a certain location or habitat (Wu et al., 2013). In addition, the list of Culture Collections Information Worldwide (CCINFO), created by the WDCM and the World Federation for Culture Collections (WFCC), provides an overview of more than 800 repositories worldwide. For more information about the history and ongoing use of culture collections, we recommend the review of Boundy-Mills and coauthors (2016). Not all culture collections index strain habitat information in a searchable way, and a reliable summary of yeast cultures from temperate forests is a current challenge.

3. Model yeast genera in temperate forests

Beginning with the genus *Saccharomyces* in the days of Louis Pasteur (Pasteur 1876), several groups of yeasts have been used as model organisms for biotechnology, cell biology, medicine, genetics, and genomics. While the impacts of these yeast groups extend well beyond yeast biology, many of them have also been isolated from temperate forest ecosystems. These include the genera *Saccharomyces*, *Komagataella*, and *Lachancea*, and yeasts that until recently were classified in the genus *Cryptococcus*, and comprise a polyphyletic group. These yeast groups have the potential to be valuable models for temperate forest ecology, and some—especially the genus *Saccharomyces*—are already well-developed ecological and evolutionary models.

3.1 The genus *Saccharomyces*

The genus *Saccharomyces* is the most frequently studied model yeast group in temperate forests. There are eight known species: *S. cerevisiae*, *S. paradoxus*, *S. mikatae*, *S. jurei*, *S. kudriavzevii*, *S. arboricola*, *S. eubayanus* and *S. uvarum* (Batschinskaya, 1914; Libkind et al., 2011; Meyen, 1839; Naseeb et al., 2017; Naumov et al., 2000; Ono et al., 2020; Pulvirenti et al., 2000; Wang et al., 2012). *S. cerevisiae* is a famous industrial and laboratory model organism, and hybrids between *Saccharomyces* species are also industrially important (Gallone et al., 2019; Langdon et al., 2019; Pérez-Torrado et al., 2018). All *Saccharomyces* species, including *S. cerevisiae*, have been isolated from temperate forest substrates, including tree bark, leaf litter, soil, fruits, insects, and leaf surfaces (Phaff et al., 1956; Libkind et al., 2011; Sampaio & Gonçalves, 2008, 2017; Naseeb et al., 2017; Naumov et al., 2000; Wang & Bai 2008; Glushakova et al., 2007). The genus includes several rare naturally-occurring hybrids and incipient hybrid species (Barbosa et al., 2016; Leducq et al., 2016; Eberlein et al., 2019) in addition to the eight *Saccharomyces* species, although most hybrids are from domesticated environments (Liti et al., 2005; Gibson et al., 2017). Interspecies *Saccharomyces* hybrids are rare outside of domesticated environments, although genomic introgressions (e.g., *S. paradoxus* open reading frames in *S. cerevisiae* genomes) have been found in temperate forest yeasts, and are evidence of past hybridization (Almeida et al., 2014; Peter et al., 2018).

Of the eight known *Saccharomyces* species, *S. paradoxus* has been most frequently isolated from temperate forests, and it is the best-studied *Saccharomyces* species outside of domesticated environments. The species was first described in 1914 based on strains isolated from oak and elm exudates in Russia (Batschinskaya, 1914). Its native range includes much of the northern hemisphere with average summer temperatures below 31 °C (Robinson et al., 2016). Its infrequent isolation in New Zealand is consistent with a recent migration of isolates with European ancestry (Zhang et al., 2010). Most isolates have been obtained from bark, sap

exudates, leaf litter, and soil; host trees are often but not always oaks (Phaff et al., 1956; Kodama, 1974; Banno & Mikata, 1981; Naumov, 1996; Naumov et al., 1998; Kowallik & Greig, 2016; Kowallik et al., 2015; Glushakova et al., 2011; Sampaio & Goncalves, 2008, 2017; Sniegowski et al., 2002). *S. paradoxus* is generally infrequent in forest habitats, but its frequency can spike seasonally or after rain events to as many as 100% of isolated colonies (Glushakova et al., 2007; Anderson et al., 2018; Charron et al., 2014); these spikes are less evident when sampling is less frequent, with a pattern of approximately even frequency year-round in at least one study (Kowallik & Greig, 2016).

Similar surveys also identified other northern hemisphere *Saccharomyces* species (Sampaio & Goncalves 2008). For instance, recent studies suggest that eastern Asia harbors several *Saccharomyces* species characterized by high genetic diversity (Duan et al., 2018; Han et al., 2021; Lee et al., 2021; Wang et al., 2012; Bing et al., 2014; Bendixen et al., 2021; Peter et al., 2018), including temperate forest *S. cerevisiae* populations that are strongly diverged from domesticated populations (Duan et al., 2018; Wang et al., 2012). Some examples of successful sampling efforts that uncovered multiple species include a study in which *S. kudriavzevii*, *S. uvarum*, and *S. cerevisiae* were isolated from oak bark in Portugal (Sampaio & Gonçalves, 2008, 2017), and another in which *S. arboricola*, *S. kudriavzevii*, and *S. cerevisiae* were isolated from mushrooms, litter and leaves in Taiwan (Naumov et al., 2013). In contrast, *S. jurei*, *S. arboricola*, and *S. mikatae* geographic ranges appear more limited. For example, the strains of *S. arboricola* were only isolated from trees of the Fagacee family in China in 2008 (Wang et al., 2008), and *S. mikatae* has only been isolated so far in soil and decayed leaves from Japan and China (Naumov et al., 2000, Sampaio & Gonçalves 2017). Along the same line, *S. jurei* has been recently isolated twice in different locations in Europe (Naseeb et al., 2017, Hutzler et al., 2021) and a metagenomic study also detected *S. jurei* in temperate forests in the Italian Alps (Alsammar et al., 2019), but it is not yet known if different populations exist outside Europe. *S. jurei* and *S. mikatae* share some genomic rearrangements, suggesting a shared evolutionary history (Naseeb et al., 2018).

Northern hemisphere *Saccharomyces* species isolated from temperate forests show generally low genetic admixture. Forest *S. paradoxus* populations have high clonality and low heterozygosity, and they experience infrequent sex with even less frequent outcrossing (3×10^{-6} to 10^{-5} per cell division) (Tsai et al., 2008; Johnson et al., 2004; Liti et al., 2006; Liti et al., 2009; Bergström et al., 2014). Researchers sampling *S. paradoxus* in temperate forests frequently find the same clonal genotype, with the probability of uncovering the same clone decreasing with distance (Koufopanou et al., 2006). The same clone can also persist over consecutive years (Koufopanou et al., 2006) (Xia et al., 2017; Anderson et al., 2018). Similarly, *S. cerevisiae* outcrossing rates are also estimated to be low at 9×10^{-5} to 2×10^{-5} per cell division (Jensen et al., 2001; Ruderfer et al., 2006). However, outcrossing rates of *S.*

cerevisiae increased at least ten-fold when yeast spores were digested by an insect dispersal vector such as *Drosophila* (Reuter et al., 2007); insect digestive systems may promote outcrossing and hybridization by bringing ascospores together (Stefanini et al., 2016).

Surveys in the southern hemisphere have further expanded our understanding of *Saccharomyces* diversity. During the past decade, *S. uvarum* and *S. eubayanus* have emerged as the most frequently isolated *Saccharomyces* species in bark, fruits and soil in the southern hemisphere, particularly in Australasian and South American temperate forests (Figure 1) (Almeida et al., 2014; Eizaguirre et al., 2018; Langdon et al., 2020; Libkind et al., 2011; Nespolo et al., 2020; Rodriguez et al., 2014). Patagonia, in particular, harbors high *S. uvarum* and *S. eubayanus* diversity (Nespolo et al., 2020; Langdon et al., 2020; Peris et al., 2016). *S. eubayanus* is widely distributed in South Patagonia and associates with cold environments, and it has also been isolated in Asia (Bing et al., 2014; Peris et al., 2016), Oceania, (Gayevskiy & Goddard, 2016) and North America (Peris et al., 2014). Interestingly, *S. eubayanus*, a parent of the hybrid lager species *S. pastorianus*, has never been isolated in Europe, challenging our understanding of the origin of lager beer yeast hybrids. While *S. eubayanus* may indeed be rare in Europe, the lack of *S. eubayanus* in Europe could also be due to insufficient sampling or absence of the species due to deforestation. Recent evidence collected from a targeted metagenomics study of *Saccharomyces* diversity in the Italian Alps suggests that *S. eubayanus* may be present in Europe, although future culture-based work is needed to confirm this (Alsammar et al., 2019).

In addition to members of the genus *Saccharomyces* isolates from throughout the globe, genome sequencing of isolates from hybrid zones have revealed admixture among partially reproductively isolated *S. paradoxus* populations (Leducq et al., 2016; Xia et al., 2017; Eberlein et al., 2019). For instance, studies on genetically diverged and partially reproductively isolated North American *S. paradoxus* lineages revealed the presence of hybrid, phenotypically diverged, and reproductively isolated lineages, which were found at intersections of the parental geographic distributions. These incipient hybrid species evolved following post glacial admixture between parental lineages (Leducq et al., 2016; Xia et al., 2017; Eberlein et al., 2019).

The ecological determinants of different *Saccharomyces* lineages' ranges are still unclear. Temperature has been suggested as an important abiotic niche determinant because experimental work found different temperature optima among *Saccharomyces* lineages (Sweeney et al., 2004; Salvado et al., 2011). In the northern hemisphere, it has been hypothesized that *S. cerevisiae* is better adapted to warmer temperatures than *S. paradoxus* (Charron et al., 2014; Robinson et al., 2016). Climatic modeling of *S. eubayanus* has also recapitulated much of its known distribution (Langdon et al., 2020). In some locations, multiple *Saccharomyces* species with different temperature optima coexist at the same site. For

example, cold-tolerant *S. kudriavzevii* and *S. uvarum* are readily isolated from the same samples that yielded warm-tolerant *S. cerevisiae* or *S. paradoxus*, when these same samples were incubated at two different temperatures (Sampaio & Goncalves 2008; Sweeney et al., 2004). Diversity in genes related to glycerol accumulation and acetaldehyde production has been implicated in temperature tolerance diversity and may therefore be important for temperature niche partitioning (Paget et al., 2014).

While temperature may indeed determine the ranges of *S. paradoxus* and *S. cerevisiae* in the northern hemisphere, this does not directly translate to a correlation between latitude and sampling success. We surveyed the literature for studies that isolated either *S. paradoxus* or *S. cerevisiae* from temperate forests and clearly stated the number of positive and negative sampling attempts (Table S3). Across these studies, rates of success in isolating either species does not correlate with latitude (Figure 2). However, *Saccharomyces cerevisiae* has been less frequently isolated at higher latitudes than *S. paradoxus* when isolation frequency is compared within the same study (Figure S2); more data are needed to further evaluate this observation. While several studies have reported isolation of one, but not both species, others have indeed isolated both *S. paradoxus* and *S. cerevisiae* from the same location (Boynton et al., 2019; Dashko et al., 2016; Kowallik et al. 2016; Sniegowski et al., 2002). This result suggests that additional factors influence these species' ranges beyond temperature.

A similar pattern of partial coexistence of *Saccharomyces* species is mediated by host tree identity in the southern hemisphere. For instance, South American *S. uvarum* and *S. eubayanus* partly share their habitat, but while *S. uvarum* is mainly found on *Nothofagus dombeyi*, *S. eubayanus* is mainly found on *Nothofagus pumilio* (Eizaguirre et al., 2018; Langdon et al., 2020; Libkind et al., 2011; Mardones et al., 2020). In spite of this apparent specialization, both species can inhabit the same tree, and elsewhere in Tibet they coexist in similar niches (Bing et al., 2014). Because dominant forest trees and species distributions change with latitude, there is a correlation between species isolated and latitude in the southern hemisphere (Figure 2, Table S3). *S. uvarum* is more frequently found at lower latitudes with warmer climates than *S. eubayanus* (Figure S2). In contrast, *S. eubayanus* is frequently found at higher latitudes and altitudes, near the treeline in extremely cold environments (Figure 2). The data used for this analysis are derived from a single study in which all samples were collected and processed using the same methods, suggesting that variability among isolation protocols across labs could add a confounding effect when estimating species ranges.

3.2 The genus *Cryptococcus* and former *Cryptococcus*

Taxonomic challenges over the past decades have drastically changed our understanding of the prevalence and frequency of the genus *Cryptococcus* in temperate

forests. Historically, sexual and asexual fungal forms were given separate names and the genus epithet “*Cryptococcus*” described asexual (i.e., mitotically reproducing) species of basidiomycetous yeasts (Vuillemin 1901; Liu et al., 2015). This system of dual nomenclature for fungi with asexual and sexual forms was abolished in 2012 (Lücking et al., 2021). At about the same time, modern DNA sequencing technology regrouped the previous polyphyletic genus *Cryptococcus* into a monophyletic genus in the former sexual *Filobasidiella* clade, plus several other basidiomycete genera. The genus *Cryptococcus* now comprises prominent human pathogens (the *C. neoformans* species complex) and a few closely related non-pathogenic yeasts with a possible link to mycoparasites (Liu et al., 2015; Passer et al., 2019). All other former *Cryptococcus* species were transferred to other genera, emended, resurrected, or newly erected (Liu et al., 2015) (Table S4). As a consequence, historic reports of *Cryptococcus* from natural substrates can be dubious because they may be reports of species no longer circumscribed in the genus *Cryptococcus* (Lücking et al., 2021).

Members of the currently delineated genus *Cryptococcus* inhabit temperate forests worldwide (Cogliati et al., 2016; Lin et al., 2021; Chowdhary et al., 2012; Kidd et al., 2007). Non-pathogenic *Cryptococcus* species are rare and only known from a very few isolates of plant-insect origin, including *C. amylorentus*, *C. wingfieldii*, and *C. floricola* (Passer et al., 2019). The limited number of reports of environmental *Cryptococcus* are restricted to selective cultivation surveys of pathogenic species and single-strain isolations of potential new species (Mittelbach et al., 2015; Yurkov et al., 2016; Passer et al., 2019). The species *Cryptococcus neoformans* and *Cryptococcus gattii* are best known as human pathogens, but they have also been isolated from soils, water, air and tree samples from temperate forests (Chowdhary et al., 2012; Kidd et al., 2007; MacDougall et al., 2007). The exchange between host niches in the forest and human hosts contributes to disease incidence (Kidd et al., 2004).

Since the taxonomic revision of the genus (Liu et al., 2015), *Cryptococcus* has not been present in the list of prevalent genera found in forests. In contrast, many yeast genera previously assigned to the genus *Cryptococcus* have been found in temperate forests, and these genera are composed of several species. They mostly inhabit soil and plant (including moss) habitats (Table S4), and many have been detected using environmental barcode sequencing, with the genera *Vishniacozyma*, *Filobasidium*, *Piskurozyma*, *Saitozyma* and *Solicoccozyma* among those with the most abundant DNA sequences recovered in environmental samples (Kemler et al., 2017). In fact, several of the species identified as most common in our global analysis of yeast sequence diversity belong to this group (Table S2). We do not know what makes these taxa so common, but speculate that some of them may be generalists, inhabiting many different global environments, or have an as-yet unexplored, but very common, ecological niche in temperate forests. For example, many members of the genera *Filobasidium*, *Papiliotrema*, and *Vishniacozyma* inhabit aboveground substrates

(Yurkov et al., 2015; Kemler et al., 2017). Moreover, *Cryptococcus* and former *Cryptococcus* species (Table S4) use a variety of nutrient acquisition and stress tolerance strategies in temperate forest environments. Many of these yeasts from temperate regions are opportunistic saprobionts, consuming the products of decomposing plant material released by other microorganisms, such as filamentous fungi (Babjeva and Golovleva, 1963; Yurkov, 2017; Mašínová et al., 2018). However, others can use complex compounds directly. For example, forest soil strains of *Solicoccozyma terricola* and *Holtermanniella wattica* (formerly *Cryptococcus terricola* and *C. watticus*) exhibit a wide assimilation spectrum of carbon sources and extracellular enzymes, including growth on oligosaccharides (Mašínová et al., 2018). Similarly, *S. terrea* and some *Apotrichum* species can grow on phenolic compounds (Botha 2006; Yurkov et al., 2017). Many former *Cryptococcus* have physiological features that can increase their chances of survival under stress as other temperate forest yeasts, including drought, nutrient limitation, and radiation in soil and phylloplane environments (Kemler et al., 2017; Yurkov, 2017). For example, members of the genera *Naganishia* and *Solicoccozyma* produce extracellular polysaccharide capsules that promote growth in low nutrient and low water activity environments (Vishniac et al., 1995, 2006; Botha et al., 2011), and some *Naganishia* species produce mycosporines as radiation protection in extreme environments (Nizovoy et al., 2021). This variety of nutrient acquisition and stress tolerance strategies illustrates the diversity and explains the heterogeneity of the former polyphyletic genus *Cryptococcus*.

3.3 The genus *Komagataella*

The seven species in the methanol-assimilating model genus *Komagataella* are all present in temperate forest ecosystems (Naumova et al., 2020). They have been isolated from tree exudates, bark, associated insects, and rotten wood of temperate forest trees including *Quercus* spp., *Populus* spp., *Ulmus* spp., *Castanea* spp. and *Betula* spp. in the northern hemisphere (Lachance et al., 1982; Naumova et al., 2020). Species in this genus have substrate preferences. For example, while most *Komagataella* species have been isolated from broad-leaved trees, *K. kurtzmanii* has been isolated only from fir flux. Additionally, *K. pseudopastoris* has higher sensitivity to the tannic acids in *Quercus* species than *K. pastoris* (Dlauchy et al., 2003; Péter et al., 2019).

Geography contributes to the taxonomic structure of the genus *Komagataella*. The genus itself includes three groups of closely related species: *K. pastoris*/*K. ulmi*, *K. kurtzmanii*/*K. phaffii*, and *K. mondaviourum*/*K. pseudopastoris*/*K. populi* (Figure S3) (Dlauchy et al., 2003; Kurtzman 2005, 2012; Naumov et al., 2013, 2018). As with *Saccharomyces*, *Komagataella* species readily hybridize in the laboratory, but hybrids are post-zygotically reproductively isolated and generate mostly unviable ascospores (0-7%) (Naumov 2015;

Naumov et al., 2016); they may also be pre-zygotically isolated by substrate or geography. We are not aware of naturally-occurring *Komagataella* hybrids in temperate forests, although the shared mating system among *Komagataella* species makes hybrids likely. *K. kurtzmanii*, *K. mondavorum*, *K. phaffii*, *K. populi*, and *K. ulmi* have frequently been isolated from North America, while *K. pastoris* and *K. pseudopastoris* are spread across Europe (Dlauchy et al., 2003; Kurtzman, 2011a, 2012; Miller et al., 1962, Naumov 2013; 2018; Phaff et al., 1972). Large-scale screening of methanol-assimilating ascomycetous yeasts failed to isolate *Komagataella* strains in Thailand and Brazil (Lamtong et al., 2013; Santos et al., 2015), suggesting that the genus might be absent from these areas. However, rare *Komagataella* isolates have been documented in Japan (Phaff et al., 1972; Kodama 1974) and Argentina (Spencer et al., 1995, 1996). Future sampling may uncover more species in this genus, which would likely expand our understanding of their ranges and ecology.

3.4 The genus *Lachancea*

The genus *Lachancea* comprises eleven species from a wide variety of ecological niches (Figure S4). Members of this genus are among the most commonly isolated yeasts worldwide and their relationship with temperate forest environments is understudied. *Lachancea* species share common physiological properties: they ferment glucose, can grow on raffinose, maltose, ethanol, and mannitol, and have similar temperature preferences and fermentation characteristics (Bandara et al., 2009; Naumova et al., 2007; S. Benito, 2018). They have been isolated from a wide variety of ecological niches, including plants (Estevez-Zarzoso et al., 2001; Gonzalez et al., 2007; Romano & Suzzi, 1993a, 1993b), tree bark (Nespolo et al., 2020), tree exudates (Varela et al., 2020), insects (Phaff et al., 1956), soil (Lee et al., 2009; Mesquita et al., 2013; Sylvester et al., 2015), water (Kenkichi. & Tadashi., 1974), and food and beverages (Magalhães et al., 2011; Marsh et al., 2014; Nova et al., 2009; Pereira et al., 2011; Tzanetakis et al., 1998; Wojtakowicz et al., 2001). While the most common species in the genus, *L. thermotolerans* and *L. fermentati*, are often associated with domesticated food and beverage environments (Aponte & Blaiotta, 2016bnt; Bagheri et al., 2016; Baleiras Couto et al., 2005; Clavijo et al., 2010; Cordero-Bueso et al., 2013; Fernández et al., 1999; Hsieh et al., 2012; Senses-Ergul et al., 2006), they are also frequently isolated from natural substrates (Sylvester et al., 2015; Spurley et al., 2021). *L. cidri*, a close relative of *L. fermentati*, has also been recently reported on *Nothofagus* bark in temperate southern Chile (Villarreal et al., 2021). Despite the large number of ecosystems where *Lachancea* is present, ranging from fermented beverages to temperate forests, there is a lack of information on why the different species inhabit such a wide variety of ecological niches (Porter et al., 2019b) and which roles they play in those ecosystems.

4. Practical suggestions for collecting yeasts from forests

Sampling and isolation steps both play a crucial role in yeast discovery and estimation of species richness (Lachance & Starmer 1998; Yurkov & Pozo 2017). When making a sampling and isolation plan, researchers should determine whether they wish to characterize the entire diversity of the environment or focus on a particular group of yeasts. Researchers' goals will ultimately inform their sampling and isolation strategies.

4.1 Collecting substrate samples from temperate forests for yeast isolation

Wild yeasts can be isolated from diverse habitats and ecological niches in temperate forests around the world, including soil, leaf litter, tree bark, rotten fruits, insect bodies, mushrooms, and other plant, animal, and fungal material. Generally, one to a few grams of material are sampled at a time (Boynton et al., 2019; Kowallik et al., 2015; Spurley et al., 2021). It is important to use proper aseptic technique to avoid contamination by wearing gloves and sterilizing tools with a solution of 70% ethanol. Substrate heterogeneity should be accounted for, either by mixing (in the case of soils), or deep and repeated sampling (in the case of difficult to mix substrates, such as phylloplane, nectar, and fluxes) (Boundy-Mills, 2006; Yurkov 2017, 2018; Yurkov & Pozo, 2017).

Because of substrate diversity and heterogeneity, researchers designing field surveys to isolate yeasts should also consider the ecological factors that might affect sampling results. These ecological factors include the geographic region of the forest, sampling substrates, and seasonal conditions. For example, *Saccharomyces* isolation success can be dependent on season, temperature, and surrounding flora; (Robinson et al., 2016; Charron et al. 2014; Leducq et al., 2014; Sylvester et al., 2015; Eizaguirre et al., 2018; Langdon et al., 2020). Additionally, genetic diversity, including levels of heterozygosity and admixture, can vary among sampled substrates: for example, wild *Saccharomyces* strains from forests have lower admixture and heterozygosity than strains isolated from fruits, flowers, insects, or human-associated substrates (Magwene et al., 2011; Hyma and Fay 2013; Tilakaratna and Bensasson 2017; Günther et al., 2019). Finally, because the number of observed species depends on the sampling intensity, species-rich substrates require more samples than species-poor ones to determine species richness (Yurkov & Pozo 2017).

Researchers should record sampling conditions carefully when conducting sampling schemes to effectively answer their research questions. Across studies, it is essential to record standard data about every yeast isolation sample, including the GPS coordinates of sampling sites; collection dates; descriptions of isolation substrates such as tree species and substrate material; descriptions of field sites; and climate data such as temperature or humidity. This metadata can be used for resampling across time and makes field collections repeatable. Metadata is also important for putting patterns of habitat preference and genetic diversity into

environmental context. Collected samples with appropriate metadata recorded can now be processed for strain isolation.

4.2 Strain isolation for diversity estimation and new species discovery

Direct streaking and dilution plating on universal media is preferable over enrichment techniques and selective media to have a broad overview of yeast biodiversity (Lachance & Starmer 1998; Boundy-Mills 2006; Yurkov et al., 2011). Yeast yield can be increased by filtration and sample concentration steps (Lachance & Starmer 1998; Boundy-Mills 2006). Media used for yeast isolation and enumeration are generally complex and nutritionally rich. Additives that suppress bacteria and filamentous fungi and special cultivation conditions increase the chances of isolating yeasts (Boundy-Mills 2006).

Inoculated tubes and plates are grown and visually inspected every two to three days for a minimum of two weeks or until growth is visualized. Isolation of slow-growing yeasts may require a prolonged incubation for more than four weeks. Purified cultures can be stored in 15-20% glycerol stocks at -80 °C, but some yeasts should be safely cryopreserved at ultra-low (-196 or -140 °C) temperatures to avoid their loss. Isolates are often identified using barcode sequencing of genomic loci, including ribosomal RNA (Robinson, Pinharanda, and Bensasson 2016; White et al., 1990; Spurley et al. 2021), using physiological tests, or other techniques such as MALDI-TOF (Boekhout et al., 2021a; Lücking et al., 2020).

4.3 Strain isolation targeting individual taxa

Strategies for isolating targeted taxa rely on phenotypic differences between target and other taxa. Selective isolation involves any unique physiological condition (e.g. fermentation ability) or tolerance to inhibitors (e.g. ethanol tolerance). These strategies include plating onto a selective medium, which has additives or environmental conditions that prevent growth of non-target taxa. Similarly, researchers can establish enrichment cultures, which have conditions that allow target taxa to grow more quickly than non-target taxa. Both strategies require some knowledge of target yeast physiology to design selective culturing conditions.

When plating, selective media can be taxon-specific (Boynton et al., 2019; Mašínová et al., 2018). When targeting yeasts, acidification and broad spectrum antibiotics reduce bacterial growth; and incubation at low temperatures, benomyl, calcium propionate and Rose Bengal may help to reduce mold growth (Boundy-Mills, 2006). Although this cultivation method can also reveal rare and minor species, selective isolation must be accompanied by an additional replicate with a complete medium when assaying diversity to ensure the complete assessment of the community (Yurkov & Pozo, 2017), and a negative control for contamination errors. Yeasts that produce forcefully ejected ballistospores can be isolated from substances such as leaves using the (ballisto)spore-fall method, (Boundy-Mills, 2006). A sample is placed

above an agar plate, and ballistospores fall and germinate onto the surface of the plate (Boundy-Mills, 2006).

Enrichment culturing, which can be especially efficient for isolation of target taxa, has been used for over a century to isolate microorganisms from mixed communities (Beijerinck 1961). It involves taking a small amount of sampled substrate, mixing the substrate with a liquid selective growth medium in which the target microbe can outcompete other non-desired microbes, and then plating enriched samples onto solid growth medium to isolate individual colonies to be tested. Enrichment media vary depending on the targeted species and multiple compositions have been developed.

Researchers have isolated a broad diversity of yeasts in the subphylum Saccharomycotina and phylum Basidiomycota using glucose-based enrichment media, and they have developed targeted enrichment strategies for more narrow yeast groups. Examples of enrichment media that target a taxonomically broad collection of yeasts include a synthetic complete (SC) based medium with a mixture of amino acids (Sylvester et al., 2015), and media containing glucose concentrations ranging from 0.8 to 8% (Spurley et al., 2021). As with selective solid media discussed above, researchers often adjust pH, add broad spectrum antibiotics, or add ethanol to reduce non-yeast microbial growth and enrich for fermentative yeast (Sylvester et al., 2015; Sampaio & Gonçalves, 2008; Fingerman et al., 2002). To increase recovered yeast diversity, enrichment cultures are incubated at different temperatures. For instance, growth at lower temperatures (< 12 °C) may favor cryotolerant yeast species (Sampaio & Gonçalves, 2008). To target specific yeast groups, such as nitrate or nitrite-utilizing fungi (Kurtzman et al., 2011a; Opulente et al., 2018; Shen et al., 2018), researchers can alter the carbon or nitrogen sources in enrichment media or adjust pH (Table S5). For example, sucrose is often used in media to enrich *Saccharomyces* and related taxa (Sniegowski et al., 2002, Robinson et al., 2016; Charron et al., 2014; Boynton et al., 2021, 2019). Negative and positive controls, to help detect contamination or media errors, can include tubes with enrichment medium only or tubes inoculated with a known culture, respectively.

Once samples are introduced into liquid enrichment media, they are usually grown for a minimum of two weeks or until growth is visualized. The presence of yeast may be visible in the form of a white sediment, bubbles, or a cloudy medium. Upon visualization of growth, a second round of enrichment is sometimes done in the same or a different liquid medium until growth is visible. Cultures are diluted and plated to solid agar plates with a medium amenable to target yeasts. Growth on plates is checked daily and separated colonies are picked for species identification. Colony preservation and identification is conducted as described above.

5. Forest yeast exploitation

Brewers, winemakers, and industrial biologists are using a variety of yeasts from temperate forests to improve their biotechnological processes. Some of the most recent and innovative attempts at wild yeast exploitation relate to the use of wild *Saccharomyces* species in the brewing industry (Cubillos et al., 2019), but diverse research and development have been carried out with a broad range of temperate forest yeasts for decades (Wegner 1983; Kurtzman 2009, 2011a). Use of wild yeasts may be a viable strategy to reintroduce diversity that was lost during modernization of the brewing process (Aquilani et al., 2015), which coincided with the use of pure brewing yeast cultures. This development greatly improved fermentation consistency and efficiency but may also have resulted in a severe drop in genotypic and phenotypic diversity among brewing yeasts. Many traditional strains have been discarded in favor of a small number of high-performing pure strains (Gallone et al., 2018). However, recently isolated wild strains have phenotypic traits useful for brewing. For instance, 353 strains of *Saccharomyces* species recently isolated from oak niches in Slovenian Sub-Mediterranean forests and nearby vineyards had useful phenotypes, including resistance to high concentrations of copper, sulfite, and ethanol (Dashko et al., 2016). Notably, *S. eubayanus* has been used in commercial beer production (Gibson et al., 2017). *Saccharomyces paradoxus* has likewise shown potential for brewing and has been used commercially (Nikulin et al., 2020b). Recent studies have also demonstrated the potential of the newly discovered *Saccharomyces jurei* for beer brewing (Giannakou et al., 2021; Hutzler et al., 2021). Though not adapted to the brewing environment, these temperate forest strains possess several traits beneficial for brewing, such as the ability to utilize the main wort sugar maltose (and in *S. jurei*'s case also maltotriose), tolerance to low temperatures, and production of desirable flavor volatiles (Gallone et al., 2016; Naseeb et al., 2017; Giannakou et al., 2021; Hutzler et al., 2021). Studies involving other fermentation systems such as winemaking and baking have likewise shown the application potential of wild yeasts (Magalhães et al., 2017a, 2017b, 2021).

A number of wild *Saccharomyces* phenotypic traits are amenable to modification by evolutionary engineering (Gibson et al., 2020). While this approach had previously been employed to enhance the potential of existing production strains, there are several recent examples of the approach being applied to the wild yeast *S. eubayanus* to improve its applicability in brewing, such as removing phenolic off-flavor, adaptation to ethanol, and utilization of the wort sugar maltotriose (Diderich et al., 2018; Mardones et al., 2021; Baker and Hittinger 2019; Brouwers et al., 2019). Increased diversity of brewing strains may be achieved through hybridization of wild and domesticated yeast strains. Several studies have shown how a lager yeast phenotype can be recapitulated by combining beneficial traits of the wild yeast *S. eubayanus* (cold tolerance) with domesticated *S. cerevisiae* strains (efficient sugar utilization) (Gibson et al., 2013; Hebly et al., 2015; Krogerus et al., 2015; Mertens et al.,

2015). Moreover, several novel approaches have been proposed to generate genetic diversity of extant or de-novo hybrids that can be exploited in industrial applications (Nikulin et al., 2018; Peris et al., 2020; Naseeb et al., 2021; Mozzachiodi et al., 2021).

Non-*Saccharomyces* yeasts from temperate forests have also been exploited in industrial applications. For example, *Lachancea* species have been used for sour beer fermentation because of their lactic acid production (Domizio et al., 2016), and for mead fermentation (Villarreal et al., 2021). Additionally, *L. thermotolerans* and *L. fermentati* produce large amounts of extracellular enzymes, which have important protease, polygalacturonase, and β -glucosidase activity for wine fermentation and aroma production (Belda et al., 2016; Porter et al., 2019a; Strauss et al., 2001). *Lachancea* spp. also could be used as biological control agents (BCAs) (Medina et al., 2017) because some members of the genus can inhibit growth or sporulation of other fungi using killer toxins or volatile organic compounds (Fiori et al., 2014; Kono & Himeno, 1997; Aponte & Blaiotta, 2016b; González-Arenzana et al., 2017). Outside of the genus *Lachancea*, *Komagataella* species isolated from broad-leaved trees have been used since the early 1970s for the production of single-cell protein from methanol, and more recently other yeasts belonging the genus *Ogataea* have been used with the same aim (Wegner 1983; Kurtzman 2009, 2011b). Subsequently, researchers developed a protein expression system in *Komagataella*, and the genus is currently used to produce biopharmaceutical proteins, recombinant enzymes, and chemicals (Cregg et al., 1985; Tschopp et al., 1987; Gasser et al., 2013; Karbalaei et al., 2020; Duman-Özdamar et al., 2021; Gao et al., 2021).

6. Future perspectives

In this review, we have presented an up-to-date overview of yeast biodiversity and abundance in temperate forests. Most of this progress is a result of technological changes that enabled more reliable yeast identification using molecular genetic information instead of phenotypic tests. Yeast researchers' next steps include investigating how temperate forest yeast diversity has been shaped by dispersal and adaptation to changing environments across a long evolutionary time frame. Temperate forest yeasts are only a subset of yeast strains available in culture collections: most of the available yeast isolates have instead been derived from fungal infections (O'Brien et al., 2021) or substrates that are related to human activities (Almeida et al., 2015; Gallone et al., 2016). Moreover, none of the yeast species described in this review has been found among the most frequent species in environmental DNA from temperate forests, as reported in the GlobalFungi database. Our understanding of their roles in ecosystem functioning, if any, is still incomplete. Recent advances in sequencing and large population genomic surveys of various yeast species revealed their possible origins, population structure, and historical admixtures between lineages (Duan et al., 2018; Peter et

al., 2018; Ropars et al., 2018). Additional applications of whole-genome resequencing include interrogating genetic diversity to reveal how species adapt in nature and using genome-wide association studies (GWASes) to pinpoint genetic variants regulating yeasts' life cycles in temperate forests (Alonso-Blanco et al. 2016; De Chiara et al., 2020). Such studies have been conducted in some model species, such as *S. cerevisiae*, and can be expanded across many other yeast species to provide a holistic view of microbial ecology.

6.1 Microbial life cycle inferred through population genomics

Population genomics will be especially useful in inferring yeast life cycle parameters. Interpreting yeasts' natural history requires the inference of various fundamental parameters in molecular evolution, such as generation time, population size, mutation rate, and frequency of sex and outcrossing (Tsai et al. 2008). From population genomic data, historical events such as hybridization (Eberlein et al., 2019) or domestication (Jeffares et al., 2015; Gallone et al., 2016) have been estimated by calculating generation numbers. However, the generation times can depend on how much time cells spend in a non-dividing state (Gray et al., 2004) or in the spore state (Freese et al., 1982) in nature. For example, generation times can vary dramatically in *Saccharomyces* between about 90 minutes per generation in laboratory culture (Herskowitz 1988) and 150 generations per year in brewing environments (Gallone et al., 2016); these clocks need to be calibrated for temperate forest conditions to be useful to understand yeast growth and evolution in natural settings. In particular, we note that laboratory calibrations disagree widely with molecular clocks calibrated against the fossil record (Shen et al., 2018). The interplay of genetic variation, mutation rate, and generation time are further complicated by other aspects of yeast biology. One example is many yeasts' predominant asexual reproduction alternates with sporulation under harsh conditions and sexual contexts. Large variations exist in the frequency of sex in yeast (for a review see Nieuwenhuis and James 2016). Population genomic methods can be used to estimate the relative frequency of asexual and sexual reproduction by comparing differences in mutation and recombination rates (Tsai et al., 2008; Lee et al., 2021; Friedrich et al., 2015) and have shown population-level differences (Tsai et al., 2008; Liti et al., 2009; Drott et al., 2020; Koufopanou et al., 2020).

6.2 Genotype-phenotype through a reverse ecology approach

With our understanding of yeasts still biased towards research conducted in laboratory settings, we need to continue sampling naturally-occurring yeasts and recording abiotic and biotic factors, which will be helpful to conduct a reverse ecology approach (Li et al., 2008) to learn how yeasts live in nature. It is especially important to identify the factors that contribute to species' fitnesses in nature and try to replicate those as closely as possible in laboratory experiments (Liti, 2015). After establishing a collection of strains, we can start addressing how

genetic diversity is shaped by the adaptation to different environments, population drift, or bursts of population expansion associated with specific substrates. Such clonal lineages are especially suitable for GWAS-based approaches. Extensive collections are already available for the yeasts *Schizosaccharomyces pombe* (Jeffares et al., 2015) and *Saccharomyces cerevisiae* (Peter et al., 2018) and have allowed development of genetic tools, such as deletion strains (Giaever and Nislow 2014; Rai et al., 2018). However, sampling biases are still present, even in these established collections, with a low number of wild isolates from sparse geographic locations. The renewed interest in yeast biodiversity from natural habitats, including from temperate forests, will close this gap to provide a holistic view of the different species biology, their interaction among them and with the environment.

Box: Outstanding questions

1. How do we resolve differences in conclusions from culture-dependent and culture-independent studies of yeast diversity?

Sequencing and culture-based studies often uncover different diversity patterns and rare species are difficult to detect with either strategy. Both culture-dependent and culture-independent assays of yeast diversity have biases, including PCR and sequencing biases, unreliable barcode and genome databases, difficult-to-culture (i.e., fastidious) taxa, and taxa that can only survive within complex microbial communities. Additionally, sequence-based methods usually only report taxon frequencies, not abundances, although abundances can be inferred by scaling frequencies to another measure of cell abundances, such as qPCR or colony counts. Future studies will explore how to combine information from both strategies to get an accurate view of yeast diversity in temperate forests.

2. What role do broader sampling biases play in influencing conclusions about yeast ecology?

Enrichment culturing introduces sampling biases. Media and other enrichment conditions influence the yeast strains recovered, as do interactions among microorganisms in enrichment cultures. How can we quantify such biases to understand the effect of sampling strategies on our understanding of culturable yeast diversity? One possible strategy might be to release yeast cells into the environment and test different recovery strategies. Year-round sampling without enrichment could also help reduce the impact of sampling biases for some targeted yeasts.

3. How do we quantify life history parameters in temperate forest yeasts?

Our current understanding of life history parameters in temperate forest yeasts relies on estimates from genome sequencing and assumptions based on laboratory experiments. Direct observations or experimental conditions that closely mimic forest environments are needed to inform genomic models. The discrepancy with molecular clocks calibrated against the fossil record also needs to be addressed and extended to non-model yeast groups.

4. What are the contributions of temperate forest yeasts to ecosystem functioning?

While we know that yeasts are common and diverse, and we often have information about their genetics and physiology, few studies have directly investigated their impacts on other members of forest communities. However, research examining microbial communities from a guild-centered point of view shows promise in helping us to understand yeasts' contributions to ecosystem functioning (Martinović et al., 2021).

5. What are the wild niches of well-studied laboratory yeasts?

Saccharomyces, *Cryptococcus*, *Komagataella*, *Lachancea*, and others are studied in the laboratory and exist in temperate forests. Understanding their natural ecologies would help laboratory researchers to put the genetics, genomics, and biotechnology of their systems into a more holistic context. These organisms have the potential to be interesting ecological models in addition to their roles in laboratory research.

6. How do we reliably identify temperate forest yeasts species and determine boundaries among yeast taxonomic units?

Mycologists often use DNA barcodes, such as ribosomal sequences, to identify yeast taxa, but barcodes do not always reliably delimit species (Schoch et al., 2012). Recent studies combined barcode sequences and physiological characteristics to describe new temperate forest yeast taxa from Bulgaria (Gouliamova et al., 2016; Gouliamova and Dimitrov, 2020). Such studies, especially when combined with genome sequencing and genetic crosses, can help determine how phenotypic characteristics shift across the boundaries of taxonomic units.

7. How will our understanding of yeast ecology in temperate forests change as we focus away from well-studied systems and towards full temperate forest yeast communities? Yeast research in general has a bias towards a few well-studied genera, such as the genus *Saccharomyces*, which are not representative of yeast diversity. Instead, temperate forest yeast ecology needs to integrate entire yeast communities.

Author contributions:

All authors participated in manuscript planning and review of the final draft. PBo, SM, and GL conceptualized the project, organized the team, and edited the final draft for submission. PBo and SM edited and synthesized all contributed material. PBo and AY wrote the introduction; PBa, and TM contributed the section on global yeast diversity, including providing new analyses for this review; KBM, PBu, BT, NC, and AY contributed the section on yeasts in culture collections; FB, GB, PBo, FC, DGr, VK, CRL, SM, ESN, and IJT contributed the section on the *Saccharomyces* genus; VK compiled data from the literature for Table S3 and Figures on *Saccharomyces cerevisiae/paradoxus* isolating success; FC provided the data on *uvarum/eubayanus*; PBu, BT, and AY contributed the section on *Cryptococcus* and former *Cryptococcus*; LH, MJ, and ESN contributed the section on *Komagataella*; FC and PV contributed the section on *Lachancea*; GB, FC, KJF, DGr, CTH, DO, JP, and AY contributed the section on yeast isolation; NC, SD, BRG, and UP contributed the section on yeast exploitation; PBo, RD, DGo, DGr, GL, SM, IJT, and AY contributed the sections on future perspectives and the contents of the outstanding questions box.

Acknowledgements:

We would like to sincerely thank members of the yeast and fungal community who helped us conceive and plan this review, and who read previous versions of this manuscript, including Douda Bensasson, Justin Fay, Matthew Goddard, Paula Gonçalves, Marizeth Groenwald, Marc-André Lachance, Paul Sniegowski, and Kathleen Treseder. Parts of this work were funded by the Czech Science Foundation (21-17749S) to PBa. Parts of this work were funded by the National Science Foundation (DEB-1442148 and DEB-2110403); the DOE Great Lakes Bioenergy Research Center (DE-SC0018409), the USDA National Institute of Food and Agriculture (Hatch Project 1020204); the Office of the Vice Chancellor for Research and Graduate Education (H. I. Romnes Faculty Fellow), University of Wisconsin-Madison; and the Wisconsin Alumni Research Foundation, to CTH. KJF is a Morgridge Metabolism Interdisciplinary Fellow of the Morgridge Institute for Research. The research of Roumen Dimitrov and Dilnora Gouliamova was supported by the grant KP-06-31/19 from Bulgarian Science Fund.

Conflict of interest: The authors declare no conflict of interest.

Data availability statement: All new data for this review is included in the supplemental materials.

Bibliography

Albarracín, M. V., Six, J., Houlton, B. Z., & Bledsoe, C. S. (2013). A nitrogen fertilization field study of carbon-13 and nitrogen-15 transfers in ectomycorrhizas of *Pinus sabiniana*. *Oecologia*, 173(4), 1439–1450. <https://doi.org/10.1007/s00442-013-2734-4>

Alfaro Reyna, T., Retana, J., & Martínez-Vilalta, J. (2018). Is there a substitution of Pinaceae by Fagaceae in temperate forests at the global scale? *Global and Planetary Change*, 166, 41–47. <https://doi.org/10.1016/j.gloplacha.2018.04.001>

Almeida, P., Barbosa, R., Zalar, P., Imanishi, Y., Shimizu, K., Turchetti, B., Legras, J.-L., Serra, M., Dequin, S., Couloux, A., Guy, J., Bensasson, D., Gonçalves, P., & Sampaio, J. P. (2015). A population genomics insight into the Mediterranean origins of wine yeast domestication. *Molecular Ecology*, 24(21), 5412–5427. <https://doi.org/10.1111/mec.13341>

Almeida, P., Gonçalves, C., Teixeira, S., Libkind, D., Bontrager, M., Masneuf-Pomarède, I., Albertin, W., Durrens, P., Sherman, D. J., Marullo, P., Todd Hittinger, C., Gonçalves, P., & Sampaio, J. P. (2014). A Gondwanan imprint on global diversity and domestication of wine and cider yeast *Saccharomyces uvarum*. *Nature Communications*, 5(1), 4044. <https://doi.org/10.1038/ncomms5044>

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M., Cao, J., Chae, E., Dezwaan, T. M., Ding, W., Ecker, J. R., Exposito-Alonso, M., Farlow, A., Fitz, J., Gan, X., Grimm, D. G., Hancock, A. M., Henz, S. R., Holm, S., ... Zhou, X. (2016). 1,135 Genomes Reveal the Global Pattern of Polymorphism in *Arabidopsis thaliana*. *Cell*, 166(2), 481–491. <https://doi.org/10.1016/j.cell.2016.05.063>

Alsammar, H. F., Naseeb, S., Brancia, L. B., Gilman, R. T., Wang, P., & Delneri, D. (2019). Targeted metagenomics approach to capture the biodiversity of *Saccharomyces* genus in wild environments. *Environmental Microbiology Reports*, 11(2), 206–214. <https://doi.org/10.1111/1758-2229.12724>

Anderson, J. B., Kasimer, D., Xia, W., Schröder, N. C. H., Cichowicz, P., Lionello, S., Chakrabarti, R., Mohan, E., & Kohn, L. M. (2018). Persistence of Resident and Transplanted Genotypes of the Undomesticated Yeast *Saccharomyces paradoxus* in Forest Soil. *MSphere*, 3(3). <https://doi.org/10.1128/mSphere.00211-18>

Aponte, M., & Blaiotta, G. (2016). Potential Role of Yeast Strains Isolated from Grapes in the Production of Taurasi DOCG. *Frontiers in Microbiology*, 7. <https://doi.org/10.3389/fmicb.2016.00809>

Aquilani, B., Laureti, T., Poponi, S., & Secondi, L. (2015). Beer choice and consumption determinants when craft beers are tasted: An exploratory study of consumer preferences. *Food Quality and Preference*, 41, 214–224. <https://doi.org/10.1016/j.foodqual.2014.12.005>

Babjeva and Golovleva, 1963 Yeast flora of the main soil types of the European part of the USSR. *Microorganisms in agriculture*

Bagheri, B., Bauer, F. F., & Setati, M. E. (2016). The Diversity and Dynamics of Indigenous Yeast Communities in Grape Must from Vineyards Employing Different Agronomic Practices and their Influence on Wine Fermentation. *South African Journal of Enology and Viticulture*, 36(2). <https://doi.org/10.21548/36-2-957>

Baker, E. P., & Hittinger, C. T. (2019). Evolution of a novel chimeric maltotriose transporter in *Saccharomyces eubayanus* from parent proteins unable to perform this function. *PLOS Genetics*, 15(4), e1007786. <https://doi.org/10.1371/journal.pgen.1007786>

Baleiras Couto, M. M., Reizinho, R. G., & Duarte, F. L. (2005). Partial 26S rDNA restriction analysis as a tool to characterise non-Saccharomyces yeasts present during red wine fermentations. *International Journal of Food Microbiology*, 102(1), 49–56. <https://doi.org/10.1016/j.ijfoodmicro.2005.01.005>

Bandara, A., Fraser, S., Chambers, P. J., & Stanley, G. A. (2009). Trehalose promotes the survival of *Saccharomyces cerevisiae* during lethal ethanol stress, but does not influence growth under sublethal ethanol stress. *FEMS Yeast Research*, 9(8), 1208–1216. <https://doi.org/10.1111/j.1567-1364.2009.00569.x>

Barbosa, R., Almeida, P., Safar, S. V. B., Santos, R. O., Morais, P. B., Nielly-Thibault, L., Leducq, J.-B., Landry, C. R., Gonçalves, P., Rosa, C. A., & Sampaio, J. P. (2016). Evidence of Natural Hybridization in Brazilian Wild Lineages of *Saccharomyces cerevisiae*. *Genome Biology and Evolution*, 8(2), 317–329. <https://doi.org/10.1093/gbe/evv263>

Banno & Mikata 1981 Ascomycetous yeasts isolated from forest materials in Japan. *IFO Res. Commun.*, 10, 10-19 (1981).

Batschinskaya A.A., (1914) *Saccharomyces paradoxus*, a new yeast species; its development and its pure culture. *J Microbiol Epidemiol Immunobiol* 1:231–250.

Beard, J. S. (1990). Temperate forests of the southern hemisphere. *Vegetatio*, 89(1), 7–10. <https://doi.org/10.1007/BF00134430>

Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., & Wood, E. F. (2018). Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5, 180214. <https://doi.org/10.1038/sdata.2018.214>

Beijerinck, M. W. (1961) Enrichment culture studies with urea bacteria (T. D. Brock, Trans.).

Belda, I., Ruiz, J., Alastruey-Izquierdo, A., Navascués, E., Marquina, D., & Santos, A. (2016). Unraveling the Enzymatic Basis of Wine “Flavorome”: A Phylo-Functional Study of Wine Related Yeast Species. *Frontiers in Microbiology*, 7. <https://doi.org/10.3389/fmicb.2016.00012>

Bendixsen, D. P., Gettle, N., Gilchrist, C., Zhang, Z., & Stelkens, R. (2021). Genomic Evidence of an Ancient East Asian Divergence Event in Wild *Saccharomyces cerevisiae*. *Genome Biology and Evolution*, 13(2), evab001. <https://doi.org/10.1093/gbe/evab001>

Benito, Á., Calderón, F., Palomero, F., & Benito, S. (2015). Combine Use of Selected *Schizosaccharomyces pombe* and *Lachancea thermotolerans* Yeast Strains as an Alternative to the Traditional Malolactic Fermentation in Red Wine Production. *Molecules*, 20(6), 9510–9523. <https://doi.org/10.3390/molecules20069510>

Bergström, A., Simpson, J. T., Salinas, F., Barré, B., Parts, L., Zia, A., Nguyen Ba, A. N., Moses, A. M., Louis, E. J., Mustonen, V., Warringer, J., Durbin, R., & Liti, G. (2014). A High-Definition View of Functional Genetic Variation from Natural Yeast Genomes. *Molecular Biology and Evolution*, 31(4), 872–888. <https://doi.org/10.1093/molbev/msu037>

Bing, J., Han, P.-J., Liu, W.-Q., Wang, Q.-M., & Bai, F.-Y. (2014). Evidence for a Far East Asian origin of lager beer yeast. *Current Biology*, 24(10), R380–R381. <https://doi.org/10.1016/j.cub.2014.04.031>

Birkhofer, K., Schöning, I., Alt, F., Herold, N., Klärner, B., Maraun, M., Marhan, S., Oelmann, Y., Wubet, T., Yurkov, A., Begerow, D., Berner, D., Buscot, F., Daniel, R., Diekötter, T., Ehnes, R. B., Erdmann, G., Fischer, C., Foesel, B., ... Schrumpf, M. (2012). General relationships between abiotic soil properties and soil biota across spatial scales and different land-use types. *PLoS One*, 7(8), e43292. <https://doi.org/10.1371/journal.pone.0043292>

Boddy, L., & Hiscox, J. (2016). Fungal Ecology: Principles and Mechanisms of Colonization and Competition by Saprotrrophic Fungi. *Microbiology Spectrum*, 4(6). <https://doi.org/10.1128/microbiolspec.FUNK-0019-2016>

Boekhout, T., Aime, M. C., Begerow, D., Gabaldón, T., Heitman, J., Kemler, M., Khayhan, K., Lachance, M.-A., Louis, E. J., Sun, S., Vu, D., & Yurkov, A. (2021). The evolving species concepts used for yeasts: From phenotypes and genomes to speciation networks. *Fungal Diversity*, 109(1), 27–55. <https://doi.org/10.1007/s13225-021-00475-9>

Botha, A. (2006). Yeasts in Soil. In G. Péter & C. Rosa (Eds.), *Biodiversity and Ecophysiology of Yeasts* (pp. 221–240). Springer-Verlag. https://doi.org/10.1007/3-540-30985-3_11

Botha, A. (2011). The importance and ecology of yeasts in soil. *Soil Biology and Biochemistry*, 43(1), 1–8. <https://doi.org/10.1016/j.soilbio.2010.10.001>

Boundy-Mills, K. L. (2006). Methods for investigating yeast biodiversity. In C. A. Rosa & G. Peter (Eds.), *Biodiversity and ecophysiology of yeasts* (pp. 67-100). Springer, Berlin, Heidelberg

Boundy-Mills, K. L., Glantschnig, E., Roberts, I. N., Yurkov, A., Casaregola, S., Daniel, H.-M., Groenewald, M., & Turchetti, B. (2016). Yeast culture collections in the twenty-first century: New opportunities and challenges: Yeast culture collections in the 21st century. *Yeast*, 33(7), 243–260. <https://doi.org/10.1002/yea.3171>

Boynton, P. J., Kowallik, V., Landermann, D., & Stukenbrock, E. H. (2019). Quantifying the efficiency and biases of forest *Saccharomyces* sampling strategies. *Yeast*, 36(11), 657–668. <https://doi.org/10.1002/yea.3435>

Boynton, P. J., Wloch-Salamon, D., Landermann, D., & Stukenbrock, E. H. (2021). Forest *Saccharomyces paradoxus* are robust to seasonal biotic and abiotic changes. *Ecology and Evolution*, 11(11), 6604–6619. <https://doi.org/10.1002/ee.3751>

Brock, J. M. R., Perry, G. L. W., Lee, W. G., & Burns, B. R. (2016). Tree fern ecology in New Zealand: A model for southern temperate rainforests. *Forest Ecology and Management*, 375, 112–126. <https://doi.org/10.1016/j.foreco.2016.05.030>

Brouwers, N., Gorter de Vries, A. R., van den Broek, M., Weening, S. M., Elink Schuurman, T. D., Kuijpers, N. G. A., Pronk, J. T., & Daran, J.-M. G. (2019). In vivo recombination of *Saccharomyces eubayanus* maltose-transporter genes yields a chimeric transporter that enables maltotriose fermentation. *PLOS Genetics*, 15(4), e1007853. <https://doi.org/10.1371/journal.pgen.1007853>

Charron, G., Leducq, J.-B., Bertin, C., Dubé, A. K., & Landry, C. R. (2014). Exploring the northern limit of the distribution of *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* in North America. *FEMS Yeast Research*, 14(2), 281–288. <https://doi.org/10.1111/1567-1364.12100>

Chernov et al., (2013) Annotated list of yeasts from Moscow region. Mikol Fitopatol

Chowdhary, A., Randhawa, H. S., Boekhout, T., Hagen, F., Klaassen, C. H., & Meis, J. F. (2012). Temperate climate niche for *Cryptococcus gattii* in Northern Europe. *Emerging Infectious Diseases*, 18(1), 172–174. <https://doi.org/10.3201/eid1801.111190>

Claudino-Sales, V. (2019). Laurisilva of Madeira, Portugal. In V. Claudino-Sales, *Coastal World Heritage Sites* (Vol. 28, pp. 243–249). Springer Netherlands. https://doi.org/10.1007/978-94-024-1528-5_36

Clavijo, A., Calderón, I. L., & Panque, P. (2010). Diversity of *Saccharomyces* and non-*Saccharomyces* yeasts in three red grape varieties cultured in the Serranía de Ronda (Spain) vine-growing region. *International Journal of Food Microbiology*, 143(3), 241–245. <https://doi.org/10.1016/j.ijfoodmicro.2010.08.010>

Cogliati, M., D'Amicis, R., Zani, A., Montagna, M. T., Caggiano, G., De Giglio, O., Balbino, S., De Donno, A., Serio, F., Susever, S., Ergin, C., Velegraki, A., Ellabib, M. S., Nardoni, S., Macci, C., Oliveri, S., Trovato, L., Dipineto, L., Rickerts, V., ... Colom, M. F. (2016). Environmental distribution of *Cryptococcus neoformans* and *C. gattii* around the Mediterranean basin. *FEMS Yeast Research*, 16(7), fow086. <https://doi.org/10.1093/femsyr/fow086>

Cordero-Bueso, G., Esteve-Zarzoso, B., Cabellos, J. M., Gil-Díaz, M., & Arroyo, T. (2013). Biotechnological potential of non-Saccharomyces yeasts isolated during spontaneous fermentations of Malvar (*Vitis vinifera* cv. L.). *European Food Research and Technology*, 236(1), 193–207. <https://doi.org/10.1007/s00217-012-1874-9>

Cowling, R. M., Rundel, P. W., Lamont, B. B., Kalin Arroyo, M., & Arianoutsou, M. (1996). Plant diversity in mediterranean-climate regions. *Trends in Ecology & Evolution*, 11(9), 362–366. [https://doi.org/10.1016/0169-5347\(96\)10044-6](https://doi.org/10.1016/0169-5347(96)10044-6)

Cregg, J. M., Barringer, K. J., Hessler, A. Y., & Madden, K. R. (1985). *Pichia pastoris* as a host system for transformations. *Molecular and Cellular Biology*, 5(12), 3376–3385. <https://doi.org/10.1128/mcb.5.12.3376-3385.1985>

Cubillos, F. A., Gibson, B., Grijalva-Vallejos, N., Krogerus, K., & Nikulin, J. (2019). Bioprospecting for brewers: Exploiting natural diversity for naturally diverse beers. *Yeast*, 36(6), 383–398. <https://doi.org/10.1002/yea.3380>

Dashko, S., Liu, P., Volk, H., Butinar, L., Piškur, J., & Fay, J. C. (2016). Changes in the Relative Abundance of Two Saccharomyces Species from Oak Forests to Wine Fermentations. *Frontiers in Microbiology*, 7. <https://doi.org/10.3389/fmicb.2016.00215>

De Chiara, M., Barré, B., Persson, K., Chioma, A. O., Irizar, A., Schacherer, J., Warringer, J., & Liti, G. (2020). Domestication reprogrammed the budding yeast life cycle [Preprint]. *Evolutionary Biology*. <https://doi.org/10.1101/2020.02.08.939314>

Diderich, J. A., Weening, S. M., van den Broek, M., Pronk, J. T., & Daran, J.-M. G. (2018). Selection of Pof-Saccharomyces eubayanus Variants for the Construction of *S. cerevisiae* × *S. eubayanus* Hybrids With Reduced 4-Vinyl Guaiacol Formation. *Frontiers in Microbiology*, 9, 1640. <https://doi.org/10.3389/fmicb.2018.01640>

Dlauchy, D. (2003). [No title found]. *Antonie van Leeuwenhoek*, 83(4), 327–332. <https://doi.org/10.1023/A:1023318829389>

Domizio, P., House, J. F., Joseph, C. M. L., Bisson, L. F., & Bamforth, C. W. (2016). *Lachancea thermotolerans* as an alternative yeast for the production of beer †. *Journal of the Institute of Brewing*, 122(4), 599–604. <https://doi.org/10.1002/jib.362>

Drott, M. T., Satterlee, T. R., Skerker, J. M., Pfannenstiel, B. T., Glass, N. L., Keller, N. P., & Milgroom, M. G. (2020). The Frequency of Sex: Population Genomics Reveals Differences in Recombination and Population Structure of the Aflatoxin-Producing Fungus *Aspergillus flavus*. *MBio*, 11(4). <https://doi.org/10.1128/mBio.00963-20>

Duan, S.-F., Han, P.-J., Wang, Q.-M., Liu, W.-Q., Shi, J.-Y., Li, K., Zhang, X.-L., & Bai, F.-Y. (2018). The origin and adaptive evolution of domesticated populations of yeast from Far East Asia. *Nature Communications*, 9(1), 2690. <https://doi.org/10.1038/s41467-018-05106-7>

Duman-Özdamar, Z. E., & Binay, B. (2021). Production of Industrial Enzymes via *Pichia pastoris* as a Cell Factory in Bioreactor: Current Status and Future Aspects. *The Protein Journal*, 40(3), 367–376. <https://doi.org/10.1007/s10930-021-09968-7>

Eberlein, C., Hénault, M., Fijarczyk, A., Charron, G., Bouvier, M., Kohn, L. M., Anderson, J. B., & Landry, C. R. (2019). Hybridization is a recurrent evolutionary stimulus in wild yeast speciation. *Nature Communications*, 10(1), 923. <https://doi.org/10.1038/s41467-019-08809-7>

Eizaguirre, J. I., Peris, D., Rodríguez, M. E., Lopes, C. A., De Los Ríos, P., Hittinger, C. T., & Libkind, D. (2018). Phylogeography of the wild Lager-brewing ancestor (*Saccharomyces eubayanus*) in Patagonia. *Environmental Microbiology*, 20(10), 3732–3743. <https://doi.org/10.1111/1462-2920.14375>

Esteve-Zarzoso, B., Peris-Torán, M. J., García-Maiquez, E., Uruburu, F., & Querol, A. (2001). Yeast Population Dynamics during the Fermentation and Biological Aging of Sherry Wines. *Applied and Environmental Microbiology*, 67(5), 2056–2061. <https://doi.org/10.1128/AEM.67.5.2056-2061.2001>

Fernandez, M. T., Ubeda, J. F., & Briones, A. I. (1999). Comparative study of non- *Saccharomyces* microflora of musts in fermentation, by physiological and molecular methods. *FEMS Microbiology Letters*, 173(1), 223–229. <https://doi.org/10.1111/j.1574-6968.1999.tb13506.x>

Fingerman, M. (Ed.). (2002). Recent Advances in Marine Biotechnology, Vol. 6 (0 ed.). CRC Press. <https://doi.org/10.1201/9781482279832>

Fiori, S., Urgeghe, P. P., Hammami, W., Razzu, S., Jaoua, S., & Migheli, Q. (2014). Biocontrol activity of four non- and low-fermenting yeast strains against *Aspergillus carbonarius* and their ability to remove ochratoxin A from grape juice. *International Journal of Food Microbiology*, 189, 45–50. <https://doi.org/10.1016/j.ijfoodmicro.2014.07.020>

França, L., Sannino, C., Turchetti, B., Buzzini, P., & Margesin, R. (2016). Seasonal and altitudinal changes of culturable bacterial and yeast diversity in Alpine forest soils. *Extremophiles*, 20(6), 855–873. <https://doi.org/10.1007/s00792-016-0874-2>

Freese, E. B., Chu, M. I., & Freese, E. (1982). Initiation of Yeast Sporulation by Partial Carbon, Nitrogen, or Phosphate Deprivation. *Journal of Bacteriology*, 149(3), 840–851. <https://doi.org/10.1128/jb.149.3.840-851.1982>

Friedrich, A., Jung, P., Reisser, C., Fischer, G., & Schacherer, J. (2015). Population Genomics Reveals Chromosome-Scale Heterogeneous Evolution in a Protoploid Yeast. *Molecular Biology and Evolution*, 32(1), 184–192. <https://doi.org/10.1093/molbev/msu295>

Gallone, B., Mertens, S., Gordon, J. L., Maere, S., Verstrepen, K. J., & Steensels, J. (2018). Origins, evolution, domestication and diversity of *Saccharomyces* beer yeasts. *Current Opinion in Biotechnology*, 49, 148–155. <https://doi.org/10.1016/j.copbio.2017.08.005>

Gallone, B., Steensels, J., Mertens, S., Dzialo, M. C., Gordon, J. L., Wauters, R., Theßeling, F. A., Bellinazzo, F., Saelens, V., Herrera-Malaver, B., Prahl, T., White, C., Hutzler, M., Meußdoerffer, F., Malcorps, P., Souffriau, B., Daenen, L., Baele, G., Maere, S., & Verstrepen, K. J. (2019). Interspecific hybridization facilitates niche adaptation in beer yeast. *Nature Ecology & Evolution*, 3(11), 1562–1575. <https://doi.org/10.1038/s41559-019-0997-9>

Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saelens, V., Herrera-Malaver, B., Merlevede, A., Roncoroni, M., Voordeckers, K., Miraglia, L., Teiling, C., Steffy, B., Taylor, M., Schwartz, A., Richardson, T., White, C., Baele, G., Maere, S., & Verstrepen, K. J. (2016). Domestication and Divergence of *Saccharomyces cerevisiae* Beer Yeasts. *Cell*, 166(6), 1397–1410.e16. <https://doi.org/10.1016/j.cell.2016.08.020>

Gao, J., Jiang, L., & Lian, J. (2021). Development of synthetic biology tools to engineer *Pichia pastoris* as a chassis for the production of natural products. *Synthetic and Systems Biotechnology*, 6(2), 110–119. <https://doi.org/10.1016/j.synbio.2021.04.005>

Gasser, B., Priehofer, R., Marx, H., Maurer, M., Nocon, J., Steiger, M., Puxbaum, V., Sauer, M., & Mattanovich, D. (2013). *Pichia pastoris*: Protein production host and model organism for biomedical research. *Future Microbiology*, 8(2), 191–208. <https://doi.org/10.2217/fmb.12.133>

Gayevskiy, V., Lee, S., & Goddard, M. R. (2016). European derived *Saccharomyces cerevisiae* colonisation of New Zealand vineyards aided by humans. *FEMS Yeast Research*, 16(7), fow091. <https://doi.org/10.1093/femsyr/fow091>

Giaever, G., & Nislow, C. (2014). The Yeast Deletion Collection: A Decade of Functional Genomics. *Genetics*, 197(2), 451–465. <https://doi.org/10.1534/genetics.114.161620>

Giannakou, K., Visinoni, F., Zhang, P., Nathoo, N., Jones, P., Cotterrell, M., Vrhovsek, U., & Delneri, D. (2021). Biotechnological exploitation of *Saccharomyces jurei* and its hybrids in craft beer fermentation uncovers new aroma combinations. *Food Microbiology*, 100, 103838. <https://doi.org/10.1016/j.fm.2021.103838>

Gibson, B., Dahabieh, M., Krogerus, K., Jouhten, P., Magalhães, F., Pereira, R., Siewers, V., & Vidgren, V. (2020). Adaptive Laboratory Evolution of Ale and Lager Yeasts for Improved Brewing Efficiency and Beer Quality. *Annual Review of Food Science and Technology*, 11(1), 23–44. <https://doi.org/10.1146/annurev-food-032519-051715>

Gibson, B., Geertman, J.-M. A., Hittinger, C. T., Krogerus, K., Libkind, D., Louis, E. J., Magalhães, F., & Sampaio, J. P. (2017). New yeasts—new brews: Modern approaches to brewing yeast design and development. *FEMS Yeast Research*, 17(4). <https://doi.org/10.1093/femsyr/fox038>

Gibson, B. R., Storgårds, E., Krogerus, K., & Vidgren, V. (2013). Comparative physiology and fermentation performance of Saaz and Frohberg lager yeast strains and the parental species *Saccharomyces eubayanus*: Fermentation performance of Saaz and Frohberg yeast. *Yeast*, 30(7), 255–266. <https://doi.org/10.1002/yea.2960>

Global Forest Resources Assessment 2020. (2020). FAO. <https://doi.org/10.4060/ca9825en>

Glushakova, A. M., Ivannikova, Yu. V., Naumova, E. S., Chernov, I. Yu., & Naumov, G. I. (2007). Massive isolation and identification of *Saccharomyces paradoxus* yeasts from plant phyllosphere. *Microbiology*, 76(2), 205–210. <https://doi.org/10.1134/S0026261707020129>

Glushakova, A. M., Kachalkin, A. V., & Chernov, I. Yu. (2011). Specific features of the dynamics of epiphytic and soil yeast communities in the thickets of Indian balsam on mucky gley soil. *Eurasian Soil Science*, 44(8), 886–892. <https://doi.org/10.1134/S1064229311080059>

Golubev and Golubeva, (2004) Yeast fungi in steppe and forest phytocenoses of the Prioksko- Terrasny Nature Reserve

González, S. S., Barrio, E., & Querol, A. (2006). Molecular identification and characterization of wine yeasts isolated from Tenerife (Canary Island, Spain). *Journal of Applied Microbiology*, 0(0), 061120055200058-???. <https://doi.org/10.1111/j.1365-2672.2006.03150.x>

González-Arenzana, L., Garijo, P., Berlanas, C., López-Alfaro, I., López, R., Santamaría, P., & Gutiérrez, A. R. (2017). Genetic and phenotypic intraspecific variability of non- *Saccharomyces* yeasts populations from La Rioja winegrowing region (Spain). *Journal of Applied Microbiology*, 122(2), 378–388. <https://doi.org/10.1111/jam.13341>

Gouliamova, D., & Dimitrov, R. (2020). *Kazachstania chrysolinae* and *Kazachstania bozae* Two New Yeast Species of the Genus *Kazachstania*. Transfer of Four *Kazachstania* Species to *Grigorovia* Gen. Nov. As New Combinations. 'Prof. Marin Drinov' Publishing House of Bulgarian Academy of Sciences. <https://doi.org/10.7546/CRABS.2020.01.06>

Gouliamova, D. E., Dimitrov, R. A., Smith, M. Th., Groenewald, M., Stoilova-Disheva, M. M., Guéorguiev, B. V., & Boekhout, T. (2016). DNA barcoding revealed *Nematodospora valgi* gen. Nov., sp. Nov. And *Candida cetoniae* sp. Nov. In the Lodderomyces clade. *Fungal Biology*, 120(2), 179–190. <https://doi.org/10.1016/j.funbio.2015.05.008>

Gray, J. V., Petsko, G. A., Johnston, G. C., Ringe, D., Singer, R. A., & Werner-Washburne, M. (2004). "Sleeping Beauty": Quiescence in *Saccharomyces cerevisiae*. *Microbiology and Molecular Biology Reviews*, 68(2), 187–206. <https://doi.org/10.1128/MMBR.68.2.187-206.2004>

Günther, C. S., & Goddard, M. R. (2019). Do yeasts and *Drosophila* interact just by chance? *Fungal Ecology*, 38, 37–43. <https://doi.org/10.1016/j.funeco.2018.04.005>

Han, D.-Y., Han, P.-J., Rumbold, K., Koricha, A. D., Duan, S.-F., Song, L., Shi, J.-Y., Li, K., Wang, Q.-M., & Bai, F.-Y. (2021). Adaptive gene content and allele distribution variations in the wild and domesticated populations of *Saccharomyces cerevisiae*. *Frontiers in Microbiology* 12, 631250. doi:10.3389/fmicb.2021.631250

Hebly, M., Brickwedde, A., Bolat, I., Driessen, M. R. M., de Hulster, E. A. F., van den Broek, M., Pronk, J. T., Geertman, J.-M., Daran, J.-M., & Daran-Lapujade, P. (2015). *S. cerevisiae* × *S. eubayanus* interspecific hybrid, the best of both worlds and beyond. *FEMS Yeast Research*, 15(3). <https://doi.org/10.1093/femsyr/fov005>

Herskowitz, I. (1988). Life cycle of the budding yeast *Saccharomyces cerevisiae*. *Microbiological Reviews*, 52(4), 536–553. <https://doi.org/10.1128/mr.52.4.536-553.1988>

Hsieh, H.-H., Wang, S.-Y., Chen, T.-L., Huang, Y.-L., & Chen, M.-J. (2012). Effects of cow's and goat's milk as fermentation media on the microbial ecology of sugary kefir grains. *International Journal of Food Microbiology*, 157(1), 73–81. <https://doi.org/10.1016/j.ijfoodmicro.2012.04.014>

Hutzler, M., Michel, M., Kunz, O., Kuusisto, T., Magalhães, F., Krogerus, K., & Gibson, B. (2021). Unique Brewing-Relevant Properties of a Strain of *Saccharomyces jurei* Isolated From Ash (*Fraxinus excelsior*). *Frontiers in Microbiology*, 12, 645271. <https://doi.org/10.3389/fmicb.2021.645271>

Hyma, K. E., & Fay, J. C. (2013). Mixing of vineyard and oak-tree ecotypes of *Saccharomyces cerevisiae* in North American vineyards. *Molecular Ecology*, 22(11), 2917–2930. <https://doi.org/10.1111/mec.12155>

Jeffares, D. C., Rallis, C., Rieux, A., Speed, D., Převorovský, M., Mourier, T., Marsella, F. X., Iqbal, Z., Lau, W., Cheng, T. M. K., Pracana, R., Mülleeder, M., Lawson, J. L. D., Chessel, A., Bala, S., Hellenthal, G., O'Fallon, B., Keane, T., Simpson, J. T., ... Bähler, J. (2015). The genomic and phenotypic diversity of *Schizosaccharomyces pombe*. *Nature Genetics*, 47(3), 235–241. <https://doi.org/10.1038/ng.3215>

Jensen, M. A., True, H. L., Chernoff, Y. O., & Lindquist, S. (2001). Molecular population genetics and evolution of a prion-like protein in *Saccharomyces cerevisiae*. *Genetics*, 159(2), 527–535. <https://doi.org/10.1093/genetics/159.2.527>

Jensen et al. (1963) Studies on the microflora of Danish beech forest soils. IV. Yeasts and yeast-like fungi. *Zentralbl Bakteriol Mikrobiol Hyg A*

Johnson, L. J., Koufopanou, V., Goddard, M. R., Hetherington, R., Schäfer, S. M., & Burt, A. (2004). Population Genetics of the Wild Yeast *Saccharomyces paradoxus*. *Genetics*, 166(1), 43–52. <https://doi.org/10.1534/genetics.166.1.43>

Kenkichi, K. & Tadashi, K. 1974 Ascosporogenous yeasts isolated from tree exudates in Japan.

Karbalaei, M., Rezaee, S. A., & Farsiani, H. (2020). *Pichia pastoris*: A highly successful expression system for optimal synthesis of heterologous proteins. *Journal of Cellular Physiology*, 235(9), 5867–5881. <https://doi.org/10.1002/jcp.29583>

Kemler, M., Witfeld, F., Begerow, D., & Yurkov, A. (2017). Phyloplane yeasts in temperate climates. In P. Buzzini, M.-A. Lachance, & A. Yurkov (Eds.), *Yeasts in Natural Ecosystems: Ecology* (pp. 171–197). Springer International Publishing. https://doi.org/10.1007/978-3-319-62683-3_6

Kidd, S. E., Hagen, F., Tscharke, R. L., Huynh, M., Bartlett, K. H., Fyfe, M., MacDougall, L., Boekhout, T., Kwon-Chung, K. J., & Meyer, W. (2004). A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). *Proceedings of the National Academy of Sciences*, 101(49), 17258–17263. <https://doi.org/10.1073/pnas.0402981101>

Kidd, S. E., Bach, P. J., Hingston, A. O., Mak, S., Chow, Y., MacDougall, L., Kronstad, J. W., & Bartlett, K. H. (2007). *Cryptococcus gattii* Dispersal Mechanisms, British Columbia, Canada. *Emerging Infectious Diseases*, 13(1), 51–57. <https://doi.org/10.3201/eid1301.060823>

Kono, I., & Himeno, K. (1997). A Novel Killer Yeast Effective on *Schizosaccharomyces pombe*. *Bioscience, Biotechnology, and Biochemistry*, 61(3), 563–564. <https://doi.org/10.1271/bbb.61.563>

Koufopanou, V., Hughes, J., Bell, G., & Burt, A. (2006). The spatial scale of genetic differentiation in a model organism: The wild yeast *Saccharomyces paradoxus*. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 361(1475), 1941–1946. <https://doi.org/10.1098/rstb.2006.1922>

Koufopanou, V., Lomas, S., Pronina, O., Almeida, P., Sampaio, J. P., Mousseau, T., Liti, G., & Burt, A. (2020). Population Size, Sex and Purifying Selection: Comparative Genomics of Two Sister Taxa of the Wild Yeast *Saccharomyces paradoxus*. *Genome Biology and Evolution*, 12(9), 1636–1645. <https://doi.org/10.1093/gbe/evaa141>

Kowallik, V., & Greig, D. (2016). A systematic forest survey showing an association of *Saccharomyces paradoxus* with oak leaf litter. *Environmental Microbiology Reports*, 8(5), 833–841. <https://doi.org/10.1111/1758-2229.12446>

Kowallik, V., Miller, E., & Greig, D. (2015). The interaction of *Saccharomyces paradoxus* with its natural competitors on oak bark. *Molecular Ecology*, 24(7), 1596–1610. <https://doi.org/10.1111/mec.13120>

Krogerus, K., Magalhães, F., Vidgren, V., & Gibson, B. (2015). New lager yeast strains generated by interspecific hybridization. *Journal of Industrial Microbiology and Biotechnology*, 42(5), 769–778. <https://doi.org/10.1007/s10295-015-1597-6>

Kurtzman, C. P. (2005). Description of *Komagataella phaffii* sp. Nov. And the transfer of *Pichia pseudopastoris* to the methylotrophic yeast genus *Komagataella*. *International Journal of Systematic and Evolutionary Microbiology*, 55(2), 973–976. <https://doi.org/10.1099/ijst.0.63491-0>

Kurtzman, C. P. (2009). Biotechnological strains of *Komagataella* (*Pichia*) *pastoris* are *Komagataella phaffii* as determined from multigene sequence analysis. *Journal of Industrial Microbiology & Biotechnology*, 36(11), 1435–1438. <https://doi.org/10.1007/s10295-009-0638-4>

Kurtzman, C. P. (2011a). *Komagataella* Y. Yamada, Matsuda, Maeda & Mikata (1995). In *The Yeasts* (pp. 491–495). Elsevier. <https://doi.org/10.1016/B978-0-444-52149-1.00037-9>

Kurtzman, C. P. (2011b). A new methanol assimilating yeast, *Ogataea parapolymorpha*, the ascosporic state of *Candida parapolymorpha*. *Antonie van Leeuwenhoek*, 100(3), 455–462. <https://doi.org/10.1007/s10482-011-9603-0>

Kurtzman, C. P. (2012). *Komagataella populi* sp. Nov. And *Komagataella ulmi* sp. Nov., two new methanol assimilating yeasts from exudates of deciduous trees. *Antonie van Leeuwenhoek*, 101(4), 859–868. <https://doi.org/10.1007/s10482-012-9702-6>

Kurtzman, C. P., Fell, J. W., & Boekhout, T. (2011). *The yeasts: A taxonomic study. Volume 1*. Elsevier Science. <http://site.ebrary.com/id/10440524>

Lachance, M.-A., & Starmer, W. T. (1998). Ecology and yeasts. In C. P. Kurtzman & J. W. Fell (Eds.), *The Yeasts, A Taxonomic Study* (pp. 21-30). Elsevier.

Lachance, M.-A., Metcalf, B. J., & Starmer, W. T. (1982). Yeasts from exudates of *Quercus*, *Ulmus*, *Populus*, and *Pseudotsuga*: New isolations and elucidation of some factors affecting ecological specificity. *Microbial Ecology*, 8(2), 191–198. <https://doi.org/10.1007/BF02010452>

Langdon, Q. K., Peris, D., Baker, E. P., Opulente, D. A., Nguyen, H.-V., Bond, U., Gonçalves, P., Sampaio, J. P., Libkind, D., & Hittinger, C. T. (2019). Fermentation innovation through complex hybridization of wild and domesticated yeasts. *Nature Ecology & Evolution*, 3(11), 1576–1586. <https://doi.org/10.1038/s41559-019-0998-8>

Langdon, Q. K., Peris, D., Eizaguirre, J. I., Opulente, D. A., Buh, K. V., Sylvester, K., Jarzyna, M., Rodríguez, M. E., Lopes, C. A., Libkind, D., & Hittinger, C. T. (2020). Postglacial migration shaped the genomic diversity and global distribution of the wild ancestor of lager-brewing hybrids. *PLOS Genetics*, 16(4), e1008680. <https://doi.org/10.1371/journal.pgen.1008680>

Leducq, J.-B., Charron, G., Samani, P., Dubé, A. K., Sylvester, K., James, B., Almeida, P., Sampaio, J. P., Hittinger, C. T., Bell, G., & Landry, C. R. (2014). Local climatic adaptation in a widespread microorganism. *Proceedings of the Royal Society B: Biological Sciences*, 281(1777), 20132472. <https://doi.org/10.1098/rspb.2013.2472>

Leducq, J.-B., Nielly-Thibault, L., Charron, G., Eberlein, C., Verta, J.-P., Samani, P., Sylvester, K., Hittinger, C. T., Bell, G., & Landry, C. R. (2016). Speciation driven by hybridization and chromosomal plasticity in a wild yeast. *Nature Microbiology*, 1(1), 15003. <https://doi.org/10.1038/nmicrobiol.2015.3>

Lee, C.-F., Yao, C.-H., Liu, Y.-R., Hsieh, C.-W., & Young, S.-S. (2009). *Lachancea dasiensis* sp. Nov., an ascosporeogenous yeast isolated from soil and leaves in Taiwan. *INTERNATIONAL JOURNAL OF SYSTEMATIC AND EVOLUTIONARY MICROBIOLOGY*, 59(7), 1818–1822. <https://doi.org/10.1099/ijns.0.008789-0>

Lee, T. J., Liu, Y.-C., Liu, W.-A., Lin, Y.-F., Lee, H.-H., Ke, H.-M., Huang, J.-P., Lu, M.-Y. J., Hsieh, C.-L., Chung, K.-F., Liti, G., & Tsai, I. J. (2021). Deep sampling of ancestral genetic diversity reveals *Saccharomyces cerevisiae* pre-domestication life histories [Preprint]. *Genomics*. <https://doi.org/10.1101/2021.09.07.459046>

Li, Y. F., Costello, J. C., Holloway, A. K., & Hahn, M. W. (2008). “REVERSE ECOLOGY” AND THE POWER OF POPULATION GENOMICS. *Evolution*, 62(12), 2984–2994. <https://doi.org/10.1111/j.1558-5646.2008.00486.x>

Libkind, D., Hittinger, C. T., Valerio, E., Goncalves, C., Dover, J., Johnston, M., Goncalves, P., & Sampaio, J. P. (2011). Microbe domestication and the identification of the wild genetic stock of lager-brewing yeast. *Proceedings of the National Academy of Sciences*, 108(35), 14539–14544. <https://doi.org/10.1073/pnas.1105430108>

Limtong, S., Kaewwichian, R., & Groenewald, M. (2013). *Ogataea kanchanaburiensis* sp. Nov. And *Ogataea wangdongensis* sp. Nov., two novel methylotrophic yeast species from phylloplane in Thailand. *Antonie van Leeuwenhoek*, 103(3), 551–558. <https://doi.org/10.1007/s10482-012-9837-5>

Lin, K., Lin, Y., Ho, M., Chen, Y., & Chung, W. (2021). Molecular epidemiology and phylogenetic analyses of environmental and clinical isolates of *Cryptococcus gattii* sensu lato in Taiwan. *Mycoses*, 64(3), 324–335. <https://doi.org/10.1111/myc.13195>

Liti, G. (2015). The fascinating and secret wild life of the budding yeast *S. cerevisiae*. *ELife*, 4, e05835. <https://doi.org/10.7554/eLife.05835>

Liti, G., Carter, D. M., Moses, A. M., Warringer, J., Parts, L., James, S. A., Davey, R. P., Roberts, I. N., Burt, A., Koufopanou, V., Tsai, I. J., Bergman, C. M., Bensasson, D., O'Kelly, M. J. T., van Oudenaarden, A., Barton, D. B. H., Bailes, E., Nguyen, A. N., Jones, M., ... Louis, E. J. (2009). Population genomics of domestic and wild yeasts. *Nature*, 458(7236), 337–341. <https://doi.org/10.1038/nature07743>

Liti, G., Peruffo, A., James, S. A., Roberts, I. N., & Louis, E. J. (2005). Inferences of evolutionary relationships from a population survey of LTR-retrotransposons and telomeric-associated sequences in the *Saccharomyces sensu stricto* complex. *Yeast*, 22(3), 177–192. <https://doi.org/10.1002/yea.1200>

Liu, X.-Z., Wang, Q.-M., Göker, M., Groenewald, M., Kachalkin, A. V., Lumbsch, H. T., Millanes, A. M., Wedin, M., Yurkov, A. M., Boekhout, T., & Bai, F.-Y. (2015). Towards an integrated phylogenetic classification of the Tremellomycetes. *Studies in Mycology*, 81, 85–147. <https://doi.org/10.1016/j.simyco.2015.12.001>

Lücking, R., Aime, M. C., Robbertse, B., Miller, A. N., Ariyawansa, H. A., Aoki, T., Cardinali, G., Crous, P. W., Druzhinina, I. S., Geiser, D. M., Hawksworth, D. L., Hyde, K. D., Irinyi, L., Jeewon, R., Johnston, P. R., Kirk, P. M., Malosso, E., May, T. W., Meyer, W., ... Schoch, C. L. (2020). Unambiguous identification of fungi: Where do we stand and how accurate and precise is fungal DNA barcoding? *IMA Fungus*, 11(1), 14. <https://doi.org/10.1186/s43008-020-00033-z>

Lücking, R., Aime, M. C., Robbertse, B., Miller, A. N., Aoki, T., Ariyawansa, H. A., Cardinali, G., Crous, P. W., Druzhinina, I. S., Geiser, D. M., Hawksworth, D. L., Hyde, K. D., Irinyi, L., Jeewon, R., Johnston, P. R., Kirk, P. M., Malosso, E., May, T. W., Meyer, W., ... Schoch, C. L. (2021). Fungal taxonomy and sequence-based nomenclature. *Nature Microbiology*, 6(5), 540–548. <https://doi.org/10.1038/s41564-021-00888-x>

MacDougall, L., Kidd, S. E., Galanis, E., Mak, S., Leslie, M. J., Cieslak, P. R., Kronstad, J. W., Morshed, M. G., & Bartlett, K. H. (2007). Spread of *Cryptococcus gattii* in British Columbia, Canada, and Detection in the Pacific Northwest, USA. *Emerging Infectious Diseases*, 13(1), 42–50. <https://doi.org/10.3201/eid1301.060827>

Magalhães, F., Calton, A., Heiniö, R.-L., & Gibson, B. (2021). Frozen-dough baking potential of psychrotolerant *Saccharomyces* species and derived hybrids. *Food Microbiology*, 94, 103640. <https://doi.org/10.1016/j.fm.2020.103640>

Magalhães, F., Krogerus, K., Castillo, S., Ortiz-Julien, A., Dequin, S., & Gibson, B. (2017a). Exploring the potential of *Saccharomyces eubayanus* as a parent for new interspecies hybrid strains in winemaking. *FEMS Yeast Research*, 17(5). <https://doi.org/10.1093/femsyr/fox049>

Magalhães, F., Krogerus, K., Vidgren, V., Sandell, M., & Gibson, B. (2017b). Improved cider fermentation performance and quality with newly generated *Saccharomyces cerevisiae* × *Saccharomyces eubayanus* hybrids. *Journal of Industrial Microbiology and Biotechnology*, 44(8), 1203–1213. <https://doi.org/10.1007/s10295-017-1947-7>

Magalhães, KT., (2011) Brazilian kefir: structure, microbial communities and chemical composition. *Braz J Microbiol*. 10.1590/S1517-838220110002000034

Magwene, P. M., Kayıkçı, Ö., Granek, J. A., Reininga, J. M., Scholl, Z., & Murray, D. (2011). Outcrossing, mitotic recombination, and life-history trade-offs shape genome evolution in *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences*, 108(5), 1987–1992. <https://doi.org/10.1073/pnas.1012544108>

Mardones, W., Villarroel, C. A., Abarca, V., Urbina, K., Peña, T. A., Molinet, J., Nespolo, R. F., & Cubillos, F. A. (2021). Rapid selection response to ethanol in *Saccharomyces eubayanus* emulates the domestication process under brewing conditions. *Microbial Biotechnology*, 1751-7915.13803. <https://doi.org/10.1111/1751-7915.13803>

Mardones, W., Villarroel, C. A., Krogerus, K., Tapia, S. M., Urbina, K., Oporto, C. I., O'Donnell, S., Minebois, R., Nespolo, R., Fischer, G., Querol, A., Gibson, B., & Cubillos, F. A. (2020). Molecular profiling of beer wort fermentation diversity across natural *Saccharomyces eubayanus* isolates. *Microbial Biotechnology*, 13(4), 1012–1025. <https://doi.org/10.1111/1751-7915.13545>

Marsh, A. J., O'Sullivan, O., Hill, C., Ross, R. P., & Cotter, P. D. (2014). Sequence-based analysis of the bacterial and fungal compositions of multiple kombucha (tea fungus) samples. *Food Microbiology*, 38, 171–178. <https://doi.org/10.1016/j.fm.2013.09.003>

Martinović, T., Odriozola, I., Mašínová, T., Doreen Bahnmann, B., Kohout, P., Sedlák, P., Merunková, K., Větrovský, T., Tomšovský, M., Ovaskainen, O., & Baldrian, P. (2021). Temporal turnover of the soil microbiome composition is guild-specific. *Ecology Letters*, 24(12), 2726–2738. <https://doi.org/10.1111/ele.13896>

Mašínová, T., Pontes, A., Carvalho, C., Sampaio, J. P., & Baldrian, P. (2017). *Libkindia masarykiana* gen. Et sp. Nov., *Yurkovia mendeliana* gen. Et sp. Nov. And *Leucosporidium krtinense* f.a. Sp. Nov., isolated from temperate forest soils. *International Journal of Systematic and Evolutionary Microbiology*, 67(4), 902–908. <https://doi.org/10.1099/ijsem.0.001707>

Mašínová, T., Yurkov, A., & Baldrian, P. (2018). Forest soil yeasts: Decomposition potential and the utilization of carbon sources. *Fungal Ecology*, 34, 10–19. <https://doi.org/10.1016/j.funeco.2018.03.005>

Medina, A., Mohale, S., Samsudin, N. I. P., Rodriguez-Sixtos, A., Rodriguez, A., & Magan, N. (2017). Biocontrol of mycotoxins: Dynamics and mechanisms of action. *Current Opinion in Food Science*, 17, 41–48. <https://doi.org/10.1016/j.cofs.2017.09.008>

Mertens, S., Steensels, J., Saels, V., De Rouck, G., Aerts, G., & Verstrepen, K. J. (2015). A Large Set of Newly Created Interspecific *Saccharomyces* Hybrids Increases Aromatic Diversity in Lager Beers. *Applied and Environmental Microbiology*, 81(23), 8202–8214. <https://doi.org/10.1128/AEM.02464-15>

Mesquita et al (2013). The molecular phylogenetic diversity of bacteria and fungi associated with the cerrado soil from different regions of Minas Gerais. Brazil. *Int. J. Microbiol. Res.*, 4, 119-131.

Meyen, F. J. F. (1839). *A Report on the Progress of Vegetable Physiology During the Year 1837*. Taylor, London

Miller, M. W., Phaff, H. J., & Snyder, H. E. (1962). On the occurrence of various species of yeast in nature. *Mycopathologia et Mycologia Applicata*, 16(1), 1–18. <https://doi.org/10.1007/BF02136176>

Mittelbach, M., Yurkov, A. M., Nocentini, D., Nepi, M., Weigend, M., & Begerow, D. (2015). Nectar sugars and bird visitation define a floral niche for basidiomycetous yeast on the Canary Islands. *BMC Ecology*, 15(1), 2. <https://doi.org/10.1186/s12898-015-0036-x>

Mozzachiodi, S., Tattini, L., Llored, A., Irizar, A., Škofljanc, N., D'Angiolo, M., De Chiara, M., Barré, B. P., Yue, J.-X., Lutazi, A., Loeillet, S., Laureau, R., Marsit, S., Stenberg, S., Albaud, B., Persson, K., Legras, J.-L., Dequin, S., Warringer, J., ... Liti, G. (2021). Aborting meiosis allows recombination in sterile diploid yeast hybrids. *Nature Communications*, 12(1), 6564. <https://doi.org/10.1038/s41467-021-26883-8>

Myster (2012). *Ecotones Between Forest and Grassland*. Springer Science+Business Media. <https://doi.org/10.1007/978-1-4614-3797-0>

Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403(6772), 853–858. <https://doi.org/10.1038/35002501>

Naseeb, S., James, S. A., Alsammar, H., Michaels, C. J., Gini, B., Nueno-Palop, C., Bond, C. J., McGhie, H., Roberts, I. N., & Delneri, D. (2017). *Saccharomyces jurei* sp. Nov., isolation and genetic identification of a novel yeast species from *Quercus robur*. *International Journal of Systematic and Evolutionary Microbiology*, 67(6), 2046–2052. <https://doi.org/10.1099/ijsem.0.002013>

Naseeb, S., Alsammar, H., Burgis, T., Donaldson, I., Knyazev, N., Knight, C., & Delneri, D. (2018). Whole genome sequencing, de novo assembly and phenotypic profiling for the new budding yeast species *Saccharomyces jurei*. *G3 Genes|Genomes|Genetics*, 8(9), 2967–2977. <https://doi.org/10.1534/g3.118.200476>

Naseeb, S., Visinoni, F., Hu, Y., Hinks Roberts, A. J., Maslowska, A., Walsh, T., Smart, K. A., Louis, E. J., & Delneri, D. (2021). Restoring fertility in yeast hybrids: Breeding and quantitative genetics of beneficial traits. *Proceedings of the National Academy of Sciences*, 118(38), e2101242118. <https://doi.org/10.1073/pnas.2101242118>

Naumov, G. (1996). Genetic identification of biological species in the *Saccharomyces* sensu stricto complex. *Journal of Industrial Microbiology & Biotechnology*, 17(3–4), 295–302. <https://doi.org/10.1007/BF01574704>

Naumov, G. I. (2015). The yeast *Komagataella*: A genetic genus in accordance with interspecies hybridization. *Microbiology*, 84(4), 538–543. <https://doi.org/10.1134/S0026261715040141>

Naumov, G. I., James, S. A., Naumova, E. S., Louis, E. J., & Roberts, I. N. (2000). Three new species in the *Saccharomyces* sensu stricto complex: *Saccharomyces cariocanus*, *Saccharomyces kudriavzevii* and *Saccharomyces mikatae*. *International Journal of Systematic and Evolutionary Microbiology*, 50(5), 1931–1942. <https://doi.org/10.1099/00207713-50-5-1931>

Naumov, G. I., Kondratieva, V. I., Meshcheryakova, E. V., & Naumova, E. S. (2016). Taxonomic genetics of methylotrophic yeast genus *Komagataella*: New biological species *K. kurtzmanii*. *Russian Journal of Genetics*, 52(4), 378–382. <https://doi.org/10.1134/S1022795416030108>

Naumov, G. I., Lee, C.-F., & Naumova, E. S. (2013). Molecular genetic diversity of the *Saccharomyces* yeasts in Taiwan: *Saccharomyces arboricola*, *Saccharomyces cerevisiae* and *Saccharomyces kudriavzevii*. *Antonie van Leeuwenhoek*, 103(1), 217–228. <https://doi.org/10.1007/s10482-012-9803-2>

Naumov, G. I., Naumova, E. S., & Boundy-Mills, K. L. (2018). Description of *Komagataella mondaviorum* sp. Nov., a new sibling species of *Komagataella* (*Pichia*) *pastoris*. *Antonie van Leeuwenhoek*, 111(7), 1197–1207. <https://doi.org/10.1007/s10482-018-1028-6>

Naumov, G. I., Naumova, E. S., & Sniegowski, P. D. (1998). *Saccharomyces paradoxus* and *Saccharomyces cerevisiae* are associated with exudates of North American oaks. *Canadian Journal of Microbiology*, 44(11), 1045–1050. <https://doi.org/10.1139/w98-104>

Naumov, G. I., Naumova, E. S., Tyurin, O. V., & Kozlov, D. G. (2013). *Komagataella kurtzmanii* sp. Nov., a new sibling species of *Komagataella* (*Pichia*) *pastoris* based on multigene sequence analysis. *Antonie van Leeuwenhoek*, 104(3), 339–347. <https://doi.org/10.1007/s10482-013-9956-7>

Naumova, E. S., Boundy-Mills, K. L., & Naumov, G. I. (2020). Phylogenetics and Biogeography of Biotechnologically Important Methylotrophic Yeasts *Komagataella*. *Microbiology*, 89(3), 294–300. <https://doi.org/10.1134/S002626172003011X>

Naumova, E. S., Serpova, E. V., Korshunova, I. V., & Naumov, G. I. (2007). Molecular genetic characterization of the yeast *Lachancea kluyveri*. *Microbiology*, 76(3), 317–323. <https://doi.org/10.1134/S0026261707030083>

Nespolo, R. F., Villarroel, C. A., Oporto, C. I., Tapia, S. M., Vega-Macaya, F., Urbina, K., De Chiara, M., Mozzachiodi, S., Mikhalev, E., Thompson, D., Larrondo, L. F., Saenz-Agudelo, P., Liti, G., & Cubillos, F. A. (2020).

An Out-of-Patagonia migration explains the worldwide diversity and distribution of *Saccharomyces eubayanus* lineages. *PLOS Genetics*, 16(5), e1008777. <https://doi.org/10.1371/journal.pgen.1008777>

Nieuwenhuis, B. P. S., & James, T. Y. (2016). The frequency of sex in fungi. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1706), 20150540. <https://doi.org/10.1098/rstb.2015.0540>

Nikulin, J., Vidgren, V., Krogerus, K., Magalhães, F., Valkeemäki, S., Kangas-Heiska, T., & Gibson, B. (2020). Brewing potential of the wild yeast species *Saccharomyces paradoxus*. *European Food Research and Technology*, 246(11), 2283–2297. <https://doi.org/10.1007/s00217-020-03572-2>

Nikulin, J., Krogerus, K., & Gibson, B. (2018). Alternative *Saccharomyces* interspecies hybrid combinations and their potential for low-temperature wort fermentation: Alternative *Saccharomyces* interspecies hybrids for brewing. *Yeast*, 35(1), 113–127. <https://doi.org/10.1002/yea.3246>

Nizovoy, P., Bellora, N., Haridas, S., Sun, H., Daum, C., Barry, K., Grigoriev, I. V., Libkind, D., Connell, L. B., & Moliné, M. (2021). Unique genomic traits for cold adaptation in *Naganishia vishniacii*, a polyextremophile yeast isolated from Antarctica. *FEMS Yeast Research*, 21(1), foaa056. <https://doi.org/10.1093/femsyr/foaa056>

Noh, J., Echeverría, C., Pauchard, A., & Cuenca, P. (2019). Extinction debt in a biodiversity hotspot: The case of the Chilean Winter Rainfall-Valdivian Forests. *Landscape and Ecological Engineering*, 15(1), 1–12. <https://doi.org/10.1007/s11355-018-0352-3>

O'Brien, C. E., Oliveira-Pacheco, J., Ó Cinnéide, E., Haase, M. A. B., Hittinger, C. T., Rogers, T. R., Zaragoza, O., Bond, U., & Butler, G. (2021). Population genomics of the pathogenic yeast *Candida tropicalis* identifies hybrid isolates in environmental samples. *PLOS Pathogens*, 17(3), e1009138. <https://doi.org/10.1371/journal.ppat.1009138>

Olson, D. M., & Dinerstein, E. (2002). The Global 200: Priority Ecoregions for Global Conservation. *Annals of the Missouri Botanical Garden*, 89(2), 199. <https://doi.org/10.2307/3298564>

Ono, J., & Greig, D. (2020). A *Saccharomyces* paradox: Chromosomes from different species are incompatible because of anti-recombination, not because of differences in number or arrangement. *Current Genetics*, 66(3), 469–474. <https://doi.org/10.1007/s00294-019-01038-x>

Ono, J., Greig, D., & Boynton, P. J. (2020). Defining and Disrupting Species Boundaries in *Saccharomyces*. *Annual Review of Microbiology*, 74(1), 477–495. <https://doi.org/10.1146/annurev-micro-021320-014036>

Opulente, D. A., Rollinson, E. J., Bernick-Roehr, C., Hulfachor, A. B., Rokas, A., Kurtzman, C. P., & Hittinger, C. T. (2018). Factors driving metabolic diversity in the budding yeast subphylum. *BMC Biology*, 16(1), 26. <https://doi.org/10.1186/s12915-018-0498-3>

Paget, C. M., Schwartz, J., & Delneri, D. (2014). Environmental systems biology of cold-tolerant phenotype in *Saccharomyces* species adapted to grow at different temperatures. *Molecular Ecology*, 23(21), 5241–5257. <https://doi.org/10.1111/mec.12930>

Passer, A. R., Coelho, M. A., Billmyre, R. B., Nowrouzian, M., Mittelbach, M., Yurkov, A. M., Averette, A. F., Cuomo, C. A., Sun, S., & Heitman, J. (2019). Genetic and Genomic Analyses Reveal Boundaries between Species Closely Related to *Cryptococcus* Pathogens. *MBio*, 10(3). <https://doi.org/10.1128/mBio.00764-19>

Pasteur, L. (1876) *Études sur la Bière, ses Maladies, Causes qui les Provoquent, Procédé pour la Rendre Inaltérable, avec une Théorie Nouvelle de la Fermentation*. Gauthier-Villars, Paris.

Pereira, L. F., Costa, C. R. L., Brasileiro, B. T. R. V., & de Moraes, M. A. (2011). *Lachancea mirantina* sp. Nov., an ascomycetous yeast isolated from the cachaça fermentation process. *International Journal of Systematic and Evolutionary Microbiology*, 61(4), 989–992. <https://doi.org/10.1099/ijss.0.020008-0>

Pérez-Torrado, R., Barrio, E., & Querol, A. (2018). Alternative yeasts for winemaking: *Saccharomyces* non-cerevisiae and its hybrids. *Critical Reviews in Food Science and Nutrition*, 58(11), 1780–1790. <https://doi.org/10.1080/10408398.2017.1285751>

Peris, D., Alexander, W. G., Fisher, K. J., Moriarty, R. V., Basuino, M. G., Ubbelohde, E. J., Wrobel, R. L., & Hittinger, C. T. (2020). Synthetic hybrids of six yeast species. *Nature Communications*, 11(1), 2085. <https://doi.org/10.1038/s41467-020-15559-4>

Peris, D., Langdon, Q. K., Moriarty, R. V., Sylvester, K., Bontrager, M., Charron, G., Leducq, J.-B., Landry, C. R., Libkind, D., & Hittinger, C. T. (2016). Complex Ancestries of Lager-Brewing Hybrids Were Shaped by Standing Variation in the Wild Yeast *Saccharomyces eubayanus*. *PLOS Genetics*, 12(7), e1006155. <https://doi.org/10.1371/journal.pgen.1006155>

Peris, D., Sylvester, K., Libkind, D., Gonçalves, P., Sampaio, J. P., Alexander, W. G., & Hittinger, C. T. (2014). Population structure and reticulate evolution of *Saccharomyces eubayanus* and its lager-brewing hybrids. *Molecular Ecology*, 23(8), 2031–2045. <https://doi.org/10.1111/mec.12702>

Péter, G., Nagy, E. S., & Dlauchy, D. (2019). Systematics, Diversity and Ecology of the Genus *Yarrowia* and the Methanol-Assimilating Yeasts. In A. Sibirny (Ed.), *Non-conventional Yeasts: From Basic Research to Application* (pp. 297–339). Springer International Publishing. https://doi.org/10.1007/978-3-030-21110-3_9

Péter, G., Takashima, M., & Čadež, N. (2017). Yeast Habitats: Different but Global. In P. Buzzini, M.-A. Lachance, & A. Yurkov (Eds.), *Yeasts in Natural Ecosystems: Ecology* (pp. 39–71). Springer International Publishing. https://doi.org/10.1007/978-3-319-61575-2_2

Peter, J., De Chiara, M., Friedrich, A., Yue, J.-X., Pflieger, D., Bergström, A., Sigwalt, A., Barre, B., Freel, K., Llored, A., Cruaud, C., Labadie, K., Aury, J.-M., Istace, B., Lebrigand, K., Barbry, P., Engelen, S., Lemainque, A., Wincker, P., ... Schacherer, J. (2018). Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature*, 556(7701), 339–344. <https://doi.org/10.1038/s41586-018-0030-5>

Phaff, H. J., Miller, M. W., & Shifrine, M. (1956). The taxonomy of yeasts isolated from *Drosophila* in the Yosemite region of California. *Antonie van Leeuwenhoek*, 22(1), 145–161. <https://doi.org/10.1007/BF02538322>

Phaff, H. J., et al. (1972) Acomparative study of the yeast flora associated with treeson the Japanese islands and on the west coast of North America. In: Terui G (ed) *Fermentation TechnologyToday*. Society of Fermentation Technology, Osaka,pp 759–774

Põlme, S., Abarenkov, K., Henrik Nilsson, R., Lindahl, B. D., Clemmensen, K. E., Kauserud, H., Nguyen, N., Kjøller, R., Bates, S. T., Baldrian, P., Frøslev, T. G., Adojaan, K., Vizzini, A., Suija, A., Pfister, D., Baral, H.-O., Järv, H., Madrid, H., Nordén, J., ... Tedersoo, L. (2020). FungalTraits: A user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Diversity*, 105(1), 1–16. <https://doi.org/10.1007/s13225-020-00466-2>

Porter, T. J., Divol, B., & Setati, M. E. (2019). *Lachancea* yeast species: Origin, biochemical characteristics and oenological significance. *Food Research International*, 119, 378–389. <https://doi.org/10.1016/j.foodres.2019.02.003>

Pulvirenti, A., Nguyen, H. V., Caggia, C., Giudici, P., Rainieri, S., Zambonelli, C., (2000). *Saccharomyces uvarum*, a proper species within *Saccharomyces* sensu stricto. *FEMS Microbiology Letters*, 192(2), 191–96.

Rai, S. K., Atwood-Moore, A., & Levin, H. L. (2018). Duplication and Transformation of the *Schizosaccharomyces pombe* Collection of Deletion Strains. In T. L. Singleton (Ed.), *Schizosaccharomyces pombe* (Vol. 1721, pp. 197–215). Springer New York. https://doi.org/10.1007/978-1-4939-7546-4_18

Reuter, M., Bell, G., & Greig, D. (2007). Increased outbreeding in yeast in response to dispersal by an insect vector. *Current Biology*, 17(3), R81–R83. <https://doi.org/10.1016/j.cub.2006.11.059>

Robinson, H. A., Pinharanda, A., & Bensasson, D. (2016). Summer temperature can predict the distribution of wild yeast populations. *Ecology and Evolution*, 6(4), 1236–1250. <https://doi.org/10.1002/ece3.1919>

Rodríguez, M. E., Pérez-Través, L., Sangorrín, M. P., Barrio, E., & Lopes, C. A. (2014). *Saccharomyces eubayanus* and *Saccharomyces uvarum* associated with the fermentation of Araucaria araucana seeds in Patagonia. *FEMS Yeast Research*, 14(6), 948–965. <https://doi.org/10.1111/1567-1364.12183>

Romano, P., & Suzzi, G. (1993). Potential use for *Zygosaccharomyces* species in winemaking. *Journal of Wine Research*, 4(2), 87–94. <https://doi.org/10.1080/09571269308717955>

Romano, P., & Suzzi, G. (1993) Higher alcohol and acetoin production by *Zygosaccharomyces* wine yeasts. *Journal of Applied Bacteriology*, 75, 541–545. <https://doi.org/10.1111/j.1365-2672.1993.tb01592.x>

Romero-Olivares, A. L., Morrison, E. W., Pringle, A., & Frey, S. D. (2021). Correction to: Linking Genes to Traits in Fungi. *Microbial Ecology*, 82(1), 156. <https://doi.org/10.1007/s00248-021-01776-x>

Ropars, J., Maufrais, C., Diogo, D., Marcet-Houben, M., Perin, A., Sertour, N., Mosca, K., Permal, E., Laval, G., Bouchier, C., Ma, L., Schwartz, K., Voelz, K., May, R. C., Poulain, J., Battail, C., Wincker, P., Borman, A. M., Chowdhary, A., ... d'Enfert, C. (2018). Gene flow contributes to diversification of the major fungal pathogen *Candida albicans*. *Nature Communications*, 9(1), 2253. <https://doi.org/10.1038/s41467-018-04787-4>

Ruderfer, D. M., Pratt, S. C., Seidel, H. S., & Kruglyak, L. (2006). Population genomic analysis of outcrossing and recombination in yeast. *Nature Genetics*, 38(9), 1077–1081. <https://doi.org/10.1038/ng1859>

Salvadó, Z., Arroyo-López, F. N., Guillamón, J. M., Salazar, G., Querol, A., & Barrio, E. (2011). Temperature Adaptation Markedly Determines Evolution within the Genus *Saccharomyces*. *Applied and Environmental Microbiology*, 77(7), 2292–2302. <https://doi.org/10.1128/AEM.01861-10>

Sampaio, J. P., & Gonçalves, P. (2008). Natural Populations of *Saccharomyces kudriavzevii* in Portugal Are Associated with Oak Bark and Are Sympatric with *S. cerevisiae* and *S. paradoxus*. *Applied and Environmental Microbiology*, 74(7), 2144–2152. <https://doi.org/10.1128/AEM.02396-07>

Sampaio, J. P., & Gonçalves, P. (2017). Biogeography and Ecology of the Genus *Saccharomyces*. In P. Buzzini, M.-A. Lachance, & A. Yurkov (Eds.), *Yeasts in Natural Ecosystems: Ecology* (pp. 131–153). Springer International Publishing. https://doi.org/10.1007/978-3-319-61575-2_5

Santos, A. R. O., Faria, E. S., Lachance, M.-A., & Rosa, C. A. (2015). *Ogataea mangiferae* sp. Nov., a methylotrophic yeast isolated from mango leaves. *International Journal of Systematic and Evolutionary Microbiology*, 65(Pt_6), 1855–1859. <https://doi.org/10.1099/ijss.0.000194>

Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W., Fungal Barcoding Consortium, Fungal Barcoding Consortium Author List, Bolchacova, E., Voigt, K., Crous, P. W., Miller, A. N., Wingfield, M. J., Aime, M. C., An, K.-D., Bai, F.-Y., Barreto, R. W., Begerow, D., ... Schindel, D. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences*, 109(16), 6241–6246. <https://doi.org/10.1073/pnas.1117018109>

Senses-Ergul, S., Ágoston, R., Belák, Á., & Deák, T. (2006). Characterization of some yeasts isolated from foods by traditional and molecular tests. *International Journal of Food Microbiology*, 108(1), 120–124. <https://doi.org/10.1016/j.ijfoodmicro.2005.10.014>

Shen, X.-X., Opulente, D. A., Kominek, J., Zhou, X., Steenwyk, J. L., Buh, K. V., Haase, M. A. B., Wisecaver, J. H., Wang, M., Doering, D. T., Boudouris, J. T., Schneider, R. M., Langdon, Q. K., Ohkuma, M., Endoh, R., Takashima, M., Manabe, R., Čadež, N., Libkind, D., ... Rokas, A. (2018). Tempo and Mode of Genome Evolution in the Budding Yeast Subphylum. *Cell*, 175(6), 1533–1545.e20. <https://doi.org/10.1016/j.cell.2018.10.023>

Sniegowski, P. D., Dombrowski, P. G., & Fingerman, E. (2002). *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* coexist in a natural woodland site in North America and display different levels of reproductive isolation from European conspecifics. *FEMS Yeast Research*, 1(4), 299–306. <https://doi.org/10.1111/j.1567-1364.2002.tb00048.x>

Spencer et al (1995) Yeasts associated with algarrobo trees (*Prosopisspp.*) in northwest Argentina: a preliminary report. *J IndustrialMicrobiol* 14:472–474

Spencer et al (1996) Yeasts associated with pods and exudates of algarrobo trees (*Prosopisspp.*) and species of columnar cacti in northwest Argentina. *App Microbiol Biotechnol* 44:736–739

Spurley, W. J., Fisher, K. J., Langdon, Q. K., Buh, K. V., Jarzyna, M., Haase, M. A. B., Sylvester, K., Moriarty, R. V., Rodriguez, D., Sheddan, A., Wright, S., Sorlie, L., Hulfachor, A. B., Opulente, D. A., & Hittinger, C. T. (2021). Substrate, temperature, and geographical patterns among nearly 2000 natural yeast isolates. *Yeast*, yea.3679. <https://doi.org/10.1002/yea.3679>

Starmer, W. T., & Lachance, M. A. (2011). Yeast ecology. In C. P. Kurtzman, J. W. Fell, & T. Boekhout (Eds.), *The yeasts, a taxonomic study*, (pp65–83). Elsevier.

Stefanini, I., Dapporto, L., Berná, L., Polzinelli, M., Turillazzi, S., & Cavalieri, D. (2016). Social wasps are a *Saccharomyces* mating nest. *Proceedings of the National Academy of Sciences*, 113(8), 2247–2251. <https://doi.org/10.1073/pnas.1516453113>

Stefanini, I. (2018). Yeast-insect associations: It takes guts. *Yeast* (Chichester, England), 35(4), 315–330. <https://doi.org/10.1002/yea.3309>

Strauss, M. L. A., Jolly, N. P., Lambrechts, M. G., & van Rensburg, P. (2001). Screening for the production of extracellular hydrolytic enzymes by non-*Saccharomyces* wine yeasts. *Journal of Applied Microbiology*, 91(1), 182–190. <https://doi.org/10.1046/j.1365-2672.2001.01379.x>

Sweeney, J., Kuehne, H., & Sniegowski, P. (2004). Sympatric natural and populations have different thermal growth profiles. *FEMS Yeast Research*, 4(4–5), 521–525. [https://doi.org/10.1016/S1567-1356\(03\)00171-5](https://doi.org/10.1016/S1567-1356(03)00171-5)

Sylvester, K., Wang, Q.-M., James, B., Mendez, R., Hulfachor, A. B., & Hittinger, C. T. (2015). Temperature and host preferences drive the diversification of *Saccharomyces* and other yeasts: A survey and the discovery of eight new yeast species. *FEMS Yeast Research*, 15(3). <https://doi.org/10.1093/femsyr/fov002>

Tedersoo, L., May, T. W., & Smith, M. E. (2010). Ectomycorrhizal lifestyle in fungi: Global diversity, distribution, and evolution of phylogenetic lineages. *Mycorrhiza*, 20(4), 217–263. <https://doi.org/10.1007/s00572-009-0274-x>

Tilakaratna, V., & Bensasson, D. (2017). Habitat Predicts Levels of Genetic Admixture in *Saccharomyces cerevisiae*. *G3 Genes|Genomes|Genetics*, 7(9), 2919–2929. <https://doi.org/10.1534/g3.117.041806>

Tláskal, V., Brabcová, V., Větrovský, T., López-Mondéjar, R., Monteiro, L.M.O., Saraiva, J.P., da Rocha, U.N., & Baldrian, P. (2021). Metagenomes, metatranscriptomes and microbiomes of naturally decomposing deadwood. *Scientific Data* 8, 198. <https://doi.org/10.1038/s41597-021-00987-8>

Treseder, K. K., & Lennon, J. T. (2015). Fungal Traits That Drive Ecosystem Dynamics on Land. *Microbiology and Molecular Biology Reviews*, 79(2), 243–262. <https://doi.org/10.1128/MMBR.00001-15>

Tsai, I. J., Bensasson, D., Burt, A., & Koufopanou, V. (2008). Population genomics of the wild yeast *Saccharomyces paradoxus*: Quantifying the life cycle. *Proceedings of the National Academy of Sciences*, 105(12), 4957–4962. <https://doi.org/10.1073/pnas.0707314105>

Tschopp, J. F., Brust, P. F., Cregg, J. M., Stillman, C. A., & Gingeras, T. R. (1987). Expression of the lacZ gene from two methanol-regulated promoters in *Pichia pastoris*. *Nucleic Acids Research*, 15(9), 3859–3876. <https://doi.org/10.1093/nar/15.9.3859>

Tzanetakis et al 1998 Yeasts of the surface microflora of Feta cheese.

Varela, C., Sundstrom, J., Cuijvers, K., Jiranek, V., & Borneman, A. (2020). Discovering the indigenous microbial communities associated with the natural fermentation of sap from the cider gum *Eucalyptus gunnii*. *Scientific Reports*, 10(1), 14716. <https://doi.org/10.1038/s41598-020-71663-x>

Větrovský, T., Kolařík, M., Žifčáková, L., Zelenka, T., & Baldrian, P. (2016). The rpb2 gene represents a viable alternative molecular marker for the analysis of environmental fungal communities. *Molecular Ecology Resources* 16, 388-401. <https://doi.org/10.1111/1755-0998.12456>

Větrovský, T., Kohout, P., Kopecký, M., Machac, A., Man, M., Bahnmann, B. D., Brabcová, V., Choi, J., Meszárošová, L., Human, Z. R., Lepinay, C., Lladó, S., López-Mondéjar, R., Martinović, T., Mašínová, T., Morais, D., Navrátilová, D., Odriozola, I., Štursová, M., ... Baldrian, P. (2019). A meta-analysis of global fungal distribution reveals climate-driven patterns. *Nature Communications*, 10(1), 5142. <https://doi.org/10.1038/s41467-019-13164-8>

Větrovský, T., Morais, D., Kohout, P., Lepinay, C., Algora, C., Awokunle Hollá, S., Bahnmann, B. D., Bílohnědá, K., Brabcová, V., D'Alò, F., Human, Z. R., Jomura, M., Kolařík, M., Kvasničková, J., Lladó, S., López-Mondéjar, R., Martinović, T., Mašínová, T., Meszárošová, L., ... Baldrian, P. (2020). GlobalFungi, a global database of fungal occurrences from high-throughput-sequencing metabarcoding studies. *Scientific Data*, 7(1), 228. <https://doi.org/10.1038/s41597-020-0567-7>

Villarreal, P., Quintrel, P. A., Olivares-Muñoz, S., Ruiz, J. J., Nespolo, R. F., & Cubillos, F. A. (2021). Identification of new ethanol-tolerant yeast strains with fermentation potential from central Patagonia. *Yeast*, yea.3662. <https://doi.org/10.1002/yea.3662>

Vishniac, H. S. (1995). Simulated in situ competitive ability and survival of a representative soil yeast, *Cryptococcus albidus*. *Microbial Ecology*, 30(3). <https://doi.org/10.1007/BF00171937>

Vishniac, H. S. (2006). A Multivariate Analysis of Soil Yeasts Isolated from a Latitudinal Gradient. *Microbial Ecology*, 52(1), 90–103. <https://doi.org/10.1007/s00248-006-9066-4>

Vuillemin, P. (1901). Les blastomycètes pathogènes. *Rev Gen Sci Pures Appl*, 12, 732-751.

Wang, Q.-M., Liu, W.-Q., Liti, G., Wang, S.-A., & Bai, F.-Y. (2012). Surprisingly diverged populations of *Saccharomyces cerevisiae* in natural environments remote from human activity. *Molecular Ecology*, 21(22), 5404–5417. <https://doi.org/10.1111/j.1365-294X.2012.05732.x>

Wang, S.-A., & Bai, F.-Y. (2008). *Saccharomyces arboricolus* sp. Nov., a yeast species from tree bark. *International Journal of Systematic and Evolutionary Microbiology*, 58(2), 510–514. <https://doi.org/10.1099/ij.s.0.65331-0>

Wegner 1983 Biochemical conversions by yeast fermentation at high cell densities. United States Pat. 4 329–414. United States Pat. 4414329.

White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, p 315–322. In Innis MA, Gelfand DH, Sninsky JJ, White TJ (ed), *PCR protocols: a guide to methods and applications*. Academic Press, Inc, New York, NY.

Wojtowicz, M., Chrzanowska, J., Juszczak, P., Skiba, A., & Gdula, A. (2001). Identification and biochemical characteristics of yeast microflora of Rokpol cheese. *International Journal of Food Microbiology*, 69(1–2), 135–140. [https://doi.org/10.1016/S0168-1605\(01\)00582-7](https://doi.org/10.1016/S0168-1605(01)00582-7)

Wu, L., Sun, Q., Sugawara, H., Yang, S., Zhou, Y., McCluskey, K., Vasilenko, A., Suzuki, K.-I., Ohkuma, M., Lee, Y., Robert, V., Ingsriswang, S., Guissart, F., Philippe, D., & Ma, J. (2013). Global catalogue of microorganisms (gcm): A comprehensive database and information retrieval, analysis, and visualization system for microbial resources. *BMC Genomics*, 14(1), 933. <https://doi.org/10.1186/1471-2164-14-933>

Xia, W., Nielly-Thibault, L., Charron, G., Landry, C. R., Kasimer, D., Anderson, J. B., & Kohn, L. M. (2017). Population genomics reveals structure at the individual, host-tree scale and persistence of genotypic variants of the undomesticated yeast *Saccharomyces paradoxus* in a natural woodland. *Molecular Ecology*, 26(4), 995–1007. <https://doi.org/10.1111/mec.13954>

Yurkov, A. M. (2005). First Isolation of the Yeast *Saccharomyces paradoxus* in Western Siberia. *Microbiology*, 74(4), 459–462. <https://doi.org/10.1007/s11021-005-0089-5>

Yurkov, A. & Pozo, M. I. (2017). Yeast community composition and structure. In P. Buzzini, M.-A. Lachance, & A. Yurkov (Eds.), *Yeasts in Natural Ecosystems: Ecology* (pp. 70–100). Springer International Publishing.

Yurkov, A. (2017). Yeasts in Forest Soils. In P. Buzzini, M.-A. Lachance, & A. Yurkov (Eds.), *Yeasts in Natural Ecosystems: Diversity* (pp. 87–116). Springer International Publishing. https://doi.org/10.1007/978-3-319-62683-3_3

Yurkov, A., Inácio, J., Chernov, I. Y., & Fonseca, Á. (2015). Yeast Biogeography and the Effects of Species Recognition Approaches: The Case Study of Widespread Basidiomycetous Species from Birch Forests in Russia. *Current Microbiology*, 70(4), 587–601. <https://doi.org/10.1007/s00284-014-0755-9>

Yurkov, A., Krüger, D., Begerow, D., Arnold, N., & Tarkka, M. T. (2012). Basidiomycetous Yeasts from Boletales Fruiting Bodies and Their Interactions with the Mycoparasite *Sepedonium chrysospermum* and the Host Fungus *Paxillus*. *Microbial Ecology*, 63(2), 295–303. <https://doi.org/10.1007/s00248-011-9923-7>

Yurkov, A. M. (2018). Yeasts of the soil—Obscure but precious. *Yeast*, 35(5), 369–378. <https://doi.org/10.1002/yea.3310>

Yurkov, A. M., Kemler, M., & Begerow, D. (2012). Assessment of yeast diversity in soils under different management regimes. *Fungal Ecology*, 5(1), 24–35. <https://doi.org/10.1016/j.funeco.2011.07.004>

Yurkov, A. M., Röhl, O., Pontes, A., Carvalho, C., Maldonado, C., & Sampaio, J. P. (2016). Local climatic conditions constrain soil yeast diversity patterns in Mediterranean forests, woodlands and scrub biome. *FEMS Yeast Research*, 16(1), fov103. <https://doi.org/10.1093/femsyr/fov103>

Yurkov, A. M., Wehde, T., Federici, J., Schäfer, A. M., Ebinghaus, M., Lotze-Engelhard, S., Mittelbach, M., Prior, R., Richter, C., Röhl, O., & Begerow, D. (2016). Yeast diversity and species recovery rates from beech forest soils. *Mycological Progress*, 15(8), 845–859. <https://doi.org/10.1007/s11557-016-1206-8>

Yurkov, A., Wehde, T., Kahl, T., & Begerow, D. (2012). Aboveground Deadwood Deposition Supports Development of Soil Yeasts. *Diversity*, 4(4), 453–474. <https://doi.org/10.3390/d4040453>


Yurkov, A. M., Kemler, M., & Begerow, D. (2011). Species Accumulation Curves and Incidence-Based Species Richness Estimators to Appraise the Diversity of Cultivable Yeasts from Beech Forest Soils. *PLoS ONE*, 6(8), e23671. <https://doi.org/10.1371/journal.pone.0023671>

Zhang, H., Skelton, A., Gardner, R. C., & Goddard, M. R. (2010). *Saccharomyces paradoxus* and *Saccharomyces cerevisiae* reside on oak trees in New Zealand: Evidence for migration from Europe and interspecies hybrids: New Zealand *Saccharomyces* ecology. *FEMS Yeast Research*, 10(7), 941–947. <https://doi.org/10.1111/j.1567-1364.2010.00681.x>

Figure legends:

Figure 1: Geographic distribution of yeast sequences within temperate forest samples in the database GlobalFungi. Points on this map were used for analyses mentioned in the text. Relative abundances represent the share of ITS yeast sequences of each species in the global fungal ITS sequence pool. Each point represents a sequenced environmental sample and blue regions represent the distribution of temperate forest biomes. Points with a relative abundance of zero represent samples with a single yeast sequence, and these were rounded down to zero for the map. Points outside blue regions are from forest environments contiguous with temperate forests; we included these in the map and analyses to avoid breaking up information from individual forests. The world map with temperate forest biomes was obtained from <https://earthobservatory.nasa.gov/biome/maptemperate.php>

Figure 2: Success rates in isolating different *Saccharomyces* species across different geographical regions. The only significant correlation regards the success rate of *S. uvarum* isolation in South America, which decreases with latitude (Spearman coefficient, $rs = -0.94286$, two-tailed p -value = 0.0048). Data from table S3.



Relative abundance

- 0
- $\leq 0.1\%$
- $0.1\text{--}1\%$
- $1\text{--}10\%$
- $> 10\%$

