New lower bounds on the size-Ramsey number of a path

Deepak Bal

Department of Marthematics Montclair State University Montclair, NJ 07043, U.S.A. deepak.bal@montclair.edu

Louis DeBiasio*

Department of Mathematics Miami University Oxford, OH 45056, U.S.A. debiasld@miamioh.edu

Submitted: Aug 21, 2020; Accepted: Nov 17, 2021; Published: Jan 28, 2022 The authors. Released under the CC BY-ND license (International 4.0).

Abstract

We prove that for all graphs with at most (3.75 - o(1))n edges there exists a 2-coloring of the edges such that every monochromatic path has order less than n. This was previously known to be true for graphs with at most 2.5n - 7.5 edges.We also improve on the best-known lower bounds in the r-color case.

Mathematics Subject Classification 5:55, 05C38

1 Introduction

Given a graph H, let $\hat{R}_r(H)$ be the minimum m such that there exists a graph G with m edges such that in every r-coloring of G, there is a monochromatic copy of H. When r = 2, we drop the subscript and just write $\hat{R}(H)$. We refer to $\hat{R}(H)$ as the size-Ramsey number of H.

Let P_n be the path with n vertices. Erdős [17] famously asked if $\hat{R}(P_n)/n \to \infty$ and $\hat{R}(P_n)/n^2 \to 0$. Beck [7] proved that, in fact, $\hat{R}(P_n)$ 6 900n (for n sufficiently large). The bound 900n was subsequently improved in [10], [11], [14], [24] and currently rests at 74n as proved by Dudek and Pra lat in [15].

As for the lower bound, it is clear that $\hat{R}(P_n) > 2n - 4$ since P_n has n - 1 edgesBeck [7] proved $\hat{R}(P_n) > (9/4-o(1))n$, Bielak [9] proved $\hat{R}(P_n) > 9n/4-3$, Bollob´ as [10] proved $\hat{R}(P_n) > (1+\frac{1}{2}-o(1))n$, and finally Dudek and Pra lat [15] proved $\hat{R}(P_n) > 5n/2-15/2$.

The closest thing there is to a conjecture about the precise value $\hat{R}(P_n)$ is Bollobás' [10] comment, "it would not be surprising if $\hat{R}(P_n)$ turned out to be about 8n." We are not aware of what insight led to this comment, but – together with the recent flurry of

^{*}Research supported in part by NSF grant DMS-1954170

activity on the upper bound – it inspired us to make a determined effort to improve the lower bound. We prove the following.

Theorem 1.For all > 0, there exists n_0 such that if $n > n_{-0}$ and G is a graph with at most (3.75 -)n edges, there exists a 2-coloring of the edges of G such that every monochromatic path has order less than $nThus \hat{R}(P_n) > (3.75 - o(1))n$.

For the general, r-color version of the problem, the best upper bound is due to Krivelevich [22] who prove $\hat{\mathbb{R}}_r(P_n) = O(r^2 \log(r)n)$ (Dudek and Pra lat [16] later gave a different proof). In fact, both [22] and [16] prove the stronger "density version" of the theorem: there exists a graph G (a binomial random graph) with $|E(G)| = O(r^2 \log(r)n)$ such that every subgraph of G with at least e(G)/r many edges contains a monochromatic path of order n (A recent paper of Balogh, Dudek, and Li [3] shows that the factor log r cannot be improved for this stronger density version in the setting of random graphs).

As for the lower bound, Dudek and Pra lat [15] proved that for any r > 2, $\hat{R}_r(P_n) > \frac{(r+3)r}{4}n - O(r^2)$ and then Krivelevich [22] proved that for any r > 3 such that r - 2 is a prime power, $\hat{R}_r(P_n) > (r-2)^2 n - o(n)$. We improve on each of these results by proving the following.

Theorem 2 Let r > 2 and let q be the largest prime power such that q 6 r – 1. Then

$$\hat{R}_r(P_n) > \max \frac{(r-1)r}{2} + 2.75 - o(1) \quad n, \ (q^2 - o(1))n$$

Note that the prime number theorem guarantees that for any $\varepsilon > 0$ and r sufficiently large, there is a prime between $(1 - \varepsilon)r$ and r, so for sufficiently large r, the second term in the maximum will dominate and we have $\hat{R}_r(P_n) > (r - 1 - o_{-r}(1))^2 n$. Determining whether $\hat{R}_r(P_n) = \Theta(r^2)n$ or not is perhaps the most interesting open problem regarding the size-Ramsey number of a path.

1.1 Outline, Notation

Our improvement in the lower bound stems from two main ideas.

1) If we can partition the graph G into sets of order at most n-1 such that the number of edges crossing the partition is at most n-2, then we can color the edges inside the sets red and the edges between the sets blue so there are no monochromatic Ps. This has some similarity to the problem of determining the bisection width of a graph 1 – in which case a result of Alon [1, Proposition 3.1] gives good bounds on the number of crossing edges in a balanced bipartition of graphs with bounded maximum degree and at most 2n-2 vertices. However, in our case, G may not have bounded maximum degree, G may have more than 2n-2 vertices, and we don't necessarily want the partition to be balanced. Nevertheless, with some extra work, we are able to use similar methods from the study of the bisection width problem (e.g. [1], [23]) in our setting.

¹the *bisection width* of a graph is the minimum number of edges crossing a balanced bipartition of the vertex set

2) From the ordinary path Ramsey problem it is known that if G has at most $\frac{3n}{2} - 2$ vertices, then there exists a 2-coloring of G such that every monochromatic path has order less than n. We show that if G has between roughly 3n/2 and 5n/3 vertices and few enough edgesthen there exists a 2-coloring of G such that every monochromatic path has order less than n. This allows us to only consider graphs with at least 5n/3 vertices.

In Section 2 we prove a number of lemmas which we will use throughout the proof. We also show how some of these lemmas imply the previously known lower bounds on the size-Ramsey number of paths. In Section 3 we prove Theorem 1. In Section 4 we prove Theorem 2. In Section 5, we list a few observations and approaches that may helpful in trying to improve the lower bounds we have provided.

Let G = (V, E) be a graph. For all $S \subseteq V$, we write G - S to mean $G[V \setminus S]$; i.e. the graph induced by $V \setminus S$. Given disjoint sets $A, B \subseteq V$, we write G[A, B] to mean the bipartite subgraph of G with vertex set $A \cup B$ with edges having one endpoint in A and the other in B and we write E[A, B] to be the edge set of G[A, B]. We sometimes write |G| to mean |V|. For any other notation we defer to [12]All logarithms are natural (base e) unless otherwise stated. Throughout the paper, if we refer to an r-coloring of G, we mean an r-coloring of the edges of G.

2 Lemmas

When proving a lower bound on the r-color size-Ramsey number of R, we are given a graph G = (V, E) and we must exhibit an r-coloring of the edges of G so that G has no monochromatic paths of order n. It is often useful to break this into cases depending the number of vertices of G. In Section 2.1 we use the examples from the ordinary path Ramsey problem to determine a lower bound on |V|. In Section 2.2 we prove a general result which allows us, when proving a lower bound on $\hat{R}_r(P_n)$, to restrict our attention to graphs with minimum degree at least r + 1, which in turn gives us an upper bound on |V|. In Section 2.3, we prove a lemma which we use in the proof of Theoremir2Section 2.4, we prove the main lemma of the paper needed for the proof of Theorem 1Finally, in Section 2.5 we show how to deal with the case when G has between roughly 3n/2 and 5n/3 vertices.

2.1 Examples from the ordinary path Ramsey problem

Proposition 3 (Gerencs'erGy'artás [19]). If G has at most $\frac{3n}{2}$ – 2 vertices, then there exists a 2-coloring of G such that every monochromatic path has order less than n.

Proof. Partition V(G) into two sets X_1 , X_2 with $|X_1|$ 6 $\frac{n}{2}$ – 1 and $|X_2|$ 6 n – 1. Color all edges incident with X_1 red and all edges inside X blue. Clearly the longest blue path has order n – 1. Any pair of consecutive vertices on a red path must contain at least one vertex of X_1 . Thus the longest red path is of order at most $2|X_1|$ + 1 6 n – 1.

Proposition 4 (Yongqi, Yuansheng, Feng, Bingxi [27]). Let r > 3. If G has at most $2(r-1)(\frac{n}{2}-1) = (r-1)(n-2)$ vertices, then there exists an r-coloring of G such that every monochromatic path has order less than n.

Proof. Partition V(G) into 2r-2 sets $X_1, X_2, \ldots, X_{r-2}$ each of order at $\operatorname{most}_{\frac{n}{2}}^n - 1$. In the following, addition is modulo 2r-2. For $i=1,\ldots,r-1$, color with color i, the edges between X_i and X_{i+1},\ldots,X_{r-2} and the edges between X_{i+r-1} and X_{i+r},\ldots,X_{r-3} . Use color r for the edges between X_i and X_{i+r-1} for $i=1,\ldots,r-1$. Color arbitrarily within the X_i 's. This coloring has no monochromatic P_i in color P_i for P_i in color P_i for the same reason as in Proposition 3. There is none in color P_i since each component of color P_i is of order less than P_i .

2.2 A reduction lemma

Fact 5. If G = (V, E) is a graph with minimum degree at least r + 1, then $|V| = 6 \frac{2|E|}{r+1}$.

The following lemma shows that in order to get a lower bound on the r-color size-Ramsey number of P, we can restrict our attention to graphs P0 with minimum degree at least P1, and consequently at mosP1 vertices. This generalizes an observation which is implicit in the proof of Beck's lower bound [7].

Lemma 6. Let r and n be positive integers with n > r + 4!f every connected graph with at most m edges and minimum degree at least r + 1 (and consequently at most 2m/(r + 1) vertices) has an r-coloring such that every monochromatic path has order less than n - 2, then every graph with at most m edges has an r-coloring such that every monochromatic path has order less than n.

Proof. Suppose that every connected graph with at most m edges and minimum degree at least r +1 has an r-coloring such that every monochromatic path has order less than n-2. Let G = (V, E) be a graph with at most m edges. Let $S = \{v \in V : d(v) \ 6 \ r\}$. We begin by describing how to color the edges of G - S so that G - S contains no monochromatic paths of order n - 2.

If G-S has fewer than n-2 vertices, then coloring the edges of G-S arbitrarily we have an r-coloring of G-S with no monochromatic paths of order n-2So suppose G-S has at least n-2>r+2 vertices. Let v be a vertex in G-S and suppose that v has exactly v+1-t neighbors in v=1 for some positive v=1. This means v=1 had at least v=1 neighbors in v=1 neighbors in v=1 for some positive v=1. This means v=1 had at least v=1 neighbors in v=1 neighbors

Now let $u_1, \dots, y_1, \dots, y_n$ be an ordering of the vertices of V where $S = \{u_1, \dots, y_n\}$. For all $i \in [s]$, let $E_i^- = \{u_1 u_1 \in E : j < i\}$, let $E_i^+ = \{u_1 u_1 \in E : j > i\}$, and let $E_i^- = E_i^- \cup E_i^+$. We now iteratively color the edges incident with each of u_1, \dots, y_n starting with u_1 . Since $\Delta(G[S])$ 6 r, we color the edges in $E_1^- = E_1^+$ with distinct colors. Now let 2 6 i 6 s and suppose that for all 1 6 i 6 i – 1, we have colored all edges in E_i^- with the property that the set of colors used on E_j^+ is disjoint from the set of colors used on E_j^+ and all the edges in E_j^+ receive distinct colors. Note that the edges in E_i^- have already been colored but since $d(u_i)$ 6 r we can color the edges in E_i^+ such that every edge receives a different color and no color which is used on an edge $i\bar{p}$ is used on an edge in E_i^+ .

We have now colored all of the edges incident with S such that every monochromatic component consisting of edges incident with S is a star with all of its leaves in We have that every monochromatic path which only uses edges from G–S has order less than n–2 and every monochromatic path which only uses edges from $E(G[S]) \cup E[S, V(G) - S]$ has order at most S. If a monochromatic, say color S, path uses an edge from S, S, then since its endpoint in S is not incident with any other edges of S color S, this edge must be a pendant edge of the path (of which there are only two) and thus the longest monochromatic path in S has order less than S and S are S are S and S are S are S and S are S and S are S and S are S and S are S are S and S are S and S are S are S and S are S are S are S and S are S and S are S are S and S are S are S and S are S and S are S and S are S are S and S are S are S and S are S and S are S and S are S are S and S are S are S are S and S are S are S and S are S and S are S and S are S are S and S are S and S are S and S are S and S are S are S and S are S and S a

Corollary 7. For all
$$n > r + 4$$
, $\hat{R}_r(P_n) > \frac{r+1}{2} \cdot R_r(P_{n-2})$.

Proof. Let G = (V, E) be a connected graph with fewer than $\frac{r+1}{2} \cdot R_r(P_{n-2})$ edges and minimum degree at least r+1. So |V| 6 $\frac{2|E|}{r+1} \le R_r(P_{n-2})$ and thus G has an r-coloring with no monochromatic P_{n-2} . So by Lemma 6, every graph with fewer that $\frac{r+1}{2} \cdot R_r(P_{n-2})$ edges has an r-coloring with no monochromatic P_n .

Remark 8. Proposition 3 and Corollary 7 imply that

$$\hat{R}(P_n) > \frac{3}{2} \cdot R(P_{n-2}) > \frac{3}{2} \cdot \frac{3}{2}(n-2) - \frac{3}{2} = \frac{9}{4}n - \frac{27}{4}.$$

Remark 9. Proposition 4 and Corollary 7 imply that for r > 3,

$$\hat{R}_r(P_n) > \frac{r+1}{2} \cdot R_r(P_{n-2}) > \frac{r+1}{2} (r-1)(n-4) = \frac{r^2-1}{2} n - 2(r^2-1).$$

Remark 10. The bound in Remark 9 is less than the bounds given in Theorer Remark 9 is the easiest way to see that $\Re_r(P_n) = \Omega(r^2 n)$.

2.3 Trimming a tree so that no long paths remain

The following is a slight generalization of the lemma used in [10] and [15] to give a lower bound on the size-Ramsey number of a path.

Lemma 11. For every tree T with |V(T)| > bn/2c, there exists a set E^0 of at most $b \frac{|V|}{bn/2c}c - 1$ edges such that $T - E^0$ has no paths of order n.

Proof. If *T* has no path of order *n* we are done, so choose a path of order *n* and delete the middle edge (or one of the two middle edges if *n* is odd). This separates *T* into two subtrees, each with at least bn/2c vertices. Now repeat on each subtree and call the set of deleted edges $\not\models^0$. When the process stops, every component of $T - E^{-0}$ has at least bn/2c vertices and no paths of order *n*. Thus $T - E^{-0}$ has at most $b \frac{|V|}{bn/2c} c$ components, which means $|E| = b \frac{|V|}{bn/2c} c - 1$.

Remark 12. Proposition 3 and Lemma 11 imply that $\hat{R}(P_n) > \frac{5}{2}n - 7$.

Proof. Let G = (V, E) be a graph with at most $\frac{5n}{2} - \frac{15}{2}$ edges. We may assume G is connected and by Proposition 3 we may assume $\frac{3n}{2} - \frac{3}{2}$ 6 |V|. Let T be a spanning tree of G (which contains at least $\frac{3n}{2} - \frac{5}{2}$ edges). Applying Lemma 11, there exists a forest F with $F \subseteq T$ such that F has at least $\frac{3n}{2} - \frac{11}{2}$ edges and no paths of order n, so we may color all of the edges of F red without creating a red P_n . There are at most $\frac{5n}{2} - \frac{15}{2} - (\frac{3n}{2} - \frac{11}{2}) = n - 2$ edges remaining in $E(G) \setminus E(F)$, all of which we may color blue without creating a blue P_n .

2.4 Main lemma

We will only use the following lemma in the case where k = 1 or k = 2, but we state it in general here. Note that for instance when k = 1, this says that if G is a graph on n - 1 < N 6 2(n - 1) vertices, then there is a bipartition of V(G) into sets of order n - 1 and N - (n - 1) such that the number of edges crossing the partition is approximately what we would get by taking a random such partition of a graph with |E(G)| - N edges.

Lemma 13. There exists n_0 such that for all $n > n_0$, if G = (V, E) is a graph on N > n vertices, then the following holds. Let k be a positive integer uniquely defined by $k(n-1) < N \ 6 \ (k+1)(n-1)$ where $k \ 6 \ n^{-1/32}$ and let $\alpha = \frac{n-1}{N}$. If every component of G has at least n vertices, $\Delta(G) \ 6 \ N^{-1/16}$, and $|E| \ 6 \ 100N \ 6 \ 100(k+1)n$, then there exists a partition of V into k+1 parts V_{-1}, \ldots, V_{k+1} such that $|V_1|, \ldots, |V_k|, |V_{k+1}| \ 6 \ n-1$ and $|V_{k+1}| \ 6 \ N - k(n-1) + N^{-15/16}$ and the number of edges crossing the partition is at most $(1-k\alpha^2-(1-k\alpha)^2)(|E|-N)+N^{-15/16}$.

The first tool needed to prove Lemma 13 is the following fact mentioned by Alon [1], stated in general and made explicit here.

Lemma 14Let G be a connected graph on p vertices with maximum degree E_Q any 1 6 ` < p, we can find a collection of connected subgraphs₁S. . . , Sof G such that (T1) $V(S_1), \ldots, V(S_t)$ form a partition of V(G) with ` < $|S_t|$ 6 1 + Δ ` for all $i \in [t-1]$ and $|S_t|$ 6 1 + Δ `

(T2)
$$P_{i=1}^{t} |E(S_{i})| > p - t$$

(T3) if $\hat{p} = b$ pc, then $\frac{1}{\Delta + 1}$ \bar{p} 6 t 6 \bar{p} + 1

Proof. Let T_0 be a rooted spanning tree of G with (arbitrary) root r. For a rooted tree T and vertex v, let s(T, v) denote the subtree of T rooted at vertex v and let C(v) denote the set of children of v. Assume T has been defined for some i > 0 and that r is still the root of T_i . Traverse down T from r until encountering a vertex v (if one exists) such that $|s(T_i, v)| >$ and $|s(T_i, u)| 6$ for all $u \in C(v)$. Then $s(T_i, v)$ satisfies

`
$$< |s(T_i, v)| = 1 + X_{u \in C(v)} |s(T_i, u)| 6 1 + \Delta$$
`. (1)

If $v \in S_i$, let $S_{i+1} = s(T_i, v)$ and $T_{i+1} = T_i - S_{i+1}$ and repeat for i + 1. If v = r or if no such vertex v exists, then set $S_{i+1} = S_i = T_i$. Each S is connected by construction. Property (T1) is satisfied by (1). Property (T2) follows since each S is connected and thus $\sum_{i=1}^{t} |E(G[S_i])| > \sum_{i=1}^{t} (|S_i| - 1) = p - t$.

Finally, if $= b \overline{p}c$ we have

$$(t-1)(b^{\sqrt{p}c+1}) \ 6 \sum_{i=1}^{X^t} |S_i| = p \ 6 \ t(1+\Delta^{\sqrt{p}c})$$

and from each of $(t-1)(\stackrel{\sqrt{p}}{\overline{p}}c+1)$ 6 p and p 6 $t(1+\stackrel{\sqrt{p}}{\overline{p}})$ 6 $t\stackrel{\sqrt{p}}{\overline{p}}(1+\stackrel{\Delta}{\Delta})$, we derive the bounds on t in (T3).

The next tool we need is the following concentration inequality of McDiarmid [25] (see also [18]). We note that McDiarmid's inequality isn't strictly necessary in this first application, but as we will use it again later in a different context, it is easiest to use it for both purposes.

Lemma 15 (McDiarmid's inequality). Let $Z = Z(X_1, \dots, X_n)$ be a random variable that depends on N independent random variables X_1, \dots, X_n . Suppose that

$$|Z(X_1,\ldots,X_n,\ldots,X_n)-Z(X_1,\ldots,X_n)| \leq c_k$$

for all k = 1, ..., N and $X_1, ..., X_k, X_k^0$. Then for any t > 0 we have

P [Z > E[Z] + t] 6 exp
$$-\frac{t^2}{2 \sum_{k \in [N]}^{t^2} c_k^2}$$
.

We are now ready to prove the main lemma.

Proof of Lemma 13. Apply Lemma 14 with $\dot{}=b$ $\sqrt{N}c$ to partition the components of G into $\frac{N}{\Delta+1}$ 6 t 6 \overline{N} + 1 connected subgraphs $S \cdot \cdot \cdot \cdot \cdot S$ each of order at most $1 + \Delta \cdot \overline{N}$. There are at least $N - (t-1) > N - \overline{N}$ edges accounted for in these subgraph Define $M = |E| - (N - \overline{N})$ to be an upper bound on the number of edges of G which are not contained in these subgraphs.

We independently at random place each such connected subgraph in one of the sets $V_1, \ldots, \not V_{k+1}$ with probability α for all V_i with $i \in [k]$ and probability $1 - k\alpha$ for V_{k+1} . Let Z_i represent the number of vertices which land in the set i for all $i \in [k+1]$.

Then E $[Z_1] = \cdots = E[Z_k] = \alpha N = n - 1$ and E $[Z_{k+1}] = (1 - k\alpha)N = N - k(n - 1)$. Note that changing the position of one of $\{S_i, \dots, S_i\}$ can change any of these variables by at most $1 + \Delta = N = 0$. Thus we may apply McDiarmid's inequality (Lemma 15) and the union bound to conclude that the probability that for some $i \in [k]$, Z_i exceeds $n - 1 + N^{7/8}$ or Z_{k+1} exceeds $N - k(n - 1) + N^{7/8}$ is at most

$$(k+1) \cdot \exp \left[-\frac{1}{2} \cdot \frac{\sqrt{N^{7/4}}}{(\overline{N}+1) \cdot (1+N^{9/16})^2}\right] = \exp -\Omega(N^{-1/8})$$

Thus at least $1 - e^{-\Omega(N^{-1/8})}$ proportion of the partitions satisfy

$$|V_1|, \dots, |M| 6 n - 1 + N^{-7/8}$$
 and $|V_{k+1}| 6 N - k(n-1) + N^{-7/8}$. (2)

Now, by linearity of expectation, the expected number of edges μ crossing the partition satisfies

$$\mu$$
 6 (1 – $k\alpha$ ² – (1 – $k\alpha$) ²) m .

So there is a partition $V \cdot \cdot \cdot \cdot \cdot V_{k+1}$ satisfying (2) with at most $(1-k\alpha^2-(1-k\alpha)^{-2})m+1$ edges crossing the partition; otherwise we would have

$$(1 - k\alpha^2 - (1 - k\alpha)^2)m > \mu > (1 - e^{-\Omega(N^{-1/8})})((1 - k\alpha^2 - (1 - k\alpha)^2)m + 1)$$

> $(1 - k\alpha^2 - (1 - k\alpha)^2)m$,

a contradiction.

Finally, in order to achieve the desired upper bounds on the sizes of $_1V \cdot \cdot \cdot \cdot \not V_{k+1}$, we potentially have to slightly modify the partition given above. When modifying the partition, we only want to move vertices which have bounded degrees olet $S = \{v \in V(G) : d(v) \in V(G)\}$ and note that

$$|S| > (1 - \frac{1}{2k})N;$$
 (3)

as otherwise there are at leas $\frac{N}{2k}$ vertices of degree greater than 400k which gives

$$200N > 2|E| = \frac{X}{v \in V(G)} d(v) > \frac{N}{2k} \cdot 400k = 200N,$$

a contradiction.

Now if |V| > n-1 for $i \in [k+1]$, there must exist $j \in [k+1] \setminus \{i\}$ such that |V| = j = n-1, so we select a vertex from MS and we move it to M. Because $|S| > (1-\frac{1}{2k})N$ and by (2), we can repeat this process for at most MN steps until we have $|V| \cdot \cdot \cdot \cdot \cdot \cdot |M| \cdot |V_{k+1}| = 6 \cdot n-1$ and $|V_{k+1}| = 6 \cdot N - k(n-1) + (k+1)N$ and $|V_{k+1}| = 6 \cdot N - k(n-1) + N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |M| \cdot |V_{k+1}| = 6 \cdot N - k(n-1) + N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |V| \cdot |V_{k+1}| = 6 \cdot N - k(n-1) + N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |V| \cdot |V_{k+1}| = 6 \cdot N - k(n-1) + N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |V| \cdot |V_{k+1}| = 6 \cdot N - k(n-1) + N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |V| \cdot |V| \cdot |V| \cdot |V| = 1 \cdot N$ steps until we have $|V| \cdot \cdot \cdot \cdot \cdot |V| \cdot |V|$

$$(1 - k\alpha^2 - (1 - k\alpha)^2)m + 1 + kN^{7/8} \cdot 400k < (1 - k\alpha^2 - (1 - k\alpha)^2)(|E| - N) + N^{15/16}$$
 as desired

2.5 Extending Proposition 3

The following observations extend Proposition 3. We note that there is a similarity between this observation and the concept of the *integrity* of a graph (see [26]).

Observation 16 *f G* has a set *S* of at most $\frac{n}{2}$ – 1 vertices such that every component of *G* – *S* has no path of order *n*, then there exists a 2-coloring of the edges of *G* such that every monochromatic path has order less than *n*.




Figure 1: Coloring the edges of *G* in Observation 16

Proof. We color all edges incident to S with blue and every other edge red. By the condition on the components of G - S, there will clearly be no red path of order n. Any pair of consecutive vertices on a blue path must contain at least one vertex of S. Thus the longest blue path is of order less than n.

We also note that there is a similarity between the following observation and the concept of the *edge integrity* of a graph (see [2]).

Observation 17. If G has a subgraph H such that H has no path of order n (in particular, if H has at most n-2 edges) and every component of $(V(G), E(G) \setminus E(H))$ has order less than n, then there exists a 2-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. Color the edges of H with red and color the remaining edges blue.

The following lemma says that if the number of vertices is not too much more than 3n/2 and the number of edges of G is small enough, we can essentially color G in a way which resembles the coloring in Proposition 3.

Given a graph H, and a positive real $\rho > 0$, we say that a graph G = (V, E) is (ρ, H) free if the graph G^0 obtained from G by adding a set of vertices U with |U| 6 ρ such that
for all $u \in U$, $N(u) = (U \cup V) \setminus \{u\}$ does not contain a copy of H. For example, every
graph with at most $n - 2d\rho e - 2$ edges is (ρ, R) -free.

Lemma 18. Let $0 < < \frac{1}{100}$, let n be sufficiently large, and let G = (V, E) be a graph with $\delta(G) > 3$ and $(\frac{3}{2} -)n < |V| < (\frac{5}{3} -2)n$. Let $0 < \sigma$ 6 $\frac{1}{6}$ be defined by $|V| = (\frac{3}{2} -+\sigma)n$ and let $d = \min\{b \frac{1/2-3}{\sigma} c + 1, 100\}$. If

$$|E| 6 \frac{3(d+1)+6\sigma}{4} - d n$$

then there exists a 2-coloring of G such that every blue component has order at most n-1 and the graph G_R induced by the red edges $is_A^n(P_n)$ -free.

Proof. First note that d is an integer with either d = 100 or d = $b \frac{1/2-3}{\sigma} c + 1 > \frac{1/2-3}{\sigma} > \frac{1/2-3}{1/6-} = 3$; either way, d > 4. To get a better feel for the statement, note that since σ

depends on |V| and d depends on σ , the upper bound on |E| is a piecewise function of |V| which increases on every interval corresponding to a fixed value of d, but which decreases as d increases (roughly, the upper bound decreases as |V| increases - see Figure 4).

The proof is based on the following claim.

Claim 19. There exists a partition {X, Y, Z} of V such that

(i) every vertex in X has at most one neighbor in Z and

(ii)
$$|Z| 6 n - 1$$
, $|Y| 6 (1/2 -)n$, and $|X| + |Y| 6 n - 1$.

Before proving the claim, note that if such a partition exists, we can color all edges inside Z and all edges inside $X \cup Y$ blue and all edges in $E[X \cup Y, Z]$ red so that clearly every blue component has at most n-1 vertices (see Figure \mathbb{Z}), see that the red graph G_R is $(n/4, P_n)$ -free, add a set V of at most n/4 vertices adjacent to every vertex in G_R and call the resulting graph G_R^0 . Let P be a longest path in G_R^0 and let $V_0^0 = V_0 \cap V(P)$, $X^0 = X \cap V(P)$, $Y^0 = Y \cap V(P)$, and $Z^0 = Z \cap V(P)$. Note that since every vertex in X has degree 1 in G_R , every vertex in X^0 which is not an endpoint of P is either preceded by or succeeded by a vertex from V so


$$|X^0| 62|V_0^0| + 2$$
 (4)

(since the same vertex from \mathscr{V} can be counted twice). Also note that every vertex in Z^o which is not the final vertex of P is succeeded by a vertex from ${}_0^{0}VX^{0} \cup Y^{0}$. So we have


$$|Z^{0}| 6 |V_{0}^{0}| + |X^{0}| + |Y^{0}| + 163 |V_{0}^{0}| + |Y^{0}| + 3$$
 (5)

and thus by (4), (5), and the fact that $|V_0|^0 = |V_0|^0 = |V_0|^0$

$$|V(P)| = |V_0^0| + |X^0| + |Y^0| + |Z^0| \cdot 62|Y^0| + 6|V_0^0| + 5 \cdot 6(1-2)n + 6 \cdot \frac{n}{4} + 5 \cdot 6n - 1.$$



(a) Finding the desired partition if Y * is small enough



(b) Finding the desired partition after moving vertices from Y*

Figure 2: Coloring the edges in Lemma 18

Proof of Claim 19. Let $X^* = \{v \in V : d(v) \ 6 \ d\}$. We first show that $|X^*|$ is significantly larger than σn . Indeed, using $\delta(G) > 3$ we have

Rearranging and using $|V| = (3/2 - + \sigma)n$ gives

$$(d-2)|X^*| > (d+1)(3/2 - + \sigma)n - \frac{3(d+1) + 6\sigma}{2} - 2d$$

$$= (d-2)\sigma n + (d-1)n,$$

and thus $|X^*| > (\sigma +)n$. So we can choose $X \subseteq X$ such that

$$|X| = b(\sigma + \frac{1}{d-1})nc.$$
 (6)

Let $Y^* = N(X) \setminus X$, and note that

$$|Y^*| 6 d|X|. (7)$$

Since $N(X) \subseteq X \cup Y$ * we would be done if $|Y^*| \in (1/2 - 2)$ by letting $Y \subseteq V \setminus X$ such that $Y^* \subseteq Y$ and |Y| = b(1/2 -)nc and letting $Z = V \setminus (X \cup Y)$ and noting that

$$|Z| = |V| - |X| - |Y| = (3/2 - + \sigma)n - |X| - |Y|$$

$$\stackrel{(6)}{6} (3/2 - + \sigma)n - (\sigma + \frac{1}{d-1})n - (1/2 - n) 6 n - 1.$$

where the last inequality holds since *n* is sufficiently large.

We now show that if $|Y^*| > (1/2 - 2)$, then we can move at least |Y'| - (1/2 - 2) vertices from Y^* to Z. We do this by showing that there exists an induced matching in the bipartite graph $G[X, Y^*]$ of size at least $|Y^*| - (1/2 - 2)$.

Let $Y_1 = \{v \in Y^* : d(v, X) = 1\}$ and $Y_2 = \{v \in Y^* : d(v, X) > 2\}$. We note that since every vertex in X sends at most d edges to Y, $G[X, Y^*]$ has an induced matching of size at least $|Y_1|/d$. We have

$$d|X| > e(X, Y^*) > |Y_1| + 2(|Y^*| - |Y_1|) = 2|Y^*| - |Y_1|$$

which implies

$$\frac{|Y_{1}|}{d} > \frac{2|Y^{*}|}{d} - |X| = |Y^{*}| - \frac{d-2}{d}|Y^{*}| - |X| \stackrel{(7)}{>} |Y^{*}| - (d-1)|X|$$

$$= |Y^{*}| - (d-1)b(\sigma + \frac{1}{d-1})nc$$

$$> |Y^{*}| - (d-1)\sigma n - n$$

$$> |Y^{*}| - \frac{1/2 - 3}{\sigma}\sigma n - n$$

$$= |Y^{*}| - (\frac{1}{2} - 2)n$$

where the last inequality holds by the upper bound on d.

Let M be a maximum induced matching in $G[X, Y^*]$ where X^0 are the vertices from X which are saturated by M and $f(X^0)$ are the vertices in Y^* which are saturated by M. Set $Y^0 = Y^* \setminus f(X^0)$. By the above we have $|Y| \in (1/2 - 2)$.

Finally, let $Y \subseteq V \setminus X$ such that $Y = 0 \subseteq Y$ and |Y| = b(1/2 -)nc. Now let $Z = V \setminus (X \cup Y)$ and note that as before

$$\begin{split} |Z| &= |V| - |X| - |Y| = (3/2 - + \sigma)n - |X| - |Y| \\ & \stackrel{(6)}{6} (3/2 - + \sigma)n - (\sigma + \frac{1}{d-1})n - (1/2 - n) \cdot 6n - 1. \end{split}$$

where the last inequality holds since *n* is sufficiently large.

This completes the proof of Claim 19.

Having established Claim 19, we have completed the proof of Lemma 18. □

3 Two colors

We are now ready to give a proof of Theorem We note that the in the following proof can be taken to be as small as $= n^{-\Theta(1)}$; however, for the sake of readability, we didn't try to optimize the value of .

Proof of Theorem 1. Let 0 < < 1/100 and let n = 0 be a sufficiently large integer (the value of which we don't explicitly compute, but we will point out which inequalities depend on n being sufficiently large). Let $G^0 = (V^0, E^0)$ be a connected graph with at most (3 + y -)n edges, where 0.6 y 6.3/2 is to be chosen later (ultimately, we will choose y = 3/4 and in fact our proof can handle larger values of y, but restricting y in this way makes it easier to apply Lemma 13 later in the proof). By Lemma 6 it suffices to assume that $\delta(G^0) > 3$ and thus $|V^0| 6.2|E^0/3.6 (2 + 2y/3 - 2/3)n 6.3(n - 1)$ (where the last inequality follows by the bounds on y and the fact that y is sufficiently large). We will exhibit a 2-coloring of y with no monochromatic y, but since we are using Lemma 6 this will prove that all graphs with at most y edges will have a 2-coloring with no monochromatic y (in other words, we will ultimately be showing that y of y edges with y of y edges will have a 2-coloring with no monochromatic y (in other words, we will ultimately be showing that y but y edges with y edges will have a 2-coloring with y edges will have a 2-coloring y edges will have a 2-coloring with y edges will have a 2-coloring y edges will have a 2-coloring y edges will have a 2-coloring with y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-coloring with no monochromatic y edges will have a 2-colorin


Let $V_0 = \{v \in V(G^0) : d(v) > n^{-1/32}\}$. We have $n^{1/32}|V_0|$ 6 2| E^0 | and thus since n is sufficiently large,

$$|V_0| 6 2(3 + y -)n^{-31/32} 6 \frac{^2n}{4}$$
 (8)


We say that a component C of $G^{-}V_{0}$ is small if |C| < n and large otherwise.

Suppose there are exactly t edges between₀V and the large components of $G - V_0$. Now let G = (V, E) be the graph obtained from G by deleting all of the vertices in V_0 , deleting all of the vertices in small components of $G - V_0$, and adding t edges inside V so that the minimum degree of G is at least 3.So |E| 6 |E| |E|.

We claim that we can 2-color the edges of G in such a way that the graph conduced by the blue edges has no components of order at least n and the graph @ induced by the red edges is $(\frac{2n}{4}, P_n)$ -free. Once we establish this claim, we complete the proof of Theorem 1 by coloring all edges between \forall and the large components of \forall V_0 red, the edges inside l/blue, the edges between Vand the small components blue, and the edges inside the small components red.







(b) Coloring the edges of G in Case 2 and Case 3

Figure 3: Coloring the edges in the proof of Theorem 1

Note that by Proposition 3, we may assume |V| > (3/2 - 2)n since any graph with at most (3/2 - 2)n vertices can be 2-colored in such a way that the red graph G_R is $(\frac{2n}{4}, P_n)$ -free and every component of the blue graph s has order at most n-1. **Case 1** $(\frac{8}{2} - 2)n < |V| < (\frac{5}{3} - 2^2)n$) In this case we are done by applying Lemma 18

(with 2 in place of).

Case 2 $(\frac{5}{8} - 2^{-2})n$ 6 |V| 6 2(n-1)) We parameterize this by introducing a variable σ such that $|V| = (3/2 - 2 + \sigma)n$ and thus $1/6 - 26 \sigma < 1/2 + 2$.

Lemma 13 (with k = 1) provides a bipartition of V into sets V_1 , V_2 of order at most n-1 such that the number of edges crossing the partition is at most

$$1 - \frac{1}{1 + (1/2 + \sigma)}^{2} - \frac{1/2 + \sigma}{1 + (1/2 + \sigma)}^{2} + 2^{2} (3/2 + y - \sigma -)n + |V|^{15/16}.$$

This quantity is less than $(1 - \frac{1}{2})n$ since n is sufficiently large and

$$y \ 6 \ \frac{3/4 + \sigma + 3\sigma^2}{1 + 2\sigma}.$$
(9)

Since we are assuming 1/6 – 2 6 σ < 1/2 + 2 , we have $\frac{3/4+\sigma+3\sigma^{-2}}{1+2\sigma}$ > 0.75 – with the minimum occurring when $\sigma = \frac{1}{6}$ – 2.

Now color the edges inside the sets $_1VV_2$ blue and the edges between the sets $_1VV_2$ red. Since the red graph $_{\mathfrak{S}}$ has at most $(1-\frac{2}{4})n$ edges, $_{\mathfrak{S}}$ is $(\frac{2n}{4}, P_n)$ -free.

Case 3 (2(n-1) < |V| 6 3(n-1)) We parameterize this by introducing a variable τ and assuming that $|V| = (2 + \tau)(n - 1)$ where $0 < \tau 6 1$. Apply Lemma 13 (with k = 2) to

get a tripartition of V into sets V_1 , V_2 , V_3 of order at most n-1 such that the number of edges crossing the partition is at most

$$1-2 \frac{1}{2+\tau}^{2} - \frac{\tau}{2+\tau}^{2!} (1+y-\tau-)n+|V|^{-15/16} < (1-\frac{\tau}{4})n,$$

where the last inequality holds provided n is sufficiently large and

$$y = 6 \frac{1 + \tau + 5\tau^2/2}{1 + 2\tau}$$

We have $\frac{1+\tau+5\tau^{-2}/2}{1+2\tau} > \frac{3}{4}(\sqrt[5]{5}-1) \approx 0.927$ with the minimum occurring when $\tau = \frac{3\sqrt[5]{5}-5}{10} \approx 0.1708$.

Now color the edges inside the sets $_1VV_2$, V_3 blue and the edges between the sets $_1VV_2$, V_3 red. Since the red graph G_1 has at most $(1 - \frac{1}{4})n$ edges, G_2 is $(\frac{2n}{4}, P_n)$ -free. \Box

One note about the previous proof. If we were to deal with the case $(\frac{3}{2}-2)n < |V|$ $\frac{6}{2}(n-1)$ by simply applying Lemma 13, the bound we obtain in (9) would be $y \in (\frac{3}{2}-1)$ which gives an overall lower bound of $\frac{1}{2}(P_n) > (2+\frac{1}{2}-o(1))n \approx 3.732n$ (this hypothetical scenario is depicted by the dashed red curve in Figure 4)So by using Lemma 18 to deal with the case $(\frac{3}{2}-2)n < |V| < (\frac{5}{3}-2^2)n$ separately, we get an improvement of about 0.018n. In light of this, if one were to attempt to improve the lower bound of (3.75-o(1))n, a good test case would be when $|V| \approx \frac{5n}{3}$, since this corresponds to the case where $|V| = \frac{3}{2}(1-2)n$ and $|V| \approx \frac{5n}{3}$, since this of the above proof.

Finally we note that by our result, one immediately obtains

$$\hat{R}(C_n, C_n) > \hat{R}(P_n, C_n) > \hat{R}(P_n, P_n) > (3.75 - o(1))n$$

for all sufficiently large n. However when n is odd, any graph G with |V(G)| 6 <math>2n - 2 can be 2-colored in a way which avoids a monochromatic R and C_n (partition the vertices into 2 sets of size at most n - 1 and note that the graph between the two sets contains no odd cycles). Thus, we can use the bounds provided by Case 3 of the proof of Theorem 1 to obtain the following.

Remark 20. For all > 0 and sufficiently large n, if G = (V, E) is a graph with |V| > 2n-1 and $|E| 6 (3 + \frac{3(-\overline{5}-1)}{4} -)n$, then there exists a 2-coloring of the edges of G such that every monochromatic path has order less than n.Consequently,for all sufficiently large odd n, $\hat{R}(C_n, C_n) > \hat{R}(P_n, C_n) > (3 + \frac{3(-\overline{5}-1)}{4} - o(1))n > 3.927n$.

The above improves a bound of $\hat{R}(P_n, C_n) > 3(n-1)$ (for odd n) noted by Dudek, Khoeini, and Pra lat in [13] and can be used to slightly improve a bound of $\hat{R}_r(C_n) > (3.75 - o(1))2^{-2} n$ (for odd n) given by Javadi and Miralaei in [21] (which uses Theorem 1 directly).




Figure 4: The value of c on the x-axis represents |V| = cn. The value of d on the y-axis represents |E| = dn. For a given value of c, the curve shows the maximum value of d so that if |E| 6 dn our proof gives a 2-coloring of G with no monochromatic P_n . The blue curve corresponds to Case 1, the solid red curve to Case 2, and the green curve to Case 3. Note that the minimum over the entire interval is 3.75 and occurs when c = 5/3.

4 More than two colors

The following statement implies the first part of Theorem 2.

Proposition 21 For all > 0, r > 2, and sufficiently large n, if G is a graph with at most $(\frac{(r-1)r}{2} + 2.75 -)n$ edges, then there exists an r-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. Let G = (V, E) be a connected graph with at most $(\frac{(r-1)r}{2} + 2.75 -)n$ edges. For r = 2, the result holds by Theorem 1. So let r > 3 and suppose the result is true in the case r - 1. If $N \in (r - 1)(n - 2)$, then we are done by Proposition 4; so suppose N > (r - 1)(n - 2) + 1. Let T be a spanning tree of G and apply Lemma 11 to get a forest F with no paths of order n and at least (r - 1)(n - 2) - 2r - 2 = (r - 1)n - 4r edges. Color the edges of the forest with color r. The number of remaining edges is at

most $(\frac{(r-1)r}{2} + 2.75 -)n - (r-1)n + 4r = (\frac{(r-2)(r-1)}{2} + 2.75 - {}^{0})n$ (where ${}^{0} = -4r/n > 0$ since n is sufficiently large) and thus we may apply induction to color the remaining edges with the remaining r-1 colors.

Definition 22. An affine plane of order q is a q-uniform hypergraph δwer tices (called points), with q(q + 1) edges (called lines) such that each pair of vertices is contained in exactly one edge.

It is well known that an affine plane of order q exists whenever q is a prime power (and it is unknown whether there exists an affine plane of non-prime power order). We collect two key properties of affine planes in the following proposition.

Proposition 23Let q > 2 be such that there exists an affine plane of order q. There exists a q + 1-coloring of the edges of k such that

- (i) every color class (called a parallelclass) consists of a collection of q vertex disjoint K_{a} 's, and
- (ii) every vertex v is contained in exactly one K_q of each color and the union of these q + 1 K_q 's incident with v covers all of $V(K_{q^2})$.

The following theorem implies the second part of Theorem We modify Krivelevich's proof [22, Theorem 8] in such a way that no color is "wasted" on the high degree vertices. This improves the lower bound from $((r-2)^2 - o(1))n$ to $((r-1)^2 - o(1))n$.

Proposition 24. Suppose that an affine plane of order q exists and suppose n is sufficiently large. For all graphs G with at most $\frac{2}{3}n-6q^4n^{0.9}=(q^2-o(1))n$ edges, there exists a q+1-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. Let G = (V, E) be a graph with $|E| 6 q^2 n - 6q^4 n^{0.9}$. Let

$$V_0 := v \in V(G)$$
 $d(v) > n^{0.1}$.

Then $q^2n > |E(G)| > \frac{1}{2}|V_0|n^{0.1}$ implies that $|V_0|$ 6 $2q^{2n^{0.9}}$. Now randomly partition $V \setminus V_0$ into q^2 parts V_1, \dots, V_d by placing each vertex into one of these sets independently with probability $1/q^2$. Let L be a line of the affine plane A_q on point set $[q^2]$. For each edge e in $G[V \setminus V_0]$, we assign color i to e if the endpoints of e are in distinct sets V_d and V_d where the unique line containing X and Y in A is in the I th parallel class of A_q . We color e arbitrarily if both of its endpoints are in V_X for some X.

For a line L of A_q , define the random variable $X := |E|_{x \in L} V_x$ |. Then

$$E[X_L] 6 \frac{1}{q^2} \cdot |E(G)| 6 n - 6q^2 n^{0.9}.$$

Since every vertex of $V \setminus V$ has degree at most h^1 , we have that moving any one vertex from V_X to V_Y can change X by at most $n^{0.1}$. Thus we may apply McDiarmid's inequality (Lemma 15) with $c_K = n^{0.1}$ for all K to conclude that

$$P X_L > n - 5q^2 n^{0.9} \ 6 \exp \ -\frac{(q^2 n^{0.9})^2}{2|V \setminus V_0| \cdot (n^{0.1})^2} = \exp -\Omega(n^{-0.6}) \ ,$$

where we used $|V \setminus V_0| \in |E| = 6 q^2 n$ in the last inequality. Thus taking a union bound over all (q + 1)q lines of A, we conclude that there exists a partition of $V \setminus V$ in which at most $n - 5q^2 n^{0.9}$ edges lie inside $V_x \in V_x$ for all lines $V_x \in V_x$. In other words, for all $V_x \in V_x$ is $(2q^2 n^{0.9}, P_n)$ -free. Suppose $V_x \in V_x \in V_x$ is such a partition.

Finally, we must color the edges incident with V_0 . We color the edges from V_0 to $V_0 \cup V_1$ arbitrarily, and for all $i \in [q^2] \setminus \{1\}$ we color the edges from V_0 to V_i the same as the color of the edges between V and V_i . By Proposition 23(ii), this accounts for all of the edges incident with V_0 , and since for all V_0 containing V_0 the graph induced by V_0 is V_0 is

5 Additional observations and conclusion

In this section we collect a few additional thoughts, none of which fit into into the main thread of the paper. The four observations below quantify the intuitive notion that if *G* is a graph having the property that every 2-coloring of the edges of *G* contains a monochromatic *R*, then *G* must be "expansive" in some sense.

For a graph G = (V, E), let S_V be the set of permutations of V. The *bandwidth*, ϕ of G is defined as

$$\phi(G) := \min_{f \in S} \max_{v} |f(u) - f(v)|.$$

Observation 25For all graphs G, if $\phi(G)$ 6 $\frac{n}{2}$ – 1, then there is a 2-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. Choose an ordering f of V(G) which witnesses the bandwidth of G; i.e.

$$\max_{uv \in E} |f(u) - f(v)| = \varphi(G).$$

Now split the vertices into sets V_1, \dots, t , with $|V_1| = \dots = |V_{t-1}| = b \frac{n}{2} - 1c$ and $|V_t| = 0$. For all odd $i \in [t]$, color the edges from V to $V_i \cup V_{i+1}$ red, and for all even $i \in [t]$ color the edges from V to $V_i \cup V_{i+1}$ blue.

A depth first search (DFS) tree (or normal tree) T rooted at x in a graph G is a subtree of G such that for all $uv \in E(G)$ with $u, v \in V(T)$, either v is on the x - v path in T or u is on the x - u path in T.

For a connected subgraph H of a graph G and vertices $u, v \in V(H)$, let $d_H(u, v)$ be the length of the shortest path between u and v in HA breadth first search (BFS) tree T rooted at x is a subtree of G such that for all $v \in V(T)$, $d_T(x, v) = d_G(x, v)$. Such a tree has the property that for all $uv \in E(G)$ with $u, v \in V(T)$, $|d_T(x, u) - d_T(x, v)|$ 6 1. The vertices at each fixed distance from the root are called the *levels* of T It is well known that for every connected graph G and every vertex $x \in V(G)$, there exists a spanning DFS tree T rooted at x and a spanning BFS tree rooted at x.

Using the notation for rooted trees from the proof of Lemma 14, we have the following observation.

Observation 26. et G be a connected graphlf there exists a vertex x and a DFS tree T rooted at x so that every child $y \in C(x)$ satisfies $|S(T, y)| 6^{\frac{5n}{4}} - 2$, then there exists a 2-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. For each sub-tree S(T, y) where $y \in C(x)$, we partition the vertices of S(T, y) into sets A_y and B_y where $|A_y| = 6 \frac{n}{4} - 1$, $y \in A_y$ and $|B_y| = 6 \frac{n}{4} - 1$. Let $A = \{x\} \cup \{x\} \cup \{y \in C(x)\} = 1$. Let $A = \{x\} \cup \{y \in C(x)\} = 1$. We color the edges of G within G blue and the edges from G to G and G includes G in

Observation 27Let G be a connected graphlf there exists a vertex x and a BFS tree T rooted at x such that every pair of consecutive levels of T have fewer than n vertices, then there exists a 2-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. For all i > 0, let $D_i = \{v : d_{\tau}(x, v) = i\}$. For all j > 0, color the edges from D_{2j} to D_{2j+1} red and the edges from D_{2j+1} to $D_{2j+1} \cup D_{2j+2}$ blue. By the property of BFS trees, this accounts for every edge in G. Since every two consecutive levels contain fewer than n vertices, there are no monochromatic paths of order n.

The following observation was inspired by Figure 2 in both [4] and [5].

Observation 28^{lf} G is a graph on N vertices with $\alpha(G) > N - (n-3)$, then there exists a 2-coloring of the edges of G such that every monochromatic path has order less than n.

Proof. Let *S* be an independent set of order at least N - (n - 3) and partition the vertices of $V(G) \setminus S$ into disjoint sets X, Y with |X|, $|Y| = 6 - \frac{n}{2} - 1$. Color all edges incident with X red and color all edges incident with Y blue (so edges between X and Y can be either color). The longest monochromatic path has order at most $2\frac{N}{2} - 1 + 1 = n - 1$.

Finally, we end with the following question which relates to the upper bound on the size-Ramsey number of a path.

Question 29.What is the largest monochromatic path one can find in an arbitrary 2-coloring of a *d*-regular graph on *N* vertices?

For instance, suppose it is always possible to 2-color the edges of 5-regular graph on N vertices (with N sufficiently large) so that there is no monochromatic path of orde $\frac{N}{30}$. This would imply that all 5-regular graphs on at most 30n vertices (which have at most 75n edges) have a 2-coloring with no monochromatic; $\frac{1}{2}$ n other words, 5-regular graphs could never improve the current best [15] upper bounce $\frac{1}{2}$ ($\frac{1}{2}$) 6 74 $\frac{1}{2}$.

Acknowledgements

We thank two very thorough referees for their careful reading of the paper and their helpful comments.

References

- [1] N. Alon. On the edge-expansion of graphs. *Combinatorics, Probability and Computing* **6**, no. 2 (1997):145–152.
- [2] N. Alon, P. Hamburger, and A. V. Kostochka. Regular honest graphs, isoperimetric numbers, and bisection of weighted graphs. *European Journal Combinatorics* **20**, no. 6 (1999): 469–481.
- [3] J. Balogh, A. Dudek, L. Li. An analogue of the Erdős-Gallai theorem for random graphs. *European Journabf Combinatorics* **91** (2021), Paper No. 103200, 11 pp.
- [4] J. Balogh, A. Kostochka, M. Lavrov, and X. Liu. Monochromatic connected matchings in 2-edge-colored multipartite graphs. arXiv:1905.04653 (2019).
- [5] J. Balogh, A. Kostochka, M. Lavrov, and X. Liu. Long monochromatic paths and cycles in 2-edge-colored multipartite graphs. Moscow Journal of Combinatorics and Number Theory **9**, no. 1 (2020):55–100.
- [6] J. Beck. On size Ramsey number of paths,trees, and circuits. I. *Journal of Graph Theory* **7**, no. 1 (1983):115–129.
- [7] J. Beck. On size Ramsey number of paths, trees and circuits. II. *In Mathematics of Ramsey theory*, pp. 34–45. Springer, Berlin, Heidelberg, 1990.
- [8] J. Bermond, J. Fouquet, M. Habib, and B. Peroche. On linear *k*-arboricity. *Discrete Mathematics* **52**, no. 2-3 (1984):123–132.
- [9] H. Bielak. Remarks on the size Ramsey number ofgraphs. *Periodica Mathematica Hungarica* **18**, no. 1 (1987)27–38.
- [10] B. Bollob´as. Extremal graph theory with emphasis on probabilistic methods. No. 62. American Mathematical Soc., 1986.
- [11] B. Bollobás. Random graphs. No. 73. Cambridge University Press, 2001.
- [12] R. Diestel. Graph theory. Fifth edition. Graduate Texts in Mathematics, 173. Springer, Berlin, 2018. xviii+428 pp.
- [13] A. Dudek, F. Khoeini, P. Pra lat. Size-Ramsey numbers of cycles versus a path. *Discrete Mathematics* **341**, no. 7 (2018)2095–2103.
- [14] A. Dudek, P. Pra lat. An alternative proof of the linearity of the size-Ramsey number of paths. *Combinatorics, Probability and Computing* **24**, no. 3 (2015)51–555.
- [15] A. Dudek, P. Pra lat. On some multicolor Ramsey properties of random graphs. *SIAM Journal on Discrete Mathematics* **31**, no. 3 (2017)2079–2092.
- [16] A. Dudek, P. Pra lat. Note on the Multicolour Size-Ramsey Number for Paths. *Electronic Journal of Combinatorics* **25**, no. 3, (2018):P3.35.

- [17] P. Erd"os.On the combinatorial problems which I would most like to see solved. *Combinatorica* **1**, no. 1 (1981):25–42.
- [18] A.M. Frieze, M. Karoński. Introduction to random graphs. Cambridge University Press, Cambridge, 2016. xvii+464 pp.
- [19] L. Gerencs'er and A. Gy'and. On Ramsey-type problems. *Ann. Univ. Sci. Budapest. E"otv"os Sect. Math* **10** (1967)67–170.
- [20] A. Gy'artás. Large monochromatic components in edge colorings of graphssurvey. In Ramsey Theory, pp. 77–96. Birkhäuser, Boston, MA, 2011.
- [21] R. Javadi, M. Miralaei. Multicolor Size-Ramsey Number of Cycles. arXiv:2106.16023 (2021).
- [22] M. Krivelevich. Long cycles in locally expanding graphs,with applications. *Combinatorica* **39**, no. 1 (2019):135–151.
- [23] A. V. Kostochka, L. S. Melnikov. On bounds of the bisection width of cubic graphs. In Fourth Czechoslovakian Symposium on CombinatoricsGraphs and Complexity, pp. 151–154. Elsevier, 1992.
- [24] S. Letzter. Path Ramsey number for random graphs. *Combinatorics, Probability and Computing* **25**, no. 4 (2016):612–622.
- [25] C. McDiarmid. On the method of bounded differences. *Surveys in combinatorics* **141**, no. 1 (1989): 148–188.
- [26] A. Vince. The integrity of a cubic graph. *Discrete Applied Mathematics* **140**, no. 1-3 (2004): 223–239.
- [27] S. Yongqi, Y. Yuansheng, X. Feng, and L. Bingxi. New lower bounds on the multicolor Ramsey numbers $R(C_{2m})$. *Graphs and Combinatorics* **22**, no. 2 (2006)283–288.