



Lior Fishman · Dmitry Kleinbock · Keith Merrill · David Simmons

Intrinsic Diophantine approximation on quadric hypersurfaces

Received April 19, 2019; revised February 21, 2021

Abstract. We consider the question of how well points in a quadric hypersurface $M \subseteq \mathbb{R}^d$ can be approximated by rational points of $\mathbb{Q}^d \cap M$. This contrasts with the more common setup of approximating points in a manifold by all rational points in \mathbb{Q}^d . We provide complete answers to major questions of Diophantine approximation in this context. Of particular interest are the impact of the real and rational ranks of the defining quadratic form, quantities whose roles in Diophantine approximation have never been previously elucidated. Our methods include a correspondence between the intrinsic Diophantine approximation theory on a rational quadric hypersurface and the dynamics of the group of projective transformations which preserve that hypersurface, similar to earlier results in the non-intrinsic setting due to Dani (1986) and Kleinbock–Margulis (1999).

Keywords. Quadratic forms, lattices, intrinsic approximation, Dirichlet’s theorem, Khintchine’s theorem

Contents

1. Introduction and motivation	1046
2. Main results	1050
3. Preliminaries on quadratic forms and lattices	1056
4. The correspondence principle	1063
5. A Dirichlet-type theorem	1068
6. Khintchine-type theorems and counting of rational points	1078
7. Proof of Theorem 6.5 modulo a volume computation	1081
8. Estimating the measure μ_{R, Λ_*}	1087
9. The exceptional quadric hypersurface	1094
References	1099

Lior Fishman: Department of Mathematics, University of North Texas, 1155 Union Circle #311430, Denton, TX 76203-5017, USA; lior.fishman@unt.edu

Dmitry Kleinbock (corresponding author): Department of Mathematics, Brandeis University, 415 South Street, Waltham, MA 02454-9110, USA; kleinboc@brandeis.edu

Keith Merrill: Department of Mathematics, Brandeis University, 415 South Street, Waltham, MA 02454-9110, USA; merrill2@brandeis.edu

David Simmons: Department of Mathematics, University of York, Heslington, York YO10 5DD, United Kingdom; david.simmons@york.ac.uk

Mathematics Subject Classification (2020): Primary 11J83; Secondary 11H55, 37A17

1. Introduction and motivation

Classical theorems in Diophantine approximation theory address questions regarding the way points $\mathbf{x} \in \mathbb{R}^d$ are approximated by rational points, considering the trade-off between the height of the rational point – the size of its denominator – and its distance to \mathbf{x} ; see [13, 49] for a general introduction. Often \mathbf{x} is assumed to lie on a certain subset of \mathbb{R}^d , for example a smooth manifold M , leading to *Diophantine approximation on manifolds*. This area of research has experienced rapid progress during the last two decades, owing much of it to methods coming from flows on homogeneous spaces.

It was observed in [11, 17, 18] that all sufficiently good rational approximants to points on certain rational varieties must in fact be *intrinsic* – that is, they are rational points lying on the variety itself. These results, in part, have motivated a new field of *intrinsic approximation*, which examines the degree to which points on a manifold or variety can be approximated by rationals lying on that same subset. Questions about the quality of these approximations were raised already by Lang [40] and Mahler [43]. Following some recent results on quadric hypersurfaces [26, 27, 50] and a comprehensive treatment of Diophantine approximation on spheres [34], this paper seeks to fully explore the topic of intrinsic approximation on quadrics. One of the most novel and important aspects of our work is an elucidation of the role of the \mathbb{Q} -rank and the \mathbb{R} -rank of the defining quadratic form (see Definition 3.3). It turns out there are qualitative differences between the intrinsic approximation theories of forms with different rank pairs, highlighting the importance of rank, rather than the dimension of the hypersurface. In particular, we will see below that our Dirichlet-type theorem, Theorem 5.1, is *independent* of the dimension d , but changes depending on whether the \mathbb{Q} -rank and \mathbb{R} -rank are equal or different. We remark that [34] considers only the case where both ranks equal 1; therefore the dependence on the ranks is not explored there, and significant new ideas have had to be developed in the present paper.

Convention 1. The symbols \lesssim , \gtrsim , and \asymp will denote asymptotics; a subscript of $+$ indicates that the asymptotic is additive, and a subscript of \times indicates that it is multiplicative. For example, $A \lesssim_{\times, K} B$ means that there exists a constant $C > 0$ (the *implied constant*), depending only on K , such that $A \leq CB$. Furthermore, $A \lesssim_{+, \times} B$ means that there exist constants $C_1, C_2 > 0$ so that $A \leq C_1 B + C_2$. In general, dependence of the implied constant(s) on universal objects such as the manifold M will be omitted from the notation.

Convention 2. For any $c \geq 0$, we let

$$\psi_c(q) := \frac{1}{q^c}.$$

Convention 3. The symbol \lhd will be used to indicate the end of a nested proof.

Glossary of notation. For the reader's convenience, we summarize a list of notations and terminology in the order that they appear in the sequel.

M	a complete metric space	(Section 1)
\mathcal{Q}	a countable subset of M	(Section 1)
H	a height function	(Section 1)
$\text{BA}(\psi, M, \mathcal{Q}, H)$	the set of badly approximable points	(Section 1)
$\text{WA}(\psi, M, \mathcal{Q}, H)$	the set of well approximable points	(Section 1)
$\text{A}(\psi, M, \mathcal{Q}, H)$	the set of ψ -approximable points	(Section 1)
H_{std}	the standard height on projective space	(Section 2)
Q	a quadratic form on \mathbb{R}^{d+1}	(Section 2)
L_Q	the light cone of Q	(Section 2)
M_Q	a nonsingular rational quadric hypersurface	(Section 2)
$p_{\mathbb{R}}$	the real rank of Q	(Section 2)
$p_{\mathbb{Q}}$	the rational rank of Q	(Section 2)
Q_{aff}	a quadratic polynomial with integer coefficients on \mathbb{R}^d	(Section 2)
$M_{Q_{\text{aff}}}$	the nonsingular rational quadric hypersurface associated to Q_{aff}	(Section 2)
$\text{A}_{M_Q}(\psi)$	the set of ψ -approximable points on M_Q	(Section 2)
BA_{M_Q}	the set of badly approximable points on M_Q	(Section 2)
Q_0	the exceptional quadratic form	(Section 2)
B_Q	the symmetric, bilinear form associated to Q	(Section 3)
\mathcal{L}_m	$\sum_{i=0}^{m-1} \mathbb{R} \mathbf{e}_i$	(Section 3)
\tilde{Q}	the remainder of the form Q after normalizing	(Section 3)
$\hat{\phi}$	the reverse of the matrix ϕ	(Section 3)
g_{ϕ}	$\begin{bmatrix} \phi & & \\ & I_{d+1-2m} & \\ & & \hat{\phi} \end{bmatrix}$	(Section 3)
$g_{\mathbf{t}}$	$g_{\text{diag}(e^{-t_0}, \dots, e^{-t_{m-1}})}$	(Section 3)
g_t	$\begin{bmatrix} e^{-t} & & \\ & I_{d-1} & \\ & & e^t \end{bmatrix}$	(Section 3)
$\delta(\Lambda)$	$\min_{\mathbf{p} \in \Lambda \setminus \{0\}} \ \mathbf{p}\ $	(Section 3)
$\delta_Q(\Lambda)$	$\min_{\mathbf{p} \in \Lambda \cap L_Q \setminus \{0\}} \ \mathbf{p}\ $	(Section 3)
$O(Q)$	$\{g \in \text{SL}_{d+1}^{\pm}(\mathbb{R}) : Q \circ g = R\}$	(Section 3)
Ω_Q	the space of Q -arithmetic lattices	(Section 3)
Ω_d	the space of all lattices in \mathbb{R}^{d+1}	(Section 3)
$O(Q; \Lambda)$	the stabilizer of Λ under the action of $O(Q)$	(Section 3)
$\Omega_{Q, \Lambda}$	the homogeneous space $O(Q)/O(Q; \Lambda)$	(Section 3)
π_1, π_2	projections $O(R) \rightarrow M_Q$ and $O(R) \rightarrow \Omega_{R, \Lambda_*}$	(Section 4)
Λ_{pr}	the set of primitive vectors of Λ	(Section 4)
$\mu_R, \mu_{R, \Lambda_*}$	Haar measures on $O(R)$ and Ω_{R, Λ_*}	(Section 4)
$\text{Codiam}(\Gamma)$	the diameter of the quotient space $\text{Span}(\Gamma)/\Gamma$	(Section 5)

$N_M(T)$	$\#([\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M : H_{\text{std}}([\mathbf{r}]) \leq T)$	(Section 6)
$S_{\Delta,z}$	$\{x \in X : \Delta(x) \geq z\}$	(Section 7)
$\Phi_{\Delta}(z)$	$\mu(S_{\Delta,z})$, the tail distribution function of Δ	(Section 7)
$\varphi^{(C)}(x)$	$\max_{\text{dist}_X(x',x) \leq C} \varphi(x')$	(Section 8)
$\varphi_C(x)$	$\min_{\text{dist}_X(x',x) \leq C} \varphi(x')$	(Section 8)
P	a parabolic subgroup of G	(Section 8)
ρ_P	the modular function of P	(Section 8)
A	a maximal \mathbb{Q} -split torus	(Section 8)
ρ	the sum of the positive roots of A , counted with multiplicity	(Section 8)

1.1. General terminology and basic problems in metric Diophantine approximation

In order to review some known facts and state our theorems, let us first introduce basic notations which we will follow throughout the paper (some of it has been introduced in a different context in [23]).

Definition 1.1. By a *Diophantine triple*, we will mean a triple (M, \mathcal{Q}, H) , where M is a closed subset of a complete metric space (X, dist) , \mathcal{Q} is a countable subset of X whose closure contains M , and H is a function from \mathcal{Q} to $(0, \infty)$.

Definition 1.2. Say that a nonincreasing¹ function $\psi: (0, \infty) \rightarrow (0, \infty)$ is a *Dirichlet function* for (M, \mathcal{Q}, H) if, for every $\mathbf{x} \in M$, there exist $C_{\mathbf{x}} > 0$ and a sequence $(\mathbf{r}_n)_{n=1}^{\infty}$ in \mathcal{Q} such that

$$\mathbf{r}_n \xrightarrow{n} \mathbf{x} \quad \text{and} \quad \text{dist}(\mathbf{r}_n, \mathbf{x}) \leq C_{\mathbf{x}} \psi(H(\mathbf{r}_n)). \quad (1.1)$$

If $C_{\mathbf{x}}$ can be chosen independent of \mathbf{x} , then we call ψ *uniformly Dirichlet*.

When ψ is a Dirichlet function, it is often important to understand whether a faster decaying function can also be Dirichlet. We formalize this thought in the next definition.

Definition 1.3. A Dirichlet function ψ is *optimal* for (M, \mathcal{Q}, H) if there is no function φ which is Dirichlet for (M, \mathcal{Q}, H) and satisfies $\frac{\varphi(x)}{\psi(x)} \rightarrow 0$ as $x \rightarrow \infty$.

It turns out that the optimality of ψ is under some fairly general assumptions equivalent to the existence of so-called *badly approximable* points. This notion deserves a special definition.

Definition 1.4. If (M, \mathcal{Q}, H) is a Diophantine triple and if $\psi: (0, \infty) \rightarrow (0, \infty)$, then a point $\mathbf{x} \in M$ is said to be *badly approximable* with respect to ψ if there exists $\varepsilon > 0$ such that, for all $\mathbf{r} \in \mathcal{Q}$,

$$\text{dist}(\mathbf{r}, \mathbf{x}) \geq \varepsilon \psi(H(\mathbf{r})).$$

The set of such points will be denoted $\text{BA}(\psi, M, \mathcal{Q}, H)$, and its complement will be denoted $\text{WA}(\psi, M, \mathcal{Q}, H)$ (the set of *well approximable* points).

¹The approximating functions ψ will be assumed to be nonincreasing throughout the paper.

If $\text{BA}(\psi, M, \mathcal{Q}, H) \neq \emptyset$, then it is easy to see that ψ is an optimal Dirichlet function for (M, \mathcal{Q}, H) .² Note also that $\mathcal{Q} \cap M$ is always contained in $\text{WA}(\psi, M, \mathcal{Q}, H)$.

Definition 1.5. Also, we will let

$$\begin{aligned} \text{A}(\psi, M, \mathcal{Q}, H) &:= \{\mathbf{x} \in M : \text{there exist infinitely many } \mathbf{r} \in \mathcal{Q} \\ &\quad \text{with } \text{dist}(\mathbf{r}, \mathbf{x}) \leq \psi(H(\mathbf{r})))\} \\ &= \limsup_{\mathbf{r} \in \mathcal{Q}} (B(\mathbf{r}, \psi(H(\mathbf{r}))) \cap M) \end{aligned}$$

be the set of ψ -approximable points. Note that

$$\text{WA}(\psi, M, \mathcal{Q}, H) = (\mathcal{Q} \cap M) \cup \bigcap_{\varepsilon > 0} \text{A}(\varepsilon\psi, M, \mathcal{Q}, H).$$

We can now list a few basic general problems one can pose, given a Diophantine triple (M, \mathcal{Q}, H) .

- (1) Find a Dirichlet function for (M, \mathcal{Q}, H) . Even better – find an optimal one; determine whether or not it is uniformly Dirichlet.
- (2) Find a function ψ such that $\text{BA}(\psi, M, \mathcal{Q}, H) \neq \emptyset$. Even better – do it for a Dirichlet function, thus proving it to be optimal. In the latter case, determine how big is the set $\text{BA}(\psi, M, \mathcal{Q}, H)$, e.g. in terms of its Hausdorff dimension.
- (3) Given a function ψ and a measure on M , what is the measure of $\text{A}(\psi, M, \mathcal{Q}, H)$? This measure could be a Riemannian volume on M if the latter is a manifold or, more generally, the Hausdorff measure relative to some dimension function. A special case of the last question is a determination of the Hausdorff dimension of $\text{A}(\psi, M, \mathcal{Q}, H)$.

Note that, since $\text{A}(\psi, M, \mathcal{Q}, H)$ is a \limsup set, the easy direction of the Borel–Cantelli lemma shows that, for any measure μ on M , if the series

$$\sum_{\mathbf{r} \in \mathcal{Q} \cap U} \mu(B(\mathbf{r}, \psi(H(\mathbf{r}))) \cap M) \tag{1.2}$$

converges whenever U is a bounded subset of X , then one has $\mu(\text{A}(\psi, M, \mathcal{Q}, H)) = 0$. The hope is that for “nice” measures the (much harder) complementary divergence case can be established. Also, in general, it is not clear how to explicitly decide for which functions ψ the sum (1.2) converges or diverges; for that, one often needs extra information concerning the number of points of \mathcal{Q} satisfying a given height bound.

1.2. Diophantine approximation in \mathbb{R}^d

In the classical Diophantine approximation setup, one has $X = M = \mathbb{R}^d$, $\mathcal{Q} = \mathbb{Q}^d$, and

$$H(\mathbf{r}) = H_{\text{std}}(\mathbf{r}) := q, \quad \text{where } \mathbf{r} = \mathbf{p}/q \text{ is written in reduced form} \tag{1.3}$$

(this will be referred to as the *standard* height).

²See [23, Theorem 2.6] where this is stated under the assumption that $M = X$; one can check that the latter assumption is not necessary for the argument. Furthermore, the converse is true assuming the σ -compactness of M ; see [23, Proposition 2.7].

Dirichlet's theorem asserts that, for all $\mathbf{x} \in \mathbb{R}^d$ and $T \geq 1$, there exists $\mathbf{p}/q \in \mathbb{Q}^d$ with $q \leq T$ satisfying

$$\text{dist}\left(\frac{\mathbf{p}}{q}, \mathbf{x}\right) \leq \frac{C}{qT^{1/d}}, \quad (1.4)$$

where $C > 0$ is a constant depending on the choice of the norm on \mathbb{R}^d . A corollary is that

$$\psi_{1+1/d} \text{ is uniformly Dirichlet for } (\mathbb{R}^d, \mathbb{Q}^d, H_{\text{std}}) \quad (1.5)$$

(see Convention 2). Note that, when the distance is given by the supremum norm on \mathbb{R}^d , one can take $C = 1$ in (1.4), and thus $C_x \equiv 1$ in (1.1). (It is clear that the property of ψ being Dirichlet or uniformly Dirichlet does not depend on the choice of the norm.)

On the other hand, it is well known that, for all d , the set

$$\text{BA}_d := \text{BA}(\psi_{1+1/d}, \mathbb{R}^d, \mathbb{Q}^d, H_{\text{std}})$$

of badly approximable vectors in \mathbb{R}^d is nonempty (see e.g. [47, 49]), implying the optimality of $\psi_{1+1/d}$ as a Dirichlet function for $(\mathbb{R}^d, \mathbb{Q}^d, H)$. Indeed, Schmidt [49] showed that

$$\text{BA}_d \text{ has full Hausdorff dimension in } \mathbb{R}^d, \quad (1.6)$$

generalizing a result of Jarník [31], who proved the case $d = 1$ of (1.6). We shall refer to (1.6) as the *Jarník–Schmidt theorem*. Note that, together, Dirichlet's theorem and the Jarník–Schmidt theorem solve problems (1) and (2) above for the case of the Diophantine triple $(\mathbb{R}^d, \mathbb{Q}^d, H_{\text{std}})$.

Resolving problem (3) gives rise to theorems of Khintchine and of Jarník–Besicovitch. For convenience, let us denote $\text{A}(\psi, \mathbb{R}^d, \mathbb{Q}^d, H_{\text{std}})$ by $\text{A}_d(\psi)$. If λ is Lebesgue measure on \mathbb{R}^d , it was proven by Khintchine [33] that, if ψ is nonincreasing³, $\text{A}_d(\psi)$ is either null or conull depending on whether the series $\sum_{q=1}^{\infty} q^{d-1} \psi(q)^d$ converges or diverges. More generally, for $0 < s < d$, one can replace λ with \mathcal{H}^s , the s -dimensional Hausdorff measure, and get the Jarník–Besicovitch theorem [5, 32]: $\mathcal{H}^s(\text{A}_d(\psi))$ is either 0 or ∞ depending on whether the series $\sum_{q=1}^{\infty} q^{d-1} \psi(q)^s$ converges or diverges.

2. Main results

Convention 4. Throughout the paper, propositions which are proven later in the paper will be numbered according to the section they are proven in.

We now consider the main setup of the paper, namely that of intrinsic approximation. One way to do it is to take $X = \mathbb{R}^d$, choose a submanifold M of \mathbb{R}^d , and let $\mathcal{Q} = \mathbb{Q}^d \cap M$ and $H = H_{\text{std}}$ as in (1.3). However, we have chosen a different approach: state and prove the main results of the paper for submanifolds of projective spaces. This way, in most cases, statements of results and their proofs become more natural and transparent; see Remark 2.1 below.

³The monotonicity assumption is not needed if $d > 1$; see [25].

Let $\mathbb{P}_{\mathbb{R}}^d$ denote the d -dimensional real projective space, and let $\pi: \mathbb{R}^{d+1} \setminus \{\mathbf{0}\} \rightarrow \mathbb{P}_{\mathbb{R}}^d$ be the quotient map $\pi(\mathbf{x}) := [\mathbf{x}]$ so that $[\mathbf{r}\mathbf{x}] = [\mathbf{x}]$. The distance on $\mathbb{P}_{\mathbb{R}}^d$ will be given by the formula $\text{dist}([\mathbf{x}], [\mathbf{y}]) = \min(\|\mathbf{y} - \mathbf{x}\|, \|\mathbf{y} + \mathbf{x}\|)$ ($\|\mathbf{x}\| = \|\mathbf{y}\| = 1$). For a subset S of \mathbb{R}^{d+1} , we let $[S] = \pi(S \setminus \{\mathbf{0}\})$. With some abuse of notation, let us define the *standard height function* $H_{\text{std}}: \mathbb{P}_{\mathbb{Q}}^d \rightarrow \mathbb{N}$ by the formula

$$H_{\text{std}}([\mathbf{p}]) = \|\mathbf{p}\|, \quad \text{where } \mathbf{p} \text{ is the unique (up to a sign) primitive integer representative of } [\mathbf{p}].$$

Here and elsewhere, $\|\cdot\|$ represents the max norm.

Remark 2.1. To see the difference between results for affine and projective spaces, note that if $\iota_d: \mathbb{R}^d \rightarrow \mathbb{P}_{\mathbb{R}}^d$ is given by the formula $\iota_d(\mathbf{x}) = [(1, \mathbf{x})]$ and if $B \subseteq \mathbb{R}^d$ is a bounded set, then $\iota_d|_B$ is bi-Lipschitz and

$$H_{\text{std}}(\iota_d(\mathbf{r})) \asymp_{\times, B} H(\mathbf{r}) \quad \text{for all } \mathbf{r} \in \mathbb{Q}^d \cap B. \quad (2.1)$$

In particular, the Diophantine triples

$$\begin{aligned} T_{\text{aff}} &:= (M, \mathbb{Q}^d \cap M, H_{\text{std}}), \\ T_{\text{proj}} &:= (\iota_d(M), \mathbb{P}_{\mathbb{Q}}^d \cap \iota_d(M), H_{\text{std}}) \end{aligned}$$

are “locally isomorphic”. However, both the bi-Lipschitz constant and the implied constant of (2.1) depend on the chosen bounded set B . Thus concepts which are robust under point-dependent multiplicative constants will not be affected by the transformation. For example, whether or not a function is Dirichlet will be the same for the triples T_{aff} and T_{proj} , but it is conceivable that a function could be uniformly Dirichlet for the triple T_{proj} but not for the triple T_{aff} .

Because of this difference, it is perhaps worthwhile to give a justification for why we are stating our results in projective space rather than affinely. The simplest answer to this question is that the projective statements are closest to how the results are actually proven. Moreover, in those cases where projective statements cannot be reformulated as affine statements, we feel it is important to keep the full strength of the projective theorem. To give a simple example, consider the classical Dirichlet theorem. By examining its proof, we can deduce that

$$\psi_{1+1/d} \text{ is uniformly Dirichlet for } (\mathbb{P}_{\mathbb{R}}^d, \mathbb{P}_{\mathbb{Q}}^d, H_{\text{std}}). \quad (2.2)$$

This result is stronger than the classical (1.5), in the sense that simply translating (1.5) to projective space along the lines indicated above does not yield (2.2), while translating (2.2) to affine space yields (1.5) at least on the unit cube $[0, 1]^d$, and applying translations recovers the full force of (1.5).

To guide the reader, we have included affine corollaries after most of the main results. Each affine corollary can be deduced from its corresponding result together with Remark 2.1. We omit those affine corollaries which would merely be restatements of the theorems with $\mathbb{P}_{\mathbb{R}}^d$ replaced by \mathbb{R}^d .

In the following theorems, we fix $d \geq 2$ and let Q be a nonsingular (see Definition 3.2) quadratic form on \mathbb{R}^{d+1} with integer coefficients. (See Remark 2.8 for a discussion of the singular case.) Denote by

$$L_Q := \{\mathbf{x} \in \mathbb{R}^{d+1} : Q(\mathbf{x}) = 0\} \quad (2.3)$$

the *light cone* of Q , and let $M_Q = [L_Q]$. Manifolds M_Q of this form are called *nonsingular rational quadric hypersurfaces*.

We will denote by $p_{\mathbb{R}}$ the \mathbb{R} -rank of Q , defined as the dimension of any maximal totally isotropic (with respect to Q) subspace of \mathbb{R}^{d+1} . Similarly, $p_{\mathbb{Q}}$ will stand for the \mathbb{Q} -rank of Q , i.e. the dimension of any maximal totally isotropic rational subspace of \mathbb{R}^{d+1} . Clearly, $p_{\mathbb{R}} \geq p_{\mathbb{Q}}$; see Section 3.2 for more details. To avoid trivialities, in our theorems, we will make the standing assumption that $p_{\mathbb{Q}} \geq 1$ or, equivalently, that

$$\mathbb{P}_{\mathbb{Q}}^d \cap M_Q \neq \emptyset. \quad (2.4)$$

Note that Meyer's theorem states that (2.4) is satisfied as soon as $d \geq 4$ and $M_Q \neq \emptyset$. Moreover, if $d = 2$ or 3 , the Hasse–Minkowski theorem (e.g. [7, Theorem 1 on p. 61]) allows one to determine computationally whether (2.4) is satisfied for any given quadratic form Q ; cf. [7, Chapter 1, Section 7, in particular the remarks on the top of page 62].

For the affine corollaries to our theorems, we consider a quadratic polynomial

$$Q_{\text{aff}}: \mathbb{R}^d \rightarrow \mathbb{R}$$

with integer coefficients, and we let $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be the projectivization of Q_{aff} , that is, the unique homogeneous quadratic polynomial (i.e. quadratic form) Q on \mathbb{R}^{d+1} such that $Q(1, \mathbf{x}) = Q_{\text{aff}}(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^d$. Then $M_{Q_{\text{aff}}}$, the zero set of Q_{aff} , is equal to $\iota_d^{-1}(M_Q)$. We call $M_{Q_{\text{aff}}}$ a nonsingular rational quadric hypersurface whenever M_Q is. Note that it may be the case that M_Q is singular due to “singularities at infinity” rather than singularities at finite points; in this case, we still consider the hypersurface $M_{Q_{\text{aff}}}$ to be singular despite its having no “singular points”.

The problem of intrinsic approximation on M_Q was implicitly considered by Druțu in [18] where the Hausdorff dimension of sets $A_{M_Q}(\psi)$ was computed. (Druțu actually studied ambient approximation on M_Q and, generalizing an earlier result of Dickinson and Dodson [17, Lemma 1], showed that it reduces to intrinsic approximation if ψ is assumed to decay fast enough.) The case $Q(\mathbf{x}) = x_1^2 + \cdots + x_d^2 - x_0^2$ was recently considered in [34].⁴ One of the theorems from the latter paper asserts⁵ that there exists $C > 0$ (possibly depending on d) such that, for all $[\mathbf{x}] \in M_Q$ and for all $T \geq T_0$, there exists $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ with

$$H_{\text{std}}([\mathbf{r}]) \leq T \quad \text{and} \quad \text{dist}([\mathbf{r}], [\mathbf{x}]) \leq \frac{C}{\sqrt{H_{\text{std}}([\mathbf{r}])T}}. \quad (2.5)$$

⁴The article [34] is written in the affine setup; specifically, the manifold $\mathbb{S}^{d-1} \subseteq \mathbb{R}^d$ is discussed. Since this set is compact, Remark 2.1 gives an exact correspondence for Diophantine results in \mathbb{S}^{d-1} and those in $\iota_d(\mathbb{S}^{d-1}) = M_Q$.

⁵Moshchevitin [46] has recently provided an elementary proof of this assertion for the case $M_{Q_{\text{aff}}} = \mathbb{S}^2$. His proof gives an explicit value for the constant C appearing in (2.5).

In particular, it follows that ψ_1 is uniformly Dirichlet for intrinsic approximation on M_Q . It was also shown in [34] that

- (i) ψ_1 is optimal – moreover, $\text{BA}_{M_Q}(\psi_1)$ has full Hausdorff dimension,
- (ii) for any $\psi: \mathbb{N} \rightarrow (0, \infty)$ such that

the function $q \mapsto q\psi(q)$ is nonincreasing,

the Lebesgue measure of $\text{A}_{M_Q}(\psi)$ is full (resp. zero) if and only if the sum

$$\sum_{q=1}^{\infty} q^{d-2} \psi(q)^{d-1}$$

diverges (resp. converges).

The last statement was also shown to imply, via the Mass Transference Principle of Beresnevich and Velani [3, Theorem 2], a similar statement for Hausdorff measures.

In the present paper, we generalize all the aforementioned results to the case of arbitrary quadric hypersurfaces.

Theorem 5.1 (Dirichlet-type theorem for quadric hypersurfaces). *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Then*

- (i) *the function ψ_1 is Dirichlet for intrinsic approximation on M_Q .*
- (ii) *ψ_1 is uniformly Dirichlet if and only if $p_{\mathbb{Q}} = p_{\mathbb{R}}$.*
- (iii) *The following are equivalent:*
 - (A) $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$.
 - (B) (“Strong Dirichlet”) *There exist $C, T_0 > 0$ such that, for all $[\mathbf{x}] \in M_Q$ and for all $T \geq T_0$, there exists $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ such that (2.5) holds.*
 - (C) *The set*

$$\{[\mathbf{x}] \in M_Q : \text{there exist } C, T_0 > 0 \text{ such that, for all } T \geq T_0, \text{ there exists } [\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q \text{ satisfying (2.5)}\}$$

has positive λ_{M_Q} -measure.

Affine Corollary. *Let $M_{Q_{\text{aff}}} \subseteq \mathbb{R}^d$ be a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Then*

- (i) *the function ψ_1 is Dirichlet for intrinsic approximation on $M_{Q_{\text{aff}}}$.*
- (ii) *If $p_{\mathbb{Q}} = p_{\mathbb{R}}$, then ψ_1 is uniformly Dirichlet on compact subsets of $M_{Q_{\text{aff}}}$.*
- (iii) *The following are equivalent:*
 - (A) $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$.
 - (B) (“Strong Dirichlet”) *For every compact set $K \subseteq M_{Q_{\text{aff}}}$, there exist $C, T_0 > 0$ such that, for all $\mathbf{x} \in K$ and for all $T \geq T_0$, there exists $\mathbf{r} \in \mathbb{Q}^d \cap M_{Q_{\text{aff}}}$ such that*

$$H_{\text{std}}(\mathbf{r}) \leq T \quad \text{and} \quad \text{dist}(\mathbf{r}, \mathbf{x}) \leq \frac{C}{\sqrt{H_{\text{std}}(\mathbf{r})T}} \quad (2.6)$$

(C) *The set*

$$\{\mathbf{x} \in M_{Q_{\text{aff}}} : \text{there exist } C, T_0 > 0 \text{ such that, for all } T \geq T_0, \\ \text{there exists } \mathbf{r} \in \mathbb{Q}^d \cap M_{Q_{\text{aff}}} \text{ satisfying (2.6)}\}$$

has positive $\lambda_{M_{Q_{\text{aff}}}}$ -measure.

As for the optimality of Theorem 5.1, as stated above, it suffices to show that the set

$$\text{BA}_{M_Q} := \text{BA}_{M_Q}(\psi_1)$$

of *intrinsically badly approximable* points of M_Q is nonempty. It follows from the Correspondence Principle below (Lemma 4.2) that points in BA_{M_Q} correspond to bounded orbits of some dynamical system (cf. Corollary 4.3). Then the results of [35] imply the following theorem.

Theorem 4.5 (Jarník–Schmidt for quadric hypersurfaces). *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface. Then $\dim(\text{BA}_{M_Q}) = \dim(M_Q)$. In particular, the Dirichlet function ψ_1 is optimal.*

(No changes needed for the affine corollary.)

Using the methods of [38], one can strengthen the conclusion of this theorem to say that BA_{M_Q} is winning (in the sense of Schmidt). This conclusion also follows from a much more general theorem in [21] which applies to *all* nondegenerate manifolds and asserts that the set of intrinsically badly approximable points is hyperplane absolute winning (see [9] for the definition).

Before stating the analogue of Khintchine's theorem for intrinsic approximation on quadric hypersurfaces, let us introduce the following definitions, which will be used in Sections 6–9.

Definition 2.4. Call a function ψ *regular* if, for every (equivalently, for some) $C_1 > 1$, there exists $C_2 > 1$ such that, for all q_1, q_2 , if $1/C_1 \leq q_2/q_1 \leq C_1$, then

$$1/C_2 \leq \psi(q_2)/\psi(q_1) \leq C_2.$$

This may be stated succinctly as follows: $q_1 \asymp q_2$ implies $\psi(q_1) \asymp \psi(q_2)$.

Definition 2.5. The *exceptional quadric hypersurface* is the hypersurface $M_{Q_0} \subseteq \mathbb{P}_{\mathbb{R}}^3$ defined by the *exceptional quadratic form*

$$Q_0(x_0, x_1, x_2, x_3) = x_0x_3 - x_1x_2. \quad (2.7)$$

If a quadratic form $Q: \mathbb{R}^4 \rightarrow \mathbb{R}$ is conjugate over \mathbb{Q} to Q_0 , we will write $Q \sim Q_0$. We remark that $Q \sim Q_0$ holds if and only if Q is a rational quadratic form in four variables for which $p_{\mathbb{Q}} = p_{\mathbb{R}} = 2$ (see Lemma 3.6 for more detail).

The hypersurface M_{Q_0} , which we study in detail in Section 9, has very interesting properties for intrinsic Diophantine approximation. Note that if $Q \sim Q_0$, then the intrinsic Diophantine theory on M_Q will be more or less the same as the intrinsic Diophantine theory on M_{Q_0} . Specifically, the rational equivalence between Q and Q_0 defines

a diffeomorphism between M_Q and M_{Q_0} which sends rational points to rational points and preserves heights up to a multiplicative constant.

Theorem 6.3 (Khintchine-type theorem for quadric hypersurfaces). *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Fix $\psi: \mathbb{N} \rightarrow (0, \infty)$, and suppose that ψ is regular and that the function $q \mapsto q\psi(q)$ is nonincreasing. Then $A_{M_Q}(\psi)$ has full Lebesgue measure if the series⁶*

$$\begin{cases} \sum_{T \in 2^{\mathbb{N}}} T^{d-1} \psi^{d-1}(T), & Q \sim Q_0, \\ \sum_{T \in 2^{\mathbb{N}}} T^2 \log \log T \psi^2(T), & Q \sim Q_0, \end{cases} \quad (6.4)$$

diverges; otherwise, $A_{M_Q}(\psi)$ is Lebesgue null.

(No changes needed for the affine corollary.)

The appearance of two cases in Theorem 6.3 is due to nontrivial relations among the collection of sets defining $A_{M_{Q_0}}$ that are not present when $Q \sim Q_0$. A discussion of these relations, and their implications, is given in Section 9 (see particularly Remark 9.3).

Using the Mass Transference Principle of Beresnevich and Velani [3, Theorem 2], one can deduce the divergence case of the Jarník–Besicovitch theorem for quadric hypersurfaces (Theorem 6.4). Combined with the convergence case (Corollary 6.2), this gives a complete analogue of the Jarník–Besicovitch theorem when $Q \sim Q_0$, and a slight discrepancy between the convergence and divergence conditions in the exceptional case. This discrepancy, however, does not affect the computation of the Hausdorff dimension of the set of intrinsically ψ_c -approximable points for all $c > 1$, $A_{M_Q}(\psi_c)$; namely, Theorem 6.4 immediately implies

$$\dim(A_{M_Q}(\psi_c)) = \frac{d-1}{c}. \quad (2.8)$$

See Section 6 for a detailed discussion.

Remark 2.7. Let \mathbb{H}^d denote the d -dimensional hyperbolic space. Given a quadric hypersurface $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ satisfying $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$, there exists a lattice $\Gamma \subseteq \text{Isom}(\mathbb{H}^d)$ and a diffeomorphism $\Phi: \partial \mathbb{H}^d \rightarrow M_Q$ such that if $P_{\Gamma} \subseteq \partial \mathbb{H}^d$ is the set of parabolic fixed points of Γ , then $\Phi(P_{\Gamma}) = \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$. This correspondence allows one to deduce the case $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$ of all the results of this subsection as consequences of known theorems about Diophantine approximation of lattices in $\text{Isom}(\mathbb{H}^d)$; see Section 3.4 for more detail.

Remark 2.8. In the above theorems, the form Q is always assumed to be nonsingular with integer coefficients. The latter assumption may be made without loss of generality since, if Q is a quadratic form which is not a scalar multiple of any quadratic form with integer coefficients, then $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ is not dense in M_Q ; cf. Remark 5.11. On the other hand, the nonsingularity assumption does involve a loss of generality. In Theorem 5.1, the singular case can be deduced from the nonsingular case; cf. Remark 5.9. However, this is not the case for Theorem 6.3. The use of the nonsingularity assumption appears unavoidable in Theorem 6.3 since, if Q is singular, then the associated algebraic group $O(Q)$ is not semisimple.

⁶Here and hereafter, $2^{\mathbb{N}}$ stands for $\{2^n : n \in \mathbb{N}\}$.

The structure of the paper. In Section 3, we recall the necessary preliminaries from the theory of quadratic forms. In Section 4, we state and prove the Correspondence Principle, which relates intrinsic Diophantine approximation on a nonsingular rational quadric hypersurface M_Q with dynamics on a certain space of arithmetic lattices. This correspondence is similar to the one developed for ambient approximation by Davenport–Schmidt and Dani; see [14–16, 36, 37] and generalizes the one used in [34]. In particular, we prove (Corollary 4.3) that $[x] \in \text{BA}_{M_Q}$ if and only if a certain trajectory on the corresponding homogeneous space is bounded.

In Section 5, we prove Theorem 5.1 (Dirichlet for quadric hypersurfaces). In Section 6, we use [37, Theorem 1.7] to reduce Theorem 6.3 (Khintchine for quadric hypersurfaces) to a statement about Haar measure on the space of Q -arithmetic lattices (Proposition 8.9). In Section 8, we use the generalized Iwasawa decomposition [39, Proposition 8.44] and the reduction theory for algebraic groups [41, Proposition 2.2] to prove Proposition 8.9, thus completing the proof of Theorem 6.3. Finally, in Section 9, we analyze in detail the exceptional quadric hypersurface M_{Q_0} and explain intuitively why the converse to (the naive application of) Borel–Cantelli does not hold for intrinsic approximation on this hypersurface.

3. Preliminaries on quadratic forms and lattices

3.1. Orthogonality and nonsingularity

Let V be a vector space over \mathbb{R} , and let $Q: V \rightarrow \mathbb{R}$ be a quadratic form. We denote by B_Q the unique symmetric bilinear form on V satisfying

$$Q(\mathbf{x}) = B_Q(\mathbf{x}, \mathbf{x}) \quad \text{for all } \mathbf{x} \in V.$$

We remark that B_Q may be written explicitly in terms of Q via the formula

$$B_Q(\mathbf{x}, \mathbf{y}) = \frac{Q(\mathbf{x} + \mathbf{y}) - Q(\mathbf{x}) - Q(\mathbf{y})}{2}.$$

Definition 3.1. Two elements $\mathbf{x}, \mathbf{y} \in V$ are Q -orthogonal if $B_Q(\mathbf{x}, \mathbf{y}) = 0$. The set of all vectors which are Q -orthogonal to a given vector \mathbf{x} will be denoted \mathbf{x}^\perp , and for any $S \subseteq V$, we let $S^\perp := \bigcap_{\mathbf{x} \in S} \mathbf{x}^\perp$.

Definition 3.2. The quadratic form Q is called *nonsingular* if, for every $\mathbf{x} \in V \setminus \{\mathbf{0}\}$, we have $\mathbf{x}^\perp \not\subseteq V$ or, equivalently, if the map $\mathbf{x} \mapsto B_Q(\mathbf{x}, \cdot)$ is an isomorphism between V and V^* .

Note that a form Q is nonsingular if and only if its corresponding hypersurface M_Q is nonsingular as a manifold. Indeed, recall that $M_Q = [L_Q]$, where L_Q is the light cone of Q defined in (2.3). Then M_Q is nonsingular if and only if $L_Q \setminus \{\mathbf{0}\}$ is nonsingular, which in turn happens if and only if $\nabla Q(\mathbf{x}) \neq 0$ for all $\mathbf{x} \in L_Q \setminus \{\mathbf{0}\}$. Since $\nabla Q(\mathbf{x}) = 2B_Q(\mathbf{x}, \cdot)$, we have $\nabla Q(\mathbf{x}) = 0$ if and only if $\mathbf{x}^\perp = \mathbb{R}^{d+1}$. Thus M_Q is nonsingular if and only if $\mathbf{x}^\perp \not\subseteq \mathbb{R}^{d+1}$ for all $\mathbf{x} \in L_Q$. Since $\mathbf{x}^\perp = \mathbb{R}^{d+1}$ implies $\mathbf{x} \in L_Q$, this proves the assertion.

3.2. Totally isotropic subspaces; rank and renormalization

Throughout this subsection, fix $\mathbb{K} \in \{\mathbb{R}, \mathbb{Q}\}$ and $d \geq 1$, and let $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be a non-singular quadratic form whose coefficients lie in \mathbb{K} . We say that a subspace $E \subseteq \mathbb{R}^{d+1}$ is a \mathbb{K} -subspace if E has a basis consisting of elements of \mathbb{K}^{d+1} or, equivalently, if E is defined by equations whose coefficients lie in \mathbb{K} . (In the literature, it is sometimes said that E is *defined over \mathbb{K}* .)

Definition 3.3. A subspace $E \subseteq \mathbb{R}^{d+1}$ is *totally isotropic* if $Q|_E = 0$. It is known (see e.g. [19, Corollary 8.12]) that any two maximal totally isotropic \mathbb{K} -subspaces of \mathbb{R}^{d+1} have the same dimension. This common dimension is called the \mathbb{K} -rank of Q and is denoted by $p_{\mathbb{K}}$.

It turns out to be convenient to conjugate totally isotropic subspaces to canonical subspaces, namely to subspaces of the form

$$\mathcal{L}_m := \sum_{i=0}^{m-1} \mathbb{R}\mathbf{e}_i. \quad (3.1)$$

By choosing the right conjugation map ϕ , we may also guarantee that the conjugated quadratic form $R = Q \circ \phi$ has a particularly nice form. We make this rigorous as follows.

Definition 3.4. For $m \leq \frac{d+1}{2}$, a quadratic form R is *m -normalized* if there exists a quadratic form \tilde{R} on \mathbb{R}^{d+1-2m} such that

$$R(\mathbf{x}) = x_0 x_d + x_1 x_{d-1} + \cdots + x_{m-1} x_{d-m+1} + \tilde{R}(x_m, \dots, x_{d-m}).$$

The quadratic form \tilde{R} will be called the *remainder* of R .

Proposition 3.5. Let $E \subseteq \mathbb{R}^{d+1}$ be a totally isotropic \mathbb{K} -subspace of dimension m . Then $m \leq \frac{d+1}{2}$, and there exists $\phi \in \mathrm{GL}_{d+1}(\mathbb{K})$ such that

- (i) $\phi^{-1}(E) = \mathcal{L}_m$ and
- (ii) $R := Q \circ \phi$ is m -normalized.

Proof. Since Q is nonsingular, we may identify E^* with \mathbb{R}^{d+1}/E^\perp via the map

$$\mathbf{x} + E^\perp \mapsto B_Q(\mathbf{x}, \cdot)|_E. \quad (3.2)$$

Let $(\mathbf{f}_i)_{i=0}^{m-1}$ be a \mathbb{K} -basis for E , and let $(\mathbf{f}'_{d-i} + E^\perp)_{i=0}^{m-1}$ be its dual basis. Inductively, define $\mathbf{f}_{d-i} \in \mathbf{f}'_{d-i} + E^\perp$ by letting

$$\mathbf{f}_{d-i} = \mathbf{f}'_{d-i} - \sum_{j=0}^{i-1} B_Q(\mathbf{f}'_{d-i}, \mathbf{f}_{d-j}) \mathbf{f}_j - \frac{1}{2} Q(\mathbf{f}'_{d-i}) \mathbf{f}_i.$$

Direct calculation shows that $B_Q(\mathbf{f}_{d-i}, \mathbf{f}_{d-j}) = 0$ for $j \leq i$. Thus $E_2 := \sum_{i=0}^{m-1} \mathbb{R}\mathbf{f}_{d-i}$ is also a totally isotropic \mathbb{K} -subspace of \mathbb{R}^{d+1} . Note that, by construction, E_2 is isomorphic to E^* via the map (3.2). Since E is totally isotropic, $E \subseteq E^\perp$ and thus $E \cap E_2 = \{\mathbf{0}\}$.

Let $E_3 = E^\perp \cap E_2^\perp = (E + E_2)^\perp$. Since $Q|_{E+E_2}$ is nonsingular, we have

$$(E + E_2) \cap E_3 = \{\mathbf{0}\}$$

and thus $\mathbb{R}^{d+1} = E \oplus E_2 \oplus E_3$. It follows that

$$\dim(E_3) = d + 1 - \dim(E) - \dim(E_2) = d + 1 - 2m,$$

and in particular $m \leq (d + 1)/2$. Let $(\mathbf{f}_i)_{i=m}^{d-m}$ be a \mathbb{K} -basis for E_3 , and let ϕ be the $(d + 1) \times (d + 1)$ matrix whose columns are given by $\mathbf{f}_0, \dots, \mathbf{f}_d$ so that $\phi(\mathbf{e}_i) = \mathbf{f}_i$ for $i = 0, \dots, d$. Then $\phi \in \mathrm{GL}_{d+1}(\mathbb{K})$ by the above-mentioned decomposition

$$\mathbb{R}^{d+1} = E \oplus E_2 \oplus E_3.$$

Parts (i) and (ii) follow immediately. \blacksquare

Note that it follows from the above proposition that $p_{\mathbb{R}}$ is always less than or equal to $\frac{d+1}{2}$. Also, if Q has coefficients in \mathbb{Q} , then $p_{\mathbb{Q}} \geq \frac{d-3}{2}$ unless $p_{\mathbb{Q}} = p_{\mathbb{R}}$. Indeed, without loss of generality, suppose that Q is $p_{\mathbb{Q}}$ -normalized, and let \tilde{Q} be the remainder of Q . If $p_{\mathbb{Q}} \neq p_{\mathbb{R}}$, then \tilde{Q} represents zero over \mathbb{R} . Since \tilde{Q} is a quadratic form in $d + 1 - 2p_{\mathbb{Q}}$ variables, if $d + 1 - 2p_{\mathbb{Q}} \geq 5$, by Meyer's theorem, \tilde{Q} represents zero over \mathbb{Q} . This would contradict the definition of $p_{\mathbb{Q}}$. So $d + 1 - 2p_{\mathbb{Q}} \leq 4$; rearranging gives $p_{\mathbb{Q}} \geq \frac{d-3}{2}$.

Another consequence of Proposition 3.5 is a nice characterization of quadratic forms rationally equivalent to the exceptional quadratic form Q_0 defined in (2.7). Recall that the determinant $\det(Q)$ of a quadratic form $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ is the determinant of the linear map $\phi_Q: \mathbb{R}^{d+1} \rightarrow (\mathbb{R}^{d+1})^* \equiv \mathbb{R}^{d+1}$ defined by $\mathbf{x} \mapsto B_Q(\mathbf{x}, \cdot)$.

Lemma 3.6. *The following are equivalent for a rational quadratic form Q in four variables with $p_{\mathbb{Q}} \geq 1$:*

- (i) $Q \sim Q_0$;
- (ii) $p_{\mathbb{Q}} = p_{\mathbb{R}} = 2$;
- (iii) $\det(Q)$ is a square of a rational number.

Proof. Note that, for any $\phi \in \mathrm{GL}_4(\mathbb{R})$, it holds that $\det(Q \circ \phi) = \det(Q) \det(\phi)^2$. In particular, if Q_1 and Q_2 are equivalent over \mathbb{Q} , then $\det(Q_1)$ is a square if and only if $\det(Q_2)$ is. Thus the implication (i) \Rightarrow (iii) follows immediately upon calculating that $\det(Q_0) = 1/16$.

For the implication (iii) \Rightarrow (ii), suppose that $\det(Q)$ is a square. By Proposition 3.5, we may without loss of generality assume that Q is 1-normalized. In this case, we have $\det(Q) = -(1/4) \det(\tilde{Q})$ where \tilde{Q} is the remainder of Q . By the well-known canonical form of quadratic forms, we may without loss of generality assume that

$$\tilde{Q}(\mathbf{x}) = a_1 x_1^2 + a_2 x_2^2 \quad \text{for some } a_1, a_2 \in \mathbb{Q}.$$

Then $-\det(\tilde{Q}) = -a_1 a_2$ is a square. Thus $\mathbf{b} := (0, a_2, \sqrt{-a_1 a_2}, 0) \in \mathbb{Q}^4$, and $\mathbb{R}\mathbf{e}_0 + \mathbb{R}\mathbf{b}$ is a totally isotropic subspace of dimension 2, proving that $p_{\mathbb{Q}} = 2$.

Finally, (ii) \Rightarrow (i) is a straightforward consequence of Proposition 3.5. \blacksquare

A convenient fact about m -normalized quadratic forms is that any element of $\mathrm{GL}_m(\mathbb{R})$ extends to an element of $\mathrm{SL}_{d+1}(\mathbb{R})$ which preserves every m -normalized quadratic form. Specifically, given a quadratic form $R: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$, let

$$\mathrm{O}(R) = \{g \in \mathrm{SL}_{d+1}^+(\mathbb{R}) : R \circ g = R\}.$$

Then a direct computation yields the following.

Observation 3.7. Fix $m \leq \frac{d+1}{2}$ and $\phi \in \mathrm{GL}_m(\mathbb{R})$. Define the *reverse* of the matrix ϕ to be the matrix whose (i, j) th entry is equal to the $(m-j, m-i)$ th entry of ϕ^{-1} , and denote this matrix by $\hat{\phi}$. Visually, $\hat{\phi}$ is ϕ^{-1} flipped along the northeast-southwest diagonal. Let

$$g_\phi = \begin{bmatrix} \phi & & \\ & I_{d+1-2m} & \\ & & \hat{\phi} \end{bmatrix}. \quad (3.3)$$

Then $g_\phi \in \mathrm{O}(R)$ for every m -normalized quadratic form R .

Next, for each $m \leq \frac{d+1}{2}$ and $\mathbf{t} \in \mathbb{R}^m$, let

$$g_{\mathbf{t}} = g_{\mathrm{diag}(e^{-t_0}, \dots, e^{-t_{m-1}})} = \begin{bmatrix} e^{-t_0} & & & & \\ & \ddots & & & \\ & & e^{-t_{m-1}} & & \\ & & & I_{d+1-2m} & e^{t_{m-1}} \\ & & & & \ddots \\ & & & & & e^{t_0} \end{bmatrix}. \quad (3.4)$$

Of particular importance will be the case $m = 1$, in which case

$$g_t = \begin{bmatrix} e^{-t} & & \\ & I_{d-1} & \\ & & e^t \end{bmatrix}. \quad (3.5)$$

A simple computation immediately yields the following observation, which will turn out to be quite useful.

Observation 3.8. For $t \geq 0$ and $\mathbf{x} \in \mathbb{R}^{d+1}$,

$$\mathrm{dist}(\mathbf{x}, \mathcal{L}_1) \leq \|g_t(\mathbf{x})\|, \quad (3.6)$$

where \mathcal{L}_1 is as in (3.1).

3.3. The space of lattices; Mahler's compactness criterion

As stated in the introduction, our main tool for proving theorems concerning intrinsic approximation on M_Q is a correspondence principle between approximations of a point in M_Q and dynamics in the space of lattices. We will describe this correspondence principle in Section 4 below, while here we introduce the space of lattices which we are interested in, namely the space of Q -*arithmetic* lattices.

Definition 3.9. Fix a quadratic form $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$. A lattice $\Lambda \subseteq \mathbb{R}^{d+1}$ is Q -arithmetic if $Q(\Lambda) \subseteq \mathbb{Z}$. (Symmetrically, we may also say that Q is Λ -arithmetic.) The set of Q -arithmetic lattices will be denoted by Ω_Q , while the set of all lattices in \mathbb{R}^{d+1} will be denoted by Ω_d .

Observation 3.10. A quadratic form is \mathbb{Z}^{d+1} -arithmetic if and only if its coefficients are integral.

Clearly, Ω_Q is preserved by the action of $O(Q)$. If $\Lambda \in \Omega_Q$ is fixed, we denote its stabilizer by $O(Q; \Lambda)$ and its orbit by $\Omega_{Q, \Lambda}$. We will implicitly identify $\Omega_{Q, \Lambda}$ with the homogeneous space $O(Q)/O(Q; \Lambda)$ via the map $gO(Q; \Lambda) \mapsto g\Lambda$. This automatically endows $\Omega_{Q, \Lambda}$ with a topological structure and, since $O(Q)$ is unimodular and $O(Q; \Lambda)$ is discrete, a Haar measure, which we will denote by $\mu_{Q, \Lambda}$.

Viewing $\Omega_{Q, \Lambda}$ as a homogeneous space could conceivably give it a different topology than viewing it as a subset of Ω_d , which has its own topology from its identification with $GL_{d+1}(\mathbb{R})/GL_{d+1}(\mathbb{Z})$ coming from the map $g GL_{d+1}(\mathbb{Z}) \mapsto g(\mathbb{Z}^{d+1})$. Fortunately, it turns out that these topologies are identical.

Proposition 3.11. *The inclusion map $\Omega_{Q, \Lambda} \rightarrow \Omega_d$ is proper and continuous when both spaces are endowed with the topologies coming from the identification with their corresponding homogeneous spaces. Consequently, the topology on $\Omega_{Q, \Lambda}$ is unambiguous.*

Proof. The continuity of the inclusion map follows directly from the continuity of the inclusion map from $O(Q)$ to $GL_{d+1}(\mathbb{R})$. Let us show that the inclusion map is proper. Let $(\Lambda_n)_1^\infty$ be a sequence in $\Omega_{Q, \Lambda}$ converging to a point $\Lambda_0 \in \Omega_d$. Then there exist

$$GL_{d+1}(\mathbb{R}) \ni g_n \rightarrow g_0 \in GL_{d+1}(\mathbb{R})$$

such that $\Lambda_n = g_n(\mathbb{Z}^{d+1})$ for all $n \geq 0$. This implies that, for all $n \geq 1$, $Q_n := Q \circ g_n$ is a \mathbb{Z}^{d+1} -arithmetic quadratic form, and $Q_n \rightarrow Q_0 := Q \circ g_0$. Since the space of \mathbb{Z}^{d+1} -arithmetic quadratic forms is discrete (being identical to the space of quadratic forms with coefficients in \mathbb{Z}), we have $Q_n = Q_0$ for all sufficiently large n . (Thus *a posteriori* Q_0 is \mathbb{Z}^{d+1} -arithmetic, or equivalently $\Lambda_0 \in \Omega_Q$.) For n satisfying $Q_n = Q_0$, we have $h_n := g_n g_0^{-1} \in O(Q)$; in particular, $\Lambda_0 = h_n^{-1}(\Lambda_n) \in \Omega_{Q, \Lambda}$. On the other hand, $\Lambda_n = h_n \Lambda_0$ and $h_n \rightarrow h_0 = id$; this implies that $\Lambda_n \rightarrow \Lambda_0$ in the topology on $\Omega_{Q, \Lambda}$ coming from its identification with the homogeneous space $O(Q)/O(Q; \Lambda)$. ■

We now recall Mahler's famous compactness criterion and deduce an analogue in the context of quadratic forms. For $\Lambda \in \Omega_d$, let

$$\delta(\Lambda) := \min_{\mathbf{p} \in \Lambda \setminus \{0\}} \|\mathbf{p}\|. \quad (3.7)$$

Theorem 3.12 (Mahler's compactness criterion [42, Theorem 2]). *A set $S \subseteq \Omega_d$ is pre-compact if and only if δ is bounded from below on S , and the covolumes of all lattices in S are uniformly bounded from above.*

For $\Lambda \in \Omega_Q$, let

$$\delta_Q(\Lambda) = \min_{\mathbf{p} \in \Lambda \cap L_Q \setminus \{0\}} \|\mathbf{p}\|.$$

We let $\delta_Q(\Lambda) = \infty$ if $\Lambda \cap L_Q \setminus \{\mathbf{0}\} = \emptyset$.

Observation 3.13. If we let

$$\|Q\| = \max_{\|\mathbf{x}\|=\|\mathbf{y}\|=1} |B_Q(\mathbf{x}, \mathbf{y})|,$$

then $\min(\delta_Q, 1/\sqrt{\|Q\|}) \leq \delta \leq \delta_Q$.

Proof. For $\mathbf{p} \in \Lambda \setminus L_Q$, $\|\mathbf{p}\| \geq \sqrt{|Q(\mathbf{p})|/\|Q\|} \geq 1/\sqrt{\|Q\|}$. ■

Corollary 3.14 (Analogue of Mahler's compactness criterion). *Fix $\Lambda \in \Omega_Q$. Then a set $S \subseteq \Omega_{Q,\Lambda}$ is precompact if and only if δ_Q is bounded from below on S .*

Proof. By Observation 3.13, δ_Q is bounded from below on S if and only if δ is bounded from below on S . But by Theorem 3.12, since the covolumes of all lattices in $\Omega_{Q,\Lambda}$ are the same, δ is bounded from below if and only if S is precompact in the topology of Ω_d . By Proposition 3.11, this occurs if and only if S is precompact in the topology of $\Omega_{Q,\Lambda}$. (Here we use not only the fact that the topology on $\Omega_{Q,\Lambda}$ is the one induced from Ω_d , but also the fact that the inclusion map is proper, and consequently $\Omega_{Q,\Lambda}$ is closed in Ω_d .) ■

3.4. Relation to Kleinian lattices

In this subsection, we describe the relation between the intrinsic Diophantine approximation of a quadric hypersurface M_Q satisfying $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$ and the approximation of points in the boundary of d -dimensional hyperbolic space \mathbb{H}^d by parabolic fixed points in a lattice $\Gamma \subseteq \text{Isom}(\mathbb{H}^d)$ which depends on the quadric hypersurface M_Q . Since the latter situation is well-studied, this correspondence can be used to immediately prove the theorems of Section 2 in the case $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$. (However, our proofs of the theorems of Section 2 in the general case are not dependent on assuming $p_{\mathbb{R}} > 1$, so this subsection can be skipped without any loss of generality.)

Let $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be a quadratic form with integer coefficients satisfying

$$p_{\mathbb{Q}} = p_{\mathbb{R}} = 1.$$

Then the signature of Q is either $(d, 1)$ or $(1, d)$. Without loss of generality, we will suppose that its signature is $(d, 1)$. The *hyperboloid model of hyperbolic geometry* is the set

$$\mathbb{H}^d := \{\mathbf{x} \in \mathbb{R}^{d+1} : Q(\mathbf{x}) = -1\}$$

with the Riemannian metric $Q|_{\mathbb{H}^d}$ (its positive definiteness is guaranteed by the fact that the signature of Q is $(d, 1)$). The *hyperbolic distance* is given by the formula

$$\cosh \text{dist}(\mathbf{x}, \mathbf{y}) = |B_Q(\mathbf{x}, \mathbf{y})|.$$

Note that, by Sylvester's law of inertia, up to isometry, the space $(\mathbb{H}^d, \text{dist})$ does not depend on Q , but only on d . For the equivalence of the hyperboloid model with other standard models of hyperbolic geometry, see e.g. [12]. The *boundary* of \mathbb{H}^d , denoted

$\partial\mathbb{H}^d$, is defined to be the boundary of $[\mathbb{H}^d]$ in $\mathbb{P}_{\mathbb{R}}^d$. Observe that $\partial\mathbb{H}^d = M_Q$. A *horoball* in \mathbb{H}^d is a set of the form

$$\{\mathbf{x} \in \mathbb{H}^d : \mathcal{B}_{[\mathbf{r}]}(\mathbf{z}, \mathbf{x}) > t\},$$

where $\mathbf{z} \in \mathbb{H}^d$, $[\mathbf{r}] \in \partial\mathbb{H}^d$, $t \in \mathbb{R}$, and $\mathcal{B}_{[\mathbf{r}]}$ denotes the *Busemann function*

$$\mathcal{B}_{[\mathbf{r}]}(\mathbf{z}, \mathbf{x}) = \lim_{[\mathbf{y}] \rightarrow [\mathbf{r}]} [\text{dist}(\mathbf{y}, \mathbf{z}) - \text{dist}(\mathbf{y}, \mathbf{x})].$$

Such a horoball is said to be *centered* at the point $[\mathbf{r}]$. The isometry group of \mathbb{H}^d is given by

$$\text{Isom}(\mathbb{H}^d) = \text{O}(Q).$$

Since Q has integer coefficients, the subgroup

$$\Gamma := \text{O}(Q; \mathbb{Z}) := \text{O}(Q) \cap \text{GL}_{d+1}(\mathbb{Z})$$

is a lattice in $\text{O}(Q)$ (see [6, Theorem 7.8]). Let $P_\Gamma \subseteq \partial\mathbb{H}^d$ denote the set of parabolic fixed points of Γ .

We now state the relation between intrinsic approximation of M_Q and approximation of $\partial\mathbb{H}^d$ by P_Γ .

Proposition 3.15. *The following statements hold.*

- (i) *There exists a Γ -invariant disjoint family of horoballs $(H_{[\mathbf{r}]})_{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}$ such that, for each $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$, $H_{[\mathbf{r}]}$ is centered at $[\mathbf{r}]$ and*

$$H_{\text{std}}([\mathbf{r}]) \asymp_{\times} e^{\text{dist}(\mathbf{z}, H_{[\mathbf{r}]})}, \quad (3.8)$$

where $\mathbf{z} \in \mathbb{H}^d$ is fixed.

- (ii) $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q = P_\Gamma$.

Using Proposition 3.15, one may translate [52, Theorems 1 and 4], [51, Theorem C], and [44, Theorem 2] (see also [24] and the references therein for subsequent generalizations) into the context of quadratic forms, yielding the results of Section 2 in the case $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$. Details are left to the reader.

Proof of (i). Fix $\varepsilon > 0$, and for each $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$, let

$$H_{[\mathbf{r}]} = \{\mathbf{x} \in \mathbb{H}^d : |B_Q(\mathbf{x}, \mathbf{r})| < \varepsilon\},$$

where \mathbf{r} is the unique primitive integral representative of $[\mathbf{r}]$. The fact that $H_{[\mathbf{r}]}$ is a horoball centered at $[\mathbf{r}]$ follows from the following well-known formula for the Busemann function in the hyperboloid model:

$$\mathcal{B}_{[\mathbf{r}]}(\mathbf{x}, \mathbf{y}) = \log \frac{|B_Q(\mathbf{x}, \mathbf{r})|}{|B_Q(\mathbf{y}, \mathbf{r})|}.$$

Since $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ and Q are both invariant under Γ , the collection $(H_{[\mathbf{r}]})_{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}$ is clearly Γ -invariant. Next, we will show that the collection $(H_{[\mathbf{r}]})_{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}$ is disjoint for ε sufficiently small. Indeed, suppose $\mathbf{x} \in H_{[\mathbf{r}_1]} \cap H_{[\mathbf{r}_2]}$, and apply $g \in \text{O}(Q)$ such

that $g(\mathbf{x}) = \mathbf{w}$, where $\mathbf{w} \in \mathbb{H}^d$ is fixed. Then $|B_Q(\mathbf{w}, g(\mathbf{r}_i))| < \varepsilon$, where \mathbf{r}_i is the primitive integral representative of $[\mathbf{r}_i]$. On the other hand, since Q has signature $(d, 1)$ and $Q(\mathbf{w}) = -1$, we have

$$|B_Q(\mathbf{w}, \mathbf{r})| \asymp_{\times} \|\mathbf{r}\| \quad \text{for all } \mathbf{r} \in L_Q. \quad (3.9)$$

Thus $\|g(\mathbf{r}_i)\| \lesssim_{\times} \varepsilon$, and so $|B_Q(\mathbf{r}_1, \mathbf{r}_2)| \lesssim_{\times} \|Q\| \varepsilon^2$. Thus $|B_Q(\mathbf{r}_1, \mathbf{r}_2)| < \frac{1}{2}$ for ε sufficiently small. On the other hand, $B_Q(\mathbf{r}_1, \mathbf{r}_2) \in \mathbb{Z}/2$ since Q has integer coefficients, so $B_Q(\mathbf{r}_1, \mathbf{r}_2) = 0$. Since $p_{\mathbb{Q}} = 1$, this implies $[\mathbf{r}_1] = [\mathbf{r}_2]$.

As the horoballs $(H_{[\mathbf{r}]})_{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}$ are disjoint open subsets of the connected set \mathbb{H}^d , there exists $\mathbf{z} \in \mathbb{H}^d \setminus \bigcup_{[\mathbf{r}]} H_{[\mathbf{r}]}$. Now fix $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$, and we will demonstrate (3.8). Letting $\mathbf{x} \in \partial H_{[\mathbf{r}]}$ be arbitrary, we calculate

$$e^{\text{dist}(\mathbf{z}, H_{[\mathbf{r}]})} = e^{\mathcal{B}_{[\mathbf{r}]}(\mathbf{z}, \mathbf{x})} = \frac{|B_Q(\mathbf{z}, \mathbf{r})|}{\varepsilon}.$$

Combining with (3.9) yields (3.8). \blacksquare

Proof of (ii). Suppose that $[\mathbf{r}]$ is a parabolic fixed point of Γ , say $g([\mathbf{r}]) = [\mathbf{r}]$ for some parabolic $g \in \Gamma$. Then the line representing $[\mathbf{r}]$ is precisely the set

$$\{\mathbf{x} \in \mathbb{R}^{d+1} : g(\mathbf{x}) = \mathbf{x}\},$$

which is a rational subspace of \mathbb{R}^{d+1} . Consequently, $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$.

Conversely, suppose that $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$. As above, we fix $\mathbf{z} \in \mathbb{H}^d \setminus \bigcup_{[\mathbf{r}]} H_{[\mathbf{r}]}$. Since the collection $(H_{[\mathbf{r}]})_{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}$ is Γ -invariant, this implies $g(\mathbf{z}) \notin H_{[\mathbf{r}]}$ for all $g \in \Gamma$. In particular, $[\mathbf{r}]$ cannot be a conical limit point of Γ (see e.g. [8, Section 3.2] for the definition). But since Γ is a lattice, every point of $\partial \mathbb{H}^d$ is either a conical limit point or a parabolic fixed point (e.g. [8, Section 4]). Thus $[\mathbf{r}] \in P_{\Gamma}$. \blacksquare

4. The correspondence principle

In this section, we introduce the correspondence principle alluded to in the introduction. It is an intrinsic approximation analogue of the so-called Dani correspondence for ambient approximation [14–16, 36, 37]. A special case can be found in [34, Theorem 1.5].

Fix $d \geq 2$, and let $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be a nonsingular quadratic form with integer coefficients. Suppose that $p_{\mathbb{Q}} \geq 1$. By Proposition 3.5, there exists a matrix $\phi \in \text{GL}_{d+1}(\mathbb{Q})$ such that $R := Q \circ \phi$ is $p_{\mathbb{Q}}$ -normalized. Let $\Lambda_* = \phi^{-1}(\mathbb{Z}^{d+1})$. Note that Λ_* is commensurable with \mathbb{Z}^{d+1} and that $\Lambda_* \subset \Omega_R$. Moreover, the \mathbb{Q} -ranks of Q and R are identical, and the same goes for the \mathbb{R} -ranks, so denoting these ranks by $p_{\mathbb{Q}}$ and $p_{\mathbb{R}}$ will not cause ambiguity.

Consider the maps $\pi_1: \text{O}(R) \rightarrow M_Q$ and $\pi_2: \text{O}(R) \rightarrow \Omega_{R, \Lambda_*}$ defined by

$$\begin{aligned} \pi_1(g) &= \phi \circ g([\mathbf{e}_0]), \\ \pi_2(g) &= g^{-1} \Lambda_* = (\phi \circ g)^{-1}(\mathbb{Z}^{d+1}). \end{aligned}$$

Now fix $g \in \mathrm{O}(R)$, and let

$$[\mathbf{x}] = \pi_1(g) \quad \text{and} \quad \Lambda = \pi_2(g). \quad (4.1)$$

The first version of the correspondence principle gives a relation between the following entities:

- (A) Rational points in $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ which are close to $[\mathbf{x}]$.
- (B) Points in $\Lambda_{\mathrm{pr}} \cap L_R$ which are close to \mathcal{L}_1 . Here Λ_{pr} denotes the set of primitive vectors of Λ , and $\mathcal{L}_1 = \mathbb{R}\mathbf{e}_0$ is as in (3.1).
- (C) Pairs (t, \mathbf{q}) , where $\mathbf{q} \in g_t \Lambda_{\mathrm{pr}} \cap L_R$ is close to $\{\mathbf{0}\}$.

Lemma 4.1 (Correspondence principle, form 1). *Let g , $[\mathbf{x}]$, and Λ be as in (4.1). Then*

- (i) *the map $\mathbf{p} \mapsto \phi \circ g([\mathbf{p}])$ is a bijection between $\Lambda_{\mathrm{pr}} \cap L_R$ and $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$.*
- (ii) *Fix $\mathbf{p} \in \Lambda_{\mathrm{pr}} \cap L_R$, and let $[\mathbf{r}] = \phi \circ g([\mathbf{p}])$. Then*

$$\mathrm{dist}([\mathbf{r}], [\mathbf{x}]) \asymp_{\times, g} \frac{\mathrm{dist}(\mathbf{p}, \mathcal{L}_1)}{\|\mathbf{p}\|} \quad \text{and} \quad H_{\mathrm{std}}([\mathbf{r}]) \asymp_{\times, g} \|\mathbf{p}\|. \quad (4.2)$$

In particular, if $\psi: (0, \infty) \rightarrow (0, \infty)$ is a regular function (cf. Definition 2.4), then

$$\frac{\mathrm{dist}([\mathbf{r}], [\mathbf{x}])}{\psi \circ H_{\mathrm{std}}([\mathbf{r}])} \asymp_{\times, g, \psi} \frac{\mathrm{dist}(\mathbf{p}, \mathcal{L}_1)}{\|\mathbf{p}\| \psi(\|\mathbf{p}\|)}. \quad (4.3)$$

In each case, the implied constant can be made independent of g if g is constrained to lie in a bounded subset of $\mathrm{O}(R)$.

- (iii) *Fix $\mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$ such that $|p_0| = \|\mathbf{p}\|$, i.e. $|p_0| \geq \mathrm{dist}(\mathbf{p}, \mathcal{L}_1)$. For $t \geq 0$,*

$$\begin{aligned} \max\left(\mathrm{dist}(\mathbf{p}, \mathcal{L}_1), \frac{\|\mathbf{p}\|}{e^t}\right) &\leq \|g_t(\mathbf{p})\| \\ &\lesssim_{\times} \max\left(\mathrm{dist}(\mathbf{p}, \mathcal{L}_1), \frac{\|\mathbf{p}\|}{e^t}, \frac{e^t \mathrm{dist}(\mathbf{p}, \mathcal{L}_1)^2}{\|\mathbf{p}\|}\right). \end{aligned} \quad (4.4)$$

In particular, letting $t(\mathbf{p}) = \log(\|\mathbf{p}\|/\mathrm{dist}(\mathbf{p}, \mathcal{L}_1))$, we have

$$\|g_{t(\mathbf{p})}(\mathbf{p})\| \asymp_{\times} \mathrm{dist}(\mathbf{p}, \mathcal{L}_1). \quad (4.5)$$

Proof. Part (i) is straightforward. Regarding part (ii), formula (4.2) is perhaps elucidated by the calculation

$$\begin{aligned} \mathrm{dist}([\mathbf{r}], [\mathbf{x}]) &= \mathrm{dist}(\phi \circ g([\mathbf{p}]), \phi \circ g([\mathbf{e}_0])) \asymp_{\times, g} \mathrm{dist}([\mathbf{p}], [\mathbf{e}_0]) \asymp_{\times} \frac{\mathrm{dist}(\mathbf{p}, \mathcal{L}_1)}{\|\mathbf{p}\|}, \\ H_{\mathrm{std}}([\mathbf{r}]) &= \|\phi \circ g(\mathbf{p})\| \asymp_{\times, g} \|\mathbf{p}\|. \end{aligned}$$

Formula (4.3) follows from (4.2) together with the regularity of ψ ; as $H_{\mathrm{std}}([\mathbf{r}]) \asymp_{\times, g} \|\mathbf{p}\|$, we have $\psi \circ H_{\mathrm{std}}([\mathbf{r}]) \asymp_{\times, g} \psi(\|\mathbf{p}\|)$, and (4.3) follows upon combining with the first part of (4.2).

We proceed to the proof of (iii). The first inequality of (4.4) is an immediate consequence of the definition of g_t . To show the second inequality of (4.4), let $\mathbf{q} = g_t(\mathbf{p})$,

and write $\mathbf{q} = (q_0, \dots, q_d)$. Then $|q_1|, \dots, |q_{d-1}| \leq \text{dist}(\mathbf{p}, \mathcal{L}_1)$, while $|q_0| = \|\mathbf{p}\|/e^t$. To bound $|q_d|$, we use the fact that $\mathbf{q} \in L_R$, which means that

$$R(\mathbf{q}) = q_0 q_d + \tilde{R}(q_1, \dots, q_{d-1}),$$

where \tilde{R} is the remainder of R . Rearranging, we have

$$\begin{aligned} |q_d| &= \frac{|\tilde{R}(q_1, \dots, q_{d-1})|}{|q_0|} \leq \frac{\|\tilde{R}\| \cdot \|(q_1, \dots, q_{d-1})\|^2}{|q_0|} \\ &\lesssim \frac{\text{dist}(\mathbf{p}, \mathcal{L}_1)^2}{|q_0|} = \frac{e^t \text{dist}(\mathbf{p}, \mathcal{L}_1)^2}{\|\mathbf{p}\|}. \end{aligned} \quad \blacksquare$$

The second version of the correspondence principle depends on $\psi: (0, \infty) \rightarrow (0, \infty)$, and may be stated as follows.

Lemma 4.2 (Correspondence principle, form 2). *Let g , $[\mathbf{x}]$, and Λ be as in (4.1), and assume that $[\mathbf{x}]$ is irrational (equivalently, that $\Lambda \cap \mathcal{L}_1 = \{\mathbf{0}\}$). Let $\psi: (0, \infty) \rightarrow (0, \infty)$ be a regular function such that the map $q \mapsto q\psi(q)$ is nonincreasing and tends to zero. Then*

$$\begin{aligned} \liminf_{\substack{[\mathbf{r}] \rightarrow [\mathbf{x}] \\ [\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q}} \frac{\text{dist}([\mathbf{r}], [\mathbf{x}])}{\psi \circ H_{\text{std}}([\mathbf{r}])} &\asymp_{\times, g, \psi} \liminf_{\substack{[\mathbf{p}] \rightarrow [\mathbf{e}_0] \\ \mathbf{p} \in \Lambda_{\text{pr}} \cap L_R}} \frac{\text{dist}(\mathbf{p}, \mathcal{L}_1)}{\|\mathbf{p}\| \psi(\|\mathbf{p}\|)} \\ &\asymp_{\times, \psi} \liminf_{t \rightarrow \infty} \frac{e^{-t}}{\psi(e^t \delta_R(g_t \Lambda))}. \end{aligned} \quad (4.6)$$

Proof. The first asymptotics follows directly from (i) and (ii) of Lemma 4.1. Using the function $\Psi(q) := q\psi(q)$, the second asymptotics can be rewritten in a more convenient form:

$$\liminf_{\substack{[\mathbf{p}] \rightarrow [\mathbf{e}_0] \\ \mathbf{p} \in \Lambda_{\text{pr}} \cap L_R}} \frac{\text{dist}(\mathbf{p}, \mathcal{L}_1)}{\Psi(\|\mathbf{p}\|)} \asymp_{\times} \liminf_{t \rightarrow \infty} \frac{\delta_R(g_t \Lambda)}{\Psi(e^t \delta_R(g_t \Lambda))}. \quad (4.7)$$

To demonstrate the \lesssim direction of (4.7), for each $t \geq 0$, choose $\mathbf{p}_t \in \Lambda_{\text{pr}} \cap L_R$ such that $\delta_R(g_t \Lambda) = \|\mathbf{p}_t\|$. Then by (4.4), we have

$$\text{dist}(\mathbf{p}_t, \mathcal{L}_1) \leq \delta_R(g_t \Lambda) \quad \text{and} \quad \|\mathbf{p}_t\| \leq e^t \delta_R(g_t \Lambda)$$

and thus

$$\frac{\text{dist}(\mathbf{p}_t, \mathcal{L}_1)}{\Psi(\|\mathbf{p}_t\|)} \leq \frac{\delta_R(g_t \Lambda)}{\Psi(e^t \delta_R(g_t \Lambda))}. \quad (4.8)$$

Here we have used the fact that the function Ψ is nonincreasing. Next, suppose we have a sequence $t_k \rightarrow \infty$ such that

$$\lim_{k \rightarrow \infty} \frac{e^{-t_k}}{\psi(e^{t_k} \delta_R(g_{t_k} \Lambda))} < \infty.$$

Since $\Psi(q) \rightarrow 0$ as $q \rightarrow \infty$, it follows that $\delta_R(g_{t_k} \Lambda) \rightarrow 0$. In particular,

$$\text{dist}(\mathbf{p}_{t_k}, \mathcal{L}_1) \rightarrow 0.$$

Since $\Lambda \cap \mathcal{L}_1 = \{\mathbf{0}\}$, this implies that the set $\{\mathbf{p}_{t_k} : k \in \mathbb{N}\}$ is infinite. Combining with (4.8) yields the \lesssim direction of (4.7).

To demonstrate the \gtrsim direction of (4.7), suppose that $\mathbf{p}_k \in \Lambda_{\text{pr}} \cap L_R$ is a sequence such that $[\mathbf{p}_k] \rightarrow [\mathbf{e}_0]$. For each k , let $t_k = t(\mathbf{p}_k)$ be defined as in (iii) of Lemma 4.1. Since $[\mathbf{p}_k] \rightarrow [\mathbf{e}_0]$, we have $t_k \rightarrow \infty$. On the other hand, by (4.5), we have

$$\begin{aligned}\delta_R(g_{t_k} \Lambda) &\leq \|g_{t_k}(\mathbf{p}_k)\| \asymp_{\times} \text{dist}(\mathbf{p}_k, \mathcal{L}_1), \\ e^{t_k} \delta_R(g_{t_k} \Lambda) &\lesssim_{\times} e^{t_k} \text{dist}(\mathbf{p}_k, \mathcal{L}_1) = \|\mathbf{p}_k\|,\end{aligned}$$

and so

$$\frac{\delta_R(g_{t_k} \Lambda)}{\Psi(e^{t_k} \delta_R(g_{t_k} \Lambda))} \lesssim_{\times} \frac{\text{dist}(\mathbf{p}_k, \mathcal{L}_1)}{\Psi(\|\mathbf{p}_k\|)}.$$

Letting $k \rightarrow \infty$ finishes the proof. \blacksquare

The next corollary is a direct analogue of Dani's correspondence between bounded orbits and badly approximable vectors/matrices [14, Theorem 2.20].

Corollary 4.3. *Let g , $[\mathbf{x}]$, and Λ be as in (4.1). Then the following are equivalent.*

- (A) $[\mathbf{x}]$ is intrinsically badly approximable, i.e. $[\mathbf{x}] \in \text{BA}_{M_Q}$.
- (B) $\inf_{\mathbf{p} \in \Lambda \cap L_Q \setminus \{\mathbf{0}\}} \text{dist}(\mathbf{p}, \mathcal{L}_1) > 0$.
- (C) The orbit $(g_t \Lambda)_{t \geq 0}$ is bounded in Ω_R .

Proof. Clearly, all the above statements are false if $[\mathbf{x}]$ is irrational. Otherwise, let \mathcal{C} be the class of all regular functions ψ such that the map $q \mapsto q\psi(q)$ is nonincreasing and tends to zero. Then (A) is equivalent to the assertion that the left-hand side of (4.6) is positive for all $\psi \in \mathcal{C}$, (B) is equivalent to the assertion that the middle of (4.6) is positive for all $\psi \in \mathcal{C}$, and (C) is equivalent (by Corollary 3.14) to the assertion that the right-hand side of (4.6) is positive for all $\psi \in \mathcal{C}$. \blacksquare

Remark 4.4. It is somewhat annoying that Lemma 4.2 requires the assumption that $q\psi(q) \rightarrow 0$ as $q \rightarrow \infty$, so that the Dirichlet function $\psi = \psi_1$ is ruled out. (If we were allowed to use $\psi = \psi_1$, then the proof of Corollary 4.3 could be made even simpler – just consider $\psi = \psi_1$ rather than all functions $\psi \in \mathcal{C}$.) However, this assumption is necessary, as can be seen as follows. Arguing as in [34, Proof of Corollary 3.5], one can show that there exists $C > 0$ such that $\delta_R(\Lambda) \leq C$ for all $\Lambda \in \Omega_{R, \Lambda_*}$. (Indeed, otherwise, one can take a sequence $\Lambda_n \in \Omega_{R, \Lambda_*}$ with $\delta_R(\Lambda_n) \rightarrow \infty$; such a sequence cannot have a convergent subsequence, yet it is precompact in view of Corollary 3.14.) This C is a uniform upper bound on the right-hand side of (4.6) when $\psi = \psi_1$. However, we know that, when $p_Q \neq p_{\mathbb{R}}$, then there is no uniform upper bound on the left-hand side of (4.6); this follows from Theorem 5.1 (ii) below. Thus the left- and right-hand sides cannot be asymptotic.⁷

⁷A closer analysis shows that, when $\psi = \psi_1$, the left- and right-hand sides of (4.6) are not necessarily asymptotic even when both of them are close to 0.

Using Corollary 4.3, we can now prove Theorem 4.5.

Theorem 4.5. *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface. Then we have $\dim(\text{BA}_{M_Q}) = \dim(M_Q)$. In particular, the Dirichlet function ψ_1 is optimal.*

Proof. First observe that $\text{BA}_{M_Q} = M_Q$ if $p_Q = 0$; thus it suffices to consider the case $p_Q \geq 1$. Let $\text{BA}_{\Omega_R} \subseteq \Omega_R$ denote the set of lattices in Ω_R whose orbit under the g_t flow is bounded. By [35, Theorem 5.2], we have $\dim(\text{BA}_{\Omega_R}) = \dim(\Omega_R)$. On the other hand, by Corollary 4.3, we have $\mathcal{B} := \pi_2^{-1}(\text{BA}_{\Omega_R}) = \pi_1^{-1}(\text{BA}_{M_Q})$.

Since π_2 is a fibration whose fibers are isomorphic to $\text{Stab}(\Lambda_*)$, the set

$$\mathcal{B} = \pi_2^{-1}(\text{BA}_{\Omega_R}) \subseteq \text{O}(R)$$

has the same local structure as the product $\text{BA}_{\Omega_R} \times \text{Stab}(\Lambda_*) \subseteq \Omega_R \times \text{Stab}(\Lambda_*)$. Now, since $\text{Stab}(\Lambda_*)$ is a manifold, its Hausdorff dimension and upper box dimension are equal. (We refer to [20, p. 38] for the definition of upper box dimension.) So, by [20, Corollary 7.4], we have $\dim(A \times \text{Stab}(\Lambda_*)) = \dim(A) + \dim(\text{Stab}(\Lambda_*))$ for all $A \subseteq \Omega_R$. Taking the cases $A = \text{BA}_{\Omega_R}$ and $A = \Omega_R$ and using the fact that Hausdorff dimension is a local property, we have

$$\begin{aligned} \dim(\mathcal{B}) &= \dim(\text{BA}_{\Omega_R}) + \dim(\text{Stab}(\Lambda_*)), \\ \dim(\text{O}(R)) &= \dim(\Omega_R) + \dim(\text{Stab}(\Lambda_*)). \end{aligned}$$

A similar argument gives

$$\begin{aligned} \dim(\mathcal{B}) &= \dim(\text{BA}_{M_Q}) + \dim(\text{Stab}([\mathbf{e}_0])), \\ \dim(\text{O}(R)) &= \dim(M_Q) + \dim(\text{Stab}([\mathbf{e}_0])). \end{aligned}$$

Thus, since $\dim(\text{BA}_{\Omega_R}) = \dim(\Omega_R)$, we have $\dim(\text{BA}_{M_Q}) = \dim(M_Q)$. \blacksquare

Under the assumption that $q\psi(q) \rightarrow 0$ as $q \rightarrow \infty$, Lemma 4.2 can be used to dynamically describe the sets $\text{A}_{M_Q}(\psi)$ and $\text{WA}_{M_Q}(\psi)$.

Corollary 4.6. *Let $\psi: (0, \infty) \rightarrow (0, \infty)$ be a regular continuous function such that the map $q \mapsto q\psi(q)$ is nonincreasing and tends to zero, let $r_\psi(t) := e^{-t}\psi^{-1}(e^{-t})$ (this is well defined for large enough t), and let*

$$\text{A}(r_\psi, \Omega_{R, \Lambda_*}) := \{\Lambda \in \Omega_{R, \Lambda_*} : \delta_R(g_t \Lambda) \leq r_\psi(t) \text{ for an unbounded set of } t \geq 0\}. \quad (4.9)$$

Then, for every compact set $\mathcal{K} \subseteq \text{O}(R)$, there exists $C > 0$ (depending on ψ and \mathcal{K}) such that

$$\pi_1^{-1}(\text{A}_{M_Q}(\psi/C)) \cap \mathcal{K} \subseteq \pi_2^{-1}(\text{A}(r_\psi, \Omega_{R, \Lambda_*})) \cap \mathcal{K} \subseteq \pi_1^{-1}(\text{A}_{M_Q}(C\psi)) \cap \mathcal{K}.$$

Consequently, if g , $[\mathbf{x}]$, and Λ are as in (4.1), then $[\mathbf{x}] \in \text{WA}_{M_Q}(\psi) \setminus (\mathbb{P}_{\mathbb{Q}}^d \cap M_Q)$ if and only if

$$\Lambda \in \text{WA}(r_\psi, \Omega_{R, \Lambda_*}) := \bigcap_{\varepsilon > 0} \text{A}(\varepsilon r_\psi, \Omega_{R, \Lambda_*})$$

Proof. Given $g \in O(R)$ and $[\mathbf{x}], \Lambda$ as in (4.1), write $C([\mathbf{x}])$ for the left-hand side of (4.6) and write $C(\Lambda)$ for the right-hand side of (4.6). Then

$$C([\mathbf{x}]) < \alpha \implies [\mathbf{x}] \in A_{M_Q}(\alpha \psi) \implies C([\mathbf{x}]) \leq \alpha$$

and

$$C(\Lambda) < 1 \implies \Lambda \in A(r_\psi, \Omega_{R, \Lambda_*}) \implies C(\Lambda) \leq 1.$$

The conclusion follows. The “consequently” part follows from the regularity of ψ and the elementary computation $r_{\varepsilon\psi}(t) = e^{-t}\psi^{-1}(e^{-t}/\varepsilon)$. \blacksquare

In applying the correspondence principle, the following observations happen to be useful.

Observation 4.7. There exists a compact set $\mathcal{K} \subseteq O(R)$ such that $\pi_1(\mathcal{K}) = M_Q$.

Proof. This follows from the facts that M_Q is compact, $O(R)$ is locally compact, and π_1 is open and surjective. \blacksquare

We remark that the corresponding assertion is not true for π_2 since Ω_{R, Λ_*} is not compact by Corollary 3.14.

Now let μ_R and μ_{R, Λ_*} denote Haar measures on $O(R)$ and Ω_{R, Λ_*} , respectively.

Observation 4.8. The measures⁸ λ_{M_Q} and $\pi_1[\mu_R]$ are mutually absolutely continuous. The measures μ_{R, Λ_*} and $\pi_2[\mu_R]$ are mutually absolutely continuous.

We remark that Corollary 4.3, the ergodicity of the g_T -action on Ω_{R, Λ_*} , and the above observation allow one to conclude that the set BA_{M_Q} is λ_{M_Q} -null. This is a special case of a more general Khintchine-type result – namely Theorem 6.3.

5. A Dirichlet-type theorem

In this section, we prove the following.

Theorem 5.1 (Dirichlet-type theorem for quadric hypersurfaces). *Fix $d \geq 2$, and let M_Q be a nonsingular rational quadric hypersurface in $\mathbb{P}_{\mathbb{R}}^d$ with $p_{\mathbb{Q}} \geq 1$. Then*

- (i) ψ_1 is Dirichlet for intrinsic approximation on M_Q .
- (ii) ψ_1 is uniformly Dirichlet if and only if $p_{\mathbb{Q}} = p_{\mathbb{R}}$.
- (iii) The following are equivalent:
 - (A) $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$.
 - (B) There exist $C, T_0 > 0$ such that, for all $[\mathbf{x}] \in M_Q$ and for all $T \geq T_0$, there exists $[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ such that

$$H_{\text{std}}([\mathbf{r}]) \leq T \quad \text{and} \quad \text{dist}([\mathbf{r}], [\mathbf{x}]) \leq \frac{C}{\sqrt{H_{\text{std}}([\mathbf{r}])T}}. \quad (5.1)$$

⁸Note that the measures $\pi_1[\mu_R]$ and $\pi_2[\mu_R]$ are not σ -finite; in fact, they are $\{0, \infty\}$ -valued.

(C) *The set*

$$\{[\mathbf{x}] \in M_Q : \text{there exist } C, T_0 > 0 \text{ such that, for all } T \geq T_0, \\ \text{there exists } [\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q \text{ satisfying (5.1)}\}$$

has positive λ_{M_Q} -measure.

Except for the forward direction of (ii) (i.e. uniformly Dirichlet implies $p_{\mathbb{Q}} = p_{\mathbb{R}}$), which we will prove separately (see p. 1075), all of these results are consequences of the following theorem together with the correspondence principle,⁹ namely Lemma 4.1 (i), (ii) and Observations 4.7 and 4.8. Details are left to the reader.

Theorem 5.2. *Fix $d \geq 2$, and let R be a nonsingular quadratic form on \mathbb{R}^{d+1} with $p_{\mathbb{Q}} \geq 1$ which is $p_{\mathbb{Q}}$ -normalized. Fix $\Lambda_* \in \Omega_R$ commensurable to \mathbb{Z}^{d+1} . Then*

- (i) *for all $\Lambda \in \Omega_{R, \Lambda_*}$, there exists $C_{\Lambda} > 0$ such that*

$$\text{dist}(\mathbf{p}, \mathcal{L}_1) \leq C_{\Lambda} \quad (5.2)$$

holds for infinitely many $\mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$.

- (ii) *If $p_{\mathbb{Q}} = p_{\mathbb{R}}$, then the constant C_{Λ} in (5.2) can be made independent of Λ .*

- (iii) *The following are equivalent:*

(A) $p_{\mathbb{Q}} = p_{\mathbb{R}} = 1$.

- (B') *There exist $C, T_0 > 0$ such that, for all $\Lambda \in \Omega_{R, \Lambda_*}$ and for all $T \geq T_0$, there exists $\mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$ with $\|\mathbf{p}\| \leq T$ such that*

$$\text{dist}(\mathbf{p}, \mathcal{L}_1) \leq C \sqrt{\frac{\|\mathbf{p}\|}{T}}. \quad (5.3)$$

(C') *The set*

$$\{\Lambda \in \Omega_{R, \Lambda_*} : \text{there exist } C, T_0 > 0 \text{ such that, for all } T \geq T_0, \\ \text{there exists } \mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\} \text{ satisfying } \|\mathbf{p}\| \leq T \text{ and (5.3)}\}$$

has positive μ_{R, Λ_*} -measure.

Proof of (i). We require the following preliminary result.

Lemma 5.3. *Let Q be a nonsingular quadratic form on \mathbb{R}^{d+1} , and fix $\Lambda \in \Omega_Q$ satisfying $\Lambda \cap L_Q \setminus \{\mathbf{0}\} \neq \emptyset$. Then*

$$\text{Span}(\Lambda \cap L_Q) = \mathbb{R}^{d+1}.$$

⁹However, the correspondence principle cannot be used to deduce Theorem 5.2 from Theorem 5.1 (or similarly, Theorem 6.5 from Theorem 6.3), due to the lack of an analogue of Observation 4.7 for π_2 . Similar considerations prevent the forwards direction of Theorem 5.1 (ii) from being deduced from an appropriate analogue in the space of lattices.

Proof. After applying a matrix (namely one whose columns form a basis of Λ), we may without loss of generality assume that $\Lambda = \mathbb{Z}^{d+1}$. The assumption $\Lambda \cap L_Q \setminus \{\mathbf{0}\} \neq \emptyset$ will then imply that $p_Q \geq 1$, and by applying Proposition 3.5, we may without loss of generality assume that Q is 1-normalized and Λ is commensurable with \mathbb{Z}^{d+1} . Then, clearly,

$$r_0 \mathbf{e}_0, r_d \mathbf{e}_d \in \Lambda \cap L_Q \quad \text{for some nonzero } r_0, r_d \in \mathbb{Q}. \quad (5.4)$$

On the other hand, for each $i = 1, \dots, d-1$, we have

$$\mathbf{e}_i + Q(\mathbf{e}_i) \mathbf{e}_0 - \mathbf{e}_d \in L_Q$$

by direct calculation. Since Λ is Q -arithmetic and commensurable with \mathbb{Z}^{d+1} , it follows that

$$r_i(\mathbf{e}_i + Q(\mathbf{e}_i) \mathbf{e}_0 - \mathbf{e}_d) \in \Lambda$$

for some nonzero $r_i \in \mathbb{Q}$; hence, in view of (5.4), $\mathbf{e}_i \in \text{Span}(\Lambda \cap L_Q)$. \triangleleft

For $t \geq 0$, let $g_t \in \text{O}(R)$ be as in equation (3.5). Applying Corollary 3.14 to the lattices $(g_t \Lambda)_{t \geq 0}$, we see that one of the following two cases holds.

Case 1: There exists a sequence $t_n \rightarrow \infty$ and a sequence $g_{t_n}(\Lambda \cap L_R) \ni g_{t_n}(\mathbf{p}_n) \rightarrow 0$. In this case, for all sufficiently large n , (3.6) implies that \mathbf{p}_n satisfies (5.2). If the set $\{\mathbf{p}_n : n \in \mathbb{N}\}$ is infinite, this completes the proof. Otherwise, there exists $\mathbf{p} \in \Lambda$ such that $\mathbf{p}_n = \mathbf{p}$ for arbitrarily large n . In particular, we have $g_{t_{n_k}}(\mathbf{p}) \rightarrow 0$ for some increasing sequence $(n_k)_1^\infty$. Comparing with (3.5), we see that $\mathbf{p} \in \mathcal{L}_1$. Since the vectors $n\mathbf{p}$ ($n \in \mathbb{Z}$) all satisfy (5.2), this completes the proof.

Case 2: There exists a sequence $t_n \rightarrow \infty$ such that $g_{t_n} \Lambda \rightarrow \tilde{\Lambda} \in \Omega_{R, \Lambda_*}$. In this case, by Lemma 5.3, we have $\tilde{\Lambda} \cap L_R \not\subseteq \mathcal{L}_1^\perp$, where \mathcal{L}_1^\perp denotes the set of vectors Q -orthogonal to \mathbf{e}_1 as in Definition 3.1. Thus we may fix $\tilde{\mathbf{p}} \in \tilde{\Lambda} \cap L_R \setminus \mathcal{L}_1^\perp$. Since $g_{t_n} \Lambda \rightarrow \tilde{\Lambda}$, there is a sequence $g_{t_n} \Lambda \ni g_{t_n}(\mathbf{p}_n) \rightarrow \tilde{\mathbf{p}}$. Let $C_\Lambda = 2\|\tilde{\mathbf{p}}\|$; then for all sufficiently large n , (3.6) implies that \mathbf{p}_n satisfies (5.2). If the set $\{\mathbf{p}_n : n \in \mathbb{N}\}$ is infinite, this completes the proof. Otherwise, there exists $\mathbf{p} \in \Lambda$ such that $\mathbf{p}_n = \mathbf{p}$ for arbitrarily large n . In particular, $e^{t_{n_k}} \text{dist}(\mathbf{p}, \mathcal{L}_1^\perp) \rightarrow \text{dist}(\tilde{\mathbf{p}}, \mathcal{L}_1^\perp) \neq 0$ for some increasing sequence $(n_k)_1^\infty$. This is clearly a contradiction. \blacksquare

Proof of (ii). We first need to define the codiameter of a discrete subgroup.

Definition 5.4. The *codiameter* of a discrete subgroup $\Gamma \subseteq \mathbb{R}^{d+1}$, written $\text{Codiam}(\Gamma)$, is the diameter of the quotient space $\text{Span}(\Gamma)/\Gamma$.

We require the following lemma.

Lemma 8.11. *There exists $C_1 > 0$ such that, for every $\Lambda \in \Omega_{R, \Lambda_*}$, there exists a totally isotropic Λ -rational¹⁰ subspace $V \subseteq \mathbb{R}^{d+1}$ of dimension p_Q satisfying*

$$\text{Codiam}(V \cap \Lambda) \leq C_1.$$

¹⁰A subspace $V \subseteq \mathbb{R}^{d+1}$ is Λ -rational if $\text{Span}(\Lambda \cap V) = V$.

The proof of Lemma 8.11 requires reduction theory, so we delay its proof until Section 8.

Let C_1 be as in Lemma 8.11. Fix $\Lambda \in \Omega_{R, \Lambda_*}$. For each $t \geq 0$, applying Lemma 8.11 to the lattice $g_t \Lambda \in \Omega_{R, \Lambda_*}$ yields a totally isotropic $g_t \Lambda$ -rational subspace $V_t \subseteq \mathbb{R}^{d+1}$ of dimension $p_{\mathbb{Q}}$ satisfying

$$\text{Codiam}(V_t \cap g_t \Lambda) \leq C_1. \quad (5.5)$$

At this point, we divide the proof into two cases.

Case 1: $\mathcal{L}_1 \subseteq V_t$ for some $t \geq 0$. In this case, since the set

$$S := \{\mathbf{x} \in g_{-t}(V_t) : \text{dist}(\mathbf{x}, \mathcal{L}_1) \leq C_1\}$$

has infinite volume in the vector space $g_{-t}(V_t)$, by Minkowski's theorem, it contains infinitely many lattice points $\mathbf{p} \in \Lambda \cap S$. Note that each such \mathbf{p} is in L_R since V_t is totally isotropic. On the other hand, (5.2) is clearly satisfied (with $C_{\Lambda} = C_1$ independent of Λ). This completes the proof.

Case 2: $\mathcal{L}_1 \not\subseteq V_t$ for all $t \geq 0$. Fix $t \geq 0$. Note that if $V_t \subseteq \mathcal{L}_1^{\perp}$, then $V_t + \mathcal{L}_1$ is a totally isotropic vector space of dimension $p_{\mathbb{Q}} + 1 = p_{\mathbb{R}} + 1 > p_{\mathbb{R}}$, a contradiction. Thus $V_t \not\subseteq \mathcal{L}_1^{\perp}$. Fix a unit vector $\mathbf{v}_t \in V_t$ which is perpendicular to $V_t \cap \mathcal{L}_1^{\perp}$ with respect to the Euclidean quadratic form $\mathcal{E}_{d+1} = \sum_0^d x_i^2$. By (5.5), there exists $g_t(\mathbf{p}_t) \in V_t \cap g_t \Lambda$ satisfying $\|g_t(\mathbf{p}_t) - 2C_1 \mathbf{v}_t\| \leq C_1$. Then (3.6) implies that \mathbf{p}_t satisfies (5.2), $C_{\Lambda} = 3C_1$ independent of Λ . If the set $\{\mathbf{p}_t : t \geq 0\}$ is infinite, this completes the proof. Otherwise, there exists $\mathbf{p} \in \Lambda$ such that $\mathbf{p}_t = \mathbf{p}$ for arbitrarily large t . However, for all t , we have $g_t(\mathbf{p}_t) \in V_t \setminus (V_t \cap \mathcal{L}_1^{\perp}) = V_t \setminus \mathcal{L}_1^{\perp}$, and thus $\mathbf{p} \notin \mathcal{L}_1^{\perp}$. This implies that $\|g_t(\mathbf{p})\| \rightarrow \infty$, a contradiction. ■

Proof of (iii). For the purpose of this proof, we introduce a new system of coordinates on \mathbb{R}^{d+1} . For $\mathbf{x} \in \mathbb{R}^{d+1}$, let

$$H(\mathbf{x}) = |x_0|, \quad W(\mathbf{x}) = \|(x_1, \dots, x_{d-1})\|, \quad L(\mathbf{x}) = |x_d|.$$

We will think of the letters H , W , and L as being short for “height”, “width”, and “length”, respectively. Note that, for $t \in \mathbb{R}$,

$$H(g_t \mathbf{x}) = e^{-t} H(\mathbf{x}), \quad W(g_t \mathbf{x}) = W(\mathbf{x}), \quad L(g_t \mathbf{x}) = e^t L(\mathbf{x}).$$

In other words, for $t \geq 0$, applying g_t decreases height and increases length while leaving width fixed. Moreover,

$$\begin{aligned} \|\mathbf{x}\| &= \max(H(\mathbf{x}), W(\mathbf{x}), L(\mathbf{x})), \\ \text{dist}(\mathbf{x}, \mathcal{L}_1) &= \max(W(\mathbf{x}), L(\mathbf{x})). \end{aligned}$$

If $\mathbf{x} \in L_R$, then

$$H(\mathbf{x})L(\mathbf{x}) = |\tilde{R}(x_1, \dots, x_{d-1})| \leq \|\tilde{R}\| W^2(\mathbf{x}), \quad (5.6)$$

where \tilde{R} is the remainder of R .

We will now rephrase the Diophantine condition on a lattice $\Lambda \in \Omega_{R, \Lambda_*}$ described in (B') and (C') of Theorem 5.2 (iii) as a dynamical condition on the same lattice Λ . Precisely, we have the following observation.

Observation 5.6. Fix $C, T_0 \geq 1$ with $T_0 > C^2$, and fix $\Lambda \in \Omega_{R, \Lambda_*}$. Then we have that $(1) \Rightarrow (2) \Rightarrow (3)$.

(1) For all $t \geq \frac{1}{2} \log(T_0)$, there exists $\mathbf{q} \in g_t \Lambda \cap L_R \setminus \{\mathbf{0}\}$ satisfying

$$\|\mathbf{q}\| \leq C \quad \text{and} \quad W(\mathbf{q}) \leq \sqrt{CH(\mathbf{q})}.$$

(2) For all $T \geq T_0$, there exists $\mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$ with $\|\mathbf{p}\| \leq T$ satisfying (5.3).

(3) For all $t \geq \log(T_0)$, there exists $\mathbf{q} \in g_t \Lambda \cap L_R \setminus \{\mathbf{0}\}$ satisfying

$$\|\mathbf{q}\| \leq C^2 \max(1, \|\tilde{R}\|) \quad \text{and} \quad W(\mathbf{q}) \leq C \sqrt{H(\mathbf{q})}.$$

(1) \Rightarrow (2). Fix $T \geq T_0$, and let $t = \log(T/C) \geq \frac{1}{2} \log(T_0)$. Let $\mathbf{q} \in g_t \Lambda \cap L_R \setminus \{\mathbf{0}\}$ be as in (1), and let $\mathbf{p} = g_{-t}(\mathbf{q}) \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$. Then

$$\|\mathbf{p}\| \leq e^t \|\mathbf{q}\| \leq \frac{T}{C} C = T.$$

To demonstrate (5.3), we bound $W(\mathbf{p})$ and $L(\mathbf{p})$. First of all,

$$W(\mathbf{p}) = W(\mathbf{q}) \leq \sqrt{CH(\mathbf{q})} = \sqrt{C \frac{H(\mathbf{p})}{T/C}} = C \sqrt{\frac{H(\mathbf{p})}{T}}. \quad (5.7)$$

On the other hand, we have

$$L(\mathbf{p}) = \frac{L(\mathbf{q})}{T/C} \leq \frac{C}{T/C} = \frac{C^2}{T},$$

which implies

$$L(\mathbf{p}) = \sqrt{L(\mathbf{p})} \sqrt{L(\mathbf{p})} \leq \sqrt{L(\mathbf{p})} \sqrt{\frac{C^2}{T}} = C \sqrt{\frac{L(\mathbf{p})}{T}}.$$

Combining with (5.7) demonstrates (5.3). \triangleleft

(2) \Rightarrow (3). Fix $t \geq \log(T_0)$, and let $T = e^t \geq T_0$. Let $\mathbf{p} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$ be as in (2), and let $\mathbf{q} = g_t(\mathbf{p}) \in g_t \Lambda \cap L_R \setminus \{\mathbf{0}\}$. Then

$$H(\mathbf{q}) = e^{-t} H(\mathbf{p}) \leq e^{-t} T = 1.$$

On the other hand, (5.3) is written in terms of height, width, and length as

$$\max(W(\mathbf{p}), L(\mathbf{p})) \leq C \sqrt{\frac{\max(H(\mathbf{p}), W(\mathbf{p}), L(\mathbf{p}))}{T}},$$

and since $T \geq T_0 > C^2$, the case where the maximum is $W(\mathbf{p})$ or $L(\mathbf{p})$ cannot occur. Thus

$$\max(W(\mathbf{p}), L(\mathbf{p})) \leq C \sqrt{\frac{H(\mathbf{p})}{T}}.$$

In particular,

$$W(\mathbf{q}) = W(\mathbf{p}) \leq C \sqrt{\frac{H(\mathbf{p})}{e^t}} = C \sqrt{H(\mathbf{q})}.$$

Since $\mathbf{q} \in L_R$, (5.6) gives

$$L(\mathbf{q}) \leq \|\tilde{R}\| \frac{W^2(\mathbf{q})}{H(\mathbf{q})} \leq \|\tilde{R}\| \frac{C^2 H(\mathbf{q})}{H(\mathbf{q})} = C^2 \|\tilde{R}\|.$$

Thus $\|\mathbf{q}\| = \max(H(\mathbf{q}), W(\mathbf{q}), L(\mathbf{q})) \leq \max(1, C, C^2 \|R\|) \leq C^2 \max(1, \|R\|)$. \triangleleft

For each $C > 0$, consider the set

$$\mathcal{F}_C := \{\Lambda \in \Omega_{R, \Lambda_*} : \text{there exists } \mathbf{q} \in \Lambda \cap L_R \setminus \{\mathbf{0}\} \text{ such that} \\ \|\mathbf{q}\| \leq C, W(\mathbf{q}) \leq \sqrt{CH(\mathbf{q})}\}.$$

Then (B') and (C') of Theorem 5.2 (iii) are equivalent to the following conditions, respectively.

- (B'') There exists $C > 0$ such that, for all $\Lambda \in \Omega_{R, \Lambda_*}$ and for all $t \geq C$, $g_t \Lambda \in \mathcal{F}_C$.
(C'') The set

$$\{\Lambda \in \Omega_{R, \Lambda_*} : \text{there exists } C > 0 \text{ such that, for all } t \geq C, g_t \Lambda \in \mathcal{F}_C\} \\ = \bigcup_{C > 0} \liminf_{t \rightarrow \infty} g_{-t}(\mathcal{F}_C)$$

has positive μ_{R, Λ_*} -measure.

Now (B'') is clearly equivalent to the following.

- (B''') There exists $C > 0$ such that $\mathcal{F}_C = \Omega_{R, \Lambda_*}$.

We claim that (C'') is also equivalent to (B'''). Indeed, it is clear that (B''') implies (C''). Conversely, if (C'') holds, then by Moore's ergodicity theorem [1, Theorem III.2.1],¹¹ the set \mathcal{F}_C has full μ_{R, Λ_*} measure, where C is large enough so that the (g_t) -invariant set $\liminf_{t \rightarrow \infty} g_{-t}(\mathcal{F}_C)$ has positive measure. But since \mathcal{F}_C is closed, this implies (B''').

To complete the proof, we must show that (B''') is equivalent to (A).

Proof of (A) \Rightarrow (B'''). Since $p_{\mathbb{R}} = 1$, the remainder \tilde{R} does not represent zero over \mathbb{R} , i.e. it is either positive definite or negative definite. Without loss of generality, suppose that it is positive definite. Then $\sqrt{\tilde{R}}$ is a norm on \mathbb{R}^{d-1} , so there exists $K > 0$ such that

$$\tilde{R}(\mathbf{x}) \geq \frac{1}{K} W^2(\mathbf{x}) \quad \text{for all } \mathbf{x} \in \mathbb{R}^{d-1}.$$

¹¹If $d = 3$ and $p_{\mathbb{R}} = 2$, then the group $G = O(Q)$ is not simple (being isomorphic to $O(2, 2)$), so one should use [1, Theorem III.2.5] rather than [1, Theorem III.2.1]. Note that the fact that the group $(g_t)_{t \in \mathbb{R}}$ is totally noncompact in G follows from the inequality $(\pi_i)'(\mathbf{z}) \neq \mathbf{0}$ proven on p. 1083 of the present paper.

Then, for all $\mathbf{x} \in L_R$,

$$W^2(\mathbf{x}) \leq K \tilde{R}(x_1, \dots, x_{d-1}) = -K x_0 x_d = K H(\mathbf{x}) L(\mathbf{x}), \quad (5.8)$$

providing an asymptotic converse to (5.6).

Let $C_1 > 0$ be as in Lemma 8.11. Fix $\Lambda \in \Omega_{R, \Lambda_*}$, and we will show that $\Lambda \in \mathcal{F}_{C_1 K}$. Indeed, by Lemma 8.11, there exists $\mathbf{q} \in \Lambda \cap L_R \setminus \{\mathbf{0}\}$ satisfying $\|\mathbf{q}\| \leq C_1$. Then (5.8) gives

$$W(\mathbf{q}) \leq \sqrt{K H(\mathbf{q}) L(\mathbf{q})} \leq \sqrt{C_1 K H(\mathbf{q})},$$

demonstrating that $\Lambda \in \mathcal{F}_{C_1 K}$. \triangleleft

Proof of (B'') \Rightarrow (A).

Claim 5.7. *We may without loss of generality¹² suppose that R is $p_{\mathbb{R}}$ -normalized and that $\Lambda_* \cap \mathcal{L}_{p_{\mathbb{Q}}} = \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$.*

Proof. Let $E_{\mathbb{Q}}$ be a Λ_* -rational totally isotropic subspace of \mathbb{R}^{d+1} of dimension $p_{\mathbb{Q}}$. Let $E_{\mathbb{R}} \supseteq E_{\mathbb{Q}}$ be a totally isotropic subspace of \mathbb{R}^{d+1} of dimension $p_{\mathbb{R}}$. By Proposition 3.5, there is a matrix $\phi_1 \in \mathrm{GL}_{d+1}(\mathbb{R})$ such that $R' := R \circ \phi_1$ is $p_{\mathbb{R}}$ -normalized and $\phi_1^{-1}(E_{\mathbb{R}}) = \mathcal{L}_{p_{\mathbb{R}}}$. In particular, $\Gamma := \phi_1^{-1}(\Lambda_* \cap E_{\mathbb{Q}}) \subseteq \mathcal{L}_{p_{\mathbb{R}}}$. Let $\phi_2 \in \mathrm{GL}_{p_{\mathbb{R}}}(\mathbb{R})$ send Γ to $\mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$. Let g_{ϕ_2} be defined by equation (3.3) so that $g_{\phi_2} \in \mathrm{O}(R')$. Then $g_{\phi_2}^{-1}(\Gamma) = \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$. Letting $\phi = \phi_1 \circ g_{\phi_2}^{-1}$, we have $\phi^{-1}(\Lambda_* \cap E_{\mathbb{Q}}) = \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$, or equivalently $\phi^{-1}(\Lambda_*) \cap \mathcal{L}_{p_{\mathbb{Q}}} = \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$. Let $\Lambda'_* = \phi^{-1}(\Lambda_*)$, and observe that $R' = R \circ \phi$. Then R' is $p_{\mathbb{R}}$ -normalized and $\Lambda'_* \cap \mathcal{L}_{p_{\mathbb{Q}}} = \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}}$. On the other hand, both conditions (A) and (B'') are unaffected by replacing R and Λ_* with R' and Λ'_* , respectively. \triangleleft

Now suppose (A) fails, i.e. $p_{\mathbb{R}} > 1$. Fix $t \geq 0$, and let $\mathbf{t} = (t, \dots, t) \in \mathbb{R}^{p_{\mathbb{Q}}}$. Then

$$\Lambda_t := g_{\mathbf{t}} \Lambda_* \in \Omega_{R, \Lambda_*}.$$

Claim 5.8. *If $\mathbf{p} \in \Lambda_t \cap L_R$ satisfies $\|\mathbf{p}\| < e^t / (2\|R\|)$, then $\mathbf{p} \in \Gamma_t := \Lambda_t \cap \mathcal{L}_{p_{\mathbb{Q}}}$.*

Proof. For each $i = 0, \dots, p_{\mathbb{Q}} - 1$, we have $\mathbf{e}_i \in \mathbb{Z}^{d+1} \cap \mathcal{L}_{p_{\mathbb{Q}}} \subseteq \Lambda_* \cap L_R$, and thus $g_{\mathbf{t}}(\mathbf{e}_i) = e^{-t} \mathbf{e}_i \in \Lambda_t \cap L_R$. Since Λ_t is R -arithmetic, we have

$$B_R(\mathbf{p}, g_{\mathbf{t}}(\mathbf{e}_i)) \in \frac{\mathbb{Z}}{2}. \quad (5.9)$$

On the other hand,

$$|B_R(\mathbf{p}, g_{\mathbf{t}}(\mathbf{e}_i))| \leq \|R\| \cdot \|\mathbf{p}\| \cdot \|g_{\mathbf{t}}(\mathbf{e}_i)\| < \|R\| \left(\frac{e^t}{2\|R\|} \right) e^{-t} = \frac{1}{2}.$$

Combining with (5.9), we see that

$$B_R(\mathbf{p}, g_{\mathbf{t}}(\mathbf{e}_i)) = e^{-t} B_R(\mathbf{p}, \mathbf{e}_i) = 0.$$

¹²Here we abandon the assumption that Λ_* is commensurable to \mathbb{Z}^{d+1} .

It follows that the Λ_t -rational subspace $\mathcal{L}_{p_{\mathbb{Q}}} + \mathbb{R}\mathbf{p}$ is totally isotropic, and so, by the maximality of $p_{\mathbb{Q}}$, we have $\mathbf{p} \in \mathcal{L}_{p_{\mathbb{Q}}}$. \triangleleft

Now let $\phi_t \in \mathrm{O}(\mathcal{E}_{p_{\mathbb{R}}})$ satisfy $\phi_t(\Gamma_t) \cap \mathcal{L}_1 = \{\mathbf{0}\}$, where $\mathcal{E}_{p_{\mathbb{R}}}$ is the Euclidean metric on $\mathbb{R}^{p_{\mathbb{R}}}$. Such a choice is possible since by assumption $p_{\mathbb{R}} > 1$. Let g_{ϕ_t} be given by (3.3) so that $g_{\phi_t} \in \mathrm{O}(R) \cap \mathrm{O}(\mathcal{E}_{d+1})$. Let $\Lambda'_t = g_{\phi_t} \Lambda_t$.

Let

$$\gamma = \begin{bmatrix} & & 1 \\ & I_{d-1} & \\ 1 & & \end{bmatrix}$$

so that

$$\begin{aligned} \gamma(\mathcal{F}_C) &= \{\Lambda \in \Omega_{R, \Lambda_*} : \text{there exists } \mathbf{q} \in \Lambda \cap L_R \setminus \{\mathbf{0}\} \text{ such that} \\ &\quad \|\mathbf{q}\| \leq C, W(\mathbf{q}) \leq \sqrt{CL(\mathbf{q})}\}. \end{aligned}$$

We claim that, for all $C > 0$, there exists $t \geq 0$ such that $\Lambda'_t \notin \gamma(\mathcal{F}_C)$; in particular, $\mathcal{F}_C \subsetneq \Omega_{R, \Lambda_*}$. Indeed, fix C and t , and suppose we have $\mathbf{q} = g_{\phi_t}(\mathbf{p}) \in \Lambda'_t \cap L_R \setminus \{\mathbf{0}\}$ with $\|\mathbf{p}\| \asymp \|\mathbf{q}\| \leq C$ and $W(\mathbf{q}) \leq \sqrt{CL(\mathbf{q})}$. If t is large enough (depending on C), then by Claim 5.8, we have $\mathbf{p} \in \Gamma_t$ and thus $\mathbf{q} \in \mathcal{L}_{p_{\mathbb{R}}} \setminus \mathcal{L}_1$. In particular, $L(\mathbf{q}) = 0$, but $W(\mathbf{q}) > 0$. This is a contradiction. Thus $\mathcal{F}_C \subsetneq \Omega_{R, \Lambda_*}$ for all $C > 0$, so (B'') fails. \triangleleft

This completes the proof of Theorem 5.2. \blacksquare

We complete the proof of Theorem 5.1 by demonstrating the forwards direction of (ii).

Proof of Theorem 5.1, forwards direction of (ii). Let $V_{\mathbb{Q}}$ be a maximal isotropic \mathbb{Q} -subspace of \mathbb{R}^{d+1} and $V_{\mathbb{R}}$ a maximal isotropic \mathbb{R} -subspace of \mathbb{R}^{d+1} such that $V_{\mathbb{Q}} \subsetneq V_{\mathbb{R}}$. Then $[V_{\mathbb{Q}}] \subsetneq [V_{\mathbb{R}}]$. By contradiction, suppose that ψ_1 is uniformly Dirichlet. This is equivalent to the existence of a constant $C > 0$ such that, for all $[\mathbf{x}] \in M_Q$, there exist infinitely many $\mathbf{r} \in \mathbb{Z}^{d+1} \cap L_Q$ satisfying

$$\mathrm{dist}(\mathbf{r}, \mathcal{L}_{[\mathbf{x}]}) \leq C, \quad (5.10)$$

where $\mathcal{L}_{[\mathbf{x}]} = \mathbb{R}\mathbf{x}$.

Fix $[\mathbf{x}] \in [V_{\mathbb{R}}] \setminus [V_{\mathbb{Q}}] \subseteq M_Q$. Since $[\mathbf{x}] \notin [V_{\mathbb{Q}}]$, only finitely many $\mathbf{r} \in V_{\mathbb{Q}} \cap \mathbb{Z}^{d+1}$ can satisfy (5.10), so there exists $\mathbf{r} \in \mathbb{Z}^{d+1} \cap L_Q \setminus V_{\mathbb{Q}}$ satisfying (5.10). Let \mathbf{x} be the projection of \mathbf{r} onto $\mathcal{L}_{[\mathbf{x}]}$ so that

$$\|\mathbf{x} - \mathbf{r}\| = \mathrm{dist}(\mathbf{r}, \mathcal{L}_{[\mathbf{x}]}) \leq C. \quad (5.11)$$

Let $\mathbf{b}_1, \dots, \mathbf{b}_{p_{\mathbb{Q}}}$ be a basis of $V_{\mathbb{Q}} \cap \mathbb{Z}^{d+1}$. Since $V_{\mathbb{R}}$ is totally isotropic and $\mathbf{x} \in V_{\mathbb{R}}$, we have $B_Q(\mathbf{x}, \mathbf{b}_i) = 0$ for all $i = 1, \dots, p_{\mathbb{Q}}$. Thus

$$\begin{aligned} |B_Q(\mathbf{r}, \mathbf{b}_i)| &= |B_Q(\mathbf{x} - \mathbf{r}, \mathbf{b}_i)| \\ &\leq \|B_Q\| \cdot \|\mathbf{x} - \mathbf{r}\| \cdot \|\mathbf{b}_i\| \leq N := \left\lceil C \|B_Q\| \max_{i=0}^{p_{\mathbb{Q}}-1} \|\mathbf{b}_i\| \right\rceil, \end{aligned}$$

and so, since Q is \mathbb{Z}^{d+1} -arithmetic,

$$\mathbf{z} := (B_Q(\mathbf{r}, \mathbf{b}_i))_{i=0}^{p_{\mathbb{Q}}-1} \in \{-N, \dots, N\}^{p_{\mathbb{Q}}}.$$

On the other hand, since $\mathbf{r} \notin V_{\mathbb{Q}}$, the maximality of $V_{\mathbb{Q}}$ implies that $V_{\mathbb{Q}} + \mathbb{R}\mathbf{r}$ is not isotropic (it is clearly a \mathbb{Q} -subspace). Thus $B_Q(\mathbf{r}, \mathbf{b}_i) \neq 0$ for some $i = 1, \dots, p_{\mathbb{Q}}$, i.e.

$$\mathbf{z} \neq \mathbf{0}.$$

Choose real numbers $c_1, \dots, c_{p_{\mathbb{Q}}}$ linearly independent over \mathbb{Q} , and let

$$\mathbf{s} = \sum_{i=1}^{p_{\mathbb{Q}}} c_i \mathbf{b}_i \in V_{\mathbb{Q}}.$$

Let

$$[\mathbf{x}_m] \xrightarrow{m} [\mathbf{s}] \quad \text{with } [\mathbf{x}_m] \in [V_{\mathbb{R}}] \setminus [V_{\mathbb{Q}}].$$

For each m , let \mathbf{r}_m , \mathbf{x}_m , and \mathbf{z}_m be defined as above, with the additional stipulation that $\|\mathbf{r}_m\| \geq m$ (this is possible since there were infinitely many possible choices for \mathbf{r}_m). Then, for each $m \in \mathbb{N}$, we have

$$|B_Q(\mathbf{r}_m, \mathbf{s})| = |\mathbf{z}_m \cdot \mathbf{c}|,$$

where $\mathbf{c} = (c_i)_{i=0}^{p_{\mathbb{Q}}-1}$. Thus

$$|B_Q(\mathbf{r}_m, \mathbf{s})| \in \{|\mathbf{z} \cdot \mathbf{c}| : \mathbf{z} \in \{-N, \dots, N\}^{p_{\mathbb{Q}}} \setminus \{\mathbf{0}\}\},$$

which implies $|B_Q(\mathbf{r}_m, \mathbf{s})| \geq \varepsilon$ for some $\varepsilon > 0$ independent of m . Let $t_m = \pm \|\mathbf{x}_m\| / \|\mathbf{s}\|$; since $[\mathbf{x}_m] \xrightarrow{m} [\mathbf{s}]$, we have

$$\left\| \mathbf{s} - \frac{\mathbf{x}_m}{t_m} \right\| \xrightarrow{m} 0$$

after choosing the appropriate \pm signs to define t_m . Now

$$\begin{aligned} \varepsilon t_m &\leq |B_Q(\mathbf{r}_m, t_m \mathbf{s})| \\ &= |B_Q(\mathbf{r}_m - \mathbf{x}_m, t_m \mathbf{s})| && \text{(since } \mathbf{x}_m, \mathbf{s} \in V_{\mathbb{R}}\text{)} \\ &\leq |B_Q(\mathbf{r}_m - \mathbf{x}_m, t_m \mathbf{s} - \mathbf{x}_m)| + |B_Q(\mathbf{r}_m - \mathbf{x}_m, \mathbf{x}_m)| \\ &= |B_Q(\mathbf{r}_m - \mathbf{x}_m, t_m \mathbf{s} - \mathbf{x}_m)| + \frac{1}{2} |Q(\mathbf{r}_m) - Q(\mathbf{x}_m) - Q(\mathbf{r}_m - \mathbf{x}_m)| \\ &= |B_Q(\mathbf{r}_m - \mathbf{x}_m, t_m \mathbf{s} - \mathbf{x}_m)| + \frac{1}{2} |Q(\mathbf{r}_m - \mathbf{x}_m)| && \text{(since } \mathbf{r}_m, \mathbf{x}_m \in L_Q\text{)} \\ &\leq \|Q\| \cdot \|\mathbf{r}_m - \mathbf{x}_m\| \left[\|t_m \mathbf{s} - \mathbf{x}_m\| + \frac{1}{2} \|\mathbf{r}_m - \mathbf{x}_m\| \right] \\ &\leq C \|Q\| \left(\frac{C}{2} + \|t_m \mathbf{s} - \mathbf{x}_m\| \right). && \text{(by (5.11))} \end{aligned}$$

Dividing by t_m , we have

$$\varepsilon \lesssim \frac{1}{t_m} + \left\| \mathbf{s} - \frac{\mathbf{x}_m}{t_m} \right\| \xrightarrow{m} 0,$$

a contradiction. ■

Remark 5.9. The hypothesis of nonsingularity can be dropped from parts (i) and (ii) of Theorem 5.1 if the hypothesis that $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q \neq \emptyset$ is replaced by the stronger hypothesis that \mathbb{Z}^{d+1} intersects $L_Q \setminus (\mathbb{R}^{d+1})^\perp$.

Proof. Any singular quadratic form is conjugate to a quadratic form $Q: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ of the form

$$Q(x_0, \dots, x_d) = \tilde{Q}(x_0, \dots, x_m),$$

where \tilde{Q} is a nonsingular quadratic form on \mathbb{R}^{m+1} for some $m < d$. In particular, we have $L_Q = L_{\tilde{Q}} \times \mathbb{R}^{d-m}$. Note that the hypothesis on Q guarantees that $\mathbb{P}_{\mathbb{Q}}^m \cap M_{\tilde{Q}} \neq \emptyset$.

Fix $[\mathbf{x}] \in M_Q$ and a representative $\mathbf{x} = (\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) \in L_Q$. Suppose first that $\mathbf{x}^{(1)} \neq 0$, and let $\mathbf{r}^{(1)} \in \mathbb{Z}^{m+1} \cap L_{\tilde{Q}}$ be such that

$$\text{dist}(\mathbf{r}^{(1)}, \mathbb{R}\mathbf{x}^{(1)}) \leq C_{[\mathbf{x}^{(1)}]}. \quad (5.12)$$

Then there exists $t \in \mathbb{R}$ so that $\|\mathbf{r}^{(1)} - t\mathbf{x}^{(1)}\| \leq C_{[\mathbf{x}^{(1)}]}$. Choose $\mathbf{r}^{(2)} \in \mathbb{Z}^{d-m}$ so that $\|\mathbf{r}^{(2)} - t\mathbf{x}^{(2)}\| \leq 1$. Then

$$\|(\mathbf{r}^{(1)}, \mathbf{r}^{(2)}) - t\mathbf{x}\| \leq C_{[\mathbf{x}^{(1)}]} + 1. \quad (5.13)$$

Now, by Theorem 5.1 (i) applied to \tilde{Q} , there exist infinitely many $\mathbf{r}^{(1)} \in \mathbb{Z}^{m+1} \cap L_{\tilde{Q}}$ satisfying (5.12); thus there exist infinitely many pairs $(\mathbf{r}^{(1)}, \mathbf{r}^{(2)})$ satisfying (5.13).

On the other hand, if $\mathbf{x}^{(1)} = 0$, let $\mathbf{r}^{(1)} = 0$, and for each $t \in \mathbb{R}$, choose $\mathbf{r}^{(2)}$ satisfying $\|\mathbf{r}^{(2)} - t\mathbf{x}^{(2)}\| \leq 1$; then (5.13) holds. Letting $t \rightarrow \infty$, there exist infinitely many pairs $(\mathbf{r}^{(1)}, \mathbf{r}^{(2)})$ satisfying (5.13).

Finally, if $p_{\mathbb{Q}} = p_{\mathbb{R}}$, then by using Theorem 5.1 (ii) in place of Theorem 5.1 (i), the above argument shows that the implied constant is independent of \mathbf{x} . ■

Remark 5.10. The same technique cannot be used to remove the nonsingularity hypothesis from Theorem 6.3 below. Indeed, if we suppose that $[\mathbf{x}^{(1)}] \in A_{\psi, M_{\tilde{Q}}}$ for some ψ , then $C_{[\mathbf{x}^{(1)}]}$ will be replaced by $CH_{\text{std}}([\mathbf{r}])\psi \circ H_{\text{std}}([\mathbf{r}])$ in (5.13), but the second term (namely 1) will not be changed. Thus the overall bound is no better than if we did not know that $[\mathbf{x}^{(1)}] \in A_{\psi, M_{\tilde{Q}}}$.

Remark 5.11. The hypothesis that M_Q is rational certainly cannot be dropped from Theorem 5.1. Indeed, Theorem 5.1 (i) implies that the set $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ is dense in M_Q whenever M_Q is a nonsingular rational quadric hypersurface in $\mathbb{P}_{\mathbb{R}}^d$ satisfying $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q \neq \emptyset$. By contrast, if Q is a quadratic form which is not a scalar multiple of any quadratic form with integer coefficients, then $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ is not dense in M_Q .

Proof. Let $\pi: \mathbb{R} \rightarrow \mathbb{Q}$ be a \mathbb{Q} -linear map, and let $R: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be the unique quadratic form so that $R = \pi \circ Q$ on \mathbb{Q}^{d+1} . Then, for $\mathbf{r} \in \mathbb{Q}^{d+1}$, $Q(\mathbf{r}) = 0$ implies $R(\mathbf{r}) = 0$; thus $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q \subseteq M_R$. If $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q$ is dense in M_Q , then $M_Q \subseteq M_R$, and so Q is a scalar multiple of R . But R has rational coefficients and is therefore a scalar multiple of a quadratic form with integer coefficients. ■

6. Khintchine-type theorems and counting of rational points

Recall that, in the classical setting, the convergence case of Khintchine's theorem follows directly from the Borel–Cantelli lemma combined with estimates for the number of rational points whose height is less than a fixed number T . So, in the case of intrinsic approximation, one must find upper bounds on expressions of the form

$$N_M(T) := \#\{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M : H_{\text{std}}([\mathbf{r}]) \leq T\},$$

where $M \subseteq \mathbb{P}_{\mathbb{R}}^d$ is an arbitrary manifold. Such bounds have been considered extensively in the case where M is algebraic in [10]. We will pay special attention to the following result due to D. R. Heath-Brown. Recall that Q is a rational quadratic form in $d+1$ variables, $\dim(M_Q) = d-1$, and Q_0 is the exceptional quadratic form on \mathbb{R}^4 defined in (2.7).

Theorem 6.1 ([30, Theorems 5, 6, 7, 8 and remarks afterwards]). *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Then*

$$N_{M_Q}(T) \asymp_{\times} \begin{cases} T^{d-1}, & Q \sim Q_0, \\ T^2 \log T, & Q \sim Q_0. \end{cases} \quad (6.1)$$

In order to clarify the relation between the above paraphrased version of Heath-Brown's results with the original theorems, we make the following comments.

(1) Theorems 5, 6, 7, and 8 in [30] provide asymptotics with an error term for the weighted sum

$$N(F, w) = N(F, w, P) := \sum_{\mathbf{x} \in \mathbb{Z}^{d+1} \cap F^{-1}(0)} w(P^{-1}\mathbf{x}),$$

where F is a rational quadratic form in $d+1$ variables, and w a function on \mathbb{R}^{d+1} which is required to be \mathcal{C}^∞ . However, to estimate $N_{M_Q}(T)$, one must let $w = \mathbb{1}_{B(0,1)}$. Since $w_0 = \mathbb{1}_{B(0,1)}$ can be approximated from above and below by \mathcal{C}^∞ functions w_n in a way such that the singular integrals $\sigma_\infty(F, w_n)$ approach $\sigma_\infty(F, w_0) \in (0, \infty)$ as $n \rightarrow \infty$, [30, Theorems 5, 6, 7, and 8] will still hold for $w_0 = \mathbb{1}_{B(0,1)}$, but without an estimate on the error term; namely, we have

$$\lim_{P \rightarrow \infty} \frac{N(F, w_0, P)}{\text{leading term}} = 1$$

for each result in [30]. In Theorem 6.1, we have stated only the weaker conclusion that the left-hand side is bounded from above and below (in \limsup and \liminf , respectively).

(2) According to [30, Theorems 5, 6, 7, and 8], the number of integer vectors on quadric hypersurfaces $Q^{-1}(0)$ of \mathbb{R}^{d+1} inside the ball of radius T is up to a multiplicative constant asymptotically equal to

$$\begin{cases} T^{d-1} & \text{if } d \geq 4 \\ T^2 & \text{if } d = 3 \text{ and } Q \sim Q_0 \\ T^2 \log T & \text{if } d = 3 \text{ and } Q \sim Q_0 \\ T \log T & \text{if } d = 2 \end{cases} \quad \begin{array}{ll} [\text{30, Theorem 5}], & \\ [\text{30, Theorem 6}], & \\ [\text{30, Theorem 7}], & \\ [\text{30, Theorem 8}]. & \end{array} \quad (6.2)$$

Note however that our goal is to count rational points on M_Q , which correspond to *primitive* integer vectors on $Q^{-1}(0)$. The relation between counting primitive vectors and counting all lattice vectors is clarified in [30] after the theorems are stated. In particular, [30, Theorems 5, 6, and 7] lead to equivalent results for counting of primitive vectors, which the only change is that the leading term is divided by a constant. However, the situation with [30, Theorem 8] is different: in view of [30, Corollary 2], for the count of primitive integer vectors, the factor $\log T$ in the last line of (6.2) disappears.

(3) In [30], it is shown that the modified singular series σ^* is positive and finite if and only if the equation $Q = 0$ has nontrivial solutions in every p -adic field. Since the forms we deal with satisfy $\mathbb{P}_{\mathbb{Q}}^d \cap M_Q \neq \emptyset$, the equation $Q = 0$ has nontrivial solutions over \mathbb{Q} , and so certainly over every p -adic field.

For any nonincreasing function $\psi: \mathbb{N} \rightarrow (0, \infty)$, we may write

$$A_{M_Q}(\psi) \subseteq \limsup_{\substack{T \rightarrow \infty \\ T \in 2^{\mathbb{N}}}} \bigcup_{\substack{[\mathbf{r}] \in \mathbb{P}_{\mathbb{Q}}^d \cap M_Q \\ H_{\text{std}}([\mathbf{r}]) \leq 2T}} B([\mathbf{r}], \psi(T)).$$

Combining with (6.1) and using the Hausdorff–Cantelli lemma [4, Lemma 3.10], one can immediately deduce the following corollary.

Corollary 6.2. *Suppose that $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ is a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Fix a positive $s \leq d - 1$, and let $\psi: \mathbb{N} \rightarrow (0, \infty)$ be nonincreasing. If the series*

$$\begin{cases} \sum_{T \in 2^{\mathbb{N}}} T^{d-1} \psi^s(T), & Q \sim Q_0, \\ \sum_{T \in 2^{\mathbb{N}}} T^2 \log T \psi^s(T), & Q \sim Q_0, \end{cases} \quad (6.3)$$

converges, then $\mathcal{H}^s(A_{M_Q}(\psi)) = 0$.

The case $s = d - 1$ corresponds to Lebesgue measure.

Based on the above, one would expect that Khintchine’s theorem for quadric hypersurfaces would state that the converse of Corollary 6.2 holds when $s = d - 1$ (possibly with some additional assumptions on ψ). However, we instead have the following.

Theorem 6.3 (Khintchine-type theorem for quadric hypersurfaces). *Let $M_Q \subseteq \mathbb{P}_{\mathbb{R}}^d$ be a nonsingular rational quadric hypersurface with $p_{\mathbb{Q}} \geq 1$. Fix $\psi: \mathbb{N} \rightarrow (0, \infty)$, and suppose that ψ is regular (see Definition 2.4) and that the function $q \mapsto q\psi(q)$ is nonincreasing. Then $A_{M_Q}(\psi)$ has full Lebesgue measure if the series*

$$\begin{cases} \sum_{T \in 2^{\mathbb{N}}} T^{d-1} \psi^{d-1}(T), & Q \sim Q_0, \\ \sum_{T \in 2^{\mathbb{N}}} T^2 \log \log T \psi^2(T), & Q \sim Q_0, \end{cases} \quad (6.4)$$

diverges; otherwise, $A_{M_Q}(\psi)$ is Lebesgue null.

In other words, whenever $Q \sim Q_0$, the above intuition is correct: Theorem 6.3 then says that, when $Q \sim Q_0$, the converse to the standard Borel–Cantelli argument holds for the collection of sets defining $A_{M_Q}(\psi)$. On the other hand, the series (6.4) does not agree with (6.3) when $Q \sim Q_0$, and so, philosophically, there is some nontrivial relation

between the sets appearing in the definition of $A_{M_{Q_0}}(\psi)$. A description of this relation is given in Section 9 (see in particular Remark 9.3), where an elementary proof of the convergence case of Theorem 6.3 for the manifold M_{Q_0} is given.

Using the Mass Transference Principle of Beresnevich and Velani [3, Theorem 2], one can immediately deduce the following.¹³

Theorem 6.4 (Jarník–Besicovitch theorem for quadric hypersurfaces). *Fix $0 < s < d - 1$. Let $\psi: \mathbb{N} \rightarrow (0, \infty)$ be regular, and suppose that $q \mapsto q^{d-1}\psi^s(q)$ is nonincreasing. If the series*

$$\begin{cases} \sum_{T \in 2^{\mathbb{N}}} T^{d-1}\psi^s(T), & Q \sim Q_0, \\ \sum_{T \in 2^{\mathbb{N}}} T^2 \log \log T \psi^s(T), & Q \sim Q_0, \end{cases} \quad (6.5)$$

diverges, then $\mathcal{H}^s(A_{M_Q}(\psi)) = \infty$.

This, in particular, computes the Hausdorff dimension of the set of ψ_c -approximable points of M_Q ; see (2.8).

It follows from Corollary 6.2 that, for $Q \sim Q_0$, convergence of (6.5) implies

$$\mathcal{H}^s(A_{M_Q}(\psi)) = 0.$$

However, in the case of the exceptional quadratic form Q_0 , there is a discrepancy between (6.5) and the series (6.3) appearing in Corollary 6.2, and the former may converge, while the latter diverges. In this case, we do not know the value of $\mathcal{H}^s(A_{M_Q}(\psi))$. However, the coarser Hausdorff dimension result (2.8) holds regardless. For reasons explained in Remark 9.3, the authors conjecture that Theorem 6.4 remains true if (6.5) is replaced by (6.3).

Note also that if $q^2\psi(q) \rightarrow 0$, then all ψ -good rational approximations of points in M_Q are intrinsic, meaning that $A_{M_Q}(\psi) = A_d(\psi) \cap M_Q$ (see [18, Lemma 4.1.1]). Consequently, for such ψ , Theorem 6.4 may be rephrased in terms of ambient approximation. The rephrased result has been proven in the case $Q_{\text{aff}}(\mathbf{x}) = x_1^2 + x_2^2$ by Dickinson and Dodson [17, Theorem 1], and in the case where $Q \sim Q_0$ by Druțu [18, Theorem 4.5.7].¹⁴

Note that Theorem 6.3 is analogous to the main result of [28], the difference being that we are considering intrinsic approximation and the authors of [28] are considering a specific type of extrinsic approximation. Also, it is likely that the techniques of Druțu [18] can be used to prove Theorem 6.3 in the case $Q \sim Q_0$ via the use of ubiquitous systems as considered in [2]. On the other hand, Druțu's methods do not apply to the exceptional quadric hypersurface M_{Q_0} (cf. footnote 14). We opt to use the machinery of Kleinbock and Margulis [37] to establish Theorem 6.3.

¹³The dimension $s > 0$ may be replaced by a dimension function f ; we omit the statement for brevity.

¹⁴Although the hypothesis $Q \sim Q_0$ does not appear explicitly in Druțu's theorem, it is required by her standing assumption that the lattice Γ is irreducible (cf. [18, Section 2.5, Section 4.5]) since, when $Q \sim Q_0$, Γ is reducible (see p. 1083).

Theorem 6.3 can be deduced directly from the following theorem together with the correspondence principle (Corollary 4.6 and Observation 4.8). As before, details are left to the reader.¹⁵

Theorem 6.5. *Fix $d \geq 2$, let R be a nonsingular $p\mathbb{Q}$ -normalized quadratic form on \mathbb{R}^{d+1} , and fix $\Lambda_* \in \Omega_R$ commensurable to \mathbb{Z}^{d+1} . Let $\psi: (0, \infty) \rightarrow (0, \infty)$ be a continuous function, and suppose that $q \mapsto q\psi(q)$ is nonincreasing. Let $r_\psi: (0, \infty) \rightarrow (0, \infty)$ and $\mathbf{A}_R(\psi) = \mathbf{A}(r_\psi, \Omega_{R, \Lambda_*})$ be defined as in Corollary 4.6; see (4.9). Then $\mathbf{A}_R(\psi)$ has full measure with respect to μ_{R, Λ_*} if (6.4) diverges; otherwise, $\mathbf{A}_R(\psi)$ is null with respect to μ_{R, Λ_*} .*

The proof of Theorem 6.5 will occupy Sections 7 and 8.

7. Proof of Theorem 6.5 modulo a volume computation

In the current section, we reduce Theorem 6.5 to a statement about the asymptotic behavior of the measure μ_{R, Λ_*} . Namely, we will deduce Theorem 6.5 as a corollary of one of the main results of [37], which we now recall.

Definition 7.1. Let (X, dist_X) be a metric space, let μ be a (finite Borel) measure on X , and let $\Delta: X \rightarrow \mathbb{R}$ be a continuous function. For each $z \in \mathbb{R}$, let

$$S_{\Delta, z} = \{x \in X : \Delta(x) \geq z\} \quad \text{and} \quad \Phi_\Delta(z) = \mu(S_{\Delta, z}),$$

where Φ_Δ is called the *tail distribution function* of Δ . We say that Δ is *distance-like* if

- (I) Δ is uniformly continuous, and
- (II) Φ_Δ is regular (see Definition 2.4).

Let G be a connected semisimple center-free Lie group without compact factors, and let $\Gamma \subseteq G$ be a lattice. By [48, Theorem 5.22], one can find connected normal subgroups $G_1, \dots, G_\ell \leq G$ such that G is the direct product of G_1, \dots, G_ℓ , $\Gamma_i := \Gamma \cap G_i$ is an irreducible lattice in G_i for each $i = 1, \dots, \ell$, and $\prod_{i=1}^\ell \Gamma_i$ has finite index in Γ . Of course, if Γ is irreducible, then we have $\ell = 1$, $G_1 = G$, and $\Gamma_1 = \Gamma$. Let π_1, \dots, π_ℓ denote the projections from G to the factors G_i .

Theorem 7.2 ([37, Theorem 1.7 (a)]). *Fix $G, \Gamma, G_1, \dots, G_\ell$ as above. Let \mathfrak{g} denote the Lie algebra of G , and let $\mathbf{z} \in \mathfrak{g}$ be an element of a Cartan subalgebra of \mathfrak{g} . Suppose that $(\pi_i)'(\mathbf{z}) \neq \mathbf{0}$ for all $i = 1, \dots, \ell$. (If G is simple, this just amounts to saying that $\mathbf{z} \neq \mathbf{0}$.) Let $X = G/\Gamma$, let μ_X be normalized Haar measure on X , let dist_G be a right-invariant*

¹⁵It is helpful to notice that the convergence/divergence of the series (6.4) is unaffected by the substitution $\psi \mapsto C\psi$, where $C > 0$ is a constant. Also, the fact that the assumption $q\psi(q) \rightarrow 0$ appears in Corollary 4.6 but not Theorem 6.3 can be remedied by the observation that BA_{M_Q} has measure zero, which follows either from applying Theorem 6.3 to any function ψ satisfying the hypotheses and such that the series (6.4) diverges, or by the argument at the end of Section 4.

Riemannian metric on G , let dist_X be the quotient of dist_G by Γ , and let $\Delta: X \rightarrow \mathbb{R}$ be a distance-like function.¹⁶ If $(z_t)_1^\infty$ is a sequence in \mathbb{R} , then

$$\begin{aligned} \mu_X(\{x \in X : e^{t\mathbf{z}}(x) \in S_{\Delta, z_t} \text{ for infinitely many } t \in \mathbb{N}\}) \\ = \begin{cases} 0 & \text{if } \sum_{t=1}^\infty \Phi_\Delta(z_t) < \infty, \\ 1 & \text{if } \sum_{t=1}^\infty \Phi_\Delta(z_t) = \infty. \end{cases} \end{aligned}$$

Remark 7.3. In [37, Theorem 1.7(a)], Γ is assumed to be irreducible, and \mathbf{z} is simply assumed to be a nonzero vector in α . However, in [37, Section 10.3], the authors of [37] describe how to modify their proof to include the case where Γ is reducible. Incorporating those modifications leads to the above theorem.

For the purposes of this paper, it will be more convenient to deal with the following “continuous” version of Theorem 7.2.

Theorem 7.4. Let $G, \Gamma, \alpha, \mathbf{z}, X, \mu_X, \Delta$ be as in Theorem 7.2. If $z: (0, \infty) \rightarrow (0, \infty)$ is nondecreasing, then

$$\begin{aligned} \mu_X(\{x \in X : e^{t\mathbf{z}}(x) \in S_{\Delta, z(t)} \text{ for arbitrarily large } t > 0\}) \\ = \begin{cases} 0 & \text{if } \sum_{t=1}^\infty \Phi_\Delta \circ z(t) < \infty, \\ 1 & \text{if } \sum_{t=1}^\infty \Phi_\Delta \circ z(t) = \infty. \end{cases} \end{aligned} \tag{7.1}$$

Proof of Theorem 7.4 using Theorem 7.2. Let $z_t^{(1)} = z(t)$, and let $z_t^{(2)} = z(t) - C$ for some $C > 0$. To complete the proof, it suffices to demonstrate the following:

- (i) $\sum_{t=1}^\infty \Phi_\Delta(z_t^{(i)}) < \infty$ if and only if $\sum_{t=1}^\infty \Phi_\Delta \circ z(t) < \infty$,
- (ii) $e^{t\mathbf{z}}(x) \in S_{\Delta, z_t^{(1)}}$ for infinitely many $t \in \mathbb{N}$ implies $e^{t\mathbf{z}}(x) \in S_{\Delta, z(t)}$ for arbitrarily large $t > 0$, and
- (iii) if C is large enough, then we have that $e^{t\mathbf{z}}(x) \in S_{\Delta, z(t)}$ for arbitrarily large $t > 0$ implies $e^{t\mathbf{z}}(x) \in S_{\Delta, z_t^{(2)}}$ for infinitely many $t \in \mathbb{N}$.

Indeed, (i) follows from the fact that Φ_Δ is regular (since Δ is assumed distance-like), and (ii) is obvious, so we turn to (iii). Suppose that $e^{t\mathbf{z}}(x) \in S_{\Delta, z(t)}$ for some t , and let $t' = \lfloor t \rfloor$. Then

$$\text{dist}_X(e^{t'\mathbf{z}}(x), e^{t\mathbf{z}}(x)) \leq C_1$$

for some constant $C_1 > 0$; since Δ is uniformly continuous, there exists $C = C_2 > 0$ independent of t so that $|\Delta(e^{t'\mathbf{z}}(x)) - \Delta(e^{t\mathbf{z}}(x))| \leq C_2$. On the other hand, since z is nondecreasing, $z_{t'}^{(2)} \leq z(t) - C$; it follows that $e^{t'\mathbf{z}}(x) \in S_{\Delta, z_t^{(2)}}$. \blacksquare

¹⁶We remark that whether or not Δ is distance-like is independent of the choice of the right-invariant Riemannian metric dist_G since any two such metrics $\text{dist}_1, \text{dist}_2$ satisfy $\text{dist}_1 \asymp_{\times} \text{dist}_2$.

Let $O(R)_0$ denote the identity component of $O(R)$. We claim that Theorem 6.5 follows from applying Theorem 7.4 with

$$\begin{aligned} G &= O(R)_0, & \Gamma &= O(R; \Lambda_*) \cap O(R)_0, \\ X &= G/\Gamma \equiv \Omega_{R, \Lambda_*}, & \Delta &= -\log \delta, \quad \text{where } \delta \text{ is as in (3.7),} \\ \mathbf{z} &= \frac{\partial}{\partial t} g_t \Big|_{t=0} = \begin{bmatrix} -1 & & & \\ & \mathbf{0}_{d-1} & & \\ & & 1 & \\ & & & \end{bmatrix}, \text{ and} \\ z(t) &= -\log r_\psi(t). \end{aligned} \tag{7.2}$$

Obviously, the verification of this claim consists of two parts: showing that the hypotheses of Theorem 7.4 are satisfied, and showing that Theorem 6.5 follows from the conclusion of Theorem 7.4.

Verification of the hypotheses. The verification of hypotheses is mostly a consequence of well-known facts; we leave the details to the reader, proving only the following statements.

(1) $(\pi_i)'(\mathbf{z}) \neq \mathbf{0}$ for all i . To see this, note that the group G is isomorphic to $O(p, q)_0$, where $p = p_{\mathbb{R}}$ and $q = d + 1 - p_{\mathbb{R}}$. Now $O(p, q)_0$ is simple as long as $p + q \geq 3$ and $(p, q) \notin \{(4, 0), (2, 2), (0, 4)\}$; if $(p, q) \in \{(4, 0), (2, 2), (0, 4)\}$, then $O(p, q)_0$ is only semisimple. In our case, we have $1 \leq p \leq q$ and $p + q = d + 1 \geq 3$, so G is simple unless $p = q = 2$. If G is simple, there is nothing to prove, so assume that $p = q = 2$. Then, by Proposition 3.5, $G \equiv O(2, 2)_0$ is conjugate in $SL_4(\mathbb{R})$ to $O(Q_0)_0$, where

$$Q_0(\mathbf{x}) = x_0 x_3 - x_1 x_2$$

is the exceptional quadratic form; moreover, it is readily seen that

$$O(Q_0)_0 = SL_2(\mathbb{R}) \times SL_2(\mathbb{R}),$$

where $G \times H$ denotes the set of all matrices of the form $g \otimes h$, where $g \in G$ and $h \in H$. (See the ‘‘product structure’’ of M_{Q_0} described in Section 9). Write

$$G = \phi(SL_2(\mathbb{R}) \times SL_2(\mathbb{R}))$$

for some matrix $\phi \in SL_4(\mathbb{R})$. Then the factors of G are given by the formulas¹⁷

$$G_1 = \phi(SL_2(\mathbb{R}) \times I), \quad G_2 = \phi(I \times SL_2(\mathbb{R})).$$

The tangent spaces are given by the formulas $g_1 = \phi(\mathfrak{sl}_2(\mathbb{R}) \times I)$, $g_2 = \phi(I \times \mathfrak{sl}_2(\mathbb{R}))$. Now any element of either of these tangent spaces has eigenvalues $\lambda, \lambda, -\lambda, -\lambda$ for some $\lambda \in \mathbb{R}$. On the other hand, the eigenvalues of \mathbf{z} are $1, 0, 0, -1$. Thus $\mathbf{z} \notin g_1, g_2$. It follows that $(\pi_i)'(\mathbf{z}) \neq \mathbf{0}$ for all i .

(2) Δ is uniformly continuous. To see this, fix $g \in G$ and $\Lambda \in X$; then for all $\mathbf{r} \in \Lambda$, we have $\|g\mathbf{r}\| \leq \|g\| \cdot \|\mathbf{r}\|$, where $\|g\|$ is the operator norm of g . Taking the minimum over $\mathbf{r} \in \Lambda \setminus \{\mathbf{0}\}$ gives $\delta(g\Lambda) \leq \|g\| \delta(\Lambda)$, or equivalently $\Delta(\Lambda) \leq \Delta(g\Lambda) + \log \|g\|$.

¹⁷If Γ is irreducible, then there will actually be only one factor, namely G , and so, as before, there is nothing to prove. (In fact, this happens if and only if $p_{\mathbb{Q}} = 1$.)

A symmetric argument gives

$$\Delta(g\Lambda) \leq \Delta(\Lambda) + \log\|g^{-1}\|.$$

Since $\log\|g\|, \log\|g^{-1}\| \leq \text{dist}_G(\text{id}, g)$ for all g , it follows that

$$\Delta \text{ is 1-Lipschitz.} \quad (7.3)$$

(3) Φ_Δ is regular. This will be a consequence of the following asymptotic formula for $\Phi_\Delta(z)$, whose proof will occupy Section 8, and which we will make further use of below.

Proposition 8.9. *For z large enough,*

$$\Phi_\Delta(z) \asymp_x \begin{cases} e^{-(d-1)z}, & R \sim Q_0, \\ e^{-2z}z, & R \sim Q_0. \end{cases}$$

This completes the verification of the hypotheses of Theorem 6.5. \blacksquare

Completion of the proof. First, we rewrite (7.1) using (7.2):

$$\begin{aligned} & \begin{cases} 0 & \text{if } \sum_{t=1}^{\infty} \Phi_\Delta(-\log r_\psi(t)) < \infty, \\ 1 & \text{if } \sum_{t=1}^{\infty} \Phi_\Delta(-\log r_\psi(t)) = \infty \end{cases} \\ &= \mu_{R, \Lambda_*}(\{\Lambda \in \Omega_{R, \Lambda_*} : g_t \Lambda \in S_{-\log \delta, -\log r_\psi(t)} \text{ for arbitrarily large } t > 0\}) \\ &= \mu_{R, \Lambda_*}(\{\Lambda \in \Omega_{R, \Lambda_*} : \delta(g_t \Lambda) \leq r_\psi(t) \text{ for arbitrarily large } t > 0\}) \\ &= \mu_{R, \Lambda_*}(\mathcal{A}_R(\psi)). \end{aligned}$$

So, to complete the proof, it suffices to show that the series

$$\sum_{t=1}^{\infty} \Phi_\Delta(-\log r_\psi(t)) \quad (7.4)$$

is asymptotic to (6.4). First of all, by Proposition 8.9, we have

$$(7.4) \asymp_x \begin{cases} \sum_{t=1}^{\infty} r_\psi(t)^{d-1}, & R \sim Q_0, \\ \sum_{t=1}^{\infty} r_\psi(t)^2 (-\log r_\psi(t)), & R \sim Q_0. \end{cases}$$

Let

$$n = \begin{cases} 0, & R \sim Q_0, \\ 1, & R \sim Q_0. \end{cases} \quad (7.5)$$

Then we can write both (6.4) and (7.4) in a uniform manner:

$$\begin{aligned} (6.4) &= \sum_{T \in 2^{\mathbb{N}}} T^{d-1} \log^n \log T \psi^{d-1}(T), \\ (7.4) &\asymp_x \sum_{t=1}^{\infty} r_\psi(t)^{d-1} (-\log r_\psi(t))^n. \end{aligned}$$

Since ψ is regular, each of these series is asymptotic to its corresponding integral, that is,

$$(6.4) \asymp_{+, \times} \int_0^\infty (2^x)^{d-1} \log^n \log(2^x) \psi^{d-1}(2^x) dx,$$

$$(7.4) \asymp_{+, \times} \int_0^\infty r_\psi(t)^{d-1} (-\log r_\psi(t))^n dt.$$

Let $\Psi(T) = T^{d-1} \log^n \log T$. In the following integrals, we omit the finite limit of integration since it is irrelevant for determining whether or not the integral converges. The reader should think of the finite limit of integration as being some arbitrarily large number.

$$\begin{aligned} (6.4) &\asymp_{+, \times} \int_0^\infty (2^x)^{d-1} \log^n \log(2^x) \psi^{d-1}(2^x) dx \\ &\asymp_{\times} \int_0^\infty T^{d-1} \log^n \log T \psi^{d-1}(T) \frac{dT}{T} \\ &= \int_0^\infty R \psi^{d-1}(T) \frac{dR}{T \Psi'(T)} \quad (\text{letting } T = \Psi^{-1}(R)) \\ &\asymp_{\times} \int_0^\infty \psi^{d-1}(\Psi^{-1}(R)) dR. \quad (\text{since } \Psi(T) \asymp_{\times} T \Psi'(T)) \end{aligned}$$

We shall now resort to the following lemma.

Lemma 7.6. *Let $f: [c, \infty) \rightarrow (0, \infty)$ be a strictly decreasing continuous function. Then*

$$\int_c^\infty f(x) dx + cf(c) = \int_0^{f(c)} f^{-1}(x) dx.$$

Proof. The regions whose areas are represented by these integrals are congruent to each other via the map $(x, y) \mapsto (y, x)$. \square

Applying this lemma with $f = \psi^{d-1} \circ \Psi^{-1}$, we continue our calculation:

$$\begin{aligned} (6.4) &\asymp_{+, \times} \int_0^\infty \Psi(\psi^{-1}(U^{\frac{1}{d-1}})) dU \quad (\text{by Lemma 7.6}) \\ &\asymp_{\times} \int_0^\infty \Psi(\psi^{-1}(e^{-t})) e^{-(d-1)t} dt \quad (\text{letting } U = e^{-(d-1)t}) \\ &= \int_0^\infty r_\psi(t)^{d-1} \log^n \log(\psi^{-1}(e^{-t})) dt. \end{aligned}$$

Comparing with (7.4), we see that we have proven Theorem 6.5 in the case $n = 0$, and also for all functions ψ satisfying

$$\log \log \psi^{-1}(e^{-t}) \asymp_{+, \times} -\log r_\psi(t). \quad (7.6)$$

Remark 7.7. For the remainder of the proof, we could require $n = 1$ and thus $d - 1 = 2$ to simplify notation somewhat. However, we prefer to keep the original notation.

For each $c > 0$, let $\psi_{1,c}$ be defined by the equation

$$r_{\psi_{1,c}}(t) = \frac{1}{t^c},$$

i.e.

$$\psi_{1,c}^{-1}(x) = \frac{1}{x(-\log x)^c}.$$

Then

$$\begin{aligned} -\log r_{\psi_{1,c}}(t) &= c \log t \asymp_{\times} \log t, \\ \log \log \psi_{1,c}^{-1}(e^{-t}) &= \log(t + c \log t) \asymp_{+} \log t. \end{aligned}$$

This yields the following.

Claim 7.8. *Fix $c_1 > c_2 > 0$. Then Theorem 6.5 holds for any function $\psi_{1,c_1} \leq \psi \leq \psi_{1,c_2}$.*

Proof. We have $\psi_{1,c_1}^{-1} \leq \psi^{-1} \leq \psi_{1,c_2}^{-1}$ and $r_{\psi_{1,c_1}} \leq r_{\psi} \leq r_{\psi_{1,c_2}}$, and thus

$$\log \log \psi^{-1}(e^{-t}) \asymp_{+} \log t \asymp_{\times} -\log r_{\psi}(t),$$

i.e. (7.6) holds. \square

Remark 7.9. This completes the proof of Theorem 6.5 for the case of most “reasonable” functions ψ , for example if ψ can be written in terms of the elementary operations together with exponents and logs. Such a ψ is always comparable to every function $\psi_{1,c}$ (see [29, Chapter III]). On the other hand, if $c_1 > \frac{1}{d-1} > c_2 > 0$, then (6.4) converges with $\psi = \psi_{1,c_1}$ but diverges with $\psi = \psi_{1,c_2}$. If $\psi \lesssim_{\times} \psi_{1,c_1}$, then $A_R(\psi) \subseteq A_R(C\psi_{1,c_1})$ for some $C > 0$, implying that $\mu_{R,\Lambda_*}(A_R(\psi)) = 0$. Similarly, if $\psi \gtrsim_{\times} \psi_{1,c_2}$, then we have $\mu_{R,\Lambda_*}(A_R(\psi)) = 1$. Finally, if $\psi_{1,c_1} \lesssim \psi \lesssim_{\times} \psi_{1,c_2}$, then Claim 7.8 gives the desired result.

We now proceed to prove the general case of Theorem 6.5, using Claim 7.8. Fix

$$c_1 > \frac{1}{d-1} > c_2 > c_3 > 0.$$

Claim 7.10. *We can without loss of generality assume $\psi \geq \psi_{1,c_1}$.*

Proof. Suppose that the theorem is true for all $\psi \geq \psi_{1,c_1}$, and let ψ be arbitrary. Let $\psi' = \max(\psi, \psi_{1,c_1})$. Note that (6.4) converges for $\psi = \psi'$ if and only if it converges for $\psi = \psi$. Applying the known case of the theorem, we have

$$\mu_{R,\Lambda_*}(A_R(\psi')) = \begin{cases} 0, & (6.4) \text{ converges,} \\ 1, & (6.4) \text{ diverges.} \end{cases}$$

On the other hand, we have

$$A_R(\psi') = A_R(\psi) \cup A_R(\psi_{1,c_1}).$$

Since the latter set has measure zero, the measures of $A_R(\psi')$ and $A_R(\psi)$ are equal. \square

So, from now on, we assume $\psi \geq \psi_{1,c_1}$. If $\psi \leq \psi_{1,c_3}$, then this completes the proof (of Theorem 6.5). So we will assume that $\psi(q) > \psi_{1,c_3}(q)$ for arbitrarily large q .

Claim 7.11. *Fix T_2 for which $\psi(T_2) > \psi_{1,c_3}(T_2)$, and let $T_1 < T_2$ be the largest value for which $\psi(T_1) \leq \psi_{1,c_2}(T_1)$. Then*

$$\begin{aligned} & \int_{T_1}^{T_2} T^{d-1} \log^n \log T \psi_{1,c_2}^{d-1}(T) \frac{dT}{T} \\ & \gtrsim_{\times} \log^{(c_2/c_3)(1-(d-1)c_2)} T_2 - C \log^{1-(d-1)c_2} T_2 \end{aligned} \quad (7.7)$$

for some constant $C > 0$.

Proof. Since $q \mapsto q\psi(q)$ is assumed to be nondecreasing, we have

$$T_1 \psi_{1,c_2}(T_1) \geq T_1 \psi(T_1) \geq T_2 \psi(T_2) > T_2 \psi_{1,c_3}(T_2).$$

On the other hand,

$$\psi_c(q) \asymp_{\times} \frac{1}{q \log^c q}, \quad \text{so} \quad \log^{c_2} T_1 \lesssim_{\times} \log^{c_3} T_2.$$

Now

$$\begin{aligned} & \int_{T_1}^{T_2} T^{d-1} \log^n \log T \psi_{1,c_2}^{d-1}(T) \frac{dT}{T} \\ & \asymp_{\times} \int_{T_1}^{T_2} \frac{\log^n \log T}{\log^{(d-1)c_2} T} \frac{dT}{T} = \int_{\log T_1}^{\log T_2} \frac{\log^n t}{t^{(d-1)c_2}} dt \geq \int_{\log T_1}^{\log T_2} t^{-(d-1)c_2} dt \\ & \asymp_{\times} \log^{1-(d-1)c_2} T_2 - \log^{1-(d-1)c_2} T_1 \\ & \gtrsim_{\times} \log^{(c_2/c_3)(1-(d-1)c_2)} T_1 - C \log^{1-(d-1)c_2} T_1. \end{aligned} \quad \triangleleft$$

Since the right-hand side of (7.7) tends to infinity as $T_2 \rightarrow \infty$, the existence of infinitely large values of T_2 for which the hypotheses of the claim are satisfied implies that

$$\int^{\infty} T^{d-1} \log^n \log T \min(\psi(T), \psi_{1,c_2}(T))^{d-1} \frac{dT}{T} = \infty,$$

i.e. (6.4) diverges for $\psi = \min(\psi, \psi_{1,c_2})$. Thus, by Claim 7.8, we have

$$\mu_{R,\Lambda_*}(\mathbf{A}_R(\min(\psi, \psi_{1,c_2}))) = 1.$$

But since $\mathbf{A}_R(\psi) \supseteq \mathbf{A}_R(\min(\psi, \psi_{1,c_2}))$, this completes the proof of Theorem 6.5. ■

8. Estimating the measure μ_{R,Λ_*}

In this section, we estimate $\int \varphi d\mu_{R,\Lambda_*}$ for any function $\varphi: \Omega_{R,\Lambda_*} \rightarrow [0, \infty)$. Our main tools will be the generalized Iwasawa decomposition (Theorem 8.1) and the reduction theory of algebraic groups (Theorem 8.4). We first prove a theorem for general algebraic groups and then specialize to the case $G = \mathrm{O}(R)_0$.

We will need the following notation: if X is a metric space with distance dist_X , φ is a nonnegative continuous function on X , and $C > 0$, we define

$$\varphi^{(C)}(x) := \max_{\text{dist}_X(x',x) \leq C} \varphi(x'), \quad \varphi_{(C)}(x) := \min_{\text{dist}_X(x',x) \leq C} \varphi(x').$$

Let G be a semisimple algebraic group. Let $P \subseteq G$ be a parabolic subgroup, and let $P = MAN$ be a Langlands decomposition of P . Let \mathfrak{g} , \mathfrak{p} , \mathfrak{m} , \mathfrak{a} , and \mathfrak{n} denote the corresponding Lie algebras. Let $K \subseteq G$ be a maximal compact subgroup whose Lie algebra \mathfrak{k} is orthogonal to \mathfrak{a} with respect to the Killing form.

Theorem 8.1 (Generalized Iwasawa decomposition [39, Proposition 8.44]). *Let ρ_P be the modular function of P . Then, given any Haar measures μ_K , μ_M , μ_A , μ_N on K , M , A , N , respectively, the measure μ_G given by*

$$\int_G \Phi d\mu_G := \int_{K \times M \times A \times N} \rho_P(a) \Phi(kman) d(\mu_K \times \mu_M \times \mu_A \times \mu_N)(k, m, a, n),$$

where Φ is a measurable function on G , is a Haar measure on G .

Now suppose that G is \mathbb{Q} -algebraic and that $P \subseteq G$ is a minimal parabolic \mathbb{Q} -subgroup. Let $\Gamma \subseteq G$ be a lattice commensurable to $G_{\mathbb{Z}}$.

Definition 8.2. A set $\mathcal{F} \subseteq G$ is a *coarse fundamental domain* for Γ if

- (I) $\mathcal{F}\Gamma = G$, and
- (II) $\#\{\gamma \in \Gamma : \mathcal{F}\gamma \cap \mathcal{F} \neq \emptyset\} < \infty$.

Consider the set

$$A^+ := \{a \in A : \text{Ad}_a|_{\mathfrak{n}} \text{ is contracting}\}. \quad (8.1)$$

Here Ad_a denotes the adjoint action of a .

Theorem 8.3 (Reduction theory for arithmetic groups, [41, Proposition 2.2] or [45, Theorem 16.9]). *There exist precompact open sets $M_0 \subseteq M$ and $N_0 \subseteq N$ and a finite set $F \subseteq G_{\mathbb{Q}}$ such that*

$$\mathcal{F} := KM_0A^+N_0F \quad (8.2)$$

is a coarse fundamental domain for Γ .

Let dist_G denote a right-invariant Riemannian metric on G . Let $X = G/\Gamma$, and consider the metric $\text{dist}_X(x, x') = \min_{g\Gamma=x, g'\Gamma=x'} \text{dist}_G(g, g')$. We note that dist_X is a Riemannian metric on X . Let μ_X denote the normalized Haar measure on X .

Theorem 8.4. *There exist $C > 0$ and a finite set $F \subseteq G_{\mathbb{Q}}$ such that, for any function $\varphi: X \rightarrow [0, \infty)$, we have*

$$\begin{aligned} \int_{A^+} \rho_P(a) \sum_{f \in F} \varphi_{(C)}(af\Gamma) d\mu_A(a) &\lesssim_X \int \varphi d\mu_X \\ &\lesssim_X \int_{A^+} \rho_P(a) \sum_{f \in F} \varphi^{(C)}(af\Gamma) d\mu_A(a). \end{aligned}$$

Proof. Let $M_0 \subseteq M$, $N_0 \subseteq N$, and $F \subseteq G_{\mathbb{Q}}$ be as in Theorem 8.3, and let \mathcal{F} be given by (8.2). Let $\mathcal{F}_0 = KM_0A^+N_0$ so that $\mathcal{F} = \mathcal{F}_0F$. Then

$$\begin{aligned} \int_{\mathcal{F}_0} \sum_{f \in F} \varphi(gf\Gamma) d\mu_G(g) &\lesssim_{\times} \int_{\mathcal{F}} \varphi(g\Gamma) d\mu_G(g) \quad (\text{since } \#(F) < \infty) \\ &\lesssim_{\times} \int \varphi d\mu_X \quad (\text{by (II) of Definition 8.2}) \\ &\leq \int_{\mathcal{F}} \varphi(g\Gamma) d\mu_G(g) \quad (\text{by (I) of Definition 8.2}) \\ &\leq \int_{\mathcal{F}_0} \sum_{f \in F} \varphi(gf\Gamma) d\mu_G(g) \end{aligned}$$

Let $\Phi(g) = \sum_{f \in F} \varphi(gf\Gamma)$ so that

$$\int \varphi d\mu_X \asymp_{\times} \int_{\mathcal{F}_0} \Phi d\mu_G. \quad (8.3)$$

Now, by Theorem 8.1,

$$\int_{\mathcal{F}_0} \Phi d\mu_G = \int_{K \times M_0 \times A^+ \times N_0} \rho_P(a) \Phi(kman) d(\mu_K \times \mu_M \times \mu_A \times \mu_N)(k, m, a, n). \quad (8.4)$$

Now let

$$C = \max\{\text{dist}_G(\text{id}, km(ana^{-1})) : k \in K, m \in M_0, a \in A^+, n \in N_0\}. \quad (8.5)$$

Since N is contracted by the adjoint action of A^+ , the set $\{ana^{-1} : a \in A^+, n \in N_0\}$ is precompact and thus $C < \infty$. For $k \in K, m \in M_0, a \in A^+$, and $n \in N_0$ fixed, we have

$$\text{dist}_G(a, kman) = \text{dist}_G(a, km(ana^{-1})a) \leq C$$

and thus

$$\Phi(kman) = \Phi(km(ana^{-1})a) \in [\Phi_{(C)}(a), \Phi^{(C)}(a)].$$

Thus, by (8.4),

$$\begin{aligned} &\int_{K \times M_0 \times A^+ \times N_0} \rho_P(a) \Phi_{(C)}(a) d(\mu_K \times \mu_M \times \mu_A \times \mu_N)(k, m, a, n) \\ &\leq \int_{\mathcal{F}_0} \Phi d\mu_G \\ &\leq \int_{K \times M_0 \times A^+ \times N_0} \rho_P(a) \Phi^{(C)}(a) d(\mu_K \times \mu_M \times \mu_A \times \mu_N)(k, m, a, n). \quad (8.6) \end{aligned}$$

Now, since K, M_0 , and N_0 are open and precompact, we have

$$\begin{aligned} &\int_{K \times M_0 \times A^+ \times N_0} \rho_P(a) \Phi^{(C)}(a) d(\mu_K \times \mu_M \times \mu_A \times \mu_N)(k, m, a, n) \\ &\asymp_{\times} \int_{A^+} \rho_P(a) \Phi^{(C)}(a) d\mu_A(a), \quad (8.7) \end{aligned}$$

and similarly for $\Phi_{(C)}$. Combining (8.3), (8.6), and (8.7) completes the proof. \blacksquare

Next we apply Theorem 8.4 to the case where $G = \mathrm{O}(R)_0$ for some quadratic form $R: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$. Suppose that Λ_* is an R -arithmetic lattice commensurable with \mathbb{Z}^{d+1} . Then $X := \Omega_{R, \Lambda_*} \cong G/\Gamma$, where $\Gamma = \mathrm{O}(R; \Lambda_*)$; see (7.2). In view of Proposition 3.11, it is properly embedded into the space Ω_d of all lattices in \mathbb{R}^{d+1} . We are going to consider functions $\varphi: \Omega_{R, \Lambda_*} \rightarrow [0, \infty)$ which are restrictions of functions on Ω_d satisfying an additional property defined below.

Definition 8.5. A function $\varphi: \Omega_d \rightarrow [0, \infty)$ is *monotonic* if

$$\Lambda_1 \subseteq \Lambda_2 \quad \text{implies} \quad \varphi(\Lambda_1) \leq \varphi(\Lambda_2).$$

Theorem 8.6. Let $R: \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ be a $p_{\mathbb{Q}}$ -normalized quadratic form, and suppose that $\Lambda_* \in \Omega_d$ is commensurable with \mathbb{Z}^{d+1} . Let

$$\mathbf{s} = \begin{bmatrix} d-1 \\ d-3 \\ \vdots \\ d+1-2p_{\mathbb{Q}} \end{bmatrix} \in \mathbb{R}^{p_{\mathbb{Q}}}.$$

There exists $C > 0$ such that, for any monotonic function $\varphi: \Omega_d \rightarrow [0, \infty)$, we have

$$\int_{\mathbf{t} \in \mathbf{a}^+} e^{-\mathbf{s} \cdot \mathbf{t}} \varphi^{(C)}(g_{\mathbf{t}} \Lambda_*) d\mathbf{t} \lesssim_{\times} \int_X \varphi d\mu_X \lesssim_{\times} \int_{\mathbf{t} \in \mathbf{a}^+} e^{-\mathbf{s} \cdot \mathbf{t}} \varphi^{(C)}(g_{\mathbf{t}} \Lambda_*) d\mathbf{t}.$$

We remark that, even though we integrate φ over $X = \Omega_{R, \Lambda_*}$, it is assumed to be a function on Ω_d ; in particular, the functions $\varphi^{(C)}$, $\varphi^{(C)}$ are defined with respect to the Riemannian distance on $\Omega_d \cong \mathrm{GL}_{d+1}(\mathbb{R})/\mathrm{GL}_{d+1}(\mathbb{Z})$.

Proof. Let $G = \mathrm{O}(R)_0$, and let $\Gamma = \mathrm{O}(R; \Lambda_*) \cap \mathrm{O}(R)_0$. Then G is a semisimple \mathbb{Q} -algebraic group, and Γ is commensurable with $G_{\mathbb{Z}}$. For $\mathbf{t} \in \mathbb{R}^{p_{\mathbb{Q}}}$, let $\Phi(\mathbf{t}) = g_{\mathbf{t}}$ be as in (3.4) so that $\Phi: \mathbb{R}^{p_{\mathbb{Q}}} \rightarrow G$ is a homomorphism. Let $A = \Phi(\mathbb{R}^{p_{\mathbb{Q}}})$. Then the Lie algebra \mathfrak{a} of A is isomorphic to $\mathbb{R}^{p_{\mathbb{Q}}}$ via the map $\Phi'(\mathbf{0})$. In our notation, we will not distinguish between \mathfrak{a} and $\mathbb{R}^{p_{\mathbb{Q}}}$.

Let $\mathfrak{a}^+ = \{\mathbf{t} \in \mathbb{R}^{p_{\mathbb{Q}}}: t_0 > t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0\} \subseteq \mathfrak{a}$, and let $A^+ = \exp(\mathfrak{a}^+)$. Then A is a maximal \mathbb{Q} -split torus, and A^+ is as in (8.1). Fix $a \in A^+$, and let $N \subseteq G$ and $P \subseteq G$ be the groups

$$N := \{g \in G: a^n g a^{-n} \rightarrow 0\},$$

$$P := \{g \in G: (a^n g a^{-n})_1^\infty \text{ is bounded}\},$$

i.e. N is the group of elements contracted by A^+ , and P is the group of elements stabilized by A^+ . Then P is a minimal parabolic \mathbb{Q} -subgroup of G whose Langlands decomposition is $P = MAN$ for some reductive group $M \subseteq P$. Moreover, A^+ is given by formula (8.1). So, by Theorem 8.4, there exist $C > 0$ and a finite set $F \subseteq G_{\mathbb{Q}}$ such that, for any $\varphi: \Omega_{R, \Lambda_*} \rightarrow [0, \infty)$, we have

$$\begin{aligned} \int_{\mathbf{t} \in \mathbf{a}^+} \rho_P(g_{\mathbf{t}}) \sum_{f \in F} \varphi^{(C)}(g_{\mathbf{t}} f \Lambda_*) d\mathbf{t} &\lesssim_{\times} \int_X \varphi d\mu_X \\ &\lesssim_{\times} \int_{\mathbf{t} \in \mathbf{a}^+} \rho_P(g_{\mathbf{t}}) \sum_{f \in F} \varphi^{(C)}(g_{\mathbf{t}} f \Lambda_*) d\mathbf{t}. \end{aligned} \quad (8.8)$$

Here we remark that formally Theorem 8.4 produces (8.8) with $\varphi_{(C)}$, $\varphi^{(C)}$ replaced by $(\varphi_X)_{(C)}$, $(\varphi_X)^{(C)}$, respectively, where the latter are defined with respect to the Riemannian distance on X . But since we clearly have $\varphi_{(C)} \leq (\varphi_X)_{(C)}$ and $\varphi^{(C)} \geq (\varphi_X)^{(C)}$, (8.8) follows.

Claim 8.7. *For some $C' > 0$,*

$$\sum_{f \in F} \varphi^{(C)}(g_t f \Lambda_*) \lesssim_{\times} \varphi^{(C')}(g_t \Lambda_*). \quad (8.9)$$

Proof. For $f \in F \subseteq G_{\mathbb{Q}}$ fixed, $f \Lambda_*$ is commensurable with Λ_* , and thus

$$\frac{1}{N_f} \Lambda_* \subseteq f \Lambda_* \subseteq N_f \Lambda_* \quad \text{for some } N_f \in \mathbb{N}.$$

In particular, since φ is monotonic,

$$\varphi^{(C)}(g_t f \Lambda_*) \leq \varphi^{(C)}(g_t N_f \Lambda_*) = \varphi^{(C)}(N_f g_t \Lambda_*) \leq \varphi^{(C + \log N_f)}(g_t \Lambda_*),$$

where the last inequality follows since the distance on Ω_d is defined via a Riemannian metric on $\mathrm{GL}_{d+1}(\mathbb{R})$. Thus (8.9) holds with $C' = C + \log \max_{f \in F} N_f$. \triangleleft

A similar argument shows that

$$\sum_{f \in F} \varphi_{(C)}(g_t f \Lambda_*) \gtrsim_{\times} \varphi_{(C')}(g_t \Lambda_*).$$

Thus (8.8) becomes

$$\begin{aligned} \int_{t \in \alpha^+} \rho_P(g_t) \varphi_{(C')}(g_t \Lambda_*) dt &\lesssim_{\times} \int \varphi d\mu_{R, \Lambda_*} \\ &\lesssim_{\times} \int_{t \in \alpha^+} \rho_P(g_t) \varphi^{(C')}(g_t \Lambda_*) dt. \end{aligned}$$

Claim 8.8. $\rho_P(g_t) = e^{-s \cdot t}$. (Here and hereafter, $s \cdot t$ denotes $\sum_{i=1}^{p_{\mathbb{Q}}-1} s_i t_i$.)

Proof. It is well known (see e.g. [39, (8.38)]¹⁸) that $\rho_P(g_t) = e^{-\rho(t)}$, where ρ is the sum of the positive roots of A , counting multiplicity.

So, to demonstrate the claim, we must show that $\rho(t) = s \cdot t$. One verifies that the positive roots of A are of the form

$$\begin{aligned} \lambda_{i,j,\pm} &:= \mathbf{e}_i^* \pm \mathbf{e}_j^*, \quad i < j < p_{\mathbb{Q}}, \\ \lambda_i &:= \mathbf{e}_i^*, \quad i < p_{\mathbb{Q}}, \end{aligned}$$

¹⁸The sign difference between [39, (8.38)] and the present formula is due to Knapp's convention of assuming that \mathfrak{n} is the union of the positive root spaces, while we assume that \mathfrak{n} is the union of the negative root spaces (cf. (8.1)).

with corresponding root spaces

$$\begin{aligned}\mathfrak{g}_{\lambda_{i,j,-}} &= \mathbb{R}(\mathbf{e}_j \cdot \mathbf{e}_i^* - \mathbf{e}_{d-i} \cdot \mathbf{e}_{d-j}^*), \\ \mathfrak{g}_{\lambda_{i,j,+}} &= \mathbb{R}(\mathbf{e}_{d-j} \cdot \mathbf{e}_i^* - \mathbf{e}_{d-i} \cdot \mathbf{e}_j^*), \\ \mathfrak{g}_{\lambda_i} &= \{\mathbf{x} \cdot \mathbf{e}_i^* - \mathbf{e}_{d-i} \cdot 2B\tilde{R}(\mathbf{x}, \cdot) : (x_{p_{\mathbb{Q}}}, \dots, x_{d-p_{\mathbb{Q}}}) \in \mathbb{R}^{d+1-2p_{\mathbb{Q}}}\}.\end{aligned}$$

In particular, the multiplicity of the root $\lambda_{i,j,\pm}$ is 1, and the multiplicity of the root λ_i is $(d+1-2p_{\mathbb{Q}})$. Thus

$$\begin{aligned}\rho &= \sum_{j=1}^{p_{\mathbb{Q}}-1} \sum_{i=0}^{j-1} [(\mathbf{e}_i^* + \mathbf{e}_j^*) + (\mathbf{e}_i^* - \mathbf{e}_j^*)] + \sum_{i=0}^{p_{\mathbb{Q}}-1} (d+1-2p_{\mathbb{Q}})\mathbf{e}_i^* \\ &= \sum_{i=0}^{p_{\mathbb{Q}}-1} [2(p_{\mathbb{Q}}-i-1) + (d+1-2p_{\mathbb{Q}})]\mathbf{e}_i^* = \sum_{i=0}^{p_{\mathbb{Q}}-1} [d-2i-1]\mathbf{e}_i^*. \quad \triangleleft\end{aligned}$$

This completes the proof of Theorem 8.6. \blacksquare

Finally, we use Theorem 8.6 to complete the proof of Theorem 6.3. Recall that Δ denotes the function $\Delta = -\log \delta: \Omega_{R,\Lambda_*} \rightarrow \mathbb{R}$ (cf. (7.2)), where δ is defined by (3.7), and that, for $z \in \mathbb{R}$,

$$S_{\Delta,z} = \{\Lambda \in \Omega_{R,\Lambda_*} : \Delta(\Lambda) \geq z\}.$$

Proposition 8.9. *For z large enough,*

$$\Phi_{\Delta}(z) := \mu_{R,\Lambda_*}(S_{\Delta,z}) \asymp_{\times} \begin{cases} e^{-(d-1)z}, & R \not\sim Q_0, \\ e^{-2z}z, & R \sim Q_0. \end{cases}$$

Proof. Clearly, $\delta(\Lambda) = \min_{\mathbf{p} \in \Lambda \setminus \{0\}} \|\mathbf{p}\|$ and $\Delta = -\log \delta$ can be extended to Ω_d using the same definition. For each $z \in \mathbb{R}$, define

$$\varphi_z := 1_{\{\Lambda \in \Omega_d : \Delta(\Lambda) \geq z\}}.$$

Then the restriction of φ_z to Ω_{R,Λ_*} is the characteristic function of $S_{\Delta,z}$ so that

$$\Phi_{\Delta}(z) = \int_{\Omega_{R,\Lambda_*}} \varphi_z \, d\mu_{R,\Lambda_*}.$$

Observe that φ_z is monotonic in the sense of Definition 8.5, with $X = \Omega_d$. Thus, by Theorem 8.6, there exists $C > 0$ independent of z such that

$$\int_{\mathbf{t} \in \alpha^+} e^{-s \cdot \mathbf{t}} (\varphi_z)_{(C)}(g_{\mathbf{t}} \Lambda_*) \, d\mathbf{t} \lesssim_{\times} \Phi_{\Delta}(z) \lesssim_{\times} \int_{\mathbf{t} \in \alpha^+} e^{-s \cdot \mathbf{t}} (\varphi_z)^{(C)}(g_{\mathbf{t}} \Lambda_*) \, d\mathbf{t}.$$

Since Δ is 1-Lipschitz (see (7.3)), we have

$$(\varphi_z)_{(C)} \geq \varphi_{z+C} \quad \text{and} \quad (\varphi_z)^{(C)} \leq \varphi_{z-C},$$

and so

$$f(z+C) \lesssim_{\times} \Phi_{\Delta}(z) \lesssim_{\times} f(z-C),$$

where

$$f(z) := \int_{\mathbf{t} \in \alpha^+} e^{-s \cdot \mathbf{t}} \varphi_z(g_{\mathbf{t}} \Lambda_*) d\mathbf{t}.$$

Thus, to complete the proof, it suffices to show that

$$f(z) \asymp_x \begin{cases} e^{-(d-1)z}, & R \sim Q_0, \\ e^{-2z} z, & R \sim Q_0. \end{cases} \quad (8.10)$$

Indeed, observe that, for $\mathbf{t} \in \alpha^+$, the smallest vector in $g_{\mathbf{t}}(\mathbb{Z}^{d+1})$ is $g_{\mathbf{t}}(\mathbf{e}_0) = e^{-t_0} \mathbf{e}_0$. Thus $\Delta(g_{\mathbf{t}} \mathbb{Z}^{d+1}) = t_0$. On the other hand, since Λ_* is commensurable with \mathbb{Z}^{d+1} , we have $\frac{1}{N} \mathbb{Z}^{d+1} \subseteq \Lambda_* \subseteq N \mathbb{Z}^{d+1}$ for some $N \in \mathbb{N}$, which implies

$$|\Delta(g_{\mathbf{t}} \Lambda_*) - \Delta(g_{\mathbf{t}} \mathbb{Z}^{d+1})| \leq \log N \quad \text{for all } \mathbf{t}.$$

It follows that $\Delta(g_{\mathbf{t}} \Lambda) \asymp_+ t_0$, and so

$$\varphi_z(g_{\mathbf{t}} \mathbb{Z}^{d+1}) \asymp_x \begin{cases} 1, & t_0 \geq z, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore,

$$f(z) \asymp_x \int_{\substack{t_0 > t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0 \\ t_0 > z}} e^{-s \cdot \mathbf{t}} d\mathbf{t}.$$

Claim 8.10. *For $x \geq 1$,*

$$\int_{x > t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0} e^{-s \cdot \mathbf{t}} d\mathbf{t} \asymp_x \begin{cases} 1, & R \sim Q_0, \\ x, & R \sim Q_0. \end{cases}$$

Proof. If $p_{\mathbb{Q}} = 1$, then the domain of integration is zero-dimensional, making the statement trivial. Thus suppose $p_{\mathbb{Q}} \geq 2$. If $d = 3$, then Proposition 3.5 implies that $R \sim Q_0$. So if $R \sim Q_0$, then $d \geq 4$ and in particular $s_1 = d - 3 > 0$. Since $s_i \geq 0$ for all i , we have

$$\begin{aligned} \int_{t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0} e^{-s \cdot \mathbf{t}} d\mathbf{t} &\leq \int_{t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0} e^{-s_1 t_1} d\mathbf{t} \\ &\leq \int_{t_1, \dots, t_{p_{\mathbb{Q}}-1} > 0} e^{-\frac{s_1}{p_{\mathbb{Q}}-1} \sum_{i=1}^{p_{\mathbb{Q}}-1} t_i} d\mathbf{t} < \infty, \end{aligned}$$

demonstrating the upper bound. The lower bound is trivial, so this completes the proof if $R \sim Q_0$.

Now suppose that $R \sim Q_0$. Then $s_1 = 0$, and

$$\int_{x > t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0} e^{-s \cdot \mathbf{t}} d\mathbf{t} = \int_{x > t_1 > 0} 1 dt_1 = x. \quad \triangleleft$$

Let n be given by (7.5) so that

$$\int_{x > t_1 > \dots > t_{p_{\mathbb{Q}}-1} > 0} e^{-s \cdot t} dt \asymp_{\times} x^n.$$

Integrating over $t_0 > z$ gives

$$f(z) \asymp_{\times} \int_{t_0 > z} e^{-s_0 t_0} t_0^n dt_0 \asymp_{\times} e^{-s_0 z} z^n = e^{-(d-1)z} z^n,$$

demonstrating (8.10). \blacksquare

We end this section by proving a lemma which was needed in the proof of Theorem 5.1 (ii), (iii). Recall the definition of codiameter given in Definition 5.4.

Lemma 8.11. *There exists $C_1 > 0$ such that, for every $\Lambda \in \Omega_{R, \Lambda_*}$, there exists a totally isotropic Λ -rational subspace $V \subseteq \mathbb{R}^{d+1}$ of dimension $p_{\mathbb{Q}}$ satisfying*

$$\text{Codiam}(V \cap \Lambda) \leq C.$$

Proof. Let G, Γ, A, A^+, N, P , and M be as in the proof of Theorem 8.6. Let $M_0 \subseteq M$, $N_0 \subseteq N$, and $F \subseteq G_{\mathbb{Q}}$ be as in Theorem 8.3, and let \mathcal{F} be given by (8.2). Then, for every $\Lambda \in \Omega_{R, \Lambda_*}$, we can write $\Lambda = g\Lambda_*$ for some $g \in \mathcal{F}$. Write

$$g = kmanf = km(ana^{-1})af,$$

where $k \in K$, $m \in M_0$, $a \in A^+$, $n \in N_0$, and $f \in F$. Write $h = km(ana^{-1})$ so that

$$\Lambda = haf\Lambda_*.$$

We recall (cf. (8.5)) that $\text{dist}_G(\text{id}, h) \leq C$ for some $C > 0$ independent of Λ .

Let $V_0 = \mathcal{L}_{p_{\mathbb{Q}}}$, and let $V = h(V_0)$. We observe that V_0 is a totally isotropic $af\Lambda_*$ -rational subspace of \mathbb{R}^{d+1} of dimension $p_{\mathbb{Q}}$, and thus V is a totally isotropic Λ -rational subspace of \mathbb{R}^{d+1} of dimension $p_{\mathbb{Q}}$.

Since a is contracting on V_0 , we have $\text{Codiam}(V_0 \cap af\Lambda_*) \leq \text{Codiam}(V_0 \cap f\Lambda_*)$. On the other hand, $\text{Codiam}(V_0 \cap f\Lambda_*) \asymp_{\times} 1$ since f ranges over a finite set. Thus

$$\text{Codiam}(V \cap \Lambda) \leq e^{\text{dist}_G(\text{id}, h)} \text{Codiam}(V_0 \cap af\Lambda_*) \lesssim_{\times} e^C.$$

This completes the proof. \blacksquare

9. The exceptional quadric hypersurface

Recall that the *exceptional quadric hypersurface* is the hypersurface M_{Q_0} defined by the exceptional quadratic form (2.7). This hypersurface occupies an interesting place in the theory of intrinsic Diophantine approximation on quadric hypersurfaces developed in this paper. To begin with, it has “more rational points than expected”. Specifically, according to Theorem 6.1,

$$N_{M_{Q_0}}(T) \asymp_{\times} T^2 \log T, \tag{9.1}$$

rather than $N_{M_Q}(T) \asymp_{\times} T^2$, which holds when Q is a quadratic form on \mathbb{R}^4 which is not equivalent to Q_0 . Nevertheless, these “extra points” do not appear to affect either the

Dirichlet- or Khintchine-type theorems of these manifolds in quite the way one would expect. With regards to the Dirichlet-type theorem, the extra points have no effect at all, and the optimal Dirichlet function for M_Q is always ψ_1 , independent of whether or not $Q \sim Q_0$. On the other hand, the extra points do affect the Khintchine-type theorem, but not as expected: they introduce a factor of $\log \log T$ into the series (6.4), rather than a factor of $\log T$ as a naive application of the Borel–Cantelli lemma would predict.

It is natural to ask whether these extraordinary properties of the exceptional quadric hypersurface are due to special algebraic properties. This turns out to be the case; in this section, we make this special structure explicit and use this explicitness to derive elementary proofs both of (9.1) and of the convergence case of Theorem 6.3 for the manifold M_{Q_0} .

We begin by describing the special algebraic property which leads to the results outlined above: the manifold M_{Q_0} is isomorphic to $\mathbb{P}_{\mathbb{R}}^1 \times \mathbb{P}_{\mathbb{R}}^1$, with the isomorphism given by the Segre embedding $\Phi: \mathbb{P}_{\mathbb{R}}^1 \times \mathbb{P}_{\mathbb{R}}^1 \rightarrow \mathbb{P}_{\mathbb{R}}^3$ defined by the formula $\Phi([\mathbf{x}], [\mathbf{y}]) = [\mathbf{x} \otimes \mathbf{y}]$, or more explicitly,

$$\Phi([(x_0, x_1)], [(y_0, y_1)]) = [(x_0 y_0, x_0 y_1, x_1 y_0, x_1 y_1)].$$

Thus M_{Q_0} has a “product structure”. This explains why the lattice $O(Q_0; \mathbb{Z}) \cap O(Q_0)_0$ factors as $SL_2(\mathbb{Z}) \times SL_2(\mathbb{Z})$; each factor of $SL_2(\mathbb{Z})$ acts on a different copy of $\mathbb{P}_{\mathbb{R}}^1$. Note that the natural metric on M_{Q_0} is compatible with the distance inherited from $\mathbb{P}_{\mathbb{R}}^3$ under the Segre embedding.

We also remark that the product structure of M_{Q_0} is consistent with its Diophantine structure. More precisely, the set of intrinsic rationals $\mathbb{P}_{\mathbb{Q}}^3 \cap M_{Q_0}$ factors as $\mathbb{P}_{\mathbb{Q}}^1 \times \mathbb{P}_{\mathbb{Q}}^1$; moreover, for $[\mathbf{p}], [\mathbf{q}] \in \mathbb{P}_{\mathbb{Q}}^1$,

$$H_{\text{std}}(\Phi([\mathbf{p}], [\mathbf{q}])) = H_{\text{std}}([\mathbf{p}]) \cdot H_{\text{std}}([\mathbf{q}]). \quad (9.2)$$

Remark 9.1. According to formula (9.2), the Diophantine triple

$$(\iota_3^{-1}(M_{Q_0}), \mathbb{Q}^3 \cap \iota_3^{-1}(M_{Q_0}), H_{\text{std}})$$

is locally isomorphic to the Diophantine triple $(\mathbb{R}^2, \mathbb{Q}^2, H_{\text{prod}})$ considered in [22]. For example, applying the affine corollary of Theorem 5.1 to the hypersurface M_{Q_0} yields an alternate proof of the case $\Theta = \text{prod}$, $d = 2$ of [22, Theorem 1.2].

We are now ready to begin proving statements about the manifold M_{Q_0} by using the decomposition $M_{Q_0} \equiv \mathbb{P}_{\mathbb{R}}^1 \times \mathbb{P}_{\mathbb{R}}^1$. We begin by computing the number of rationals up to a given height.

An elementary proof of (9.1). It is well known that

$$\#\{[\mathbf{p}] \in \mathbb{P}_{\mathbb{Q}}^1 : T/2 < H_{\text{std}}([\mathbf{p}]) \leq T\} \asymp_{\times} \#\{[\mathbf{p}] \in \mathbb{P}_{\mathbb{Q}}^1 : H_{\text{std}}([\mathbf{p}]) \leq T\} \asymp_{\times} T^2. \quad (9.3)$$

Now, by (9.2),

$$N_{M_Q}(2^N) = \#\{([\mathbf{p}], [\mathbf{q}]) \in (\mathbb{P}_{\mathbb{Q}}^1)^2 : H_{\text{std}}([\mathbf{p}]) \cdot H_{\text{std}}([\mathbf{q}]) \leq 2^N\}$$

$$\begin{aligned}
&= \sum_{n=0}^N \# \left\{ ([\mathbf{p}], [\mathbf{q}]) \in (\mathbb{P}_{\mathbb{Q}}^1)^2 : 2^{n-1} < H_{\text{std}}([\mathbf{p}]) \leq 2^n, \right. \\
&\quad \left. H_{\text{std}}([\mathbf{q}]) \leq \frac{2^N}{H_{\text{std}}([\mathbf{p}])} \right\} \\
&\asymp \sum_{n=0}^N \sum_{\substack{[\mathbf{p}] \in \mathbb{P}_{\mathbb{Q}}^1 \\ 2^{n-1} < H_{\text{std}}([\mathbf{p}]) \leq 2^n}} \left(\frac{2^N}{H_{\text{std}}([\mathbf{p}])} \right)^2 \quad (\text{by (9.3)}) \\
&\asymp \sum_{n=0}^N (2^{N-n})^2 \# \{ [\mathbf{p}] \in \mathbb{P}_{\mathbb{Q}}^1 : 2^{n-1} < H_{\text{std}}([\mathbf{p}]) \leq 2^n \} \\
&\asymp \sum_{n=0}^N (2^N)^2 \quad (\text{by (9.3)}) \\
&= (2^N)^2 (N+1) \asymp (2^N)^2 \log(2^N),
\end{aligned}$$

demonstrating (9.1) in the case $T \in 2^{\mathbb{N}}$. The general case follows from a standard approximation argument. \blacksquare

Next, we give an elementary proof of the convergence case of Theorem 6.3 for the manifold M_{Q_0} . This proof will give insight as to why in this case Theorem 6.3 does not simply state the converse of the (naive) Borel–Cantelli lemma; cf. Remark 9.3.

Remark 9.2. In the following proof, we will assume that ψ is regular, but we do not need to assume that $q \mapsto q\psi(q)$ is nonincreasing, as was assumed in the proof of Theorem 6.3.

Proof of the convergence case of Theorem 6.3 assuming $Q = Q_0$. Let λ denote normalized Lebesgue measure on $\mathbb{P}_{\mathbb{R}}^1$, and note that $\lambda_{M_Q} \asymp \Phi(\lambda \times \lambda)$. Let

$$A_{\psi} = \{([\mathbf{x}], [\mathbf{y}]) \in (\mathbb{P}_{\mathbb{R}}^1)^2 : \text{there exist infinitely many } ([\mathbf{p}], [\mathbf{q}]) \in (\mathbb{P}_{\mathbb{Q}}^1)^2 \text{ such that} \\ \text{dist}([\mathbf{p}], [\mathbf{x}]), \text{dist}([\mathbf{q}], [\mathbf{y}]) \leq \psi(H_{\text{std}}([\mathbf{p}]) \cdot H_{\text{std}}([\mathbf{q}]))\}.$$

Then $A_{M_{Q_0}}(\psi) = \Phi(A_{\psi})$. So, to prove the convergence case of Theorem 6.3, we should show that $\lambda \times \lambda(A_{\psi}) = 0$, assuming that the series

$$\sum_{T \in 2^{\mathbb{N}}} T^2 \log \log T \psi^2(T) \tag{9.4}$$

converges.

For each $n \geq 0$, let

$$\mathcal{Z}_n = \{[\mathbf{p}] \in \mathbb{P}_{\mathbb{Q}}^1 : 2^n \leq H_{\text{std}}([\mathbf{p}]) < 2^{n+1}\}.$$

By (9.3), we have $\#(\mathcal{Z}_n) \asymp (2^n)^2$. Now fix $0 \leq n \leq N$, and let

$$A_{n,N} = B(\mathcal{Z}_n, C\psi(2^N)) \times B(\mathcal{Z}_{N-n}, C\psi(2^N)),$$

where $C > 0$ is a large constant. Since ψ is regular (as assumed in Theorem 6.3), if C is large enough, then

$$A_\psi \subseteq \limsup_{N \rightarrow \infty} \bigcup_{0 \leq n \leq N} A_{n,N},$$

and so, by the Borel–Cantelli lemma, if the series

$$\sum_{N=0}^{\infty} \sum_{n=0}^N (\lambda \times \lambda)(A_{n,N}) \quad (9.5)$$

converges, then $(\lambda \times \lambda)(A_\psi) = 0$. So, to complete the proof, it suffices to show that (9.5) $\lesssim_{\times} (9.4)$.

Fix $0 \leq n \leq N$. We have

$$(\lambda \times \lambda)(A_{n,N}) = \lambda(B(\mathcal{Z}_n, \psi(2^N))) \cdot \lambda(B(\mathcal{Z}_{N-n}, \psi(2^N))).$$

Since $\lambda(B([\mathbf{x}], \rho)) \asymp_{\times} r$ for all $[\mathbf{x}] \in \mathbb{P}_{\mathbb{R}}^1$ and $0 < \rho \leq 1$, subadditivity gives

$$\lambda(B(\mathcal{Z}_n, \psi(2^N))) \lesssim_{\times} \#(\mathcal{Z}_n) \psi(2^N).$$

However, in some cases, it may be better to simply estimate from above by $\lambda(\mathbb{P}_{\mathbb{R}}^1) = 1$:

$$\lambda(B(\mathcal{Z}_n, \psi(2^N))) \leq 1.$$

Similar bounds hold for $\lambda(B(\mathcal{Z}_{N-n}, \psi(2^N)))$. Thus

$$\begin{aligned} (\lambda \times \lambda)(A_{n,N}) &\lesssim_{\times} \min(1, \#(\mathcal{Z}_n) \psi(2^N)) \min(1, \#(\mathcal{Z}_{N-n}) \psi(2^N)) \\ &\asymp_{\times} \min(1, (2^n)^2 \psi(2^N)) \min(1, (2^{N-n})^2 \psi(2^N)) \\ &= \begin{cases} (2^n)^2 \psi(2^N), & n \leq N + \log_2 \sqrt{\psi(2^N)}, \\ (2^{N-n})^2 \psi(2^N), & n \geq -\log_2 \sqrt{\psi(2^N)}, \\ (2^N)^2 \psi^2(2^N) & \text{otherwise.} \end{cases} \end{aligned} \quad (9.6)$$

The case $N + \log_2 \sqrt{\psi(2^N)} \geq n \geq -\log_2 \sqrt{\psi(2^N)}$ cannot occur (for all but finitely many N) since $\psi(2^N)$ is less than $1/2^N$ for all sufficiently large N (otherwise, the series (9.4) would diverge).

Geometrically, note that the first two cases correspond to the bounds on $(\lambda \times \lambda)(A_{n,N})$ which result from covering $A_{n,N}$ by vertical and horizontal rectangles, respectively, while the third case corresponds to covering $A_{n,N}$ by squares.

Now fix N , and vary $0 \leq n \leq N$. We have

$$\begin{aligned} \sum_{n=0}^N (\lambda \times \lambda)(A_{n,N}) &\asymp_{\times} \sum_{n=0}^{\lfloor N/2 \rfloor} (\lambda \times \lambda)(A_{n,N}) \quad (\text{by symmetry}) \\ &\lesssim_{\times} \sum_{n=0}^{\lfloor N + \log_2 \sqrt{\psi(2^N)} \rfloor} (2^n)^2 \psi(2^N) \\ &\quad + \sum_{n=\lfloor N + \log_2 \sqrt{\psi(2^N)} \rfloor + 1}^{\lfloor N/2 \rfloor} (2^N)^2 \psi^2(2^N) \end{aligned}$$

$$\begin{aligned}
& \asymp_{\times} (2^{N+\log_2 \sqrt{\psi(2^N)}})^2 \psi(2^N) \\
& \quad + (2^N)^2 \psi^2(2^N) \left(\frac{N}{2} - (N + \log_2 \sqrt{\psi(2^N)}) \right) \\
& = (2^N)^2 \psi^2(2^N) + (2^N)^2 \psi^2(2^N) \frac{1}{2} \log_2 \left(\frac{1}{2^N \psi(2^N)} \right) \\
& \asymp_{\times} (2^N)^2 \psi^2(2^N) \log \left(\frac{1}{2^N \psi(2^N)} \right).
\end{aligned}$$

Thus, for any function ψ satisfying

$$\log \left(\frac{1}{q \psi(q)} \right) \lesssim_{\times} \log \log q, \quad (9.7)$$

we have (9.5) \lesssim_{\times} (9.4), and thus the conclusion of Theorem 6.3 holds in the convergence case for such ψ .

To complete the proof, fix $\varepsilon > 0$, and let

$$\psi_*(q) = \frac{1}{q \log^{1/2+\varepsilon} q}.$$

Then ψ_* satisfies (9.7); moreover, (9.4) converges at $\psi = \psi_*$. Given any function ψ , let

$$\psi' = \max(\psi_*, \psi).$$

Then, if (9.4) converges at ψ , it also converges at ψ' . Moreover, ψ' satisfies (9.7), so if (9.4) converges at ψ , then $A_{\psi'}$ is a nullset. But since $\psi' \geq \psi$, we have $A_{\psi} \subseteq A_{\psi'}$, so this completes the proof. \blacksquare

Remark 9.3. There are two important points to be made about the above proof. The first point is that the calculation (9.6) indicates what the nontrivial relation is which causes the series (9.4) to differ from (6.3). Indeed, (9.6) shows that if $n \leq N + \log_2 \sqrt{\psi(2^N)}$ or $n \geq -\log_2 \sqrt{\psi(2^N)}$, then we are better off computing $(\lambda \times \lambda)(A_{n,N})$ not by simply adding the measures of the squares

$$B(\cdot, C\psi(2^N)) \times B(\cdot, C\psi(2^N))$$

which define $A_{n,N}$, but by estimating the measure of $A_{n,N}$ in terms of the rectangles

$$B(\cdot, \psi(2^N)) \times \mathbb{P}_{\mathbb{R}}^1 \quad \text{or} \quad \mathbb{P}_{\mathbb{R}}^1 \times B(\cdot, \psi(2^N)),$$

respectively. Inside each rectangle, there are many overlapping squares, and this overlap is what causes the difference in the series.

The second point is that we should not expect there to be a difference in series for the Jarník–Besicovitch theorem if $s < d - 1$. Indeed, the same argument would work up until the point where inequality (9.7) is required. But when $s < d - 1$, then the ψ which we “expect to see” (i.e. those which are near the boundary of convergence/divergence) will satisfy

$$\log \left(\frac{1}{q \psi(q)} \right) \asymp_{\times} \log q$$

rather than (9.7). Thus the “refined argument” for the convergence case produces in this case the same series (6.3).

Acknowledgments. The authors would like to thank Victor Beresnevich, Cornelia Druțu, and Sanju Velani for helpful discussions, and an anonymous referee for useful comments.

Funding. The first-named author was supported in part by the Simons Foundation grant #245708. The second-named author was supported in part by the NSF grants DMS-1101320 and DMS-1600814. The fourth-named author was supported in part by the EPSRC Programme Grant EP/J018260/1.

References

- [1] Bekka, M. B., Mayer, M.: Ergodic Theory and Topological Dynamics of Group Actions on Homogeneous Spaces. London Math. Soc. Lecture Note Ser. 269, Cambridge University Press, Cambridge (2000) Zbl 0961.37001 MR 1781937
- [2] Beresnevich, V., Dickinson, D., Velani, S.: Measure theoretic laws for \limsup sets. Mem. Amer. Math. Soc. **179**, x+91 (2006) Zbl 1129.11031 MR 2184760
- [3] Beresnevich, V., Velani, S.: A mass transference principle and the Duffin–Schaeffer conjecture for Hausdorff measures. Ann. of Math. (2) **164**, 971–992 (2006) Zbl 1148.11033 MR 2259250
- [4] Bernik, V. I., Dodson, M. M.: Metric Diophantine Approximation on Manifolds. Cambridge Tracts in Math. 137, Cambridge University Press, Cambridge (1999) Zbl 0933.11040 MR 1727177
- [5] Besicovitch, A. S.: Sets of fractional dimensions (IV): On rational approximation to real numbers. J. Lond. Math. Soc. **9**, 126–131 (1934) Zbl 0009.05301 MR 1574327
- [6] Borel, A., Harish-Chandra: Arithmetic subgroups of algebraic groups. Ann. of Math. (2) **75**, 485–535 (1962) Zbl 0107.14804 MR 147566
- [7] Borevich, A. I., Shafarevich, I. R.: Number Theory. Pure Appl. Math. 20, Academic Press, New York (1966) MR 0195803
- [8] Bowditch, B. H.: Geometrical finiteness for hyperbolic groups. J. Funct. Anal. **113**, 245–317 (1993) Zbl 0789.57007 MR 1218098
- [9] Broderick, R., Fishman, L., Kleinbock, D., Reich, A., Weiss, B.: The set of badly approximable vectors is strongly C^1 incompressible. Math. Proc. Cambridge Philos. Soc. **153**, 319–339 (2012) Zbl 1316.11064 MR 2981929
- [10] Browning, T. D.: Quantitative Arithmetic of Projective Varieties. Progr. Math. 277, Birkhäuser, Basel (2009) Zbl 1188.14001 MR 2559866
- [11] Budarina, N., Dickinson, D., Levesley, J.: Simultaneous Diophantine approximation on polynomial curves. Mathematika **56**, 77–85 (2010) Zbl 1279.11076 MR 2604984
- [12] Cannon, J. W., Floyd, W. J., Kenyon, R., Parry, W. R.: Hyperbolic geometry. In: Flavors of Geometry, Math. Sci. Res. Inst. Publ. 31, Cambridge University Press, Cambridge, 59–115 (1997) Zbl 0899.51012 MR 1491098
- [13] Cassels, J. W. S.: An Introduction to Diophantine Approximation. Cambridge Tracts Math. Math. Phys. 45, Cambridge University Press, New York (1957) Zbl 0077.04801 MR 0087708

- [14] Dani, S. G.: Divergent trajectories of flows on homogeneous spaces and Diophantine approximation. *J. Reine Angew. Math.* **359**, 55–89 (1985) Zbl 0578.22012 MR 794799
- [15] Davenport, H., Schmidt, W. M.: Approximation to real numbers by quadratic irrationals. *Acta Arith.* **13**, 169–176 (1967/68) Zbl 0155.09503 MR 219476
- [16] Davenport, H., Schmidt, W. M.: A theorem on linear forms. *Acta Arith.* **14**, 209–223 (1967/68) Zbl 0179.07303 MR 225728
- [17] Dickinson, H., Dodson, M. M.: Simultaneous Diophantine approximation on the circle and Hausdorff dimension. *Math. Proc. Cambridge Philos. Soc.* **130**, 515–522 (2001) Zbl 0992.11046 MR 1816807
- [18] Druțu, C.: Diophantine approximation on rational quadrics. *Math. Ann.* **333**, 405–469 (2005) Zbl 1082.11047 MR 2195121
- [19] Elman, R., Karpenko, N., Merkurjev, A.: The Algebraic and Geometric Theory of Quadratic Forms. Amer. Math. Soc. Colloq. Publ. 56, American Mathematical Society, Providence (2008) Zbl 1165.11042 MR 2427530
- [20] Falconer, K.: Fractal Geometry. John Wiley & Sons, Chichester (1990) Zbl 0689.28003 MR 1102677
- [21] Fishman, L., Kleinbock, D., Merrill, K., Simmons, D.: Intrinsic Diophantine approximation on manifolds: General theory. *Trans. Amer. Math. Soc.* **370**, 577–599 (2018) Zbl 1422.11149 MR 3717990
- [22] Fishman, L., Simmons, D.: Unconventional height functions in simultaneous Diophantine approximation. *Monatsh. Math.* **182**, 577–618 (2017) Zbl 1367.11060 MR 3607503
- [23] Fishman, L., Simmons, D., Urbański, M.: Diophantine approximation in Banach spaces. *J. Théor. Nombres Bordeaux* **26**, 363–384 (2014) Zbl 1370.11076 MR 3320484
- [24] Fishman, L., Simmons, D., Urbański, M.: Diophantine approximation and the geometry of limit sets in Gromov hyperbolic metric spaces. *Mem. Amer. Math. Soc.* **254**, v+137 (2018) MR 3826896
- [25] Gallagher, P. X.: Metric simultaneous diophantine approximation. II. *Mathematika* **12**, 123–127 (1965) Zbl 0142.01504 MR 188154
- [26] Ghosh, A., Gorodnik, A., Nevo, A.: Diophantine approximation and automorphic spectrum. *Int. Math. Res. Not. IMRN* **2013**, 5002–5058 (2013) Zbl 1370.11077 MR 3123673
- [27] Ghosh, A., Gorodnik, A., Nevo, A.: Metric Diophantine approximation on homogeneous varieties. *Compos. Math.* **150**, 1435–1456 (2014) Zbl 1309.37005 MR 3252026
- [28] Gorodnik, A., Shah, N. A.: Khinchin’s theorem for approximation by integral points on quadratic varieties. *Math. Ann.* **350**, 357–380 (2011) Zbl 1260.11049 MR 2794914
- [29] Hardy, G. H.: Orders of infinity. The *Infinitärcalcül* of Paul du Bois-Reymond. Cambridge Tracts Math. Math. Phys. 12, Hafner Publishing, New York (1971) MR 0349922
- [30] Heath-Brown, D. R.: A new form of the circle method, and its application to quadratic forms. *J. Reine Angew. Math.* **481**, 149–206 (1996) Zbl 0857.11049 MR 1421949
- [31] Jarník, V.: Zur metrischen Theorie der diophantischen Approximationen, *Prace Mat. Fiz.* **36** (1928), 91–106
- [32] Jarník, V.: Über die simultanen diophantischen Approximationen. *Math. Z.* **33**, 505–543 (1931) MR 1545226
- [33] Khintchine, A.: Zur metrischen Theorie der diophantischen Approximationen. *Math. Z.* **24**, 706–714 (1926) MR 1544787

- [34] Kleinbock, D., Merrill, K.: Rational approximation on spheres. *Israel J. Math.* **209**, 293–322 (2015) Zbl 1332.11070 MR 3430242
- [35] Kleinbock, D. Y., Margulis, G. A.: Bounded orbits of nonquasiunipotent flows on homogeneous spaces. In: Sinai's Moscow Seminar on Dynamical Systems, Amer. Math. Soc. Transl. Ser. 2 171, American Mathematical Society, Providence, 141–172 (1996) Zbl 0843.22027 MR 1359098
- [36] Kleinbock, D. Y., Margulis, G. A.: Flows on homogeneous spaces and Diophantine approximation on manifolds. *Ann. of Math.* (2) **148**, 339–360 (1998) Zbl 0922.11061 MR 1652916
- [37] Kleinbock, D. Y., Margulis, G. A.: Logarithm laws for flows on homogeneous spaces. *Invent. Math.* **138**, 451–494 (1999) Zbl 0934.22016 MR 1719827
- [38] Kleinbock, D., Weiss, B.: Modified Schmidt games and a conjecture of Margulis. *J. Mod. Dyn.* **7**, 429–460 (2013) Zbl 1286.11112 MR 3296561
- [39] Knapp, A. W.: Lie Groups Beyond an Introduction. Progr. Math. 140, Birkhäuser, Boston, 2nd ed. (2002) Zbl 1075.22501 MR 1920389
- [40] Lang, S.: Report on diophantine approximations. *Bull. Soc. Math. France* **93**, 177–192 (1965) Zbl 0135.10802 MR 193064
- [41] Leuzinger, E.: Tits geometry, arithmetic groups, and the proof of a conjecture of Siegel. *J. Lie Theory* **14**, 317–338 (2004) Zbl 1086.53073 MR 2066859
- [42] Mahler, K.: On lattice points in n -dimensional star bodies. I. Existence theorems. *Proc. Roy. Soc. London Ser. A* **187**, 151–187 (1946) Zbl 0060.11710 MR 17753
- [43] Mahler, K.: Some suggestions for further research. *Bull. Austral. Math. Soc.* **29**, 101–108 (1984) Zbl 0517.10001 MR 732177
- [44] Melián, M. V., Pestana, D.: Geodesic excursions into cusps in finite-volume hyperbolic manifolds. *Michigan Math. J.* **40**, 77–93 (1993) Zbl 0793.53052 MR 1214056
- [45] Morris, D. W.: Introduction to Arithmetic Groups. Deductive Press (2015) Zbl 1319.22007 MR 3307755
- [46] Moshchevitin, N.: Über die rationalen Punkte auf der Sphäre. *Monatsh. Math.* **179**, 105–112 (2016) Zbl 1331.11051 MR 3439274
- [47] Perron, O.: Über diophantische Approximationen. *Math. Ann.* **83**, 77–84 (1921) MR 1512000
- [48] Raghunathan, M. S.: Discrete Subgroups of Lie Groups. Ergeb. Math. Grenzgeb. (3) 68, Springer, Heidelberg (1972) Zbl 0254.22005 MR 0507234
- [49] Schmidt, W. M.: Badly approximable systems of linear forms. *J. Number Theory* **1**, 139–154 (1969) Zbl 0172.06401 MR 248090
- [50] Schmutz, E.: Rational points on the unit sphere. *Cent. Eur. J. Math.* **6**, 482–487 (2008) Zbl 1176.11037 MR 2425007
- [51] Stratmann, B.: Diophantine approximation in Kleinian groups. *Math. Proc. Cambridge Philos. Soc.* **116**, 57–78 (1994) Zbl 0809.30036 MR 1274159
- [52] Stratmann, B., Velani, S. L.: The Patterson measure for geometrically finite groups with parabolic elements, new and old. *Proc. Lond. Math. Soc.* (3) **71**, 197–220 (1995) Zbl 0821.58026 MR 1327939