



Hessenberg varieties associated to ad-nilpotent ideals

Caleb Ji & Martha Precup

To cite this article: Caleb Ji & Martha Precup (2021): Hessenberg varieties associated to ad-nilpotent ideals, Communications in Algebra, DOI: [10.1080/00927872.2021.1988629](https://doi.org/10.1080/00927872.2021.1988629)

To link to this article: <https://doi.org/10.1080/00927872.2021.1988629>



Published online: 05 Nov 2021.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)



Hessenberg varieties associated to ad-nilpotent ideals

Caleb Ji^a and Martha Precup^b

^aDepartment of Mathematics, Columbia University in the City of New York, New York, New York, USA;

^bDepartment of Mathematics and Statistics, Washington University in St. Louis, St. Louis, Missouri, USA;

ABSTRACT

We consider Hessenberg varieties in the flag variety of $GL_n(\mathbb{C})$ with the property that the corresponding Hessenberg function defines an ad-nilpotent ideal. Each such Hessenberg variety is contained in a Springer fiber. We extend a theorem of Tymoczko to this setting, showing that these varieties have an affine paving obtained by intersecting with Schubert cells. Our method of proof constructs an affine paving for each Springer fiber that restricts to an affine paving of the Hessenberg variety. We use the combinatorial properties of this paving to prove that Hessenberg varieties of this kind are connected.

ARTICLE HISTORY

Received 3 September 2019

Revised 27 September 2021

Communicated by K. Misra

KEYWORDS

Affine paving; flag variety;
Hessenberg variety;
Springer fiber;
Young diagram

2020 MATHEMATICS SUBJECT

CLASSIFICATION

14M15; 14L35; 05A05

1. Introduction

This article studies topological and combinatorial properties of a certain class of Hessenberg varieties. Hessenberg varieties, as introduced in ref. [3], are subvarieties of the flag variety. They are important examples of varieties whose geometry and topology can be characterized using combinatorial techniques (see, e.g. [1, 10, 11]). The Hessenberg variety $\mathcal{H}\text{ess}(X, h)$ is parametrized by two pieces of data: a matrix $X \in \mathfrak{gl}_n(\mathbb{C})$ and a non-decreasing function $h: [n] \rightarrow [n]$, known as a Hessenberg function.

Most of the existing literature on Hessenberg varieties considers only Hessenberg functions with the property that $h(i) \geq i$ for all i . Tymoczko has shown that the Hessenberg varieties corresponding to such Hessenberg functions have a paving by affines [17]. This article investigates Hessenberg varieties corresponding to Hessenberg functions such that $h(i) < i$ for all i . In this case, the Hessenberg space of the function h is an ad-nilpotent ideal and $\mathcal{H}\text{ess}(X, h)$ is a subvariety of the Springer fiber for X . We construct an affine paving for these varieties and explore additional geometric and combinatorial properties.

The fact that Hessenberg varieties of this kind are paved by affines is not new; Fresse proves this statement for a more general class of Hessenberg varieties in ref. [6]. While the arguments used in that article are broader in scope, they do not compute the dimension of each affine cell in the paving. Our methods are constructive and we obtain combinatorial formulas for the dimension of the cells, recovering Tymoczko's results in this setting. In Section 5 below, we define explicit coordinates for an affine paving of the Springer fiber. We then obtain a paving of the Hessenberg variety $\mathcal{H}\text{ess}(X, h)$ by setting certain coordinates equal to zero; this is recorded in Theorem 5.7. Our arguments are of a similar flavor as those given by Spaltenstein in ref. [16].

We give two applications of our results in [Section 6](#). Recall that the irreducible components of the Springer fibers are in bijection with standard tableaux. This is one of the key conclusions of Springer theory. The Hessenberg varieties we consider here may not be equidimensional, so the cells in the affine paving of maximal dimension are not in bijection with irreducible components. However, [Theorem 6.3](#) below shows that these cells are still indexed by standard tableaux. The second main result of [Section 6](#), namely [Theorem 6.5](#), proves that the Hessenberg varieties we consider are always connected (in the type A case). [Example 6.7](#) shows that this property may not be true for analogous Hessenberg varieties defined using other classical groups.

The constructions in this article are motivated by the goal of better understanding the geometry of the affine paving. Determining the closure relations between cells in the paving and identifying singularities of the irreducible components of $\mathcal{H}\text{ess}(X, h)$ are both interesting open questions. Even in the case of the Springer fiber, the answer to these questions is unknown, although progress has been made in special cases [\[5, 8, 9\]](#). Since the Hessenberg varieties considered here are all subvarieties of a Springer fiber, a thorough study of their geometry has the potential to shed new light on these subjects.

The organization of this article is as follows. In [Section 2](#), we review necessary definitions and prior results. In [Section 3](#) we study the notion of Hessenberg inversions, originally introduced by Tymoczko in [ref. \[17\]](#). We define certain subgroups of matrices crucial to the construction of our paving in [Section 4](#). Our affine paving is defined in [Section 5](#) and we prove our main result, which is [Theorem 5.7](#). Finally, we explore some combinatorial properties of our construction in [Section 6](#).

2. Preliminaries

Let n be a positive integer and $[n]$ denote the set of positive integers $\{1, 2, \dots, n\}$. We work in type A throughout (except for [Example 6.7](#) in [Section 6](#)), where $GL_n(\mathbb{C})$ is the group of invertible $n \times n$ complex matrices and $\mathfrak{gl}_n(\mathbb{C})$ is the Lie algebra of all $n \times n$ complex matrices. Let B be the Borel subgroup of $GL_n(\mathbb{C})$ consisting of upper triangular matrices and U be the subgroup of upper triangular matrices with diagonal entries equal to 1.

The Weyl group of $GL_n(\mathbb{C})$ is S_n , which we identify with the subgroup of permutation matrices in $GL_n(\mathbb{C})$. Given $w \in S_n$, let

$$\text{inv}(w) := \{(i, j) \mid i > j \text{ and } w(i) < w(j)\}$$

denote the set of inversions of w . Note that we adopt the nonstandard notation of listing the larger number in the pair $(i, j) \in \text{inv}(w)$ first; this simplifies our exposition below. The Bruhat length of a permutation $w \in S_n$ is $\ell(w) := |\text{inv}(w)|$.

2.1. Hessenberg varieties

The flag variety is the collection of all full flags in $\mathbb{C}^n$,

$$\text{Flags}(\mathbb{C}^n) := \{V_\bullet = (\{0\} \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq \mathbb{C}^n) \mid \dim_{\mathbb{C}}(V_i) = i \text{ for all } i \in [n]\}.$$

Given a full flag $V_\bullet$, let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis of $\mathbb{C}^n$ such that for each i , $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_i\}$ is a basis for V_i . We denote the flag $V_\bullet$ by $V_\bullet = (\mathbf{v}_1 \mid \mathbf{v}_2 \mid \dots \mid \mathbf{v}_n)$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ be the standard basis of $\mathbb{C}^n$. The standard flag $E_\bullet$ is the full flag $E_\bullet = (\mathbf{e}_1 \mid \mathbf{e}_2 \mid \dots \mid \mathbf{e}_n)$. Every flag $V_\bullet$ is of the form $gE_\bullet$ where $g \in GL_n(\mathbb{C})$ such that $g\mathbf{e}_k = \mathbf{v}_k$ and $gE_\bullet := (g\mathbf{e}_1 \mid g\mathbf{e}_2 \mid \dots \mid g\mathbf{e}_n)$.

Remark 2.1. The flag variety identifies with the homogeneous space $GL_n(\mathbb{C})/B$ via the map $gB \mapsto gE_\bullet$. In this article, we interchange notation for the flag $gE_\bullet$ and coset gB whenever it is convenient.

A Hessenberg variety in $\text{Flags}(\mathbb{C}^n)$ is specified by two pieces of data: a Hessenberg function and an element of $\mathfrak{gl}_n(\mathbb{C})$. A Hessenberg function is a function $h : [n] \rightarrow [n]$ such that $h(i) \leq h(i+1)$ for all $i \in [n-1]$. We frequently write a Hessenberg function by listing its values in sequence, that is, $h = (h(1), h(2), \dots, h(n))$. We now define the main objects of interest in this article.

Definition 2.2. Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $X \in \mathfrak{gl}_n(\mathbb{C})$. The Hessenberg variety associated to h and X is

$$\mathcal{H}\text{ess}(X, h) := \{V_\bullet \in \text{Flags}(\mathbb{C}^n) \mid X(V_i) \subseteq V_{h(i)} \text{ for all } i \in [n]\}.$$

If $V_\bullet = (\mathbf{v}_1 \mid \mathbf{v}_2 \mid \dots \mid \mathbf{v}_n)$ then $V_\bullet \in \mathcal{H}\text{ess}(X, h)$ if and only if $X\mathbf{v}_i \in \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_{h(i)}\}$ for all $i \in [n]$. When $X \in \mathfrak{g}$ is a nilpotent matrix and $h = (0, 1, \dots, n-2, n-1)$, the variety $\mathcal{B}(X, h)$ is the Springer fiber of X , which we denote by $\mathcal{B}^X$.

The following remark indicates that we may choose any matrix within a given conjugacy class for our computations without alternating the geometric invariants of the corresponding Hessenberg variety.

Remark 2.3. Given a fixed Hessenberg function h , we have $\mathcal{H}\text{ess}(X, h) \simeq \mathcal{H}\text{ess}(g^{-1}Xg, h)$ for all $g \in GL_n(\mathbb{C})$.

Most of the existing literature on Hessenberg varieties assumes that the Hessenberg function also satisfies the condition that $h(i) \geq i$. The main reason is that this condition on the Hessenberg function ensures $\mathcal{H}\text{ess}(X, h) \neq \emptyset$ for all $X \in \mathfrak{gl}_n(\mathbb{C})$. One of the main purposes of this article is to explore Hessenberg varieties corresponding to Hessenberg functions with the property that $h(i) < i$ for all i . This is exactly the case in which the corresponding Hessenberg space, defined by:

$$H(h) := \text{span}\{E_{ij} \mid i \leq h(j)\} \subseteq \mathfrak{gl}_n(\mathbb{C}),$$

is an ad-nilpotent ideal (that is, its lower central series is finite). Thus, for the remainder of this manuscript we assume that any Hessenberg function $h : [n] \rightarrow [n]$ satisfies $h(i) < i$ for all $i \in [n]$.

Let $\mathcal{H} := \{h : [n] \rightarrow [n] \mid h(i+1) \geq h(i) \text{ and } h(i) < i\}$ denote the set of all Hessenberg functions satisfying the condition that $h(i) < i$. There is a partial ordering on this set defined by

$$h_1 \preceq h_2 \iff h_1(i) \leq h_2(i) \text{ for all } i$$

for $h_1, h_2 \in \mathcal{H}$. A partial order like this one is studied by Drellich in ref. [4]. It follows directly from the definition that if $h_1 \preceq h_2$ then $\mathcal{H}\text{ess}(X, h_1) \subseteq \mathcal{H}\text{ess}(X, h_2)$ for all $X \in \mathfrak{gl}_n(\mathbb{C})$. Note that our set of Hessenberg functions contains a unique maximal element with respect to $\preceq$, namely the Hessenberg function $h = (0, 1, \dots, n-1)$. Thus $\mathcal{H}\text{ess}(X, h) \subseteq \mathcal{B}^X$ for any nilpotent matrix $X \in \mathfrak{gl}_n(\mathbb{C})$ and $h \in \mathcal{H}$.

2.2. Affine pavings

The first main goal of this manuscript is to demonstrate an affine paving of the Hessenberg variety $\mathcal{H}\text{ess}(X, h)$ obtained by intersecting with the Schubert cells. We do so by first constructing an explicit affine paving of the Springer fiber $\mathcal{B}^X$. We then prove that this paving restricts to a paving of $\mathcal{H}\text{ess}(X, h)$ in a natural way. Note that it is very well known that Springer fibers are paved by affines [5, 16], and Tymoczko's results prove that such a paving can be obtained by intersecting with the Schubert cells [17], so our result in that case is not new.

Definition 2.4. A paving of an algebraic variety Y is a filtration by closed subvarieties

$$Y_0 \subset Y_1 \subset \cdots \subset Y_i \subset \cdots \subset Y_d = Y.$$

A paving is affine if every $Y_i - Y_{i-1}$ is isomorphic to a finite disjoint union of affine spaces; we call these spaces the affine cells of the paving.

An affine paving allows us to compute the Betti numbers of an algebraic variety Y , as shown in ref. [7, 1.9.1, 19.1.11]. In the statement below, $H_c^*(Y)$ denotes cohomology with compact support of the algebraic variety Y .

Lemma 2.5. *Let Y be an algebraic variety with an affine paving, $Y_0 \subset Y_1 \subset \cdots \subset Y_i \subset \cdots \subset Y_d = Y$. Then the nonzero cohomology groups of Y are given by $H_c^{2k}(Y) = \mathbb{Z}^{n_k}$ where n_k denotes the number of affine components of dimension k .*

In this article we apply the lemma for Y a complex projective variety, so $H^*(Y) = H_c^*(Y)$. There is a well-known affine paving of $\text{Flags}(\mathbb{C}^n)$ induced by the **Bruhat decomposition**:

$$\text{Flags}(\mathbb{C}^n) = \bigsqcup_{w \in S_n} C_w \text{ where } C_w = BwE_\bullet.$$

The B -orbit C_w is called the Schubert cell indexed by $w \in S_n$. It is well known that each Schubert cell is isomorphic to the subgroup $U^w := U \cap wU^-w^{-1}$ where U^- is the subgroup of lower triangular matrices with diagonal entries equal to 1. In other words, each flag $bwE_\bullet \in C_w$ can be written uniquely as $bwE_\bullet = uwE_\bullet$ for some $u \in U^w$. Since U^w is a unipotent subgroup we have $U^w \simeq \text{Lie}(U^w)$, where $\text{Lie}(U^w)$ an affine space of dimension $\ell(w)$. Therefore $C_w \simeq \mathbb{C}^{\ell(w)}$. It is well known that $\overline{C_w} = \bigcup_{y \leq w} C_y$, where $\leq$ denotes the Bruhat order on S_n . Thus, the Schubert cells are affine cells for the paving of $\text{Flags}(\mathbb{C}^n)$ defined by

$$\mathcal{B}_0 \subset \mathcal{B}_1 \subset \cdots \subset \mathcal{B}_{\frac{n(n-1)}{2}} = \text{Flags}(\mathbb{C}^n) \text{ where } \mathcal{B}_i := \bigcup_{\substack{w \in S_n \\ \ell(w)=i}} \overline{C_w}.$$

We prove $\mathcal{H}\text{ess}(X, h)$ has an affine paving by considering the intersections $C_w \cap \mathcal{H}\text{ess}(X, h)$.

Remark 2.6. It follows from the discussion in ref. [14, §2.2] that in order to prove $\mathcal{H}\text{ess}(X, h)$ is paved by affines, it suffices to prove $\mathcal{H}\text{ess}(X, h) \cap C_w$ is isomorphic to affine space $\mathbb{C}^d$ with $d \in \mathbb{Z}_{\geq 0}$.

2.3. Factorization

We now describe a method for identifying a portion of any Schubert cell in $\text{Flags}(\mathbb{C}^n)$ with a Schubert cell in the flag variety associated to $GL_{n-1}(\mathbb{C})$, namely $\text{Flags}(\mathbb{C}^{n-1})$. We view $GL_{n-1}(\mathbb{C})$ as a subgroup of $GL_n(\mathbb{C})$ by identifying it with its image under the map:

$$GL_{n-1}(\mathbb{C}) \rightarrow GL_n(\mathbb{C}); a \mapsto \begin{bmatrix} a & 0 \\ 0 & 1 \end{bmatrix} \text{ for all } a \in GL_{n-1}(\mathbb{C}). \quad (2.1)$$

Let U_0 be the unipotent subgroup of $GL_{n-1}(\mathbb{C})$ of upper triangular matrices with diagonal entries equal to 1. We view U_0 as a subgroup of $GL_n(\mathbb{C})$ via $U_0 = \{u \in U \mid u_{in} = 0 \text{ if } i \neq n\}$. Similarly, we identify S_{n-1} with the subgroup $\{\sigma \in S_n \mid \sigma(n) = n\}$ of S_n .

Each permutation $w \in S_n$ can be factorized uniquely as

$$w = \nu y \text{ where } \nu = s_i s_{i+1} \cdots s_{n-2} s_{n-1}, \text{ for } i = w(n) \text{ and } y \in S_{n-1}. \quad (2.2)$$

Here s_j denotes the simple transposition swapping j and $j+1$. Note that ν in the factorization above is called the shortest left coset representative for $w = \nu y$ with respect to the Young subgroup $S_{n-1} = \langle s_1, \dots, s_{n-2} \rangle$, see [2, Proposition 2.4.4]. In one-line notation, we have that ν is the permutation with the property that $\nu(n) = i$ and all remaining values are placed in positions $1, 2, \dots, n-1$ of the one-line notation for ν in increasing order; y is the unique permutation with

the property that $y(n) = n$ and the rest of the entries in the one-line notation for y are in the same relative order as the entries of w .

The factorization given in (2.2) satisfies the condition that $\ell(w) = \ell(v) + \ell(y)$ and:

$$\text{inv}(w) = \text{inv}(y) \sqcup y^{-1}(\text{inv}(v)). \quad (2.3)$$

Example 2.7. Let $w = [3, 4, 1, 2] = s_2 s_3 s_1 s_2 \in S_4$. Then $w(4) = 2$ and we see that

$$w = vy \quad \text{where} \quad v = s_2 s_3 \quad \text{and} \quad y = s_1 s_2.$$

In one-line notation, $v = [1, 3, 4, 2]$ and $y = [2, 3, 1, 4]$. We have that

$$\text{inv}(w) = \{(3, 2), (3, 1), (4, 2), (4, 1)\}$$

where $\text{inv}(y) = \{(3, 2), (3, 1)\}$ and $y^{-1}(\text{inv}(v)) = \{(4, 2), (4, 1)\}$, confirming (2.3).

Recall that $U^w := U \cap wU^{-1}w^{-1}$. In the special case where $v = s_i s_{i+1} \cdots s_{n-2} s_{n-1}$ for some $i \in [n]$ we have $U^v = \{u \in U \mid u_{kj} = 0 \text{ for all } k \neq i, k < j\}$, that is, U^v is the i -th row of U , which we denote by U_i . The next lemma tells us there is a factorization of the elements of U^w that is compatible with the factorization of permutations given in (2.2) above. This is a special case of [12, Proposition 28.1].

Lemma 2.8. Suppose $w \in S_n$ and let $w = vy$ be the factorization given in (2.2) with $i = w(n) = v(n)$. For each $u \in U^w$ the product uw can be written uniquely as

$$uw = u_i v u_0 y \quad \text{for some } u_i \in U_i = U^v \text{ and } u_0 \in U^y.$$

Lemma 2.8 gives us an inductive decomposition of each Schubert cell, as we now explain. Let $w \in S_n$ and $w = vy$ be the factorization from (2.2). Given $uwE_\bullet$ with $u \in U$, by **Lemma 2.8** we may write $uwE_\bullet = u_i v u_0 y E_\bullet$ with $u_i \in U_i$ and $u_0 \in U_0$. (Note that since $y(n) = n$ we have $U^y \subseteq U_0$.) We obtain an isomorphism,

$$\rho : C_w \rightarrow \mathbb{C}^{\ell(v)} \times C_y, \quad uwE_\bullet = u_i v u_0 y E_\bullet \mapsto (u_i, u_0 y E'_\bullet) \quad (2.4)$$

where $E'_\bullet := (\mathbf{e}_1 \mid \mathbf{e}_2 \mid \dots \mid \mathbf{e}_{n-1})$ is the standard flag in $\mathbb{C}^{n-1} = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n-1}\}$ and we identify U_i with affine space via $u_i \mapsto (u_{ij})_{j>i}$, where $u_i = I_n + \sum_{j>i} u_{ij} E_{ij}$. We make the identification $U_i \simeq \mathbb{C}^{\ell(v)}$ implicitly throughout this article. Let

$$\rho_1 : C_w \rightarrow \mathbb{C}^{\ell(v)}, \quad uwE_\bullet = u_i v u_0 E_\bullet \mapsto u_i \quad (2.5)$$

be the map obtained from ρ via composition with projection to the first factor. Denote by P the maximal standard parabolic subgroup with Levi subgroup equal to the image of (2.1). That is, $P = \{g \in G \mid g_{nj} = 0 \text{ for all } j < n\}$. The map ρ_1 can be identified with the restriction to C_w of the canonical projection $G/B \rightarrow G/P$; in the notation of cosets we have $\rho_1(uwB) = uwP = u_i vP$. Thus ρ_1 is a morphism of varieties and commutes with the action of B .

3. Hessenberg inversions

Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ be a weak composition of n and $\text{Par}(\lambda)$ be the partition we obtain from λ by rearranging the parts of λ in decreasing order. We begin by fixing an element X_λ in the conjugacy class $\mathcal{O}_{\text{Par}(\lambda)}$ of nilpotent matrices of Jordan type λ .

Definition 3.1. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ be a weak composition of n drawn as a diagram, namely with k rows of boxes so that the i th row from the top has λ_i boxes. The base filling of λ is obtained as follows. Fill the boxes of λ with integers 1 to n starting at the bottom of the leftmost column and moving up the column by increments of one. Then move to the lowest box of the

next column and so on. Denote the base filling of λ by $R(e)$. We now define:

$$X_\lambda := \sum_{(\ell, r)} E_{\ell r} \quad (3.1)$$

where the sum is taken over the set of all pairs (ℓ, r) such that r labels the box directly to the right of ℓ in the base filling of λ .

Example 3.2. If $n=7$ and $\lambda = (2, 3, 1, 1)$ then the base filling of λ is

4	6	
3	5	7
2		
1		

and we have $X_{(2, 3, 1, 1)} =$

0	0	0	0	0	0	0
0	0	0	0	1	0	0
0	0	0	0	0	1	0
0	0	0	0	0	0	1
0	0	0	0	0	0	0
0	0	0	0	0	0	0

For each $w \in S_n$, let $R(w)$ denote the tableau of composition shape λ obtained by labeling the i -th box in the base filling of λ by $w^{-1}(i)$. We say that $R(w)$ is h -strict if $\ell \leq h(r)$ whenever ℓ labels a box directly to the left of r in $R(w)$. Let $\text{RS}_h(\lambda)$ denote the set of all h -strict tableaux of composition shape λ . The set of h -strict tableaux determines which Schubert cells intersect the Hessenberg variety. This is proved by Tymoczko in ref. [17, Theorem 7.1] for Hessenberg varieties associated to Hessenberg functions such that $h(i) \geq i$ for all i . The proof below is the same; we give a sketch using our notation for the reader's convenience.

Lemma 3.3. *Let $w \in S_n$ and $h \in \mathcal{H}$. Then $C_w \cap \mathcal{H}\text{ess}(X_\lambda, h) \neq \emptyset$ if and only if $R(w) \in \text{RS}_h(\lambda)$.*

Sketch of proof. By definition, $wE_\bullet \in \mathcal{H}\text{ess}(X_\lambda, h)$ if and only if

$$X_\lambda \mathbf{e}_{w(r)} \in \text{span}\{\mathbf{e}_{w(1)}, \dots, \mathbf{e}_{w(h(r))}\}$$

for all $r \in [n]$. Suppose ℓ labels the box directly to the left of r in $R(w)$. The $w(\ell)$ labels the box directly to the left of $w(r)$ in $R(e)$. Since $X_\lambda \mathbf{e}_{w(r)} = \mathbf{e}_{w(\ell)}$ we therefore have $wE_\bullet \in \mathcal{H}\text{ess}(X_\lambda, h)$ if and only if $\ell \leq h(r)$ for any such pair (ℓ, r) .

To complete the proof we have only to show that $\mathcal{H}\text{ess}(X_\lambda, h) \cap C_w \neq \emptyset$ implies $wE_\bullet \in \mathcal{H}\text{ess}(X_\lambda, h)$. Assume $u w E_\bullet \in \mathcal{H}\text{ess}(X_\lambda, h)$ for some $u \in U^w$. Then

$$X_\lambda u \mathbf{e}_{w(r)} \in \text{span}\{u \mathbf{e}_{w(1)}, \dots, u \mathbf{e}_{w(h(r))}\} \iff (u^{-1} X_\lambda u) \mathbf{e}_{w(r)} \in \text{span}\{\mathbf{e}_{w(1)}, \dots, \mathbf{e}_{w(h(r))}\}$$

for all $r \in [n]$. The desired statement now follows immediately from the fact that the pivots of $u^{-1} X_\lambda u$ are in the same position as the pivots of X_λ (as proved by Tymoczko in ref. [17, Proposition 4.6]). $\square$

When $h = (0, 1, \dots, n-1)$ we have that $\text{RS}_h(\lambda) =: \text{RS}(\lambda)$ is the set of tableaux of composition shape λ which are row-strict, that is, increasing across rows. By definition, $\text{RS}_h(\lambda) \subseteq \text{RS}(\lambda)$ for all Hessenberg functions $h \in \mathcal{H}$. Our next definition comes from [17, Theorem 7.1], see [15] also.

Definition 3.4 . Let λ be a weak composition of n and $k, \ell \in [n]$. We say (k, ℓ) is a Hessenberg inversion of $R(w)$ for $w \in S_n$ if $k > \ell$ and:

- (1) k occurs in a box below ℓ and in the same column or in any column strictly to the left of the column containing ℓ in $R(w)$, and
- (2) if the box directly to the right of ℓ in $R(w)$ is labeled by r , then $k \leq h(r)$.

Denote the set of Hessenberg inversions in $R(w)$ by $\text{inv}_{h, \lambda}(w)$.

Note that if the pair (k, ℓ) satisfies condition (1), then $(k, \ell) \in \text{inv}(w)$; so Hessenberg inversions are a subset of the inversions of w .

Remark 3.5. If $(k, \ell_1), (k, \ell_2) \in \text{inv}_{h, \lambda}(w)$ then k, ℓ_1 , and ℓ_2 are all in different rows of $R(w)$, or equivalently, $w(k)$, $w(\ell_1)$, and $nw(\ell_2)$ are all in different rows of $R(e)$. Indeed if ℓ_1 fills a box to the left of ℓ_2 and in the same row, the assumption that (k, ℓ_1) is a Hessenberg inversion implies that k is less than every entry to the right of ℓ_1 , implying (k, ℓ_2) cannot be an inversion.

Now suppose ℓ_1 and k occur in the same row. Since (k, ℓ_1) is a Hessenberg inversion, k must occur to the left of ℓ_1 . On the other hand, $R(w)$ must be row-strict and $k > \ell_1$ so we obtain a contradiction.

Example 3.6. Let $n=7$ and $\lambda = (2, 3, 1, 1)$. The tableau $R(w)$ for $w = [4, 3, 1, 6, 5, 7, 2]$ is shown below.

1	4	
2	5	6
7		
3		

If $h = (0, 1, 2, 3, 4, 5, 6)$ then $R(w)$ has inversion set $\text{inv}_{\lambda, h}(w) = \{(7, 6), (7, 4), (5, 4), (3, 2), (3, 1), (2, 1)\}$. Note that $(7, 5) \in \text{inv}(w)$ but $(7, 5)$ is not a Hessenberg inversion since $7 \not\leq 5 = h(6)$. If $h = (0, 0, 1, 2, 3, 3, 3)$ then the inversion set becomes $\text{inv}_{\lambda, h}(w) = \{(7, 6), (7, 4), (5, 4), (3, 2), (2, 1)\}$ since $3 \not\leq 2 = h(4)$ now.

When $h = (0, 1, 2, \dots, n-1)$, we called the pairs in Definition 3.4 above Springer inversions, denoted $\text{inv}_{\lambda}(w)$ in this case. We let

$$\text{inv}_{\lambda}^k(w) := \{(k, \ell) \mid 1 \leq \ell < k \text{ and } (k, \ell) \in \text{inv}_{\lambda}(w)\}$$

so $\text{inv}_{\lambda}(w) = \sqcup_{k=2}^n \text{inv}_{\lambda}^k(w)$. We set $d_k := |\text{inv}_{\lambda}^k(w)|$ for all $2 \leq k \leq n$.

Let $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$. Since $R(w)$ is row-strict, the box labeled by n must appear at the end of a row. Let λ' be the composition of $n-1$ we obtain from λ by deleting the box labeled by n in $R(w)$, or equivalently, deleting the box labeled by $i = w(n)$ in $R(e)$. Our next lemma shows that the Hessenberg inversions of w are well-behaved with respect to the decomposition of $\text{inv}(w)$ given in (2.3).

Lemma 3.7. Suppose $R(w) \in \text{RS}(\lambda)$ and $w = vy$ is the factorization from (2.2) with $i = w(n)$. Then $R(y) \in \text{RS}(\lambda')$ and

$$\text{inv}_{\lambda'}(y) = \text{inv}_{\lambda}(w) \setminus \text{inv}_{\lambda}^n(w).$$

where λ' is the composition of $n-1$ we obtain from λ by deleting the box labeled by n in $R(w)$.

Proof. Recall that v is the permutation whose one-line notation has i in the n -th position and all remaining entries are placed in positions $1, 2, \dots, n-1$ in increasing order. In particular, we obtain the base filling of the composition λ' from the base filling of λ by deleting the box containing i and applying v^{-1} to the remaining entries. It follows immediately that $R(y)$ is the tableau of composition shape λ' we obtain by deleting the box containing n from $R(w)$ so $R(y) \in \text{RS}(\lambda')$. Thus $\text{inv}_{\lambda'}(y) = \sqcup_{k=2}^{n-1} \text{inv}_{\lambda}^k(w)$ as desired. $\square$

Motivated by the inductive formula from Lemma 3.7, we let $X_{\lambda'} \in \mathfrak{gl}_{n-1}(\mathbb{C})$ be the nilpotent matrix defined as in (3.1) for the composition λ' of $n-1$. The proof of the lemma implies $X_{\lambda'}$ is the matrix corresponding to the linear transformation obtained by restricting $v^{-1}X_{\lambda}v$ to $\mathbb{C}^{n-1} \simeq \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n-1}\}$.

Remark 3.8. Let $\mathcal{B}^{X_{\lambda'}}$ denote the Spring fiber in $\mathcal{F}\text{lags}(\mathbb{C}^{n-1})$ corresponding to $X_{\lambda'} \in \mathfrak{gl}_{n-1}(\mathbb{C})$. The discussion above implies that for all $u_0 \in U_0$, $\nu u_0 y E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$ if and only if $u_0 y E'_{\bullet} \in \mathcal{B}^{X_{\lambda'}}$.

4. The $B_k(w)$ -subgroups

We now introduce a collection of subgroups of U associated to each $w \in S_n$ with $R(w) \in \text{RS}(\lambda)$. We use these subgroups in the next section to construct an affine paving of the Springer fiber $\mathcal{B}^X$ that restricts to a paving of the subvariety $\mathcal{H}\text{ess}(X, h)$. Throughout this section, let λ be a fixed weak composition of n and $X_{\lambda} \in \mathcal{O}_{\text{Par}(\lambda)}$ the matrix from Definition 3.1 above.

Suppose $w E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$, or equivalently by Lemma 3.3, that $R(w) \in \text{RS}(\lambda)$. Using the factorization from (2.2) we write $w = \nu \gamma$ for $\nu = s_i s_{i+1} \cdots s_{n-2} s_{n-1}$ where $i = w(n) = \nu(n)$. Recall the morphism $\rho_1 : C_w \rightarrow \mathbb{C}^{\ell(\nu)}$ defined in (2.5). The next lemma tells us that if $uw E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$, then certain entries of $u_i = \rho_1(uw E_{\bullet})$ must be zero.

Lemma 4.1. *Suppose $uw E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$ and $\rho_1(uw E_{\bullet}) = u_i \in U_i$ for $i = w(n)$. Let u_{ij} for $j > i$ denote the entry in the i -th row and j -th column of u_i . Then $u_{ij} = 0$ unless j appears at the end of a row in the base filling $R(e)$. In particular, the morphism of varieties $\rho_1 : C_w \rightarrow \mathbb{C}^{\ell(\nu)}$ defined in (2.5) restricts to a morphism*

$$\rho_1^{(\lambda)} : C_w \cap \mathcal{B}^{X_{\lambda}} \rightarrow \mathbb{C}^{d_n} \quad (4.1)$$

where $d_n := |\text{inv}_{\lambda}^n(w)|$.

Proof. By Lemma 2.8 we may write $uw E_{\bullet} = u_i \nu u_0 y E_{\bullet}$ for some $u_0 \in U_0$ and $u_i = \rho_1(uw E_{\bullet})$. Suppose $j > i$ does not fill a box at the end of a row in the base filling $R(e)$ of λ . This implies $\mathbf{e}_j \in \text{im}(X_{\lambda})$. Let $V_{\bullet} = (\mathbf{v}_1 | \mathbf{v}_2 | \cdots | \mathbf{v}_n)$ where $\mathbf{v}_k = uw(\mathbf{e}_k)$. Since $V_{\bullet} \in \mathcal{B}^{X_{\lambda}}$ we must have $\text{im}(X_{\lambda}) \subseteq V_{n-1}$ and so $\mathbf{e}_j \in \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_{n-1}\}$. Thus $\mathbf{e}_j = \sum_{k=1}^{n-1} c_k \mathbf{v}_k$ for some $c_1, \dots, c_{n-1} \in \mathbb{C}$. Applying $(u_i \nu)^{-1}$ to both sides we obtain

$$\begin{aligned} \nu^{-1}(\mathbf{e}_j - u_{ij} \mathbf{e}_i) &= \sum_{k=1}^{n-1} c_k u_0 y(\mathbf{e}_k) \Rightarrow \mathbf{e}_{\nu^{-1}(j)} - u_{ij} \mathbf{e}_n = \sum_{k=1}^{n-1} c_k u_0 \mathbf{e}_{y(k)} \\ &\Rightarrow -u_{ij} \mathbf{e}_n = -\mathbf{e}_{\nu^{-1}(j)} + \sum_{k=1}^{n-1} c_k u_0 \mathbf{e}_{y(k)} \end{aligned}$$

The RHS of the above equation is in $\text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_{n-1}\}$, implying $u_{ij} = 0$ as desired. Finally, we note that $j > i$ and j appears at the end of a row in the base filling $R(e)$ if and only if $(w^{-1}(i), w^{-1}(j)) = (n, w^{-1}(j))$ is a Springer inversion of $R(w)$ in $\text{inv}_{\lambda}^n(w)$. This shows

$$\text{inv}_{\lambda}^n(w) = \{(n, w^{-1}(j)) \mid i < j \text{ and } j \text{ labels a box at the end of a row in } R(e)\}$$

Thus if $uw E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$ we get

$$\rho_1(uw B) = u_i = I_n + \sum_{\substack{i < j \\ (n, w^{-1}(j)) \in \text{inv}_{\lambda}^n(w)}} u_{ij} E_{ij}.$$

This yields the description of the restriction of ρ_1 to $C_w \cap \mathcal{B}^{X_{\lambda}}$ in (4.1). □

The goal of the next section is to construct a generic element of $C_w \cap \mathcal{B}^{X_{\lambda}}$ whenever this intersection is nonempty. We do so by introducing a collection of subgroups of U associated to each $R(w) \in \text{RS}(\lambda)$. Recall that $\text{inv}_{\lambda}(w)$ denotes the set of Hessenberg inversions corresponding to $h = (0, 1, 2, \dots, n-1)$, namely the Springer inversions.

Definition 4.2. Let $2 \leq k \leq n$. We define $B_k(w)$ to be the set of all matrices g_k such that:

(1) if $(k, \ell) \in \text{inv}_\lambda^k(w)$ and $\mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(i)}$ for some $m \in \mathbb{Z}_{\geq 0}$ then

$$g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)} + x_{w(k)w(\ell)} X_\lambda^m \mathbf{e}_{w(k)} \text{ for some } x_{w(k)w(\ell)} \in \mathbb{C},$$

(2) and $g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$ otherwise.

From the definition above, we see that each element g_k of $B_k(w)$ is uniquely determined by the values of $(x_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)})$ for $\text{inv}_\lambda^k(w) = \{(k, \ell_1), \dots, (k, \ell_d)\}$ and $d = d_k$. To emphasize this, we sometimes write $g_k = g_k(x_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)})$ and say that $(x_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)})$ are the coordinates of g_k .

Example 4.3. Let $n = 7$ and $\lambda = (3, 2, 2)$. We consider $w = [3, 2, 6, 1, 7, 4, 5]$ with corresponding tableau $R(w) \in \text{RS}(\lambda)$ as shown below; the base filling $R(e)$ is also below.

$$R(w) = \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 7 & \\ \hline 4 & 6 & \\ \hline \end{array} \quad R(e) = \begin{array}{|c|c|c|} \hline 3 & 6 & 7 \\ \hline 2 & 5 & \\ \hline 1 & 4 & \\ \hline \end{array}$$

In this case, $\text{inv}_\lambda(w) = \{(7, 5), (6, 5), (4, 2), (4, 3), (2, 1)\}$ so, in particular, $\text{inv}_\lambda^4(w) = \{(4, 2), (4, 3)\}$. We have $(w(4), w(2)) = (1, 2)$ and $(w(4), w(3)) = (1, 6)$. Since $X_\lambda \mathbf{e}_{w(4)} = \mathbf{0}$, the elements of $B_4(w)$ are matrices of the form

$$I_7 + x_{12}E_{12} + x_{16}E_{16} \text{ where } x_{12}, x_{16} \in \mathbb{C}.$$

For another example, consider $\text{inv}_\lambda^6(w) = \{(6, 5)\}$; we have $(w(6), w(5)) = (4, 7)$. In this case, $X_\lambda \mathbf{e}_{w(6)} = X_\lambda \mathbf{e}_4 = \mathbf{e}_1$ and $X_\lambda^2 \mathbf{e}_{w(6)} = X^2 \mathbf{e}_4 = \mathbf{0}$. Therefore the elements of $B_6(w)$ are matrices of the form

$$I_7 + x_{47}E_{47} + x_{47}E_{16} \text{ where } x_{47} \in \mathbb{C}.$$

Let $2 \leq k \leq n$ and $g_k \in B_k(w)$. Then $(g_k)_{aa} = 1$ for all $a \in [n]$. Suppose $(g_k)_{ab} \neq 0$ for $a \neq b$. By definition, $(a, b) = (w(\ell'), w(k'))$ where $\ell', k' \in [n]$ such that $X^m \mathbf{e}_{w(\ell)} = \mathbf{e}_{w(\ell')}$ and $X^m \mathbf{e}_{w(k)} = \mathbf{e}_{w(k')}$ for some $m \in \mathbb{Z}_{\geq 0}$. Since the action of X_λ on the standard basis vectors is determined by the base filling $R(e) \in \text{RS}(\lambda)$, it follows that $w(k')$ fills the m -th box to the left of $w(k)$ in $R(e)$ and $w(\ell')$ fills the m -th box to the left of $w(\ell)$ in $R(e)$. Equivalently, k' fills the m -th box to the left of k in $R(w)$ and ℓ' fills the m -th box to the left of ℓ in $R(w)$.

Since $(k, \ell) \in \text{inv}_\lambda(w)$, we know that the box labeled by k appears below the box labeled by ℓ and in the same column or in any column strictly to the left of ℓ in $R(w)$. Therefore the same must be true of the pair (k', ℓ') , that is, the box labeled by k' appears below the box labeled by ℓ' and in the same column or in any column strictly to the left of ℓ' in $R(w)$; and similarly for the boxes labeled by $w(k')$ and $w(\ell')$ in $R(e)$. By definition of the base filling $R(e)$, we conclude that $a = w(k') < w(\ell') = b$ so $g_k \in U$.

We summarize the discussion above in the following remark.

Remark 4.4 . Let $2 \leq k \leq n$ and $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$. Then $B_k(w) \subseteq U$ and furthermore, given $g_k \in B_k$ we have $(g_k)_{ab} \neq 0$ if and only if $a = b$ (in which case $(g_k)_{aa} = 1$) or $(a, b) = (w(\ell'), w(k'))$ where k' fills the m -th box to the left of k in $R(w)$ and ℓ' fills the m -th box to the left of ℓ in $R(w)$ for some $m \in \mathbb{Z}_{\geq 0}$ and $(k, \ell) \in \text{inv}_\lambda^k(w)$.

The remainder of this section contains results describing the structure of the matrices in $B_k(w)$; in most cases, our proofs consist of straightforward computations using linear algebra.

Lemma 4.5. Let $2 \leq k \leq n$ and $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$. The set of matrices $B_k(w)$ from Definition 4.2 is an abelian subgroup of U of dimension $|\text{inv}_\lambda^k(w)|$.

Proof. Let $g_k, h_k \in B_k(w)$, where g_k has coordinates $(x_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)})$ and h_k has coordinates $(y_{w(k)w(\ell_1)}, \dots, y_{w(k)w(\ell_d)})$ for $d = d_k$. We will prove $g_k h_k$ is the matrix in $B_k(w)$ with coordinates

$(x_{w(k)w(\ell_1)} + y_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)} + y_{w(k)w(\ell_d)})$. In other words, we prove that if $(k, \ell) \in \text{inv}_\lambda^k(w)$ and $\mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(\ell)}$ for some $m \in \mathbb{Z}_{\geq 0}$, then

$$(g_k h_k) \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)} + (x_{w(k)w(\ell)} + y_{w(k)w(\ell)}) X_\lambda^m \mathbf{e}_{w(k)} \quad (4.2)$$

and $(g_k h_k) \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$ otherwise. It follows directly from this formula that $B_k(w)$ is an abelian subgroup of U .

Now consider the action of $g_k h_k$ on any basis element $\mathbf{e}_{w(j)}$. If $w(j)$ does not appear in $R(e)$ to the left and in the same row as any $w(\ell)$ for $(k, \ell) \in \text{inv}_\lambda^k(w)$, that is, if $\mathbf{e}_{w(j)} \neq X_\lambda^m \mathbf{e}_{w(\ell)}$ for some $m \in \mathbb{Z}_{\geq 0}$, then both g_k and h_k map $\mathbf{e}_{w(j)}$ to itself and $(g_k h_k) \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$.

If $w(j)$ is to the left and in the same row of $R(e)$ as $w(\ell)$ for some $(k, \ell) \in \text{inv}_\lambda^k(w)$, then $g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)} + x_{w(k)w(\ell)} X_\lambda^m \mathbf{e}_{w(k)}$ and $h_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)} + y_{w(k)w(\ell)} X_\lambda^m \mathbf{e}_{w(k)}$ for some $m \in \mathbb{Z}_{\geq 0}$. Write $X_\lambda^m \mathbf{e}_{w(k)} = \mathbf{e}_{w(k')}$ where $w(k')$ labels the m -th box to the left of $w(k)$ in $R(e)$. Then k' labels the m -th box to the left of k in $R(w)$ and hence k' is not in the same row as ℓ for any $(k, \ell) \in \text{inv}_\lambda^k(w)$ by Remark 3.5. This implies $g_k \mathbf{e}_{w(k')} = \mathbf{e}_{w(k')} = h_k \mathbf{e}_{w(k')}$ and formula (4.2) now follows. The assertion that the dimension of this subgroup is $|\text{inv}_\lambda^k(w)|$ is clear from the definition. $\square$

Lemma 4.6. *Let $2 \leq k \leq n$ and $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$. For all $g_k \in B_k(w)$ we have*

$$g_k(\text{span}\{\mathbf{e}_{w(1)}, \mathbf{e}_{w(2)}, \dots, \mathbf{e}_{w(k)}\}) = \text{span}\{\mathbf{e}_{w(1)}, \mathbf{e}_{w(2)}, \dots, \mathbf{e}_{w(k)}\}$$

and $g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$ for all $j \geq k$. Furthermore, $g_k(\ker(X_\lambda)) \subseteq \ker(X_\lambda)$.

Proof. By definition, if $\mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(\ell)}$ for some $m \in \mathbb{Z}_{\geq 0}$ and $(k, \ell) \in \text{inv}_\lambda^k(w)$ then $w(j)$ fills the m -th box to the left of $w(\ell)$ in $R(e)$. Therefore j fills the m -th box to the left of ℓ in $R(w)$ and since $R(w) \in \text{RS}(\lambda)$ we have $j \leq \ell < k$. This shows $g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$ for all $j \geq k$. Similarly, if $X_\lambda^m \mathbf{e}_{w(k)} = \mathbf{e}_{w(k')}$ then k' fills the m -th box to the left of k in $R(w)$ and $k' \leq k$. The first assertion of the lemma now follows.

To prove the second, suppose $\mathbf{e}_{w(j)} \in \ker(X_\lambda)$ with $\mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(\ell)}$ for some $m \in \mathbb{Z}_{\geq 0}$ and $(k, \ell) \in \text{inv}_\lambda^k(w)$. We must show $g_k \mathbf{e}_{w(j)} \in \ker(X_\lambda)$, or equivalently, if $X_\lambda^m \mathbf{e}_{w(k)} \neq \mathbf{0}$ then $X_\lambda^m \mathbf{e}_{w(k)} \in \ker(X_\lambda)$. Since $g_k \in U$, if $X_\lambda^m \mathbf{e}_{w(k)} \neq \mathbf{0}$ then $X_\lambda^m \mathbf{e}_{w(k)} = \mathbf{e}_{w(k')}$ where $w(k') < w(j)$. The assumption that $\mathbf{e}_{w(j)} \in \ker(X_\lambda)$ implies $w(j)$ fills a box in the first column of $R(e)$ and the fact that $w(k') < w(j)$ implies that $w(k')$ fills a box below $w(j)$ in the first column of $R(e)$ so $X_\lambda \mathbf{e}_{w(k')} = \mathbf{0}$ as desired. $\square$

The next lemma shows that the matrices in $B_k(w)$ for $k < n$ almost commute with X_λ and the elements of $B_n(w)$ always commute with X_λ .

Lemma 4.7. *Suppose $R(w) \in \text{RS}(\lambda)$ and let $2 \leq k \leq n$. Given $\ell \in [n]$, let r_ℓ denote the label of the box directly to the right of the box labeled by ℓ in $R(w)$, if such a box exists. Then for all $g_k \in B_k(w)$,*

$$(g_k X_\lambda - X_\lambda g_k) \mathbf{e}_{w(j)} = \begin{cases} x_{w(k)w(\ell)} \mathbf{e}_{w(k)} & \text{if } j = r_\ell, (k, \ell) \in \text{inv}_\lambda^k(w) \\ 0 & \text{otherwise} \end{cases} \quad (4.3)$$

where $x_{w(k)w(\ell)} \in \mathbb{C}$ for $(k, \ell) \in \text{inv}_\lambda^k(w)$ is a coordinate of $g_k \in B_k(w)$. In particular, if $k = n$ this formula shows that X_λ and g_n commute.

Proof. First, suppose $j = r_\ell$ for some $\ell \in [n]$ such that $(k, \ell) \in \text{inv}_\lambda^k(w)$. By definition, this is the case if and only if $X_\lambda \mathbf{e}_{w(j)} = \mathbf{e}_{w(\ell)}$ and $k \leq j$. In particular, Lemma 4.6 tells us $g_k \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$. We now have:

$$(g_k X_\lambda - X_\lambda g_k) \mathbf{e}_{w(j)} = g_k \mathbf{e}_{w(\ell)} - \mathbf{e}_{w(\ell)} = x_{w(k)w(\ell)} \mathbf{e}_{w(k)}$$

where the last equation follows directly from Definition 4.2.

It remains to show that if $j \neq r_\ell$ for any $\ell \in [n]$ such that $(k, \ell) \in \text{inv}_\lambda^k(w)$, then $g_k X_\lambda \mathbf{e}_{w(j)} = X_\lambda g_k \mathbf{e}_{w(j)}$. This is straightforward to prove using Definition 4.2, so we omit the details.

Finally, when $k = n$ we have $(n, \ell) \in \text{inv}_\lambda(w)$ only if ℓ labels a box at the end of a row in $R(e)$. Thus (4.3) tells us X_λ and g_n commute. $\square$

Our goal is to use induction to analyze the flags in $C_w \cap \mathcal{B}^{X_\lambda}$. In particular, the next statement shows that we can naturally identify the group $B_k(w)$ for $k < n$ with a subgroup of the same type in $GL_{n-1}(\mathbb{C})$ using the map from (2.1).

Proposition 4.8. *Suppose $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$ and $w = vy$ is the factorization defined in (2.2), with $i = w(n)$.*

- (1) For all $2 \leq k \leq n-1$, we have $v^{-1}B_k(w)v = B_k(y)$ where $B_k(y)$ is the subgroup of $U_0 \subseteq GL_{n-1}(\mathbb{C})$ corresponding to $R(y) \in \text{RS}(\lambda')$.
- (2) Each $g_n \in B_n(w)$ can be factored uniquely as $g_n = u_i b_n$ where $u_i \in U_i$, $b_n \in vU_0 v^{-1}$, and the i -th row of u_i is equal to the i -th row of g_n . In particular, $u_i = \rho_1^{(\lambda)}(g_n w E_\bullet)$ and there exists an isomorphism of varieties $\mathbb{C}^{d_n} \rightarrow B_n(w)$ with inverse given by $g_n \mapsto \rho_1^{(\lambda)}(g_n w E_\bullet)$.

Proof. The proof of Lemma 3.7 shows that $R(y)$ is the row-strict tableau we obtain by deleting the box labeled by n from $R(w)$. This lemma also tells us that $R(y) \in \text{RS}(\lambda')$ where λ' is the composition of n corresponding to $R(y)$, and for all $k \leq n-1$, $(k, \ell) \in \text{inv}_{\lambda'}(y)$ if and only if $(k, \ell) \in \text{inv}_\lambda(w)$. Recall that $X_{\lambda'}$ is given by the restriction of $v^{-1}X_\lambda v$ to $\mathbb{C}^{n-1} = \text{span}\{\mathbf{e}_1, \dots, \mathbf{e}_{n-1}\}$.

Let $g_k \in B_k(w)$ have coordinates $(x_{w(k)w(\ell_1)}, \dots, x_{w(k)w(\ell_d)})$ where $d = d_k$. We prove $v^{-1}g_k v \in B_k(y)$ by showing $v^{-1}g_k v$ acts on the basis vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_{n-1}\}$ by the formulas given in Definition 4.2 where λ is replaced by λ' and w is replaced by y . Suppose $(k, \ell) \in \text{inv}_{\lambda'}(y)$ and $\mathbf{e}_{y(j)} = X_{\lambda'}^m \mathbf{e}_{y(\ell)}$ for some $m \in \mathbb{Z}_{\geq 0}$. By definition of $X_{\lambda'}$, this is the case if and only if $\mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(\ell)}$ for $(k, \ell) \in \text{inv}_\lambda^k(w)$. Thus $v^{-1}g_k v$ is the matrix such that:

$$\begin{aligned} (v^{-1}g_k v)(\mathbf{e}_{y(j)}) &= v^{-1}(g_k \mathbf{e}_{w(j)}) = v^{-1}(\mathbf{e}_{w(j)} + x_{w(k)w(\ell)} X_\lambda^m \mathbf{e}_{w(k)}) \\ &= \mathbf{e}_{y(j)} + x_{w(k)w(\ell)} X_{\lambda'}^m \mathbf{e}_{y(k)}. \end{aligned}$$

By similar reasoning, we have $(v^{-1}g_k v)\mathbf{e}_{y(j)} = \mathbf{e}_{y(j)}$ if $\mathbf{e}_{y(j)} \neq X_{\lambda'}^m \mathbf{e}_{y(\ell)}$ for any $(k, \ell) \in \text{inv}_{\lambda'}(y)$ and $m \in \mathbb{Z}_{\geq 0}$. This proves $v^{-1}B_k(w)v \subseteq B_k(y)$, after relabeling the coordinate $x_{w(k)w(\ell)}$ by $x_{y(k)y(\ell)}$. The proof that $B_k(y) \subseteq v^{-1}B_k(w)v$ follows in exactly the same way.

To prove statement (2), suppose $g_n \in B_n(w)$ with coordinates $(x_{w(n)w(\ell_1)}, \dots, x_{w(n)w(\ell_d)}) = (x_{iw(\ell_1)}, \dots, x_{iw(\ell_d)})$ where $d = d_n$. Let u_i be the matrix defined uniquely by the equation

$$u_i(\mathbf{e}_{w(\ell)}) = \begin{cases} \mathbf{e}_{w(\ell)} + x_{iw(\ell)} X_\lambda^m \mathbf{e}_i & \text{if } (n, \ell) \in \text{inv}_\lambda^n(w) \\ \mathbf{e}_{w(\ell)} & \text{otherwise.} \end{cases}$$

Then $u_i \in U_i$ and by definition u_i has i -th row equal to g_n . Furthermore, b_n is uniquely determined by the formula

$$b_n(\mathbf{e}_{w(j)}) = \begin{cases} \mathbf{e}_{w(j)} + x_{iw(\ell)} X_\lambda^m \mathbf{e}_i & \text{if } (n, \ell) \in \text{inv}_\lambda^n(w), \mathbf{e}_{w(j)} = X_\lambda^m \mathbf{e}_{w(\ell)} \text{ for some } m \geq 1 \\ \mathbf{e}_{w(j)} & \text{otherwise.} \end{cases}$$

for all $m \geq 1$.

By the previous paragraph, the matrix u_i is uniquely determined by, and uniquely determines, g_n . We now show that $b_n := u_i^{-1}g_n \in vU_0 v^{-1}$. It is straightforward to see from the definition of u_i and g_n that $u_i^{-1}g_n$ has the same entries as g_n except for the i -th row, which is 0 in all off-diagonal entries. Thus $u_i^{-1}g_n$ may only have a nonzero off-diagonal entry in positions $(w(n'), w(\ell'))$ with $w(n') < w(\ell')$ where n' labels the m -th box to the left of n in $R(w)$ and ℓ' labels the m -th box to the left of ℓ in $R(w)$ for some $m \in \mathbb{Z}_{\geq 1}$. In particular, $n' \neq n$. It follows that $v^{-1}(u_i^{-1}g_n)v$ may

only have nonzero off-diagonal entries in positions $(y(n'), y(\ell'))$ where $y(n') \neq n$, $y(\ell') \neq n$, and $y(n') < y(\ell')$ so $b_n \in vU_0v^{-1}$ as desired.

Finally, the map $\mathbb{C}^{d_n} \rightarrow B_n(w)$ defined by $(x_{iw(\ell_1)}, \dots, x_{iw(\ell_d)}) \mapsto g_n(x_{iw(\ell_1)}, \dots, x_{iw(\ell_d)})$ is clearly a morphism of varieties. By the above, we may write $g_n = u_i b_n$. Since g_n commutes with X_λ we get $g_n w E_\bullet \in \mathcal{B}^{X_\lambda}$ and:

$$g_n w E_\bullet = u_i v u'_0 y E_\bullet \text{ where } u'_0 = v^{-1} b_n v \in U_0, u_i \in U_i.$$

This shows that $\rho_1^{(\lambda)}(g_n w E_\bullet) = u_i$, where $\rho_1^{(\lambda)}$ is the morphism defined in [Lemma 4.1](#). The inverse map $B_n(w) \rightarrow \mathbb{C}^{d_n}$ is precisely $g_n \mapsto u_i = \rho_1^{(\lambda)}(g_n w E_\bullet)$. This proves $\mathbb{C}^{d_n} \rightarrow B_n(w)$ is an isomorphism of varieties. $\square$

Example 4.9. We illustrate the previous proposition with an example. Let $n=6$ and $\lambda = (2, 2, 2)$ and $w = [3, 6, 2, 1, 5, 4]$. In this case, $v = [1, 2, 3, 5, 6, 4]$ and $y = [3, 5, 2, 1, 4, 6]$.

$$R(w) = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & 6 \\ \hline \end{array} \quad R(e) = \begin{array}{|c|c|} \hline 3 & 6 \\ \hline 2 & 5 \\ \hline 1 & 4 \\ \hline \end{array}$$

We have $\text{inv}_\lambda^6(w) = \{(6, 5), (6, 2)\}$ where $(w(6), w(5)) = (4, 5)$ and $(w(6), w(2)) = (4, 6)$. An arbitrary element of $B_6(w)$ is of the form:

$$\begin{aligned} g_6 &= I_6 + x_{45}(E_{45} + E_{12}) + x_{46}(E_{46} + E_{13}) \\ &= (I_6 + x_{45}E_{45} + x_{46}E_{46})(I_6 + x_{45}E_{12} + x_{46}E_{13}) = u_4 b_6 \end{aligned}$$

where $x_{45}, x_{46} \in \mathbb{C}$ are the coordinates of g_6 , $u_4 = I_6 + x_{45}E_{45} + x_{46}E_{46}$ and $b_6 = I_6 + x_{45}E_{12} + x_{46}E_{13}$. Note that $v^{-1}b_6v = b_6 \in U_0$, confirming statement (2) of [Proposition 4.8](#). The isomorphism $\mathbb{C}^2 \rightarrow B_6(w)$ is defined by

$$(x_{45}, x_{46}) \mapsto (I_6 + x_{45}E_{45} + x_{46}E_{46})(I_6 + x_{45}E_{12} + x_{46}E_{13}) = g_6(x_{45}, x_{46}).$$

Now consider $\text{inv}_\lambda^4(w) = \{(4, 2), (4, 3)\}$; we have $(w(4), w(2)) = (1, 6)$ and $(w(4), w(3)) = (1, 2)$. An arbitrary element of $B_4(w)$ is of the form:

$$g_4 = I_6 + x_{12}E_{12} + x_{16}E_{16}$$

where x_{12}, x_{16} are the coordinates of g_4 . We have $v^{-1}g_4v = I_6 + x_{12}E_{12} + x_{16}E_{15} \in B_4(y) \subset U_0$. Since $B_4(y) = \{I_5 + x_{12}E_{12} + x_{15}E_{15} \mid x_{12}, x_{15}\}$ we obtain $v^{-1}B_4(w)v = B_4(y)$ via the identification from (2.1), confirming statement (1) of [Proposition 4.8](#).

5. An affine paving

In this section, we prove our first main theorem. In particular, [Theorem 5.7](#) below tells us that $\mathcal{H}\text{ess}(X_\lambda, h)$ is paved by affines for all $h \in \mathcal{H}$. As noted above, this result is not new, but our constructive methods are elementary and provide more insight into the structure of the paving. We obtain our paving of $\mathcal{H}\text{ess}(X_\lambda, h)$ by restricting an affine paving for the Springer fiber $\mathcal{B}^{X_\lambda}$. As a consequence, we recover Tymoczko's formulas for the dimension of each affine cell, originally proved in [17, Theorem 7.1] for Hessenberg varieties in the flag variety of $GL_n(\mathbb{C})$ corresponding to Hessenberg functions such that $h(i) \geq i$ for all i .

We begin by constructing an affine paving of the Springer fiber $\mathcal{B}^{X_\lambda}$ using the subgroups $B_k(w)$ introduced above. Throughout this section, λ denotes a fixed composition of n . The next proposition is the lynchpin of our inductive arguments; our proof resembles that of ref. [16, Lemma 1] (see also [13, §11.3]).

Proposition 5.1. Suppose $R(w) \in \text{RS}(\lambda)$ and $w = vy$ is the factorization from (2.2) with $i = w(n)$. Let λ' be the composition of $n - 1$ obtained from λ by deleting the box labeled by n in $R(w)$ and $X_{\lambda'}$ denote the restriction of $v^{-1}X_{\lambda}v$ to $\mathbb{C}^{n-1} = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n-1}\}$. There exists an isomorphism of varieties

$$\varphi_w : \mathbb{C}^{d_n} \times (C_y \cap \mathcal{B}^{X_{\lambda'}}) \rightarrow C_w \cap \mathcal{B}^{X_{\lambda}}$$

where $d_n = |\text{inv}_{\lambda}^n(w)|$.

Proof. We use the isomorphism $\mathbb{C}^{d_n} \rightarrow B_n(w)$ from Proposition 4.8(2) throughout the proof. Given a flag $V'_{\bullet} \in \mathcal{B}^{X_{\lambda'}}$ we write $V'_{\bullet} = u_0 y E'_{\bullet}$ for some $u_0 \in U_0$. It suffices to show that

$$\varphi_w : B_n(w) \times (C_y \cap \mathcal{B}^{X_{\lambda'}}) \rightarrow C_w \cap \mathcal{B}^{X_{\lambda}}, (g_n, u_0 y E'_{\bullet}) \mapsto g_n v u_0 y E_{\bullet}$$

is an isomorphism of varieties. Note first that φ_w is well defined since g_n commutes with X_{λ} by Lemma 4.7 and $v u_0 y E_{\bullet} \in \mathcal{B}^{X_{\lambda}}$ by Remark 3.8.

To begin, we argue that φ_w is a bijection. Suppose first that $g_n v u_0 y E_{\bullet} = h_n v u'_0 y E_{\bullet}$ for some $g_n, h_n \in B_n(w)$ and $u_0, u'_0 \in U_0$. By Proposition 4.8(2), we may write $g_n = u_i b_n$ for $u_i \in U_i$ such that the i -th row of u_i is equal to the i -th row of g_n and $b_n \in v U_0 v^{-1}$. Similarly, we have $h_n = u'_i b'_n$ for $u'_i \in U_i$ such that the i -th row of u'_i is equal to the i -th row of h_n and $b'_n \in v U_0 v^{-1}$. Thus

$$u_i = \rho_1^{(\lambda)}(g_n v u_0 E_{\bullet}) = \rho_1^{(\lambda)}(h_n v u'_0 E_{\bullet}) = u'_i$$

where $\rho_1^{(\lambda)}$ is the map defined in (4.1) above. Proposition 4.8(2) now implies $g_n = h_n$ and consequently $u_0 y E_{\bullet} = u'_0 y E_{\bullet}$ as well. This shows that φ_w is injective.

To prove surjectivity, let $u w E_{\bullet} \in C_w \cap \mathcal{B}^{X_{\lambda}}$ for some $u \in U^w$. Using Lemma 2.8 we may write $u w E_{\bullet} = u_i v u_0 y E_{\bullet}$ for $u_0 \in U_0$ and $u_i \in U_i$. By Lemma 4.1, we have that $\rho_1^{(\lambda)}(u w E_{\bullet}) = u_i \in \mathbb{C}^{d_n}$. Let $g_n \in B_n(w)$ have coordinates determined by $\rho_1^{(\lambda)}(u w E_{\bullet})$ via the isomorphism from Proposition 4.8(2). Thus we have $g_n = u_i b_n$ where $u_i = \rho_1^{(\lambda)}(u w E_{\bullet})$ and $b_n \in v U_0 v^{-1}$. Now

$$g_n^{-1} u w E_{\bullet} = v u'_0 y E_{\bullet} \in \mathcal{B}^{X_{\lambda}} \quad \text{where} \quad u'_0 = v^{-1} b_n v u_0 \in U_0.$$

This shows that $\varphi_w(g_n, u'_0 y E'_{\bullet}) = u w E_{\bullet}$, so φ_w is also surjective.

The bijection φ_w is in fact a morphism of varieties since $\mathbb{C}^{d_n} \rightarrow B_n(w)$ and the action map $GL_{n-1}(\mathbb{C}) \times \text{Flags}(\mathbb{C}^n) \rightarrow \text{Flags}(\mathbb{C}^n)$ are morphisms of varieties. The inverse map, defined by $\varphi_w^{-1}(u w E_{\bullet}) = (g_n, v^{-1} g_n^{-1} u w E'_{\bullet})$ where $g_n \in B_n(w)$ has coordinates determined by $\rho_1^{(\lambda)}(u w E_{\bullet})$, is also a morphism of varieties. This concludes the proof. $\square$

We now introduce flags constructed using the subgroups $B_k(w)$ of the previous section. We use these flags to give an explicit description of our affine paving of the Springer fibers below.

Definition 5.2. Let $w \in S_n$ such that $R(w) \in \text{RS}(\lambda)$. We define a subset $\mathcal{D}_w \subseteq C_w$ inductively as follows. First, set $\mathcal{D}_w^1 = \{w(E_{\bullet})\}$. Then for each $2 \leq k \leq n$ let

$$\mathcal{D}_w^k = \{g_k(V_{\bullet}) \mid g_k \in B_k(w) \text{ and } V_{\bullet} \in \mathcal{D}_w^{k-1}\}.$$

Let $\mathcal{D}_w := \mathcal{D}_w^n$. In other words, $b w E_{\bullet} \in \mathcal{D}_w$ if and only if $b = g_n g_{n-1} \dots g_2$ with $g_k \in B_k(w)$ for all k .

The definition of $\mathcal{D}_w$ depends on our choice of composition λ (since $R(w) \in \text{RS}(\lambda)$ and the definition of $B_k(w)$ depends on $R(w)$), but we suppress this dependence in the notation for $\mathcal{D}_w$. Note that for any composition, we have $\mathcal{D}_e = \{E_{\bullet}\}$.

Example 5.3. Continuing Example 4.9, let $n = 6$ and $\lambda = (2, 2, 2)$ and $w = [3, 6, 2, 1, 5, 4]$.

$$R(w) = \begin{bmatrix} 1 & 2 \\ 3 & 5 \\ 4 & 6 \end{bmatrix} \quad R(e) = \begin{bmatrix} 3 & 6 \\ 2 & 5 \\ 1 & 4 \end{bmatrix}$$

In this case, $\text{inv}_\lambda(w) = \{(6, 5), (6, 2), (5, 2), (4, 3), (4, 2), (3, 2)\}$. Computing the $B_k(w)$ subgroups, we have that $B_2(w) = \{I_6\}$ and:

$$B_3(w) = \left\{ \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & x_{26} \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right\}, \quad B_4(w) = \left\{ \begin{bmatrix} 1 & x_{12} & 0 & 0 & 0 & x_{16} \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right\},$$

$$B_5(w) = \left\{ \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & x_{56} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & x_{56} \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right\}, \quad B_6(w) = \left\{ \begin{bmatrix} 1 & x_{45} & x_{46} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & x_{45} & x_{46} \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right\}$$

where $x_{26}, x_{12}, x_{16}, x_{56}, x_{45}, x_{46} \in \mathbb{C}$. This gives:

$$\begin{aligned} \mathcal{D}_w^1 &= \mathcal{D}_w^2 = \{(\mathbf{e}_3 | \mathbf{e}_6 | \mathbf{e}_2 | \mathbf{e}_1 | \mathbf{e}_5 | \mathbf{e}_4)\} \\ \mathcal{D}_w^3 &= \{(\mathbf{e}_3 | \mathbf{e}_6 + x_{26}\mathbf{e}_2 | \mathbf{e}_2 | \mathbf{e}_1 | \mathbf{e}_5 | \mathbf{e}_4)\} \\ \mathcal{D}_w^4 &= \{(\mathbf{e}_3 | \mathbf{e}_6 + x_{16}\mathbf{e}_1 + x_{26}(\mathbf{e}_2 + x_{12}\mathbf{e}_1) | \mathbf{e}_2 + x_{12}\mathbf{e}_1 | \mathbf{e}_1 | \mathbf{e}_5 | \mathbf{e}_4)\} \\ \mathcal{D}_w^5 &= \{(\mathbf{e}_3 + x_{56}\mathbf{e}_2 | \mathbf{e}_6 + x_{56}\mathbf{e}_5 + x_{16}\mathbf{e}_1 + x_{26}(\mathbf{e}_2 + x_{12}\mathbf{e}_1) | \mathbf{e}_2 + x_{12}\mathbf{e}_1 | \mathbf{e}_1 | \mathbf{e}_5 | \mathbf{e}_4)\}, \\ \mathcal{D}_w^6 &= \{(\mathbf{e}_3 + x_{46}\mathbf{e}_1 + x_{56}(\mathbf{e}_2 + x_{45}\mathbf{e}_1) | \mathbf{e}_6 + x_{46}\mathbf{e}_4 + x_{56}(\mathbf{e}_5 + x_{45}\mathbf{e}_4) + x_{16}\mathbf{e}_1 \cdots \\ &\quad \cdots + x_{26}(\mathbf{e}_2 + x_{45}\mathbf{e}_1 + x_{12}\mathbf{e}_1) | \mathbf{e}_2 + x_{45}\mathbf{e}_1 + x_{12}\mathbf{e}_1 | \mathbf{e}_1 | \mathbf{e}_5 + x_{45}\mathbf{e}_4 | \mathbf{e}_4)\}. \end{aligned}$$

The set $\mathcal{D}_w$ has an inductive structure. Indeed, if $w(n) = i$ write $w = \nu y$ where $y \in S_{n-1}$ and $\nu = s_i s_{i+1} \cdots s_{n-2} s_{n-1}$ as in (2.2). Let $V_\bullet = g_n g_{n-1} \cdots g_2 w E_\bullet \in \mathcal{D}_w$. Statement (1) of Proposition 4.8 implies $g'_k := \nu^{-1} g_k \nu \in B_k(y)$ for all $2 \leq k \leq n-1$, so $V'_\bullet := g'_{n-1} \cdots g'_2 y(E'_\bullet) \in \mathcal{D}_y \subset \text{Flags}(\mathbb{C}^{n-1})$. Here $\mathcal{D}_y$ is the collection of flags determined as in Definition 5.2 for $y \in S_{n-1}$ with $R(y) \in \text{RS}(\lambda')$. We obtain a surjective map:

$$B_n(w) \times \mathcal{D}_y \rightarrow \mathcal{D}_w; (g_n, V'_\bullet) \mapsto V_\bullet. \quad (5.1)$$

The next result gives our affine paving of $\mathcal{B}^{X_\lambda}$.

Theorem 5.4. *Let $w \in S_n$ and λ a composition of n such that $R(w) \in \text{RS}(\lambda)$. There exists an isomorphism of varieties $\tilde{\varphi}_w : \mathbb{C}^{d_w} \rightarrow C_w \cap \mathcal{B}^{X_\lambda}$ with image equal to $\mathcal{D}_w$.*

Proof. We argue using induction on $n \geq 1$. The base case of $n = 1$ is trivial since $\text{Flags}(\mathbb{C}) = \{E_\bullet\}$ where $E_\bullet = \text{span}\{\mathbf{e}_1\}$ in this case. The required isomorphism $\tilde{\varphi}_e$ is the map $\mathbf{0} \mapsto E_\bullet$.

Now suppose $n \geq 2$. Let $w = \nu y$ be the factorization of w from (2.2), λ' the composition of $n-1$ obtained from λ by deleting the box labeled by n in $R(w)$, and $X_{\lambda'}$ the restriction of $\nu^{-1} X_\lambda \nu$ to $\mathbb{C}^{n-1} = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n-1}\}$. Consider $C_y \cap \mathcal{B}^{X_{\lambda'}} \subseteq \text{Flags}(\mathbb{C}^{n-1})$. By induction, there exists an isomorphism of varieties $\tilde{\varphi}_y : \mathbb{C}^{d_y} \rightarrow C_y \cap \mathcal{B}^{X_{\lambda'}}$ with image equal to $\mathcal{D}_y$. Here $d_y = |\text{inv}_{\lambda'}(y)|$ and $\mathcal{D}_y$ is the collection of flags determined as in Definition 5.2 for $y \in S_{n-1}$ with $R(y) \in \text{RS}(\lambda')$. By

Lemma 3.7 we have $d_w = d_n + d_y$ and **Proposition 4.8(2)** gives us an isomorphism $\mathbb{C}^{d_w} \simeq B_n(w) \times \mathbb{C}^{d_y}$. Using this identification we define

$$\tilde{\varphi}_w : B_n(w) \times \mathbb{C}^{d_y} \rightarrow C_w \cap \mathcal{B}^{X_\lambda}, \quad \tilde{\varphi}_w(g_n, \mathbf{x}) = \varphi_w(g_n, \tilde{\varphi}_y(\mathbf{x}))$$

where φ_w is the isomorphism of **Proposition 5.1**. This is an isomorphism of varieties since both φ_w and $\tilde{\varphi}_y$ are isomorphisms. Using the description of the image of $\tilde{\varphi}_y$ and definition of φ_w , it follows that the image of $\tilde{\varphi}_w$ is equal to the image of the map (5.1), which is $\mathcal{D}_w$. $\square$

By the theorem, $C_w \cap \mathcal{B}^{X_\lambda} = \mathcal{D}_w$ for all $R(w) \in \text{RS}(\lambda)$. The main theorem of this section analyzes the structure of the intersections $\mathcal{D}_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$ in greater detail. We begin with a technical statement generalizing **Lemma 4.7**; it shows that the flags in $\mathcal{D}_w$ satisfy strong linear conditions.

Lemma 5.5. *Let $R(w) \in \text{RS}(\lambda)$ and $g_k \in B_k(w)$ for each k . Furthermore, for each $2 \leq k \leq n$ let*

$$\mathbf{v}_j^{(k)} = g_k g_{k-1} \cdots g_2 \mathbf{e}_{w(j)} \text{ for all } 1 \leq j \leq n.$$

Given $\ell \in [n]$, let r_ℓ denote the label of the box directly to the right of the box labeled by ℓ in $R(w)$, if such a box exists. Then

$$(g_k X_\lambda - X_\lambda g_k) \mathbf{v}_j^{(k-1)} = \begin{cases} x_{w(k)w(\ell)} \mathbf{v}_k^{(k-1)} & \text{if } j = r_\ell, (k, \ell) \in \text{inv}_\lambda^k(w) \\ 0 & \text{otherwise} \end{cases} \quad (5.2)$$

where $x_{w(k)w(\ell)} \in \mathbb{C}$ for $(k, \ell) \in \text{inv}_\lambda^k(w)$ is a coordinate of $g_k \in B_k(w)$.

Proof. Since $\mathbf{v}_j^{(k-1)} = g_{k-1} \cdots g_2 \mathbf{e}_{w(j)}$ and $\mathbf{v}_k^{(k-1)} = g_{k-1} \cdots g_2 \mathbf{e}_{w(k)}$, by **Lemma 4.7** it suffices to show that the matrices $A = g_{k-1} \cdots g_2$ and $g_k X_\lambda - X_\lambda g_k$ commute.

In order to establish this claim, note that **Lemma 4.7** also implies

$$\ker(g_k X_\lambda - X_\lambda g_k) = \text{span}\{\mathbf{e}_{w(j)} \mid j \neq r_\ell \text{ for } (k, \ell) \in \text{inv}_\lambda^k(w)\}.$$

In particular, we have $\text{span}\{\mathbf{e}_{w(1)}, \dots, \mathbf{e}_{w(k)}\} \subseteq \ker(g_k X_\lambda - X_\lambda g_k)$.

If $\mathbf{e}_{w(j)} \notin \ker(g_k X_\lambda - X_\lambda g_k)$ then $j = r_\ell$ for some $(k, \ell) \in \text{inv}_\lambda^k(w)$ and $j \geq k$. **Lemma 4.6** implies $A \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$ and $A \mathbf{e}_{w(k)} = \mathbf{e}_{w(k)}$ so:

$$(g_k X_\lambda - X_\lambda g_k) A \mathbf{e}_{w(j)} = x_{w(k)w(\ell)} \mathbf{e}_{w(k)} = A(g_k X_\lambda - X_\lambda g_k) \mathbf{e}_{w(j)}$$

by **Lemma 4.7**. Now suppose $\mathbf{e}_{w(j)} \in \ker(g_k X_\lambda - X_\lambda g_k)$. **Lemma 4.6** implies that for all $j \leq k$, $A \mathbf{e}_{w(j)} \in \text{span}\{\mathbf{e}_{w(1)}, \dots, \mathbf{e}_{w(k)}\}$ and if $j \geq k$, $A \mathbf{e}_{w(j)} = \mathbf{e}_{w(j)}$. Thus $A \mathbf{e}_{w(j)} \in \ker(g_k X_\lambda - X_\lambda g_k)$ so

$$(g_k X_\lambda - X_\lambda g_k) A \mathbf{e}_{w(j)} = \mathbf{0} = A(g_k X_\lambda - X_\lambda g_k) \mathbf{e}_{w(j)}$$

in this case. $\square$

The following proposition records a key property satisfied by flags in $\mathcal{D}_w$. This statement is the technical heart of the proof of our main theorem below.

Proposition 5.6. *Suppose $R(w) \in \text{RS}(\lambda)$ and $V_\bullet = (\mathbf{v}_1 \mid \mathbf{v}_2 \mid \cdots \mid \mathbf{v}_n) \in \mathcal{D}_w$ so $V_\bullet = g_n g_{n-1} \cdots g_2 w(E_\bullet)$ for $g_k \in B_k(w)$, $2 \leq k \leq n$. Let $\ell \in [n]$. Suppose ℓ does not label a box at the end of a row in $R(w)$ and let $r = r_\ell$ denote the label of the box directly to the right of ℓ in $R(w)$. Then*

$$\mathbf{v}_\ell = X_\lambda \mathbf{v}_r + \sum_{\substack{\ell < t \leq n \\ (t, \ell) \in \text{inv}_\lambda(w)}} x_{w(t)w(\ell)} \mathbf{v}_t. \quad (5.3)$$

We assume the sum appearing above is zero whenever the index set is empty and $x_{w(t)w(\ell)} \in \mathbb{C}$ for $(t, \ell) \in \text{inv}_\lambda(w)$ are coordinates of $g_t \in B_t(w)$ for each t .

Proof. Let $V_\bullet^1 := w(E_\bullet)$ and

$$V_\bullet^k := g_k g_{k-1} \cdots g_2 w(E_\bullet) = (\mathbf{v}_1^{(k)} \mid \mathbf{v}_2^{(k)} \mid \cdots \mid \mathbf{v}_n^{(k)})$$

with $\mathbf{v}_j^{(k)}$ as in the statement of [Lemma 5.5](#). By definition, $V_\bullet = V_\bullet^n$ and $\mathbf{v}_k = \mathbf{v}_k^{(n)}$ for all $k \in [n]$. Note that [Lemma 4.6](#) implies $\mathbf{v}_k^{(k-1)} = \mathbf{e}_{w(k)} = \mathbf{v}_k^{(k)}$ for all $k \in [n]$.

Since r appears in the box directly to the right of ℓ in $R(w)$, we know $X_\lambda \mathbf{e}_{w(r)} = \mathbf{e}_{w(\ell)}$. We will show

$$\mathbf{v}_\ell^{(k)} - X_\lambda \mathbf{v}_r^{(k)} = \sum_{\substack{\ell < t \leq k \\ (t, \ell) \in \text{inv}_\lambda(w)}} x_{w(t)w(\ell)} \mathbf{v}_t^{(k)} \quad (5.4)$$

for every $2 \leq k \leq n$. This gives us (5.3) when $k = n$, proving the desired result.

We proceed by induction on k . When $k = 2$ the formula in [Lemma 4.7](#) implies

$$(g_2 X_\lambda - X_\lambda g_2) \mathbf{e}_{w(r)} = \sum_{(2, \ell) \in \text{inv}_\lambda(w)} x_{w(2)w(\ell)} \mathbf{e}_{w(2)}$$

where the RHS is zero whenever the index set is empty, that is, whenever $(2, \ell) \notin \text{inv}_\lambda^2(w)$. Applying the identities $\mathbf{v}_\ell^{(2)} = g_2 \mathbf{e}_{w(\ell)}$, $\mathbf{v}_r^{(2)} = g_2 \mathbf{e}_{w(r)}$, and $\mathbf{e}_{w(2)} = \mathbf{v}_2^{(2)}$ we obtain

$$\mathbf{v}_\ell^{(2)} - X_\lambda \mathbf{v}_r^{(2)} = \sum_{(2, \ell) \in \text{inv}_\lambda(w)} x_{w(2)w(\ell)} \mathbf{v}_2^{(2)}.$$

as desired.

Next, assume $k > 2$ and (5.4) holds for $k - 1$. Applying g_k to both sides of this equation, we obtain

$$\mathbf{v}_\ell^{(k)} - g_k X_\lambda \mathbf{v}_r^{(k-1)} = \sum_{\substack{\ell < t \leq k-1 \\ (t, \ell) \in \text{inv}_\lambda(w)}} x_{w(t)w(\ell)} \mathbf{v}_t^{(k)}. \quad (5.5)$$

The formula of [Lemma 5.5](#) shows that

$$g_k X_\lambda \mathbf{v}_r^{(k-1)} = \begin{cases} X_\lambda g_k \mathbf{v}_r^{(k-1)} + x_{w(k)w(\ell)} \mathbf{v}_k^{(k-1)} & \text{if } (k, \ell) \in \text{inv}_\lambda^k(w) \\ X_\lambda g_k \mathbf{v}_r^{(k-1)} & \text{otherwise.} \end{cases}$$

Substitute this formula for $g_k X_\lambda \mathbf{v}_r^{(k-1)}$ into the LHS of (5.5). Since $g_k \mathbf{v}_r^{(k-1)} = \mathbf{v}_r^{(k)}$ and $\mathbf{v}_k^{(k-1)} = \mathbf{v}_k^{(k)}$, this substitution yields the desired result. $\square$

We can now prove our main result.

Theorem 5.7. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function such that $h(i) < i$ for all i , λ be a composition of n , and $w \in S_n$ such that $R(w) \in \text{RS}_h(\lambda)$. Suppose $V_\bullet = g_n g_{n-1} \cdots g_2 w E_\bullet \in \mathcal{D}_w$ where $g_k \in B_k(w)$ for all $2 \leq k \leq n$.*

- (1) We have $V_\bullet \in \mathcal{H}\text{ess}(X_\lambda, h)$ if and only if $x_{w(k)w(\ell)} = 0$ for all $(k, \ell) \in \text{inv}_\lambda(w) \setminus \text{inv}_{\lambda, h}(w)$, where $x_{w(k)w(\ell)} \in \mathbb{C}$ is a coordinate of g_k .
- (2) The isomorphism of [Theorem 5.4](#) restricts to an isomorphism $\mathbb{C}^{d_{w, h}} \rightarrow C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$ where $d_{w, h} = |\text{inv}_{\lambda, h}(w)|$.

In particular, there is an affine paving of $\mathcal{H}\text{ess}(X_\lambda, h)$ with affine cells obtained by intersecting $\mathcal{H}\text{ess}(X_\lambda, h)$ with the Schubert cells.

Proof. By [Lemma 4.6](#), each $g_k \in B_k(w)$ preserves the kernel of X_λ . It follows immediately that if r labels a box in the first column of $R(w)$ (that is, if $\mathbf{e}_{w(r)} \in \ker(X_\lambda)$), then $\mathbf{v}_r = g_n g_{n-1} \cdots g_2 \mathbf{e}_{w(r)} \in \ker(X_\lambda)$. Thus, in this case, the condition that $X_\lambda \mathbf{v}_r \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{h(r)}\}$ is vacuously true.

Next, we consider the case in which r is not in the first column of $R(w)$. Rewriting formula (5.3) from [Proposition 5.6](#) we have,

$$X_\lambda \mathbf{v}_r = \mathbf{v}_\ell - \sum_{\substack{\ell < t \leq n \\ (t, \ell) \in \text{inv}_\lambda(w)}} x_{w(t)w(\ell)} \mathbf{v}_t$$

where ℓ is the label of the box immediately to the left of r in $R(w)$. Since $(t, \ell) \in \text{inv}_\lambda(w) \setminus \text{inv}_{\lambda, h}(w)$ if and only if $t > \ell$ and $h(r) < t < r$ it follows immediately that $X_\lambda \mathbf{v}_r \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{h(r)}\}$ if and only if $x_{w(t)w(\ell)} = 0$ for all $(t, \ell) \in \text{inv}_\lambda(w) \setminus \text{inv}_{\lambda, h}(w)$. This proves statement (1).

Let $\mathbf{x} = (x_{w(k)w(\ell)})_{(k, \ell) \in \text{inv}_\lambda(w)}$. By definition, $\tilde{\varphi}_w(\mathbf{x}) = g_n g_{n-1} \dots g_2 w E_\bullet$ where g_k has coordinates $(x_{w(k)w(\ell_1)}, \dots, x_{w(k), w(\ell_d)})$ for $d = d_k$ and $\text{inv}_\lambda^k(w) = \{(k, \ell_1), \dots, (k, \ell_d)\}$. Statement (1) implies $\tilde{\varphi}_w(\mathbf{x}) \in C_w \cap \mathcal{H}\text{ess}(X_\lambda, h) = \mathcal{D}_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$ if and only if $x_{w(k)w(\ell)} = 0$ for all $(k, \ell) \notin \text{inv}_{\lambda, h}(w)$. Thus, the restriction of $\tilde{\varphi}_w$ obtained by setting $x_{w(k)w(\ell)} = 0$ for all $(k, \ell) \notin \text{inv}_{\lambda, h}(w)$ yields an isomorphism $\mathbb{C}^{d_{w, h}} \rightarrow C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$. This proves (2) and the final assertion of the theorem follows from Remark 2.6. $\square$

Remark 5.8. One can also recover the results of Theorem 5.7 using similar methods as the second author in ref. [14]. It is an exercise to show that the formula for $\dim(C_w \cap \mathcal{H}\text{ess}(X_\lambda, h))$ given in Proposition 3.7 of ref. [14] is equal to $|\text{inv}_{\lambda, h}(w)|$.

Our next example continues the work of Examples 4.9 and 5.3.

Example 5.9 . Let $n = 6$ and $\lambda = (2, 2, 2)$ and $w = [3, 6, 2, 1, 5, 4]$. Consider the Hessenberg function $h = (0, 0, 1, 1, 3, 4)$. Then

$$\begin{aligned} \mathcal{D}_w \cap \mathcal{H}\text{ess}(X_\lambda, h) = \{ & (\mathbf{e}_3 + x_{46}\mathbf{e}_1 + x_{56}(\mathbf{e}_2 + x_{45}\mathbf{e}_1) \mid \mathbf{e}_6 + x_{46}\mathbf{e}_4 + x_{56}(\mathbf{e}_5 + x_{45}\mathbf{e}_4) + x_{16}\mathbf{e}_1 \cdots \\ & \cdots + x_{26}(\mathbf{e}_2 + x_{45}\mathbf{e}_1) \mid \mathbf{e}_2 + x_{45}\mathbf{e}_1 \mid \mathbf{e}_1 \mid \mathbf{e}_5 + x_{45}\mathbf{e}_4 \mid \mathbf{e}_4) \}. \end{aligned}$$

and

$$\begin{aligned} \mathcal{D}_w = \{ & (\mathbf{e}_3 + x_{46}\mathbf{e}_1 + x_{56}(\mathbf{e}_2 + x_{45}\mathbf{e}_1) \mid \mathbf{e}_6 + x_{46}\mathbf{e}_4 + x_{56}(\mathbf{e}_5 + x_{45}\mathbf{e}_4) + x_{16}\mathbf{e}_1 \cdots \\ & \cdots + x_{26}(\mathbf{e}_2 + x_{45}\mathbf{e}_1 + x_{12}\mathbf{e}_1) \mid \mathbf{e}_2 + x_{45}\mathbf{e}_1 + x_{12}\mathbf{e}_1 \mid \mathbf{e}_1 \mid \mathbf{e}_5 + x_{45}\mathbf{e}_4 \mid \mathbf{e}_4) \}. \end{aligned}$$

Since $(4, 3) \in \text{inv}_\lambda(w) \setminus \text{inv}_{\lambda, h}(w)$ and $(w(4), w(3)) = (1, 2)$, to get the former paving from the latter we set $x_{12} = 0$.

6. Geometric and combinatorial properties

In this section, we use the affine paving from above to study the geometry of $\mathcal{H}\text{ess}(X_\lambda, h)$ using the combinatorics of h -strict tableaux. We prove two main results, generalizing known facts about the Springer fiber to the Hessenberg varieties studied in this article. The first is that, when λ is a partition and the cell $C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$ has maximal dimension, then $R(w) \in \text{RS}_h(\lambda)$ is a standard tableau. Our second result proves that $\mathcal{H}\text{ess}(X_\lambda, h)$ is connected.

Assume for now that λ is a partition. If R is a row-strict tableau of shape λ , we let $\text{std}(R)$ denote the standard tableau obtained by reordering the entries in each column so that they increase from top to bottom. The next result generalizes Theorem 3.5 in ref. [15].

Lemma 6.1. *Let λ be a partition of n . If $R \in \text{RS}_h(\lambda)$ then $\text{std}(R) \in \text{RS}_h(\lambda)$.*

Proof. Suppose a_i is the entry in row i and column $k > 1$ of $\text{std}(R)$. Then a_i is greater than precisely $i - 1$ other entries of the k -th column in R . Since $R \in \text{RS}_h(\lambda)$, $h(a_i)$ is greater than or equal to at least i distinct entries in column $k - 1$. Thus $h(a_i)$ is greater than or equal to the entry in the box to the immediate left of a_i in $\text{std}(R)$. This implies that $\text{std}(R) \in \text{RS}_h(\lambda)$. $\square$

We now prove the first of the main results in this section. In general, the varieties $\mathcal{H}\text{ess}(X_\lambda, h)$ need not be equidimensional, but the following theorem shows that the maximal dimension cells in our affine paving correspond to standard tableaux. First we require the following definition.

Definition 6.2. Suppose $R \in \text{RS}_h(\lambda)$ has m columns. Let $d_R(i, j)$ for $1 \leq i \leq j \leq m$ be the number of Hessenberg inversions (k, ℓ) with k in column i and ℓ in column j of R .

Note that in the definition above, i and j may be equal. In the proof of the theorem below, we will consider a process that alters the entries of a tableau R . We use the notation $d(i, j)$ to refer to the number of Hessenberg inversions in columns i and j at each step of the process in our proof.

Theorem 6.3. *If $w \in S_n$ such that $\dim(C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)) = |\text{inv}_{\lambda, h}(w)|$ is maximal, then $R(w)$ is a standard tableau.*

Proof. Suppose $R \in \text{RS}_h(\lambda)$ is not a standard tableau. Write $R = R(y)$ and $\text{std}(R) = R(w)$ for some $y, w \in S_n$. We claim that $d_R(i, j) \leq d_{\text{std}(R)}(i, j)$ for all pairs (i, j) , and furthermore, at least one inequality is strict. This implies $|\text{inv}_{\lambda, h}(y)| < |\text{inv}_{\lambda, h}(w)|$, proving the theorem.

We prove this claim by describing a process which, when applied to any three columns $i, j, j+1$, will sort them. Furthermore, each step in this process will not increase the value of $d(i, j)$. (If $i = j$, then there are only two columns.) This will imply that $d_R(i, j) \leq d_{\text{std}(R)}(i, j)$.

We now describe this process; the idea is to “bubble sort columns.” If there is an inversion between two adjacent boxes in column i , we exchange them as well as the entries in that same row for column j and column $j+1$. We consider all rows to have the same number of squares and count blank squares as having a value of $+\infty$. Note that this means that, in the middle of this process, we may end up with a composition of n that is not a partition. Continue this until column i is increasing. Then do this for column j ; bubble sort out the inversions while simultaneously exchanging the corresponding entries in column $j+1$. Finally, bubble sort column $j+1$. This results in the entries of columns i, j , and $j+1$ being reordered so that they increase from top to bottom (i.e. we obtain the corresponding columns of $\text{std}(R)$).

First we analyze what happens when $i = j$. Our goal is to prove that this process does not increase the number of Hessenberg inversions (k, ℓ) with k and ℓ in column i . In this case, any such pair (k, ℓ) will still be a pair after sorting column i because $k > \ell$ so k will still label a box below ℓ in column i , and the number in the box to the right of ℓ does not change. Note that sorting column i does not change the fact that columns i and $i+1$ are h -strict.

Now consider sorting column $j+1 = i+1$. Suppose $\ell > s$ are adjacent entries in column i (so ℓ appears directly below s) and that we swap r_ℓ and r_s (the entries to the right of ℓ and s , respectively) as we sort column $i+1$, so $r_\ell < r_s$. We first prove that the h -strictness of columns i and $i+1$ is preserved. Indeed, if $s \leq h(r_s)$ and $\ell \leq h(r_\ell)$ then

$$\ell \leq h(r_\ell) \leq h(r_s) \text{ and } s < \ell \leq h(r_\ell)$$

Since $r_\ell < r_s$ implies $h(r_\ell) \leq h(r_s)$. Now, if we lose any pair counted by $d(i, j)$, it must be of the form (k, s) , where $k > s$ appears below s in column i and $h(r_\ell) < k \leq h(r_s)$. Note that $k \neq \ell$ in this case, since $\ell \leq h(r_s)$. Since $k > s$ and ℓ appears immediately below s , we get that $k > \ell$ labels a box below ℓ in column i . Since k is below ℓ and r_s now labels the box directly to the right of ℓ after the swap, we gain the Hessenberg inversion (k, ℓ) . Thus $d(i, j)$ does not decrease.

Now suppose $i < j$ and consider the operations on column i first. Because every move swaps entire rows, bubble sorting column i will not result in the loss of any Hessenberg inversions, and the columns j and $j+1$ remain h -strict. Now consider sorting column j . This clearly does not affect the inversions in $d(i, j)$ for the same reason, namely the columns j and $j+1$ are being simultaneously swapped. Columns j and $j+1$ remain h -strict in this case also.

Finally, consider the operations on column $j+1$. Say s is directly above ℓ in column j (so $s < \ell$) and $r_s > r_\ell$, as above. Swapping r_s and r_ℓ preserves the h -strictness of columns j and $j+1$; the argument is the same as above. If we lose any Hessenberg inversion (k, s) where k appears in

column i , then we must have $h(r_\ell) < k \leq h(r_s)$, just as above. But we gain the pair (k, ℓ) after swapping r_s and r_ℓ since $\ell \leq h(r_\ell) < k$ and $k \leq h(r_s)$. Once again, we see that $d(i, j)$ does not decrease.

Now we show that $d_{\text{std}(R)}(i, j) > d_R(i, j)$ for some (i, j) . Let column k be the last column of R whose entries are not already increasing from top to bottom. Consider the first instance the bubble sorting process swaps two elements when $i = j = k$. Say we swap ℓ and s with $\ell > s$. Since column $j + 1$ begins sorted, we have $r_\ell < r_s$. Now $\ell \leq h(r_\ell)$, so we gain the Hessenberg inversion (ℓ, s) . Moreover, as we showed in the general case where $i = j$, we do not lose any pairs counted by $d(i, i)$ when sorting column i , and every pair we lose when sorting column $i + 1$ matches with one we gain upon doing so. Thus $d_{\text{std}(R)}(k, k) > d_R(k, k)$, as desired. $\square$

We demonstrate the algorithm in the previous proof with an example.

Example 6.4. Let $n = 12$ and $h : [12] \rightarrow [12]$ the Hessenberg function defined by $h(i) = \max\{0, i - 2\}$. Consider beginning with the following h -strict tableaux R and $\text{std}(R)$:

$$R = \begin{array}{|c|c|c|c|} \hline 2 & 4 & 8 & 10 \\ \hline 1 & 5 & 7 & 11 \\ \hline 3 & 9 & 12 & \\ \hline 6 & & & \\ \hline \end{array} \rightarrow \text{std}(R) = \begin{array}{|c|c|c|c|} \hline 1 & 4 & 7 & 10 \\ \hline 2 & 5 & 8 & 11 \\ \hline 3 & 9 & 12 & \\ \hline 6 & & & \\ \hline \end{array}$$

When $i = j = 1$, we check that $d_R(i, j)$ does not decrease during the operations described in the proof. Each step of the process is displayed below; in this case there are only two steps.

$$\begin{array}{|c|c|} \hline 2 & 4 \\ \hline 1 & 5 \\ \hline 3 & 9 \\ \hline 6 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 1 & 5 \\ \hline 2 & 4 \\ \hline 3 & 9 \\ \hline 6 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & 9 \\ \hline 6 & \\ \hline \end{array}$$

We see $d_R(1, 1) = 2 \leq 3 = d_{\text{std}(R)}(1, 1)$, as desired. Next, when $i = 1$ and $j = 2$:

$$\begin{array}{|c|c|c|} \hline 2 & 4 & 8 \\ \hline 1 & 5 & 7 \\ \hline 3 & 9 & 12 \\ \hline 6 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 5 & 7 \\ \hline 2 & 4 & 8 \\ \hline 3 & 9 & 12 \\ \hline 6 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 4 & 8 \\ \hline 2 & 5 & 7 \\ \hline 3 & 9 & 12 \\ \hline 6 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 3 & 9 & 12 \\ \hline 6 & & \\ \hline \end{array}$$

and we again have $d_R(1, 2) = 1 \leq 1 = d_{\text{std}(R)}(1, 2)$. Since column 3 is the last column of R in which the values do not increase from top to bottom, we check that when $i = j = 3$, $d_R(i, j) < d_{\text{std}(R)}(i, j)$.

$$\begin{array}{|c|c|} \hline 8 & 10 \\ \hline 7 & 11 \\ \hline 12 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 7 & 10 \\ \hline 8 & 11 \\ \hline 12 & \\ \hline \end{array}$$

Indeed, $d_R(3, 3) = 0$ and $d_{\text{std}(R)}(3, 3) = 1$.

We now return to the setting in which λ is any composition of n . We prove that each Hessenberg variety $\mathcal{H}\text{ess}(X_\lambda, h)$ is connected. Recall that the dimension of the 0-cohomology group of any algebraic variety is equal to the number of connected components of that space. Thus, by Lemma 2.5 and Theorem 5.7 it suffices to show that there is a unique permutation $w \in S_n$ such that $\dim(C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)) = 0$.

Theorem 6.5. *For any Hessenberg function $h \in \mathcal{H}$ and composition λ , the Hessenberg variety $\mathcal{H}\text{ess}(X_\lambda, h)$ is connected whenever it is nonempty.*

Proof. By Remark 2.3, we may assume without loss of generality that λ is a partition of n . It suffices to prove that there exists a unique h -strict tableau that has no Hessenberg dimension pairs. Consider the following algorithm. Begin at the right-most column of λ . Label the boxes in this column from top to bottom, assigning n to the first box and decreasing by one at a time. Then move to the next right-most column, filling the boxes from top to bottom with the largest available number subject to the constraint that doing so does not violate h -strictness. Continue in this way and denote the resulting tableau by R_0 . We first show that this algorithm results in an h -strict tableau whenever $\mathcal{H}\text{ess}(X_\lambda, h) \neq \emptyset$; then we will show that R_0 is the unique tableaux satisfying the desired properties.

If $\mathcal{H}\text{ess}(X_\lambda, h)$ is nonempty, then at least one intersection $C_w \cap \mathcal{H}\text{ess}(X_\lambda, h)$ must be nonempty. Let $R = R(w)$ be the corresponding h -strict tableau. We show that R can be transformed into an h -strict tableau obtained via the algorithm above by a sequence of swaps. Given $\ell \in [n]$, recall that we denote by r_ℓ the label of the box directly to the right of ℓ , if it exists.

Going in order from top to bottom in each column, starting the right-most column and moving left, take the first square in R whose label differs from that dictated by the algorithm. In other words, we consider the first square in R with label ℓ such that there exists $k > \ell$ with $k \leq r_\ell$ and such that k appears below the box containing ℓ and in the same column or in any column strictly to the left of the column containing ℓ . We may assume k is the maximal with respect to these properties. Let ℓ_m and k_m be the values m boxes to the left of ℓ and k in tableau R , respectively.

Define R' to be the following h -strict tableau obtained from R . Exchange the positions of k and ℓ in R . If this is an h -strict tableau, then stop. Otherwise, the only problem that can arise is that $k_1 > h(\ell)$. In that case, we must also have $k_1 > \ell_1$. Now exchange the positions of k_1 and ℓ_1 ; if this results in an h -strict tableau then stop. Otherwise note that swapping k_1 into its new row does not change the fact that that row is h -strict. The only problem that can occur is if $k_2 > h(\ell_1)$ (and therefore $k_2 > \ell_2$). Continue this process of swapping until the tableau is h -strict. Indeed, because k originally appeared in a ‘later’ box than ℓ , we will always be able to perform this swap until the entire tableau is h -strict. The resulting R' is h -strict and has labels as dictated by the algorithm up through the box which now contains k .

Repeating the operation of the previous two paragraphs yields an h -strict tableau whose entries are dictated by the algorithm. This shows that the algorithm does indeed produce a completed h -strict tableau, R_0 . Now consider any pair (k, ℓ) with $k > \ell$ such that k labels a box in R_0 below ℓ and in the same column or in any column to the left of the column containing ℓ . By construction, placing k in the box containing ℓ would lead to a violation of h -strictness, that is, $k > h(r_\ell)$. This proves that R_0 has no Hessenberg inversions.

Finally, we show there are no other h -strict tableau $R(w)$ with $|\text{inv}_{\lambda, h}(w)| = 0$. Suppose $R(w) = R \in \text{RS}_h(\lambda)$ such that $R \neq R_0$ and that in the first box (using the same ordering as above) in which R differs from R_0 , R is labeled by ℓ rather than a k . Then $k > \ell$ since our algorithm for constructing R_0 always chooses the largest available number. The value k must be placed in a ‘later’ box of R than the box containing ℓ , so (k, ℓ) is an inversion of w . Since k could have been placed in the box containing ℓ (as it is in R_0), we have that $k \leq h(r_\ell)$, so (k, ℓ) is a Hessenberg inversion pair. This shows that all other h -strict tableaux have at least one Hessenberg inversion pair. Thus R_0 is the unique h -strict tableau with the desired properties. $\square$

In the Springer fiber case, the unique row-strict tableau without any Springer inversions is the base filling from Definition 3.1 above. However, in general the base filling may not be h -strict, as the next example demonstrates.

Example 6.6. Let $n=12$ and $h : [12] \rightarrow [12]$ the Hessenberg function defined by $h(i) = \max\{0, i - 3\}$. The base filling for $\lambda = (4, 4, 3, 1)$ and the h -strict tableau R_0 constructed in the proof of Theorem 6.5 are displayed below.

$$R(e) = \begin{array}{|c|c|c|c|} \hline 4 & 7 & 10 & 12 \\ \hline 3 & 6 & 9 & 11 \\ \hline 2 & 5 & 8 & \\ \hline 1 & & & \\ \hline \end{array} \quad R_0 = \begin{array}{|c|c|c|c|} \hline 3 & 6 & 9 & 12 \\ \hline 2 & 5 & 8 & 11 \\ \hline 4 & 7 & 10 & \\ \hline 1 & & & \\ \hline \end{array}$$

Note that $R(e)$ is not h -strict, since 10 and 12 appear as consecutive entries in the first row, but $10 \not\preceq 9 = h(12)$. The unique row-strict tableau with no Hessenberg inversions is R_0 .

Finally, we conclude with an example which shows that analogous Hessenberg varieties defined for other classical groups need not be connected.

Example 6.7. Consider the symplectic group $SP_4(\mathbb{C})$. Our convention is that the inner product on $\mathbb{C}^4 = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4\}$ is determined by the following matrix

$$s_J = \left[\begin{array}{c|c} 0 & J \\ \hline -J & 0 \end{array} \right] \text{ where } J = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

In other words, given $g \in GL_4(\mathbb{C})$ we have $g \in SP_4(\mathbb{C})$ if and only if $g^T s_J g = s_J$ where g^T denotes the transpose of g . The symplectic group has Lie algebra $\mathfrak{sp}_4(\mathbb{C}) = \{X \in \mathfrak{gl}_4(\mathbb{C}) \mid s_J X = -X^T s_J\}$.

The flag variety of $SP_4(\mathbb{C})$ consists of full flags of isotropic subspaces. Let

$$X = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \in \mathfrak{sp}_4(\mathbb{C})$$

and h be the Hessenberg function $(0, 1, 1, 3)$. This Hessenberg function gives a well-defined subvariety of the flag variety for $SP_4(\mathbb{C})$ using the Hessenberg space $H = H(h) \cap \mathfrak{sp}_4(\mathbb{C})$; see, for example, [14, Section 2.2] for the definition of the Hessenberg variety in an arbitrary flag variety. The variety of isotropic flags $V_\bullet = (\mathbf{v}_1 \mid \mathbf{v}_2 \mid \mathbf{v}_3 \mid \mathbf{v}_4)$ such that:

$$X\mathbf{v}_1 = \mathbf{0}; \quad X\mathbf{v}_2, X\mathbf{v}_3 \in V_1 = \text{span}\{\mathbf{v}_1\}; \quad \text{and} \quad X\mathbf{v}_4 \in V_3 = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} \quad (6.1)$$

is the Hessenberg variety $\mathcal{H}\text{ess}(X, H)$ in the flag variety of $SP_4(\mathbb{C})$. It is a subvariety of the Springer fiber corresponding to X . In this small dimensional case, it is easy to show that $\mathcal{H}\text{ess}(X, H)$ is paved by affines, and this paving is obtained by intersecting $\mathcal{H}\text{ess}(X, H)$ with each Schubert cell. Let s_1 be the element of the Weyl group of $SP_4(\mathbb{C})$ such that $s_1 \mathbf{e}_1 = \mathbf{e}_2$, $s_1 \mathbf{e}_2 = \mathbf{e}_1$, $s_1 \mathbf{e}_3 = \mathbf{e}_4$, $s_1 \mathbf{e}_4 = \mathbf{e}_3$, and let e denote the identity element. The Schubert cell C_{s_1} consists of all flags of the form:

$$(\mathbf{e}_2 + c \mathbf{e}_1 \mid \mathbf{e}_1 \mid \mathbf{e}_4 - c \mathbf{e}_3 \mid \mathbf{e}_3) \text{ for some } c \in \mathbb{C}$$

and $C_e = \{E_\bullet\}$. The conditions from (6.1) now imply:

$$C_e \cap \mathcal{H}\text{ess}(X, H) = \{E_\bullet\} \text{ and } C_{s_1} \cap \mathcal{H}\text{ess}(X, H) = \{s_1 E_\bullet\}$$

so $\dim(C_e \cap \mathcal{H}\text{ess}(X, H)) = \dim(C_{s_1} \cap \mathcal{H}\text{ess}(X, H)) = 0$. Since the 0-cohomology group of $\mathcal{H}\text{ess}(X, H)$ has dimension 2, $\mathcal{H}\text{ess}(X, H)$ is not connected.

Acknowledgements

The authors are thankful to the anonymous referee for their feedback, which improved the exposition of Section 5.

Funding

The second author is supported in part by NSF DMS 1954001.

References

- [1] Abe, H., Harada, M., Horiguchi, T., Masuda, M. (2014). The equivariant cohomology rings of regular nilpotent Hessenberg varieties in Lie type A: Research announcement. *Morfismos*. 18(2):51–65.
- [2] Björner, A., Brenti, F. (2005). *Combinatorics of Coxeter Groups, Volume 231 of Graduate Texts in Mathematics*. New York: Springer.
- [3] De Mari, F., Procesi, C., Shayman, M. A. (1992). Hessenberg varieties. *Trans. Amer. Math. Soc.* 332(2): 529–534. DOI: [10.1090/S0002-9947-1992-1043857-6](https://doi.org/10.1090/S0002-9947-1992-1043857-6).
- [4] Drellich, E. (2017). The containment poset of type A Hessenberg varieties. arXiv: 1710.05412.
- [5] Fresse, L. (2010). A unified approach on Springer fibers in the hook, two-row and two-column cases. *Transform. Groups*. 15(2):285–331. DOI: [10.1007/s00031-010-9096-3](https://doi.org/10.1007/s00031-010-9096-3).
- [6] Fresse, L. (2016). *Existence of affine pavings for varieties of partial flags associated to nilpotent elements*. International Mathematics Research Notices, Oxford University Press, pp. 418–472.
- [7] Fulton, W. (1984). *Intersection Theory*. Berlin: Springer-Verlag.
- [8] Fung, F. Y.C. (2003). On the topology of components of some Springer fibers and their relation to Kazhdan–Lusztig theory. *Adv. Math.* 178(2):244–276. DOI: [10.1016/S0001-8708\(02\)00072-5](https://doi.org/10.1016/S0001-8708(02)00072-5).
- [9] Graham, W., Zierau, R. (2011). Smooth components of Springer fibers. *Ann. Inst. Fourier*. 61(5):2139–2182. DOI: [10.5802/aif.2669](https://doi.org/10.5802/aif.2669).
- [10] Harada, M., Precup, M. (2017). The cohomology of abelian Hessenberg varieties and the Stanley–Stembridge conjecture. arXiv: 1709.06736.
- [11] Harada, M., Tymoczko, J. (2017). Poset pinball, GKM-compatible subspaces, and Hessenberg varieties. *J. Math. Soc. Japan*. 69(3):945–994. DOI: [10.2969/jmsj/06930945](https://doi.org/10.2969/jmsj/06930945).
- [12] Humphreys, J. E. (1964). *Linear Algebraic Groups*. New York: Springer-Verlag.
- [13] Jantzen, J. C. (2004). Nilpotent orbits in representation theory. In *Lie Theory*, vol. 228 of *Progr. Math.* Boston, MA: Birkhäuser Boston, pp. 1–211.
- [14] Precup, M. (2013). Affine pavings of Hessenberg varieties for semisimple groups. *Sel. Math. New. Ser.* 19(4):903–922. DOI: [10.1007/s00029-012-0109-z](https://doi.org/10.1007/s00029-012-0109-z).
- [15] Precup, M., Tymoczko, J. (2019). Springer fibers and Schubert points. *Eur. J. Combinatorics*. 76:10–26. DOI: [10.1016/j.ejc.2018.08.010](https://doi.org/10.1016/j.ejc.2018.08.010).
- [16] Spaltenstein, N. (1976). The fixed point set of a unipotent transformation on the flag manifold. *Nederl. Akad. Wetensch. Proc. Ser. A Indag. Math.* 79(5):452–456. DOI: [10.1016/S1385-7258\(76\)80008-X](https://doi.org/10.1016/S1385-7258(76)80008-X).
- [17] Tymoczko, J. S. (2006). Linear conditions imposed on flag varieties. *Amer. J. Math.* 128(6):1587–1604. DOI: [10.1353/ajm.2006.0050](https://doi.org/10.1353/ajm.2006.0050).