

On the Almeida-Thouless transition line in the Sherrington-Kirkpatrick model with centered Gaussian external field*

Wei-Kuo Chen[†]

Abstract

We study the phase transition of the free energy in the Sherrington-Kirkpatrick mean-field spin glass model with centered Gaussian external field. We show that the corresponding Almeida-Thouless line is the correct transition curve that distinguishes between the replica symmetric and replica symmetry breaking solutions in the Parisi formula.

Keywords: Almeida-Thouless line; Parisi formula; Sherrington-Kirkpatrick model.

MSC2020 subject classifications: 60K35; 82B44.

Submitted to ECP on March 11, 2021, final version accepted on November 3, 2021.

1 Introduction and main results

The famous Sherrington-Kirkpatrick (SK) mean-field spin glass model was introduced in [12] aiming to explain some unusual magnetic behavior of certain alloys. By means of the replica method, it was also proposed in [12] that the thermodynamic limit of the free energy in the SK model can be solved by the replica symmetric ansatz at very high temperature. A complete picture was later settled in the seminal works [10, 11] of Parisi, in which he adapted an ultrametric ansatz and deduced a variational formula for the limiting free energy at all temperature, known as the Parisi formula. This formula was rigorously established by Talagrand [13] utilizing the replica symmetry breaking bound discovered by Guerra [5]. See [8] for physicists' studies of the SK model as well as [9, 14, 15] for the recent mathematical progress.

For any $N \geq 1$, the Hamiltonian of the SK model is defined as

$$-H_N(\sigma) = \frac{\beta}{\sqrt{N}} \sum_{1 \leq i < j \leq N} g_{ij} \sigma_i \sigma_j + h \sum_{i=1}^N \sigma_i, \quad \forall \sigma \in \{-1, 1\}^N, \quad (1.1)$$

where g_{ij} 's are i.i.d. standard Gaussian. The parameters $\beta > 0$ and $h \in \mathbb{R}$ are the (inverse) temperature and external field, respectively. Define the free energy as

$$F_N(\beta, h) = \frac{1}{N} \log \sum_{\sigma \in \{-1, 1\}^N} e^{H_N(\sigma)}.$$

*Partly supported by NSF grant DMS-17-52184.

[†]University of Minnesota, United States of America. E-mail: wkchen@umn.edu

The famous Parisi formula [9, 15] asserts that almost surely,

$$\lim_{N \rightarrow \infty} F_N(\beta, h) = \min_{\alpha \in \text{Pr}([0,1])} P(\alpha). \quad (1.2)$$

Here, $\text{Pr}([0, 1])$ is the collection of all probability distribution functions on $[0, 1]$ equipped with the $L^1(dx)$ distance and P is a functional on $\text{Pr}([0, 1])$ defined as

$$P(\alpha) = \log 2 + \mathbb{E} \Phi_\alpha(0, h) - \frac{\beta^2}{2} \int_0^1 s \alpha(s) ds,$$

where Φ_α is the weak solution [6] to

$$\partial_s \Phi_\alpha(s, x) = -\frac{\beta^2}{2} (\partial_{xx} \Phi_\alpha(s, x) + \alpha(s) (\partial_x \Phi_\alpha(s, x))^2), \quad (s, x) \in [0, 1] \times \mathbb{R} \quad (1.3)$$

with boundary condition $\Phi(1, x) = \log \cosh x$. It is known [2] that the Parisi formula has a unique minimizer denoted by α_P . We say that the Parisi formula is solved by the replica symmetric solution if $\alpha_P = 1_{[q,1]}$ for some $q \in [0, 1]$ and is solved by the replica symmetry breaking solution if otherwise.

For any $\beta, h > 0$, let $q = q(\beta, h)$ be the unique solution (see [4] and [14, Proposition A.14.1]) to

$$q = \mathbb{E} \tanh^2(h + \beta z \sqrt{q}), \quad (1.4)$$

where z is standard Gaussian. In [11], it was conjectured that for $\beta, h > 0$, the SK model is solved by the replica symmetric solution if and only if (β, h) satisfies

$$\beta^2 \mathbb{E} \frac{1}{\cosh^4(h + \beta z \sqrt{q})} \leq 1.$$

In other words, for $\beta, h > 0$, the following equation, known as the Almeida-Thouless line,

$$\beta^2 \mathbb{E} \frac{1}{\cosh^4(h + \beta z \sqrt{q})} = 1, \quad (1.5)$$

characterizes the transition between the replica symmetric and replica symmetry breaking solutions. Toninelli [16] proved that above the AT line, i.e. $\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{q}) > 1$, the solution to the Parisi formula is replica symmetry breaking. Later Talagrand [15] and Jagannath-Tobasco [7] showed that inside the AT line, $\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{q}) \leq 1$, there exist fairly large regimes in which the Parisi formula is solved by replica symmetric solution. As parts of their regimes are not up to the AT line, verifying the exactness of the AT line remains open.

In this short note, we investigate the SK model with centered Gaussian external field and show that the corresponding AT line is indeed the transition line distinguishing between the replica symmetric and replica symmetry breaking solutions in the Parisi formula. For $\beta, h > 0$, the Hamiltonian of the SK model with centered Gaussian external field is defined as

$$-\mathcal{H}_N(\sigma) = \frac{\beta}{\sqrt{N}} \sum_{1 \leq i < j \leq N} g_{ij} \sigma_i \sigma_j + h \sum_{i=1}^N \xi_i \sigma_i, \quad \forall \sigma \in \{-1, 1\}^N,$$

where $(g_{ij})_{1 \leq i < j \leq N}$ and $(\xi_i)_{1 \leq i \leq N}$ are i.i.d. standard normal and are independent of each other. The free energy associated to this Hamiltonian is given by

$$\mathcal{F}_N(\beta, h) = \frac{1}{N} \mathbb{E} \log \sum_{\sigma \in \{-1, 1\}^N} \exp(-\mathcal{H}_N(\sigma)).$$




Figure 1: This figure describes the phase transition in the SK model with centered Gaussian external field. The gray area is the regime of replica symmetric solutions, while the white area is the regime of replica symmetry breaking solutions. The boundary between these two regimes is the AT line (1.7) corresponding to our model. The black-dash line is the AT line (1.5) corresponding to the original SK model with deterministic external field.

In a similar manner, the limiting free energy can be expressed by the Parisi formula as in (1.3) with a replacement of h by $h\xi$ for ξ a standard Gaussian random variable, more precisely, we have that almost surely,

$$\lim_{N \rightarrow \infty} \mathcal{F}_N(\beta, h) = \min_{\alpha \in \text{Pr}([0,1])} \mathcal{P}(\alpha),$$

where

$$\mathcal{P}(\alpha) = \log 2 + \mathbb{E} \Phi_\alpha(0, h\xi) - \frac{\beta^2}{2} \int_0^1 s \alpha(s) ds,$$

and Φ_α is defined as (1.3). Note that the proof of this formula is identically the same as those in [9, 15] with no essential modifications. Moreover, as in [2], the Parisi formula here also has a unique minimizer denoted by α_P . We again say that the Parisi formula is solved by the replica symmetric solution if $\alpha_P = 1_{[q,1]}$ for some $q \in [0, 1]$ and is solved by the replica symmetry breaking solution if otherwise.

The AT line corresponds to our model is formulated analogously. Let z, ξ be i.i.d. standard Gaussian random variables. For $\beta, h > 0$, let $q = q(\beta, h)$ be the unique fixed point of

$$q = \mathbb{E} \tanh^2(h\xi + \beta z \sqrt{q}), \quad (1.6)$$

where the existence and uniqueness of q are guaranteed thanks to the Latala-Guerra lemma.¹ The AT line associated to the SK model with centered Gaussian external field is the collection of all $(\beta, h) \in (0, \infty) \times (0, \infty)$ satisfying

$$\beta^2 \mathbb{E} \frac{1}{\cosh^4(h\xi + \beta z \sqrt{q})} = 1. \quad (1.7)$$

The following is our main result.

¹ In [4] and [14, Proposition A.14.1], it is known that for any $r \in \mathbb{R}$, $x^{-1} \mathbb{E} \tanh^2(r + \beta z \sqrt{x})$ is a strictly decreasing function in $x > 0$, which implies that $f(x) := x^{-1} \mathbb{E} \tanh^2(h\xi + \beta z \sqrt{x})$ is also strictly decreasing on $(0, \infty)$. Since $f(\infty) = 0$ and $f(0) = \infty$, there exists a unique x_0 such that $f(x_0) = 1$. This ensures that (1.6) has a unique solution.

Theorem 1.1. *Consider the SK model with centered Gaussian external field. For any $\beta > 0$ and $h > 0$, the Parisi formula exhibits the replica symmetric solution if and only if (β, h) lies inside the AT line, i.e.,*

$$\beta^2 \mathbb{E} \frac{1}{\cosh^4(h\xi + \beta z \sqrt{q})} \leq 1. \quad (1.8)$$

The rest of the paper is organized as follows. In Section 2, we gather some results regarding the directional derivative of the functional $\mathcal{P}$ and some consequences following the first order optimality of the Parisi variational formula. The proof of Theorem 1.1 is presented in Section 3.

2 Preliminary results

Let ξ be standard Gaussian. Let $\alpha \in \text{Pr}[0, 1]$. Conditionally on ξ , let $(X_\alpha^\xi(s))_{0 \leq s \leq 1}$ be the solution to the following stochastic differential equation with boundary condition $X_\alpha^\xi(0) = h\xi$,

$$dX_\alpha^\xi(s) = \beta^2 \alpha(s) \partial_x \Phi_\alpha(s, X_\alpha^\xi(s)) ds + \beta dW(s),$$

where $(W(s))_{s \in [0, 1]}$ is a standard Brownian motion independent of ξ . Set

$$\phi_\alpha(s) = \mathbb{E}(\partial_x \Phi_\alpha(s, X_\alpha^\xi(s)))^2, \quad 0 \leq s \leq 1.$$

The following proposition establishes the directional derivative of $\mathcal{P}$ in terms of ϕ_α .

Proposition 2.1. *For $\alpha_0, \alpha_1 \in \text{Pr}[0, 1]$ and $\theta \in [0, 1]$, set $\alpha_\theta = (1 - \theta)\alpha_0 + \theta\alpha_1$. The right-derivative of $\theta \mapsto \mathcal{P}(\alpha_\theta)$ at zero is given by*

$$\left. \frac{d\mathcal{P}(\alpha_\theta)}{d\theta^+} \right|_{\theta=0} = \frac{\beta^2}{2} \int_0^1 (\alpha_1(s) - \alpha_0(s)) (s - \phi_{\alpha_0}(s)) ds.$$

Furthermore, α_0 is the minimizer of $\min_{\alpha \in \text{Pr}[0, 1]} \mathcal{P}(\alpha)$ if and only if $\left. \frac{d\mathcal{P}(\alpha_\theta)}{d\theta^+} \right|_{\theta=0} \geq 0$ for all $\alpha_1 \in \text{Pr}[0, 1]$.

Proposition 2.2. *If α_0 is the minimizer of $\mathcal{P}$, then every point in the support of α_0 must satisfy $\phi_{\alpha_0}(s) = s$.*

In the case of the SK model with deterministic external field (1.1), the two propositions above were established in Theorem 2 and Proposition 1 in [3], respectively, where the statements are essentially the same as Propositions 2.1 and 2.2 with a replacement of $h\xi$ by h . As the proofs in [3] can be directly applied to Propositions 2.1 and 2.2 with no major changes, we do not reproduce the details here. The following lemma provides a simpler expression for ϕ_α .

Lemma 2.3. *Let $\alpha \in \text{Pr}[0, 1]$. For any $s \in [0, 1]$, we have that*

$$\phi_\alpha(s) = \mathbb{E} \partial_x \Phi_\alpha(s, M(s))^2 \exp \left(\int_0^s (\Phi_\alpha(s, M(s)) - \Phi_\alpha(t, M(t))) \alpha(dt) \right),$$

where $M(s) := h\xi + \beta W(s)$.

Proof. Conditionally on ξ , let $\mathbb{E}^\xi$ be the expectation associated to the probability measure

$$d\mathbb{P}^\xi = \exp \left(- \int_0^1 \beta \alpha(r) \partial_x \Phi(r, X_\alpha^\xi(r)) dW(r) - \frac{1}{2} \int_0^1 \beta^2 \alpha(r)^2 \partial_x \Phi_\alpha(r, X_\alpha^\xi(r))^2 dr \right) d\mathbb{P}.$$

From the Girsanov theorem, under $\mathbb{E}^\xi$,

$$\left(\int_0^s \beta \alpha(r) \partial_x \Phi_\alpha(r, X_\alpha^\xi(r)) dr + W(s) \right)_{0 \leq s \leq 1}$$

is a standard Brownian motion for which we denote by $(W^\xi(s))_{0 \leq s \leq 1}$. Set $M^\xi(s) = h\xi + \beta W^\xi(s)$. Under $\mathbb{E}^\xi$, write

$$\begin{aligned} & \int_0^1 \beta \alpha(r) \partial_x \Phi(r, X_\alpha^\xi(r)) dW(r) + \frac{1}{2} \int_0^1 \beta^2 \alpha(r)^2 \partial_x \Phi_\alpha(r, X_\alpha^\xi(r))^2 dr \\ &= \int_0^1 \beta \alpha(r) \partial_x \Phi(r, X_\alpha^\xi(r)) \left(\beta \alpha(r) \partial_x \Phi_\alpha(r, X_\alpha^\xi(r)) dr + dW(r) \right) \\ & \quad - \frac{1}{2} \int_0^1 \beta^2 \alpha(r)^2 \partial_x \Phi_\alpha(r, X_\alpha^\xi(r))^2 dr \\ &= \int_0^1 \beta \alpha(r) \partial_x \Phi(r, M^\xi(r)) dW^\xi(r) - \frac{1}{2} \int_0^1 \beta^2 \alpha(r)^2 \partial_x \Phi_\alpha(r, M^\xi(r))^2 dr. \end{aligned}$$

Consequently,

$$\begin{aligned} \mathbb{E}[(\partial_x \Phi_\alpha(s, X_\alpha^\xi(s)))^2 | \xi] &= \mathbb{E}^\xi(\partial_x \Phi_\alpha(s, M^\xi(s)))^2 \exp\left(\int_0^1 \beta \alpha(r) \partial_x \Phi_\alpha(r, M^\xi(r)) dW^\xi(r) \right. \\ & \quad \left. - \frac{1}{2} \int_0^1 \beta^2 \alpha(r)^2 \partial_x \Phi_\alpha(r, M^\xi(r))^2 dr\right). \end{aligned} \quad (2.1)$$

Next, note that from Itô's formula and (1.3), for $0 \leq t < s \leq 1$,

$$\begin{aligned} & \Phi_\alpha(s, M^\xi(s)) - \Phi_\alpha(t, M^\xi(t)) \\ &= -\frac{\beta^2}{2} \int_t^s \alpha(r) (\partial_x \Phi_\alpha(r, M^\xi(r)))^2 dr + \beta \int_t^s \partial_x \Phi_\alpha(r, M^\xi(r)) W^\xi(dr). \end{aligned}$$

and then from the Fubini theorem,

$$\begin{aligned} & \int_0^s (\Phi_\alpha(s, M^\xi(s)) - \Phi_\alpha(t, M^\xi(t))) \alpha(dt) \\ &= -\frac{\beta^2}{2} \int_0^s \alpha(r)^2 (\partial_x \Phi_\alpha(r, M^\xi(r)))^2 dr + \beta \int_0^s \alpha(r) \partial_x \Phi_\alpha(r, M^\xi(r)) W^\xi(dr). \end{aligned}$$

Plugging this into (2.1) yields that

$$\begin{aligned} \mathbb{E}[(\partial_x \Phi_\alpha(s, X_\alpha^\xi(s)))^2 | \xi] &= \mathbb{E}^\xi \left[(\partial_x \Phi_\alpha(s, M^\xi(s)))^2 \right. \\ & \quad \left. \exp\left(\int_0^s (\Phi_\alpha(s, M^\xi(s)) - \Phi_\alpha(t, M^\xi(t))) \alpha(dt)\right) \right] \\ &= \mathbb{E} \left[(\partial_x \Phi_\alpha(s, M(s)))^2 \right. \\ & \quad \left. \exp\left(\int_0^s (\Phi_\alpha(s, M(t)) - \Phi_\alpha(t, M(t))) \alpha(dt)\right) \right] | \xi, \end{aligned}$$

where we used that $(M(s))_{0 \leq s \leq 1} \stackrel{d}{=} (M^\xi(s))_{0 \leq s \leq 1}$ conditionally on ξ . Finally, taking expectation in ξ completes our proof. $\square$

While Lemma 2.3 holds for any α , the important case that we shall use in our main proof is when $\alpha = 1_{[q,1]}$ for some $q \in [0,1]$. In this case, one can compute ϕ_α more explicitly. Let z, z' be i.i.d. standard Gaussian independent of ξ . Denote by $\mathbb{E}'$ the expectation with respect to z' only.

Lemma 2.4. Let $\alpha = 1_{[q,1]}$ for some $q \in [0, 1]$. We have that

$$\phi_\alpha(s) = \begin{cases} \mathbb{E} \left(\frac{\mathbb{E}' \tanh^2(h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q}) \cosh(h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q})}{\mathbb{E}' \cosh(h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q})} \right), & \text{if } s \in [q, 1], \\ \mathbb{E} (\mathbb{E}' \tanh(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}))^2, & \text{if } s \in [0, q) \end{cases} \quad (2.2)$$

and

$$\phi'_\alpha(s) = \begin{cases} \beta^2 \mathbb{E} \left(\frac{\mathbb{E}' \cosh^{-3}(h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q})}{\mathbb{E}' \cosh(h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q})} \right), & \text{if } s \in [q, 1], \\ \beta^2 \mathbb{E} (\mathbb{E}' \cosh^{-2}(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}))^2, & \text{if } s \in [0, q). \end{cases} \quad (2.3)$$

Proof. A direct computation gives that

$$\Phi_\alpha(s, x) = \begin{cases} \frac{\beta^2}{2} (1-s) + \log \cosh x, & \text{if } s \in [q, 1], \\ \frac{\beta^2}{2} (1-q) + \mathbb{E} \log \cosh(x + \beta z \sqrt{q-s}), & \text{if } s \in [0, q) \end{cases}$$

and that

$$\partial_x \Phi_\alpha(s, x) = \begin{cases} \tanh x, & \text{if } s \in [q, 1], \\ \mathbb{E} \tanh(x + \beta z \sqrt{q-s}), & \text{if } s \in [0, q). \end{cases}$$

Plugging these along with the assumption $\alpha = 1_{[q,1]}$ into Lemma 2.3 establishes (2.2).

As for (2.3), note that for any twice differentiable function f with $\|f''\|_\infty < \infty$, we can compute by using the Gaussian integration by parts² to obtain

$$\frac{d}{ds} \mathbb{E}' f(z' \sqrt{s-q}) = \frac{1}{2\sqrt{s-q}} \mathbb{E}' f'(z' \sqrt{s-q}) = \frac{1}{2} \mathbb{E}' f''(z' \sqrt{s-q}), \quad \forall s \in (q, 1]. \quad (2.4)$$

Applying this equation and

$$\begin{aligned} (\tanh^2(x) \cosh(x))'' &= \frac{2}{\cosh^3(x)} + \tanh^2(x) \cosh(x), \\ (\cosh(x))'' &= \cosh(x) \end{aligned}$$

to the $\mathbb{E}'$ -expectations in the first equation of (2.2), the first equation of (2.3) follows by a straightforward computation. To obtain the second equation in (2.3), write

$$\mathbb{E} (\mathbb{E}' \tanh(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}))^2 = \mathbb{E} [\tanh(h\xi + \beta z \sqrt{s} + \beta z_1 \sqrt{q-s}) \tanh(h\xi + \beta z \sqrt{s} + \beta z_2 \sqrt{q-s})]$$

for z_1, z_2 i.i.d. standard Gaussian independent of z . From the last equation and $(\tanh(x))' = \cosh^{-2}(x)$, we have

$$\begin{aligned} & \frac{d}{ds} \mathbb{E} (\mathbb{E}' \tanh(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}))^2 \\ &= \frac{\beta}{2} \mathbb{E} \frac{\tanh(h\xi + \beta z \sqrt{s} + \beta z_1 \sqrt{q-s})}{\cosh^2(h\xi + \beta z \sqrt{s} + \beta z_2 \sqrt{q-s})} \left(\frac{z}{\sqrt{s}} - \frac{z_2}{\sqrt{q-s}} \right) \\ &+ \frac{\beta}{2} \mathbb{E} \frac{\tanh(h\xi + \beta z \sqrt{s} + \beta z_2 \sqrt{q-s})}{\cosh^2(h\xi + \beta z \sqrt{s} + \beta z_1 \sqrt{q-s})} \left(\frac{z}{\sqrt{s}} - \frac{z_1}{\sqrt{q-s}} \right). \end{aligned}$$

²Let $Z = (z_1, \dots, z_n)$ be a centered Gaussian random vector and let F be a differentiable function on $\mathbb{R}^n$ with $\sum_{i=1}^n \|\partial_{x_i} F\|_\infty < \infty$. We have that $\mathbb{E} z_1 f(Z) = \sum_{i=1}^n \mathbb{E}[z_1 z_i] \mathbb{E}[\partial_{x_i} f(Z)]$.

Using Gaussian integration by parts yields that

$$\begin{aligned} & \frac{d}{ds} \mathbb{E} \left(\mathbb{E}' \tanh(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}) \right)^2 \\ &= \beta^2 \mathbb{E} \frac{1}{\cosh^2(h\xi + \beta z \sqrt{s} + \beta z_1 \sqrt{q-s}) \cosh^2(h\xi + \beta z \sqrt{s} + \beta z_2 \sqrt{q-s})} \\ &= \beta^2 \mathbb{E} \left(\mathbb{E}' \cosh^{-2}(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}) \right)^2. \end{aligned} \quad \square$$

3 Proof of Theorem 1.1

Throughout the entire proof, we let q be the unique solution to (1.6). First, we assume that (β, h) lies above the AT line, i.e.,

$$\beta^2 \mathbb{E} \frac{1}{\cosh^4(h\xi + \beta z \sqrt{q})} > 1.$$

We show that α_P can not be replica symmetric. If not, then we must have that $\alpha_P = 1_{[q',1]}$ for some $q' \in [0, 1]$ and from Proposition 2.2 and Lemma 2.4, q' must satisfy

$$\mathbb{E} \tanh^2(h\xi + \beta z \sqrt{q'}) = \phi_{\alpha_P}(q') = q'.$$

Since this equation has only one unique solution (see (1.6)), we must have that $q = q'$. Consequently, $\phi_{\alpha_P}(q) = q$ and from (2.3),

$$\phi'_{\alpha_P}(q) = \beta^2 \mathbb{E} \frac{1}{\cosh^4(h\xi + \beta z \sqrt{q})} > 1.$$

If we let $\varepsilon > 0$ be small enough and $\alpha_1 = 1_{[q+\varepsilon,1]}$, then $\phi_{\alpha_P}(s) > s$ for $s \in [q, q+\varepsilon]$ so that from Proposition 2.1,

$$\frac{d}{d\theta^+} \mathcal{P}(\alpha_\theta) \Big|_{\theta=0} = -\frac{\beta^2}{2} \int_q^{q+\varepsilon} (\phi_{\alpha_P}(s) - s) ds < 0$$

and consequently, $1_{[q,1]}$ can not be the optimizer. Hence, outside the AT line, the SK model is not replica symmetric.

Next, we assume that (β, h) lies inside the AT line, i.e., (1.8) holds. We proceed to show that $\alpha_P = 1_{[q,1]}$. To this end, let $\alpha_0 = 1_{[q,1]}$ and we claim that (1.8) implies that $\phi_{\alpha_0}(s) \geq s$ if $s < q$ and $\phi_{\alpha_0}(s) \leq s$ if $s > q$. If this claim is valid, then for any $\alpha_1 \in \text{Pr}[0, 1]$,

$$\frac{d}{d\theta} \mathcal{P}(\alpha_\theta) \Big|_{\theta=0^+} = \frac{\beta^2}{2} \left(\int_0^q \alpha_1(s) (\phi_{\alpha_0}(s) - s) ds + \int_q^1 (\alpha_1(s) - 1) (\phi_{\alpha_0}(s) - s) ds \right) \geq 0.$$

Hence, from Proposition 2.1, $\alpha_0 = 1_{[q,1]}$ is the minimizer of the Parisi formula and this will complete our proof.

We now turn to the proof of our claim. First of all, from (2.3) and the Jensen inequality with respect to $\mathbb{E}'$, for $s \in [0, q]$,

$$\phi'_{\alpha_0}(s) \leq \beta^2 \mathbb{E} \mathbb{E}' \cosh^{-4}(h\xi + \beta z \sqrt{s} + \beta z' \sqrt{q-s}) = \beta^2 \mathbb{E} \cosh^{-4}(h\xi + \beta z \sqrt{q}) \leq 1.$$

Since $\phi_{\alpha_0}(q) = q$ by (2.2), we see that $\phi_{\alpha_0}(s) \geq s$ for $s \in [0, q]$. Next, for $s \in (q, 1)$, recall from the first equation in (2.3) that by denoting $Y = h\xi + \beta z \sqrt{q} + \beta z' \sqrt{s-q}$,

$$\phi'_{\alpha_0}(s) = \beta^2 \mathbb{E} \frac{\mathbb{E}' \cosh^{-3} Y}{\mathbb{E}' \cosh Y}$$




Figure 2: This is a simulation of $s \in [q, 1] \mapsto \beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{s})$ with $\beta = 4.05$, $h = 5$, and $q \approx 0.909$. The blue line is the level $\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{q})$ and the red line is q . It can be seen that $\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{s})$ lies above $\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{q})$ for $s \in [q, 1]$ and above 1 for s large enough.

Write $\cosh^{-3} x = (\cosh^{-4} x)(\cosh x)$. Since $\cosh^{-4} x$ is decreasing and $\cosh x$ is increasing for $x > 0$, applying the FKG inequality³ implies

$$\begin{aligned} \mathbb{E}' \cosh^{-3} Y &= \mathbb{E}' \cosh^{-3} |Y| \\ &\leq \mathbb{E}' \cosh^{-4} |Y| \cdot \mathbb{E}' \cosh |Y| \\ &= \mathbb{E}' \cosh^{-4} Y \cdot \mathbb{E}' \cosh Y. \end{aligned}$$

Thus, for any $q < s \leq 1$,

$$\begin{aligned} \phi'_{\alpha_0}(s) &\leq \beta^2 \mathbb{E} \cosh^{-4} Y \\ &= \beta^2 \mathbb{E} \cosh^{-4}(h\xi + \beta z \sqrt{s}) \\ &= \beta^2 \mathbb{E} \cosh^{-4}(z \sqrt{h^2 + \beta^2 s}) \\ &\leq \beta^2 \mathbb{E} \cosh^{-4}(z \sqrt{h^2 + \beta^2 q}) \\ &= \beta^2 \mathbb{E} \cosh^{-4}(h\xi + \beta z \sqrt{q}) \leq 1, \end{aligned}$$

where the second inequality is valid since $\cosh^{-4} x$ is even and decreasing in $x > 0$. Consequently, $\phi'_{\alpha_0}(s) - 1 \leq 0$ for any $q < s \leq 1$. Since $\phi_{\alpha_0}(q) = q$, we deduce that $\phi_{\alpha_0}(s) \leq s$ for $q < s \leq 1$. This finishes the proof of our claim.

Remark 3.1. In the case of the SK model with non-random external field (1.1), the corresponding $\phi_\alpha(s)$ for $\alpha = 1_{[q,1]}$ and q satisfying (1.4) is the same as in Lemma 2.4 except that $h\xi$ is replaced by h . The same argument in Theorem 1.1 enables us to show that the Parisi formula can not be solved by the replica symmetric solution if (β, h) lies outside the AT line. However, if (β, h) lies inside the AT line, while it can still be shown that $\phi_\alpha(s) \geq s$ for all $s \in [0, q]$, it is unclear why $\phi_\alpha(s) \leq s$ for all $s \in [q, 1]$. In this case, one can still use the FKG inequality to obtain that for $q \leq s \leq 1$,

$$\phi'_\alpha(s) \leq \beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{s}),$$

but a numerical simulation, see Figure 2, suggests that the following does not always hold:

$$\beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{s}) \leq \beta^2 \mathbb{E} \cosh^{-4}(h + \beta z \sqrt{q})$$

for all $q \leq s \leq 1$.

³The FKG inequality states that if X is a random variable and f, g are both nondecreasing functions with $\text{Var}(f(X)) < \infty$ and $\text{Var}(g(X)) < \infty$, then $\mathbb{E}f(X)g(X) \geq \mathbb{E}f(X)\mathbb{E}g(X)$.

References

- [1] J. F. L. Almeida and D. J. Thouless, *Stability of the Sherrington-Kirkpatrick solution of a spin glass model*, J. Phys. A: Math. Gen. **11** (1978), 983–990.
- [2] A. Auffinger and W.-K. Chen, *The Parisi formula has a unique minimizer*, Comm. Math. Phys. **335** (2015), no. 3, 1429–1444. MR3320318
- [3] W.-K. Chen, *Variational representations for the Parisi functional and the two-dimensional Guerra-Talagrand bound*, Ann. Probab. **45** (2017), no. 6A, 3929–3966. MR3729619
- [4] F. Guerra, *Sum rules for the free energy in the mean field spin glass model*, Fields Institute Communications **30** (2001), no. 161. MR1867553
- [5] F. Guerra, *Broken replica symmetry bounds in the mean field spin glass model*, Comm. Math. Phys. **233** (2003), no. 1, 1–12. MR1957729
- [6] A. Jagannath and I. Tobasco, *A dynamic programming approach to the Parisi functional*, Proc. Amer. Math. Soc. **144** (2016), no. 7, 3135–3150. MR3487243
- [7] A. Jagannath and I. Tobasco, *Some properties of the phase diagram for mixed p -spin glasses*, Probab. Theory Related Fields **167** (2017), no. 3–4, 615–672. MR3627426
- [8] M. Mézard, G. Parisi, and M. A. Virasoro, *Spin glass theory and beyond*, World Scientific Lecture Notes in Physics, vol. 9, World Scientific Publishing Co., Inc., Teaneck, NJ, 1987. MR1026102
- [9] D. Panchenko, *The Sherrington-Kirkpatrick model*, Springer Monographs in Mathematics, Springer, New York, 2013. MR3052333
- [10] G. Parisi, *Infinite number of order parameters for spin-glasses*, Phys. Rev. Lett. **43** (1979), 1754–1756. MR0702601
- [11] G. Parisi, *A sequence of approximate solutions to the SK model for spin glasses*, J. Phys. A. **13** (1980), no. L-115.
- [12] D. Sherrington and S. Kirkpatrick, *Solvable model of a spin glass*, Phys. Rev. Lett. **35** (1972), 1792–1796.
- [13] M. Talagrand, *The Parisi formula*, Ann. of Math. (2) **163** (2006), no. 1, 221–263. MR2195134
- [14] M. Talagrand, *Mean field models for spin glasses. Volume I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 54, Springer-Verlag, Berlin, 2011, Basic examples. MR2731561
- [15] M. Talagrand, *Mean field models for spin glasses. Volume II*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 55, Springer, Heidelberg, 2011, Advanced replica-symmetry and low temperature. MR3024566
- [16] F. Toninelli, *About the Almeida-Thouless transition line in the Sherrington-Kirkpatrick mean-field spin glass model*, Europhys. Lett. **60** (2020), 764–767.

Acknowledgments. The author thanks the anonymous referees for some helpful comments regarding the presentation of this paper. Special thanks are due to Si Tang for running the simulations in Figures 1 and 2.

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>