

# Total Synthesis of the Chlorinated Pentacyclic Indole Alkaloid (+)-Ambiguine G

Lingbowei Hu and Viresh H. Rawal\*

Department of Chemistry, University of Chicago, Chicago, 5735 South Ellis Avenue, Chicago, Illinois 60637, United States

**ABSTRACT:** Reported herein is the total synthesis of (+)-ambiguine G, the first member of the chlorinated pentacyclic ambiguines to yield to chemical synthesis. The synthesis is accomplished through a convergent strategy that proceeds in 10 steps from (S)-carvone oxide. Pivotal to the concise route is the successful realization of a [4+3] cycloaddition that conjoins two easily synthesized components of the carbon framework of the natural product. Also featured in the synthesis is the efficient, diastereoselective construction of a key vinylated chloroketone and the unprecedented, one-pot reduction-elimination-oxidation sequence that transforms an enone to an advanced hydroxylated-diene intermediate.

The ambiguines are a subset of the large halapindole family of more than 80 cyanobacteria metabolites that also includes the fischerindoles and welwitindolinones.<sup>1,2</sup> The first of the ambiguines were identified by Smitka and Moore in 1992 while screening fungicidal extracts primarily from the terrestrial cyanophytes *Fischerella ambiguua*.<sup>3</sup> Although the full bioactivity profiles of these alkaloids have yet to be fully assessed, several members have displayed useful properties. Of note, ambiguine I isonitrile (4) is not only a stronger antibacterial and antifungal agent than established clinical agents, but it is also a potent NF- $\kappa$ B inhibitor ( $IC_{50}=30$  nM), with cytotoxic activity against HT-29 colon cancer and MCF-7 breast cancer cells (Figure 1).<sup>4,5</sup> Structurally, all ambiguines contain the tetracyclic core of the halapindoles, but thirteen of the eighteen members possess an additional, seven-membered ring that connects the indole to the distal six-membered ring. Furthermore, over half of the ambiguines possess a chlorine atom at C13, rendering them significantly more difficult as targets for synthesis.<sup>6</sup> The intricate polycyclic architecture and the unpredictable reactivity of the pentacyclic ambiguines present a significant challenge to the state-of-the-art of synthesis, one that went unmet despite numerous efforts over many years.<sup>7,8</sup> It was only in 2019 that the first pentacyclic member of this family of natural products succumbed to synthesis. Two contemporaneous publications, one by Sarpong and coworkers and the other us, presented distinctly different strategies for the synthesis of ambiguine P (7).<sup>9</sup> We now report the total synthesis of (+)-ambiguine G (8), the first member of the chlorinated pentacyclic ambiguines to yield to chemical synthesis.<sup>10</sup>

Our strategy to ambiguine G (8) is intimated in the retrosynthesis shown in Scheme 1 and is enabled by three key insights. First, the chlorine atom at C13 would be installed early in the synthesis to avoid potential rearrangements induced by the adjacent vinyl group in a rigid, advanced intermediate, as observed in the welwitindolinones.<sup>11</sup> Second, through advanced model studies, we determined that the desired [4+3] cycloaddition reaction, which was unrealized in our ambiguine synthesis,<sup>9b</sup> could be rendered efficacious by using an alkoxy diene instead of a siloxyl diene. Lastly, a removable functionality at C15 with low tendency to leave as a cation was deemed





Figure 1. Selected members of the ambiguine natural products.



essential for the late-stage functionalization of C23. Otherwise, installation of the nitrile group at that position, whether through site-selective, direct electrophilic cyanation or via halogenation followed by transition metal catalyzed coupling with cyanide, was expected to be complicated by untoward reactions (e.g., proton loss from C15).

## Scheme 1. Retrosynthetic Analysis



Our synthesis of ambiguine G (**8**) commenced with the preparation of chloroketone **17**, a functionalized six-membered ring unit common to numerous members of the halapindole family. While seemingly simple, ketone **17** presents unique challenges, and the only reported synthesis of it requires 10 steps.<sup>12</sup> In devising an alternate route to **17**, we planned to install the chloride via a stereoinvertive-deoxychlorination of a hydroxy ketone precursor **16**, which mapped nicely over (*S*)-carvone oxide, provided a vinyl group could be introduced from the side opposite to that of the isopropenyl unit. In simple carvone derivatives, however, it is well documented that carbon electrophiles are introduced at C2 *cis* to the isopropenyl unit due to stereoelectronic factors. Therefore, installation of a substituent *trans* to the isopropenyl unit would require harnessing the chirality of a preexisting functionality on a carvone derivative, thereby overriding the intrinsic diastereoselectivity. With this recognition, we examined different strategies with the goal of preinstalling a hydroxyl group at C3 and using it to direct a vinylation reaction. Success was achieved through the method reported by Coltrart and coworkers.<sup>13</sup> Addition of vinylmagnesium bromide to tosylhydrazone **15**, which was easily prepared from commercially available (*S*)-carvone oxide,<sup>14</sup> followed directly by copper (II) chloride mediated hydrolysis of the hydrazone provided ketone **16** with nearly complete diastereoselectivity (Scheme 2).<sup>15</sup> The vinyl addition took place as desired from the side away from the isopropenyl unit, ostensibly directed by coordination of the Grignard reagent with the alkoxide intermediate.<sup>13a,c</sup> To our knowledge, this epoxyhydrazone-mediated directed introduction of a carbon substituent *α*-to a carbonyl group has not been utilized in natural product synthesis.<sup>16</sup> Conversion of the vinyl-alcohol product **16** to chloroketone **17** was accomplished using N-chlorosuccinimide and PPh<sub>3</sub> with complete stereoinversion at the chlorine attaching carbon. The stereoretentive chlorination product was not observed.


### Scheme 2. Synthesis of Chloroketone **17**



With a practical, two-step synthesis of chloroketone **17** in hand, we focused our attention on assembling the carbon framework of the natural product via the [4+3] cycloaddition reaction.<sup>17</sup> Although our published route to ambiguine P (**7**) was inspired by this cycloaddition as the key step, in practice it proved unsuccessful.<sup>9b</sup> Rather than forging two C-C bonds to form the seven-membered ring, the reaction gave what is effectively the Friedel-Crafts alkylation product of the silyl enol ether and the benzylic cation (*cf.* **11** + **12**, Scheme 1). We reasoned that the reaction may proceed in a stepwise manner, wherein the labile silyl group falls off after formation of the first C-C bond to give an “interrupted” [4+3] product. Based on this hypothesis, we examined the cycloaddition reaction of ethoxy diene **18**, which was easily synthesized from ketone **17** via triflation followed by Stille cross-coupling. To our delight, treatment of diene **18** and indolic silyl ether **19**<sup>18</sup> with TMSOTf promoted the desired [4+3] reaction to afford tetracycle **20** cleanly, with no evidence of the Friedel-Crafts reaction product (Scheme 3).

An efficient, two-pot reaction sequence was developed to transform tetracycle **20** to pentacyclic alcohol **23**. First, the [4+3] cycloadduct was treated with BF<sub>3</sub>•OEt<sub>2</sub> to annulate the final ring through a Friedel-Crafts reaction. Subsequent addition of TBAF to quench the Lewis acid followed by DDQ oxidized the intermediate ketone to enone **10** in good yield. In the next protocol, DIBAL reduction of the carbonyl group and elimination of the resulting alkoxide using Et<sub>2</sub>AlCl produced a mixture of the conjugated diene **21** and the cross-conjugated diene **22**, greatly favoring the latter. The high regioselectivity for diene **22** likely reflects the stereoelectronic preference for elimination of the axially-oriented C15 proton over the C23 proton in the conformationally rigid pentacyclic framework. The facile deprotonation at C15 also complicates the required electrophilic functionalization at C23 on diene **21** and necessitates the installation of a blocking group at C15. Fortunately, the C15 carbon of diene **22** was found to be unexpectedly electron rich, making it susceptible to air oxidation. Based on this realization, we developed a highly efficient procedure wherein after DIBAL reduction and Et<sub>2</sub>AlCl-mediated elimination, KHMDS was added to deprotonate the indole nitrogen and then the reaction mixture was exposed to air. Gratifyingly, the intermediate indole anion reacted with oxygen at C15 and the resulting hydroperoxide was reduced by the P(OMe)<sub>3</sub> present to provide alcohol **23** in good yield and excellent diastereoselectivity (13:1).

**Scheme 3. Total Synthesis of (+)-Ambiguine G**



Having installed the hydroxyl group at C15, and thereby fore stalled side reactions arising from proton loss from that position, the next task was to introduce the nitrile group at C23. Although methods for the direct introduction the nitrile group proved unsuccessful, treatment of diene **23** with *N*-bromosuccinimide selectively brominated the distal carbon of the conjugated diene without touching the vinyl group. A subsequent tautomerization in the presence of pyridinium *p*-toluenesulfonate produced alkenyl bromide **24** in high yield. The nitrile group was then introduced in good yield by a palladium catalyzed coupling reaction.<sup>19</sup> Having served its function, the hydroxyl group was removed under ionic hydrogenation conditions ( $\text{BF}_3\text{-OEt}_2$  and  $\text{Et}_3\text{SiH}$ ) to afford (+)-ambiguine G (**8**), which was formed as a single diastereomer.

In summary, we have completed the enantiospecific synthesis of (+)-ambiguine G (**8**), a chlorinated member of the ambiguine family of indole alkaloids. The synthesis is accomplished through a convergent strategy that proceeds in 10 synthetic operations from (*S*)-carvone oxide and demonstrates: (1) the construction of a key chlorine substituted cyclohexanone precursor through an alkoxide directed vinylation reaction, (2) the rapid assembly of the core skeleton of the natural product by a [4+3] cycloaddition reaction, and (3) the unprecedented, one-pot reduction-elimination-oxidation sequence that transforms an enone intermediate to a pivotal hydroxy diene. The efficiency of the route is expected to provide ready access to more intricate members of the pentacyclic ambiguines, as well as their analogs.

## ASSOCIATED CONTENT

### SUPPORTING INFORMATION

This supporting information is available free of charge via the Internet at <http://pubs.acs.org>.

Experimental procedures and spectroscopic data for all new compounds (PDF).

### AUTHOR INFORMATION

#### Corresponding Author

Viresh H. Rawal – Department of Chemistry, University of Chicago, Chicago, Illinois 60637, United States; orcid.org/0000-0003-4606-0239; Email: vrawal@uchicago.edu

#### Authors

Lingbowei Hu – Department of Chemistry, University of Chicago, Chicago, Illinois 60637, United States; orcid.org/0000-0001-8525-9983

### Author Contributions

All authors have given approval to the final version of the manuscript.

### Funding Sources

National Science Foundation (NSF-1566402).

### Notes

The authors declare no competing financial interest.

### ACKNOWLEDGMENT

Financial support from the National Science Foundation (NSF-1566402) is gratefully acknowledged.

### REFERENCES

(1) Bhat, V.; Davé, A.; MacKay, J. A.; Rawal, V. H. The chemistry of halapindoles, fischerindoles, ambiguines and welwitindolinones. in *The Alkaloids, Chem. Biol.* **2014**, 73, 65-160.

(2) (a) Smith, C. D.; Zilfou, J. T.; Stratmann, K.; Patterson, G. M. L.; Moore, R. E. Welwitindolinone analogues that reverse P-glycoprotein mediated multiple drug resistance. *Mol. Pharmacol.* **1995**, 47, 241–247. (b) Zhang, X.; Smith, C. D. Microtubule effects of welwistatin, a cyanobacterial indolinone that circumvents multiple drug resistance. *Mol. Pharmacol.* **1996**, 49, 288–294. (c) Acuña, U. M.; Mo, S.; Zi, J.; Orjala, J.; Carcache de Blanco, E. J. Hapalindole H induces apoptosis as an inhibitor of NF-κB and affects the intrinsic mitochondrial pathway in PC-3 androgen-insensitive prostate cancer cells. *Anticancer Res.* **2018**, 38, 3299-3307. (d) Chilczuk, T.; Steinborn, C.; Breinlinger, S.; Zimmermann-Klemd, A. M.; Huber, R.; Enke, H.; Enke, D.; Nidermeyer, T. H. J.; Gründemann, C. Hapalindoles from the cyanobacterium *Hapalsiphon* sp. inhibit T cell proliferation. *Planta Med.* **2020**, 86, 96-103. (e) Kim, H.; Lantvit, D.; Hwang, C. H.; Kroll, D. J.; Swanson, S. M.; Franzblau, S. G.; Orjala, J. Indole alkaloids from two cultured cyanobacteria, *Westiellopsis* sp. and *Fischerella muscicola*. *Bioorg. Med. Chem.* **2012**, 20, S290-5295.

(3) Smitka, T. A.; Bonjouklian, R.; Doolin, L.; Jones, N. D.; Deeter, J. B.; Yoshida, W. Y.; Prinsep, M. R.; Moore, R. E.; Patterson, G. M. L. Ambiguine isonitriles, fungicidal hapalindole-type alkaloids from three genera of blue-green algae belonging to the Stigonemataceae. *J. Org. Chem.* **1992**, 57, 857-861.

(4) Raveh, A.; Carmeli, S. Antimicrobial ambiguines from the cyanobacterium *Fischerella* sp. collected in Israel. *J. Nat. Prod.* **2007**, 70, 196–201.

(5) Acuña, U. M.; Zi, J.; Orjala, J.; Carcache de Blanco, E. J.. Ambiguine I isonitrile from *Fischerella ambigua* induces caspase-independent cell death in MCF-7 hormone dependent breast cancer cells. *Int. J. Cancer Res.* **2015**, 49, 1655-1662.

(6) Walton, K.; Gantar, M.; Gibbs, P. D. L.; Schmale, M. C.; Berry, J. P. Indole alkaloids from *Fischerella* inhibit vertebrate development in the zebrafish (*Danio rerio*) embryo model. *Toxins* **2014**, 6, 3568-3581.

(7) Synthetic efforts toward seven-membered ring containing ambiguines: (a) Chandra, A.; Viswanathan, R.; Johnston, J. N. Synthesis of the ABC-and D-ring systems of the indole alkaloid ambiguine G. *Org. Lett.* **2007**, 9, 5027-5029. (b) Rafferty, R. J.; Williams, R. M. Synthetic studies on the ambiguine family of alkaloids: Construction of the ABCD ring system. *Tetrahedron Lett.* **2011**, 52, 2037-2040.

(8) Synthesis of tetracyclic ambiguines lacking the seven-membered ring: (a) Baran, P. S.; Maimone, T. J.; Richter, J. M. Total synthesis of marine natural products without using protecting groups. *Nature* **2007**, 446, 404-408.

(b) Rafferty, R. J.; Williams, R. M. Formal synthesis of hapalindole O and synthetic efforts towards hapalindole K and ambiguine A. *Heterocycles* **2012**, 86, 219-231. (c) Maimone, T. J.; Ishihara, Y.; Baran, P. S. Scalable total syntheses of (-)-hapalindole U and (+)-ambiguine H. *Tetrahedron* **2015**, 71, 3652-3665. (d) Sahu, S.; Das, B.; Maji, M. S. Stereodivergent total synthesis of hapalindoles, fischerindoles, hapalonamide H, and ambiguine H alkaloids by developing a biomimetic, redox-neutral, cascade Prins-type cyclization. *Org. Lett.* **2018**, 20, 6485-6489.

(9) (a) Johnson, R. E.; Ree, H.; Hartmann, M.; Lang, L.; Sawano, S.; Sarpong, R. Total synthesis of pentacyclic (-)-ambiguine P using sequential indole functionalizations. *J. Am. Chem. Soc.* **2019**, 141, 2233-2236. (b) Xu, J.; Rawal, V. H. Total synthesis of (-)-ambiguine P. *J. Am. Chem. Soc.* **2019**, 141, 4820-4823.

(10) Huber, U.; Moore, R. E.; Patterson, G. M. L. Isolation of a nitrile-containing indole alkaloid from the terrestrial blue-green alga *Hapalosiphon delicatulus*. *J. Nat. Prod.* **1998**, 61, 1304-1306.

(11) (a) Bhat, V.; MacKay, J. A.; Rawal, V. H. Lessons learned while traversing the welwitindolinone alkaloids obstacle course. *Tetrahedron* **2011**, 67, 10097-10104. (b) Reyes, J. R.; Xu, J.; Kobayashi, K.; Bhat, V.; Rawal, V. H. Total synthesis of (-)-N-methylwelwitindolinone B isothiocyanate. *Angew. Chem., Int. Ed.* **2017**, 56, 9962-9966.

(12) Fukuyama, T.; Chen, X. Stereocontrolled synthesis of (-)-hapalindole G. *J. Am. Chem. Soc.* **1994**, 116, 3125-3126.

(13) (a) Uteuliyev, M. M.; Nguyen, T. T.; Colart, D. M. Diastereoselective addition of Grignard reagents to  $\alpha$ -epoxy N-sulfonyl hydrazones. *Nat. Chem.* **2015**, 7, 1024-1027. For similar transformations using organocuprates, see: (b) Corey, E. J.; Melvin, L. S. Jr.; Haslanger, M. F. A novel  $\alpha$ -alkylation of  $\alpha,\beta$ -epoxy ketones. *Tetrahedron Lett.* **1975**, 36, 3117-3120. (c) Fuchs, P. L.  $\alpha$ -Arylation of  $\alpha,\beta$ -unsaturated ketones. Utilization of the  $\alpha$ -epoxytosylhydrazone functional group as a  $\Delta^2$ -enonium synthon. *J. Org. Chem.* **1976**, 41, 2935-2937. (d) Stork, G.; Ponaras, A. A.  $\alpha$ -Alkylation and arylation of  $\alpha,\beta$ -unsaturated ketones. *J. Org. Chem.* **1976**, 41, 2937-2939.

(14) (*S*)-Carvone oxide is commercially available from Ambeed, Toronto Research Chemicals etc. It is also easily prepared from (*S*)-carvone through epoxidation: Weinstabl, H.; Gaich, T.; Mulzer, J. Application of the Rodriguez-Pattenden photo-ring contraction: Total synthesis and configurational reassignment of 11-gorgiacerol and 11-epigorgiacerol. *Org. Lett.* **2012**, 14, 2834-2837.

(15) Attempted isolation of the intermediate tosylhydrazone **15**, which degraded slowly at room temperature, gave irreproducible yields, whereas direct addition of vinylmagnesium bromide to a solution of the hydrazone

(19) Böttcher, C.; Zeyat, G.; Ahmed, S. A.; Irran, E.; Cordes, T.; Elsner, C.; Zinth, W.; Ruech-Braun, K. Synthesis of novel photochromic pyrans via palladium-mediated reactions. *Beilstein J. Org. Chem.* **2009**, 5, No. 25.

condensation mixture provided the vinyl addition product in more consistent yields. Additionally, it was observed that the hydrazone could be hydrolyzed in the same pot in good yield.

(16) For related chemistry using oximes, see: Sawada, D.; Shibasaki, M. Enantioselective total synthesis of epothilone A using multifunctional asymmetric catalyses. *Angew. Chem. Int. Ed.* **2000**, 39, 209-213.

(17) Selected examples of [4+3] cycloaddition reactions in total synthesis: (a) Zhang, J.; Li, L.; Wang, Y.; Wang, W.; Xue, J.; Li, Y. A novel, facile approach to frondosin B and 5-*epi*-liphagal via a new [4 + 3]-cycloaddition. *Org. Lett.* **2012**, 14, 4528-4530. (b) Granger, B. A.; Jewett, I. T.; Butler, J. D.; Hua, B.; Knezevic, C. E.; Parkinson, E. I.; Hergenrother, P. J.; Martin, S. F. Synthesis of ( $\pm$ )-actinophyllic acid and analogs: Applications of cascade reactions and diverted total synthesis. *J. Am. Chem. Soc.* **2013**, 135, 12984-12986. (c) Granger, B. A.; Jewett, I. T.; Butler, J. D.; Martin, S. F. Concise total synthesis of ( $\pm$ )-actinophyllic acid. *Tetrahedron* **2014**, 70, 4094-4104. (d) Laplace, D. R.; Verbraeken, B.; Hecke, K. V.; Winne, J. M. Total synthesis of (+/-)-frondosin B and (+/-)-5-*epi*-liphagal by using a concise (4+3) cycloaddition approach. *Chem. Eur. J.* **2014**, 20, 253-262.

(18) Compound **19** was prepared in two steps from commercially available 2-ethoxycarbonylindole. See supporting information for synthetic procedures and characterizations.

**Table of content:**

