



# Chromosome Evolution in the Family Solanaceae

Rocío Deanna<sup>1,2†</sup>, María Cristina Acosta<sup>1</sup>, Marisel Scaldaferrero<sup>1</sup> and Franco Chiarini<sup>1\*†</sup>

<sup>1</sup> Instituto Multidisciplinario de Biología Vegetal (CONICET-UNC), Córdoba, Argentina, <sup>2</sup> Department of Ecology and Evolutionary Biology, University of Colorado at Boulder, Boulder, CO, United States

This review summarizes and discusses the knowledge of cytogenetics in Solanaceae, the tomato family, its current applications, and prospects for making progress in fundamental systematic botany and plant evolution. We compile information on basic chromosome features (number, size, morphology) and molecular cytogenetics (chromosome banding and rDNA patterns). These data were mapped onto the Solanaceae family tree to better visualize the changes in chromosome features and evaluate them in a phylogenetic context. We conclude that chromosomal features are important in understanding the evolution of the family, especially in delimiting clades, and therefore it is necessary to continue producing this type of data. The potential for future applications in plant biology is outlined. Finally, we provide insights into understanding the mechanisms underlying Solanaceae's diversification that could substantially contribute to developing new approaches for future research.

**Keywords:** comparative methods, cytogenetics, karyotype, nightshades, phylogeny

## OPEN ACCESS

### Edited by:

Peter Poczai,  
University of Helsinki, Finland

### Reviewed by:

Sonia Garcia,  
Botanical Institute of Barcelona,  
CSIC, Spain  
Jaume Pellicer,  
Spanish National Research Council  
(CSIC), Spain

### \*Correspondence:

Franco Chiarini  
chiarini@imbiv.unc.edu.ar;  
franco.e.chiarini@gmail.com

<sup>†</sup> These authors have contributed  
equally to this work and share first  
authorship

### Specialty section:

This article was submitted to  
Plant Systematics and Evolution,  
a section of the journal  
Frontiers in Plant Science

**Received:** 30 September 2021

**Accepted:** 22 December 2021

**Published:** 28 January 2022

### Citation:

Deanna R, Acosta MC,  
Scaldaferrero M and Chiarini F (2022)  
Chromosome Evolution in the Family  
Solanaceae.  
Front. Plant Sci. 12:787590.  
doi: 10.3389/fpls.2021.787590

## INTRODUCTION

Chromosomes provide information for inferring phylogenetic relationships, since they are hereditary elements of the whole nuclear genome and discrete hereditary units of mutation. Within taxa, they may vary in number, size, morphology, and staining properties (Weiss-Schneeweiss and Schneeweiss, 2013). Chromosome number has always been a common character employed, since it is the most easily obtained information and the only one that is known for most plant groups. Karyotypes represent an important aspect in plant speciation since chromosomal differences establish immediate postzygotic crossing barriers. They provide diagnostic characters for plant systematics and evolution that are usually expected to be congruent with clade divergence (e.g., Baltisberger and Hörandl, 2016). In Solanaceae, chromosomes have played a relevant role: they have been useful to delimit taxa (Camadro et al., 2012), for genetic studies with model species (Blakeslee et al., 1920; Avery et al., 1959), and for the selection of cultivars in economically important species (Pringle and Murray, 1992). With the advent of molecular phylogenetic techniques, chromosome data acquired a new value, to the point that within the family, there is a major lineage named as the “ $x = 12$  clade,” whose members share such cytological synapomorphy (Olmstead et al., 2008). This review summarizes and discusses the knowledge of cytogenetics in Solanaceae, its current applications and prospects for making progress in fundamental systematic botany and plant evolution. Available chromosome data were reassessed by employing them in ancestral state reconstruction and thus revealing how many times traits changed over evolutionary time, inferring the degree of convergence, and suggesting relationships between chromosomes and other Solanaceae characteristics. Continuous cytogenetic characters (C, mean chromosome length; r,

mean arm ratio or  $r$  index; TL, total haploid chromosome length; and  $m\_ratio$ , the ratio between the number of metacentric to the total number of chromosomes) were mapped and plotted onto the family phylogeny of Särkinen et al. (2013) with modifications by Dupin et al. (2017), after removing those species without data using the `drop.tip` function of the package `{ape}` in R (Paradis et al., 2004). The ancestral states were estimated assuming that the species evolved under a Brownian model and the mapping was performed using the `ContMap` function of the package `{phytools}` in R version 3.4.2 (Revell, 2012; **Supplementary Figures 1–4**). All the methodologies to gather the data are in **Supplementary Table 2**.

## CHROMOSOME COUNTS

Knowing the basic number  $x$  for all Angiosperms and its clades has been a recurring question (Darlington, 1956). Initially, it was considered of little taxonomic utility at high hierarchical levels (orders, families) due to homoplasy. For Solanales, Raven (1975) proposed  $x = 7$  while  $x = 12$  may have derived from the tetraploid level ( $n = 14$ ) by aneuploid reduction in the early history of Solanaceae. Acosta et al. (2006) suggested  $x = 11$  as the ancestral number that best fits the available phylogenetic hypotheses (Olmstead et al., 1999). Recently, models of chromosome evolution were customized using statistical frameworks, allowing to leverage other tools, like Ancestral State Reconstruction (ASR, FitzJohn, 2012), and to infer character evolution across phylogenetic trees (Revell, 2012). The compilation and analysis of data in a phylogenetic context allow us to recalculate the first estimates and reconstruct the history of the changes. Although the basic number  $x$  is homoplastic in Solanaceae, it is more conserved than other karyotype features, which makes it useful to delimit taxa at lower hierarchical levels: tribes, subtribes, genera (e.g., Rodríguez et al., 2020). The availability of chromosome data is essential to delve into these analyses, but the information is still scattered and fragmentary (**Supplementary Tables 1, 2**). Among the five largest genera of Solanaceae, the best studied (*Solanum*) has 48% of its species counted. There is a bias toward groups of species with economically valuable representatives (e.g., *Capsicum*, *Nicotiana*), while small clades and monotypic genera are karyologically unknown (**Supplementary Tables 1, 2**). Applying model approaches to chromosome number would clarify this topic (Glick and Mayrose, 2014), although such methodologies have difficulties given the size of the matrix for the entire family (**Supplementary Table 1**) and discussing such downsides is beyond the scope of this work.

Solanaceae have undergone whole genome duplications occurring near the time of its origin, which could have contributed to the rise of key traits and drove species diversification (Schranz et al., 2012). Polyploidy, a phenomenon constantly reviewed (e.g., Van de Peer et al., 2021), because of the advantages it may confer, has been a subject since the beginning of cytogenetics in Solanaceae (e.g., Blakeslee et al., 1920) up to the present (e.g., Hijmans et al., 2007; Chiarini et al., 2018). Nowadays, the availability of data makes it possible to analyze polyploidy throughout the evolutionary history of the family and


relate it to colonization of new habitats, speciation events and to the rising of adaptive traits (Hijmans et al., 2007; Scaldaferrero et al., 2012; Zenil-Ferguson et al., 2018, 2019; McCarthy et al., 2019).

## KARYOTYPES

Blakeslee's pioneering work on *Datura* mutants (Blakeslee et al., 1920, and subsequent papers) allowed to identify pairs of homologs by their shape, and correlate alterations in meiotic behavior to exomorphology. The chromosome morphology observable with classical technique is still essential to detect rearrangements involved in speciation, for example, detection of allopolyploids in *Nicotiana* (Goodspeed, 1954), Robertsonian translocations in *Solanum* (Chiarini and Bernardello, 2006) and in *Chamaesaracha* (Rodríguez et al., 2020), and centric fission in *Capsicum* (Jarret et al., 2019). In contrast, uniform chromosome morphology may be the explanation for crossability among Iochrominae species (Deanna et al., 2018) and suggests the phenomenon of "karyotype orthoselection" in *Lycieae* (Bernardello et al., 2008). Karyotype data integrated with molecular phylogenies provide characters for Solanaceae systematics and evolution (e.g., *Nierembergia*: Acosta et al., 2016; *Jaborosa*, Chiarini et al., 2017; *Iochroma*, Deanna et al., 2018; *Physalis*, Rodríguez et al., 2020). However, the distribution of cytological studies is uneven among the genera of the family (**Supplementary Table 2**). We registered 420 species of Solanaceae with some kind of data (karyotype formulae, idiograms, asymmetry indices), which makes up a percentage of ca. 15% of the family. The most frequently informed characters are the following.

## Chromosome Size

Chromosome size (usually expressed as length in  $\mu\text{m}$ ) is useful to single out individuals, samples, populations or species. DNA content is directly related to it, so the alleged factors affecting the former may indirectly affect the latter as the speed of DNA replication and the duration of the life cycle. Also, long chromosomes would undergo a higher number of chiasmata than small ones (Turney et al., 2004) and low recombination due to small chromosomes would be compensated by high ploidy levels (Nakazato et al., 2006). In Solanaceae, average chromosome size varies from ca. 1  $\mu\text{m}$  (in *Solanum* and *Atropa*) to 6.5–11.5  $\mu\text{m}$  (in *Cestrum* and the Cyphomandra clade of *Solanum*, **Figure 1**, **Supplementary Figure 1**, and **Supplementary Table 3**) although most species have small to medium-sized chromosomes, with an overall mean chromosome length of  $2.95 \pm 1.78$  (**Supplementary Table 3**). In order to better visualize the changes in chromosome size and evaluate them in a phylogenetic context, we mapped C and TL onto the Solanaceae family tree (**Supplementary Tables 3, 4**). Although these two variables are homoplastic (**Supplementary Figures 1, 2**) and have no significant phylogenetic signal, it is still useful to characterize some clades which are equivalent to medium-hierarchy taxonomic levels. *Cestreae*, *Cyphomandra* and *Capsicum* clades are outstanding because they have high C, conspicuously different from the remaining



**FIGURE 1** | Photomicrographs of mitotic metaphases of Solanaceae, illustrating the chromosome variety found in the family. **(A)** *Cestrum buxifolium* Kunth ( $2n = 16$ ,  $c = 11.5 \mu\text{m}$ ,  $r = 1.4$ ). **(B)** *Solanum lidii* Sunding ( $2n = 24$ ,  $c = 1.3 \mu\text{m}$ ,  $r = 1.6$ ). **(C)** *Lochroma edule* S. Leiva ( $2n = 24$ ,  $c = 4.0$ ,  $r = 1.2$ ). **(D)** *Physalis lagascae* Roem. & Schult. ( $2n = 24$ ,  $c = 2.5 \mu\text{m}$ ,  $r = 2.8$ ). Panels **(A,B)** with classical staining, panels **(C,D)** fluorescence *in situ* hybridization with 5S (red) and 18-5.8-26S (green) rDNA probes. All at the same scale. Bar =  $10 \mu\text{m}$ .

clades (**Supplementary Figure 1**). These synapomorphic large chromosomes arose three times separately and are an interesting case to study phenomena such as “genome obesity” (Bennetzen and Kellogg, 1997), increases in DNA content through the incorporation of heterochromatin, repetitive sequences, and retrotransposons (Fregonezi et al., 2007; Miguel et al., 2012; Scaldaferrero et al., 2013).

## Chromosome Morphology

### Asymmetry

There are a variety of indices to estimate chromosomal asymmetry and their accuracy is arguable (Peruzzi and Eroglu, 2013). Here we have considered the  $r$  index because it is the most widespread, intuitive and easy to calculate, and also the most frequently informed (**Supplementary Table 5**). Asymmetry indices try to provide an idea of the general morphology of the karyotype through a single number, but logically not regarding the morphology of each chromosome. When all pairs of chromosomes undergo similar changes at the same time (like in *Nicotiana*, Kovarik et al., 2004), other measurements are

needed to distinguish individual chromosome rearrangements. In a general survey of Solanaceae, Badr et al. (1997) reported values of  $r$  from 1.17 to 2.78, whereas in *Solanum* it ranges from 1.19 to 3.71 with an overall mean of 1.64 (Chiarini et al., 2018). Thus, karyotype asymmetry is not uniform in the family. According to ASR, symmetrical karyotypes are plesiomorphic (**Supplementary Figure 3**), being asymmetry a synapomorphy of some clades (*Physalis*, *Nicotiana*), while others would have conserved symmetry (*Lycium*). This supports the idea that in Solanaceae, shifts in chromosome morphology are frequent when considering broad frames of time (Wu and Tanksley, 2010; Chiarini et al., 2018).

### Karyotype Formulae

The karyotype formula is a mean of expressing the result of a measurement process of the chromosome complement, i.e., the whole set of chromosomes in a nucleus, and it gives an overall idea of its morphology. Morphology provides the first clue to single out chromosome pairs (e.g., heteromorphic sex chromosomes). Chromosome morphology can be related to the

plant mating system: in *Rumex*, a species with sex chromosomes, the suppression of recombination varies between metacentric and acrocentric chromosomes (e.g., Rifkin et al., 2021). Also, bimodal karyotypes visible with classical technique evidence the presence of two genomes (Brandham and Doherty, 1998). According to the widespread Levan's classification (Levan et al., 1964), chromosomes can be, in increasing order of asymmetry, *m*, *sm*, *st*, and *t*. On this basis, karyotypes of Solanaceae are symmetrical, since only 17 out of the total species (4%) with known karyotypes contain 1–3 *t* chromosomes, while 100 (24%) present 1–7 *sm* chromosomes. Most Solanaceae have more than half of chromosomes per complement in the *m* and/or *sm* categories. These features are useful to characterize entire clades within the family: there are groups with a majority of *m* chromosomes, e.g., *Lycium* (Stiefkens et al., 2020), but there are also groups in which complements include *st* and *t* chromosomes, e.g., *Nicotiana*, *Capsicum*, *Jaborosa*, *Physalis* (Figure 1D), *Hyoscyamus*, *Nierembergia* (Sheidai et al., 1999; Moscone et al., 2006; Scaldaferrero et al., 2013; Acosta et al., 2016; Chiarini et al., 2017; Rodríguez et al., 2020) and in some clades of *Solanum* (Bernardello et al., 1994; Acosta et al., 2005). Some *Nicotiana* species have karyotypes mostly with *st* chromosomes (Villa, 1984) and *Leptoglossis linifolia* has mostly *sm* chromosomes (Acosta et al., 2016).

In Solanaceae, establishing the direction of karyotype evolution is unlikely, as many reversals might have occurred (Stace, 2000), and karyotype asymmetry might be a temporary state rather than an evolutionary end (Mandáková and Lysak, 2008). Comparing karyotype formulae is arguable because it is difficult to establish homologies of chromosome pairs among different species. Here, we calculated the proportion of *m* chromosomes for each nightshade with the available karyotype formula and mapped this ratio onto the phylogeny (Figure 2, Supplementary Figure 4, and Supplementary Table 6). This reconstruction suggests the most probable ancestral karyotypes had  $\pm$  80% of *m* chromosomes (i.e., symmetric karyotype). Formulae with few or no *m* chromosomes are autapomorphic (e.g., *Nicotiana plumbaginifolia*) while complete formulae with *m* chromosomes is a synapomorphy for certain clades (*Lycium*, Iochrominae, Figure 1C). Transitions between karyotype formulae can be interpreted as evidence of chromosome rearrangements. Particularly, formulae with *st* or *t* chromosomes are seemingly the result of a deletion or translocation of the entire or part of one arm (Weiss-Schneeweiss and Schneeweiss, 2013) or centric fissions (Moscone et al., 2006; Jarret et al., 2019).

## Chromosome Banding


Banding techniques are the next step in identifying chromosome pairs in a complement. They leverage on the physical properties of chromatin, using dyes that differentially stain euchromatin and fractions within heterochromatin (e.g., AT- or GC-rich), which are then visualized as bands across the chromatids. In Solanaceae, 218 spp. out of approx. 2,800 (ca. 7.7%) have been studied with any banding technique. The largest number of species studied belong to *Solanum* (69 out of 1,238 species, 5.6%), followed by *Capsicum* (25 out of 41, 60.9%) and *Lycium*

(24 out of 105, 22.9%) (Supplementary Table 2). These last two genera are also the best studied in relative terms, together with *Nierembergia* (71.4% of 21 spp.), *Jaborosa* (65.2% of 23 spp.) and *Iochroma* (22.6% of 31 spp.) (Supplementary Table 2). A variety of banding techniques have been essayed in Solanaceae to answer a range of questions: silver impregnation detected active nucleolar organizing regions (NORs) (e.g., in *Cestrum*, Berg and Greilhuber, 1993; *Capsicum*, Scaldaferrero et al., 2016; *Solanum*, Miguel et al., 2012; *Deprea*, Deanna et al., 2014); Giemsa C-banding revealed differences between taxa and contributed to taxonomic grouping of *Capsicum* (Moscone et al., 1993); Fluorescent banding, mostly with the double staining CMA/DAPI technique (Schweizer, 1976) demonstrated that chromosomes are mostly composed of non-coding chromatin, and also revealed strong phylogenetic signal, defining specific patterns in different clades (Acosta et al., 2016). CMA/DAPI technique helped to detect variability despite the morphological uniformity of the chromosomes, provided information on genetic variation at a population level regarding speciation, and revealed AT- and GC-rich regions of B chromosomes. Together with FISH (see below), this technique showed that B chromosomes possess nucleolar activity and nucleolar competition with the A chromosomes (Acosta and Moscone, 2011; Acosta et al., 2016; Montechiari et al., 2020).

Heterochromatin amount (H.a.) is a value often reported (usually as a percentage). It exhibits a remarkable variation among species of the same genus, for example in *Solanum* it varies from 1.86 to 35.43% (Brasileiro-Vidal et al., 2009; Chiarini et al., 2018). The data on H.a. have served to discuss whether the increases in genome size (either measured visually as TL, or as DNA content through flow cytometry) are due to increases in the amount of one or another fraction of chromatin (e.g., Karsburg et al., 2009). In some clades, H.a. is positively correlated with karyotype length (Moscone et al., 2006; Scaldaferrero et al., 2013), while in others there is negative or no correlation, which is unexpected because heterochromatin is considered as an additional component of the genome (Pringle and Murray, 1993; Acosta et al., 2012, 2016). Heterochromatin patterns are more variable in some genera (*Jaborosa*, *Solanum*) than in other members of the  $x = 12$  clade (e.g., *Lycium* and *Sclerophylax*), where heterochromatin is scarce and restricted to the NORs (Lujea and Chiarini, 2017; Stiefkens et al., 2020). Variable patterns might be evidence of intense chromosomal rearrangements (Evtushenko et al., 2016) associated with diversification and colonization of new habitats, since they function as species barriers (Hughes and Hawley, 2009).

## rDNA and Repetitive DNA Patterns

Ribosomal DNAs are fundamental components of all cell types. In most plants, 5S and 18S-5.8S-26S rDNAs are present in high copy numbers to satisfy the cellular requirement for ribosomes. These rDNAs, highly conserved in plants, are the commonest markers for FISH (Heslop-Harrison and Schwarzacher, 2011). They are repetitive, tandemly arranged and generally clustered at different loci, either with a spatial separation or also linked in a single unit. FISH enables



direct visualization of rDNA loci on the chromosomes and determination of their rearrangements and organization (e.g., Srebniak et al., 2002; Melo and Guerra, 2003; Lou et al., 2010). Thus, homologous chromosomes in a complement can be identified and related species can be compared, answering evolutionary questions. Number, position and organization of rDNAs have been suitably employed in Solanaceae as systematic and evolutionary approaches (Figures 1C,D and Supplementary Table 2), for example in *Nierembergia* (Acosta et al., 2016), *Jaborosa* (Chiarini et al., 2017) and *Iochroma* (Deanna et al., 2018). Genera with economically important species (*Capsicum*, *Nicotiana*) have received special attention (Kitamura et al., 2005; Scaldaferrero et al., 2016). Within the largest genera, *Lycium* is the most studied (Stiefkens et al., 2020), revealing uniformity of patterns, but little is known about the rest of the family. Chromosomal rearrangements between *Solanum* crops and several related species have been studied using tomato and potato bacterial artificial chromosomes (BACs) in multiple FISH essays, providing support for grouping of species into sections and suggesting interspecific hybridization events (Doganlar et al., 2002; Frary et al., 2016). Another type of repetitive DNA revealed by FISH are the telomeric sequences. Most plants present the Arabidopsis-type telomeres, but some Solanaceae are exceptional in lacking such sequences: the characterization of unusual telomeres in the three genera of Cestreae has shed light on patterns of telomere

evolution, maintenance and function (Sýkorová et al., 2003; Peška et al., 2015).

## CYTOGENETIC AND PHYLOGENETIC GAPS

There is a growing interdependence between chromosomal data and molecular phylogeny. Cytogenetics and molecular phylogeny feed into each other, providing evidence in both ways. To some extent, Solanaceae ASRs allow reconstructing the cytogenetic data when there is molecular data (e.g., Rodríguez et al., 2020), but the reverse is obviously not possible. Thus, the largest information gaps are found in *Cestrum*, (165 species without any data of a total of 233, neither counts nor sequences), *Lycianthes* (116 species without data of about 161), *Jaltomata* (32 species of 73), *Physalis* (47 of 109) and *Browallia* (18 species without data of about 22). Of the ca. 2,900 species in the family, 494 (17%) possess chromosome counts and are included in the phylogeny of Särkinen et al. (2013), while of these, only 207 (7% of the family) also possess karyotypes. At the same time, there are 137 species for which there are counts but no sequences, and of these, 40 also have karyotypes, but these data cannot be analyzed in a molecular phylogenetic context (Supplementary Tables 7, 8). Important gaps in both cytogenetics and phylogeny still wait to be filled: in Goetzeoideae, a clade of which nothing is known;

in Solandreae, only chromosome studies in one *Trianaea* species and two *Dyssochroma*; *Cuatresia*, genus of Physalideae with an important number of species about which nothing is known (Supplementary Table 8).

## CYTOGENETICS AND GENOMICS IN SOLANACEAE

Extensive pairwise comparative mapping studies have been performed for several major solanaceous crops relative to tomato (Tanksley et al., 1988, 1992; Livingstone et al., 1999; Doganlar et al., 2002; Wu et al., 2009; Wu and Tanksley, 2010). By using single-copy conserved ortholog sets (COSII), the outcomes of chromosomal evolution in the Solanaceae over 30 million years were deduced, estimating the rates and timing of chromosomal rearrangements as well as calculating the age of ancestral species and predicting their genome features (Wu and Tanksley, 2010). Thus, studies integrating cytogenetics and genomics, though focused on main crops and close relatives, generate knowledge that can be extrapolated to the rest of the species of the family.

Recently, Cytogenetics developed feedback with Genomics: the availability of species with complete chloroplast genome sequenced and some with complete nuclear genome, allows the finding of new markers for hybridization techniques. In return, these cytogenetic data become valuable information on chromosome identification and genome organization, enabling the spatial location of specific, single, or low-copy sequences. Next-generation sequencing speeds up cytogenetic research on Solanaceae. In species with available transcriptomes, like *S. dulcamara*, new markers like SNP SSR, AFLP and CAPS were developed, allowing the construction of a genome-wide genetic linkage map (D'Agostino et al., 2013). Based on gene orthologs, the markers were anchored to the genome of related *Solanum* species (tomato, potato and eggplant), revealing both conserved and novel chromosomal rearrangements. Estimates of the evolutionary moment of rearrangements were possible and showed that chromosomal breakpoints are regularly re-used.

Also, advances in NGS, such as the study of repetitive DNA, contributed to the design of more specific oligo-probes for FISH (Buggs et al., 2012), and hence bringing new chances to improve our understanding of systematics and genome organization at different taxonomic hierarchies. More recently, there has been progress in the phylogenomics of Solanaceae, particularly in some clades within the family, through the use of nuclear target capture and high-throughput sequencing of transcriptomic data (Gates et al., 2018; Poczai et al., 2018; Gagnon et al., 2021). These approaches provide more comprehensive and better resolved phylogenies which can be used in the study of chromosome evolution.

## REFERENCES

Acosta, M. C., Bernardello, G., Guerra, M., and Moscone, E. A. (2005). Karyotype analysis in several South American species of *Solanum* and *Lycianthes rantonnei* (Solanaceae). *Taxon* 54, 713–723. doi: 10.2307/25065428

## CONCLUDING REMARKS

- Evolutionary trends: The ancestor of the family had a karyotype formula with 80% of m chromosomes, each about 2  $\mu\text{m}$  in size. Increased asymmetry and size are synapomorphies of some clades that arose in independent events.
- Although homoplastic at a higher hierarchical level, karyotype traits are useful to delimit taxa at lower levels: tribes, subtribes, genera.
- Important gaps in both cytogenetics and phylogeny still wait to be filled.
- Potential for future applications: once there is a genomic phylogenetic backbone of Solanaceae, including divergence time estimates, adding chromosomes to it would be illuminating, because relationships between major clades of the family and Solanoideae are still poorly resolved using Sanger sequence data alone.

## AUTHOR CONTRIBUTIONS

FC and RD conceived the ideas, designed the methodology, analyzed the data, and wrote the manuscript. FC, RD, MS, and MA collected the data. All authors contributed to the article and approved the submitted version.

## FUNDING

This work was supported by National Agency for Scientific Promotion and Technological grant PICT 2017-2370 and PICT 2016-1525 (FONCyT), the National Science Foundation (NSF-DEB 1902797), and National Council for Scientific and Technological Research (CONICET). Publication of this article was funded by the University of Colorado Boulder Libraries Open Access Fund.

## ACKNOWLEDGMENTS

We would like to thank Stacey Smith for advice during the comparative analyses performed.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2021.787590/full#supplementary-material>

Acosta, M. C., Guerra, M., and Moscone, E. A. (2012). Karyological relationships among some South American species of *Solanum* (Solanaceae) based on fluorochrome banding and nuclear DNA amount. *Plant Syst. Evol.* 298, 1547–1556. doi: 10.1007/s00606-012-0657-x

- Acosta, M. C., and Moscone, E. A. (2011). B Chromosomes in *Nierembergia aristata* (Solanaceae): Nucleolar Activity and Competition with the A Chromosomes. *Cytogenet. Gen. Res.* 132, 105–112. doi: 10.1159/000320705
- Acosta, M. C., Moscone, E. A., and Cocucci, A. A. (2016). Using chromosomal data in the phylogenetic and molecular dating framework: karyotype evolution and diversification in *Nierembergia* (Solanaceae) influenced by historical changes in sea level. *Plant Biol.* 18, 514–526. doi: 10.1111/plb.12430
- Acosta, M. C., Ordoñez, A. D. V., Cocucci, A. A., and Moscone, E. A. (2006). Chromosome reports in South American Nicotianeae (Solanaceae) with particular reference to *Nierembergia*. *Ann. Mo. Bot. Gard.* 93, 634–646. doi: 10.3417/0026-6493(2006)93[634:crisan]2.0.co;2
- Avery, A. G., Satina, S., and Rietsema, J. (1959). *Blakeslee: The Genus Datura*. New York: Ronald Press Co.
- Badr, A., Khalifa, S. F., Aboel-Atta, A. I., and Abou-El-Enain, M. M. (1997). Chromosomal criteria and taxonomic relationships in the Solanaceae. *Cytologia* 62, 103–113. doi: 10.1508/cytologia.62.103
- Baltisberger, M., and Hörandl, E. (2016). Karyotype evolution supports the molecular phylogeny in the genus *Ranunculus* (Ranunculaceae). *Perspect. Plant Ecol. Evol. Syst.* 18, 1–14. doi: 10.1016/j.ppees.2015.11.001
- Bennetzen, J. L., and Kellogg, E. A. (1997). Do plants have a one-way ticket to genomic obesity? *Plant Cell* 9, 1509–1514. doi: 10.2307/3870439
- Berg, C., and Greilhuber, J. (1993). Cold-sensitive chromosome regions and heterochromatin in *Cestrum* (Solanaceae): *C. strigillatum*, *C. fasciculatum*, and *C. elegans*. *Plant Syst. Evol.* 185, 133–151. doi: 10.1007/BF00937725
- Bernardello, G., Stiefkens, L., and Las Peñas, M. L. (2008). Karyotype studies in *Grabowskia* and *Phrodus* (Solanaceae). *Plant Syst. Evol.* 275, 265–269. doi: 10.1007/s00606-008-0060-9
- Bernardello, L. M., Heiser, C. B., and Piazzano, M. (1994). Karyotypic studies in *Solanum* section *Lasiocarpa* (Solanaceae). *Am. J. Bot.* 81, 95–103. doi: 10.1002/j.1537-2197.1994.tb15414.x
- Blakeslee, A. F., Belling, J., and Farnham, M. E. (1920). Chromosomal duplication and Mendelian phenomena in *Datura* mutants. *Science* 52, 388–390. doi: 10.1126/science.52.1347.388
- Brandham, P. E., and Doherty, M. J. (1998). Genome size variation in the Alooaceae, an angiosperm family displaying karyotypic orthoselection. *Ann. Bot.* 82, 67–73. doi: 10.1006/anbo.1998.0742
- Brasileiro-Vidal, A. C., Melo-Oliveira, M. B., Carvalho, G. M. G., and Guerra, M. (2009). Different chromatin fractions of tomato (*Solanum lycopersicum* L.) and related species. *Micron* 40, 851–859. doi: 10.1016/j.micron.2009.06.004
- Buggs, R. J., Renny-Byfield, S., Chester, M., Jordon-Thaden, I. E., Viccini, L. F., Chamala, S., et al. (2012). Next-generation sequencing and genome evolution in allopolyploids. *Amer. J. Bot.* 99, 372–382. doi: 10.3732/ajb.1100395
- Camadro, E. L., Erazzu, L. E., Maune, J. F., and Bedogni, M. C. (2012). A genetic approach to the species problem in wild potato. *Plant Biol.* 14, 543–554. doi: 10.1111/j.1438-8677.2012.00563.x
- Chiarini, F., and Bernardello, G. (2006). Karyotype studies in South American species of *Solanum* subgen. *Leptostemonum* (Solanaceae). *Plant Biol.* 8, 486–493. doi: 10.1055/s-2006-923859
- Chiarini, F., Moreno, N., Moré, M., and Barboza, G. (2017). Chromosomal changes and recent diversification in the Andean genus *Jaborosa* (Solanaceae). *Bot. J. Linn. Soc.* 183, 57–74. doi: 10.1111/boj.12493
- Chiarini, F., Sazatornil, F., and Bernardello, G. (2018). Data reassessment in a phylogenetic context gives insight into chromosome evolution in the giant genus *Solanum* (Solanaceae). *Syst. Biodivers.* 16, 397–416. doi: 10.1080/14772000.2018.1431320
- D'Agostino, N., Golas, T., Van de Geest, H., Bombarely, A., Dawood, T., Zethof, J., et al. (2013). Genomic analysis of the native European *Solanum* species, *S. dulcamara*. *BMC Gen.* 14:356. doi: 10.1186/1471-2164-14-356
- Darlington, C. D. (1956). *Chromosome Botany*. London, United Kingdom: George Allen & Unwin
- Deanna, R., Barboza, G. E., and Scaldaferrero, M. A. (2014). First karyological report in *Larnax* and *Deprea* (Solanaceae). *Austral. J. Bot.* 62, 251–261. doi: 10.1071/BT14041
- Deanna, R., Smith, S. D., Särkinen, T., and Chiarini, F. (2018). Patterns of chromosomal evolution in the florally diverse Andean clade Iochrominae (Solanaceae). *Perspect. Plant Ecol. Evol. Syst.* 35, 31–43. doi: 10.1016/j.ppees.2018.09.004
- Doganlar, S., Frary, A., Daunay, M. C., Lester, R. N., and Tanksley, S. D. (2002). A comparative genetic linkage map of eggplant (*Solanum melongena*) and its implications for genome evolution in the Solanaceae. *Genetics* 161, 1697–1711. doi: 10.1093/genetics/161.4.1697
- Dupin, J., Matzke, N. J., Särkinen, T., Knapp, S., Olmstead, R. G., Bohs, L., et al. (2017). Bayesian estimation of the global biogeographical history of the Solanaceae. *J. Biogeogr.* 44, 887–899. doi: 10.1111/jbi.12898
- Evtushenko, E. V., Levitsky, V. G., Elisafenko, E. A., Gunbin, K. V., Belousov, A. I., Šafař, J., et al. (2016). The expansion of heterochromatin blocks in rye reflects the co-amplification of tandem repeats and adjacent transposable elements. *BMC Gen.* 17:337. doi: 10.1186/s12864-016-2667-5
- FitzJohn, R. G. (2012). Diversitree: comparative phylogenetic analyses of diversification in R. *Methods Ecol. Evol.* 3, 1084–1092. doi: 10.1111/j.2041-210X.2012.00234.x
- Frary, A., Doganlar, S., and Frary, A. (2016). “Synteny among solanaceae genomes,” in *The Tomato Genome*, eds M. Causse, J. Giovannoni, M. Bouzayen and M. Zouine (Berlin, Heidelberg: Springer), 217–243. doi: 10.1007/978-3-662-53389-5\_12
- Fregonezi, J. N., Vilas-Boas, L. A., Fungaro, M. H. P., Gaeta, M. L., and Vanzela, A. L. L. (2007). Distribution of a Ty3/gypsy-like retroelement on the A and B-chromosomes of *Cestrum strigilatum* Ruiz & Pav. and *Cestrum intermedium* Sendtn. (Solanaceae). *Genet. Mol. Biol.* 30, 599–604. doi: 10.1590/S1415-4752007000400017
- Gagnon, E., Hilgenhof, R., Orejuela, A., McDonnell, A., Sablok, G., Aubriot, X., et al. (2021). Phylogenomic data reveal hard polytomies across the backbone of the large genus *Solanum* (Solanaceae). *bioRxiv* [Preprint]. doi: 10.1101/2021.03.25.436973
- Gates, D. J., Pilson, D., and Smith, S. D. (2018). Filtering of target sequence capture individuals facilitates species tree construction in the plant subtribe Iochrominae (Solanaceae). *Mol. Phylogenet. Evol.* 123, 26–34. doi: 10.1016/j.ympev.2018.02.002
- Glick, L., and Mayrose, I. (2014). ChromEvol: assessing the pattern of chromosome number evolution and the inference of polyploidy along a phylogeny. *Mol. Biol. Evol.* 31, 1914–1922. doi: 10.1093/molbev/msu122
- Goodspeed, T. H. (1954). *The Genus Nicotiana; Origins, Relationships And Evolution Of Its Species In The Light Of Their Distribution, Morphology And Cytogenetics*. Waltham, Mass: Chronica Botanica Co.
- Heslop-Harrison, J. S., and Schwarzscher, T. (2011). Organisation of the plant genome in chromosomes. *Plant J.* 66, 18–33. doi: 10.1111/j.1365-313X.2011.04544.x
- Hijmans, R. J., Gavrilenko, T., Stephenson, S., Bamberg, J., Salas, A., and Spooner, D. M. (2007). Geographical and environmental range expansion through polyploidy in wild potatoes (*Solanum* section *Petota*). *Glob. Ecol. Biogeogr.* 16, 485–495. doi: 10.1111/j.1466-8238.2007.00308.x
- Hughes, S. E., and Hawley, R. S. (2009). Heterochromatin: a rapidly evolving species barrier. *PLoS Biol.* 7:e1000233. doi: 10.1371/journal.pbio.1000233
- Jarret, R. L., Barboza, G. E., da Costa Batista, F. R., Berke, T., Chou, Y. Y., Hulse-Kemp, A., et al. (2019). *Capsicum*—An abbreviated compendium. *J. Am. Soc. Hortic. Sci.* 144, 3–22. doi: 10.21273/JASHS04446-18
- Karsburg, I. V., Carvalho, C. R., and Clarindo, W. R. (2009). Identification of chromosomal deficiency by flow cytometry and cytogenetics in mutant tomato (*Solanum lycopersicum*, Solanaceae) plants. *Aust. J. Bot.* 57, 444–449. doi: 10.1071/BT08223
- Kitamura, S., Tanaka, A., and Inoue, M. (2005). Genomic relationships among *Nicotiana* species with different ploidy levels revealed by 5S rDNA spacer sequences and FISH/GISH. *Genes Genet. Syst.* 80, 251–260. doi: 10.1266/ggs.80.251
- Kovarik, A., Matyasek, R., Lim, K. Y., Skalická, K., Koukalova, B., Knapp, S., et al. (2004). Concerted evolution of 18–5.8–26S rDNA repeats in *Nicotiana* allotetraploids. *Biol. J. Linn. Soc.* 8, 615–625. doi: 10.1111/j.1095-8312.2004.00345.x
- Levan, A., Fredga, L., and Sandberg, A. (1964). Nomenclature for centromeric position on chromosomes. *Hereditas* 52, 201–220. doi: 10.1111/j.1601-5223.1964.tb01953.x
- Livingstone, K. D., Lackney, V. K., Blauth, J. R., Van Wijk, R. I. K., and Jahn, M. K. (1999). Genome mapping in *Capsicum* and the evolution of genome structure in the Solanaceae. *Genetics* 152, 1183–1202. doi: 10.1093/genetics/152.3.1183

- Lou, Q., Iovene, M., Spooner, D. M., Buell, C. R., and Jiang, J. (2010). Evolution of chromosome 6 of *Solanum* species revealed by comparative fluorescence in situ hybridization mapping. *Chromosoma* 119, 435–442. doi: 10.1007/s00412-010-0269-6
- Lujea, N. C., and Chiarini, F. E. (2017). Differentiation of *Nolana* and *Sclerophylax* (Solanaceae) by means of heterochromatin and rDNA patterns. *N. Z. J. Bot.* 55, 163–177. doi: 10.1080/0028825X.2016.1269812
- Mandáková, T., and Lysak, M. A. (2008). Chromosomal phylogeny and karyotype evolution in  $x = 7$  crucifer species (Brassicaceae). *Plant Cell* 20, 2559–2570. doi: 10.1105/tpc.108.062166
- McCarthy, E. W., Landis, J. B., Kurti, A., Lawhorn, A. J., Chase, M. W., Knapp, S., et al. (2019). Early consequences of allopolyploidy alter floral evolution in *Nicotiana* (Solanaceae). *BMC Plant Biol.* 19:162. doi: 10.1186/s12870-019-1771-5
- Melo, N. F., and Guerra, M. (2003). Variability of the 5S and 45S rDNA sites in *Passiflora* L. species with distinct base chromosome numbers. *Ann. Bot.* 92, 309–316. doi: 10.1093/aob/mcg138
- Miguel, V., Acosta, M. C., and Moscone, E. A. (2012). Karyotype analysis in two species of *Solanum* (Solanaceae) Sect. *Cyphomandropsis* based on chromosome banding. *N. Z. J. Bot.* 50, 217–225. doi: 10.1080/0028825X.2012.662904
- Montechiari, K. A., González, M. L., Yañez Santos, A. M., Hajduczyk Rutz, J. L., and Urdampilleta, J. D. (2020). Structure, behaviour and repetitive DNA of B-chromosomes in *Cestrum nocturnum* (Solanaceae). *Plant Biosyst.*, 29–37. doi: 10.1080/11263504.2018.1559249
- Moscone, E. A., Lambrou, M., Hunziker, A. T., and Ehrendorfer, F. (1993). Giemsa C-banded karyotypes in *Capsicum* (Solanaceae). *Plant Syst. Evol.* 186, 213–229. doi: 10.1007/BF00940799
- Moscone, E. A., Scaldaferrro, M. A., Grabile, M., Cecchini, N. M., Sánchez García, Y., Jarret, R., et al. (2006). “The evolution of chili peppers (*Capsicum*-Solanaceae): a cytogenetic perspective,” in *VI International Solanaceae Conference: Genomics Meets Biodiversity*, (Leuven: International Society for Horticultural Science) 745, 137–170. doi: 10.17660/ActaHortic.2007.745.5
- Nakazato, T., Jung, M. K., Housworth, E. A., Rieseberg, L. H., and Gastony, G. J. (2006). Genetic map-based analysis of genome structure in the homosporous fern *Ceratopteris richardii*. *Genetics* 173, 1585–1597. doi: 10.1534/genetics.106.055624
- Olmstead, R. G., Bohs, L., Migid, H. A., Santiago-Valentin, E., Garcia, V. F., and Collier, S. M. (2008). A molecular phylogeny of the Solanaceae. *Taxon* 57, 1159–1181. doi: 10.1002/tax.574010
- Olmstead, R. G., Sweere, J. A., Spangler, R. E., Bohs, L., and Palmer, J. D. (1999). “Phylogeny and provisional classification of the Solanaceae based on chloroplast DNA,” in *Solanaceae IV: advances in biology and utilization*, eds M. Nee, D. E. Symon, R. N. Lester, and J. P. Jessop (Kew: Royal Botanic Gardens), 257–274.
- Paradis, E., Claude, J., and Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289–290. doi: 10.1093/bioinformatics/btg412
- Peruzzi, L., and Eroglu, H. E. (2013). Karyotype asymmetry: again, how to measure and what to measure? *Comp. Cytogenet.* 7, 1–9. doi: 10.3897/compcytogen.v7i1.4431
- Peška, V., Fajkus, P., Fojtová, M., Dvořáčková, M., Hapala, J., Dvořáček, V., et al. (2015). Characterisation of an unusual telomere motif (TTTTTTAGGG)n in the plant *Cestrum elegans* (Solanaceae), a species with a large genome. *Plant J.* 82, 644–654. doi: 10.1111/tj.12839
- Poczai, P., Amirouze, A., Hyvönen, J. T., and Sablok, G. (2018). “Chloroplast and nuclear phylogenomics reveals concordant phylogenetic structure in wild tomatoes (*Solanum* sect. *Lycopersicon*, Solanaceae),” in *The 15th Solanaceae Conference: Applied Genomics, Accelerated Breeding, Gene Targeting*, (Chiang Mai: University of Helsinki)
- Pringle, G. J., and Murray, B. G. (1992). Polyploidy and aneuploidy in the tamarillo, *Cyphomandra betacea* (Cav.) Sendt. (Solanaceae). I. Spontaneous polyploidy and features of the euploids. *Plant Breed.* 108, 132–138. doi: 10.1111/j.1439-0523.1992.tb00112.x
- Pringle, G. J., and Murray, B. G. (1993). Karyotypes and C banding patterns in species of *Cyphomandra* Mart. ex Sendtn. (Solanaceae). *Bot. J. Linn. Soc.* 111, 331–342. doi: 10.1006/bojil.1993.1023
- Raven, P. H. (1975). The bases of angiosperm phylogeny: cytology. *Ann. Mo. Bot. Gard.* 62, 724–764. doi: 10.2307/2395272
- Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223. doi: 10.1111/j.2041-210X.2011.00169.x
- Rifkin, J. L., Beaudry, F. E., Humphries, Z., Choudhury, B. I., Barrett, S. C., and Wright, S. I. (2021). Widespread recombination suppression facilitates plant sex chromosome evolution. *Mol. Biol. Evol.* 38, 1018–1030. doi: 10.1093/molbev/msaa271
- Rodríguez, J., Deanna, R., and Chiarini, F. (2020). Karyotype asymmetry shapes diversity within the physaloids (Physalidinae, Physalidae, Solanaceae). *Syst. Biodivers.* 19, 168–185. doi: 10.1080/14772000.2020.1832156
- Särkinen, T., Bohs, L., Olmstead, R. G., and Knapp, S. (2013). A phylogenetic framework for evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC Evol. Biol.* 13, 1–15. doi: 10.1186/1471-2148-13-214
- Scaldaferrro, M. A., Grabile, M., and Moscone, E. A. (2013). Heterochromatin type, amount and distribution in wild species of chili peppers (*Capsicum*-Solanaceae). *Genet. Res. Crop Evol.* 60, 693–709. doi: 10.1007/s10722-012-9867-x
- Scaldaferrro, M. A., Romero da Cruz, M. V., Cecchini, N. M., and Moscone, E. A. (2016). FISH and AgNor-mapping of the 45S and 5S rRNA genes in wild and cultivated *Capsicum* species (Solanaceae). *Genome* 59, 95–113. doi: 10.1139/gen-2015-0099
- Scaldaferrro, M., Chiarini, F., Santiñaque, F. F., Bernardello, G., and Moscone, E. A. (2012). Geographical pattern and ploidy levels of the weed *Solanum elaeagnifolium* (Solanaceae) from Argentina. *Genet. Res. Crop Evol.* 59, 1833–1847. doi: 10.1007/s10722-012-9807-9
- Schranz, M. E., Mohammadin, S., and Edger, P. P. (2012). Ancient whole genome duplications, novelty and diversification: the WGD Radiation Lag-Time Model. *Curr. Opin. Plant Biol.* 15, 147–153. doi: 10.1016/j.pbi.2012.03.011
- Schweizer, D. (1976). Reverse fluorescent chromosome banding with chromomycin and DAPI. *Chromosoma* 58, 307–324. doi: 10.1007/BF00292840
- Sheidai, M., Mosallanejad, M., and Khatamsaz, M. (1999). Karyological studies in *Hyoscyamus* species of Iran. *Nord. J. Bot.* 19, 369–374. doi: 10.1111/j.1756-1051.1999.tb01130.x
- Srebnia, M., Rasmussen, O., and Maluszynska, J. (2002). Cytogenetic analysis of an asymmetric potato hybrid. *J. Appl. Genet.* 43, 19–32.
- Stace, C. A. (2000). Cytology and cytogenetics as a fundamental taxonomic resource for the 20th and 21st centuries. *Taxon* 49, 451–477. doi: 10.2307/1224344
- Stiefkens, L., Las Peñas, M. L., Levin, R. A., Miller, J. S., and Bernardello, G. (2020). Chromosome evolution in the cosmopolitan genus *Lycium* (Solanaceae). *Taxon* 69, 124–141. doi: 10.1002/tax.12205
- Sýkorová, E., Lim, K. Y., Chase, M. W., Knapp, S., Leitch, I. J., Leitch, A. R., et al. (2003). The absence of Arabidopsis-type telomeres in *Cestrum* and closely related genera *Vestia* and *Sessea* (Solanaceae): first evidence from eudicots. *Plant J.* 34, 283–291. doi: 10.1046/j.1365-313x.2003.01731.x
- Tanksley, S. D., Bernatzky, R., Lapitan, N. L., and Prince, J. P. (1988). Conservation of gene repertoire but not gene order in pepper and tomato. *Proc. Natl. Acad. Sci.* 85, 6419–6423. doi: 10.1073/pnas.85.17.6419
- Tanksley, S. D., Ganai, M. W., Price, J. P., de Vicente, M. C., Bonierbale, M. W., Broun, P., et al. (1992). High density molecular linkage maps of the tomato and potato genomes; biological inferences and practical applications. *Genetics* 132, 1141–1160. doi: 10.1093/genetics/132.4.1141
- Turney, D., de los Santos, T., and Hollingsworth, N. M. (2004). Does chromosome size affect map distance and genetic interference in budding yeast? *Genetics* 168, 2421–2424. doi: 10.1534/genetics.104.033555
- Van de Peer, Y., Ashman, T. L., Soltis, P. S., and Soltis, D. E. (2021). Polyploidy: an evolutionary and ecological force in stressful times. *Plant Cell* 33, 11–26. doi: 10.1093/plcell/koaa015
- Villa, A. (1984). The chromosome idiogram of *Nicotiana plumbaginifolia*. *Genetica* 64, 145–148. doi: 10.1007/BF00120266
- Weiss-Schneeweiss, H., and Schneeweiss, G. M. (2013). “Karyotype diversity and evolutionary trends in angiosperms,” in *Plant Genome Diversity Volume 2*, ed. I. J. Leitch (Vienna: Springer), 209–230. doi: 10.1007/978-3-7091-1160-4\_13
- Wu, F., Eannetta, N. T., Xu, Y., and Tanksley, S. D. (2009). A detailed synteny map of the eggplant genome based on conserved ortholog set II (COSII) markers. *Theor. Appl. Genet.* 118, 927–935. doi: 10.1007/s00122-008-0950-9
- Wu, F., and Tanksley, S. D. (2010). Chromosomal evolution in the plant family Solanaceae. *BMC Genomics* 11:182. doi: 10.1186/1471-2164-11-182



- Zenil-Ferguson, R., Burleigh, J. G., Freyman, W. A., Igić, B., Mayrose, I., and Goldberg, E. E. (2019). Interaction among ploidy, breeding system and lineage diversification. *New Phytol.* 224, 1252–1265. doi: 10.1111/nph.16184
- Zenil-Ferguson, R., Burleigh, J. G., and Ponciano, J. M. (2018). Chromploid: An R package for chromosome number evolution across the plant tree of life. *Appl. Plant Sci.* 6:e1037. doi: 10.1002/aps3.1037

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Deanna, Acosta, Scaldaferrero and Chiarini. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.