

Gluon quasiparticles and the CGC density matrix

Haowu Duan *North Carolina State University, Raleigh, North Carolina 27695, USA*

Alex Kovner

Physics Department, University of Connecticut, 2152 Hillside Road, Storrs, Connecticut 06269, USA

Vladimir V. Skokov*

RIKEN-BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973, USA

(Received 31 December 2021; accepted 25 February 2022; published 10 March 2022)

We revisit and extend the calculation of the density matrix and entanglement entropy of a color glass condensate (CGC) by including the leading saturation corrections in the calculation. We show that the density matrix is diagonal in the quasiparticle basis, where it has the Boltzmann form. The quasiparticles in a wide interval of momenta behave as massless two-dimensional bosons with the temperature proportional to the typical semihard scale $T = Q_s / \sqrt{\alpha_s N_c}$. Thus, the semihard momentum region $Q_s < k < Q_s / \sqrt{\alpha_s N_c}$ arises as a well-defined intermediate regime between the perturbatively hard momenta and the non-perturbative soft momenta $k < Q_s$ in the CGC description of a hadronic wave function.

DOI: [10.1103/PhysRevD.105.056009](https://doi.org/10.1103/PhysRevD.105.056009)

I. INTRODUCTION

Recently, there has been an increased interest in incorporating the concepts and methods of quantum information theory into nuclear and particle physics [1]. In particular, various aspects of entanglement in application to hadronic collisions have been considered in [2–8]. A new intriguing connection between the physics of black holes and high energy hadrons was formulated in Ref. [9]. In the context of high energy collisions, various ideas about possible relevance of entanglement to thermalization and parton distributions have been discussed in [10–13]. The entanglement entropy between strongly coupled nonperturbative modes and partonic components of a hadronic wave function was conjectured to be the origin of the Boltzmann entropy of particles produced in the collisions.

It was pointed out in [14] that the color glass condensate (CGC) effective theory provides an explicit and calculable model of entanglement in a high energy hadronic wave function. The concept of entanglement implies a partition of a system into the system of interest and its complement. In [14], the soft gluon degrees of freedom were considered as the system of interest while the valence degrees of

freedom (larger x partons) in the hadron wave function were treated as the complement. This partition is directly relevant to measurements of observables at midrapidity which reflect the properties of the soft gluons. Integrating out the valence degrees of freedom in the original (pure state) hadron wave function produces a mixed state density matrix of the soft gluons. The entropy associated with this mixed state is the entanglement entropy in question.

Originally, the entanglement entropy between valence degrees of freedom and soft gluons was calculated in [14] in the dilute limit, $Q_s^2/k^2 \ll 1$. In the present paper, we extend this calculation by accounting for saturation effects in a “mean field” approximation. In addition, we discuss some properties of the CGC reduced density matrix and the associated entropy which were not investigated in earlier work. In particular, we notice that the entropy has a Boltzmann form. This implies that the associated reduced density matrix not only can be diagonalized (this, of course, can always be done, in principle) but also that the eigenvalues of the density matrix are given by the powers of the same single number (which in our case is a function of transverse momentum). In other words, in the appropriate basis, the reduced density matrix has a Boltzmann form albeit with a momentum-dependent effective temperature. We explicitly find this basis by performing a Bogoliubov transformation from the original free gluon basis and discuss some interesting properties of the corresponding quasiparticles. With the benefit of hindsight, we do not find our results on the Boltzmann form of the entanglement entropy surprising. At the leading order, the

*VSkokov@ncsu.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

hadron wave function is a coherent operator acting on the soft vacuum. It is given by the exponential of the argument linear in the creation-annihilation operators. Integrating the valence degrees of freedom in a Gaussian/McLerran-Venugopalan leads to a reduced density matrix for the soft sector, which has a general Gaussian form. A proper Bogoliubov transformation would thus always lead to a reduced density matrix with a Boltzmann structure.

We start in Sec. II by reviewing the derivation of [14] while also providing some technical details. In Sec. III, we discuss the extension of this calculation by including the saturation effects. In Sec. IV, we diagonalize the density matrix and give explicitly its Boltzmann form in the quasiparticle basis. We conclude with a short discussion in Sec. V.

II. REVIEW: THE ENTANGLEMENT IN THE COLOR GLASS CONDENSATE

A. The CGC hadron wave function

For an ultrarelativistic proton, a large fraction of the longitudinal momentum is carried by the valence degrees of freedom. When boosted, the valence partons radiate gluons with lower longitudinal momentum which have a relatively short lifetime. It is then natural to separate the degrees of freedom according to their longitudinal momentum: the large longitudinal momentum partons can be treated as static sources of soft, low longitudinal momentum, gluons. These degrees of freedom are of course strongly correlated, and these correlations play an important role for the phenomenology. For example, triggering on high multiplicities at forward rapidities also selects events with high multiplicity at midrapidity. A less conventional quantity which measures the correlation strength is the entanglement entropy—the main focus of this paper.

In the CGC approach, the hadron state vector can be written in the form

$$|\psi\rangle = |s\rangle \otimes |v\rangle, \quad (1)$$

where $|v\rangle$ is the state vector characterizing the valence degrees of freedom and $|s\rangle$ the state in the soft gluon Hilbert space. The direct product in this equation is not mathematically precise, as we alluded to before, since the soft gluons are sourced by valence degrees of freedom. In CGC, this is encoded by

$$|s\rangle = \mathcal{C}|0\rangle, \quad (2)$$

with the coherent operator

$$\mathcal{C} = \exp \left\{ 2i \int_{\underline{k}} \text{tr} b^i(\underline{k}) [a_i^+(\underline{k}) + a_i(-\underline{k})] \right\}, \quad (3)$$

where the summation over all color is implied. Here, $|0\rangle$ is the Fock vacuum of the soft gluon Hilbert space. The

“background” Weizsäcker-Williams gluon field b_a^i is the solution of the static Yang-Mills equation

$$\partial_i b_i^a(x) = g\rho^a(x), \quad (4)$$

where $\rho^a(x)$ is the color charge density of the valence gluons.

Equation (2) strictly speaking is valid in the regime when the color charge density is weak where one can perform the perturbative diagonalization of the QCD Hamiltonian in the soft gluon sector. Nevertheless, in principle, the solution to Eq. (2) contains nonlinearities in ρ which reflect certain gluon saturation effects. Often, for the sake of simplification, one considers the limit $k \gg Q_s$ where these nonlinearities are small. In this case, the solution of the Yang-Mills equation can be expressed as¹

$$b_a^i(\underline{k}) = g\rho_a(\underline{k}) \frac{ik_i}{k^2} + c_a^i(\underline{k}), \quad (5)$$

where $c_a^i(\underline{k})$ is at least $\mathcal{O}(\rho^2)$. In this section, following the original derivation of Ref. [14], we neglect the contribution due to $c_a^i(\underline{k})$. We come back to consider nonlinear terms in the next section. Note that the first term in Eq. (5) is longitudinal (in the two dimensional sense) while c^i is transverse; that is, $c^i k^i = 0$. Thus, neglecting c_a^i leads to excitation of only longitudinal gluon degrees of freedom in the soft gluon wave function.

We use the McLerran-Venugopalan model to model the valence state [15,16]. This corresponds to treating the color charges as the only relevant valence degrees of freedom with the distribution

$$\langle \rho | v \rangle \langle v | \rho \rangle = e^{-\int_{\underline{k}} \rho_a(\underline{k}) \frac{1}{2\mu^2(\underline{k})} \rho_a^*(\underline{k})}. \quad (6)$$

The density matrix of the complete system is thus given by

$$\hat{\rho} = |s\rangle \langle s| \otimes |v\rangle \langle v|, \quad (7)$$

and corresponds to a pure state, as it should.

¹For completeness, we provide the conventions for the Fourier transformation in two dimensions,

$$f(\underline{x}) = \int \frac{d^2 \underline{k}}{(2\pi)^2} e^{ik \cdot \underline{x}} f(\underline{k}) \equiv \int_{\underline{k}} e^{ivk \cdot \underline{x}} f(\underline{k})$$

$$f(\underline{k}) = \int d^2 x e^{-ik \cdot \underline{x}} f(\underline{x}).$$

In this paper, $\underline{k}$ stands for 2-d vector, and k stands for its magnitude.

B. The reduced density matrix of soft gluons

To proceed with evaluation of the entanglement entropy, we integrate out the valence modes. The resulting reduced density matrix for the soft modes is defined as

$$\hat{\rho}_r = \sum_v \langle v | \hat{\rho} | v \rangle = \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} \rho_a(k) \rho_a^*(k)} \mathcal{C} | 0 \rangle \langle 0 | \mathcal{C}^\dagger. \quad (8)$$

Our next step is to compute the reduced density matrix explicitly. For this, it is convenient to introduce the notation

$$\begin{aligned} \phi(\underline{k}) &= a(\underline{k}) + a^+(-\underline{k}), \\ \phi(x) &= a(x) + a^+(x). \end{aligned}$$

The *matrix element* of the reduced density matrix in the field basis is

$$\begin{aligned} \langle \phi_1 | \hat{\rho}_r | \phi_2 \rangle &= \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} \rho_a(k) \rho_a^*(k)} \\ &\times \langle \phi_1 | \mathcal{C}(\rho, \phi) | 0 \rangle \langle 0 | \mathcal{C}^\dagger(\rho, \phi) | \phi_2 \rangle. \quad (9) \end{aligned}$$

In the above,

$$\begin{aligned} \langle \phi_1 | \mathcal{C}(\rho, \phi) | 0 \rangle &= \langle \phi_1 | \exp \left\{ i \int_k b_a^i(\underline{k}) \phi_a^*(\underline{k}) \right\} | 0 \rangle \\ &= \exp \left\{ i \int_k b_a^i(\underline{k}) \phi_{1a}^*(\underline{k}) \right\} \langle \phi_1 | 0 \rangle. \quad (10) \end{aligned}$$

The wave function for the coherent vacuum $\langle \phi_1 | 0 \rangle$ is known, and we use its explicit form at a later stage of the derivation. Equation (9) then becomes

$$\begin{aligned} \langle \phi_1 | \hat{\rho} | \phi_2 \rangle &= \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} \rho_c(k) \rho_c^*(k)} e^{i \int_k b_a^i(\underline{k}) \phi_{1a}^*(\underline{k})} \\ &\times \langle \phi_1 | 0 \rangle \langle 0 | \phi_2 \rangle e^{-i \int_k b_b^j(\underline{k}) \phi_{2b}^j(\underline{k})}. \quad (11) \end{aligned}$$

The integral over the color charge density ρ is Gaussian and can be evaluated in a straightforward manner. Using Eq. (5) and neglecting nonlinear terms in the solution, we obtain

$$\begin{aligned} \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} \rho_c(k) \rho_c^*(k) + i \int_k b_a^i(\underline{k}) \phi_{1a}^*(\underline{k}) - i \int_k b_b^j(\underline{k}) \phi_{2b}^j(\underline{k})} &= \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} \rho_c(k) \rho_c^*(k) + i \int_k b_a^i(\underline{k}) (\phi_{1a}^i(-\underline{k}) - \phi_{2a}^i(-\underline{k}))} \\ &= \mathcal{N} \int D\rho e^{-\int \frac{1}{k^2 \mu^2(k)} [\rho_a(\underline{k}) - g\mu^2 \frac{k_i}{k^2} (\phi_{a1i}(\underline{k}) - \phi_{a2i}(\underline{k}))] [\rho_a(-\underline{k}) - g\mu^2 \frac{-k_i}{k^2} (\phi_{a1i}(-\underline{k}) - \phi_{a2i}(-\underline{k}))]} \\ &\times e^{-\int_k \frac{g^2 \mu^2 k_i k_j}{k^4} (\phi_{a1j}(\underline{k}) - \phi_{a2j}(\underline{k})) (\phi_{a1i}(-\underline{k}) - \phi_{a2i}(-\underline{k}))} \\ &= e^{-\int_k \frac{g^2 \mu^2 k_i k_j}{k^4} (\phi_{a1j}(\underline{k}) - \phi_{a2j}(\underline{k})) (\phi_{a1i}(-\underline{k}) - \phi_{a2i}(-\underline{k}))}. \quad (12) \end{aligned}$$

Therefore, the matrix element reads

$$\begin{aligned} \langle \phi_1 | \hat{\rho} | \phi_2 \rangle &= \langle \phi_1 | 0 \rangle \langle 0 | \phi_2 \rangle e^{-\int_k \frac{1}{2} M_{ij}^{ab}(\underline{k}) (\phi_{1i}^a(\underline{k}) - \phi_{2i}^a(\underline{k})) (\phi_{1j}^b(-\underline{k}) - \phi_{2j}^b(-\underline{k}))}, \quad (13) \end{aligned}$$

with

$$M_{ij}^{ab}(\underline{k}) \equiv g^2 \mu^2(k) \delta^{ab} \frac{k_i k_j}{k^4}. \quad (14)$$

This has to be supplemented by the vacuum wave function (in terms of fields in the coordinate space representation),

$$\langle \phi | 0 \rangle = N_{vac} e^{-\frac{1}{4} \int_x \phi_i^a(x) \phi_i^a(x)}, \quad (15)$$

where N is defined by the condition²

$$1 = \langle 0 | 0 \rangle = \int D\phi \langle 0 | \phi \rangle \langle \phi | 0 \rangle = N_{vac}^2 \int D\phi e^{-\frac{1}{2} \int_x \phi_i^a(x) \phi_i^a(x)}.$$

We finally obtain that the matrix element that has the following form:

$$\langle \phi_1 | \hat{\rho} | \phi_2 \rangle = N_{vac}^2 e^{-\int_k \left[\frac{1}{2} M_{ij}^{ab}(\underline{k}) (\phi_{1i}^a(\underline{k}) - \phi_{2i}^a(\underline{k})) (\phi_{1j}^b(-\underline{k}) - \phi_{2j}^b(-\underline{k})) + \frac{1}{4} \phi_{1i}^a(\underline{k}) \phi_{1i}^a(-\underline{k}) + \frac{1}{4} \phi_{2i}^a(\underline{k}) \phi_{2i}^a(-\underline{k}) \right]}. \quad (16)$$

²The argument of the exponential of the vacuum wave function is normalized to yield the same result for the density matrix as we previously obtained in [14], see also [17,18].

The representation (16) is the most convenient for computing the von Neumann entropy.

C. The von Neumann entropy

The von Neumann entropy of a density matrix is defined as³

$$S^E = -\text{tr}[\hat{\rho} \ln \hat{\rho}]. \quad (17)$$

The calculation is facilitated by using the replica trick. Using the identity

$$\ln \hat{\rho} = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (\hat{\rho}^\epsilon - 1), \quad (18)$$

we have

$$S^E = -\lim_{\epsilon \rightarrow 0} \text{tr} \left(\frac{\hat{\rho}^N - \hat{\rho}}{\epsilon} \right) = -\lim_{\epsilon \rightarrow 0} \frac{\text{tr} \hat{\rho}^N - 1}{\epsilon}, \quad (19)$$

where $N = \epsilon + 1$. Thus, the problem of evaluating the von Neumann entropy reduces to computation of $\text{tr} \hat{\rho}^N$ for integer N and subsequent analytic continuation to arbitrary N . Note that $\text{tr} \hat{\rho}^N$ is related to the N th Renyi entropy

$$S_N = \frac{1}{1-N} \ln [\text{Tr} \hat{\rho}^N]. \quad (20)$$

It is straightforward to proceed with

$$\begin{aligned} \text{Tr} \hat{\rho}^N &= \int D\phi_1 \langle \phi_1 | \hat{\rho}^N | \phi_1 \rangle \\ &= \int D\phi_1 D\phi_2 \langle \phi_1 | \hat{\rho} | \phi_2 \rangle \langle \phi_2 | \hat{\rho}^{N-1} | \phi_1 \rangle = \dots \\ &= \int \prod_{n=1}^N D\phi_n \langle \phi_n | \hat{\rho} | \phi_{n+1} \rangle, \end{aligned}$$

where the fields satisfy periodic boundary conditions in replica space $\phi_{N+1} = \phi_1$. We use boldface font to denote the field index. With the help of Eq. (16), we can explicitly write

$$\begin{aligned} \text{Tr} \hat{\rho}^N &= N_{vac}^N \int D\phi_1 D\phi_2 \dots D\phi_N \\ &\times \exp \left\{ -\frac{1}{2} \sum_{\mathbf{n}=1}^N \int_{\underline{k}} \phi_{\mathbf{n}i}^a(\underline{k}) \phi_{\mathbf{n}i}^a(-\underline{k}) \right. \\ &- \frac{1}{2} \sum_{\mathbf{n}=1}^N \int_{\underline{k}} [\phi_{\mathbf{n}i}^a(\underline{k}) - \phi_{(\mathbf{n}+1)i}^a(\underline{k})] M_{ij}^{ab}(\underline{k}) [\phi_{\mathbf{n}j}^b(-\underline{k}) \\ &- \phi_{(\mathbf{n}+1)j}^b(-\underline{k})] \left. \right\}. \quad (21) \end{aligned}$$

³For a thorough discussion of various definitions of entropy, see [19].

The integrand involves mixing terms between different replica fields; however, it can be diagonalized by performing the Fourier transformation with respect to the replica index (we suppress other indices for simplicity),

$$\tilde{\phi}_\eta = \frac{1}{N} \sum_{\mathbf{n}=1}^N e^{i\frac{2\pi}{N}\mathbf{n}\eta} \phi_{\mathbf{n}}, \quad (22)$$

$$\phi_{\mathbf{n}} = \sum_{\eta=0}^{N-1} e^{-i\frac{2\pi}{N}\mathbf{n}\eta} \tilde{\phi}_\eta. \quad (23)$$

This yields

$$\begin{aligned} &\sum_{\mathbf{n}=1}^N (\phi_{\mathbf{n}i}^a - \phi_{(\mathbf{n}+1)i}^a) (\phi_{\mathbf{n}j}^b - \phi_{(\mathbf{n}+1)j}^b) \\ &= N \sum_{\eta=0}^{N-1} \left(1 - e^{-i\frac{2\pi\eta}{N}} \right) \left(1 - e^{i\frac{2\pi\eta}{N}} \right) \tilde{\phi}_{\eta i}^a \tilde{\phi}_{(-\eta)j}^b \quad (24) \end{aligned}$$

$$= 4N \sum_{\eta=0}^{N-1} \sin^2 \left(\frac{\pi}{N} \eta \right) \tilde{\phi}_{\eta i}^a \tilde{\phi}_{(-\eta)j}^b, \quad (25)$$

and the problem is reduced to a standard Gaussian integral,

$$\begin{aligned} \text{Tr} \hat{\rho}^N &= \mathcal{N} \int D\tilde{\phi}_0 D\tilde{\phi}_1 \dots D\tilde{\phi}_{N-1} \\ &\times \exp \left\{ -N \sum_{\eta=0}^{N-1} \tilde{\phi}_{\eta i}^a \left[\frac{1}{2} \delta_{ij} \delta^{ab} + 2M_{ij}^{ab} \sin^2 \left(\frac{\pi}{N} \eta \right) \right] \right. \\ &\times \left. \tilde{\phi}_{(-n)j}^b \right\}, \quad (26) \end{aligned}$$

where we have absorbed the Jacobian into the normalization factor which we establish below. The Gaussian integral yields

$$\text{Tr} \hat{\rho}^N = \mathcal{N} \det \left[\prod_{\eta=0}^{N-1} \left(\frac{1}{2} + 2M \sin^2 \left(\frac{\pi}{N} \eta \right) \right)^{-\frac{1}{2}} \right] \quad (27)$$

$$= \mathcal{N} \det \left[\prod_{\eta=0}^{N-1} \left(\frac{1}{2} + M \left(1 - \cos \left(\frac{2\pi}{N} \eta \right) \right) \right)^{-\frac{1}{2}} \right]. \quad (28)$$

The matrix M in Eq. (28) is diagonal in color, momentum, and the replica space. Its polarization structure is purely longitudinal, so that the eigenvalues are $M_- = 0$ and $M_+ = \frac{q^2 \mu^2}{k^2}$. We therefore get

$$\begin{aligned}
 \text{Tr } \hat{\rho}^N &= \mathcal{N} \det \left[\prod_{\eta=0}^{N-1} \left(\frac{1}{2} + M_- \left(1 - \cos \left(\frac{2\pi}{N} \eta \right) \right) \right) \right]^{-\frac{1}{2}} \Big]^{N_a} \\
 &\times \det \left[\prod_{\eta=0}^{N-1} \left(\frac{1}{2} + M_+ \left(1 - \cos \left(\frac{2\pi}{N} \eta \right) \right) \right) \right]^{-\frac{1}{2}} \Big]^{N_a} \\
 &= \mathcal{N} \det \left[\prod_{\eta=0}^{N-1} \left(1 + 2M_+ \left(1 - \cos \left(\frac{2\pi}{N} \eta \right) \right) \right) \right]^{-N_a/2}, \quad (29)
 \end{aligned}$$

where we again absorbed irrelevant constants into $\mathcal{N}$.

To perform summation over η , we adapt the formula (1.394) from [20],

$$\prod_{l=0}^{N-1} \left[x^2 - 2xy \cos \frac{2l\pi}{N} + y^2 \right] = (x^N - y^N)^2, \quad (30)$$

using the following mapping:

$$x^2 + y^2 = 1 + 2M_+, 2xy = 2M_+.$$

The result is

$$\begin{aligned}
 &\prod_{\eta=0}^{N-1} \left(1 + 2M_+ \left(1 - \cos \left(\frac{2\pi}{N} \eta \right) \right) \right) \\
 &= \frac{1}{2^{2N}} \left[(\sqrt{1+4M_+} + 1)^N - (\sqrt{1+4M_+} - 1)^N \right]^2. \quad (31)
 \end{aligned}$$

Thus, we arrive at

$$\begin{aligned}
 \text{Tr } \hat{\rho}^N &= \mathcal{N} \det_k \{ 2^{NN_a} [(\sqrt{1+4M_+} + 1)^N \\
 &\quad - (\sqrt{1+4M_+} - 1)^N]^{-N_a} \}, \quad (32)
 \end{aligned}$$

where $\det_k$ denotes a determinant in momentum space (product over momenta) only, and $N_a = N_c^2 - 1$. The normalization factor $\mathcal{N}$ can now be determined by requiring that the reduced density matrix is properly normalized. Setting $N = 1$, we have $\text{Tr } \hat{\rho} = \mathcal{N} \det_k 1$, and thus, $\mathcal{N} = 1/\det_k 1$.

Performing analytical continuation $N = 1 + \epsilon$ and expanding in ϵ , we obtain

$$\begin{aligned}
 \text{Tr } \hat{\rho}^N &\approx \mathcal{N} \det \left\{ 1 - \epsilon \frac{N_a}{2} \left(\ln M_+ + \sqrt{1+4M_+} \right. \right. \\
 &\quad \times \ln \left[1 + \frac{1}{2M_+} + \frac{1}{2M_+} \sqrt{1+4M_+} \right] \Big) \Big\} \\
 &\approx 1 - \epsilon \frac{N_a}{2} S_\perp \int_{\underline{k}} \left(\ln M_+ + \sqrt{1+4M_+} \right. \\
 &\quad \times \ln \left[1 + \frac{1}{2M_+} + \frac{1}{2M_+} \sqrt{1+4M_+} \right] \Big). \quad (33)
 \end{aligned}$$

Therefore, the entanglement entropy is given by

$$\begin{aligned}
 S_E &= \frac{N_a}{2} S_\perp \int_{\underline{k}} \left(\ln M_+ + \sqrt{1+4M_+} \right. \\
 &\quad \times \ln \left[1 + \frac{1}{2M_+} + \frac{1}{2M_+} \sqrt{1+4M_+} \right] \Big). \quad (34)
 \end{aligned}$$

With the definition $M_+ = \frac{g^2 \mu^2}{k^2}$, this reproduces the result of Ref. [14].⁴

III. BEYOND THE DILUTE APPROXIMATION

In the previous section, we derived the entanglement entropy under the dilute approximation,

$$\begin{aligned}
 \langle b_a^i(\underline{q}) b_b^j(-\underline{p}) \rangle &\equiv \mathcal{N} \int D\rho e^{-\int_{\underline{k}^2 \mu^2(\underline{k})} \rho_c(\underline{k}) \rho_c^*(\underline{k})} b_a^i(\underline{q}) b_b^j(-\underline{p}) \\
 &\approx \delta^2(\underline{q} - \underline{p}) (2\pi)^2 g^2 \mu^2 \delta^{ab} \frac{p^i p^j}{p^4}. \quad (35)
 \end{aligned}$$

As we pointed out before, this approximation neglects any saturation corrections in the wave function, and as a result, only the longitudinally polarized gluons contribute to the entanglement entropy. Here, our goal is to take into account the saturation corrections.

In general, the solution of the static Yang-Mills equation for the Weizsäcker-Williams field is given by

$$\begin{aligned}
 b_i^a(\underline{x}) &= \frac{1}{igN_c} \text{Tr} [T^a U^+(\underline{x}) \partial_i U(\underline{x})] \\
 &= \frac{2}{ig} \text{Tr} [t^a V^+(\underline{x}) \partial_i V(\underline{x})]. \quad (36)
 \end{aligned}$$

Given that the color charge distribution must be globally color invariant, the field correlator must have the form $\langle b_a^i(\underline{x}) b_b^j(\underline{y}) \rangle \propto \delta^{ab}$. Summing with respect to the colors, we have

⁴Reference [14] used a different normalization of μ , see also Ref. [17].

$$\begin{aligned}
\langle b_i^a(\underline{x}) b_j^a(\underline{y}) \rangle &= -\frac{4}{g^2} \langle \text{Tr}[t^a V^+(\underline{x}) \partial_i V(\underline{x})] \text{Tr}[t^a V^+(\underline{y}) \partial_j V(\underline{y})] \rangle \\
&= -\frac{2}{g^2} \langle \text{Tr}[V^+(\underline{x}) \partial_i V(\underline{x}) V^+(\underline{y}) \partial_j V(\underline{y})] \rangle \\
&= \frac{(2\pi)^3}{2} x G_{ij}^{\text{WW}}(x, \underline{x} - \underline{y}).
\end{aligned} \quad (37)$$

Here, $x G_{ij}^{\text{WW}}(x, \underline{x} - \underline{y})$ is the Weizsäcker-Williams gluon distribution in the coordinate space. Performing Fourier transformation and taking into account the translational invariance of the Weizsäcker-Williams gluon distribution function, we arrive at

$$\begin{aligned}
\langle b_i^a(\underline{k}) b_j^a(\underline{q}) \rangle &= \frac{(2\pi)^5}{2} \delta^{(2)}(\underline{k} + \underline{q}) \int d^2 r e^{-i \underline{r} \cdot \underline{k}} x G_{ij}^{\text{WW}}(x, \underline{r}) \\
&= \frac{(2\pi)^5}{2} \frac{\delta^{(2)}(\underline{k} + \underline{q})}{S_\perp} x G_{ij}^{\text{WW}}(x, k).
\end{aligned} \quad (38)$$

The factor of the transverse area $S_\perp$ in the denominator originates from the commonly accepted definition of $x G_{ij}^{\text{WW}}(x, k)$,

$$\begin{aligned}
x G_{ij}^{\text{WW}}(x, k) &= \int d^2 x d^2 y e^{-i(x-y)k} x G_{ij}^{\text{WW}}(x, \underline{x} - \underline{y}) \\
&= S_\perp \int d^2 r e^{-i \underline{r} \cdot \underline{k}} x G_{ij}^{\text{WW}}(x, \underline{r}).
\end{aligned} \quad (39)$$

Thus,

$$\langle b_i^a(\underline{k}) b_j^b(\underline{q}) \rangle = \frac{(2\pi)^5}{2(N_c^2 - 1)S_\perp} \delta^{(2)}(\underline{k} + \underline{q}) \delta^{ab} x G_{ij}^{\text{WW}}(x, k). \quad (40)$$

The tensor $x G_{ij}^{\text{WW}}(x, k)$ is conventionally split into two independent components,

$$x G_{ij}^{\text{WW}}(x, k) = \frac{1}{2} \delta_{ij} x G^{(1)}(x, k) - \frac{1}{2} \left(\delta_{ij} - 2 \frac{k_i k_j}{k^2} \right) x h^{(1)}(x, k), \quad (41)$$

where $x h^{(1)}$ is the linearly polarized gluon distribution. Thus, in general, the Weizsäcker-Williams field correlator contains both longitudinal and transverse components, with the transverse component proportional to $x G^{(1)} - x h^{(1)}$.

In the MV model, both components can be computed semianalytically [21] to yield

$$\begin{aligned}
x h^{(1)}(x, q_\perp) &= \frac{S_\perp}{2\pi^3 \alpha_s} \frac{N_c^2 - 1}{N_c} \int_0^\infty dr_\perp \frac{r_\perp J_2(q_\perp r_\perp)}{r_\perp^2 \ln\left(\frac{1}{r_\perp^2 \Lambda^2}\right)} \\
&\times \left[1 - e^{-\frac{1}{4} r_\perp^2 Q_s^2 \ln\left(\frac{1}{r_\perp^2 \Lambda^2}\right)} \right],
\end{aligned} \quad (42)$$

$$\begin{aligned}
x G^{(1)}(x, q_\perp) &= \frac{S_\perp}{2\pi^3 \alpha_s} \frac{N_c^2 - 1}{N_c} \int_0^\infty dr_\perp \frac{r_\perp J_0(q_\perp r_\perp)}{r_\perp^2} \\
&\times \left[1 - e^{-\frac{1}{4} r_\perp^2 Q_s^2 \ln\left(\frac{1}{r_\perp^2 \Lambda^2}\right)} \right],
\end{aligned} \quad (43)$$

where Λ is a nonperturbative IR scale, and the saturation momentum Q_s is given by

$$Q_s^2 = N_c \alpha_s g^2 \mu^2. \quad (44)$$

In this paper, we use the expression in the MV model for the Weizsäcker-Williams gluon field distributions; our results, however, can be straightforwardly extended to account for the small- x evolution [22].

Now, we are ready to revise the derivation of the reduced matrix to account for the saturation corrections. We go back to the integration over the valence degrees of freedom in Eq. (12),

$$\begin{aligned}
&\left\langle e^{-i \int_{\underline{k}} b_a^i(\underline{k}) (\phi_{1,a}^i(-\underline{k}) - \phi_{2,a}^i(-\underline{k}))} \right\rangle \\
&= 1 + \sum_{q=1}^\infty \frac{1}{q!} \left\langle \left[-i \int_{\underline{k}} b_a^i(\underline{k}) (\phi_{1,a}^i(-\underline{k}) - \phi_{2,a}^i(-\underline{k})) \right]^q \right\rangle.
\end{aligned}$$

The right-hand side contains all higher order correlators of the Weizsäcker-Williams field. We will however invoke a simple minded mean field Gaussian approximation in which all higher correlators factorize into products of the two point function. In this approximation, we have

$$\begin{aligned}
&\left\langle e^{-i \int_{\underline{k}} b_a^i(\underline{k}) (\phi_{1,a}^i(-\underline{k}) - \phi_{2,a}^i(-\underline{k}))} \right\rangle_{\text{MV}} \\
&= e^{-\int_{\underline{k}} \tilde{M}_{ij}^{ab}(\underline{k}) (\phi_{b1j}(\underline{k}) - \phi_{b2j}(\underline{k})) (\phi_{a1i}(-\underline{k}) - \phi_{a2i}(-\underline{k}))},
\end{aligned} \quad (45)$$

with

$$\tilde{M}_{ij}^{ab}(\underline{k}) = \frac{(2\pi)^3 \delta^{ab}}{2(N_c^2 - 1)S_\perp} x G_{ij}^{\text{WW}}(x, k). \quad (46)$$

This approximation, albeit simple, allows us to incorporate the main saturation effects in the CGC density matrix. While deriving Eq. (46), we took into account that $\delta^{(2)}(\underline{k} = 0) = S_\perp / (2\pi)^2$.

In the limit $k \gg Q_s$, we recover the dilute approximation of the previous section, i.e., $\tilde{M}_{ij}^{ab}(k) \rightarrow M_{ij}^{ab}(k)$. To show this explicitly, consider the eigenvalues of $x G_{ij}^{\text{WW}}(x, k)$.

Those are $(xG^{(1)} \pm xh^{(1)})/2$. At large momentum, one can consider small r in the integrals (42) and (43). Expanding the exponentials, we obtain

$$\begin{aligned} xh^{(1)}(x, q_\perp) &\approx \frac{S_\perp}{2\pi^3\alpha_s} \frac{N_c^2 - 1}{N_c} \frac{Q_s^2}{4} \int_0^\infty dr_\perp r_\perp J_2(q_\perp r_\perp) \\ &\approx \frac{S_\perp}{4\pi^3\alpha_s} \frac{N_c^2 - 1}{N_c} \frac{Q_s^2}{q_\perp^2}, \\ xG^{(1)}(x, q_\perp) &\approx \frac{S_\perp}{2\pi^3\alpha_s} \frac{N_c^2 - 1}{N_c} \frac{Q_s^2}{4} \int_0^\infty dr_\perp r_\perp J_0(q_\perp r_\perp) \ln\left(\frac{1}{r_\perp^2 \Lambda^2}\right) \\ &\approx \frac{S_\perp}{4\pi^3\alpha_s} \frac{N_c^2 - 1}{N_c} \frac{Q_s^2}{q_\perp^2}. \end{aligned} \quad (47)$$

This reproduces the limits discussed in Ref. [23]. We thus obtain one zero eigenvalue and the other one given by

$$\lim_{q \gg Q_s} \frac{xG^{(1)} + xh^{(1)}}{2} = \frac{S_\perp}{4\pi^3\alpha_s} \frac{N_c^2 - 1}{N_c} \frac{Q_s^2}{q_\perp^2}. \quad (48)$$

The nontrivial eigenvalue of $\tilde{M}$ becomes

$$\frac{S_\perp}{4\pi^3\alpha_s} \frac{1}{N_c} \frac{Q_s^2}{q_\perp^2} \times \frac{(2\pi)^3 \delta^{ab}}{2S_\perp} = \frac{g^2 \mu^2 \delta^{ab}}{q_\perp^2}. \quad (49)$$

This indeed reduces to the expression in the dilute limit used in the previous section.

Repeating the derivation of the previous section separately for each eigenvalue of the matrix $\tilde{M}$, we obtain

$$\begin{aligned} S^E &= \frac{N_c^2 - 1}{2} \sum_{\nu=\pm} \int_{\underline{k}} \left[\ln \tilde{M}_\nu(\underline{k}) + \sqrt{1 + 4\tilde{M}_\nu(\underline{k})} \right. \\ &\quad \left. \times \ln \left(1 + \frac{1}{2\tilde{M}_\nu(\underline{k})} + \frac{\sqrt{1 + 4\tilde{M}_\nu(\underline{k})}}{2\tilde{M}_\nu(\underline{k})} \right) \right], \end{aligned} \quad (50)$$

where

$$\tilde{M}_\pm = \frac{(2\pi)^3}{2S_\perp(N_c^2 - 1)} \frac{xG^{(1)} \pm xh^{(1)}}{2}. \quad (51)$$

Comparing to Eq. (34), there are two major differences. First, Eq. (50) contains a nontrivial contribution of the transverse mode, and second, the small momentum behavior of the larger eigenvalue is very different from that of M_+ . We return to the discussion of this point later on.

IV. DIAGONALIZATION OF THE REDUCED DENSITY MATRIX

A. The Boltzmann density matrix

As we alluded to in the Introduction, it is anticipated that the result of the previous section can be rewritten in a Boltzmann form,⁵

$$S_E = (N_c^2 - 1) S_\perp \sum_{\nu=\pm} \int \frac{d^2k}{(2\pi)^2} \left[(1 + f_\nu) \ln(1 + f_\nu) - f_\nu \ln f_\nu \right]. \quad (52)$$

Here, we defined the distribution functions

$$f_\pm(k) = \frac{1}{\exp(\beta\omega_\pm(k)) - 1}, \quad (53)$$

with

$$\beta\omega_\pm(\underline{k}) = 2 \ln \left(\frac{1}{2\sqrt{\tilde{M}_\pm(\underline{k})}} + \sqrt{1 + \frac{1}{4\tilde{M}_\pm(\underline{k})}} \right). \quad (54)$$

This suggests that in a basis in which the density matrix is diagonal, it must have the Boltzmann form,

$$\hat{\rho} = N e^{-\beta\omega_+ \hat{n}_+} e^{-\beta\omega_- \hat{n}_-}, \quad (55)$$

where $\hat{n}_\pm$ is the corresponding number density of the quasiparticles. We refer to this basis as the quasiparticle basis.

Our purpose in this section is to find the quasiparticle basis explicitly. Before turning to this problem, we ask what is the dispersion relation of the quasiparticles provided our interpretation of Eq. (52) is correct.

Let us first examine the dilute case of Sec. II. For the only nontrivial polarization $M_+ = g^2 \mu^2 / k^2$ and at small momenta $k \ll g^2 \mu^2$, Eq. (54) gives $\beta\omega_+ \approx k / (g\mu)$. Interestingly, this looks like a dispersion relation of a massless particle. Although only the product of the frequency and the inverse temperature is determined by Eq. (54), assuming the velocity of quasiparticles is the speed of light, the inverse temperature is $\beta = (g\mu)^{-1}$. At large momentum, $\beta\omega_+ \approx \ln(k^2 / g^2 \mu^2)$ or $f_+ \approx g^2 \mu^2 / k^2$. This perturbativelike behavior is then interpreted as a logarithmic dispersion relation for quasiparticles at high momenta.

One interesting point to note is that the transition between the “low momentum” and “high momentum” regimes in the present context is given by the scale $g^2 \mu^2$ which is *parametrically larger* than the saturation momentum Q_s , i.e., $g^2 \mu^2 = Q_s^2 / \alpha_s N_c$. Physically, this is easy to understand. While saturation effects become important at the momentum scale at which the gluon occupation number

⁵A similar result was obtained in a general Gaussian density matrix [10].

is large, of order $1/\alpha_s$, the perturbative regime, i.e., momenta for which the eigenvalue M_+ becomes small, requires the occupation number to be smaller than unity. Thus, there is a whole range of momenta, at which the gluon occupation number is greater than one, but saturation effects are still unimportant. We refer to this range of momenta, $Q_s^2 < k^2 < Q_s^2/\alpha_s N_c$ as “semihard”. At small 'tHooft coupling $\alpha_s N_c$, the semihard region is parametrically large.

The saturation corrections discussed in the previous section have a strong effect on the “dispersion relation” at low momenta. Beyond the dilute limit, $xG^{(1)}$ dominates over $xh^{(1)}$ at $k < Q_s$, so that both eigenvalues $\tilde{M}_\pm$ behave as $xG^{(1)} \propto \ln Q_s/k$ in the infrared. This is a well-known effect where the infrared $1/k^2$ behavior of the transverse momentum dependent which formally leads to a power like infrared divergence in the produced particle spectrum is tamed by the saturation corrections and becomes logarithmic. This correction results in the logarithmic “dispersion relation” $\beta\omega_\pm \propto 1/\sqrt{\ln Q_s/k}$. This modification however is only important at very low momenta $k^2 < Q_s^2$. In the semihard regime $Q_s^2 < k^2 < Q_s^2/\alpha_s N_c$, the quasiparticles corresponding to the eigenvalue M_+ still behave approximately as massless bosons. For these momenta, the eigenvalue M_+ is large. For large M_+ , we can expand the logarithm in Eq. (54), which leads to $\beta\omega_+ \approx 1/\sqrt{\tilde{M}_+}$. Then, taking the high momentum approximation, that is $\tilde{M}_+ \propto Q_s^2/\alpha_s k^2$, we obtain $\beta\omega_+ \propto \sqrt{\alpha_s} k/Q_s$, i.e., thermal spectrum for massless particles. The temperature is of the order $T \sim Q_s/\sqrt{\alpha_s N_c}$ which is parametrically larger than the saturation momentum. The reason, as we explained above, is that the gluon occupation number is of order unity for momenta which are much higher than Q_s , and it is the value of the momentum of the highest occupied state that determines the effective temperature.

B. Construction of the eigenvalue problem

We now perform the explicit diagonalization of the reduced density matrix.

First, we note that the density matrix has a product form in the momentum space due to the factorization of the momentum modes. Using Eq. (16), we can write the density matrix operator as

$$\begin{aligned} \hat{\rho} &= \prod_{\underline{k}} \hat{\rho}_\nu(\underline{k}) = \prod_{\nu=\pm} \prod_{\underline{k}} \mathcal{N} \int D\phi(\underline{k}) D\Phi(\underline{k}) \\ &\times e^{-\frac{1}{4}(\phi(\underline{k})\phi(-\underline{k})+\Phi(\underline{k})\Phi(-\underline{k}))} e^{-\frac{1}{2}\tilde{M}(\phi(\underline{k})\phi(-\underline{k})-\frac{1}{2}\tilde{M}\Phi(\underline{k})\Phi(-\underline{k}))} \\ &\times e^{\tilde{M}\phi(\underline{k})\Phi(-\underline{k})} \times |\phi(\underline{k})\rangle\langle\Phi(\underline{k})|, \end{aligned} \quad (56)$$

where $\tilde{M}$ is an eigenvalue of $\tilde{M}^{ij}$

$$\tilde{M} = \tilde{M}_+ \text{ or } \tilde{M}_-. \quad (57)$$

Equation (56) should also contain a product over the index $\nu = \pm$, which we do not indicate explicitly.

Since $\hat{\rho}$ is a product over momentum (and polarization), we consider only a single momentum (and polarization) mode for the sake of simplifying the notations.

The Boltzmann form of the entanglement entropy suggests that all eigenvalues of the density matrix are given by integer powers of the same number. Our goal is to find these eigenvalues explicitly by solving the eigenvalue equation,

$$\hat{\rho}|\Psi_i\rangle = \lambda_i|\Psi_i\rangle. \quad (58)$$

We write the eigenstate in the field basis as

$$|\Psi_i(\underline{k})\rangle = \int d\phi(\underline{k}) f_i(\phi(\underline{k})) |\phi(\underline{k})\rangle. \quad (59)$$

The eigenvalue equation for the wave function f_i becomes

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int d\Phi e^{-\frac{(1+2M)}{4}(\phi(-\underline{k})\phi(\underline{k})+\Phi(-\underline{k})\Phi(\underline{k}))} e^{M\phi(-\underline{k})\Phi(\underline{k})} f_i(\Phi) \\ = \lambda_i f_i(\phi). \end{aligned} \quad (60)$$

The form of the integrand suggests to look for the ground state in the form of the Gaussian

$$f_0(\phi(\underline{k})) = N \exp[-\alpha\phi(-\underline{k})\phi(\underline{k})]. \quad (61)$$

Once we find the constant α by solving Eq. (60), we generate exciting states by acting with the creation operators, which are defined analogously to the quantum oscillator problem

$$c(\underline{k}) = \frac{1}{\sqrt{2}} \left(\sqrt{\alpha}\phi(\underline{k}) + \frac{(2\pi)^2}{\sqrt{\alpha}} \frac{\delta}{\delta\phi(-\underline{k})} \right), \quad (62)$$

$$c^\dagger(\underline{k}) = \frac{1}{\sqrt{2}} \left(\sqrt{\alpha}\phi(-\underline{k}) - \frac{(2\pi)^2}{\sqrt{\alpha}} \frac{\delta}{\delta\phi(\underline{k})} \right). \quad (63)$$

The operators satisfy

$$[c(\underline{p}), c^\dagger(\underline{k})] = (2\pi)^2 \delta^{(2)}(\underline{p} - \underline{k}), \quad (64)$$

$$c(\underline{k}) f_0(\phi(\underline{k})) = 0. \quad (65)$$

It can be checked straightforwardly that once the appropriate α is found, the states obtained by repeated action of $c^\dagger(\underline{k})$ on the Gaussian state Eq. (61),

$$f_n(\phi(\underline{k})) = \left(\frac{\alpha}{2} \right)^{\frac{n}{2}} \phi(-\underline{k})^n \exp[-\alpha\phi(\underline{k})\phi(-\underline{k})], \quad (66)$$

are in fact eigenstates of the density matrix.

Substituting the Gaussian function Eq. (61) into Eq. (60) and after a little algebra, we find

$$4\alpha^\pm = \sqrt{1 + 4\tilde{M}_\pm}, \quad (67)$$

$$\lambda_0^\pm = \frac{\sqrt{2}}{\sqrt{1 + 2\tilde{M}_\pm + 4\alpha^\pm}} = \frac{2}{1 + \sqrt{1 + 4\tilde{M}_\pm}}. \quad (68)$$

Proceeding similarly for the excited states, we find

$$\lambda_n^\pm = \left[\frac{2\tilde{M}_\pm}{(1 + \sqrt{1 + 4\tilde{M}_\pm})^2} \right]^n \lambda_0^\pm. \quad (69)$$

This result confirms our earlier expectation that the density matrix has the Boltzmann form. In terms of the operators c and $c^\dagger$, it can be written as

$$\hat{\rho}(\underline{k}) = N \left[\frac{2\tilde{M}_\pm}{(1 + \sqrt{1 + 4\tilde{M}_\pm})^2} \right]^{c^\dagger c} = N e^{-\beta\omega(\underline{k})c^\dagger(\underline{k})c(\underline{k})}, \quad (70)$$

with

$$\beta\omega = \ln \left[\frac{(1 + \sqrt{1 + 4\tilde{M}_\pm})^2}{2\tilde{M}_\pm} \right], \quad (71)$$

and N the appropriate normalization factor. This coincides with Eq. (54).

C. The Bogoliubov transformation

We have thus explicitly established that in the basis of the quasiparticles defined by the creation and annihilation operators $c^\dagger(k)$ and $c(k)$, the density matrix is diagonal and has a Boltzmann form. It is easy to show that the quasiparticle basis is related to the perturbative gluon Fock space basis by a simple Bogoliubov transformation,

$$c_\pm(k) = \cosh(B_\pm)a_\pm(k) + \sinh(B_\pm)a_\pm^\dagger(-k), \quad (72)$$

$$c_\pm^\dagger(k) = \cosh(B_\pm)a_\pm^\dagger(k) + \sinh(B_\pm)a_\pm(-k), \quad (73)$$

where $B_\pm = \ln 2\sqrt{\alpha_\pm} = \frac{1}{4}\ln(1 + 4\tilde{M}_\pm)$. Here, we have restored the polarization indices and have defined $a_+(k) \equiv \hat{k}_i a_i(k)$ and $a_- \equiv \epsilon_{ij}\hat{k}^i a_j(k)$ with $\hat{k}$ a unit vector in the direction of the transverse momentum k .

How “far” is the quasiparticle Fock space removed from the perturbative gluon Fock space? The answer clearly depends on the value of the transverse momentum. Let us consider the two simple limiting cases:

[$k \gg Q_s$:] here, we have $\tilde{M}_- \propto k^{-4}$, while $\tilde{M}_+ \propto k^{-2}$.

For either polarization, $B_\pm \approx 0$ so that the quasiparticle basis practically coincides with the perturbative gluon basis $c_\pm(k) \approx a_\pm(k)$. This is natural since for large momenta the occupation number of gluons

vanishes, and the density matrix in the first approximation is just given by the perturbative vacuum.

[$k \ll Q_s$:] the situation is quite different in this limit. Here, $\tilde{M}_\pm$ is large, and $\sinh(B_\pm) \simeq \cosh(B_\pm) \simeq e^{B_\pm}$. The transformation in this case corresponds to maximal mixing. Interestingly, this maximal mixing regime only requires that $\tilde{M} \gg 1$. Thus, it is not only valid for very small momenta but also for a considerably large range of “semihard” momenta, $k < 2\pi Q_s$. This is the same momenta for which the dispersion relation of the quasiparticles is approximately linear.

V. DISCUSSION

In this paper, we have extended the approach to the CGC density matrix pioneered in [14] by including saturation corrections in a mean-field approximation. The effect of these corrections is two prong. First, they result in excitation of the transverse gluon mode, so that both gluon polarizations now contribute to entanglement entropy. Second, the infrared behavior of the Weizsäcker-Williams field propagator is softened.

We have also pointed out that the reduced soft gluon density matrix can be explicitly diagonalized by a Bogoliubov transformation. In the quasiparticle basis, it has a Boltzmann form; i.e., for a given transverse momentum and polarization, its eigenvalues are powers of one number. If interpreted as a thermal density matrix, this determines the product of the quasiparticle energy and the inverse temperature. We found that for semihard momenta, $Q_s^2 < k^2 < Q_s^2/\alpha_s N_c$, the dispersion relation of the quasiparticles is approximately linear with momentum, while at very low momenta $k^2 < Q_s^2$ the saturation effects lead logarithmic dispersion relation. The saturation then induces an effective mass for the quasiparticles, albeit this mass is not fixed but rather runs with the momentum into the infrared.

We also noted that the effective temperature for the quasiparticle system is not given by the saturation momentum Q_s but rather by a parametrically greater scale $T \sim Q_s/\sqrt{\alpha_s N_c}$. The physical reason is that the temperature is determined by the momentum of those levels for which the occupation number is of order unity rather than much larger than unity.

The two distinct scales that arise in the physics of saturated wave function is reminiscent of the two scales present at high temperature in weakly interacting quark-gluon plasma. The softest scale is where the saturation momentum is closely analogous to the so called “magnetic mass”. Both arise due to self-interaction of very soft modes, and both are parametrically $m_{\text{soft}} \propto \alpha_s \Lambda_{\text{hard}}$, where in the case of plasma $\Lambda_{\text{hard}} = T$, while in CGC $\Lambda_{\text{hard}} = \mu$. The semihard scale, parametrically $m_{\text{semi hard}} \propto g\Lambda_{\text{hard}}$ is identified with the “electric mass” in plasma and the effective temperature in CGC. This scale arises in both cases via the interaction of semihard modes with the hard ones. In the case of plasma, the relevant mechanism is “hard thermal loops”,

while in the case of CGC it is the eikonal emissions by the valence charges that populate the CGC wave function in the semihard region. Amusingly, in both cases, only “electric” modes are affected by this scale, in the sense that in the plasma the electric mass produces finite correlation length only for the chromoelectric field, while in the CGC wave function only the electric (longitudinal in the two dimensional sense) modes are populated in the semihard region.

Note that the value of the effective temperature in this paper is very different from the one obtained in [14]. The reason is that in [14] the inverse temperature was defined as the derivative of the entropy with respect to transverse energy. The total transverse energy is dominated by the UV modes despite the low occupancy of each energy level. Thus, the temperature calculated this way in [14] was dominated by the contribution of the UV modes and came out proportional to the UV cutoff. Conversely, when considering properties of the density matrix of *produced gluons*, which is dominated by momenta of order Q_s , the temperature in [14] came out to be of order Q_s . In the present paper however, we discussed the effective temperature of those modes which have an approximate Boltzmann distribution of massless bosons. These turned out to be semihard modes, and the temperature accordingly turned out to be tied to the appropriate semihard scale.

We conclude by noting that the concept of gluon quasiparticles inside a hadron is a very interesting concept. It is especially worth stressing, that while at high momenta the quasiparticles coincide with perturbative gluons, at

semihard momenta the quasiparticle operators are very different from the perturbative gluon creation and annihilation operators as is clear from our discussion in the previous section. Thus, the semihard momentum region arises here as a well defined transition region between the perturbative hard regime and a genuinely nonperturbative soft ($k < Q_s$) regime. One is reminded of the Landau liquid theory, where quasiparticles indeed arise via possibly strong dressing of original particles while still retaining the same quantum numbers and a particle identity.

It would be extremely interesting to understand how to probe experimentally the properties of such gluon quasiparticles. This is of course a very difficult question, since the quasiparticles discussed here exist inside the wave function of a hadron, while any final state is described (modulo hadronization corrections) in terms of original perturbative gluons. Nevertheless, it seems to us that this problem is well worth thinking about.

ACKNOWLEDGMENTS

We acknowledge useful conversations with E. Levin, M. Li, and M. Lublinsky. We are especially thankful to R. Venugopalan for illuminating discussions. A. K. is supported by the NSF Nuclear Theory Grant No. 1913890. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics through the Contract No. DE-SC0020081 (H. D. and V. V. S.).

-
- [1] N. Klcio, A. Roggero, and M. J. Savage, [arXiv:2107.04769](#).
 - [2] C. Robin, M. J. Savage, and N. Pillet, *Phys. Rev. C* **103**, 034325 (2021).
 - [3] S. R. Beane and P. Ehlers, *Mod. Phys. Lett. A* **35**, 2050048 (2020).
 - [4] N. Armesto, F. Dominguez, A. Kovner, M. Lublinsky, and V. Skokov, *J. High Energy Phys.* **05** (2019) 025.
 - [5] D. Neill and W. J. Waalewijn, *Phys. Rev. Lett.* **123**, 142001 (2019).
 - [6] A. Kovner, M. Lublinsky, and M. Serino, *Phys. Lett. B* **792**, 4 (2019).
 - [7] Y. Hagiwara, Y. Hatta, B.-W. Xiao, and F. Yuan, *Phys. Rev. D* **97**, 094029 (2018).
 - [8] Y. Hatta and Y. Hagiwara, *EPJ Web Conf.* **112**, 01010 (2016).
 - [9] G. Dvali and R. Venugopalan, [arXiv:2106.11989](#).
 - [10] J. Berges, S. Floerchinger, and R. Venugopalan, *J. High Energy Phys.* **04** (2018) 145.
 - [11] J. Berges, S. Floerchinger, and R. Venugopalan, *Phys. Lett. B* **778**, 442 (2018).
 - [12] D. E. Kharzeev and E. M. Levin, *Phys. Rev. D* **95**, 114008 (2017).
 - [13] E. Gotsman and E. Levin, *Phys. Rev. D* **102**, 074008 (2020).
 - [14] A. Kovner and M. Lublinsky, *Phys. Rev. D* **92**, 034016 (2015).
 - [15] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **49**, 2233 (1994).
 - [16] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **49**, 3352 (1994).
 - [17] H. Duan, C. Akkaya, A. Kovner, and V. V. Skokov, *Phys. Rev. D* **101**, 036017 (2020).
 - [18] H. Duan, in *Proceedings of the 28th International Workshop on Deep Inelastic Scattering and Related Subjects* (2021), [arXiv:2107.10812](#).
 - [19] F. I. Ijpelaar, Master’s thesis, University Groningen, VSI, 2021.
 - [20] J. Alan and Z. Daniel, *Table of Integrals, Series, and Products* (Academic Press, Cambridge, 2007).
 - [21] F. Dominguez, J.-W. Qiu, B.-W. Xiao, and F. Yuan, *Phys. Rev. D* **85**, 045003 (2012).
 - [22] A. Dumitru, T. Lappi, and V. Skokov, *Phys. Rev. Lett.* **115**, 252301 (2015).
 - [23] A. Metz and J. Zhou, *Phys. Rev. D* **84**, 051503 (2011).