

Entropy Admissibility of the Limit Solution for a Nonlocal Model of Traffic Flow

Alberto Bressan and Wen Shen

Department of Mathematics, Penn State University.
University Park, PA 16802, USA.

e-mails: axb62@psu.edu, wxs27@psu.edu

February 11, 2021

Abstract

We consider a conservation law model of traffic flow, where the velocity of each car depends on a weighted average of the traffic density ρ ahead. The averaging kernel is of exponential type: $w_\varepsilon(s) = \varepsilon^{-1}e^{-s/\varepsilon}$. For any decreasing velocity function v , we prove that, as $\varepsilon \rightarrow 0$, the limit of solutions to the nonlocal equation coincides with the unique entropy-admissible solution to the scalar conservation law $\rho_t + (\rho v(\rho))_x = 0$.

1 Introduction

We consider a nonlocal PDE model for traffic flow, where the traffic density $\rho = \rho(t, x)$ satisfies a scalar conservation law with nonlocal flux

$$\rho_t + (\rho v(q))_x = 0. \quad (1.1)$$

Here $\rho \mapsto v(\rho)$ is a decreasing function, modeling the velocity of cars depending on the traffic density, while the integral

$$q(x) = \int_x^{+\infty} \varepsilon^{-1} e^{(x-y)/\varepsilon} \rho(y) ds \quad (1.2)$$

computes a weighted average of the density of cars ahead. As in [2], the assumption that the averaging kernel is of exponential type plays a crucial role in our analysis. For more general averaging kernels, the limiting behavior of the solutions as $\varepsilon \rightarrow 0$ is not yet fully understood. Following [2] we also assume

(A1) *The velocity function $v : [0, \rho_{jam}] \mapsto \mathbb{R}_+$ is \mathcal{C}^2 , and satisfies*

$$v(\rho_{jam}) = 0, \quad v'(\rho) \leq -\delta_* < 0, \quad \text{for all } \rho \in [0, \rho_{jam}]. \quad (1.3)$$

One can think of ρ_{jam} as the maximum possible density of cars along the road, when all cars are packed bumper-to-bumper and nobody moves. The conservation equation (1.1) will be solved with initial data

$$\rho(0, x) = \bar{\rho}(x) \in [0, \rho_{jam}]. \quad (1.4)$$

As $\varepsilon \rightarrow 0+$, the weight function $w_\varepsilon(s) = \varepsilon^{-1}e^{-s/\varepsilon}$ converges to a Dirac mass at the origin, and the nonlocal equation (1.1)-(1.2) formally converges to the scalar conservation law

$$\rho_t + (\rho v(\rho))_x = 0. \quad (1.5)$$

Assuming that the initial datum $\bar{\rho}$ has bounded total variation and takes uniformly positive values, the recent analysis in [2] has established:

- (i) For every $\varepsilon > 0$, the Cauchy problem with non-local flux (1.1), (1.2), (1.4), has a unique solution $\rho = \rho_\varepsilon(t, x)$. Its total variation satisfies a uniform bound

$$\text{Tot.Var.}\{\rho_\varepsilon(t, \cdot)\} \leq M \quad (1.6)$$

where the constant M is independent of t, ε .

- (ii) As $\varepsilon \rightarrow 0$, by possibly taking a subsequence, one obtains the convergence $\rho_\varepsilon \rightarrow \rho$ in \mathbf{L}_{loc}^1 . The limit function $\rho = \rho(t, x)$ provides a weak solution to the Cauchy problem (1.4)-(1.5).

A major issue, which was not fully resolved in [2], is the entropy admissibility of the limit solution ρ . Aim of the present note is to resolve this question in the affirmative. Namely, we prove:

Theorem. *Let v satisfy the assumptions **(A1)**, and let ρ_ε be a sequence of solutions to the nonlocal Cauchy problem (1.1), (1.2) and (1.4), satisfying the uniform BV bounds (1.6). Assume that, as $\varepsilon \rightarrow 0$, we have the convergence $\rho_\varepsilon \rightarrow \rho$ in \mathbf{L}_{loc}^1 . Then ρ is the unique entropy admissible solution to the Cauchy problem (1.4)-(1.5).*

The above result was proved in [2] in the special case where the velocity is affine: $v(\rho) = a - b\rho$. The earlier proof was based on the Hardy-Littlewood inequality. In the next section we give a simpler proof, valid for a general class of velocity functions v .

For a more general class of averaging kernels, assuming that the initial datum $\bar{\rho}$ satisfies a one-sided Lipschitz condition, the convergence to the unique entropy admissible solution was recently proved in [3]. Our result requires an exponential kernel, but it applies to any BV initial data. In particular, $\bar{\rho}$ can be piecewise constant.

For the general theory of conservation laws we refer to [1, 5, 6]. A brief review of literature on hyperbolic conservation laws with nonlocal flux can be found in [2].

2 Proof of the theorem

1. According to [4, 7], to prove uniqueness it suffices to prove that the limit solution dissipates one single strictly convex entropy. We thus consider the entropy and entropy

flux pair

$$\eta(\rho) = \frac{\rho^2}{2}, \quad \psi(\rho) = \int_0^\rho [sv(s) + s^2v'(s)] ds. \quad (2.1)$$

For future use, we observe that (1.2) implies

$$\rho = q - \varepsilon q_x. \quad (2.2)$$

Moreover, we introduce the function

$$W(\rho) \doteq \int_0^\rho s^2v'(s) ds. \quad (2.3)$$

The equation (1.1) can now be written as

$$\rho_t + (\rho v(\rho))_x = \left(\rho(v(\rho) - v(q)) \right)_x.$$

Multiplying both sides by $\eta'(\rho) = \rho$, we obtain

$$\eta(\rho)_t + \psi(\rho)_x = \rho \left(\rho(v(\rho) - v(q)) \right)_x. \quad (2.4)$$

2. Given a test function $\varphi \in \mathcal{C}_c^1(\mathbb{R})$, $\varphi \geq 0$, we estimate the quantity

$$\begin{aligned} J &\doteq 2 \int \rho \left(\rho(v(\rho) - v(q)) \right)_x \varphi dx \\ &= \int (\rho^2)_x (v(\rho) - v(q)) \varphi dx + \int 2\rho^2 (v(\rho) - v(q))_x \varphi dx \\ &= - \int \rho^2 (v(\rho) - v(q)) \varphi_x dx + \int \rho^2 (v(\rho) - v(q))_x \varphi dx \\ &\doteq J_1 + J_2. \end{aligned} \quad (2.5)$$

Concerning the second integral, using (2.2) we obtain

$$\begin{aligned} J_2 &= \int \rho^2 v'(\rho) \rho_x \varphi dx - \int \rho q v'(q) q_x \varphi dx + \int \rho \varepsilon (q_x)^2 v'(q) \varphi dx \\ &\doteq J_{21} + J_{22} + J_{23}. \end{aligned} \quad (2.6)$$

Using (2.2) once again, we now compute

$$\begin{aligned} J_{21} + J_{22} &= \int \rho^2 v'(\rho) \rho_x \varphi dx - \int q^2 v'(q) q_x \varphi dx + \int q \varepsilon (q_x)^2 v'(q) \varphi dx \\ &\doteq J_3 + J_4 + J_5. \end{aligned} \quad (2.7)$$

Since $\rho, q, \varphi \geq 0$ while $v' \leq 0$, from (2.6) and (2.7) we immediately see that

$$J_{23} \leq 0, \quad J_5 \leq 0. \quad (2.8)$$

On the other hand, integrating by parts and recalling (2.3), we obtain

$$\begin{aligned} J_3 + J_4 &= \int [W(\rho)]_x \varphi dx - \int [W(q)]_x \varphi dx \\ &= - \int [W(\rho) - W(q)] \varphi_x dx. \end{aligned} \quad (2.9)$$

3. To conclude, consider a sequence of solutions ρ_ε to (1.1)-(1.2), (1.4). Assume that, as $\varepsilon \rightarrow 0$, we have the convergence $\rho_\varepsilon \rightarrow \rho$ in \mathbf{L}_{loc}^1 . Notice that this implies $q_\varepsilon \rightarrow \rho$ in \mathbf{L}_{loc}^1 as well. Hence, the integrals J_1 and $J_3 + J_4$ both approach zero. By the previous analysis,

$$\begin{aligned} & 2 \iint \{ \eta(\rho_\varepsilon) \varphi_t + \psi(\rho_\varepsilon) \varphi_x \} dxdt \\ & \geq \iint \rho_\varepsilon^2 (v(\rho_\varepsilon) - v(q_\varepsilon)) \varphi_x dxdt + \iint [W(\rho_\varepsilon) - W(q_\varepsilon)] \varphi_x dxdt. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, since the right hand side converges to zero, we obtain

$$\iint \{ \eta(\rho) \varphi_t + \psi(\rho) \varphi_x \} dxdt \geq 0.$$

This proves that the limit solution ρ is entropy admissible. In particular, by [4, 7], ρ is the unique entropy weak solution to the Cauchy problem (1.4)-(1.5). \square

Acknowledgment. This research was partially supported by NSF with grant DMS-2006884, ‘‘Singularity and error bounds for hyperbolic equations’’.

References

- [1] A. Bressan, *Hyperbolic Systems of Conservation Laws. The One Dimensional Cauchy Problem*, Oxford University Press, 2000.
- [2] A. Bressan and W. Shen, On traffic flow with nonlocal flux: a relaxation representation, *Arch. Rational Mech. Anal.* **237** (2020), 1213–1236.
- [3] M. Colombo, G. Crippa, E. Marconi, and L.V. Spinolo, Local limit of nonlocal traffic models: convergence results and total variation blow-up. Preprint 2019, arXiv:1808.03529v2.
- [4] C. De Lellis, F. Otto, and M. Westdickenberg, Minimal entropy conditions for Burgers equation. *Quart. Appl. Math.* **62** (2004), 687–700.
- [5] C. Dafermos, *Hyperbolic Conservation Laws in Continuum Physics*, Fourth edition. Springer-Verlag, Berlin, 2016.
- [6] H. Holden and N. Risebro, *Front Tracking for Hyperbolic Conservation Laws*. Springer-Verlag, Berlin, 2002.
- [7] E. Y. Panov. Uniqueness of the solution of the Cauchy problem for a first order quasi-linear equation with one admissible strictly convex entropy. (Russian) *Mat. Zametki* **55** (1994), 116–129; translation in *Math. Notes* **55** (1994), 517–525.