

Learning Low-Degree Functions from a Logarithmic Number of Random Queries

Alexandros Eskenazis
University of Cambridge
Cambridge, UK
ae466@cam.ac.uk

Paata Ivanisvili
University of California, Irvine
Irvine, CA, USA
pivanisv@uci.edu

ABSTRACT

We prove that every bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d can be learned with L_2 -accuracy ε and confidence $1 - \delta$ from $\log(\frac{n}{\delta}) \varepsilon^{-d-1} C^{d^{3/2}} \sqrt{\log d}$ random queries, where $C > 1$ is a universal finite constant.

CCS CONCEPTS

• **Theory of computation** → **Sample complexity and generalization bounds**; **Boolean function learning**; • **Mathematics of computing** → **Probabilistic algorithms**; **Approximation**.

KEYWORDS

Discrete hypercube, learning theory, Bohnenblust–Hille inequality

ACM Reference Format:

Alexandros Eskenazis and Paata Ivanisvili. 2022. Learning Low-Degree Functions from a Logarithmic Number of Random Queries. In *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing (STOC '22)*, June 20–24, 2022, Rome, Italy. ACM, New York, NY, USA, 5 pages. <https://doi.org/10.1145/3519935.3519981>

1 INTRODUCTION

Every function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ admits a unique Fourier–Walsh expansion of the form

$$\forall x \in \{-1, 1\}^n, \quad f(x) = \sum_{S \subseteq \{1, \dots, n\}} \hat{f}(S) w_S(x), \quad (1)$$

where $w_S(x) = \prod_{i \in S} x_i$ and the Fourier coefficients $\hat{f}(S)$ are given by

$$\forall S \subseteq \{1, \dots, n\}, \quad \hat{f}(S) = \frac{1}{2^n} \sum_{y \in \{-1, 1\}^n} f(y) w_S(y). \quad (2)$$

We say that f has degree at most $d \in \{1, \dots, n\}$ if $\hat{f}(S) = 0$ for every subset S with $|S| > d$.

1.1 Learning Functions on the Hypercube

Let $\mathcal{C}$ be a class of functions $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ on the n -dimensional discrete hypercube. The problem of learning the class $\mathcal{C}$ can be described as follows: given a source of examples $(x, f(x))$, where $x \in \{-1, 1\}^n$, for an unknown function $f \in \mathcal{C}$, compute a hypothesis

function $h : \{-1, 1\}^n \rightarrow \mathbb{R}$ which is a good approximation of f up to a given error in some prescribed metric. In this paper we will be interested in the *random query model* with L_2 -error, in which we are given N independent examples $(x, f(x))$, each chosen uniformly at random from the discrete hypercube $\{-1, 1\}^n$, and we want to efficiently construct a (random) function $h : \{-1, 1\}^n \rightarrow \mathbb{R}$ such that $\|h - f\|_{L_2}^2 < \varepsilon$ with probability at least $1 - \delta$, where $\varepsilon, \delta \in (0, 1)$ are given accuracy and confidence parameters. The goal is to construct a randomized algorithm which produces the hypothesis function h from a minimal number N of examples.

The above very general problem has been studied for decades in computational learning theory and many results are known¹, primarily for various classes $\mathcal{C}$ of structured Boolean functions $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$. Already since the late 1980s, researchers used the Fourier–Walsh expansion [1] to design such learning algorithms (see the survey [14]). Perhaps the most classical of these is the *Low-Degree Algorithm* of Linial, Mansour and Nisan [12] who showed that for the class $\mathcal{C}_b^d$ of all bounded functions $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d there exists an algorithm which produces an ε -approximation of f with probability at least $1 - \delta$ using $N = \frac{2n^d}{\varepsilon} \log(\frac{2n^d}{\delta})$ samples. In this generality, the $O_{\varepsilon, \delta, d}(n^d \log n)$ estimate of [12] was the state of the art until the recent work [11] of Iyer, Rao, Reis, Rothvoss and Yehudayoff who employed analytic techniques to derive new bounds on the ℓ_1 -size of the Fourier spectrum of bounded functions (see also Section 3) and used these estimates to show that $N = O_{\varepsilon, \delta, d}(n^{d-1} \log n)$ examples suffice to learn $\mathcal{C}_b^d$. The goal of the present paper is to further improve this result and show that in fact $N = O_{\varepsilon, \delta, d}(\log n)$ samples suffice for this purpose.

THEOREM 1. Fix $\varepsilon, \delta \in (0, 1)$, $n \in \mathbb{N}$, $d \in \{1, \dots, n\}$ and a bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d . If $N \in \mathbb{N}$ satisfies

$$N \geq \min \left\{ \frac{\exp(Cd^{3/2} \sqrt{\log d})}{\varepsilon^{d+1}}, \frac{4dn^d}{\varepsilon} \right\} \log \left(\frac{n}{\delta} \right), \quad (3)$$

where $C \in (0, \infty)$ is a large numerical constant, then N uniformly random independent queries of pairs $(x, f(x))$, where $x \in \{-1, 1\}^n$, suffice for the construction of a random function $h : \{-1, 1\}^n \rightarrow \mathbb{R}$ satisfying the condition $\|h - f\|_{L_2}^2 < \varepsilon$ with probability at least $1 - \delta$.

The proof of Theorem 1 relies on some important approximation theoretic estimates going back to the 1930s which we shall now describe (see also [9]). To the best of our knowledge, these tools had not yet been exploited in the computational learning theory literature.

¹We will by no means attempt to survey this (vast) field, so we refer the interested reader to the relevant chapters of O’Donnell’s book [15] and the references therein.



This work is licensed under a Creative Commons Attribution 4.0 International License.

STOC '22, June 20–24, 2022, Rome, Italy

© 2022 Copyright held by the owner/author(s).

ACM ISBN 978-1-4503-9264-8/22/06.

<https://doi.org/10.1145/3519935.3519981>

1.2 The Fourier Growth of Walsh Polynomials in $\ell^{\frac{2d}{d+1}}$

Estimates for the growth of coefficients of polynomials as a function of their degree and their maximum on compact sets go back to the early days of approximation theory (see [5]). A seminal result of this nature is Littlewood's celebrated $\frac{4}{3}$ -inequality [13] for bilinear forms which was later generalized by Bohnenblust and Hille [4] for multilinear forms on the torus $\mathbb{T}^n$ or the unit square $[-1, 1]^n$. By means of polarization, one can use this multilinear estimate to derive an inequality for polynomials which reads as follows². For every $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ and $d \in \mathbb{N}$, there exists $B_d^{\mathbb{K}} \in (0, \infty)$ such that for every $n \in \mathbb{N}$ and every coefficients $c_\alpha \in \mathbb{K}$, where $\alpha \in (\mathbb{N} \cup \{0\})^n$ with $|\alpha| \leq d$, we have

$$\left(\sum_{|\alpha| \leq d} |c_\alpha|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \leq B_d^{\mathbb{K}} \max \left\{ \left| \sum_{|\alpha| \leq d} c_\alpha x^\alpha \right| : \|x\|_{\ell_\infty^n(\mathbb{K})} \leq 1 \right\}. \quad (4)$$

Moreover, $\frac{2d}{d+1}$ is the smallest exponent for which the optimal constant in (4) is independent of the number of variables n of the polynomial. The exact asymptotics of the constants $B_d^{\mathbb{R}}$ and $B_d^{\mathbb{C}}$ remain unknown, however it is known that there is a significant gap between $B_d^{\mathbb{R}}$ and $B_d^{\mathbb{C}}$, namely that $\limsup_{d \rightarrow \infty} (B_d^{\mathbb{R}})^{1/d} = 1 + \sqrt{2}$ whereas $B_d^{\mathbb{C}} \leq C^{\sqrt{d \ln d}}$ for a finite constant $C > 1$ (see [1, 6–9] for these and other important advances of the last decade). Restricting inequality (4) to real multilinear polynomials, convexity shows that the maximum on the right-hand side is attained at a point $x \in \{-1, 1\}^n$, which, in view of (1), makes (4) an estimate for the Fourier–Walsh growth of functions on the discrete hypercube. We shall denote by $B_d^{\{\pm 1\}}$ the corresponding optimal constant (first explicitly investigated by Blei in [3, p. 175]), that is, the least constant such that for every $n \in \mathbb{N}$ and every function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ of degree at most d ,

$$\left(\sum_{S \subseteq \{1, \dots, n\}} |\hat{f}(S)|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \leq B_d^{\{\pm 1\}} \|f\|_{L_\infty}. \quad (5)$$

The best known quantitative result in this setting is due to Defant, Mastylo and Pérez [8] who showed that $B_d^{\{\pm 1\}} \leq \exp(\kappa \sqrt{d \log d})$ for a universal constant $\kappa \in (0, \infty)$. The main contribution of this work is the following theorem relating the growth of the constant $B_d^{\{\pm 1\}}$ and learning.

THEOREM 2. Fix $\varepsilon, \delta \in (0, 1)$, $n \in \mathbb{N}$, $d \in \{1, \dots, n\}$ and a bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d . If $N \in \mathbb{N}$ satisfies

$$N \geq \frac{e^8 d^2}{\varepsilon^{d+1}} (B_d^{\{\pm 1\}})^{2d} \log \left(\frac{n}{\delta} \right), \quad (6)$$

then given N uniformly random independent queries of pairs $(x, f(x))$, where $x \in \{-1, 1\}^n$, one can construct a function $h : \{-1, 1\}^n \rightarrow \mathbb{R}$ satisfying $\|h - f\|_{L_2}^2 < \varepsilon$ with probability at least $1 - \delta$.

In Section 2 we will prove Theorem 2 and use it to derive Theorem 1. In Section 3 we will present some additional remarks on

²For $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{N} \cup \{0\})^n$, we use the standard notations $|\alpha| = \alpha_1 + \dots + \alpha_n$ and $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$.

Boolean analysis and learning, in particular showing that the dependence on n in Theorem 1 is optimal for $\delta \asymp \frac{1}{n}$. Moreover, we shall improve the recent bounds of [11] on the ℓ_1 -Fourier growth of bounded functions of low degree.

2 PROOFS

PROOF OF THEOREM 2. Fix a parameter $b \in (0, \infty)$ and denote by

$$N_b \stackrel{\text{def}}{=} \left\lceil \frac{2}{b^2} \log \left(\frac{2}{\delta} \sum_{k=0}^d \binom{n}{k} \right) \right\rceil. \quad (7)$$

Let $X_1, \dots, X_{N_b}$ be independent random vectors, each uniformly distributed on $\{-1, 1\}^n$. For a subset $S \subseteq \{1, \dots, n\}$ with $|S| \leq d$ consider the empirical Walsh coefficient of f , given by

$$\alpha_S = \frac{1}{N_b} \sum_{j=1}^{N_b} f(X_j) w_S(X_j). \quad (8)$$

As α_S is a sum of bounded i.i.d. random variables and $\mathbb{E}[\alpha_S] = \hat{f}(S)$, the Chernoff bound gives

$$\forall S \subseteq \{1, \dots, n\}, \quad \mathbb{P}\{|\alpha_S - \hat{f}(S)| > b\} \leq 2 \exp(-N_b b^2 / 2). \quad (9)$$

Therefore, using the union bound and taking into account that f has degree at most d , we get

$$\begin{aligned} \mathbb{P}\{|\alpha_S - \hat{f}(S)| \leq b, \text{ for every } S \subseteq \{1, \dots, n\} \text{ with } |S| \leq d\} \\ \geq 1 - 2 \sum_{k=0}^d \binom{n}{k} \exp(-N_b b^2 / 2) \stackrel{(7)}{\geq} 1 - \delta. \end{aligned} \quad (10)$$

Denote the event on the LHS by G_b . Fix an additional parameter $a \in (b, \infty)$ and consider the random collection of sets given by

$$\mathcal{S}_a \stackrel{\text{def}}{=} \{S \subseteq \{1, \dots, n\} : |\alpha_S| \geq a\}. \quad (11)$$

Observe that if the event G_b of equation (10) holds, then

$$\forall S \notin \mathcal{S}_a, \quad |\hat{f}(S)| \leq |\alpha_S - \hat{f}(S)| + |\alpha_S| < a + b \quad (12)$$

and

$$\forall S \in \mathcal{S}_a, \quad |\hat{f}(S)| \geq |\alpha_S| - |\alpha_S - \hat{f}(S)| \geq a - b. \quad (13)$$

Finally, consider the random function $h_{a,b} : \{-1, 1\}^n \rightarrow \mathbb{R}$ given by

$$\forall x \in \{-1, 1\}^n, \quad h_{a,b}(x) \stackrel{\text{def}}{=} \sum_{S \in \mathcal{S}_a} \alpha_S w_S(x). \quad (14)$$

Combining (13) with inequality (5), we deduce that

$$\begin{aligned} |\mathcal{S}_a| &\stackrel{(13)}{\leq} (a - b)^{-\frac{2d}{d+1}} \sum_{S \in \mathcal{S}_a} |\hat{f}(S)|^{\frac{2d}{d+1}} \\ &\leq (a - b)^{-\frac{2d}{d+1}} \sum_{S \subseteq \{1, \dots, n\}} |\hat{f}(S)|^{\frac{2d}{d+1}} \stackrel{(5)}{\leq} (a - b)^{-\frac{2d}{d+1}} (B_d^{\{\pm 1\}})^{\frac{2d}{d+1}}. \end{aligned} \quad (15)$$

Therefore, on the event G_b we have

$$\begin{aligned}
\|h_{a,b} - f\|_{L_2}^2 &= \sum_{S \subseteq \{1, \dots, n\}} |\hat{h}_{a,b}(S) - \hat{f}(S)|^2 \\
&= \sum_{S \in \mathcal{S}_a} |\alpha_S - \hat{f}(S)|^2 + \sum_{S \notin \mathcal{S}_a} |\hat{f}(S)|^2 \\
&\stackrel{(12)}{<} |\mathcal{S}_a| b^2 + (a+b)^{\frac{2}{d+1}} \sum_{S \notin \mathcal{S}_a} |\hat{f}(S)|^{\frac{2d}{d+1}} \\
&\stackrel{(5) \wedge (15)}{\leq} (B_d^{\{\pm 1\}})^{\frac{2d}{d+1}} ((a-b)^{-\frac{2d}{d+1}} b^2 + (a+b)^{\frac{2}{d+1}}).
\end{aligned} \tag{16}$$

Choosing $a = b(1 + \sqrt{d+1})$, we deduce that

$$\begin{aligned}
&\|h_{b(1+\sqrt{d+1}), b} - f\|_{L_2}^2 \\
&< (B_d^{\{\pm 1\}})^{\frac{2d}{d+1}} b^{\frac{2}{d+1}} ((d+1)^{-\frac{d}{d+1}} + (2 + \sqrt{d+1})^{\frac{2}{d+1}}).
\end{aligned} \tag{17}$$

Next, we need the technical inequality: for all $d \geq 1$,

$$(d+1)^{-\frac{d}{d+1}} + (2 + \sqrt{d+1})^{\frac{2}{d+1}} \leq (e^4(d+1))^{\frac{1}{d+1}}. \tag{18}$$

Rearranging the terms, it suffices to show that

$$(2 + \sqrt{d+1})^{\frac{2}{d+1}} \leq (d+1)^{\frac{1}{d+1}} (e^{\frac{4}{d+1}} - \frac{1}{d+1}), \tag{19}$$

which is equivalent to $(\frac{2}{\sqrt{d+1}} + 1)^{\frac{2}{d+1}} \leq e^{\frac{4}{d+1}} - \frac{1}{d+1}$. We have

$$\left(\frac{2}{\sqrt{d+1}} + 1\right)^{\frac{2}{d+1}} \leq (\sqrt{2} + 1)^{\frac{2}{d+1}} \stackrel{(*)}{\leq} 1 + \frac{3}{d+1} \leq e^{\frac{4}{d+1}} - \frac{1}{d+1},$$

where inequality $(*)$ holds because the left hand side is convex in the variable $\lambda \stackrel{\text{def}}{=} \frac{2}{d+1}$ whereas the right hand side is linear and since $(*)$ holds at the endpoints $\lambda = 0, 1$.

Combining (17) and (18) we see that $\|h_{b(1+\sqrt{d+1}), b} - f\|_{L_2}^2 < \varepsilon$ holds for $b^2 \leq e^{-5} d^{-1} \varepsilon^{d+1} (B_d^{\{\pm 1\}})^{-2d}$. Plugging this choice of b in (7) shows that given N random queries, where

$$N = \left\lceil \frac{e^6 d (B_d^{\{\pm 1\}})^{2d}}{\varepsilon^{d+1}} \log \left(\frac{2}{\delta} \sum_{k=0}^d \binom{n}{k} \right) \right\rceil, \tag{20}$$

the function $h_{b(1+\sqrt{d+1}), b}$ satisfies $\|h_{b(1+\sqrt{d+1}), b} - f\|_{L_2}^2 < \varepsilon$ with probability at least $1 - \delta$ and the conclusion of the theorem follows from elementary estimates, such as

$$\sum_{k=0}^d \binom{n}{k} \leq \sum_{k=0}^d \frac{n^k}{k!} = \sum_{k=0}^d \frac{d^k}{k!} \left(\frac{n}{d}\right)^k \leq \left(\frac{en}{d}\right)^d. \quad \square$$

Theorem 1 is a straightforward consequence of Theorem 2.

PROOF OF THEOREM 1. Our Theorem 2 combined with the bound $B_d^{\{\pm 1\}} \leq \exp(\kappa \sqrt{d \log d})$ of [8] imply the conclusion of Theorem 1 for $\varepsilon \geq \frac{\exp(C \sqrt{d \log d})}{n}$, where $C \in (0, \infty)$ is a large universal constant. The case $\varepsilon < \frac{\exp(C \sqrt{d \log d})}{n}$ follows from the Low-Degree Algorithm of [12]. $\square$

3 CONCLUDING REMARKS

We conclude with a few additional remarks on the spectrum of bounded functions defined on the hypercube and learning algorithms. For a function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$, its Rademacher projection on level $\ell \in \{1, \dots, n\}$ is defined as

$$\forall x \in \{-1, 1\}^n, \quad \text{Rad}_\ell f(x) = \sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S| = \ell}} \hat{f}(S) w_S(x). \tag{21}$$

1. The first main theorem of [11] asserts that if $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ is a function of degree d , then for $\ell \in \{1, \dots, d\}$,

$$\|\text{Rad}_\ell f\|_{L_\infty} \leq \begin{cases} \frac{|T_d^{(\ell)}(0)|}{\ell!} \cdot \|f\|_{L_\infty}, & \text{if } (d - \ell) \text{ is even} \\ \frac{|T_{d-1}^{(\ell)}(0)|}{\ell!} \cdot \|f\|_{L_\infty}, & \text{if } (d - \ell) \text{ is odd} \end{cases}, \tag{22}$$

where $T_d(t)$ is the d -th Chebyshev polynomial of the first kind, that is, the unique real polynomial of degree d such that $\cos(d\theta) = T_d(\cos \theta)$ for every $\theta \in \mathbb{R}$. Moreover, Iyer, Rao, Reis, Rothvoss and Yehudayoff observed in [11, Proposition 2] that this estimate is asymptotically sharp. We present a simple proof of their inequality (22) (see also [10] for related arguments).

PROOF OF (22). For any $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ consider its harmonic extension on $[-1, 1]^n$,

$$\forall x \in [-1, 1]^n, \quad \tilde{f}(x_1, \dots, x_n) = \sum_{S \subseteq \{1, \dots, n\}} \hat{f}(S) \prod_{j \in S} x_j. \tag{23}$$

By convexity $\|\tilde{f}\|_{L_\infty([-1, 1]^n)} = \|f\|_{L_\infty(\{-1, 1\}^n)}$. In particular, the restriction of $\tilde{f}$ on the ray $t(x_1, \dots, x_n)$, $t \in [-1, 1]$, i.e.

$$\forall t \in \mathbb{R}, \quad h_x(t) \stackrel{\text{def}}{=} \sum_{S \subseteq \{1, \dots, n\}} \hat{f}(S) w_S(x) t^{|S|} \tag{24}$$

satisfies $\max_{t \in [-1, 1]} |h_x(t)| \leq \|f\|_{L_\infty}$ for all $(x_1, \dots, x_n) \in \{-1, 1\}^n$. Therefore, since $\deg h_x \leq d$, a classical inequality of Markov (see e.g. [5, p. 248]) gives

$$|\text{Rad}_\ell f(x)| = \frac{|h_x^{(\ell)}(0)|}{\ell!} \leq \begin{cases} \frac{|h_x^{(\ell)}(0)|}{\ell!} \cdot \|f\|_{L_\infty}, & \text{if } (d - \ell) \text{ is even} \\ \frac{|h_x^{(\ell)}(0)|}{\ell!} \cdot \|f\|_{L_\infty}, & \text{if } (d - \ell) \text{ is odd} \end{cases}$$

and (22) follows by taking a maximum over all $x \in \{-1, 1\}^n$. $\square$

In particular, as observed in [11], inequality (22) implies that if f has degree at most d then

$$\forall \ell \in \{1, \dots, d\}, \quad \|\text{Rad}_\ell f\|_{L_\infty} \leq \frac{d^\ell}{\ell!} \cdot \|f\|_{L_\infty}. \tag{25}$$

2. The second main theorem of [11] asserts that if a function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ has degree at most d , then for every $\ell \in \{1, \dots, d\}$ we have

$$\sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S| = \ell}} |\widehat{\text{Rad}_\ell f}(S)| = \sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S| = \ell}} |\hat{f}(S)| \leq n^{\frac{\ell-1}{2}} d^\ell e^{\binom{\ell+1}{2}}. \tag{26}$$

The Bohnenblust–Hille-type inequality of [8] implies the following improved bound.

COROLLARY 3. *Let $n \in \mathbb{N}$ and $d \in \{1, \dots, n\}$. Then, every bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d satisfies*

$$\sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S| = \ell}} |\hat{f}(S)| \leq \binom{n}{\ell}^{\frac{\ell-1}{2\ell}} e^{\kappa \sqrt{\ell \log \ell}} \frac{d^\ell}{\ell!} = O(n^{\frac{\ell-1}{2}} d^\ell \ell^{-c\ell}), \quad (27)$$

for some universal constant $c \in (0, 1)$ and all $\ell \in \{1, \dots, d\}$.

PROOF. Combining Hölder's inequality with the estimate of [8] and (25) we get

$$\begin{aligned} \sum_{\substack{S \subseteq \{1, \dots, n\} \\ |S| = \ell}} |\hat{f}(S)| &\leq \binom{n}{\ell}^{\frac{\ell-1}{2\ell}} \left(\sum_{S \subseteq \{1, \dots, n\}} |\widehat{\text{Rad}_\ell f}(S)|^{\frac{2\ell}{\ell+1}} \right)^{\frac{\ell+1}{2\ell}} \\ &\leq \binom{n}{\ell}^{\frac{\ell-1}{2\ell}} \exp(\kappa \sqrt{\ell \log \ell}) \|\text{Rad}_\ell f\|_{L_\infty} \\ &\stackrel{(25)}{\leq} \binom{n}{\ell}^{\frac{\ell-1}{2\ell}} \exp(\kappa \sqrt{\ell \log \ell}) \frac{d^\ell}{\ell!}. \end{aligned} \quad (28)$$

The last inequality of (27) follows from (22) and the elementary bound $\binom{n}{\ell} \leq \left(\frac{ne}{\ell}\right)^\ell$. $\square$

We refer to the recent work [2] for a systematic study of inequalities relating the Fourier growth with various well-studied properties of Boolean functions.

3. It is simple to observe (see also [15, Proposition 3.31]) that if $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ is a Boolean function and $h : \{-1, 1\}^n \rightarrow \mathbb{R}$ is an arbitrary function, then

$$\begin{aligned} \|\text{sign}(h) - f\|_{L_2}^2 &= 4\mathbb{P}\{\text{sign}(h) \neq f\} \\ &\leq 4\mathbb{P}\{|h - f| \geq 1\} \leq 4\|h - f\|_{L_2}^2, \end{aligned} \quad (29)$$

where we define $\text{sign}(0)$ as ± 1 arbitrarily. Therefore, applying Theorem 1 to a Boolean function, the above algorithm produces a Boolean function $\tilde{h} = \text{sign}(h)$ which is a 4ϵ -approximation of f .

4. In Theorem 1 we showed that functions $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d can be learned with accuracy at most ϵ and confidence at least $1 - \delta$ from $N = O_{\epsilon, d}(\log(n/\delta))$ random queries. We will now show that this estimate is sharp for small enough values of δ .

PROPOSITION 4. *Suppose that all the bounded linear functions $\ell : \{-1, 1\}^n \rightarrow [-1, 1]$ can be learned with accuracy at most $\frac{1}{2}$ and confidence at least $1 - \frac{1}{2n}$ from N random queries. Then $N > \log_2 n$.*

PROOF. By the assumption, for any input $(X_1, y_1), \dots, (X_N, y_N) \in \{-1, 1\}^n \times \{-1, 1\}$, there exists a function

$$h_{(X_1, y_1), \dots, (X_N, y_N)} : \{-1, 1\}^n \rightarrow \mathbb{R} \quad (30)$$

such that if the random vectors $X_1, \dots, X_N$ are chosen independently and uniformly from $\{-1, 1\}^n$ and there exists a linear function $\ell : \{-1, 1\}^n \rightarrow [-1, 1]$ such that $y_j = \ell(X_j)$ for every j , then $\mathbb{P}(\Omega_\ell) > 1 - \frac{1}{2n}$, where Ω_ℓ is the event

$$\Omega_\ell \stackrel{\text{def}}{=} \left\{ \mathbb{E}(h_{(X_1, \ell(X_1)), \dots, (X_N, \ell(X_N))}) - \ell \right\}^2 < \frac{1}{2}. \quad (31)$$

Let $X_j = (X_j(1), \dots, X_j(n))$ for $j \in \{1, \dots, N\}$ and consider the event

$$\mathcal{W} = \{X_j(1) = X_j(2), \forall j \in \{1, \dots, N\}\}. \quad (32)$$

By the independence of the samples, we have $\mathbb{P}(\mathcal{W}) = \frac{1}{2^N}$. Therefore, if $N \leq \log_2 n$ and we consider the standard linear functions $r_i : \{-1, 1\}^n \rightarrow \{-1, 1\}$ given by $r_i(x) = x_i$, then

$$\mathbb{P}(\Omega_{r_1} \cap \Omega_{r_2}) > 1 - \frac{1}{n} \geq 1 - \frac{1}{2^N} = 1 - \mathbb{P}(\mathcal{W}), \quad (33)$$

which implies that $\Omega_{r_1} \cap \Omega_{r_2} \cap \mathcal{W} \neq \emptyset$. Choosing $X_1, \dots, X_N$ from this event and denoting by

$$h = h_{(X_1, X_1(1)), \dots, (X_N, X_N(1))} = h_{(X_1, X_1(2)), \dots, (X_N, X_N(2))}, \quad (34)$$

we deduce from the triangle inequality that

$$2 = \mathbb{E}(r_1 - r_2)^2 \leq 2\mathbb{E}(h - r_1)^2 + 2\mathbb{E}(h - r_2)^2 \stackrel{(31)}{<} 2 \quad (35)$$

which is clearly a contradiction. Therefore $N > \log_2 n$. $\square$

ACKNOWLEDGMENTS

We are very grateful to Assaf Naor for constructive feedback and to Lauritz Streck for useful discussions which led to Proposition 4. A. E. was supported by a Junior Research Fellowship from Trinity College, Cambridge. P. I. was partially supported by the NSF grants DMS-2152346 and CAREER-DMS-2152401.

REFERENCES

- [1] Frédéric Bayart, Daniel Pellegrino, and Juan Benigno Seoane-Sepúlveda. 2014. The Bohr radius of the n -dimensional polydisk is equivalent to $\sqrt{(\log n)/n}$. *Adv. Math.* 264 (2014), 726–746. <https://doi.org/10.1016/j.aim.2014.07.029>
- [2] Jarosław Blasiok, Peter Ivanov, Yaonan Jin, Chin Ho Lee, Rocco A. Servedio, and Emanuele Viola. 2021. Fourier growth of structured $\mathbb{F}_2$ -polynomials and applications. (2021). To appear in RANDOM 2021. Preprint available at <https://arxiv.org/abs/2107.10797>.
- [3] Ron Blei. 2001. *Analysis in integer and fractional dimensions*. Cambridge Studies in Advanced Mathematics, Vol. 71. Cambridge University Press, Cambridge. xx+556 pages. <https://doi.org/10.1017/CBO9780511543012>
- [4] Henri Frederic Bohnenblust and Einar Hille. 1931. On the absolute convergence of Dirichlet series. *Ann. of Math.* (2) 32, 3 (1931), 600–622. <https://doi.org/10.2307/1968255>
- [5] Peter Borwein and Tamás Erdélyi. 1995. *Polynomials and polynomial inequalities*. Graduate Texts in Mathematics, Vol. 161. Springer-Verlag, New York. x+480 pages. <https://doi.org/10.1007/978-1-4612-0793-1>
- [6] Jamilson R. Campos, Pablo Jiménez-Rodríguez, Gustavo A. Muñoz Fernández, Daniel Pellegrino, and Juan Benigno Seoane-Sepúlveda. 2015. On the real polynomial Bohnenblust-Hille inequality. *Linear Algebra Appl.* 465 (2015), 391–400. <https://doi.org/10.1016/j.laa.2014.09.040>
- [7] Andreas Defant, Leonhard Frerick, Joaquim Ortega-Cerdà, Myriam Ounaies, and Kristian Seip. 2011. The Bohnenblust-Hille inequality for homogeneous polynomials is hypercontractive. *Ann. of Math.* (2) 174, 1 (2011), 485–497. <https://doi.org/10.4007/annals.2011.174.1.13>
- [8] Andreas Defant, Mirosław Mastyło, and Antonio Pérez. 2019. On the Fourier spectrum of functions on Boolean cubes. *Math. Ann.* 374, 1-2 (2019), 653–680. <https://doi.org/10.1007/s00208-018-1756-y>
- [9] Andreas Defant and Pablo Sevilla-Peris. 2014. The Bohnenblust-Hille cycle of ideas from a modern point of view. *Funct. Approx. Comment. Math.* 50, 1, [2013 on table of contents] (2014), 55–127. <https://doi.org/10.7169/facm/2014.50.1.2>
- [10] Alexandros Eskenazis and Paata Ivanisvili. 2020. Polynomial inequalities on the Hamming cube. *Probab. Theory Related Fields* 178, 1-2 (2020), 235–287. <https://doi.org/10.1007/s00440-020-00973-y>
- [11] Siddharth Iyer, Anup Rao, Victor Reis, Thomas Rothvoss, and Amir Yehudayoff. 2021. Tight bounds on the Fourier growth of bounded functions on the hypercube. (2021). To appear in ECCO 2021. Preprint available at <https://arxiv.org/abs/2107.06309>.
- [12] Nathan Linial, Yishay Mansour, and Noam Nisan. 1993. Constant depth circuits, Fourier transform, and learnability. *J. Assoc. Comput. Mach.* 40, 3 (1993), 607–620. <https://doi.org/10.1145/174130.174138>

- [13] John Edensor Littlewood. 1930. On bounded bilinear forms in an infinite number of variables. *Q. J. Math.* os-1, 1 (01 1930), 164–174. <https://doi.org/10.1093/qmath/os-1.1.164> arXiv:<https://academic.oup.com/qjmath/article-pdf/os-1/1/164/4482719/os-1-1-164.pdf>
- [14] Yishay Mansour. 1994. *Learning Boolean Functions via the Fourier Transform*. Springer US, Boston, MA, 391–424. https://doi.org/10.1007/978-1-4615-2696-4_11
- [15] Ryan O'Donnell. 2014. *Analysis of Boolean functions*. Cambridge University Press, New York. xx+423 pages. <https://doi.org/10.1017/CBO9781139814782>