



Threshold solutions for the nonlinear Schrödinger equation

Luccas Campos, Luiz Gustavo Farah and Svetlana Roudenko

Abstract. We study the focusing NLS equation in \mathbb{R}^N in the mass-supercritical and energy-subcritical (or *intercritical*) regime, with H^1 data at the mass-energy threshold $\mathcal{ME}(u_0) = \mathcal{ME}(Q)$, where Q is the ground state. Previously, Duyckaerts–Merle studied the behavior of threshold solutions in the H^1 -critical case, in dimensions $N = 3, 4, 5$, later generalized by Li–Zhang for higher dimensions. In the intercritical case, Duyckaerts–Roudenko studied the threshold problem for the 3d cubic NLS equation.

In this paper, we generalize the results of Duyckaerts–Roudenko for any dimension and any power of the nonlinearity for the *entire* intercritical range. We show the existence of special solutions, Q^\pm , besides the standing wave $e^{it}Q$, which exponentially approach the standing wave in the positive time direction, but differ in its behavior for negative time. We classify solutions at the threshold level, showing either blow-up occurs in finite (positive and negative) time, or scattering in both time directions, or the solution is equal to one of the three special solutions above, up to symmetries. Our proof extends to the H^1 -critical case, thus, giving an alternative proof of the Li–Zhang result and unifying the critical and intercritical cases.

These results are obtained by studying the linearized equation around the standing wave and some tailored approximate solutions to the NLS equation. We establish important decay properties of functions associated to the spectrum of the linearized Schrödinger operator, which, in combination with modulational stability and coercivity for the linearized operator on special subspaces, allows us to use a fixed-point argument to show the existence of special solutions. Finally, we prove the uniqueness by studying exponentially decaying solutions to a sequence of linearized equations.

Contents

1. Introduction	1638
2. Notation and preliminaries	1644
3. The linearized equation	1647
4. Construction of special solutions	1664
5. Modulation	1669

6. Convergence to the standing wave above the ground state 1673
 7. Convergence to the standing wave below the ground state 1674
 8. Estimates on exponentially decaying solutions 1676
 9. Closure of the main theorems 1682
 A. Appendix 1684
 References 1706

1. Introduction

We consider the focusing energy-subcritical nonlinear Schrödinger (NLS) equation in $H^1(\mathbb{R}^N)$, $N \geq 1$:

$$(1.1) \quad \begin{cases} i \partial_t u + \Delta u + |u|^{p-1} u = 0, \\ u(x, 0) = u_0(x) \in H^1(\mathbb{R}^N), \end{cases}$$

where $1 + 4/N < p < 2^* - 1 := \begin{cases} 1 + 4/(N - 2), & N \geq 3, \\ +\infty, & N = 1, 2. \end{cases}$

When $N \geq 3$, we also consider the focusing energy-critical nonlinear Schrödinger equation in $\dot{H}^1(\mathbb{R}^N)$ with nonlinearity power $p_c := (N + 2)/(N - 2)$:

$$(1.2) \quad \begin{cases} i \partial_t u + \Delta u + |u|^{p_c-1} u = 0, \\ u(x, 0) = u_0(x) \in \dot{H}^1(\mathbb{R}^N). \end{cases}$$

The equation (1.1) is considered in the inhomogeneous space $H^1(\mathbb{R}^N)$ with the norm $\|f\|_{H^1} := \|f\|_{L^2} + \|\nabla f\|_{L^2}$, while (1.2) is studied in the homogeneous Sobolev space $\dot{H}^1(\mathbb{R}^N)$ with the norm $\|f\|_{\dot{H}^1} := \|\nabla f\|_{L^2}$.

Note that (1.1) and (1.2) are invariant under scaling. Namely, if u is a solution, then

$$u_\delta(x, t) = \delta^{2/(p-1)} u(\delta x, \delta^2 t)$$

is also a solution to the same equation, for any $\delta > 0$. Computing the homogeneous $\dot{H}^s(\mathbb{R}^N)$ norm yields

$$\|u_\delta(\cdot, 0)\|_{\dot{H}^s} = \delta^{s-(N/2-2/(p-1))} \|u(\cdot, 0)\|_{\dot{H}^s}.$$

Hence, the scale-invariant norm is $\dot{H}^{s_c}(\mathbb{R}^N)$, where

$$s_c = \frac{N}{2} - \frac{2}{p-1}.$$

The conditions on p are equivalent to $0 < s_c < 1$ in (1.1), and to $s_c = 1$ in (1.2).

In addition to scaling invariance, the equations (1.1) and (1.2) exhibit several symmetries, such as, space translation, time translation, phase rotation and time-reversal. Indeed, if $u(x, t)$ is such a solution, so is

$$e^{i\theta_0} u(x + x_0, t + t_0) \quad \text{or} \quad e^{i\theta_0} \bar{u}(x + x_0, t + t_0),$$

with $(\theta_0, x_0, t_0) \in [0, 2\pi) \times \mathbb{R}^N \times \mathbb{R}$.

All these symmetries leave the \dot{H}^{s_c} norm invariant. Another symmetry that does not have this characteristic is the *Galilean boost*, given by

$$e^{ix \cdot \xi_0} e^{-it|\xi_0|^2} u(x - 2\xi_0 t, t), \quad \xi_0 \in \mathbb{R}^N.$$

Moreover, if $u_0 \in \dot{H}^1(\mathbb{R}^N)$, then solutions to (1.1) and (1.2) conserve the *energy*

$$E(u(t)) := \frac{1}{2} \int |\nabla u(t)|^2 - \frac{1}{p+1} \int |u(t)|^{p+1} = E(u_0),$$

and whenever u_0 belongs to $L^2(\mathbb{R}^N)$, the *mass*

$$M(u(t)) := \int |u(t)|^2 = M(u_0)$$

and the *momentum*

$$P(u(t)) := \text{Im} \int \bar{u}(t) \nabla u(t) = P(u_0)$$

are also conserved.

The Cauchy problem for (1.1) was first studied by Ginibre and Velo [18]. Namely, for initial data $u_0 \in H^1(\mathbb{R}^N)$, there exists a non-empty maximal interval I and a unique local-in-time solution $u: \mathbb{R}^N \times I \rightarrow \mathbb{C}$ that belongs to $C_t^0 H_x^1(\mathbb{R}^N \times J)$ for every compact interval $J \subset I$. Moreover, the map $u_0 \mapsto u$ is uniformly continuous and u satisfies the Duhamel formula

$$u(t) = e^{it\Delta} u_0 + i \int_0^t e^{i(t-s)\Delta} |u|^{p-1} u(s) ds$$

for all $t \in I$. The solution is also known to be in $L_t^q W_x^{1,r}(\mathbb{R}^N \times J)$ for any Strichartz pair (q, r) (see Section 2).

In the energy-critical case, the Cauchy problem for (1.2) was first considered by Cazenave and Weissler [7]. They proved that, for initial data $u_0 \in \dot{H}^1$, there exists a unique solution defined on a maximal non-empty interval I , satisfying the corresponding Duhamel formula and belonging to $C_t^0 \dot{H}_x^1(\mathbb{R}^N \times J) \cap L_{t,x}^{2(N+2)/(N-2)}(\mathbb{R}^N \times J)$ for every compact interval $J \subset I$. Later works, see [10, 35, 40], proved that the map from the initial data to the solution is also uniformly continuous.

The impossibility to extend the solution to all times is related to the concept of *finite-time blow-up*. We say that a solution to (1.1) blows up in finite positive time $T > 0$ if

$$\lim_{t \nearrow T} \|\nabla u(t)\|_{L^2} = +\infty.$$

For the energy-critical case, given that the scale-invariant norm is $\dot{H}^1(\mathbb{R}^N)$, this criterion is not enough to preclude the possibility of continuing the solution. Rather, we say that a solution to (1.2) blows up in finite positive time $T > 0$ if

$$\int_0^T \int |u(x, t)|^{\frac{2(N+2)}{N-2}} dx dt = +\infty.$$

In a similar way, blow-up in finite negative time is defined.

Solutions to (1.1) and (1.2) can also exhibit a *scattering* behavior in the energy space. We say that a solution to (1.1) scatters forward in time if it is defined for any $t \in [0, +\infty)$ and there exists $\psi \in H^1(\mathbb{R}^N)$ such that

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} \psi\|_{H^1} = 0.$$

In the energy-critical case, the definition is the same, except that the $\dot{H}^1(\mathbb{R}^N)$ norm is used instead. Scattering backward in time is defined analogously.

The $L_{t,x}^{2(N+2)/(N-2s_c)}$ norm also plays an important role in the scattering theory (see, for instance, Cazenave [6], Chapter 7): solutions to either (1.1) or (1.2) defined on the time interval $[0, +\infty)$ scatter forward in time if

$$\int_0^{+\infty} \int |u(x, t)|^{\frac{2(N+2)}{N-2s_c}} dx dt < +\infty.$$

By the time-reversal symmetry, a similar scattering criterion backward in time can be obtained.

Besides finite-time blow-up and scattering, there is the concept of *standing waves*. Consider the elliptic equation

$$(1.3) \quad \Delta \psi - (1 - s_c) \psi + |\psi|^{p-1} \psi = 0.$$

It is known that, for $s_c < 1$, this equation admits the unique radial, positive solution in $H^1(\mathbb{R}^N)$, which we call the *ground state* and denote by $Q = Q_{p,N}$ (see Strauss [37], Berestycki, Lions and Peletier [4], Kwong [29] and also Tao [39], Appendix B, for a textbook exposition). If Q solves (1.3), then the standing wave

$$u(x, t) = e^{i(1-s_c)t} Q(x)$$

is a solution to (1.1) that neither blows up in finite time, nor scatters, in any time direction. On the other hand, if $s_c = 1$, since the equation (1.3) is invariant by scaling, the radial positive solution to (1.3) is not unique. In this case, an explicit solution is given by

$$Q_{\frac{2N}{N-2}, N}(x) := \frac{1}{\left(1 + \frac{|x|^2}{N(N-2)}\right)^{(N-2)/2}}.$$

This solution is commonly denoted by W , and we shall often do so.

A simple calculation shows that $W \in \dot{H}^1(\mathbb{R}^N)$ for any $N \geq 3$, and that $W \in L^2(\mathbb{R}^N)$ if and only if $N \geq 5$. As its subcritical counterpart, the static solution $u(x, t) = W(x)$ to (1.2) neither blows up in finite time, nor scatters, in any time direction.

Also, the following *Pohozaev* identities follow from (1.3):

$$(1.4) \quad \begin{aligned} \int |Q|^{p+1} &= \frac{2(p+1)}{N(p-1)} \int |\nabla Q|^2, \\ \int |Q|^2 &= \frac{(N-2)(p+1)-4}{(1-s_c)N(p-1)} \int |\nabla Q|^2, \quad \text{if } 0 < s_c < 1. \end{aligned}$$

Remark 1.1. The choice of the constant $(1-s_c)$ in (1.3) is only for convenience. If $s_c < 1$, we can modify Q and replace $(1-s_c)$ by any positive constant by scaling. Similarly, if $s_c = 1$, the choice of $Q_{2N/(N-2),N} = W$ is arbitrary, and we could have used any rescaled version of W . Since we will state our results up to a constant scaling (among other symmetries), there is no loss of generality.

The works of Weinstein [42] in the case $0 < s_c < 1$ and of Aubin [2] and Talenti [38] for $s_c = 1$ give the characterization of the ground state as the minimizer of

$$(1.5) \quad \|f\|_{L^{p+1}}^{p+1} \leq C_{N,p} \|\nabla f\|_{L^2}^{N(p-1)/2} \|f\|_{L^2}^{2-(N-2)(p-1)/2},$$

with equality if and only if $f(x) = e^{i\theta_0} Q(x + x_0)$, in the case $0 < s_c < 1$, or $f(x) = e^{i\theta_0} \lambda_0^{(N-2)/2} W(\lambda_0 x + x_0)$, if $s_c = 1$, for some $\theta_0 \in [0, 2\pi)$, $x_0 \in \mathbb{R}^N$ and $\lambda_0 > 0$. Here, $C_{N,p}$ is the sharp constant of the inequality (1.5). In the subcritical case, the inequality (1.5) is known as (one version of) the Gagliardo–Nirenberg inequality, and in the energy-critical case, it reduces to the classical Sobolev inequality.

The ground state is also associated with the *threshold* for a dichotomy between finite-time blow-up and scattering. The behavior of solutions below the ground state level is now well understood for both the focusing energy-critical and the energy-subcritical nonlinear Schrödinger equations. Indeed, for the Cauchy problem (1.2), solutions with $E(u_0) < E(W)$ were first considered by Kenig and Merle [27] in the radial setting for $N = 3, 4$ and 5, introducing the concentration-compactness and rigidity approach for dispersive models. They showed that if $\|\nabla u_0\|_{L^2} < \|\nabla W\|_{L^2}$, then the corresponding solution exists globally in time and scatters in both time directions. On the other hand, if $\|\nabla u_0\|_{L^2} > \|\nabla W\|_{L^2}$, then the corresponding solution blows up in finite positive and negative times (provided it is radial or of finite variance). Later, by exploiting a double Duhamel trick, Killip and Visan [28] proved the corresponding result for $N \geq 5$, removing the radial assumption. More recently, Dodson [11] proved the non-radial scattering in dimension $N = 4$ using a delicate control on the scaling parameter, given by long time Strichartz estimates and an interaction Morawetz estimate. It is worth noting that there are important differences regarding the dimension in the energy-critical problem: for example, in dimensions $N \geq 5$, the lack of regularity of the nonlinearity poses technical difficulties, which are handled using generalizations of classical estimates. However, the faster time decay of the linear Schrödinger operator helps obtaining extra space-time bounds on the solution. The linear decay is only double integrable in dimensions $N \geq 5$, diverges logarithmically in dimension $N = 4$, and diverges polynomially in dimension $N = 3$. The same situation occurs for the ground state, which belongs to L_x^2 in higher dimensions, but misses it by a logarithm divergence in dimension $N = 4$ and by a polynomial divergence in dimension $N = 3$. The non-radial scattering below the ground state for the focusing energy-critical nonlinear Schrödinger equation in dimension $N = 3$ still remains an open problem.

For the Cauchy problem (1.1), it was first studied by Holmer and Roudenko [23] in the 3d cubic radial case. In [22], for $0 < s_c < 1$, they consider the following scale-invariant quantities:

$$\mathcal{ME}(u(t)) = \frac{M(u(t))^{(1-s_c)/s_c} E(u(t))}{M(Q)^{(1-s_c)/s_c} E(Q)} = \mathcal{ME}(u_0),$$

and

$$\mathcal{M}\mathcal{G}(u(t)) = \frac{\|u_0\|_{L^2}^{(1-s_c)/s_c} \|\nabla u(t)\|_{L^2}}{\|Q\|_{L^2}^{(1-s_c)/s_c} \|\nabla Q\|_{L^2}}.$$

Based on the concentration-compactness and rigidity approach, in [22] they proved that if $\mathcal{M}\mathcal{E}(u_0) < 1$ and $\mathcal{M}\mathcal{G}(u_0) < 1$, then the corresponding solution exists globally in time and scatters in both time directions. In [22] they also proved that if $\mathcal{M}\mathcal{E}(u_0) < 1$, $\mathcal{M}\mathcal{G}(u_0) > 1$ and either u_0 is radial or $|x|u_0 \in L^2(\mathbb{R}^N)$, then the corresponding solution blows up in both finite positive and negative times¹, establishing the dichotomy result in the 3d cubic radial case. Later Duyckaerts, Holmer and Roudenko [23] removed the radial assumption in the scattering result. Fang, Xie and Cazenave [15] and Guevara [20] (see also Guevara and Carreon [19]) extended this result to all intercritical ranges and dimensions.

This dichotomy does not hold above the ground state mass-energy threshold. In [21], Holmer, Platte and Roudenko proved blow-up criteria that included solutions above the mass-energy threshold. In [14], Duyckaerts and Roudenko showed, for $0 < s_c \leq 1$, the existence of asymmetric behavior in time of solutions to the NLS equation that are above the threshold and that scatter in one time direction and blow-up in finite time in the other time direction (in fact, they showed that it suffices to multiply the ground state by a quadratic phase to produce such a result). Moreover, they proved a dichotomy-type result above the mass-energy threshold with some conditions on the variance of the initial data.

At the threshold level, there exists a richer dynamics for the asymptotic behavior of solutions. Indeed, for the focusing energy-critical NLS equation, this problem was first considered by Duyckaerts and Merle [12], in the radial case for $N = 3, 4$ and 5 . In particular, they proved the existence of special solutions that approach W as $t \rightarrow +\infty$ and either blow-up or scatter as $t \rightarrow -\infty$. Later, Li and Zhang [30] studied the case $N \geq 6$. For the focusing energy-subcritical nonlinear Schrödinger equation, Duyckaerts and Roudenko [13] treated the 3d cubic case. The main goal of this paper is to generalize the results in [13] to the entire intercritical range $0 < s_c < 1$. More precisely, we prove the following.

Theorem 1.2 (Energy-subcritical case). *For $N \geq 1$, there exist two radial solutions Q^+ and Q^- to (1.1) in $H^1(\mathbb{R}^N)$ such that*

- $M[Q^\pm] = M[Q]$, $E[Q^\pm] = E[Q]$, Q^\pm is defined at least on $[0, +\infty)$ and there exist $C, e_0 > 0$ such that

$$\|Q^\pm(t) - e^{it}Q\|_{H^1} \leq Ce^{-e_0t} \quad \text{for all } t \geq 0,$$

- $\|\nabla Q_0^+\|_2 > \|\nabla Q\|_2$, and Q^+ blows-up in finite negative time,
- $\|\nabla Q_0^-\|_2 < \|\nabla Q\|_2$, and Q^- is globally defined and scatters backward in time.

Theorem 1.3 (Energy-subcritical case). *For $N \geq 1$, let u be a solution to (1.1) such that $\mathcal{M}\mathcal{E}(u_0) = 1$. Then, the following holds.*

- If $\mathcal{M}\mathcal{G}(u_0) < 1$, then u is defined for all times. Moreover, either u scatters in both time directions, or $u = Q^-$ up to the symmetries of the equation.

¹If u_0 is nonradial and has infinite variance, then there exists a sequence of times $\{t_n\}$ such that $\|\nabla u(t_n)\|_{L^2} \rightarrow +\infty$, as shown by Holmer and Roudenko in [24].

- If $\mathcal{ME}(u_0) = 1$, then $u = Q$ up to the symmetries of the equation.
- If $\mathcal{ME}(u_0) > 1$ and u_0 is either radial or $|x|u_0 \in L^2(\mathbb{R}^N)$, then either u blows up in finite positive and negative time, or $u = Q^\pm$ up to the symmetries of the equation.

Remark 1.4. In the 3d cubic setting and for $\mathcal{ME}(u_0) < 1$, the infinite-variance case was treated by Holmer and Roudenko in [24]. It was shown that, in this case, the H^1 norm of the solution is unbounded, even if the solution exists for all $t > 0$. That proof works the same way if one considers any $0 < s_c < 1$ and $\mathcal{ME}(u_0) \leq 1$.

There are two major difficulties in extending the previous results. The first one is to deal with low powers of the parameter p . If $p < 3$, then the nonlinearity $|u|^{p-1}u$ is not a smooth function of (u, \bar{u}) . Moreover, as the power of the nonlinearity is not an odd integer, the difference $|u|^{p-1}u - |v|^{p-1}v$ cannot be written as a polynomial. Therefore, we cannot use the same estimates as in [13], as they rely heavily on $H^s(\mathbb{R}^N)$ estimates, for large values² of s . Moreover, if $p \leq 2$, then the nonlinearity is not twice real-differentiable. In order to perform the necessary estimates, we employ the fractional calculus tools introduced by Christ and Weinstein [9] and Visan [41].

Another problem arises from the fast decay of the ground state Q . When constructing the solutions Q^\pm , we must deal with some estimates that involve terms of the form $\|Q^{-1}f\|_{L^\infty}$. Even though $(Q^{-1}f)(x)$ is pointwise defined for any function f , the exponential decay of Q makes it harder to obtain good estimates. Therefore, we have to carefully study the desired functions f to ensure that they have the necessary decay. We establish the decay via several bootstrap arguments, and by making use of resolvent convolution kernels associated to the corresponding elliptic equations.

It is worth mentioning that, in order to prove the classification in Theorems 1.2 and 1.3, one has to change the orthogonality conditions that were used in [13], as in the lower dimensions they would not necessarily ensure coercivity. See Remark 3.4 and the proof of Lemma 3.5 for details.

Since our proof can be readily applied to the energy-critical case, we also state and prove similar results for $s_c = 1$ and $N \geq 6$. Note that $p_c \leq 2$ happens exactly when $N \geq 6$, which again implies that the nonlinearity is not twice real-differentiable. We have the following.

Theorem 1.5 (Energy-critical case). *Let $N \geq 6$. There exist two radial solutions W^+ and W^- to (1.2) in $\dot{H}^1(\mathbb{R}^N)$ such that*

- $E[W^\pm] = E[W]$, W^\pm is defined at least in $[0, +\infty)$ and there exist $C, e_0 > 0$ such that

$$\|W^\pm(t) - W\|_{H^1} \leq Ce^{-e_0t} \quad \text{for all } t \geq 0,$$

- $\|\nabla W_0^+\|_2 > \|\nabla W\|_2$, and W^+ blows-up in finite negative time,
- $\|\nabla W_0^-\|_2 < \|\nabla W\|_2$, and W^- is globally defined and scatters backward in time.

Theorem 1.6 (Energy-critical case). *For $N \geq 6$, let u be a radial solution to (1.2) such that $E(u_0) = E(W)$. Then, the following holds.*

²To be precise, at least $s > N/2$, to make use of the fact that $H^s(\mathbb{R}^N)$ is an algebra.

- If $\|u_0\|_{\dot{H}^1} < \|W\|_{\dot{H}^1}$, then u is defined for all times. Moreover, either u scatters in both time directions, or $u = W^-$ up to the symmetries of the equation.
- If $\|u_0\|_{\dot{H}^1} = \|W\|_{\dot{H}^1}$, then $u = W$ up to the symmetries of the equation.
- If $\|u_0\|_{\dot{H}^1} > \|W\|_{\dot{H}^1}$ and $u_0 \in L^2$, then either u blows-up in finite positive and negative time, or $u = W^+$ up to the symmetries of the equation.

The last two theorems were originally proved by Li and Zhang [30], by means of weighted Sobolev estimates. Since our approach is considerably different from [30], we include our proofs in this paper.

Remark 1.7. By scaling, the condition $\mathcal{ME}(u_0) = 1$ can be read, without loss of generality, as

$$\begin{cases} M(u_0) = M(Q), \\ E(u_0) = E(Q). \end{cases}$$

Indeed, considering $u_{0,\delta}(x) = \delta^{2/(p-1)}u_0(\delta x)$, with $\delta = (M(u_0)/M(Q))^{1/(2s_c)}$, gives the above condition for $u_{0,\delta}$. Similarly, the condition

$$\mathcal{ME}(u_0) < 1$$

(resp. “=”, “>”) can be read as

$$\|\nabla u_0\|_{L^2} < \|\nabla Q\|_{L^2}$$

(resp. “=”, “>”). Unless stated otherwise, we shall adopt this simplification throughout the whole paper.

2. Notation and preliminaries

We will need the following tools from harmonic analysis.

Lemma 2.1 (Sobolev inequality, see Stein [36]). *If $0 < \rho - \sigma < N$, $1 < q < p < \infty$, and*

$$\frac{1}{p} = \frac{1}{q} - \frac{\rho - \sigma}{N},$$

then the following estimate holds:

$$\|D^\sigma u\|_{L^p(\mathbb{R}^N)} \leq C \|D^\rho u\|_{L^q(\mathbb{R}^N)}.$$

Lemma 2.2 (Leibniz rule, see Christ and Weinstein [9]). *Let $s \in (0, 1)$, $p_j, q_j \in (1, \infty)$, with $1/p = 1/p_j + 1/q_j$, $j = 1, 2$. Then*

$$(2.1) \quad \|D^s(fg)\|_{L^p(\mathbb{R}^N)} \leq C (\|D^s f\|_{L^{p_1}(\mathbb{R}^N)} \|g\|_{L^{q_1}(\mathbb{R}^N)} + \|f\|_{L^{p_2}(\mathbb{R}^N)} \|D^s g\|_{L^{q_2}(\mathbb{R}^N)}).$$

Lemma 2.3 (Fractional chain rule for Hölder continuous functions, see Visan [41]). *Let F be a Hölder continuous function of order $0 < \alpha < 1$. Then, for every $0 < s < \alpha$, $1 < p < \infty$, and $s/\alpha < v < 1$, we have*

$$(2.2) \quad \|D^s F(u)\|_{L^p(\mathbb{R}^N)} \leq C \| |u|^{\alpha-s/v} \|_{L^{p_1}(\mathbb{R}^N)} \|D^v u\|_{L^{\frac{p}{v}q_1}(\mathbb{R}^N)}^{s/v},$$

provided $1/p = 1/p_1 + 1/q_1$ and $(1 - s/(v\alpha))p_1 > 1$.

Lemma 2.4 (Kato–Strichartz inequalities, see Cazenave [6], Kato [25], Foschi [16], Keel and Tao [26]). *Let $N \geq 1$, $I \subset \mathbb{R}$ and $1 \leq q_i, r_i \leq \infty$, $i = 1, 2$. Assume the pairs (q_1, r_1) and (q_2, r_2) satisfy*

$$\frac{1}{q_i} < N \left(\frac{1}{2} - \frac{1}{r_i} \right) \quad \text{or} \quad (q_i, r_i) = (\infty, 2), \quad i = 1, 2, \quad \frac{1}{q_1} + \frac{1}{q_2} = N \left(1 - \frac{1}{r_1} - \frac{1}{r_2} \right),$$

and

- if $N = 2$, we require that $r_1, r_2 < +\infty$,
- if $N > 2$, we consider two subcases,
 - non sharp case:

$$\frac{1}{q_1} + \frac{1}{q_2} < 1, \quad \frac{N-2}{r_1} \leq \frac{N}{r_2}, \quad \frac{N-2}{r_1} \leq \frac{N}{r_2},$$

- sharp case:

$$\frac{1}{q_1} + \frac{1}{q_2} = 1, \quad \frac{N-2}{r_1} < \frac{N}{r_2}, \quad \frac{N-2}{r_1} < \frac{N}{r_2}, \quad \frac{1}{r_1} \leq \frac{1}{q_1}, \quad \frac{1}{r_2} \leq \frac{1}{q_2}.$$

Then the following estimate holds:

$$(2.3) \quad \left\| \int_{s>t} e^{i(t-s)\Delta} F(s) ds \right\|_{L_I^{q_1} L_x^{r_1}} + \left\| \int_{s<t} e^{i(t-s)\Delta} F(s) ds \right\|_{L_I^{q_1} L_x^{r_1}} \lesssim \|F\|_{L_I^{q_2} L_x^{r_2}}.$$

Definition 2.5. We say that the pair (q, r) is \dot{H}^s -admissible if $2 \leq q, r \leq +\infty$, $(q, r, N) \neq (2, \infty, 2)$, and

$$\frac{2}{q} + \frac{N}{r} = \frac{N}{2} - s.$$

If $s = 0$, we say that the pair (q, r) is L^2 -admissible.

We define Strichartz norms for the energy-critical and intercritical cases separately.

Definition 2.6 (Critical case). Let I be a (possibly unbounded) time interval. Given $0 < \varepsilon \ll \frac{4}{N-2}$, $N \geq 6$, define the spaces

$$\begin{aligned} S(\dot{H}^1, I) &= L_I^\infty L_x^{\frac{2N}{N-2}} \\ S(\dot{H}^{1-\varepsilon}, I) &= L_I^\infty L_x^{\frac{2N}{N-2+2\varepsilon}} \cap L_I^{4/\varepsilon} L_x^{\frac{2N}{N-2+\varepsilon}} \cap L_I^{\frac{2(N-2)}{\varepsilon(N-4)}} L_x^{\frac{2N(N-2)}{(N-2)^2+4\varepsilon}}, \\ S'(\dot{H}^{-(1-\varepsilon)}, I) &= L_I^{2/\varepsilon} L_x^{\frac{2N}{N+2}}, \\ S(L^2, I) &= \{L_I^q L_x^r \mid (q, r) \text{ is } L^2\text{-admissible}\}, \\ S'(L^2, I) &= L_I^2 L_x^{\frac{2N}{N+2}}. \end{aligned}$$

Remark 2.7. In particular, we make use of the following spaces in $S(L^2, I)$: $L_I^\infty L_x^2$,

$$L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{N-\varepsilon}}, L_I^{\frac{2(N-2)}{\varepsilon(N-4)}} L_x^{\frac{2N(N-2)}{N(N-2)-2\varepsilon(N-4)}}, L_I^2 L_x^{\frac{2N}{N-2}}, L_I^{\frac{2(N+2)}{N-2}} L_x^{\frac{2(N+2)}{N^2+4}}, \text{ and } L_I^{\frac{16}{\varepsilon(N-2)}} L_x^{\frac{8N}{4N-\varepsilon(N-2)}}.$$

Remark 2.8. By the Sobolev embedding, if $f \in S(\dot{H}^1, I) \cap \nabla^{-1}S(L^2, I)$,

$$\|f\|_{S(\dot{H}^1, I)} + \|D^\varepsilon f\|_{S(\dot{H}^{1-\varepsilon}, I)} \lesssim \|\nabla f\|_{S(L^2, I)},$$

and by the Kato–Strichartz estimates, non sharp case,

$$\begin{aligned} \left\| \int_{s>t} e^{i(t-s)\Delta} F(s) ds \right\|_{S(L^2, I)} &\lesssim \|F\|_{S'(L^2, I)}, \\ \left\| \int_{s>t} e^{i(t-s)\Delta} F(s) ds \right\|_{S(\dot{H}^{1-\varepsilon}, I)} &\lesssim \|F\|_{S'(\dot{H}^{-(1-\varepsilon)}, I)}. \end{aligned}$$

Remark 2.9. Note that the pair in $S(\dot{H}^1, I)$ is \dot{H}^1 -admissible, the pairs in $S(\dot{H}^{1-\varepsilon}, I)$ are $\dot{H}^{1-\varepsilon}$ -admissible, the pairs in $S(L^2, I)$ and in the dual space of $S'(L^2, I)$ are L^2 -admissible, and the pair corresponding to the dual space of $S'(\dot{H}^{-(1-\varepsilon)}, I)$ is $\dot{H}^{-(1-\varepsilon)}$ -admissible.

Definition 2.10 (Intercritical case). Define the set

$$\mathcal{A}_0 = \{(q, r) | (q, r) \text{ is } L^2\text{-admissible}\}.$$

For $s \in (0, 1)$, define \mathcal{A}_s as the \dot{H}^s -admissible pairs that satisfy

$$\begin{cases} \frac{2N}{N-2s} \leq r \leq \left(\frac{2N}{N-2}\right)^-, & N \geq 3, \\ \frac{2}{1-s} \leq r \leq \left(\left(\frac{2}{1-s}\right)^+\right)', & N = 2, \\ \frac{2}{1-2s} \leq r \leq \infty, & N = 1, \end{cases}$$

and \mathcal{A}_{-s} as the \dot{H}^{-s} -admissible pairs that satisfy

$$\begin{cases} \left(\frac{2N}{N-2s}\right)^+ \leq r \leq \left(\frac{2N}{N-2}\right)^-, & N \geq 3, \\ \left(\frac{2}{1-s}\right)^+ \leq r \leq \left(\left(\frac{2}{1-s}\right)^+\right)', & N = 2, \\ \left(\frac{2}{1-2s}\right)^+ \leq r \leq \infty & N = 1, \end{cases}$$

where $(a^+)'$ is the number such that

$$\frac{1}{a} = \frac{1}{a^+} + \frac{1}{(a^+)'}.$$

Let I be a (possibly unbounded) time interval. For $s \in [0, 1)$, we define the following Strichartz norms:

$$\|u\|_{S(L^2, I)} = \sup_{(q, r) \in \mathcal{A}_0} \|u\|_{L_t^q L_x^r}, \quad \text{and} \quad \|u\|_{S(\dot{H}^s, I)} = \sup_{(q, r) \in \mathcal{A}_s} \|u\|_{L_t^q L_x^r},$$

and the dual Strichartz norms

$$\|u\|_{S'(L^2, I)} = \inf_{(q, r) \in \mathcal{A}_0} \|u\|_{L_t^{q'} L_x^{r'}}, \quad \text{and} \quad \|u\|_{S'(\dot{H}^{-s}, I)} = \inf_{(q, r) \in \mathcal{A}_{-s}} \|u\|_{L_t^{q'} L_x^{r'}}.$$

Remark 2.11. By the Sobolev embedding, if $f \in S(\dot{H}^{s_c}, I) \cap \langle \nabla \rangle^{-1} S(L^2, I)$,

$$\|f\|_{S(\dot{H}^{s_c}, I)} \lesssim \| |\nabla|^{s_c} f \|_{S(L^2, I)} \lesssim \|\langle \nabla \rangle f\|_{S(L^2, I)},$$

and by the Kato–Strichartz estimates, non sharp case,

$$\begin{aligned} \left\| \int_{s>t} e^{i(t-s)\Delta} F(s) ds \right\|_{S(L^2, I)} &\lesssim \|F\|_{S'(L^2, I)}, \\ \left\| \int_{s>t} e^{i(t-s)\Delta} F(s) ds \right\|_{S(\dot{H}^{s_c}, I)} &\lesssim \|F\|_{S'(\dot{H}^{-s_c}, I)}. \end{aligned}$$

3. The linearized equation

In order to prove the main theorems of this paper, we need to carefully study (1.1) and (1.2) around the ground state. We identify the complex number $a + bi$ with the vector $\begin{pmatrix} a \\ b \end{pmatrix}$. For a complex-valued function f , we write $f = f_1 + if_2$. We next introduce the following definition.

Definition 3.1. For $0 < s_c \leq 1$, we define

$$\begin{aligned} L_+ &= (1 - s_c) - \Delta - pQ^{p-1}, \quad L_- = (1 - s_c) - \Delta - Q^{p-1}, \quad \mathcal{L} := \begin{pmatrix} 0 & -L_- \\ L_+ & 0 \end{pmatrix}, \\ R(f) &= |Q + f|^{p-1}(Q + f) - Q^p - pQ^{p-1}f_1 - iQ^{p-1}f_2, \\ K(f) &= pQ^{p-1}f_1 + iQ^{p-1}f_2. \end{aligned}$$

If u is a solution to (1.1), write $u = e^{i(1-s_c)t}(Q + v)$. Then v must satisfy

$$(3.1) \quad \partial_t v + \mathcal{L}v = iR(v),$$

or, writing it as a Schrödinger equation,

$$(3.2) \quad i\partial_t v + \Delta v - (1 - s_c)v + K(v) = -R(v).$$

In the next two sections we recall some properties of the operator \mathcal{L} .

3.1. The linearized operator

For $0 < s_c \leq 1$, we have, by a direct calculation,

$$L_-(Q) = 0 \quad \text{and} \quad L_+(\partial_k Q) = 0, \quad 1 \leq k \leq N.$$

This implies

$$\mathcal{L}(\partial_k Q) = \mathcal{L}(iQ) = 0, \quad 1 \leq k \leq N.$$

Also, defining Λf as the scaling generator $\frac{2}{p-1}f + x \cdot \nabla f$, we have

$$L_+(\Lambda Q) = -2(1 - s_c)Q, \quad \text{if } 0 < s_c < 1, \quad \text{and} \quad L_+(\Lambda W) = 0, \quad \text{if } s_c = 1.$$

The above directions are obtained from Q by the symmetries of the NLS equation. Indeed, defining

$$f_{[x_0, \lambda_0, \theta_0]}(x) = e^{i\theta_0} \frac{1}{\lambda_0^{2/(p-1)}} f\left(\frac{x}{\lambda_0} + x_0\right),$$

we have

$$(\nabla Q, -\Lambda Q, iQ) = \frac{\partial Q_{[x_0, \lambda_0, \theta_0]}}{\partial (x_0, \lambda_0, \theta_0)} \Big|_{(x_0, \lambda_0, \theta_0) = (0, 1, 0)}.$$

We also define the bilinear form

$$\begin{aligned} B(f, g) &:= \frac{1}{2} (L_+ f_1, g_1)_{L^2} + \frac{1}{2} (L_- f_2, g_2)_{L^2} \\ &= \frac{1-s_c}{2} \int f_1 \cdot g_1 + \frac{1}{2} \int \nabla f_1 \cdot \nabla g_1 - \frac{p}{2} \int Q^{p-1} f_1 g_1 \\ &\quad + \frac{1-s_c}{2} \int f_2 \cdot g_2 + \frac{1}{2} \int \nabla f_2 \cdot \nabla g_2 - \frac{1}{2} \int Q^{p-1} f_2 g_2, \end{aligned}$$

and the linearized energy

$$\begin{aligned} \Phi(f) &:= B(f, f) = \frac{1}{2} (L_+ f_1, f_1)_{L^2} + \frac{1}{2} (L_- f_2, f_2)_{L^2} \\ &= \frac{1-s_c}{2} \int |f|^2 + \frac{1}{2} \int |\nabla f|^2 - \frac{1}{2} \int Q^{p-1} (p|f_1|^2 + |f_2|^2). \end{aligned}$$

If $0 < s_c \leq 1$, one can check directly that, for any $f, g \in S(\mathbb{R}^N)$,

$$\begin{aligned} (3.3) \quad &B(f, g) = B(g, f), \quad B(\mathcal{L}f, g) = -B(f, \mathcal{L}g), \\ &B(iQ, f) = 0, \quad B(\partial_k Q, f) = 0, \quad k \leq N, \\ &B(\Lambda Q, f) = -(1-s_c) \int Q f_1, \quad \Phi(\mathcal{Y}_+) = \Phi(\mathcal{Y}_-) = 0. \end{aligned}$$

In the energy-critical case, we have

$$B(\Lambda W, f) = 0.$$

The following result is well known, but for completeness we provide its proof here.

Lemma 3.2 ([8, 12]). *Let $\sigma(\mathcal{L})$ be the spectrum of the operator \mathcal{L} , defined in $L^2(\mathbb{R}^N) \times L^2(\mathbb{R}^N)$ with domain $H^2(\mathbb{R}^N) \times H^2(\mathbb{R}^N)$, and let $\sigma_{\text{ess}}(\mathcal{L})$ be its essential spectrum. Then*

$$\sigma_{\text{ess}}(\mathcal{L}) = \{iy : y \in \mathbb{R}, |y| \geq 1 - s_c\}, \quad \sigma \cap \mathbb{R} = \{-e_0, 0, e_0\} \quad \text{with } e_0 > 0.$$

Moreover, e_0 and $-e_0$ are simple eigenvalues of \mathcal{L} with eigenfunctions \mathcal{Y}_+ and $\mathcal{Y}_- = \overline{\mathcal{Y}_+} \in \mathcal{S}$, respectively. The null space of \mathcal{L} is spanned by iQ and $\partial_k Q$, $1 \leq k \leq N$ (and, in the energy-critical case, also by ΛW).

Remark 3.3. By Lemma 3.2, if $\mathcal{Y}_1 = \text{Re}(\mathcal{Y}_+)$ and $\mathcal{Y}_2 = \text{Im}(\mathcal{Y}_+)$, then

$$L_+ \mathcal{Y}_1 = e_0 \mathcal{Y}_2 \quad \text{and} \quad L_- \mathcal{Y}_2 = -e_0 \mathcal{Y}_1.$$

Furthermore, the null space of L_+ is spanned by the vectors $\partial_k Q$, $1 \leq k \leq N$ (and by ΛW , if $s_c = 1$), and the null space of L_- is spanned by Q .

Proof of Lemma 3.2. In this proof, for brevity, we write $V = Q^{p-1}$ for $0 < s_c \leq 1$. Note that V defines a compact operator from H^1 to L^2 .

Intercritical case. For convenience (for example, as in Holmer–Roudenko [22] or Weinstein [42]), from now on, we rescale Q in the intercritical case to have unit coefficients in the ground state equation (1.3), i.e.,

$$(3.4) \quad \Delta Q - Q + Q^p = 0.$$

This rescaling is done in order to keep the exposition more transparent in computations. The term $(1 - s_c)$ is thus replaced by 1 in the standing wave solution $e^{i(1-s_c)t} Q$ and in the definitions of \mathcal{L} , B and Φ as well.

The operator \mathcal{L} is a relatively compact perturbation of $i(1 - \Delta)$, and therefore, has the same essential spectrum. We now prove the existence of exactly one negative eigenvalue to \mathcal{L} .

From the proof of Lemma 3.5, we see that L_- on L^2 with domain H^2 is non-negative. Since it is also self-adjoint, it has a unique square root $L_-^{1/2}$ with domain H^1 . It is equivalent to show that the self-adjoint operator $P := L_-^{1/2} L_+ L_-^{1/2}$ on L^2 with domain H^4 has a unique negative eigenvalue. Indeed, consider the function

$$Z = \Lambda Q - \frac{(\Lambda Q, Q)_{L^2}}{(Q, Q)_{L^2}} Q.$$

One can check that $Z \in H^2$, $Z \in \{Q\}^\perp$ and, for $0 < s_c \leq 1$,

$$(3.5) \quad (L_+ Z, Z)_{L^2} = -\frac{N^2(p-1)}{4(p+1)} \left[p - \left(1 + \frac{4}{N} \right) \right] \int Q^{p+1} < 0.$$

Defining $h := L_-^{-1/2} Z \in Q^\perp$, one also has

$$h = (L_-^{1/2} L_-^{-1})(L_-^{-1} L_-) Z = L_-^{-1} L_-^{-1/2} L_- Z \in H^3.$$

For $\varepsilon > 0$, choose $\tilde{h}_\varepsilon \in H^4$ such that $\tilde{h}_\varepsilon \perp Q$ and $\|h - \tilde{h}_\varepsilon\|_{H^3} < \varepsilon$. We have

$$\inf_{f \in H^4} \frac{(Pf, f)_{L^2}}{\|f\|_{L^2}^2} \leq \frac{(L_+ L_-^{1/2} \tilde{h}_\varepsilon, L_-^{1/2} \tilde{h}_\varepsilon)_{L^2}}{\|\tilde{h}_\varepsilon\|_{L^2}^2} < 0,$$

if ε is small enough.

Hence, by the minimax principle, P has a negative eigenvalue $-e_0^2$ and an associated eigenfunction g . Defining $\mathcal{Y}_1 := L_-^{1/2} g$, $\mathcal{Y}_2 := \frac{1}{e_0} L_+ \mathcal{Y}_1$, and $\mathcal{Y}_\pm := \mathcal{Y}_1 \pm i \mathcal{Y}_2$, we have $\mathcal{L} \mathcal{Y}_\pm = \pm e_0 \mathcal{Y}_\pm$. Uniqueness of the eigenfunction corresponding to the negative eigenvalue of P follows from the non-negativity of L_+ on $\{Q^p\}^\perp$, since

$$(Pf, f)_{L^2} = (L_+ L_-^{1/2} f, L_-^{1/2} f) \geq 0$$

for $f \in H^4$ such that $(L_-^{1/2} f, Q^p) = (f, L_-^{1/2} Q^p) = 0$. The assertions about the kernel of \mathcal{L} follow from the coercivity given by Lemma 3.5.

It remains to prove that $\mathcal{Y}_\pm \in S(\mathbb{R}^N)$. It suffices to prove this assertion for $\mathcal{Y}_1 = \text{Re } \mathcal{Y}_+$. The differential equation for \mathcal{Y}_1 is

$$(3.6) \quad [(1 - \Delta)^2 + e_0^2] \mathcal{Y}_1 = [pV^2 + V(1 - \Delta)] \mathcal{Y}_1 - p(1 - \Delta)[V \mathcal{Y}_1].$$

Since the Fourier symbol of $(1 - \Delta)^2 + e_0^2$ is $(1 + |\xi|^2)^2 + e_0^2 \approx (1 + |\xi|^2)^2$, and $V, \mathcal{Y}_1 \in H^2(\mathbb{R}^N)$, we have that $\mathcal{Y}_1 \in H^s$ for all $s \geq 0$. As in [12], we show that for all non-negative integers k, s and all $\varphi \in C_c^\infty(\mathbb{R}^N)$, we have

$$(3.7) \quad \|\varphi(x/R)\mathcal{Y}_1\|_{H^s} \leq \frac{C(\varphi, s, k)}{R^k} \quad \text{for all } R \geq 1.$$

Indeed, the inequality (3.7) holds if $k = 0$, for any $s \geq 0$. By induction, we show that if it holds for (k, s) , it also holds for $(k + 1, s + 1)$. Given φ , consider $\tilde{\varphi} \in C_c^\infty(\mathbb{R}^N)$ such that $\tilde{\varphi}$ is 1 on the support of φ , so that we have $\tilde{\varphi} \partial^\alpha \varphi = \partial^\alpha \varphi$ for any multi-index α . Since Q and its derivatives decay (more than) polynomially, (3.6) gives, for $s \geq 3$,

$$\|\varphi(x/R)[(1 - \Delta)^2 + e_0^2]\mathcal{Y}_1\|_{H^{s-3}} \leq \frac{C}{R} \|\tilde{\varphi}(x/R)\mathcal{Y}_1\|_{H^{s-1}} \leq \frac{C}{R} \|\tilde{\varphi}(x/R)\mathcal{Y}_1\|_{H^s}.$$

Using the trivial commutator estimate $\|[(1 - \Delta)^2 + e_0^2, \phi(x/R)]\|_{H^{s-3} \rightarrow H^s} \leq C/R$, we get

$$\|\varphi(x/R)\mathcal{Y}_1\|_{H^{s+1}} \approx \|[(1 - \Delta)^2 + e_0^2](\varphi(x/R)\mathcal{Y}_1)\|_{H^{s-3}} \leq \frac{C}{R} \|\tilde{\varphi}(x/R)\mathcal{Y}_1\|_{H^s}.$$

By the induction hypothesis, we get $\|\varphi(x/R)\mathcal{Y}_1\|_{H^{s+1}} \leq C/R^{k+1}$, as desired. The same argument shows that, if $\lambda \in \mathbb{R} \setminus \sigma(\mathcal{L})$, then $(\lambda - \mathcal{L})^{-1}S(\mathbb{R}^N) \subset S(\mathbb{R}^N)$.

Critical case. The range of the operator L_- is no longer closed, but the operator $1 + L_-$ is invertible on $\{Q\}^\perp$. Therefore, for any $\varepsilon > 0$, one can take $G_\varepsilon \in H^2$ such that

$$\|L_-G_\varepsilon - (1 + L_-)Z\|_{L^2} < \varepsilon.$$

Letting

$$\begin{aligned} h_\varepsilon &:= (1 + L_-)^{-1}L_-^{1/2}G_\varepsilon = L_-^{1/2}(1 + L_-)^{-1}G_\varepsilon \\ &= (1 + L_-)^{-1}(1 + L_-)^{-1/2}L_-^{1/2}(1 + L_-)^{-1/2}G_\varepsilon \in H^3, \end{aligned}$$

we have

$$\begin{aligned} \|L_-^{1/2}h_\varepsilon - Z\|_{H^2} &= \|(1 - \Delta)(1 + L_-)^{-1}[L_-G_\varepsilon - (1 + L_-)Z]\|_{L^2} \\ &\leq \varepsilon \| [1 - V(1 - \Delta)^{-1}]^{-1} \|_{L^2 \rightarrow L^2}. \end{aligned}$$

Choosing $\tilde{h}_\varepsilon \in H^4$ such that $\tilde{h}_\varepsilon \perp Q$ and $\|h_\varepsilon - \tilde{h}_\varepsilon\|_{H^3} < \varepsilon$, and recalling $P = L_-^{1/2}L_+L_-^{1/2}$ and (3.5), we get

$$(P\tilde{h}_\varepsilon, \tilde{h}_\varepsilon)_{L^2} = (L_+Z, Z)_{L^2} + O(\varepsilon).$$

Thus, if ε is small enough, the conclusion follows. The regularity and the decay of \mathcal{Y}_\pm follow analogously from the argument for the intercritical case. ■

If $0 < s_c < 1$, consider the following orthogonality relations:

$$(3.8) \quad \int Qv_2 = \int \partial_k Qv_1 = 0, \quad 1 \leq k \leq N,$$

$$(3.9) \quad \int Q^p v_1 = 0,$$

$$(3.10) \quad \int \mathcal{Y}_1 v_2 = \int \mathcal{Y}_2 v_1 = 0.$$

Denote by G^\perp the set of $v \in H^1$ satisfying (3.8) and (3.9), and \tilde{G}^\perp the set of $v \in H^1$ satisfying (3.8) and (3.10).

Remark 3.4. Differently from [13], we use the orthogonality condition (3.9) instead of $\int \Delta Q v_1 = 0$. We make this choice in order to be able to prove coercivity in all dimensions, specially in dimension $N = 1$.

By direct calculations, one sees that

$$\Phi|_{\text{span}\{\nabla Q, iQ\}} = 0$$

and

$$(3.11) \quad \Phi(Q) = -\frac{p+1}{2} \int Q^{p+1} < 0.$$

If $s_c = 1$, consider the directions $W, iW, \Lambda W = \frac{N-2}{2}W + x \cdot \nabla W$ and $\partial_k W$, for $1 \leq k \leq N$, in the Hilbert space $\dot{H}^1 = \dot{H}^1(\mathbb{R}^N, \mathbb{C})$ with real inner product defined in (3.12). Denote by $G := \text{span}\{W, \nabla W, iW, \Lambda W\}$ and by G^\perp its orthogonal complement in \dot{H}^1 with

$$(3.12) \quad (f, g)_{\dot{H}^1} = \int \nabla f_1 \cdot \nabla g_1 + \int \nabla f_2 \cdot \nabla g_2 = \text{Re} \int \nabla f \cdot \nabla \bar{g}.$$

Let \tilde{G}^\perp be the set $\{v \in \dot{H}^1; v \perp \text{span}\{\nabla W, iW, \Lambda W\}, B(\mathcal{Y}_+, v) = B(\mathcal{Y}_-, v) = 0\}$.

By direct calculations, one sees that

$$\Phi|_{\text{span}\{\nabla W, iW, \Lambda W\}} = 0$$

and

$$(3.13) \quad \Phi(W) = -\frac{2}{(N-2)C_N^N} < 0,$$

where C_N is the sharp constant for Sobolev inequality for the embedding $\dot{H}^1(\mathbb{R}^N) \hookrightarrow L^{\frac{2N}{N-2}}(\mathbb{R}^N)$. The following lemma shows that Φ is coercive in $G^\perp \cup \tilde{G}^\perp$.

Lemma 3.5. *For $0 < s_c \leq 1$, there exists a constant $\tilde{c} > 0$ such that, for any $f \in G^\perp \cup \tilde{G}^\perp$,*

$$\Phi(f) \geq \tilde{c} \|f\|_{\dot{H}^1}^2.$$

This result was proved in a different context in [34], in the energy-critical case, and in [13] in the 3d cubic case. We give here the proof for all $0 < s_c \leq 1$, in any dimension.

Proof of Lemma 3.5, energy-critical case.

Step 1. Coercivity in G^\perp .

We adapt here the proof in [34] to our context. Let $\Pi: S^N \rightarrow \mathbb{R}^N$ be the “stretched” stereographic projection of the sphere S^N onto \mathbb{R}^N , with respect to the North pole, defined by

$$y_i = \frac{1}{N(N-2)} \frac{x_i}{1-x_{N+1}}, \quad 1 \leq i \leq N.$$

If $y = \Pi x$ and v is a real function is defined on \mathbb{R}^N , we define a function u on S^N by

$$u(x) = W^{-1}(y)v(y).$$

By integration by parts, one can check that

$$\int_{S^N} |\nabla_{S^N} u|^2 d\sigma = 2^{N-2} \int_{\mathbb{R}^N} (|\nabla v|^2 - W^{p_c} v^2) dy,$$

and

$$\int_{S^N} u^2 d\sigma = \frac{2^N}{N(N-2)} \int_{\mathbb{R}^N} W^{p_c} v^2 dy.$$

The spectrum of Δ_{S^N} is well known [5]. Namely, for the first eigenvalues λ_k , with multiplicity n_k and associated eigenfunctions $u_{k,\alpha}$, with $\alpha = (\alpha_1, \dots, \alpha_N) \in \mathbb{Z}_{\geq 0}^N$, we have

$$\begin{aligned} \lambda_0 &= 0, & n_0 &= 1, & u_0 &= 1, \\ \lambda_1 &= N, & n_1 &= N + 1, & u_{1,j} &= x_i, \quad 1 \leq j \leq N + 1, \\ \lambda_2 &= 2(N + 1), & n_2 &= 2N + 3. \end{aligned}$$

Therefore, if $v \perp W$ in \dot{H}^1 , then u is orthogonal to u_0 , and we have

$$\int_{\mathbb{R}^N} (|\nabla v|^2 - W^{p_c} v^2) dy \geq \frac{4\lambda_1}{N(N-2)} \int_{\mathbb{R}^N} W^{p_c} v^2 dy,$$

which is equivalent to

$$\int_{\mathbb{R}^N} (|\nabla v|^2 - W^{p_c} v^2) dy \geq \frac{4}{N+2} \int_{\mathbb{R}^N} |\nabla v|^2 dy.$$

Similarly, if $v \perp \text{span}\{W, \nabla W, \Lambda W\}$ in \dot{H}^1 , then u is orthogonal to $u_0, u_{1,i}, 1 \leq i \leq N + 1$, and thus

$$\int_{\mathbb{R}^N} (|\nabla v|^2 - W^{p_c} v^2) dy \geq \frac{4\lambda_2}{N(N-2)} \int_{\mathbb{R}^N} W^{p_c} v^2 dy,$$

which is equivalent to

$$\int_{\mathbb{R}^N} (|\nabla v|^2 - p_c W^{p_c} v^2) dy \geq \frac{4}{N+2} \int_{\mathbb{R}^N} |\nabla v|^2 dy.$$

Therefore, we proved that, for $h \in G^\perp$,

$$\Phi(h) \geq \frac{4}{N+2} \|h\|_{\dot{H}^1}.$$

Step 2. Coerciveness of Φ in \tilde{G}^\perp .

We first claim that $B(\mathcal{Y}_+, \mathcal{Y}_-) \neq 0$. If $B(\mathcal{Y}_+, \mathcal{Y}_-)$ were 0, then Φ would be identically 0 on $\text{span}\{\nabla W, iW, \Lambda W, \mathcal{Y}_+, \mathcal{Y}_-\}$, a subspace of dimension $N + 4$. But this cannot happen, given Φ is positive definite on G^\perp , which is of codimension $N + 3$.

We now show that $\Phi(h) > 0$ on $\tilde{G}^\perp \setminus \{0\}$. Assume, by contradiction, that there exists $h \in \tilde{G}^\perp \setminus \{0\}$ such that $\Phi(h) \leq 0$. Recall that $\ker \mathcal{L} = \text{span}\{\nabla W, iW, \Lambda W\}$, and that, by definition of $\tilde{G}^\perp \setminus \{0\}$, $B(\mathcal{Y}_+, h) = 0$. Hence, the vectors $\partial_k W$, $1 \leq k \leq N$, iW , ΛW , \mathcal{Y}_+ and h are mutually orthogonal under the symmetric form B . Since

$$\Phi(\partial_k W) = \Phi(iW) = \Phi(\Lambda W) = \Phi(\mathcal{Y}_+) = 0,$$

we get

$$\Phi|_{\text{span}\{\nabla W, iW, \Lambda W, \mathcal{Y}_+, h\}} \leq 0.$$

We claim that these vectors are independent. Indeed, if

$$\sum_k \alpha_k \partial_k W + \beta iW + \gamma \Lambda W + \delta \mathcal{Y}_+ + \epsilon h = 0,$$

then

$$\delta B(\mathcal{Y}_+, \mathcal{Y}_-) = 0,$$

and since $B(\mathcal{Y}_+, \mathcal{Y}_-) \neq 0$, $\delta = 0$. Therefore, the claim is proven, since $\partial_k W$, iW , ΛW and h are orthogonal in the real Hilbert space \dot{H}^1 .

To prove coercivity, we rely on a compactness argument. Suppose, by contradiction, that there exists $\{h_n\} \subset \tilde{G}^\perp$ such that

$$\lim \Phi(h_n) = 0, \quad \|h_n\|_{\dot{H}^1} = 1.$$

Up to a subsequence, we may assume $h_n \rightharpoonup h^*$ weakly in \dot{H}^1 . This implies $h^* \in \tilde{G}^\perp$. Since the operator $\int W^{p_c-1} |\cdot|^2$ is compact, we have $\int W^{p_c-1} |h_n|^2 > 0$ and

$$\Phi(h^*) \leq \liminf \Phi(h_n) = 0.$$

This contradicts the strict positivity of Φ on $\tilde{G}^\perp \setminus \{0\}$. ■

Proof of Lemma 3.5, intercritical case. Since the explicit formula for Q in the intercritical case is not available, we cannot proceed as in the energy-critical case. We follow here [43] and [13].

Step 1. Non-negativity on G^\perp . Define the functional

$$J(u) = \frac{\left(\int |\nabla u|^2\right)^a \left(\int |u|^2\right)^b}{\int |u|^{p+1}},$$

where

$$a = \frac{N(p-1)}{4}, \quad b = \frac{2p+2-N(p-1)}{4}.$$

By the sharp Gagliardo–Nirenberg inequality, one can see that this functional achieves an absolute minimum at Q . Therefore, the minimization condition $\frac{d^2}{d\varepsilon^2} J(Q + \varepsilon h)|_{\varepsilon=0} \geq 0$ for all functions $h \in H^1$ gives

$$\Phi(h) \geq \frac{b}{\left(\int Q^2\right)} \left[\frac{1}{a} \left(\int \Delta Q h_1\right)^2 + \frac{1}{b} \left(\int Q h_1\right)^2 - \left(\int Q^p h_1\right)^2 \right].$$

Since a and b are positive if $0 < s_c < 1$, we have that $\Phi(h) \geq 0$ if $\int Q^p h_1 = 0$. Therefore, Φ must be non-negative on G^\perp .

Step 2. Coercivity on G^\perp . We now employ compactness to show that, for every real function $h \in G^\perp$,

$$(L_\pm h, h)_{L^2} \gtrsim \|h\|_{L^2}.$$

Assuming the above inequality, multiplying the immediate bound

$$(L_\pm h, h)_{L^2} \geq \frac{1}{2} \|\nabla h\|_{L^2}^2 - C \|h\|_{L^2}^2,$$

by a small constant, and summing both inequalities, the coercivity follows. Suppose that there is a sequence of real H^1 functions $\{h_n\}$ in G^\perp such that

$$\lim_{n \rightarrow \infty} (L_+ h_n, h_n)_{L^2} = \Phi(h_n) = 0 \quad \text{and} \quad \|h_n\|_{L^2} = 1.$$

This implies

$$0 \leq \frac{1}{2} \int |\nabla h_n|^2 = -\frac{1}{2} + \frac{p}{2} \int Q^{p-1} h_n^2 + \Phi(h_n) \lesssim 1.$$

Therefore, $\|\nabla h_n\| \lesssim 1$ and, for large n , $\int Q^{p-1} h_n^2 \gtrsim 1$. Passing to a subsequence, and recalling that Q decays at infinity, we get that there exists $h_* \in G^\perp$ such that

$$h_n \rightharpoonup h_* \text{ weakly in } H^1, \quad \text{and} \quad \int Q^{p-1} h_*^2 > 0.$$

In particular, $h_* \neq 0$. Moreover,

$$\Phi(h_*) \leq \frac{1}{2} \liminf_{n \rightarrow \infty} \|h_n\|_{H^1}^2 - \frac{p}{2} \lim_{n \rightarrow \infty} \int Q^{p-1} h_n^2 = \liminf_{n \rightarrow \infty} \Phi(h_n) = 0.$$

Recall that $\Phi(h_*) \geq 0$ by Step 1. Therefore, $\Phi(h_*) = 0$ and h_* is the solution to the minimization problem

$$0 = (L_+ h_*, h_*)_{L^2} = \min_{f \in E} (L_+ f, f)_{L^2},$$

$$\text{where } E := \{h \in H^1 : \|h\|_{L^2} = \|h_*\|_{L^2} \text{ and } h \in G^\perp\}.$$

Thus, there exist Lagrange multipliers $\lambda_0, \dots, \lambda_{N+1}$ such that

$$L_+ h_* = \lambda_0 Q^p + \sum_{j=1}^N \lambda_j \partial_j Q + \lambda_{N+1} h_*.$$

Since $h_* \in G^\perp \setminus \{0\}$ and $(L_+ h_*, h_*)_{L^2} = 0$, we have $\lambda_{N+1} = 0$. By testing the last equation against $\partial_j Q$ and using that $L_+(\partial_k Q) = 0$, for all $k \leq N$, we conclude that

$$L_+ h_* = \lambda_0 Q^p.$$

Recalling that $L_+Q = -\frac{p-1}{2}Q^p$ and $\ker(L_+) = \text{span}\{\nabla Q\}$, we conclude that there exist μ_1, \dots, μ_N such that

$$h_* = -\frac{2\lambda_0}{p-1}Q + \sum_{j=1}^N \mu_j \partial_j Q.$$

Noting that $\int Q \partial_j Q = \frac{1}{2} \int \partial_j(Q^2) = 0$, and recalling that $h_* \in G^\perp$, gives $\mu_j = 0$ for all j . Therefore,

$$h_* = \frac{2\lambda_0}{p-1}Q.$$

And, by direct calculation,

$$(L_+h_*, h_*)_{L^2} = -\left(\frac{2\lambda_0}{p-1}\right)^2 \int Q^{p+1} < 0.$$

This contradicts $(L_+h_*, h_*)_{L^2} \geq 0$ and $h_* \neq 0$, and proves that

$$(L_+h, h)_{L^2} \gtrsim \|h\|_{L^2}^2$$

for any real function $h \in G^\perp$. The proof for L_- is analogous. In particular, we have strict positivity of Φ on $G^\perp \setminus \{0\}$ and, by compactness, the coercivity follows on G^\perp .

Step 3. Coercivity on \tilde{G}^\perp .

We have, as in the critical case, $B(\mathcal{Y}_+, \mathcal{Y}_-) \neq 0$, otherwise, the restriction of Φ to the $(N + 3)$ -dimensional space $\text{span}\{\nabla Q, iQ, \mathcal{Y}_+, \mathcal{Y}_+\}$ would be identically zero, contradicting the coercivity on G^\perp , which is of codimension $N + 2$.

Now, if there exists $h \in \tilde{G}^\perp \setminus \{0\}$ such that $\Phi(h) \leq 0$, we have $\Phi|_{\text{span}\{\nabla Q, iQ, \mathcal{Y}_+, h\}} \leq 0$. But if

$$\sum_k \alpha_k \partial_k + \beta iQ + \gamma \mathcal{Y}_+ + \delta h = 0,$$

then $\gamma B(\mathcal{Y}_+, \mathcal{Y}_-) = 0$, which implies $\gamma = 0$. Therefore, since $h \in \tilde{G}^\perp \setminus \{0\}$, the subspace spanned by $\{\nabla Q, iQ, \mathcal{Y}_+, h\}$ has dimension $N + 3$, contradicting the coercivity on G^\perp . It remains to show that the coercivity follows from the strict positivity.

Indeed, consider a sequence $\{h_n\} \subset \tilde{G}^\perp$ such that $\|h_n\|_{H^1} = 1$ for all n and such that

$$\Phi(h_n) \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Up to a subsequence, $h_n \rightharpoonup h^* \in \tilde{G}^\perp$ weakly on H^1 . But since Q decays at infinity, this implies $\Phi(h^*) \leq \liminf \Phi(h_n) = 0$, which contradicts the strict positivity of Φ on \tilde{G}^\perp . ■

Unlike the energy-critical case, the ground state decays exponentially if $s_c < 1$. In the next sections, we need sharp bounds on the decay of Q and its derivatives. We start recalling the following result, proved by Gidas, Ni and Nirenberg [17]. Recall that $2^* = 2N/(N - 2)$, if $N \geq 3$, and $2^* = +\infty$, if $N = 1, 2$.

Lemma 3.6 (Gidas et al., [17], Theorem 2, p. 370). *For $1 + 4/N \leq p < 2^* - 1$, let $Q \in \mathcal{S}(\mathbb{R}^N)$, Schwartz space, be the unique radial positive solution of the equation (3.4). Then there exists $C > 0$ such that*

$$\lim_{|x| \rightarrow +\infty} |x|^{(N-1)/2} e^{|x|} Q(x) = C.$$

We next study the decay of solutions to the equation (3.4). The next lemma, as well as its corollary, might be known in the theory of elliptic equations. However, we could not find a specific reference, and for convenience provide a proof here.

Lemma 3.7. *Let $f \in S(\mathbb{R}^N)$ and $\lambda \in \mathbb{R}$. If f solves*

$$(1 - \Delta + \lambda i)f = G, \quad \text{with } |G(x)| \lesssim \frac{e^{-a|x|}}{(1 + |x|^{(N-1)/2})^b},$$

for $0 < a \neq \operatorname{Re} \sqrt{1 + \lambda i}$, $0 < b \neq 1$, then

$$|f(x)| \lesssim \frac{1}{(1 + |x|^{(N-1)/2})^{\min\{b,1\}}} (e^{-|x|})^{\min\{a, \operatorname{Re} \sqrt{1+\lambda i}\}}.$$

Proof. Let $c = \operatorname{Re} \sqrt{1 + \lambda i} \geq 1$. We recall the integral form of the resolvent (see [1])

$$(1 - \Delta + \lambda i)^{-1}G = K * G,$$

where $K \in L^1(\mathbb{R}^N)$ is such that, for $|x| \gg 1$,

$$(3.14) \quad K(x) \lesssim \frac{e^{-c|x|}}{1 + |x|^{(N-1)/2}},$$

and, for $|x| \ll 1$,

$$(3.15) \quad K(x) \lesssim \begin{cases} \frac{1}{|x|^{(N-1)/2}} & \text{for } N > 2, \\ \ln \frac{1}{|x|} & \text{for } N = 2, \\ 1 & \text{for } N < 2. \end{cases}$$

Consider first the case $0 < a < c$. We estimate

$$\begin{aligned} |K * G(x)| &\lesssim \int K(y) \frac{e^{-a|x-y|}}{(1 + |x - y|^{(N-1)/2})^b} dy \\ &\lesssim \frac{e^{-a|x|}}{(1 + |x|^{(N-1)/2})^{\min\{b,1\}}} \int K(y) e^{a|y|} \frac{(1 + |x - y|^{(N-1)/2} + |y|^{(N-1)/2})^{\min\{b,1\}}}{(1 + |x - y|^{(N-1)/2})^b} dy \\ &\lesssim \frac{e^{-a|x|}}{(1 + |x|^{(N-1)/2})^{\min\{b,1\}}} \int K(y) e^{a|y|} (1 + |y|^{(N-1)/2})^{\min\{b,1\}} dy. \end{aligned}$$

By (3.14) and (3.15), the integral in the last inequality is $O(1)$. For $a > c$, the estimate is

$$\begin{aligned} |K * G(x)| &\lesssim \frac{e^{-c|x|}}{(1 + |x|^{(N-1)/2})^{\min\{1,b\}}} \int K(y) e^{c|y|} (1 + |y|^{(N-1)/2})^{\min\{1,b\}} e^{-(a-c)|x-y|} dy \\ &\lesssim \frac{e^{-c|x|}}{(1 + |x|^{(N-1)/2})^{\min\{1,b\}}} \left[\int e^{-(a-c)|x-y|} dy + \int_{|y| \leq 1} K(y) e^{c|y|} (1 + |y|^{(N-1)/2}) dy \right]. \end{aligned}$$

Since the first integral in the last inequality is bounded uniformly in x , the lemma is proved. ■

Corollary 3.8. *For any multi-index $\alpha \in \mathbb{Z}_+^N$, the following estimates hold:*

- (i) $\|Q^{-1}\partial^\alpha Q\|_{L^\infty} < +\infty$,
- (ii) $\|Q^{-1}e^{\eta|x|}\partial^\alpha \mathcal{Y}_\pm\|_{L^\infty} < +\infty$, for some $0 < \eta \ll 1$,
- (iii) $\|Q^{-1}e^{\eta|x|}\partial^\alpha [(\mathcal{L} - \lambda)^{-1}f]\|_{L^\infty} < +\infty$, for every $\lambda \in \mathbb{R} \setminus \sigma(\mathcal{L})$ and every $f \in S(\mathbb{R}^N)$ such that $\|Q^{-1}e^{\eta|x|}\partial^\beta f\|_{L^\infty} < +\infty$ for some $0 < \eta < \text{Re}(\sqrt{1 + \lambda i})$ and any $\beta \in \mathbb{Z}_+^N$.

Proof. We first remark that Q is strictly positive, and thus, Q^{-1} is well-defined. Recalling Lemma 3.6, we have, for all x ,

$$Q(x) \approx \frac{e^{-|x|}}{1 + |x|^{(N-1)/2}}.$$

We differentiate (1.3) to obtain

$$(1 - \Delta)\nabla Q = pQ^{p-1}\nabla Q.$$

Since $Q \in S$, by Lemma 3.7 and a bootstrap argument, we conclude (i) for $|\alpha| = 1$. Differentiating (1.3) and repeating the argument, we conclude (i), by induction, for any multi-index α .

To prove (ii), recall the differential equation for $\mathcal{Y}_1 = \text{Re}(\mathcal{Y}_+)$:

$$(1 - \Delta - pQ^{p-1})(1 - \Delta - Q^{p-1})\mathcal{Y}_1 = -e_0^2 \mathcal{Y}_1.$$

By factoring $[(1 - \Delta)^2 + e_0^2] = (1 - \Delta + ie_0)(1 - \Delta - ie_0)$ and using item (i), this equation can be rewritten as

$$(1 - \Delta + ie_0)(1 - \Delta - ie_0)\mathcal{Y}_1 = G_2(\mathcal{Y}_1),$$

where we define $G_k(f)$ as a linear function on f and its derivatives up to order k , that satisfies, for any $k \geq 1$,

$$|G_k(f)| \lesssim Q^{p-1} \sum_{|\alpha| \leq k} |\partial^\alpha f|.$$

Writing $g = (1 - \Delta - ie_0)\mathcal{Y}_1$, we have, for any multi-indices α, β ,

$$\begin{cases} (1 - \Delta + ie_0)\partial^\alpha g = G_{|\alpha|+2}(\mathcal{Y}_1) \\ (1 - \Delta - ie_0)\partial^\beta \mathcal{Y}_1 = \partial^\beta g. \end{cases}$$

Therefore, using Lemma 3.7 and bootstrapping, we prove that $Q^{-1}e^{\eta|x|}\partial^\alpha \mathcal{Y}_1 \in L^\infty$ for any multi-index α , where $0 < \eta \ll \text{Re}(\sqrt{1 + ie_0}) - 1$. The estimate on \mathcal{Y}_2 is analogous, and hence (ii) holds. We now turn to estimate (iii). If $g = (\mathcal{L} - \lambda)^{-1}f$, then, for any α ,

$$\begin{cases} -\partial^\alpha(1 - \Delta - Q^{p-1})g_2 - \lambda \partial^\alpha g_1 = \partial^\alpha f_1, \\ \partial^\alpha(1 - \Delta - pQ^{p-1})g_1 - \lambda \partial^\alpha g_2 = \partial^\alpha f_2. \end{cases}$$

We can rewrite this system as

$$[(1 - \Delta)^2 + \lambda^2]\partial^\alpha g_1 = G_{|\alpha|+2}(g_1) + H_{|\alpha|+2}(f),$$

where we define $H_k(f)$ as a linear function on f and its derivatives up to order k , that satisfies, for any $k \geq 1$,

$$|H_k(f)| \lesssim \sum_{|\alpha| \leq k} |\partial^\alpha f|.$$

Bootstrapping similarly to the previous items, and noting that the argument for g_2 is analogous, completes the proof of Lemma 3.7. ■

3.2. Estimates on the linearized equation

We now prove some estimates that will be used in the next sections. We start with estimates for the energy-critical case.

Lemma 3.9 (Preliminary estimates). *Let $s_c = 1$, $N \geq 6$ (and then $p_c - 1 = \frac{4}{N-2} \leq 1$), $0 < \epsilon \ll \frac{4}{N-2}$, and let I be a bounded time interval with $|I| \leq 1$. Consider $f, g \in S(\dot{H}^1, I)$ such that $\nabla f, \nabla g \in S(L^2, I)$. The following estimates hold.*

- (i) $\|\nabla K(f)\|_{S'(L^2, I)} \lesssim |I|^{1/2} \|\nabla f\|_{S(L^2, I)}$,
- (ii) $\|\nabla(R(f) - R(g))\|_{S'(L^2, I)}$
 $\lesssim \|\nabla(f - g)\|_{S(L^2, I)} (\|\nabla f\|_{S(L^2, I)}^{p_c-1} + \|\nabla g\|_{S(L^2, I)}^{p_c-1})$
 $+ \|D^\epsilon(f - g)\|_{S(\dot{H}^{1-\epsilon}, I)}^{p_c-1} (\|\nabla f\|_{S(L^2, I)} + \|\nabla g\|_{S(L^2, I)}).$

If $N > 6$, then also

- (iii) $\|D^\epsilon K(f)\|_{S'(\dot{H}^{-(1-\epsilon)}, I)} \lesssim |I|^{\epsilon/(N-2)} \|D^\epsilon f\|_{S(\dot{H}^{1-\epsilon}, I)}$,
- (iv) $\|D^\epsilon(R(f) - R(g))\|_{S'(\dot{H}^{-(1-\epsilon)}, I)}$
 $\lesssim \|D^\epsilon(f - g)\|_{S(\dot{H}^{1-\epsilon}, I)} (\|\nabla f\|_{S(L^2, I)}^{p_c-1} + \|\nabla g\|_{S(L^2, I)}^{p_c-1}).$

Remark 3.10. It is necessary to treat the case $N > 6$ differently due to the low power of the nonlinearity. If $N \leq 6$, then it is possible to estimate $\|\nabla(R(f) - R(g))\|_{S'(L^2, I)}$ at least linearly in terms of $\|\nabla(f - g)\|_{S(L^2, I)}$. In higher dimensions, one of the terms must be in the form $\|\nabla(f - g)\|_{S(L^2, I)}^{p_c-1}$, which is not good enough for the fixed-point argument carried on in the next section. The use of less than one derivative enables us to keep the desired linearity.

Proof of Lemma 3.9. We start by proving the following claim.

Claim 3.11. *Let H be a map such that $H(0) = 0$ and $|H(f) - H(g)| \leq C|f - g|^{\frac{4}{N-2}}$ for all functions $f, g: \mathbb{R}^N \rightarrow \mathbb{C}$, $N > 6$. Then, for all $f, g \in S(\dot{H}^1, I) \cap \nabla^{-1}S(L^2, I)$,*

$$\begin{aligned} & \|D^\epsilon(H(f)g)\|_{L_t^{2/\epsilon} L_x^{\frac{2N}{N+2}}} \\ & \lesssim \|\nabla f\|_{\frac{4}{\epsilon(N-2)}, \frac{8N}{4N-\epsilon(N-2)}}^{\frac{4}{N-2}} \|D^\epsilon g\|_{\frac{4}{\epsilon}, \frac{2N}{N-2+\epsilon}} + \|\nabla f\|_{\frac{4}{\epsilon}, \frac{2N}{N-\epsilon}}^{\frac{4}{N-2}} \|D^\epsilon g\|_{\frac{2(N-2)}{\epsilon(N-4)}, \frac{2N(N-2)}{(N-2)^2+4\epsilon}}. \end{aligned}$$

In other words,

$$\|D^\epsilon(H(f)g)\|_{S'(\dot{H}^{-(1-\epsilon)}, I)} \lesssim \|\nabla f\|_{S(L^2, I)}^{4/(N-2)} \|D^\epsilon g\|_{S(\dot{H}^{1-\epsilon}, I)}.$$

Proof of Claim 3.11. By the Leibniz rule (2.1) and Holder’s inequality, we can write

$$\begin{aligned} & \|D^\varepsilon(H(f)g)\|_{L_I^{\frac{2}{\varepsilon}} L_x^{\frac{2N}{N+2}}} \\ & \lesssim \left\| \|D^\varepsilon H(f)\|_{L_x^{\frac{2N}{4+\varepsilon}}} \|g\|_{L_x^{\frac{2N}{N-2-\varepsilon}}} \right\|_{L_I^{2/\varepsilon}} + \left\| \|H(f)\|_{L_x^{\frac{N(N-2)}{2(N-2-\varepsilon)}}} \|D^\varepsilon g\|_{L_x^{\frac{2N(N-2)}{(N-2)^2+4\varepsilon}}} \right\|_{L_I^{2/\varepsilon}} \\ & \lesssim \|D^\varepsilon H(f)\|_{L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{4+\varepsilon}}} \|g\|_{L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{N-2-\varepsilon}}} + \|H(f)\|_{L_I^{\frac{N-2}{\varepsilon}} L_x^{\frac{N(N-2)}{2(N-2-\varepsilon)}}} \|D^\varepsilon g\|_{L_I^{\frac{2(N-2)}{\varepsilon(N-4)} L_x^{\frac{2N(N-2)}{(N-2)^2+4\varepsilon}}}. \end{aligned}$$

By the Sobolev inequality,

$$\|g\|_{L_I^{4/\varepsilon} L_x^{\frac{2N}{N-2-\varepsilon}}} \lesssim \|D^\varepsilon g\|_{L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{N-2+\varepsilon}}},$$

and since, by the assumption on H , $|H(f)| \lesssim |f|^{4/(N-2)}$, we have, by Sobolev,

$$\|H(f)\|_{L_I^{\frac{N-2}{\varepsilon}} L_x^{\frac{N(N-2)}{2(N-2-\varepsilon)}}} \lesssim \|f\|_{L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{N-2-\varepsilon}}}^{\frac{4}{\varepsilon}} \lesssim \|\nabla f\|_{L_I^{\frac{4}{\varepsilon}} L_x^{\frac{2N}{N-2-\varepsilon}}}^{\frac{4}{\varepsilon}}.$$

It remains to estimate $\|D^\varepsilon H(f)\|_{L_x^{2N/(4+\varepsilon)}}$. Choosing ν such that $(N - 2)\varepsilon/4 < \nu < 1$, by fractional chain rule³ (2.2) and the Sobolev embeddings, we have

$$\|D^\varepsilon H(f)\|_{L_x^{\frac{2N}{4+\varepsilon}}} \lesssim \|f\|_{L_x^{\frac{4}{(N-2-\frac{\varepsilon}{\nu})p_1}}}^{\frac{4}{N-2-\frac{\varepsilon}{\nu}}} \|D^\nu f\|_{L_x^{\frac{\varepsilon/\nu}{q_1}}}^{\varepsilon/\nu} \lesssim \|\nabla f\|_{L_x^{\frac{4}{N-2-\frac{\varepsilon}{\nu}}}}^{\frac{4}{N-2-\frac{\varepsilon}{\nu}}} \|\nabla f\|_{L_x^{q_2}}^{\varepsilon/\nu},$$

where we choose $p_2 = q_2 = \frac{8N}{4N-\varepsilon(N-2)} \in (1, +\infty)$, and p_1 and q_1 must satisfy

$$1 < p_1, q_1 < \infty, \quad \frac{1}{p_2} = \frac{1}{\left(\frac{4}{N-2} - \frac{\varepsilon}{\nu}\right) p_1} + \frac{1}{N}, \quad \frac{1}{q_2} = \frac{1}{\frac{\varepsilon}{\nu} q_1} + \frac{1-\nu}{N},$$

and

$$\left(1 - \frac{\varepsilon(N-2)}{4\nu}\right) p_1 > 1.$$

These conditions can be easily satisfied if ε is small enough (depending only on the dimension). The claim is now proved. ■

Estimate (iii) of Lemma 3.9 follows directly from Sobolev’s inequality and Claim 3.11, by taking $H(W) = |W|^{4/(N-2)}$, and from the fact that $|\nabla W| \in L_x^2 \cap L_x^\infty$, if $N > 6$.

To prove (iv), note that $R(f) = W^{p_c} J(W^{-1} f)$, where

$$J(z) = |1 + z|^{p_c-1} (1 + z) - 1 - \frac{p_c + 1}{2} z - \frac{p_c - 1}{2} \bar{z}$$

is $C^1(\mathbb{C})$. Its derivatives J_z and $J_{\bar{z}}$ satisfy $J_z(0) = J_{\bar{z}}(0) = 0$ and, if $N > 6$, are Hölder continuous of order $p_c - 1 < 1$. Therefore, writing

$$\begin{aligned} (3.16) \quad & R(f) - R(g) \\ & = W^{p_c-1} \int_0^1 J_z(W^{-1}(sf + (1-s)g))(f-g) + J_{\bar{z}}(W^{-1}(sf + (1-s)g))(\overline{f-g}) \, ds, \end{aligned}$$

we can apply Claim 3.11 to estimate each term in (3.16), taking $H(f) = W^{p_c-1} J_z(W^{-1} f)$ or $H(f) = W^{p_c-1} J_{\bar{z}}(W^{-1} f)$. Estimate (iv) then follows directly.

³This is where the hypothesis $N > 6$ is used, as the fractional chain rule requires $0 < 4/(N - 2) < 1$.

To prove (i), we write

$$|\nabla K(f)| \lesssim |W|^{p_c-2} |\nabla W| |f| + |W|^{p_c-1} |\nabla f|.$$

Using the fact that $|\partial^\alpha W(x)| \leq C_\alpha |W(x)|$ for every multi-index $\alpha \in \mathbb{Z}_+^N$ and all x , we have, by Hölder’s inequality,

$$\begin{aligned} \|\nabla K(f)\|_{L_t^2 L_x^{\frac{2N}{N+2}}} &\lesssim \left\| \|W\|_{L_x^{\frac{4(N-2)}{2N/(N-2)}}} \|f\|_{L_x^{2N/(N-2)}} \right\|_{L_t^2} + \left\| \|W\|_{L_x^{\frac{4(N-2)}{4N/(N-2)}}} \|\nabla f\|_{L_x^2} \right\|_{L_t^2} \\ &\lesssim |I|^{1/2} (\|f\|_{L_t^\infty L_x^{2N/(N-2)}} + \|\nabla f\|_{L_t^\infty L_x^2}). \end{aligned}$$

Note that we used that $W \in L_x^{2N/(N-2)} \cap L_x^{4N/(N-2)}$, which follows from the fact that $W \in L_x^2 \cap L_x^\infty$, if $N > 6$. The inequality follows from the Sobolev embedding.

We finally turn to estimate (ii). Write

$$\begin{aligned} \nabla(R(f) - R(g)) &= \underbrace{p_c W^{p_c-1} \nabla W (J(W^{-1} f) - J(W^{-1} g))}_{(a)} \\ &\quad + \underbrace{W^{p_c-1} J_z(W^{-1} f) \nabla f - W^{p_c-1} J_z(W^{-1} g) \nabla g}_{(b)} \\ &\quad + \underbrace{W^{p_c-1} J_{\bar{z}}(W^{-1} f) \nabla \bar{f} - W^{p_c-1} J_{\bar{z}}(W^{-1} g) \nabla \bar{g}}_{(c)} \\ &\quad + \underbrace{W^{p_c-2} \nabla W J_z(W^{-1} f) f - W^{p_c-2} \nabla W J_z(W^{-1} g) g}_{(d)} \\ &\quad + \underbrace{W^{p_c-2} \nabla W J_{\bar{z}}(W^{-1} f) \bar{f} - W^{p_c-2} \nabla W J_{\bar{z}}(W^{-1} g) \bar{g}}_{(e)}. \end{aligned}$$

To estimate (a), note that

$$\begin{aligned} |(a)| &\lesssim W^{p_c-1} \int_0^1 |J_z(W^{-1}(sf + (1-s)g))(f-g) + J_{\bar{z}}(W^{-1}(sf + (1-s)g))(\overline{f-g})| ds \\ &\lesssim (|f|^{p_c-1} + |g|^{p_c-1}) |f - g|. \end{aligned}$$

Thus, by the Hölder and Sobolev inequalities,

$$\begin{aligned} \|(a)\|_{L_t^2 L_x^{\frac{2N}{N+2}}} &\lesssim |I|^{1/2} (\|f\|_{L_t^\infty L_x^{2N/(N-2)}}^{p_c-1} + \|g\|_{L_t^\infty L_x^{2N/(N-2)}}^{p_c-1}) \|f - g\|_{L_t^\infty L_x^{2N/(N-2)}} \\ &\lesssim |I|^{1/2} (\|\nabla f\|_{L_t^\infty L_x^2}^{p_c-1} + \|\nabla g\|_{L_t^\infty L_x^2}^{p_c-1}) \|\nabla(f - g)\|_{L_t^\infty L_x^2}. \end{aligned}$$

We now estimate (b). By the triangle inequality,

$$\begin{aligned} |(b)| &\leq W^{p_c-1} |J_z(W^{-1} f)| |\nabla f - \nabla g| + W^{p_c-1} |J_z(W^{-1} f) - J_z(W^{-1} g)| |\nabla g| \\ &\leq |f|^{p_c-1} |\nabla(f - g)| + |f - g|^{p_c-1} |\nabla g|. \end{aligned}$$

So that, by the Hölder and Sobolev inequalities,

$$\begin{aligned} \|(b)\|_{L^2_I L^{\frac{2N}{N+2}}_x} &\lesssim \|f\|_{L^\infty_I L^{\frac{2N}{N-2}}_x}^{p_c-1} \|\nabla(f-g)\|_{L^2_I L^{\frac{2N}{N-2}}_x} + \|f-g\|_{L^\infty_I L^{\frac{2N}{N-2}}_x}^{p_c-1} \|\nabla g\|_{L^2_I L^{\frac{2N}{N-2}}_x} \\ &\lesssim \|\nabla f\|_{L^\infty_I L^2_x}^{p_c-1} \|\nabla(f-g)\|_{L^2_I L^{\frac{2N}{N-2}}_x} + \|D^\varepsilon(f-g)\|_{L^\infty_I L^{\frac{2N}{N-2+2\varepsilon}}_x}^{p_c-1} \|\nabla g\|_{L^2_I L^{\frac{2N}{N-2}}_x}. \end{aligned}$$

The estimate for (c) is analogous. To estimate (d), we write

$$\begin{aligned} |(d)| &\leq W^{p_c-1} |J_z(W^{-1}f)| |f-g| + W^{p_c-1} |J_z(W^{-1}f) - J_z(W^{-1}g)| |g| \\ &\leq |f|^{p_c-1} |f-g| + |f-g|^{p_c-1} |g|. \end{aligned}$$

Therefore, by the Hölder and Sobolev inequalities,

$$\begin{aligned} \|(d)\|_{L^2_I L^{\frac{2N}{N+2}}_x} &\lesssim |I|^{\frac{1}{2}} \|f\|_{L^\infty_I L^{\frac{2N}{N-2}}_x}^{p_c-1} \|f-g\|_{L^2_I L^{\frac{2N}{N-2}}_x} + |I|^{\frac{1}{2}} \|f-g\|_{L^\infty_I L^{\frac{2N}{N-2}}_x}^{p_c-1} \|g\|_{L^2_I L^{\frac{2N}{N-2}}_x} \\ &\lesssim |I|^{\frac{1}{2}} \|\nabla f\|_{L^\infty_I L^2_x}^{p_c-1} \|\nabla(f-g)\|_{L^2_I L^2_x} + |I|^{\frac{1}{2}} \|D^\varepsilon(f-g)\|_{L^\infty_I L^{\frac{2N}{N-2+2\varepsilon}}_x}^{p_c-1} \|\nabla g\|_{L^2_I L^2_x}. \end{aligned}$$

Since the estimate for (e) is analogous, the proof of Lemma 3.9 is complete. ■

The following Strichartz-type continuity argument follows from Lemma 3.9 and will be useful in proving the main theorems of this paper.

Lemma 3.12. *Let h be a solution to (3.2). If, for some $c > 0$, and all $t > 0$,*

$$(3.17) \quad \|h(t)\|_{\dot{H}^1} \lesssim e^{-ct},$$

then, for all $t > 0$,

$$(3.18) \quad \|\nabla h\|_{S(L^2, [t, +\infty))} \lesssim e^{-ct}.$$

Proof. Differentiating (3.2), we get

$$i \partial_t (\nabla h) + \Delta(\nabla h) + \nabla(K(h) + R(h)) = 0.$$

By Duhamel’s formula, the Strichartz estimates and items (i) and (ii) of Lemma 3.9, if $0 < \tau < 1$,

$$\|\nabla h\|_{S(L^2, [t, t+\tau])} \lesssim \|h(t)\|_{\dot{H}^1} + \tau^{1/2} \|\nabla h\|_{S(L^2, [t, t+\tau])} + \|\nabla h\|_{S(L^2, [t, t+\tau])}^{p_c}.$$

By (3.17), we get, for some $K > 0$,

$$(3.19) \quad \|\nabla h\|_{S(L^2, [t, t+\tau])} \leq K (e^{-ct} + \tau^{1/2} \|\nabla h\|_{S(L^2, [t, t+\tau])} + \|\nabla h\|_{S(L^2, [t, t+\tau])}^{p_c}).$$

This implies, for large t ,

$$\|\nabla h\|_{S(L^2, [t, t+\tau_0])} < 2Ke^{-ct}, \quad \text{with } \tau_0 = \frac{1}{9K^2}.$$

Indeed, assume by contradiction that there exists $\tau \in (0, \tau_0]$ such that $\|h\|_{S(L^2, [t, t+\tau])} = 2Ke^{-ct}$, for fixed $t > 0$. Then, by (3.19),

$$2Ke^{-ct} \leq Ke^{-ct} + 2K^2\tau^{1/2}e^{-ct} + (2K)^{p_c}Ke^{-c p_c t} \leq \frac{5}{3}Ke^{-ct} + (2K)^{p_c}Ke^{-c p_c t},$$

which is a contradiction if t is large. Therefore, by decomposing $[t, +\infty) = \bigcup_{j=0}^\infty [t + j\tau_0, t + (j + 1)\tau_0]$ and using the triangle inequality, we see that (3.18) holds. ■

The following lemma is the intercritical version of Lemma 3.9, and its proof is analogous.

Lemma 3.13 (Preliminary estimates, subcritical case). *Let $0 < s_c < 1$, and let I be a bounded time interval such that $|I| \leq 1$. Consider $f, g \in S(L^2, I)$ such that $\nabla f, \nabla g \in S(L^2, I)$. There exists $\alpha > 0$ such that the following estimates hold.*

For $p > 1$:

- (i) $\|\langle \nabla \rangle K(f)\|_{S'(L^2, I)} \lesssim |I|^\alpha \|\langle \nabla \rangle f\|_{S(L^2, I)},$
- (ii) $\|K(f)\|_{S'(\dot{H}^{-s_c}, I)} \lesssim |I|^\alpha \|f\|_{S(\dot{H}^{s_c}, I)}.$

For $p > 2$:

- (iii) $\|\langle \nabla \rangle (R(f) - R(g))\|_{S'(L^2, I)}$
 $\lesssim \|\langle \nabla \rangle (f - g)\|_{S(L^2, I)} [\|\langle \nabla \rangle f\|_{S(L^2, I)} + \|\langle \nabla \rangle g\|_{S(L^2, I)}]$
 $+ \|\langle \nabla \rangle f\|_{S(L^2, I)}^{p-1} + \|\langle \nabla \rangle g\|_{S(L^2, I)}^{p-1},$
- (iv) $\|R(f) - R(g)\|_{S'(\dot{H}^{-s_c}, I)}$
 $\lesssim \|f - g\|_{S(\dot{H}^{s_c}, I)} [\|f\|_{S(\dot{H}^{s_c}, I)} + \|g\|_{S(\dot{H}^{s_c}, I)} + \|f\|_{S(\dot{H}^{s_c}, I)}^{p-1} + \|g\|_{S(\dot{H}^{s_c}, I)}^{p-1}],$

For $1 < p \leq 2$:

- (v) $\|\langle \nabla \rangle (R(f) - R(g))\|_{S'(L^2, I)}$
 $\lesssim \|\langle \nabla \rangle (f - g)\|_{S(L^2, I)} (\|f\|_{S(\dot{H}^{s_c}, I)}^{p-1} + \|g\|_{S(\dot{H}^{s_c}, I)}^{p-1})$
 $+ \|f - g\|_{S(\dot{H}^{s_c}, I)}^{p-1} (\|\langle \nabla \rangle f\|_{S(L^2, I)} + \|\langle \nabla \rangle g\|_{S(L^2, I)}),$
- (vi) $\|R(f) - R(g)\|_{S'(\dot{H}^{-s_c}, I)} \lesssim \|f - g\|_{S(\dot{H}^{s_c}, I)} (\|f\|_{S(\dot{H}^{s_c}, I)}^{p-1} + \|g\|_{S(\dot{H}^{s_c}, I)}^{p-1}).$

Proof. The estimates are very similar as the ones in the proof of the energy-critical case. We use the following classical inequalities:

$$\| |a|^{p-1} b \|_{S'(L^2)} \leq \|a\|_{S(\dot{H}^{s_c})}^{p-1} \|b\|_{S(L^2)} \lesssim \|\langle \nabla \rangle a\|_{S(L^2)}^{p-1} \|b\|_{S(L^2)}$$

and

$$\| |a|^{p-1} b \|_{S'(\dot{H}^{-s_c})} \leq \|a\|_{S(\dot{H}^{s_c})}^{p-1} \|b\|_{S(\dot{H}^{s_c})} \lesssim \|\langle \nabla \rangle a\|_{S(L^2)}^{p-1} \|\langle \nabla \rangle b\|_{S(L^2)},$$

which can be verified using the pairs $(\frac{4(p+1)}{N(p-1)}, p+1) \in \mathcal{A}_0, (\frac{2(p-1)(p+1)}{4-(N-2)(p-1)}, p+1) \in \mathcal{A}_{s_c},$ and $(\frac{2(p-1)(p+1)}{(p-1)(Np-2)-4}, p+1) \in \mathcal{A}_{-s_c},$ together with the Sobolev inequality. Let us estimate,

for example, $\|\nabla(R(f) - R(g))\|_{S'(L^2, I)}$. Write

$$\begin{aligned} \nabla(R(f) - R(g)) &= \underbrace{pQ^{p-1}\nabla Q(J(Q^{-1}f) - J(Q^{-1}g))}_{(a)} \\ &\quad + \underbrace{Q^{p-1}J_z(Q^{-1}f)\nabla f - Q^{p-1}J_z(Q^{-1}g)\nabla g}_{(b)} \\ &\quad + \underbrace{Q^{p-1}J_{\bar{z}}(Q^{-1}f)\nabla \bar{f} - Q^{p-1}J_{\bar{z}}(Q^{-1}g)\nabla \bar{g}}_{(c)} \\ &\quad + \underbrace{Q^{p-2}\nabla QJ_z(Q^{-1}f)f - Q^{p-2}\nabla QJ_z(Q^{-1}g)g}_{(d)} \\ &\quad + \underbrace{Q^{p-2}\nabla QJ_{\bar{z}}(Q^{-1}f)\bar{f} - Q^{p-2}\nabla QJ_{\bar{z}}(Q^{-1}g)\bar{g}}_{(e)}. \end{aligned}$$

Making use of $|\nabla Q| \lesssim Q$ (which follows from Corollary 3.8), we bound (a) as

$$|(a)| \lesssim Q^{p-1} \int_0^1 |J_z(Q^{-1}(sf + (1-s)g))(f-g) + J_{\bar{z}}(Q^{-1}(sf + (1-s)g))(\overline{f-g})| ds.$$

Now, since

$$|J_z(z_1) - J_z(z_2)| + |J_{\bar{z}}(z_1) - J_{\bar{z}}(z_2)| \lesssim \begin{cases} |z_1 - z_2|(1 + |z_1|^{p-2} + |z_2|^{p-2}), & p \geq 2, \\ |z_1 - z_2|^{p-1}, & 1 < p < 2, \end{cases}$$

we have

$$|(a)| \lesssim \begin{cases} (Q^{p-2}|f| + Q^{p-2}|g| + |f|^{p-1} + |g|^{p-1})|f - g|, & p \geq 2, \\ (|f|^{p-1} + |g|^{p-1})|f - g|, & 1 < p < 2. \end{cases}$$

Thus, since $Q \in \mathcal{S}(\mathbb{R}^N)$ and $|I| \leq 1$,

$$\begin{aligned} &\|(a)\|_{S'(L^2, I)} \\ &\lesssim \begin{cases} (\|f\|_{S(\dot{H}^{s_c, I})} + \|g\|_{S(\dot{H}^{s_c, I})} + \|f\|_{S(\dot{H}^{s_c, I})}^{p-1} + \|g\|_{S(\dot{H}^{s_c, I})}^{p-1})\|f - g\|_{S'(L^2, I)}, & p \geq 2, \\ (\|f\|_{S(\dot{H}^{s_c, I})}^{p-1} + \|g\|_{S(\dot{H}^{s_c, I})}^{p-1})\|f - g\|_{S'(L^2, I)}, & 1 < p < 2. \end{cases} \end{aligned}$$

We also have

$$\begin{aligned} &\|(b)\|_{S'(L^2, I)} + \|(d)\|_{S'(L^2, I)} \\ &\lesssim \begin{cases} \|\nabla(f - g)\|_{S(L^2, I)}(\|\langle \nabla \rangle f\|_{S(L^2, I)} + \|\langle \nabla \rangle f\|_{S(L^2, I)}^{p-1} \\ \quad + \|\langle \nabla \rangle g\|_{S(L^2, I)} + \|\langle \nabla \rangle g\|_{S(L^2, I)}^{p-1}), & p \geq 2, \\ \|\langle \nabla \rangle(f - g)\|_{S(L^2, I)}(\|f\|_{S(\dot{H}^{s_c, I})}^{p-1} + \|g\|_{S(\dot{H}^{s_c, I})}^{p-1}) \\ \quad + \|f - g\|_{S(\dot{H}^{s_c, I})}^{p-1}(\|\langle \nabla \rangle f\|_{S(L^2, I)} + \|\langle \nabla \rangle g\|_{S(L^2, I)}), & 1 < p < 2, \end{cases} \end{aligned}$$

with the same bounds for (c) and (e). ■

Remark 3.14. We do not employ the same estimates as in [13], since the nonlinearity $|u|^{p-1}u$ is not a polynomial in (u, \bar{u}) if p is not an odd integer. Therefore, instead of using H^s estimates, we rely on $S(L^2)$ and $S(\dot{H}^{s_c})$ estimates, that are more suitable for generalizing the result to all dimensions and powers of the nonlinearity.

Remark 3.15. We employ a different approach than Li and Zhang [30], that divide all estimates in regions where $|f| > W$ or $|f| \leq W$. Instead, we use fractional derivatives to avoid some sublinear estimates.

4. Construction of special solutions

In this section, we construct *special* solutions to the NLS equation (1.2), in the sense that they are on the same energy level of the ground state, converge to the standing wave in \dot{H}^1 as $t \rightarrow +\infty$, but have kinetic energy different from $\|\nabla Q\|_{L^2}$.

4.1. Construction of a family of approximate solutions

We start with a result that was first proved by Duyckaerts and Merle in [12], for $s_c = 1$, and later in [13] for the 3d cubic case. We extend here those proofs to the intercritical case. The main difference from the energy-critical case is that Q decays exponentially if $0 < s_c < 1$, so we need to be careful with its spatial decay, as we make use of the estimates of the type $\|Q^{-1}f\|_{L^\infty}$. To this end, we make use of the sharp decay estimate for Q given by (3.6) and of the control on the spatial decay given by Corollary 3.8.

Proposition 4.1. *Let $0 < s_c \leq 1$ and $A \in \mathbb{R}$. There exists a sequence $(Z_k^A)_{k \geq 1}$ of functions in $\mathcal{S}(\mathbb{R}^N)$ such that $Z_1^A = AY_+$ and, if $k \geq 1$ and $\mathcal{V}_k^A = \sum_{j=1}^k e^{-je_0t} Z_j^A$, then as $t \rightarrow +\infty$ we have*

$$(4.1) \quad \partial_t \mathcal{V}_k^A + \mathcal{L} \mathcal{V}_k^A = iR(\mathcal{V}_k^A) + O(e^{-(k+1)e_0t}) \quad \text{in } \mathcal{S}(\mathbb{R}^N),$$

where \mathcal{L} and R are given in Definition 3.1.

Proof. We prove this proposition by induction. For simplicity, we often omit the super-script A .

Define $Z_1 = AY_+$ and $\mathcal{V}_1 = e^{-e_0t} Z_1$. Thus,

$$\partial_t \mathcal{V}_1 + \mathcal{L} \mathcal{V}_1 - iR(\mathcal{V}_1) = -iR(\mathcal{V}_1).$$

Note now that $R(f) = Q^p J(Q^{-1}f)$, where

$$J(z) = |1 + z|^{p-1}(1 + z) - 1 - \frac{p + 1}{2} z - \frac{p - 1}{2} \bar{z}$$

is real-analytic in the disc $\{z : |z| < 1\}$, and satisfies $J(0) = \partial_z J(0) = \partial_{\bar{z}} J(0) = 0$. Write its Taylor expansion as

$$(4.2) \quad J(z) = \sum_{i+j \geq 2} a_{ij} z^i \bar{z}^j;$$

this series and all of its derivatives converge uniformly in the compact disc $\{z : |z| \leq 1/2\}$.

Now, if $s_c = 1$, since $Z_1 \in \mathcal{S}(\mathbb{R}^N)$ and W decays polynomially, we have $\|W^{-1}Z_1\|_{L^\infty} < +\infty$. For $0 < s_c < 1$, we make use of Corollary 3.8 (ii), to conclude that $\|Q^{-1}Z_1\|_{L^\infty} < +\infty$. In any case, we can choose t_0 such that $|\mathcal{V}_1(t)| \leq \frac{1}{2}Q$, for any $t \geq t_0$. Therefore, for large t , we have

$$|R(\mathcal{V}_1)| \leq \|Q\|_{L^\infty}^p \left(\sum_{i+j \geq 0} |a_{ij}| \frac{1}{2^{i+j}} \right) |Q^{-1}\mathcal{V}_1|^2 = C|Q^{-1}\mathcal{V}_1|^2.$$

In the same fashion, we can use the Leibniz rule, equation (4.2) and items (i) and (ii) of Corollary 3.8 to bound all the derivatives of $R(\mathcal{V}_1)$. Using that $\mathcal{V}_1 = e^{-e_0 t} Z_1$, we conclude that $R(\mathcal{V}_1) = O(e^{-2e_0 t})$ in $\mathcal{S}(\mathbb{R}^N)$. Moreover, by Corollary 3.8 (ii), we have $\|Q^{-1}e^{\eta|x|}\partial^\alpha Z_1\|_{L^\infty} < +\infty$.

Now let $k \geq 1$ and assume that the \mathcal{V}_i are defined and satisfy (4.1) for all $i \leq k$. For $0 < s_c < 1$, assume furthermore that, for all $i \leq k$, and all multi-indices α ,

$$(4.3) \quad \|Q^{-1}e^{\eta|x|}\partial^\alpha Z_i\|_{L^\infty} < +\infty.$$

Define

$$(4.4) \quad \epsilon_k = \partial_t \mathcal{V}_k + \mathcal{L}\mathcal{V}_k - iR(\mathcal{V}_k),$$

and note that

$$\partial_t \mathcal{V}_k = \sum_{j=1}^k (-je_0) e^{-je_0 t} Z_k,$$

so that (4.4) can be written as

$$(4.5) \quad \epsilon_k(x, t) = \sum_{j=1}^k e^{-je_0 t} (-je_0 Z_k(x) + \mathcal{L}Z_k(x)) - iR(\mathcal{V}_k(x, t)).$$

Recall that, for all k , $Z_k \in \mathcal{S}(\mathbb{R}^N)$. If $0 < s_c < 1$, we also have (4.3). Therefore, for large t , and all x , $|\mathcal{V}_k(x, t)| \leq \frac{1}{2}Q(x)$. Writing $R(\mathcal{V}_k) = Q^p J(Q^{-1}\mathcal{V}_k)$ and using again the expansion (4.2), we get by (4.5) that there exist functions $F_j \in \mathcal{S}(\mathbb{R}^N)$ such that for large t ,

$$\epsilon_k(x, t) = \sum_{j=1}^{k+1} e^{-je_0 t} F_j(x) + O(e^{-e_0(k+2)t}) \quad \text{in } \mathcal{S}(\mathbb{R}^N).$$

By (4.1), we conclude that $F_j = 0$ for $j \leq k$, which shows

$$(4.6) \quad \epsilon_k(x, t) = e^{-(k+1)e_0 t} F_{k+1} + O(e^{-(k+2)e_0 t}).$$

Note that $(k + 1)e_0$ is not in the spectrum of \mathcal{L} , and define

$$Z_{k+1} = -(\mathcal{L} + (k + 1)e_0)^{-1} F_{k+1},$$

which belongs to \mathcal{S} (see Appendix A). Moreover, if $0 < s_c < 1$, Z_{k+1} satisfies (4.3) with k replaced by $k + 1$. By definition, we have $\mathcal{V}_{k+1} = \mathcal{V}_k + e^{-(k+1)e_0 t} Z_{k+1}$. Furthermore,

$$\epsilon_{k+1} = \epsilon_k - e^{-(k+1)e_0 t} F_{k+1} - i(R(\mathcal{V}_{k+1}) - R(\mathcal{V}_k)).$$

By (4.6), $\epsilon_k - e^{-(k+1)e_0 t} F_{k+1} = O(e^{-(k+2)e_0 t})$. Writing again $R(f) = Q^p J(Q^{-1}f)$, and using the expansion (4.2), we conclude that $R(\mathcal{V}_{k+1}) - R(\mathcal{V}_k) = O(e^{-(k+2)e_0 t})$, completing the proof. ■

4.2. Contraction argument near an approximate solution

We now prove the key result of this section. The propositions for the energy-critical and for the intercritical cases are stated separately.

4.2.1. Energy-critical case. We only treat here the case $N \geq 6$, as in the lower-dimensional cases this result is proved in [12]. The main difference here from [12] is that $0 < p_c - 1 < 1$ if $N > 6$, so that the nonlinearity is no longer C^2 , and its derivative is only Hölder-continuous of order $p_c - 1$. This introduces difficulties, as the control of the convergence of ∇U^A to ∇W is not enough to close the contraction argument, and we need to ensure that the higher order terms $D^\varepsilon(U^A - W - \mathcal{V}_k)$ converge faster to 0, for small $\varepsilon > 0$. The fractional derivative D^ε is needed here to avoid certain end-point Strichartz estimates, which are not available for any combination of \dot{H}^1 -admissible and \dot{H}^{-1} -admissible pairs.

Proposition 4.2. *Let $N \geq 6$. There exists $k_0 > 0$ such that for any $k \geq k_0$, there exist $t_k \geq 0$ and a solution U^A to (1.2) such that for $t \geq t_k$ and $l(k) = \lceil \frac{N-2}{4}k + \frac{N-6}{4} \rceil$ (where $\lceil x \rceil$ denotes the least integer bigger or equal to x),*

$$(4.7) \quad \begin{aligned} \|D^\varepsilon(U^A - W - \mathcal{V}_{l(k)}^A)\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))} &\leq e^{-(k+\frac{1}{2})\frac{N-2}{4}e_0t} \quad \text{and} \\ \|\nabla(U^A - W - \mathcal{V}_{l(k)}^A)\|_{S(L^2, [t, +\infty))} &\leq e^{-(k+\frac{1}{2})e_0t}. \end{aligned}$$

Furthermore, U^A is the unique solution to (1.2) satisfying (4.7) for large t . Finally, U^A is independent of k and satisfies, for large t ,

$$(4.8) \quad \|U^A(t) - W - Ae^{-e_0t}y_+\|_{\dot{H}^1} \leq e^{-2e_0t}.$$

Proof. Since $A \in \mathbb{R}$ is fixed in the proof, we omit the superscripts A . Define

$$h = U^A - W - \mathcal{V}_{l(k)}^A,$$

so that U^A is a solution to (1.2) if and only if h satisfies

$$i \partial_t h + \Delta h = -K(h) - (R(\mathcal{V}_{l(k)} + h) - R(\mathcal{V}_{l(k)})) + i\epsilon_{l(k)},$$

where $\epsilon_{l(k)} = O(e^{-(l(k)+1)e_0t})$ in $\mathcal{S}(\mathbb{R}^N)$ for all $k \geq 0$. Therefore, the existence of U^A can be written as the fixed-point problem

$$h(t) = \mathcal{M}(h)(t),$$

where

$$\mathcal{M}(h)(t) = -i \int_t^{+\infty} e^{i(t-s)\Delta} [-K(h) - (R(\mathcal{V}_{l(k)} + h) - R(\mathcal{V}_{l(k)})) + i\epsilon_{l(k)}] ds.$$

Let first $N > 6$. We will show that \mathcal{M} is a contraction on B defined by

$$B = B(k, t_k) := \{h \in E : \|h\|_E \leq 1\},$$

$$E = E(k, t_k) := \{h \in S(\dot{H}^1, [t_k, +\infty)), D^\varepsilon h \in S(\dot{H}^{1-\varepsilon}, [t_k, +\infty)), \nabla h \in S(L^2, [t_k, +\infty)) : \|h\|_E < +\infty\},$$

$$\|h\|_E := \sup_{t \geq t_k} e^{(k+\frac{1}{2})\frac{N-2}{4}e_0t} \|D^\varepsilon h\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))} + \sup_{t \geq t_k} e^{(k+\frac{1}{2})e_0t} \|\nabla h\|_{S(L^2, [t, +\infty))},$$

equipped with the metric

$$\rho(u, v) = \sup_{t \geq t_k} e^{(k+\frac{1}{2})\frac{N-2}{4}e_0 t} \|D^\varepsilon(u - v)\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))}.$$

Let $\{h_n\} \subset B$ and $h \in S(\dot{H}^{1-\varepsilon}, I)$ with $\rho(h_n, h) \rightarrow 0$. By reflexivity and uniqueness between the weak and strong limits, $h \in B$. This shows that (B, ρ) is a complete metric space. We will show that $\mathcal{M}(B) \subset B$ and that \mathcal{M} is a contraction.

By Strichartz estimates, there exists $C^* > 0$ such that

$$(4.9) \quad \|\nabla \mathcal{M}(h)\|_{S(L^2, [t, +\infty))} \leq C^* [\|\nabla K(h)\|_{S'(L^2, [t, +\infty))} + \|\nabla(R(\mathcal{V}_l(k) + h) - R(\mathcal{V}_l(k)))\|_{S'(L^2, [t, +\infty))} + \|\nabla \epsilon_l(k)\|_{S'(L^2, [t, +\infty))}],$$

$$(4.10) \quad \|D^\varepsilon \mathcal{M}(h)\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))} \leq C^* [\|D^\varepsilon K(h)\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} + \|D^\varepsilon(R(\mathcal{V}_l(k) + h) - R(\mathcal{V}_l(k)))\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} + \|D^\varepsilon \epsilon_l(k)\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))}],$$

and

$$(4.11) \quad \|D^\varepsilon(\mathcal{M}(g) - \mathcal{M}(h))\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))} \leq C^* [\|D^\varepsilon K(g - h)\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} + \|D^\varepsilon(R(\mathcal{V}_l(k) + g) - R(\mathcal{V}_l(k) + h))\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))}].$$

To finish the argument, we need the following estimates.

Lemma 4.3. *For every $\eta > 0$, there exists $\tilde{k}(\eta) > 0$ such that, if $k \geq \tilde{k}(\eta)$, then for any $g, h \in B$, the following inequalities hold for all $t \geq t_k$:*

- (i) $\|\nabla K(h)\|_{S'(L^2, [t, +\infty))} \leq \eta e^{-(k+\frac{1}{2})e_0 t} \|h\|_E,$
- (ii) $\|\nabla(R(\mathcal{V}_l(k) + h) - R(\mathcal{V}_l(k)))\|_{S'(L^2, [t, +\infty))} \leq C_k e^{-(k+\frac{1}{2}+\frac{4}{N-2})e_0 t},$
- (iii) $\|D^\varepsilon K(h)\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} \leq \eta e^{-(k+\frac{1}{2})\frac{N-2}{4}e_0 t} \rho(h, 0),$
- (iv) $\|D^\varepsilon(R(\mathcal{V}_l(k) + g) - R(\mathcal{V}_l(k) + h))\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} \leq C_k e^{-(k+\frac{1}{2}+\frac{16}{(N-2)^2})\frac{N-2}{4}e_0 t} \rho(g, h)$
- (v) $\|\nabla \epsilon_l(k)\|_{S'(L^2, [t, +\infty))} + \|D^\varepsilon \epsilon_l(k)\|_{S'(\dot{H}^{-(1-\varepsilon)}, [t, +\infty))} \leq C_k e^{-(k+1)\frac{N-2}{4}e_0 t}.$

Indeed, assuming this lemma, choosing first $\eta > 0$ small enough, and then a large enough t_k , we see by (4.9) and (4.10) that $\mathcal{M}(B) \subset B$. Moreover, by (4.11), \mathcal{M} is a contraction on B . Thus, for every $k \geq k_0 = \tilde{k}(\eta)$, there is a unique solution U^A to (1.2) satisfying (4.7) for $t \geq t_k$. Note that the uniqueness still holds in the class of solutions to (1.2) satisfying (4.7) for $t \geq t'_k$, where $t'_k \geq t_k$. Thus, uniqueness of the solutions to (1.2) shows that U^A does not depend on k .

It remains to show Lemma 4.3. By Lemma 3.9 (i), if $\tau_0 > 0$ and $t \geq t_k$, then

$$\|\nabla K(h)\|_{S'(L^2, [t, t+\tau_0])} \leq C \tau_0^{1/2} e^{-(k+\frac{1}{2})e_0 t} \|h\|_E.$$

Summing up this equation at times $t_j = t + j\tau_0$, and using the triangle inequality, we get a geometric series, whose sum is

$$\|\nabla K(h)\|_{S'(L^2, [t, +\infty))} \leq C \frac{\tau_0^{1/2}}{1 - e^{-(k+\frac{1}{2})\frac{4}{N-2}e_0 \tau_0}} e^{-(k+\frac{1}{2})e_0 t} \|h\|_E.$$

Choosing τ_0 and k_0 such that $\tau_0^{1/2} = \eta/(2C)$ and $e^{-(k+\frac{1}{2})e_0\tau_0} \leq 1/2$, we get the estimate (i) of Lemma 4.3. The estimate (iii) follows analogously from Lemma 3.9 (iii).

We now turn to the item (ii). By Lemma 3.9 (ii), we have

$$\begin{aligned} & \|\nabla(R(\mathcal{V}_{l(k)} + h) - R(\mathcal{V}_{l(k)}))\|_{S'(L^2, [t, t+1])} \\ & \leq (A) \cdot \|\nabla h\|_{S(L^2, [t, t+1])} + (B) \cdot \|D^\varepsilon h\|_{S(\dot{H}^{1-\varepsilon}, [t, t+1])}^{\frac{4}{N-2}}, \end{aligned}$$

where

$$(A) \lesssim \|\nabla \mathcal{V}_{l(k)}\|_{S(L^2, [t, t+1])}^{\frac{4}{N-2}} + \|\nabla h\|_{S(L^2, [t, t+1])}^{\frac{4}{N-2}}$$

and

$$(B) \lesssim \|\nabla \mathcal{V}_{l(k)}\|_{S(L^2, [t, t+1])} + \|\nabla h\|_{S(L^2, [t, t+1])}.$$

By the explicit form of \mathcal{V}_k and the fact that $h \in B$, we get

$$(A) + (B) \leq C_k e^{-e_0 \frac{4}{N-2} t}.$$

Therefore,

$$\begin{aligned} & \|(R(\mathcal{V}_{l(k)} + h) - R(\mathcal{V}_{l(k)}))\|_{S'(L^2, [t, t+1])} \\ & \leq C_k e^{-e_0 \frac{4}{N-2} t} (\|\nabla h\|_{S(L^2, [t, t+1])} + \|D^\varepsilon h\|_{S(\dot{H}^{1-\varepsilon}, [t, t+1])}^{\frac{4}{N-2}}) \leq C_k e^{-(k+\frac{1}{2}+\frac{4}{N-2})e_0 t}. \end{aligned}$$

Since $h \in B$, the triangle inequality and the sum of the resulting geometric series gives us (ii). As for the item (iv), it follows analogously from Lemma 3.9 (iv). The estimate (v) of Lemma 4.3 follows immediately from (4.6) and from the bound

$$l(k) + 1 \geq (k + 1) \frac{N - 2}{4}.$$

Finally, given that $U^A = W + \mathcal{V}_k + h$ with $h \in B$, we see that, for large k ,

$$\|\nabla h(t)\|_{L^2} \leq e^{-\frac{5}{2}e_0 t} \|h\|_E \leq e^{-\frac{5}{2}e_0 t},$$

and recalling the definition of \mathcal{V}_k given in Proposition 4.1, we have, for all k ,

$$\mathcal{V}_{l(k)} = Ae^{-e_0 t} y_+ + O(e^{-2e_0 t}) \quad \text{in } \mathcal{S}(\mathbb{R}^N),$$

which proves (4.8), and finishes the case $N > 6$.

For the case $N = 6$, note that $(N - 2)/4 = 1$, so that, by the Sobolev embedding, only the space $S(L^2, I)$ is enough for the contraction argument. Therefore, defining the space B as

$$\begin{aligned} B &= B(k, t_k) := \{h \in E : \|h\|_E \leq 1\}, \\ E &= E(k, t_k) := \{h \in C_t \dot{H}^1([t_k, +\infty)) \cap S(\dot{H}^1, [t_k, +\infty)), \\ & \quad \nabla h \in S(L^2, [t_k, +\infty)) : \|h\|_E < +\infty\}, \\ \|h\|_E &:= \sup_{t \geq t_k} e^{(k+\frac{1}{2})e_0 t} \|\nabla h\|_{S(L^2, [t, +\infty))}, \end{aligned}$$

equipped with the metric

$$\rho(u, v) = \sup_{t \geq t_k} e^{(k+\frac{1}{2})e_0 t} \|\nabla(u - v)\|_{S(L^2, [t, +\infty))},$$

we observe that, by Lemma 3.9, the analogous estimates of Lemma 4.3 hold. Hence, the conclusion of Proposition 4.2 also holds for $N = 6$, finishing its proof. ■

4.2.2. Intercritical case.

Proposition 4.4. *There exists $k_0 > 0$ such that for any $k \geq k_0$, there exist $t_k \geq 0$ and a solution U^A to (1.1) such that for $t \geq t_k$ and $l(k) = \max\{\lceil(k + 1)/(p - 1) - 1\rceil, k\}$, we have*

$$(4.12) \quad \|U^A - e^{it} Q - e^{it} \mathcal{V}_{l(k)}^A\|_{S(\dot{H}^{s_c}, [t, +\infty))} \leq e^{-(k+\frac{1}{2}) \max\{\frac{1}{p-1}, 1\} e_0 t}$$

and

$$(4.13) \quad \|(\nabla)(U^A - e^{it} Q - e^{it} \mathcal{V}_{l(k)}^A)\|_{Z(t, +\infty)} \leq e^{-(k+\frac{1}{2}) e_0 t}.$$

Furthermore, U^A is the unique solution to the NLS equation (1.1) satisfying (4.13) for large t . Finally, U^A is independent of k and satisfies, for large t ,

$$(4.14) \quad \|U^A(t) - e^{it} Q - Ae^{-e_0 t + it} \mathcal{Y}_+\|_{H^1} \leq e^{-2e_0 t}.$$

In view of Lemma 3.13, the proof of Proposition 4.4 is essentially the same as in the energy-critical case, and we state Lemma 4.5 below for completeness. Note that (4.12) is a consequence of (4.13) in the case $p \geq 2$, due to the Sobolev inequalities.

Lemma 4.5. *For every $\eta > 0$, there exists $\tilde{k}(\eta) > 0$ such that, if $k \geq \tilde{k}(\eta)$, then for any $g, h \in B$ the following inequalities hold for all $t \geq t_k$.*

- (i) $\|\nabla K(h)\|_{S'(L^2, [t, +\infty))} \leq \eta e^{-(k+\frac{1}{2}) e_0 t} \|h\|_E,$
- (ii) $\|\nabla(R(\mathcal{V}_{l(k)} + h) - R(\mathcal{V}_{l(k)}))\|_{S'(L^2, [t, +\infty))} \leq C_k e^{-\min\{p-1, 1\} e_0 t} e^{-(k+\frac{1}{2}) e_0 t},$
- (iii) $\|K(h)\|_{S'(\dot{H}^{-s_c}, [t, +\infty))} \leq \eta e^{-(k+\frac{1}{2}) \max\{\frac{1}{p-1}, 1\} e_0 t} \rho(h, 0),$
- (iv) $\|R(\mathcal{V}_{l(k)} + g) - R(\mathcal{V}_{l(k)} + h)\|_{S'(\dot{H}^{-s_c}, [t, +\infty))} \leq C_k e^{-\min\{p-1, 1\} e_0 t} e^{-(k+\frac{1}{2}) \max\{\frac{1}{p-1}, 1\} e_0 t} \rho(g, h),$
- (v) $\|\nabla \epsilon_{l(k)}\|_{S'(L^2, [t, +\infty))} + \|\epsilon_{l(k)}\|_{S'(\dot{H}^{-s_c}, [t, +\infty))} \leq C_k e^{-(k+1) \max\{\frac{1}{p-1}, 1\} e_0 t}.$

5. Modulation

Throughout the rest of the paper, we write, for $0 < s_c \leq 1$,

$$d(f) = \left| \|\nabla f\|_{L^2} - \|\nabla Q\|_{L^2} \right|.$$

If u is a solution to (1.1) or to (1.2) and if there is no risk of ambiguity, we also write

$$d(t) = d(u(t)).$$

5.1. Energy-critical case

The variational characterization of W , see [2, 32, 38], shows that, if $E(f) = E(W)$, then

$$\inf_{\substack{x \in \mathbb{R}^N \\ \lambda > 0 \\ \theta \in \mathbb{R}}} \|f_{[x,\lambda,\theta]} - W\|_{\dot{H}^1} \leq \epsilon(d(f))$$

with $\epsilon = \epsilon(\delta)$ such that

$$\lim_{\delta \rightarrow 0^+} \epsilon(\delta) = 0.$$

The goal of this section is to construct modulation parameters x_0, λ_0 and θ_0 such that the quantity $d(f)$ controls linearly $\|f_{[x_0,\lambda_0,\theta_0]} - W\|_{\dot{H}^1}$ as well as the parameters and its derivatives.

By making use of the implicit function theorem, we can construct appropriate modulation parameters. The proof of the next two lemmas is very similar to the ones in [12], with the introduction of a translation parameter, and will be given in Appendix A.

Lemma 5.1. *There exist $\delta_0 > 0$ and a positive function $\epsilon(d)$ defined for $0 < d < \delta_0$, with $\epsilon(d) \rightarrow 0$ as $d \rightarrow 0$, such that, for all $f \in \dot{H}^1$ satisfying $E(f) = E(W)$ and $d(f) < \delta_0$, there exist (x, λ, θ) such that*

$$\|f_{[x,\lambda,\theta]} - W\|_{\dot{H}^1} \leq \epsilon(d(u)) \quad \text{and} \quad f_{[x,\lambda,\theta]} \perp \text{span}\{\nabla W, iW, \Lambda W\}.$$

The parameters (x, λ, θ) are unique in $\mathbb{R}^N \times \mathbb{R}/2\pi\mathbb{Z} \times \mathbb{R}_+$, and the mapping $u \mapsto (x, \lambda, \theta)$ is C^1 .

Let u be a solution to (1.2) and let I be a time interval such that

$$d(t) < \delta_0 \quad \text{for all } t \in I.$$

For each $t \in I$, choose the parameters $(x(t), \lambda(t), \theta(t))$ according to Lemma 5.1. Write

$$(5.1) \quad u_{[x(t),\lambda(t),\theta(t)]}(t) = (1 + \alpha(t))W + h(t),$$

where

$$\alpha(t) = \frac{1}{\|W\|_{\dot{H}^1}} (u_{[x(t),\lambda(t),\theta(t)]}, W)_{\dot{H}^1} - 1.$$

Note that $\alpha(t)$ is chosen so that $h(t) \in G^\perp$. By Lemma 5.1 and a standard regularization argument (see, for instance Lemma 4 in [33] for details in a similar context), the map $t \mapsto (x(t), \lambda(t), \theta(t))$ is C^1 . We are now able to prove estimates on the modulation.

Lemma 5.2. *Let u be a solution to (1.2) satisfying $E(u_0) = E(W)$. Taking a smaller δ_0 , if necessary, the following estimates hold on I :*

$$(5.2) \quad |\alpha(t)| \approx \|h(t)\|_{\dot{H}^1} \approx d(t) < \delta_0,$$

$$(5.3) \quad |\alpha'(t)| + |x'(t)| + |\theta'(t)| + \left| \frac{\lambda'(t)}{\lambda(t)} \right| \lesssim \lambda^2(t)d(t).$$

In the next two sections, we mainly consider *radial* solutions to (1.2). Since the translation parameter is not needed, we write

$$f_{[\lambda_0, \theta_0]}(x) = e^{i\theta_0} \frac{1}{\lambda_0^{(N-2)/2}} f\left(\frac{x}{\lambda_0}\right).$$

In some results regarding compactness, the parameter θ_0 can also be omitted. In this case, we write

$$f_{[\lambda_0]}(x) = \frac{1}{\lambda_0^{(N-2)/2}} f\left(\frac{x}{\lambda_0}\right).$$

For solutions to (1.1), since the scaling parameter is fixed (see Remark 1.7), we use the notation

$$f_{[x_0, \theta_0]}(x) = e^{i\theta_0} f(x + x_0).$$

When θ_0 can be omitted, we write

$$f_{[x_0]}(x) = f(x + x_0).$$

5.2. Intercritical case

For $0 < s_c < 1$, the variational characterization of Q , [31], shows that, if $M(f) = M(Q)$ and $E(f) = E(Q)$, then

$$(5.4) \quad \inf_{\substack{x \in \mathbb{R}^N \\ \theta \in \mathbb{R}}} \|f_{[x, \theta]} - Q\|_{H^1} \leq \epsilon(d(f)),$$

with

$$\lim_{\delta \rightarrow 0^+} \epsilon(\delta) = 0.$$

As before, the goal here is to construct modulation parameters x_0 and θ_0 such that the quantity $d(f)$ controls linearly $\|f_{[x_0, \theta_0]} - Q\|_{H^1}$, as well as the parameters and its derivatives. We follow mainly [13] here, and the proofs for the next two lemmas are almost identical; thus, we omit them.

Lemma 5.3. *There exist $\delta_0 > 0$ and a positive function $\epsilon(d)$ defined for $0 < d < \delta_0$, with $\epsilon(d) \rightarrow 0$ as $d \rightarrow 0$, such that, for all $f \in H^1$ satisfying $M(f) = M(Q)$, $E(f) = E(Q)$ and $d(f) < \delta_0$, there exist (x, θ) such that*

$$\|f_{[x, \theta]} - Q\|_{H^1} \leq \epsilon(d(f)) \quad \text{and} \quad f_{[x, \theta]} \perp \text{span}\{\nabla Q, iQ\}.$$

The parameters (x, θ) are unique in $\mathbb{R}^N \times \mathbb{R}/2\pi\mathbb{Z}$, and the mapping $u \mapsto (x, \theta)$ is C^1 .

Let u be a solution to (1.1) and let I be a time interval such that $d(t) := d(u(t)) < \delta_0$ for all $t \in I$. For each $t \in I$, choose the parameters $(x(t), \theta(t))$ according to Lemma 5.3. Write

$$(5.5) \quad e^{-it} u_{[x(t), \theta(t)]}(t) = (1 + \alpha(t))Q + h(t),$$

where

$$\alpha(t) = \frac{\text{Re}(e^{-it} \int Q^p u_{[x(t), \theta(t)]})}{\|Q\|_{L^{p+1}}^{p+1}} - 1.$$

Note that $\alpha(t)$ is chosen so that $h(t) \in G^\perp$. Recall that the parameters (x, θ) are C^1 . We are now able to prove estimates on the modulation.

Lemma 5.4. *Let u be a solution to (1.1) satisfying $M(u_0) = M(Q)$ and $E(u_0) = E(Q)$. Taking a smaller δ_0 , if necessary, the following estimates hold on I :*

$$(5.6) \quad |\alpha(t)| \approx \|h(t)\|_{H^1} \approx \left| \int Qh_1 \right| \approx d(t),$$

$$(5.7) \quad |\alpha'(t)| \approx |x'(t)| \approx |\theta'(t)| \lesssim d(t).$$

We finish this section with a lemma that will be useful in later sections.

Lemma 5.5. *Let u be a solution to (1.1) such that $M(u_0) = M(Q)$ and $E(u_0) = E(Q)$. Assume that u is defined on $[0, +\infty)$ and that there exists $c > 0$ such that*

$$(5.8) \quad \int_t^{+\infty} d(s) ds \lesssim e^{-ct}.$$

Then there exist (x_0, θ_0) such that

$$\|u_{[x_0, \theta_0]} - e^{it}Q\|_{H^1} \lesssim e^{-ct}.$$

Proof. Step 1. Convergence of $\delta(t)$. We claim that

$$(5.9) \quad \lim_{t \rightarrow +\infty} d(t) = 0.$$

To prove this, first note that (5.8) implies that there exists a sequence $\{t_n\}$ with $t_n \rightarrow +\infty$ such that

$$(5.10) \quad \lim_{n \rightarrow +\infty} d(t_n) = 0.$$

Suppose now, by contradiction, that (5.9) does not hold. In this case, we can find another sequence $\{t'_n\}$ and $0 < \epsilon_1 < \delta_0$ such that

$$(5.11) \quad t_n < t'_n \quad \forall n, \quad d(t'_n) = \epsilon_1 \quad \text{and} \quad d(t) < \epsilon_1 \quad \forall t \in [t_n, t'_n].$$

Since $\epsilon_1 < \delta_0$, the parameter $\alpha(t)$ is well-defined on $[t_n, t'_n]$. By Lemma 5.4, $|\alpha'(t)| \lesssim d(t)$, so that $\int_{t_n}^{t'_n} |\alpha'(t)| dt \lesssim e^{-ct_n}$, by (5.8). Therefore,

$$\lim_{n \rightarrow \infty} |\alpha(t_n) - \alpha(t'_n)| = 0.$$

Since, by Lemma (5.4), $|\alpha(t)| \approx d(t)$, we get that $\alpha(t_n)$ tends to 0, which contradicts (5.11) and proves (5.9). Recalling the decomposition (5.5), to conclude the proof of Lemma 5.5, it is sufficient to prove that there exists (x_0, θ_0) such that

$$d(t) + |\alpha(t)| + \|h(t)\|_{\dot{H}^1} + |x(t) - x_0| + |\theta(t) - \theta_0| \lesssim e^{-ct}.$$

By Lemma 5.4, $|\alpha(t)| \approx d(t) \rightarrow 0$ as $t \rightarrow +\infty$, and hence,

$$|\alpha(t)| \leq \int_t^{+\infty} |\alpha'(s)| ds \lesssim \int_t^{+\infty} d(s) ds \lesssim e^{-ct},$$

since $|\alpha'(t)| \approx d(t)$. Again by Lemma 5.4, $d(t) \approx \|h(t)\|_{\dot{H}^1} \approx |\alpha(t)|$, and we get the bounds on $d(t)$ and $\|h(t)\|_{\dot{H}^1}$. To obtain the bounds on $x(t)$ and $\theta(t)$, it is sufficient to recall that Lemma 5.4 says that $|x'(t)| + |\theta'(t)| \lesssim d(t) \lesssim e^{-ct}$. ■

6. Convergence to the standing wave above the ground state

6.1. Energy-critical case

In this and the next sections, we prove that radial solutions to (1.2) on the same energy level as W that do not blow-up in finite positive time (and have finite mass), and that do not scatter forward in time, must converge exponentially to W as $t \rightarrow +\infty$. We follow closely [12], and give the proofs in Appendix A.

Proposition 6.1. *Let u be a solution to (1.2) defined on $[0, +\infty)$ satisfying*

$$(6.1) \quad E(u_0) = E(W) \quad \text{and} \quad \|u_0\|_{\dot{H}^1} > \|W\|_{\dot{H}^1}.$$

Assume, furthermore, that u_0 is radial and belongs to $L^2(\mathbb{R}^N)$. Then there exist (λ_0, θ_0) and $c > 0$ such that

$$(6.2) \quad \|u - W_{[\lambda_0, \theta_0]}\|_{\dot{H}^1} \lesssim e^{-ct}.$$

Moreover, u blows up in finite negative time.

We will work with a truncated variance. Consider a radial function $\phi \in C_0^\infty(\mathbb{R}^N)$ such that

$$\phi(r) \geq 0 \quad \forall r \geq 0, \quad \phi(r) = \begin{cases} r^2, & r \leq 1, \\ 0, & r \geq 3, \end{cases} \quad \text{and} \quad \frac{d^2\phi}{dr^2}(r) \leq 2, r \geq 0.$$

Define $\phi_R(x) = R^2\phi(x/R)$ and

$$F_R(t) = \int \phi_R |u(t)|^2.$$

By virial identities, if $E(u_0) = E(W)$, we have

$$(6.3) \quad F'_R(t) = 2 \operatorname{Im} \int \nabla \phi \cdot \nabla u \bar{u}$$

$$(6.4) \quad F''_R(t) = -\frac{16}{N-2} d(t) + A_R(u(t)),$$

where

$$(6.5) \quad A_R(u(t)) = \int_{|x| \geq R} |\nabla u(t)|^2 (4\partial_r^2 \phi_R - 8) + \int_{|x| \geq R} |u|^{2^*} \left(8 - \frac{4}{N} \Delta \phi_R\right) - \int_{|x| \geq R} |u|^2 \Delta^2 \phi_R.$$

Recall that, if $\|\nabla u_0\|_{\dot{H}^1} > \|\nabla W\|_{\dot{H}^1}$, then, for all t in the interval of definition of u ,

$$d(t) = \left| \|\nabla u(t)\|_{\dot{H}^1} - \|\nabla W\|_{\dot{H}^1} \right| = \|\nabla u(t)\|_{\dot{H}^1} - \|\nabla W\|_{\dot{H}^1}.$$

In order to prove Proposition 6.1, we need the following lemma, which is also proved in Appendix A.

Lemma 6.2. *Let u be a radial solution to (1.2) defined on $[0, +\infty)$ and satisfying (6.1). Assume, furthermore, that the mass $M(u_0)$ is finite. Then, there exists a constant $R_0 > 0$ such that, for all t in the interval of existence of u and all $R \geq R_0$,*

$$(6.6) \quad F'_R(t) > 0,$$

and there exists $c > 0$ such that

$$(6.7) \quad \int_t^{+\infty} d(s) ds \lesssim e^{-ct}, \quad \forall t \geq 0.$$

6.2. Intercritical case

Here, we state the corresponding result for the intercritical case. Since the proof is very similar to the energy-critical case, we mainly sketch it in Appendix A.

Proposition 6.3. *Let u be a solution to (1.1) defined on $[0, +\infty)$ satisfying*

$$(6.8) \quad M(u_0) = M(Q), \quad E(u_0) = E(Q) \quad \text{and} \quad \|\nabla u_0\|_{L^2} > \|\nabla Q\|_{L^2}.$$

Assume, furthermore, that either u_0 is radial or $|x|u_0 \in L^2(\mathbb{R}^N)$. Then there exist (x_0, θ_0) and $c > 0$ such that

$$(6.9) \quad \|u - e^{it} Q_{[x_0, \theta_0]}\|_{H^1} \lesssim e^{-ct}.$$

Moreover, u blows up in finite negative time.

7. Convergence to the standing wave below the ground state

7.1. Energy-critical case

In this section, we consider solutions such that

$$(7.1) \quad E(u_0) = E(W) \quad \text{and} \quad \|u_0\|_{\dot{H}^1} < \|W\|_{\dot{H}^1}.$$

Definition 7.1. A solution u to (1.2) with lifespan I is said to be *almost periodic modulo symmetries* on $J \subset I$ if there exist functions $x: J \rightarrow \mathbb{R}^N$, $\lambda: J \rightarrow \mathbb{R}_+^*$ and $C: \mathbb{R}_*^+ \rightarrow \mathbb{R}_*^+$ such that for all $t \in I$ and all $\eta > 0$,

$$\int_{|x-x(t)| \geq C(\eta)/\lambda(t)} |\nabla u(x, t)|^2 dx \leq \eta \quad \text{and} \quad \int_{|\xi| \geq C(\eta)\lambda(t)} |\xi|^2 |\hat{u}(\xi, t)|^2 d\xi \leq \eta.$$

Remark 7.2. By the Arzelà–Ascoli theorem, almost periodicity modulo symmetries is equivalent to the set $\{u_{[x(t), \lambda(t), 0]} : t \in J\}$ being precompact in \dot{H}^1 .

Remark 7.3. If the solution is radial, $x(t)$ can be chosen to be zero.

Proposition 7.4. *Let u be a solution to (1.2) and let $I = (T^-, T^+)$ be its maximal interval of existence. If u satisfies (7.1), then $I = \mathbb{R}$.*

Furthermore, if

$$(7.2) \quad \int_0^{+\infty} \int_{\mathbb{R}^N} |u(x, t)|^{\frac{2(N+2)}{N-2}} dx dt = +\infty,$$

then u is almost periodic modulo symmetries on $[0, +\infty)$.

The proof of Proposition 7.4 is essentially contained in the proof of Proposition 3.1 in [28], which extended the work in [27] to dimensions $N \geq 6$.

Remark 7.5. By time-reversal symmetry, the analogous version of (7.2) for the interval $(-\infty, 0]$ holds.

The next theorem is the main result proved in [28] (Theorem 1.7).

Theorem 7.6. For $N \geq 5$, let $u: I \times \mathbb{R}^N \rightarrow \mathbb{C}$ be a solution to (1.2) satisfying

$$E_* := \sup_{t \in I} \|u(t)\|_{\dot{H}^1} < \|W\|_{\dot{H}^1}.$$

Then,

$$\int_I \int_{\mathbb{R}^N} |u(x, t)|^{\frac{2(N+2)}{N-2}} dx dt = C(E_*) < +\infty.$$

In particular, by uniqueness of solutions and continuity of the flow of (1.2), we have the following consequence.

Corollary 7.7. For $N \geq 5$, let u be a solution to (1.2) satisfying (7.1) and (7.2). Then there exists a sequence $t_n \rightarrow +\infty$ such that

$$\lim_{n \rightarrow +\infty} d(u(t_n)) = 0.$$

The main aim of this section is to prove the following proposition.

Proposition 7.8. Let u be a radial solution to (1.2) satisfying (7.1) and (7.2). Then there exist (λ_0, θ_0) and $c > 0$ such that, for all $t \geq 0$,

$$(7.3) \quad \|u - W_{[\lambda_0, \theta_0]}\|_{\dot{H}^1} \lesssim e^{-ct}.$$

Moreover, u scatters backward in time.

The proof follows the same lines as in [12], and is given in Appendix A.

7.2. Intercritical case

Here, we consider solutions such that

$$(7.4) \quad M(u_0) = M(Q), \quad E(u_0) = E(Q) \quad \text{and} \quad \|\nabla u_0\|_{L^2} < \|\nabla Q\|_{L^2}.$$

Since the scaling parameter is fixed a priori in the intercritical regime, controlling scaling is no longer an issue. We can then use the fact that the solution has finite mass, together with information given by the virial-type and compactness arguments, to control the translation parameter, allowing us to prove results in the non-radial setting. We start with a definition.

Definition 7.9. A solution u to (1.1) with lifespan I is said to be *almost periodic modulo symmetries* on $J \subset I$ if there exist functions $x: J \rightarrow \mathbb{R}^N$ and $C: \mathbb{R}_*^+ \rightarrow \mathbb{R}_*^+$ such that for all $t \in J$ and all $\eta > 0$,

$$\int_{|x-x(t)| \geq C(\eta)} |\nabla u(x, t)|^2 + |u(x, t)|^2 dx \leq \eta.$$

Remark 7.10. By the Arzelà–Ascoli theorem, almost periodicity modulo symmetries is equivalent to the set $\{u_{[x(t)]} : t \in J\}$ being precompact in H^1 .

Proposition 7.11. *Let u be a solution to (1.1) and let $I = (T^-, T^+)$ be its maximal interval of existence. If u satisfies (7.4), then $I = \mathbb{R}$.*

Furthermore, if

$$(7.5) \quad \|u\|_{S(0, +\infty)} = +\infty,$$

then u is almost periodic modulo symmetries on $[0, +\infty)$, and we have

$$P(u) = \text{Im} \int \bar{u} \nabla u = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{x(t)}{t} = 0.$$

The proof of Proposition 7.11 is now classical, and it is essentially the same as in [13], Lemma 6.2, Corollary 6.3 and Lemma 6.4.

Remark 7.12. By time-reversal symmetry, the analogous version of (7.5) for the interval $(-\infty, 0]$ holds.

Remark 7.13. As in Lemma 6.2 of [13], the function $x(t)$ can be chosen to be continuous on \mathbb{R} and the same as the one given in Lemmas 5.3 and 5.4, if $d(t) < \delta_0$.

Proposition 7.14. *Let u be a solution to (1.1) satisfying (7.4) and (7.5). Then there exist (x_0, θ_0) and $c > 0$ such that, for all $t \geq 0$,*

$$(7.6) \quad \|u - e^{it} Q_{[x_0, \theta_0]}\|_{H^1} \lesssim e^{-ct}.$$

Moreover, u scatters backward in time.

8. Estimates on exponentially decaying solutions

According to the previous sections, we must study the behavior of solutions approaching $e^{it} Q$ exponentially fast in time. We start with the energy-critical setting.

8.1. Energy-critical case

In contrast to the previous two sections, the radially assumption is not needed to prove the results in this subsection. We consider the linearized approximate equation

$$(8.1) \quad \partial_t h + \mathcal{L}h = \epsilon$$

with h and ϵ such that, for $t \geq 0$,

$$(8.2) \quad \|h(t)\|_{\dot{H}^1} \lesssim e^{-c_0 t} \quad \text{and} \quad \|\epsilon(t)\|_{L^{2N/(N+2)}} + \|\nabla \epsilon\|_{S'(L^2, [t, +\infty))} \lesssim e^{-c_1 t},$$

where $c_1 > c_0 > 0$. The following self-improving estimate was proved for radial data in [12]. We give the proof without the radial assumption in Appendix A.

Using the notation a^- for $a - \delta$ with arbitrarily small $\delta > 0$, we have the following lemma.

Lemma 8.1. *Under the assumptions (8.2),*

(i) *if $e_0 \notin [c_0, c_1)$, then*

$$(8.3) \quad \|h(t)\|_{\dot{H}^1} \lesssim e^{-c_1^- t},$$

(ii) *if $e_0 \in [c_0, c_1)$, then there exists $A \in \mathbb{R}$ such that*

$$(8.4) \quad \|h(t) - Ae^{-e_0 t} y_+\|_{\dot{H}^1} \lesssim e^{-c_1^- t}.$$

To further improve the convergence in the case $N \geq 6$, we study the linearized equation around $W + \mathcal{V}_k^A$, for $A \neq 0$, which was defined in (4.1). For simplicity, we omit the superscript A . Defining, for every k ,

$$\begin{aligned} \tilde{\mathcal{L}}_k &= \begin{pmatrix} 0 & \Delta \\ -\Delta & 0 \end{pmatrix} + \frac{(p_c + 1)}{2} |W + \mathcal{V}_k|^{p_c - 1} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ &\quad + \frac{(p_c - 1)}{2} |W + \mathcal{V}_k|^{p_c - 3} \begin{pmatrix} \text{Im}(W + \mathcal{V}_k)^2 & -\text{Re}(W + \mathcal{V}_k)^2 \\ -\text{Re}(W + \mathcal{V}_k)^2 & -\text{Im}(W + \mathcal{V}_k)^2 \end{pmatrix}, \\ \tilde{K}_k(h) &= \frac{(p_c + 1)}{2} |W + \mathcal{V}_k|^{p_c - 1} h + \frac{(p_c - 1)}{2} |W + \mathcal{V}_k|^{p_c - 3} (W + \mathcal{V}_k)^2 \bar{h}, \end{aligned}$$

and

$$\tilde{R}_k(h) = |W + \mathcal{V}_k|^{p_c - 1} (W + \mathcal{V}_k) J((W + \mathcal{V}_k)^{-1} h),$$

where

$$J(z) = |1 + z|^{p_c - 1} (1 + z) - 1 - \frac{(p_c + 1)}{2} z - \frac{(p_c - 1)}{2} \bar{z},$$

we have that if $u = W + \mathcal{V}_k + h$ satisfies (1.2), then h satisfies

$$(8.5) \quad \partial_t h + \tilde{\mathcal{L}}_k h = i \tilde{R}_k(h) + \epsilon_k,$$

or in the form of a Schrödinger equation,

$$i \partial_t h + \Delta h + \tilde{K}_k h = -\tilde{R}_k(h) + i \epsilon_k,$$

where ϵ_k are $O(e^{-(k+1)e_0 t})$ in $\mathcal{S}(\mathbb{R}^N)$. Note that the operator $\tilde{\mathcal{L}}_k$ is time-dependent and that, by the construction of \mathcal{V}_k , we have, for all $t \geq 0$,

$$|\mathcal{V}_k(t)| \lesssim e^{-e_0 t} |W| \quad \text{and} \quad |\nabla \mathcal{V}_k(t)| \lesssim e^{-e_0 t} |\nabla W| \lesssim e^{-e_0 t} |W|.$$

This implies that the estimates in Lemmas 3.9 and 3.12 hold with the same proof if we replace K by \tilde{K}_k and R by \tilde{R}_k . Therefore, we have the following results.

Lemma 8.2. *Let $N \geq 6$, $k \geq 1$ and let I be a bounded time interval. Consider $f \in S(\dot{H}^1, I)$ such that $\nabla f \in S(L^2, I)$. The following estimates hold.*

- (i) $\|\nabla \tilde{K}_k(f)\|_{S'(L^2, I)} \lesssim |I|^{1/2} \|\nabla f\|_{S(L^2, I)},$
- (ii) $\|\nabla \tilde{R}_k(f)\|_{S'(L^2, I)} + \|\tilde{R}_k(f)\|_{L^{2N/(N+2)}} \lesssim (1 + |I|^{1/2}) \|\nabla f\|_{S(L^2, I)}^{p_c}.$

Lemma 8.3. *Let h be a solution to (8.5). If, for some $c > 0$ and for any $t \geq 0$,*

$$\|h(t)\|_{\dot{H}^1} \lesssim e^{-ct},$$

then

$$(8.6) \quad \|\nabla h\|_{S(L^2, [t, +\infty))} \lesssim e^{-\min\{c, (k+1^-)e_0\}t}.$$

In the spirit of Lemma 8.1, we prove the following estimate.

Lemma 8.4. *For $N \geq 6$, let h be a solution to*

$$(8.7) \quad \partial_t h + \tilde{\mathcal{L}}_k h = \epsilon,$$

with h and ϵ such that, for $t \geq 0$,

$$(8.8) \quad \|h(t)\|_{\dot{H}^1} \lesssim e^{-c_0 t} \quad \text{and} \quad \|\epsilon(t)\|_{L^{2N/(N+2)}} + \|\nabla \epsilon\|_{S'(L^2, [t, +\infty))} \lesssim e^{-c_1 t},$$

where $(k + 1)e_0 > c_1 > c_0 > e_0$. Then,

$$(8.9) \quad \|h(t)\|_{\dot{H}^1} \lesssim e^{-c_1^- t}.$$

Proof of Lemma 8.4. Since the subscript k will be fixed in this proof, we omit it. By Lemma 8.3, we have

$$\|\nabla h(t)\|_{S(L^2, [t, +\infty))} \lesssim e^{-c_0 t}.$$

We first note that (8.7) can be written as

$$\partial_t h + \mathcal{L}h = \epsilon + (\mathcal{L} - \tilde{\mathcal{L}}_k)h.$$

Now, if $N > 6$ and $h \in \dot{H}^1$,

$$|(\mathcal{L} - \tilde{\mathcal{L}})h| \lesssim |\mathcal{V}(t)|^{p_c-1} |h| \lesssim e^{-(p_c-1)e_0 t} |W|^{p_c-1} |h|,$$

and

$$\begin{aligned} |\nabla[(\mathcal{L} - \tilde{\mathcal{L}})h]| &\lesssim |W|^{p_c-2} [e^{-e_0 t} |\nabla W| + |\nabla \mathcal{V}(t)|] |h| + |\mathcal{V}(t)|^{p_c-1} |\nabla h| \\ &\lesssim e^{-(p_c-1)e_0 t} [|W|^{p_c-1} |h| + |\nabla h|], \end{aligned}$$

where we used the fact that $\mathcal{V}(t) \in \mathcal{S}(\mathbb{R}^N)$, $\|\mathcal{V}(t)\|_{L^\infty} \lesssim e^{-e_0 t}$ and $|\nabla W| \lesssim |W|$. Thus,

$$\|(\mathcal{L} - \tilde{\mathcal{L}})h\|_{L^{2N/(N+2)}} + \|\nabla[(\mathcal{L} - \tilde{\mathcal{L}})h]\|_{S'(L^2, [t, +\infty))} \lesssim e^{-[\min\{c_0, (k+1^-)e_0\} + (p_c-1)e_0]t}.$$

Therefore, by Lemma 8.1, since $c_0 > e_0$ by hypothesis,

$$\|h\|_{\dot{H}^1} \lesssim e^{-t \min\{[c_0 + (p_c-1)e_0], c_1\}^-}.$$

By iterating this argument, we get (8.9). ■

We now improve the convergence given by Propositions 6.1 and 7.8.

Lemma 8.5. *For $N \geq 6$, if u is a solution to (1.2) satisfying, for all $t \geq 0$,*

$$(8.10) \quad \|u(t) - W\|_{\dot{H}^1} \lesssim e^{-ct} \quad \text{and} \quad E(u_0) = E(W),$$

then there exists a unique $A \in \mathbb{R}$ such that $u = U^A$.

Proof. *Step 1. Linearization around W to improve the decay on time.*

If u is a solution to (1.2), write $u = h + W$. Recall that h is a solution to (3.1). We first show the bound

$$(8.11) \quad \|\nabla(R(h))\|_{S'(L^2, [t, +\infty))} + \|R(h(t))\|_{L^{2N/(N+2)}} \lesssim e^{-cp_c t} \quad \text{for all } t \geq 0.$$

Indeed, by Lemmas 3.12 and 3.9 (ii),

$$\|\nabla(R(h))\|_{S'(L^2, [t, t+1])} \lesssim e^{-cp_c t}.$$

Therefore, the triangle inequality gives

$$\|\nabla(R(h))\|_{S'(L^2, [t, +\infty))} \lesssim e^{-cp_c t}.$$

Now, by (3.16), we have

$$|R(h(t))| \leq |h(t)|^{p_c},$$

so that, by the Sobolev inequality,

$$\|R(h(t))\|_{L^{2N/(N+2)}} \lesssim \|h(t)\|_{L^{2^*}}^{p_c} \lesssim e^{-cp_c t}.$$

Therefore, the bound (8.11) is proved.

We are now under the hypotheses of Lemma 8.1, with $c_0 = c$ and $c_1 = cp_c > c$. The conclusion of this lemma gives

$$\|h(t)\|_{\dot{H}^1} \lesssim e^{-e_0 t} + e^{-cp_c^- t}.$$

If $c > e_0/p_c$, we get

$$\|h(t)\|_{\dot{H}^1} \lesssim e^{-e_0 t},$$

and, by the same argument used to prove (8.11),

$$\|\nabla(R(h))\|_{S'(L^2, [t, +\infty))} + \|R(h(t))\|_{L^{2N/(N+2)}} \lesssim e^{-e_0 p_c t}.$$

Thus, (8.4) gives

$$(8.12) \quad \|h(t) - Ae^{-e_0 t} \mathbf{y}_+\|_{\dot{H}^1} \lesssim e^{-p_c^- e_0 t}.$$

If, however, $c \leq e_0/p_c$, then assumption (8.10) holds with $\frac{1+p_c}{2}c > c$ instead of c . By iteration, we get (8.12).

Step 2. Linearize around $W + \mathcal{V}_k$ to improve higher order convergence to U^A .

For $k \geq 2$ to be chosen later, write $\tilde{h} = h - \mathcal{V}_k$, so that \tilde{h} is a solution to (8.5). Since k is fixed throughout the proof, it will be omitted. By Lemma 8.2, we have

$$\|\nabla(\tilde{R}(\tilde{h}))\|_{S'(L^2, [t, +\infty))} + \|\tilde{R}(\tilde{h}(t))\|_{L^{2N/(N+2)}} \lesssim_k \|\nabla\tilde{h}\|_{S(L^2, [t, +\infty))}^{p_c}.$$

Therefore, by (8.5),

$$\partial_t h + \tilde{\mathcal{L}}_k h = \eta,$$

with

$$(8.13) \quad \|\nabla \eta\|_{S'(L^2, [t, +\infty))} + \|\eta\|_{L^{2N/(N+2)}} \lesssim_k \|\nabla \tilde{h}\|_{S(L^2, [t, +\infty))}^{p_c} + e^{-(k+1)e_0 t}.$$

By (8.12) and the definition of \mathcal{V}_k , we have

$$(8.14) \quad \|\tilde{h}\|_{\dot{H}^1} \lesssim \|h - Ae^{-e_0 t} \mathbf{y}_+\|_{\dot{H}^1} + O(e^{-2e_0 t}) \lesssim_k e^{-p_c^- e_0 t}.$$

By iteration, starting with (8.14), and repeatedly applying Lemmas 8.4 and 8.6, as well as estimate (8.13), we have, for any $k \geq 2$,

$$\|\tilde{h}\|_{\dot{H}^1} \lesssim_k e^{-(k+1^-)e_0 t}.$$

Therefore, choosing $k = l(k_0)$, where k_0 and l are defined in Proposition 4.2, we have, for $N \geq 6$ and $t \geq 0$,

$$\begin{aligned} \|D^\varepsilon(u - W - \mathcal{V}_{l(k_0)})\|_{S(\dot{H}^{1-\varepsilon}, [t, +\infty))} &\lesssim \|\nabla(u - W - \mathcal{V}_{l(k_0)})\|_{S(L^2, [t, +\infty))} \\ &\lesssim e^{-(k_0 + \frac{3}{4})\frac{N-2}{4}e_0 t}. \end{aligned}$$

Hence, by the uniqueness in Proposition 4.2, we get that $u = U^A$. ■

Corollary 8.6. *Let $N \geq 6$. For any $A \neq 0$, there exists $T_A \in \mathbb{R}$ such that*

$$\text{either } U^A(t) = W^+(t + T_A), \text{ if } A > 0, \text{ or } U^A(t) = W^-(t + T_A), \text{ if } A < 0.$$

Proof. Choose T_A such that $|A|e^{-e_0 T_A} = 1$. We have, by (4.8),

$$(8.15) \quad \|U^A(t + T_A) - W \mp e^{-e_0 t} \mathbf{y}_+\|_{\dot{H}^1} \lesssim e^{-2e_0 t}.$$

Note that $U^A(t + T_A)$ satisfies the hypotheses of Lemma 8.5. Thus, there exists $\bar{A} \in \mathbb{R}$ such that $U^A(t + T_A) = U^{\bar{A}}$. But (8.15) implies that $\bar{A} = 1$, if $A > 0$, and $\bar{A} = -1$, if $A < 0$, finishing the proof of the corollary. ■

8.2. Intercritical case

For $0 < s_c < 1$, we study the linearized approximate equation

$$(8.16) \quad \partial_t h + \mathcal{L}h = \epsilon$$

with h and ϵ such that, for $t \geq 0$,

$$(8.17) \quad \|h(t)\|_{H^1} \lesssim e^{-c_0 t} \quad \text{and} \quad \|\langle \nabla \rangle \epsilon\|_{S'(L^2, [t, +\infty))} \lesssim e^{-c_1 t},$$

where $c_1 > c_0 > 0$. We merely state the results in this case, as their proofs are very close to the energy-critical case (in fact, some proofs are easier, since the L^2 norm of the solution is finite).

Lemma 8.7. *Under the assumptions (8.17),*

(i) *if $e_0 \notin [c_0, c_1)$, then*

$$(8.18) \quad \|h(t)\|_{H^1} \lesssim e^{-c_1^{-1}t},$$

(ii) *if $e_0 \in [c_0, c_1)$, then there exists $A \in \mathbb{R}$ such that*

$$(8.19) \quad \|h(t) - Ae^{-e_0t}y_+\|_{H^1} \lesssim e^{-c_1^{-1}t}.$$

Omitting A for simplicity and defining, for every k ,

$$\begin{aligned} \tilde{\mathcal{L}}_k &= \begin{pmatrix} 0 & 1 - \Delta \\ -(1 - \Delta) & 0 \end{pmatrix} + \frac{(p+1)}{2} |Q + \mathcal{V}_k|^{p-1} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ &\quad + \frac{(p-1)}{2} |Q + \mathcal{V}_k|^{p-3} \begin{pmatrix} \text{Im}(Q + \mathcal{V}_k)^2 & -\text{Re}(Q + \mathcal{V}_k)^2 \\ -\text{Re}(Q + \mathcal{V}_k)^2 & -\text{Im}(Q + \mathcal{V}_k)^2 \end{pmatrix}, \\ \tilde{K}_k(h) &= \frac{(p+1)}{2} |Q + \mathcal{V}_k|^{p-1}h + \frac{(p-1)}{2} |Q + \mathcal{V}_k|^{p-3} (Q + \mathcal{V}_k)^2 \bar{h}, \end{aligned}$$

and

$$\tilde{R}_k(h) = |Q + \mathcal{V}_k|^{p-1} (Q + \mathcal{V}_k) J((Q + \mathcal{V}_k)^{-1}h),$$

where

$$J(z) = |1 + z|^{p-1} (1 + z) - 1 - \frac{(p+1)}{2} z - \frac{(p-1)}{2} \bar{z},$$

we have that if $u = e^{it}(Q + \mathcal{V}_k + h)$ is a solution to the NLS equation (1.1), then h satisfies

$$(8.20) \quad \partial_t h + \tilde{\mathcal{L}}_k h = i \tilde{R}_k(h) + \epsilon_k,$$

or in the form of a Schrödinger equation,

$$i \partial_t h + \Delta h - h + \tilde{K}_k h = -\tilde{R}_k(h) + i \epsilon_k,$$

where ϵ_k are $O(e^{-(k+1)e_0t})$ in $\mathcal{S}(\mathbb{R}^N)$. By the construction of \mathcal{V}_k , we have, for all $t \geq 0$,

$$|\mathcal{V}_k(t)| \lesssim e^{-e_0t} |Q| \quad \text{and} \quad |\nabla \mathcal{V}_k(t)| \lesssim e^{-e_0t} |\nabla Q| \lesssim e^{-e_0t} |Q|.$$

Therefore, as in the energy-critical case, we have the following results.

Lemma 8.8. *Let $p > 1$, $k \geq 1$ and I be a bounded time interval, and consider $f \in S(L^2, I)$ such that $\nabla f \in S(L^2, I)$. There exists $\alpha > 0$ such that the following estimates hold.*

For all $p > 1$:

(i) $\|\langle \nabla \rangle \tilde{K}_k(f)\|_{S'(L^2, I)} \lesssim |I|^\alpha \|\langle \nabla \rangle f\|_{S(L^2, I)}.$

For $p > 2$:

(ii) $\|\langle \nabla \rangle \tilde{R}_k(f)\|_{S'(L^2, I)} \lesssim \|\langle \nabla \rangle f\|_{S(L^2, I)} (|I|^\alpha \|\langle \nabla \rangle f\|_{S(L^2, I)} + \|\langle \nabla \rangle f\|_{S(L^2, I)}^{p-1}).$

For $1 < p \leq 2$:

(iii) $\|\langle \nabla \rangle \tilde{R}_k(f)\|_{S'(L^2, I)} \lesssim (1 + |I|^\alpha) \|\langle \nabla \rangle f\|_{S(L^2, I)}^p.$

Lemma 8.9. *Let h be a solution to (8.20). If, for some $c > 0$ and for any $t \geq 0$,*

$$\|h(t)\|_{H^1} \lesssim e^{-ct},$$

then

$$(8.21) \quad \|\langle \nabla \rangle h\|_{S(L^2, [t, +\infty))} \lesssim e^{-\min\{c, (k+1^-)e_0\}t}.$$

Lemma 8.10. *Let h be a solution to*

$$(8.22) \quad \partial_t h + \tilde{\mathcal{L}}_k h = \epsilon,$$

with h and ϵ such that, for $t \geq 0$,

$$(8.23) \quad \|h(t)\|_{H^1} \lesssim e^{-c_0 t} \quad \text{and} \quad \|\langle \nabla \rangle \epsilon\|_{S'(L^2, [t, +\infty))} \lesssim e^{-c_1 t},$$

where $(k + 1)e_0 > c_1 > c_0 > e_0$. Then,

$$(8.24) \quad \|h(t)\|_{H^1} \lesssim e^{-c_1^- t}.$$

Lemma 8.11. *If u is a solution to (1.1) satisfying*

$$(8.25) \quad \|u(t) - e^{it} Q\|_{H^1} \lesssim e^{-ct}, \quad M(u_0) = M(Q), \quad E(u_0) = E(Q),$$

then there exists a unique $A \in \mathbb{R}$ such that $u = U^A$.

Corollary 8.12. *Let $1 + 4/N < p < 2^* - 1$. For any $A \neq 0$, there exists $T_A \in \mathbb{R}$ such that*

$$\text{either } U^A(t) = Q^+(t + T_A), \text{ if } A > 0, \text{ or } U^A(t) = Q^-(t + T_A), \text{ if } A < 0.$$

9. Closure of the main theorems

Having proved Propositions 6.1 and 7.8, and Lemma 8.5, we can proceed as in [12].

Proof of Theorem 1.5. Recall the notation $\mathcal{Y}_1 = \text{Re } \mathcal{Y}_+ = \text{Re } \mathcal{Y}_-$. We claim that $(W, \mathcal{Y}_1)_{\dot{H}^1} \neq 0$. If not, since W solves the equation $\Delta W = -W^{p_c}$, we would have

$$B(W, \mathcal{Y}_\pm) = \frac{1}{2} \int \nabla W \cdot \nabla \mathcal{Y}_1 - \frac{p_c}{2} \int W^{p_c} \mathcal{Y}_1 = \frac{p_c}{2} \int \Delta W \mathcal{Y}_1 = 0,$$

so that $W \in G^\perp$. But, by Lemma 3.5, Φ is nonnegative (in fact, it is coercive) on G^\perp , which contradicts (3.13).

Replacing \mathcal{Y}_\pm , if necessary, we may assume

$$(W, \mathcal{Y}_1)_{\dot{H}^1} > 0.$$

Defining

$$W^+ := U^1 \quad \text{and} \quad W^- := U^{-1},$$

we claim that the conclusions of Theorem 1.5 hold.

By the strong convergence $U^A(t) \rightarrow W$ in \dot{H}^1 and energy conservation, we conclude $E(W^\pm) = E(W)$. Moreover, by (4.7),

$$\|U^A(t)\|_{\dot{H}^1}^2 = \|W\|_{\dot{H}^1}^2 + 2A e^{-2e_0 t} (W, \mathcal{Y}_1)_{\dot{H}^1} + O(e^{-3e_0 t}),$$

which shows that $\|U^A(t)\|_{\dot{H}^1} - \|W\|_{\dot{H}^1}$ has the same sign as A , for large t . By uniqueness and continuity of the flow, this sign must remain the same for every t in the intervals of existence of W^\pm . By Proposition 7.4, W^- is defined on \mathbb{R} , and by Proposition 7.8, W^- scatters backward in time.

We now show that U^A has finite mass for $N \geq 5$. As in the proof of Proposition 7.4, let ϕ be a smooth, positive, radial cutoff of the set $\{|x| \leq 1\}$. Define, for $R > 0$ and large t ,

$$F_R(t) = \int |U^A(x, t)|^2 \phi\left(\frac{x}{R}\right) dx.$$

Since U^A is a solution to (1.2), by Lemma A.2 and Hardy’s inequality, we have

$$|F'_R(t)| \lesssim \|U^A(t) - W\|_{\dot{H}^1} \left(\int \frac{1}{|x|^2} |U^A(t)|^2 \right)^{1/2} \lesssim \|U^A(t) - W\|_{\dot{H}^1} \|U^A(t)\|_{\dot{H}^1} \lesssim e^{-e_0 t}.$$

Hence, integrating from a large t to $+\infty$, we get

$$\left| F_R(t) - \int |W|^2 \phi\left(\frac{x}{R}\right) dx \right| \lesssim e^{-e_0 t}.$$

Recalling that $W \in L^2(\mathbb{R}^N)$ if $N \geq 5$, we can make $R \rightarrow +\infty$ to obtain $M(U^A) = M(W) < +\infty$. In particular, $W^\pm \in L^2(\mathbb{R}^N)$ and, by Proposition 6.1, W^+ blows up in finite negative time. This finishes the proof of Theorem 1.5. ■

Proof of Theorem 1.6. The case $\|u_0\|_{\dot{H}^1} < \|W\|_{\dot{H}^1}$ follows immediately from Proposition 7.8, Lemma 8.5 and Corollary 8.6. The case $\|u_0\|_{\dot{H}^1} = \|W\|_{\dot{H}^1}$ is a consequence of the variational characterization of W . Finally, $\|u_0\|_{\dot{H}^1} > \|W\|_{\dot{H}^1}$ follows from Proposition 6.1, Lemma 8.5 and Corollary 8.6. ■

Proof of Theorem 1.2. Recall the notation $\mathcal{Y}_1 = \text{Re } \mathcal{Y}_+ = \text{Re } \mathcal{Y}_-$. We claim that $(Q, \mathcal{Y}_1)_{H^1} \neq 0$. If not, since Q solves the equation $Q - \Delta Q = -Q^p$, we would have

$$B(Q, \mathcal{Y}_\pm) = \frac{1}{2} \int Q \mathcal{Y}_1 + \frac{1}{2} \int \nabla Q \cdot \nabla \mathcal{Y}_1 - \frac{p}{2} \int Q^{p_c} \mathcal{Y}_1 = \frac{p+1}{2} (Q, \mathcal{Y}_1)_{H^1} = 0,$$

so that $Q \in \tilde{G}^\perp$. But, by Lemma 3.5, Φ is nonnegative (in fact, it is coercive) on \tilde{G}^\perp , which contradicts (3.11).

Replacing \mathcal{Y}_\pm , if necessary, we may assume

$$(Q, \mathcal{Y}_1)_{H^1} > 0.$$

Defining

$$Q^+ := U^1 \quad \text{and} \quad Q^- := U^{-1},$$

we claim that the conclusions of Theorem 1.2 hold.

By the strong convergence $e^{-it}U^A(t) \rightarrow Q$ in H^1 and energy conservation, we conclude $M(Q^\pm) = M(Q)$ and $E(Q^\pm) = E(Q)$. Moreover, by (4.13),

$$\|U^A(t)\|_{H^1}^2 = \|Q\|_{H^1}^2 + 2A e^{-2e_0t} (Q, \mathcal{Y}_1)_{H^1} + O(e^{-2p_c e_0t}),$$

which shows that $\|U^A(t)\|_{H^1} - \|Q\|_{H^1}$ has the same sign as A , for large t . By uniqueness and continuity of the flow, this sign must remain the same for every t in the intervals of existence of Q^\pm . By Proposition 7.11, Q^- is defined on \mathbb{R} , and by Proposition 7.14, Q^- scatters backward in time. Finally, by Proposition 6.3, Q^+ blows up in finite negative time. This finishes the proof of Theorem 1.2. ■

Proof of Theorem 1.3. The case $\|\nabla u_0\|_{L^2} < \|\nabla Q\|_{L^2}$ follows immediately from Proposition 7.14, Lemma 8.11 and Corollary 8.12. The case $\|\nabla u_0\|_{L^2} = \|\nabla Q\|_{L^2}$ is a consequence of the variational characterization of Q . Finally, $\|\nabla u_0\|_{L^2} > \|\nabla Q\|_{L^2}$ follows from Proposition 6.3, Lemma 8.11 and Corollary 8.12. ■

A. Appendix

A.1. Proof of modulation results

Proof of Lemma 5.1. The idea of the proof is already well-known (see, for instance, Section 7.1 of [13] for the 3d cubic NLS equation, Section 3 of [13] for the energy-critical NLS equation, in the radial case, for dimensions $N = 3, 4$ and 5 , and Section 2 of [33] in the context of the Korteweg–de Vries equation), and we extend the proofs here to any dimension, any $0 < s_c \leq 1$, not assuming radially of the solution. The case $s_c = 1$ is Lemma 5.1, and the case $0 < s_c < 1$ is Lemma 5.3. We first show Lemma 5.1 when u is close to W . Define the functional $J = (J_0, \dots, J_{N+1})$ on $\mathbb{R}^N \times \mathbb{R}_+ \times \mathbb{R} \times \dot{H}^1$ as

$$\begin{aligned} J_0 &: (\theta, x, \lambda, u) \mapsto (f_{[x,\lambda,\theta]}, iW)_{\dot{H}^1}, \\ J_k &: (\theta, x, \lambda, u) \mapsto (f_{[x,\lambda,\theta]}, \partial_k W)_{\dot{H}^1}, \quad 1 \leq k \leq N, \\ J_{N+1} &: (\theta, x, \lambda, u) \mapsto (f_{[x,\lambda,\theta]}, \Lambda W)_{\dot{H}^1}. \end{aligned}$$

By direct calculation, one can check that

$$\det \left(\frac{\partial J}{\partial(\theta, x, \lambda)} \right) = \left(\int |W|^2 \right) \left(\prod_k \int |\partial_k W|^2 \right) \left(- \int |\Lambda W|^2 \right) \neq 0,$$

and that $J(0, 1, 0, W) = 0$. Hence, by the implicit function theorem, there exist ϵ_0 and η_0 such that, if $f \in \dot{H}^1$ and $\|f - W\|_{\dot{H}^1} < \epsilon_0$, then there exists a unique n -tuple (x, λ, θ) such that

$$|x| + |\lambda| + |\theta - 1| \leq \eta_0 \quad \text{and} \quad J(\theta, x, \lambda, f) = 0.$$

Now, if u is as in the lemma, by the variational characterization of W , if $d(u)$ is small, then there exists $(x_0, \lambda_0, \theta_0)$ such that $u_{[x_0,\lambda_0,\theta_0]} = W + f$, with $\|f\|_{\dot{H}^1} \leq \epsilon(d(f))$. We are thus back to the preceding case. Existence, local uniqueness and regularity follow again from the implicit function theorem. ■

Proof of Lemma 5.2. For a fixed t , write $v = u_{[x(t), \lambda(t), \theta(t)]}(t) - W = \alpha(t)W + h(t)$ as in (5.1). Since $h \in G^\perp$, we have

$$(A.1) \quad \|v\|_{\dot{H}^1}^2 = \alpha^2 \|W\|_{\dot{H}^1}^2 + \|h\|_{\dot{H}^1}^2.$$

Since $h \in G^\perp$, and W satisfies the equation $\Delta W + W^{p_c} = 0$, we have

$$B(W, h) = \frac{1}{2} \int \nabla W \cdot \nabla h_1 + \frac{p}{2} \int \Delta W h_1 = 0.$$

Therefore, $\Phi(v) = \Phi(\alpha W + h) = \alpha^2 \Phi(W) + \Phi(h)$. Recalling that W is a critical point for the energy functional E , we have $E(W + v) = E(W) + \Phi(v) + O(\|v\|_{\dot{H}^1}^3)$. Since $E(W + v) = E(W)$, and by the coercivity given by Lemma 3.5, we have $\Phi(h) \approx \|h\|_{\dot{H}^1}^2$. Thus, we have

$$(A.2) \quad |\alpha^2 \Phi(W) + \Phi(h)| = O(\|v\|_{\dot{H}^1}^3)$$

Since $\|v\|_{\dot{H}^1}$ is small when $d(u)$ is small, estimates (A.1) and (A.2) give $|\alpha| \approx \|h\|_{\dot{H}^1} \approx \|v\|_{\dot{H}^1}$. Finally, since

$$d(u) = \left| \|W + v\|_{\dot{H}^1}^2 - \|W\|_{\dot{H}^1}^2 \right| = \left| \|v\|_{\dot{H}^1}^2 + 2\alpha \|W\|_{\dot{H}^1}^2 \right|,$$

we have $d(u) \approx |\alpha|$, and (5.2) is proved.

It remains to prove (5.3). Consider the variables y and s given by

$$y = \frac{x}{\mu(t)} \quad \text{and} \quad dt = \frac{1}{\mu^2(t)} ds.$$

In view of (5.2) and the decomposition (5.1), we can rewrite (1.2) as

$$(A.3) \quad i \partial_s h + \Delta h - i \alpha_s W + \theta_s W - i x_s \cdot \nabla W + i \frac{\lambda_s}{\lambda} \Lambda W = O(\epsilon(s)) \quad \text{in } \dot{H}^1,$$

where $\epsilon(s) := d(u(t(s))) (d(u(t(s))) + |\theta_s| + |x_s| + |\lambda_s/\lambda|)$. Since $h \in G^\perp$, projecting (A.3) in \dot{H}^1 onto W , iW , ∇W and ΛW and integrating by parts (possible due to a standard regularization argument) yields

$$|\alpha_s| + |\theta_s| + |x_s| + \left| \frac{\lambda_s}{\lambda} \right| = O(d + \epsilon(s)),$$

which is enough to conclude (5.3) and to finish the proof of Lemma 5.2. ■

Proof of Lemma 5.3. The proof is analogous to that of Lemma 5.1, and is omitted. ■

Proof of Lemma 5.4. The orthogonality condition (3.9) implies $B(Q, h) = 0$. Since $E(u) = E(Q)$, and Q is a critical point for E , we have

$$\alpha^2 \Phi(Q) + \Phi(h) = \Phi(\alpha Q + h) = O(|\alpha|^3 + \|h\|_{\dot{H}^1}^3).$$

By coercivity, $\Phi(h) \approx \|h\|_{H^1}^2$, and hence, $|\alpha| \approx \|h\|_{H^1}$. The relation $M(u) = M(Q)$ gives

$$(A.4) \quad \left| \alpha \int Q^2 + \int Qh_1 \right| = \frac{1}{2} \int |\alpha Q + h|^2 = O(|\alpha|^2),$$

and thus,

$$|\alpha| \approx \left| \int Qh_1 \right|.$$

Now, using (3.9),

$$d(u) = \left| \int |\nabla u|^2 - \int |\nabla Q|^2 \right| = \left| 2\left(\alpha \int |\nabla Q|^2 - \int Qh_1 \right) + O(|\alpha|^2) \right|,$$

which, together with (A.4), gives

$$d(u) = \left| 2\alpha \left(\int |\nabla Q|^2 - \int Q^2 \right) + O(|\alpha|^2) \right|.$$

Since, by the Pohozaev identities (1.4), $\|\nabla Q\|_{L^2} \neq \|Q\|_{L^2}$ for any N and any p in the intercritical range, we conclude $d(u) \approx |\alpha|$, and hence, (5.6) holds. The rest of the proof goes along the same lines of the proof of Lemma 5.2 (without the need of self-similar variables), and is omitted. ■

A.2. Convergence to the standing wave above the ground state

Proof of Lemma 6.2 (Critical case).

Step 1. A general bound on A_R (recall (6.5)).

By the definition of ϕ_R , we have the bounds $4\partial_r^2\phi_R \leq 8$, $|\Delta^2\phi_R| \lesssim 1$ and $|\Delta^2\phi_R(r)| \lesssim 1/r^2$. Therefore,

$$(A.5) \quad A_R(u(t)) \lesssim \int_{|x| \geq R} |u(x, t)|^{2^*} + \frac{1}{R^2} |u(x, t)|^2 dx.$$

We now recall Strauss' lemma, [37], and make use of the decay given by radiality in H^1 .

Lemma A.1 ([37]). *There is a constant $C > 0$ such that, for any radial function f in $H^1(\mathbb{R}^N)$ and any $R > 0$,*

$$\|f\|_{L^\infty_{\{|x| \geq R\}}} \leq \frac{C}{R^{(N-1)/2}} \|f\|_{L^2}^{1/2} \|\nabla f\|_{L^2}^{1/2}.$$

We can now bound

$$\int_{|x| \geq R} |u(x, t)|^{2^*} dx \leq \|u(t)\|_{L^\infty_{\{|x| \geq R\}}}^{\frac{4}{N-2}} \|u_0\|_{L^2}^2 \leq \frac{C}{R^{\frac{2N-2}{N-2}}} \|\nabla u(t)\|_{L^2}^{\frac{2}{N-2}} \|u_0\|_{L^2}^{\frac{2N-2}{N-2}},$$

to obtain

$$(A.6) \quad A_R(u(t)) \leq C_0 \left[\frac{1}{R^2} + \frac{1}{R^{\frac{2N-2}{N-2}}} (d(t) + \|W\|_{\dot{H}^1})^{\frac{1}{N-2}} \right],$$

where C_0 depends only on $M(u_0)$.

Step 2. A bound on A_R when $d(t)$ is small.

Taking a small δ_1 , write the decomposition (5.1) as $u_{[x(t),\lambda(t),\theta(t)]} = W + v$, with $\|v\|_{\dot{H}^1} \lesssim d(t)$, by Lemma 5.2. We first prove

$$(A.7) \quad \lambda_- := \inf\{\lambda(t), t \geq 0, d(t) \leq \delta_1\} > 0.$$

Indeed, by mass conservation,

$$\|u_0\|_{L^2} \geq \int_{|x| \leq \lambda(t)} |u(t)|^2 = \frac{1}{\lambda^2(t)} \left(\int_{|x| \leq 1} W^2(x) dx - C d^2(t) \right).$$

If $d(t) \leq \delta_1$ and δ_1 is small enough, then (A.7) holds. We now give an estimate on A_R when $d(t)$ is small. Since W is a static solution to (1.2), $\frac{d}{dt} \int \phi_R |W|^2 = 0$, so that $A_R(W) = 0$, for all $R > 0$, by (6.4). If we assume $R \geq 1$, by a change of variables, the Hölder, Hardy and Sobolev inequalities, we can write (6.5) as

$$(A.8) \quad |A_R(u(t))| = |A_{R\lambda(t)}(W + v)| = |A_{R\lambda(t)}(W + v) - A_{R\lambda(t)}(W)| \\ \leq C \left[\int_{|x| \geq R\lambda(t)} |\nabla v|^2 + |\nabla W \cdot \nabla v| + W^{2^*-1}|v| + |v|^{2^*} + \frac{1}{(R\lambda(t))^2} (W|v| + |v|^2) \right] \\ \leq C \left[\|v\|_{\dot{H}^1}^2 + \frac{1}{(R\lambda(t))^{\frac{N-2}{2}}} \|v\|_{\dot{H}^1} + \|v\|_{\dot{H}^1}^{2^*} + \frac{1}{(R\lambda(t))^{\frac{N+2}{2}}} \|v\|_{\dot{H}^1} + \frac{1}{\lambda^2(t)} \|v\|_{\dot{H}^1}^2 \right] \\ \leq C_1 \left[d^2(t) + \frac{1}{R^{(N-2)/2}} d(t) \right],$$

where we used the fact that $\|\nabla W\|_{L^2_{\{|x| \geq r\}}} \approx \|W\|_{L^{2^*}_{\{|x| \geq r\}}} \approx 1/r^{(N-2)/2}$, which can be verified by explicit computation. Note that the constant C_1 depends only on λ_- , which in turn depends only on $M(u_0)$.

Step 3. Bounds on A_R prove bounds on $d(t)$.

We now claim the bound

$$(A.9) \quad A_R(t) \leq \frac{8}{N-2} d(t).$$

This follows from the bound (A.8), if $d(t) \leq \delta_1$ and $R \geq R_1$, where R_1 is a large constant depending only on $M(u_0)$. Now, if $d(t) > \delta_1$, consider the function

$$\varphi_R(\delta) = \frac{C_0}{R^2} + \frac{C_0}{R^{(2N-2)/(N-2)}} (\delta + \|W\|_{\dot{H}^1})^{1/(N-2)} - \frac{8}{N-2} \delta.$$

By direct computation, we see that $\varphi_R''(\delta) < 0$ for any $\delta > 0$. We can choose a large $R_2 \geq R_1$ (depending again only on $M(u_0)$) such that $\varphi_{R_2}(\delta_1) \leq 0$ and $\varphi'_{R_2}(\delta_1) \leq 0$, so that $\varphi_{R_2}(\delta) \leq 0$ for all $\delta \geq \delta_1$. The bound (A.9) is now proved.

The bound (A.9), together with (6.4), gives, for $R \geq R_2$ and any $t \geq 0$,

$$(A.10) \quad F_R''(t) \leq -\frac{8}{N-2} d(t) < 0.$$

Note that we must have $F_R'(t) > 0$, for all $t \geq 0$, as (A.10) would otherwise contradict the positivity of F_R , thus, proving (6.6).

We now make use of the following claim, in the spirit of Lemma 2.1 in [3] and Claim 5.4 in [13].

Claim A.2. *Let $\phi \in C^1(\mathbb{R}^N)$ and $f \in H^1(\mathbb{R}^N)$. Assume that $\int |\nabla\phi|^2 |f|^2 < +\infty$ and $E(f) = E(W)$. Then*

$$\left(\operatorname{Im} \int \nabla\phi \cdot \nabla f \bar{f} \right)^2 \lesssim d^2(f) \int |\nabla\phi|^2 |f|^2.$$

By Claim A.2 and the fact that $F'_R(t) > 0$ and $F''(t) < 0$, we can write

$$\frac{F'_R(t)}{\sqrt{F_R(t)}} \lesssim -F''_R(t),$$

so that

$$\int_t^{+\infty} d(s) ds \lesssim e^{-ct},$$

which proves (6.7) and finishes the proof of Lemma 6.2. ■

Proof of Claim A.2. Let $\delta(f) = \int |\nabla W|^2 - \int |\nabla f|^2$ and $\lambda \in \mathbb{R}$. By Sobolev’s inequality,

$$\|\nabla(e^{i\lambda\phi} f)\|_{L^2} \geq \frac{\|\nabla W\|_{L^2}}{\|W\|_{L^{2^*}}} \|f\|_{L^{2^*}}.$$

Squaring the last inequality and expanding the term $\|\nabla(e^{i\lambda\phi} f)\|_{L^2}$, we get

$$\lambda^2 \int |\nabla\phi|^2 |\nabla f|^2 + 2\lambda \operatorname{Im} \int (\nabla\phi \cdot \nabla f) \bar{f} + \int |\nabla f|^2 - \frac{\|\nabla W\|_{L^2}^2}{\|W\|_{L^{2^*}}^2} \|f\|_{L^{2^*}}^2 \geq 0.$$

The discriminant of this quadratic form must be non-positive, thus,

$$\left(\operatorname{Im} \int (\nabla\phi \cdot \nabla f) \bar{f} \right)^2 \leq \left(\int |\nabla f|^2 - \frac{\|\nabla W\|_{L^2}^2}{\|W\|_{L^{2^*}}^2} \|f\|_{L^{2^*}}^2 \right) \left(\int |\nabla\phi|^2 |\nabla f|^2 \right).$$

Since

$$\int |\nabla f|^2 = \int |\nabla W|^2 - \delta(f),$$

by $E(f) = E(W)$, we have,

$$0 < \int |f|^{2^*} = \int |W|^{2^*} - \frac{N}{N-2} \delta(f).$$

Therefore,

$$\begin{aligned} \int |\nabla f|^2 - \frac{\|\nabla W\|_{L^2}^2}{\|W\|_{L^{2^*}}^2} \|f\|_{L^{2^*}}^2 &= \int |\nabla W|^2 - \delta(f) - \frac{\|\nabla W\|_{L^2}^2}{\|W\|_{L^{2^*}}^2} \left(\int |W|^{2^*} - \frac{N}{N-2} \delta(f) \right)^{\frac{N-2}{N}} \\ &= \int |\nabla W|^2 - \delta(f) - \frac{\|\nabla W\|_{L^2}^2}{\|W\|_{L^{2^*}}^2} \left(\|W\|_{L^{2^*}}^2 - \|W\|_{L^{2^*}}^{-\frac{4}{N-2}} \delta(f) + O(\delta(f)^2) \right) = O(\delta(f)^2), \end{aligned}$$

and Claim A.2 is proved. ■

Proof of Proposition 6.1. We first prove that

$$(A.11) \quad \lim_{t \rightarrow +\infty} d(t) = 0.$$

Indeed, by Lemma 6.2, there exists $t_n \rightarrow +\infty$ such that $d(t_n) \rightarrow 0$. Assume, by contradiction, that there exists a sequence $t'_n > t_n$ such that

$$(A.12) \quad d(t'_n) = \delta_0 \quad \text{and} \quad 0 < d(t) < \delta_0 \quad \forall t \in (t_n, t'_n),$$

where δ_0 is given by Lemmas 5.1 and 5.2. Recall the decomposition (5.1):

$$u_{[\lambda(t), \theta(t)]}(t) = (1 + \alpha(t))W + h(t), \quad \text{with } h \in G^\perp.$$

By taking subsequences, if necessary, we can assume

$$\lim \lambda(t_n) = \lambda_\infty \in (0, +\infty].$$

We now prove that $\lambda_\infty < +\infty$.

If $\lambda_\infty = +\infty$, as $u_{[\lambda(t_n), \theta(t_n)]}$ converges to W in \dot{H}^1 , we have, for any $C > 0$,

$$\int_{|x| \geq C} |u(t_n)|^{2^*} \rightarrow 0.$$

For any $\epsilon > 0$ we have, by Hölder's inequality,

$$|F_R(t_n)| \lesssim \epsilon \|u(t_n)\|_{\dot{H}^1} + \int_{|x| \geq C_\epsilon} |u(t_n)|^{2^*},$$

so that

$$\lim F_R(t_n) = 0.$$

However, by (6.6), $F'_R(t) > 0$. This implies $F_R(t) < 0$ for all $t \geq 0$, contradicting the fact that F_R is positive. Therefore, $\lambda(t_n)$ must be bounded.

Now, by Lemma 5.2, we have $|\lambda'(t)/\lambda^3(t)| \lesssim d(t)$. This implies, if $t \in (t_n, t'_n)$,

$$\left| \frac{1}{\lambda^2(t)} - \frac{1}{\lambda^2(t_n)} \right| \lesssim e^{-ct_n}.$$

Therefore, $\lambda(t) \leq 2\lambda_\infty$ on $\cup_n (t_n, t'_n)$, for large t . Turning to the bound on α' in Lemma 5.2,

$$|\alpha'(t)| \lesssim \lambda^2(t) d(t) \lesssim d(t).$$

This implies $|\alpha(t_n) - \alpha(t'_n)| \rightarrow 0$. Moreover, again by Lemma 5.2, $|\alpha(t)| \approx d(t)$, which contradicts (A.12) and proves (A.11).

To finish the proof of Proposition 6.1, we must refine the estimate on $d(t)$. Since $d(t) \rightarrow 0$ as $t \rightarrow +\infty$, the decomposition (5.1) is well-defined for all large times. Therefore, by (A.11) and (A.7), we have

$$\lim_{t \rightarrow +\infty} \lambda(t) = \lambda_\infty \in (0, +\infty), \quad \lim_{t \rightarrow +\infty} \alpha(t) = \lim_{t \rightarrow +\infty} d(t) = 0,$$

and

$$\|h(t)\|_{\dot{H}^1} \approx |\alpha(t)| = \int_t^{+\infty} |\alpha'(s)| ds \lesssim \int_t^{+\infty} \lambda^2(s) d(s) ds \lesssim e^{-ct}.$$

Furthermore, the bound $|\theta'(t)| \lesssim \lambda^2(t)d(t)$ implies that there exists θ_∞ such that

$$\lim_{t \rightarrow +\infty} |\theta(t) - \theta_\infty| = 0.$$

Therefore, (6.2) is proven.

It remains to prove the finite-time blow-up. This is a corollary of Lemma 6.2, applied to the time-reversed solution $v(t) := \bar{u}(-t)$. If u is defined on \mathbb{R} , by (6.6), we have

$$\operatorname{Im} \int \nabla \phi \cdot \nabla u_0 \bar{u}_0 > 0, \quad \text{and} \quad \operatorname{Im} \int \nabla \phi \cdot \nabla v_0 \bar{v}_0 > 0,$$

which clearly contradicts the fact that

$$\operatorname{Im} \int \nabla \phi \cdot \nabla u_0 \bar{u}_0 = -\operatorname{Im} \int \nabla \phi \cdot \nabla v_0 \bar{v}_0.$$

This finishes the proof of Proposition 6.1. ■

Proof of Proposition 6.3 (intercritical case). We divide the proof in two cases: the finite-variance case, and the radial case. Using the same argument of the finite-time blow-up as in the energy-critical case, and in view of Lemmas 5.5, A.3 and A.5 in the next subsections, Proposition 6.3 follows. ■

A.2.1. Finite-variance solutions.

Lemma A.3. *Let u be a solution to (1.1) defined on $[0, +\infty)$ and satisfying (6.8) and $\| |x| u_0 \|_{L^2} < +\infty$. Then, for all t in the interval of existence of u ,*

$$(A.13) \quad \operatorname{Im} \int x \cdot \nabla u(t) \overline{u(t)} > 0,$$

and there exists $c > 0$ such that

$$(A.14) \quad \int_t^{+\infty} d(s) ds \lesssim e^{-ct}, \quad \forall t \geq 0.$$

Proof. Let $F(t) = \int |x|^2 |u(x, t)|^2 dx$. Then, by the virial identities, we have, for all $t \geq 0$,

$$F'(t) = 4 \operatorname{Im} \int x \cdot \nabla u(t) \overline{u(t)}.$$

Note that, by Cauchy–Schwarz, $F'(t)$ is well-defined. Furthermore, since $E(u) = E(Q)$,

$$F''(t) = -[2N(p - 1) - 8] \left(\int |\nabla u|^2 - \int |u|^{p+1} \right) = -[2N(p - 1) - 8] d(u(t)).$$

Now, if (A.13) does not hold, there exists t_1 such that $F'(t_1) \leq 0$. Since $F'' \leq 0$, for any $t_0 > t_1$,

$$F'(t) \leq F'(t_0) < 0, \quad \forall t \geq t_0.$$

This implies that $F(t) < 0$ for large t , contradicting the definition of F .

We now claim that

$$(A.15) \quad [F'(t)]^2 \lesssim F(t) [F''(t)]^2,$$

which is a consequence of the following claim, which is proved similarly to Claim A.2.

Claim A.4. Let $\phi \in C^1(\mathbb{R}^N)$ and let $f \in H^1(\mathbb{R}^N)$. Assume that $\int |\nabla\phi|^2 |f|^2 < +\infty$, $M(f) = M(Q)$ and $E(f) = E(Q)$. Then

$$\left(\operatorname{Im} \int \nabla\phi \cdot \nabla f \bar{f} \right)^2 \lesssim d^2(f) \int |\nabla\phi|^2 |f|^2.$$

Taking $\phi(x) = |x|^2$ yields (A.15).

Since $F'(t) > 0$ and $F''(t) < 0$ for all $t \geq 0$, equation (A.15) can be rewritten as

$$\frac{F'(t)}{\sqrt{F(t)}} \lesssim -F''(t).$$

Integrating from 0 to $t \geq 0$,

$$\sqrt{F(t)} - \sqrt{F(0)} \lesssim -(F'(t) - F'(0)) \leq F'(0).$$

From (A.15), we deduce

$$F'(t) \lesssim -(\sqrt{F(0)} + F'(0)) F''(t) \lesssim -F''(t),$$

which shows

$$F'(t) \lesssim e^{-ct}.$$

Finally,

$$F'(t) = - \int_t^{+\infty} F''(s) ds = 4 \int_t^{+\infty} d(s) ds,$$

producing (A.14), which proves Lemma A.3. ■

A.2.2. Radial solutions. We now work with a truncated variance. Consider a radial function $\phi \in C_0^\infty(\mathbb{R}^N)$ such that

$$\phi(r) \geq 0 \quad \forall r \geq 0, \quad \phi(r) = \begin{cases} r^2, & r \leq 1, \\ 0, & r \geq 3, \end{cases} \quad \text{and} \quad \frac{d^2\phi}{dr^2}(r) \leq 2, \quad r \geq 0.$$

Define $\phi_R(x) = R^2\phi(x/R)$ and

$$F_R(t) = \int \phi_R |u(t)|^2.$$

By the virial identities, if $M(u_0) = M(Q)$ and $E(u_0) = E(Q)$, we have

$$(A.16) \quad F'_R(t) = 2 \operatorname{Im} \int \nabla\phi_R \cdot \nabla u \bar{u},$$

$$(A.17) \quad F''_R(t) = -[2N(p-1) - 8]d(t) + A_R(t),$$

where

$$(A.18) \quad \begin{aligned} A_R(u(t)) &= \int_{|x| \geq R} |\nabla u(t)|^2 (4\partial_r^2\phi_R - 8) \\ &+ \frac{2(p-1)}{(p+1)} \int_{|x| \geq R} |u|^{p+1} (2N - \Delta\phi_R) - \int_{|x| \geq R} |u|^2 \Delta^2\phi_R. \end{aligned}$$

The following lemma holds.

Lemma A.5. *Let u be radial a solution to (1.1) defined on $[0, +\infty)$ and satisfying (6.8). Then, there exists a constant $R_0 > 0$ such that, for all t in the interval of existence of u and all $R \geq R_0$,*

$$(A.19) \quad F'_R(t) > 0,$$

and there exists $c > 0$ such that

$$(A.20) \quad \int_t^{+\infty} d(s) ds \lesssim e^{-ct}, \quad \forall t \geq 0.$$

Moreover, u_0 has finite variance.

Proof. The proof of Lemma A.5 is essentially the same as in the energy-critical case, and is omitted, except for the finite-variance part.

By hypothesis, there is a sequence $t_n \rightarrow +\infty$ such that $d(t_n) \rightarrow 0$. By (5.4), extracting a subsequence, if necessary, we have $u_n \rightarrow e^{i\theta_0} Q$ in H^1 for some $\theta_0 \in \mathbb{R}$. Since F_R is increasing by (A.19), we have

$$\int \phi_R |u_0|^2 = F_R(0) \leq \int \phi_R Q^2.$$

Thus, we can make $R \rightarrow +\infty$, which proves the finite variance of u_0 . ■

Proof of Lemma A.4. The proof is analogous to that of Lemma A.2 and, thus, omitted. ■

A.3. Convergence to the standing wave below the ground state

As in the proof of Proposition 6.1, we need to show that

$$d(t) = \|W\|_{\dot{H}^1} - \|u(t)\|_{\dot{H}^1} \rightarrow 0 \quad \text{as } t \rightarrow +\infty.$$

We start by stating the following monotonicity results.

Lemma A.6. *Consider two real sequences $\{t_n\}_n$ and $\{t'_n\}_n$, $t_n < t'_n$, and a sequence $\{u_n\}_n$ of radial solutions to (1.2) on $[t_n, t'_n]$ satisfying (7.1). Assume there exists $\{\lambda_n(t)\}_n \subset \mathbb{R}_+^*$ such that the set*

$$K = \{(u_n(t))_{[\lambda_n(t)]} : n \in \mathbb{N}, t \in [t_n, t'_n]\}$$

is relatively compact in \dot{H}^1 . If

$$\lim_n d(u_n(t_n)) + d(u_n(t'_n)) = 0,$$

then

$$(A.21) \quad \lim_n \left\{ \sup_{t \in [t_n, t'_n]} d(u_n(t)) \right\} = 0.$$

Lemma A.7. *Under the assumptions of Lemma A.6, if n is large enough so that $d(u_n(t)) \leq \delta_0$ on $[t_n, t'_n]$ and if θ_n, μ_n and α_n are the parameters of the decomposition (5.1), then*

$$(A.22) \quad \lim_n \frac{\sup_{t \in [t_n, t'_n]} \mu_n(t)}{\inf_{t \in [t_n, t'_n]} \mu_n(t)} = 1.$$

Remark A.8. In Lemmas A.6 and A.7, it is sufficient to assume

$$\inf_{t \in [t_n, t'_n]} \lambda(t) = 1 \quad \forall n \in \mathbb{N}.$$

In fact, if $\tilde{\lambda}_n := \inf_{t \in [t_n, t'_n]} \lambda(t)$, then

$$u_n^*(x, t) := \frac{1}{\tilde{\lambda}_n^{(N-2)/2}} u_n\left(\frac{x}{\tilde{\lambda}_n}, \frac{t}{\tilde{\lambda}_n^2}\right), \quad \lambda_n^*(t) := \frac{\lambda_n(t)}{\tilde{\lambda}_n}, \quad t_n^* := \frac{t_n}{\tilde{\lambda}_n^2}, \quad t_n'^* := \frac{t_n'}{\tilde{\lambda}_n^2},$$

$$K^* := \{(u_n^*(t))_{[\lambda_n^*(t)]} : n \in \mathbb{N}, t \in [t_n^*, t_n'^*]\}$$

satisfy the assumptions of Lemma A.6. Moreover, the conclusions of the lemmas are unchanged under these transformations.

Before proving Lemmas A.6 and A.7, we prove two auxiliary lemmas.

Lemma A.9. Consider two real sequences $\{t_n\}_n$ and $\{t'_n\}_n$, $t_n < t'_n$, and a sequence $\{u_n\}_n$ of radial solutions to (1.2) on $[t_n, t'_n]$ satisfying (7.1). Assume there exists $\{\lambda_n(t)\}_n \subset \mathbb{R}_+^*$ such that the set

$$K = \{(u_n(t))_{[\lambda_n(t)]} : n \in \mathbb{N}, t \in [t_n, t'_n]\}$$

is relatively compact in \dot{H}^1 . Assume furthermore that

$$(A.23) \quad \inf_{t \in [t_n, t'_n]} \lambda(t) = 1 \quad \forall n \in \mathbb{N}.$$

Then, for all $n \in \mathbb{N}$,

$$(A.24) \quad \int_{t_n}^{t'_n} d(u(t)) \lesssim d(u(t_n)) + d(u(t'_n)).$$

Proof of Lemma A.9. For $R > 0$, consider the function

$$F_{R,n}(t) = \int \phi_R |u_n(t)|^2.$$

By the Hölder and Sobolev inequalities, and recalling that $\|u(t)\|_{\dot{H}^1} \leq \|W\|_{\dot{H}^1}$, we have

$$F_{R,n}(t) \lesssim R^4.$$

By Lemma A.2,

$$(A.25) \quad |F'_{R,n}(t)| \lesssim d(u_n(t)) \sqrt{F_{R,n}(t)} \lesssim R^2 d(u_n(t)).$$

By (A.23), $\lambda(t) \geq 1$ on $[t_n, t'_n]$. We claim that, whenever defined, μ_n is bounded away from zero. In fact, by the precompactness of K and decomposition (5.1), we have

$$(u_n(t))_{[\lambda_n(t)]} = (1 + \alpha_n(t)) W_{[\lambda_n(t)/\mu_n(t)]} + (h_n(t))_{[\lambda_n(t)/\mu_n(t)]}$$

with $(h_n(t))_{[\lambda_n(t)/\mu_n(t)]} \perp W_{[\lambda_n(t)/\mu_n(t)]}$ and $\alpha_n(t) \leq \|u_{[\lambda_n(t)]}\|_{\dot{H}^1} + 1$. Therefore, the set

$$(A.26) \quad \bigcup_n \{W_{[\lambda_n(t)/\mu_n(t)]} : t \in [t_n, t'_n], d(u_n(t)) \leq \delta_0\}$$

must be precompact. Since W does not depend on time, we get

$$(A.27) \quad \lambda_n(t) \approx \mu_n(t) \text{ on } \{t \in [t_n, t'_n], d(u_n(t)) \leq \delta_0\}.$$

(Note that the constant does not depend on n). Thus, we set

$$\mu_- := \inf_{\substack{t \in [t_n, t'_n], \\ d(u_n(t)) \leq \delta_0}} \mu_n(t) \gtrsim 1.$$

We will now give a lower bound to $F''_{R,n}(t)$. Recalling (A.8), if $d(u_n(t)) \leq \delta_0$ and $R \geq 1/\mu_-$, we have

$$|A_R(u_n(t))| \lesssim \left[d^2(u_n(t)) + \frac{1}{(R\mu_-)^{(N-2)/2}} d(u_n(t)) \right].$$

Therefore, there exist $\delta_1 > 0$ and $R_1 > 0$ such that, if $d(u_n(t)) \leq \delta_1$, then

$$|A_R(u_n(t))| \leq \frac{8}{N-2} d(u_n(t)).$$

Now, by almost periodicity modulo symmetries and (A.23), if $\eta > 0$ and $R \geq C(\eta)$, then

$$|A_R(u_n(t))| \lesssim \eta.$$

Thus, we can choose $\eta_1 = \eta_1(\delta_1)$ such that, if $d(u_n(t)) \geq \delta_1$ and $R \geq C(\eta_1)$, then

$$|A_R(u_n(t))| \leq \frac{8}{N-2} \delta_1 \leq \frac{8}{N-2} d(u_n(t)).$$

Finally, since

$$F''_{R,n}(t) = \frac{16}{N-2} d(u_n(t)) + A_R(u_n(t)),$$

we get, if $R \geq \max\{R_1, C(\eta_1)\}$,

$$(A.28) \quad F''_{R,n}(t) \geq \frac{8}{N-2} d(u_n(t)).$$

Integrating (A.28) and using (A.25), we obtain (A.24). ■

Lemma A.10. *Under the assumptions of Lemma A.6 and Remark A.8, if $s_n \in [t_n, t'_n]$ and the sequence $\lambda_n(s_n)$ is bounded, then*

$$(A.29) \quad \lim_n d(u_n(s_n)) = 0.$$

Proof. By Remark A.8, we have $\lambda_n(s_n) \approx 1$, hence, we can assume that the sequence $\{u_n(s_n)\}_n$ converges to some $v_0 \in \dot{H}^1$. If (A.29) does not hold, then

$$(A.30) \quad d(v_0) > 0 \quad \text{and} \quad \|v_0\|_{\dot{H}^1} < \|W\|_{\dot{H}^1}.$$

By strong convergence, we have $E(v_0) = E(W)$. Let v be the solution to (1.2) with initial condition v_0 . By Proposition 7.4, $v(t)$ is defined for any $t \in \mathbb{R}$.

We claim that, for large n , $s_n + 1 \leq t'_n$. Indeed, if $t'_n \in (s_n, s_n + 1)$ for an infinite number of n , after extracting a subsequence, we have that $t'_n - s_n$ converges to some $\tau \in [0, 1]$. By continuity of the flow, $u_n(t'_n) \rightarrow v(\tau)$. But since $d(u_n(t'_n)) \rightarrow 0$, we have $d(v(\tau)) = 0$, which implies that $v = W_{[\lambda_0, \theta_0]}$, for some fixed λ_0, θ_0 . Uniqueness of solutions to (1.2) then contradicts (A.30). Therefore, for large n , $t_n \leq s_n \leq s_n + 1 \leq t'_n$. Again by continuity of the flow,

$$\lim_n \int_{s_n}^{s_n+1} d(u_n(t)) dt = \int_0^1 d(v(t)) dt > 0.$$

However, Lemma A.9 gives

$$\lim_n \int_{s_n}^{s_n+1} d(u_n(t)) dt \leq \lim_n \int_{t_n}^{t'_n} d(u_n(t)) dt \lesssim \lim_n d(u_n(t_n)) + d(u_n(t'_n)) = 0,$$

completing the proof of Lemma A.10. ■

We now prove Lemmas A.6 and A.7.

Proof of Lemma A.6. By Remark A.8, we can choose, for every n , $b_n \in [t_n, t'_n]$ such that

$$\lim_n \lambda_n(b_n) = 1.$$

This implies, by Lemma A.10, that

$$\lim_n d(u_n(b_n)) = 0.$$

Assume, by contradiction, that (A.21) does not hold. Without loss of generality, there exists $\delta_1 > 0$ such that

$$(A.31) \quad \sup_{t \in [t_n, b_n]} d(u_n(t)) \geq \delta_1, \quad \forall n \in \mathbb{N}.$$

Choosing $\delta_2 < \min\{\delta_0, \delta_1\}$, by continuity there exists $a_n \in [t_n, b_n]$ such that

$$d(u_n(t)) < \delta_2 \quad \text{on } (a_n, b_n), \quad \text{and} \quad d(u_n(a_n)) = \delta_2.$$

Since $\delta_2 < \delta_0$, the modulation parameter μ_n is well-defined. Recalling that the set defined by (A.26) is precompact, we must have $\lambda_n \approx \mu_n$, where the constants do not depend on n . Thus, up to a subsequence, we can assume

$$\mu_n(b_n) \rightarrow \mu_0 \in (0, +\infty) \quad \text{as } n \rightarrow +\infty.$$

We will now show that the μ_n are uniformly bounded on $\cup_n [a_n, b_n]$. Suppose, by contradiction, that there exists $c_n \in [a_n, b_n]$ such that, for large n ,

$$(A.32) \quad \mu_n(t) < 2\mu_0 \quad \text{on } (c_n, b_n), \quad \text{and} \quad \mu_n(c_n) = 2\mu_0.$$

Since $\mu_n(c_n)$ is bounded, so is $\lambda_n(c_n)$. Therefore, by Lemma A.10, $\lim_{n \rightarrow \infty} d(u_n(c_n)) = 0$. Recalling Lemma 5.2, we have

$$\left| \frac{1}{\mu_n^2(c_n)} - \frac{1}{\mu_n^2(b_n)} \right| \leq \int_{c_n}^{b_n} \left| \frac{\mu'_n(t)}{\mu_n^3(t)} \right| \lesssim \int_{c_n}^{b_n} d(u_n(t)) dt.$$

By Lemma A.9, the last integral converges to 0, contradicting (A.32). Therefore,

$$\sup_{\substack{t \in [a_n, b_n] \\ n \in \mathbb{N}}} \mu_n(t) < +\infty.$$

We conclude that $\mu_n(a_n)$ must be bounded, and so must be $\lambda_n(a_n)$. Invoking again Lemma A.10, we have $\lim_{n \rightarrow \infty} d(u_n(a_n)) = 0$, contradicting (A.31). Lemma A.6 is proven. ■

Proof of Lemma A.7. As in the proof of the previous lemma, we have, by Remark A.8 and Lemmas A.9 and A.10, that $\mu_n \approx 1$, where the constant does not depend on n . Let a_n and b_n be such that

$$\mu_n(a_n) = \inf_{t \in [t_n, t'_n]} \mu_n(t) \quad \text{and} \quad \mu_n(b_n) = \sup_{t \in [t_n, t'_n]} \mu_n(t).$$

Let $\bar{a}_n = \min\{a_n, b_n\}$ and $\bar{b}_n = \max\{a_n, b_n\}$. Then,

$$\left| \frac{1}{\mu_n^2(a_n)} - \frac{1}{\mu_n^2(b_n)} \right| \leq \int_{\bar{a}_n}^{\bar{b}_n} \left| \frac{\mu'_n(t)}{\mu_n^3(t)} \right| \lesssim \int_{\bar{a}_n}^{\bar{b}_n} d(u_n(t)) dt \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Since $\mu_n(b_n)$ is bounded, we get (A.22), proving Lemma A.7. ■

We now have all the tools to prove Proposition 7.8.

Proof of Proposition 7.8. By Corollary 7.7, there exists a sequence $t_n \rightarrow +\infty$ such that

$$\lim_n d(u(t_n)) = 0.$$

By Lemma (A.6), with $u_n = u$, $\lambda_n = \lambda$ (where λ is the frequency scale obtained from Proposition 7.4) and $t'_n = t_{n+1}$, this implies

$$(A.33) \quad \lim_{t \rightarrow +\infty} d(t) = 0.$$

Therefore, the modulation parameters $\alpha(t)$, $\mu(t)$ and $\theta(t)$ are defined for large t . We now prove that

$$(A.34) \quad \lim_{t \rightarrow +\infty} \mu(t) = \mu_\infty \in (0, +\infty).$$

Indeed, if not, then as $t \rightarrow +\infty$, $\log(\mu(t))$ does not satisfy the Cauchy criterion. Therefore, there must exist sequences $\{T_n\}$ and $\{T'_n\}$ such that $T_n < T'_n$ and

$$(A.35) \quad \lim_n \frac{\mu(T'_n)}{\mu(T_n)} \neq 1.$$

But $d(T_n) + d(T'_n) \rightarrow 0$ by (A.33). By Lemma (A.7), with $u_n = u$, $\lambda_n = \lambda$, $t_n = T_n$ and $t'_n = T'_n$, we have

$$\lim_n \frac{\sup_{t \in [T_n, T'_n]} \mu(t)}{\inf_{t \in [T_n, T'_n]} \mu(t)} = 1,$$

contradicting (A.35). Turning to the proof of (7.3), we claim the following inequality:

$$(A.36) \quad \int_t^{+\infty} d(u(s)) \, ds \lesssim d(u(t)).$$

Suppose by contradiction that (A.36) does not hold. Then there exists a sequence $T_n \rightarrow +\infty$ such that

$$\int_{T_n}^{+\infty} d(u(s)) \, ds \geq 2n d(u(T_n)).$$

Moreover, there exists a sequence $\{S_n\}_n$ such that $S_n > T_n$ for all n , and

$$\int_{T_n}^{S_n} d(u(s)) \, ds \geq n d(u(T_n)).$$

By (A.34), for any sequence $\{T'_n\}_n$ such that $T'_n \geq S_n$ for all n , we are under the assumptions of Lemma A.9, with $u_n = u$, $\lambda_n = \lambda$, $T_n = t_n$ and $t'_n = T'_n$. Hence,

$$n d(u(T_n)) \leq \int_{T_n}^{T'_n} d(u(s)) \, ds \lesssim d(u(T_n)) + d(u(T'_n)).$$

Since T'_n can be taken arbitrarily large, and the implicit constant is independent of the choice of a particular $\{T'_n\}_n$ (given the function u itself does not change), we have a contradiction.

Note that (A.36) is equivalent to the existence of $c > 0$ such that

$$\int_t^{+\infty} d(u(s)) \, ds \lesssim e^{-ct}.$$

By Lemma 5.2, since $|\alpha(t)| \approx d(u(t))$ and μ is bounded, there exists θ_∞ such that

$$|\alpha(t)| + |\theta(t) - \theta_\infty| + \|h(t)\|_{\dot{H}^1} \lesssim e^{-ct}.$$

Therefore, the bound (7.3) is proven. The assertion about scattering for negative times is a corollary of Lemma A.9. Indeed, if

$$\|u\|_{S(0,+\infty)} = \|u\|_{S(-\infty,0)} = +\infty,$$

by time-reversal and (7.3), we see that the set $\{u(t) : t \in \mathbb{R}\}$ is relatively compact and that

$$\lim_{t \rightarrow \pm\infty} d(t) = 0.$$

Therefore, by Lemma A.9, with $u_n = u$, $\lambda_n = 1$, $t_n = -n$ and $t'_n = n$, we have

$$\int_{-\infty}^{+\infty} d(t) \, dt = \lim_{n \rightarrow +\infty} \int_{-n}^n d(t) \, dt \lesssim d(-n) + d(n) = 0.$$

Therefore, $d(u_0) = 0$, contradicting (7.1). Proposition 7.8 is proven. ■

For the intercritical case, as in the proof of Proposition 6.3, we need to show that

$$(A.37) \quad \int_t^{+\infty} d(s) ds \lesssim e^{-ct}, \quad \forall t \geq 0.$$

We start with the following lemmas.

Lemma A.11. *Let u be a solution to (1.1) satisfying (7.4) and (7.5). Then*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T d(t) dt = 0.$$

We next state a key result to prove Proposition A.37.

Lemma A.12. *Let u be a solution to (1.1) satisfying (7.4) and (7.5), and let $x(t)$ be as in Proposition 7.11 and Remark 7.13. Then, for any $0 \leq \sigma \leq \tau$,*

$$(A.38) \quad \int_\sigma^\tau d(u(t)) \lesssim \left[1 + \sup_{\sigma \leq t \leq \tau} |x(t)| \right] (d(u(\sigma)) + d(u(\tau))),$$

and, if $\tau \geq \sigma + 1$,

$$(A.39) \quad |x(\tau) - x(\sigma)| \lesssim \int_\sigma^\tau d(u(t)).$$

The proof of (A.38) is similar to the energy-critical setting (it is in fact easier, since there is no scaling involved). We refer to Lemma 6.7 in [13] for the argument in the 3d cubic case. The proof of (A.39) follows verbatim the proof of Lemma 6.8 in [13].

Proof of Lemma A.11. Let $R > 0$ to be chosen later and let ϕ_R and F_R be as in the previous section. Then, by Hölder’s inequality,

$$(A.40) \quad |F'_R(t)| \lesssim R.$$

Moreover, we have

$$(A.41) \quad F''_R(t) = [2N(p - 1) - 8]d(t) + A_R(u(t)),$$

where A_R is given by (A.18).

Fix $\eta > 0$. By definition of ϕ_R and almost periodicity modulo symmetries, if $R \geq C(\eta)$, we have

$$(A.42) \quad |A_R(u(t))| \lesssim \int_{|x| \geq R} |\nabla u(x, t)|^2 + |u(x, t)|^{p+1} + \frac{1}{|x|^2} |u(x, t)|^2 dx.$$

Choose $T_0(\eta) \geq 0$ such that, for any $t \geq T_0$,

$$|x(t)| \leq \eta t.$$

For $T \geq T_0$, choose $R = \eta T + C(\eta) + 1$. With this choice of R , we have

$$\begin{aligned} |A_R(u(t))| &\lesssim \int_{|x-x(t)|+|x(t)|\geq R} |\nabla u(x, t)|^2 + |u(x, t)|^{p+1} + |u(x, t)|^2 dx \\ &\lesssim \int_{|x-x(t)|\geq C(\eta)} |\nabla u(x, t)|^2 + |u(x, t)|^{p+1} + |u(x, t)|^2 dx \lesssim \eta. \end{aligned}$$

By (A.40), (A.41), and (A.42),

$$\begin{aligned} [2N(p - 1) - 8] \int_{T_0}^T d(t) dt &\lesssim |F'_R(T)| + |F'_R(T_0)| + \eta(T - T_0) \\ &\lesssim R + \eta(T - T_0) = \eta T + \eta(T - T_0). \end{aligned}$$

Letting $T \rightarrow +\infty$, we deduce

$$\limsup_{T \rightarrow +\infty} \frac{1}{T} \int_0^T d(t) dt \lesssim \eta.$$

Since η is arbitrary, we conclude the proof of Lemma A.11. ■

We are now able to prove Proposition 7.14, following the proof in [13].

Proof of Proposition 7.14. We first show that $x(t)$ is bounded. By Lemma A.11, there exists a sequence $\{t_n\}_n$ such that $t_{n+1} \geq t_n + 1$ for all n , and $d(u(t_n)) \rightarrow 0$. Now, by Lemma A.12, there exists $C_0 > 0$ such that, if $n > n_0$ and $1 + t_{n_0} \leq t \leq t_n$, then

$$|x(t) - x(t_{n_0})| \leq C_0 \left[1 + \sup_{t_{n_0} \leq s \leq t_n} |x(s)| \right] (d(u(t_n)) + d(u(t_{n_0}))).$$

If n_0 is large enough so that $d(u(t_n)) + d(u(t_{n_0})) \leq 1/(2C_0)$, and t is chosen in $[t_{n_0} + 1, t_n]$ so that $\sup_{t_{n_0} + 1 \leq s \leq t_n} |x(s)| = |x(t)|$, then

$$\sup_{t_{n_0} + 1 \leq s \leq t_n} |x(s)| \leq C(n_0) + \frac{1}{2} \sup_{t_{n_0} + 1 \leq s \leq t_n} |x(s)|,$$

where $C(n_0) = |x(t_{n_0})| + \frac{1}{2} \sup_{t_{n_0} \leq s \leq t_{n_0} + 1} |x(t)| + 1/2$. Therefore, $x(t)$ is bounded on $[t_{n_0} + 1, +\infty)$, and hence, by continuity, on $[0, +\infty)$.

By the boundedness of $x(t)$ and (A.38), we have

$$\int_{\sigma}^{\tau} d(u(t)) \lesssim d(u(\sigma)) + d(u(\tau)).$$

For a fixed $\sigma \geq 0$ and choosing $\tau = t_n$, we let $n \rightarrow +\infty$ to obtain

$$\int_{\sigma}^{\infty} d(u(t)) \lesssim d(u(\sigma)).$$

By Gronwall's lemma, we have (A.37) and, using Lemma 5.5, we finish the proof of Proposition 7.14. ■

A.4. Results for the linearized equation

Proof of Lemma 8.1. By Lemma 3.12, we can assume that

$$(A.43) \quad \|h(t)\|_{S(L^2, [t, +\infty))} \lesssim e^{-c_0 t}.$$

We first normalize the eigenfunctions of \mathcal{L} . Define

$$f_0 = \frac{iW}{\|W\|_{\dot{H}^1}}, \quad f_k = \frac{\partial_k W}{\|\partial_k W\|_{\dot{H}^1}}, \quad 1 \leq k \leq N, \quad \text{and} \quad f_{N+1} = \frac{\Lambda W}{\|\Lambda W\|_{\dot{H}^1}}.$$

We have

$$B(f_k, h) = 0, \quad \|f_k\|_{\dot{H}^1}, \quad \forall 0 \leq k \leq N + 1, \quad \forall h \in \dot{H}^1.$$

Recall that $B(\mathcal{Y}_+, \mathcal{Y}_-) \neq 0$. Normalize $\mathcal{Y}_+, \mathcal{Y}_-$ such that $B(\mathcal{Y}_+, \mathcal{Y}_-) = 1$. Next, write

$$(A.44) \quad h(t) = \alpha_+(t)\mathcal{Y}_+ + \alpha_-(t)\mathcal{Y}_- + \sum_k \beta_k(t)f_k + g(t), \quad g(t) \in \tilde{G}^\perp,$$

where, recalling that $\mathcal{L}_{|\text{span}\{f_k, k \leq N+1\}} = 0$ and that $\Phi(\mathcal{Y}_+) = \Phi(\mathcal{Y}_-) = 0$,

$$(A.45) \quad \alpha_+(t) = B(h(t), \mathcal{Y}_-), \quad \alpha_-(t) = B(h(t), \mathcal{Y}_+),$$

$$(A.46) \quad \beta_k(t) = (h(t), f_k)_{\dot{H}^1} - \alpha_+(t)(\mathcal{Y}_+, f_k)_{\dot{H}^1} - \alpha_-(t)(\mathcal{Y}_-, f_k)_{\dot{H}^1}, \quad \forall k \leq N + 1.$$

Step 1. Differential inequalities on the coefficients. We will show

$$(A.47) \quad \frac{d}{dt}(e^{e_0 t} \alpha_+(t)) = e^{e_0 t} B(\mathcal{Y}_-, \epsilon), \quad \frac{d}{dt}(e^{-e_0 t} \alpha_-(t)) = e^{-e_0 t} B(\mathcal{Y}_+, \epsilon),$$

$$(A.48) \quad \frac{d}{dt}(e^{-e_0 t} \beta_k(t)) = (f_k, \epsilon)_{\dot{H}^1} - (\mathcal{Y}_+, f_k)_{\dot{H}^1} B(\mathcal{Y}_-, \epsilon) - (\mathcal{Y}_-, f_k)_{\dot{H}^1} B(\mathcal{Y}_+, \epsilon) - (\mathcal{L}g, f_k)_{\dot{H}^1},$$

$$(A.49) \quad \frac{d\Phi(h(t))}{dt} = 2B(h, \epsilon).$$

By equation (8.1),

$$(A.50) \quad \begin{aligned} \alpha'_+(t) &= B(\partial_t h, \mathcal{Y}_-) = B(-\mathcal{L}h + \epsilon, \mathcal{Y}_-) \\ &= B(h, \mathcal{L}\mathcal{Y}_-) + B(\epsilon, \mathcal{Y}_-) = -e_0 \alpha_+(t) + B(\epsilon, \mathcal{Y}_-). \end{aligned}$$

This yields the first equation in (A.47). The second equation follows similarly.

Now, differentiating (A.46), we obtain

$$\beta'_k = (-\mathcal{L}h + \epsilon - \alpha'_+ \mathcal{Y}_+ - \alpha'_- \mathcal{Y}_-, f_k)_{\dot{H}^1}.$$

Note that $\mathcal{L}h = e_0 \alpha_+ \mathcal{Y}_+ - e_0 \alpha_- \mathcal{Y}_- + \mathcal{L}g$, by (A.44), which proves (A.48), in view of (A.47).

Finally, differentiating $\Phi(h(t))$,

$$\frac{d}{dt}\Phi(t) = 2B(h, \partial_t h) = -2B(h, \mathcal{L}h) + 2B(h, \epsilon) = 2B(h, \epsilon),$$

by the skew-symmetry of \mathcal{L} in (3.3). Equation (A.49) is then proved.

Step 2. Estimates on α_{\pm} . We claim that

$$(A.51) \quad |\alpha_{-}(t)| \lesssim e^{-c_1 t},$$

$$(A.52) \quad |\alpha_{+}(t)| \lesssim \begin{cases} e^{-c_1 t} & \text{if } e_0 < c_0, \\ e^{-e_0 t} + e^{-c_1^{-} t} & \text{if } e_0 \geq c_0. \end{cases}$$

We need the following claim, which is an immediate application of Hölder’s inequality.

Claim A.13. *If I is a finite time interval, and $f \in L_I^{\infty} L_x^{2N/(N-2)}$, $g \in L_I^{\infty} L_x^{2N/(N+2)}$ are such that $\nabla f \in L_I^2 L_x^{2N/(N-2)}$, $\nabla g \in L_I^2 L_x^{2N/(N+2)}$, then*

$$\int_I |B(f(t), g(t))| dt \lesssim \|\nabla f\|_{L_I^2 L_x^{2N/(N-2)}} \|\nabla g\|_{L_I^2 L_x^{2N/(N+2)}} + \|I\| \|f\|_{L_I^{\infty} L_x^{2N/(N-2)}} \|g\|_{L_I^{\infty} L_x^{2N/(N+2)}}.$$

The above claim, (A.43) and (A.47) yield

$$\int_t^{t+1} |e^{-e_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \leq e^{-e_0 t} \int_t^{t+1} |B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{-(e_0+c_1)t}.$$

By the triangle inequality, integrating the second equation in (A.47) gives

$$|\alpha_{-}(t)| \lesssim e^{e_0 t} \int_t^{+\infty} |e^{-e_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{-c_1 t},$$

which proves (A.51).

To prove (A.52), consider first the case $c_0 > e_0$. Then, by (A.43) and (A.45), we have that $e^{e_0 t} \alpha_{+}(t)$ vanishes as $t \rightarrow +\infty$. By Claim A.13,

$$\int_t^{t+1} |e^{e_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{e_0 t} \int_t^{t+1} |B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{(e_0-c_1)t}.$$

Integrating the equation on α_{+} in (A.47), recalling that $c_1 > c_0$, and using triangle inequality, we get (A.52) if $c_0 > e_0$.

Assume now that $c_0 \leq e_0$. Integrating (A.47),

$$|\alpha_{+}(t) - e^{-e_0 t} \alpha_{+}(0)| \leq e^{-e_0 t} \int_0^t e^{e_0 s} |B(\mathcal{Y}_-, \epsilon(s))| ds \lesssim e^{-c_1^{-} t},$$

and the proof of (A.52) is finished.

Step 3. Bounds on g and β_k . We will prove

$$(A.53) \quad \|g(t)\|_{\dot{H}^1} + \sum_k |\beta_k(t)| \lesssim e^{-(c_0+c_1)t/2}.$$

Again by Claim A.13, $\int_t^{t+1} |B(h(s), \epsilon(s))| ds \lesssim e^{-(c_0+c_1)t}$. By the triangle inequality, integrating (A.49), we get

$$\Phi(h(t)) \lesssim e^{-(c_0+c_1)t}.$$

Therefore,

$$|2\alpha_{+}\alpha_{-}B(\mathcal{Y}_+, \mathcal{Y}_-) + \Phi(g)| = |\Phi(h)| \lesssim e^{-(c_0+c_1)t}.$$

By Step 2,

$$|\Phi(g)| \lesssim \begin{cases} e^{-(c_0+c_1)t} + e^{-2c_1t} & \text{if } c_0 > e_0, \\ e^{-(c_0+c_1)t} + e^{-c_1t}(e^{-e_0t} + e^{-c_1^{-1}t}) & \text{if } c_0 \leq e_0. \end{cases}$$

In any case, $|\Phi(g)| \leq e^{-(c_0+c_1)t}$. Using the coercivity of Φ , given by Lemma 3.5, the estimate for g in (A.53) is proven.

Consider now (A.48). By (A.43),

$$\begin{aligned} \beta_k(t + 1) - \beta_k(t) &\lesssim e^{-c_1t} + \int_t^{t+1} |(f_k, \mathcal{L}g(s))_{\dot{H}^1}| ds \\ &= e^{-c_1t} + \int_t^{t+1} \left| \operatorname{Re} \int \mathcal{L}^*(\Delta f_k) \bar{g}(s) \right| ds, \end{aligned}$$

where $\mathcal{L}^* = \begin{pmatrix} 0 & L_+ \\ -L_- & 0 \end{pmatrix}$ is the L^2 -adjoint of \mathcal{L} .

One can check explicitly that, for any $0 \leq k \leq N + 1$, $|\mathcal{L}^*(\Delta f_k)| \lesssim \frac{1}{1+|x|^{N+4}}$. Therefore, $\mathcal{L}^*(\Delta f_k) \in L^{2N/(N+2)}(\mathbb{R}^N)$, so that, by the estimate on g in (A.53), we obtain

$$\left| \operatorname{Re} \int \mathcal{L}^*(\Delta f_k) \bar{g}(t) \right| \lesssim \|g(t)\|_{2N/(N-2)} \lesssim \|g\|_{\dot{H}^1} \lesssim e^{-(c_0+c_1)t/2}.$$

Step 4. Closure. By the decomposition (A.44), as well as Steps 2 and 3, so far we have

$$\|h(t)\|_{\dot{H}^1} \lesssim \begin{cases} e^{-(c_0+c_1)t/2} & \text{if } c_0 > e_0, \\ e^{-e_0t} + e^{-(c_0+c_1)t/2} & \text{if } c_0 \leq e_0. \end{cases}$$

Now, if $e_0 \notin [c_0, c_1]$, by iterating the argument, we obtain

$$\|h(t)\|_{\dot{H}^1} \lesssim e^{-c_1^{-1}t},$$

which proves (8.3).

Assume now $e_0 \in [c_0, c_1]$. Then, the estimate (A.47) on α_+ ensures the existence of a limit A to $e^{e_0t}\alpha_+(t)$ as $t \rightarrow +\infty$. Integrating (A.47) from t to $+\infty$, we get

$$|A - e^{e_0t}\alpha_+| \leq e^{e_0t} \int_t^{+\infty} |B(y_+, \epsilon(s))| ds \lesssim e^{(e_0-c_1)t}.$$

In view of the decomposition (A.44) and estimates (A.51), (A.52) and (A.53), we get

$$\|h(t) - Ae^{-e_0t}y_+\|_{\dot{H}^1} \lesssim e^{-(c_0+c_1)t/2}.$$

Since $\mathcal{L}y_+ = e_0y_+$, we see that $\tilde{h}(t) := h(t) - Ae^{-e_0t}y_+$ satisfies the differential equation (8.1) with the same ϵ , and with c_0 replaced by $(c_0 + c_1)/2 > c_0$ in the condition (8.2). By iterating the argument a finite number of times, we end up under the condition (8.3), which implies condition (8.4) for the original h , and finishes the proof of Lemma 8.1. ■

Proof of Lemma 8.7. We first normalize the eigenfunctions of \mathcal{L} . Define

$$f_0 := \frac{iQ}{\|Q\|_{L^2}}, \quad f_k := \frac{\partial_k Q}{\|\partial_k Q\|_{L^2}}, \quad 1 \leq k \leq N.$$

We have

$$B(f_k, h) = 0, \quad \|f_k\|_{L^2} = 1, \quad \forall 0 \leq k \leq N, \forall h \in H^1.$$

Recall that $B(\mathcal{Y}_+, \mathcal{Y}_-) \neq 0$. Normalize $\mathcal{Y}_+, \mathcal{Y}_-$ such that $B(\mathcal{Y}_+, \mathcal{Y}_-) = 1$. Next, write

$$(A.54) \quad h(t) = \alpha_+(t)\mathcal{Y}_+ + \alpha_-(t)\mathcal{Y}_- + \sum_k \beta_k(t)f_k + g(t), \quad g(t) \in \tilde{G}^\perp,$$

where, recalling that $\mathcal{L}|_{\text{span}\{f_k, k \leq N\}} = 0$ and that $\Phi(\mathcal{Y}_+) = \Phi(\mathcal{Y}_-) = 0$, we have

$$(A.55) \quad \alpha_+(t) = B(h(t), \mathcal{Y}_-), \quad \alpha_-(t) = B(h(t), \mathcal{Y}_+),$$

$$(A.56) \quad \beta_k(t) = (h(t), f_k)_{L^2} - \alpha_+(t)(\mathcal{Y}_+, f_k)_{L^2} - \alpha_-(t)(\mathcal{Y}_-, f_k)_{L^2}, \quad \forall k \leq N.$$

Step 1. Differential inequalities on the coefficients. We show

$$(A.57) \quad \frac{d}{dt}(e^{e_0 t} \alpha_+(t)) = e^{e_0 t} B(\mathcal{Y}_-, \epsilon), \quad \frac{d}{dt}(e^{-e_0 t} \alpha_-(t)) = e^{-e_0 t} B(\mathcal{Y}_+, \epsilon),$$

$$(A.58) \quad \frac{d}{dt}(e^{-e_0 t} \beta_k(t)) = (f_k, \epsilon)_{L^2} - (\mathcal{Y}_+, f_k)_{L^2} B(\mathcal{Y}_-, \epsilon) - (\mathcal{Y}_-, f_k)_{L^2} B(\mathcal{Y}_+, \epsilon) - (\mathcal{L}g, f_k)_{L^2},$$

$$(A.59) \quad \frac{d\Phi(h(t))}{dt} = 2B(h, \epsilon).$$

By equation (8.16),

$$(A.60) \quad \begin{aligned} \alpha'_+(t) &= B(\partial_t h, \mathcal{Y}_-) = B(-\mathcal{L}h + \epsilon, \mathcal{Y}_-) \\ &= B(h, \mathcal{L}\mathcal{Y}_-) + B(\epsilon, \mathcal{Y}_-) = -e_0 \alpha_+(t) + B(\epsilon, \mathcal{Y}_-). \end{aligned}$$

This yields the first equation in (A.57). The second equation follows similarly.

Now, differentiating (A.56), we obtain

$$\beta'_k = (\mathcal{L}h + \epsilon - \alpha'_+ \mathcal{Y}_+ - \alpha'_- \mathcal{Y}_-, f_k)_{L^2}.$$

Note that $\mathcal{L}h = e_0 \alpha_+ \mathcal{Y}_+ - e_0 \alpha_- \mathcal{Y}_- + \mathcal{L}g$, by (A.54), which proves (A.58), in view of (A.57).

Finally, differentiating $\Phi(h(t))$,

$$\frac{d}{dt}\Phi(t) = 2B(h, \partial_t h) = -2B(h, \mathcal{L}h) + 2B(h, \epsilon) = 2B(h, \epsilon),$$

by the skew-symmetry of \mathcal{L} in (3.3). Equation (A.59) is then proved.

Step 2. Estimates on α_\pm . We claim that

$$(A.61) \quad |\alpha_-(t)| \lesssim e^{-c_1 t},$$

$$(A.62) \quad |\alpha_+(t)| \lesssim \begin{cases} e^{-c_1 t} & \text{if } c_0 \leq e_0, \\ e^{-e_0 t} + e^{-c_1^- t} & \text{if } c_0 > e_0. \end{cases}$$

We will need the following inequality, which is an immediate application of Hölder’s inequality:

$$(A.63) \quad \int_I |B(f(t), g(t))| dt \lesssim \| \langle \nabla \rangle f \|_{S(L^2, I)} \| \langle \nabla \rangle g \|_{S'(L^2, I)}.$$

The above inequality, assumption (8.17) and (A.57) yield

$$\int_t^{+\infty} |e^{-\epsilon_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \leq e^{-\epsilon_0 t} \int_t^{+\infty} |B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{-(\epsilon_0+c_1)t}.$$

By integrating the second equation in (A.57), we get

$$|\alpha_-(t)| \lesssim e^{\epsilon_0 t} \int_t^{+\infty} |e^{-\epsilon_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{-c_1 t},$$

which proves (A.61).

To prove (A.62), consider first the case $c_1 > c_0 > \epsilon_0$. Then, by assumption (8.17) and (A.55), $e^{\epsilon_0 t} \alpha_+(t)$ vanishes as $t \rightarrow +\infty$. By (A.63), integrating the equation on α_+ in (A.57),

$$|e^{\epsilon_0 t} \alpha_+(t)| \lesssim \int_t^{+\infty} |e^{\epsilon_0 s} B(\mathcal{Y}_+, \epsilon(s))| ds \lesssim e^{(\epsilon_0-c_1)t},$$

and we get (A.62) if $c_0 > \epsilon_0$.

Assume now that $c_0 \leq \epsilon_0$. By (A.57), we have

$$|\alpha_+(t) - e^{-\epsilon_0 t} \alpha_+(0)| \leq e^{-\epsilon_0 t} \int_0^t e^{\epsilon_0 s} |B(\mathcal{Y}_-, \epsilon(s))| ds \lesssim e^{-c_1^- t},$$

and the proof of (A.62) is finished.

Step 3. Bounds on g and β_k . We prove

$$(A.64) \quad \|g(t)\|_{L^2} + \sum_k |\beta_k(t)| \lesssim e^{-(c_0+c_1)t/2}.$$

Again by (A.63), $\int_t^{+\infty} |B(h(s), \epsilon(s))| ds \lesssim e^{-(c_0+c_1)t}$. By integrating (A.59), we get

$$\Phi(h(t)) \lesssim e^{-(c_0+c_1)t}.$$

Therefore,

$$|2\alpha_+ \alpha_- B(\mathcal{Y}_+, \mathcal{Y}_-) + \Phi(g)| = |\Phi(h)| \lesssim e^{-(c_0+c_1)t}.$$

By Step 2,

$$|\Phi(g)| \lesssim \begin{cases} e^{-(c_0+c_1)t} + e^{-2c_1 t} & \text{if } c_0 > \epsilon_0, \\ e^{-(c_0+c_1)t} + e^{-c_1 t} (e^{-\epsilon_0 t} + e^{-c_1^- t}) & \text{if } c_0 \leq \epsilon_0. \end{cases}$$

In any case, $|\Phi(g)| \leq e^{-(c_0+c_1)t}$. Using the coercivity of Φ , given by Lemma 3.5, the estimate for Φ in (A.64) is proven.

Consider now (A.58). By the assumption (8.17),

$$\beta_k(t + 1) - \beta_k(t) \lesssim e^{-c_1 t} + \int_t^{t+1} |(f_k, \mathcal{L}g(s))_{L^2}| ds = e^{-c_1 t} + \int_t^{t+1} \left| \operatorname{Re} \int \mathcal{L}^*(f_k) \bar{g}(s) \right| ds,$$

where $\mathcal{L}^* = \begin{pmatrix} 0 & L_+ \\ -L_- & 0 \end{pmatrix}$ is the L^2 -adjoint of \mathcal{L} .

One can check that, for any $0 \leq k \leq N$, $|\mathcal{L}^*(f_k)| \lesssim e^{-|x|}$. Therefore, $\mathcal{L}^*(f_k) \in L^2$ so that, by the estimate on g in (A.64),

$$\left| \operatorname{Re} \int \mathcal{L}^*(f_k) \bar{g}(t) \right| \lesssim \|g(t)\|_{L^2} \leq \|g\|_{H^1} \lesssim e^{-(c_0+c_1)t/2}.$$

Step 4. Closure. By the decomposition (A.54), as well as Steps 2 and 3, so far we have

$$\|h(t)\|_{H^1} \lesssim \begin{cases} e^{-(c_0+c_1)t/2} & \text{if } c_0 > e_0, \\ e^{-e_0 t} + e^{-(c_0+c_1)t/2} & \text{if } c_0 \leq e_0. \end{cases}$$

Now, if $e_0 \notin [c_0, c_1]$, by iterating the argument, we obtain

$$\|h(t)\|_{H^1} \lesssim e^{-c_1^- t},$$

which proves (8.18).

Assume now $e_0 \in [c_0, c_1]$. Then, the estimate (A.57) on α_+ ensures the existence of a limit A to $e^{e_0 t} \alpha_+(t)$, as $t \rightarrow +\infty$. Integrating (A.57) from t to $+\infty$,

$$|A - e^{e_0 t} \alpha_+| \leq e^{e_0 t} \int_t^{+\infty} |B(y_+, \epsilon(s))| ds \lesssim e^{(e_0-c_1)t}.$$

In view of the decomposition (A.54) and the estimates (A.61), (A.62) and (A.64), we get

$$\|h(t) - Ae^{-e_0 t} y_+\|_{H^1} \lesssim e^{-(c_0+c_1)t/2}.$$

Since $\mathcal{L}y_+ = e_0 y_+$, we see that $\tilde{h}(t) := h(t) - Ae^{-e_0 t} y_+$ satisfies the differential equation (8.16) with the same ϵ , and with c_0 replaced by $(c_0 + c_1)/2 > c_0$ in condition (8.17). By iterating the argument a finite number of times, we end up under condition (8.18), which implies condition (8.19) for the original h , and finishes the proof of Lemma 8.7. ■

Acknowledgements. This project was mostly done while L. C. was visiting the Department of Mathematics and Statistics at Florida International University, Miami, FL, during his third year of PhD training. He thanks the Department and the University for the hospitality and support. He would like to thank Thomas Duyckaerts and Nicola Visciglia for their valuable comments and suggestions, which helped improve the manuscript.

Funding. L. C. was financed by grant #2020/10185-1, São Paulo Research Foundation (FAPESP) and in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior -Brasil (CAPES)-, Finance Code 001. S. R. was partially supported by the NSF DMS grant 1927258, and part of research travel of L. C. was funded by the same grant DMS-1927258 (PI: Roudenko). L. G. F. was partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior -Brasil (CAPES)-, Finance Code 001, Conselho Nacional de Desenvolvimento Científico e Tecnológico-Brasil (CNPq) and Fundação de Amparo a Pesquisa do Estado de Minas Gerais- Brasil (FAPEMIG).

References

- [1] Aronszajn, N. and Smith, K. T.: Theory of Bessel potentials. I. *Ann. Inst. Fourier (Grenoble)* **11** (1961), 385–475.
- [2] Aubin, T.: Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire. *J. Math. Pures Appl. (9)* **55** (1976), no. 3, 269–296.
- [3] Banica, V.: Remarks on the blow-up for the Schrödinger equation with critical mass on a plane domain. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **3** (2004), no. 1, 139–170.
- [4] Berestycki, H., Lions, P.-L. and Peletier, L. A.: An ODE approach to the existence of positive solutions for semilinear problems in \mathbb{R}^N . *Indiana Univ. Math. J.* **30** (1981), no. 1, 141–157.
- [5] Berger, M., Gauduchon, P. and Mazet, E.: *Le spectre d'une variété riemannienne*. Lecture Notes in Mathematics 194, Springer-Verlag, Berlin-New York, 1971.
- [6] Cazenave, T.: *Semilinear Schrödinger equations*. Courant Lecture Notes in Mathematics 10, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- [7] Cazenave, T. and Weissler, F. B.: The Cauchy problem for the critical nonlinear Schrödinger equation in H^s . *Nonlinear Anal.* **14** (1990), no. 10, 807–836.
- [8] Chang, S.-M., Gustafson, S., Nakanishi, K. and Tsai, T.-P.: Spectra of linearized operators for NLS solitary waves. *SIAM J. Math. Anal.* **39** (2007/08), no. 4, 1070–1111.
- [9] Christ, F. M. and Weinstein, M. I.: Dispersion of small amplitude solutions of the generalized Korteweg–de Vries equation. *J. Funct. Anal.* **100** (1991), no. 1, 87–109.
- [10] Colliander, J., Keel, M., Staffilani, G., Takaoka, H. and Tao, T.: Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in \mathbb{R}^3 . *Ann. of Math. (2)* **167** (2008), no. 3, 767–865.
- [11] Dodson, B.: Global well-posedness and scattering for the focusing, cubic Schrödinger equation in dimension $d = 4$. *Ann. Scient. Éc. Norm. Sup.* **52** (2019), no. 1, 139–180.
- [12] Duyckaerts, T. and Merle, F.: Dynamic of threshold solutions for energy-critical NLS. *Geom. Funct. Anal.* **18** (2009), no. 6, 1787–1840.
- [13] Duyckaerts, T. and Roudenko, S.: Threshold solutions for the focusing 3D cubic Schrödinger equation. *Rev. Mat. Iberoam.* **26** (2010), no. 1, 1–56.
- [14] Duyckaerts, T. and Roudenko, S.: Going beyond the threshold: scattering and blow-up in the focusing NLS equation. *Comm. Math. Phys.* **334** (2015), no. 3, 1573–1615.
- [15] Fang, D. Y., Xie, J., and Cazenave, T.: Scattering for the focusing energy-subcritical nonlinear Schrödinger equation. *Sci. China Math.* **54** (2011), no. 10, 2037–2062.
- [16] Foschi, D.: Inhomogeneous Strichartz estimates. *J. Hyperbolic Differ. Equ.* **2** (2005), no. 1, 1–24.
- [17] Gidas, B., Ni, W. M. and Nirenberg, L.: Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n . In *Mathematical analysis and applications, Part A*, pp. 369–402. Adv. in Math. Suppl. Stud. 7, Academic Press, New York-London, 1981.
- [18] Ginibre, J. and Velo, G.: On a class of nonlinear Schrödinger equations. I. The Cauchy problem, general case. *J. Funct. Anal.* **32** (1979), no. 1, 1–32.
- [19] Guevara, C. and Carreon, F.: Scattering and blow up for the two-dimensional focusing quintic nonlinear Schrödinger equation. In *Recent advances in harmonic analysis and partial differential equations*, pp. 117–153. Contemp. Math. 581, Amer. Math. Soc., Providence, RI, 2012.

- [20] Guevara, C.: Global behavior of finite energy solutions to the d -dimensional focusing nonlinear Schrödinger equation. *Appl. Math. Res. Express. AMRX* (2014), no. 2, 177–243.
- [21] Holmer, J., Platte, R. and Roudenko, S.: Blow-up criteria for the 3D cubic nonlinear Schrödinger equation. *Nonlinearity* **23** (2010), no. 4, 977–1030.
- [22] Holmer, J. and Roudenko, S.: On blow-up solutions to the 3D cubic nonlinear Schrödinger equation. *Appl. Math. Res. Express. AMRX* (2007), Art. ID abm004, 31 pp.
- [23] Holmer, J. and Roudenko, S.: A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation. *Comm. Math. Phys.* **282** (2008), no. 2, 435–467.
- [24] Holmer, J. and Roudenko, S.: Divergence of infinite-variance nonradial solutions to the 3D NLS equation. *Commun. Partial Differential Equations* **35** (2010), no. 5, 878–905.
- [25] Kato, T.: An $L^{q,r}$ -theory for nonlinear Schrödinger equations. In *Spectral and scattering theory and applications*, pp. 223–238. Adv. Stud. Pure Math. 23, Math. Soc. Japan, Tokyo, 1994.
- [26] Keel, M. and Tao, T.: Endpoint Strichartz estimates. *Amer. J. Math.* **120** (1998), no. 5, 955–980.
- [27] Kenig, C. E. and Merle, F.: Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. *Invent. Math.* **166** (2006), no. 3, 645–675.
- [28] Killip, R. and Visan, M.: The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher. *Amer. J. Math.* **132** (2010), no. 2, 361–424.
- [29] Kwong, M. K.: Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbb{R}^n . *Arch. Rational Mech. Anal.* **105** (1989), no. 3, 243–266.
- [30] Li, D. and Zhang, X.: Dynamics for the energy critical nonlinear Schrödinger equation in high dimensions. *J. Funct. Anal.* **256** (2009), no. 6, 1928–1961.
- [31] Lions, P.-L.: The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1** (1984), no. 4, 223–283.
- [32] Lions, P.-L.: The concentration-compactness principle in the calculus of variations. The limit case. II. *Rev. Mat. Iberoamericana* **1** (1985), no. 2, 45–121.
- [33] Martel, Y. and Merle, F.: Instability of solitons for the critical generalized Korteweg–de Vries equation. *Geom. Funct. Anal.* **11** (2001), no. 1, 74–123.
- [34] Rey, O.: The role of the Green’s function in a nonlinear elliptic equation involving the critical Sobolev exponent. *J. Funct. Anal.* **89** (1990), no. 1, 1–52.
- [35] Ryckman, E. and Visan, M.: Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in \mathbb{R}^{1+4} . *Amer. J. Math.* **129** (2007), no. 1, 1–60.
- [36] Stein, E. M.: *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series 30, Princeton University Press, Princeton, NJ, 1970.
- [37] Strauss, W. A.: Existence of solitary waves in higher dimensions. *Comm. Math. Phys.* **55** (1977), no. 2, 149–162.
- [38] Talenti, G.: Best constant in Sobolev inequality. *Ann. Mat. Pura Appl. (4)* **110** (1976), 353–372.
- [39] Tao, T.: *Nonlinear dispersive equations*. CBMS Regional Conference Series in Mathematics 106, published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2006.
- [40] Tao, T. and Visan, M.: Stability of energy-critical nonlinear Schrödinger equations in high dimensions. *Electron. J. Differential Equations* (2005), no. 118, 28 pp.

- [41] Visan, M.: The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions. *Duke Math. J.* **138** (2007), no. 2, 281–374.
- [42] Weinstein, M. I.: Nonlinear Schrödinger equations and sharp interpolation estimates. *Comm. Math. Phys.* **87** (1982/83), no. 4, 567–576.
- [43] Weinstein, M. I.: The nonlinear Schrödinger equation – singularity formation, stability and dispersion. In *The connection between infinite-dimensional and finite-dimensional dynamical systems (Boulder, CO, 1987)*, pp. 213–232. Contemp. Math. 99, Amer. Math. Soc., Providence, RI, 1989.

Received January 12, 2021; revised December 1, 2021. Published online February 9, 2022.

Luccas Campos

Department of Mathematics, Universidade Federal de Minas Gerais (UFMG),
Belo Horizonte, MG 31270-901; and IMECC, Universidade Estadual de Campinas
(UNICAMP), Campinas, SP 13083-970, Brazil;
luccas@ime.unicamp.br

Luiz Gustavo Farah

Department of Mathematics, Universidade Federal de Minas Gerais (UFMG),
Belo Horizonte, MG 31270-901, Brazil;
farah@mat.ufmg.br

Svetlana Roudenko

Department of Mathematics and Statistics, Florida International University (FIU),
Miami, FL 33199, USA;
sroudenko@fiu.edu