

UP- AND DOWN-OPERATORS ON YOUNG'S LATTICE

RICKY INI LIU AND CHRISTIAN SMITH

ABSTRACT. The up-operators u_i and down-operators d_i (introduced as Schur operators by Fomin) act on partitions by adding/removing a box to/from the i th column if possible. It is well known that the u_i alone satisfy the relations of the (local) plactic monoid, and the present authors recently showed that relations of degree at most 4 suffice to describe all relations between the up-operators. Here we characterize the algebra generated by the up- and down-operators together, showing that it can be presented using only quadratic relations.

1. INTRODUCTION

The *up-operators* u_i for $i \in \mathbb{N}$ act on a partition λ by adding a box to the i th column of λ if the result is a partition and by sending λ to 0 otherwise. Similarly, the *down-operators* d_i act on λ by subtracting a box from the i th column if the result is a partition and by sending it to 0 otherwise. These operators were introduced as *Schur operators* by Fomin [1] and further discussed by Fomin and Greene [2] in the context of noncommutative Schur functions. They can also be seen as refinements of the raising and lowering operators U and D acting on Young's lattice as defined by Stanley [7] in his study of differential posets.

It was noted in [2] that the u_i give a representation for the *local plactic monoid* as they satisfy the relations:

$$\begin{aligned} u_i u_j &= u_j u_i & \text{for } |i - j| \geq 2, \\ u_i u_{i+1} u_i &= u_{i+1} u_i u_i, \\ u_{i+1} u_i u_{i+1} &= u_{i+1} u_{i+1} u_i. \end{aligned}$$

(In particular, the u_i satisfy the classical Knuth relations of the plactic monoid—see for instance [4].) The current authors proved in [5] (see also Meinel [6]) that the u_i also satisfy the additional degree 4 relation

$$u_{i+1} u_{i+2} u_{i+1} u_i = u_{i+1} u_{i+2} u_i u_{i+1}$$

and that this relation along with the local plactic relations characterize the algebra generated by the u_i , therein called the *algebra of Schur operators*.

It was also noted in [1] (using the fact that the down-operators can be thought of as transposes of the up-operators) that the d_i satisfy:

$$\begin{aligned} d_j d_i &= d_i d_j & \text{for } |i - j| \geq 2, \\ d_i d_{i+1} d_i &= d_i d_i d_{i+1}, \\ d_{i+1} d_i d_{i+1} &= d_i d_{i+1} d_{i+1}, \end{aligned}$$

Date: June 29, 2021.

Keywords: Schur operator, plactic monoid, Young's lattice, up- and down-operators.

R. I. Liu and C. Smith were partially supported by National Science Foundation grant DMS-1700302. R. I. Liu was also partially supported by National Science Foundation grant CCF-1900460.

and that together the u_i and d_i satisfy:

$$\begin{aligned} d_i u_j &= u_j d_i && \text{for } i \neq j, \\ d_1 u_1 &= id, \\ d_{i+1} u_{i+1} &= u_i d_i. \end{aligned}$$

In this paper we give a complete description of the algebra generated by the u_i and d_i , which we call the *algebra of up- and down-operators for Young's lattice*. Surprisingly, the following theorem shows that quadratic relations suffice to give a presentation of this algebra.

Theorem 1.1. *The algebra of up- and down-operators for Young's lattice is defined by the relations:*

$$\begin{aligned} u_i u_j &= u_j u_i && \text{for } |i - j| \geq 2, \\ d_i d_j &= d_j d_i && \text{for } |i - j| \geq 2, \\ d_i u_j &= u_j d_i && \text{for } i \neq j, \\ d_1 u_1 &= id, \\ d_{i+1} u_{i+1} &= u_i d_i. \end{aligned}$$

In particular, it follows that the local plactic relations are implied by the quadratic relations in Theorem 1.1.

In contrast, we also give a complete description of the subalgebra generated by u_t and d_t for a fixed $t > 1$ and show that it cannot be presented using relations of bounded degree.

We provide necessary background information about partitions and the up- and down-operators in Section 2. The characterization of the algebra of up- and down- operators is given in Section 3, and a discussion of subalgebras can be found in Section 4.

2. PRELIMINARIES

In this section, we discuss some background on partitions and up- and down-operators.

2.1. Partitions. A *partition* $\lambda = (\lambda_1, \lambda_2, \dots)$ of $|\lambda| = \sum_{i=1} \lambda_i$ is a sequence of nonincreasing nonnegative integers. We associate to each partition a collection of left-aligned boxes with λ_i boxes in the i th row called the *Young diagram* of λ . We define the *conjugate partition* λ' to be the partition whose Young diagram is obtained by reflecting the Young diagram of λ across the main diagonal.

We consider the partial order on partitions λ and μ such that $\mu \leq \lambda$ if and only if the Young diagram of μ fits inside the Young diagram of λ , that is, $\mu_i \leq \lambda_i$ for all i . Note that this means that if $\mu \leq \lambda$, then λ covers μ (denoted $\mu \lessdot \lambda$) if and only if λ/μ is a single box, where λ/μ is the *skew Young diagram* consisting of all boxes in λ that are not in μ . We take *Young's lattice* (\mathbf{Y}, \leq) to be the partially ordered set of partitions with the above partial order.

2.2. Words in the alphabet. Let $\mathbf{N} = \{1, 2, \dots\}$, $\overline{\mathbf{N}} = \{\overline{1}, \overline{2}, \dots\}$, and $\Gamma = \mathbf{N} \cup \overline{\mathbf{N}}$. We refer to elements $1, 2, \dots$ of \mathbf{N} as unbarred letters and elements $\overline{1}, \overline{2}, \dots$ of $\overline{\mathbf{N}}$ as barred letters.

Let $x = x_1 \cdots x_\ell$ be a word of *length* ℓ in the alphabet Γ . The *weight* of x is the vector $w(x) = (w_1(x), w_2(x), \dots)$ where

$$w_i(x) = (\text{the number of times } i \text{ appears in } x) - (\text{the number of times } \overline{i} \text{ appears in } x).$$

We also define the α -vector of x to be $\alpha(x) = (\alpha_1(x), \alpha_2(x), \dots)$ where

$$\alpha_i(x) = \max\{w_{i+1}(\tilde{x}) - w_i(\tilde{x}) \mid \tilde{x} \text{ is a suffix subword of } x\}.$$

Here a suffix subword \tilde{x} is a word of the form $\tilde{x} = x_j x_{j+1} \cdots x_\ell$ for some $1 \leq j \leq \ell + 1$. When $j = \ell + 1$, \tilde{x} is the empty word, in which case $w_{i+1}(\tilde{x}) = w_i(\tilde{x}) = 0$, so it follows that $\alpha_i(x) \geq 0$ for all i .

Example 2.1. Let $x = 11\bar{3}3\bar{2}32\bar{1}21$. Then $w(x) = (2, 1, -1, 0, \dots)$ and $\alpha(x) = (2, 0, 1, 0, \dots)$. For instance, for $\alpha_1(x) = 2$, the maximum value of $w_2(\tilde{x}) - w_1(\tilde{x})$ first occurs when $\tilde{x} = 2\bar{1}21$.

2.3. Up-operators and down-operators. Let \mathcal{U} be the free associative algebra over the complex field \mathbf{C} generated by elements u_i for $i \in \Gamma$. We will write $d_i = u_{\bar{i}}$ for all barred letters \bar{i} . For any word $x = x_1 x_2 \dots x_\ell$ in the alphabet Γ , we define $u_x = u_{x_1} u_{x_2} \cdots u_{x_\ell}$. We also use the alternate notation $i = u_i$ for $i \in \Gamma$. To avoid potential confusion in the future, we note now that $(i+j)$ denotes u_{i+j} and *not* the sum $u_i + u_j$.

Let $\mathbf{C}[\mathbf{Y}]$ be the complex vector space with basis \mathbf{Y} . We define an action of \mathcal{U} on $\mathbf{C}[\mathbf{Y}]$ in the following way. For $\lambda \in \mathbf{Y}$ and $i \in \mathbf{N}$, we let

$$u_i(\lambda) = \begin{cases} \mu & \text{if } \mu \in \mathbf{Y} \text{ and } \mu/\lambda \text{ is a single box in column } i, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$d_i(\lambda) = \begin{cases} \mu & \text{if } \mu \in \mathbf{Y} \text{ and } \lambda/\mu \text{ is a single box in column } i, \\ 0 & \text{otherwise.} \end{cases}$$

Example 2.2. Let $\lambda = (3, 1)$. Then $u_2(\lambda) = (3, 2)$, $d_3 u_2(\lambda) = (2, 2)$, but $d_1 d_3 u_2(\lambda) = 0$ since subtracting a box from the first column does not yield a partition.

$$\begin{array}{|c|c|c|} \hline & \square & \square \\ \hline \square & & \square \\ \hline \end{array} \xrightarrow{u_2} \begin{array}{|c|c|c|} \hline & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \xrightarrow{d_3} \begin{array}{|c|c|} \hline & \square \\ \hline \square & \square \\ \hline \end{array} \xrightarrow{d_1} 0$$

Note that $u_i(\lambda)$ is either 0 or a partition that covers λ in \mathbf{Y} , so we refer to u_i as an *up-operator*, and similarly we call d_i a *down-operator*. These operators were introduced by Fomin [1] under the name *Schur operators*.

The action of u_x on partitions is determined by the weight and α -vector of x as follows.

Proposition 2.3. *Let x be a word and $\lambda \in \mathbf{Y}$. Then*

$$u_x(\lambda) = \begin{cases} (\lambda'_1 + w_1(x), \lambda'_2 + w_2(x), \dots)' & \text{if } \lambda'_i - \lambda'_{i+1} \geq \alpha_i(x) \text{ for all } i, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We have $u_x(\lambda) \neq 0$ if and only if $u_{\tilde{x}}(\lambda)$ is a partition for each suffix subword \tilde{x} of x . Fix some \tilde{x} and suppose $\mu = u_{\tilde{x}}(\lambda) \neq 0$. We then have $\mu'_i = \lambda'_i + w_i(\tilde{x})$ for all i . The condition for μ to be a partition is that $\mu'_i \geq \mu'_{i+1}$ for all i , or equivalently

$$\lambda'_i + w_i(\tilde{x}) \geq \lambda'_{i+1} + w_{i+1}(\tilde{x}).$$

Rearranging this gives

$$\lambda'_i - \lambda'_{i+1} \geq w_{i+1}(\tilde{x}) - w_i(\tilde{x}).$$

By the definition of $\alpha_i(x)$, these inequalities hold for all suffix subwords \tilde{x} if and only if $\lambda'_i - \lambda'_{i+1} \geq \alpha_i(x)$. \square

The following corollary then follows from Proposition 2.3.

Corollary 2.4. *Let x and y be words. Then u_x and u_y act identically on \mathbf{Y} if and only if $\alpha(x) = \alpha(y)$ and $w(x) = w(y)$.*

Proof. The backwards implication is immediate from Proposition 2.3. For the forward direction, suppose $\alpha(x) \neq \alpha(y)$. Then we may assume without loss of generality that $\alpha_j(x) < \alpha_j(y)$ for some j . Taking λ such that $\lambda'_i - \lambda'_{i+1} = \alpha_i(x)$, we have $u_x(\lambda) \neq 0 = u_y(\lambda)$, so u_x and u_y do not act identically. If instead $\alpha(x) = \alpha(y)$ but $w(x) \neq w(y)$, then for this same choice of λ , $u_x(\lambda) \neq u_y(\lambda)$ by Proposition 2.3. \square

It was noted in [1] that u_i and d_i are transposes with respect to the basis \mathbf{Y} , which we may write as $u_i^T = d_i$. Also in [1], various relations among the u_i and d_i were described, including the local plactic relations and various quadratic relations (as described in Section 3). Our main result will be to show that in fact these quadratic relations generate all relations between these operators.

3. THE ALGEBRA OF UP- AND DOWN-OPERATORS

Let I be the two-sided ideal consisting of all elements of \mathcal{U} that annihilate $\mathbf{C}[\mathbf{Y}]$. We call the algebra \mathcal{U}/I the *algebra of up- and down-operators for Young's lattice*. Let J be the two-sided ideal generated by the following relations.

- (1) $u_i u_j \equiv u_j u_i \quad \text{for } |i - j| \geq 2,$
- (2) $d_i d_j \equiv d_j d_i \quad \text{for } |i - j| \geq 2,$
- (3) $d_i u_j \equiv u_j d_i \quad \text{for } i \neq j,$
- (4) $d_1 u_1 \equiv id,$
- (5) $d_{i+1} u_{i+1} \equiv u_i d_i.$

Our main result (Theorem 1.1) will be to show that $I = J$. We first verify that $J \subseteq I$.

Proposition 3.1. *The inclusion of ideals $J \subseteq I$ holds.*

Proof. It suffices to show that for each of (1)–(5), the two terms in the relation are in fact equivalent modulo I . We show this for relation (5); the other relations are similar. By Corollary 2.4 we need only show $\alpha(x) = \alpha(y)$ and $w(x) = w(y)$ where $x = i\bar{i}$ and $y = (i+1)\bar{(i+1)}$. Indeed, $w(x) = (0, 0, \dots) = w(y)$, while $\alpha(x) = (0, \dots, 0, 1, 0, \dots) = \alpha(y)$, where the 1 occurs in the i th position. \square

It therefore remains only to show that $I \subseteq J$. The next proposition proves that I is a *binomial ideal*, that is, I is generated by elements of the form $u_x - u_y$. The proof of this proposition is very similar to that of Proposition 3.3 in [5], but we include it here for completeness.

Proposition 3.2. *The ideal I is a binomial ideal.*

Proof. Let I' be the two-sided ideal generated by all binomials $u_x - u_y$ such that $u_x \equiv u_y \pmod{I}$, and suppose $R \in I$. Since \mathcal{U} is graded by weight and I is homogeneous with respect to weight, we may assume that all terms appearing in R have weight w for some $w = (w_1, w_2, \dots)$. We can then find $R' \equiv R \pmod{I'}$ for some

$$R' = \sum_{i=1}^n c_{x^{(i)}} u_{x^{(i)}},$$

where $x^{(i)}$ is a word in Γ of weight w , $u_{x^{(i)}} \not\equiv u_{x^{(j)}} \pmod{I}$ whenever $i \neq j$, and $0 \neq c_{x^{(i)}} \in \mathbf{C}$ for all $i \in [n]$. In particular, by Corollary 2.4, the $\alpha(x^{(i)})$ are distinct, so suppose without loss of generality that they occur in lexicographic order.

If $n \geq 1$, let $\lambda \in \mathbf{Y}$ be such that $\lambda'_k - \lambda'_{k+1} = \alpha_k(x^{(1)})$ for all k . By Proposition 2.3, $u_{x^{(1)}}(\lambda) \neq 0$. For each $i > 1$, by the lexicographic ordering, there exists some s such that $\alpha_s(x^{(1)}) < \alpha_s(x^{(i)})$. Then by Proposition 2.3, $u_{x^{(i)}}(\lambda) = 0$. Thus $0 = R'(\lambda) = c_{x^{(1)}}u_{x^{(1)}}(\lambda)$, which implies $c_{x^{(1)}} = 0$. This is a contradiction, so we must have $R' = 0$. Thus $I = I'$. \square

Our goal for the rest of this section is to show that if $u_x \equiv u_y \pmod{I}$, then $u_x \equiv u_y \pmod{J}$. Our general strategy is as follows. Let $[u_x]_I$ be the equivalence class of u_x modulo I . We will construct a representative word $[x]$ such that $u_{[x]} \in [u_x]_I$. This representative will only depend on $\alpha(x)$ and $w(x)$, so if $u_x \equiv u_y \pmod{I}$, then $[x] = [y]$. We will then show that $u_x \equiv u_{[x]} \pmod{J}$ and similarly for y , which will complete the proof.

Definition 3.3. For a word x , define

$$m(x) = \max_{i \in \mathbf{N}} \{-(\alpha_i(x) + w_i(x))\} \geq 0,$$

$$n(x) = \max \{t \in \mathbf{N} \mid t \text{ or } \bar{t} \text{ appears in } x\}.$$

For any $m \geq m(x), n \geq n(x)$, we let

$$[x]_{m,n} = (\bar{1}^m \cdots \bar{n}^m)(n^{\beta_n^m(x)} \bar{n}^{\alpha_n(x)} \cdots 1^{\beta_1^m(x)} \bar{1}^{\alpha_1(x)})$$

where $\beta_i^m(x) = \alpha_i(x) + w_i(x) + m$.

Note that the definition of m ensures that all of the exponents appearing in the definition of $[x]_{m,n}$ are nonnegative. We will often abbreviate $[x] = [x]_{m,n}$. We now show that indeed $u_{[x]} \in [u_x]_I$.

Proposition 3.4. For any word x , $u_x \equiv u_{[x]} \pmod{I}$.

Proof. Let $i \in \mathbf{N}$. Then $w_i([x]) = -m + \beta_i^m(x) - \alpha_i(x) = w_i(x)$. We now show that $\alpha_i([x]) = \alpha_i(x)$. For ease of notation, we will write $\alpha_i = \alpha_i(x)$, $w_i = w_i(x)$, and $\beta_i = \beta_i^m(x)$. Since α_i relies only upon the appearances of i , \bar{i} , $(i+1)$, and $(\bar{i}+1)$ in x , we need only consider the subword

$$\bar{i}^m \overline{(i+1)}^m (i+1)^{\beta_{i+1}} \overline{(i+1)}^{\alpha_{i+1}} i^{\beta_i} \bar{i}^{\alpha_i}.$$

To calculate $\alpha_i([x])$, we need to find the maximum value of $w_{i+1}(\tilde{x}) - w_i(\tilde{x})$ for each suffix subword \tilde{x} . This value only increases when adding an occurrence of \bar{i} or $(i+1)$ to \tilde{x} . Thus we need only verify a few choices of \tilde{x} :

$$\begin{aligned} \tilde{x} &= \bar{i}^{\alpha_i} : & w_{i+1}(\tilde{x}) - w_i(\tilde{x}) &= \alpha_i, \\ \tilde{x} &= (i+1)^{\beta_{i+1}} \overline{(i+1)}^{\alpha_{i+1}} i^{\beta_i} \bar{i}^{\alpha_i} : & w_{i+1}(\tilde{x}) - w_i(\tilde{x}) &= \beta_{i+1} - \alpha_{i+1} - \beta_i + \alpha_i \\ &&&= w_{i+1} - w_i, \\ \tilde{x} &= [x] : & w_{i+1}(\tilde{x}) - w_i(\tilde{x}) &= w_{i+1} - w_i. \end{aligned}$$

The maximum of these is just α_i . \square

We now wish to show that $u_x \equiv u_{[x]_{m,n}} \pmod{J}$ for sufficiently large m and n . To that end we will make use of the following two lemmas. As a reminder, we will use i and \bar{i} to represent u_i and d_i , respectively.

Lemma 3.5. Let $x = \bar{1} \cdots \bar{n} n \cdots 1$ for any $n \in \mathbf{N}$. Then $u_x \equiv id \pmod{J}$.

Proof. First note the equivalence

$$(6) \quad \bar{n}n(n-1)\cdots 1 \equiv (n-1)(n-2)\cdots 1 \pmod{J},$$

which holds by repeated application of (5) and a single use of (4). Then

$$\begin{aligned} \bar{1}\cdots\bar{n}n\cdots 1 &\equiv \bar{1}\cdots\overline{(n-1)}(n-1)\cdots 1 \\ &\equiv \bar{1}\cdots\overline{(n-2)}(n-2)\cdots 1 \\ &\quad \vdots \\ &\equiv id \end{aligned}$$

by repeated application of (6). \square

Lemma 3.6. *The following equivalences hold modulo J :*

$$(7) \quad u_i \equiv u_i d_i u_i,$$

$$(8) \quad d_i \equiv d_i u_i d_i,$$

$$(9) \quad u_i u_{i+1} u_i \equiv u_{i+1} u_i u_i,$$

$$(10) \quad d_i d_{i+1} d_i \equiv d_i d_i d_{i+1},$$

$$(11) \quad u_{i+1} u_i u_{i+1} \equiv u_{i+1} u_{i+1} u_i,$$

$$(12) \quad d_{i+1} d_i d_{i+1} \equiv d_i d_{i+1} d_{i+1}.$$

Proof. For (7), we have

$$\begin{aligned} n &\equiv n\bar{1}\cdots\bar{n}n\cdots 1 && \text{(Lemma 3.5)} \\ &\equiv \bar{1}\cdots\overline{(n-1)}n\bar{n}n\cdots 1 && (3) \\ &\equiv \bar{1}\cdots\overline{(n-1)}\overline{(n+1)}(n+1)n\cdots 1 && (5) \\ &\equiv \overline{(n+1)}(n+1)n\bar{1}\cdots\overline{(n-1)}(n-1)\cdots 1 && (2), (3) \\ &\equiv \overline{(n+1)}(n+1)n && \text{(Lemma 3.5)} \\ &\equiv n\bar{n}n. && (5) \end{aligned}$$

For (9) we have

$$\begin{aligned} n(n+1)n &\equiv n(n+1)\overline{(n+1)}(n+1)n && (7) \\ &\equiv n\overline{(n+2)}(n+2)(n+1)n && (5) \\ &\equiv \overline{(n+2)}(n+2)n(n+1)n && (1), (3) \\ &\equiv (n+1)\overline{(n+1)}n(n+1)n && (5) \\ &\equiv (n+1)n\overline{(n+1)}(n+1)n && (3) \\ &\equiv (n+1)n\bar{n}n\bar{n}n && (5) \\ &\equiv (n+1)n. && (7) \end{aligned}$$

For (11), assume $n \geq 2$ (the case when $n = 1$ is similar). Then

$$(\mathbf{n} + 1)\mathbf{n}(\mathbf{n} + 1) \equiv (\mathbf{n} + 1)\mathbf{n}\bar{\mathbf{n}}\mathbf{n}(\mathbf{n} + 1) \quad (7)$$

$$\equiv (\mathbf{n} + 1)\mathbf{n}(\mathbf{n} - 1)\overline{(\mathbf{n} - 1)}(\mathbf{n} + 1) \quad (5)$$

$$\equiv (\mathbf{n} + 1)\mathbf{n}(\mathbf{n} + 1)(\mathbf{n} - 1)\overline{(\mathbf{n} - 1)} \quad (1), (3)$$

$$\equiv (\mathbf{n} + 1)\mathbf{n}(\mathbf{n} + 1)\bar{\mathbf{n}}\mathbf{n} \quad (5)$$

$$\equiv (\mathbf{n} + 1)\mathbf{n}\bar{\mathbf{n}}(\mathbf{n} + 1)\mathbf{n} \quad (3)$$

$$\equiv (\mathbf{n} + 1)\overline{(\mathbf{n} + 1)}(\mathbf{n} + 1)(\mathbf{n} + 1)\mathbf{n} \quad (5)$$

$$\equiv (\mathbf{n} + 1)(\mathbf{n} + 1)\mathbf{n}. \quad (7)$$

The proofs of (8), (10), and (12) are similar to the proofs of (7), (9), and (11), respectively. \square

In particular, one can observe that (9) and (11) are *Knuth relations*, which, together with (1), verify that the quadratic relations imply that J contains all the relations of the *local plactic monoid* generated by the u_i (see [1]).

We are now ready to prove the heart of our main theorem.

Proposition 3.7. *Let $x = x_1 \cdots x_\ell$ be a word. Then there exist $M, N \in \mathbf{N}$ such that $u_x \equiv u_{[x]_{m,n}} \pmod{J}$ for all $m \geq M, n \geq N$.*

Proof. As before, we will abbreviate $[x] = [x]_{m,n}$ and $[y] = [y]_{m,n}$. We proceed by induction on the length of x . First suppose $\ell = 0$ (that is, x is the empty word), and take any $m, n \geq 0$. Then we have $[x] = \bar{1}^m \cdots \bar{n}^m n^m \cdots 1^m$ and we wish to show that $u_{[x]} \equiv id \pmod{J}$. By (1) and (11),

$$\mathbf{n} \cdots 1\mathbf{n} \equiv \mathbf{n}(\mathbf{n} - 1)\mathbf{n}(\mathbf{n} - 2) \cdots 1 \equiv \mathbf{n}\mathbf{n}(\mathbf{n} - 1) \cdots 1.$$

In other words, \mathbf{n} and $\mathbf{n} \cdots 1$ commute. Therefore

$$\begin{aligned} (\mathbf{n} \cdots 1)\mathbf{n}^{m-1} \cdots 1^{m-1} &\equiv \mathbf{n}^m((\mathbf{n} - 1) \cdots 1)(\mathbf{n} - 1)^{m-1} \cdots 1^{m-1} \\ &\equiv \mathbf{n}^m(\mathbf{n} - 1)^m((\mathbf{n} - 2) \cdots 1)(\mathbf{n} - 2)^{m-1} \cdots 1^{m-1} \\ &\quad \vdots \\ &\equiv \mathbf{n}^m \cdots 1^m, \end{aligned}$$

so $\mathbf{n}^m \cdots 1^m \equiv (\mathbf{n} \cdots 1)^m$. Similarly by (2) and (12), $\bar{1}^m \cdots \bar{n}^m \equiv (\bar{1} \cdots \bar{n})^m$. Then applying Lemma 3.5 repeatedly to

$$\bar{1}^m \cdots \bar{n}^m \mathbf{n}^m \cdots 1^m \equiv (\bar{1} \cdots \bar{n})^m (\mathbf{n} \cdots 1)^m$$

gives the claim.

Now suppose the proposition statement is true for all words of length less than ℓ . Let $x = x_1 \cdots x_\ell$ and $y = x_1 \cdots x_{\ell-1}$. By induction we know the statement holds for y for some $N', M' \in \mathbf{N}$. Then take $M = \max\{m(x), M'\}$ and $N = \max\{n(x), N'\}$ and let $m \geq M$ and $n \geq N$. By induction we have $u_x = u_y u_{x_\ell} \equiv u_{[y]} u_{x_\ell} \pmod{J}$. From this we see that it suffices to show $u_{[y]} u_{x_\ell} \equiv u_{[x]} \pmod{J}$. For ease of notation we let $\alpha_i = \alpha_i(y)$, $\beta_i = \beta_i^m(y)$, $w_i = w_i(y)$, and $\beta_i(x) = \beta_i^m(x)$ for all i .

We now split the argument into four cases depending on x_ℓ and α_i . Note that if $x_\ell = t$ or \bar{t} for $t \geq 1$, then $\alpha_i(x) = \alpha_i$, $w_i(x) = w_i$, and $\beta_i(x) = \beta_i$ for all $i \neq t, t-1$.

Case 1. Suppose $x_\ell = t$ and $\alpha_t = 0$. We have $\alpha_{t-1}(x) = \alpha_{t-1} + 1$, $w_{t-1}(x) = w_{t-1}$, $\alpha_t(x) = 0$, and $w_t(x) = w_t + 1$, so that $\beta_{t-1}(x) = \beta_{t-1} + 1$ and $\beta_t(x) = \beta_t + 1$. Then

$$\begin{aligned}
u_{[y]} u_t &\equiv \cdots \mathbf{t}^{\beta_t} (\mathbf{t}-1)^{\beta_{t-1}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots \mathbf{t} \\
&\equiv \cdots \mathbf{t}^{\beta_t} (\mathbf{t}-1)^{\beta_{t-1}} \mathbf{t} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots & (1), (3) \\
&\equiv \cdots \mathbf{t}^{\beta_t} (\mathbf{t}-1)^{\beta_{t-1}} \mathbf{t} \bar{\mathbf{t}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots & (7) \\
&\equiv \cdots \mathbf{t}^{\beta_t} (\mathbf{t}-1)^{\beta_{t-1}} \mathbf{t} (\mathbf{t}-1) \overline{(\mathbf{t}-1)}^{\alpha_{t-1}+1} \cdots & (5) \\
&\equiv \cdots \mathbf{t}^{\beta_t+1} (\mathbf{t}-1)^{\beta_{t-1}+1} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}+1} \cdots & (9) \\
&= u_{[x]}.
\end{aligned}$$

Case 2. Suppose that $x_\ell = t$ and $\alpha_t \neq 0$. We have $\alpha_{t-1}(x) = \alpha_{t-1} + 1$, $w_{t-1}(x) = w_{t-1}$, $\alpha_t(x) = \alpha_t - 1$, and $w_t(x) = w_t + 1$, so that $\beta_{t-1}(x) = \beta_{t-1} + 1$ and $\beta_t(x) = \beta_t$. Then

$$\begin{aligned}
u_{[y]} u_t &\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots \mathbf{t} \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t-1} (\mathbf{t}-1)^{\beta_{t-1}} \bar{\mathbf{t}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots & (1), (3) \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t-1} (\mathbf{t}-1)^{\beta_{t-1}+1} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}+1} \cdots & (5) \\
&= u_{[x]}.
\end{aligned}$$

Case 3. Suppose that $x_\ell = \bar{t}$ and $\alpha_{t-1} = 0$. We have $\alpha_{t-1}(x) = 0$, $w_{t-1}(x) = w_{t-1}$, $\alpha_t(x) = \alpha_t + 1$, and $w_t(x) = w_t - 1$, so that $\beta_{t-1}(x) = \beta_{t-1}$ and $\beta_t(x) = \beta_t$. Then

$$\begin{aligned}
u_{[y]} d_t &\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}} \cdots \bar{\mathbf{t}} \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t+1} (\mathbf{t}-1)^{\beta_{t-1}} \cdots & (2), (3) \\
&= u_{[x]}.
\end{aligned}$$

Case 4. Finally, suppose that $x_\ell = \bar{t}$ and $\alpha_{t-1} \neq 0$. We have $\alpha_{t-1}(x) = \alpha_{t-1} - 1$, $w_{t-1}(x) = w_{t-1}$, $\alpha_t(x) = \alpha_t + 1$, and $w_t(x) = w_t - 1$, so that $\beta_{t-1}(x) = \beta_{t-1} - 1$ and $\beta_t(x) = \beta_t$. Then

$$\begin{aligned}
u_{[y]} d_t &\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \cdots \bar{\mathbf{t}} \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}} \bar{\mathbf{t}} \cdots & (2), (3) \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}-1} (\mathbf{t}-1) \overline{(\mathbf{t}-1)} \bar{\mathbf{t}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}-1} \cdots & (10) \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}-1} \bar{\mathbf{t}} \bar{\mathbf{t}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}-1} \cdots & (5) \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t} (\mathbf{t}-1)^{\beta_{t-1}-1} \bar{\mathbf{t}} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}-1} \cdots & (8) \\
&\equiv \cdots \mathbf{t}^{\beta_t} \bar{\mathbf{t}}^{\alpha_t+1} (\mathbf{t}-1)^{\beta_{t-1}-1} \overline{(\mathbf{t}-1)}^{\alpha_{t-1}-1} \cdots & (3) \\
&= u_{[x]}.
\end{aligned}$$

This completes the proof. □

It is now easy to complete the proof of our main theorem.

Theorem 3.8. *The ideals I and J are equal. Equivalently, the algebra of up- and down-operators for Young's lattice (generated by the u_i and d_i) is determined by relations (1)–(5).*

Proof. The inclusion $J \subseteq I$ follows from Proposition 3.2. For the other direction, note that by Proposition 3.1 we need only prove that $u_x \equiv u_y \pmod{I}$ implies $u_x \equiv u_y \pmod{J}$ for words x and y . By Proposition 3.7 there exist nonnegative integers m and n sufficiently large such that $u_x \equiv u_{[x]_{m,n}} = u_{[y]_{m,n}} \equiv u_y \pmod{J}$. \square

Remark 3.9. The up- and down-operators u_i and d_i refine the raising and lowering operators U and D on Young's lattice introduced by Stanley [7] in his study of differential posets via $U = \sum_i u_i$ and $D = \sum_i d_i$. (Although these sums are infinite, only finitely many terms yield a nonzero result when applied to a particular element of $\mathbf{C}[\mathbf{Y}]$.) Therefore the main relation $DU - UD = id$ defining differential posets must be a consequence of relations (1)–(5). Indeed, we see that

$$DU - UD = \sum_{i \neq j} (d_i u_j - u_j d_i) + \sum_{i \geq 1} (d_{i+1} u_{i+1} - u_i d_i) + d_1 u_1 = id.$$

4. SUBALGEBRAS

We now turn our attention to various subalgebras generated by up- and down-operators. We briefly discuss a subalgebra studied by the authors in [5], and we introduce two other subalgebras of interest, giving a complete list of relations for each of them.

4.1. Up-operators and down-operators. We first consider the subalgebra generated by the up-operators u_i . Let \mathcal{U}' be the subalgebra of \mathcal{U} generated by u_i for $i \in \mathbf{N}$. Furthermore, let $I_{\mathcal{U}'} = I \cap \mathcal{U}'$ be the ideal of \mathcal{U}' consisting of all elements of \mathcal{U}' that annihilate \mathbf{Y} . We call $\mathcal{U}'/I_{\mathcal{U}'}$ the *subalgebra of up-operators for Young's lattice*. In [5], the present authors described this as the *algebra of Schur operators* and proved the following theorem. (See also Meinel [6].)

Theorem 4.1. *The ideal $I_{\mathcal{U}'}$ is generated by the following relations:*

$$\begin{aligned} u_i u_j &\equiv u_j u_i & \text{for } |i - j| \geq 2, \\ u_i u_{i+1} u_i &\equiv u_{i+1} u_i u_i, \\ u_{i+1} u_i u_{i+1} &\equiv u_{i+1} u_{i+1} u_i, \\ u_{i+1} u_{i+2} u_{i+1} u_i &\equiv u_{i+1} u_{i+2} u_i u_{i+1}. \end{aligned}$$

Note that most of these relations do not appear in the list of relations for the algebra of up- and down-operators, as they are implied by the quadratic relations (1)–(5) when the down-operators are included.

It is natural to also consider the subalgebra generated by the down-operators. Let \mathcal{D} be the subalgebra of \mathcal{U} generated by d_i for $i \in \mathbf{N}$, and let $I_{\mathcal{D}} = I \cap \mathcal{D}$. The *subalgebra of down-operators for Young's lattice* is then $\mathcal{D}/I_{\mathcal{D}}$. Recall that with respect to the basis \mathbf{Y} , we have $u_i^T = d_i$. Applying this transpose property to the relations in Theorem 4.1 gives the following characterization of $\mathcal{D}/I_{\mathcal{D}}$.

Theorem 4.2. *The ideal $I_{\mathcal{D}}$ is generated by the following relations:*

$$\begin{aligned} d_i d_j &\equiv d_j d_i & \text{for } |i - j| \geq 2, \\ d_i d_{i+1} d_i &\equiv d_i d_i d_{i+1}, \\ d_{i+1} d_i d_{i+1} &\equiv d_i d_{i+1} d_{i+1}, \\ d_i d_{i+1} d_{i+2} d_{i+1} &\equiv d_{i+1} d_i d_{i+2} d_{i+1}. \end{aligned}$$

4.2. u_t and d_t for fixed t . Fix some $1 < t \in \mathbf{N}$. Let \mathcal{B} be the subalgebra of \mathcal{U} generated by u_t and d_t , and consider the subalgebra $\mathcal{B}/I_{\mathcal{B}} = \mathcal{B}/(I_{\mathcal{U}} \cap \mathcal{B}) \subseteq \mathcal{U}/I_{\mathcal{U}}$. We will show that its ideal of relations $I_{\mathcal{B}}$ is generated by

$$(13) \quad u_t^{i+1} d_t^i \equiv u_t^{i+1} d_t^{i+1} u_t$$

$$(14) \quad u_t^i d_t^{i+1} \equiv d_t u_t^{i+1} d_t^{i+1}$$


for all $i \in \mathbf{N}$. Let $J_{\mathcal{B}}$ be the ideal generated by relations (13) and (14), so that we wish to show $J_{\mathcal{B}} = I_{\mathcal{B}}$.

(When $t = 1$, it is straightforward to verify that the only relation between u_1 and d_1 is (4), namely $d_1 u_1 \equiv id$, as this relation can be used to rewrite any monomial in the form $u_1^i d_1^j$, and all such monomials act independently on \mathbf{Y} .)

4.2.1. *Peaks and valleys.* One convenient way to interpret a word consisting only of the letters t and \bar{t} is as a graph of diagonal steps. More precisely, we construct a graph corresponding to a word x in the following way. Starting at the origin in the plane we read x from right to left. When we encounter a t we take a diagonal step up and to the left by adding $(-1, 1)$, and when we encounter a \bar{t} we take a diagonal step down and to the left by adding $(-1, -1)$. One must be careful since we are reading both the word and its graph from right to left.

We call a point of the graph with maximal height a *peak* and a point with minimal height a *valley*. (Peaks and valleys need not be unique.) It is straightforward to see that if (a, b) is a peak and (c, d) is a valley, then $\alpha_{t-1}(x) = b$ and $\alpha_t(x) = -d$. Also note that if (e, f) is the (leftmost) endpoint of the graph, then $w_t(x) = f$. Therefore by Corollary 2.4, the action of x on \mathbf{Y} is determined entirely by the heights of its peaks, valleys, and endpoint.

Example 4.3. The word $x = t^2 \bar{t}^4 t^3$ has the graph shown below.



This graph has a peak at $(-3, 3)$ and a valley at $(-7, -1)$. Correspondingly, $\alpha_{t-1}(x) = 3$ and $\alpha_t(x) = 1$. The leftmost point of the graph is $(-9, 1)$, so $w_t(x) = 1$.

Note that relations (13) and (14) are not bounded in degree since the only condition on i is that it be a nonnegative integer. This differs from the previous algebras that we examined in that the largest degree needed in those cases was 4 (as in the subalgebra of up-operators $\mathcal{U}'/I_{\mathcal{U}'}$). Indeed, relations of unbounded degree are required due to the following proposition.

Proposition 4.4. *The ideal $I_{\mathcal{B}}$ cannot be generated by elements of bounded degree.*

Proof. Suppose for contradiction that the largest degree appearing among the generators of $I_{\mathcal{B}}$ is $h \in \mathbf{N}$. Choose an integer $k > h$, and let $x = t^k$ and $y = t^k \bar{t}^k t^k$. Observe that $w(x) = w(y) = (0, \dots, 0, k, 0, \dots)$ and $\alpha(x) = \alpha(y) = (0, \dots, 0, k, 0, \dots)$, and so $u_x \equiv u_y \pmod{I_{\mathcal{B}}}$ by Corollary 2.4.

Note that in the graph of x , there is never a peak occurring to the right of a valley. In other words, if $x = x_1 \dots x_k$, then there do not exist $i < j$ such that $(-i, \alpha_{t-1}(x))$ and $(-j, -\alpha_t(x))$

appear in the graph of x . We will call an instance of a peak occurring to the right of a valley a *peak/valley pair*. For instance, x has no peak/valley pair but y does, corresponding to the suffix subwords t^k and $\bar{t}^k t^k$, respectively.

We now show that for words z satisfying $\alpha_{t-1}(z) + \alpha_t(z) > h$, our degree boundedness assumption implies that the existence of a peak/valley pair is invariant modulo $I_{\mathcal{B}}$. This will then lead to an immediate contradiction when applied to x and y . Let $u_m - u_{m'}$ be a generator of $I_{\mathcal{B}}$ of degree at most h . It suffices to show that if the word $z = m_1 m m_2$ has a peak/valley pair, then so does $z' = m_1 m' m_2$.

Since $u_m \equiv u_{m'} \pmod{I_{\mathcal{B}}}$, the graphs of m and m' must have their peaks, valleys, and endpoints at the same heights. Therefore z has a peak or valley within m if and only if z' has a peak or valley within m' . If z has a peak/valley pair with neither peak nor valley occurring within m , then z' has a peak/valley pair at the same locations. If at most one of the peak or valley occurs within m , say the peak, then the valley must occur within m_1 , so z' will have a peak within m' and a valley within m_1 and hence a peak/valley pair. (The other case is similar.) The only remaining possibility is if both the peak and valley occur within m (for they might switch order in m'). However, since $\alpha_{t-1}(z) + \alpha_t(z) > h$, the difference in height between the peak and valley is more than h , so they cannot both appear within m , which has length at most h . This completes the proof. \square

4.2.2. Proof of relations. We now prove that relations (13) and (14) suffice. The proofs for the following two propositions are essentially the same as the proofs of the analogous propositions in Section 3.

Proposition 4.5. *The ideal $I_{\mathcal{B}}$ is a binomial ideal.*

Proposition 4.6. *The inclusion of ideals $J_{\mathcal{B}} \subseteq I_{\mathcal{B}}$ holds.*

As in Section 3, our approach is to construct a standard equivalence class representative $u_{[x]}$ (modulo $I_{\mathcal{B}}$) for every monomial u_x and to then show that $u_x \equiv u_{[x]} \pmod{J_{\mathcal{B}}}$.

Definition 4.7. For any word x be a word in t and \bar{t} , define

$$[x] = t^{w_t(x) + \alpha_t(x)} \bar{t}^{\alpha_{t-1}(x) + \alpha_t(x)} t^{\alpha_{t-1}(x)}.$$

We say that such a word $[x]$ is the *standard representative* for x , or alternatively that it is in *standard form*.

Note that all the exponents appearing in $[x]$ are nonnegative: in particular, by the definition of $\alpha_t(x)$ we have $\alpha_t(x) \geq -w_t(x)$, and so $w_t(x) + \alpha_t(x) \geq 0$. It is straightforward to check that $w(x) = (0, \dots, 0, w_t(x), 0, \dots) = w([x])$ and $\alpha(x) = (0, \dots, 0, \alpha_{t-1}(x), \alpha_t(x), 0, \dots) = \alpha([x])$, so Corollary 2.4 implies that $[x]$ is the unique word in standard form such that $u_x \equiv u_{[x]} \pmod{I_{\mathcal{B}}}$.

Proposition 4.8. *Let $x = x_1 \cdots x_{\ell}$ be a word in t and \bar{t} . We have $u_x \equiv u_{[x]} \pmod{J_{\mathcal{B}}}$.*

Proof. We prove this by induction on the length of x . If $\ell = 0$ or if $x = \bar{t}$, then $[x] = x$, so there is nothing to prove. If $x = t$, then $[x] = t\bar{t}t$, and $t \equiv \bar{t}t$ by (13) for $i = 0$.

Now suppose the statement holds for all words shorter than x . We have that $u_x = u_{x_1} u_y$ where $y = x_2 \cdots x_{\ell}$. By induction, $u_x = u_{x_1} u_y \equiv u_{x_1} u_{[y]} \pmod{J_{\mathcal{B}}}$, so we need to show $u_{x_1} u_{[y]} \equiv u_{[x]}$.

If $x_1 = t$ and $w_t(y) < \alpha_{t-1}(y)$, then $w_t(x) = w_t(y) + 1$ while $\alpha(x) = \alpha(y)$. Hence $[x] = t[y]$, so there is nothing to show. Similarly if $x_1 = \bar{t}$ and $\alpha_t(y) = -w_t(y)$, then

$$[x] = \bar{t}^{\alpha_{t-1}(y)+\alpha_t(y)+1} t^{\alpha_{t-1}(y)} = \bar{t}[y],$$

so again there is nothing to show.

Suppose $x_1 = t$ and $w_t(y) = \alpha_{t-1}(y)$. Then $w_t(x) = w_t(y) + 1$, $\alpha_{t-1}(x) = \alpha_{t-1}(y) + 1$, and $\alpha_t(x) = \alpha_t(y)$. Here the graph of x has a new peak at its leftmost point, so $t[y]$ is not in standard form. Applying (13) with $i = w_t(y) + \alpha_t(y)$ gives

$$\begin{aligned} u_t u_{[y]} &= t^{w_t(y)+\alpha_t(y)+1} \bar{t}^{w_t(y)+\alpha_t(y)} t^{\alpha_{t-1}(y)} \\ &\equiv t^{w_t(y)+\alpha_t(y)+1} \bar{t}^{w_t(y)+\alpha_t(y)+1} t^{\alpha_{t-1}(y)+1} = u_{[x]}. \end{aligned}$$

Finally, suppose $x_1 = \bar{t}$ and $\alpha_t(y) > -w_t(y)$. We then have $w_t(x) = w_t(y) - 1$ and $\alpha(x) = \alpha(y)$. Again $\bar{t}[y]$ is not in standard form since it begins with \bar{t} . Note that by definition $\alpha_{t-1}(y) \geq w_t(y)$, so $\alpha_{t-1}(y) + \alpha_t(y) \geq w_t(y) + \alpha_t(y)$. Therefore we can apply (14) with $i = w_t(y) + \alpha_t(y) - 1 = w_t(x) + \alpha_t(x)$ to get

$$\begin{aligned} d_t u_{[y]} &= \bar{t} t^{w_t(y)+\alpha_t(y)} \bar{t}^{\alpha_{t-1}(y)+\alpha_t(y)} t^{\alpha_{t-1}(y)} \\ &\equiv t^{w_t(y)+\alpha_t(y)-1} \bar{t}^{\alpha_{t-1}(y)+\alpha_t(y)} t^{\alpha_{t-1}(y)} = u_{[x]}. \end{aligned}$$

□

Theorem 4.9. *The ideals $I_{\mathcal{B}}$ and $J_{\mathcal{B}}$ are equal.*

Proof. This follows from Propositions 4.5, 4.6, and 4.8. □

4.2.3. Up- and down-operators on finite chains. Consider again the operators u_t and d_t for some fixed $t > 1$. The action of these operators on \mathbf{Y} splits up as a direct sum of the action on chains C , where C is a set of partitions λ that have fixed values for λ'_i for all $i \neq t$. The action is then determined entirely by $\rho = \lambda'_{t-1} - \lambda'_{t+1}$, the difference between the $(t-1)$ st and $(t+1)$ st columns. (Equivalently, C is a chain with $\rho+1$ elements, and u_t and d_t act as up- and down-operators on this chain.)

Fix ρ , and let I_C be the two-sided ideal of \mathcal{B} containing all elements which annihilate C , a chain with $\rho+1$ elements. We characterize the algebra \mathcal{B}/I_C by showing that I_C is generated by the following relations:

$$(15) \quad u_t^{i+1} d_t^i \equiv u_t^{i+1} d_t^{i+1} u_t \quad \text{for } 0 \leq i \leq \rho-1,$$

$$(16) \quad u_t^i d_t^{i+1} \equiv d_t u_t^{i+1} d_t^{i+1} \quad \text{for } 0 \leq i \leq \rho-1,$$

$$(17) \quad u_t^{\rho+1} \equiv 0,$$

$$(18) \quad d_t^{\rho+1} \equiv 0.$$

Let J_C be the ideal generated by relations (15)–(18). We will show that $J_C = I_C$ by exploiting the close relationship between these ideals and $I_{\mathcal{B}}$.

Theorem 4.10. *The ideals I_C and J_C are equal.*

Proof. Recall that $I_{\mathcal{B}}$ is the two-sided ideal of \mathcal{B} containing all elements which annihilate \mathbf{Y} . Let P be the two-sided ideal of \mathcal{B} which is generated by relations (17) and (18). It is straightforward to see that $J_C = I_{\mathcal{B}} + P$ (since (13) and (14) for $i \geq \rho$ are implied by (17) and (18)), so we need to show that $I_C = I_{\mathcal{B}} + P$.

The inclusion $I_{\mathcal{B}} + P \subseteq I_{\mathcal{C}}$ holds since both (17) and (18) annihilate C . For the reverse direction, note that by Proposition 4.8, $\mathcal{B}/I_{\mathcal{B}}$ has a basis consisting of the standard representatives $u_{[x]}$. A basis element $u_{[x]}$ annihilates C if and only if the power of \bar{t} appearing in it is larger than ρ , which occurs if and only if it lies in P . The other basis elements act independently on C as in the proof of Proposition 3.2. It follows that $I_{\mathcal{C}} \subseteq I_{\mathcal{B}} + P$. \square

5. CONCLUSION

While the results of this paper and [5] have answered various questions about up- and down-operators, there still remain directions to explore on this subject. For instance, recall that the ideal of relations among the up- and down-operators is generated by relations of bounded degree (in fact, of degree 2), while some subalgebras such as $\mathcal{B}/I_{\mathcal{B}}$ cannot be presented by relations of bounded degree. It would be interesting to determine for which subalgebras this is true. In other words, can one characterize when the generating relations among a subset of operators are bounded versus unbounded in degree?

More generally, it would be interesting to explore these up- and down-operators for posets other than Young's lattice. Let P be a poset with an edge labeling from an index set I . We can define up-operators u_i for $i \in I$ such that, for $p \in P$, $u_i(p) = q$ if $p \lessdot q$ and the edge between p and q is labeled i , and otherwise $u_i(p) = 0$ if no such q exists. Note that for Young's lattice as considered above, the label between λ and μ where $\lambda \lessdot \mu$ is the column i in which the unique box of μ/λ appears. One can consider the algebras generated by these operators (or the analogously defined d_i) and try to describe their relations for other posets of interest, such as Bruhat order or absolute order on a Coxeter group. (The case of weak order leads to the study of nil-Coxeter algebras [3].) It would also be interesting if it were possible to relate structural properties of these algebras to the structure of the corresponding posets in some way.

REFERENCES

- [1] Sergey Fomin. Schur operators and Knuth correspondences. *J. Combin. Theory Ser. A*, 72(2):277–292, 1995.
- [2] Sergey Fomin and Curtis Greene. Noncommutative Schur functions and their applications. *Discrete Math.*, 193(1-3):179–200, 1998. Selected papers in honor of Adriano Garsia (Taormina, 1994).
- [3] Sergey Fomin and Richard P. Stanley. Schubert polynomials and the nil-Coxeter algebra. *Adv. Math.*, 103(2):196–207, 1994.
- [4] Alain Lascoux and Marcel-P. Schützenberger. Le monoïde plaxique. In *Noncommutative structures in algebra and geometric combinatorics (Naples, 1978)*, volume 109 of *Quad. “Ricerca Sci.”*, pages 129–156. CNR, Rome, 1981.
- [5] Ricky Ini Liu and Christian Smith. The algebra of Schur operators. *European J. Combin.*, 87:103130, 9, 2020.
- [6] Joanna Meinel. A plactic algebra action on bosonic particle configurations: The classical case. *arXiv e-prints*, January 2019. Available at arXiv:1901.00847.
- [7] Richard P. Stanley. Differential posets. *J. Amer. Math. Soc.*, 1(4):919–961, 1988.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC
Email address: `riliu@ncsu.edu`

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC
Email address: `casmit34@ncsu.edu`