

Research Article

Cite this article: Alnaqeb H, Galbreath KE, Koehler AV, Campbell ML, Jiménez FA (2022). *Citellinema* (Nematoda: Heligmosomidae) from North America with descriptions of 2 new species from the red squirrel *Tamiasciurus hudsonicus* and 1 from the Canadian woodchuck, *Marmota monax*. *Parasitology* **149**, 1199–1218. <https://doi.org/10.1017/S0031182022000737>

Received: 25 March 2022
Revised: 10 May 2022
Accepted: 12 May 2022
First published online: 27 May 2022

Keywords:

Citellinema; Heligmosomoidea; Nematoda; scientific collections; *Tamiasciurus hudsonicus*

Author for correspondence:

F. Agustín Jiménez,
Email: agustinjz@zoology.siu.edu

Citellinema (Nematoda: Heligmosomidae) from North America with descriptions of 2 new species from the red squirrel *Tamiasciurus hudsonicus* and 1 from the Canadian woodchuck, *Marmota monax*

Haitham Alnaqeb¹ , Kurt E. Galbreath² , Anson V. Koehler³,
Mariel L. Campbell⁴ and F. Agustín Jiménez¹ 

¹School of Biological Sciences, Southern Illinois University, Carbondale, Illinois 62901-6501, USA; ²Department of Biology, Northern Michigan University, Marquette, Michigan 49855, USA; ³Department of Veterinary Biosciences, Melbourne Veterinary School, The University of Melbourne, Parkville, Victoria 3010, Australia and ⁴Division of Genomic Resources, Museum of Southwestern Biology, University of New Mexico, Albuquerque, New Mexico 87131, USA

Abstract

Citellinema Hall, 1918 includes 6 valid species of gastrointestinal nematodes of sciurids. Two species occur in the Palearctic and 4 in the Nearctic, 3 of which occur minimally across Colorado, Idaho and Oregon and 1, *Citellinema bifurcatum*, has a wide distribution across North America. Members of the genus are didelphic, possess a cephalic vesicle, a terminal spine-like process in females and feature robust spicules, consisting of a proximal end fused and semicylindrical shaft connected to a lamina supported by 2 terminal filiform processes. Typically, the size of the spicules is used to differentiate species. As part of the Beringian Coevolution Project, specimens provisionally identified as *C. bifurcatum* were collected through intensive field sampling of mammals and associated parasites from across localities spanning the Holarctic. These specimens revealed considerable genetic variability at both mitochondrial and nuclear loci, supporting the identification of deeply divergent clades. Examination of these new specimens, along with the holotypes of *C. bifurcatum* and *Citellinema quadrivittati* indicates that *Citellinema monacis* (previously synonymized with *C. bifurcatum*) should be resurrected and 3 additional species described. We suggest that the apparent bifurcated nature of the spicule should be considered a generic diagnostic trait, while the proportional length of the lamina relative to that of the spicule is used as a specific character. We demonstrate the critical need for continued inventory of often poorly known assemblages of hosts and parasites, contributing to a growing baseline of archival specimens, collections and information that make explorations of faunal structure and diversity possible.

Introduction

Citellinema Hall, 1916 includes 6 species of gastrointestinal nematodes that infect squirrels. They are distributed across the Holarctic with 4 species present in North America and 2 in northeastern Eurasia. The species present in North America include *Citellinema bifurcatum* Hall, 1916; *Citellinema quadrivittati* Hall, 1916; *Citellinema columbianum* Dikmans, 1938 and *Citellinema grisei* Lichtenfels, 1971. The species present in Asia include *Citellinema nipponicum* Yamaguti, 1941 and *Citellinema orientale* Schulz, 1933, known from Japan and Siberia, respectively. No other species in the genus are known from southeast Asia, Africa, Central America and South America despite the presence of sciurids in those regions.

The lack of detailed comparative morphological and molecular studies of *Citellinema* created a gap that led taxonomists to identify most of the specimens of *Citellinema* collected across North America as *C. bifurcatum*. In an initial revision of the specimens included in the genus, Dikmans (1938) concluded that meristic similarities, especially spicule length, warranted the placement of *Citellinema sleggsi* Manter, 1930 and *Citellinema monacis* Manter, 1930 as junior synonyms of *C. bifurcatum*. Further examination of other specimens collected across North America, including individuals from 6 different sciurid species from localities in Illinois, Maryland, Michigan, Maine, Minnesota and New York in the USA, and Saskatchewan in Canada allowed Dikmans (1938) to conclude that *C. bifurcatum* has a wide geographic range and diverse host spectrum. Notably, he reported the presence of prominent prebursal papillae observed in specimens collected from a fox squirrel (*Sciurus niger* L.) from Maryland. Though these features were absent in other North American *Citellinema*, he did not propose that the specimens represented a distinct species because they were almost identical to *C. bifurcatum* in meristic traits such as the size of spicules.

In a subsequent evaluation of the specimens that served as the basis for description of *C. sleggsi* and *C. monacis* Manter (1930), and then Durette-Desset (1969) described conspicuous

variation in the number of ridges in the synlophe, which led them to consider those 2 species to be valid. Subsequently, Lichtenfels (1971) emphasized the relevance of the length of the genital cone, the size of the shaft and the configuration of the tip of the lamina as important sources of informative characters. He also emphasized the relevance of characterizing the variability of the number of ridges along the body of individuals before proposing this number as a viable character to differentiate species. Because this number could not be determined for species described based on sole individuals, he rejected the validity of *C. sleggsi* and *C. monacis* until a proper characterization of the variation of the synlophe becomes available for these species.

In part, the practice of assigning most specimens to *C. bifurcatum* reflects the emphasis of the description on the spicules and on the fact that this description was based on a single specimen mounted on a permanent slide. As a consequence, there is neither documentation of the variation in the number of ridges for the synlophe, nor there is information regarding variability in the size of spicules or in the distance from the anterior end to the deirids, nerve ring and excretory pore.

The purpose of this study is to re-examine and re-evaluate the available specimens of *Citellinema* along the Pacific coast of North America and document the species diversity using morphological and molecular approaches. This character reassessment will permit the proper identification of intra and interspecific variation, leading to increasingly robust documentation of the biodiversity present in the continent. Furthermore, we reassess these characters based on specimens with properly vouchered hosts, which is critical for ensuring that the scope of host diversity is correctly described (Galbreath *et al.*, 2019). As a result of this analysis, 2 new species of *Citellinema* from Alaska and British Columbia, collected from *Tamiasciurus hudsonicus* (red squirrel) from British Columbia, Canada and Alaska are herein described. The description of these species and the characterization of their variability and diversity are important in understanding the interactions of these parasites with their hosts, and with their environment. In the Pacific Northwest the distribution of 3 species of *Tamiasciurus* Trouessart, 1880 shows evidence of isolation across a historical biogeographic barrier in the Nass River valley (Hope *et al.*, 2016). We therefore apply a molecular phylogenetic approach to investigate if sciurid-dwelling parasites feature a similar pattern of historic isolation coincident with that of their hosts.

Materials and methods

Specimen sampling

One hundred and forty-two specimens of *Citellinema* including *C. bifurcatum*, *Citellinema quadrivittati*, *C. sleggsi* and *C. monacis* were used in this study (Table 1). Specimens were borrowed from the Museum of Southwestern Biology (MSB, University of New Mexico, Albuquerque, New Mexico) and National Museum of Natural History (USNM, Smithsonian Institution, Washington D.C., USA), Northern Michigan University (Marquette, Michigan, USA), Harold W. Manter Laboratory of Parasitology (HWML, University of Nebraska State Museum, Lincoln, Nebraska, USA) and The Cornell University Museum of Vertebrates (Ithaca, New York, USA). Most of these specimens were preserved in 70% ethanol, though some, such as holotypes, were preserved in formalin or mounted on permanent slides after being cleared with lactophenol or glycerin. All of the worms were examined under the dissection and compound microscope to confirm their identity based on appropriate characters. Upon obtaining permission for destructive sampling, a small number of specimens from each lot was cleared and dissected to observe the morphological variability of characters.

Molecular systematics

A list of the nematode specimens used for DNA extraction and sequencing is detailed in Table 2. DNeasy Blood and Tissue spin columns (Qiagen Inc., Madison, Wisconsin, USA) were used for tissues excised between the mid-body and posterior end of the worm. The rest of the worm was saved as a voucher to be preserved in the Parasite Collection of the MSB. Amplification of the mitochondrial gene cytochrome b (*CYTB*), cytochrome c oxidase subunit 1 (*COI*) and a continuous region of rDNA including internal transcribed spacer 1 (*ITS1*), 5.8S and *ITS2* (hereafter, *ITS*) was completed using primers and protocols described elsewhere (Jiménez *et al.*, 2012; Alnaqeb *et al.*, 2022).

Raw sequences were assembled in Sequencer v.3.5 (Gene Codes Corporation, Ann Arbor, Michigan, USA). Annotated sequences were complemented with sequences from relevant sister taxa published elsewhere (Zaleśny *et al.*, 2014) (Table 2), aligned and trimmed in Mesquite v.3.40 (Maddison, 2018) equipped with Clustal v.2.1 (Sievers *et al.*, 2011) applying default settings for coding genes where the invertebrate genetic code was used to identify the coding regions of mitochondrial genes. Non-coding DNA regions were aligned enforcing a gap open penalty of 15, a gap extension penalty of 6 and re-aligned by hand. Selection of the models of evolution for each dataset, description of the parameters used for the phylogenetic reconstruction using maximum likelihood (ML) and estimation of both branch support and posterior probability are described elsewhere (Alnaqeb *et al.*, 2022). A complete list of sequences generated in this study is presented in Table 3 including their accession numbers. The aligned matrixes are universally available at Open SIUC (https://opensiuc.lib.siu.edu/zoal_data/17).

Genetic distance was calculated separately for each gene as implemented in Phylogenetic Analysis Using Parsimony (PAUP*) v.4.0a (Swofford, 2002). To assess the monophyly of the new species both mtDNA and ribosomal nuclear DNA genes were analysed separately. These analyses included a reduced dataset, in which only operational taxonomic units with all 3 genes (*ITS*, *Cytb* and *COI*) were included. Species limits were established using the diagnosability of clades. These limits were compared against results of a tree-based Poisson Tree Process (PTP) model using the tree topology that included the specimens of *Citellinema* herein described and other species in the family (Alnaqeb *et al.*, 2022).

Morphological examination

Nematodes were cleared using lactophenol and compared against type and voucher specimens described in Table 1. For examination of the variation of the synlophe, transversal sections were prepared for 2 males and 2 females at mid-body, and at the anterior and posterior ends. Drawings were made using an Olympus BX50 microscope (Olympus Co. Ltd., Tokyo, Japan) equipped with a drawing tube. For morphometrical characters, the range is given first followed by the mean and coefficient of variation.

Results

Phylogeny

In the phylogram (Fig. 1) based on a combination of ribosomal nuclear DNA (*ITS*) and mtDNA (*COI* and *CYTB*), the clade of *Citellinema* is placed as the putative sister group to *Heligmosomoides* and *Heligmosomum* with a bootstrap support value of 100%. Within the clade of *Citellinema*, a western (yellow/light rectangle) and an eastern (blue/dark rectangle) clade form reciprocally monophyletic groups. Monophyly of these 2 clades is consistent in the phylogenetic trees of nDNA (Fig. 2) and mtDNA (Fig. 3). The topologies of both ML and Bayesian

Table 1. List of lots of worms examined for the completion of this study indicating their original catalogue numbers from the MSB, National Parasite Collection (USNM), HWML and Northern Michigan University (NMI)

Species	Host	Locality	Museums records
<i>Citellinema kinsellai</i>	<i>Tamiasciurus hudsonicus</i>	South Kinaskan Lake, Stewart Cassiar Highway 37, Line 5, British Columbia, Canada	MSB:Para:19047; 24859; 24858 and 24860
<i>C. kinsellai</i>	<i>T. hudsonicus</i>	South Kinaskan Lake, Stewart Cassiar Highway 37, Line 1, British Columbia, Canada	MSB:Para:19050 and 24866
<i>C. kinsellai</i>	<i>T. hudsonicus</i>	Bob Quinn Lake, Stewart Cassiar Highway 37, Line 1A, British Columbia, Canada	MSB:Para:24826
<i>C. kinsellai</i>	<i>T. hudsonicus</i>	South Kinaskan Lake, Stewart Cassiar Highway 37, Line 7, British Columbia, Canada	MSB:Para:24863
<i>C. kinsellai</i>	<i>T. hudsonicus</i>	North Bob Quinn Lake, Stewart Cassiar Highway 37, Line 1B, British Columbia, Canada	MSB:Para:24864
<i>C. kinsellai</i>	<i>Glaucomys sabrinus</i>	Huntington Wildlife Forest, Essex County, New York	USNM:1397705
<i>C. kinsellai</i>	<i>G. sabrinus</i>	Douglas Lake, Michigan	USNM:1330957
<i>Citellinema meagheri</i>	<i>T. hudsonicus</i>	St. John's Harbor, Zarembo Island, Alexander Archipelago, Petersburg Quad, Alaska	MSB:Para:25568
<i>C. meagheri</i>	<i>T. hudsonicus</i>	Salmon Run Campground, Haines, Line 1, Alaska	MSB:Para:24829, MSB:Para:24828, MSB:Para:24830
<i>C. meagheri</i>	<i>Tamias amoenus</i>	W Nez Perce Peak, Idaho County, Idaho	NMU:Para:16
<i>Citellinema</i> sp.	<i>Tamias minimus</i>	La Plata County, Colorado, USA	MSB:Para:24527
<i>Citellinema quadrivittati</i>	<i>Tamias quadrivittatus</i>	Crested Butte, Gunnison County, Colorado	USNM1324374
<i>Citellinema sleggsi</i>	<i>Sciurus carolinensis</i>	Prince George's County, Maryland	USNM:1329987
<i>Citellinema manteri</i>	<i>Marmota monax</i>	Thunder Bay, Ontario, Canada	USNM:1374038
<i>Citellinema</i> sp.	<i>T. minimus</i>	La Plata County, Colorado	MSB:Para:24527
<i>Citellinema</i> sp.	<i>Sciurus niger</i>	Alachua County, Florida, USA	USNM:1378695
<i>Citellinema</i> sp.	<i>Spermophilus armatus</i>	Jackson Hole, Teton County, Wyoming, USA	USNM:1391614
<i>Heligmosomoides bibullosus</i>	<i>Peromyscus maniculatus</i>	Klondike Highway, 22 km NW of Stewart Crossing, at mouth of Moose Creek Yukon Territory, Canada	MSB:Para:24578; 24579; 24580; 24581; 24583; 24584; 24585; 24595; 24596; 24597; 24598 and 24599
<i>H. bibullosus</i>	<i>P. maniculatus</i>	South Kinaskan Lake, Stewart Cassiar Highway 37, Line 5, British Columbia, Canada	MSB:Para:24819 and 24862
<i>H. bibullosus</i>	<i>P. maniculatus</i>	South side Dettaki Creek, British Columbia, Canada	MSB:Para:26332
<i>Heligmosomoides americanus</i>	<i>Phenacomys intermedius</i>	Cassiar Highway (Hwy 37), Burrage River Crossing, British Columbia, Canada	MSB:Para:24546
<i>H. americanus</i>	<i>P. intermedius</i>	Unknown, British Columbia, Canada	MSB:Para:26333
<i>H. americanus</i>	<i>P. intermedius</i>	1.22 km N of Tee's Creek, British Columbia, Canada	MSB:Para:26334
<i>H. americanus</i>	<i>P. intermedius</i>	N of Stikine River (E side of road), British Columbia, Canada	MSB:Para:26335
<i>Heligmosomoides bakeri</i>	<i>Mus musculus</i>	Woodland, Conway Ranch, California, USA	USNM:1367761 and 1397110
<i>Heligmosomoides johnsoni</i>	<i>P. intermedius</i>	Olympic Peninsula, Mt. Angelus Trail, Hurricane Ridge, Washington, USA	USNM:1368236
<i>Heligmosomoides thomomyos</i>	<i>Thomomys bottae</i>	Humboldt County, California, USA	USNM:1372129
<i>H. americanus</i>	<i>P. intermedius</i>	Lubrecht Forest, Missoula County, Montana, USA	USNM:1367553
<i>Heligmosomoides vandegrifti</i>	<i>Peromyscus leucopus</i>	Pennsylvania State Experimental Forest, Centre County, Pennsylvania, USA	HWML
<i>Heligmosomoides</i> sp.	<i>Peromyscus</i> sp.	NW of Helena City, Montana	HWML RZ1; RZ56; RZ62; RZ78 and RZ92
<i>Heligmosomum mixtum</i>	<i>Alticola lemminus</i>	40 km W Magadan, Magadan, Oblast, Russia	MSB:Para:26336; 26337; 26339 and 26338

(Continued)

Table 1. (Continued.)

Species	Host	Locality	Museums records
<i>H. mixtum</i>	<i>Myodes rufocanus</i>	Contact Station, 150 km NW Ust-Omchug, Magadan Oblast, Russia	MSB:Para:26341
<i>H. mixtum</i>	<i>Myodes rutilus</i>	40 km W Magadan, Magadan Oblast, Russia, Asia	MSB:Para:26343; 26344; 26347 and 26346
<i>H. mixtum</i>	<i>M. rutilus</i>	Elegan River, 5 km S Seymchan, Magadan Oblast, Russia	MSB:Para:26348
<i>H. mixtum</i>	<i>M. rutilus</i>	Kolymskaya Highway, 15 km E Susuman, Magadan Oblast, Russia	MSB:Para:26349; 26350 and 26352
<i>H. mixtum</i>	<i>M. rutilus</i>	Contact Station, 150 km NW Ust-Omchug, Magadan Oblast, Russia	MSB:Para:26357; 26358; 26359 and 26360
<i>H. mixtum</i>	<i>M. rutilus</i>	Alekchen Lakes, Magadan Oblast, Russia	MSB:Para:26361
<i>H. mixtum</i>	<i>M. rutilus</i>	Delyankir River, Sakha Republic, Russia	MSB:Para:25528 and 25531
<i>H. mixtum</i>	<i>M. rutilus</i>	Elgi River, Sakha Republic, Russia	MSB:Para:25529; 25533 and 25538
<i>H. mixtum</i>	<i>M. rutilus</i>	Indigirka River basin, 20 km NW from Ust-Nera, Sakha Republic, Russia	MSB:Para:25518; 25523; 25524 and 25532
<i>H. mixtum</i>	<i>M. rutilus</i>	Taryn-Yuryakh River, Sakha Republic, Russia	MSB:Para:25525; 25534; 25547; 25553; 25554 and 25555

Table 2. Comparative presentation of diagnostic features used to identify species of *Citellinema*

Male	<i>Citellinema bifurcatum</i> Hall, 1916	<i>Citellinema quadrivittati</i> Hall, 1916	<i>Citellinema manteri</i>	<i>Citellinema kinsellai</i>	<i>Citellinema meagheri</i>
Sources	This study	This study	This study	This study	This study
Type host	<i>Urocitellus elegans</i>	<i>Tamias quadrivittatus</i>	<i>Marmota monax</i>	<i>Tamiasciurus hudsonicus</i>	<i>Tamiasciurus hudsonicus</i>
Type locality	Waldon, Colorado	Crested Butte, Gunnison County, Colorado	Thunder Bay, Ontario, Canada	British Colombia	Alaska
Body length (mm)	–	7.24	7.42–10.45	9.35–12.81	6.83–7.42
Excretory pore	328	425	251–295	321–488	251–288
Nerve ring	165	130	200–263	214–364	150–200
Oesophagus length	695	215	669–787	576–717	503–669
Spicule length	334 and 392	623 and 627	427–507	436–489	437–507
Proportion spicules/body length (%)	–	8.6	4.85–5.75	3.8–4.6	6.3–6.8
Bifurcation from proximal end	70 and 87	73 and 81	53–84	60–97	48–76
Width of spicules (R/L)	30/33	25/29	19–20/20–22	19/18	17/14
Anterior ridges	–	–	33	31	26–29
Mid-body ridges	–	–	39	25–29	26–27
Posterior ridges	–	–	38	25	26–27

Male	<i>Citellinema bifurcatum</i>	<i>Citellinema bifurcatum</i>	<i>Citellinema orientale</i>	<i>Citellinema quadrivittati</i>	<i>Citellinema quadrivittati</i>	<i>Citellinema columbianum</i>	<i>Citellinema nipponicum</i>	<i>Citellinema grisei</i>
Sources	Hall, 1916	Sleggs, 1925	Schults, 1933	Hall, 1916	Durette-Desset, 1969	Dikmans, 1938	Yamaguti, 1941	Lichtenfels, 1971
Type host	<i>Citellus elegans</i>	<i>Citellus richardsonii</i>	<i>Eutamias asiaticus orientalis</i>	<i>Eutamias quadrivittatus</i>	<i>Glaucomys sabrinus</i>	<i>Citellus columbianus</i>	<i>Sciurus lis</i>	<i>Sciurus griseus</i>
Type locality	Waldon, Colorado	Saskatchewan, Canada	Ferner Osten, Umgebung von Nikolajewsk am Amur und Birschewaja, Russia	Crested Butte Mountain	Mill Canyon, Montana	Soda Spring, Idaho, USA	Kiso, Nagano Prefecture	Bald Mountain, Jackson County, Oregon, USA
Body length (mm)	6.8	7–12	11.2	6.21	11.5	–	12–14	8.6–14.1
Excretory pore		–	–	–	530	–	–	340–654
Nerve ring	170	–	–	130	280	–	240	203–292
Oesophagus length	535	600	–	215	735	–	540–600	480–590
Spicules	360	–	747–896	695	575	3.6 mm	940–1100	831–1060
Spicules proportion to body (%)	5.2	5–8.5	6.6–8	11.1	5	–	7.8	7.5–9.6
Anterior ridges	–	–	–	–	–	–	–	24
Mid-body ridges	–	24	20–22	–	29	–	17	23–27
Posterior ridges	–	–	–	–	–	–	–	19

Table 3. Correlational table including accession numbers available in the data repository of the National Center for Biotechnology Information, species identities and scientific collection numbers

Species	Sources	Host	Locality	ITS	COI	Cytb	28S
<i>Citellinema kinsellai</i> (MSB19047F1)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865429	MN961337	MN953025	
<i>C. kinsellai</i> (MSB19047F2)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865430	MN961338	MN953026	
<i>C. kinsellai</i> (MSB19047F3)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865431	MN961339	MN953027	
<i>C. kinsellai</i> (MSB19050F1)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865432	MN961340	MN953028	
<i>C. kinsellai</i> (MSB19050F2)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865433	MN961341	MN953029	
<i>C. kinsellai</i> (MSB24859F2)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	British Columbia, Canada	MN865434	MN961342	MN953030	
<i>Citellinema meagheri</i> (MSB25568F1)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	Alaska, Zarembo Island	MN865435	MN961343	MN953031	
<i>C. meagheri</i> (MSB25568F2)	Alnaqeb et al. (2022)	<i>Tamiasciurus hudsonicus</i>	Alaska, Zarembo Island	MN865436	MN961344	MN953032	
<i>Heligmosomoides americanus</i>	This study	<i>Phenacomys intermedius</i>	British Columbia, Canada	MT023510	MT122764	MT138452	
<i>H. americanus</i>	This study	<i>Phenacomys intermedius</i>	British Columbia, Canada	MT023511	MT122765	MT138453	
<i>H. americanus</i>	Alnaqeb et al. (2022)	<i>Phenacomys intermedius</i>	British Columbia, Canada	MN865426	MN927212	MN953021	
<i>H. americanus</i>	Alnaqeb et al. (2022)	<i>Phenacomys intermedius</i>	British Columbia, Canada	MN865427	MN927213	MN953022	
<i>H. americanus</i>	Alnaqeb et al. (2022)	<i>Phenacomys intermedius</i>	USA	KF921074	KF921077	KF921078	
<i>Heligmosomoides bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865417	MN939001	MN953012	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865418	MN939002	MN953013	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865419	MN939003	MN953014	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865420	MN939004	MN953015	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865421	MN939005	MN953016	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865422	MN939006	MN953017	
<i>H. bibullosus</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Yukon, Canada	MN865423	MN939007	MN953018	
<i>Heligmosomoides vandegrifti</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Pennsylvania	MN865424	MN927210	MN953019	
<i>H. vandegrifti</i>	Alnaqeb et al. (2022)	<i>Peromyscus maniculatus</i>	Pennsylvania	MN865425	MN927211	MN953020	
<i>Heligmosomoides bakeri</i>	GenBank	<i>Mus musculus</i>	UK	DQ408624	DQ408627	KJ994554	
<i>Heligmosomoides kurilensis kobayashii</i>	GenBank	<i>Apodemus speciosus</i>	Japan	AM409077	–	AJ971146	
<i>Heligmosomoides glareoli</i>	GenBank	<i>Myodes glareolus</i>	UK	KF765460	KF765457	KF765453	
<i>Heligmosomoides neopolygyrus</i>	GenBank	<i>Apodemus agrarius</i>	Poland	KF765458	KF765455	KJ994551	
<i>Heligmosomoides p corsicum</i>	GenBank	<i>Apodemus mystacinus</i>	Turkey	KJ994556	KJ994540	KJ994547	
<i>Heligmosomoides polygyrus</i>	GenBank	<i>Apodemus sylvaticus</i>	Belgium	DQ408618	DQ408632	AJ608863	
<i>H. polygyrus</i>	GenBank	<i>Apodemus sylvaticus</i>	Norway	KJ994557	KJ994543	KJ994553	
<i>H. polygyrus</i>	GenBank	<i>Apodemus sylvaticus</i>	Poland	KF765459	KF765456	KF765452	
<i>Heligmosomoides</i> sp.	Alnaqeb et al. (2022)	<i>Phenacomys</i> sp.	Montana, USA	MN865428	MN952981	MN953023	
<i>Heligmosomoides</i> sp.	Alnaqeb et al. (2022)	<i>Phenacomys</i> sp.	Montana, USA	–	MN952982	MN953024	

(Continued)

Table 3. (Continued.)

Species	Sources	Host	Locality	ITS	COI	Cytb	28S
<i>Heligmosomum mixtum</i>	Alnaqeb et al. (2022)	<i>Alticola lemminus</i>	Asia, Russia, Magadan Oblast	MN865437	MN939008	MN953033	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Alticola lemminus</i>	Asia, Russia, Magadan Oblast	MN865438	MN939009	MN953034	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Alticola lemminus</i>	Asia, Russia, Magadan Oblast	MN865439	MN939010	MN953035	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Alticola lemminus</i>	Asia, Russia, Magadan Oblast	MN865440	MN939011	MN953036	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Magadan Oblast	MN865441	MN939012	MN953037	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Magadan Oblast	MN865442	MN939013	MN953038	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Taryn-Yuryakh River	MN865443	MN939014	MN953039	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Taryn-Yuryakh River	MN865444	MN939015	MN953040	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Taryn-Yuryakh River	MN865445	MN939016	MN953041	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Taryn-Yuryakh River	MN865446	MN939017	MN953042	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Indigirka River basin	MN865447	MN939018	MN953043	
<i>H. mixtum</i>	Alnaqeb et al. (2022)	<i>Myodes rutilus</i>	Asia, Russia, Sakha Republic, Indigirka River basin	MN865448	MN939019	MN953044	
<i>H. mixtum</i>	GenBank	<i>Myodes glareolus</i>	Poland	KF765461	DQ408635	KF765454	
<i>Viannaia didelphis</i>	GenBank	<i>Didelphis marsupialis</i>	Panama	JX877689	JX877731	JX877722	
<i>V. didelphis</i>	GenBank	<i>Didelphis marsupialis</i>	Costa Rica	JX877688	JX877728	JX877717	
<i>Viannaia viannai</i>	GenBank	<i>Didelphis marsupialis</i>	Panama	JX877693	JX877726	JX877716	
<i>V. viannai</i>	GenBank	<i>Didelphis marsupialis</i>	Panama	JX877687	JX877725	JX877715	
<i>Hassalstrongylus</i> sp.	GenBank	<i>Calomys</i> sp.	Argentina	JX877694	JX877723	JX877713	




Fig. 1. ML phylogeny based on concatenated nDNA (*ITS1*, 5.8 and *ITS2*) and mtDNA (*CYTB* and *COI*) sequences. Numbers on branches indicate bootstrap support values (ML) followed by posterior probabilities (Bayesian) for major nodes. Tips are labelled with species names, followed by museum catalogue numbers and GenBank accession numbers as appropriate (Table 3).

inference are congruent. The average uncorrected *p*-distances and Jukes–Cantor genetic distance between both clades is 7% for mtDNA while it is 4% for *ITS*. Intraspecific genetic distance for mtDNA is 0.07% in the eastern clade (blue/dark rectangle) and

1.6% in the western clade (yellow/light rectangle); for nDNA, this distance is 0.04% in the eastern clade and 0% in the western clade. The PTP analysis revealed a support value of 0.887 for the eastern clade and 0.759 for the western clade.




Fig. 2. ML phylogeny based on the ribosomal nuclear DNA (*ITS*) sequences. Numbers on branches indicate bootstrap support values (ML) followed by posterior probabilities (Bayesian) for major nodes. Tips are labelled with species names, followed by museum catalogue numbers and GenBank accession numbers as appropriate (Table 3).




Fig. 3. ML phylogeny based on mtDNA (*CYTB* and *COI*) sequences. Numbers on branches indicate bootstrap support values (ML) followed by posterior probabilities (Bayesian) for major nodes. Tips are labelled with species names, followed by museum catalogue numbers and GenBank accession numbers as appropriate (Table 3).


Description

Citellinema kinsellai n. sp. (Figs 4–17 and 40B)

Synlophe based on 2 males and 3 females: Cuticle with longitudinal ridges; ventral ridges are larger than dorsal ridges. Male with 31 ridges (18 dorsal/13 ventral) at anterior end, 25–29 at mid-

body and 25 (15 dorsal/10 ventral) at posterior end (Figs 4–6). Females with 32 ridges (18 dorsal/14 ventral) at anterior end, 33–37 at mid-body and 36–37 at posterior end (Figs 7–9).

Holotype: Length 10.67 mm. Width at mid-body 113. Cephalic vesicle 116 long × 48 wide. Body with 5 coils. Excretory pore,



Figs 4–17. *Citellinema kinsellai* n. sp. from the red squirrel, *Tamiasciurus hudsonicus*. Synopse sections at anterior end (4), mid-body (5) and posterior end (6) of a male. Synopse sections of a female at level of oesophagus (7), mid-body (8) and posterior end (9). Scale bar for Figs 5–10 = 50 μ m. (10) Anterior end of male showing cephalic vesicle, oesophagus and nerve ring; scale bar = 100 μ m. (11) Detail of a caudal bursa; scale bar = 50 μ m. (12) Detail of the spicule; scale bar = 50 μ m. (13) Configuration of the dorsal ray; scale bar = 50 μ m. (14) Detail of the genital cone; scale bar = 50 μ m. (15) Anterior end of female showing cephalic vesicle, oesophagus and nerve ring; scale bar = 100 μ m. (16) Posterior end of female showing vulva and ovejector and 2 uterine branches; scale bar = 100 μ m. (17) Detail of the tail and terminal spine; scale bar = 50 μ m.

nerve ring and deirids located at 448, 182 and 282, from anterior end, respectively. Oesophagus 576 long and 46 wide at posterior end. Inflection of testis 350 from posterior end. Bursa is asymmetrical with 2–3 arrangement. Spicules sub-equal, proximal end vase-shaped transitions into a cylindrical shaft, lamina relatively wide near the shaft with walls that run very close from one another tapering towards the end (Fig. 12). Right spicule 468 long, 19 wide at proximal end, bifurcation 60 from proximal

end representing 12.8% of body length. Left spicule 484 long, 18 wide. Spicule length represents 4.4% of body length. Dorsal ray divided at about mid-length into 2 branches. Genital cone with 2 prominent lyre-shaped papillae.

Male paratypes based on whole mounts of 7 individuals unless otherwise noted: Length 9.35–12.81 mm (10.14 mm, 10.93%). Width at mid-body 108–125 (117, 5.38%). Cephalic vesicle 102–120 (110, 5.17%) long $\times$ 48–57 (51, 5.2%) wide. Body with

2–7 (5, 38.3%, $n = 12$) coils. Excretory pore, nerve ring and deirids located at 321–488 (432, 14.32%), 214–364 (292, 19.97%, $n = 5$) and 330 ($n = 1$), from anterior end, respectively. Oesophagus 580–717 (643, 6.46%) long and 45–55 (50, 7.46%) wide at posterior end (Fig. 10). Inflection of testis 307–389 (248, 11.78%, $n = 2$) from posterior end. Bursa asymmetrical with 2–3 arrangement (Fig. 11). Spicules subequal, with features seen in holotype; 436–489 (466, 3.95%, $n = 8$) long (Figs 12 and 40B), representing 3.8–4.6% of length of body. Width at proximal end 17–27 (20, 14.67%, $n = 20$) wide. Bifurcation 65–97 (73, 11.28%, $n = 15$) from proximal end, representing 14.9–19.8% of spicule. Dorsal ray divided at about mid-length into 2 branches (Fig. 13). Genital cone with 2 prominent lyre-shaped papillae (Fig. 14).

Allotype: 19.24 mm long and 162 wide. Body with 9 coils. Cephalic vesicle 123 long $\times$ 55 wide. Excretory pore, nerve ring and deirids located at 539, 303 and 241 from anterior end, respectively. Oesophagus 836 long, 42 wide at posterior end. Vulva 2.87 mm from posterior end. Vagina vera 63 long, vestibule 872 long, anterior sphincter 55 long and 65 wide, posterior sphincter 41 long, 55 wide. Anterior infundibulum 168 long, posterior infundibulum 171 long. Ovejector containing mature eggs 62–78 long (69, 5.1%, $n = 10$) and 38–58 wide (44, 6.3%, $n = 10$). Tail 60 long with caudal spine 16 long.

Female paratypes based on whole mounts of 14 females: Length 10.97–21.1 mm (18.47 mm, 14.41%, $n = 12$). Maximum width at mid-body 155–205 (170, 8.5%, $n = 11$). Cephalic vesicle 71–134 (109, 23%, $n = 6$) long $\times$ 54–69 (61, 10.5%, $n = 6$) wide. Body with 4–15 (9, 37%, $n = 14$) coils. Excretory pore, nerve ring and deirids located 371–544 (451, 17.5%, $n = 6$), 224–373 (287, 19.3%, $n = 5$) and 392–580 (474, 20.2%, $n = 3$) from anterior end, respectively. Oesophagus 755–866 (823, 4.8%, $n = 6$) long $\times$ 47–75 (64, 14.7%, $n = 6$) wide (Fig. 15). Vulva 2.87–3.29 mm (3.12 mm, 5.7%, $n = 6$) long from the posterior end. Vagina vera 42–76 (64, 18.8%, $n = 6$) long, vestibule 664–1059 (870, 16%, $n = 5$), anterior sphincter 45–77 (58, 19.6%, $n = 6$) long and 47–86 (67, 18.8%, $n = 6$) wide, posterior sphincter 41–71 (57, 17.3%, $n = 7$) long and 50–73 (58, 14.4%, $n = 7$) wide, anterior infundibulum 146–195 (177, 10.5%, $n = 6$) long, posterior infundibulum 136–263 (179, 22.6%, $n = 7$) long (Fig. 16). Ovejector containing mature eggs 52–78 (67, 8.4%, $n = 68$) long, 32–58 (41, 10.2%, $n = 68$) wide. Immature eggs 37–78 (55, 16.8%, $n = 231$) long, 22–51 (39, 10.53%, $n = 231$) wide. Tail 60–134 (101, 26.8%, $n = 6$) long; caudal spine 14–22 (18, 18.92%, $n = 5$) (Fig. 17).

Taxonomic summary

Type host: *Tamiasciurus hudsonicus* (red squirrel) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Symbiotype: MSB:Mamm:266934 (NK232761, collected 31 July 2013).

Type locality: Bob Quinn Lake, Stewart Cassiar Highway 37, British Columbia, Canada (56°44′37.122″N, 129°47′31.1568″W).

Other localities: South Kinaskan Lake, Stewart Cassiar Highway 37, British Columbia, Canada (57°26′33.9432″N, 130°14′23.2548″W); Bob Quinn Lake, Stewart Cassiar Highway 37, British Columbia, Canada (56°44′37.122″N, 129°47′31.1568″W); South Kinaskan Lake, Stewart Cassiar Highway 37, British Columbia, Canada (57°27′51.48″N, 130°13′51.5748″W).

Specimens deposited: Holotype (MSB:Para:32215) and 7 paratypes (MSB:Para:19047); allotype (MSB:Para:32216) and 9 paratype females (MSB:Para:19050).

Specimens examined: MSB:Para:19047, MSB:Para:19050, MSB:Para:24826, MSB:Para:24858, MSB:Para:24859, MSB:Para:24860, MSB:Para:24863, MSB:Para:24864, MSB:Para:24866, MSB:Para:24597, MSB:Para:24599, MSB:Para:24819, MSB:Para:24862; MSB:Para:26332.

Etymology: This species is named in honour of John M. Kinsella, who intensively contributes to the field of nematode taxonomy. **Zoobank Registration:** urn:lsid:zoobank.org:pub:F6C33748-B11F-4E64-8CE1-A0229F63201E.

Remarks

Citellinema kinsellai n. sp. can be immediately separated from 4 Nearctic species on the basis of the size of the spicules, the ratio of these spicules relative to the body length and the number and distribution of ridges of the male and female (Table 2). The most similar species is *C. columbianum*; however, the spicules of the latter are over 3 mm in length whereas the spicules of *C. kinsellai* are 436–529. Furthermore, the spicules in *C. columbianum* bifurcate at 150 from the proximal end, which represents 4% of the spicule length, while the spicules of *C. kinsellai* bifurcate at 64–97, representing 14.7–18.7% of the spicule length. The spicules in *C. kinsellai* are narrower compared to those in *C. bifurcatum* and in *C. quadrivittati* (Table 2). In *C. kinsellai*, the vellum that produces the pattern of bifurcation starts at the anterior conical base and runs to about the middle of the length of the spicules (Figs 12 and 40B). Conversely, the lamina between the filiform processes or walls of the spicule in *C. bifurcatum* extends further through the length of the spicules (Figs 39 and 40A). These morphological differences support the separation between *C. kinsellai* and other described species including *C. bifurcatum*.

Intraspecific genetic distance based on mtDNA is 0.07% and for nDNA it is 0.04%. Because of the morphological differences of the size, proportion and bifurcation of the spicules as well as the characterization of the synlophe and the length of body, we consider *C. kinsellai* to be a new species.


Description

Citellinema meagheri n. sp. (Figs 18–30 and 40C)

Synlophe (based on 2 males and 3 females): Cuticle with longitudinal ridges; right ventral ridges are large. Male with 26–29 ridges at anterior end, 26–27 at mid-body and 26–27 at posterior end (Figs 18–20). Females with 26–31 ridges at anterior end, 26–28 at mid-body and 27–33 at posterior end (Figs 21–23).

Holotype: Body coiled, length 6.83 mm. Width at mid-body 117. Cephalic vesicle 75 long $\times$ 47 wide. Excretory pore, nerve ring and deirids located at 288, 150 and 216 from anterior end, respectively. Oesophagus 503 long and 36 wide at posterior end. Inflection of testis 218 from posterior end. Bursa is asymmetrical with 2–3 arrangement. Spicules sub-equal, proximal end of uniform edge, shaft cylindrical, wider than it is long; lamina fusiform, limited by thick walls, walls approach each other tapering towards the end (Fig. 26). Spicules 485 and 468 long, representing 7.1% of length of body. Width at proximal end 17 and 14 for right and left spicules, respectively. Spicules bifurcate 76 from proximal end, representing 15.6% of length of spicule. Dorsal ray divided at about mid-length into 2 branches. Genital cone with 2 prominent lyre-shaped papillae.

Male paratypes based on whole mounts of 3 individuals unless otherwise noted: Length 6.83–7.42 mm (7.04 mm, 3.88%). Width at mid-body 105–139 (120, 11.81%). Cephalic vesicle 47–55 (51, 12.87%, $n = 2$) long $\times$ 47–55 (51, 7.44%, $n = 2$) wide. Body with 1–5 (3, 42.16%, $n = 5$) coils. Excretory pore and nerve ring located at 251–288 (266, 5.94%) and 150–200 (168, 13.37%) from anterior end, respectively. Oesophagus 503–669 (572, 12.27%) long and 36–48 (44, 12.09%) wide at posterior end (Fig. 24). Inflection of testis 261–277 (269, 3.07%, $n = 2$) from posterior end. Bursa is asymmetrical with 2–3 arrangement (Fig. 25). Spicules 437–507 (478, 5.34%, $n = 4$) long (Figs 26 and 40C), representing 6.3–6.8% of length of body. Width at proximal end 14–19 (16, 11.32%, $n = 7$) wide. Spicules bifurcate 48–76 (65, 18.65%) from



Figs 18–30. *Citellinema meagheri* n. sp. from the red squirrel, *T. hudsonicus*. Synlophe sections at anterior end (18), mid-body (19) and posterior ends of a male (20). Synlophe sections at level of nerve ring (21), mid-body (22) and posterior end (23) of a female; scale bar for Figs 19–24 = 50 μ m. (24) Anterior end of male showing cephalic vesicle, oesophagus, nerve ring and excretory pore; scale bar = 100 μ m. (25) Caudal bursa; scale bar = 100 μ m. (26) Detail of the spicule; scale bar = 50 μ m. (27) Genital cone; scale bar = 25 μ m. (28) Anterior end of female showing cephalic vesicle, oesophagus, position of the nerve ring; scale bar = 100 μ m. (29) Detail of the vulva and ovejector; scale bar = 100 μ m. (30) Detail of the caudal terminus; scale bar = 25 μ m.

proximal end, representing 10.9–14.9% of length of spicule. Genital cone with 2 prominent lyre-shaped papillae (Fig. 27).

Allotype: Body coiled, length 16.23 mm; maximum width 121. Cephalic vesicle 94 long $\times$ 60 wide. Excretory pore, nerve ring and deirids located at 262, 224 and 333 from anterior end, respectively. Oesophagus 682 long, 50 wide at posterior end. Vulva located at 2682 from posterior end. Vagina vera 62 long, vestibule 768, anterior sphincter 66 $\times$ 68, posterior sphincter 44 $\times$ 45,

anterior infundibulum 152 long, posterior infundibulum 125 long. Ovejector containing mature eggs 61–68 long (64, 4.16%, $n = 6$), 31–43 wide (38, 11.81%, $n = 6$). Immature eggs 47–86 (59, 17.4%, $n = 11$) long, 35–43 (39, 7.41%, $n = 11$) wide. Tail 72 long; caudal spine 21.

Female paratypes based on whole mounts of 6 individuals: Length 11.17–17.48 mm (15.36 mm, 11.3%). Width at mid-body 114–175 (145, 17.25%). Cephalic vesicle 63–114 (92, 24.5%, $n = 5$)

long $\times$ 44–58 (52, 9.77%) wide. Body with 5–12 (7, 40%, $n = 4$) coils. Excretory pore, nerve ring and deirids located at 262–414 (314, 17.5%, $n = 3$); 161–216 (197, 15.64%, $n = 1$) and 468 ($n = 1$) from anterior end, respectively. Oesophagus 639–822 (680, 11.4%, $n = 5$) long $\times$ 47–72 (56, 16.97%, $n = 5$) wide (Fig. 28). Vulva located at 2.47–2.98 mm (2.63 mm, 8.94%, $n = 4$) from posterior end. Vagina vera 50–71 (62, 12.95%, $n = 5$) long, vestibule 740–1355 (955, 24.72%, $n = 5$), anterior sphincter 48–66 (59, 10.67%, $n = 4$) long $\times$ 30–47 (41, 16%, $n = 5$), posterior sphincter 35–58 (48, 17.6%, $n = 5$) long $\times$ 40–65 (50, 20.17%, $n = 5$), anterior infundibulum 120–122 (121, 1.3%, $n = 2$) long, posterior infundibulum 136–164 (153, 7.4%, $n = 4$) long (Fig. 29). Ovejector containing mature eggs 61–101 (69, 13.88%, $n = 26$) long and 25–46 (38, 11.4%, $n = 23$) wide. Immature eggs 37–69 (57, 13.8%, $n = 64$) long and 25–47 (38, 13.6%, $n = 64$) wide. Tail 118 ($n = 1$) long; caudal spine 15–21 (19, 16.5%, $n = 2$) long (Fig. 30).

Taxonomic summary

Type host: *Tamiasciurus hudsonicus* (red squirrel) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Symbiotype: MSB:Mamm:195556 (NK152789, collected 28 July 2009).

Type locality: St. John's Harbor, Zarembo Island, Alexander Archipelago, Petersburg Quad, Alaska, USA (56°25'49.4"N, 132°58'26.8"W).

Other localities: Salmon Run Campground, Haines, Alaska (59°17'51.8"N, 135°30'54.4"W and 59°17'52.6"N, 135°30'52.8"W).

Specimens deposited: Holotype (MSB:Para:32212) and 4 paratypes (MSB:Para:25568); allotype (MSB:Para:32212) and 2 paratype females (MSB:Para:24829). Additional paratypes MSB:Para:24828; MSB:Para:24830.

Etymology: This species is named in honour of Shawn Meagher, colleague dedicated to the study of nematode evolution. *Zoobank Registration:* urn:lsid:zoobank.org:pub:F6C33748-B11F-4E64-8CE1-A0229F63201E.

Remarks

Citellinema meagheri n. sp. can be separated from the 5 Nearctic species of *Citellinema* known mainly on the basis of the variation in the number of ridges of the synlophe, their overall smaller size, the size/ratio/width of spicules and their width at the shaft. Additionally, the female can be separated from other species based on the larger length of the oesophagus and the location of the vulva from the posterior end and the size of infundibulum (Table 2). The bifurcation pattern in the spicules among *C. meagheri* is different from *C. bifurcatum* in that it is less pronounced. In *C. meagheri*, the lamina is conspicuously different from the shaft and it extends until about the middle of the length of spicules (Figs 26 and 40C). From that level and toward the posterior end, the two filiform processes that limit the lamina come very close to each other. In contrast, the lamina limited by the filiform processes in *C. bifurcatum* is more conspicuous in that it is evident towards the posterior end of the spicules (Figs 39 and 40A).

Differences between *C. meagheri* and *C. kinsellai* include the range in the number of ridges in synlophe at the anterior end of the male, which is 26–29 in *C. meagheri* and 31 in *C. kinsellai*. Additionally, the range in body size is conspicuously different (6.83–8.57 mm in *C. meagheri* compared to 9.35–12.81 mm in *C. kinsellai*). The size and ratio of spicules in *C. meagheri* (437–510) represent approximately 5.9% of the body length whereas it represents approximately 4.5% of the body length in *C. kinsellai* (436–519). The shape and configuration of the lamina in the spicules are different between *C. meagheri* and *C. kinsellai*.




Fig. 31. *Citellinema manteri* n. sp. from the Canadian woodchuck, *Marmota monax*. Micrograph of the transversal section at mid-body of a male

In females, the length of the oesophagus also differentiates specimens of *C. meagheri* from *C. grisei*, since these are 668–822 and 575–676 long, respectively. The ovejector represents approximately 7.8% of body length in *C. meagheri* while it is about 11.2% of the body length in *C. kinsellai*. These proposed morphological differences support the proposal of *C. meagheri* as a valid species.


Description

Citellinema manteri n. sp. (Figs 31–38)

Synlophe (based on 1 male and 1 female): Cuticle with longitudinal ridges. Male with 33 ridges (14 dorsal/19 ventral) at anterior end, 39 (19 dorsal/20 ventral) at mid-body and 38 (19 dorsal/19 ventral) at posterior end (Figs 31–33). Females with 24 ridges (11 dorsal/13 ventral) at anterior end, 39 (18 dorsal/21 ventral) at mid-body and 42 (21 dorsal/21 ventral) at posterior end (Figs 34–36).

Males based on 5 worms unless otherwise noted: Length 7.42–10.45 mm (8.47 mm, 13.9%); width at mid-body 139–175 (157, 13.8%). Cephalic vesicle 97–128 (112, 13.4%) long $\times$ 55–63 (59, 11.7%) wide. Body with 4–5 (4.5, 17.6%) coils. Excretory pore and nerve ring located at 251–295 (273, 34.7%, $n = 4$) and 200–263 (231, 10.7%), from anterior end, respectively. Deirids not seen. Oesophagus 669–787 (728, 10.7%, $n = 4$) long and 48–55 (51, 11.9%, $n = 4$) wide at posterior end (Fig. 37). Inflection of testis 277 ($n = 1$) from posterior end. Bursa is asymmetrical with 2–3 arrangement (Fig. 38). Spicules sub-equal, conical proximal end and tapering shaft; lamina features conspicuous filiform processes that run posteriad gradually approaching each other. Spicules 427–507 (467, 13%, $n = 7$) long, 19–22 wide at proximal end; spicular length represents 4.85–5.75% of length of body. Spicules bifurcated at 53–84 (71, 20%, $n = 4$) from proximal end, representing 12.4–16.5% of length of spicule. Dorsal rays and genital cone unknown.

Females based on 5 individuals unless otherwise noted: Length 16.65–19.88 mm (17.76 mm, 7.43%), maximum width 162–194 (180, 7.6%). Cephalic vesicle 89–124 (102, 12%, $n = 3$) long $\times$ 64–69 (66, 4.5%, $n = 3$) wide. Body with 9–12 (9, 13.3%) coils. Excretory pore and nerve ring located at 544–582 (564, 2.9%, $n = 4$) and 232–288 (256, 9.2%) from anterior end, respectively.



Figs 32–38. *Citellinema manteri* n. sp. synlophe sections at anterior (32) and posterior (33) ends of a male; scale bar = 50 μ m. Sections of the synlophe at anterior end, close to the level of excretory pore (34), mid-body (35) and posterior end (36) of a female; scale bar = 50 μ m. (37) Anterior end of a male showing cephalic vesicle, oesophagus and nerve ring; scale bar = 100 μ m. (38) Detail of the bursa; scale bar = 200 μ m.

Deirids not seen. Oesophagus 684–876 (755, 13.8%, $n = 3$) long $\times$ 63–77 (71, 7.7%, $n = 4$) wide. Vulva located at 3.41 mm ($n = 1$) from posterior end. Vagina vera 43–45 (58, 2.5%, $n = 3$) long, vestibule 299 ($n = 1$), anterior sphincter 66 ($n = 1$) long and 89–90 (90, 0.74%, $n = 2$) wide, posterior sphincter 64–82 (73, 14.3%, $n = 2$) long and 56 ($n = 1$) wide, anterior infundibulum 90–122 (106, 17.6%, $n = 2$) long, posterior infundibulum 65 ($n = 1$) long. Ovejector containing mature eggs 62–84 (73, 8.9%, $n = 27$) long and 34–48 (41, 8.4%, $n = 27$) wide. Immature eggs 39–48 (43, 9.8%, $n = 3$) long and 33–36 (35, 4.7%, $n = 3$) wide. Tail 153–225 (189, 18.8%, $n = 3$) long with caudal spine 18–21 (13, 18.3%, $n = 2$) long.

Taxonomic summary

Host: *Marmota monax* (Canadian woodchuck) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Locality: Thunder Bay, Ontario, Canada.

Specimens examined: 10 syntypes (USNM:1374038 = USNPC:078608.01).

Etymology: This species is named in honour of Dr Harold W. Manter, founder of the parasitological collection of the University of Nebraska State Museum and pioneer in the taxonomic assessment of *Citellinema*. **Zoobank Registration:** urn:lsid:zoobank.org:pub:F6C33748-B11F-4E64-8CE1-A0229F63201E.

Remarks

Based on the number of cuticular ridges in the synlophe the species is different from all other species in the genus (Table 2). At mid-body the number of ridges is 39 for both male and female; the species with greater similarities to *C. manteri* would include

C. monacis, which features 41 ridges at mid-body. However, the spicules of the latter range between 280 and 330, whereas the spicules in *C. manteri* range between 427 and 507. Furthermore, the distances of the anterior end to excretory pore are other morphological characters that can be used to differentiate *C. manteri* from the other species of the genus as shown in Table 2. Finally, the right spicule of *C. manteri* appears narrower than all the examined species (Table 2).

Dikmans (1938) considered *C. monacis* to be a junior synonym of *C. bifurcatum*. His interpretation was based on the limited morphological characteristics offered by Manter (1930). Subsequent researchers supported the use of the name *C. bifurcatum* for parasites found in the woodchuck, citing the lack of clarity in the solution of this taxonomic problem (Fleming et al., 1979). Our characterization allows the comparison of intraspecific morphological variability. It is apparent that parasites infecting woodchucks feature a greater number of ridges in the synlophe than any of the other species thus far characterized. We posit that the differences in the number of ridges and spicule size among worms collected in Maine and Thunder Bay, Ontario represent different species. Size variation of the spicule and total body length resulting from long-term preservation has been documented even for species in this genus; nevertheless, the differences are seldom equivalent to 1/3 of the length of the spicule as observed in this case. Consequently, we reject the proposed hypothesis of *C. monacis* being a junior synonym of *C. bifurcatum* posited by Dikmans (1938).

Redescriptions

Citellinema bifurcatum Hall, 1916 (Figs 39 and 40A).

Synlophe: Unknown.




Fig. 39. Spicule of *Citellinema bifurcatum* from the Wyoming ground squirrel, *Urocitellus elegans*.

Redescription of holotype based on slide consisting of anterior and posterior ends: Length unknown. Approximate width at mid-body 166. Cephalic vesicle 38 wide. Excretory pore and nerve ring situated at 328 and 165, from anterior end, respectively. Deirids not seen. Oesophagus 695 long and 70 wide at posterior end. Inflection of testis not evident. Bursa is asymmetrical with 2–3 arrangement. Spicules unequal, manubrium or proximal end cup-shaped; proximal end fused with cylindrical shaft; lamina starts immediately after shaft, characterized by the presence of 2 bifurcating filiform processes or walls. Lamina divided into 2 parts: the first part is as long as the proximal end and it is limited by conspicuous filiform processes or walls; the second part is characterized by these filiform processes or walls gradually approaching each other towards the distal end (Fig. 39). Right spicule 334 long, bifurcation starting 70 from proximal end, representing 20.9% of length of spicule (Figs 39 and 40A). Left spicule 392 long, with bifurcation starting 87 from proximal end, 22.1% of length of spicule. Width of right and left spicules at proximal end 30 and 33, respectively. Dorsal rays and genital cone not seen.

Female: Unknown.

Taxonomic summary

Type host: *Spermophilus elegans* (= *Urocitellus elegans*) (Wyoming ground squirrel) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Type locality: Waldon, Colorado, USA.

Specimens examined: A single slide of holotype (USNM:1324365 = USNPC:016176.01).

Remarks

The holotype is the single specimen available from the type series as designated by Hall (1916). The present re-description indicated slight meristic differences in the body size, length of the oesophagus and size of spicules. Hall (1916) provided measurements for body size, length of the oesophagus and spicules size, which are 6.8 mm, 535 and 360 (5.2% of the body length), respectively. Since the specimen is fragmented, it was not possible to measure the length of the body, yet we measured a slightly longer oesophagus, 695, and subequal spicules of sizes 334 and 392, respectively. The combined length of the body for the 2 fragments (parts) of the anterior and posterior ends mounted on the slide was 8.21 mm. It is uncertain if the 2 fragments mounted on the slide represent the complete parts and size of the total length of the

holotype, or if there is another dissected part that was missing from the specimen before it was mounted.

This specimen is different from other species in the size and width of spicules as shown in Table 2. Because no additional specimens are available other than the holotype we were unable to examine the characteristics of the synlophe in specimens collected from the host and type locality. However, Durette-Desset (1969) listed the number of ridges in the synlophe in the mid-body of specimens identified as *C. bifurcatum* that originated from the typical host, these range from 16 to 20 ridges, while it is 20–22 in *C. orientale*, 29 in *C. quadrivittati*, 17 in *C. nipponicum* and 23–27 in *C. grisei*. Because of the lysis of the holotype specimens of *C. bifurcatum* slide, we were unable to observe any ridges along the body of the holotype. One should bear in mind that the characterization of the synlophe of *C. bifurcatum* presented by Durette-Desset (1969) was conducted at mid-body, but it was not clearly stated if these sections were made using type material. Finally, reconstructing and describing the characteristics of the posterior end, including the bursa, was difficult, since the specimen is mounted in a permanent medium.

Redescription

Citellinema quadrivittati (Hall, 1916) Manter, 1930

Synlophe: Unknown.

Redescription based on holotype slide: Length 7.24 mm. Width at mid-body 169. Cephalic vesicle 60 wide. Excretory pore and nerve ring situated at 425 and 130 from anterior end, respectively. Oesophagus 215 long and 17 wide at posterior end. Inflection of testis not observed. Bursa is asymmetrical with 2–3 arrangement. Spicules, sub-equal; right and left spicules 623 and 627 long, representing 8.6% of length of body. Proximal end vase shaped followed by cylindrical shaft; lamina bifurcates forming 2 long filiform processes that run parallel to each other towards the distal end of the lamina; one of these processes ends slightly before the other. Bifurcation of lamina 73 from proximal end, representing 11.7% of length of right spicule; in left spicule bifurcation starts 81 from proximal end, representing 12.9% of length. Width of right and left proximal processes 25 and 29, respectively. Dorsal rays and genital cone not seen.

Females: Unknown.

Taxonomic summary

Synonyms: *Warrenius quadrivittati* Hall, 1916.

Type host: *Tamias quadrivittatus* (Colorado chipmunk) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Type locality: Crested Butte, Gunnison County, Colorado, USA.

Specimens examined: A single slide of holotype (USNM:1324374 = USNPC:016185.01).

Remarks

Hall (1916) proposed *Warrenius* Hall, 1916 based on *Warrenius quadrivittati* Hall, 1916. The morphological traits used to establish the genus included the size of spicules and the presence of unilateral cervical membrane at the anterior end of the single male examined. The nomenclatural history of *Citellinema* and *Warrenius* was efficiently summarized by Dikmans (1938). Relative to the diagnostic feature, the presence of the cervical ala is perceived as an artefact due to the width of the cephalic vesicle, which makes it apparently thicker on 1 of the sides, yet thin on the opposite side.

Based on our observations, the type of *C. quadrivittati* shows slight morphological differences from the data presented in the original description of Hall (1916); in particular in body size, width and spicules' length. Body length, width and spicules are



Fig. 40. Comparison of the spicules of (A) *C. bifurcatum*, (B) *C. kinsellai* and (C) *C. meagheri*; scale bar = 50 μ m.

7.24 mm; 169 and 623–627 in our re-description compared to 6.21 mm, 112 and 695 reported by Hall (1916).

Furthermore, Durette-Desset (1969) examined a specimen of *Citellinema* sp. isolated in *Glaucomys sabrinus* from Montana; this specimen is very similar to *C. quadrivittati*, yet the specimen from Montana differs from the holotype on being larger (11.5 mm) and possessing shorter spicules (570 and 575); the proportion of spicule/body length of 5% is however, relatively close to the range of 8–11% established for the holotype of *C. quadrivittati* (Table 2). The examination of fresher material that allows the establishment of the size variation of the body, the spicules and their proportions should provide an additional morphological element to separate *C. quadrivittati* from *C. bifurcatum* as by Hall (1916).

Reidentification

Citellinema sleggsi Manter, 1930

The specimens of this lot were determined by a donor to the USNPC. The examination of the material allowed us to determine that they do not belong to *Citellinema* because the females lack the characteristic cephalic cap and spine; furthermore, the spicules do not feature thick walls limiting the lamina. As a consequence, we cannot make the relevant comparisons against the rest of the species in the genus, precluding the assessment of the independence of this species relative to *C. bifurcatum*.

Taxonomic summary

Synonyms: *Warrenius bifurcatus* Sleggs, 1925.

Host: *Sciurus carolinensis* (eastern grey squirrel) (Rodentia, Sciuridae).

Site of infection: Small intestine.

Locality: Bowie, Prince George's County, Maryland, USA.

Specimens studied: 3 males and 3 females deposited in (USNM:1329987 = USNPC:027851.01).

Discussion

The origins of *C. kinsellai* and *C. meagheri* may be closely tied to geographic isolation that also drove speciation in their sciurid hosts. Some of the specimens used in this analysis were collected from localities that are separated by the Nass River, which cuts through the coastal mountains of British Columbia and was shown to be a significant barrier limiting the distribution of species of red squirrels (Hope *et al.*, 2016). It appears that the parasites of these squirrels may also be limited by this geographical barrier given the strong divergence between the *C. kinsellai* and *C. meagheri* clades (Figs 1–3) at both mtDNA and nDNA loci.

The lack of *Citellinema* specimens available for DNA isolation limited the scope of our phylogenetic analysis. Inclusion of specimens of *C. bifurcatum* and other species such as *C. orientale*, *C. quadrivittati*, *C. columbianum*, *C. nipponicum* and *C. grisei* is important for reconstructing the phylogeny of the genus and family; this phylogeny will enable us to better understand the history

of diversification in this group. Most specimens deposited in scientific museums are old and either preserved as permanent slide mounts or in formalin. Furthermore, those collections that do exist commonly are represented by a few individual specimens that have been archived, and geographic sampling is poor. Therefore, fresh specimens collected through spatially extensive and site-intensive sampling are required to resolve the relationships among species of the genus *Citellinema*.

Following the standards of the times, the original description of the type species is incomplete, lacking important morphological characteristics such as the synoppe that were later applied in diagnoses of these strongyles. In the earlier part of the 20th century taxonomists relied on the size of the spicules as the main morphological characteristic to differentiate species. This practice was changed by Durette-Desset (1969) and Lichtenfels (1971) who attempted to standardize morphological descriptions by adding information about variability of the synoppe along the body and other structures omitted in the original descriptions.

In addition to discrete synoppe morphologies, we demonstrated considerable levels of genetic divergence between *C. kinsellai* and *C. meagheri*. Although the divergence in the nDNA genes was substantially lower than that in the mtDNA genes, it is consistent with the patterns of genetic divergence observed in other groups of nematodes, and genetic distances in the range of 3–4% for mtDNA have been suggested as rough benchmarks for interspecific divergence (Hebert *et al.*, 2003; Mayer *et al.*, 2007; Pyziel *et al.*, 2017).

In the recent past, scientists underestimated the diversity of strongyles by not recognizing the predictive value of some portions of the spicules. The spicules in *Citellinema* appear to be uniform in that there is no conspicuous separation between the proximal end, shaft and lamina, and the narrow lamina is surrounded by walls that confer it the appearance of being bifurcated. Our observations revealed that the proportion of the lamina relative to the proximal end and shaft may be a useful trait to diagnose species in this genus. This determination was made possible by the proper preservation of specimens deposited in scientific collections, even when these specimens were expected to belong to *C. bifurcatum*, which was historically assumed to be a geographically widespread taxon. We suspect that this expectation of finding an extremely common parasite in squirrels from across North America caused many researchers to make assumptions regarding the identity of nematodes collected from squirrels. Reports of *C. bifurcatum* that are not supported by properly archived voucher specimens should therefore be viewed with caution. This phenomenon, dubbed the ‘fallacy of expected identification’ (Hoberg and Soudachanh, 2021) prevents an accurate documentation of diversity, since no species identity can be tested nor assumed in the absence of the materialistic evidence of the voucher.

Biological collections and specimen archives are the foundation for documenting and understanding patterns and distribution of diversity for complex parasite–host assemblages and for tracking the outcomes accelerating climate and environmental changes in the biosphere (Dunnum *et al.*, 2017; Colella *et al.*, 2021). Our results were possible because primary materials for species of *Citellinema* were largely derived from extensive field collections linking Alaska, Siberia and regions of the continental USA under the Beringian Coevolution Project over the past 2 decades (Cook *et al.*, 2005, 2016; Hoberg *et al.*, 2012), which include vouchered records for both parasite and host specimens. These sampling efforts provide holistic information regarding mammal–parasite assemblages across spatial and temporal scales, and have been instrumental in building an increasingly nuanced picture and synthesis for the history of diversity and faunal assembly at northern latitudes during the Late Pliocene

and Quaternary (Hoberg *et al.*, 2012; Haas *et al.*, 2020). Especially critical are archives and repositories with sufficient scope and depth to reveal the limits and occurrence of cryptic diversity as demonstrated in our explorations of *Citellinema* (e.g. Hoberg *et al.*, 2015).

Data. The aligned matrices are universally available at Open SIUC (https://opensiuc.lib.siu.edu/zoool_data/17). Specimens were returned to the MSB and the USNM. Sequences were deposited in GenBank.

Acknowledgements. We are grateful to Drs J. A. Cook and E. P. Hoberg at the MSB who organized and facilitated extensive field inventories in Alaska, Siberia and the continental USA under the Beringian Coevolution Project. Dr Sara Brant facilitated access to the materials in MSB. Dr Jerzy Behnke provided specimens collected in the UK. Collectors whose tears, sweat and blood made these collections possible include J. W. Brunt, J. A. Cook, N. Dawson, L. Delehanty, N. Dokuchaev, A. Hope, A. Lahzuhtkin, S. O. MacDonald, A. Runck, A. Tsvetkova and M. Westover. Drs K. Neubig, E. A. Zieman and Mr C. Williams provided technical assistance and feedback.

Author contributions. H. A. and F. A. J. conceived and designed the study. K. E. G. and F. A. J. helped securing funding. K. E. G., A. V. K. and M. L. C. conducted fieldwork, archived specimens and documented parasite presence. H. A. conducted data gathering. H. A. and F. A. J. performed phylogenetic analyses. H. A., K. E. G., A. V. K., M. L. C. and F. A. J. wrote the article.

Financial support. Alnaqeb was funded by the Saudi Arabian Cultural Mission (SACM). Funding for this research was possible through a grant in aid by the Annual Midwestern Conference of Parasitologists (AMCOP), collection was enabled by the Beringian Coevolution Project funded by NSF-DEB 0196095, 0415668 and 1258010; USDA Forest Service and US Fish and Wildlife Service contracts to J. A. Cook and NSF-DEB 1256943 to K. E. Galbreath. Writing supported in part by NSF-DUE1564969.

Conflict of interest. The authors declare there are no conflicts of interest.

Ethical standards. Collecting and handling of mammals was done in accordance with the approved Animal Care and Use Committee of Southern Illinois University, USA (Protocol 21-017, Assurance Number D16-00044).

References

- Alnaqeb H, Greiman S, Vandegrift KJ, Campbell ML, Meagher S and Jiménez FA (2022) A molecular reconstruction of Holarctic Heligmosomidae reveals a new species of *Heligmosomoides* (Nematoda: Heligmosomidae) in *Peromyscus maniculatus* (Neotominae) from Canada. *Systematics and Biodiversity* 20, 1–19. doi: 10.1080/14772000.2022.2046199.
- Colella JP, Bates J, Burneo SF, Camacho MA, Carrion Bonilla C, Constable I, D’Elia G, Dunnum JL, Greiman S and Hoberg EP (2021) Leveraging natural history biorepositories as a global, decentralized, pathogen surveillance network. *PLoS Pathogens* 17, e1009583.
- Cook JA, Hoberg EP, Koehler A, Henttonen H, Wickström L, Haukisalmi V, Galbreath K, Chernyavski F, Dokuchaev N and Lahzuhtkin A (2005) Beringia: intercontinental exchange and diversification of high latitude mammals and their parasites during the Pliocene and Quaternary. *Mammal Study* 30, S33–S44.
- Cook JA, Galbreath KE, Bell KC, Campbell ML, Carrière S, Colella JP, Dawson NG, Dunnum JL, Eckerlin RP and Fedorov V (2016) The Beringian coevolution project: holistic collections of mammals and associated parasites reveal novel perspectives on evolutionary and environmental change in the North. *Arctic Science* 3, 585–617.
- Dikmans G (1938) A consideration of the nematode genus *Citellinema* with description of a new species, *Citellinema columbianum*. *Proceedings of the Helminthological Society of Washington* 5, 55–58.
- Dunnum JL, Yanagihara R, Johnson KM, Armien B, Batsaikhan N, Morgan L and Cook JA (2017) Biospecimen repositories and integrated databases as critical infrastructure for pathogen discovery and pathobiology research. *PLoS Neglected Tropical Diseases* 11, e0005133.
- Durette-Desset MC (1969) Remarques sur un *Citellinema* sp., Nématode Trichostrongylide parasite d’un *Glaucomys sabrinus* en Californie. *Bulletin du Muséum National d’Histoire Naturelle* 41, 940–945.

- Fleming W, Georgi J and Caslick J (1979) Parasites of the woodchuck (*Marmota monax*) in central New York State. *Proceedings of the Helminthological Society of Washington* **46**, 115–127.
- Galbreath KE, Hoberg EP, Cook JA, Armien B, Bell KC, Campbell ML, Dunnun JL, Dursahinhan AT, Eckerlin RP, Gardner SL, Greiman SE, Henttonen H, Jiménez FA, Koehler AVA, Nyamsuren B, Tkach VV, Torres-Pérez F, Tsvetkova A and Hope AG (2019) Building an integrated infrastructure for exploring biodiversity: field collections and archives of mammals and parasites. *Journal of Mammalogy* **100**, 382–393.
- Haas GM, Hoberg EP, Cook JA, Henttonen H, Makarikov AA, Gallagher SR, Dokuchaev NE and Galbreath KE (2020) Taxon pulse dynamics, episodic dispersal and host colonization across Beringia drive diversification of a Holarctic tapeworm assemblage. *Journal of Biogeography* **47**, 2457–2471.
- Hall MC (1916) Nematode parasites of mammals of the orders Rodentia, Lagomorpha, and Hyracoidea. *Proceedings of the United States National Museum* **50**, 1–258.
- Hebert PDN, Cywinska A, Ball SL and DeWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London, Series B: Biological Sciences* **270**, 313–321.
- Hoberg EP and Soudachanh KM (2021) Diversity of Tetrabothriidae (Eucestoda) among Holarctic Alcidae (Charadriiformes): resolution of the *Tetrabothrius jagerskioldi* cryptic species complex – cestodes of Alcinae – provides insights on the dynamic nature of tapeworm and marine bird faunas under the Stockholm paradigm. *MANTER: Journal of Parasite Biodiversity* **16**, 1–76. doi: 10.32873/unl.dc.manter16
- Hoberg EP, Galbreath KE, Cook JA, Kutz SJ and Polley L (2012) Northern host–parasite assemblages: history and biogeography on the borderlands of episodic climate and environmental transition. *Advances in Parasitology* **79**, 1–97.
- Hoberg EP, Agosta SJ, Boeger WA and Brooks DR (2015) An integrated parasitology: revealing the elephant through tradition and invention. *Trends in Parasitology* **31**, 128–133.
- Hope AG, Malaney JL, Bell KC, Salazar-Miralles F, Chavez AS, Barber BR and Cook JA (2016) Revision of widespread red squirrels (genus: *Tamiasciurus*) highlights the complexity of speciation within North American forests. *Molecular Phylogenetics and Evolution* **100**, 170–182.
- Jiménez FA, Gardner SL, Navone GT and Ortí G (2012) Four events of host-switching in Aspidoderidae (Nematoda) involve convergent lineages of mammals. *Journal of Parasitology* **98**, 1166–1175.
- Lichtenfels JR (1971) *Citellinema grisei* sp. n. (Nematoda: Trichostrongylidae) from the western gray squirrel, *Sciurus griseus*. *Proceedings of the Helminthological Society of Washington* **38**, 257–261.
- Maddison W (2018) Mesquite: a modular system for evolutionary analysis. Version 3.51. Available at <http://www.mesquiteproject.org>.
- Manter HW (1930) Two new nematodes from the woodchuck, *Marmota monax canadensis*. *Transactions of the American Microscopical Society* **49**, 26–33.
- Mayer F, Dietz C and Kiefer A (2007) Molecular species identification boosts bat diversity. *Frontiers in Zoology* **4**, 4.
- Pyziel AM, Laskowski Z, Demiaszkiewicz AW and Höglund J (2017) Interrelationships of *Dictyocaulus* spp. in wild ruminants with morphological description of *Dictyocaulus cervi* n. sp. (Nematoda: Trichostrongyloidea) from red deer, *Cervus elaphus*. *Journal of Parasitology* **103**, 506–518.
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M and Söding J (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, 7, 539.
- Swofford D (2002) PAUP* [Phylogenetic Analysis Using Parsimony (and other methods)]. Version 4. Sunderland, Sinauer Associates, 4.
- Zaleśny G, Hildebrand J, Paziewska-Harris A, Behnke JM and Harris PD (2014) *Heligmosomoides neopolygyrus* Asakawa & Ohbayashi, 1986, a cryptic Asian nematode infecting the striped field mouse *Apodemus agrarius* in Central Europe. *Parasites & Vectors* **7**, 457–466.