

DOMAINS OF ATTRACTION OF INVARIANT DISTRIBUTIONS OF THE INFINITE ATLAS MODEL

BY SAYAN BANERJEE^a AND AMARJIT BUDHIRAJA^b

*Department of Statistics and Operations Research, University of North Carolina, ^asayan@email.unc.edu,
^bbudhiraj@email.unc.edu*

The infinite Atlas model describes a countable system of competing Brownian particles where the lowest particle gets a unit upward drift and the rest evolve as standard Brownian motions. The stochastic process of gaps between the particles in the infinite Atlas model does not have a unique stationary distribution and in fact for every $a \geq 0$, $\pi_a := \bigotimes_{i=1}^{\infty} \text{Exp}(2 + ia)$ is a stationary measure for the gap process. We say that an initial distribution of gaps is in the weak domain of attraction of the stationary measure π_a if the time averaged laws of the stochastic process of the gaps, when initialized using that distribution, converge to π_a weakly in the large time limit. We provide general sufficient conditions on the initial gap distribution of the Atlas particles for it to lie in the weak domain of attraction of π_a for each $a \geq 0$. The cases $a = 0$ and $a > 0$ are qualitatively different as is seen from the analysis and the sufficient conditions that we provide. Proofs are based on the analysis of synchronous couplings, namely, couplings of the ranked particle systems started from different initial configurations, but driven using the same set of Brownian motions.

1. Introduction. Rank-based diffusions comprise systems of Brownian particles where the drift and diffusivity of each particle depends on its relative rank in the system. Such diffusions are given by a system of stochastic differential equations (SDE) for $m \in \mathbb{N}_0 \cup \{\infty\}$, of the following form:

$$(1.1) \quad dY_i(s) = \sum_{j=0}^m a_j \mathbf{1}(Y_i(s) = Y_{(j)}(s)) ds + \sum_{j=0}^m b_j \mathbf{1}(Y_i(s) = Y_{(j)}(s)) dW_i(s),$$

for $s \geq 0, i \in \{0, \dots, m\}$. Here, a_j and b_j , respectively, denote the drift and diffusion coefficients of the rank j particle, $\{W_i(\cdot) : i \in \{0, \dots, m\}\}$ are independent standard Brownian motions, $\mathbf{Y}(0) = \{Y_i(0)\}_{i=0}^m$ is a $\mathbb{R}^m$ -valued random variable independent of the above Brownian motions, and $\{Y_{(i)}(\cdot) : i \in \{0, \dots, m\}\}$ are the associated ranked (ordered) processes defined in an appropriate way. Rank-based diffusions have gained significant attention in recent years as models in Stochastic Portfolio Theory [11, 12], scaling limits of interacting particle systems like the simple exclusion process [20], and their connection with nonlinear diffusion processes [9, 19]. Rank-based diffusions also have close connections with Aldous’ “Up the river” stochastic control problem [1], which was recently solved in [29].

The finite-dimensional setting $m < \infty$ is reasonably well-understood as in that case the system of gaps between the ranked particles can be described by a finite-dimensional reflected Brownian motion (RBM). One can then apply the general theory of RBM [14, 15, 32] to study the process in a pathwise sense. In particular, it is known that the system in (1.1) has a unique weak solution [4]. Moreover, the solution exists in a strong sense and is pathwise unique until the first time of a triple collision (three particles at the same location) [17], and

Received March 2021; revised October 2021.

MSC2020 subject classifications. Primary 60K35, 82C22, 60J60; secondary 60J55.

Key words and phrases. Atlas model, interacting diffusions, infinite dimensional diffusions, time averaged occupancy measures, synchronous couplings, local time, ergodicity.

general criteria are known for the almost sure absence of triple collisions [16, 18, 26]. In particular, for the finite Atlas model, that is, the case where $a_j = \mathbf{1}(j = 0)$, $b_j \equiv 1$ for $j = 0, 1, \dots, m$, these criteria give unique strong solutions for (1.1). Further, under very general stability and nondegeneracy conditions on the drift and diffusion coefficients $\{a_j\}$, $\{b_j\}$, the RBM describing gaps between the ranked particles has a unique stationary measure [14]. Geometric ergodicity results can be found in [6] and rates of convergence to stationarity, depending explicitly on $\{a_j\}$, $\{b_j\}$ have recently been obtained [2, 3].

Much less is known for the case $m = \infty$. Owing to the nonstandard structure of the drift and diffusion coefficients, the infinite-dimensional rank-based diffusions are technically much more challenging to analyze and even some basic questions like uniqueness in law of the reflected process describing gaps between the ranked particles remain open in full generality. Moreover, unlike its finite-dimensional analogue, the gap process for infinite rank-based diffusions can have multiple stationary measures (like the model considered below) and the domains of attraction of these stationary measures are far from being well-understood. See [7, 8, 10, 27] and references therein for comprehensive surveys of the known results in the $m = \infty$ case.

In this article, we study the long time behavior of the infinite Atlas model which, like its finite-dimensional analogue, is a rank-based diffusion where, at any point of time, the lowest particle has a unit upward drift and the remaining particles perform independent standard Brownian motions. This is a special case of the $\mathbb{R}^\infty$ -valued process $\{Y_i(\cdot) : i \in \mathbb{N}_0\}$ defined in (1.1) with $m = \infty$ and $a_j = \mathbf{1}(j = 0)$, $b_j \equiv 1$ for $j \in \mathbb{N}_0$. Hence, $\{Y_i(\cdot) : i \in \mathbb{N}_0\}$ satisfies the following system of SDE:

$$(1.2) \quad dY_i(s) = \mathbf{1}(Y_i(s) = Y_{(0)}(s)) ds + dW_i(s), \quad s \geq 0, i \in \mathbb{N}_0.$$

It was shown in [27], Theorem 3.2, that, if the starting points of the infinite Atlas model almost surely satisfy

$$(1.3) \quad \sum_{i=0}^{\infty} e^{-\alpha Y_i(0)^2} < \infty \quad \text{for all } \alpha > 0,$$

then the system (1.2) has a weak solution that is unique in law. If (1.3) is satisfied, it can be shown that, almost surely, the system is locally finite in the sense that for any $T > 0$, $u > 0$, there exist only finitely many $i \in \mathbb{N}_0$ such that $\min_{s \in [0, T]} Y_i(s) \leq u$ [27], Lemma 3.4. Thus, almost surely, the ranking $(Y_0(t), Y_1(t), \dots) \mapsto (Y_{(0)}(t), Y_{(1)}(t), \dots)$ is well defined for all $t \geq 0$ by prescribing the following convention for ties: if there exist $t \geq 0$ and $i < j$ such that $Y_i(t) = Y_j(t)$, we assign a lower rank to Y_i and higher rank to Y_j at time t . We will assume throughout this article that $0 \leq Y_0(0) \leq Y_1(0) \leq Y_2(0) \leq \dots$.

By [27], Lemma 3.5, the processes defined by

$$(1.4) \quad B_i^*(t) := \sum_{j=0}^{\infty} \int_0^t \mathbf{1}(Y_j(s) = Y_{(i)}(s)) dW_j(s), \quad i \in \mathbb{N}_0, t \geq 0,$$

are independent standard Brownian motions which can be used to write down the following stochastic differential equation for $\{Y_{(i)}(\cdot) : i \in \mathbb{N}_0\}$:

$$(1.5) \quad dY_{(i)}(t) = \mathbf{1}(i = 0) dt + dB_i^*(t) - \frac{1}{2} dL_{i+1}^*(t) + \frac{1}{2} dL_i^*(t), \quad t \geq 0, i \in \mathbb{N}_0.$$

Here, $L_0^*(\cdot) \equiv 0$ and for $i \in \mathbb{N}$, $L_i^*(\cdot)$ denotes the local time of collision between the $(i-1)$ th and i th particles, that is, the unique nondecreasing continuous process satisfying $L_i^*(0) = 0$ and $L_i^*(t) = \int_0^t \mathbf{1}(Y_{(i-1)}(s) = Y_{(i)}(s)) dL_i^*(s)$ for all $t \geq 0$. The gap process in the infinite Atlas model is the $\mathbb{R}_+^\infty$ -valued process $\mathbf{Z}(\cdot) = (Z_1(\cdot), Z_2(\cdot), \dots)$ defined by

$$(1.6) \quad Z_i(\cdot) := Y_{(i)}(\cdot) - Y_{(i-1)}(\cdot), \quad i \in \mathbb{N}.$$

We will be primarily interested in the long time behavior of the gap process. Thus, without loss of generality, we will assume $Y_{(0)}(0) = 0$. Denote by $\mathcal{S}_0$ the class of probability measures on $\mathbb{R}_+^\infty$ such that the corresponding $\mathbb{R}_+^\infty$ -valued random variable $\mathbf{Y}(0) = (Y_i(0))_{i \in \mathbb{N}_0}$ satisfies (1.3) and

$$0 = Y_0(0) \leq Y_1(0) \leq \cdots \quad \text{a.s.}$$

Given a measure $\gamma \in \mathcal{S}_0$, from weak existence and uniqueness, we can construct a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ on which are given mutually independent $\mathcal{F}_t$ -Brownian motions $\{W_i(\cdot), i \in \mathbb{N}_0\}$ and $\mathcal{F}_t$ adapted continuous processes $\{Y_i(t), t \geq 0, i \in \mathbb{N}_0\}$ such that Y_i solve (1.2) with $\mathbb{P} \circ \mathbf{Y}(0)^{-1} = \gamma$. On this space the processes $Y_{(i)}$ and Z_i are well defined and the distribution $\mathbb{P} \circ \mathbf{Z}(\cdot)^{-1}$ is uniquely determined from γ . Furthermore, $\mathbf{Z}$ is a Markov process with values in (a subset of) $\mathbb{R}_+^\infty$. Let

$$\mathcal{S} = \{\mathbb{P} \circ \mathbf{Z}(0)^{-1} : \mathbb{P} \circ \mathbf{Y}(0)^{-1} \in \mathcal{S}_0\}.$$

Note that $\mu \in \mathcal{S}$ if for some $\mathbb{R}_+^\infty$ -valued random variable $\mathbf{Y}(0)$ with probability law in $\mathcal{S}_0$, the vector $(Y_1(0) - Y_0(0), Y_2(0) - Y_1(0), \dots)$ has probability law μ .

It was shown in [25] that the distribution $\pi := \bigotimes_{i=1}^\infty \text{Exp}(2)$ (which is clearly an element of $\mathcal{S}$) is a stationary distribution of the gap process $\mathbf{Z}(\cdot)$. It was also conjectured there that this is the unique stationary distribution of the gap process. Surprisingly, this was shown to be not true in [28] who gave an uncountable collection of stationary distributions in $\mathcal{S}$ for the gaps in the infinite Atlas model defined as

$$(1.7) \quad \pi_a := \bigotimes_{i=1}^\infty \text{Exp}(2 + ia), \quad a \geq 0.$$

The ‘maximal’ stationary distribution π_0 (in the sense of stochastic domination) is somewhat special in this collection as described below. For this reason, in what follows, we will continue using the notation π for the measure π_0 .

For a $\nu \in \mathcal{S}$, we will denote by $\hat{\nu}_t$ the probability law of $\mathbf{Z}(t)$, when $\mathbb{P} \circ \mathbf{Z}(0)^{-1} = \nu$. Obviously, when $\nu = \pi_a$, $\hat{\nu}_t = \pi_a$ for all $t \geq 0$. In general, one expects that under suitable conditions on $\nu \in \mathcal{S}$, $\hat{\nu}_t$ converges (say in the weak convergence topology on the space $\mathcal{P}(\mathbb{R}_+^\infty)$ of probability measures on $\mathbb{R}_+^\infty$) to one of the stationary measures π_a , $a \geq 0$. When that happens (i.e., $\hat{\nu}_t \rightarrow \pi_a$ weakly as $t \rightarrow \infty$ for some $a \geq 0$), we say that ν is in the *Domain of Attraction* (DoA) of π_a . Characterizing the domain of attraction of the above collection of stationary measures has been a longstanding open problem. It was shown in [27], Theorem 4.7, using comparison techniques with finite-dimensional Atlas models that if the law ν of the initial gaps $\mathbf{Z}(0)$ stochastically dominates $\pi = \pi_0$ (in a coordinatewise sense), then ν is in DoA of π , that is, the law of $\mathbf{Z}(t)$ converges weakly to π as $t \rightarrow \infty$ (see also Lemma 6.1). Recently, [8] established a significantly larger domain of attraction for π using relative entropy and Dirichlet form techniques. They showed that a $\nu \in \mathcal{S}$ is in DoA of π if the random vector $\mathbf{Z}(0)$ with distribution ν almost surely satisfies the following conditions for some $\beta \in [1, 2)$ and eventually nondecreasing sequence $\{\theta(m) : m \geq 1\}$ with $\inf_m \{\theta(m-1)/\theta(m)\} > 0$:

$$(1.8) \quad \limsup_{m \rightarrow \infty} \frac{1}{m^{\beta\theta(m)}} \sum_{j=1}^m Z_j(0) < \infty,$$

$$(1.9) \quad \limsup_{m \rightarrow \infty} \frac{1}{m^{\beta\theta(m)}} \sum_{j=1}^m (\log Z_j(0))_- < \infty,$$

$$(1.10) \quad \liminf_{m \rightarrow \infty} \frac{1}{m^{\beta^2/(1+\beta)\theta(m)}} \sum_{j=1}^m Z_j(0) = \infty,$$

with the additional requirement that $\theta(m) \geq \log m$, $m \geq 1$, if $\beta = 1$. This is a big leap in our understanding of the domain of attraction properties of the infinite Atlas model. However, the conditions (1.8)–(1.10) involve upper and lower bounding the growth rate of the starting points $Y_m(0) = \sum_{j=1}^m Z_j(0)$ with respect to m in terms of a common parameter $\beta \in [1, 2)$, which partially restricts its applicability. In particular, they do not cover all initial gap distributions that stochastically dominate π (e.g., $Z_j(0) \sim e^{j^2}$), which were shown to be in DoA of π in [27]. Moreover, (1.9) is not satisfied if even one of the gaps is zero, which intuitively should not drastically influence ergodic properties of the model. This condition arises from the relative entropy methods used in [8] (see, e.g., the estimate (3.5) therein) which make an important use of the fact that the gaps are nonzero.

Beyond the results presented above, little is known about the domain of attraction of stationary measures of the infinite Atlas model. Especially, for the other stationary measures π_a , $a > 0$, nothing is known about the domain of attraction. To investigate the latter question, the techniques of [8] can no longer be applied in a straightforward manner, as we now explain, and one needs new ideas. The methods in [8] involve approximating the infinite Atlas model by the finite Atlas model with $d + 1$ particles, given by the solution to the SDE (1.2) with $\mathbb{N}_0$ replaced with $\{0, 1, \dots, d\}$. A key step in the proof is to argue that the law of the first k gaps, as $t \rightarrow \infty$ and $d \rightarrow \infty$ simultaneously, in a suitable fashion, converges to the k -marginal of π (i.e., $\bigotimes_{i=1}^k \text{Exp}(2)$). It is then shown by a natural coupling argument that, on any compact time interval $[0, T]$, there is a $d_T \in \mathbb{N}$ such that the ranked particles in the d_T -dimensional Atlas model stay uniformly close to the lowest $d_T + 1$ particles in the infinite Atlas model on $[0, T]$. A crucial fact that is exploited in the proof is that the *unique* stationary measure of the finite Atlas model with $d + 1$ particles, given by $\pi^{(d)} := \bigotimes_{i=1}^d \text{Exp}(2(1 - i/(d + 1)))$, converges to π as $d \rightarrow \infty$. Since the finite Atlas model has a unique stationary distribution, a finite-dimensional approximation approach of the form used in [8] cannot work in addressing convergence to π_a for $a \neq 0$, as this approximation is *designed* to select π_0 . One may consider other types of finite-dimensional approximations to the infinite Atlas model (see Remark 2.7), however they present different challenges with implementing the approach as in [8] for showing convergence to π_a for $a \neq 0$. These are further discussed in Remark 2.7.

In the current work, we consider a somewhat weaker formulation of domain of attraction of the stationary measures π_a , $a \geq 0$. Specifically, for $\nu \in \mathcal{S}$, define for $t > 0$

$$\nu_t \doteq \frac{1}{t} \int_0^t \hat{\nu}_s ds,$$

where $\hat{\nu}_t$ is as introduced in the paragraph below (1.7). We say that a $\nu \in \mathcal{S}$ is in the *Weak Domain of Attraction* (WDoA) of π_a for $a \geq 0$, if $\nu_t \rightarrow \pi_a$ weakly as $t \rightarrow \infty$. Note that, although the convergence of the time-averaged laws ν_t to π_a (ergodic limit) is implied by the convergence of $\hat{\nu}_t$ to π_a (marginal time limit), the converse is not clear. However, if $\nu \in \mathcal{S}$ is in the WDoA of π_a for some $a > 0$, it is *not* in the DoA of $\pi = \pi_0$. In this article we will provide sufficient conditions for a measure $\nu \in \mathcal{S}$ to be in the WDoA of π_a for a general $a \geq 0$.

In Theorem 2.1 and Corollary 2.2, we describe a large set of measures $\nu \in \mathcal{S}$ that are in the weak domain of attraction of $\pi = \pi_0$. The sufficient condition for ν to be in the WDoA of π is a condition similar to (1.10) with $\beta = 1$ (see (2.1)). In particular, the sufficient condition in (2.1) is implied by (1.10) if the initial gap sequence $\{Z_j(0)\}$ satisfies $\limsup_{n \rightarrow \infty} Z_n(0) < \infty$ a.s. (see Remark 2.3). We do not require upper bounds on growth rates of $m \mapsto Y_m(0) = \sum_{j=1}^m Z_j(0)$ or $m \mapsto \sum_{j=1}^m (\log Z_j(0))_-$ of the form in (1.8), (1.9). In particular, unlike the condition in (1.9), the sufficient condition we provide does not require $Z_j(0)$ to be nonzero. Furthermore, the sufficient condition in (2.1) is satisfied by all gap distributions which stochastically dominate π . In Corollary 2.2 and Remark 2.3, we provide

additional examples where the sufficient condition (2.1) is satisfied and make some comparisons with the sufficient conditions given in [8].

In Theorem 2.4 and Corollary 2.5, we provide sufficient conditions for $\nu \in \mathcal{S}$ to be in the weak domain of attraction of the alternate stationary measures π_a , $a > 0$. This condition is formulated in terms of growth rates of the L^1 distance between the first m coordinates of the initial gap process and that of a coupled random variable $\mathbf{V}_a$, with distribution π_a , as $m \rightarrow \infty$. In particular, this sufficient condition holds if the initial gap sequence is given as a perturbation $\Theta = (\Theta_i)_{i \in \mathbb{N}}$ of $\mathbf{V}_a$ which satisfies

$$(1.11) \quad \sum_{i=1}^d |\Theta_i| = o\left(\frac{\log d}{\log \log d}\right) \quad \text{and} \quad \limsup_{d \rightarrow \infty} \frac{|\Theta_d|}{dV_{a,d}} < \infty.$$

Note that, for $a, a' > 0$, the L^1 -distance between the first d coordinates of the stationary measures π_a and $\pi_{a'}$ grows at rate $O(|a - a'| \log d)$ as $d \rightarrow \infty$ and thus the size of the perturbation allowed in (1.11) is not very far from what one expects. See Remark 2.6 for further discussion of this point. To the best of our knowledge, this is the first result on convergence properties for these alternate stationary measures. In Corollary 2.5 and Remark 2.6, we provide some examples of measures $\nu \in \mathcal{S}$ in WDoA of π_a for $a > 0$.

Our proofs are based on a pathwise approach to studying the long time behavior of the infinite Atlas model using *synchronous couplings*, namely, two versions of the infinite Atlas model described in terms of the ordered particles via (1.5) and started from different initial conditions but driven by the same collection of Brownian motions. Central ingredients in the proofs are suitable estimates on the decay rate of the L^1 distance between the gaps in synchronously coupled ordered infinite Atlas models. These estimates are obtained by analyzing certain excursions of the difference of the coupled processes where each excursion ensures the contraction of the L^1 distance by a fixed deterministic amount. The key is to appropriately control the number and the lengths of such excursions. Certain monotonicity properties of synchronous couplings (see Proposition 3.1), and a quantification of the influence of far away coordinates on the first few gaps (see Section 4), also play a crucial role. Using these tools, the discrepancy between the gap processes of two synchronously coupled infinite Atlas models can be controlled in terms of associated gap processes when the starting configurations differ only in finitely many coordinates. The latter are more convenient to work with as for them the initial L^1 distance between the gap processes is finite, and the aforementioned excursion analysis can be applied to obtain our main results.

This article is organized as follows. In Section 2, we state the main results. In Section 3, we collect some crucial monotonicity properties of synchronous couplings including the monotonicity and quantitative control of the L^1 distance between them. In Section 4, we quantify the influence of far away coordinates on the first k gaps of the infinite Atlas model. This is crucial in reducing the problem of convergence to stationarity from arbitrary starting gap configurations to those that are perturbations of the stationary gaps at finitely many coordinates. In Section 5, we identify excursions in the paths of the synchronously coupled gap processes that result in reductions of the L^1 distance between them by fixed amounts. Finally, in Section 6, the main results are proved by analyzing these excursions of synchronously coupled gap processes from carefully chosen starting configurations.

Notation. For a vector $\mathbf{v} = (v_1, v_2, \dots)^T \in \mathbb{R}_+^\infty$ and $m \in \mathbb{N}$, we will write $\mathbf{v}|_m := (v_1, \dots, v_m)^T$. We will write $s(\mathbf{v})$ for the new vector in $\mathbb{R}_+^\infty$ whose i th coordinate is $v_1 + \dots + v_i$, for $i \in \mathbb{N}$. For two vectors $\mathbf{v}_1$ and $\mathbf{v}_2$ in $\mathbb{R}_+^m$, where $m \in \mathbb{N} \cup \{\infty\}$, $\mathbf{v}_1 \leq \mathbf{v}_2$ will be used to denote that each coordinate of $\mathbf{v}_1$ is less than or equal to the corresponding coordinate of $\mathbf{v}_2$. Similarly, $\mathbf{v}_1 \wedge \mathbf{v}_2$ and $\mathbf{v}_1 \vee \mathbf{v}_2$ will, respectively, denote the coordinatewise minimum and maximum.

We will denote by $C([0, \infty) : \mathbb{R}_+^\infty)$ the space of all continuous $\mathbb{R}_+^\infty$ -valued functions defined on $[0, \infty)$. This space is equipped with the usual local uniform topology, namely a sequence $f^{(m)} \in C([0, \infty) : \mathbb{R}_+^\infty)$ converges to $f \in C([0, \infty) : \mathbb{R}_+^\infty)$ if

$$\lim_{m \rightarrow \infty} \sup_{t \in [0, T]} |f_j^{(m)}(t) - f_j(t)| = 0 \quad \text{for any } T \in (0, \infty) \text{ and } j \in \mathbb{N}.$$

In the sequel, for a collection of events $\{E_t : t \in [0, \infty)\}$, we will often write “ E_t holds for all $t \geq 0$ ” to mean “almost surely, E_t holds for all $t \geq 0$.” $\Phi(\cdot)$ will denote the normal CDF and $\bar{\Phi}(\cdot) := 1 - \Phi(\cdot)$. We will write $X \sim \mu$ to denote that the random variable X has distribution μ . For any measure ν on $\mathbb{R}_+^\infty$ and any $k \in \mathbb{N}$, we call the measure $\nu^{(k)}$ on $\mathbb{R}_+^k$ defined by $\nu^{(k)}(A) := \nu(A \times \mathbb{R}_+^\infty)$, $A \in \mathcal{B}(\mathbb{R}_+^k)$, as the k -marginal of ν . By a coupling of two probability measures θ_1 and θ_2 on some Polish space S , we mean S -valued random variables X_1, X_2 on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mathbb{P} \circ X_i^{-1} = \theta_i$ for $i = 1, 2$. Coupling of more than two probability measures is defined in a similar manner. For $m \in \mathbb{N} \cup \{\infty\}$ and probability measures θ_1 and θ_2 on $\mathbb{R}_+^m$ we say θ_2 stochastically dominates θ_1 if $\theta_2(-\infty, \mathbf{x}] \leq \theta_1(-\infty, \mathbf{x}]$ for all $\mathbf{x} \in \mathbb{R}_+^m$. In this case, by Strassen’s theorem [23], there is a coupling (X, Y) of (θ_1, θ_2) such that $X_i \leq Y_i$ a.s. for all i .

2. Main results. Our first theorem gives a sufficient condition for a measure on $\mathbb{R}_+^\infty$ to be in the WDoA of π .

THEOREM 2.1. *Suppose that the probability measure μ on $\mathbb{R}_+^\infty$ satisfies the following: there exists a coupling $(\mathbf{U}, \mathbf{V})$ of μ and π such that, almost surely,*

$$(2.1) \quad \liminf_{d \rightarrow \infty} \frac{1}{\sqrt{d}(\log d)} \sum_{i=1}^d U_i \wedge V_i = \infty.$$

Then $\mu \in \mathcal{S}$ and it belongs to the WDoA of π .

The following corollary gives some natural situations in which (2.1) holds.

COROLLARY 2.2. *Let μ be a probability measure on $\mathbb{R}_+^\infty$ and $\mathbf{U} \sim \mu$. Suppose one of the following conditions hold:*

(i) *μ is stochastically dominated by π and, almost surely,*

$$(2.2) \quad \liminf_{d \rightarrow \infty} \frac{1}{\sqrt{d}(\log d)} \sum_{i=1}^d U_i = \infty.$$

(ii) *For some a.s. finite random variable M*

$$(2.3) \quad \liminf_{d \rightarrow \infty} \frac{1}{\sqrt{d}(\log d)} \sum_{i=1}^d (U_i \wedge M) = \infty, \quad \text{a.s.}$$

(iii) *$U_i = \lambda_i \Theta_i$, $i \in \mathbb{N}$, where $\{\Theta_i\}$ are i.i.d. nonnegative random variables satisfying $\mathbb{P}(\Theta_1 > 0) > 0$, and $\{\lambda_i\}$ are positive deterministic real numbers satisfying one of the following:*

- (a) $\liminf_{j \rightarrow \infty} \lambda_j > 0$,
- (b) $\limsup_{j \rightarrow \infty} \lambda_j < \infty$ and

$$(2.4) \quad \liminf_{d \rightarrow \infty} \frac{1}{\sqrt{d}(\log d)} \sum_{i=1}^d \lambda_i = \infty.$$

Then (2.1) holds. Hence, in all the above cases, $\mu \in \mathcal{S}$ and is in the WDoA of π .

REMARK 2.3. We make the following observations:

(a) Suppose that π is stochastically dominated by μ . Then (2.1) holds and so $\mu \in \mathcal{S}$ and is in the WDoA of π . In fact in this case [27], Theorem 4.7 (cf. Lemma 6.1) shows that μ is in the DoA of π .

(b) Note that for any $C \in (0, \infty)$,

$$((2C) \wedge 1)[U_i \wedge (1/2)] \leq U_i \wedge C \leq ((2C) \vee 1)[U_i \wedge (1/2)].$$

Hence, if (2.3) holds for one a.s. finite random variable M then it holds for every such random variable. In particular, if $\limsup_{n \rightarrow \infty} U_n < \infty$ a.s. and (2.2) is satisfied then (2.1) holds and so $\mu \in \mathcal{S}$ and is in the WDoA of π .

(c) The paper [8] notes two important settings where conditions (1.8)–(1.10) are satisfied. These are: (i) for some $c \in [1, \infty)$, $\lambda_j \in [c^{-1}, c]$ for all $j \in \mathbb{N}$ and $Z_j(0) = \lambda_j \Theta_j$ where Θ_j are i.i.d. with finite mean and such that $E \log(\Theta_1)_- < \infty$, and (ii) $Z_j(0) = \lambda_j \Theta_j$ where Θ_j are i.i.d. exponential with mean 1 and, either $\lambda_d \downarrow 0$ and $\frac{1}{\sqrt{d} \log d} \sum_{i=1}^d \lambda_i \rightarrow \infty$, as $d \rightarrow \infty$, or $\lambda_d \uparrow \infty$ and $\limsup_{d \rightarrow \infty} \frac{1}{d^\beta} \sum_{i=1}^d \lambda_i < \infty$ for some $\beta < 2$. We note that conditions assumed in the above settings are substantially stronger than the one assumed in part (iii) of Corollary 2.2. However, [8] establishes the stronger marginal time convergence as opposed to an ergodic limit considered in the present paper. See part (g) on a related open problem.

(d) In the setting of (iii) the sequence $\{\lambda_i\}$ satisfies (2.4) if $\lambda_i \geq \varphi_i \log i / \sqrt{i}$ for sufficiently large i , for any nonnegative sequence $\varphi_i \rightarrow \infty$ as $i \rightarrow \infty$. Indeed, note that, for any $C > 0$, there exists $i_C \in \mathbb{N}$ such that for all $i \geq i_C$,

$$\sum_{j=i_C}^i \lambda_j \geq C \sum_{j=i_C}^i \frac{\log j}{\sqrt{j}} \geq C \int_{i_C}^{i+1} \frac{\log z}{\sqrt{z}} dz.$$

Using the fact that the primitive of $\log z / \sqrt{z}$ is $2\sqrt{z}(\log z - 2)$, $\{\lambda_j\}$ is seen to satisfy (2.4) and so, if in addition $\limsup_{j \rightarrow \infty} \lambda_j < \infty$, we have that the conditions of (iii)(b) are satisfied.

(e) We note that the conditions prescribed in Theorem 2.1 and Corollary 2.2 all involve quantifying how close together the particles can be on the average in the initial configuration of the Atlas model. In particular, we do not require any upper bounds on the sizes of U_i . This is in contrast with condition (1.8) assumed in [8] which requires the particles to be not too far apart on the average. Intuitively one expects convergence to π to hold when initially the particles are not too densely packed and upper bounds on the average rate of growth of the initial spacings are somewhat unnatural. The result in [27], Theorem 4.7, also points to this heuristic by showing weak convergence to π from all initial gap configurations that stochastically dominate π . We also note that we do not require any condition analogous to (1.9) and can, in particular, allow gaps to be zero.

(f) We note that the conditions (1.8)–(1.10) of [8] do not imply our conditions. To see this, consider the deterministic sequence of initial gaps: $U_i = i^{-2/3}$ for $n^3 < i < (n+1)^3$ and $U_{n^3} = n$, for any $n \in \mathbb{N}$. It can be checked that, $\sum_{i=1}^d U_i$ grows like $d^{2/3}$ and $\sum_{i=1}^d (\log U_i)_-$ grows like $d \log d$ as $d \rightarrow \infty$. Thus, (1.8)–(1.10) of [8] hold with $\beta = 1$ and $\theta(d) = \log d$. However, almost surely, $\sum_{i=1}^d U_i \wedge V_i$ grows like $d^{1/3}$, and therefore, (2.1) is violated.

(g) It will be interesting to investigate whether the condition in Theorem 2.1 in fact implies the stronger property that μ is in the DoA of π . We leave this as an open problem.

Our second theorem provides sufficient conditions for a measure on $\mathbb{R}_+^\infty$ to be in the WDoA of one of the other stationary measures π_a , $a > 0$.

THEOREM 2.4. Fix $a > 0$. Suppose that the probability measure μ on $\mathbb{R}_+^\infty$ satisfies the following: There exists a coupling $(\mathbf{U}, \mathbf{V}_a)$ of μ and π_a such that, almost surely,

$$(2.5) \quad \limsup_{d \rightarrow \infty} \frac{\log \log d}{\log d} \sum_{i=1}^d |V_{a,i} - U_i| = 0 \quad \text{and} \quad \limsup_{d \rightarrow \infty} \frac{U_d}{dV_{a,d}} < \infty.$$

Then $\mu \in \mathcal{S}$ and it belongs to the WDoA of π_a .

We remark that Theorem 2.4 also holds for the case $a = 0$. Indeed, suppose that the condition (2.5) in the theorem holds for $a = 0$. Denoting the corresponding $\mathbf{V}_0$ as $\mathbf{V}$, we have that

$$\sum_{i=1}^d U_i \wedge V_i = \sum_{i=1}^d U_i \vee V_i - \sum_{i=1}^d |V_i - U_i| \geq \sum_{i=1}^d V_i - \sum_{i=1}^d |V_i - U_i|.$$

Using this and the law of large numbers for $\{V_i\}$, it follows that (2.1) holds and thus from Theorem 2.1 the conclusion of Theorem 2.4 holds with $a = 0$. The reason we do not note the case $a = 0$ in the statement of Theorem 2.4 is because (2.1) is a strictly weaker assumption than (2.5) when $a = 0$. This is expected as for any initial gap distribution that stochastically dominates π , convergence to π holds (see Remark 2.3(a)). However, for convergence to π_a for some $a > 0$, the initial gap distribution should be appropriately close to π_a (see Remark 2.6(b) below).

The following corollary gives a set of random initial conditions for which (2.5) holds.

COROLLARY 2.5. Fix $a > 0$. Let

$$\mu := \bigotimes_{i=1}^{\infty} \text{Exp}(2 + ia + \lambda_i),$$

where $\{\lambda_i\}$ are real numbers satisfying $\lambda_i \geq -\beta(2 + ia)$ for some $\beta < 1$, and $\lambda_i \leq C(2 + ia)$ for some $C > 0$, for all i . Moreover, assume

$$(2.6) \quad \limsup_{d \rightarrow \infty} \frac{\log \log d}{\log d} \sum_{i=1}^d \frac{|\lambda_i|}{i^2} = 0.$$

Then (2.5) holds, and hence, $\mu \in \mathcal{S}$ and belongs to the WDoA of π_a .

REMARK 2.6. We note the following.

(a) Clearly, there are measures μ of the form described in Corollary 2.5 for which $\liminf_{i \rightarrow \infty} \frac{|\lambda_i|}{i} > 0$ and which do not lie in the WDoA of π_a (e.g., $\pi_{a'}$ for any $a' \neq a$). However, the corollary says that if $|\lambda_i|$ grows slightly slower than i then μ is indeed in the WDoA of π_a . More precisely, the convergence in (2.6) holds in particular if $|\lambda_i| \leq i\delta_i/(\log \log i)$ for sufficiently large i , for some nonnegative sequence $\delta_i \rightarrow 0$ as $i \rightarrow \infty$. Indeed, note that for any $\delta > 0$, there exists sufficiently large $i_\delta \in \mathbb{N}$ such that for $i \geq i_\delta$,

$$(2.7) \quad \sum_{j=i_\delta}^i \frac{|\lambda_j|}{j^2} \leq \sum_{j=i_\delta}^i \frac{\delta_j}{j \log \log j} \leq \delta \int_{i_\delta-1}^i \frac{1}{z \log \log z} dz = \delta \int_{\log(i_\delta-1)}^{\log i} \frac{1}{\log w} dw.$$

As $w \mapsto \frac{1}{\log w}$ is a slowly varying function, by Karamata's theorem [24], Theorem 1.2.6 (a),

$$\lim_{i \rightarrow \infty} \frac{\log \log i}{\log i} \int_{\log(i_0-1)}^{\log i} \frac{1}{\log w} dw = 1.$$

This can also be directly shown by noting that the primitive of $1/\log w$ is $\text{Li}(w)$, and applying the l'Hôpital's rule. Thus, we conclude from (2.7) that, for any $\delta > 0$,

$$\limsup_{i \rightarrow \infty} \frac{\log \log i}{\log i} \sum_{j=1}^i \frac{|\lambda_j|}{j^2} \leq \delta.$$

As $\delta > 0$ is arbitrary, (2.6) holds.

(b) Note that if $\mathbf{U} \sim \pi$, then for $a, a' > 0$,

$$V_{a,i} = \frac{2}{2+ia} U_i, \quad V_{a',i} = \frac{2}{2+ia'} U_i, \quad i \in \mathbb{N}$$

defines a coupling of π_a and $\pi_{a'}$. For this coupling,

$$(2.8) \quad \sum_{i=1}^d |V_{a,i} - V_{a',i}| \sim O(|a - a'| \log d) \quad \text{and} \quad \limsup_{d \rightarrow \infty} \frac{V_{a',d}}{d V_{a,d}} = 0.$$

Since $\pi_{a'}$ is obviously not in WDoA of π_a for $a \neq a'$, the first property in (2.8) says that the first requirement in (2.5) is not far from what one expects. We conjecture that for μ to be in the WDoA of π_a it is necessary that $(\log d)^{-1} \sum_{i=1}^d |V_{a,i} - U_i| \rightarrow 0$ as $d \rightarrow \infty$ for some coupling $(\mathbf{U}, \mathbf{V}_a)$ of μ and π_a . Theorem 2.4 says that if the first convergence holds at a rate faster than $1/(\log \log d)$ for some coupling of μ and π_a then that (together with the second condition in (2.5)) is sufficient for μ to be in the WDoA of π_a .

Further, observe that if the second condition in (2.5) does not hold, then there are infinitely many $d \in \mathbb{N}$ such that the lowest $d + 1$ particles are separated from the rest by a large initial gap. If one such gap is very large, then it could happen that the distribution of gaps between the $d + 1$ particles stabilizes toward the unique stationary gap distribution $\pi^{(d)}$ of the corresponding finite Atlas model before the remaining particles have interacted with them. As $\pi^{(d)}$ converges weakly to π as $d \rightarrow \infty$, one does not expect in such situations the convergence (of time averaged laws) to π_a to hold for any $a > 0$.

REMARK 2.7. As noted in the [Introduction](#), the approach of [8] using finite-dimensional approximations does not seem to have an easy extension when investigating the DoA (or WDoA) of π_a for $a > 0$. This is because the stationary distribution of the finite-dimensional Atlas model approaches $\pi = \pi_0$ as the dimension approaches infinity. Nevertheless, there is an alternative natural finite-dimensional rank-based diffusion model that one may consider as an approximation to the infinite Atlas model which is better suited for proving convergence to π_a for $a > 0$. Consider the SDE with $s \geq 0$ and $i \in \{0, 1, \dots, d\}$,

$$(2.9) \quad \begin{aligned} dY_i(s) &= \left(1 - \frac{a}{d+1}\right) \mathbf{1}(Y_i(s) = Y_{(0)}(s)) ds - \sum_{j=1}^d \frac{ja}{d+1} \mathbf{1}(Y_i(s) \\ &= Y_{(j)}(s)) ds + dW_i(s). \end{aligned}$$

It can be checked using the form of the generator of the above diffusion that the corresponding gap process has the unique stationary distribution $\pi^{a,(d)} := \bigotimes_{i=1}^d \text{Exp}((2+ia)(1-i/(d+1)))$, which clearly converges to π_a as $d \rightarrow \infty$. There is also a natural analog of the right-anchored rank-based diffusion of the form in [8], Section 2 (see equation (2.12) therein) which is associated with the above dynamics and the stationary distribution π_a . In this model, the bottom d particles evolve as a finite Atlas model reflected off the top particle, which moves deterministically with constant velocity $-a/2$. Both of these Markov processes (as in the setting of [8]) are reversible with respect to the corresponding stationary distributions and are therefore well suited for Dirichlet form methods developed in [8]. In particular, one can

obtain an estimate analogous to [8], Proposition 1.3, for these finite-dimensional diffusions. However, because the drift and reflection structure of the above finite rank-based diffusions is substantially different from that of the infinite Atlas model, it is not clear how to devise comparison techniques between these diffusions and the infinite Atlas model, that are crucial to both [8] and our work.

Nevertheless, the above diffusions do capture some key qualitative features that are exhibited by the bottom $d + 1$ particles in the infinite Atlas model starting from configurations close (in distribution) to π_a . In particular, as proved in [30], if the initial gaps are distributed as π_a , the Atlas processes $\{Y_{(i)}(t) + at/2 : t \geq 0\}_{i \in \mathbb{N}_0}$ (where $Y_{(i)}(\cdot)$ satisfy (1.5)) form a tight collection. The drift $-aj/(d + 1)$ of the j th ordered particle, for $j = 1, \dots, d$, in the diffusion satisfying (2.9) implies an analogous property for the ordered processes in this diffusion for large d . To see this, observe that the center of mass of the system has drift $(d + 1)^{-1}[(1 - \frac{a}{d+1}) - \sum_{j=1}^d \frac{ja}{d+1}]$ which is asymptotically $-a/2$ as $d \rightarrow \infty$. A similar property is true for the anchored dynamics as the top particle moves as a linear barrier with slope $-a/2$. Understanding relationships between these finite-dimensional rank-based diffusions and the infinite Atlas model, particularly over long time intervals, is key to extending the techniques of [8] to study the DoA of π_a for $a > 0$. We hope to explore this in future work.

3. Synchronous couplings of the infinite ordered Atlas model. As mentioned earlier, a synchronous coupling of the ranked particles in the infinite Atlas model refers to coupling two copies of the order statistics processes starting from different initial configurations but driven by the same set of Brownian motions via the SDE (1.5). The finite-dimensional analogue of the SDE (1.5) is known to have a unique strong solution [13], which implies the existence of synchronous couplings for the ranked processes of finite Atlas models starting from any two initial configurations. For the infinite Atlas model, devising synchronous couplings is a more delicate issue as described below.

In [27], a systematic method for constructing solutions of the infinite-dimensional SDE (1.5), from a given collection of Brownian motions, has been devised under the name of *approximative versions*. It is based on taking appropriate limits of solutions of the corresponding finite-dimensional systems. This, in turn, can be used to construct synchronous couplings in an appropriate sense. We collect some results from [27] in the following proposition.

PROPOSITION 3.1. *Suppose $\mathbf{x}$ is a $\mathbb{R}_+^\infty$ -valued random variable satisfying $0 \leq x_0 \leq x_1 \leq x_2 \leq \dots$ and, almost surely,*

$$(3.1) \quad \sum_{i=0}^{\infty} e^{-\alpha x_i^2} < \infty \quad \text{for all } \alpha > 0.$$

Consider a collection of independent standard Brownian motions $B_0(\cdot), B_1(\cdot), B_2(\cdot), \dots$, independent of $\mathbf{x}$. Consider the collection of SDE, for $i \in \mathbb{N}_0$,

$$(3.2) \quad dX_i(t) = \mathbf{1}(i = 0)dt + dB_i(t) - \frac{1}{2}dL_{i+1}(t) + \frac{1}{2}dL_i(t), \quad X_i(0) = x_i, t \geq 0,$$

where $L_0(\cdot) \equiv 0$, and for $i \in \mathbb{N}$, $L_i(\cdot)$ is the associated local time of collision between the $(i - 1)$ th and i th particle, namely

$$(3.3) \quad L_i, X_i - X_{i-1} \text{ are nonnegative, continuous process, } L_i \text{ is nondecreasing,}$$

$$L_i(0) = 0 \quad \text{and} \quad L_i(t) = \int_0^t \mathbf{1}(X_{i-1}(s) = X_i(s)) dL_i(s) \quad \text{for all } t \geq 0.$$

Also consider for fixed $m \in \mathbb{N}$, the system of SDE in (3.2) for $i = 0, 1, \dots, m$, with starting configuration $X_i(0) = x_i$, $0 \leq i \leq m$, and with L_i satisfying (3.3) for $1 \leq i \leq$

m and $L_0(\cdot) \equiv L_{m+1}(\cdot) \equiv 0$. Denote by $\mathbf{X}^{(m)}(\cdot) = (X_0^{(m)}(\cdot), \dots, X_m^{(m)}(\cdot))$ and $\mathbf{L}^{(m)}(\cdot) = (L_0^{(m)}(\cdot), \dots, L_m^{(m)}(\cdot))$ the unique strong solution to this finite-dimensional system of reflected SDE which we call the ‘finite ordered Atlas model with $m + 1$ particles’ with driving Brownian motion $(B_0(\cdot), \dots, B_m(\cdot))$. Then, the following hold:

(i) There exist continuous $\mathbb{R}^\infty$ -valued processes $\mathbf{X}(\cdot) := (X_i(\cdot) : i \in \mathbb{N}_0)$, $\mathbf{L}(\cdot) := (L_i(\cdot) : i \in \mathbb{N}_0)$, adapted to $\mathcal{F}_t \doteq \sigma\{B_i(s) : s \leq t, i \in \mathbb{N}_0\}$, such that, a.s. $(\mathbf{X}, \mathbf{L})$ satisfy (3.2) and (3.3) and for any $T \in (0, \infty)$,

$$\lim_{m \rightarrow \infty} \sup_{t \in [0, T]} [|X_i^{(m)}(t) - X_i(t)| + |L_i^{(m)}(t) - L_i(t)|] = 0 \quad \text{a.s., for all } i \in \mathbb{N}_0.$$

We will call $\mathbf{X}(\cdot)$ the ‘infinite ordered Atlas model’ with driving Brownian motions $B_0(\cdot), B_1(\cdot), B_2(\cdot), \dots$, started from $\mathbf{x} = (x_0, x_1, x_2, \dots)$.

(ii) Suppose $\mathbf{u}, \mathbf{v}$ are $\mathbb{R}_+^\infty$ -valued random variables independent of $\{B_i, i \in \mathbb{N}_0\}$, such that $u_i \leq u_{i+1}$ and $v_i \leq v_{i+1}$ a.s. for all $i \in \mathbb{N}_0$, and (3.1) holds a.s. with $\mathbf{x}$ replaced with $\mathbf{u}, \mathbf{v}$. Then, on $(\Omega, \mathcal{F}, \mathbb{P})$ there are continuous, $\mathbb{R}^\infty$ -valued processes $\mathbf{X}^{\mathbf{u}}, \mathbf{L}^{\mathbf{u}}$ and $\mathbf{X}^{\mathbf{v}}, \mathbf{L}^{\mathbf{v}}$ that are adapted to $\hat{\mathcal{F}}_t = \mathcal{F}_t \vee \sigma\{\mathbf{u}, \mathbf{v}\}$ and solve (3.2), (3.3) with $\mathbf{X}^{\mathbf{u}}(0) = \mathbf{u}$ and $\mathbf{X}^{\mathbf{v}}(0) = \mathbf{v}$, respectively. These processes are given as follows. Let $\mathbf{X}^{(m), \mathbf{u}}(\cdot)$ and $\mathbf{X}^{(m), \mathbf{v}}(\cdot)$ be the unique strong solutions of the finite ordered Atlas model with $m + 1$ particles, with $\mathbf{X}^{(m), \mathbf{u}}(0) = \mathbf{u}|_{m+1}$, $\mathbf{X}^{(m), \mathbf{v}}(0) = \mathbf{v}|_{m+1}$. Then $\mathbf{X}^{\mathbf{u}}, \mathbf{X}^{\mathbf{v}}$ are defined as the a.s. limits of $\mathbf{X}^{(m), \mathbf{u}}(\cdot)$ and $\mathbf{X}^{(m), \mathbf{v}}(\cdot)$ as $C([0, \infty) : \mathbb{R}^\infty)$ -valued random variables. We will refer to these limit processes as a synchronous coupling of the infinite ordered Atlas model driven by Brownian motions $\{B_i, i \in \mathbb{N}_0\}$ with initial configuration $\mathbf{X}^{\mathbf{u}}(0) = \mathbf{u}$ and $\mathbf{X}^{\mathbf{v}}(0) = \mathbf{v}$.

(iii) If $(\mathbf{u}, \mathbf{v})$ in (ii) are such that $\mathbf{u} \leq \mathbf{v}$, then the synchronous coupling defined in (ii) almost surely satisfies $\mathbf{X}^{\mathbf{u}}(t) \leq \mathbf{X}^{\mathbf{v}}(t)$ for all $t \geq 0$.

Define the corresponding ‘gap processes’ $\mathbf{Z}^{\mathbf{u}}(\cdot)$ and $\mathbf{Z}^{\mathbf{v}}(\cdot)$ by $Z_i^{\mathbf{u}}(\cdot) := X_i^{\mathbf{u}}(\cdot) - X_{i-1}^{\mathbf{u}}(\cdot)$ and $Z_i^{\mathbf{v}}(\cdot) := X_i^{\mathbf{v}}(\cdot) - X_{i-1}^{\mathbf{v}}(\cdot)$ for $i \in \mathbb{N}$. Then $\mathbf{Z}^{\mathbf{u}}(0) \leq \mathbf{Z}^{\mathbf{v}}(0)$ almost surely implies the following hold almost surely:

$$\mathbf{Z}^{\mathbf{u}}(t) \leq \mathbf{Z}^{\mathbf{v}}(t) \quad \text{for all } t \geq 0 \quad \text{and} \quad \mathbf{L}^{\mathbf{v}}(t) - \mathbf{L}^{\mathbf{u}}(t) \leq \mathbf{L}^{\mathbf{v}}(s) - \mathbf{L}^{\mathbf{u}}(s) \quad \text{for all } 0 \leq s \leq t.$$

(iv) Analogous statements hold with $\mathbf{1}(i = 0)$ in (3.2) replaced with $\gamma \mathbf{1}(i = 0)$ where $\gamma \in \mathbb{R}$. Specifically, for $\gamma \in \mathbb{R}$, define the ‘infinite ordered Atlas model with drift γ ’ to be the $\mathbb{R}^\infty$ -valued stochastic process $\mathbf{X}^{[\gamma]}(\cdot) := (X_0^{[\gamma]}(\cdot), X_1^{[\gamma]}(\cdot), \dots)$ satisfying the SDE

$$(3.4) \quad dX_i^{[\gamma]}(t) = \gamma \mathbf{1}(i = 0) dt + dB_i(t) - \frac{1}{2} dL_{i+1}^{[\gamma]}(t) + \frac{1}{2} dL_i^{[\gamma]}(t), \quad t \geq 0, i \in \mathbb{N}_0,$$

where $L_0^{[\gamma]}(\cdot) \equiv 0$ and $(L_1^{[\gamma]}(\cdot), L_2^{[\gamma]}(\cdot), \dots)$ are the associated local times as in (3.3). Then $\mathbf{X}^{[\gamma]}(\cdot)$ can be obtained as a limit of the solutions to the finite-dimensional analogues of the SDE (3.4) in the sense of (i). In particular, one can construct a synchronous coupling of $\mathbf{X}(\cdot)$ and $\mathbf{X}^{[\gamma]}(\cdot)$ for same or different (coupled) initial configurations as in (ii). Moreover, if $\gamma \leq 1$ and $\mathbf{Z}(\cdot)$ and $\mathbf{Z}^{[\gamma]}(\cdot)$ are the associated gap processes under synchronous coupling and $\mathbf{Z}(0) = \mathbf{Z}^{[\gamma]}(0)$, then, almost surely,

$$\mathbf{Z}(t) \leq \mathbf{Z}^{[\gamma]}(t) \quad \text{for all } t \geq 0.$$

PROOF. (i) follows from [27], Definition 7 and Theorem 3.7, [27], Lemma 6.4, and the discussion following it. Part (ii) is immediate from (i). (iii) follows from [27], Corollary 3.10. The assertion in (iv) that $\mathbf{X}^\gamma(\cdot)$ can be obtained as a limit of finite systems again follows from [27], Definition 7 and Theorem 3.7. The pathwise domination assertion under synchronous coupling follows from [27], Corollary 3.12(ii). $\square$

Proposition 3.1 constructs the infinite ordered Atlas model via approximation by finite ordered Atlas models. As the solution to the finite-dimensional analogue of (3.2) is unique in law (which follows from pathwise uniqueness), the finite ordered Atlas model has the same law as the ranked processes in the finite Atlas model (which is a weak solution to the finite-dimensional analogue of (1.2)). The following lemma says that the infinite ordered Atlas model, constructed in Proposition 3.1(i), indeed has the same law as the ranked trajectories of the infinite Atlas model satisfying (1.2). A similar result was discussed in [27], Remark 2. We provide a proof for completeness.

LEMMA 3.2. *Let $\mathbf{x}$ be a $\mathbb{R}_+^\infty$ -valued random variable satisfying $0 \leq x_0 \leq x_1 \leq x_2 \leq \dots$ such that, almost surely, (3.1) holds, and let $\{B_i, i \in \mathbb{N}_0\}$ be a sequence of independent standard Brownian motions, independent of $\mathbf{x}$. Let $\mathbf{X}$ be the process constructed in Proposition 3.1(i) with $\mathbf{X}(0) = \mathbf{x}$. Let μ be the probability law of $\mathbf{x}$ and $\mathbf{Y}(\cdot) := (Y_0(\cdot), Y_1(\cdot), \dots)$ be the unique weak solution of (1.2) with $\mathbf{Y}(0) \sim \mu$. Then $\mathbf{X}$ has the same law as the ranked processes $\mathbf{Y}_r(\cdot) := (Y_{(0)}(\cdot), Y_{(1)}(\cdot), \dots)$.*

PROOF. Let $\mathbf{Y}$ be as in the statement of the lemma and for any $m \in \mathbb{N}$, let $\mathbf{Y}^{(m)}(\cdot) = (Y_0^{(m)}(\cdot), \dots, Y_m^{(m)}(\cdot))$ be the unique weak solution of the finite Atlas model (1.2) with $m+1$ particles (i.e., with $i \in \mathbb{N}_0$ in (1.2) replaced by $0 \leq i \leq m$) and with $\mathbf{Y}^{(m)}(0)$ distributed same as $(Y_0(0), Y_1(0), \dots, Y_m(0))$. Let $\mathbf{Y}_r^{(m)}(\cdot) := (Y_{(0)}^{(m)}(\cdot), \dots, Y_{(m)}^{(m)}(\cdot))$ be the corresponding ranked trajectories. By [27], Theorem 3.3, there exists a sequence $m_j \rightarrow \infty$ such that for any $T > 0$ and $k \in \mathbb{N}$,

$$(3.5) \quad (\mathbf{Y}^{(m_j)}(\cdot)|_k, \mathbf{Y}_r^{(m_j)}(\cdot)|_k) \xrightarrow{d} (\mathbf{Y}(\cdot)|_k, \mathbf{Y}_r(\cdot)|_k) \quad \text{on } \mathbb{C}([0, T] : \mathbb{R}^{2k}).$$

Next, let $\mathbf{x}, \{B_i, i \in \mathbb{N}_0\}$ be as in the statement of the lemma and let $(\mathbf{X}^{(m)}, \mathbf{L}^{(m)})$ be as in Proposition 3.1 (i) given as the unique strong solution of the corresponding $(m+1)$ -dimensional reflected SDE, with $\mathbf{X}^{(m)}(0) = \mathbf{x}|_{m+1}$. From the same argument used to derive (1.5) from (1.2), it follows that $\mathbf{Y}_r^{(m_j)}(\cdot)$ satisfies the m_j -dimensional version of the SDE (1.5) for each $j \geq 1$. By pathwise uniqueness of solutions to the finite-dimensional reflected SDE [13], for any $k \in \mathbb{N}$,

$$(3.6) \quad \mathbf{Y}_r^{(m_j)}(\cdot)|_k \stackrel{d}{=} \mathbf{X}^{(m_j)}(\cdot)|_k \quad \text{for all } j \geq 1.$$

Moreover, by Proposition 3.1(ii), $\mathbf{X}^{(m)}|_k \rightarrow \mathbf{X}|_k$, almost surely, in $C([0, \infty) : \mathbb{R}^{k+1})$, as $m \rightarrow \infty$. This observation, along with (3.5) and (3.6), gives

$$\mathbf{Y}_r(\cdot)|_k \stackrel{d}{=} \mathbf{X}(\cdot)|_k \quad \text{for any } k \in \mathbb{N},$$

which proves the lemma. $\square$

A pathwise analysis of synchronous couplings will be the main tool in proving our main results. A key observation is the following control on the L^1 distance between the gap processes corresponding to two synchronously coupled infinite ordered Atlas models. In particular, it shows (using Proposition 3.1(iii)) that if the initial gap distributions satisfy a coordinatewise ordering, then this L^1 distance is nonincreasing in time and its decay rate can be precisely estimated from the local times of collision between the bottom two particles. We note that the proof for an analogous result for the finite ordered Atlas model is easier and can be deduced from standard properties of reflected Brownian motions [21], Theorem 6.

LEMMA 3.3. Let $B_0(\cdot), B_1(\cdot), B_2(\cdot), \dots$ be standard Brownian motions given on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\mathbf{X}(\cdot)$ and $\tilde{\mathbf{X}}(\cdot)$ be synchronously coupled infinite ordered Atlas models driven by $\{B_i, i \in \mathbb{N}_0\}$ as constructed in Proposition 3.1(ii), with the initial configurations $\mathbf{X}(0)$ and $\tilde{\mathbf{X}}(0)$ such that the corresponding probability laws are in S_0 . Let $\mathbf{Z}(\cdot)$ and $\tilde{\mathbf{Z}}(\cdot)$ denote the corresponding gap processes. Suppose $\tilde{\mathbf{Z}}(0) \geq \mathbf{Z}(0)$. Write $\Delta \mathbf{Z}(\cdot) := \tilde{\mathbf{Z}}(\cdot) - \mathbf{Z}(\cdot)$ and assume $\sum_{j=1}^{\infty} j \Delta Z_j(0) < \infty$. Let $\mathbf{L}(\cdot)$ and $\tilde{\mathbf{L}}(\cdot)$ denote the local times of collision for the respective Atlas models and write $\Delta \mathbf{L}(\cdot) := \tilde{\mathbf{L}}(\cdot) - \mathbf{L}(\cdot)$. Then, almost surely,

$$(3.7) \quad \sum_{j=1}^{\infty} \Delta Z_j(t) = \sum_{j=1}^{\infty} \Delta Z_j(0) + \frac{1}{2} \Delta L_1(t), \quad t \geq 0.$$

PROOF. Observe from (3.2) that for any $d \in \mathbb{N}, t \geq 0$,

$$(3.8) \quad \begin{aligned} \sum_{j=1}^d \Delta Z_j(t) &= \sum_{j=1}^d \Delta Z_j(0) + \sum_{j=1}^d \left(\Delta L_j(t) - \frac{1}{2} \Delta L_{j-1}(t) - \frac{1}{2} \Delta L_{j+1}(t) \right) \\ &= \sum_{j=1}^d \Delta Z_j(0) + \frac{1}{2} \Delta L_1(t) + \frac{1}{2} \Delta L_d(t) - \frac{1}{2} \Delta L_{d+1}(t). \end{aligned}$$

Thus, to show (3.7), it suffices to show that, for every $t \geq 0$,

$$(3.9) \quad \lim_{d \rightarrow \infty} \frac{1}{2} \Delta L_d(t) - \frac{1}{2} \Delta L_{d+1}(t) = 0.$$

This will be shown in several steps as follows.

For $m \in \mathbb{N}$, denote by $\mathbf{Z}^{(m)}(\cdot)$ and $\tilde{\mathbf{Z}}^{(m)}(\cdot)$ the m -dimensional gap processes associated with ranked processes $\mathbf{X}^{(m)}(\cdot)$ and $\tilde{\mathbf{X}}^{(m)}(\cdot)$ for the finite ordered Atlas models with $m+1$ particles, respectively, starting from $\mathbf{X}(0)|_{m+1}$ and $\tilde{\mathbf{X}}(0)|_{m+1}$ and driven by $\{B_i\}_{i=0}^m$. Recall from Proposition 3.1 that $\mathbf{Z}^{(m)}(\cdot)$ and $\tilde{\mathbf{Z}}^{(m)}(\cdot)$, respectively, converge to $\mathbf{Z}(\cdot)$ and $\tilde{\mathbf{Z}}(\cdot)$ a.s. in $C([0, \infty) : \mathbb{R}_+^{\infty})$ as $m \rightarrow \infty$. Write $\Delta \mathbf{Z}^{(m)}(\cdot) := \tilde{\mathbf{Z}}^{(m)}(\cdot) - \mathbf{Z}^{(m)}(\cdot)$. Similarly, let $\Delta \mathbf{L}^{(m)}(\cdot) := \tilde{\mathbf{L}}^{(m)}(\cdot) - \mathbf{L}^{(m)}(\cdot)$ denote the difference vector between the respective local times. Then, from the SDE for $\mathbf{X}^{(m)}(\cdot)$ and $\tilde{\mathbf{X}}^{(m)}(\cdot)$ (i.e., the finite-dimensional version of (3.2)), we obtain

$$(3.10) \quad \Delta \mathbf{Z}^{(m)}(t) = \Delta \mathbf{Z}^{(m)}(0) + R^{(m)} \Delta \mathbf{L}^{(m)}(t), \quad t \geq 0,$$

where $R^{(m)}$ is an $m \times m$ matrix given by $R_{ii}^{(m)} = 1$ for all $1 \leq i \leq m$, $R_{i(i-1)}^{(m)} = -1/2$ for $2 \leq i \leq m$, $R_{i(i+1)}^{(m)} = -1/2$ for $1 \leq i \leq m-1$ and $R_{ij}^{(m)} = 0$ if $|i-j| \geq 2$. It can be checked (see, e.g., proof of [3], Theorem 4) that $R^{(m)}$ is invertible and

$$(3.11) \quad (R^{(m)})_{ij}^{-1} = 2(i \wedge j) \left(1 - \frac{i \vee j}{m+1} \right), \quad 1 \leq i, j \leq m.$$

Also, from [21], Theorem 6,

$$(3.12) \quad \begin{aligned} &\text{For every } t \geq 0, \tilde{\mathbf{Z}}^{(m)}(t) \geq \mathbf{Z}^{(m)}(t) \text{ and } -\Delta \mathbf{L}^{(m)}(\cdot) \\ &\text{is nonnegative and nondecreasing, a.s.} \end{aligned}$$

Write $\zeta := \sum_{j=1}^{\infty} j \Delta Z_j(0) < \infty$. Then, it follows from (3.10) that, for any $1 \leq d \leq m, t \geq 0$,

$$\sum_{j=1}^d (R^{(m)})_{dj}^{-1} \Delta Z_j^{(m)}(t)$$

$$\begin{aligned}
 (3.13) \quad & \leq \sum_{j=1}^m (R^{(m)})_{dj}^{-1} \Delta Z_j^{(m)}(t) = \sum_{j=1}^m (R^{(m)})_{dj}^{-1} \Delta Z_j^{(m)}(0) + \Delta L_d^{(m)}(t) \\
 & \leq \sum_{j=1}^m (R^{(m)})_{dj}^{-1} \Delta Z_j^{(m)}(0) \leq 2 \sum_{j=1}^m j \Delta Z_j^{(m)}(0) = 2 \sum_{j=1}^m j \Delta Z_j(0) \leq 2\zeta,
 \end{aligned}$$

where we used the nonnegativity of $(R^{(m)})_{dj}^{-1}$ (see (3.11)) and the first part of (3.12) for the first inequality, the second part of (3.12) for the second inequality and the explicit form of $(R^{(m)})^{-1}$ given in (3.11) in the third inequality. Recall that, for any $t \geq 0$, $j \in \mathbb{N}$, $\Delta Z_j^{(m)}(t) \rightarrow \Delta Z_j(t)$ as $m \rightarrow \infty$. Moreover, using (3.11), for any fixed $j, d \in \mathbb{N}$ with $j \leq d$, $(R^{(m)})_{dj}^{-1} \rightarrow 2j$ as $m \rightarrow \infty$. Thus, fixing $d \in \mathbb{N}$, we obtain by letting $m \rightarrow \infty$ on the left-hand side of (3.13), for all $d \in \mathbb{N}$,

$$(3.14) \quad 2 \sum_{j=1}^d j \Delta Z_j(t) \leq 2\zeta, \quad t \geq 0.$$

By Proposition 3.1(iii), the left-hand side of (3.14) is nondecreasing in d . Hence, taking limit as $d \rightarrow \infty$,

$$(3.15) \quad \sum_{j=1}^{\infty} j \Delta Z_j(t) \leq \zeta, \quad t \geq 0.$$

Next, we claim that for any $d \in \mathbb{N}$, $t \geq 0$,

$$(3.16) \quad 0 \leq -\Delta L_d(t) \leq 2\zeta.$$

To show this, we note that by (3.12) and (3.13), for any $m \geq d$,

$$-\Delta L_d^{(m)}(t) \leq \sum_{j=1}^m (R^{(m)})_{dj}^{-1} \Delta Z_j^{(m)}(0) \leq 2\zeta.$$

The upper bound in (3.16) now follows upon taking a limit as $m \rightarrow \infty$ on the left-hand side above and using Proposition 3.1(i). The lower bound follows from Proposition 3.1(iii).

Finally, we claim that for any $d \in \mathbb{N}$, $t \geq 0$,

$$(3.17) \quad -\zeta \leq \frac{d+1}{2} \Delta L_d(t) - \frac{d}{2} \Delta L_{d+1}(t) \leq \zeta.$$

To see this, observe that

$$\begin{aligned}
 \sum_{j=1}^d j \Delta Z_j(t) &= \sum_{j=1}^d j \Delta Z_j(0) + \sum_{j=1}^d j \left(\Delta L_j(t) - \frac{1}{2} \Delta L_{j-1}(t) - \frac{1}{2} \Delta L_{j+1}(t) \right) \\
 &= \sum_{j=1}^d j \Delta Z_j(0) + \frac{d+1}{2} \Delta L_d(t) - \frac{d}{2} \Delta L_{d+1}(t).
 \end{aligned}$$

Hence,

$$\frac{d+1}{2} \Delta L_d(t) - \frac{d}{2} \Delta L_{d+1}(t) = \sum_{j=1}^d j \Delta Z_j(t) - \sum_{j=1}^d j \Delta Z_j(0).$$

The inequality in (3.17) is now immediate from the above upon using Proposition 3.1(iii) and (3.15) and noting that

$$-\zeta = -\sum_{j=1}^{\infty} j \Delta Z_j(0) \leq \sum_{j=1}^d j \Delta Z_j(t) - \sum_{j=1}^d j \Delta Z_j(0) \leq \sum_{j=1}^{\infty} j \Delta Z_j(t) \leq \zeta.$$

The convergence in (3.9) now follows from (3.16) and (3.17) as for every $t \geq 0$,

$$\lim_{d \rightarrow \infty} \frac{1}{2} \Delta L_d(t) - \frac{1}{2} \Delta L_{d+1}(t) = \lim_{d \rightarrow \infty} \left[\frac{1}{d} \left(\frac{d+1}{2} \Delta L_d(t) - \frac{d}{2} \Delta L_{d+1}(t) \right) - \frac{1}{2d} \Delta L_d(t) \right] = 0.$$

This proves (3.7) and hence the lemma. $\square$

4. Influence of far away coordinates. A crucial component of our approach will be to obtain a quantitative understanding of the time taken for the first k gaps between the ranked particles in the infinite Atlas model to ‘feel the effect’ of the unordered processes $Y_i(\cdot)$ for $i \gg k$. In this section, we derive such estimates. We recall that the infinite Atlas model is given as the unique weak solution of (1.2) when the law of $Y(0)$ is an element of $\mathcal{S}_0$. Note that the latter property, which will be assumed throughout this section, says that $Y_{(0)}(0) = 0$ and $Y_{(i)}(0) = Y_i(0)$ for all $i \in \mathbb{N}_0$.

We will rely on the following estimates derived in [8].

LEMMA 4.1. (i) ([8], Lemma 3.2). For any $k \in \mathbb{N}$, $t > 0$, $l \in \mathbb{N}$, $\Gamma \in \mathbb{R}$, a.s.,

$$(4.1) \quad \mathbb{P} \left(\sup_{s \in [0, t]} Y_{(k)}(s) \geq \Gamma | \mathbf{Y}(0) \right) \leq 2 \bar{\Phi} \left(\frac{l(\Gamma - Y_k(0))/3 - t - \sum_{j=0}^{l-1} Y_j(0)}{\sqrt{lt}} \right) + 4(k+1) \bar{\Phi} \left(\frac{\Gamma - Y_k(0)}{3\sqrt{t}} \right).$$

(ii) ([8], Lemma 3.3). For any $d \in \mathbb{N}$, $t > 0$, $\Gamma > 0$, a.s.,

$$(4.2) \quad \mathbb{P} \left(\inf_{s \in [0, t]} \inf_{i \geq d} Y_i(s) \leq \Gamma | \mathbf{Y}(0) \right) \leq 2 \sum_{i \geq d} \bar{\Phi} \left(\frac{Y_i(0) - \Gamma}{\sqrt{t}} \right).$$

The following lemma will be used in Section 6 (see proof of Lemma 6.3) to quantify influence of far away coordinates when the starting configuration satisfies (2.1).

LEMMA 4.2. Suppose that the distribution of $Y_i(0) - Y_{i-1}(0)$ is stochastically dominated by the $\text{Exp}(2)$ distribution for each $i \in \mathbb{N}$. Moreover, assume

$$(4.3) \quad \liminf_{d \rightarrow \infty} \frac{Y_d(0)}{\sqrt{d}(\log d)} = \infty \quad \text{almost surely.}$$

Then for any $A \geq 4$, $k \in \mathbb{N}$, writing $t_d^A := Ad(\log d)$ and $\Gamma_d^A := 6A\sqrt{d}(\log d)$ for $d \in \mathbb{N}$,

$$\begin{aligned} \mathbb{P} \left(\inf_{s \in [0, t_d^A]} \inf_{i \geq d} Y_i(s) \leq \Gamma_d^A \right) &\rightarrow 0 \quad \text{as } d \rightarrow \infty, \\ \mathbb{P} \left(\sup_{s \in [0, t_d^A]} Y_{(k)}(s) \geq \Gamma_d^A \right) &\rightarrow 0 \quad \text{as } d \rightarrow \infty. \end{aligned}$$

PROOF. For $d \in \mathbb{N}$, let $l_d := \lceil \sqrt{d} \rceil$. Define the event

$$\mathcal{A}(n) := \{Y_i(0) - Y_{i-1}(0) \leq \log i \text{ for all } i > n\}, \quad n \in \mathbb{N}.$$

By the stochastic domination assumption,

$$(4.4) \quad \begin{aligned} \mathbb{P}(\mathcal{A}(n)^c) &\leq \sum_{i=n+1}^{\infty} \mathbb{P}(Y_i(0) - Y_{i-1}(0) > \log i) \\ &\leq \sum_{i=n+1}^{\infty} i^{-2} \leq n^{-1} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

For fixed $n \in \mathbb{N}$ and d sufficiently large, we have

$$\begin{aligned} \sum_{j=0}^{l_d-1} Y_j(0) &= \sum_{j=0}^{n-1} Y_j(0) + \sum_{j=n}^{l_d-1} Y_j(0) = \sum_{j=0}^{n-1} Y_j(0) + Y_n(0)(l_d - n) + \sum_{j=n}^{l_d-1} (Y_j(0) - Y_n(0)) \\ &\leq Y_n(0)l_d + \sum_{j=n}^{l_d-1} (Y_j(0) - Y_n(0)). \end{aligned}$$

Also, for $n \leq j \leq l_d - 1$, on the event $\mathcal{A}(n)$,

$$Y_j(0) - Y_n(0) = \sum_{k=n+1}^j (Y_k(0) - Y_{k-1}(0)) \leq \sum_{k=n+1}^j \log k.$$

Thus, on $\mathcal{A}(n)$, for d sufficiently large,

$$\sum_{j=0}^{l_d-1} Y_j(0) \leq Y_n(0)l_d + \sum_{k=n+1}^{l_d-1} (k-1) \log k \leq (1 + \sqrt{d})Y_n(0) + (1 + \sqrt{d})^2 \log(1 + \sqrt{d}).$$

Thus, for fixed $n \in \mathbb{N}$, there exists deterministic $d_0 \in \mathbb{N}$ such that, on the event $\mathcal{A}(n)$, for all $d \geq d_0$,

$$\begin{aligned} &\frac{l_d(\Gamma_d^A - Y_k(0))/3 - t_d^A - \sum_{j=0}^{l_d-1} Y_j(0)}{\sqrt{l_d t_d^A}} \\ &\geq \frac{2Ad(\log d) - Ad(\log d) - (1 + \sqrt{d})^2 \log(1 + \sqrt{d})}{[(1 + \sqrt{d})Ad(\log d)]^{1/2}} - \frac{(1 + \sqrt{d})(Y_k(0)/3 + Y_n(0))}{[\sqrt{d}Ad(\log d)]^{1/2}} \\ &\geq \frac{Ad(\log d)}{4\sqrt{Ad}^{3/4}(\log d)^{1/2}} - \frac{2(Y_k(0)/3 + Y_n(0))d^{1/4}}{\sqrt{Ad} \log d} \\ &= \frac{\sqrt{A}}{4} d^{1/4} (\log d)^{1/2} - \frac{2(Y_k(0)/3 + Y_n(0))}{\sqrt{Ad}^{1/4} (\log d)^{1/2}}. \end{aligned}$$

Also, for any $d \in \mathbb{N}$,

$$\frac{\Gamma_d^A - Y_k(0)}{3\sqrt{t_d^A}} = \frac{2A\sqrt{d}(\log d)}{\sqrt{Ad}(\log d)} - \frac{Y_k(0)}{3\sqrt{Ad}(\log d)} = 2\sqrt{A \log d} - \frac{Y_k(0)}{3\sqrt{Ad}(\log d)}.$$

Thus, using (4.1), for any $d \geq d_0$, on the event $\mathcal{A}(n)$,

$$\begin{aligned} \mathbb{P}\left(\sup_{s \in [0, t_d^A]} Y_{(k)}(s) \geq \Gamma_d^A \mid \mathbf{Y}(0)\right) &\leq 2\bar{\Phi}\left(\frac{\sqrt{A}}{4} d^{1/4} (\log d)^{1/2} - \frac{2(Y_k(0)/3 + Y_n(0))}{\sqrt{Ad}^{1/4} (\log d)^{1/2}}\right) \\ &\quad + 4(k+1)\bar{\Phi}\left(2\sqrt{A \log d} - \frac{Y_k(0)}{3\sqrt{Ad}(\log d)}\right). \end{aligned}$$

Note that the right-hand side converges to 0 as $d \rightarrow \infty$. Hence, for any $n \in \mathbb{N}$,

$$\limsup_{d \rightarrow \infty} \mathbb{P}\left(\sup_{s \in [0, t_d^A]} Y_{(k)}(s) \geq \Gamma_d^A\right) \leq \mathbb{P}(\mathcal{A}(n)^c).$$

As $n \in \mathbb{N}$ is arbitrary, we conclude from (4.4) that

$$(4.5) \quad \limsup_{d \rightarrow \infty} \mathbb{P}\left(\sup_{s \in [0, t_d^A]} Y_{(k)}(s) \geq \Gamma_d^A\right) = 0.$$

This proves the second convergence statement in the lemma. For the first statement, define the events $\mathcal{A}^*(d) := \{Y_i(0) \geq 7A\sqrt{i}(\log i) \text{ for all } i \geq d\}$, $d \in \mathbb{N}$. By assumption (4.3),

$$(4.6) \quad \mathbb{P}(\mathcal{A}^*(d)) \rightarrow 1 \quad \text{as } d \rightarrow \infty.$$

For any $d \in \mathbb{N}$, on the set $\mathcal{A}^*(d)$, for all $i \geq d$,

$$\frac{Y_i(0) - \Gamma_d^A}{\sqrt{t_d^A}} \geq \frac{7A\sqrt{i}(\log i) - 6A\sqrt{d}(\log d)}{\sqrt{Ad(\log d)}} \geq \sqrt{A \log i}.$$

Hence, using (4.2), for all $d \geq 2$, on the event $\mathcal{A}^*(d)$,

$$(4.7) \quad \begin{aligned} \mathbb{P}\left(\inf_{s \in [0, t_d^A]} \inf_{i \geq d} Y_i(s) \leq \Gamma_d^A \mid \mathbf{Y}(0)\right) &\leq 2 \sum_{i \geq d} \bar{\Phi}(\sqrt{A \log i}) \\ &\leq \sum_{i \geq d} \frac{2}{\sqrt{2\pi A \log i}} e^{-A(\log i)/2} \leq \sum_{i \geq d} i^{-A/2} \leq \frac{1}{d-1}. \end{aligned}$$

Hence, from (4.6) and (4.7),

$$(4.8) \quad \limsup_{d \rightarrow \infty} \mathbb{P}\left(\inf_{s \in [0, t_d^A]} \inf_{i \geq d} Y_i(s) \leq \Gamma_d^A\right) = 0.$$

This proves the first convergence statement in the lemma and completes the proof of the lemma. $\square$

The next few lemmas will be used to study the case when the starting configuration satisfies (2.5).

LEMMA 4.3. *Fix $\theta > 0$. Suppose there exists $i_\theta \geq 2$, such that $Y_i(0) \geq \theta \log i$ for all $i \geq i_\theta$. Then there exists $\delta_\theta \in (0, 1)$ and positive constants C_1, C_2 (depending on θ) such that, a.s.,*

$$\mathbb{P}\left(\inf_{s \in [0, \delta_\theta t]} \inf_{i \geq e^t} Y_i(s) \leq \theta t/2 \mid \mathbf{Y}(0)\right) \leq C_1 e^{-C_2 t} \quad \text{for all } t \geq 0.$$

PROOF. For any $i \geq e^t \vee i_\theta$, $\delta \in (0, 1)$, by the assumption on the starting configuration,

$$\begin{aligned} \bar{\Phi}\left(\frac{Y_i(0) - \theta t/2}{\sqrt{\delta t}}\right) &\leq \bar{\Phi}\left(\frac{\theta \log i - \theta t/2}{\sqrt{\delta t}}\right) \\ &\leq \bar{\Phi}\left(\frac{\theta \log i}{2\sqrt{\delta t}}\right) \leq \frac{1}{\sqrt{2\pi}} \frac{2\sqrt{\delta t}}{\theta \log i} \exp\left\{-\frac{(\log i)^2 \theta^2}{8\delta t}\right\}. \end{aligned}$$

Hence, using (4.2) with $d = \lceil e^t \rceil$, $\Gamma = \theta t/2$ and $t = \delta t$, we obtain $t_0 > 0$ such that for all $t \geq t_0$,

$$\begin{aligned} \mathbb{P}\left(\inf_{s \in [0, \delta t]} \inf_{i \geq e^t} Y_i(s) \leq \theta t/2 \mid \mathbf{Y}(0)\right) &\leq 2 \sum_{i \geq e^t} \bar{\Phi}\left(\frac{Y_i(0) - \theta t/2}{\sqrt{\delta t}}\right) \leq 2 \sum_{i \geq e^t} \exp\left\{-\frac{(\log i)^2 \theta^2}{8\delta t}\right\} \\ &= 2 \sum_{j=1}^{\infty} \sum_{e^{jt} \leq i < e^{(j+1)t}} \exp\left\{-\frac{(\log i)^2 \theta^2}{8\delta t}\right\} \\ &\leq 2 \sum_{j=1}^{\infty} e^{(j+1)t} \exp\left\{-\frac{\theta^2 j^2 t}{8\delta}\right\}. \end{aligned}$$

By the above bound, we can obtain $\delta_\theta \in (0, 1)$ and $C_1, C_2 > 0$ such that for all $t \geq 0$,

$$\mathbb{P}\left(\inf_{s \in [0, \delta_\theta t]} \inf_{i \geq e^t} Y_i(s) \leq \theta t/2 | \mathbf{Y}(0)\right) \leq C_1 e^{-C_2 t},$$

proving the lemma. $\square$

LEMMA 4.4. *For any $k \in \mathbb{N}, \alpha > 0$,*

$$\lim_{t \rightarrow \infty} \mathbb{P}\left(\sup_{s \in [0, t]} Y_{(k)}(s) \geq \alpha t\right) = 0.$$

PROOF. For any $A \geq 1, t > 0$, using (4.1) with $\Gamma = \alpha t$, we obtain for any $l \geq k$,

$$\mathbb{P}\left(\sup_{s \in [0, t]} Y_{(k)}(s) \geq \alpha t, Y_l(0) \leq A\right) \leq 2\bar{\Phi}\left(\frac{l(\alpha t - A)/3 - t - Al}{\sqrt{lt}}\right) + 4(k+1)\bar{\Phi}\left(\frac{\alpha t - A}{3\sqrt{t}}\right).$$

Hence, choosing and fixing any $l \geq k \vee (18/\alpha)$, for all $t \geq 16A/(3\alpha)$,

$$\mathbb{P}\left(\sup_{s \in [0, t]} Y_{(k)}(s) \geq \alpha t, Y_l(0) \leq A\right) \leq 2\bar{\Phi}\left(\frac{\sqrt{t}}{2\sqrt{l}}\right) + 4(k+1)\bar{\Phi}\left(\frac{13\alpha\sqrt{t}}{48}\right).$$

Hence,

$$\limsup_{t \rightarrow \infty} \mathbb{P}\left(\sup_{s \in [0, t]} Y_{(k)}(s) \geq \alpha t\right) \leq \mathbb{P}(Y_l(0) > A).$$

As $A \geq 1$ is arbitrary, the lemma follows upon taking a limit as $A \rightarrow \infty$ in the above bound. $\square$

Recall that, for $\mathbf{v} = (v_1, v_2, \dots)^T \in \mathbb{R}_+^\infty$, $s(\mathbf{v})$ denotes the vector in $\mathbb{R}_+^\infty$ whose i th coordinate is $v_1 + \dots + v_i$, $i \in \mathbb{N}$. In the following, for a $\mathbb{R}_+^\infty$ -valued random vector $\mathbf{v}$, we will say $s(\mathbf{v})$ satisfies (3.1) if, almost surely, (3.1) holds with $x_0 = 0$ and $x_i = (s(\mathbf{v}))_i$ for $i \geq 1$.

LEMMA 4.5. *Fix $a > 0$. Let μ be a probability measure on $\mathbb{R}_+^\infty$ such that there exists a coupling $(\mathbf{U}, \mathbf{V}_a)$ of μ and π_a such that, almost surely, (2.5) holds. Then, almost surely, there exists $i_0 \geq 2$ such that $s(\mathbf{U})_i \geq (4a)^{-1} \log i$ for all $i \geq i_0$. In particular, $s(\mathbf{U})$ satisfies (3.1), and consequently $\mu \in \mathcal{S}$. Furthermore, $s(\mathbf{U} \wedge \mathbf{V}_a)$ and $s(\mathbf{V}_a)$ satisfy (3.1) as well.*

PROOF. We will show that, almost surely, there exists $i'_0 \geq 2$ such that

$$(4.9) \quad \sum_{j=1}^i V_{a,j} \geq \frac{1}{2a} \log i \quad \text{for all } i \geq i'_0.$$

It will then follow, by (2.5), that there is a $i_0 \geq i'_0$ such that for all $i \geq i_0$,

$$(4.10) \quad s(\mathbf{U})_i = \sum_{j=1}^i U_j \geq \sum_{j=1}^i V_{a,j} - \sum_{j=1}^i |V_{a,j} - U_j| \geq \frac{1}{2a} \log i - \frac{\log i}{\log \log i} \geq \frac{1}{4a} \log i.$$

The same argument shows that $s(\mathbf{U} \wedge \mathbf{V}_a)_i \geq (4a)^{-1} \log i$ for $i \geq i_0$ as well. Thus, in order to prove the lemma, it suffices to prove (4.9). Note that $E_j := (2 + ja)V_{a,j}$, $j \in \mathbb{N}$, are i.i.d. exponential random variables with mean one. Define

$$S_i := \sum_{j=1}^i V_{a,j} = \sum_{j=1}^i \frac{E_j}{2 + ja}, \quad i \in \mathbb{N}.$$

Note that

$$\mathbb{E}(S_i) = \sum_{j=1}^i \frac{1}{2+ja} \geq \int_1^{i+1} \frac{1}{2+as} ds = \frac{1}{a} \log\left(\frac{2+a(i+1)}{2+a}\right).$$

Moreover,

$$\text{Var}(S_i) = \sum_{j=1}^i \frac{1}{(2+ja)^2} \leq \sum_{j=1}^{\infty} \frac{1}{(2+ja)^2} < \infty.$$

Hence, $M_i := S_i - \mathbb{E}(S_i)$, $i \in \mathbb{N}$, is an L^2 -bounded martingale. Therefore, by the martingale convergence theorem [22], Theorem 11.10, there exists a random variable M_∞ with $\mathbb{E}(M_\infty^2) < \infty$ such that $M_i \rightarrow M_\infty$ as $i \rightarrow \infty$, a.s. and in L^2 . Thus, almost surely, there exists $\underline{M} \in (-\infty, \infty)$ and $i_0'' \geq 2$ such that

$$S_i \geq \frac{1}{a} \log\left(\frac{2+a(i+1)}{2+a}\right) - \underline{M} \quad \text{for all } i \geq i_0''.$$

The estimate in (4.9), and hence the lemma, is immediate from the above. $\square$

The following lemma uses the above three lemmas to quantify the influence of far away coordinates when the starting configuration satisfies (2.5). Note that Lemma 4.5 shows that, if $\mathbf{U}$ is as in that lemma, then the distribution of $(0, s(\mathbf{U})^T)^T$ is in $\mathcal{S}_0$.

LEMMA 4.6. *Fix $a > 0$. Let μ be a probability measure on $\mathbb{R}_+^\infty$ such that there exists a coupling $(\mathbf{U}, \mathbf{V}_a)$ of μ and π_a such that, almost surely, (2.5) holds. Let $\mathbf{Y}(\cdot)$ be the infinite Atlas model given as the unique weak solution of (1.2) with $\mathbf{Y}(0)$ distributed as $(0, s(\mathbf{U})^T)^T$. Then there exists $\delta_0 \in (0, 1)$ such that,*

$$\limsup_{d \rightarrow \infty} \mathbb{P}\left(\inf_{s \in [0, \delta_0 \log d]} \inf_{i \geq d} Y_i(s) \leq \frac{1}{8a} \log d\right) = 0,$$

and for any $k \in \mathbb{N}$

$$\limsup_{d \rightarrow \infty} \mathbb{P}\left(\sup_{s \in [0, \delta_0 \log d]} Y_{(k)}(s) \geq \frac{1}{8a} \log d\right) = 0.$$

PROOF. Take $\delta_0 := \delta_{(4a)^{-1}}$ as defined in Lemma 4.3. The second limit above follows by Lemma 4.4. To prove the first limit, define the event $\tilde{\mathcal{E}} := \{\exists i_0 \geq 2 \text{ such that } s(\mathbf{U})_i \geq (4a)^{-1} \log i \text{ for all } i \geq i_0\}$. By Lemma 4.3, with $\theta = (4a)^{-1}$,

$$\limsup_{d \rightarrow \infty} \mathbb{P}\left(\inf_{s \in [0, \delta_0 \log d]} \inf_{i \geq d} Y_i(s) \leq \frac{1}{8a} \log d\right) \leq \mathbb{P}(\tilde{\mathcal{E}}^c).$$

The right-hand side above is zero by Lemma 4.5. This proves the lemma. $\square$

5. Analyzing excursions of synchronous couplings. In this section, we will estimate the rate of decay of the L^1 distance between synchronously coupled infinite ordered Atlas processes by defining and analyzing suitable excursions of the processes. Each such excursion will capture an event which ensures that the L^1 distance decreases by a fixed amount. Theorems 2.1 and 2.4 will then be proved by estimating the number of such excursions on the time interval $[0, t]$ for large t .

Let $(\mathbf{U}_1, \mathbf{U}_2)$ be an $\mathbb{R}_+^\infty \times \mathbb{R}_+^\infty$ -valued random variable such that $\mathbf{U}_1 \leq \mathbf{U}_2$ and $s(\mathbf{U}_1)$ (and thus $s(\mathbf{U}_2)$) satisfies (3.1). Consider the synchronously coupled copies $\mathbf{X}^{\mathbf{U}_1}(\cdot)$ and

$\mathbf{X}^{\mathbf{U}_2}(\cdot)$ of the infinite ordered Atlas model (in the sense of Proposition 3.1(ii)) starting from $(0, s(\mathbf{U}_1)^T)^T$ and $(0, s(\mathbf{U}_2)^T)^T$, respectively, driven by Brownian motions $\{B_i\}_{i \in \mathbb{N}_0}$ (that are independent of $\mathbf{U}_1, \mathbf{U}_2$). Let $\mathbf{Z}^{\mathbf{U}_1}(\cdot)$ and $\mathbf{Z}^{\mathbf{U}_2}(\cdot)$ be the associated gap processes as introduced in Proposition 3.1(iii), and let $\Delta \mathbf{Z}(\cdot) := \mathbf{Z}^{\mathbf{U}_2}(\cdot) - \mathbf{Z}^{\mathbf{U}_1}(\cdot)$.

Fix $k \in \mathbb{N}$. For $s \geq 0$, define the following stopping times: $T_0^k(s) := s$, and

$$(5.1) \quad T_j^k(s) := \inf\{u \geq T_{j-1}^k(s) : \Delta Z_{k-j+1}(u) = 0\}, \quad 1 \leq j \leq k.$$

Set $\mathcal{T}_k(s) := T_k^k(s)$. We will analyze excursions of $\Delta \mathbf{Z}$ the coupled processes defined by the following stopping times. Fix $\epsilon > 0$. Define $\sigma_0 := 0$, $\sigma_1 := \inf\{s \geq 0 : \Delta Z_k(s) \geq \epsilon\}$, and for $j \geq 0$,

$$\begin{aligned} \sigma_{2j+2} &:= \inf\{s \geq \mathcal{T}_k(\sigma_{2j+1}) : \Delta Z_k(s) = 0\}, \\ \sigma_{2j+3} &:= \inf\{s \geq \sigma_{2j+2} : \Delta Z_k(s) \geq \epsilon\}, \end{aligned}$$

where the infimum in (5.1) and in the above two definitions is taken to be ∞ if the corresponding sets are empty. Although the stopping times above depend on k , we suppress this dependence for notational convenience. For $T \geq 0$, define the number of excursions until time T by

$$\mathcal{N}_T := \begin{cases} \sup\{j \geq 0 : \sigma_{2j+1} \leq T\} + 1 & \text{if } \sigma_1 \leq T, \\ 0 & \text{if } \sigma_1 > T. \end{cases}$$

In the next two lemmas, we work with a fixed $k \in \mathbb{N}$ and stopping times $\{\sigma_j : j \geq 0\}$ defined for this fixed k .

Each of the above excursions ensure a decrease of the L^1 norm of $\Delta \mathbf{Z}(\cdot)$ by a fixed amount as described by the following lemma.

LEMMA 5.1. *Assume $\sum_{j=1}^{\infty} j \Delta Z_j(0) < \infty$. Then, for any $j \geq 0$, when $\sigma_{2j+2} < \infty$,*

$$\sum_{i=1}^{\infty} \Delta Z_i(\sigma_{2j+2}) - \sum_{i=1}^{\infty} \Delta Z_i(\sigma_{2j+1}) \leq -\epsilon/2^k.$$

PROOF. By Lemma 3.3,

$$\sum_{i=1}^{\infty} \Delta Z_i(t) = \sum_{i=1}^{\infty} \Delta Z_i(0) + \frac{1}{2} \Delta L_1(t), \quad t \geq 0.$$

Thus, it suffices to show that

$$(5.2) \quad \Delta L_1(\sigma_{2j+2}) - \Delta L_1(\sigma_{2j+1}) \leq -\epsilon/2^{k-1}.$$

Recalling the stopping times $\{T_l^k(\cdot)\}_{0 \leq l \leq k}$ from (5.1), note that for any $1 \leq i \leq k$, $j \in \mathbb{N}_0$,

$$\begin{aligned} (5.3) \quad 0 &= \Delta Z_i(T_{k-i+1}^k(\sigma_{2j+1})) \\ &= \Delta Z_i(\sigma_{2j+1}) + (\Delta L_i(T_{k-i+1}^k(\sigma_{2j+1})) - \Delta L_i(\sigma_{2j+1})) \\ &\quad - \frac{1}{2}(\Delta L_{i-1}(T_{k-i+1}^k(\sigma_{2j+1})) - \Delta L_{i-1}(\sigma_{2j+1})) \\ &\quad - \frac{1}{2}(\Delta L_{i+1}(T_{k-i+1}^k(\sigma_{2j+1})) - \Delta L_{i+1}(\sigma_{2j+1})). \end{aligned}$$

Recalling that $\mathbf{U}_1 \leq \mathbf{U}_2$, by Proposition 3.1(iii), for each $i \in \mathbb{N}$, $j \in \mathbb{N}_0$, $\Delta Z_i(\sigma_{2j+1}) \geq 0$ and $\Delta L_i(\cdot)$ is nonpositive and nonincreasing. Thus, from (5.3) we obtain for any $1 \leq i \leq k$,

$$\begin{aligned} -(\Delta L_i(T_{k-i+1}^k(\sigma_{2j+1})) - \Delta L_i(\sigma_{2j+1})) &\geq -\frac{1}{2}(\Delta L_{i+1}(T_{k-i+1}^k(\sigma_{2j+1})) - \Delta L_{i+1}(\sigma_{2j+1})) \\ &\geq -\frac{1}{2}(\Delta L_{i+1}(T_{k-i}^k(\sigma_{2j+1})) - \Delta L_{i+1}(\sigma_{2j+1})). \end{aligned}$$

Using the above bound recursively for $i = 1, \dots, k-1$, we get, for any $j \in \mathbb{N}_0$,

$$\begin{aligned} &-(\Delta L_1(T_k^k(\sigma_{2j+1})) - \Delta L_1(\sigma_{2j+1})) \\ (5.4) \quad &\geq -\frac{1}{2}(\Delta L_2(T_{k-1}^k(\sigma_{2j+1})) - \Delta L_2(\sigma_{2j+1})) \\ &\geq \dots \geq -\frac{1}{2^{k-1}}(\Delta L_k(T_1^k(\sigma_{2j+1})) - \Delta L_k(\sigma_{2j+1})). \end{aligned}$$

Moreover, from (5.3) with $i = k$, again using Proposition 3.1(iii),

$$(5.5) \quad -(\Delta L_k(T_1^k(\sigma_{2j+1})) - \Delta L_k(\sigma_{2j+1})) \geq \Delta Z_k(\sigma_{2j+1}) \geq \epsilon.$$

Recalling $\sigma_{2j+2} \geq \mathcal{T}_k(\sigma_{2j+1}) = T_k^k(\sigma_{2j+1})$ and $\Delta L_1(\cdot)$ is nonincreasing, (5.2) follows from (5.4) and (5.5). This proves the lemma. $\square$

The following lemma will be used to control the maximum length of excursions in a large time interval.

LEMMA 5.2. *Suppose there exists $D \geq 1$ such that the distribution of $\mathbf{U}_2/D$ is stochastically dominated by π . Let $T_k := \inf\{S \geq e : 48D^2k(k+1)^2 \log T \leq T \text{ for all } T \geq S\}$. Then, for any $n \in \mathbb{N}$, $T \geq T_k$,*

$$\begin{aligned} &\mathbb{P}(\sigma_1 \leq T, \sigma_{2j+2} - \sigma_{2j+1} > 48D^2k(k+1)^2 \log T \text{ for some } 0 \leq j \leq \mathcal{N}_T - 1) \\ &\leq \mathbb{P}(\mathcal{N}_T > n) + 5knT^{-2}. \end{aligned}$$

PROOF. We can assume without loss of generality that, in addition to $\mathbf{U}_1, \mathbf{U}_2$, we are given another $\mathbb{R}_+^\infty$ -valued random variable $\mathbf{V}$ (on the same probability space) such that $\mathbf{U}_1 \leq \mathbf{U}_2 \leq D\mathbf{V}$. In addition to infinite ordered Atlas models $\mathbf{X}^{\mathbf{U}_i}$, with $\mathbf{X}^{\mathbf{U}_i}(0) = \mathbf{U}_i$ for $i = 1, 2$, driven by Brownian motions $\{B_i\}$, we also construct infinite ordered Atlas models $\mathbf{X}^{D\mathbf{V}}$ and $\mathbf{X}^{[\gamma]}$ with drifts 1 and $\gamma = 1/D$, respectively (the latter defined as in Proposition 3.1(iv)), satisfying $\mathbf{X}^{D\mathbf{V}}(0) = \mathbf{X}^{[\gamma]}(0) = D\mathbf{V}$, and driven by the same Brownian motions $\{B_i\}$ (that are independent of $\mathbf{U}_1, \mathbf{U}_2, \mathbf{V}$). The associated gap processes are denoted as $\mathbf{Z}^{D\mathbf{V}}$ and $\mathbf{Z}^{[\gamma]}$. In order to note the dependence of γ on D , we will denote the processes $\mathbf{X}^{[\gamma]}, \mathbf{Z}^{[\gamma]}$ as $\mathbf{X}^{[D]}, \mathbf{Z}^{[D]}$, respectively.

By Proposition 3.1(iii) and (iv),

$$(5.6) \quad \mathbf{Z}^{\mathbf{U}_2}(t) \leq \mathbf{Z}^{D\mathbf{V}}(t) \leq \mathbf{Z}^{[D]}(t) \quad \text{for all } t \geq 0.$$

For $u \geq 0$, define

$$S_i(u) := \inf\{s \geq u : Z_i^{[D]}(s) = 0\} \quad \text{for } 1 \leq i \leq k, \quad S(u) := \max_{1 \leq i \leq k} S_i(u).$$

Write $A := 24D^2k(k+1)$. Note that $2A(k+1) \log T \leq T$ for all $T \geq T_k$. Suppose for some $j \geq 0$, $T \geq T_k$, the following event holds:

$$\mathcal{E}_{T,j} := \{\sigma_{2j+1} \leq T, \sigma_{2j+2} - \sigma_{2j+1} > 2A(k+1) \log T\}.$$

Then, either

$$(5.7) \quad \text{there is } 1 \leq l \leq k \text{ such that } T_l^k(\sigma_{2j+1}) - T_{l-1}^k(\sigma_{2j+1}) > 2A \log T,$$

or

$$(5.8) \quad \sigma_{2j+2} - \mathcal{T}_k(\sigma_{2j+1}) = T_1^k(\mathcal{T}_k(\sigma_{2j+1})) - \mathcal{T}_k(\sigma_{2j+1}) > 2A \log T.$$

Suppose (5.7) holds. Let l be the minimum integer in $\{1, \dots, k\}$ for which (5.7) holds. Then recalling $\sigma_{2j+1} \leq T$,

$$T_{l-1}^k(\sigma_{2j+1}) \leq \sigma_{2j+1} + 2A(l-1) \log T \leq T + 2A(k-1) \log T \leq 2T - A \log T,$$

where the last inequality is true because $T \geq T_k$. Similarly, if (5.8) holds and (5.7) does not hold for any $1 \leq l \leq k$, then

$$\mathcal{T}_k(\sigma_{2j+1}) \leq \sigma_{2j+1} + 2Ak \log T \leq 2T - A \log T.$$

Observe that for any $1 \leq l \leq k$, $j \geq 0$, by (5.6),

$$\begin{aligned} T_l^k(\sigma_{2j+1}) &:= \inf\{u \geq T_{l-1}^k(\sigma_{2j+1}) : \Delta Z_{k-l+1}(u) = 0\} \\ &\leq \inf\{u \geq T_{l-1}^k(\sigma_{2j+1}) : Z_{k-l+1}^{(D)}(u) = 0\} = S_{k-l+1}(T_{l-1}^k(\sigma_{2j+1})), \end{aligned}$$

and similarly,

$$\sigma_{2j+2} = T_1^k(\mathcal{T}_k(\sigma_{2j+1})) \leq S_k(\mathcal{T}_k(\sigma_{2j+1})).$$

Hence, for any $T \geq T_k$,

$$\mathcal{E}_{T,j} \subseteq \mathcal{E}'_T := \{\exists t \in [0, 2T - A \log T] \text{ such that } S(t) - t > 2A \log T\}.$$

From this observation and the union bound, for any $n \in \mathbb{N}$, $T \geq T_k$,

$$\begin{aligned} (5.9) \quad &\mathbb{P}(\sigma_1 \leq T, \sigma_{2j+2} - \sigma_{2j+1} > 2A(k+1) \log T \text{ for some } 0 \leq j \leq \mathcal{N}_T - 1) \\ &\leq \mathbb{P}(\mathcal{N}_T > n) + \sum_{j=0}^{n-1} \mathbb{P}(\mathcal{E}_{T,j}) \leq \mathbb{P}(\mathcal{N}_T > n) + n\mathbb{P}(\mathcal{E}'_T). \end{aligned}$$

In the rest of the proof, we will estimate $\mathbb{P}(\mathcal{E}'_T)$.

For any $T > 0$, define $n(T) := \lceil 2T/(A \log T) \rceil \vee 2$ and

$$t_l := Al \log T, \quad 0 \leq l \leq n(T).$$

Then $\bigcup_{l=0}^{n(T)} [t_l, t_{l+1}] \supset [0, 2T]$ and each interval $[t_l, t_{l+1}]$ is of length $A \log T$.

Suppose $S(t) - t > 2A \log T$ for some $t \in [t_l, t_{l+1}]$, $0 \leq l \leq n(T) - 2$. Then there exists $1 \leq i \leq k$ such that $S_i(t_{l+1}) - t_{l+1} > A \log T$. Hence, for any $T \geq T_k$,

$$(5.10) \quad \mathbb{P}(\mathcal{E}'_T) \leq \mathbb{P}(\exists 0 \leq l \leq n(T) - 2, 1 \leq i \leq k \text{ such that } S_i(t_{l+1}) - t_{l+1} > A \log T).$$

Note that, by scaling properties of the Atlas model, the process $\mathbf{Z}^{(D)}(\cdot)$ is stationary with $\mathbf{Z}^{(D)}(t) \sim \bigotimes_{i=1}^{\infty} \text{Exp}(2/D)$ for all $t \geq 0$. Define for any $T \geq e$ the event

$$\mathcal{A}_T := \{Z_i^{(D)}(t_l) \leq 24D \log T \text{ for all } 1 \leq i \leq k, 0 \leq l \leq n(T) - 1\}.$$

By the stationarity of $\mathbf{Z}^{(D)}(\cdot)$ and the union bound, for any $T \geq e$,

$$(5.11) \quad \mathbb{P}(\mathcal{A}_T^c) \leq kn(T)e^{-48 \log T} \leq k \left(2 + \frac{2T}{A \log T}\right) T^{-48} \leq 4kT^{-47}.$$

Denote by $\mathbf{L}^{\{D\}} = (L_0^{\{D\}}(\cdot), L_1^{\{D\}}(\cdot), \dots)$ the collision local times associated with the infinite ordered Atlas model $\mathbf{X}^{\{D\}}$ (with drift $\gamma = 1/D$). Then, for any $0 \leq l \leq n(T) - 1$, $1 \leq i \leq k$, $t \geq 0$, $T \geq e$,

$$(5.12) \quad \{S_i(t_l) - t_l > t\} = \{X_{i-1}^{\{D\}}(s) < X_i^{\{D\}}(s) \text{ for all } s \in [t_l, t_l + t]\}.$$

If the above event holds, then $L_i(t_l + s) - L_i(t_l) = 0$ for all $0 \leq s \leq t$. Hence, using the fact that $X_j^{\{D\}}(\cdot)$'s are ranked and applying (3.4), we obtain for any $0 \leq s \leq t$, on the event in (5.12),

$$\begin{aligned} X_{i-1}^{\{D\}}(s + t_l) &\geq \frac{1}{i} \sum_{q=0}^{i-1} X_q^{\{D\}}(s + t_l) \\ &= \frac{s}{Di} + \frac{1}{i} \sum_{q=0}^{i-1} X_q^{\{D\}}(t_l) + \frac{1}{i} \sum_{q=0}^{i-1} (B_q(s + t_l) - B_q(t_l)), \end{aligned}$$

and

$$X_i^{\{D\}}(s + t_l) \leq X_i^{\{D\}}(t_l) + (B_i(s + t_l) - B_i(t_l)).$$

Moreover, if $\mathcal{A}_T$ holds, then for any $0 \leq l \leq n(T) - 1$, $1 \leq q \leq i - 1$,

$$X_i^{\{D\}}(t_l) - X_q^{\{D\}}(t_l) = \sum_{p=q+1}^i Z_p^{\{D\}}(t_l) \leq 24D(i - q) \log T.$$

Hence, defining the new Brownian motions

$$\tilde{B}_1(s) := \frac{1}{\sqrt{i}} \sum_{q=0}^{i-1} (B_q(s + t_l) - B_q(t_l)), \quad \tilde{B}_2(s) := (B_i(s + t_l) - B_i(t_l)), \quad 0 \leq s \leq t,$$

and using the above observations in (5.12), we obtain for any $0 \leq l \leq n(T) - 1$, $1 \leq i \leq k$, $t \geq 0$, $T \geq e$,

$$\begin{aligned} &\mathbb{P}(S_i(t_l) - t_l > t, \mathcal{A}_T) \\ &\leq \mathbb{P}\left(\frac{t}{Di} + \frac{1}{i} \sum_{q=0}^{i-1} X_q^{\{D\}}(t_l) + \frac{1}{\sqrt{i}} \tilde{B}_1(t) < X_i^{\{D\}}(t_l) + \tilde{B}_2(t), \mathcal{A}_T\right) \\ &= \mathbb{P}\left(\frac{1}{\sqrt{i}} \tilde{B}_1(t) - \tilde{B}_2(t) < -\frac{t}{Di} + \frac{1}{i} \sum_{q=0}^{i-1} (X_i^{\{D\}}(t_l) - X_q^{\{D\}}(t_l)), \mathcal{A}_T\right) \\ &\leq \mathbb{P}\left(\frac{1}{\sqrt{i}} \tilde{B}_1(t) - \tilde{B}_2(t) < -\frac{t}{Di} + \frac{1}{i} \sum_{q=0}^{i-1} 24D(i - q) \log T\right) \\ &= \mathbb{P}\left(\frac{1}{\sqrt{i}} \tilde{B}_1(t) - \tilde{B}_2(t) < -\frac{t}{Di} + 12D(i + 1) \log T\right). \end{aligned}$$

Hence, taking $t = 24D^2i(i + 1) \log T$ and using the fact that $\frac{1}{\sqrt{i}} \tilde{B}_1(24D^2i(i + 1) \log T) - \tilde{B}_2(24D^2i(i + 1) \log T)$ is a mean zero normal random variable with variance $24D^2(i +$

$1)^2 \log T$, we obtain for any $0 \leq l \leq n(T) - 1$, $1 \leq i \leq k$, $T \geq e$,

$$\begin{aligned}
 & \mathbb{P}(S_i(t_l) - t_l > 24D^2 i(i+1) \log T, \mathcal{A}_T) \\
 (5.13) \quad & \leq \bar{\Phi}\left(-\frac{\sqrt{24 \log T}}{2}\right) \\
 & \leq \frac{2}{\sqrt{48\pi \log T}} \exp\{-24 \log T/8\} \leq \frac{1}{2\sqrt{3\pi}} T^{-3}.
 \end{aligned}$$

Thus, recalling that $A = 24D^2 k(k+1)$, using (5.11) and (5.13) in (5.10), we obtain for any $T \geq T_k$,

$$\begin{aligned}
 \mathbb{P}(\mathcal{E}'_T) & \leq \mathbb{P}(\exists 0 \leq l \leq n(T) - 2, 1 \leq i \leq k \text{ such that } S_i(t_{l+1}) - t_{l+1} > A \log T) \\
 & \leq \sum_{l=0}^{n(T)-2} \sum_{i=1}^k \mathbb{P}(S_i(t_{l+1}) - t_{l+1} > 24D^2 i(i+1) \log T, \mathcal{A}_T) + \mathbb{P}(\mathcal{A}_T^c) \\
 & \leq n(T)k \frac{1}{2\sqrt{3\pi}} T^{-3} + 4kT^{-47} \leq \frac{2k}{\sqrt{3\pi}} T^{-2} + 4kT^{-47}.
 \end{aligned}$$

Finally, using the above bound in (5.9), for any $n \in \mathbb{N}$, $T \geq T_k$,

$$\begin{aligned}
 & \mathbb{P}(\sigma_1 \leq T, \sigma_{2j+2} - \sigma_{2j+1} > 2A(k+1) \log T \text{ for some } 0 \leq j \leq \mathcal{N}_T - 1) \\
 & \leq \mathbb{P}(\mathcal{N}_T > n) + \frac{2kn}{\sqrt{3\pi}} T^{-2} + 4knT^{-47} \leq \mathbb{P}(\mathcal{N}_T > n) + 5knT^{-2},
 \end{aligned}$$

which proves the lemma. $\square$

6. Proofs of main results. In this section we will prove Theorem 2.1 and Theorem 2.4, and their corollaries (i.e., Corollary 2.2 and Corollary 2.5).

6.1. Proof of Theorem 2.1. The key idea in the proof of Theorem 2.1 is to sandwich the initial gap vectors $\mathbf{U}$ and $\mathbf{V}$ between the $\mathbb{R}_+^\infty$ -valued random vectors $\mathbf{U} \wedge \mathbf{V}$ and $\mathbf{U} \vee \mathbf{V}$. Using monotonicity properties of synchronous couplings described in Proposition 3.1, one can bound the gap process $\mathbf{Z}^{\mathbf{U}}(\cdot)$ from above and below in a pathwise fashion by the synchronously coupled processes $\mathbf{Z}^{\mathbf{U} \vee \mathbf{V}}(\cdot)$ and $\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(\cdot)$, respectively. Lemma 6.2 uses the excursion estimates in Section 5 to analyze synchronously coupled gap processes, respectively, started from $\mathbf{V}$ and a perturbation of $\mathbf{V}$ with the first d entries replaced by those of $\mathbf{U} \wedge \mathbf{V}$. It shows that, for fixed $k \in \mathbb{N}$, $\epsilon > 0$, the average time spent ϵ distance apart by the k th coordinates of these two processes on a suitably large d -dependent time interval converges to zero in probability as $d \rightarrow \infty$.

Lemma 6.3 uses Lemma 6.2 and the quantitative estimates for the influence of far away coordinates obtained in Section 4 to show that the averaged occupancy measure for the gap process started from $\mathbf{U} \wedge \mathbf{V}$ weakly converges to π . The convergence of the averaged occupancy measure for the gap process started from $\mathbf{U} \vee \mathbf{V}$ is a consequence of the following lemma, which is a direct corollary of [27], Theorem 4.7. Theorem 2.1 follows from these observations.

LEMMA 6.1 (Corollary of Theorem 4.7 of [27]). *Let $(\mathbf{U}_\circ, \mathbf{V})$ be an $\mathbb{R}_+^\infty \times \mathbb{R}_+^\infty$ -valued random variable such that $\mathbf{U}_\circ \geq \mathbf{V}$ and $\mathbf{V} \sim \pi$. Consider the synchronously coupled copies $\mathbf{X}^{\mathbf{U}_\circ}(\cdot)$ and $\mathbf{X}^{\mathbf{V}}(\cdot)$ of the infinite ordered Atlas model (in the sense of Proposition 3.1(ii)) starting from $(0, s(\mathbf{U}_\circ)^T)^T$ and $(0, s(\mathbf{V})^T)^T$, respectively, and let $\mathbf{Z}^{\mathbf{U}_\circ}(\cdot)$ and $\mathbf{Z}^{\mathbf{V}}(\cdot)$ denote the associated gap processes as in Proposition 3.1(iii). Then the law of $\mathbf{Z}^{\mathbf{U}_\circ}(t)$ converges weakly to π as $t \rightarrow \infty$.*

LEMMA 6.2. Suppose that $\mathbf{U}$ and $\mathbf{V}$ are as in the statement of Theorem 2.1. For $d \in \mathbb{N}$, define the vector $\mathbf{U}_{(d)}$ by

$$U_{(d),i} := (U_i \wedge V_i)\mathbf{1}(i \leq d) + V_i\mathbf{1}(i > d), \quad i \in \mathbb{N}.$$

Then, $s(\mathbf{U} \wedge \mathbf{V}), s(\mathbf{U}), s(\mathbf{V}), s(\mathbf{U}_{(d)})$ all satisfy (3.1). Define infinite ordered Atlas models $\mathbf{X}^{\mathbf{U}}, \mathbf{X}^{\mathbf{V}}$ and $\mathbf{X}^{\mathbf{U}_{(d)}}$, starting from $(0, s(\mathbf{U})^T)^T, (0, s(\mathbf{V})^T)^T$ and $(0, s(\mathbf{U}_{(d)})^T)^T$, respectively, as in Proposition 3.1(i) using an infinite sequence of Brownian motions $\{B_i\}_{i \in \mathbb{N}_0}$ (independent of $\mathbf{U}, \mathbf{V}$). Let $\mathbf{Z}^{\mathbf{U}}, \mathbf{Z}^{\mathbf{V}}, \mathbf{Z}^{\mathbf{U}_{(d)}}$ be the associated gap processes. Then, for any $k \in \mathbb{N}, \epsilon \in (0, 1)$, there exists $A_0 \geq 1$ such that for any $\vartheta > 0, 1 \leq j \leq k$,

$$\mathbb{P}\left(\frac{1}{A_0 d \log d} \int_0^{A_0 \vartheta d \log d} \mathbf{1}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}_{(d)}}(s)| > \epsilon) ds \geq 1\right) \rightarrow 0 \quad \text{as } d \rightarrow \infty.$$

PROOF. First note that, by (2.1), almost surely, there exists $i' \in \mathbb{N}$ such that $s(\mathbf{U} \wedge \mathbf{V})_i \geq \sqrt{i} \log i$ for all $i \geq i'$. In particular, $s(\mathbf{U} \wedge \mathbf{V})$, and hence $s(\mathbf{U}), s(\mathbf{V})$, satisfies (3.1). As $s(\mathbf{U}_{(d)}) \geq s(\mathbf{U} \wedge \mathbf{V})$ and $s(\mathbf{U} \wedge \mathbf{V})$ satisfies (3.1), we have that $s(\mathbf{U}_{(d)})$ also satisfies (3.1).

Fix any $k \in \mathbb{N}, \epsilon \in (0, 1), \vartheta > 0$. Write $\Delta \mathbf{Z}'(\cdot) := \mathbf{Z}^{\mathbf{V}}(\cdot) - \mathbf{Z}^{\mathbf{U}_{(d)}}(\cdot)$.

Recall the stopping times $\{\sigma_j : j \geq 0\}$ and the random variables $\{\mathcal{N}_T : T \geq 0\}$ from Section 5 with our choice of $\epsilon, \mathbf{U}_1 = \mathbf{U}_{(d)}$ and $\mathbf{U}_2 = \mathbf{V}$. For $A \geq 4, d \in \mathbb{N}$, recall from Lemma 4.2 that $t_d^A := Ad \log d$, and write $\mathcal{N}_{A, \vartheta, d} := \mathcal{N}_{\vartheta t_d^A}$.

As $\sum_{j=1}^{\infty} j \Delta Z'_j(0) \leq \sum_{j=1}^d j V_j < \infty$, by Lemma 5.1, for any $j \geq 0$, when $\sigma_{2j+2} < \infty$,

$$\sum_{i=1}^{\infty} \Delta Z'_i(\sigma_{2j+2}) - \sum_{i=1}^{\infty} \Delta Z'_i(\sigma_{2j+1}) \leq -\epsilon/2^k.$$

Recalling the monotonicity property in Proposition 3.1(iii), for any $A > 0$,

$$(6.1) \quad \mathbb{P}(\mathcal{N}_{A, \vartheta, d} > 2^k \epsilon^{-1} d + 1) \leq \mathbb{P}\left(\sum_{i=1}^d \Delta Z'_i(0) > d\right) \leq \mathbb{P}\left(\sum_{i=1}^d V_i > d\right) \leq e^{-d/8}.$$

For $d \in \mathbb{N}, A \geq 4$, define the event

$$\mathcal{E}_{d,A} := \{\sigma_1 \leq \vartheta t_d^A, \sigma_{2j+2} - \sigma_{2j+1} \leq 48k(k+1)^2 \log(\vartheta t_d^A) \text{ for all } 0 \leq j \leq \mathcal{N}_{A, \vartheta, d} - 1\}.$$

By Lemma 5.2 (with $D = 1, T = \vartheta t_d^A$ and $n = 2^k \epsilon^{-1} d + 1$) and (6.1), there exists $d_0 \in \mathbb{N}$ such that for any $d \geq d_0, A \geq 4$,

$$(6.2) \quad \mathbb{P}(\mathcal{E}_{d,A}^c \cap \{\sigma_1 \leq \vartheta t_d^A\}) \leq e^{-d/8} + \frac{10k2^k \epsilon^{-1}}{A^2 \vartheta^2 d (\log d)^2}.$$

For $d \geq A \vee \vartheta \geq 4$, on the event $\mathcal{E}_{d,A}$,

$$\begin{aligned} & \frac{1}{t_d^A} \int_0^{\vartheta t_d^A} \mathbf{1}(|\Delta Z'_k(s)| \geq \epsilon) ds \\ (6.3) \quad & \leq \frac{1}{t_d^A} \sum_{j=0}^{\mathcal{N}_{A, \vartheta, d}-1} \int_{\sigma_{2j+1}}^{\sigma_{2j+2}} \mathbf{1}(|\Delta Z'_k(s)| \geq \epsilon) ds \\ & \leq 48k(k+1)^2 \left(\frac{\log(\vartheta t_d^A)}{t_d^A}\right) \mathcal{N}_{A, \vartheta, d} \leq \frac{192k(k+1)^2}{Ad} \mathcal{N}_{A, \vartheta, d}. \end{aligned}$$

Fix $A = A_0(k) := 385\epsilon^{-1}2^k k(k+1)^2$ (suppressing dependence on ϵ for notational convenience). Note that this does not depend on ϑ . Writing $t_{d,k} := t_d^{A_0(k)}$ and $\mathcal{E}_{d,k} := \mathcal{E}_{d,A_0(k)}$, and using (6.1) and (6.3), we obtain for $d \geq A_0(k) \vee \vartheta$,

$$\begin{aligned} (6.4) \quad & \mathbb{P}\left(\frac{1}{t_{d,k}} \int_0^{\vartheta t_{d,k}} \mathbf{1}(|\Delta Z'_k(s)| \geq \epsilon) ds \geq 1, \mathcal{E}_{d,k}\right) \\ & \leq \mathbb{P}\left(\frac{192k(k+1)^2}{A_0(k)d} \mathcal{N}_{A_0(k), \vartheta, d} \geq 1\right) \\ & \leq \mathbb{P}(\mathcal{N}_{A_0(k), \vartheta, d} > 2^k \epsilon^{-1} d + 1) \leq e^{-d/8}. \end{aligned}$$

Thus, from (6.2) and (6.4), for any $\vartheta > 0$, and d sufficiently large,

$$\begin{aligned} (6.5) \quad & \mathbb{P}\left(\frac{1}{t_{d,k}} \int_0^{\vartheta t_{d,k}} \mathbf{1}(|\Delta Z'_k(s)| \geq \epsilon) ds \geq 1\right) \\ & \leq 2e^{-d/8} + \frac{10k2^k \epsilon^{-1}}{A_0(k)^2 \vartheta^2 d (\log d)^2} \rightarrow 0 \quad \text{as } d \rightarrow \infty. \end{aligned}$$

Note that the above holds for any $1 \leq j \leq k$, upon replacing k with j . Thus, for any $1 \leq j \leq k$, writing $\vartheta_k := \vartheta t_{d,k}/t_{d,1} = \vartheta A_0(k)/A_0(1) = \vartheta 2^{k-3} k(k+1)^2$,

$$\mathbb{P}\left(\frac{1}{t_{d,k}} \int_0^{\vartheta t_{d,k}} \mathbf{1}(|\Delta Z'_j(s)| \geq \epsilon) ds \geq 1\right) \leq \mathbb{P}\left(\frac{1}{t_{d,j}} \int_0^{\vartheta_k t_{d,j}} \mathbf{1}(|\Delta Z'_j(s)| \geq \epsilon) ds \geq 1\right),$$

which converges to zero as $d \rightarrow \infty$. This proves the lemma with $A_0 := A_0(k)$. $\square$

LEMMA 6.3. *Suppose that $\mathbf{U}$ and $\mathbf{V}$ are as in Theorem 2.1. Define the infinite ordered Atlas model $\mathbf{X}^{\mathbf{U} \wedge \mathbf{V}}$ with $\mathbf{X}^{\mathbf{U} \wedge \mathbf{V}}(0) = (0, s(\mathbf{U} \wedge \mathbf{V})^T)^T$ and the same sequence of Brownian motions used to define $\mathbf{X}^{\mathbf{U}}$, $\mathbf{X}^{\mathbf{V}}$ and $\mathbf{X}^{\mathbf{U}^{(d)}}$ in Lemma 6.2. Let $\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}$ be the corresponding gap process. Define the measure*

$$(6.6) \quad \underline{\mu}_t(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty), t > 0.$$

Then $\underline{\mu}_t$ converges weakly to π as $t \rightarrow \infty$.

PROOF. Fix any $\epsilon \in (0, 1)$, $\delta \in (0, 1)$, $k \in \mathbb{N}$. Recall A_0 from Lemma 6.2 for this choice of ϵ, k . For $t > \delta^{-1} A_0 e$, write $d(t) := \lfloor \delta A_0^{-1} t / \log(\delta A_0^{-1} t) \rfloor$. Choose $t_0 > \delta^{-1} A_0 e$ large enough such that $d(t) \geq \delta A_0^{-1} t / [2 \log(\delta A_0^{-1} t)] > k$ and $\log(\delta A_0^{-1} t) \geq 2 \log(2 \log(\delta A_0^{-1} t))$ for all $t \geq t_0$. Also, set $\vartheta := 4\delta^{-1}$. Note that for all $t \geq t_0$,

$$A_0 d(t) \log d(t) \leq \delta t, \quad A_0 \vartheta d(t) \log d(t) \geq t.$$

Recall the gap processes $\mathbf{Z}^{\mathbf{V}}, \mathbf{Z}^{\mathbf{U}^{(d)}}$ from Lemma 6.2. Then, for all $t \geq t_0$, $1 \leq j \leq k$,

$$\begin{aligned} & \mathbb{P}\left(\frac{1}{t} \int_0^t \mathbf{1}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}^{(d(t))}}(s)| > \epsilon) ds \geq \delta\right) \\ & \leq \mathbb{P}\left(\frac{1}{A_0 d(t) \log d(t)} \int_0^{A_0 \vartheta d(t) \log d(t)} \mathbf{1}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}^{(d(t))}}(s)| > \epsilon) ds \geq 1\right). \end{aligned}$$

Hence, applying Lemma 6.2 with $d(t)$ in place of d and the above choice of ϑ , we obtain for any $1 \leq j \leq k$,

$$(6.7) \quad \mathbb{P}\left(\frac{1}{t} \int_0^t \mathbf{1}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}^{(d(t))}}(s)| > \epsilon) ds \geq \delta\right) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Define the measures $\{\tilde{\mu}_t^{(k)}, t > 0\}$ on $\mathbb{R}_+^k$ by

$$\tilde{\mu}_t^{(k)}(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U}_{(d(t))}}(s)|_k \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^k), t > 0.$$

Take any closed set $F \in \mathcal{B}(\mathbb{R}_+^k)$ and, for $\eta > 0$, let $F^\eta := \{y \in \mathbb{R}_+^k : \|y - x\|_1 \leq \eta \text{ for some } x \in F\}$ be the closed η -neighborhood of F with respect to L^1 distance. Observe that

$$\tilde{\mu}_t^{(k)}(F) \leq \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{V}}(s)|_k \in F^{k\epsilon}) ds + \sum_{j=1}^k \frac{1}{t} \int_0^t \mathbb{P}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}_{(d(t))}}(s)| > \epsilon) ds.$$

Using the inequality

$$\frac{1}{t} \int_0^t \mathbb{P}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}_{(d(t))}}(s)| > \epsilon) ds \leq \delta + \mathbb{P}\left(\frac{1}{t} \int_0^t \mathbf{1}(|Z_j^{\mathbf{V}}(s) - Z_j^{\mathbf{U}_{(d(t))}}(s)| > \epsilon) ds \geq \delta\right)$$

for $1 \leq j \leq k$, together with (6.7), and the stationarity of $\mathbf{Z}^{\mathbf{V}}(\cdot)$, we now obtain for any closed set $F \in \mathcal{B}(\mathbb{R}_+^k)$,

$$(6.8) \quad \limsup_{t \rightarrow \infty} \tilde{\mu}_t^{(k)}(F) \leq \pi^{(k)}(F^{k\epsilon}) + k\delta,$$

where $\pi^{(k)} = \bigotimes_{i=1}^k \text{Exp}(2)$.

Our next step is to show the above bound (6.8) with $\tilde{\mu}_t^{(k)}$ replaced by the measure $\underline{\mu}_t^{(k)}$ (the k -marginal of $\underline{\mu}_t$) by showing that, for any closed set $F \in \mathcal{B}(\mathbb{R}_+^k)$,

$$(6.9) \quad \underline{\mu}_t^{(k)}(F) - \tilde{\mu}_t^{(k)}(F) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

To see this, take any $t \geq t_0$. Let $\gamma(t)$ be the probability law of $(0, s(\mathbf{U}_{(d(t))})^T)^T$ and denote by $\mathbf{Y}^{\gamma(t)}$ the unique weak solution of (1.1) with $\mathbf{Y}^{\gamma(t)}(0)$ distributed as $\gamma(t)$, given on some probability space with driving Brownian motions $\{W_i\}_{i \in \mathbb{N}_0}$. Denote the corresponding process of gaps (as defined in (1.6)) as $\hat{\mathbf{Z}}^{\gamma(t)}$.

Recall $d(t) > k$. For any $\zeta \in [0, t]$,

$$(6.10) \quad \begin{aligned} \mathbb{P}(\mathbf{Z}^{\mathbf{U}_{(d(t))}}(\zeta)|_k \in F) &= \mathbb{P}(\hat{\mathbf{Z}}^{\gamma(t)}(\zeta)|_k \in F) \\ &= \mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t)}(\zeta)|_k \in F, \inf_{s \in [0, t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) > \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\right) \\ &\quad + \mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t)}(\zeta)|_k \in F, \inf_{s \in [0, t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\right). \end{aligned}$$

Recall from the [Introduction](#) that a finite-dimensional Atlas model has a unique strong solution. We denote by $\hat{\mathbf{Y}}^{\gamma(t), d(t)}$ such a process with $d(t) + 1$ particles, driven by Brownian motions $\{W_i\}_{i=0}^{d(t)}$ and initial conditions $\hat{Y}_i^{\gamma(t), d(t)}(0) = Y_i^{\gamma(t)}(0)$, $i = 0, 1, \dots, d(t)$. Denote by $\hat{\mathbf{X}}^{\gamma(t), d(t)}$ the corresponding process of ordered particles and denote the associated gap process by $\hat{\mathbf{Z}}^{\gamma(t), d(t)}$. Then, on the set, $\{\inf_{s \in [0, t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) > \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\}$, $\hat{\mathbf{Z}}^{\gamma(t)}(s)|_k = \hat{\mathbf{Z}}^{\gamma(t), d(t)}(s)|_k$ for all $s \in [0, t]$ and so, using (6.10), for $\zeta \in [0, t]$,

$$\begin{aligned} \mathbb{P}(\mathbf{Z}^{\mathbf{U}_{(d(t))}}(\zeta)|_k \in F) &= \mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t), d(t)}(\zeta)|_k \in F, \inf_{s \in [0, t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) > \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\right) \\ &\quad + \mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t)}(\zeta)|_k \in F, \inf_{s \in [0, t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\right) \\ &= \mathbb{P}(\hat{\mathbf{Z}}^{\gamma(t), d(t)}(\zeta)|_k \in F) \end{aligned}$$

$$\begin{aligned}
& -\mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t),d(t)}(\zeta)|_k \in F, \inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma(t)}(s)\right) \\
& + \mathbb{P}\left(\hat{\mathbf{Z}}^{\gamma(t)}(\zeta)|_k \in F, \inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma(t)}(s)\right),
\end{aligned}$$

which shows that

$$\begin{aligned}
(6.11) \quad & |\mathbb{P}(\mathbf{Z}^{\mathbf{U}(d(t))}(\zeta)|_k \in F) - \mathbb{P}(\hat{\mathbf{Z}}^{\gamma(t),d(t)}(\zeta)|_k \in F)| \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma(t)}(s)\right).
\end{aligned}$$

Now let γ^* be the probability law of $(0, s(\mathbf{U} \wedge \mathbf{V})^T)^T$ and denote by $\mathbf{Y}^{\gamma^*}$ the unique weak solution of (1.1) with $\mathbf{Y}^{\gamma^*}(0)$ distributed as γ^* . Denote the corresponding process of gaps as $\hat{\mathbf{Z}}^{\gamma^*}$. The process of ordered particles $\hat{\mathbf{X}}^{\gamma^*,d(t)}$ with $d(t) + 1$ particles and initial values $\mathbf{Y}^{\gamma^*}(0)|_{d(t)}$, and the associated gap process $\hat{\mathbf{Z}}^{\gamma^*,d(t)}$, are defined in a similar manner as $\hat{\mathbf{X}}^{\gamma(t),d(t)}$ and $\hat{\mathbf{Z}}^{\gamma(t),d(t)}$. By a similar argument as above

$$\begin{aligned}
(6.12) \quad & |\mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(\zeta)|_k \in F) - \mathbb{P}(\hat{\mathbf{Z}}^{\gamma^*,d(t)}(\zeta)|_k \in F)| \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma^*}(s)\right).
\end{aligned}$$

Since $\mathbf{U} \wedge \mathbf{V}|_{d(t)} = \mathbf{U}_{(d(t))}|_{d(t)}$ we have that $\hat{\mathbf{Z}}^{\gamma^*,d(t)}$ and $\hat{\mathbf{Z}}^{\gamma(t),d(t)}$ have the same distribution and so from (6.11) and (6.12) we have that

$$\begin{aligned}
(6.13) \quad & |\mathbb{P}(\mathbf{Z}^{\mathbf{U}(d(t))}(\zeta)|_k \in F) - \mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(\zeta)|_k \in F)| \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma(t)}(s)\right) \\
& + \mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma^*}(s)\right).
\end{aligned}$$

Now, recalling $t_{d(t)}^{A_0\vartheta} = A_0\vartheta d(t) \log d(t) \geq t$ and that $\Gamma_{d(t)}^{A_0\vartheta} = 6A_0\vartheta \sqrt{d(t)}(\log d(t))$ from Lemma 4.2, we see that

$$\begin{aligned}
& \mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma(t)}(s)\right) \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t_{d(t)}^{A_0\vartheta}]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0,t_{d(t)}^{A_0\vartheta}]} Y_{(k)}^{\gamma(t)}(s)\right) \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t_{d(t)}^{A_0\vartheta}]} \inf_{i \geq d(t)} Y_i^{\gamma(t)}(s) \leq \Gamma_{d(t)}^{A_0\vartheta}\right) + \mathbb{P}\left(\sup_{s \in [0,t_{d(t)}^{A_0\vartheta}]} Y_{(k)}^{\gamma(t)}(s) \geq \Gamma_{d(t)}^{A_0\vartheta}\right) \\
& \leq \mathbb{P}\left(\inf_{s \in [0,t_{d(t)}^{A_0\vartheta}]} \inf_{i \geq d(t)} Y_i^{\gamma^*}(s) \leq \Gamma_{d(t)}^{A_0\vartheta}\right) + \mathbb{P}\left(\sup_{s \in [0,t_{d(t)}^{A_0\vartheta}]} Y_{(k)}^{\pi}(s) \geq \Gamma_{d(t)}^{A_0\vartheta}\right) \rightarrow 0 \quad \text{as } t \rightarrow \infty,
\end{aligned}$$

where $\mathbf{Y}^{\pi}(\cdot)$ is the infinite Atlas model with $\mathbf{Y}^{\pi}(0)$ distributed as $(0, s(\mathbf{V})^T)^T$. The last inequality above uses the observation $\mathbf{U} \wedge \mathbf{V} \leq \mathbf{U}_{(d(t))} \leq \mathbf{V}$, $\inf_{i \geq d(t)} Y_i^{\gamma^*}(s) = Y_{(d(t))}^{\gamma^*}(s)$ for all $s \geq 0$, and [27], Corollary 3.10(i). The limit follows from Lemma 4.2 with $A = A_0\vartheta$, $d = d(t)$ upon using $\mathbf{U} \wedge \mathbf{V} \leq \mathbf{V}$ and noting that $\mathbf{Y}^{\gamma^*}(0)$ satisfies condition (4.3) by virtue of assumption (2.1) and $\mathbf{Y}^{\pi}(0)$ satisfies (4.3) by the law of large numbers.

Similarly,

$$\mathbb{P}\left(\inf_{s \in [0,t]} \inf_{i \geq d(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0,t]} Y_{(k)}^{\gamma^*}(s)\right) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Using the above observations along with (6.13), we obtain

$$\sup_{\zeta \in [0, t]} |\mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(\zeta)|_k \in F) - \mathbb{P}(\mathbf{Z}^{\mathbf{U}(d(t))}(\zeta)|_k \in F)| \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

(6.9) follows from this. From (6.8) and (6.9), we obtain for any closed set $F \in \mathcal{B}(\mathbb{R}_+^k)$,

$$\limsup_{t \rightarrow \infty} \underline{\mu}_t^{(k)}(F) \leq \pi^{(k)}(F^{k\epsilon}) + k\delta.$$

As the left-hand side above does not depend on ϵ, δ , we take a limit as ϵ, δ go to zero in the above to obtain for any $k \in \mathbb{N}$,

$$\limsup_{t \rightarrow \infty} \underline{\mu}_t^{(k)}(F) \leq \pi^{(k)}(F) \quad \text{for any closed set } F \in \mathcal{B}(\mathbb{R}_+^k).$$

The lemma now follows from the Portmanteau theorem [5], Chapter 1, Theorem 2.1. $\square$

Now, we are ready to prove Theorem 2.1.

PROOF OF THEOREM 2.1. Let $(\mathbf{U}, \mathbf{V})$ be as in the statement of Theorem 2.1. Let $\mathbf{Z}^{\mathbf{U}}, \mathbf{Z}^{\mathbf{V}}$ and $\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}$ be as in Lemma 6.2 and Lemma 6.3. By the monotonicity property in Proposition 3.1(iii),

$$\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}}(\cdot) \leq \mathbf{Z}^{\mathbf{U}}(\cdot) \leq \mathbf{Z}^{\mathbf{U} \vee \mathbf{V}}(\cdot).$$

Define the measure

$$\bar{\mu}_t(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U} \vee \mathbf{V}}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty), t > 0,$$

and recall $\underline{\mu}_t$ defined in (6.6). Also define

$$\mu_t(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U}}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty), t > 0.$$

Then, for any $k \in \mathbb{N}, \mathbf{z} \in \mathbb{R}_+^k$,

$$(6.14) \quad \bar{\mu}_t^{(k)}((-\infty, \mathbf{z}]) \leq \mu_t^{(k)}((-\infty, \mathbf{z}]) \leq \underline{\mu}_t^{(k)}((-\infty, \mathbf{z}]).$$

By Lemma 6.3, $\underline{\mu}_t^{(k)}((-\infty, \mathbf{z}]) \rightarrow \pi^{(k)}((-\infty, \mathbf{z}])$ as $t \rightarrow \infty$. Moreover, as $\mathbf{U} \vee \mathbf{V} \geq \mathbf{V}$, by Lemma 6.1, the law of $\mathbf{Z}^{\mathbf{U} \vee \mathbf{V}}(t)$ converges weakly to π . In particular, $\bar{\mu}_t^{(k)}((-\infty, \mathbf{z}]) \rightarrow \pi^{(k)}((-\infty, \mathbf{z}])$ as $t \rightarrow \infty$. Hence, by (6.14), for any $k \in \mathbb{N}, \mathbf{z} \in \mathbb{R}_+^k$,

$$\mu_t^{(k)}((-\infty, \mathbf{z}]) \rightarrow \pi^{(k)}((-\infty, \mathbf{z}]) \quad \text{as } t \rightarrow \infty.$$

This proves the theorem. $\square$

PROOF OF COROLLARY 2.2. The fact that (i) implies (2.1) is immediate on observing that we can find a coupling $(\mathbf{U}, \mathbf{V})$ of μ and π such that $\mathbf{U} \leq \mathbf{V}$ a.s.

For the statement in (ii), we will consider the independent coupling of $(\mathbf{U}, \mathbf{V})$, that is, under the coupling, the marginals $\mathbf{U}$ and $\mathbf{V}$ are independent. Recall from Remark 2.3(b) that the statement in (2.3) is equivalent to

$$(6.15) \quad \liminf_{d \rightarrow \infty} \frac{1}{\sqrt{d}(\log d)} \sum_{i=1}^d \left(U_i \wedge \frac{1}{2} \right) = \infty, \quad \text{a.s.}$$

It suffices to show that, conditional on a realization of $\mathbf{U} = \mathbf{u}$ such that (6.15) holds with $\mathbf{U} = \mathbf{u}$, the independent random variables $\Psi_i := V_i \wedge u_i$ satisfy

$$(6.16) \quad \liminf_{m \rightarrow \infty} \frac{1}{\sqrt{m} \log m} \sum_{i=1}^m \Psi_i = \infty \quad \text{almost surely.}$$

Note that

$$\mu_i := \mathbb{E}(\Psi_i) = \int_0^{u_i} 2ve^{-2v} dv + \int_{u_i}^{\infty} 2u_i e^{-2v} dv = \frac{1}{2}(1 - e^{-2u_i}).$$

As $x \wedge (1/2) \leq 1 - e^{-2x} \leq 2(x \wedge (1/2))$ for all $x \geq 0$, (6.15) with $\mathbf{U} = \mathbf{u}$ is equivalent to the condition

$$(6.17) \quad \lim_{m \rightarrow \infty} \frac{1}{\sqrt{m} \log m} \sum_{i=1}^m \mu_i = \infty.$$

Take any $C > 0$. By (6.17), there exists $m_C \in \mathbb{N}$ such that $\sum_{i=1}^m \mu_i \geq 2C\sqrt{m} \log m$ for all $m \geq m_C$. Note that for any $k \in \mathbb{N}$,

$$\mathbb{E}(|\Psi_i - \mu_i|^k) \leq 2^k \mathbb{E}(V_i^k + \mu_i^k) \leq 2^k(k!2^{-k} + 2^{-k}) \leq 2k!.$$

Therefore, Ψ_i satisfies condition (2.16) in [31], Chapter 2, with $\sigma = 2$, $b = 1$. Hence, by [31], Chapter 2, Proposition 2.3, $\mathbb{E}(e^{\lambda(\Psi_i - \mu_i)}) \leq \exp\{2\lambda^2/(1 - |\lambda|)\} \leq \exp\{4\lambda^2\}$ for all $|\lambda| < 1/2$. Hence, using [31], Chapter 2, Theorem 2.3, with $D_k = \Psi_k$, $b_k = 2$ and $v_k = 2\sqrt{2}$, for all $m \geq m_C$ satisfying $C\sqrt{m} \log m \leq 4m$,

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^m \Psi_i \leq C\sqrt{m} \log m\right) &\leq \mathbb{P}\left(\sum_{i=1}^m (\Psi_i - \mu_i) \leq -C\sqrt{m} \log m\right) \\ &\leq 2 \exp\left\{-\frac{(C\sqrt{m} \log m)^2}{16m}\right\} = 2 \exp\{-C^2(\log m)^2/16\}. \end{aligned}$$

As the bound above is summable in m , we conclude from the Borel–Cantelli lemma that for any $C > 0$, almost surely, $\liminf_{m \rightarrow \infty} (\sqrt{m} \log m)^{-1} \sum_{i=1}^m \Psi_i \geq C$. This implies (6.16) and thus proves that the statement in (ii) implies (2.1).

Consider now (iii). Thus $U_i = \lambda_i \Theta_i$, $i \in \mathbb{N}$, where $\{\Theta_i\}$ are i.i.d. nonnegative random variables satisfying $\mathbb{P}(\Theta_1 > 0) > 0$, and $\{\lambda_i\}$ are positive deterministic real numbers. If $\{\lambda_j\}$ satisfy condition (a), that is, $\liminf_{j \rightarrow \infty} \lambda_j > 0$, there exists $\eta > 0$ such that $U_i \wedge (1/2) \geq (\eta \Theta_i) \wedge (1/2)$ for all sufficiently large i . Note that $\{(\eta \Theta_i) \wedge (1/2)\}$ are i.i.d. with positive mean as $\mathbb{P}(\Theta_1 > 0) > 0$. Therefore, (6.15) holds by the strong law of large numbers, implying (2.1).

If $\{\lambda_j\}$ satisfy condition (b), namely $\limsup_{j \rightarrow \infty} \lambda_j < \infty$ and (2.4) holds, there exists $M \in [1, \infty)$ such that $\lambda_j \leq M$ for all $j \in \mathbb{N}$. Note that $\mathbb{E}(U_i \wedge (1/2)) \geq \lambda_j \mathbb{E}(\Theta_i \wedge (1/2M)) \geq \lambda_j M^{-1} \mathbb{E}(\Theta_1 \wedge (1/2))$. As $\mathbb{P}(\Theta_1 > 0) > 0$, $\mathbb{E}(\Theta_1 \wedge (1/2)) > 0$. Hence, by (2.4), for any $C > 0$, there exists $m_C \in \mathbb{N}$ such that for all $m \geq m_C$,

$$\sum_{i=1}^m \mathbb{E}(U_i \wedge (1/2)) \geq M^{-1} \mathbb{E}(\Theta_1 \wedge (1/2)) \sum_{i=1}^m \lambda_i \geq 2C\sqrt{m} \log m.$$

As $|U_i \wedge (1/2) - \mathbb{E}(U_i \wedge (1/2))| \leq 1$ for all $i \in \mathbb{N}$, by the Azuma–Hoeffding inequality [31], Chapter 2, Corollary 2.1, for all $m \geq m_C$,

$$\begin{aligned} &\mathbb{P}\left(\sum_{i=1}^m U_i \wedge (1/2) \leq C\sqrt{m} \log m\right) \\ &\leq \mathbb{P}\left(\sum_{i=1}^m (U_i \wedge (1/2) - \mathbb{E}(U_i \wedge (1/2))) \leq -C\sqrt{m} \log m\right) \leq 2 \exp\{-2C^2(\log m)^2\}, \end{aligned}$$

which is summable in m . Hence, we conclude from the Borel–Cantelli lemma that

$$\liminf_{m \rightarrow \infty} (\sqrt{m} \log m)^{-1} \sum_{i=1}^m U_i \wedge (1/2) = \infty$$

almost surely. Thus, (6.15) holds which shows (2.1). $\square$

6.2. Proof of Theorem 2.4. We now prove Theorem 2.4. As in the proof of Theorem 2.1, we will proceed by defining new starting configurations for the gap process in terms of $\mathbf{U}$ and $\mathbf{V}_a$ and exploiting the monotonicity properties of the synchronous coupling described in Proposition 3.1.

PROOF OF THEOREM 2.4. Let $(\mathbf{U}, \mathbf{V}_a)$ be as in the statement of Theorem 2.4. First note that, by Lemma 4.5, $s(\mathbf{U})$, $s(\mathbf{V}_a)$ and $s(\mathbf{U} \wedge \mathbf{V}_a)$ all satisfy (3.1). Define the vector $\mathbf{U}_{a,(d)}$ by

$$U_{a,(d),i} := (U_i \wedge V_{a,i})\mathbf{1}(i \leq d) + (U_i \vee V_{a,i})\mathbf{1}(i > d), \quad i \in \mathbb{N}, d \geq 2.$$

As $\mathbf{U}_{a,(d)} \geq \mathbf{U} \wedge \mathbf{V}_a$ and $s(\mathbf{U} \wedge \mathbf{V}_a)$ satisfies (3.1), we have that $s(\mathbf{U}_{a,(d)})$ satisfies (3.1).

Observe that, with $V_i := (1 + ia/2)V_{a,i}$, $i \in \mathbb{N}$, $(\mathbf{U}_{a,(d)}, \mathbf{U} \vee \mathbf{V}_a, \mathbf{V})$ defines a coupling of (μ, π_a, π) . By the second condition in (2.5), $\limsup_{d \rightarrow \infty} \frac{U_d}{V_d} < \infty$ a.s. and so the events

$$\mathcal{E}^{(D)} := \{\mathbf{U} \vee \mathbf{V}_a \leq D\mathbf{V}\}, \quad D \geq 1,$$

satisfy

$$(6.18) \quad \mathbb{P}(\mathcal{E}^{(D)}) \rightarrow 1 \quad \text{as } D \rightarrow \infty.$$

Fix $D \geq 1$. Note that, for $i \in \mathbb{N}$,

$$s(\mathbf{U}_{a,(d)} \wedge D\mathbf{V})_i = \sum_{j=1}^i (\mathbf{U}_{a,(d)})_j \wedge D\mathbf{V}_j \geq \sum_{j=1}^i V_{a,j} - \sum_{j=1}^i |V_{a,j} - U_j|.$$

By a similar calculation as in (4.10) it then follows that, for some $i' \in \mathbb{N}$ and all $i \geq i'$, $s(\mathbf{U}_{a,(d)} \wedge D\mathbf{V})_i \geq (4a)^{-1} \log i$. This shows that $s(\mathbf{U}_{a,(d)} \wedge D\mathbf{V})$ satisfies (3.1). Similarly, $s((\mathbf{U} \vee \mathbf{V}_a) \wedge D\mathbf{V})$ satisfies (3.1).

In the rest of the proof we will consider several random variables $\tilde{\mathbf{U}}$ such that $s(\tilde{\mathbf{U}})$ satisfies (3.1) and are given on the same probability space. For all such random variables we use Proposition 3.1(ii) to construct infinite ordered Atlas models $\mathbf{X}^{\tilde{\mathbf{U}}}(\cdot)$ starting from $(0, s(\tilde{\mathbf{U}})^T)^T$ driven by a common collection of Brownian motions $\{B_i\}_{i \in \mathbb{N}_0}$ (that are independent of all the $\tilde{\mathbf{U}}$). We denote by $\mathbf{Z}^{\tilde{\mathbf{U}}}(\cdot)$ the associated gap processes.

Let $\Delta \mathbf{Z}^{*,D}(\cdot) := \mathbf{Z}^{(\mathbf{U} \vee \mathbf{V}_a) \wedge D\mathbf{V}}(\cdot) - \mathbf{Z}^{\mathbf{U}_{a,(d)} \wedge D\mathbf{V}}(\cdot)$.

Fix $\epsilon, \delta > 0$, $k \in \mathbb{N}$. Recall the stopping times $\{\sigma_j : j \geq 0\}$ and the random variables $\{\mathcal{N}_T : T \geq 0\}$ from Section 5 with $\mathbf{U}_1 = \mathbf{U}_{a,(d)} \wedge D\mathbf{V}$ and $\mathbf{U}_2 = (\mathbf{U} \vee \mathbf{V}_a) \wedge D\mathbf{V}$. Write $t'_d := \log d$, and $\mathcal{N}'_d := \mathcal{N}'_{t'_d}$. We suppress the dependence of $\{\sigma_j\}$ and $\mathcal{N}'_d$ on D for notational convenience.

As $\sum_{j=1}^{\infty} j \Delta Z_j^{*,D}(0) \leq \sum_{j=1}^d j |V_{a,j} - U_j| < \infty$, by Lemma 5.1, for any $j \geq 0$, when $\sigma_{2j+2} < \infty$,

$$\sum_{i=1}^{\infty} \Delta Z_i^{*,D}(\sigma_{2j+2}) - \sum_{i=1}^{\infty} \Delta Z_i^{*,D}(\sigma_{2j+1}) \leq -\epsilon/2^k.$$

Hence, as in the proof of (6.1), for any $A > 0$,

$$\begin{aligned}
 & \mathbb{P}\left(\mathcal{N}'_d \geq 2^k A \frac{\log d}{\log \log d}\right) \\
 (6.19) \quad & \leq \mathbb{P}\left(\sum_{i=1}^d \Delta Z_i^{*,D}(0) \geq A\epsilon \frac{\log d}{\log \log d} - 2^{-k}\epsilon\right) \\
 & \leq \mathbb{P}\left(\frac{\log \log d}{\log d} \sum_{i=1}^d |V_{a,i} - U_i| \geq A\epsilon - 2^{-k}\epsilon \frac{\log \log d}{\log d}\right) \rightarrow 0 \quad \text{as } d \rightarrow \infty,
 \end{aligned}$$

where the claimed limit is a consequence of (2.5).

For $d \geq 2$, define the events

$$\mathcal{E}'_{d,D} := \{\sigma_1 \leq \log d, \sigma_{2j+2} - \sigma_{2j+1} \leq 48D^2k(k+1)^2 \log \log d \text{ for all } 0 \leq j \leq \mathcal{N}'_d - 1\}.$$

By Lemma 5.2 with $T = \log d$, $n = \lfloor 2^k \log d / \log \log d \rfloor$, for sufficiently large d ,

$$\mathbb{P}((\mathcal{E}'_{d,D})^c \cap \{\sigma_1 \leq \log d\}) \leq \mathbb{P}\left(\mathcal{N}'_d > 2^k \frac{\log d}{\log \log d}\right) + 5k \left(2^k \frac{\log d}{\log \log d}\right) (\log d)^{-2},$$

which converges to zero as $d \rightarrow \infty$ using (6.19) with $A = 1$. On the event $\mathcal{E}'_{d,D}$,

$$\begin{aligned}
 \frac{1}{t'_d} \int_0^{t'_d} \mathbf{1}(|\Delta Z_k^{*,D}(s)| \geq \epsilon) ds & \leq \frac{1}{t'_d} \sum_{j=0}^{\mathcal{N}'_d-1} \int_{\sigma_{2j+1}}^{\sigma_{2j+2}} \mathbf{1}(|\Delta Z_k^{*,D}(s)| \geq \epsilon) ds \\
 & \leq 48D^2k(k+1)^2 \left(\frac{\log \log d}{\log d}\right) \mathcal{N}'_d.
 \end{aligned}$$

Hence, using (6.19) with $A = A_{D,k,\delta} := \delta[48D^2 \times 2^k k(k+1)^2]^{-1}$,

$$\begin{aligned}
 & \mathbb{P}\left(\frac{1}{t'_d} \int_0^{t'_d} \mathbf{1}(|\Delta Z_k^{*,D}(s)| \geq \epsilon) ds > \delta\right) \\
 (6.20) \quad & \leq \mathbb{P}\left(\mathcal{N}'_d > 2^k A_{D,k,\delta} \frac{\log d}{\log \log d}\right) + \mathbb{P}((\mathcal{E}'_{d,D})^c \cap \{\sigma_1 \leq \log d\}) \rightarrow 0 \\
 & \quad \text{as } d \rightarrow \infty.
 \end{aligned}$$

Recalling the event $\mathcal{E}^{(D)}$ and using (6.20),

$$\begin{aligned}
 \limsup_{d \rightarrow \infty} \mathbb{P}\left(\frac{1}{t'_d} \int_0^{t'_d} \mathbf{1}(|Z_k^{\mathbf{U} \vee \mathbf{V}_a}(s) - Z_k^{\mathbf{U}_{a,(d)}}(s)| \geq \epsilon) ds > \delta\right) \\
 \leq \limsup_{d \rightarrow \infty} \mathbb{P}\left(\frac{1}{t'_d} \int_0^{t'_d} \mathbf{1}(|\Delta Z_k^{*,D}(s)| \geq \epsilon) ds > \delta\right) + \mathbb{P}((\mathcal{E}^{(D)})^c) = \mathbb{P}((\mathcal{E}^{(D)})^c).
 \end{aligned}$$

As $D \geq 1$ is arbitrary, we obtain from (6.18), for any $\epsilon, \delta > 0$, $k \in \mathbb{N}$,

$$(6.21) \quad \limsup_{d \rightarrow \infty} \mathbb{P}\left(\frac{1}{t'_d} \int_0^{t'_d} \mathbf{1}(|Z_k^{\mathbf{U} \vee \mathbf{V}_a}(s) - Z_k^{\mathbf{U}_{a,(d)}}(s)| \geq \epsilon) ds \geq \delta\right) = 0.$$

Now, from the monotonicity property in Proposition 3.1(iii),

$$(6.22) \quad \mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{U}}(\cdot) \leq \mathbf{Z}^{\mathbf{U} \vee \mathbf{V}_a}(\cdot), \quad \mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{U} \vee \mathbf{V}_a}(\cdot).$$

Let $d'(t) := \lceil e^{t/\delta_0} \rceil$, $t > 0$, where δ_0 is the constant appearing in Lemma 4.6. Define the following measures:

$$\bar{\mu}_{a,t}(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U} \vee \mathbf{V}_a}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty),$$

$$\underline{\mu}_{a,t}(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty),$$

for $t > 0$. For $k \in \mathbb{N}$, $t > 0$, define the measure on $\mathbb{R}_+^k$,

$$\tilde{\mu}_{a,t}^{(k)}(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U}_{a,(d'(t))}}(s)|_k \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^k).$$

Take any $k \in \mathbb{N}$ and $\mathbf{z} \in \mathbb{R}_+^k$. For $\eta \in \mathbb{R}$, write $\mathbf{z}^\eta$ for the vector obtained by adding η to each coordinate. Note that, for any $\epsilon > 0$,

$$(6.23) \quad \begin{aligned} \tilde{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) &\leq \bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}^\epsilon]) \\ &+ \sum_{j=1}^k \frac{1}{t} \int_0^t \mathbb{P}(|Z_j^{\mathbf{U} \vee \mathbf{V}_a}(s) - Z_j^{\mathbf{U}_{a,(d'(t))}}(s)| > \epsilon) ds. \end{aligned}$$

Hence, using (6.21),

$$\limsup_{t \rightarrow \infty} \tilde{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) \leq \limsup_{t \rightarrow \infty} \bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}^\epsilon]) \leq \pi_a^{(k)}((-\infty, \mathbf{z}^\epsilon]),$$

where the last inequality uses $\mathbf{Z}^{\mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{U} \vee \mathbf{V}_a}(\cdot)$ (cf. Proposition 3.1(iii)) and the stationarity of the process $\mathbf{Z}^{\mathbf{V}_a}(\cdot)$. Taking $\epsilon \downarrow 0$ above, we obtain for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$(6.24) \quad \limsup_{t \rightarrow \infty} \tilde{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) \leq \pi_a^{(k)}((-\infty, \mathbf{z}]).$$

As in the proof of Theorem 2.1, we will show that for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$(6.25) \quad \mu_{a,t}^{(k)}((-\infty, \mathbf{z}]) - \tilde{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Fix $t_0 > 0$ such that $d'(t_0) > k$. For $t \geq t_0$, let $\gamma(t)$ be the probability law of $(0, s(\mathbf{U}_{a,(d'(t))})^T)^T$ and denote by $\mathbf{Y}^{\gamma(t)}$ the unique weak solution of (1.1) with $\mathbf{Y}^{\gamma(t)}(0)$ distributed as $\gamma(t)$, given on some probability space with driving Brownian motions $\{W_i\}_{i \in \mathbb{N}_0}$. Denote the corresponding process of gaps (as defined in (1.6)) as $\hat{\mathbf{Z}}^{\gamma(t)}$. As in the proof of Lemma 6.3, we use the fact that a finite-dimensional Atlas model has a unique strong solution. We denote by $\hat{\mathbf{Y}}^{\gamma(t), d'(t)}$ such a process with $d'(t) + 1$ particles, driven by Brownian motions $\{W_i\}_{i=0}^{d'(t)}$ and initial conditions $\hat{Y}_i^{\gamma(t), d'(t)}(0) = Y_i^{\gamma(t)}(0)$, $i = 0, 1, \dots, d'(t)$. Denote by $\hat{\mathbf{X}}^{\gamma(t), d'(t)}$ the corresponding process of ordered particles and denote the associated gap process by $\hat{\mathbf{Z}}^{\gamma(t), d'(t)}$. Exactly as in the proof of (6.11) we now see that, for any $\zeta \in [0, t]$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$(6.26) \quad \begin{aligned} &|\mathbb{P}(\mathbf{Z}^{\mathbf{U}_{a,(d'(t))}}(\zeta)|_k \in (-\infty, \mathbf{z}]) - \mathbb{P}(\hat{\mathbf{Z}}^{\gamma(t), d'(t)}(\zeta)|_k \in (-\infty, \mathbf{z}])| \\ &\leq \mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0, t]} Y^{\gamma(t)}(s)\right). \end{aligned}$$

Now let γ^* be the probability law of $(0, s(\mathbf{U} \wedge \mathbf{V}_a)^T)^T$ and denote by $\mathbf{Y}^{\gamma^*}$ the unique weak solution of (1.1) with $\mathbf{Y}^{\gamma^*}(0)$ distributed as γ^* . Denote the corresponding process of gaps as $\hat{\mathbf{Z}}^{\gamma^*}$. The process of ordered particles $\hat{\mathbf{X}}^{\gamma^*, d'(t)}$ with $d'(t) + 1$ particles and initial values

$\mathbf{Y}^{\gamma^*}(0)|_{d'(t)}$, and the associated gap process $\hat{\mathbf{Z}}^{\gamma^*, d'(t)}$, are defined in a similar manner as $\hat{\mathbf{X}}^{\gamma(t), d'(t)}$ and $\hat{\mathbf{Z}}^{\gamma(t), d'(t)}$. By a similar argument as above,

$$(6.27) \quad \begin{aligned} & |\mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\zeta)|_k \in (-\infty, \mathbf{z}]) - \mathbb{P}(\hat{\mathbf{Z}}^{\gamma^*, d'(t)}(\zeta)|_k \in (-\infty, \mathbf{z}])| \\ & \leq \mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma^*}(s)\right). \end{aligned}$$

Since $\mathbf{U} \wedge \mathbf{V}_a|_{d'(t)} = \mathbf{U}_{a, (d'(t))}|_{d'(t)}$ we have that $\hat{\mathbf{Z}}^{\gamma^*, d'(t)}$ and $\hat{\mathbf{Z}}^{\gamma(t), d'(t)}$ have the same distribution and so from (6.26) and (6.27) we have that

$$(6.28) \quad \begin{aligned} & |\mathbb{P}(\mathbf{Z}^{\mathbf{U}_{a, (d'(t))}}(\zeta)|_k \in (-\infty, \mathbf{z}]) - \mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\zeta)|_k \in (-\infty, \mathbf{z}])| \\ & \leq \mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0, t]} Y^{\gamma(t)}(s)\right) \\ & \quad + \mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma^*}(s)\right). \end{aligned}$$

As in the proof of Theorem 2.1,

$$\begin{aligned} & \mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma(t)}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma(t)}(s)\right) \\ & \leq \mathbb{P}\left(\inf_{s \in [0, \delta_0 \log d'(t)]} \inf_{i \geq d'(t)} Y_i^{\gamma^*}(s) \leq \frac{1}{8a} \log d'(t)\right) \\ & \quad + \mathbb{P}\left(\sup_{s \in [0, \delta_0 \log d'(t)]} Y_{(k)}^{\tilde{\gamma}}(s) \geq \frac{1}{8a} \log d'(t)\right), \end{aligned}$$

where $\mathbf{Y}^{\tilde{\gamma}}(\cdot)$ is the infinite Atlas model with $\mathbf{Y}^{\tilde{\gamma}}(0)$ distributed as $(0, s(\mathbf{U} \vee \mathbf{V}_a)^T)^T$. The last inequality above uses the observation $\mathbf{U} \wedge \mathbf{V}_a \leq \mathbf{U}_{(d'(t))} \leq \mathbf{U} \vee \mathbf{V}_a$ and [27], Corollary 3.10(i). By Lemma 4.6, the right-hand side above converges to zero as $d \rightarrow \infty$. Similarly,

$$\mathbb{P}\left(\inf_{s \in [0, t]} \inf_{i \geq d'(t)} Y_i^{\gamma^*}(s) \leq \sup_{s \in [0, t]} Y_{(k)}^{\gamma^*}(s)\right) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Hence, from (6.28), for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$\sup_{\zeta \in [0, t]} |\mathbb{P}(\mathbf{Z}^{\mathbf{U}_{a, (d'(t))}}(\zeta)|_k \in (-\infty, \mathbf{z}]) - \mathbb{P}(\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\zeta)|_k \in (-\infty, \mathbf{z}])| \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

which implies (6.25). By (6.24) and (6.25), for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$\limsup_{t \rightarrow \infty} \underline{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}]) \leq \pi_a^{(k)}((-\infty, \mathbf{z}]).$$

Moreover, as $\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{V}_a}(\cdot)$, $\liminf_{t \rightarrow \infty} \underline{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}]) \geq \pi_a^{(k)}((-\infty, \mathbf{z}])$. Thus, we conclude that for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$(6.29) \quad \lim_{t \rightarrow \infty} \underline{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}]) = \pi_a^{(k)}((-\infty, \mathbf{z}]).$$

Now note that by (6.23) and (6.21), for any $\epsilon > 0$, $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$

$$\liminf_{t \rightarrow \infty} \tilde{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}^{-\epsilon}]) \leq \liminf_{t \rightarrow \infty} \bar{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}]).$$

Also, by (6.25), $\liminf_{t \rightarrow \infty} \tilde{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}^{-\epsilon}]) = \liminf_{t \rightarrow \infty} \underline{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}^{-\epsilon}])$. Hence, using (6.29),

$$\pi_a^{(k)}((-\infty, \mathbf{z}^{-\epsilon}]) = \liminf_{t \rightarrow \infty} \underline{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}^{-\epsilon}]) \leq \liminf_{t \rightarrow \infty} \bar{\mu}_{a, t}^{(k)}((-\infty, \mathbf{z}]).$$

As $\epsilon > 0$ is arbitrary, we conclude $\pi_a^{(k)}((-\infty, \mathbf{z}]) \leq \liminf_{t \rightarrow \infty} \bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}])$. Moreover, as $\mathbf{U} \vee \mathbf{V}_a \geq \mathbf{V}_a$, we have $\limsup_{t \rightarrow \infty} \bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) \leq \pi_a^{(k)}((-\infty, \mathbf{z}])$. Hence, for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$(6.30) \quad \lim_{t \rightarrow \infty} \bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) = \pi_a^{(k)}((-\infty, \mathbf{z}]).$$

Finally, as $\mathbf{Z}^{\mathbf{U} \wedge \mathbf{V}_a}(\cdot) \leq \mathbf{Z}^{\mathbf{U}}(\cdot) \leq \mathbf{Z}^{\mathbf{U} \vee \mathbf{V}_a}(\cdot)$, for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$, $t > 0$,

$$\bar{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]) \leq \mu_t^{(k)}((-\infty, \mathbf{z}]) \leq \underline{\mu}_{a,t}^{(k)}((-\infty, \mathbf{z}]),$$

where

$$\mu_t(F) := \frac{1}{t} \int_0^t \mathbb{P}(\mathbf{Z}^{\mathbf{U}}(s) \in F) ds, \quad F \in \mathcal{B}(\mathbb{R}_+^\infty), t > 0.$$

Thus, using (6.29) and (6.30), we obtain for any $k \in \mathbb{N}$, $\mathbf{z} \in \mathbb{R}_+^k$,

$$\lim_{t \rightarrow \infty} \mu_t^{(k)}((-\infty, \mathbf{z}]) = \pi_a^{(k)}((-\infty, \mathbf{z}]),$$

which proves the theorem. $\square$

PROOF OF COROLLARY 2.5. Suppose $\mu := \bigotimes_{i=1}^\infty \text{Exp}(2 + ia + \lambda_i)$. Let $\mathbf{V} \sim \pi$. Consider the coupling $(\mathbf{U}, \mathbf{V}_a)$ of μ and π_a defined as

$$U_i := \frac{2V_i}{2 + ia + \lambda_i}, \quad V_{a,i} := \frac{2V_i}{2 + ia}, \quad i \in \mathbb{N}.$$

For sufficiently large $i \in \mathbb{N}$,

$$(6.31) \quad |V_{a,i} - U_i| = \frac{2|\lambda_i|}{(2 + ia)(2 + ia + \lambda_i)} V_i \leq \frac{2|\lambda_i|}{(1 - \beta)(2 + ia)^2} V_i \leq \frac{2}{a^2(1 - \beta)} \frac{|\lambda_i|}{i^2} V_i.$$

By (2.6),

$$(6.32) \quad \limsup_{d \rightarrow \infty} \frac{\log \log d}{\log d} \sum_{i=1}^d \mathbb{E} \left(\frac{|\lambda_i| V_i}{i^2} \right) = 0.$$

Moreover, $M_i^* := \sum_{j=1}^i \left(\frac{|\lambda_j| V_j}{j^2} - \mathbb{E} \left(\frac{|\lambda_j| V_j}{j^2} \right) \right)$, $i \in \mathbb{N}$, is a martingale with uniformly bounded second moment. Indeed, for $i \in \mathbb{N}$,

$$\mathbb{E}[(M_i^*)^2] = \sum_{j=1}^i \frac{|\lambda_j|^2}{4j^4} \leq \sum_{j=1}^\infty \frac{(C^2 + 1)(2 + ja)^2}{4j^4} < \infty.$$

Hence, by the martingale convergence theorem [22], Theorem 11.10, M_i^* almost surely converges to a finite limit as $i \rightarrow \infty$. This, along with (6.32), implies that, almost surely,

$$\limsup_{d \rightarrow \infty} \frac{\log \log d}{\log d} \sum_{i=1}^d \frac{|\lambda_i| V_i}{i^2} = 0.$$

By (6.31), we conclude that the first statement in (2.5) holds. Also note that

$$\frac{U_d}{dV_{a,d}} = \frac{2 + da}{d(2 + da + \lambda_d)} \leq \frac{1}{(1 - \beta)d},$$

which shows that the second statement in (2.5) holds as well. $\square$

Acknowledgments. We thank two anonymous referees whose valuable inputs significantly improved the article.

Funding. The research of SB was supported in part by the NSF CAREER award DMS-2141621. The research of AB was supported in part by the NSF (DMS-1814894 and DMS-1853968).

REFERENCES

- [1] ALDOUS, D. (2002). “Up the River” game story. Available at <http://www.stat.berkeley.edu/~aldous/Research/OP/river.pdf>.
- [2] BANERJEE, S. and BROWN, B. (2020). Dimension-free local convergence and perturbations for reflected Brownian motions. *Ann. Appl. Probab.* To appear. Available at [arXiv:2009.12937](https://arxiv.org/abs/2009.12937).
- [3] BANERJEE, S. and BUDHIRAJA, A. (2020). Parameter and dimension dependence of convergence rates to stationarity for reflecting Brownian motions. *Ann. Appl. Probab.* **30** 2005–2029. [MR4149521](https://doi.org/10.1214/19-AAP1550) <https://doi.org/10.1214/19-AAP1550>
- [4] BASS, R. F. and PARDOUX, É. (1987). Uniqueness for diffusions with piecewise constant coefficients. *Probab. Theory Related Fields* **76** 557–572. [MR0917679](https://doi.org/10.1007/BF00960074) <https://doi.org/10.1007/BF00960074>
- [5] BILLINGSLEY, P. (2013). *Convergence of Probability Measures*. Wiley, New York.
- [6] BUDHIRAJA, A. and LEE, C. (2007). Long time asymptotics for constrained diffusions in polyhedral domains. *Stochastic Process. Appl.* **117** 1014–1036. [MR2340877](https://doi.org/10.1016/j.spa.2006.11.007) <https://doi.org/10.1016/j.spa.2006.11.007>
- [7] CABEZAS, M., DEMBO, A., SARANTSEV, A. and SIDORAVICIUS, V. (2019). Brownian particles with rank-dependent drifts: Out-of-equilibrium behavior. *Comm. Pure Appl. Math.* **72** 1424–1458. [MR3957396](https://doi.org/10.1002/cpa.21825) <https://doi.org/10.1002/cpa.21825>
- [8] DEMBO, A., JARA, M. and OLLA, S. (2019). The infinite Atlas process: Convergence to equilibrium. *Ann. Inst. Henri Poincaré Probab. Stat.* **55** 607–619. [MR3949947](https://doi.org/10.1214/17-aihp875) <https://doi.org/10.1214/17-aihp875>
- [9] DEMBO, A., SHKOLNIKOV, M., VARADHAN, S. R. S. and ZEITOUNI, O. (2016). Large deviations for diffusions interacting through their ranks. *Comm. Pure Appl. Math.* **69** 1259–1313. [MR3503022](https://doi.org/10.1002/cpa.21640) <https://doi.org/10.1002/cpa.21640>
- [10] DEMBO, A. and TSAI, L.-C. (2017). Equilibrium fluctuation of the Atlas model. *Ann. Probab.* **45** 4529–4560. [MR3737917](https://doi.org/10.1214/16-AOP1171) <https://doi.org/10.1214/16-AOP1171>
- [11] FERNHOLZ, E. R. (2002). Stochastic portfolio theory. In *Stochastic Portfolio Theory* 1–24. Springer.
- [12] FERNHOLZ, R. and KARATZAS, I. (2009). Stochastic portfolio theory: An overview. *Handb. Numer. Anal.* **15** 89–167.
- [13] HARRISON, J. M. and REIMAN, M. I. (1981). Reflected Brownian motion on an orthant. *Ann. Probab.* **9** 302–308. [MR0606992](https://doi.org/10.1214/16-AOP1171)
- [14] HARRISON, J. M. and WILLIAMS, R. J. (1987). Brownian models of open queueing networks with homogeneous customer populations. *Stochastics* **22** 77–115. [MR0912049](https://doi.org/10.1080/17442508708833469) <https://doi.org/10.1080/17442508708833469>
- [15] HARRISON, J. M. and WILLIAMS, R. J. (1987). Multidimensional reflected Brownian motions having exponential stationary distributions. *Ann. Probab.* **15** 115–137. [MR0877593](https://doi.org/10.1214/09-AAP641)
- [16] ICHIBA, T. and KARATZAS, I. (2010). On collisions of Brownian particles. *Ann. Appl. Probab.* **20** 951–977. [MR2680554](https://doi.org/10.1214/09-AAP641) <https://doi.org/10.1214/09-AAP641>
- [17] ICHIBA, T., KARATZAS, I. and SHKOLNIKOV, M. (2013). Strong solutions of stochastic equations with rank-based coefficients. *Probab. Theory Related Fields* **156** 229–248. [MR3055258](https://doi.org/10.1007/s00440-012-0426-3) <https://doi.org/10.1007/s00440-012-0426-3>
- [18] ICHIBA, T. and SARANTSEV, A. (2017). Yet another condition for absence of collisions for competing Brownian particles. *Electron. Commun. Probab.* **22** Paper No. 8. [MR3607803](https://doi.org/10.1214/17-ECP41) <https://doi.org/10.1214/17-ECP41>
- [19] JOURDAIN, B. and MALRIEU, F. (2008). Propagation of chaos and Poincaré inequalities for a system of particles interacting through their CDF. *Ann. Appl. Probab.* **18** 1706–1736. [MR2462546](https://doi.org/10.1214/07-AAP513) <https://doi.org/10.1214/07-AAP513>
- [20] KARATZAS, I., PAL, S. and SHKOLNIKOV, M. (2016). Systems of Brownian particles with asymmetric collisions. *Ann. Inst. Henri Poincaré Probab. Stat.* **52** 323–354. [MR3449305](https://doi.org/10.1214/14-AIHP646) <https://doi.org/10.1214/14-AIHP646>
- [21] KELLA, O. and WHITT, W. (1996). Stability and structural properties of stochastic storage networks. *J. Appl. Probab.* **33** 1169–1180. [MR1416235](https://doi.org/10.1017/s0021900200100567) <https://doi.org/10.1017/s0021900200100567>

- [22] KLENKE, A. (2014). *Probability Theory: A Comprehensive Course*, 2nd ed. *Universitext*. Springer, London. MR3112259 <https://doi.org/10.1007/978-1-4471-5361-0>
- [23] LINDVALL, T. (1999). On Strassen's theorem on stochastic domination. *Electron. Commun. Probab.* **4** 51–59. MR1711599 <https://doi.org/10.1214/ECP.v4-1005>
- [24] MIKOSCH, T. (1999). *Regular Variation, Subexponentiality and Their Applications in Probability Theory* **99**. Eindhoven Univ. Technology Eindhoven, The Netherlands.
- [25] PAL, S. and PITMAN, J. (2008). One-dimensional Brownian particle systems with rank-dependent drifts. *Ann. Appl. Probab.* **18** 2179–2207. MR2473654 <https://doi.org/10.1214/08-AAP516>
- [26] SARANTSEV, A. (2015). Triple and simultaneous collisions of competing Brownian particles. *Electron. J. Probab.* **20** no. 29. MR3325099 <https://doi.org/10.1214/EJP.v20-3279>
- [27] SARANTSEV, A. (2017). Infinite systems of competing Brownian particles. *Ann. Inst. Henri Poincaré Probab. Stat.* **53** 2279–2315. MR3729655 <https://doi.org/10.1214/16-AIHP791>
- [28] SARANTSEV, A. and TSAI, L.-C. (2017). Stationary gap distributions for infinite systems of competing Brownian particles. *Electron. J. Probab.* **22** Paper No. 56. MR3672832 <https://doi.org/10.1214/17-EJP78>
- [29] TANG, W. and TSAI, L.-C. (2018). Optimal surviving strategy for drifted Brownian motions with absorption. *Ann. Probab.* **46** 1597–1650. MR3785596 <https://doi.org/10.1214/17-AOP1211>
- [30] TSAI, L.-C. (2018). Stationary distributions of the Atlas model. *Electron. Commun. Probab.* **23** Paper No. 10. MR3771768 <https://doi.org/10.1214/18-ECP112>
- [31] WAINWRIGHT, M. J. (2019). *High-Dimensional Statistics: A Non-asymptotic Viewpoint*. *Cambridge Series in Statistical and Probabilistic Mathematics* **48**. Cambridge Univ. Press, Cambridge. MR3967104 <https://doi.org/10.1017/9781108627771>
- [32] WILLIAMS, R. J. (1995). Semimartingale reflecting Brownian motions in the orthant. In *Stochastic Networks*. *IMA Vol. Math. Appl.* **71** 125–137. Springer, New York. MR1381009 https://doi.org/10.1007/978-1-4757-2418-9_7