



David Loeffler · Christopher Skinner · Sarah Livia Zerbes

Euler systems for $\mathrm{GSp}(4)$

Received March 18, 2019

Abstract. We construct an Euler system for Galois representations associated to cohomological cuspidal automorphic representations of GSp_4 , using the pushforwards of Eisenstein classes for $\mathrm{GL}_2 \times \mathrm{GL}_2$.

Keywords. Euler systems, Siegel modular forms

Contents

1. Introduction	669
2. General notation	673
3. Preliminaries I: Local representation theory	673
4. Preliminaries II: Algebraic representations and Lie theory	689
5. Modular varieties	692
6. Coefficient sheaves on modular varieties	696
7. Eisenstein classes for GL_2	699
8. Construction of Lemma–Eisenstein classes	702
9. Moment maps and p -adic interpolation	710
10. Mapping to Galois cohomology	717
11. Selmer groups and p -adic L -functions	727
References	730

1. Introduction

The theory of Euler systems is one of the most powerful tools available for studying the arithmetic of global Galois representations. However, constructing Euler systems is a difficult problem, and the list of known constructions is accordingly rather short.

David Loeffler: Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK;
d.a.loeffler@warwick.ac.uk

Christopher Skinner: Department of Mathematics, Princeton University, Princeton, NJ
 08544-1000, USA; cmcls@princeton.edu

Sarah Livia Zerbes (corresponding author): Department of Mathematics, University College
 London, London WC1E 6BT, UK; s.zerbes@ucl.ac.uk

Mathematics Subject Classification (2020): 11F46, 11F67, 11F80, 11R23

In this paper, we construct a new example of an Euler system, for the 4-dimensional Galois representations associated to cohomological cuspidal automorphic representations of $\mathrm{GSp}_4/\mathbb{Q}$, and apply this to studying the Selmer groups of these Galois representations. Our construction relies crucially on an unexpected relation with branching problems in smooth representation theory, which is the key input in proving the norm-compatibility relations for our Euler system classes.

We construct this Euler system in the étale cohomology of the Shimura variety of GSp_4 . The strategy that we use for this construction is also applicable to many other examples of Shimura varieties, including those associated to the groups $\mathrm{GU}(2, 1)$, GSp_6 , and $\mathrm{GSp}_4 \times \mathrm{GL}_2$, which will be explored in forthcoming work.

The starting point for our construction is a family of motivic cohomology classes for Siegel threefolds, which were introduced and studied by Francesco Lemma [20–22]. Lemma’s classes are constructed by using the subgroup $H = \mathrm{GL}_2 \times_{\mathrm{GL}_1} \mathrm{GL}_2$ inside GSp_4 . Beilinson’s Eisenstein symbol gives a supply of motivic cohomology classes for the Shimura varieties attached to H , and pushing these forward to GSp_4 gives motivic cohomology classes for the Siegel threefold. By applying the étale realisation map and projecting to an appropriate Hecke eigenspace, Lemma’s motivic classes give rise to elements of the groups $H^1(\mathbb{Q}, W_\Pi^*(-q))$, where Π is a suitable automorphic representation of GSp_4 , W_Π^* the dual of the associated p -adic Galois representation, and q is an integer in a certain range depending on the weight of Π .

To build an Euler system for these representations W_Π^* , we need to modify this construction in order to obtain classes defined over cyclotomic fields $\mathbb{Q}(\zeta_m)$. These classes are required to satisfy an appropriate norm-compatibility relation as m changes, and to take values in a $\mathbb{Z}_p$ -lattice in W_Π^* independent of m . We define these classes by translating the natural embedding of H in G via appropriately chosen elements of $G(\mathbb{A}_f)$, following a strategy that has been successfully used in several earlier Euler-system constructions [18, 19].

Using the theory of Λ -adic Eisenstein classes initiated by Kings, we show that these Euler system classes can be interpolated p -adically as the parameters (including the Tate twist q) vary. This leads to a definition of a “motivic p -adic L -function” for Π , which is a p -adic measure on $\mathbb{Z}_p^\times$ interpolating the images of the Euler system classes under the Bloch–Kato logarithm and dual-exponential maps at p . Assuming various technical hypotheses, we prove in §11 that if this motivic L -function is non-vanishing for a value of q such that $W_\Pi^*(-q)$ is critical (in the sense of Deligne), then the corresponding Bloch–Kato Selmer groups $H_f^1(\mathbb{Q}, W_\Pi^*(-q))$ and $H_f^1(\mathbb{Q}, W_\Pi(1+q))$ are zero. Our motivic p -adic L -function should interpolate the critical values of the spin L -function of Π (that is, we expect an “explicit reciprocity law”, analogous to those that have been proved for the Beilinson–Kato and Beilinson–Flach Euler systems). If such an explicit reciprocity law holds, then our bounds for H_f^1 would give new cases of the Bloch–Kato conjecture. The construction of a spin p -adic L -function and the proof of an explicit reciprocity law are the topics of forthcoming joint work with Vincent Pilloni.

One of the chief novelties of our construction is in the proofs of the norm-compatibility relations for the Euler system classes. In place of the (exceedingly laborious) double-coset computations used in [18] for example, we use methods of smooth representation

theory to reduce the norm-compatibility statement to a far easier, purely local statement involving Bessel models of unramified representations of $\mathrm{GSp}_4(\mathbb{Q}_\ell)$. This reduction is possible thanks to a case of the local Gan–Gross–Prasad conjecture due to Kato, Murase and Sugano, showing that the space of $\mathrm{SO}_4(\mathbb{Q}_\ell)$ -invariant linear functionals on an irreducible spherical representation of $\mathrm{SO}_4(\mathbb{Q}_\ell) \times \mathrm{SO}_5(\mathbb{Q}_\ell)$ has dimension ≤ 1 . This technique promises to be applicable in many other settings where local multiplicity 1 results of this type are known; for instance, in [23] we use a similar approach to prove norm-compatibility relations in an Euler system for the Shimura variety of the unitary group $\mathrm{GU}(2, 1)$, using the local Gan–Gross–Prasad conjecture for the pair $(\mathrm{U}(2), \mathrm{U}(3))$.

Outline of the paper

For the benefit of the reader, we give a brief outline of how our Euler system classes are constructed, and how the norm-compatibility relations for these are proved.

Construction of the elements. Let $G = \mathrm{GSp}_4$, and for each open compact $U \subset G(\mathbb{A}_f)$, let $Y_G(U)$ be the Siegel three-fold of level U (a Shimura variety for G). We construct a map of $G(\mathbb{A}_f)$ -representations, the *Lemma–Eisenstein map*,

$$\mathcal{LE} : I \otimes_{\mathcal{H}(H(\mathbb{A}_f))} \mathcal{H}(G(\mathbb{A}_f)) \rightarrow \varinjlim_U H_{\mathrm{mot}}^4(Y_G(U), \mathcal{D})$$

where $\mathcal{H}(-)$ denotes the Hecke algebra, $\mathcal{D}$ is a relative Chow motive (a “motivic sheaf”) over Y_G arising from some algebraic representation of G , and I is a certain explicit representation of $H(\mathbb{A}_f)$. This construction depends on parameters a, b, q, r , specifying weights for G and for H , but we shall suppress this for now. The construction of $\mathcal{LE}$, given in §8.3, is essentially formal: the representation I records the data needed to define an Eisenstein class in the motivic cohomology of Y_H , and $\mathcal{LE}$ maps this Eisenstein class to a linear combination of $G(\mathbb{A}_f)$ -translates of its pushforward to Y_G , with the $\mathcal{H}(G(\mathbb{A}_f))$ term recording which translations to apply.

Let K be a level, unramified outside $S \cup \{p\}$ (where $S \not\ni p$ is a finite set of primes) and having a certain specific form at p . Then for any n coprime to S , the base-extension $Y_G(K) \times_{\mathrm{Spec} \mathbb{Q}} \mathrm{Spec} \mathbb{Q}(\mu_n)$ is itself a Shimura variety for some subgroup $K' \subseteq K$. In §8.4 we use these isomorphisms, and certain explicit choices of test data as input to $\mathcal{LE}$, to define a collection of classes

$$z_{M,m} \in H_{\mathrm{mot}}^4(Y_G(K) \times \mathrm{Spec} \mathbb{Q}(\mu_{Mp^m}), \mathcal{D})$$

for $m \geq 0$ and M square-free and coprime to pS . These are our Euler system classes.

Norm-compatibility. The “ideal” norm-compatibility result for these classes would be an identity of the form

$$\mathrm{norm}(z_{\ell M,m}) = \mathcal{P}_\ell(\sigma_\ell^{-1}) \cdot z_{M,m}$$

for primes $\ell \nmid MpS$. Here “norm” denotes the Galois norm map from $\mathbb{Q}(\mu_{M\ell p^m})$ to $\mathbb{Q}(\mu_{Mp^m})$, σ_ℓ is the Frobenius at ℓ , and $\mathcal{P}_\ell(X)$ is a degree-4 polynomial with coefficients in the spherical Hecke algebra at ℓ , which acts on each irreducible representation as the corresponding spin L -factor. However, we cannot prove the full strength of this statement here (we hope to return to this issue in a later paper). Instead, we prove a version of this result after mapping to Galois cohomology. We choose Π a suitably nice cohomological automorphic representation of GSp_4 such that $\Pi_f^K \neq 0$. (We need Π_ℓ to be generic for almost all ℓ , which excludes certain “endoscopic” representations such as Saito–Kurokawa lifts.) Then $\Pi_f^* \otimes W_\Pi^*$ appears with multiplicity 1 as a direct summand of $\lim_{\rightarrow U} H_{\text{ét}}^3(Y_G(U)_{\overline{\mathbb{Q}}}, \mathcal{D})$, and does not contribute to cohomology outside degree 3. Choosing a vector $\varphi \in \Pi_f$ thus gives a homomorphism of Galois representations

$$\Pi_f^* \otimes W_\Pi^* \rightarrow W_\Pi^*,$$

which factors through $(\Pi_f^*)^K$ if φ is K -invariant. Combining this with the Hochschild–Serre spectral sequence gives a map of vector spaces

$$H_{\text{ét}}^4(Y_G(K) \times \mathrm{Spec} \mathbb{Q}(\mu_{Mp^m}), \mathcal{D}) \rightarrow H^1(\mathbb{Q}(\mu_{Mp^m}), W_\Pi^*).$$

We thus obtain a collection of cohomology classes $z_{M,m}^\Pi \in H^1(\mathbb{Q}(\mu_{Mp^m}), W_\Pi^*)$, depending on the choice of φ , and we shall prove the norm-compatibility relations for these instead.

For simplicity, assume that $M = 1$ and $m = 0$, so we are trying to compare $z_{1,0}^\Pi$ with $\mathrm{norm}(z_{\ell,0}^\Pi)$ (the general case can be reduced to this by twisting). We have constructed a $G(\mathbb{A}_f)$ -equivariant bilinear pairing

$$\left(I \otimes_{\mathcal{H}(H(\mathbb{A}_f))} \mathcal{H}(G(\mathbb{A}_f)) \right) \otimes \Pi_f \rightarrow H^1(\mathbb{Q}, W_\Pi^*)$$

or equivalently (via Frobenius reciprocity) an $H(\mathbb{A}_f)$ -equivariant pairing

$$\mathfrak{z} : I \otimes \Pi_f \rightarrow H^1(\mathbb{Q}, W_\Pi^*).$$

By construction the classes $z_{1,0}^\Pi$ and $\mathrm{norm}(z_{\ell,0}^\Pi)$ are values of this pairing, at different choices of test data $v, v' \in I \otimes \Pi_f$. In most cases (away from a few small weights) the representation I is a direct sum of principal series representations τ , each of which factors as $\bigotimes_{w \text{ prime}} \tau_w$; and by construction the projections of v and v' to $\tau_w \otimes \Pi_w$ coincide for $w \neq \ell$.

It is at this point that the decisive input from local representation theory appears: known cases of the Gan–Gross–Prasad conjecture imply that $\mathrm{Hom}_{H(\mathbb{Q}_\ell)}(\tau_\ell \otimes \Pi_\ell, 1_H)$ is 1-dimensional, and we can construct a canonical basis $\mathfrak{z}_\ell$ of this space using zeta-integrals. So it suffices to show that $\mathfrak{z}_\ell(v_\ell) = P_\ell(1)\mathfrak{z}_\ell(v'_\ell)$, which is a simple, purely local computation which we carry out in §3. It then follows that $\mathfrak{z}'_\ell(v_\ell) = P_\ell(1)\mathfrak{z}'_\ell(v'_\ell)$ for every H -equivariant homomorphism $\mathfrak{z}'_\ell$ from $\tau_\ell \otimes \Pi_\ell$ to a space with trivial H -action, and the desired norm relation follows (see Proposition 10.5.2).

2. General notation

- Let J be the skew-symmetric 4×4 matrix over $\mathbb{Z}$ given by $\begin{pmatrix} & & & 1 \\ & & 1 & \\ & -1 & & \\ -1 & & & \end{pmatrix}$. We let $G = \mathrm{GSp}_4$ be the group scheme over $\mathbb{Z}$ defined by

$$G(R) := \mathrm{GSp}_4(R) = \{(g, \mu) \in \mathrm{GL}_4(R) \times \mathrm{GL}_1(R) : g^t \cdot J \cdot g = \mu J\}$$

for any commutative unital ring R . We write $\mu : G \rightarrow \mathrm{GL}_1$ for the symplectic multiplier map.

- We define the *standard Borel subgroup* $B \subseteq G$ to be the subgroup $\{(g, \mu) : g \text{ is upper-triangular}\}$.
- We define a *standard parabolic subgroup* to be a subgroup of G containing B ; there are exactly four of these, namely B , G , the *Siegel parabolic* P_S and the *Klingen parabolic* P_{Kl} , where

$$P_S = \begin{pmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ * & * & & \end{pmatrix}, \quad P_{\mathrm{Kl}} = \begin{pmatrix} * & * & * & * \\ * & * & * & * \\ * & * & * & * \\ & * & * & \end{pmatrix}.$$

- We write T for the diagonal torus of G , which is equal to the product $A \times T'$, where A, T' are the tori defined by

$$A = \begin{pmatrix} x & & & \\ & y & & \\ & & x & \\ & & & y \end{pmatrix}, \quad T' = \begin{pmatrix} x & & & \\ & x & & \\ & & 1 & \\ & & & 1 \end{pmatrix}.$$

- Let $H = \mathrm{GL}_2 \times_{\mathrm{GL}_1} \mathrm{GL}_2$ (fibre product over the determinant map), and let ι denote the embedding $H \hookrightarrow G$ given by

$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \right) \mapsto \begin{pmatrix} a & & & b \\ & a' & b' & \\ & c' & d' & \\ c & & & d \end{pmatrix}.$$

We write $B_H = \iota^{-1}(B) = \iota^{-1}(P_S)$ for the standard Borel subgroup of H .

Remark 2.1. The quotient of G by its centre Z_G is the split form of the orthogonal group SO_5 . We have $Z_G \subset \iota(H)$, and the image of H/Z_G in G/Z_G via ι is the split form of SO_4 , embedded as the stabiliser of an anisotropic vector in the defining 5-dimensional representation. This will be used in §3.7 below, in order to make use of the results of the Gan–Gross–Prasad theory of restriction of representations of SO_5 to SO_4 . $\diamond$

3. Preliminaries I: Local representation theory

In this section, we fix a prime ℓ and collect some definitions and results regarding smooth representations of the groups $\mathrm{GL}_2(\mathbb{Q}_\ell)$, $G(\mathbb{Q}_\ell)$, and $H(\mathbb{Q}_\ell)$ on complex vector spaces. For brevity we shall write $G_\ell = G(\mathbb{Q}_\ell)$ and similarly for H_ℓ .

3.1. Principal series representations of $\mathrm{GL}_2(\mathbb{Q}_\ell)$

Notation 3.1.1. We write $\mathrm{d}x$ and $\mathrm{d}^\times x$ for the Haar measures on $\mathbb{Q}_\ell$ and $\mathbb{Q}_\ell^\times$ normalised so that $\mathbb{Z}_\ell$ (resp. $\mathbb{Z}_\ell^\times$) has volume 1. The norm $|\cdot|$ is normalised so that $|\ell| = 1/\ell$. If χ is a smooth character of $\mathbb{Q}_\ell^\times$, we write $L(\chi, s)$ for its local L -factor, which is

$$L(\chi, s) = L(\chi|\cdot|^s, 0) = \begin{cases} (1 - \chi(\ell)\ell^{-s})^{-1} & \text{if } \chi|_{\mathbb{Z}_\ell^\times} = 1, \\ 1 & \text{otherwise.} \end{cases}$$

Definition 3.1.2. Given two smooth characters χ and ψ of $\mathbb{Q}_\ell^\times$, we let $I(\chi, \psi)$ be the space of smooth functions $f : \mathrm{GL}_2(\mathbb{Q}_\ell) \rightarrow \mathbb{C}$ such that

$$f\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} g\right) = \chi(a)\psi(d)|a/d|^{1/2} f(g),$$

equipped with a $\mathrm{GL}_2(\mathbb{Q}_\ell)$ -action via right translation of the argument.

As is well-known, the pairing $I(\chi, \psi) \times I(\chi^{-1}, \psi^{-1}) \rightarrow \mathbb{C}$ defined by

$$\langle f_1, f_2 \rangle = \int_{\mathrm{GL}_2(\mathbb{Z}_\ell)} f_1(g) f_2(g) \mathrm{d}g,$$

where we normalise the measure so that $\mathrm{GL}_2(\mathbb{Z}_\ell)$ has volume 1, identifies $I(\chi^{-1}, \psi^{-1})$ with the dual of $I(\chi, \psi)$. Moreover, if $\chi/\psi \neq |\cdot|^{\pm 1}$, then $I(\chi, \psi)$ is an irreducible representation.

We will frequently need to use analytic continuation in an auxiliary parameter s . The following construction will be helpful:

Definition 3.1.3. A *polynomial section* of the family of representations $I(\chi|\cdot|^s, \psi|\cdot|^{-s})$ is a function on $\mathrm{GL}_2(\mathbb{Q}_\ell) \times \mathbb{C}$, $(g, s) \mapsto f_s(g)$, such that $g \mapsto f_s(g)$ is in $I(\chi|\cdot|^s, \psi|\cdot|^{-s})$ for each $s \in \mathbb{C}$, and $s \mapsto f_s(g)$ lies in $\mathbb{C}[\ell^s, \ell^{-s}]$ for every $g \in \mathrm{GL}_2(\mathbb{Q}_\ell)$. A section is *flat* if its restriction to $\mathrm{GL}_2(\mathbb{Z}_\ell)$ is independent of s .

From the Iwasawa decomposition, one sees that every $f \in I(\chi, \psi)$ extends to a unique flat section. The space of polynomial sections is stable under the action of $\mathrm{GL}_2(\mathbb{Q}_\ell)$ (while the space of flat sections clearly is not).

Definition 3.1.4. Let $M : I(\chi, \psi) \rightarrow I(\psi, \chi)$ be the normalised standard intertwining operator, defined by analytic continuation to $s = 0$ of the integral

$$M(f_s; g) := L(\chi/\psi, 2s)^{-1} \int_{\mathbb{Q}_\ell} f_s(w \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} g) \mathrm{d}n$$

where $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ is the long Weyl element.

More precisely, if $|\chi/\psi| = |\cdot|^r$, and $(f_s)_{s \in \mathbb{C}}$ is any polynomial section, then the intertwining integral for $M(f_s; g)$ is absolutely convergent for $r + 2\Re(s) > 0$ and defines a polynomial section of $I(\psi|\cdot|^{-s}, \chi|\cdot|^s)$ (see e.g. [3, Proposition 4.5.7] for further details). The specialisation of this section at $s = 0$ depends only on $f_0 \in I(\chi, \psi)$, not on the choice of section passing through f_0 , and this defines a non-zero intertwiner between $I(\chi, \psi)$ and $I(\psi, \chi)$ (even in the exceptional case $\chi = \psi$).

Proposition 3.1.5. *Suppose χ and ψ are unramified. Then for all $f_1 \in I(\chi, \psi)$ and all $f_2 \in I(\psi^{-1}, \chi^{-1})$ we have*

$$\langle M(f_1), f_2 \rangle = \langle f_1, M(f_2) \rangle.$$

Proof. By choosing polynomial sections passing through the f_i , we may assume without loss of generality that $\chi/\psi \neq |\cdot|^{\pm 1}$, so that both $I(\chi, \psi)$ and $I(\psi^{-1}, \chi^{-1})$ are irreducible. Hence it suffices to check the equality when f_1 and f_2 are the respective spherical vectors (normalised so that $f_i(1) = 1$). With our conventions, M sends the normalised spherical vector of $I(\chi, \psi)$ to the normalised spherical vector of $I(\psi, \chi)$, and these normalised vectors pair to 1 under the duality pairing. ■

3.2. Siegel sections

Notation 3.2.1. Let $\mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$ denote the space of Schwartz functions (locally constant, compactly supported functions) on $\mathbb{Q}_\ell^2$. We let $\mathrm{GL}_2(\mathbb{Q}_\ell)$ act on this space from the left by $(g \cdot \phi)(x, y) = \phi((x, y) \cdot g)$ for $g \in \mathrm{GL}_2(\mathbb{Q}_\ell)$ and $\phi \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$. For $\phi \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$, we define its Fourier transform $\hat{\phi}$ by

$$\hat{\phi}(x, y) = \iint e_\ell(xv - yu) \phi(u, v) \, du \, dv,$$

where $e_\ell(x)$ is the standard additive character of $\mathbb{Q}_\ell$, mapping $1/\ell^n$ to $\exp(2\pi i/\ell^n)$.

Proposition 3.2.2. *Let $\phi \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$, and let χ, ψ be characters of $\mathbb{Q}_\ell^\times$ with $|\chi/\psi| = |\cdot|^r$. Then the integral*

$$f_{\phi, \chi, \psi}(g, s) := \frac{\chi(\det g) |\det g|^{s+1/2}}{L(\chi/\psi, 2s+1)} \int_{\mathbb{Q}_\ell^\times} \phi((0, x)g) (\chi/\psi)(x) |x|^{2s+1} \, d^\times x$$

converges for $r + 2\Re(s) > -1$, and defines a polynomial section of $I(|\chi| \cdot |\cdot|^s, |\psi| \cdot |\cdot|^{-s})$, so $f_{\phi, \chi, \psi}(g) := f_{\phi, \chi, \psi}(g, 0) \in I(\chi, \psi)$ is well-defined. These elements satisfy

$$\begin{aligned} f_{g \cdot \phi, \chi, \psi}(h) &= \chi(\det g)^{-1} |\det g|^{-1/2} f_{\phi, \chi, \psi}(hg), \\ \widehat{f_{g \cdot \phi, \chi, \psi}}(h) &= \psi(\det g)^{-1} |\det g|^{-1/2} \widehat{f_{\phi, \chi, \psi}}(hg) \end{aligned} \tag{1}$$

for all $g, h \in \mathrm{GL}_2(\mathbb{Q}_\ell)$.

Proof. The convergence of the integral, and its analytic continuation as a function of s , form part of Tate's theory of local zeta integrals for GL_1 . The fact that $f_{\phi, \chi, \psi}(-, s)$ lies in $I(|\chi| \cdot |\cdot|^s, |\psi| \cdot |\cdot|^{-s})$ is immediate from the definition in the region of convergence of the integral, and follows for all s by analytic continuation. The first of the transformation formulae (1) is obvious from the definition, and the second follows from the identity

$$\widehat{(g \cdot \phi)} = \frac{1}{|\det g|} ({}^t g \cdot \hat{\phi}), \quad \text{where } {}^t g = (\det g)^{-1} g. \quad \blacksquare$$

Proposition 3.2.3. *We have*

$$M(f_{\phi, \chi, \psi}) = \frac{\varepsilon(\psi/\chi)}{L(\chi/\psi, 1)} f_{\hat{\phi}, \psi, \chi},$$

where $\varepsilon(\psi/\chi)$ is the local ε -factor (a non-zero scalar, equal to 1 if χ/ψ is unramified).

Proof. This is a straightforward consequence of the functional equation for Tate's GL_1 zeta integral. ■

If $\chi/\psi = |\cdot|^{-1}$ we interpret the right-hand side as 0, so the elements $f_{\phi, \chi, \psi}$ all land in the 1-dimensional subrepresentation. Let us evaluate these integrals explicitly for some specific choices of ϕ , assuming now that χ and ψ are *unramified* characters.

Definition 3.2.4. We define functions $\phi_t \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$ for integers $t \geq 0$ as follows.

- For $t = 0$, we let $\phi_0 := \mathrm{ch}(\mathbb{Z}_\ell \times \mathbb{Z}_\ell)$.
- For $t > 0$, we let $\phi_t := \mathrm{ch}(\ell^t \mathbb{Z}_\ell \times \mathbb{Z}_\ell^\times)$.

Note that ϕ_t is preserved by the action of the group $K_0(\ell^t) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}_\ell) : c = 0 \bmod \ell^t \right\}$.

Lemma 3.2.5. *We have*

$$f_{\phi_t, \chi, \psi}(1) = \begin{cases} 1 & \text{if } t = 0, \\ L(\chi/\psi, 1)^{-1} & \text{if } t > 0. \end{cases}$$

Moreover, the function $f_{\phi_t, \chi, \psi}$ is supported on $B(\mathbb{Q}_\ell)K_0(\ell^t)$.

Proof. The computation of the value at the identity is immediate. The assertion regarding the support of the function is vacuous for $t = 0$, and for $t > 1$ we have

$$\phi_t = \begin{pmatrix} \ell^{1-t} & 0 \\ 0 & 1 \end{pmatrix} \phi_1,$$

so in fact it suffices to prove the assertion for $t = 1$; in this case, we simply observe that the function $f_{\phi_1, \chi, \psi}$ vanishes on the long Weyl element $w = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, since $\phi_1((0, x)w) = \phi_1(-x, 0) = 0$ for all x . ■

3.3. Notation: subgroups of G_ℓ and H_ℓ

We now define an assortment of open compact subgroups of G_ℓ and H_ℓ . We represent elements of G_ℓ in block form $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where A, B, C, D are 2×2 matrices.

- $K_{G_\ell} = G(\mathbb{Z}_\ell)$.
- $K_{G_\ell, 0}(\ell^n) = \{g \in G(\mathbb{Z}_\ell) : g = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \bmod \ell^n\}$ for $n \geq 0$.
- $K_{G_\ell, 1}(\ell^n) = \{g \in G(\mathbb{Z}_\ell) : g = \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \bmod \ell^n\}$ for $n \geq 0$.
- $K_{G_\ell}(\ell^m, \ell^n) = \{g \in K_{G_\ell, 1}(\ell^n) : \mu(g) = 1 \bmod \ell^m\}$ for $m, n \geq 0$.
- $K'_{G_\ell}(\ell^m, \ell^n) = \{g \in K_{G_\ell, 1}(\ell^n) : g = 1 \bmod \ell^m\}$ for $n \geq m \geq 0$.

(The subgroups $K'_G(\ell^m, \ell^n)$ will not be used until §8.4.) We define subgroups K_{H_ℓ} , $K_{H_\ell, 0}(\ell^n)$, etc. of H_ℓ as the preimages (via ι) of the corresponding subgroups of G_ℓ . We write dg and dh for the Haar measures on G_ℓ and H_ℓ normalised so that K_{G_ℓ} (resp. K_{H_ℓ}) has volume 1.

3.4. Induced representations of H_ℓ

Given two pairs of characters $\underline{\chi} = (\chi_1, \chi_2)$ and $\underline{\psi} = (\psi_1, \psi_2)$, we define $I_H(\underline{\chi}, \underline{\psi})$ as the representation of H_ℓ given by the normalised induction from $B_H(\mathbb{Q}_\ell)$ of the character

$$\begin{pmatrix} a & & & \\ & a' & & \\ & & b' & \\ & & & b \end{pmatrix} \mapsto \chi_1(a)\psi_1(b)\chi_2(a')\psi_2(b') \quad (\text{where } ab = a'b').$$

Since H_ℓ acts transitively on $\mathbb{P}^1(\mathbb{Q}_\ell) \times \mathbb{P}^1(\mathbb{Q}_\ell)$, restriction of functions from $\mathrm{GL}_2(\mathbb{Q}_\ell) \times \mathrm{GL}_2(\mathbb{Q}_\ell)$ to H_ℓ defines an isomorphism of H_ℓ -representations $I(\chi_1, \psi_1) \otimes I(\chi_2, \psi_2) \rightarrow I_H(\underline{\chi}, \underline{\psi})$, where H_ℓ acts on the source via its inclusion in $\mathrm{GL}_2(\mathbb{Q}_\ell) \times \mathrm{GL}_2(\mathbb{Q}_\ell)$. In particular, there is an intertwining operator $M : I_H(\underline{\chi}, \underline{\psi}) \rightarrow I_H(\underline{\psi}, \underline{\chi})$ given by the tensor product of the two GL_2 intertwining operators.

Proposition 3.4.1. *If there is no quadratic character η such that $\chi_1/\psi_1 = \chi_2/\psi_2 = \eta$, then every irreducible subquotient of $I(\chi_1, \psi_1) \otimes I(\chi_2, \psi_2)$ as a representation of $\mathrm{GL}_2(\mathbb{Q}_\ell) \times \mathrm{GL}_2(\mathbb{Q}_\ell)$ remains irreducible as a representation of H_ℓ .*

Proof. This is an instance of [10, Lemma 2.1]. ■

Given $\underline{\phi} = \sum \phi_{i,1} \otimes \phi_{i,2} \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})^{\otimes 2}$, we let $f_{\underline{\phi}, \underline{\chi}, \underline{\psi}}$ be the image in $I_H(\underline{\chi}, \underline{\psi})$ of the element

$$\sum_i f_{\phi_{i,1}, \chi_1, \psi_1} \otimes f_{\phi_{i,2}, \chi_2, \psi_2} \in I(\chi_1, \psi_1) \otimes I(\chi_2, \psi_2).$$

For a non-negative integer t , let $\underline{\phi}_t = \phi_t \otimes \phi_t$.

Proposition 3.4.2. *Let $t \geq 0$. Then:*

(a) *We have*

$$f_{\underline{\phi}_t, \underline{\chi}, \underline{\psi}}(1) = \begin{cases} 1 & \text{if } t = 0, \\ L(\chi_1/\psi_1, 1)^{-1} L(\chi_2/\psi_2, 1)^{-1} & \text{if } t \geq 1. \end{cases}$$

(b) *$f_{\underline{\phi}_t, \underline{\chi}, \underline{\psi}}$ is supported on $B_H(\mathbb{Q}_\ell)K_{H_\ell, 0}(\ell^t)$.*

Proof. Follows from Lemma 3.2.5. ■

3.5. Representations of G_ℓ

3.5.1. Principal series representations of G_ℓ . We follow the notations of [34] for representations of G_ℓ . See *op. cit.* for further details (in particular §2.2 and the tables in Appendix A).

Definition 3.5.1. Let χ_1, χ_2, ρ be smooth characters of $\mathbb{Q}_\ell^\times$ such that

$$|\cdot|^{\pm 1} \notin \{\chi_1, \chi_2, \chi_1\chi_2, \chi_1/\chi_2\}. \quad (2)$$

We let $\chi_1 \times \chi_2 \rtimes \rho$ denote the representation of G_ℓ afforded by the space of smooth functions $f : G_\ell \rightarrow \mathbb{C}$ satisfying

$$f\left(\begin{pmatrix} a & * & * & * \\ & b & * & * \\ & & cb^{-1} & * \\ & & & ca^{-1} \end{pmatrix} g\right) = \frac{|a^2b|}{|c|^{3/2}} \chi_1(a) \chi_2(b) \rho(c) f(g),$$

with G_ℓ acting by right translation. We refer to such representations as *irreducible principal series*.

The above representation has central character $\chi_1\chi_2\rho^2$; the condition (2) implies that it is irreducible and generic. If η is a smooth character of $\mathbb{Q}_\ell^\times$, then twisting $\chi_1 \times \chi_2 \rtimes \rho$ by η (regarded as a character of G_ℓ via the multiplier map) gives the representation $\chi_1 \times \chi_2 \rtimes \rho\eta$.

Definition 3.5.2. Let $\sigma = \chi_1 \times \chi_2 \rtimes \rho$ be an irreducible principal series representation. The local (spin) L -factor of σ is the function

$$L(\sigma, s) = L(\sigma \otimes |\cdot|^s, 0) = L(\rho, s) L(\rho\chi_1, s) L(\rho\chi_2, s) L(\rho\chi_1\chi_2, s).$$

Proposition 3.5.3. If $\sigma = \chi_1 \times \chi_2 \rtimes \rho$ is an irreducible principal series representation, then σ is unramified if and only if all three characters χ_1, χ_2, ρ are unramified. Moreover, every irreducible, generic, unramified representation of G_ℓ is isomorphic to $\chi_1 \times \chi_2 \rtimes \rho$, for a unique Weyl-group orbit of unramified characters (χ_1, χ_2, ρ) satisfying (2).

Proof. See [34, §2.2]. ■

3.5.2. Hecke operators. Firstly, we consider the action of the spherical Hecke algebra $\mathcal{H}(K_{G_\ell} \backslash G_\ell / K_{G_\ell})$ on $\sigma^{K_{G_\ell}}$ when σ is an unramified principal series representation.

Lemma 3.5.4. Consider the following elements in the Hecke algebra $\mathcal{H}(K \backslash G_\ell / K)$:

$$\begin{aligned} T(\ell) &= K_{G_\ell} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ell & \\ & & & \ell \end{pmatrix} K_{G_\ell}, & T_1(\ell^2) &= K_{G_\ell} \begin{pmatrix} 1 & & & \\ & \ell & & \\ & & \ell & \\ & & & \ell^2 \end{pmatrix} K_{G_\ell}, \\ R(\ell) &= K_{G_\ell} \begin{pmatrix} \ell & & & \\ & \ell & & \\ & & \ell & \\ & & & \ell \end{pmatrix} K_{G_\ell}. \end{aligned}$$

If $\mathcal{P}_\ell(X)$ is the polynomial over $\mathcal{H}(K_{G_\ell} \backslash G_\ell / K_{G_\ell})$ defined by

$$1 - T(\ell)X + \ell(T_1(\ell^2) + (\ell^2 + 1)R(\ell))X^2 - \ell^3 T(\ell)R(\ell)X^3 + \ell^6 R(\ell)^2 X^4,$$

then for any unramified principal series representation $\sigma = \chi_1 \times \chi_2 \rtimes \rho$, $\mathcal{P}_\ell(\ell^{-s})$ acts on $\sigma^{K_{G_\ell}}$ as $L(\sigma, s - 3/2)^{-1}$.

Proof. See [41, §2.4]; our $\mathcal{P}_\ell(X)$ is $X^4 Q_\ell(1/X)$ in Taylor's notation. ■

Secondly, we consider the larger space of invariants under the Siegel parahoric subgroup $K_{G_\ell,0}(\ell)$. We let $U(\ell)$ denote the $K_{G_\ell,0}(\ell)$ -double coset of $\begin{pmatrix} \ell & & & \\ & \ell & & \\ & & 1 & \\ & & & 1 \end{pmatrix}$, which acts on $\sigma^{K_{G_\ell,0}(\ell)}$ via

$$x \mapsto \sum_{u,v,w \in \mathbb{Z}/\ell} \begin{pmatrix} \ell & u & v \\ & \ell & w \\ & & 1 & \\ & & & 1 \end{pmatrix} x.$$

Proposition 3.5.5. *If σ is an unramified principal series representation, then $\sigma^{K_{G_\ell,0}(\ell)}$ is 4-dimensional, and*

$$\det(1 - U(\ell)\ell^{-s} \mid \sigma^{K_{G_\ell,0}(\ell)}) = L(\sigma, s - 3/2)^{-1}.$$

Proof. See [41, Lemma 2.4]. ■

3.6. Zeta integrals for G

In this section we isolate the key local zeta integral calculations used in our proofs of the tame norm relations.

3.6.1. The Bessel model.

Definition 3.6.1. Let A be the torus $\left\{ \begin{pmatrix} x & & & \\ & y & & \\ & & x & \\ & & & y \end{pmatrix} : x, y \in \mathbb{G}_m \right\}$. The *Bessel subgroup* R of G is the semidirect product $A \ltimes N_S$, where N_S is the unipotent radical of the Siegel parabolic P_S .

Definition 3.6.2. Let λ be a character of A . A (split) λ -*Bessel functional* on a representation σ of G_ℓ is a linear functional $\mu : \sigma \rightarrow \mathbb{C}$ transforming under left-translation by $R(\mathbb{Q}_\ell)$ via the formula

$$\mu \left(\begin{pmatrix} 1 & u & v \\ & 1 & w \\ & & 1 & \\ & & & 1 \end{pmatrix} a \cdot \varphi \right) = e_\ell(u)\lambda(a)\mu(\varphi) \quad (3)$$

for all $\varphi \in \sigma$, $a \in A$, and $u, v, w \in \mathbb{Q}_\ell$.

If σ is irreducible, then the space of λ -Bessel functionals on σ has dimension ≤ 1 , by [35, Theorem 6.3.2]. It is clearly zero unless $\lambda|_{Z(G_\ell)}$ coincides with the central character of σ .

Theorem 3.6.3 (Roberts–Schmidt). *If σ is an irreducible generic representation of G_ℓ (such as an irreducible principal series representation), then σ admits a non-zero λ -Bessel functional μ_λ for every λ whose restriction to $Z(G_\ell)$ agrees with the central character of σ . If both σ and λ are unramified, then we may normalise μ_λ so that $\mu_\lambda(\varphi_0) = 1$, where φ_0 is the spherical vector of σ .*

Proof. For the existence of the Bessel functional see [35, Proposition 3.4.2]. It is shown in *op. cit.* that the Bessel functional can be explicitly constructed by integrating functions in the Whittaker model of σ ; and the assertion that in the unramified case the spherical vector maps to 1 under this functional follows, for example, from the computations of [34, §7.1]. ■

If σ is any irreducible representation admitting some λ -Bessel functional μ_λ , then for any $\varphi \in \sigma$ we may define a function $B_{\varphi,\lambda}$ on G_ℓ by $B_{\varphi,\lambda}(g) = \mu_\lambda(g \cdot \varphi)$. The space of functions $\{B_{\varphi,\lambda} : \varphi \in \sigma\}$ is the λ -Bessel model of σ .

Proposition 3.6.4. *Let σ be an irreducible representation of G_ℓ admitting a λ -Bessel model, and let $\varphi \in \sigma$ be invariant under $N_S(\mathbb{Z}_\ell)$. Define*

$$U(\ell^k)\varphi := \sum_{u,v,w \in \mathbb{Z}/\ell^k} \begin{pmatrix} 1 & u & v \\ & 1 & w \\ & & 1 \end{pmatrix} \begin{pmatrix} \ell^k & & \\ & \ell^k & \\ & & 1 \end{pmatrix} \varphi.$$

Then

$$B_{U(\ell^k)\varphi,\lambda} \left(\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix} \right) = \begin{cases} 0 & \text{if } |x| > 1, \\ \ell^{3k} B_{\varphi,\lambda} \left(\begin{pmatrix} \ell^k x & & \\ & \ell^k x & \\ & & 1 \end{pmatrix} \right) & \text{if } |x| \leq 1. \end{cases}$$

Proof. We first note that the assumption that the Bessel function $B_{\varphi,\lambda}$ is fixed by right-translation by $N_S(\mathbb{Z}_\ell)$, and transforms on the left via (3), implies that $B_{\varphi,\lambda} \left(\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix} \right)$ is zero if $|x| > 1$. We now compute

$$\begin{aligned} B_{U(\ell^k)\varphi,\lambda} \left(\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix} \right) &= \sum_{u,v,w} B_{\varphi,\lambda} \left(\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & u & v \\ & 1 & w \\ & & 1 \end{pmatrix} \begin{pmatrix} \ell^k & & \\ & \ell^k & \\ & & 1 \end{pmatrix} \right) \\ &= \ell^{2k} \sum_{u \bmod \ell^k} e_\ell(xu) B_{\varphi,\lambda} \left(\begin{pmatrix} \ell^k x & & \\ & \ell^k x & \\ & & 1 \end{pmatrix} \right). \end{aligned}$$

If $|x| > \ell^k$ then the Bessel function is zero; and if $\ell^k \geq |x| > 1$, then the sum of the e_ℓ terms vanishes. This leaves the cases $|x| \leq 1$, in which case the terms $e_\ell(xv)$ are all equal to 1 and we obtain the result. ■

3.6.2. Novodvorsky's integral. In order to construct an intertwining operator between σ and a principal-series H -representation, we shall use an integral involving a choice of vector in the Bessel model of σ , for some choice of character λ as above. For $\varphi \in \sigma$, η an unramified character of $\mathbb{Q}_\ell^\times$, and $s \in \mathbb{C}$, we define

$$Z(\varphi, \eta, \lambda, s) := L(\sigma \otimes \eta, s)^{-1} \int_{\mathbb{Q}_\ell^\times} B_{\varphi,\lambda} \left(\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix} \right) \eta(x) |x|^{s-3/2} d^\times x.$$

Here $L(\sigma \otimes \eta, s)$ is the spin L -factor of $\sigma \otimes \eta$, as in Definition 3.5.2.

Remark 3.6.5. This integral apparently first appears in [28, Equation 2.7]. In an earlier draft of the present paper, we mistakenly ascribed this construction to Sugano; in fact Sugano's paper [40] considers a related but slightly different integral – see Remark 3.7.4 below. ◇

Proposition 3.6.6. *Suppose σ is an irreducible unramified principal series representation, with central character χ_σ , and let η be an unramified character. Let λ be given by $\begin{pmatrix} x & & & \\ & y & & \\ & & x & \\ & & & y \end{pmatrix} \mapsto \lambda_1(x)\lambda_2(y)$ for unramified characters λ_1, λ_2 of $\mathbb{Q}_\ell^\times$.*

- (a) *The integral defining $Z(\varphi, \eta, \lambda, s)$ is absolutely convergent for $\Re(s) \gg 0$, and it has analytic continuation to all $s \in \mathbb{C}$ as an element of $\mathbb{C}[\ell^s, \ell^{-s}]$.*
 (b) *If φ_0 is the spherical vector (normalised so that $B_{\varphi_0, \lambda}(1) = 1$) then*

$$Z(\varphi_0, \eta, \lambda, s) = [L(\lambda_1 \eta, s + 1/2)L(\lambda_2 \eta, s + 1/2)]^{-1}.$$

- (c) *We have*

$$Z\left(\begin{pmatrix} ta & & & \\ & tb & w & \\ & & a & \\ & & & b \end{pmatrix} \varphi, \eta, \lambda, s\right) = \frac{\lambda_1(a)\lambda_2(b)}{\eta(t)|t|^{s-3/2}} Z(\varphi, \eta, \lambda, s)$$

for any $v, w \in \mathbb{Q}_\ell$ and $a, b, t \in \mathbb{Q}_\ell^\times$.

Proof. Replacing σ with $\sigma \otimes \eta$, and (λ_1, λ_2) with $(\eta\lambda_1, \eta\lambda_2)$, we may assume η is trivial. It suffices to prove (a) under the assumption that $\varphi = g \cdot \varphi_0$ for some $g \in G_\ell$ (since these vectors span σ). The validity of (a) for this vector will only depend on the class of g in the double coset space $R(\mathbb{Q}_\ell) \backslash G_\ell / K_{G_\ell}$. A set of coset representatives for this double quotient, and a formula for the values of $B_{\varphi_0, \lambda}$ on these representatives, is given in [40, Proposition 2-5]; see also [4, Corollary 1.9] for an alternative, slightly more concrete formulation. The result now follows by an explicit calculation, which also gives (b) as a special case (compare also [31, §3.2]).

Finally, part (c) is obvious from the integral formula if $\Re(s) \gg 0$, and follows for all s by analytic continuation. ■

From Proposition 3.6.4 above, we see that

$$Z(U(\ell)\varphi_0, \eta, \lambda, s) = \frac{\ell^{s+3/2}}{\eta(\ell)} [L(\lambda_1 \eta, s + 1/2)^{-1} L(\lambda_2 \eta, s + 1/2)^{-1} - L(\sigma \otimes \eta, s)^{-1}]. \quad (4)$$

This formula will be fundamental to the proof of our Euler system norm relations later in the paper.

Remark 3.6.7. It is not always true that the ideal of $\mathbb{C}[q^s, q^{-s}]$ given by $\{Z(\varphi, \eta, \lambda, s) : \varphi \in \sigma\}$ is the unit ideal. A sufficient condition is that $L(\lambda_1 \eta, s + 1/2)L(\lambda_2 \eta, s + 1/2)$ and $L(\sigma \otimes \eta, s)$ should have no poles in common, since then at least one of $Z(\varphi_0, \eta, \lambda, s)$ and $Z(U(\ell)\varphi_0, \eta, \lambda, s)$ is non-vanishing for every s . In fact, this condition is also necessary, as shown by the computations of [36], although we do not need this here. ◇

3.7. A local bilinear form

As in the preceding section, let σ be an irreducible unramified principal series representation of G_ℓ , with central character χ_σ . Let $\underline{\chi} = (\chi_1, \chi_2)$ and $\underline{\psi} = (\psi_1, \psi_2)$ be pairs of unramified characters of $\mathbb{Q}_\ell^\times$ satisfying

$$\chi_1 \chi_2 \cdot \psi_1 \psi_2 \cdot \chi_\sigma = 1,$$

and suppose that neither χ_1/ψ_1 nor χ_2/ψ_2 is quadratic or equal to $|\cdot|^{-1}$ (but we do allow either or both to equal $|\cdot|$). In the notation of the above section, we define a character λ of A by $\lambda_1 = (\psi_1\chi_2)^{-1}$, $\lambda_2 = (\chi_1\psi_2)^{-1}$; and we take for η the character $\psi_1\psi_2$. For brevity, we write $\underline{\chi}_s$ for the pair $(\chi_1|\cdot|^{-s}, \chi_2|\cdot|^{-s})$, and similarly $\underline{\psi}_s = (\psi_1|\cdot|^s, \psi_2|\cdot|^s)$.

Proposition 3.7.1. *Mapping $\varphi \in \sigma$ to the function $z_s(\varphi)$ on H_ℓ defined by*

$$z_s(\varphi)(h) = Z(h \cdot \varphi, \eta, \lambda, 2s + 1/2)$$

is an H_ℓ -equivariant map from σ to the space of polynomial sections of $I_H(\underline{\psi}_s^{-1}, \underline{\chi}_s^{-1})$. For the spherical vector φ_0 , normalised as in Proposition 3.6.6(b), we have

$$\begin{aligned} z_s(\varphi_0)(1) &= L(\psi_1/\chi_1, 2s + 1)^{-1} L(\psi_2/\chi_2, 2s + 1)^{-1}, \\ z_s(U(\ell)\varphi_0)(1) &= \frac{\ell^{2+2s}}{\psi_1\psi_2(\ell)} \\ &\quad \times \left[L\left(\frac{\psi_1}{\chi_1}, 2s + 1\right)^{-1} L\left(\frac{\psi_2}{\chi_2}, 2s + 1\right)^{-1} - L\left(\sigma \otimes \psi_1\psi_2, 2s + \frac{1}{2}\right)^{-1} \right]. \end{aligned}$$

Proof. Follows from Proposition 3.6.6 and equation (4). ■

We impose the following assumption:

- The functions $L(\sigma \otimes \psi_1\psi_2, s)$ and $L(\psi_1/\chi_1, s + 1/2)L(\psi_2/\chi_2, s + 1/2)$ do not both have a pole at $s = 1/2$.

It follows that the homomorphism $z \in \text{Hom}_{H_\ell}(\sigma, I_H(\underline{\psi}^{-1}, \underline{\chi}^{-1}))$ given by specialising z_s at $s = 0$ is not zero, since at least one of $z(\varphi_0)$ and $z(U(\ell)\varphi_0)$ is non-vanishing at $1 \in H_\ell$. Our assumptions on $\underline{\chi}$, $\underline{\psi}$ imply that, although $I_H(\underline{\psi}^{-1}, \underline{\chi}^{-1})$ may be reducible, it has a unique irreducible subrepresentation, and this subrepresentation is generic.

Lemma 3.7.2. *The image of the homomorphism z is contained in the unique irreducible subrepresentation of $I_H(\underline{\psi}^{-1}, \underline{\chi}^{-1})$.*

Proof. If $L(\psi_1/\chi_1, s + 1/2)L(\psi_2/\chi_2, s + 1/2)$ is finite at $s = 1/2$, then $I_H(\underline{\psi}^{-1}, \underline{\chi}^{-1})$ is irreducible and there is nothing to prove. So it suffices to treat the case when one or both of χ_i/ψ_i is $|\cdot|$, assuming that $L(\sigma \otimes \psi_1\psi_2, s)$ has no pole at $s = 1/2$. We shall not give the details of this computation, as it is somewhat technical, and it will only be relevant in a few boundary cases. We write σ as an induced representation from the Siegel parabolic $P_S(\mathbb{Q}_\ell)$. There are exactly two orbits of H_ℓ on the flag variety $G_\ell/P_S(\mathbb{Q}_\ell)$, and an application of Mackey theory allows us to compute $\text{Hom}_{H_\ell}(\sigma, \tau)$ for each non-generic quotient τ of $I_H(\underline{\psi}^{-1}, \underline{\chi}^{-1})$ in terms of the inducing data for σ . These Hom-spaces all turn out to be zero unless $L(\sigma \otimes \psi_1\psi_2, s)$ has a pole at $s = 1/2$. ■

Corollary 3.7.3. *Let $\langle \cdot, \cdot \rangle$ denote the canonical duality pairing*

$$I_H(\underline{\psi}_s, \underline{\chi}_s) \times I_H(\underline{\psi}_s^{-1}, \underline{\chi}_s^{-1}) \rightarrow \mathbb{C}.$$

Then the bilinear form $\mathfrak{z}_{\underline{\chi}, \underline{\psi}} \in \text{Hom}_{H_\ell}(I(\underline{\chi}, \underline{\psi}) \otimes \sigma, \mathbb{C})$ defined by

$$\mathfrak{z}_{\underline{\chi}, \underline{\psi}}(f \otimes \varphi) := \lim_{s \rightarrow 0} L(\psi_1/\chi_1, 2s + 1)L(\psi_2/\chi_2, 2s + 1)\langle M(f_s), z_s(\varphi) \rangle,$$

for f_s any polynomial section of $I(\underline{\chi}_s, \underline{\psi}_s)$ passing through f , is well-defined and non-zero.

Proof. We have $\langle M(f_s), z_s(\varphi) \rangle = \langle f_s, M(z_s(\varphi)) \rangle$ by Proposition 3.1.5. From the previous lemma, $M(z_s(\varphi))$ vanishes at $s = 0$ to order equal to the order of the pole of the Euler factor, so the limit is well-defined and depends only on f . ■

Remark 3.7.4. In the paper [29] (published in 1997, but circulated as a preprint many years before), Piatetski-Shapiro defines a zeta-integral $Z(\varphi, \underline{\phi}, \lambda, \eta, s)$, for $\varphi \in \sigma$ and $\underline{\phi} \in \mathcal{S}(\mathbb{Q}_\ell^2 \times \mathbb{Q}_\ell^2)$, by

$$Z(\varphi, \underline{\phi}, \lambda, \eta, s) := \int_{N_{H_\ell} \backslash H_\ell} B_{\varphi, \lambda}(h) \underline{\phi}((0, 1) \cdot h_1, (0, 1) \cdot h_2) \eta(\det h) |\det h|^{s+1/2} dh,$$

where N_{H_ℓ} is the unipotent radical of $B_H(\mathbb{Q}_\ell)$. This integral also appears in Sugano's work [40]. If η and λ are chosen as above, then one checks that $\delta_{\underline{\chi}, \underline{\psi}}(f_{\hat{\underline{\phi}}, \underline{\chi}, \underline{\psi}} \otimes \varphi)$ is equal to the leading term of $Z(\varphi, \underline{\phi}, \Lambda, \eta, s)$ at $s = 1/2$, up to a non-zero scalar factor. However, we cannot simply take this as the definition of $\delta_{\underline{\chi}, \underline{\psi}}$, since it is not *a priori* clear that this leading term depends only on the vector $f_{\hat{\underline{\phi}}, \underline{\chi}, \underline{\psi}} \in I_H(\underline{\chi}, \underline{\psi})$ rather than on $\underline{\phi}$ itself. ◊

Vitally, the linear functional $\delta_{\underline{\chi}, \underline{\psi}}$ of Proposition 3.7.3 is *unique*:

Theorem 3.7.5 (Kato–Murase–Sugano). *For $\underline{\chi}$ and $\underline{\psi}$ satisfying the above assumptions, we have*

$$\dim \mathrm{Hom}_{H_\ell}(I_H(\underline{\chi}, \underline{\psi}) \otimes \sigma, \mathbb{C}) \leq 1.$$

Proof. Let τ be the representation $I_H(\underline{\chi}, \underline{\psi})$. Since the group of unramified characters of $\mathbb{Q}_\ell^\times$ is 2-divisible, we may replace σ and τ with $\sigma \otimes \omega^{-1}$ and $\tau \otimes \omega$, where ω is any square root of χ_σ , and therefore assume that both σ and τ are trivial on $Z(G_\ell)$. Thus σ factors through $\overline{G} = \mathrm{PGSp}_4(\mathbb{Q}_\ell) = \mathrm{SO}_5(\mathbb{Q}_\ell)$; and τ factors through the image $\overline{H}$ of H in $\mathrm{SO}_5(\mathbb{Q}_\ell)$, which is a copy of $\mathrm{SO}_4(\mathbb{Q}_\ell)$, embedded as the stabiliser of an anisotropic vector in the defining 5-dimensional representation.

We now apply the main theorem of [13], which shows that for any representations σ of SO_5 and τ of SO_4 which are generated by a spherical vector, the space $\mathrm{Hom}(\tau \otimes \sigma, \mathbb{C})$ has dimension ≤ 1 . ■

Remark 3.7.6. Alternatively, the proof of Lemma 3.7.2 shows that this Hom-space injects into $\mathrm{Hom}_{H_\ell}(\sigma \otimes \tau_0, \mathbb{C})$ where τ_0 is the unique irreducible subrepresentation of $I_H(\underline{\chi}, \underline{\psi})$. We can now invoke a very general result, which forms part of the Gan–Gross–Prasad conjecture for special orthogonal groups: for any $n \geq 0$ and any irreducible smooth representations σ of $\mathrm{SO}_{n+1}(\mathbb{Q}_\ell)$ and ρ of $\mathrm{SO}_n(\mathbb{Q}_\ell)$, one has $\dim \mathrm{Hom}_{\mathrm{SO}_n}(\sigma \otimes \rho, \mathbb{C}) \leq 1$, by [45, Théorème 1]. ◊

Corollary 3.7.7. *In the situation of Theorem 3.7.5, the bilinear form $\delta_{\underline{\chi}, \underline{\psi}}$ is a basis of $\mathrm{Hom}_{H_\ell}(I_H(\underline{\chi}, \underline{\psi}) \otimes \sigma, \mathbb{C})$.*

Proof. Clear. ■

3.8. Explicit formulae for the unramified local pairing

We record the following formulae for the values of $\mathfrak{z}_{\underline{\chi}, \underline{\psi}}$. We assume, as before, that σ is an irreducible unramified principal series representation of G_ℓ . We choose our characters $\underline{\chi}$ and $\underline{\psi}$ as follows:

- $\psi_1 = \psi_2 = |\cdot|^{-1/2}$,
- $\chi_i = |\cdot|^{(1/2+k_i)} \tau_i$, where $\underline{\tau} = (\tau_1, \tau_2)$ is a pair of finite-order unramified characters, and $k_i \geq 0$ are integers.

If one or both of the k_i is zero, we also assume that σ is essentially tempered (a twist of a tempered representation); since χ_σ is $|\cdot|^{-(k_1+k_2)}$ up to a finite-order character, all poles of $L(\sigma, s)$ therefore have real part $(k_1 + k_2)/2 \geq 0$, so that $L(\sigma \otimes \psi_1 \psi_2, 1/2) = L(\sigma, -1/2)$ is finite and the assumptions of the previous section are satisfied.

For $\underline{\phi} \in \mathcal{S}(\mathbb{Q}_\ell^2)^{\otimes 2}$, we write $F_{\underline{\phi}}$ for the Siegel section $f_{\hat{\underline{\phi}}, \underline{\chi}, \underline{\psi}} \in I_H(\underline{\chi}, \underline{\psi})$; from (1), this depends H_ℓ -equivariantly on $\underline{\phi}$. We shall apply this to the particular Schwartz functions ϕ_t introduced in §3.4 above.

Theorem 3.8.1. *Let $\mathfrak{z} \in \text{Hom}_{H_\ell}(I(\underline{\chi}, \underline{\psi}) \otimes \sigma, \mathbb{C})$. Then, for any $t \geq 1$, we have*

$$\mathfrak{z}(F_{\underline{\phi}_t}, \varphi_0) = \frac{1}{\ell^{2t-2}(\ell+1)^2} \left(1 - \frac{\ell^{k_1}}{\tau_1(\ell)}\right) \left(1 - \frac{\ell^{k_2}}{\tau_2(\ell)}\right) \cdot \mathfrak{z}(F_{\underline{\phi}_0}, \varphi_0)$$

and

$$\mathfrak{z}(F_{\underline{\phi}_1}, U(\ell)\varphi_0) = \frac{\ell}{(\ell+1)^2} \left[\left(1 - \frac{\ell^{k_1}}{\tau_1(\ell)}\right) \left(1 - \frac{\ell^{k_2}}{\tau_2(\ell)}\right) - L(\sigma, -1/2)^{-1} \right] \mathfrak{z}(F_{\underline{\phi}_0}, \varphi_0).$$

Proof. We know that $\text{Hom}_{H_\ell}(I(\underline{\chi}, \underline{\psi}) \otimes \sigma, \mathbb{C})$ is 1-dimensional and spanned by the specific bilinear form $\mathfrak{z}_{\underline{\chi}, \underline{\psi}}$ constructed above, so it suffices to assume that $\mathfrak{z} = \mathfrak{z}_{\underline{\chi}, \underline{\psi}}$. By construction $F_{\underline{\phi}_t}$ is the value at $s = 0$ of the Siegel section $f_{\hat{\underline{\phi}}, \underline{\chi}_s, \underline{\psi}_s}$, and we have

$$M(f_{\hat{\underline{\phi}}, \underline{\chi}_s, \underline{\psi}_s}) = L(\chi_1/\psi_1, 1-2s)^{-1} L(\chi_2/\psi_2, 1-2s)^{-1} f_{\hat{\underline{\phi}}, \underline{\psi}_s, \underline{\chi}_s},$$

by the functional equation for Siegel sections (Proposition 3.2.3). As we have seen above, the restriction of $f_{\hat{\underline{\phi}}, \underline{\psi}_s, \underline{\chi}_s}$ to $H(\hat{\mathbb{Z}})$ is a scalar multiple of the characteristic function of $K_{H_\ell, 0}(\ell^t)$, so we have

$$\begin{aligned} \mathfrak{z}_{\underline{\chi}, \underline{\psi}}(F_{\underline{\phi}_t}, \varphi) &= \frac{\text{Vol } K_{H_\ell, 0}(\ell^t) f_{\hat{\underline{\phi}}, \underline{\psi}, \underline{\chi}}(1)}{L(\chi_1/\psi_1, 1) L(\chi_2/\psi_2, 1)} \\ &\quad \times \lim_{s \rightarrow 0} [L(\psi_1/\chi_1, 1+2s) L(\psi_2/\chi_2, 1+2s) z_s(\varphi)(1)] \end{aligned}$$

for any $\varphi \in \sigma$ invariant under $K_{H_\ell, 0}(\ell^t)$. In particular, if $\varphi = \varphi_0$ then the bracketed term is identically 1, and from the formula for $f_{\hat{\underline{\phi}}, \underline{\chi}, \underline{\psi}}(1)$ given in Lemma 3.2.5 we see that for $t \geq 1$ we have

$$\mathfrak{z}_{\underline{\chi}, \underline{\psi}}(F_{\underline{\phi}_t}, \varphi_0) = \text{Vol } K_{H_\ell, 0}(\ell^t) \cdot L(\psi_1/\chi_1, 1)^{-1} L(\psi_2/\chi_2, 1)^{-1} \mathfrak{z}_{\underline{\chi}, \underline{\psi}, s}(F_{\underline{\phi}_0}, \varphi_0),$$

which is the first formula claimed. The second formula is similar, using the formula for $z_s(U(\ell)\varphi_0)(1)$ given in Proposition 3.7.1. $\blacksquare$

3.9. An application of Frobenius reciprocity

Proposition 3.9.1. *Let τ (resp. σ) be smooth representations of H_ℓ and G_ℓ respectively. Then there are canonical bijections of $\mathbb{C}$ -vector spaces*

$$\mathrm{Hom}_{G_\ell}(\mathrm{c}\text{-}\mathrm{Ind}_{H_\ell}^{G_\ell}(\tau), \sigma^\vee) \cong \mathrm{Hom}_{G_\ell}(\mathrm{c}\text{-}\mathrm{Ind}_{H_\ell}^{G_\ell}(\tau) \otimes \sigma, \mathbb{C}) \cong \mathrm{Hom}_{H_\ell}(\tau \otimes (\sigma|_{H_\ell}), \mathbb{C}).$$

Proof. The first isomorphism is standard, and interchanging the roles of σ and the compactly induced representation also shows that

$$\mathrm{Hom}_{G_\ell}(\mathrm{c}\text{-}\mathrm{Ind}_{H_\ell}^{G_\ell}(\tau) \otimes \sigma, \mathbb{C}) \cong \mathrm{Hom}_{G_\ell}(\sigma, (\mathrm{c}\text{-}\mathrm{Ind}_{H_\ell}^{G_\ell} \tau)^\vee).$$

One has a canonical isomorphism $(\mathrm{c}\text{-}\mathrm{Ind}_{H_\ell}^{G_\ell} \tau)^\vee = \mathrm{Ind}_{H_\ell}^{G_\ell}(\tau^\vee)$ [33, §III.2.7]. (Care must be taken here since the contragredient on the left-hand side denotes G_ℓ -smooth vectors in the abstract vector-space dual, while on the right-hand side it denotes H_ℓ -smooth vectors.) We then apply Frobenius reciprocity for the non-compact induction [33, §III.2.5] to obtain

$$\begin{aligned} \mathrm{Hom}_{G_\ell}(\sigma, \mathrm{Ind}_{H_\ell}^{G_\ell}(\tau^\vee)) &\cong \mathrm{Hom}_{H_\ell}((\sigma|_{H_\ell}), \tau^\vee) \\ &\cong \mathrm{Hom}_{H_\ell}(\tau \otimes (\sigma|_{H_\ell}), \mathbb{C}), \end{aligned}$$

as required. ■

Remark 3.9.2. The Hom-spaces in Proposition 3.9.1 will *not* in general be isomorphic to $\mathrm{Hom}_{H_\ell}(\tau, (\sigma^\vee)|_{H_\ell})$. The problem is that $(\sigma^\vee)|_{H_\ell}$ is in general much smaller than $(\sigma|_{H_\ell})^\vee$, since the two notions of contragredient do not match – an H_ℓ -smooth linear functional on σ may not be G_ℓ -smooth. ◇

For later use it will be important to have an explicit form for this bijection. Let $\mathcal{H}(G_\ell)$ denote the Hecke algebra of locally constant, compactly supported $\mathbb{C}$ -valued functions on G_ℓ , with the algebra structure defined by convolution (normalising Haar measure as in §3.3). We regard σ as a left $\mathcal{H}(G_\ell)$ -module via the usual formula

$$\xi \cdot \varphi = \int_{G_\ell} \xi(g)(g \cdot \varphi) \, dg,$$

so that $g_1 \cdot (\xi \cdot (g_2 \cdot \varphi)) = \xi(g_1^{-1}(-)g_2^{-1}) \cdot \varphi$.

Definition 3.9.3. For smooth representations τ, σ as above, let $\mathfrak{X}(\tau, \sigma^\vee)$ denote the space of linear maps

$$\mathfrak{Z} : \tau \otimes_{\mathbb{C}} \mathcal{H}(G_\ell) \rightarrow \sigma^\vee$$

which are $H_\ell \times G_\ell$ -equivariant, where the actions are defined as follows:

- The H_ℓ factor acts trivially on $\sigma^\vee$, and on $\tau \otimes_{\mathbb{C}} \mathcal{H}(G_\ell)$ it acts via the formula

$$h \cdot (v \otimes \xi) = (h \cdot v) \otimes \xi(h^{-1}(-)).$$

- The G_ℓ factor acts trivially on τ , and on $\mathcal{H}(G_\ell)$ it acts via $g \cdot \xi = \xi((-)g)$.

Unwinding the definitions, we reach the following formula:

Proposition 3.9.4. *There is a canonical bijection $\mathfrak{X}(\tau, \sigma^\vee) \cong \text{Hom}_{H_\ell}(\tau \otimes (\sigma|_{H_\ell}), \mathbb{C})$, characterised as follows: if $\mathfrak{Z} \in \mathfrak{X}(\tau, \sigma^\vee)$ corresponds to $\mathfrak{z} \in \text{Hom}_{H_\ell}(\tau \otimes (\sigma|_{H_\ell}), \mathbb{C})$, then*

$$\mathfrak{Z}(f \otimes \xi)(\varphi) = \mathfrak{z}(f \otimes (\xi \cdot \varphi))$$

for all $f \in \tau$, $\xi \in \mathcal{H}(G_\ell)$, and $\varphi \in \sigma$.

Proof. Immediate. ■

Corollary 3.9.5. *Suppose $\mathfrak{z} \leftrightarrow \mathfrak{Z}$ as in the above proposition; and let $U_0 \geq U_1$ be two open compact subgroups of G_ℓ , $f_0, f_1 \in \tau$, and $g_0, g_1 \in G_\ell$. Suppose that*

$$\mathfrak{z}(f_1, g_1 \cdot \varphi) = \mathfrak{z}(f_0, g_0 \cdot R \cdot \varphi)$$

for some $R \in \mathcal{H}(U_0 \backslash G_\ell / U_0)$ and all $\varphi \in \sigma^{U_0}$. Then the elements $\mathfrak{Z}_i = \mathfrak{Z}(f_i \otimes \text{ch}(g_i U_i)) \in (\sigma^\vee)^{U_i}$, $i = 0, 1$, are related by

$$\sum_{u \in U_0 / U_1} u \cdot \mathfrak{Z}_1 = R' \cdot \mathfrak{Z}_0$$

as elements of $(\sigma^\vee)^{U_0}$, where $R'(g) = R(g^{-1})$.

Proof. Since both sides of the desired equality are in $(\sigma^\vee)^{U_0} = (\sigma^{U_0})^\vee$, it suffices to check that they pair to the same value with φ for every $\varphi \in \sigma^{U_0}$. This follows from the above description of $\mathfrak{Z}(-)(\varphi)$. ■

3.10. Results for deeper levels

In order to prove norm-compatibility relations in the “ p -direction” for our Euler system, we shall also need a few supplementary results which are proved directly (rather than using the uniqueness result of Theorem 3.7.5). In this section, W denotes an arbitrary smooth complex representation of G_ℓ (not necessarily irreducible or even admissible), and we let $\mathfrak{X}(W)$ denote the space of homomorphisms

$$\mathfrak{Z} : \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})^{\otimes 2} \otimes_{\mathbb{C}} \mathcal{H}(G_\ell) \rightarrow W$$

satisfying the same equivariance property under $H_\ell \times G_\ell$ as in Definition 3.9.3.

Notation 3.10.1. For $t \geq 1$, let $\phi_{1,t} \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})$ denote the characteristic function of the set $\ell^t \mathbb{Z}_\ell \times (1 + \ell^t \mathbb{Z}_\ell)$, and $\underline{\phi}_{1,t} = \phi_{1,t} \otimes \phi_{1,t} \in \mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})^{\otimes 2}$. This is stable under the group $K_{H_\ell, 1}(\ell^t)$ (see §3.3).

Lemma 3.10.2. *Let $\xi \in \mathcal{H}(G_\ell)$ be invariant under left-translation by the principal congruence subgroup of level ℓ^T in $H(\mathbb{Z}_\ell)$, for some $T \geq 1$. Then, for any $\mathfrak{Z} \in \mathfrak{X}(W)$, the expression*

$$\frac{1}{\text{Vol } K_{H_\ell, 1}(\ell^t)} \mathfrak{Z}(\underline{\phi}_{1,t} \otimes \xi)$$

is independent of $t \geq T$, where $\text{Vol}(-)$ denotes volume with respect to our fixed Haar measure on H_ℓ .

Proof. For any integers $t \geq T \geq 1$, let J be a set of coset representatives for the quotient $K_{H_\ell,1}(\ell^T)/K_{H_\ell,1}(\ell^t)$. Then $\phi_{1,T} = \sum_{\gamma \in J} \gamma \cdot \phi_{1,t}$, so we have

$$\mathfrak{Z}(\phi_{1,T} \otimes \xi) = \sum_{\gamma \in J} \mathfrak{Z}((\gamma \cdot \phi_{1,t}) \otimes \xi) = \sum_{\gamma \in J} \mathfrak{Z}(\phi_{1,t} \otimes \xi(\gamma(-))).$$

We can (and do) assume that J is a subset of the principal congruence subgroup of level ℓ^T in H . By assumption, all such elements will act trivially on ξ from the left, so the above equality becomes

$$\mathfrak{Z}(\phi_{1,T} \otimes \xi) = (\#J) \cdot \mathfrak{Z}(\phi_{1,t} \otimes \xi) = \frac{\mathrm{Vol} K_{H_\ell,1}(\ell^T)}{\mathrm{Vol} K_{H_\ell,1}(\ell^t)} \mathfrak{Z}(\phi_{1,t} \otimes \xi)$$

as required. ■

Notation 3.10.3. We write $\mathfrak{Z}(\phi_{1,\infty} \otimes \xi)$ for this limiting value.

The case that interests us is the following. Let m, n be integers with $m \geq 0$ and $n \geq \max(m, 1)$, and let $\eta_m = \begin{pmatrix} 1 & \ell^{-m} & & \\ & 1 & \ell^{-m} & \\ & & 1 & \\ & & & 1 \end{pmatrix}$. Recall the subgroup $K_{G_\ell}(\ell^m, \ell^n)$ of §3.3; we consider the Hecke operator

$$U'(\ell) = \frac{1}{\mathrm{Vol} K_{G_\ell}(\ell^m, \ell^n)} \mathrm{ch} \left(K_{G_\ell}(\ell^m, \ell^n) \begin{pmatrix} \ell^{-1} & & & \\ & \ell^{-1} & & \\ & & 1 & \\ & & & 1 \end{pmatrix} K_{G_\ell}(\ell^m, \ell^n) \right).$$

Proposition 3.10.4. For any $\mathfrak{Z} \in \mathfrak{X}(W)$ we have

$$\begin{aligned} & \mathfrak{Z}(\phi_{1,\infty} \otimes \mathrm{ch}(\eta_{m+1} K_{G_\ell}(\ell^m, \ell^n))) \\ &= \begin{cases} \frac{1}{\ell} U'(\ell) & \text{if } m \geq 1 \\ \frac{1}{\ell-1} [U'(\ell) - 1] & \text{if } m = 0 \end{cases} \cdot \mathfrak{Z}(\phi_{1,\infty} \otimes \mathrm{ch}(\eta_m K_{G_\ell}(\ell^m, \ell^n))). \end{aligned}$$

Proof. Writing $K = K_{G_\ell}(\ell^m, \ell^n)$ for brevity, we have

$$\begin{aligned} U'(\ell) \cdot \mathfrak{Z}(\phi_{1,\infty} \otimes \mathrm{ch}(\eta_m K)) &= \sum_{u,v,w \in \mathbb{Z}/\ell} \mathfrak{Z} \left(\phi_{1,\infty} \otimes \mathrm{ch} \left(\eta_m \begin{pmatrix} \ell & u & v \\ & \ell & w \\ & & 1 & u \\ & & & 1 \end{pmatrix} K_{m,n} \right) \right) \\ &= \frac{1}{\mathrm{Vol} K_{H_\ell,1}(\ell^n)} \sum_{u,v,w} \mathfrak{Z} \left(\left(\begin{pmatrix} \ell & v \\ & \ell & w \\ & & 1 \end{pmatrix} \right)^{-1} \phi_{1,n} \otimes \mathrm{ch}(\eta_{m+1}^{(1+\ell^m u)} K) \right) \\ &= \frac{\ell^2}{\mathrm{Vol} K_{H_\ell,1}(\ell^n)} \sum_u \mathfrak{Z}(\mathrm{ch}(\ell^{n+1} \mathbb{Z}_\ell \times (1 + \ell^n \mathbb{Z}_\ell))^{\otimes 2} \otimes \mathrm{ch}(\eta_{m+1}^{(1+\ell^m u)} K)) \\ &= \sum_u \mathfrak{Z}(\phi_{1,\infty} \otimes \mathrm{ch}(\eta_{m+1}^{(1+\ell^m u)} K)). \end{aligned}$$

There are now two cases to consider. If $m \geq 1$ then all terms in this sum are actually equal, since the powers of η_{m+1} are conjugate via elements of the form $\begin{pmatrix} a & & \\ & a & \\ & & 1 & \\ & & & 1 \end{pmatrix}$ (with $a \in 1 + \ell^m \mathbb{Z}_\ell$), which are in K and act trivially on the Schwartz function $\phi_{1,n}$; so the

sum is simply $\ell \mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(\eta_m K))$ as required. If $m = 0$, then $\ell - 1$ of the terms are conjugate, but the term for $u = -1$ requires special consideration since $1 + \ell^m u = 0$; thus we obtain

$$[U'(\ell) - 1] \cdot \mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(K)) = (\ell - 1) \mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(\eta_1 K)),$$

which proves the formula in this case. $\blacksquare$

We also have an analogous result for $n = 0$, under rather stricter hypotheses. We take for W the smooth dual $\sigma^\vee$ of an essentially tempered, unramified principal series representation of G_ℓ . We shall suppose that $\mathfrak{Z} \in \mathfrak{X}(\sigma^\vee)$ factors through a certain induced representation of H : more precisely, we shall take pairs of characters $\underline{\chi} = (\chi_1, \chi_2)$ and $\underline{\psi} = (\psi_1, \psi_2)$ of $\mathbb{Q}_\ell^\times$ with $\psi_1 = \psi_2 = |\cdot|^{-1/2}$ and $\chi_i = |\cdot|^{k_i+1/2} \tau_i$ for finite-order characters τ_i and positive integers k_i , so our setup is similar to Theorem 3.8.1 except that we do *not* assume that the τ_i are unramified. We then have a natural map

$$\mathcal{S}(\mathbb{Q}_\ell^2, \mathbb{C})^{\otimes 2} \rightarrow I_H(\underline{\chi}, \underline{\psi}), \quad \underline{\phi} \mapsto F_{\underline{\phi}},$$

and we suppose that $\mathfrak{Z}$ factors through this map.

Corollary 3.10.5. *In this situation, we have*

$$\mathfrak{Z}(\phi_{1,\infty} \otimes (\text{ch}(K_{G_\ell}) - \text{ch}(\eta_1 K_{G_\ell}))) = \frac{\ell}{\ell-1} L(\sigma, -1/2)^{-1} \cdot \mathfrak{Z}(\phi_0, \text{ch}(K_{G_\ell})).$$

In particular, if the τ_i are not both unramified, then this holds vacuously (both sides of the formula are zero).

Proof. Since $K_{G_\ell,0}(\ell)$ fixes $\text{ch}(K_{G_\ell})$ on the left, we have

$$\mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(K_{G_\ell})) = \frac{1}{\text{Vol } K_{H_\ell,0}(\ell)} \mathfrak{Z}(\phi_1 \otimes \text{ch}(K_{G_\ell})) = (\ell + 1)^2 \mathfrak{Z}(\phi_1 \otimes \text{ch}(K_{G_\ell})).$$

where (as in §3.4) $\phi_1 = \text{ch}(\ell \mathbb{Z}_\ell \times \mathbb{Z}_\ell^\times)^{\otimes 2}$. On the other hand, taking $m = 0$ and $n = 1$ in Proposition 3.10.4, we have

$$\mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(\eta_1 K_{G_{\ell,1}}(\ell))) = \frac{U'(\ell) - 1}{\ell - 1} \mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(K_{G_{\ell,1}}(\ell))).$$

Since the action of the quotient $K_{G_\ell,0}(\ell)/K_{G_{\ell,1}}(\ell) \cong \text{GL}_2(\mathbb{Z}/\ell)$ commutes with the Hecke operator $U'(\ell)$, we can sum over representatives for the quotient to deduce that

$$\mathfrak{Z}(\phi_{1,\infty} \otimes \text{ch}(\eta_1 K_{G_\ell})) = \frac{(\ell + 1)^2}{\ell - 1} \sum_{\gamma \in K_{G_\ell}/K_{G_{\ell,0}}(\ell)} \gamma \cdot (U'(\ell) - 1) \mathfrak{Z}(\phi_1 \otimes \text{ch}(K_{G_{\ell,0}}(\ell))).$$

Combining these two formulae we have

$$\begin{aligned} \mathfrak{Z}(\phi_{1,\infty} \otimes (\text{ch}(K_{G_\ell}) - \text{ch}(\eta_1 K_{G_\ell}))) &= (\ell + 1)^2 \left(1 + \frac{1}{\ell-1}\right) \mathfrak{Z}(\phi_1 \otimes \text{ch}(K_{G_\ell})) \\ &\quad - \frac{(\ell+1)^2}{\ell-1} \sum_{\gamma \in K/K_{G_{\ell,0}}(\ell)} \gamma U'(\ell) \cdot \mathfrak{Z}(\phi_1 \otimes \text{ch}(K_{G_{\ell,0}}(\ell))). \end{aligned}$$

We can now quickly dispose of the ramified cases. The map $\phi \mapsto F_\phi$ is a morphism of $(\mathrm{GL}_2 \times \mathrm{GL}_2)$ -representations (not only of H -representations). Moreover, the elements ϕ_0 and ϕ_1 are the characteristic functions of subsets of $\mathbb{Q}_\ell^2 \times \mathbb{Q}_\ell^2$ invariant under $\mathbb{Z}_\ell^\times \times \mathbb{Z}_\ell^\times$; hence their images in any representation of $\mathrm{GL}_2 \times \mathrm{GL}_2$ with ramified central character must be zero. Hence F_{ϕ_0} and F_{ϕ_1} are both zero, and the desired formula becomes $0 = 0$, if either of the characters τ_i is ramified.

Let us now assume that the τ_i are unramified, which means we may apply Theorem 3.8.1. Translating to the homomorphism $\mathfrak{Z}$ from its corresponding bilinear form $\mathfrak{z}$, the first statement in the theorem (for $t = 1$) becomes

$$(\ell + 1)^2 \left(1 + \frac{1}{\ell - 1}\right) \mathfrak{Z}(\phi_1 \otimes \mathrm{ch}(K_{G_\ell})) = \frac{\ell}{\ell - 1} \left(1 - \frac{\ell^{k_1}}{\tau_1(\ell)}\right) \left(1 - \frac{\ell^{k_2}}{\tau_2(\ell)}\right) \cdot \mathfrak{Z}(\phi_0 \otimes \mathrm{ch}(K_{G_\ell})).$$

On the other hand, the second statement of Theorem 3.8.1 gives us

$$\begin{aligned} \frac{(\ell + 1)^2}{\ell - 1} \sum_{\gamma \in K_{G_\ell} / K_{G_\ell, 0}(\ell)} \gamma U'(\ell) \cdot \mathfrak{Z}(\phi_1 \otimes \mathrm{ch}(K_{G_\ell, 0}(\ell))) \\ = \frac{\ell}{\ell - 1} \left[\left(1 - \frac{\ell^{k_1}}{\tau_1(\ell)}\right) \left(1 - \frac{\ell^{k_2}}{\tau_2(\ell)}\right) - L(\sigma, -1/2)^{-1} \right] \mathfrak{Z}(\phi_0 \otimes \mathrm{ch}(K_{G_\ell})). \end{aligned}$$

Combining these two formulae, the “extra” Euler factors coming from the τ_i cancel out, and we are left with the desired formula. ■

4. Preliminaries II: Algebraic representations and Lie theory

4.1. Representations of G

We recall the parametrisation of algebraic representations of the group GSp_4 .

Notation 4.1.1. We write T for the diagonal torus of G (as in §2 above), and we write $\chi_1, \dots, \chi_4$ for characters of T given by projection onto the four entries. Thus $\chi_1 + \chi_4 = \chi_2 + \chi_3$ is the restriction to T of the symplectic multiplier μ , and $\{\chi_1, \chi_2, \mu\}$ is a basis of the character group $X^\bullet(T)$.

Definition 4.1.2. Let $a, b \geq 0$ be integers. We denote by $V^{a,b}$ the unique (up to isomorphism) irreducible algebraic representation of G whose highest weight, with respect to B , is the character $(a + b)\chi_1 + a\chi_2$.

This representation has dimension $\frac{1}{6}(a + 1)(b + 1)(a + b + 2)(2a + b + 3)$. Its central character is $x \mapsto x^{2a+b}$, and it satisfies

$$(V^{a,b})^* \cong V^{a,b} \otimes \mu^{-(2a+b)}.$$

Note that $V^{0,1}$ is the 4-dimensional defining representation of GSp_4 , and $V^{1,0}$ is the 5-dimensional direct summand of $\bigwedge^2 V^{0,1}$. The representation $V^{1,0} \otimes \mu^{-1}$ has trivial central character, and is the defining representation of $G/Z_G \cong \mathrm{SO}_5$.

4.2. Integral models

Let

$$\lambda = (a + b)\chi_1 + a\chi_2 + c\mu, \quad \text{with } a, b \geq 0,$$

be a dominant integral weight, V_λ the corresponding representation, and v_λ a highest weight vector in V_λ . The pair (V_λ, v_λ) is then unique up to unique isomorphism.

An *admissible lattice* in V_λ is a $\mathbb{Z}$ -lattice L with the following properties:

- the homomorphism $\mathrm{GSp}_4 \rightarrow \mathrm{GL}(V_\lambda)$ extends to a homomorphism $\mathrm{GSp}_4 \rightarrow \mathrm{GL}(L)$ of group schemes over $\mathbb{Z}$;
- the intersection of L with the highest weight space of V_λ is $\mathbb{Z} \cdot v_\lambda$.

It is known that there are finitely many such lattices, each of which is the direct sum of its intersections with the weight spaces; and we set $V_{\lambda, \mathbb{Z}}$ to be the *maximal* such lattice.

Proposition 4.2.1. *Let λ, λ' be dominant integral weights. Then there is a unique G -equivariant homomorphism, the Cartan product,*

$$V_{\lambda, \mathbb{Z}} \otimes V_{\lambda', \mathbb{Z}} \rightarrow V_{\lambda + \lambda', \mathbb{Z}}, \quad v \otimes w \mapsto v \cdot w,$$

such that $v_\lambda \cdot v_{\lambda'} = v_{\lambda + \lambda'}$. Moreover, for any non-zero $v \in V_\lambda$ and $v' \in V_{\lambda'}$, we have $v \cdot v' \neq 0$.

Proof. After tensoring with $\mathbb{Q}$ the existence and uniqueness of this homomorphism is obvious from highest-weight theory. Hence the image of $V_{\lambda, \mathbb{Z}} \otimes V_{\lambda', \mathbb{Z}}$ is a $\mathbb{Z}$ -lattice in $V_{\lambda + \lambda'}$, which is clearly admissible; so it must be contained in the maximal one, which is $V_{\lambda + \lambda', \mathbb{Z}}$.

This product gives the ring $\bigoplus_\lambda V_{\lambda, \mathbb{Z}}$ the structure of a graded ring. The Borel–Weil theorem shows that this ring injects into $\mathcal{O}(G)$, which is an integral domain; so the Cartan product of non-zero vectors is non-zero. $\blacksquare$

4.3. Branching laws

We are interested in the restriction of $V^{a,b}$ to H via the embedding $\iota : H \hookrightarrow G$, which we shall denote by $\iota^*(V^{a,b})$. Computing the weights of these representations (and their multiplicities), one deduces the following branching law describing $\iota^*(V^{a,b})$:

Proposition 4.3.1. *The restriction of $V^{a,b}$ to $H = \mathrm{GL}_2 \times_{\mathrm{GL}_1} \mathrm{GL}_2$ via ι is given by*

$$\iota^*(V^{a,b}) = \bigoplus_{0 \leq q \leq a} \bigoplus_{0 \leq r \leq b} W^{a+b-q-r, a-q+r} \otimes \det^q,$$

where $W^{c,d}$ denotes the representation $\mathrm{Sym}^c \boxtimes \mathrm{Sym}^d$ of H .

Remark 4.3.2. Compare [22, §1] for an equivalent, although less explicit, statement. In Lemma’s notations the highest weight of our representation $V^{a,b}$ is $\lambda(a + b, a, 2a + b)$. $\diamond$

For the constructions below it will be useful to fix choices of highest-weight vectors in each of these subrepresentations. For $0 \leq q \leq a$ and $0 \leq r \leq b$ we define a vector $v^{a,b,q,r} \in V_{\mathbb{Z}}^{a,b}$ as follows:

$$v^{a,b,q,r} = w^{a-q} \cdot v^{b-r} \cdot (w')^q \cdot (v')^r$$

where the product operation is the Cartan product, and:

- $v \in V^{0,1}$ is the highest-weight vector;
- $v' = X_{21} \cdot v$ is a basis of the χ_2 weight space, where $X_{21} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix} \in \mathrm{Lie} G$.
- w is the highest-weight vector of $V^{1,0}$.
- $w' = Z \cdot w$ is a basis of the μ weight space of $V^{1,0}$, where $Z = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \in \mathrm{Lie} G$.

Remark 4.3.3. We can identify $V^{0,1}$ with the standard representation of $\mathrm{GSp}_4 \subseteq \mathrm{GL}_4$, with basis $(e_1, \dots, e_4)$, by choosing the highest-weight vector $v = e_1$; of course we then have $v' = e_2$. Moreover, we can identify $V^{1,0}$ with a subspace of $\bigwedge^2 V^{0,1}$, by choosing $e_1 \wedge e_2$ for the highest-weight vector w ; and it follows that w' is the vector $e_1 \wedge e_4 - e_2 \wedge e_3$. $\diamond$

Proposition 4.3.4. *For all integers $0 \leq q \leq a$ and $0 \leq r \leq b$, the vector $v^{a,b,q,r}$ thus defined is a non-zero highest-weight vector for the unique irreducible H -summand of $\iota^*(V^{a,b})$ isomorphic to $W^{a+b-q-r, a-q+r} \otimes \det^q$.*

Proof. Since $v^{a,b,q,r}$ is a Cartan product of non-zero H -highest weight vectors (i.e. vectors fixed by the action of the unipotent radical of the Borel of H), it is itself a non-zero H -highest weight vector, and thus generates an irreducible H -subrepresentation of $V^{a,b}$. The result now follows by comparing weights. $\blacksquare$

Since the representation $W^{c,d}$ of H has a canonical highest-weight vector (namely $e_1^c \boxtimes f_1^d$, where (e_1, e_2) and (f_1, f_2) are bases of the standard representations of the two GL_2 factors), we therefore have a canonical homomorphism of H -representations

$$\mathrm{br}^{[a,b,q,r]} : W^{a+b-q-r, a-q+r} \otimes \det^q \hookrightarrow \iota^*(V^{a,b}) \quad (5)$$

mapping the highest-weight vector to $v^{a,b,q,r}$. We refer to these maps as *branching maps*.

Proposition 4.3.5. *The maps $\mathrm{br}^{[a,b,q,r]}$ restrict to maps*

$$W_{\mathbb{Z}}^{a+b-q-r, a-q+r} \otimes \det^q \hookrightarrow V_{\mathbb{Z}}^{a,b},$$

where $W_{\mathbb{Z}}^{a+b-q-r, a-q+r}$ is the minimal¹ admissible lattice in $W^{a+b-q-r, a-q+r}$.

¹Note that the minimal admissible lattice in the representation Sym^k of GL_2 is isomorphic to the module $\mathrm{TSym}^k \mathbb{Z}^2$ of symmetric tensors, while $\mathrm{Sym}^k \mathbb{Z}^2$ is the maximal lattice.

Proof. It is clear that $(\mathrm{br}^{[a,b,q,r]})^{-1}(V_{\mathbb{Z}}^{a,b})$ is a lattice in $W^{a+b-q-r, a-q+r} \otimes \det^q$ stable under the action of H , and since $v^{a,b,q,r} \in V_{\mathbb{Z}}^{a,b}$, the intersection of this lattice with the highest-weight subspace contains the highest-weight vector $e_1^{a+b-q-r} \boxtimes f_1^{a-q+r}$. Hence this lattice must contain the minimal admissible lattice in $W^{a+b-q-r, a-q+r}$. ■

4.4. A Lie-theoretic computation

As in §2 above, let $T' \subset T$ be the rank-1 split torus $\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix}$; and let u be the element $\begin{pmatrix} 1 & & 1 \\ & 1 & \\ & & 1 \end{pmatrix}$ of $G(\mathbb{Z})$.

Since T' is split, the representations $V^{a,b}$ are the direct sums of their weight spaces relative to T' , with weights between 0 and $2a + b$; and the T' -weight of $v^{a,b,q,r}$ is $2a + b - q$. The purpose of this section is to prove the following result, which will be used in §9.5:

Lemma 4.4.1. *Let $v = v^{a,b,q,r} \in V^{a,b}$ be one of the above H -highest weight vectors. Then for any non-zero integer h , the projection of $u^h(v)$ to the highest- T' -weight space is given by $(2h)^q v^{a,b,0,r}$.*

Proof. Recall that $v^{a,b,q,r} = v^{b-r} \cdot (v')^r \cdot w^{a-q} \cdot (w')^q$. The vectors v , v' , and w all lie in the highest- T' -weight subspaces of their parent representations, so they are fixed by u . Hence it suffices to check that the projection of $u^h(w')$ to the highest- T' -weight subspace of $V^{1,0}$ is non-trivial; and one computes easily that $u^h(w') = w' + 2hw$. ■

5. Modular varieties

5.1. Modular curves

We fix conventions for modular curves.

Definition 5.1.1.

- (a) For $N \geq 3$, we let $Y(N)$ be the $\mathbb{Q}$ -variety parametrising triples (E, e_1, e_2) , where E is an elliptic curve (over some $\mathbb{Q}$ -algebra R) and $e_1, e_2 \in E(R)[N]$ are a basis of the N -torsion of E .
- (b) If Σ is a finite set of primes containing all those dividing N , we write $Y(N)_{\Sigma}$ for the natural model of $Y(N)$ over $\mathbb{Z}[1/\Sigma]$ (representing the same functor on $\mathbb{Z}[1/\Sigma]$ -algebras).

We identify $Y(N)(\mathbb{C})$ with the double quotient

$$\mathrm{GL}_2^+(\mathbb{Q}) \backslash (\mathrm{GL}_2(\mathbb{A}_f) \times \mathcal{H}) / U(N),$$

where $U(N) \subset \mathrm{GL}_2(\hat{\mathbb{Z}})$ is the principal congruence subgroup of level N and $\mathcal{H}$ is the upper half-plane, in such a way that:

- The double coset of $(1, \tau)$, for $\tau \in \mathcal{H}$, corresponds to the triple

$$\left(\frac{\mathbb{C}}{\mathbb{Z}\tau + \mathbb{Z}}, \frac{\tau}{N}, \frac{1}{N} \right).$$

- The right-translation action of $g \in \mathrm{GL}_2(\hat{\mathbb{Z}})$ on the double quotient corresponds to the action on $Y(N)(\mathbb{C})$ given by

$$(E, e_1, e_2) \cdot g = (E, e'_1, e'_2), \quad \begin{pmatrix} e'_1 \\ e'_2 \end{pmatrix} = g^{-1} \cdot \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}.$$

If $g \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ then the above action of g on $Y(N)(\mathbb{C})$ coincides with the action of γ^{-1} on $\mathcal{H}$, for any $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ congruent to g modulo N . The components of $Y(N)(\mathbb{C})$ are indexed by the set μ_N° of primitive N -th roots of unity, via the Weil pairing $(E, e_1, e_2) \mapsto \langle e_1, e_2 \rangle_N$; and the induced action of $g \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ on μ_N° is given by $g \cdot \zeta = \zeta^{1/\det(g)}$.

Remark 5.1.2. Note that our model is *not* the Deligne–Shimura canonical model of the Shimura variety for GL_2 with its standard Shimura datum [25, Example 5.6]. Rather, it is the canonical model for the twisted Shimura datum defined by

$$(a + ib) \in \mathrm{Res}_{\mathbb{C}/\mathbb{R}}(\mathbb{G}_m) \mapsto \frac{1}{a^2 + b^2} \begin{pmatrix} a & b \\ -b & a \end{pmatrix},$$

which has the effect of flipping the sign of the Galois action on the connected components. $\diamond$

By passage to the quotient, we define similarly algebraic varieties $Y(U)$ over $\mathbb{Q}$, for every open compact subgroup $U \subset \mathrm{GL}_2(\mathbb{A}_f)$; if U is unramified outside the finite set Σ , then $Y(U)$ has a model over $\mathbb{Z}[1/\Sigma]$ which we denote $Y(U)_\Sigma$.

The right-translation action gives isomorphisms $\eta : Y(U) \rightarrow Y(\eta^{-1}U\eta)$ for every $\eta \in \mathrm{GL}_2(\mathbb{A}_f)$, which are compatible with the action of η^{-1} on $\mathcal{H}$ if $\eta \in \mathrm{GL}_2^+(\mathbb{Q})$; in particular, scalar matrices $\begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}$ with $A \in \mathbb{Q}^\times$ act trivially. This structure allows us to view the inverse limit $Y = \varprojlim_U Y(U)$ as a pro-variety over $\mathbb{Q}$ with a right action of $\mathrm{GL}_2(\mathbb{A}_f)$, whose $\mathbb{C}$ -points are $\mathrm{GL}_2^+(\mathbb{Q}) \backslash (\mathrm{GL}_2(\mathbb{A}_f) \times \mathcal{H})$.

Definition 5.1.3. We say $U \subset \mathrm{GL}_2(\mathbb{A}_f)$ is *sufficiently small* if every non-identity element of U acts without fixed points on the set $\mathrm{GL}_2^+(\mathbb{Q}) \backslash (\mathrm{GL}_2(\mathbb{A}_f) \times \mathcal{H})$.

This condition is equivalent to requiring that for all $g \in \mathrm{GL}_2(\mathbb{A}_f)$, every non-identity element of the discrete group $\Gamma = \mathrm{GL}_2^+(\mathbb{Q}) \cap gUg^{-1}$ acts without fixed points on $\mathcal{H}$. For instance, $U(N)$ is sufficiently small if $N \geq 3$. If U is sufficiently small, then $Y(U)$ is the solution to a moduli problem (classifying elliptic curves with appropriate level structure), and therefore has an associated universal elliptic curve $\mathcal{E}(U) \rightarrow Y(U)$.

We define similarly algebraic surfaces $Y_H(U)$, where U is an open compact subgroup of $H(\mathbb{A}_f)$, and by passage to the limit a pro-variety $Y_H = \varprojlim_U Y_H(U)$ with a right action of $H(\mathbb{A}_f)$. Of course, if U is a fibre product $U_1 \times U_2$ of subgroups of $\mathrm{GL}_2(\mathbb{A}_f)$ such that $\det(U_1) = \det(U_2)$, then $Y_H(U)$ is the fibre product of the modular curves $Y(U_1)$ and $Y(U_2)$ over their common component set $\hat{\mathbb{Z}}^\times / \det(U_1) = \hat{\mathbb{Z}}^\times / \det(U_2)$.

5.2. Siegel modular varieties

Definition 5.2.1. Let $N \geq 3$. There exists a $\mathbb{Q}$ -variety $Y_G(N)$, smooth and quasiprojective of dimension 3, parametrisng 6-tuples $(A, \lambda, e_1, \dots, e_4)$ where

- A is an abelian surface (over some $\mathbb{Q}$ -algebra R);
- λ is a principal polarization $A \cong A^\vee$;
- $e_1, \dots, e_4$ are N -torsion sections of A giving an isomorphism $A[N] \cong (\mathbb{Z}/N\mathbb{Z})^4$;
- the matrix of the Weil pairing (induced by the polarization λ), with respect to the basis $e_1, \dots, e_4$, is $J \cdot \zeta$ for some $\zeta \in R^\times$.

Moreover, for any finite set of primes Σ containing the primes dividing N , $Y_G(N)$ has a model $Y_G(N)_\Sigma$ over $\mathbb{Z}[1/\Sigma]$, representing the same functor for $\mathbb{Z}[1/N]$ -algebras.

For the existence of this scheme see e.g. [17, Corollary 3.3]. The complex manifold $Y_G(N)(\mathbb{C})$ can be identified with the double quotient

$$\mathrm{GSp}_4^+(\mathbb{Q}) \backslash (\mathrm{GSp}_4(\mathbb{A}_f) \times \mathcal{H}_2) / U_G(N),$$

where $\mathcal{H}_2$ denotes the genus-2 Siegel space of symmetric complex 2×2 matrices with positive-definite imaginary part, and $U_G(N)$ is the principal congruence subgroup of level N (the kernel of reduction $\mathrm{GSp}_4(\widehat{\mathbb{Z}}) \rightarrow \mathrm{GSp}_4(\mathbb{Z}/N\mathbb{Z})$).

The right-translation action of $\mathrm{GSp}_4(\mathbb{Z}/N\mathbb{Z})$ on $Y_G(N)$ corresponds to the action on the moduli problem given by $g : (A, \lambda, e_1, \dots, e_4) \mapsto (A, \lambda, e'_1, \dots, e'_4)$, where

$$\begin{pmatrix} e'_1 \\ \vdots \\ e'_4 \end{pmatrix} = g^{-1} \cdot \begin{pmatrix} e_1 \\ \vdots \\ e_4 \end{pmatrix}.$$

More generally, if U is any open compact subgroup of $G(\mathbb{A}_f)$, we define a $\mathbb{Q}$ -model for $Y_G(U)$ by taking the quotient of $Y_G(N)$ by the action of $U/U_G(N)$, for any $N \geq 3$ such that $U_G(N) \subseteq U$. The same procedure gives $\mathbb{Z}[1/\Sigma]$ -models $Y_G(U)_\Sigma$ if U is unramified outside Σ . If U is sufficiently small (in the same sense as for GL_2), then $Y_G(U)$ is smooth, and can be interpreted as a moduli space for abelian surfaces with level U structure. As before, the inverse limit $\varprojlim_U Y_G(U)$ acquires a right action of $G(\mathbb{A}_f)$.

5.3. The embedding of Y_H in Y_G

For any open compact subgroup U of $G(\mathbb{A}_f)$, we have a natural morphism of $\mathbb{Q}$ -varieties

$$\iota_U : Y_H(U \cap H) \rightarrow Y_G(U).$$

The map ι_U is not always injective (even if U is sufficiently small). However, we have the following criterion:

Proposition 5.3.1. *Suppose there is an open compact subgroup $\tilde{U}$ containing U and wUw , where $w = \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}$, with $\tilde{U}$ sufficiently small. Then ι_U is injective.*

Proof. As above, write Y_G for the infinite-level Shimura variety $G(\mathbb{Q})_+ \backslash (G(\mathbb{A}_f) \times \mathcal{H}_2)$, and similarly for Y_H . It is clear that ι gives an injection $Y_H \hookrightarrow Y_G$. If $Q, Q' \in Y_H$ have the same image in $Y_G(U)$, then $Q' = Qu$ for some $u \in U$; we want this to imply that u lies in $U \cap H$. So it suffices to prove that, for any element of $U - (U \cap H)$, we have $Y_H u \cap Y_H = \emptyset$ as subsets of Y_G .

Since w is central in H , its action on Y_G fixes Y_H pointwise. Thus, if $Q \in Y_H$ and $Qu \in Y_H$, we have $Quw = Qu = Qwu$, so $\tilde{u} = u \cdot wu^{-1}w$ fixes Q . This element $\tilde{u}$ lies in $\tilde{U}$, by hypothesis, and since $\tilde{U}$ is sufficiently small, we conclude that $\tilde{u} = 1$. Thus u lies in the centraliser of w in $G(\mathbb{A}_f)$, which is exactly $H(\mathbb{A}_f)$. ■

We shall say a subgroup U is *H-small* if it satisfies the hypotheses of the above proposition. For instance, if U is contained in the principal congruence subgroup $U_G(N)$ for some $N \geq 3$, then U is *H-small* (since $U_G(N)$ is normal in $G(\widehat{\mathbb{Z}})$, and sufficiently small by [30, §0.6]).

5.4. Component groups and base extension

Via strong approximation for Sp_4 , we have an isomorphism of component sets

$$\pi_0(Y_G(\mathbb{C})) = \mathbb{Q}_+^\times \backslash \mathbb{A}_f^\times \cong \widehat{\mathbb{Z}}^\times.$$

Our moduli-space description of Y_G determines a Galois action on these components as follows.

Definition 5.4.1. We write

$$\mathrm{Art} : \mathbb{Q}_+^\times \backslash \mathbb{A}_f^\times \rightarrow \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})^{\mathrm{ab}}$$

for the Artin reciprocity map of class field theory, normalised so that for $x \in \widehat{\mathbb{Z}}^\times \subset \mathbb{A}_f^\times$, $\mathrm{Art}(x)$ acts on roots of unity as $\zeta \mapsto \zeta^x$ (and hence uniformizers map to geometric Frobenius elements).

Proposition 5.4.2. *All components of $Y_G(\mathbb{C})$ are defined over the cyclotomic extension $\mathbb{Q}^{\mathrm{ab}} = \mathbb{Q}(\zeta_n : n \geq 1)$, and the right-translation action of $u \in G(\mathbb{A}_f)$ on $\pi_0(Y_G(\mathbb{C}))$ coincides with the action of the Galois automorphism $\mathrm{Art}(\mu(u)^{-1})$.*

Proof. This is an instance of Deligne's reciprocity law for the action of Galois on the connected components of any Shimura variety; see e.g. [25, §13]. ■

We will be particularly interested in the following special case. If $U \subset G(\mathbb{A}_f)$ is an open compact subgroup, and V_N is the subgroup of $\widehat{\mathbb{Z}}^\times$ defined by $\{x : x \equiv 1 \pmod{N}\}$ for some integer N , then there is an embedding of $\mathbb{Q}$ -varieties

$$Y_G(U \cap \mu^{-1}(V_N)) \hookrightarrow Y_G(U) \times_{\mathrm{Spec} \mathbb{Q}} \mathrm{Spec} \mathbb{Q}(\zeta_N),$$

which is an isomorphism if $\mu(U)$ surjects onto $(\mathbb{Z}/N\mathbb{Z})^\times$. This map intertwines the action of $g \in G(\mathbb{A}_f)$ on the left-hand side with that of (g, σ) on the right-hand side, where σ is the image of $\mathrm{Art}(\mu(g)^{-1})$ in $\mathrm{Gal}(\mathbb{Q}(\zeta_N)/\mathbb{Q})$.

6. Coefficient sheaves on modular varieties

6.1. Étale coefficient sheaves

Let $U \subset \mathrm{GL}_2(\hat{\mathbb{Z}})$ be a sufficiently small open compact subgroup, and S a finite set with a continuous left action of U . Then we may construct a finite étale covering of $Y(U)$ as follows: we take any open normal subgroup $V \trianglelefteq U$ acting trivially on S , and we let $\mathcal{S}$ be the quotient of $Y(V) \times S$ by the left action of U/V given by

$$h \cdot (y, s) = (yh^{-1}, hs).$$

If S is a $\mathbb{Z}[U]$ -module, then $\mathcal{S}$ can be considered as a locally constant étale sheaf of abelian groups over $Y(U)$. Note that the sections of $\mathcal{S}$ over $Y(V)$, for any $V \trianglelefteq U$ open, are canonically isomorphic to S^V , and the pullback action of $u \in U/V$ on $H^0(Y(V), \mathcal{S})$ is identified with the natural left action of U/V on S^V . This construction extends in the obvious fashion to profinite modules S , and in particular to continuous representations of U on finite-rank $\mathbb{Z}_p$ -modules; via passage to the isogeny category we may also allow S to be a $\mathbb{Q}_p$ -vector space.

If the action of U on S extends to some larger monoid $\mathcal{M} \subseteq \mathrm{GL}_2(\mathbb{A}_f)$ containing U , then the sheaf $\mathcal{S}$ naturally becomes $\mathcal{M}$ -equivariant. That is, for every $\sigma \in \mathcal{M}$, given a morphism of varieties $Y(U) \xrightarrow{\sigma} Y(\sigma^{-1}U\sigma)$, we have morphisms $\sigma^*(\mathcal{S}') \rightarrow \mathcal{S}$, where $\mathcal{S}$ and $\mathcal{S}'$ are the sheaves on $Y(U)$ and $Y(\sigma^{-1}U\sigma)$, respectively, corresponding to S ; and these morphisms satisfy an appropriate cocycle condition. This construction equips the cohomology groups $H^*(Y(U), \mathcal{S})$ with an action of the Hecke algebra $\mathcal{H}(U \backslash \mathcal{M} / U)$.

Remark 6.1.1. Compare [24, Proposition 4.4.3]; our conventions here are a little different as we are considering right, rather than left, actions on our Shimura varieties. $\diamond$

Exactly the same theory applies, of course, to the modular varieties $Y_G(U)$ and $Y_H(U)$, and these constructions are compatible via ι : the pullback functor ι^* on étale sheaves corresponds to restriction of representations from G to H .

6.2. Sheaves corresponding to algebraic representations

As we have seen above, the modular curves $Y(U)$, for U sufficiently small, are moduli spaces: $Y(U)$ parametrises elliptic curves E equipped with a U -orbit of isomorphisms $E_{\mathrm{tors}} \cong (\mathbb{Q}/\mathbb{Z})^2$. Thus $Y(U)$ comes equipped with a universal elliptic curve $\mathcal{E}$. From the description of the action of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ on the moduli problem, one deduces the following compatibility:

Lemma 6.2.1. *Suppose $U \subseteq \mathrm{GL}_2(\hat{\mathbb{Z}})$. For $N \geq 1$, the sheaf $\mathcal{E}[N]$ of N -torsion points of $\mathcal{E}$ is canonically isomorphic to the sheaf associated to the **dual** of the standard representation of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$.*

Let p be a prime. Taking $N = p^r$ and passing to the limit over r shows that the relative Tate module $T_p \mathcal{E}$ corresponds to the dual of the standard representation of $\mathrm{GL}_2(\mathbb{Z}_p)$. On

the other hand, $T_p \mathcal{E}$ is a lattice in the p -adic étale realisation of a “motivic sheaf” – a relative Chow motive – over $Y(U)$, namely $h^1(\mathcal{E})^\vee$.

This is the first instance of a more general phenomenon. Let $\mathcal{G}$ temporarily denote any of the three groups $\{\mathrm{GL}_2, \mathrm{GL}_2 \times_{\mathrm{GL}_1} \mathrm{GL}_2, \mathrm{GSp}_4\}$; and let U be a sufficiently small open compact in $\mathcal{G}(\mathbb{A}_f)$, so we have an associated Shimura variety $Y_{\mathcal{G}}(U)$.

Lemma 6.2.2 ([1, Theorem 8.6]). *There is an additive functor*

$$\mathrm{Anc}_{\mathcal{G},U} : \mathrm{Rep}(\mathcal{G}) \rightarrow \mathrm{CHM}(Y_{\mathcal{G}}(U))$$

from the category of representations of $\mathcal{G}$ over $\mathbb{Q}$ to the category of relative Chow motives over $Y_{\mathcal{G}}(U)$ with the following properties:

- $\mathrm{Anc}_{\mathcal{G},U}$ preserves tensor products and duals;
- if μ denotes the multiplier map $\mathcal{G} \rightarrow \mathbb{G}_m$, then $\mathrm{Anc}_{\mathcal{G},U}(\mu)$ is the Lefschetz motive $\mathbb{Q}(-1)$;
- if V denotes the defining representation of $\mathcal{G}$, then $\mathrm{Anc}_{\mathcal{G},U}(V) = h^1(\mathcal{A}_U)$, where $\mathcal{A}_U$ is the universal PEL abelian variety over $Y_{\mathcal{G}}(U)$;
- for any prime p and $\mathcal{G}$ -representation V , the p -adic realisation of $\mathrm{Anc}_{\mathcal{G},U}(V)$ is the étale sheaf associated to $V \otimes \mathbb{Q}_p$, regarded as a left U -representation via $U \hookrightarrow \mathcal{G}(\mathbb{A}_f) \twoheadrightarrow \mathcal{G}(\mathbb{Q}_p)$.

Remark 6.2.3. In fact Ancona’s construction is much more general, applying to arbitrary PEL Shimura varieties, but we shall only need the above three groups here. The theorem stated in *op. cit.* is slightly different from ours, since he normalises his functor to send the multiplier representation to $\mathbb{Q}(1)$, and the defining representation to $h^1(\mathcal{A})^\vee$; our functor is obtained from his by composing with the automorphism of $\mathrm{Rep}(\mathcal{G})$ induced by the map $g \mapsto \mu(g)^{-1}g$ on $\mathcal{G}$. $\diamond$

We shall need some “naturality” properties of Ancona’s construction, which we now recall.

Proposition 6.2.4. *Suppose U, U' are open subgroups of $\mathcal{G}(\mathbb{A}_f)$ and $\sigma \in \mathcal{G}(\mathbb{A}_f)$ is such that $\sigma^{-1}U\sigma \subseteq U'$, so that right translation defines a map $\sigma : Y_{\mathcal{G}}(U) \rightarrow Y_{\mathcal{G}}(U')$. Then we have isomorphisms of functors*

$$\sigma^* \circ \mathrm{Anc}_{\mathcal{G},U'} \cong \mathrm{Anc}_{\mathcal{G},U},$$

satisfying a suitable compatibility condition under composition.

Proof. Since $\mathrm{Anc}_{\mathcal{G},U}(V)$ for a general V is defined as a direct summand of a tensor power of $h^1(\mathcal{A}_U)$, one reduces easily to checking this functoriality property for the specific relative motives $h^1(\mathcal{A}_U)$.

By a standard argument (see e.g. [6, Prop. 3.3]) one can interpret $Y_{\mathcal{G}}(U)$ as a moduli space for abelian varieties *up to isogeny*, from which we can deduce that there is a canonical isomorphism

$$\lambda_\sigma : \mathcal{A}_U \otimes \mathbb{Q} \xrightarrow{\cong} \sigma^*(\mathcal{A}_{U'}) \otimes \mathbb{Q}$$

in the isogeny category of abelian varieties over $Y_{\mathcal{G}}(U)$. The functor $h^1(-)$ extends to the isogeny category of abelian varieties, so λ_{σ} induces an isomorphism of relative Chow motives. Moreover, the λ_{σ} satisfy a cocycle condition for varying σ (which we leave it to the reader to formulate explicitly). ■

Remark 6.2.5. Note that λ_{σ} comes from a “genuine” isogeny if and only if the matrix of σ (in the defining matrix representation of $\mathcal{G}$) has entries in $\hat{\mathbb{Z}}$. To fix conventions, note that if $U = U'$ and $\sigma = \text{diag}(x, \dots, x)$ for some $x \in \mathbb{Q}^{\times+}$, then the map λ_{σ} is given by multiplication by x . ◇

An equivalent way of stating Proposition 6.2.4 is as follows. We define a $\mathcal{G}(\mathbb{A}_f)$ -equivariant relative Chow motive over $Y_{\mathcal{G}}$ to be the data of a relative Chow motive $\mathcal{V}_U$ over $Y_{\mathcal{G}}(U)$ for each sufficiently small open $U \subset G(\mathbb{A}_f)$, together with a collection of isomorphisms $\sigma^*(\mathcal{V}_{U'}) \cong \mathcal{V}_U$ for each inclusion $\sigma^{-1}U\sigma \subseteq U'$, compatible with composition. These objects form a category $\text{CHM}(Y_{\mathcal{G}})^{\mathcal{G}(\mathbb{A}_f)}$, and the proposition states that the functors $\text{Anc}_{\mathcal{G},U}$ for varying U assemble into a functor

$$\text{Anc}_{\mathcal{G}} : \text{Rep}(\mathcal{G}) \rightarrow \text{CHM}(Y_{\mathcal{G}})^{\mathcal{G}(\mathbb{A}_f)}.$$

(Note that the isomorphisms $\text{Anc}_{\mathcal{G},U}(\mu) \cong \mathbb{Q}(-1)$ are not compatible with the equivariant structure, since the isogenies λ_{σ} only preserve the polarisation up to a scalar; as equivariant motives we have $\text{Anc}_{\mathcal{G}}(\mu) = \mathbb{Q}(-1)[-1]$, where $[-1]$ denotes that the equivariant structure is twisted by the character $\|\mu\|^{-1}$ of $G(\mathbb{A}_f)$.)

If $\mathcal{V}$ is a $\mathcal{G}(\mathbb{A}_f)$ -equivariant relative Chow motive over $Y_{\mathcal{G}}$, we can define its motivic cohomology by

$$H_{\text{mot}}^*(Y_{\mathcal{G}}, \mathcal{V}) := \lim_{\substack{\longrightarrow \\ U}} H_{\text{mot}}^*(Y_{\mathcal{G}}(U), \mathcal{V}_U), \quad (6)$$

and this is naturally a smooth representation of $\mathcal{G}(\mathbb{A}_f)$. As motivic cohomology with rational coefficients satisfies Galois descent (see e.g. [7, §1.3]), for each sufficiently small U we can recover $H_{\text{mot}}^*(Y_{\mathcal{G}}(U), \mathcal{V}_U)$ as the U -invariants of the direct limit (6).

Finally, we shall need to show a compatibility with respect to changing $\mathcal{G}$:

Proposition 6.2.6 (“Branching” for motivic sheaves). *Let $G = \text{GSp}_4$ and $H = \text{GL}_2 \times_{\text{GL}_1} \text{GL}_2$, as in §2 above. Then there is a commutative diagram of functors*

$$\begin{array}{ccc} \text{Rep}(G) & \xrightarrow{\text{Anc}_G} & \text{CHM}(Y_G)^{G(\mathbb{A}_f)} \\ \iota^* \downarrow & & \downarrow \iota^* \\ \text{Rep}(H) & \xrightarrow{\text{Anc}_H} & \text{CHM}(Y_H)^{H(\mathbb{A}_f)} \end{array}$$

where the left-hand ι^* denotes restriction of representations, and the right-hand ι^* denotes pullback of relative motives.

Proof. This is an instance of a general theorem due to Torzewski [43, Corollary 9.8], which verifies the above naturality property for a wide class of homomorphisms of PEL-type Shimura data. ■

7. Eisenstein classes for GL_2

7.1. Modular units

Let $\mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Q}) \subseteq \mathcal{S}(\mathbb{A}_f^2, \mathbb{Q})$ denote the subspace of functions satisfying $\phi(0, 0) = 0$. Recall that Y denotes the infinite-level modular curve, so that $\mathcal{O}(Y) = \varinjlim_U \mathcal{O}(Y(U))$.

Proposition 7.1.1. *There is a canonical, $\mathrm{GL}_2(\mathbb{A}_f)$ -equivariant map $\mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Q}) \rightarrow \mathcal{O}(Y)^\times \otimes \mathbb{Q}$, $\phi \mapsto g_\phi$, with the following characterising property: if ϕ is the characteristic function of $(a, b) + N\hat{\mathbb{Z}}^2$ for some $N \geq 1$ and $(a, b) \in \mathbb{Q}^2 - N\mathbb{Z}^2$, then g_ϕ is the Siegel unit $g_{a/N, b/N}$ in the notation of [12, §1.4].*

Proof. See e.g. [5, Théorème 1.8]. ■

In order to work integrally, we need to modify the construction somewhat. Let $c > 1$ be an integer. We let ${}_c\mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Z})$ denote the subgroup of $\mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Q})$ consisting of all functions of the form $\phi = \phi^{(c)} \cdot \mathrm{ch}(\mathbb{Z}_c^2)$, where $\phi^{(c)}$ is a $\mathbb{Z}$ -valued Schwartz function on $(\mathbb{A}_f^{(c)})^2$, and $\mathbb{Z}_c = \prod_{\ell|c} \mathbb{Z}_\ell$. Then we have the following refinement:

Proposition 7.1.2. *If c is coprime to 6, there is a map ${}_c\mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Z}) \rightarrow \mathcal{O}(Y)^\times$, $\phi \mapsto {}_cg_\phi$, which is equivariant for the action of $\mathrm{GL}_2(\mathbb{A}_f^{(c)})$ and satisfies*

$${}_cg_\phi \otimes 1 = (c^2 - \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}^{-1})g_\phi \quad \text{as elements of } \mathcal{O}(Y)^\times \otimes \mathbb{Q},$$

where $\begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}$ is understood as an element of $\mathrm{GL}_2(\mathbb{A}_f^{(c)})$.

Proof. See [12, §1.4]. ■

7.2. Higher Eisenstein classes

Definition 7.2.1. For $k \geq 0$, let $\mathcal{H}_{\mathbb{Q}}^k$ denote the $\mathrm{GL}_2(\mathbb{A}_f)$ -equivariant relative Chow motive over Y associated to the representation $\mathrm{Sym}^k(\mathrm{std}) \otimes \det^{-k}$ of $\mathrm{GL}_2/\mathbb{Q}$.

Theorem 7.2.2 (Beilinson). *Let $k \geq 1$. There is a $\mathrm{GL}_2(\mathbb{A}_f)$ -equivariant map $\mathcal{S}(\mathbb{A}_f^2, \mathbb{Q}) \rightarrow H_{\mathrm{mot}}^1(Y, \mathcal{H}_{\mathbb{Q}}^k(1))$, $\phi \mapsto \mathrm{Eis}_{\mathrm{mot}, \phi}^k$, the motivic Eisenstein symbol, with the following property: the pullback of the de Rham realization $r_{\mathrm{dR}}(\mathrm{Eis}_{\mathrm{mot}, \phi}^k)$ to the upper half-plane is the $\mathcal{H}^k$ -valued differential 1-form*

$$-F_\phi^{(k+2)}(\tau)(2\pi i \, \mathrm{d}z)^k(2\pi i \, \mathrm{d}\tau),$$

where $F_\phi^{(k+2)}$ is the Eisenstein series defined by

$$F_\phi^{(k+2)}(\tau) = \frac{(k+1)!}{(-2\pi i)^{k+2}} \sum_{\substack{x, y \in \mathbb{Q} \\ (x, y) \neq (0, 0)}} \frac{\hat{\phi}(x, y)}{(x\tau + y)^{k+2}}.$$

Proof. See [2]. ■

Remark 7.2.3. Note that if ϕ is the characteristic function of $(0, b) + N\hat{\mathbb{Z}}^2$, then $\text{Eis}_{\text{mot}, \phi}^k$ is the class $\text{Eis}_{\text{mot}, b, N}^k$ defined in [16, Theorem 4.1.1]. If $k = 0$, then we need to assume $\phi \in \mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Q})$ in order for the series defining $F_\phi^{(2)}$ to be absolutely convergent. With this assumption, we may define $\text{Eis}_{\text{mot}, \phi}^0$ to be the unit g_ϕ , since $H_{\text{mot}}^1(Y, \mathbb{Q}(1)) = \mathcal{O}^\times(Y) \otimes \mathbb{Q}$; the de Rham realisation of this class is then $\text{dlog } g_\phi = -F_\phi^{(2)} \cdot (2\pi i \, \text{d}\tau)$, so our statements are consistent. $\diamond$

Since we lack a good theory of relative Chow motives with coefficients in $\mathbb{Z}$, we do not have an integral version of the motivic Eisenstein classes for $k > 0$. However, we can obtain a $\mathbb{Z}_p$ -structure (for a fixed p) using étale cohomology instead. For each U we have an *étale realisation* map

$$r_{\text{ét}} : H_{\text{mot}}^1(Y(U), \mathcal{H}_{\mathbb{Q}}^k(1)) \rightarrow H_{\text{ét}}^1(Y(U), \mathcal{H}_{\mathbb{Q}_p}^k(1))$$

for any prime p , where $H_{\text{ét}}^*$ denotes continuous étale cohomology in the sense of [11], and $\mathcal{H}_{\mathbb{Q}_p}^k$ is the lisse étale $\mathbb{Q}_p$ -sheaf which is the p -adic realisation of $\mathcal{H}_{\mathbb{Q}}$. This is naturally the base extension to $\mathbb{Q}_p$ of the étale $\mathbb{Z}_p$ -sheaf $\mathcal{H}_{\mathbb{Z}_p}^k$ associated to the *minimal* admissible lattice in the GL_2 -representation $\text{Sym}^k(\text{std}) \otimes \det^{-k}$.

Proposition 7.2.4. *Let $k \geq 0$. If c is coprime to $6p$, then for each sufficiently small open compact $U \subset \text{GL}_2(\mathbb{A}_f^{(pc)} \times \mathbb{Z}_{pc})$ there is a map*

$${}_c\mathcal{S}((\mathbb{A}_f^{(p)} \times \mathbb{Z}_p)^2, \mathbb{Z}_p)^U \rightarrow H_{\text{ét}}^1(Y(U), \mathcal{H}_{\mathbb{Z}_p}^k(1)), \quad \phi \mapsto {}_c\text{Eis}_{\text{ét}, \phi}^k,$$

which is equivariant for the action of $\text{GL}_2(\mathbb{A}_f^{(pc)} \times \mathbb{Z}_{pc})$, and satisfies

$${}_c\text{Eis}_{\text{ét}, \phi}^k \otimes 1 = (c^2 - c^{-k} \begin{pmatrix} c & 0 \\ 0 & c \end{pmatrix}^{-1}) r_{\text{ét}}({}_c\text{Eis}_{\text{mot}, \phi}^k)$$

as elements of $H_{\text{ét}}^1(Y(U), \mathcal{H}_{\mathbb{Q}_p}^k(1))$.

Note that étale cohomology with $\mathbb{Z}_p$ coefficients does not satisfy Galois descent, so it is important to formulate Proposition 7.2.4 for each individual level, rather than simply passing to the direct limit.

Proof of Proposition 7.2.4. For levels of the form $U(N)$ this is explained in [14], and the arguments apply without change to a general U . $\blacksquare$

Remark 7.2.5. For $k = 0$, $\mathcal{H}_{\mathbb{Z}_p}^k$ is the constant sheaf $\mathbb{Z}_p$, and of course ${}_c\text{Eis}_{\text{ét}, \phi}^0$ is the image of ${}_c g_\phi$ under the Kummer map, so the $k = 0$ case of Proposition 7.2.4 is consistent with Proposition 7.1.2. $\diamond$

7.3. The modular unit representation

We will need a description of the modular units $\mathcal{O}^\times(Y) \otimes \mathbb{C}$ as a $\text{GL}_2(\mathbb{A}_f)$ -representation.

Definition 7.3.1. For $k \geq 0$, and η a finite-order character of $\mathbb{A}_f^\times / \mathbb{Q}^{\times+}$ satisfying $\eta(-1) = (-1)^k$, let $I_k(\eta)$ denote the space of functions $f : \mathrm{GL}_2(\mathbb{A}_f) \rightarrow \mathbb{C}$ satisfying

$$f\left(\begin{pmatrix} a & b \\ & d \end{pmatrix} g\right) = \|a\|^{k+1} \|d\|^{-1} \eta(a) f(g) \quad \text{for all } a, b, d \in \mathrm{GL}_2(\mathbb{A}_f),$$

regarded as a representation of $\mathrm{GL}_2(\mathbb{A}_f)$ by right translation. For $k = 0$ and $\eta = 1$, let $I_0^0(1)$ denote the subrepresentation which is the kernel of the natural map $I_0(1) \rightarrow \mathbb{C}$ given by integration over $\mathrm{GL}_2(\mathbb{A}_f)/B(\mathbb{A}_f)$.

Theorem 7.3.2. *There is a $\mathrm{GL}_2(\mathbb{A}_f)$ -equivariant isomorphism*

$$\partial_0 : \frac{\mathcal{O}^\times(Y)}{(\mathbb{Q}^{\mathrm{ab}})^\times} \otimes \mathbb{C} \xrightarrow{\cong} I_0^0(1) \oplus \bigoplus_{\eta \neq 1} I_0(\eta),$$

characterised by the statement that if $g \in \mathcal{O}^\times(Y)$, then $\partial_0(g)(1)$ is the order of vanishing of g at the cusp ∞ .

Proof. See [39, Theorem 3]. (Scholl's normalisations are slightly different from ours, as he uses the canonical model of Y for a different choice of Shimura datum; see Remark 5.1.2. The above formulation is correct for our choice of model.) ■

For $k \geq 1$ we have an analogous statement for the image of the Eisenstein symbol, although we do not know if this image is the whole of the motivic cohomology:

Theorem 7.3.3. *For $k \geq 1$, there is a surjective $\mathrm{GL}_2(\mathbb{A}_f)$ -equivariant map*

$$\partial_k : H_{\mathrm{mot}}^1(Y, \mathcal{H}_{\mathbb{Q}}^k(1)) \otimes \mathbb{C} \twoheadrightarrow \bigoplus_{\eta} I_k(\eta),$$

such that $\partial_k(x)(1)$ is the residue at ∞ of the 1-form $r_{\mathrm{dR}}(x)$. This map is an isomorphism on the image of the Eisenstein symbol $\phi \mapsto \mathrm{Eis}_{\mathrm{mot}, \phi}^k$.

Proof. It is shown in [38, Theorem 7.4] that the residue map ∂_k gives an isomorphism between the image of the Eisenstein symbol and a certain vector space denoted $\mathcal{B}_k$. For the description of this $\mathcal{B}_k$ as a sum of induced representations, see [22, Lemma 4.3]. ■

For η a character of $\mathbb{A}_f^\times / \mathbb{Q}^{\times+}$ as above, let us write $\mathcal{S}(\mathbb{A}_f^2, \mathbb{C})^\eta$ for the subspace of $\mathcal{S}(\mathbb{A}_f^2, \mathbb{C})$ on which $\widehat{\mathbb{Z}}^\times$ acts via the character η .

Proposition 7.3.4. *Let $\phi \in \mathcal{S}(\mathbb{A}_f^2, \mathbb{C})^\eta$ be of the form $\prod_{\ell \text{ prime}} \phi_\ell$. If $k = 0$ and $\eta = 1$ then assume that $\phi(0, 0) = 0$. Then*

$$\partial_k(\mathrm{Eis}_{\mathrm{mot}, \phi}^k) = \frac{2(k+1)!L(k+2, \eta)}{(-2\pi i)^{k+2}} \prod_{\ell} f_{\phi_\ell, \eta_\ell, |\cdot|^{k+1/2}, |\cdot|^{-1/2}}.$$

Proof. This follows from a computation of the constant term of the Eisenstein series $F_\phi^{(k+2)}$ at the cusp ∞ , using the formulae in [12, Proposition 3.10]. ■

8. Construction of Lemma–Eisenstein classes

8.1. Coefficients

Let $(a, b) \geq 0$ be a pair of non-negative integers, defining an algebraic representation $V^{a,b}$ of G . We write $D^{a,b}$ for the twist $V^{a,b} \otimes \mu^{-(2a+b)}$. We then have an equivariant relative Chow motive $\mathcal{D}_{\mathbb{Q}}^{a,b} = \text{Anc}_G(D^{a,b})$ over the Shimura variety Y_G .

Notation 8.1.1. Let us choose integers (q, r) with $0 \leq q \leq a$ and $0 \leq r \leq b$, and set $c = (a - q) + (b - r)$, $d = (a - q) + r$ (so $c, d \geq 0$).

Then there is a branching map

$$\text{br}^{[a,b,q,r]} : (\text{Sym}^c \boxtimes \text{Sym}^d) \otimes \det^{-(c+d)} \rightarrow D^{a,b} \otimes \mu^q$$

as in (5). Via the commutative diagram of functors in Proposition 6.2.6, we have a homomorphism of equivariant Chow motives over Y_H ,

$$\text{br}^{[a,b,q,r]} : \mathcal{H}_{\mathbb{Q}}^{c,d} \rightarrow \iota^*(\mathcal{D}_{\mathbb{Q}}^{a,b}(-q)[-q]),$$

where $\mathcal{H}_{\mathbb{Q}}^{c,d}$ is the relative motive associated to the H -representation $(\text{Sym}^c \boxtimes \text{Sym}^d) \otimes \det^{-(c+d)}$. (Recall that $[m]$ denotes twisting by the character $\|\mu(-)\|^m$ of $G(\mathbb{A}_f)$.)

8.2. Pushforwards in motivic cohomology

Let (a, b, q, r) and (c, d) be as in the previous section. We shall define in this section a homomorphism of left $H(\mathbb{A}_f) \times G(\mathbb{A}_f)$ -representations

$$\iota_*^{[a,b,q,r]} : H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) \otimes \mathcal{H}(G(\mathbb{A}_f); \mathbb{Q}) \rightarrow H_{\text{mot}}^4(Y_G, \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))[-q]. \quad (7)$$

The actions of $H(\mathbb{A}_f) \times G(\mathbb{A}_f)$ for which this map is equivariant are given as follows:

- The $H(\mathbb{A}_f)$ factor acts trivially on the right-hand side of (8), and on the left-hand side it acts via the formula

$$h \cdot (x \otimes \xi) = (h \cdot x) \otimes \xi(h^{-1}(-)).$$

- On the left-hand side, the $G(\mathbb{A}_f)$ factor acts trivially on $H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2))$, and on $\mathcal{H}(G(\mathbb{A}_f); \mathbb{Z})$ it acts via $g \cdot \xi = \xi((-)g)$.
- On the right-hand side, $G(\mathbb{A}_f)$ via its natural action on $H_{\text{mot}}^4(Y_G(U), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))$ (deduced from Proposition 6.2.4) twisted by the character $\|\mu(-)\|^{-q}$.

For ease of reading we shall drop the superscripts $[a, b, q, r]$ for the rest of this section.

Lemma 8.2.1. *The space $H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) \otimes \mathcal{H}(G(\mathbb{A}_f); \mathbb{Q})$ is spanned by vectors of the form $x \otimes \text{ch}(gU)$, where $x \in H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2))$, $g \in G(\mathbb{A}_f)$, and U is an open compact subgroup of $G(\mathbb{A}_f)$ such that $H(\mathbb{A}_f) \cap gUg^{-1}$ fixes x and gUg^{-1} is H -small in the sense of §5.3.*

Proof. This is immediate from the fact that the principal congruence subgroup $U_G(N)$, for any $N \geq 3$, is H -small, and these are cofinal among open compact subgroups of $G(\mathbb{A}_f)$. ■

For an element $x \otimes \mathrm{ch}(Ug)$ as in Lemma 8.2.1, we have a closed immersion $\iota_{gU} : Y_H(V) \hookrightarrow Y_G(U)$, where $V = U \cap H(\mathbb{A}_f)$, given by the composite

$$Y_H(V) \xrightarrow{\iota_{gUg^{-1}}} Y_G(gUg^{-1}) \xrightarrow{g^{-1}} Y_G(U).$$

Combining this with the morphism of sheaves $\mathrm{br}^{[a,b,q,r]}$ gives a map

$$\iota_{gU,*} : H_{\mathrm{mot}}^2(Y_H(V), \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) \rightarrow H_{\mathrm{mot}}^4(Y_G(U), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))[-q].$$

We define $\iota_*(x \otimes \mathrm{ch}(gU))$ as the image of the element

$$\mathrm{Vol}(V) \cdot \iota_{gU,*}(x) \in H_{\mathrm{mot}}^4(Y_G(U), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))[-q]$$

in the direct limit (6). (Here $\mathrm{Vol}(V)$ is volume with respect to Haar measure, normalised such that $\mathrm{Vol}(H(\hat{\mathbb{Z}})) = 1$.)

Now, suppose $U' \subset U$ is another H -small open compact subgroup, so that $\mathrm{ch}(gU) = \sum_{\gamma \in U/U'} \mathrm{ch}(g\gamma U')$. We want to show that $\iota_*(x \otimes \mathrm{ch}(gU)) = \sum_{\gamma \in U/U'} \iota_*(x \otimes \mathrm{ch}(g\gamma U'))$ for any x invariant under V . It suffices to prove this when $U' \trianglelefteq U$ (since otherwise we may compare both U and U' with a third open compact U'' normal in both U and U'); we may clearly also assume $g = 1$.

Let $V' = U' \cap H(\mathbb{A}_f)$; we then have degeneracy maps $\mathrm{pr}_U^{U'} : Y_G(U') \rightarrow Y_G(U)$ and $\mathrm{pr}_V^{V'} : Y_H(V') \rightarrow Y_H(V)$, fitting into a commutative diagram

$$\begin{array}{ccc} Y_H(V') & \xrightarrow{\iota_{U'}} & Y_G(U') \\ \mathrm{pr}_V^{V'} \downarrow & & \downarrow \mathrm{pr}_U^{U'} \\ Y_H(V) & \xrightarrow{\iota_U} & Y_G(U) \end{array}$$

By the functoriality of the pushforward maps, we have

$$(\mathrm{pr}_U^{U'})^* \circ (\mathrm{pr}_U^{U'})_* \circ \iota_{U',*} = (\mathrm{pr}_U^{U'})^* \circ \iota_{U,*} \circ (\mathrm{pr}_V^{V'})_*.$$

The composite $(\mathrm{pr}_U^{U'})^* (\mathrm{pr}_U^{U'})_*$ is given by $\sum_{\gamma \in U/U'} \gamma^*$; and as x is invariant under V (not only V'), then $(\mathrm{pr}_V^{V'})_*(x) = [V : V']x$. So we can write this as

$$\mathrm{Vol}(V') \cdot \sum_{\gamma \in U/U'} \gamma^* \iota_{U',*}(x) = \mathrm{Vol}(V) \cdot (\mathrm{pr}_U^{U'})^* \iota_{U,*}(x).$$

Pulling back from level U' to the direct limit over all levels, we can write this as

$$\iota_*(x \otimes \mathrm{ch}(U)) = \sum_{\gamma \in U/U'} \iota_*(x \otimes \mathrm{ch}(\gamma U')).$$

It follows that ι_* is well-defined on $H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) \otimes \mathcal{H}(G(\mathbb{A}_f); \mathbb{Q})$. It is obvious that this map is $G(\mathbb{A}_f)$ -equivariant; and the $H(\mathbb{A}_f)$ -equivariance follows from the obvious compatibility

$$\iota_{U,*}(h \cdot x) = h \cdot \iota_{hUh^{-1},*}(x)$$

of pushforward maps at finite level.

8.3. The Lemma–Eisenstein map

The cup-product of the Eisenstein symbols for the two factors of H defines an $H(\mathbb{A}_f)$ -equivariant map

$$\mathcal{S}_0(\mathbb{A}_f^2; \mathbb{Q})^{\otimes 2} \rightarrow H_{\text{mot}}^2(Y_H, \mathcal{H}_{\mathbb{Q}}^{c,d}(2)), \quad \underline{\phi} \mapsto \text{Eis}_{\text{mot}, \underline{\phi}}^{c,d}.$$

Definition 8.3.1. We define the *Lemma–Eisenstein map*

$$\mathcal{LE}^{[a,b,q,r]} : \mathcal{S}_0(\mathbb{A}_f^2; \mathbb{Q})^{\otimes 2} \otimes \mathcal{H}(G(\mathbb{A}_f); \mathbb{Q}) \rightarrow H_{\text{mot}}^4(Y_G, \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))[-q] \quad (8)$$

by $\mathcal{LE}^{[a,b,q,r]}(\underline{\phi} \otimes \xi) = \iota_*^{[a,b,q,r]}(\text{Eis}_{\text{mot}, \underline{\phi}}^{c,d} \otimes \xi)$, where $\iota_*^{[a,b,q,r]}$ is as in (7).

Remark 8.3.2. When ξ is the characteristic function of an open compact subgroup $U \subseteq G(\mathbb{A}_f)$, our class $\mathcal{LE}_U^{[a,b,q,r]}(\underline{\phi} \otimes \xi)$ coincides with the motivic cohomology class $\text{Eis}_{\mathcal{M}}^{m,n,W}(h)$ considered in [21] for $S = Y_G(U)$, $(m, n) = (a - q + b - r, a - q + r)$, W the representation $V^{a,b}$, and h an appropriate element of Lemma’s space $\mathcal{B}_m \otimes \mathcal{B}_n$ depending on $\underline{\phi}$. In particular, it follows from the regulator computations of [22, §7] that the Lemma–Eisenstein map is non-zero under fairly mild hypotheses on a, b, q, r . $\diamond$

8.4. Choices of the local data

We shall now fix choices of the input data to the above map $\mathcal{LE}^{[a,b,q,r]}$, in order to define a collection of motivic cohomology classes satisfying appropriate norm relations (a “motivic Euler system”). We shall work with arbitrary (but fixed) choices of local data at the bad primes; it is the local data at good primes which we shall vary, according to the values of three parameters M, m, n .

8.4.1. Subgroups of tame level 1. We fix a prime p , a finite set of primes S not containing p , and an arbitrary open compact subgroup $K_S \subset G(\mathbb{Q}_S) = \prod_{\ell \in S} G(\mathbb{Q}_{\ell})$. By enlarging S and shrinking K_S if necessary, we may assume that the open compact subgroup

$$K_G = K_S \times \prod_{\ell \notin S} G(\mathbb{Z}_{\ell}) \subseteq G(\mathbb{A}_f)$$

is *sufficiently small* in the sense of §5.2. For each $n \geq 0$, we define a compact subgroup of $G(\mathbb{A}_f)$ by

$$K_{G,0}(p^n) := K_S \times K_{G_p,0}(p^n) \times \prod_{\ell \notin S \cup \{p\}} G(\mathbb{Z}_{\ell}),$$

where the subgroup $K_{G_p,0}(p^n)$ of $G_p = G(\mathbb{Q}_p)$ is as defined in §3.3. We define similarly subgroups $K_{G,1}(p^n)$ for $n \geq 0$, and $K_G(p^m, p^n)$ for $m, n \geq 0$, using the other local subgroups at p defined in §3.3. All of these groups are contained in K_G , and hence are sufficiently small.

Notation 8.4.1. We adopt the notational convention that if $K_{G,\star}(\square)$ denotes some open subgroup of $G(\mathbb{A}_f)$, then $Y_{G,\star}(\square)$ denotes the corresponding Shimura variety, so e.g. $Y_{G,1}(p^n)$ is an abbreviation for $Y_G(K_{G,1}(p^n))$.

8.4.2. *Local data at the bad primes.* We choose the following “test data” at S :

- A vector $\phi_S \in \mathcal{S}(\mathbb{Q}_S^2, \mathbb{Z})^{\otimes 2}$.
- An open compact subgroup $W_S \subset H(\mathbb{Q}_S)$ such that $W_S \subseteq H(\mathbb{Q}_S) \cap K_S$, and W_S acts trivially on ϕ_S .

Whenever we deal with norm-compatibility relations we shall assume that the local data K_S , W_S , ϕ_S remain fixed (i.e. we shall not attempt to formulate any non-trivial norm-compatibilities at the bad primes). Regarding the choice of ϕ_S , see Remark 10.6.4 below.

8.4.3. *Subgroups of higher tame level.* Now let us choose a square-free integer $M \geq 1$ coprime to $S \cup \{p\}$ (which we shall refer to as a “tame level”). For $m \geq 0$ and $n \geq 1$, we define a subgroup $K_G(M, p^m, p^n) \subseteq K_G(p^m, p^n)$ by

$$K_G(M, p^m, p^n) = \{k \in K_G(p^m, p^n) : \mu(k) = 1 \bmod M\}.$$

As explained in §5.4, we have isomorphisms

$$J_{Mp^m} : Y_G(M, p^m, p^n) \xrightarrow{\cong} Y_{G,1}(p^n) \times_{\mathrm{Spec} \mathbb{Q}} \mathrm{Spec} \mathbb{Q}(\zeta_{Mp^m}). \quad (9)$$

Assuming $n \geq m$, we also define $K'_G(M, p^m, p^n) = \{k \in K'_G(p^m, p^n) : \mu(k) = 1 \bmod M\}$; note that the difference between this and $K_G(M, p^m, p^n)$ is only at p – we do not impose stronger congruences at M .

8.4.4. *Test data of higher level.* Let (M, m, n) be integers as above. For each such triple, we shall define the following data:

- an element $\xi_{M,m,n} \in \mathcal{H}(G(\mathbb{A}_f), \mathbb{Z})$, fixed by the right-translation action of $K_G(M, p^m, p^n)$;
- a subgroup W of $H(\mathbb{A}_f)$ such that for all x in the support of $\xi_{M,m,n}$, we have $W \subseteq H(\mathbb{A}_f) \cap xK_G(M, p^m, p^n)x^{-1}$;
- an element $\phi_{M,m,n} \in \mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Z})^{\otimes 2}$ stable under W .

We shall define these as products

$$\xi_{M,m,n} = \mathrm{ch}(K_S) \otimes \bigotimes_{\ell \notin S} \xi_\ell, \quad W = W_S \times \prod_{\ell \notin S} W_\ell, \quad \phi_{M,m,n} = \phi_S \otimes \bigotimes_{\ell \notin S} \phi_\ell,$$

where the local data K_S, W_S, ϕ_S at the bad places are the ones chosen above (independently of M, m, n) and the local data at primes $\ell \notin S$ are as follows. As in §3, we let $\eta_{\ell,r} \in G(\mathbb{Q}_\ell)$ denote the element

$$\begin{pmatrix} 1 & \ell^{-r} & & \\ & 1 & \ell^{-r} & \\ & & 1 & \\ & & & 1 \end{pmatrix}.$$

- If $\ell \nmid Mp$, we set $\xi_\ell = \text{ch}(G(\mathbb{Z}_\ell))$, $W_\ell = H(\mathbb{Z}_\ell)$, and $\phi_\ell = \text{ch}(\mathbb{Z}_\ell^2)^{\otimes 2}$.
- If $\ell \mid M$, we set $\xi_\ell = \text{ch}(K_{G_\ell}(\ell, 1)) - \text{ch}(\eta_{\ell,1} \cdot K_{G_\ell}(\ell, 1))$, and $W_\ell = K_{H_\ell}(\ell, \ell^2)$. We take $\phi_\ell = \text{ch}(\ell^2 \mathbb{Z}_\ell \times (1 + \ell^2 \mathbb{Z}_\ell))^{\otimes 2}$.
- For $\ell = p$, we set $\xi_p = \text{ch}(\eta_{p,m} \cdot K_{G_p}(p^m, p^n))$. We choose an integer $t \geq 1$ sufficiently large² that $K_{H_p}(p^m, p^t)$ is contained in $\eta_{p,m} \cdot K_{G_p}(p^m, p^n) \cdot \eta_{p,m}^{-1}$; we let W_p be this subgroup, and we define

$$\phi_p = \text{ch}(p^t \mathbb{Z}_p \times (1 + p^t \mathbb{Z}_p))^{\otimes 2}.$$

Note that $\phi_{M,m,n} \in \mathcal{S}_0(\mathbb{A}_f^2, \mathbb{Z})^{\otimes 2} \subset \mathcal{S}(\mathbb{A}_f^2, \mathbb{Z})^{\otimes 2}$, since our local Schwartz functions at p vanish at $(0, 0)$. Both the element $\phi_{M,m,n}$, and the group W , depend on the auxiliary choice of t ; but if we let $t^\circ > t$ be another choice, and $\phi_{M,m,n}^\circ, W^\circ$ the objects defined using t° in place of t , then we have

$$\phi_{M,m,n} = \sum_{w \in W/W^\circ} w \cdot \phi_{M,m,n}^\circ. \quad (10)$$

We also define a version mildly modified at p , assuming that $n \geq 1$ and $m \leq n$. Recall the subgroup $K'_{G_p}(p^m, p^n)$ defined in §3.3. We define $\xi'_p = \text{ch}(\eta_{p,0} K'_{G_p}(p^m, p^n)) = \text{ch}(K'_{G_p}(p^m, p^n) \eta_{p,0})$. Thus ξ'_p is preserved under left-translation by $W'_p = K'_{H_p}(p^m, p^n)$; and we choose $\phi'_p = \text{ch}(p^n \mathbb{Z}_p \times (1 + p^n \mathbb{Z}_p))^{\otimes 2}$. We let $K'_G(M, p^m, p^n)$, $\xi'_{M,m,n}$, $\phi'_{M,m,n}$ and W' be the adélic objects defined using these modified choices at p , and the same choices as before at all other primes.

Remark 8.4.2. These alternative local choices will give elements related to the “non-dashed” versions in the same way as the elements $\mathbb{Z}_{\dots}$ relate to the elements $\Xi_{\dots}$ in [18]. As in *op. cit.*, it is the non-dashed versions that are of interest for applications, but the dashed versions are convenient for certain calculations, in particular for studying p -adic integrality and interpolation properties. $\diamond$

8.4.5. The Lemma–Eisenstein classes and their norm relations at p . With the above notations and choices, let us define

$$\begin{aligned} & z_{M,m,n}^{[a,b,q,r]} \\ &= \frac{1}{\text{Vol}(W)} \mathcal{L}\mathcal{E}_{K_G(M,p^m,p^n)}^{[a,b,q,r]}(\phi_{M,m,n} \otimes \xi_{M,m,n}) \in H_{\text{mot}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q)). \end{aligned}$$

²One can check that $t = n + 2m$ suffices.

We refer to these elements as *Lemma–Eisenstein classes*. *A priori* this element depends on the auxiliary integer t , but it follows readily from (10) that it is in fact independent of this choice (this is essentially the same computation as in Lemma 3.10.2). It can be written concretely as follows: letting U be the subgroup $K_G(M, p^m, p^n)$, we can write our Hecke-algebra element $\xi_{M,m,n}$ as a finite $\mathbb{Z}$ -linear combination of characteristic functions $\mathrm{ch}(x_i U)$. For each of these terms, if we set $U_i = x_i U x_i^{-1}$, then by hypothesis we have $W \subseteq V_i := H \cap U_i$, and we can consider the composition of maps

$$\begin{aligned} H_{\mathrm{mot}}^2(Y_H(W), \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) &\xrightarrow{(\mathrm{pr}_{V_i}^W)_*} H_{\mathrm{mot}}^2(Y_H(V_i), \mathcal{H}_{\mathbb{Q}}^{c,d}(2)) \\ &\xrightarrow{\iota_{U_i,*}^{[a,b,q,r]}} H_{\mathrm{mot}}^4(Y_G(U_i), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q)) \xrightarrow{x_i} H_{\mathrm{mot}}^4(Y_G(U), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q)). \end{aligned} \quad (11)$$

(Note that U_i may not be H -small, so $\iota_{U_i} : Y_H(V_i) \rightarrow Y_G(U_i)$ may not be a closed immersion, but it is still a finite morphism of smooth varieties and this suffices to define the pushforward map $\iota_{U_i,*}^{[a,b,q,r]}$.)

Theorem 8.4.3. *The Lemma–Eisenstein classes satisfy the following norm-compatibility relations as m and n vary:*

(i) *For $n \geq 1$, we have*

$$(\mathrm{pr}_{K_G(M,p^m,p^n)}^{K_G(M,p^m,p^{n+1})})_* (z_{M,m,n+1}^{[a,b,q,r]}) = z_{M,m,n}^{[a,b,q,r]}.$$

(ii) *For $m \geq 0$, we have*

$$(\mathrm{pr}_{K_G(M,p^m,p^n)}^{K_G(M,p^{m+1},p^n)})_* (z_{M,m+1,n}^{[a,b,q,r]}) = \begin{cases} \frac{U'(p)}{p^q} & \text{if } m \geq 1 \\ \frac{U'(p)}{p^q} - 1 & \text{if } m = 0 \end{cases} \cdot z_{M,m,n}^{[a,b,q,r]}.$$

Here $U'(p) \in \mathcal{H}(G_p)$ is given by the $K_{G_p}(p^m, p^n)$ -double coset of $\begin{pmatrix} p^{-1} & & & \\ & p^{-1} & & \\ & & 1 & \\ & & & 1 \end{pmatrix}$.

Proof. Part (i) is immediate from the definition of the classes, since the sum of the translates of $\xi_{M,m,n+1}$ over $K_G(M, p^m, p^n)/K_G(M, p^m, p^{n+1})$ is $\xi_{M,m,n}$.

For part (ii), we note that the Hecke-algebra elements $\xi_{M,m,n}$ and $\xi_{M,m+1,n}$ are identical outside p , as are the Schwartz functions $\phi_{M,m,n}$ and $\phi_{M,m+1,n}$. So we need to compare two values of a map on $\mathcal{S}(\mathbb{Q}_p^2, \mathbb{Q})^{\otimes 2} \otimes \mathcal{H}(G(\mathbb{Q}_p))$ (given by tensoring with the common away-from- p parts and applying $\mathcal{L}\mathcal{E}$). It clearly suffices to check the equality after tensoring with $\mathbb{C}$, which puts us in a position where we may apply Proposition 3.10.4 (for $\ell = p$). If we assume that the parameters t are chosen identically for the two elements, then the proposition shows that

$$\begin{aligned} (\mathrm{pr}_{K_G(M,p^m,p^n)}^{K_G(M,p^{m+1},p^n)})_* \mathcal{L}\mathcal{E}(\phi_{M,m+1,n} \otimes \xi_{M,m+1,n}) \\ = \left\{ \begin{array}{l} \frac{1}{p} U'(p) \\ \frac{1}{p-1} (U'(p) - 1) \end{array} \right\} \cdot \mathcal{L}\mathcal{E}(\phi_{M,m,n} \otimes \xi_{M,m,n}) \end{aligned}$$

as elements of $H_{\text{mot}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))[-q]$. The factor of $\frac{1}{p}$ (resp. $\frac{1}{p-1}$) is cancelled out by the factors $\frac{1}{\text{Vol}(W)}$, since the subgroups W corresponding to the classes at levels Mp^{m+1} and Mp^m differ in volume by exactly this factor. Finally, the twist $[-q]$ gives a factor of p^q . ■

8.4.6. Integral p -adic étale classes. We now treat questions of integrality. We choose integers $c_1, c_2 > 1$ satisfying the following list of conditions:

- The c_i are coprime to $6p \prod_{\ell \in S} \ell$.
- Our chosen vector $\underline{\phi}_S \in \mathcal{S}(\mathbb{Q}_S^2, \mathbb{Z})^{\otimes 2}$ is preserved by the action of the elements $((c_1 \ 1), (c_1 \ 1)^{-1})$ and $((c_2 \ 1), (c_2 \ 1)^{-1})$ of $(\text{GL}_2 \times \text{GL}_2)(\mathbb{Q}_S)$. (Note that these elements are not in H .)
- For each $\ell \in S$, the subgroup K_ℓ is normalised by the elements $\begin{pmatrix} 1 & & & \\ & c_1 & & \\ & & 1 & \\ & & & c_1 \end{pmatrix}$ and $\begin{pmatrix} c_2 & & & \\ & 1 & & \\ & & c_2 & \\ & & & 1 \end{pmatrix}$ of $G(\mathbb{Q}_\ell)$.

(The last two conditions can, of course, always be achieved by taking c_1 and c_2 to be sufficiently close ℓ -adically to 1, for all $\ell \in S$.)

Recall the alternative local data $\xi'_{M,m,n}, \underline{\phi}'_{M,m,n}, W'$ introduced at the end of §8.4.

Definition 8.4.4. For $n \geq \max(m, 1)$, let

$$c_{1,c_2} \mathcal{Z}_{\text{ét}, M, m, n}^{[a,b,q,r]} \in H_{\text{ét}}^4(Y'_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q))$$

be the class defined using the alternative local data $\xi'_{M,m,n}, \underline{\phi}'_{M,m,n}, W'$ in place of their non-dashed versions, and substituting for $\text{Eis}_{\text{mot}, \underline{\phi}}^{c,d}$ the integral étale Eisenstein classes $c_{1,c_2} \text{Eis}_{\text{ét}, \underline{\phi}}^{c,d}$.

To see that this is well-defined, we use the explicit description of the Lemma–Eisenstein map as a sum of pushforward maps as in (11). Since $\xi'_{M,m,n}$ is a $\mathbb{Z}$ -linear combination of cosets $x_i U$ (where $U = K'_G(M, p^m, p^n)$) with $x_i \in G(\mathbb{A}_f^{(p)} \times \mathbb{Z}_p)$, the maps $\iota_{x_i U x_i^{-1}, *}^{[a,b,q,r]}$ are well-defined on étale cohomology with coefficients in the integral sheaf $\mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)$. (Note that $\xi_{M,m,n}$ is not supported in $G(\mathbb{A}_f^{(p)} \times \mathbb{Z}_p)$, which is why we need to introduce the alternative data.)

Definition 8.4.5. For $n \geq \max(m, 1)$, let $s_m : Y'_G(M, p^m, p^n) \rightarrow Y_G(M, p^m, p^n)$ be the map given by the action of $\begin{pmatrix} p^m & & & \\ & p^m & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \in G(\mathbb{Q}_p)$; and let $s_{m,\#} : \mathcal{D}_{\mathbb{Z}_p}^{a,b} \rightarrow s_m^*(\mathcal{D}_{\mathbb{Z}_p}^{a,b})$ be the morphism of sheaves given by the action of $\begin{pmatrix} p & & & \\ & p & & \\ & & 1 & \\ & & & 1 \end{pmatrix}^{-m}$ on the representation $D_{\mathbb{Z}_p}^{a,b}$.

See [15, §6.1]. The morphism s_m is well-defined on the Shimura varieties, since

$$\begin{pmatrix} p^m & & & \\ & p^m & & \\ & & 1 & \\ & & & 1 \end{pmatrix}^{-1} K'_{G_p}(p^m, p^n) \begin{pmatrix} p^m & & & \\ & p^m & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \subseteq K_{G_p}(p^m, p^n).$$

To construct the morphism $s_{m,\#}$ of integral coefficient sheaves, we note that the representation $D^{a,b}$ of G has all weights ≤ 0 for the torus T' of §2, so the action of $G(\mathbb{Z}_p)$ on $D_{\mathbb{Z}_p}^{a,b}$ extends to an action of the monoid generated by $G(\mathbb{Z}_p)$ and $\begin{pmatrix} p & & \\ & p & \\ & & 1 \end{pmatrix}^{-1}$.

Proposition 8.4.6. *If (c_1, c_2) satisfy the above conditions, then for any m, n, M as in the previous section with M chosen coprime to c_1 and c_2 , there is a class*

$$c_1, c_2 z_{\text{ét}, M, m, n}^{[a, b, q, r]} \in H_{\text{ét}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a, b}(3 - q))$$

whose image in the cohomology of $\mathcal{D}_{\mathbb{Q}_p}^{a, b}(3 - q)$ is

$$p^{mq} \left(c_1^2 - c_1^{-(a-q+b-r)} \begin{pmatrix} 1 & & \\ & c_1 & \\ & & 1 \end{pmatrix}^{-1} \right) \left(c_2^2 - c_2^{-(a-q+r)} \begin{pmatrix} c_2 & & \\ & 1 & \\ & & c_2 \end{pmatrix}^{-1} \right) r_{\text{ét}}(z_{M, m, n}^{[a, b, q, r]}),$$

where the matrices on the left-hand side are understood as elements of $\prod_{\ell | MpS} G(\widehat{\mathbb{Z}})$ acting on $Y_G(M, p^m, p^n)$ by right translation.

Remark 8.4.7. If $a = b = 0$, then $c_1, c_2 z_{\text{ét}, M, m, n}^{[0, 0, 0, 0]}$ is the image of a class $c_1, c_2 z_{M, m, n}^{[0, 0, 0, 0]}$ in the motivic cohomology with $\mathbb{Z}$ coefficients. However, we do not know how to define this group for $a, b > 0$. $\diamond$

Proof of Proposition 8.4.6. Assume for the moment that $m \leq n$, and define

$$c_1, c_2 z_{\text{ét}, M, m, n}^{[a, b, q, r]} := s_{m,*}(c_1, c_2 \mathcal{Z}_{\text{ét}, M, m, n}^{[a, b, q, r]}),$$

where the action of s_m on the coefficients is given by $s_{m,\#}$. By definition, this has coefficients in $\mathcal{D}_{\mathbb{Z}_p}^{a, b}(3 - q)$, so it remains to verify that is related to $z_{M, m, n}^{[a, b, q, r]}$ in the stated manner.

A simple check shows that

$$c_1, c_2 \tilde{z}_{\text{ét}, M, m, n}^{[a, b, q, r]} = p^{mq} r_{\text{ét}}(\tilde{z}_{M, m, n}^{[a, b, q, r]})$$

where $\tilde{z}$ is the class obtained in the same way as z , with the Schwartz function ϕ replaced by

$$(c_1^2 - c_1^{-(a-q+b-r)} ((\begin{smallmatrix} c_1 & 0 \\ 0 & c_1 \end{smallmatrix}), \text{id})^{-1}) (c_2^2 - c_2^{-(a-q+r)} (\text{id}, (\begin{smallmatrix} c_2 & 0 \\ 0 & c_2 \end{smallmatrix}))^{-1}) \underline{\phi}.$$

(The factor p^{mq} appears because we ignored the twist $[-q]$ in the definition of $s_{m,\#}$.) However, our assumptions on the c_i imply that

$$\begin{aligned} & (c_1^2 - c_1^{-(a-q+b-r)} ((\begin{smallmatrix} c_1 & 0 \\ 0 & c_1 \end{smallmatrix}), \text{id})^{-1}) (c_2^2 - c_2^{-(a-q+r)} (\text{id}, (\begin{smallmatrix} c_2 & 0 \\ 0 & c_2 \end{smallmatrix}))^{-1}) \underline{\phi} \\ &= (c_1^2 - c_1^{-(a-q+b-r)} ((\begin{smallmatrix} 1 & 0 \\ 0 & c_1 \end{smallmatrix}), (\begin{smallmatrix} c_1 & 0 \\ 0 & 1 \end{smallmatrix}))^{-1}) (c_2^2 - c_2^{-(a-q+r)} ((\begin{smallmatrix} c_2 & 0 \\ 0 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & c_2 \end{smallmatrix}))^{-1}) \underline{\phi} \end{aligned}$$

and in this second formula, all the elements acting are in H , normalise our level groups, and commute with η , so we can pull them through the equivariance properties of the Lemma–Eisenstein map to obtain the result.

Finally, we remove the restriction $m \leq n$: if $n < m$, then we simply define $c_{1,c_2} z_{\text{ét},M,m,n}^{[a,b,q,r]}$ to be the pushforward of $c_{1,c_2} z_{\text{ét},M,m,n'}^{[a,b,q,r]}$ for any integer $n' \geq m$. This is independent of the choice of n' , as is easily seen, and using Theorem 8.4.3(i) and the preceding argument with n' in place of n , we see that it has the required properties. ■

Remark 8.4.8. It follows from Theorem 8.4.3 that the Lemma–Eisenstein classes $c_{1,c_2} z_{\text{ét},M,m,n}^{[a,b,q,r]}$ and their variants $c_{1,c_2} \mathcal{Z}_{\text{ét},M,m,n}^{[a,b,q,r]}$ satisfy norm-compatibility relations in both m and n after tensoring with $\mathbb{Q}_p$. However, one can check that these norm relations actually hold *integrally*, without needing to quotient out by the torsion subgroup of the étale cohomology group. This is not obvious from the proofs we have given, but can easily be verified after carefully unwinding the normalisation factors. ◇

9. Moment maps and p -adic interpolation

We now study the interpolation of the étale Euler system classes, for varying values of the parameters (a, b, q, r) . Our goal is Theorem 9.6.4, which shows that these classes can all be obtained as specialisations of a single class “at infinite level”.

9.1. Interpolation of the GL_2 Eisenstein classes

We begin by recalling a theorem of Kings [14], which will be the fundamental input for our p -adic interpolation results. In this section, let us fix an arbitrary open compact subgroup $K^{(p)} \subset \text{GL}_2(\mathbb{A}_f^{(p)})$, and for $n \geq 1$, write $K_n = K^{(p)} \times \{g \in \text{GL}_2(\mathbb{Z}_p) : g \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{p^n}\}$. Let us assume, by shrinking $K^{(p)}$ if necessary, that K_1 is sufficiently small (and hence so is K_n for all $n \geq 1$).

We also choose a finite set of primes Σ containing p and all primes where $K^{(p)}$ is ramified, so the modular curves have models $Y(K_n)_\Sigma$ over $\mathbb{Z}[1/\Sigma]$ for all n .

Remark 9.1.1. Working with integral models is necessary here, because continuous étale cohomology for $\mathbb{Q}$ -varieties does not necessarily commute with inverse limits, but this problem does not arise for finite-type $\mathbb{Z}$ -schemes such as the $Y(K_n)_\Sigma$. ◇

Definition 9.1.2. We define

$$H_{\text{Iw}}^1(Y(K_\infty)_\Sigma, \mathbb{Z}_p(1)) := \varprojlim_{s \geq 1} H_{\text{ét}}^1(Y(K_s)_\Sigma, \mathbb{Z}_p(1))$$

where the inverse limit is with respect to the pushforward maps.

If $\mathcal{H}_n^k$ denotes the mod p^n reduction of the sheaf $\mathcal{H}_{\mathbb{Z}_p}^k$ on $Y(K_n)$ (cf. §7.2), then we have a canonical section

$$e_{k,n} = (e_1)^k \in H_{\text{ét}}^0(Y(K_n)_\Sigma, \mathcal{H}_n^k).$$

Hence, for any $n \geq 1$, we have a map

$$\text{mom}_n^k : H_{\text{Iw}}^1(Y(K_\infty)_\Sigma, \mathbb{Z}_p(1)) \rightarrow H_{\text{ét}}^1(Y(K_n), \mathcal{H}_{\mathbb{Z}_p}^k(1)),$$

mapping $(g_s)_{s \geq 1}$ to the restriction to the generic fibre $Y(K_n) \subset Y(K_n)_\Sigma$ of the element

$$(\mathrm{pr}_{K_n}^{K_s})_*(g_s \cup e_{k,s})_{s \geq n} \in \varprojlim_{s \geq n} H_{\mathrm{\acute{e}t}}^1(Y(K_n)_\Sigma, \mathcal{H}_s^k(1)) = H_{\mathrm{\acute{e}t}}^1(Y(K_n)_\Sigma, \mathcal{H}_{\mathbb{Z}_p}^k(1)).$$

Definition 9.1.3. Let ϕ be a $\mathbb{Z}_p$ -valued Schwartz function on $(\mathbb{A}_f^{(p)})^2$, stable under $K^{(p)}$; and let $\phi_s = \phi \otimes \mathrm{ch}(p^s \mathbb{Z}_p \times (1 + p^s \mathbb{Z}_p))$. For $n \geq 1$, and $c > 1$ coprime to $6p$ and to all primes where ϕ is ramified, we define

$${}_c \mathcal{E} \mathcal{I}_\phi = ({}_c g_{\phi_s})_{s \geq 1} \in H_{\mathrm{Iw}}^1(Y(K_\infty)_\Sigma, \mathbb{Z}_p(1)).$$

The following theorem, which will be fundamental for our p -adic interpolation results later in this paper, shows that the Siegel units interpolate Eisenstein classes of all weights via these moment maps:

Theorem 9.1.4 (Kings). *For all integers $k \geq 0$ and $n \geq 1$, we have*

$$\mathrm{mom}_n^k({}_c \mathcal{E} \mathcal{I}_{\phi,n}) = {}_c \mathrm{Eis}_{\mathrm{\acute{e}t}, \phi_n}^k$$

as elements of $H_{\mathrm{\acute{e}t}}^1(Y(K_n), \mathcal{H}_{\mathbb{Z}_p}^k(1))$.

Proof. This is a generalisation of [15, Theorems 4.4.4 & 4.5.1], which is the case where ϕ is the characteristic function of $(0, 1) + N(\widehat{\mathbb{Z}}^{(p)})^2$ for some integer N . The general case can be recovered from this using the action of the group $J = \prod_{\ell \in \Sigma - \{p\}} \mathrm{GL}_2(\mathbb{Q}_\ell)$, since both the Siegel units and Eisenstein classes depend J -equivariantly on ϕ , and the moment map clearly commutes with the action of J (as it acts trivially on $e_{k,s}$). ■

Of course, this argument carries over readily to the modular varieties for H : if we fix a small enough prime-to- p level $K_H^{(p)}$ and let $K_{H,n} = K_H^{(p)} \times K_{H,p,1}(p^n)$, then we obtain moment maps

$$\mathrm{mom}_{H,n}^{c,d} : H_{\mathrm{Iw}}^2(Y_H(K_{H,\infty})_\Sigma, \mathbb{Z}_p(2)) \rightarrow H_{\mathrm{\acute{e}t}}^2(Y_H(K_{H,n}), \mathcal{H}_{\mathbb{Z}_p}^{c,d}(2)),$$

for any $n \geq 1$ and integers $c, d \geq 0$; and there is a class ${}_{c_1, c_2} \mathcal{E} \mathcal{I}_\phi$ in H_{Iw}^2 , for any $\phi \in \mathcal{S}(\mathbb{A}_f^{(p)2}, \mathbb{Z}_p)^{\otimes 2}$ stable under $K_H^{(p)}$ and unramified at the primes dividing $c_1 c_2$, whose images under $\mathrm{mom}_n^{c,d}$ are the Eisenstein classes ${}_{c_1, c_2} \mathrm{Eis}_{\mathrm{\acute{e}t}, \phi_n}^{c,d}$.

9.2. Moment maps for G

For the group $G = \mathrm{GSp}_4$ we have analogous moment maps, as we shall now explain. As in the GL_2 case, we fix an arbitrary subgroup $K_G^{(p)} \subset G(\mathbb{A}_f^{(p)})$ unramified outside Σ , and write $K_{G,n} = K_G^{(p)} \times K_{G,p,1}(p^n)$. We assume that $K_{G,n}$ is sufficiently small for all $n \geq 1$.

Proposition 9.2.1. *Let $d^{[a,b,q,r]}$ be the image of $v^{[a,b,q,r]} \in V_{\mathbb{Z}}^{a,b}$ in $D_{\mathbb{Z}}^{a,b} = V_{\mathbb{Z}}^{a,b} \otimes \mu^{-(2a+b)}$; and let $d_n^{[a,b,q,r]}$ be its reduction modulo p^n . Then the vectors $d_n^{[a,b,0,r]}$, for $0 \leq r \leq b$, are stable under $K_{G,p,1}(p^n)$.*

Proof. We recall that the vectors $v^{[a,b,0,r]}$ lie in the highest- T' -weight subspace of $V^{a,b}$, where T' is the torus $\begin{pmatrix} x & & \\ & x & \\ & & 1 \end{pmatrix}$. Hence they are fixed by the unipotent radical N_S of the Siegel parabolic, and T' acts on them as $x \mapsto x^{2a+b}$. Thus the twists $d^{[a,b,0,r]}$ are fixed by T' and by N_S , and the same holds for their reductions modulo p . ■

Definition 9.2.2. For $n \geq 1$, and any integers $0 \leq q \leq a$ and $0 \leq r \leq b$, we define the moment map $\text{mom}_{G,n}^{[a,b,q,r]}$ as the following composition of maps:

$$\begin{aligned} H_{\text{lw}}^*(Y_G(K_{G,\infty})_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3)) &\xrightarrow{\cong} \varprojlim_s H_{\text{ét}}^*(Y_G(K_{G,s})_\Sigma, \mathcal{D}_s^{q,0}(3)) \\ &\rightarrow \varprojlim_s H_{\text{ét}}^*(Y_G(K_{G,s})_\Sigma, (\mathcal{D}_s^{q,0} \otimes \mathcal{D}_s^{a-q,b})(3)) \\ &\rightarrow \varprojlim_s H_{\text{ét}}^*(Y_G(K_{G,s})_\Sigma, \mathcal{D}_s^{a,b}(3)) \\ &\rightarrow \varprojlim_s H_{\text{ét}}^*(Y_G(K_{G,n})_\Sigma, \mathcal{D}_s^{a,b}(3)) \\ &\xrightarrow{\cong} H_{\text{ét}}^*(Y_G(K_{G,n})_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3)) \\ &\rightarrow H_{\text{ét}}^*(Y_G(K_{G,n})_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3)). \end{aligned}$$

Here the second arrow is cup-product with $d_s^{[a-q,b,0,r]} \in H_{\text{ét}}^0(Y_G(K_{G,s})_\Sigma, \mathcal{D}_s^{a-q,0})$; the third arrow is given by the Cartan product; the fourth by projection to level n ; and the final one by restriction to the generic fibre.

Remark 9.2.3. This construction also has an interpretation in terms of sheaves of measures as in [14]. Suppose $q = 0$ for simplicity. One finds that $H_{\text{lw}}^*(Y_G(K_{G,\infty})_\Sigma, \mathbb{Z}_p(3)) = H_{\text{ét}}^*(Y_G(K_{G,n})_\Sigma, \Lambda(3))$, where Λ is the sheaf on $Y_G(K_{G,n})_\Sigma$ given by $\varprojlim_s (\text{pr}_{K_{G,n}}^{K_{G,s}})_*(\mathbb{Z}_p)$. We can interpret Λ as the sheaf corresponding to the profinite $\mathbb{Z}_p[K_{G,n}]$ -module of $\mathbb{Z}_p$ -valued measures on the quotient $X_n = K_{G,n}/K_{G,\infty}$; in this optic, the moment map is given by the morphism of sheaves

$$\Lambda(X_n) \rightarrow D_{\mathbb{Z}_p}^{a,b}, \quad \mu \mapsto \int_{X_n} g \cdot d^{[a,b,0,r]} d\mu(g). \quad \diamond$$

Proposition 9.2.4. The Hecke operator $U'(p)$ is well-defined as an endomorphism of the inverse limit $H_{\text{lw}}^*(Y_G(K_{G,\infty})_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3))$, and the moment map $\text{mom}_{G,n}^{[a,b,q,r]}$ is compatible with the actions of $U'(p)$ and of $\prod_{\ell \in \Sigma - \{p\}} G(\mathbb{Q}_\ell)$ on both sides.

Proof. Easy check, compare e.g. [15, Remark 4.5.3]. ■

9.3. Compatibility with the moment maps for H

We shall now consider compatibility of these constructions between G and H . We take $K_H^{(p)} = K_G^{(p)} \cap H(\mathbb{A}_f^{(p)})$, so we have maps $\iota_n : Y_H(K_{H,n}) \rightarrow Y_G(K_{G,n})$ for all $n \geq 1$, and write ι_∞ for the collection $(\iota_s)_{s \geq 1}$. As before, let $(c, d) = (a + b - q - r, a - q + r)$.

Proposition 9.3.1. *There is a commutative diagram*

$$\begin{array}{ccc} H_{\mathrm{Iw}}^2(Y_H(K_{H,\infty})_\Sigma, \mathbb{Z}_p(2)) & \xrightarrow{\iota_{\infty,*} \circ \mathrm{br}^{[q,0,q,0]}} & H_{\mathrm{Iw}}^4(Y_G(K_{G,\infty})_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3-q)) \\ \mathrm{mom}_{H,n}^{c,d} \downarrow & & \downarrow \mathrm{mom}_{G,n}^{[a,b,q,r]} \\ H_{\mathrm{\acute{e}t}}^2(Y_H(K_{H,n}), \mathcal{H}_{\mathbb{Z}_p}^{c,d}(2)) & \xrightarrow{\iota_{n,*} \circ \mathrm{br}^{[a,b,q,r]}} & H_{\mathrm{\acute{e}t}}^4(Y_G(K_{G,n}), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)) \end{array}$$

Proof. After much unwinding, this reduces to the assertion that the modulo p^s reduction of $\mathrm{br}^{[a,b,q,r]} : \mathcal{H}_{\mathbb{Z}_p}^{c,d} \rightarrow \iota^* \mathcal{D}_{\mathbb{Z}_p}^{a,b}$ maps the section $(e_{1,s})^c \boxtimes (f_{1,s})^d$ over $K_{H,s}$ to the pullback of $d_s^{[q,0,q,0]} \cdot d_s^{[a-q,b,0,r]} = d_s^{[a,b,q,r]}$, which is true by the construction of the branching maps. $\blacksquare$

Remark 9.3.2. In [15], the analogous statement for the $\mathrm{GL}_2 \times \mathrm{GL}_2$ -moment maps (Lemma 6.3.1) gives rise to a binomial factor; so using the Cartan product simplifies matters considerably. $\diamond$

9.4. Application to Lemma–Eisenstein classes

We now return to the situation considered in §8.4. We can apply the machinery of the previous section with $K_G^{(p)}$ taken to be the product $K_S \times \prod_{\ell \notin S \cup \{p\}} G(\mathbb{Z}_\ell)$, so the $K_{G,n}$ of the previous section is $K_{G,1}(p^n)$, and we obtain moment maps

$$\mathrm{mom}_{G,n}^{[a,b,q,r]} : H_{\mathrm{Iw}}^4(Y_{G,1}(p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3-q)) \rightarrow H_{\mathrm{\acute{e}t}}^4(Y_{G,1}(p^n), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)),$$

for any $\Sigma \supseteq S \cup \{p\}$. More generally, if we take any $m \geq 0$ and any squarefree $M \geq 1$ whose prime divisors lie in $\Sigma - (S \cup \{p\})$, the same construction also gives maps

$$H_{\mathrm{Iw}}^4(Y_G(M, p^m, p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3-q)) \rightarrow H_{\mathrm{\acute{e}t}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)) \quad (12a)$$

for any $n \geq 1$, and

$$H_{\mathrm{Iw}}^4(Y'_G(M, p^m, p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3-q)) \rightarrow H_{\mathrm{\acute{e}t}}^4(Y'_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)) \quad (12b)$$

for any $n \geq \max(m, 1)$, which we also denote by $\mathrm{mom}_{G,n}^{[a,b,q,r]}$.

Proposition 9.4.1. *For each $q \geq 0$, there exists a class*

$${}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q \in H_{\mathrm{Iw}}^4(Y'_G(M, p^m, p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q, 0}(3 - q))$$

such that

$$\mathrm{mom}_{G, n}^{[a, b, q, r]}({}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q) = {}_{c_1, c_2} \mathcal{Z}_{\mathrm{\acute{e}t}, M, m, n}^{[a, b, q, r]}$$

for all integers a, b, r, n with $a \geq q$, $0 \leq r \leq b$ and $n \geq \max(m, 1)$.

Remark 9.4.2. Strictly speaking the elements ${}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q$ depend also on Σ , but they are easily seen to be compatible with the natural maps given by enlarging Σ , so we will suppress this from the notation. $\diamond$

Proof of Proposition 9.4.1. We define ${}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q$ to be $({}_{c_1, c_2} \mathcal{Z}_{\mathrm{\acute{e}t}, M, m, n}^{[q, 0, q, 0]})_{n \geq \max(1, m)}$, which is norm-compatible by the same argument as in Theorem 8.4.3(i) (and Remark 8.4.8).

By construction, there is a finite set of integers a_i and $x_i \in G(\mathbb{A}_f)$, independent of m and n , such that

$$\xi'_{M, m, n} = \sum_i a_i \mathrm{ch}(x_i K'_G(M, p^m, p^n)).$$

We can therefore write

$$\begin{aligned} {}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q &= \sum_i a_i (\iota_{x_i} K'_G(M, p^m, p^\infty)_* \circ \mathrm{br}^{[q, 0, q, 0]})_{c_1, c_2} \mathcal{E} \mathcal{I}_\phi, \\ {}_{c_1, c_2} \mathcal{Z}_{\mathrm{\acute{e}t}, M, m, n}^{[a, b, q, r]} &= \sum_i a_i (\iota_{x_i} K'_G(M, p^m, p^n)_* \circ \mathrm{br}^{[a, b, q, r]})_{c_1, c_2} \mathrm{Eis}_{\phi_n}^{[c, d]}. \end{aligned}$$

All the x_i have the same p -component, namely $\eta_{p, 0}$; this acts trivially on the vector $d^{[a-q, b, 0, r]}$, and hence commutes with the moment map. We know that $\mathrm{Eis}_{\phi_n}^{[c, d]}$ is the image of $\mathcal{E} \mathcal{I}_\phi$ under $\mathrm{mom}_{H, n}^{c, d}$ by Theorem 9.1.4, and the commutative diagram of Proposition 9.3.1 (taking for $K^{(p)}$ the prime-to- p part of $x_i^{-1} K'_G(M, p^m, p^n) x_i$, for each i) shows that the image of each summand at level ∞ under $\mathrm{mom}_{G, n}^{[a, b, q, r]}$ coincides with the corresponding summand at level n . $\blacksquare$

Corollary 9.4.3. *For each $q \geq 0$, there exists a class*

$${}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q \in H_{\mathrm{Iw}}^4(Y_G(M, p^m, p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{q, 0}(3 - q))$$

such that

$$\mathrm{mom}_{G, n}^{[a, b, q, r]}({}_{c_1, c_2} \mathcal{Z}_{\mathrm{Iw}, M, m}^q) = {}_{c_1, c_2} \mathcal{Z}_{\mathrm{\acute{e}t}, M, m, n}^{[a, b, q, r]}$$

for all integers a, b, r, n with $a \geq q$, $0 \leq r \leq b$ and $n \geq 1$.

Proof. The moment maps of (12a) and (12b) are compatible with respect to the push-forwards $(s_m)_*$, because the action of $\mathrm{diag}(p^{-1}, p^{-1}, 1, 1)$ on $D_{\mathbb{Z}_p}^{a-q, b}$ fixes the vector $d^{[a-q, b, 0, r]}$. So for $n \geq m$ the corollary follows from Proposition 9.4.1 by applying $(s_m)_*$ to both sides; and since both sides of the desired formula are norm-compatible in n , the result follows for $n < m$ also. $\blacksquare$

9.5. Cyclotomic twists

We now consider the more difficult problem of interpolating the classes of the previous sections as the parameter q varies, analogously to [15, Theorem 6.2.4] in the Rankin–Selberg setting. Our main technical result will be the following:

Proposition 9.5.1. *For each $m \geq 1$ and $q \geq 0$, we have*

$$c_{1,c_2} z_{\mathrm{Iw},M,m}^q = (-2)^q \cdot c_{1,c_2} z_{\mathrm{Iw},M,m}^0 \cup (d_m^{[q,0,0]} \otimes \zeta_{p^m}^{-q}) \bmod p^m.$$

Here $\zeta_{p^m} \in H_{\mathrm{et}}^0(Y_G(1, p^m, p^m), \mathbb{Z}/p^m(1))$ is the canonical p^m -th root of unity given by the isomorphism (9). In an attempt to restrain the excessive proliferation of indices in our notations, we shall give the arguments assuming $M = 1$, and drop M from the subscripts throughout the remainder of the section; the case of general M can be handled similarly (using the decomposition of $\xi'_{M,m,n}$ as a finite sum of characteristic functions of cosets, as in the proof of Proposition 9.4.1). In this case, we have

$$c_{1,c_2} \mathcal{Z}_{\mathrm{Iw},m}^q := (\eta_* \circ \iota_{\infty,*} \circ \mathrm{br}^{[q,0,q,0]})(c_{1,c_2} \mathcal{E} \mathcal{I}_{\phi'}) \in H_{\mathrm{Iw}}^4(Y'_G(p^m, p^\infty), \mathcal{D}_{\mathbb{Z}_p}^{q,0}(3-q)).$$

The branching map $\mathrm{br}^{[q,0,q,0]}$ appearing in the above constructions is given by mapping $1 \in \mathbb{Z}_p$ to the $H(\mathbb{Z}_p)$ -invariant element $d^{[q,0,q,0]} \otimes \zeta^{-q} \in D_{\mathbb{Z}_p}^{q,0}(-q)$, where ζ denotes a basis of the multiplier representation μ of G . After reducing modulo p^m , this element is invariant under a larger group:

Proposition 9.5.2. *The modulo p^m reduction $d_m^{[q,0,q,0]}$ is stable under $K'_{G_p}(p^m, p^\infty) \subset G(\mathbb{Z}_p)$.*

Proof. Since $K'_{G_p}(p^m, p^\infty)$ is contained in the principal congruence subgroup modulo p^m , it acts trivially on $\mathcal{D}_m^{a,b}$ for any a, b . $\blacksquare$

It follows that we may write

$$c_{1,c_2} \mathcal{Z}_{\mathrm{Iw},m}^q = [(\eta_* \circ \iota_{\infty,*})(c_{1,c_2} \mathcal{E} \mathcal{I}_{\phi'})] \cup (\eta_* d_m^{[q,0,q,0]} \otimes \zeta^{-q}), \quad (13)$$

where $\eta_* d_m^{[q,0,q,0]} \in H^0(Y'_G(p^m, p^\infty), \mathcal{D}_m^{q,0})$.

Proposition 9.5.3. *We have*

$$s_{m,\#}(\eta_* d_m^{[q,0,q,0]}) = (-2)^q s_m^*(d_m^{[q,0,0,0]})$$

as sections of $s_m^*(\mathcal{D}_m^{q,0})$.

Proof. We may decompose $D_{\mathbb{Z}_p}^{q,0}$ as a direct sum of its eigenspaces for the action of the torus T' , which all have weights ≤ 0 . On all eigenspaces other than the weight-0 eigenspace, the map $s_{m,\#}$ is zero, since $\mathrm{diag}(p, p, 1, 1)^{-m}$ acts as a positive power of p^m , which annihilates the module $D_m^{q,0}$. Hence $s_{m,\#}$ factors through projection to the highest-weight space relative to T' . So we need to compute the projection of $\eta_*(d_m^{[q,0,q,0]}) = (\eta^{-1})^*(d_m^{[q,0,q,0]})$ to this weight space. This is precisely the situation of Lemma 4.4.1 (with $h = -1$ in the notation of the lemma), which gives the result above. $\blacksquare$

Proposition 9.5.1 follows immediately from this, by applying $s_{m,*}$ to both sides of (13).

9.6. Projection to the ordinary part

We now define a limiting element in which m (as well as n) goes to ∞ . We set

$$H_{\text{Iw}}^4(Y_G(M, p^\infty, p^\infty)_\Sigma, \mathbb{Z}_p(3)) = \varprojlim_m H_{\text{Iw}}^4(Y_G(M, p^m, p^\infty)_\Sigma, \mathbb{Z}_p(3)).$$

On this module, there is an action of the ordinary idempotent $e'_{\text{ord}} = \lim_{k \rightarrow \infty} U'(p)^{k!}$.

Remark 9.6.1. The fact that this limit exists, and is an idempotent, follows from the corresponding statements for étale cohomology at finite levels, for which see [42]. Note that Tilouine and Urban define multiple ordinary idempotents, one for each standard parabolic subgroup; ours is the one associated to the Siegel parabolic P_S . $\diamond$

Definition 9.6.2. We set

$$c_1, c_2 z_{\text{Iw}, M} = (U'(p)^{-t} e'_{\text{ord}} \cdot c_1, c_2 z_{\text{Iw}, M, m}^0)_{m \geq 1}.$$

This is a well-defined element of $H_{\text{Iw}}^4(Y_G(M, p^\infty, p^\infty)_\Sigma, \mathbb{Z}_p(3))$, since $U'(p)$ is invertible on the image of e'_{ord} , and the terms in the limit are norm-compatible by Theorem 8.4.3(ii).

Definition 9.6.3. For integers $m \geq 0, n \geq 1, 0 \leq r \leq b, a \geq 0$, and $q \in \mathbb{Z}$, we define

$$\text{mom}_{G, m, n}^{[a, b, q, r]} : H_{\text{Iw}}^4(Y_G(M, p^\infty, p^\infty)_\Sigma, \mathbb{Z}_p(3)) \rightarrow H_{\text{ét}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a, b}(3 - q))$$

by cup-product with $d^{[a, b, 0, r]} \otimes \zeta^{-q} \in H_{\text{ét}}^0(Y_G(M, p^\infty, p^\infty)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{a, b}(-q))$, where ζ is the canonical basis of $\mathbb{Z}_p(1)$ over $\text{Spec } \mathbb{Z}[\Sigma^{-1}, \zeta_{Mp^\infty}]$.

Note that we do *not* need to assume that q lies in the interval $\{0, \dots, a\}$ in order to define this moment map. However, when we do impose this additional assumption, we obtain compatibility with the preceding constructions:

Theorem 9.6.4. For integers $m \geq 0, n \geq 1, 0 \leq q \leq a$ and $0 \leq r \leq b$, we have

$$\text{mom}_{G, m, n}^{[a, b, q, r]}(c_1, c_2 z_{\text{Iw}, M}) = \frac{1}{(-2)^q} \begin{cases} U'(p)^{-m} & \text{if } m \geq 1 \\ 1 - \frac{p^q}{U'(p)} & \text{if } m = 0 \end{cases} \cdot e'_{\text{ord}}(c_1, c_2 z_{\text{ét}, M, m, n}^{[a, b, q, r]}).$$

Proof. We factor $d^{[a, b, 0, r]}$ as the Cartan product of $d^{[a-q, b, 0, r]}$ and $d^{[q, 0, 0, 0]}$. Proposition 9.5.1 shows that cup-product with $d^{[q, 0, 0, 0]} \otimes \zeta^{-q}$ sends $c_1, c_2 z_{\text{Iw}, M}$ to the inverse system

$$(-2)^{-q} (U'(p)^{-t} e'_{\text{ord}} \cdot c_1, c_2 z_{\text{Iw}, M, t}^q)_{t \geq 1}.$$

Projecting this to level m gives the element

$$\frac{1}{(-2)^q} \begin{cases} U'(p)^{-m} & \text{if } m \geq 1 \\ 1 - \frac{p^q}{U'(p)} & \text{if } m = 0 \end{cases} \cdot e'_{\text{ord}}(c_1, c_2 z_{\text{Iw}, M, m}^q).$$

(This is true by definition for $m \geq 1$, and the case $m = 0$ follows by computing the norm of the $m = 1$ element using the appropriate case of Theorem 8.4.3(ii).) Computing the image of this element under $\mathrm{mom}_{G,n}^{[a,b,q,r]}$ using Proposition 9.4.3 gives the result. ■

Remark 9.6.5. The module $e'_{\mathrm{ord}} H_{\mathrm{Iw}}^4(Y_G(M, p^\infty, p^\infty)_\Sigma, \mathbb{Z}_p(3))$ can be regarded as an interpolation of the Iwasawa cohomology of the Galois representations attached to p -ordinary Siegel modular forms with weights varying in a Hida family. Thus Theorem 9.6.4 can be interpreted as stating that our Euler system classes interpolate in Hida families. We have not pursued this viewpoint in the present paper for reasons of space, but we intend to revisit the topic of Hida-family variation in a future project. ◇

Corollary 9.6.6 (Cohomological triviality). *If $m \geq 1$ or $q \geq 1$, then $e'_{\mathrm{ord}}(c_1, c_2 z_{\mathrm{ét}, M, m, n}^{[a,b,q,r]})$ is in the kernel of the base-extension map*

$$H_{\mathrm{ét}}^4(Y_G(M, p^m, p^n), \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q)) \rightarrow H_{\mathrm{ét}}^4(Y_G(M, p^m, p^n)_{\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q))^{\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})}.$$

Proof. Using (9), for each M and each $n \geq 1$ we have

$$\begin{aligned} \lim_{\leftarrow m} H^0(\mathbb{Q}, H_{\mathrm{ét}}^4(Y_G(M, p^m, p^n)_{\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q))) \\ = \lim_{\leftarrow m} H^0(\mathbb{Q}(\zeta_{Mp^m}), H_{\mathrm{ét}}^4(Y_{G,1}(p^n)_{\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3-q))). \end{aligned}$$

This inverse limit is zero, by standard properties of Iwasawa cohomology (see e.g. [26, Proposition 8.3.5]).

For $m \geq 1$, it is immediate from Theorem 9.6.4 that the image of $e'_{\mathrm{ord}} \cdot c_1, c_2 z_{\mathrm{ét}, M, m, n}^{[a,b,q,r]}$ under the edge map lies in the image of this inverse limit, and hence is also zero. For $m = 0$, this argument shows that the class becomes cohomologically trivial after applying $1 - \frac{p^q}{U^r(p)}$, and for $q \geq 1$ this operator acts invertibly on the image of e'_{ord} . ■

10. Mapping to Galois cohomology

In this section, we will use the motivic and étale classes we have constructed above in order to define Galois cohomology classes in automorphic Galois representations. We begin by recalling a number of results (due to various authors) on Galois representations appearing in the cohomology of the Siegel varieties Y_G .

10.1. Automorphic representations of GSp_4

Let (k_1, k_2) be integers with $k_1 \geq k_2 \geq 3$, and let $(a, b) = (k_2 - 3, k_1 - k_2)$. There are exactly two unitary discrete series representations of $G(\mathbb{R})$ which are cohomological with coefficients in the algebraic representation $V^{a,b}$: the holomorphic discrete series representation Π_{k_1, k_2}^H , and a non-holomorphic generic discrete series representation Π_{k_1, k_2}^W . We

refer to these as the *discrete series representations of weight* (k_1, k_2) . The cuspidal automorphic representations with infinite component Π_{k_1, k_2}^H are precisely those generated by classical holomorphic Siegel modular forms of weight (k_1, k_2) .

Remark 10.1.1. The pair $\{\Pi_{k_1, k_2}^H, \Pi_{k_1, k_2}^W\}$ is an example of a local L -packet. $\diamond$

Definition 10.1.2. Let $\Pi = \Pi_f \otimes \Pi_\infty$ be a cuspidal automorphic representation of $G(\mathbb{A}_f)$ with Π_∞ a discrete series representation of weight (k_1, k_2) .

- We say Π is of *Saito–Kurokawa type* if $k_1 = k_2$ and there exists a Dirichlet character χ such that $\chi^2 = \omega_\Pi$, and a cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A})$ of central character ω_Π attached to some holomorphic newform of weight $2k_1 - 2$, such that for all but finitely many places v we have

$$L(\Pi_v, s) = L(\pi_v, s)L(\chi_v, s - 1/2)L(\chi_v, s + 1/2).$$

- We say Π is of *Yoshida type* if there is a pair (π_1, π_2) of cuspidal automorphic representations of $\mathrm{GL}_2(\mathbb{A})$, both with central character ω_Π , corresponding to two elliptic modular newforms of weights $r_1 = k_1 + k_2 - 2$ and $r_2 = k_1 - k_2 + 2$, such that for all but finitely many places v we have

$$L(\Pi_v, s) = L(\pi_{1,v}, s)L(\pi_{2,v}, s).$$

- Otherwise, we say Π is *non-endoscopic*.

Theorem 10.1.3 (Taylor, Weissauer, Urban, Xu). *Let Π be as in the previous definition, and suppose Π is non-endoscopic. Let S be the set of primes at which Π ramifies, and let $w = k_1 + k_2 - 3$.*

- (1) *The representation Π is one of a pair $\{\Pi^H, \Pi^W\} = \{\Pi_f \otimes \Pi_{k_1, k_2}^H, \Pi_f \otimes \Pi_{k_1, k_2}^W\}$ of cuspidal automorphic representations having the same finite part, both of which have multiplicity 1 in $L_{\mathrm{cusp}}^2(G(\mathbb{Q}) \backslash G(\mathbb{A}), \omega_\Pi)$.*
- (2) *For any prime $\ell \notin S$, the local representation Π_ℓ is an unramified principal series representation.*
- (3) *For $\ell \notin S$, let $P(X) \in \mathbb{C}[X]$ denote the quartic polynomial such that*

$$L(\Pi_\ell, s - w/2) = P_\ell(\ell^{-s})^{-1}.$$

Then the subfield $E \subset \mathbb{C}$ generated by the coefficients of the $P_\ell(X)$, for all $\ell \notin S$, is a finite extension of $\mathbb{Q}$.

- (4) *For any prime p and choice of embedding $E \hookrightarrow \overline{\mathbb{Q}}_p$, there is a semisimple Galois representation*

$$\rho_{\Pi, p} : \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathrm{GL}_4(\overline{\mathbb{Q}}_p)$$

characterised (up to isomorphism) by the property that, for all primes $\ell \notin S \cup \{p\}$,

$$\det(1 - X\rho_{\Pi, p}(\mathrm{Frob}_\ell^{-1})) = P_\ell(X),$$

where Frob_ℓ is the arithmetic Frobenius.

- (5) The representation $\rho_{\Pi,p}$ is either irreducible or the direct sum of two distinct irreducible 2-dimensional representations. In particular,

$$H^0(\mathbb{Q}^{\mathrm{ab}}, \rho_{\Pi,p}) = 0.$$

- (6) The restriction of $\rho_{\Pi,p}$ to a decomposition group at p is de Rham, and its Hodge numbers³ are $\{0, k_2 - 2, k_1 - 1, k_1 + k_2 - 3\}$. If Π_p is unramified, then $\rho_{\Pi,p}$ is crystalline, and

$$\det(1 - X\varphi \mid \mathbb{D}_{\mathrm{cris}}(\rho_{\Pi,p})) = P_p(X).$$

Remark 10.1.4. Note that $P_\ell(X)$ in (3) is given by the action of the Hecke-operator-valued polynomial $\mathcal{P}_\ell(X)$ of §3.5.2 on the spherical vector of $\Pi_\ell \otimes |\cdot|^{(3-w)/2}$. $\diamond$

Proof of Theorem 10.1.3. Parts (3) and (4) are [46, Theorem I]. Part (2) is also implicit in that theorem, since the “purity” statement on $\rho_{\Pi,p}$ implies that the local L -factor $L(\Pi_\ell, s)$ has the form $\prod_{i=1}^4 (1 - \alpha_i \ell^{-s})^{-1}$ with $|\alpha_i| = 1$, which rules out all of the other classes of unramified representations of $G(\mathbb{Q}_\ell)$ (all of which have $\alpha_i/\alpha_j = \ell$ for some i, j). Part (5) is contained in Theorem II of *op. cit.*

For parts (1) and (6), the fact that Π^H and Π^W have the same multiplicity, $m(\Pi^H) = m(\Pi^W)$, and the characterisation of the Hodge numbers of $\rho_{\Pi,p}$, was proved in [46, Proposition 1.5 & Theorem III] under the assumption that Π is weakly equivalent to a globally generic representation; and in fact all such Π have this property by the main theorem of [47]. The assertion regarding $\mathbb{D}_{\mathrm{cris}}$ is [44, Theorem 1]. Finally, Xu has shown in [48, §3.5] that the common multiplicity of Π^H and Π^W is equal to 1. $\blacksquare$

It is expected that $\rho_{\Pi,p}$ is always irreducible, but this is only known for large p :

Theorem 10.1.5 (Ramakrishnan). *If Π is unramified at p and $p > 2w + 1$, then the representation $\rho_{\Pi,p}$ is irreducible.*

Proof. By [9], the automorphic representation Π lifts to a cuspidal automorphic representation of GL_4 . This lifted representation is regular at ∞ (since its local parameter at ∞ is determined by that of Π_∞ , via the compatibility of the local and global liftings). If Π_p is unramified, then the corresponding Galois representation is crystalline at p , and hence Theorem B of [32] shows that it is irreducible as long as $p - 1$ is greater than the largest difference between the Hodge–Tate weights of $\rho_{\Pi,p}$, which translates into the condition on p stated above. $\blacksquare$

10.2. Automorphic cohomology

As before, given integers $k_1 \geq k_2 \geq 3$ as above, we let $(a, b) = (k_2 - 3, k_1 - k_2)$, so that $w = 2a + b + 3$. Choosing a (sufficiently small) level group K , we thus have a Shimura

³Here “Hodge numbers” are the jumps in the Hodge filtration of $\mathbb{D}_{\mathrm{dR}}$, which are the negatives of Hodge–Tate weights, so the cyclotomic character has Hodge number -1 .

variety $Y_G(K)$, and a relative Chow motive $\mathcal{D}_{\mathbb{Q}}^{a,b} = \text{Anc}_G(D^{a,b})$ over this variety. We are interested in the parabolic cohomology of the p -adic étale realisation

$$H_{\text{ét},!}^3(Y_G(K)_{\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Q}_p}^{a,b}) = \text{image}(H_{\text{ét},c}^3 \rightarrow H_{\text{ét}}^3).$$

Since $\mathcal{D}_{\mathbb{Q}_p}^{a,b}$ has an equivariant structure, the direct limit $H_{\text{ét},!}^3(Y_{G,\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Q}_p}^{a,b})$ is an admissible smooth representation of $G(\mathbb{A}_f)$, with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ commuting with the $G(\mathbb{A}_f)$ -action.

Notation 10.2.1. Let $\Sigma(k_1, k_2)$ denote the set of isomorphism classes of representations Π_f of $G(\mathbb{A}_f)$ which are the finite part of a cuspidal automorphic representation $\Pi = \Pi_f \otimes \Pi_{\infty}$ in which Π_{∞} is one of the two discrete series representations of weight (k_1, k_2) .

Theorem 10.2.2 (Taylor, Weissauer). *There is a $G(\mathbb{A}_f) \times \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -equivariant direct sum decomposition*

$$H_{\text{ét},!}^3(Y_{G,\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Q}_p}^{a,b}(3)) \otimes \overline{\mathbb{Q}}_p \cong \bigoplus_{\Pi_f \in \Sigma(k_1, k_2)} (\Pi_f^*[\frac{w-3}{2}] \otimes W_{\Pi_f}^*)$$

where W_{Π_f} is a finite-dimensional p -adic representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. If Π is non-endoscopic, then the term corresponding to Π is a direct summand of the full cohomology $H_{\text{ét}}^3(Y_{G,\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Q}_p}^{a,b})$, and the semisimplification of W_{Π_f} is isomorphic to $\rho_{\Pi,p}$.

Here $\Pi_f[r] = \Pi_f \otimes \|\mu(-)\|^r$, as above.

Proof. See [46, §1]. ■

10.3. Arithmetic étale cohomology

We now consider the étale cohomology of $Y_G(K)$ as a $\mathbb{Q}$ -variety (not as a $\overline{\mathbb{Q}}$ -variety); more precisely, we work with continuous étale cohomology in the sense of Jannsen [11]. Note that this space is not finite-dimensional in general. Nonetheless, there is a Hochschild–Serre spectral sequence associated to the structure map $Y_G(K) \rightarrow \text{Spec } \mathbb{Q}$, for any lisse étale $\mathbb{Z}_p$ -sheaf or $\mathbb{Q}_p$ -sheaf $\mathcal{F}$,

$$E_2^{rs} = H^r(\mathbb{Q}, H_{\text{ét}}^s(Y_G(K)_{\overline{\mathbb{Q}}}, \mathcal{F})) \Rightarrow H_{\text{ét}}^{r+s}(Y_G(K), \mathcal{F}),$$

for any integer n . Consequently, if we define

$$H_{\text{ét}}^4(Y_G(K), \mathcal{F})_0 := \ker(H_{\text{ét}}^4(Y_G(K), \mathcal{F}) \rightarrow H_{\text{ét}}^4(Y_G(K)_{\overline{\mathbb{Q}}}, \mathcal{F}))$$

(the cohomologically trivial classes), then we have a natural “Abel–Jacobi” map

$$H_{\text{ét}}^4(Y_G(K), \mathcal{F})_0 \rightarrow H^1(\mathbb{Q}, H_{\text{ét}}^3(Y_G(K)_{\overline{\mathbb{Q}}}, \mathcal{F})).$$

These maps are compatible with changing K , and therefore assemble into a map of $G(\mathbb{A}_f)$ -representations. More generally, this also applies with Y_G replaced by its base-extension $Y_G \times \text{Spec } \mathbb{Q}(\zeta_N)$, for any integer N .

Definition 10.3.1. Suppose Π is non-endoscopic. For any $N \geq 1$ and $q \in \mathbb{Z}$, we write

$$\mathrm{pr}_{\Pi^*} : H_{\mathrm{et}}^4(Y_{G, \mathbb{Q}(\zeta_N)}, \mathcal{D}_{\mathbb{Q}_p}^{a,b}(3-q))_0 \rightarrow \Pi_f^* \left[\frac{w-3}{2} \right] \otimes H^1(\mathbb{Q}(\zeta_N), W_{\Pi_f}^*(-q))$$

for the composite of the Abel–Jacobi map and projection onto the Π_f^* -isotypical component.

10.4. Ordinarity

Let us now choose a non-endoscopic Π with Π_∞ a discrete series representation of weight (k_1, k_2) , as above, and let E be a number field as in Theorem 10.1.3(3), so we have polynomials $P_\ell(X) \in E[X]$ for all unramified primes ℓ .

Proposition 10.4.1. *Let p be a prime such that Π_p is unramified, and write*

$$P_p(X) = 1 + a_1 X + \cdots + a_4 X^4.$$

Then, for any embedding $E \hookrightarrow \overline{\mathbb{Q}_p}$, we have the relations

$$\begin{aligned} v_p(a_1) &\geq 0, & v_p(a_2) &\geq k_2 - 2, \\ v_p(a_3) &\geq k_1 + k_2 - 3, & v_p(a_4) &= 2k_1 + 2k_2 - 6, \end{aligned}$$

where the valuation is normalised so that $v_p(p) = 1$.

Proof. By Theorem 10.1.3(6), we know that the crystalline $G_{\mathbb{Q}_p}$ -representation $\rho_{\Pi, p}|_{G_{\mathbb{Q}_p}}$ has Hodge numbers $\{0, k_2 - 2, k_1 - 1, k_1 + k_2 - 3\}$ and its Frobenius has characteristic polynomial $P_p(X)$. The above relations are now precisely the assertion that the Newton polygon associated to this representation lies on or above its Hodge polygon (with the same endpoints), which is a general property of crystalline representations: see e.g. [8, Proposition 5.4.2]. ■

Remark 10.4.2. These inequalities can also be proved directly (without using p -adic Hodge theory), by expressing the coefficients of $P_p(X)$ as Hecke eigenvalues using the formula of Lemma 3.5.4, and showing that suitable scalar multiples of the Hecke operators preserve an integral lattice in Betti cohomology. ◇

Definition 10.4.3. We say that Π is *good ordinary* at p , with respect to some choice of embedding $E \hookrightarrow \overline{\mathbb{Q}_p}$, if Π is unramified at p and the eigenvalue of $T(p)$ acting on $\Pi_p \left[\frac{3-w}{2} \right]^{G(\mathbb{Z}_p)}$ is a p -adic unit.

Since this $T(p)$ -eigenvalue is a_1 in the notation of the previous proposition, it follows easily from the theory of Newton polygons that Π is good ordinary at p if and only if it is unramified at p and $P_p(X)$ has a factor in $\overline{\mathbb{Q}_p}[X]$ of the form $1 - \alpha X$ with α a p -adic unit. Moreover, this α is unique if it exists. By Proposition 3.5.5, α is also an eigenvalue of $U(p)$ acting on the 4-dimensional space $\Pi_p \left[\frac{3-w}{2} \right]^{K_{G_p, 0}(p)}$, or dually of $U'(p)$ acting on $\Pi_p^* \left[\frac{w-3}{2} \right]^{K_{G_p, 0}(p)}$.

We assume henceforth that Π is good ordinary at p ; and we choose one final piece of data needed to define our Euler system. Having fixed a set S as in §8.4 above, we obtain a level group

$$K_{G,0}(p) = K_S \times \prod_{\ell \nmid pS} G(\mathbb{Z}_\ell) \times K_{G_p,0}(p).$$

Enlarging S and shrinking K_S if necessary, we suppose that Π_f has non-zero invariants under $K_{G,0}(p)$.

Definition 10.4.4. We choose a vector $v_\alpha \in \Pi_f$ invariant under the subgroup $K_{G,0}(p)$ and lying in the $U(p) = \alpha$ eigenspace.

This choice gives a linear functional $\Pi_f^* \rightarrow \overline{\mathbb{Q}}_p$, and hence a homomorphism of Galois representations

$$H_{\text{ét}}^3(Y_{G,\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Q}_p}^{a,b}(3)) \rightarrow W_{\Pi_f}^*,$$

which we also denote by v_α . It seems reasonable to interpret this as a “modular parametrisation” of the Galois representation $W_{\Pi_f}^*$. Composing with the map pr_{Π^*} of the previous section, for each $N \geq 1$ and $0 \leq q \leq a$ we have a map

$$\text{pr}(\Pi^*, v_\alpha) : H_{\text{ét}}^4(Y_{G,\mathbb{Q}(\zeta_N)}, \mathcal{D}_{\mathbb{Q}_p}^{a,b}(3-q)) \rightarrow H^1(\mathbb{Q}(\zeta_N), W_{\Pi_f}^*(-q)).$$

Our local hypothesis at p implies that this homomorphism factors through projection to the $U'(p) = \alpha$ eigenspace at level $K_{G,0}(p)$, and in particular through the ordinary idempotent e'_{ord} .

Remark 10.4.5. It will come as no great surprise to the well-informed reader to learn that the ordinarity condition can be relaxed somewhat, to allow sufficiently small positive values of the “slope” $v_p(t(p))$, where $t(p)$ is the eigenvalue of $T(p)$ on $\Pi_f \left[\frac{3-w}{2} \right]$. (Slope < 1 is easy; slope $< 1 + a$ may be accessible, using the methods of [24].) However, we stick with $v_p(t(p)) = 0$ in the present paper for simplicity. $\diamond$

10.5. Lemma–Eisenstein classes in Galois cohomology

We apply the maps of the previous section to the classes $z_{M,m,1}^{[a,b,q,r]}$, or more precisely to their images in $H_{\text{mot}}^4(Y_{G,1}(p)_{\mathbb{Q}(\zeta_{Mp^m})}, \mathcal{D}_{\mathbb{Q}}^{a,b}(3-q))$ via the map J_{Mp^m} of (9).

Definition 10.5.1. For $m \geq 0$, we define a class

$$z_{M,m}^{[\Pi,q,r]} \in H^1(\mathbb{Q}(\zeta_{Mp^m}), W_{\Pi_f}^*(-q))$$

by

$$\frac{1}{M} \cdot (\text{pr}_{\Pi^*, v_\alpha} \circ J_{Mp^m}) \left\{ \begin{array}{ll} \left(\frac{p^q}{U'(p)} \right)^m e'_{\text{ord}} & \text{if } m \geq 1 \\ \left(1 - \frac{p^q}{U'(p)} \right) e'_{\text{ord}} & \text{if } m = 0 \end{array} \right\} \cdot r_{\text{ét}}(z_{M,m,1}^{[a,b,q,r]}).$$

Note that this is well-defined, since the classes concerned are cohomologically trivial by Corollary 9.6.6. It follows from Theorem 8.4.3(ii) that the classes $z_{M,m}^{[\Pi,q,r]}$ are compatible under the Galois corestriction maps as m varies. More importantly, they also satisfy a compatibility with respect to M :

Proposition 10.5.2. *Let $\ell \nmid Mp$ be a prime, with $\ell \notin S$. Then*

$$\mathrm{cores}_{\mathbb{Q}(\zeta_{Mp^m})}^{\mathbb{Q}(\zeta_{\ell Mp^m})}(z_{\ell M,m}^{[\Pi,q,r]}) = P_\ell(\ell^{-1-q}\sigma_\ell^{-1})z_{M,m}^{[\Pi,q,r]} \quad (14)$$

where $\sigma_\ell \in \mathrm{Gal}(\mathbb{Q}(\zeta_{Mp^m})/\mathbb{Q})$ is the arithmetic Frobenius at ℓ .

Proof. This will (eventually) turn out to be a mildly disguised form of Corollary 3.10.5. As in the proofs of the “ p -direction” norm relations, we are comparing the images of two elements of $\mathcal{S}(\mathbb{A}_f^2)^{\otimes 2} \otimes \mathcal{H}(G(\mathbb{A}_f))$ which are pure tensors having the same local components at all primes away from ℓ , and at ℓ are

$$\frac{\ell-1}{\ell} \cdot \underline{\phi}_{1,1} \otimes \mathrm{ch}(\eta_{\ell,1}G(\mathbb{Z}_\ell)) \quad \text{and} \quad \underline{\phi}_0 \otimes \mathrm{ch}(G(\mathbb{Z}_\ell)).$$

(The factor $\ell - 1$ arises by comparing the volumes of the subgroups W , and the ℓ from the $1/M$ in the definition of $z_{M,m}^{[\Pi,q,r]}$.)

Moreover, the map which we are applying to these elements factors through the Eisenstein symbol map $\mathrm{Eis}^{(c,d)}$, where $(c,d) = (a+q+b-r, a-q+r)$. We first give the proof assuming $c, d \geq 1$. In this case the divisor map ∂ is injective on the image of the Eisenstein symbol, so the local version of our map factors through the map

$$\mathcal{S}(\mathbb{Q}_\ell^2)^{\otimes 2} \rightarrow \bigoplus I(\underline{\eta}), \quad \underline{\phi} \mapsto F_{\underline{\phi}},$$

where the sum is over some set of pairs $\eta = (\eta_1, \eta_2)$ of finite-order characters of $\mathbb{Q}_\ell^\times$, and $I(\eta)$ is an irreducible principal series representation of $H(\mathbb{Q}_\ell)$. So it suffices to check that for any $(G \times H)(\mathbb{Q}_\ell)$ -equivariant homomorphism $I(\eta) \otimes \mathcal{H}(G(\mathbb{Q}_\ell)) \rightarrow W$, where W is an irreducible principal series representation of $G(\mathbb{Q}_\ell)$, the images of $\underline{\phi}_{1,1} \otimes \mathrm{ch}(\eta_{\ell,1}G(\mathbb{Z}_\ell))$ and $F_{\underline{\phi}_0} \otimes \mathrm{ch}(G(\mathbb{Z}_\ell))$ in $W^{G(\mathbb{Z}_\ell)}$ are related via an Euler factor. Corollary 3.10.5 gives a relation of exactly this form, with $\frac{\ell}{\ell-1}L(W^\vee, -1/2)^{-1}$ as the correction factor. If $M = 1$ and $m = 0$, applying this with $W = \Pi_\ell^*[\frac{w-3}{2} - q]$ gives the result, noting that $L(W^\vee, -1/2)^{-1} = L(\Pi_\ell, 1 + q - w/2)^{-1} = P_\ell(\ell^{-1-q})$.

To obtain the result in general, we apply this to each of the twists of W by Dirichlet characters modulo Mp^m ; the twist of course modifies the Euler factor P_ℓ , which corresponds to the appearance of σ_ℓ in the statement of the theorem.

If either c or d (or both) is zero, then the divisor map has a kernel (consisting of modular units which are constant along one of the factors of Y_H). However, the kernel of this map is a sum of non-generic representations of $H(\mathbb{Q}_\ell)$, and Lemma 3.7.2 shows that if τ_ℓ is any of these representations, then every $G \times H$ -equivariant map $\tau_\ell \otimes \mathcal{H}(G(\mathbb{Q}_\ell)) \rightarrow W$ is zero. ■

This Euler system norm relation is not terribly useful on its own – we need to combine it with some uniform control over the denominators of these classes. Let $T_{\Pi_f}^*$ denote the lattice in $W_{\Pi_f}^*$ generated by the image of $H_{\text{ét}}^3(Y_{G,1}(p)_{\overline{\mathbb{Q}}}, \mathcal{D}_{\mathbb{Z}_p}^{a,b})$ under v_α . We choose a pair (c_1, c_2) of integers > 1 as before, and we impose the additional assumption that the elements $\text{diag}(1, c_1, 1, c_1)$ and $\text{diag}(c_2, 1, c_2, 1) \in G(\mathbb{A}_f)$ act trivially on the vector v_α (which can be arranged, as usual, by assuming that the c_i are sufficiently close to 1 modulo the primes in S).

Definition 10.5.3. For M coprime to $c_1 c_2$, and $m \geq 0$, let us define

$$c_{1,c_2} z_{M,m}^{[\Pi,q,r]} \in H^1(\mathbb{Q}(\zeta_{Mp^m}), T_{\Pi_f}^*(-q))$$

by

$$\frac{1}{M} \cdot (\text{pr}_{\Pi^*, v_\alpha} \circ JM_{p^m}) \left\{ \begin{array}{ll} U'(p)^{-m} & \text{if } m \geq 1 \\ 1 - \frac{p^q}{U'(p)} & \text{if } m = 0 \end{array} \right\} \cdot e'_{\text{ord}}(c_{1,c_2} z_{M,m,1}^{[a,b,q,r]}).$$

From Proposition 8.4.6 (and the interaction between JM_{p^m} and the $G(\mathbb{A}_f)$ action described in §5.4), we see that the image of $c_{1,c_2} z_{M,m}^{[\Pi,q,r]}$ after inverting p is

$$(c_1^2 - c_1^{-(a-q+b-r)} \sigma_{c_1})(c_2^2 - c_2^{-(a-q+r)} \sigma_{c_2}) \cdot c_{1,c_2} z_{M,m}^{[\Pi,q,r]},$$

where $\sigma_{c_i} \in \text{Gal}(\mathbb{Q}(\zeta_{Mp^m})/\mathbb{Q})$ maps to $c_i \bmod Mp^m$ under the mod Mp^m cyclotomic character.

Theorem 10.5.4. Suppose M is coprime to $c_1 c_2$. Then:

(a) For each integer r with $0 \leq r \leq b$, there is a class

$$c_{1,c_2} z_{\text{Iw},M}^{[\Pi,r]} \in H_{\text{Iw}}^1(\mathbb{Q}(\zeta_{Mp^\infty}), T_{\Pi_f}^*)$$

uniquely determined by the following property: for each $m \geq 0$ and $q \in \{0, \dots, a\}$, the image of $(-2)^q c_{1,c_2} z_{\text{Iw},M}^{[\Pi,r]}$ in $H^1(\mathbb{Q}(\zeta_{Mp^m}), T_{\Pi_f}^*(-q))$ is $c_{1,c_2} z_{M,m}^{[\Pi,q,r]}$.

(b) If $\ell \nmid Mpc_1 c_2$ and $\ell \notin S$, then we have the norm relation

$$\text{cores}_{\mathbb{Q}(\zeta_{Mp^\infty})}^{\mathbb{Q}(\zeta_{\ell Mp^\infty})}(c_{1,c_2} z_{\text{Iw},\ell M}^{[\Pi,r]}) = P_\ell(\ell^{-1} \sigma_\ell^{-1}) \cdot c_{1,c_2} z_{\text{Iw},M}^{[\Pi,r]}.$$

Proof. The image of the class $c_{1,c_2} z_{\text{Iw},M}$ under the maps $\text{mom}_{G,m,1}^{[a,b,0,r]}$, for $m \geq 0$, define a class in $H_{\text{Iw}}^4(Y_G(M, p^\infty, p)_\Sigma, \mathcal{D}_{\mathbb{Z}_p}^{a,b}(3))$ where Σ is the set of primes dividing MpS . This class is cohomologically trivial by Corollary 9.6.6. By Theorem 9.6.4, the image of this class under tensor product with ζ^{-q} and projection to level m agrees, up to an appropriate correction factor, with the étale class $c_{1,c_2} z_{\text{ét},M,m,1}^{[a,b,q,r]}$. Applying the maps $\text{pr}_{(\Pi^*, v_\alpha)} \circ JM_{p^m}$ and dividing by M gives an Iwasawa cohomology class with the required properties.

Let us now prove (b). We claim that, for any given q , the map

$$H_{\text{Iw}}^1(\mathbb{Q}(\zeta_{Mp^m}), T_{\Pi_f}^*) \rightarrow \varprojlim_m H^1(\mathbb{Q}(\zeta_{Mp^m}), W_{\Pi_f}^*(-q))$$

is injective (i.e. there is no non-zero Iwasawa cohomology class whose image at every finite level is p -torsion). This is the map denoted by λ_q in [18, Appendix A]. By Proposition A.2.6 of *op. cit.*, its kernel is contained in the $\Lambda(\Gamma)$ -torsion submodule of the Iwasawa cohomology; and this torsion submodule is in fact zero, by Theorem 10.1.3(5). With this injectivity in hand, the norm relation (b) follows from the norm relation of Proposition 10.5.2 for the non-integral classes at finite level. ■

We conclude this section with a local property of these Galois cohomology classes.

Proposition 10.5.5. *For each prime $\lambda \nmid p$ of $\mathbb{Q}(\zeta_{Mp^m})$, the image of $c_{1,c_2} z_{M,m}^{[\Pi,q,r]}$ in $H^1(I_\lambda, T_{\Pi_f}^*(-q))$ is zero, where $I_\lambda \subset \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_{Mp^m}))$ is any choice of inertia group at λ . For the primes above p , the localisation lies in the Bloch–Kato crystalline subspace $H_f^1(\mathbb{Q}(\zeta_{Mp^m})_\lambda, T_{\Pi_f}^*(-q))$.*

Proof. It is a standard result that Galois cohomology classes which are universal norms in the p -cyclotomic extension are always unramified outside the primes above p (see e.g. [37, Corollary B.3.5]). The fact that our classes satisfy the Bloch–Kato condition at p is deeper. The comparison between étale cohomology and the syntomic cohomology of Nekovář–Nizioł [27] shows that the localisations lie in the possibly larger Bloch–Kato space $H_g^1 \supseteq H_f^1$. It suffices to check that p^{-1} is not an eigenvalue of crystalline Frobenius on $\mathbb{D}_{\mathrm{cris}}(W_{\Pi_f}^*(-q))$, since this implies that the H_f^1 and H_g^1 spaces coincide. However, the eigenvalues of Frobenius on this space are exactly the quantities $p^q \alpha_i^{-1}$, where $L(\Pi_p, s - w/2) = \prod_{i=1}^4 (1 - \alpha_i p^{-s})^{-1}$. Since the α_i are Weil numbers of weight w , the $p^q \alpha_i^{-1}$ have weight $2q - w \leq -3$; so none of these quantities may be equal to p^{-1} , and H_f^1 and H_g^1 coincide for this representation. ■

Equivalently, we have shown that

$$c_{1,c_2} z_{M,m}^{[\Pi,q,r]} \in H_f^1(\mathbb{Q}(\zeta_{Mp^m}), T_{\Pi_f}^*(-q)),$$

where the right-hand side is the global Bloch–Kato Selmer group.

10.6. The Euler-system map

In this short section we give a slicker reinterpretation of the above results. Let L be a finite extension of $\mathbb{Q}_p$ with ring of integers $\mathcal{O}$; and Σ a finite set of primes including p . We let $\mathcal{R}$ denote the set of square-free products of primes not in Σ . If T is a finite free $\mathcal{O}$ -module with a continuous action of $\mathrm{Gal}(\mathbb{Q}^\Sigma/\mathbb{Q})$, and $\ell \notin \Sigma$, we let $P_\ell(T; X) = \det(1 - X \mathrm{Frob}_\ell^{-1} : T)$.

Definition 10.6.1. For (T, Σ) as above, we define $\mathrm{ES}(T, \Sigma)$ to be the set of families of cohomology classes $(c_M)_{M \in \mathcal{R}}$, with $c_M \in H_{\mathrm{Iw}}^1(\mathbb{Q}(\zeta_{Mp^\infty}), T)$, satisfying

$$\mathrm{cores}_{\mathbb{Q}(\zeta_{Mp^\infty})}^{\mathbb{Q}(\zeta_{\ell Mp^\infty})}(c_{M\ell}) = P_\ell(T^*(1); \sigma_\ell^{-1})c_M$$

for ℓ prime with $\ell \nmid M$, $\ell \notin \Sigma$. We refer to such families as *Euler systems* for (T, Σ) .

(In the notation of [37, Definition 2.1.1], these are Euler systems for $(T, \mathcal{K}, \mathcal{N})$ where $\mathcal{K}$ is the compositum of the $\mathbb{Q}(\zeta_{Mp^\infty})$ for $M \in \mathcal{R}$, and $\mathcal{N} = \prod_{\ell \in \Sigma} \ell$.)

Theorem 10.5.4 shows that the classes $(c_1, c_2 z_{\text{Iw}, M}^{[\Pi, r]})_{M \in \mathcal{R}}$ are an Euler system for $T = T_{\Pi_f}^*$, with Σ the set of primes not dividing pc_1c_2S . This Euler system, of course, depends on choices of local data at the primes in S , namely the Schwartz function ϕ_S and the group $W_S = \prod_{\ell \in S} W_\ell \subset H(\mathbb{Q}_S)$. The goal of this section is to make this dependence precise.

Let $K^{S,p} = \prod_{\ell \nmid pS} G(\mathbb{Z}_\ell) \times K_{G_p, 0}(p)$. Then our modular parametrisation v_α is an element of the space

$$\sigma_S := \Pi_f \left[\frac{3-w}{2} \right]^{(K^{S,p}, U(p)=\alpha)},$$

which is an irreducible representation of $G(\mathbb{Q}_S)$, isomorphic to $\Pi_S \left[\frac{3-w}{2} \right]$. The choice of subgroups K_S and W_S at the bad primes only affects the construction of the Euler system through the volume factor $\text{Vol}(W)$, so we have in fact defined a canonical bilinear map

$$\mathcal{L}\mathcal{E}_{\text{ES}, S}^{[\Pi, r]} : \mathcal{S}(\mathbb{Q}_S^2, L)^{\otimes 2} \otimes \sigma_S \rightarrow \text{ES}(T_{\Pi_f}^*, \Sigma) \otimes_{\mathcal{O}} L,$$

mapping $\phi_S \otimes v_\alpha$ to $\text{Vol}(W_S) \cdot (c_1, c_2 z_{\text{Iw}, M}^{[\Pi, r]})_{M \in \mathcal{R}}$.

Remark 10.6.2. The map $\mathcal{L}\mathcal{E}_{\text{ES}, S}^{[\Pi, r]}$ does still depend on (c_1, c_2) . In fact, we assumed above that the c_i were close to 1 locally above S – where the meaning of “close” depended on the local data chosen – but this is not needed in order to define the classes $c_1, c_2 z_{\text{Iw}, M}^{[\Pi, r]}$ or to prove their norm relations, only to state simply their relation to the non-integral classes. $\diamond$

Proposition 10.6.3. *This map satisfies the equivariance property*

$$\mathcal{L}\mathcal{E}_{\text{ES}, S}^{[\Pi, r]}(h\phi, hv) = \text{Art}(\det h)^{-1} \cdot \mathcal{L}\mathcal{E}_{\text{ES}, S}^{[\Pi, r]}(\phi, v)$$

for all $h \in H(\mathbb{Q}_S)$, where we let $\text{Gal}(\mathbb{Q}^{\text{ab}}/\mathbb{Q})$ act on $\text{ES}(T_{\Pi_f}^*, \Sigma)$ via its natural map to $\text{Gal}(\mathbb{Q}(\zeta_{Mp^\infty})/\mathbb{Q})$ for all $M \in \mathcal{R}$.

Proof. This follows from the $H(\mathbb{A}_f) \times G(\mathbb{A}_f)$ -equivariance of the Lemma–Eisenstein map. $\blacksquare$

Remark 10.6.4. Similarly, mapping $\phi_S \otimes v_\alpha$ to $\text{Vol}(W_S) \cdot z_{1,0}^{[\Pi, q, r]}$ defines an $H(\mathbb{Q}_S)$ -invariant bilinear form on $\mathcal{S}(\mathbb{Q}_S^2, L)^{\otimes 2} \otimes \sigma_S[q]$. If we fix characters ν_1, ν_2 of $\mathbb{Z}_S^\times$ and restrict to Schwartz functions on which the centre of $\text{GL}_2(\mathbb{Z}_S) \times \text{GL}_2(\mathbb{Z}_S)$ acts via $\nu_1 \times \nu_2$, then this bilinear form is forced to factor through $\tau_S \otimes \sigma_S[q]$ for some irreducible principal series representation τ_S of $H(\mathbb{Q}_S)$. Of course, the restriction is zero unless $(\nu_1 \nu_2)^{-1}$ coincides with the restriction to $\mathbb{Z}_S^\times$ of the central character of σ_S .

We expect that $\dim \text{Hom}_{H(\mathbb{Q}_S)}(\tau_S \otimes \sigma_S[q], L)$ should have L -dimension ≤ 1 . This follows from the Gan–Gross–Prasad conjecture for $\text{SO}_4 \times \text{SO}_5$ if $\nu_1 \nu_2$ is a square in the group of characters of $\mathbb{Z}_S^\times$, and should probably be true more generally, but we do not know a reference; let us assume this for the duration of this remark.

If this dimension is 0, then the cohomology class $z_{1,0}^{[\Pi,q,r]}$ is zero for every choice of the local data in the (v_1, v_2) eigenspace. If this dimension is 1 – which is the case if Π_ℓ is generic for all $\ell \in S$ – then the choice of local data only affects this class up to a scaling factor, which is essentially the local zeta integral of Piatetski-Shapiro appearing in [22, §5.2]. We expect, but cannot prove, that if there exists a prime ℓ such that Π_ℓ is not generic, then the classes $z_{M,m}^{[\Pi,q,r]}$ are zero for all choices of the local data and the parameters (q, r, M, m) . $\diamond$

11. Selmer groups and p -adic L -functions

We conclude by showing that if the above Euler system is non-zero, it gives bounds on Selmer groups.

11.1. Assumptions on Π

In this section, Π will denote a non-endoscopic automorphic representation of GSp_4 , discrete series at ∞ of some weight (k_1, k_2) , as before. We suppose that $p \neq 2$, and fix an embedding $E \hookrightarrow \overline{\mathbb{Q}}_p$, where E is a number field as in Theorem 10.1.3(3). We let L be a sufficiently large finite extension of $\mathbb{Q}_p$, with ring of integers $\mathcal{O}_L$ and residue field k_L , such that L contains the image of E and $W_{\Pi_f}^*$ is definable over L as a quotient of the cohomology of $Y_{G,\overline{\mathbb{Q}}}$.

We also impose the following extra hypotheses:

Assumption 11.1.1 (“no exceptional zero”). None of the roots of the polynomial $P_p(X)$ are of the form $p^n \zeta$ with $n \in \mathbb{Z}$ and ζ a root of unity.

Assumption 11.1.2 (“big image”).

- (i) The representation $T_{\Pi_f}^* \otimes k_L$ is irreducible as a $k_L[\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_{p^\infty}))]$ -module.
- (ii) There exists $\tau \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_{p^\infty}))$ such that $T_{\Pi_f}^*/(\tau - 1)T_{\Pi_f}^*$ is free of rank 1 over $\mathcal{O}$. (This is precisely Hyp(K_∞, T) of [37].)

Assumption 11.1.3 (“Siegel ordinarity”). Π is good ordinary at p in the sense of Definition 10.4.3.

Remark 11.1.4. Note that the “big image” assumption is clearly satisfied if the image of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ in $\mathrm{Aut} W_{\Pi_f}^*$ contains a conjugate of $\mathrm{Sp}_4(\mathbb{Z}_p)$. This is expected to hold for all but finitely many p if Π is “sufficiently general” (i.e. not a functorial lift from a proper subgroup of GSp_4). However, the big image assumption is also satisfied in certain other cases, such as twisted Yoshida lifts of suitable Hilbert modular forms. $\diamond$

11.2. Ordinary submodules at p

Recall that the ordinarity property implies that there is a unique reciprocal root α of $P_p(X)$ which is a p -adic unit.

Proposition 11.2.1 (Urban; [44, Corollaire 1(i)]). *The representation $W_{\Pi_f}|_{G_{\mathbb{Q}_p}}$ has a 1-dimensional unramified subrepresentation on which geometric Frobenius acts as α .*

Equivalently, $W_{\Pi_f}^*$ has a decreasing filtration by $G_{\mathbb{Q}_p}$ -stable subspaces,

$$W_{\Pi_f}^* \supsetneq \mathcal{F}^1 W_{\Pi_f}^* \supsetneq \mathcal{F}^3 W_{\Pi_f}^* \supsetneq \{0\},$$

with $\mathcal{F}^i$ having codimension i , and the quotient $W_{\Pi_f}^*/\mathcal{F}^1$ is unramified. We let $\mathcal{F}^i T_{\Pi_f}^*$ be the intersection of $T_{\Pi_f}^*$ with $\mathcal{F}^i W_{\Pi_f}^*$.

Proposition 11.2.2. *For any $0 \leq r \leq b$, and any $M \in \mathcal{R}$, the image of the element $c_1, c_2 z_{\text{Iw}, M}^{[\Pi, r]}$ in $H_{\text{Iw}}^1(\mathbb{Q}_p(\zeta_{Mp^\infty}), T_{\Pi_f}^*/\mathcal{F}^1)$ is zero.*

Proof. From the ordinarity of Π_p it follows that $H^0(\mathbb{Q}_p(\zeta_{Mp^\infty}), T_{\Pi_f}^*/\mathcal{F}^1) = 0$, and hence the natural map

$$H_{\text{Iw}}^1(\mathbb{Q}_p(\zeta_{Mp^\infty}), T_{\Pi_f}^*/\mathcal{F}^1) \rightarrow \varprojlim_m H_{\text{Iw}}^1(\mathbb{Q}_p(\zeta_{Mp^m}), W_{\Pi_f}^*(-q)/\mathcal{F}^1)$$

is injective for any integer q . So it suffices to show that the image of $c_1, c_2 z_{M, m}^{[\Pi, q, r]}$ in $W_{\Pi_f}^*(-q)/\mathcal{F}^1$ is zero for all m (for any choice of q). However, if q is chosen such that $0 \leq q \leq a$, then this element lands in the Bloch–Kato subspace $H_f^1(\mathbb{Q}_p(\zeta_{Mp^m}), W_{\Pi_f}^*(-q)/\mathcal{F}^1)$, by Proposition 10.5.5; and this subspace is zero, since $W_{\Pi_f}^*(-q)/\mathcal{F}^1$ has all Hodge–Tate weights ≤ 0 (and is not the trivial representation). ■

Let us write $\mathbb{Q}_\infty = \mathbb{Q}(\zeta_{p^\infty})$. We can use the submodule $\mathcal{F}^1 T_{\Pi_f}^*$ to define Selmer groups $\tilde{H}_{\text{Iw}}^i(\mathbb{Q}_\infty, T_{\Pi_f}^*)$, via Nekovář’s formalism of Selmer complexes [26]; cf. [15, §11.2]. These are finitely generated Λ -modules, where Λ denotes the Iwasawa algebra $\mathcal{O}[[\Gamma]]$ of $\Gamma = \text{Gal}(\mathbb{Q}_\infty/\mathbb{Q}) \cong \mathbb{Z}_p^\times$. They admit the following somewhat concrete descriptions:

- $\tilde{H}_{\text{Iw}}^i(\mathbb{Q}_\infty, T_{\Pi_f}^*) = 0$ unless $i = 1$ or $i = 2$.
- $\tilde{H}_{\text{Iw}}^1(\mathbb{Q}_\infty, T_{\Pi_f}^*)$ is the kernel of the map

$$H_{\text{Iw}}^1(\mathbb{Q}_\infty, T_{\Pi_f}^*) \rightarrow H_{\text{Iw}}^1(\mathbb{Q}_{p, \infty}, T_{\Pi_f}^*/\mathcal{F}^1).$$

- Let S_∞ denote the set of primes of $\mathbb{Q}_\infty$ above S , and $A = T_{\Pi_f}(1) \otimes \mathbb{Q}_p/\mathbb{Z}_p$. If we define $\mathcal{S}(\mathbb{Q}_\infty, A)$ to be the p -torsion group

$$\ker\left(H^1(\mathbb{Z}[\zeta_{p^\infty}, 1/pS], A) \rightarrow \bigoplus_{v \in S_\infty} H^1(\mathbb{Q}_{\infty, v}, A) \oplus H^1(\mathbb{Q}_{p, \infty}, A/\mathcal{F}^3 A)\right),$$

then there is an exact sequence

$$0 \rightarrow \mathcal{S}(\mathbb{Q}_\infty, A)^\vee \rightarrow \tilde{H}_{\text{Iw}}^2(\mathbb{Q}_\infty, T_{\Pi_f}^*) \rightarrow H_{\text{Iw}}^2(\mathbb{Q}_{p, \infty}, \mathcal{F}^1 T_{\Pi_f}^*)$$

where the last module is a finite group (cf. [15, Proposition 11.2.8]).

Moreover, an Euler characteristic computation (using the fact that complex conjugation acts on W_{Π_f} with two $+1$ eigenvalues and two -1 eigenvalues) shows that $\mathrm{rank}_{\Lambda} \tilde{H}^1 - \mathrm{rank}_{\Lambda} \tilde{H}^2 = 1$.

Theorem 11.2.3. *Suppose that there exists an r such that ${}_{c_1, c_2} z_{\mathrm{Iw}, 1}^{[\Pi, r]}$ is non-torsion. Then $\tilde{H}_{\mathrm{Iw}}^2(\mathbb{Q}_{\infty}, T_{\Pi_f}^*)$ is a torsion Λ -module, and we have the divisibility of characteristic ideals*

$$\mathrm{char}_{\Lambda}(\tilde{H}_{\mathrm{Iw}}^2) \mid \mathrm{char}_{\Lambda}\left(\frac{\tilde{H}_{\mathrm{Iw}}^1}{\Lambda \cdot {}_{c_1, c_2} z_{\mathrm{Iw}, 1}^{[\Pi, r]}}\right).$$

Proof. This follows by applying the “Euler system machine” to the Euler system ${}_{c_1, c_2} z_{\mathrm{Iw}, M}^{[\Pi, r]}$. Compare [15, Theorem 11.4.3]. ■

Corollary 11.2.4. *Suppose q is an integer ≥ 0 , and assume that there is an $r \in \{0, \dots, b\}$ such that the cohomology class $z_{1,0}^{[\Pi, q, r]} \in H^1(\mathbb{Q}, W_{\Pi_f}^*(-q))$ is non-zero.*

- *If $\mathrm{loc}_p(z_{1,0}^{[\Pi, q, r]})$ lies in the subspace $H_{\mathrm{f}}^1(\mathbb{Q}_p, W_{\Pi_f}^*(-q))$, then the Bloch–Kato Selmer group $H_{\mathrm{f}}^1(\mathbb{Q}, W_{\Pi_f}^*(-q))$ has dimension 1 over L , and $z_{1,0}^{[\Pi, q, r]}$ is a basis of this space.*
- *If $\mathrm{loc}_p(z_{1,0}^{[\Pi, q, r]})$ does not lie in $H_{\mathrm{f}}^1(\mathbb{Q}_p, W_{\Pi_f}^*(-q))$, then $H_{\mathrm{f}}^1(\mathbb{Q}, W_{\Pi_f}^*(-q))$ is zero.*

Note that the second case cannot occur if $0 \leq q \leq a$, by Proposition 10.5.5.

Proof of Corollary 11.2.4. This follows from the previous theorem by descent; compare [15, Proposition 11.5.1]. ■

11.3. The motivic p -adic L -function

We can also interpret the above results in terms of p -adic L -functions. For simplicity, we assume that Π_p is ordinary for the Borel subgroup (not just for the Siegel parabolic) in the sense of [44]. In this case, Corollary 1 of *op. cit.* shows that there is a 2-dimensional $G_{\mathbb{Q}_p}$ -stable subspace $\mathcal{F}^2 W_{\Pi_f}^*$, with $\mathcal{F}^1 \supsetneq \mathcal{F}^2 \supsetneq \mathcal{F}^3$. Then the graded piece $\mathcal{F}^1 W_{\Pi_f}^* / \mathcal{F}^2$ has Hodge–Tate weight $a + 1$, and Perrin-Riou’s “big logarithm” map gives a canonical isomorphism

$$\mathcal{L} : H_{\mathrm{Iw}}^1(\mathbb{Q}_{p, \infty}, \mathcal{F}^1 W_{\Pi_f}^* / \mathcal{F}^2) \rightarrow \Lambda_L(\Gamma) \otimes \mathbb{D}_{\mathrm{cris}}(\mathcal{F}^1 W_{\Pi_f}^* / \mathcal{F}^2).$$

Composing $\mathcal{L}$ with evaluation at χ^q , where χ is the cyclotomic character, interpolates the Bloch–Kato logarithm (for $q \leq a$) or dual exponential (for $q \geq a + 1$) of the image of z in $H^1(\mathbb{Q}_p, \mathcal{F}^1 W_{\Pi_f}^*(-q) / \mathcal{F}^2)$.

Definition 11.3.1. We let ${}_{c_1, c_2} L_p^{\mathrm{mot}, r}(\Pi_f)$ be the image of ${}_{c_1, c_2} z_{\mathrm{Iw}, 1}^{[\Pi, r]}$ under the map $\mathcal{L}$.

By choosing a basis of the 1-dimensional L -vector space $\mathbb{D}_{\mathrm{cris}}(\mathcal{F}^1 W_{\Pi_f}^* / \mathcal{F}^2)$, we may regard this as an element of $\Lambda_L(\Gamma)$, well-defined up to non-zero scalars. We call this measure the *motivic p -adic L -function*.

Theorem 11.3.2. *Let $a + 1 \leq q \leq a + b + 1$, and suppose that ${}_{c_1, c_2} L_p^{\text{mot}, r}(\Pi_f)$ is non-vanishing at χ^q for some r . Then $H_f^1(\mathbb{Q}, W_{\Pi_f}^*(-q)) = H_f^1(\mathbb{Q}, W_{\Pi_f}(1+q)) = 0$.*

Proof. By construction, if the motivic L -function does not vanish, then the image of ${}_{c_1, c_2} z_{\text{lw}, 1}^{[\Pi, r]}$ in $H^1(\mathbb{Q}_p, \mathcal{F}^1 W_{\Pi_f}^*(-q)/\mathcal{F}^2)$ is non-zero. The hypotheses on q imply that this subquotient has vanishing H_f^1 , so we conclude that ${}_{c_1, c_2} z_{\text{lw}, 1}^{[\Pi, r]}$ cannot lie in H_f^1 locally at p . By Corollary 11.2.4 it follows that the global H_f^1 is zero. ■

Remark 11.3.3. Note that $a + 1 \leq q \leq a + b + 1$ is precisely the range such that $L(\Pi, 1 + q - w/2)$ is a critical value of the spin L -function. We conjecture that, for an appropriate r and suitably chosen test data ϕ, v_α , the value at χ^q of ${}_{c_1, c_2} L_p^{\text{mot}, r}(\Pi_f)$ should be non-zero if and only if the critical L -value does not vanish. ◇

Acknowledgments. This paper owes its existence to a question raised by Francesco Lemma, who asked us whether the techniques used to build the Euler system of Beilinson–Flach elements for $\text{GL}_2 \times \text{GL}_2$ could be adapted to the setting of GSp_4 . We are very grateful to Francesco for this inspiration, and for several interesting conversations during the writing of the paper. We would also like to thank several others for helpful advice and comments, notably Giuseppe Ancona, Martin Dickson, Dimitar Jetchev, Jacques Tilouine, and Bin Xu; and the referee for numerous helpful corrections and comments.

Substantial parts of this paper were written during a visit by the first and third authors to the Institute for Advanced Study in Princeton in the spring of 2016, and we are grateful to the Institute for its support and hospitality.

Funding. Supported by: Royal Society University Research Fellowship “ L -functions and Iwasawa theory” (Loeffler); Simons Investigator Grant #376203 from the Simons Foundation and NSF grant DMS-1501064 (Skinner); ERC Consolidator Grant “Euler systems and the Birch–Swinnerton-Dyer conjecture” (Zerbes).

References

- [1] Ancona, G.: Décomposition de motifs abéliens. *Manuscripta Math.* **146**, 307–328 (2015) Zbl [1453.14017](#) MR [3312448](#)
- [2] Beilinson, A. A.: Higher regulators of modular curves. In: *Applications of Algebraic K-Theory to Algebraic Geometry and Number Theory, Part I, II* (Boulder, CO, 1983), *Contemp. Math.* 55, Amer. Math. Soc., Providence, RI, 1–34 (1986) Zbl [0609.14006](#) MR [862627](#)
- [3] Bump, D.: *Automorphic Forms and Representations*. *Cambridge Stud. Adv. Math.* 55, Cambridge Univ. Press, Cambridge (1997) Zbl [0911.11022](#) MR [1431508](#)
- [4] Bump, D., Friedberg, S., Furusawa, M.: Explicit formulas for the Waldspurger and Bessel models. *Israel J. Math.* **102**, 125–177 (1997) Zbl [1073.11513](#) MR [1489103](#)
- [5] Colmez, P.: La conjecture de Birch et Swinnerton-Dyer p -adique. *Astérisque* **294**, ix, 251–319 (2004) Zbl [1094.11025](#) MR [2111647](#)
- [6] Deligne, P.: Formes modulaires et représentations l -adiques. In: *Séminaire Bourbaki. Vol. 1968/69: Exposés 347–363, Lecture Notes in Math.* 175, Springer, Berlin, Exp. 355, 139–172 (1971) Zbl [0206.49901](#) MR [3077124](#)

- [7] Deninger, C., Scholl, A. J.: The Beilinson conjectures. In: *L-functions and Arithmetic* (Durham, 1989), London Math. Soc. Lecture Note Ser. 153, Cambridge Univ. Press, Cambridge, 173–209 (1991) Zbl [0729.14002](#) MR [1110393](#)
- [8] Fontaine, J.-M.: Représentations p -adiques semi-stables. *Astérisque* **223**, 113–184 (1994) Zbl [0865.14009](#) MR [1293972](#)
- [9] Gan, W. T., Takeda, S.: The local Langlands conjecture for $\mathrm{GSp}(4)$. *Ann. of Math. (2)* **173**, 1841–1882 (2011) Zbl [1230.11063](#) MR [2800725](#)
- [10] Gelbart, S. S., Knapp, A. W.: L -indistinguishability and R groups for the special linear group. *Adv. Math.* **43**, 101–121 (1982) Zbl [0493.22005](#) MR [644669](#)
- [11] Jannsen, U.: Continuous étale cohomology. *Math. Ann.* **280**, 207–245 (1988) Zbl [0649.14011](#) MR [929536](#)
- [12] Kato, K.: p -adic Hodge theory and values of zeta functions of modular forms. *Astérisque* **295**, ix, 117–290 (2004) Zbl [1142.11336](#) MR [2104361](#)
- [13] Kato, S., Murase, A., Sugano, T.: Whittaker–Shintani functions for orthogonal groups. *Tohoku Math. J. (2)* **55**, 1–64 (2003) Zbl [1037.22034](#) MR [1956080](#)
- [14] Kings, G.: Eisenstein classes, elliptic Soulé elements and the ℓ -adic elliptic polylogarithm. In: *The Bloch–Kato Conjecture for the Riemann Zeta Function*, London Math. Soc. Lecture Note Ser. 418, Cambridge Univ. Press, Cambridge, 239–296 (2015) Zbl [1353.14029](#) MR [3497682](#)
- [15] Kings, G., Loeffler, D., Zerbes, S. L.: Rankin–Eisenstein classes and explicit reciprocity laws. *Cambridge J. Math.* **5**, 1–122 (2017) Zbl [1428.11103](#) MR [3637653](#)
- [16] Kings, G., Loeffler, D., Zerbes, S. L.: Rankin–Eisenstein classes for modular forms. *Amer. J. Math.* **142**, 79–138 (2020) Zbl [07176031](#) MR [4060872](#)
- [17] Laumon, G.: Fonctions zêtas des variétés de Siegel de dimension trois. *Astérisque* **302**, 1–66 (2005) Zbl [1097.11021](#) MR [2234859](#)
- [18] Lei, A., Loeffler, D., Zerbes, S. L.: Euler systems for Rankin–Selberg convolutions of modular forms. *Ann. of Math. (2)* **180**, 653–771 (2014) Zbl [1315.11044](#) MR [3224721](#)
- [19] Lei, A., Loeffler, D., Zerbes, S. L.: Euler systems for Hilbert modular surfaces. *Forum Math. Sigma* **6**, art. e23, 67 pp. (2018) Zbl [1437.11073](#) MR [3876763](#)
- [20] Lemma, F.: A norm compatible system of Galois cohomology classes for $\mathrm{GSp}(4)$. *Comment. Math. Helv.* **85**, 885–905 (2010) Zbl [1250.11095](#) MR [2718141](#)
- [21] Lemma, F.: On higher regulators of Siegel threefolds I: The vanishing on the boundary. *Asian J. Math.* **19**, 83–120 (2015) Zbl [1375.19010](#) MR [3318014](#)
- [22] Lemma, F.: On higher regulators of Siegel threefolds II: the connection to the special value. *Compos. Math.* **153**, 889–946 (2017) Zbl [1430.11091](#) MR [3705247](#)
- [23] Loeffler, D., Skinner, C., Zerbes, S. L.: An Euler system for $\mathrm{GU}(2, 1)$, *Math. Ann.* (online, 2021); DOI: [10.1007/s00208-021-02224-4](#)
- [24] Loeffler, D., Zerbes, S. L.: Rankin–Eisenstein classes in Coleman families. *Res. Math. Sci.* **3**, art. 29, 53 pp. (2016) Zbl [1414.11070](#) MR [3552987](#)
- [25] Milne, J. S.: Introduction to Shimura varieties. In: *Harmonic Analysis, the Trace Formula, and Shimura Varieties*, Clay Math. Proc. 4, Amer. Math. Soc., Providence, RI, 265–378 (2005) Zbl [1148.14011](#) MR [2192012](#)
- [26] Nekovář, J.: Selmer complexes. *Astérisque* **310**, viii+559 pp. (2006) Zbl [1211.11120](#) MR [2333680](#)

- [27] Nekovář, J., Nizioł, W.: Syntomic cohomology and p -adic regulators for varieties over p -adic fields. *Algebra Number Theory* **10**, 1695–1790 (2016) Zbl [1375.14081](#) MR [3556797](#)
- [28] Novodvorsky, M. E.: Automorphic L -functions for symplectic group $\mathrm{GSp}(4)$. In: *Automorphic Forms, Representations and L -functions* (Corvallis, OR, 1977), Part 2, Proc. Sympos. Pure Math. 33, Amer. Math. Soc., Providence, RI, 87–95 (1979) Zbl [0408.12013](#) MR [546610](#)
- [29] Piatetski-Shapiro, I. I.: L -functions for GSp_4 . *Pacific J. Math.*, Special Issue, 259–275 (1997) Zbl [1001.11020](#) MR [1610879](#)
- [30] Pink, R.: Arithmetical compactification of mixed Shimura varieties. *Bonner Math. Schriften* 209, Univ. Bonn, Math. Institut, Bonn (1990) Zbl [0748.14007](#) MR [1128753](#)
- [31] Pitale, A., Schmidt, R.: Integral representation for L -functions for $\mathrm{GSp}_4 \times \mathrm{GL}_2$. *J. Number Theory* **129**, 1272–1324 (2009) Zbl [1258.11059](#) MR [2521475](#)
- [32] Ramakrishnan, D.: Decomposition and parity of Galois representations attached to $\mathrm{GL}(4)$. In: *Automorphic Representations and L -functions*, Tata Inst. Fund. Res. Stud. Math. 22, Tata Inst. Fund. Res., Mumbai, 427–454 (2013) Zbl [1316.11040](#) MR [3156860](#)
- [33] Renard, D.: Représentations des groupes réductifs p -adiques. *Cours Spécialisés* 17, Soc. Math. France, Paris (2010) Zbl [1186.22020](#) MR [2567785](#)
- [34] Roberts, B., Schmidt, R.: Local Newforms for $\mathrm{GSp}(4)$. *Lecture Notes in Math.* 1918, Springer, Berlin (2007) Zbl [1126.11027](#) MR [2344630](#)
- [35] Roberts, B., Schmidt, R.: Some results on Bessel functionals for $\mathrm{GSp}(4)$. *Doc. Math.* **21**, 467–553 (2016) Zbl [1417.11088](#) MR [3522249](#)
- [36] Rösner, M., Weissauer, R.: Regular poles for spinor L -series attached to split Bessel models of $\mathrm{GSp}(4)$. *arXiv:1711.07409* (2017)
- [37] Rubin, K.: Euler Systems. *Ann. of Math. Stud.* 147, Princeton Univ. Press, Princeton, NJ (2000) Zbl [0977.11001](#) MR [1749177](#)
- [38] Schappacher, N., Scholl, A. J.: The boundary of the Eisenstein symbol. *Math. Ann.* **290**, 303–321 (1991) Zbl [0729.11027](#) MR [1109636](#)
- [39] Scholl, A. J.: On modular units. *Math. Ann.* **285**, 503–510 (1989) Zbl [0663.10025](#) MR [1019715](#)
- [40] Sugano, T.: On holomorphic cusp forms on quaternion unitary groups of degree 2. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **31**, 521–568 (1985) Zbl [0559.10020](#) MR [776691](#)
- [41] Taylor, R. L.: On congruences between modular forms. PhD thesis, Princeton Univ. (1988) MR [2636500](#)
- [42] Tilouine, J., Urban, E.: Several-variable p -adic families of Siegel–Hilbert cusp eigensystems and their Galois representations. *Ann. Sci. École Norm. Sup. (4)* **32**, 499–574 (1999) Zbl [0991.11016](#) MR [1693583](#)
- [43] Torzewski, A.: Functoriality of motivic lifts of the canonical construction. *Manuscripta Math.* **163**, 27–56 (2020) Zbl [1454.14012](#) MR [4131990](#)
- [44] Urban, E.: Sur les représentations p -adiques associées aux représentations cuspidales de $\mathrm{GSp}_4/\mathbb{Q}$. *Astérisque* **302**, 151–176 (2005) Zbl [1100.11017](#) MR [2234861](#)
- [45] Waldspurger, J.-L.: Une variante d’un résultat de Aizenbud, Gourevitch, Rallis et Schiffmann. *Astérisque* **346**, 313–318 (2012) Zbl [1308.22008](#) MR [3202559](#)
- [46] Weissauer, R.: Four dimensional Galois representations. *Astérisque* **302**, 67–150 (2005) Zbl [1097.11027](#) MR [2234860](#)

-
- [47] Weissauer, R.: Existence of Whittaker models related to four dimensional symplectic Galois representations. In: *Modular Forms on Schiermonnikoog*, Cambridge Univ. Press, Cambridge, 285–310 (2008) Zbl [1228.11080](#) MR [2530981](#)
- [48] Xu, B.: L -packets of quasisplit $\mathrm{GSp}(2n)$ and $\mathrm{GO}(2n)$. *Math. Ann.* **370**, 71–189 (2018) Zbl [1384.22008](#) MR [3747484](#)