

ARTICLE

Warmer and wetter climate promotes net primary production in C₄ grassland with additional enhancement by hay harvesting

Chang Gyo Jung^{1,2,3}  | Xia Xu⁴ | Zheng Shi⁵ | Shuli Niu^{6,7} | Jianyang Xia^{8,9} | Rebecca Sherry¹⁰ | Lifan Jiang² | Kai Zhu¹¹  | Enqing Hou²  | Yiqi Luo^{1,2}

¹Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona, USA

²Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, Arizona, USA

³Department of Biology, University of Central Florida, Orlando, Florida, USA

⁴College of Biology and the Environment, Nanjing Forestry University, Nanjing, China

⁵Department of Ecology and Evolutionary Biology, University of California Irvine, Irvine, California, USA

⁶Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, China

⁷Department of Resources and Environment, University of Chinese Academy of Sciences, Beijing, China

⁸Tiantong National Forest Ecosystem Observation and Research Station, School of Ecological and Environmental Sciences, East China Normal University, Shanghai, China

⁹Research Center for Global Change and Ecological Forecasting, East China Normal University, Shanghai, China

¹⁰Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma, USA

¹¹Department of Environmental Studies, University of California, Santa Cruz, California, USA

Correspondence

Chang Gyo Jung

Email: cg.jung86@gmail.com

Funding information

US National Science Foundation, Grant/Award Numbers: DBI 0850290, DEB 0444518, DEB 0743778, DEB 0840964, EPS 0919466, OIA-1301789

Handling Editor: Andrew Felton

Abstract

Grassland ecosystems provide essential services to society. To maintain ecosystem functions and services of grasslands under changing environments, it is critical to understand how grasslands respond and feedback to climate change. Here, we present results from a long-term (16 years) warming and clipping (to mimic hay harvesting or grazing) experiment conducted in a grassland ecosystem dominated by C₄ grasses in the Great Plains, USA. We analyzed responses of net primary production (NPP), aboveground NPP (ANPP), and belowground NPP (BNPP) to the expanded ranges of climate conditions observed in the experiment. NPP, ANPP, and BNPP all responded to the climatic variables nonlinearly, with higher degrees of nonlinearity for BNPP than that for ANPP. BNPP peaked at the optimum air temperature of 17.5–18.5°C while ANPP peaked at the optimum air temperature of 18.5–20.0°C. The optimum air temperatures for both ANPP and BNPP were higher than mean annual temperature (16.3°C) at the site. The optimum precipitation for BNPP was intermediate levels (775–1250 mm) while ANPP was maximized in wetter conditions (1250–1605 mm) compared with BNPP. Clipping significantly

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Author(s). *Ecosphere* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

reduced ANPP and shifted the optimum conditions for BNPP to warmer and wetter conditions. In summary, C₄-grass dominant ecosystems have the potential to increase NPP in future warmer and wetter conditions, and clipping may amplify this positive effect in this grassland ecosystem.

KEYWORDS

climate change, clipping, grassland, hay harvesting, net primary production, warming experiment

INTRODUCTION

Net primary production (NPP) is a carbon (C) flux that links terrestrial ecosystems with the atmosphere and thus it is a major process that can trigger ecosystem C cycle feedbacks to climate change (Field et al., 2007; Luo, 2007). As many models have widely projected increasing temperature trends along with altered precipitation regimes in future, it is vital to understand how NPP responds to changes in temperature and precipitation and feedbacks to climate change (Franklin et al., 2016; IPCC, 2013a; Sala et al., 2017). Yet, predicting responses of NPP to climate change is still challenging due to the nonlinearity and the complex interactions between climatic factors (Huston, 1997; Yachi & Loreau, 1999).

The roles of temperature and precipitation in regulating ecosystem production varied across ecosystems as well as different components of NPP, that is, aboveground NPP (ANPP) and belowground NPP (BNPP) (Field et al., 2007; Luo, 2007; Mooney & Gulmon, 1979). For example, warming stimulated ANPP in colder ecosystems (Rustad et al., 2001), but reduced ANPP in warmer ecosystems (Lin et al., 2010). Similarly, above-average temperature was beneficial to sagebrush productions in colder areas but detrimental in hotter areas (Kleinhesselink & Adler, 2018). Warmer climate enhanced foliage biomass while root biomass was inhibited by warmer climate in forests (Lilley et al., 2001). In grass-legume ecosystems, BNPP was also found to have a negative relationship with temperature (Lilley et al., 2001; Reich et al., 2014).

In addition to temperature, changes in precipitation can also have significant impacts on ecosystem production. BNPP increased from drought to wetter treatments in a mixed-grass prairie (Byrne et al., 2013). Similarly, BNPP in a tallgrass prairie had a positive relationship with increased precipitation (Xu et al., 2012). In a synthesis study, Wu et al. (2011) concluded that increases in precipitation stimulated ANPP and BNPP. In contrast, other studies reported no effects of increased precipitation on ecosystem production (Kleinhesselink & Adler, 2018; Rustad et al., 2001). Temperature and precipitation could also interactively affect ecosystem production, but the interactive effects were

generally smaller than the single effect of temperature or precipitation (Wu et al., 2011). Importantly, studies on interactive effects of temperature and precipitation are still inadequate to fully understand when and where climate change would stimulate or depress ecosystem production.

Due to the existence of nonlinear responses of ecosystem production to temperature and precipitation, identifying optimum climatic conditions for ecosystem production is critical to assess whether climate change will increase or decrease ecosystem production. Numerous studies have observed nonlinear responses of ecosystem production (i.e., NPP, ANPP, and BNPP) to temperature and precipitation (Knapp et al., 2016; Luo et al., 2017; Schlenker & Roberts, 2009; Zhu et al., 2016). For example, according to Zhu et al. (2016), ecosystem production generally exhibited saturated or symmetrical responses to changing climatic conditions. Wilcox et al. (2017) also found that the sensitivity of BNPP to altered precipitation declined with mean annual precipitation. With increasing frequency of climate extremes (IPCC, 2013a, 2013b), it is important to understand whether nonlinear responses of ecosystem production to temperature and precipitation are likely to become more common.

Moreover, land-use management, such as hay harvesting and grazing, can not only directly influence plant production but also have indirect effects on plant production through its interactions with climate variables (Canadell & Schulze, 2014; Gao et al., 2008). As hay harvesting or grazing physically removes aboveground parts of plants, it usually leads to higher temperature and lower soil moisture by changing light availability (Collins et al., 1998; Wan et al., 2002). Alternatively, clipping may help conserve soil moisture by reducing transpiration of leaves (Frank et al., 2018). Clipping decreased ANPP but increased carbon allocation to belowground production (Gao et al., 2011; Shi et al., 2016; Xu et al., 2012). Furthermore, clipping was found to modify precipitation effects on ANPP and BNPP in a tallgrass prairie, depending on levels of precipitation: negative and positive precipitation effects under normal and lower precipitation conditions, respectively (Xu et al., 2013). Therefore, the regulations of clipping on the responses of ecosystem production to climate change need to be further examined.

In this study, we examined how ecosystem production (i.e., NPP, ANPP, and BNPP) of a grassland dominated by C_4 plants responded to climate condition (i.e., temperature and precipitation) and how the responses were modified by land management practice in a long-term (16 years) warming and clipping experiment in the Great Plains, USA. We tested three hypotheses: (1) ecosystem production (i.e., NPP, ANPP, and BNPP) nonlinearly responds to temperature and precipitation; (2) interactions between temperature and precipitation positively affect ecosystem production; and (3) clipping reduces ecosystem production in higher levels of climatic conditions (i.e., warmer and wetter conditions). We further projected NPP, ANPP, and BNPP in a temperature-precipitation space to determine the optimum climate conditions for ecosystem production using the relationship between climate variables and ecosystem production.

MATERIALS AND METHODS

Descriptions of the long-term experiment

The long-term experimental site was located in the Kessler Atmospheric and Ecological Field Station in central Oklahoma in the Great Plains, USA (34°58'31.8"N, 97°31'19.6"W), which had remained uncultivated and ungrazed for 40 years before the experiment began in 1999. The site was mostly dominated by C_4 grasses (*Schizachyrium scoparium* and *Sorghastrum nutans*) and few C_3 forbs (*Ambrosia psilostachya*, *Solidago nemoralis*, and *Solidago rigida*). During the experimental period from 1999 to 2015, mean annual temperature for this site was 16.0°C, ranging from 15.3 to 20.0°C. Mean annual precipitation of the study period was 837 mm, ranging from 515 to 1605 mm. The driest year (2005) and wettest year (2015) were classified by the Standardized Precipitation Evapotranspiration Index (SPEI), with SPEI being < -2 and > 2 , respectively (Jung et al., 2019).

In 1999, 12 square plots of 2 m in length were established. Each plot was subdivided into four equal-sized subplots (1 m $\times$ 1 m). The experimental design was a nested design with warming as the main factor and clipping as a nested treatment with six replicates. To treat the plots for continuous warming year-round, we suspended infrared (IR) heaters (Kalglo Electronics, Bethlehem, PA, USA) 1.5 m above the ground with a radiation output of 100 W/m². The control plots with ambient temperature had dummy heaters to ensure a similar shading effect. There was a buffer area of 3 m between the warmed and control plots to avoid heating the control plots. Plants in two diagonal subplots were

clipped at a height of 10 cm above the ground once a year in peak biomass season, usually in July or August to mimic hay harvesting. The other two subplots remained unclipped. After clipping the plants, the materials were removed from the plots, used for measurements of ANPP, and not returned back to the plots.

IR warming can cause dry-down of shallow soil layers, especially during those periods with less rainfall as artifact effects (Kimball et al., 2018; Sherwood & Fu, 2014). However, our measurements showed that soil moisture content in the warmed plots was not significantly lower than that in the control plots in most of the time (except winter time) (Jung et al., 2019; Wan et al., 2002). Therefore, this warming method simulated the environmental conditions under future climate warming (1.4–5.8°C in the 21st century) (IPCC, 2013a; Wan et al., 2002). On average, daily mean air temperature was increased by 1.4°C in our warming experiment (Jung et al., 2019).

Measurements of climatic data

Air temperature was measured by sheltered thermocouples at a height of 25 cm above the ground in the center of each control and warming plots. Detailed information on the air temperature measurements has been described previously (Luo et al., 2009). Missing data of temperature due to mechanical issues of the data logger or the probes were estimated through regressions between available data and reference air temperatures from the Washington station of Oklahoma Mesonet, which was located 200 m away from the study site ($R^2 > 0.98$ and $p < 0.01$) (Brock et al., 1995; McPherson et al., 2007). The annual precipitation data during the study period from 2000 to 2015 were obtained from the Mesonet station.

The radiative energy emitted by the IR heating system will be converted to thermal energy when the radiation hits any objects, such as leaves and soil surface. Therefore, the air temperature measured at 25 cm above the ground within a canopy, which was approximately 1 m tall, over the growing season potentially reflected aerodynamic mixing of heat dissipated from the leaves, soil surface, and ambient air. In general, the measured air temperature in this study could closely represent leaf temperature because boundary layer conductance was high for the narrow grass leaves and wind speed was high in this Great Plains area (Brock et al., 1995; Gates et al., 1965; McPherson et al., 2007; Sinolair, 1970; Vogel, 2009). Thus, as a grass-dominant ecosystem with high wind speed, air temperature in this ecosystem was a suitable metric for assessments of temperature effects on plant production.

Plant production measurements

NPP and its components, ANPP and BNPP, were measured at peak plant biomass each year. For ANPP, we weighed oven-dried (65°C for 72 h) clipped plant biomass as described earlier for the clipping treatment. Plant biomass, that is, ANPP, in unclipped plots was estimated indirectly with a pin-contact method (Frank & McNaughton, 1990). Detailed procedure for estimating aboveground biomass in unclipped plots was described previously in Sherry et al. (2008). Briefly, we counted total hits of C_3 and C_4 plants in both unclipped and clipped plots in peak biomass season, including four directions that the pin frame faced. Then we established correlations between total hits and biomass of both C_3 and C_4 plants in the clipped plots, using a regression method. These correlations were applied to the unclipped plots to estimate the biomass in unclipped plots. The coefficients of the regressions ranged from 0.51 in August of 2002 to 0.84 in August of 2003 (Luo et al., 2009). BNPP was estimated with the root ingrowth-core method (Gao et al., 2008). We took soil samples sequentially from the following three depths: 0–15, 15–30, and 30–45 cm, with soil cores of 4.05 cm in diameter in both unclipped and clipped subplots within each plot in the fall of each year. The sequential soil cores were taken from the same spot in each subplot to estimate annual root growth, that is, BNPP. We backfilled the holes using soils in similar layers from adjacent area; that is, we took soils from shallow (0–30 cm) and deep (30–90 cm) layers and sieved soils to remove roots before putting into the holes. Root samples were gently washed, oven-dried at 70°C for 48 h, and weighed to calculate BNPP. For this analysis, root biomass from the entire depth of 0–45 cm was used. No root samples were taken in 2000–2004 and root samples were not processed in 2011, so data of BNPP and NPP were missing for these years.

Data analysis

We used the linear-mixed effect model in this study. This model has been applied to the Jasper Ridge Global Change Experiment and its assumptions (i.e., linear and nonlinear terms as main effects and interactive term as interactive effect) have been described in detail (Zhu et al., 2016). In this study, we used two environmental variables, annual mean air temperature (T) and annual precipitation (P), as the continuous variables and clipping treatment (C) as the categorical variable to represent whether or not a plot was clipped.

To conduct the analysis, we first tested time-dependent effects, that is, the progressive effect, of treatments

(i.e., warming, clipping, and interaction of warming and clipping) using standard analysis of variance (ANOVA). Despite significant treatment effects in some years, coefficients of ANOVA did not show significant temporal trends (i.e., $p > 0.05$ for the linear regressions between years and significant coefficients of ANOVA) (Appendix S1: Figure S1).

To test how the ecosystem production responded to the environmental factors, in the linear mixed-effects model, the main and interactive environmental factors were set as fixed effects and the plots were set as random effects (Bates et al., 2015). Linear functions for T , P , and C , as well as quadratic functions for T^2 and P^2 were used. Since there were only two environmental factors, temperature–precipitation interaction ($T:P$) was used to test their interactive effect. Ecosystem production, that is, NPP, ANPP, and BNPP, was log-transformed to satisfy normality before any statistical analysis. To ensure the environmental variables were statistically comparable, T and P were standardized by subtracting mean and divided by standard deviation (SD). Because of the log-scaled ecosystem production and standardized environmental variables, the estimated climate effects were interpreted as proportional changes in plant production ($d \log y$, where d was the proportional change and y was NPP, ANPP, or BNPP) with respect to the change of a SD in environmental parameters (dx , where x was a SD of T or P), $d \log y/dx = (dy/y)/dx$. The estimated clipping effects were interpreted as changes in log plant production with respect to the clipping treatment (C). The estimated coefficients for environmental factors (T and P) and clipping treatment (C) were summarized in Table 1.

The nonlinear model with a quadratic function was selected since it fit our data better than the linear model according to Akaike information criterion (AIC). AIC is an index of the goodness of fit of a model, including a penalty for overfitting. We used a model having lower AIC value. Although delta AIC (difference between nonlinear and linear models) of ANPP was relatively smaller (i.e., 3.0) than delta AIC of NPP (i.e., 44.9) and BNPP (i.e., 56.3), nonlinear model for ANPP was the best model and substantially different from the linear model since delta AIC of ANPP was larger than 2 (Burnham & Anderson, 2004). The selected models explained 29% (NPP), 31% (ANPP), and 29% (BNPP) of the observed variations. AIC values of linear and nonlinear models for NPP, ANPP, and BNPP were shown in Appendix S1: Table S1.

The nonlinear model, which was selected as the best one, was then used to predict the plant production in the response surfaces. We calculated expected plant production using the environmental variables for each treatment. Continuous temperature and precipitation used for

TABLE 1 Model coefficients (means and 95% confidence intervals)

Coefficient	NPP	ANPP	BNPP
T^2	−0.039 (−0.075, −0.003)	0.005 (−0.029, 0.039)	−0.107 (−0.176, −0.039)
P^2	−0.131 (−0.166, −0.095)	−0.036 (−0.062, −0.009)	−0.274 (−0.342, −0.207)
T	0.118 (0.065, 0.171)	0.096 (0.044, 0.148)	0.092 (−0.01, 0.194)
P	0.148 (0.088, 0.208)	0.153 (0.095, 0.211)	0.111 (−0.004, 0.225)
C	−0.003 (−0.075, 0.069)	−0.167 (−0.237, −0.098)	0.056 (−0.082, 0.193)
$T:P$	0.034 (−0.03, 0.098)	0.06 (0.002, 0.119)	−0.035 (−0.157, 0.088)
$T:C$	−0.011 (−0.085, 0.063)	−0.009 (−0.081, 0.062)	0.037 (−0.104, 0.178)
$P:C$	−0.01 (−0.082, 0.062)	0.063 (−0.006, 0.133)	−0.043 (−0.18, 0.094)
$T:P:C$	0.07 (−0.018, 0.159)	−0.046 (−0.125, 0.033)	0.193 (0.025, 0.362)

Note: Model structure: Plant production (net primary production [NPP], aboveground NPP [ANPP], or belowground NPP [BNPP]) $\sim T^2 + P^2 + T + P + C + T:P + T:C + P:C + T:P:C + (1|\text{Plot})$. T : Temperature; P : Precipitation; C : Clipping; $T:P$: Temperature–precipitation interaction; $T:C$: Temperature–clipping interaction; $P:C$: Precipitation–clipping interaction; $T:P:C$: Temperature–precipitation–clipping interaction.

the predictions were 15.2–20.1°C and 485–1650 mm, respectively. Categorical factors for clipping treatment were 0 for unclipped control and 1 for clipped treatment. We then visualized the prediction results by temperature and precipitation under unclipped or clipped conditions.

To evaluate model performance, we did in-sample predictions to compare the modeled plant production with the observed plant production. Overall, the model predicted plant production well ($r_{\text{NPP}} = 0.65$, $r_{\text{ANPP}} = 0.55$, and $r_{\text{BNPP}} = 0.58$; Appendix S1: Figure S1a). Modeled and observed NPP, ANPP, and BNPP were close to the 1:1 reference line.

To test model assumptions of normality, residual diagnostics regarding progressive (year-dependent) effects were performed. If progressive year effects were present, the residual (i.e., unexplained component of observed data) would be correlated with year. However, the residuals in diagnostics did not show such correlations (Appendix S1: Figure S2b). The histogram in Appendix S1: Figure S2c validated the assumption of normality of the model residuals. All analyses were performed with the packages “lme4” (Bates et al., 2015) and “ggplot2” (Wickham, 2009) in R (version 3.3.1) (R Core Team, 2016).

RESULTS

Nonlinear responses of plant production to temperature and precipitation

In the study period, annual air temperature ranged from 15.3 to 20.0°C with a threefold difference in annual precipitation (505–1605 mm) (Figure 1). The ranges of precipitation in this study fell beyond from 10th to 90th percentiles over the past 120 years of historical precipitation (Jung




FIGURE 1 Nonlinear responses of plant production to air temperature and precipitation. Hollow and filled circles indicate ambient and warming treatment, respectively. Black, red, and blue colors represent net primary production (NPP) (a), aboveground NPP (ANPP) (b), and belowground NPP (BNPP) (c), respectively. Gray shading shows 95% confidence interval

et al., 2019). Both relationships between NPP and BNPP and climatic variables, that is, air temperature and precipitation, showed unimodal shapes (hump-shape) (Figure 1). Negative warming effects on BNPP were larger than ANPP, resulting in decreases in NPP with higher




FIGURE 2 Standardized model coefficients. Circles indicate model's coefficients within 95% confidence interval. Sample sizes of net primary production (NPP), aboveground NPP (ANPP), and belowground NPP (BNPP) are 240, 384, and 240, respectively. T : Temperature; P : Precipitation; C : Clipping; $T:P$: Temperature–precipitation interaction; $T:C$: Temperature–clipping interaction; $P:C$: Precipitation–clipping interaction; $T:P:C$: Temperature–precipitation–clipping interaction

temperature (Figure 1). For example, warming caused hump-shaped responses of NPP and BNPP, both of which peaked at the intermediate temperature range (17.5–18.5°C) that was slightly higher than the long-term averaged temperature (17.2°C). However, ANPP did not exhibit a clear hump-shaped relationship in comparison to NPP and BNPP. Similar to the responses of plant production to air temperature, clear hump-shaped relationships of NPP and BNPP with precipitation were observed, having the peak production at the intermediate level of precipitation (900–1100 mm; Figure 1), which was slightly higher than long-term averaged precipitation (873 mm). ANPP also exhibited a nonlinear response to precipitation but followed a saturating rather than hump-shaped pattern (Figure 1).

Coefficients for the linear mixed effect model

Except for ANPP response to temperature, the standardized quadratic coefficients (T^2 and P^2) in the linear mixed effect model for fitting the relationship between plant production and temperature or precipitation were all significantly negative (Table 1 and Figure 2). That is, the response curves of plant production to climatic factors had concave down hump-shapes (Figure 1). The only exception to this pattern was the relationship between ANPP and precipitation, which showed a saturated pattern at the high level of precipitation and could be explained by the slightly significantly negative standardized quadratic coefficient (Figures 1 and 2).

The coefficients for the linear terms of temperature (T) and precipitation (P) in the mixed effect

model were all positive, indicating increases in plant production linearly with temperature and precipitation, peaking at higher values of temperature and precipitation with exception of BNPP. Clipping (C) significantly decreased ANPP while it had no significant effects on either NPP or BNPP. The temperature–precipitation interaction was significantly positive for ANPP, suggesting that the interaction between temperature and precipitation would be an additional positive effect on ANPP since the single effects of temperature and precipitation were both positive. In addition, interactive effect among temperature, precipitation and clipping ($T:P:C$) on BNPP was significantly positive, meaning that the positive effects were intensified when temperature, precipitation, and clipping effects were combined.

Response of plant production to temperature and precipitation in response surfaces

Modeled plant production and its respective contour lines were codetermined by the main effects of temperature, precipitation, and clipping and their interactive effects. Within observed climate ranges, the response surfaces of NPP, ANPP, and BNPP generally showed a rising ridge pattern under unclipped and clipped conditions, that is, a simple maximum pattern, with exception of BNPP under unclipped condition (Figure 3). Temperature had a stronger effect on NPP and ANPP at higher precipitation levels, that is, between the long-term average (873 mm, horizontal gray dashed line in Figure 3) and the maximum precipitation (1650 mm). The maximum NPP and




FIGURE 3 Predicted and observed net primary production (NPP) (a), aboveground NPP (ANPP) (b), and belowground NPP (BNPP) (c). The x- and y-axes are continuous air temperature (15.2–20.1°C) and annual precipitation (485–1650 mm), respectively. Gradient colors with contour lines indicate modeled results. Bubbles represent observed data collected from each plot. Vertical and horizontal gray dash lines indicate long-term average of temperature (17.2°C) and annual precipitation (873 mm), respectively

ANPP both occurred at high levels of temperature and precipitation (Figure 3a,b). In contrast, within the precipitation range between the long-term average (horizontal gray dashed line in Figure 3c) and 1490 mm, temperature increases strongly stimulated BNPP (Figure 3c). BNPP under the unclipped condition first increased with increase in temperature when precipitation was at relatively low level but started to decline with increased temperature when precipitation exceeded 1490 mm. Alternatively, BNPP can be decreased when precipitation level was drier conditions (i.e., lower than long-term average), but stimulated by decreases in precipitation from 1650 to 1490 mm. Together,

BNPP reached its maximum values at intermediate level of precipitation (1000–1250 mm) and relatively high temperature (Figure 3c). Overall, clipping decreased ANPP in the response surface, but the clipping treatment shifted the optimum environmental conditions for NPP and BNPP to higher levels of temperature and precipitation.

DISCUSSION

By analyzing NPP, ANPP, and BNPP from a long-term warming and clipping experiment of 16 years, we tested

three hypotheses regarding (1) nonlinear responses of ecosystem production to climatic variables, (2) interactive effects of climatic variables on ecosystem production, and (3) clipping impacts on the interaction of climate variables in regulating ecosystem production. Firstly, we found nonlinear relationships between ecosystem production and climatic variables (i.e., temperature and precipitation), except for the relationship between ANPP and temperature. Secondly, we detected a positive interactive effect of temperature and precipitation on ANPP. Lastly, clipping only amplified the interactive effect of temperature and precipitation on BNPP. Projections of NPP, ANPP, and BNPP based on those relationships revealed that warmer and wetter conditions were the optimum conditions for NPP and ANPP while BNPP reached peak values under warmer temperature but intermediate precipitation conditions. In addition, clipping shifted peak NPP and BNPP to higher temperature and precipitation levels while ANPP was significantly reduced under clipping.

Warmer and wetter climate conditions stimulate plant production

Previous studies have demonstrated that response curves of ecosystem production against climate gradients were nonlinear, that is, symmetrical and asymmetrical responses of NPP to temperature or precipitation (Knapp et al., 2017; Luo et al., 2017; Wilcox et al., 2017; Zhu et al., 2016). Consistently, our results exhibited nonlinear responses—plant production increased with climatic variables until it reached a peak at optimal condition, and then slightly decreased or saturated under higher levels of climatic conditions possibly due to life-history constraints (Hsu et al., 2012; Zhu et al., 2016). Similar patterns were reported previously in a C_3 -dominated grassland in California (Zhu et al., 2016) and they found long-term average climate condition was the optimum condition for NPP, showing concave down responses to temperature and precipitation. When temperature was high and soil was moist, plants naturally tended to minimize heat stress by cooling their leaves through increased transpiration and thus more water loss (Crawford et al., 2012), which might exert a negative effect for plant production. In addition, growth suppression under high precipitation might also be attributed to more frequent cloudy days (thus less radiation for photosynthesis) and nutrient leaching (Reichstein et al., 2013; Wang et al., 2020; Zhu et al., 2016). Contrary to our results, ANPP of semiarid ecosystems showed a positive asymmetry response to precipitation: the magnitude of stimulation under extreme wet conditions was greater than the

magnitude in decrease under extreme dry conditions (Felton et al., 2019). The contrasting results might be caused by different plant community compositions and ecoregion as Felton et al. (2019) have observed varied sensitivities of plant production to climate among different plant functional types in a semiarid area, that is, higher positive asymmetry of forbs than C_4 grasses.

Interestingly, while single factor of temperature or precipitation played a negative role in their higher levels in our observations (Figure 1) as well as previous studies (Schlenker & Roberts, 2009; Zhu et al., 2016), in our study site NPP would peak at warmer and wetter conditions due to combined effects of temperature and precipitation as well as their interaction (Figure 3). The negative effect of temperature or precipitation on NPP in their higher levels eliminated when the single-factor effects of temperature and precipitation were combined (Figure 3), as reflected in the response space without interaction effects (e.g., absence of $T:P$ and $T:P:C$ in the model; Appendix S1: Figure S3). Both empirical and modeling studies across different ecosystems (e.g., tropical, temperate, and boreal forests and grasslands) suggested that the single positive effects of temperature or precipitation on NPP would be intensified when warming was combined with additional precipitation inputs (Luo et al., 2008; Schuur, 2003; Wu et al., 2011). Since our study site was dominated by C_4 grasses, this ecosystem was expected to have higher optimum temperature and water use efficiency (Way et al., 2014; Yamori et al., 2014). While warm years were usually not wet at this site during the study period (Appendix S1: Figure S4), warmer and wetter conditions would favor production of C_4 plant species (Lundgren & Christin, 2016). In addition, more rainfall during warm months could benefit C_4 grasses (Knapp et al., 2020). As anomalies of temperature and precipitation in the Great Plains kept increasing over time (Kunkel et al., 2013), our results imply that this grassland ecosystem has a potential to increase NPP under the possible warmer and wetter climate (Allan & Soden, 2008; Greve et al., 2014; Wentz et al., 2007).

Contrasting responses of ANPP and BNPP to climatic variables

Response surfaces of ANPP and BNPP exhibited different patterns in this study as a result of their different responses to temperature and precipitation. Temperature had a significantly positive effect on ANPP and marginally positive effect on BNPP (i.e., positive coefficients of T in Figure 2). These results were consistent with previous studies documenting positive temperature effects on NPP, ANPP, and BNPP within a single

ecosystem (Litton & Giardina, 2008; Reich et al., 2014; Xu et al., 2012; Xu et al., 2013) as well as across different ecosystems based on a meta-analysis (Wu et al., 2011). In fact, an earlier analysis from the same experiment as the present study concluded that increased temperature exerted positive impacts on ANPP and BNPP due to high rain use efficiency in this ecosystem (Xu et al., 2012). In contrast, in a C_3 -grass dominant ecosystem, negative temperature effects on NPP, ANPP, and BNPP were detected (Zhu et al., 2016), which was distinct from the present study in a C_4 -grass dominant ecosystem. It has been well understood that the optimum temperature for photosynthesis of C_4 species was generally higher than that for C_3 species (Yamori et al., 2014). Although ANPP and BNPP had similar positive responses to temperature, BNPP had a greater degree of nonlinearity than ANPP (more negative coefficient of T^2 in BNPP than ANPP in Figure 2), which might be due to higher temperature sensitivity of BNPP than ANPP (Gibson, 2009; Xu et al., 2013).

Increased precipitation enhanced ANPP before saturating at the high level of precipitation in this study (Figure 1 and significant positive and negative coefficients of P and P^2 for ANPP, respectively, in Figure 2), which was partly consistent with previous studies (Hsu et al., 2012; Huxman et al., 2004; Sala et al., 2012; Ye et al., 2017). Similarly, BNPP showed marginally significant positive response to precipitation (coefficient of P for BNPP in Figure 2) but presented greater nonlinearity than ANPP (Figure 1 and more negative coefficient of P^2 for BNPP than ANPP in Figure 2). Due to the limited number of studies about BNPP response to a wide range of precipitation, fully comparing our results with previous findings is difficult, but previous water addition treatments were found to stimulate BNPP (Wilcox et al., 2015; Xu et al., 2013). A recent synthesis demonstrated that extremely wet conditions triggered negative responses of BNPP to precipitation while ANPP kept increasing with wetter conditions (Wilcox et al., 2017), which supported our nonlinear response of BNPP to precipitation and an increasing pattern of ANPP under high precipitation despite slightly nonlinear responses. The decline of BNPP under an extremely wet condition would be attributed to decrease in root lifespan due to water logging of soil (Kozlowski, 1997).

Given these various responses of ANPP and BNPP to climate variables, different optimal conditions for ANPP and BNPP were predicted in the response surfaces. As for ANPP, the warmer and wetter conditions were optimal conditions, which was in line with previous studies: A positive warming effect on ANPP could be intensified by additional precipitation (Dukes et al., 2005; Wu et al., 2011). This effect of precipitation on ANPP was clear within intermediate–high temperature while a nonlinear effect of precipitation on BNPP was identified

in the same temperature range. The nonlinear effects of precipitation on BNPP were exhibited under both drier and wetter conditions, which has been observed from previous studies (Kozlowski, 1997; Xu et al., 2013).

The regulatory role of clipping on the climate change effects

Clipping-mediated reduction in ANPP found in this study was consistent with previous studies (Carlyle et al., 2014; Shi et al., 2016). Grazing-mediated reductions in ANPP were linked to soil nutrient loss in other ecosystems (Giese et al., 2013; Liu et al., 2015); this was a potential cause of the observed reduction in ANPP in clipped plots even though we did not measure it in this study. In addition, there were less organic material inputs by removing plants (Luo et al., 2009). While clipping reduced ANPP in the overall response space, ANPP was still maximized in the warmer and wetter condition, same as under the unclipped condition (Figure 3b), indicating that clipping would not change the directions of the effects of temperature and precipitation and their interaction on ANPP. Previous studies illustrated that clipping increased temperature but decreased soil moisture by reducing shading, leading to negative effects on plant production under normal precipitation condition, but positive effects on plant production under the limited precipitation condition due to less water demands by removing plants (Li et al., 2011; Xu et al., 2012, 2013). Yet, our analysis did not detect such patterns of clipping effects. The combination of warming and clipping might have exacerbated not only soil nutrient loss via soil erosion (Xue et al., 2011), but also soil water loss, especially under water addition experiments (Zhou et al., 2006). This might explain clipping-mediated suppression of ANPP under warming via nutrient loss and stress of temperature and soil moisture (Xue et al., 2011).

On the other hand, clipping significantly altered the response surface of BNPP from simple maximum to rising ridge pattern due to its interactions with climatic variables. Clipping enhanced warming effects on BNPP, especially under higher precipitation levels due to increased rain use efficiency (Xu et al., 2012). Furthermore, BNPP was increased by warming and clipping treatments (Xu et al., 2012, 2013, 2014), which supported our results, that is, higher BNPP under clipped warmer and wetter conditions. These increases led to the pattern of stimulated NPP under warmer and wetter conditions.

Our results from a long-term experimental study suggested that warmer and wetter climate conditions, one of the climate scenarios predicted for the future in

this region, could further enhance NPP in this C₄-grass dominant ecosystem. Plant production in this ecosystem responded nonlinearly to temperature or precipitation individually, with slightly negative effects under warmer or wetter conditions. However, the overall positive effects of temperature or precipitation and their interaction would offset the negative effects at high temperature and high level of precipitation. Importantly, clipping stimulated BNPP under warmer and wetter conditions but decreased ANPP with no shift in the optimum condition. Another factor that might contribute to the changes in plant production was a gradual increase in atmospheric CO₂ concentration from 369 ppm in 2000 to 401 ppm in 2015 (Keeling et al., 2005). Especially under warmer and wetter conditions, rising CO₂ concentration has been reported to enhance production of C₄ plants (Augustine et al., 2018) and suppress production of C₃ plants (Zhu et al., 2016). However, forage quality in grassland ecosystems may decrease as a result of increased production, which is a challenge to sustain high livestock yields (Augustine et al., 2018). Future research should include field experiments manipulating both temperature and precipitation as well as elevated CO₂ in order to explore not only effects of climate change on plant production under warmer and wetter conditions beyond what we have observed in this study on plant production to confirm our results, but also the potential interactive effects between climate change and rising atmospheric CO₂ concentration. Overall, our 16-year experiment highlights that possible future climate in this region, that is, high temperature and high precipitation, would favor the increase of NPP, and clipping may amplify this positive effect in this grassland ecosystem.

ACKNOWLEDGMENTS

The authors thank many laboratory members for their help with field works and root washing and weighting tasks. This project was financially supported by US National Science Foundation (NSF) grants DEB 0444518, DEB 0743778, DEB 0840964, DBI 0850290, EPS 0919466, and OIA-1301789.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

Chang Gyo Jung and Yiqi Luo contributed equally to this work. Yiqi Luo designed the experiments. Chang Gyo Jung, Xia Xu, Zheng Shi, Shuli Niu, Jianyang Xia, Rebecca Sherry, and Lifan Jiang collected the data. Chang Gyo Jung, Lifan Jiang, Kai Zhu, Enqing Hou, and Yiqi Luo performed data analyses. All authors discussed the results and wrote the manuscript.

ORCID

Chang Gyo Jung  <https://orcid.org/0000-0002-9845-7732>

Kai Zhu  <https://orcid.org/0000-0003-1587-3317>

Enqing Hou  <https://orcid.org/0000-0003-4864-2347>

REFERENCES

- Allan, R.P., and B.J. Soden. 2008. "Atmospheric Warming and the Amplification of Precipitation Extremes." *Science* 321: 1481–4.
- Augustine, D.J., D.M. Blumenthal, T.L. Springer, D.R. LeCain, S.A. Gunter, and J.D. Derner. 2018. "Elevated CO₂ Induces Substantial and Persistent Declines in Forage Quality Irrespective of Warming in Mixedgrass Prairie." *Ecological Applications* 28: 721–35.
- Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. "Fitting Linear Mixed-Effects Models Using lme4." *Journal of Statistical Software* 67: 1–48.
- Brock, F.V., K.C. Crawford, R.L. Elliott, G.W. Cuperus, S.J. Stadler, H.L. Johnson, and M.D. Eilts. 1995. "The Oklahoma Mesonet: A Technical Overview." *Journal of Atmospheric and Oceanic Technology* 12: 5–19.
- Burnham, K.P., and D.R. Anderson. 2004. "Multimodel Inference: Understanding AIC and BIC in Model Selection." *Sociological Methods & Research* 33: 261–304.
- Byrne, K.M., W.K. Lauenroth, and P.B. Adler. 2013. "Contrasting effects of precipitation manipulations on production in two sites within the central grassland region, USA." *Ecosystems* 16: 1039–1051.
- Canadell, J.G., and E.D. Schulze. 2014. "Global Potential of Biospheric Carbon Management for Climate Mitigation." *Nature Communications* 5: 5282.
- Carlyle, C.N., L.H. Fraser, and R. Turkington. 2014. "Response of Grassland Biomass Production to Simulated Climate Change and Clipping Along an Elevation Gradient." *Oecologia* 174: 1065–73.
- Collins, S.L., A.K. Knapp, J.M. Briggs, J.M. Blair, and E.M. Steinauer. 1998. "Modulation of Diversity by Grazing and Mowing in Native Tallgrass Prairie." *Science* 280: 745–7.
- Crawford, A.J., D.H. McLachlan, A.M. Hetherington, and K.A. Franklin. 2012. "High Temperature Exposure Increases Plant Cooling Capacity." *Current Biology* 22: R396–7.
- Dukes, J.S., N.R. Chiariello, E.E. Cleland, L.A. Moore, M.R. Shaw, S. Thayer, T. Tobeck, H.A. Mooney, and C.B. Field. 2005. "Responses of Grassland Production to Single and Multiple Global Environmental Changes." *PLoS Biology* 3: e319.
- Felton, A.J., S. Zavislan-Pullaro, and M.D. Smith. 2019. "Semiarid Ecosystem Sensitivity to Precipitation Extremes: Weak Evidence for Vegetation Constraints." *Ecology* 100: e02572.
- Field, C.B., D.B. Lobell, H.A. Peters, and N.R. Chiariello. 2007. "Feedbacks of Terrestrial Ecosystems to Climate Change." *Annual Review of Environment and Resources* 32: 1–29.
- Frank, D.A., and S.J. McNaughton. 1990. "Aboveground Biomass Estimation with the Canopy Intercept Method: A Plant Growth Form Caveat." *Oikos* 57: 57–60.
- Frank, D.A., R.L. Wallen, E.W. Hamilton, III, P.J. White, and J.D. Fridley. 2018. "Manipulating the System: How Large Herbivores Control Bottom-Up Regulation of Grasslands." *Journal of Ecology* 106: 434–43.
- Franklin, J., J.M. Serra-Diaz, A.D. Syphard, and H.M. Regan. 2016. "Global Change and Terrestrial Plant Community Dynamics."

- Proceedings of the National Academy of Sciences of the United States of America* 113: 3725–34.
- Gao, Y.Z., M. Giese, S. Lin, B. Sattelmacher, Y. Zhao, and H. Brueck. 2008. “Belowground Net Primary Productivity and Biomass Allocation of a Grassland in Inner Mongolia Is Affected by Grazing Intensity.” *Plant and Soil* 307: 41–50.
- Gao, Y.Z., Q. Chen, S. Lin, M. Giese, and H. Brueck. 2011. “Resource Manipulation Effects on Net Primary Production, Biomass Allocation and Rain-Use Efficiency of Two Semiarid Grassland Sites in Inner Mongolia, China.” *Oecologia* 165: 855–64.
- Gates, D.M., E. Tibbals, and F. Kreith. 1965. “Radiation and Convection for Ponderosa Pine.” *American Journal of Botany* 52: 66–71.
- Gibson, D.J. 2009. *Grasses and Grassland Ecology*. UK: Oxford University Press.
- Giese, M., H. Brueck, Y. Gao, S. Lin, M. Steffens, I. Kögel-Knabner, T. Glindemann, A. Susenbeth, F. Taube, and K. Butterbach-Bahl. 2013. “N Balance and Cycling of Inner Mongolia Typical Steppe: A Comprehensive Case Study of Grazing Effects.” *Ecological Monographs* 83: 195–219.
- Greve, P., B. Orlowsky, B. Mueller, J. Sheffield, M. Reichstein, and S.I. Seneviratne. 2014. “Global Assessment of Trends in Wetting and Drying over Land.” *Nature Geoscience* 7: 716–21.
- Hsu, J.S., J. Powell, and P.B. Adler. 2012. “Sensitivity of Mean Annual Primary Production to Precipitation.” *Global Change Biology* 18: 2246–55.
- Huston, M.A. 1997. “Hidden Treatments in Ecological Experiments: Re-Evaluating the Ecosystem Function of Biodiversity.” *Oecologia* 110: 449–60.
- Huxman, T.E., M.D. Smith, P.A. Fay, A.K. Knapp, M.R. Shaw, M.E. Loik, S.D. Smith, D.T. Tissue, J.C. Zak, and J.F. Weltzin. 2004. “Convergence across Biomes to a Common Rain-Use Efficiency.” *Nature* 429: 651–4.
- IPCC. 2013a. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge and New York: Cambridge University Press.
- IPCC. 2013b. “Summary for Policymakers.” In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley, 1–30. Cambridge and New York: Cambridge University Press.
- Jung, C.G., X. Xu, S. Niu, J. Liang, X. Chen, Z. Shi, L. Jiang, and Y. Luo. 2019. “Experimental Warming Amplified Opposite Impacts of Drought vs. Wet Extremes on Ecosystem Carbon Cycle in a Tallgrass Prairie.” *Agricultural and Forest Meteorology* 276–277: 107635.
- Keeling, C.D., S.C. Piper, R.B. Bacastow, M. Wahlen, T.P. Whorf, M. Heimann, and H.A. Meijer. 2005. “Atmospheric CO₂ and ¹³CO₂ Exchange with the Terrestrial Biosphere and Oceans from 1978 to 2000: Observations and Carbon Cycle Implications.” In *A History of Atmospheric CO₂ and Its Effects on Plants, Animals, and Ecosystems* (vol. 177, pp. 83–113). New York: Springer.
- Kimball, B.A., A.M. Alonso-Rodríguez, M.A. Cavaleri, S.C. Reed, G. González, and T.E. Wood. 2018. “Infrared Heater System for Warming Tropical Forest Understory Plants and Soils.” *Ecology and Evolution* 8: 1932–44.
- Kleinheselink, A.R., and P.B. Adler. 2018. “The Response of Big Sagebrush (*Artemisia tridentata*) to Interannual Climate Variation Changes across its Range.” *Ecology* 99: 1139–49.
- Knapp, A.K., P. Ciais, and M.D. Smith. 2016. “Reconciling Inconsistencies in Precipitation–Productivity Relationships: Implications for Climate Change.” *New Phytologist* 214: 41–7.
- Knapp, A.K., P. Ciais, and M.D. Smith. 2017. “Reconciling Inconsistencies in Precipitation–Productivity Relationships: Implications for Climate Change.” *The New Phytologist* 214: 41–7.
- Knapp, A.K., A. Chen, R.J. Griffin-Nolan, L.E. Baur, C.J.W. Carroll, J.E. Gray, A.M. Hoffman, et al. 2020. “Resolving the Dust Bowl Paradox of Grassland Responses to Extreme Drought.” *Proceedings of the National Academy of Sciences of the United States of America* 117: 22249.
- Kozłowski, T. 1997. “Responses of Woody Plants to Flooding and Salinity.” *Tree Physiology* 17: 490.
- Kunkel, K. E., L. E. Stevens, S. E. Stevens, L. Sun, E. Janssen, D. Wuebbles, M. C. Kruk, D. Thomas, M. Shulski, and N. A. Umphlett. 2013. *Regional Climate Trends and Scenarios for the US National Climate Assessment Part 4. Climate of the US Great Plains*. Washington, DC: U.S. Department of Commerce National Oceanic and Atmospheric Administration.
- Li, J., S. Lin, F. Taube, Q. Pan, and K. Dittert. 2011. “Above and Belowground Net Primary Productivity of Grassland Influenced by Supplemental Water and Nitrogen in Inner Mongolia.” *Plant and Soil* 340: 253–64.
- Lilley, J., T. Bolger, and R. Gifford. 2001. “Productivity of Trifolium Subterraneum and Phalaris Aquatica under Warmer, High CO₂ Conditions.” *New Phytologist* 150: 371–83.
- Lin, D., J. Xia, and S. Wan. 2010. “Climate Warming and Biomass Accumulation of Terrestrial Plants: A Meta-Analysis.” *New Phytologist* 188: 187–98.
- Litton, C., and C. Giardina. 2008. “Below-Ground Carbon Flux and Partitioning: Global Patterns and Response to Temperature.” *Functional Ecology* 22: 941–54.
- Liu, N., H. Kan, G. Yang, and Y. Zhang. 2015. “Changes in Plant, Soil, and Microbes in a Typical Steppe from Simulated Grazing: Explaining Potential Change in Soil C.” *Ecological Monographs* 85: 269–86.
- Lundgren, M.R., and P.-A. Christin. 2016. “Despite Phylogenetic Effects, C3–C4 Lineages Bridge the Ecological Gap to C4 Photosynthesis.” *Journal of Experimental Botany* 68: 241–54.
- Luo, Y. 2007. “Terrestrial Carbon-Cycle Feedback to Climate Warming.” *Annual Review of Ecology, Evolution, and Systematics* 38: 683–712.
- Luo, Y., D. Gerten, G. Le Maire, W.J. Parton, E. Weng, X. Zhou, C. Keough, C. Beier, P. Ciais, and W. Cramer. 2008. “Modeled Interactive Effects of Precipitation, Temperature, and [CO₂] on Ecosystem Carbon and Water Dynamics in Different Climatic Zones.” *Global Change Biology* 14: 1986–99.
- Luo, Y., R. Sherry, X. Zhou, and S. Wan. 2009. “Terrestrial Carbon-Cycle Feedback to Climate Warming: Experimental Evidence on Plant Regulation and Impacts of Biofuel Feedstock Harvest.” *GCB Bioenergy* 1: 62–74.
- Luo, Y., L. Jiang, S. Niu, and X. Zhou. 2017. “Nonlinear Responses of Land Ecosystems to Variation in Precipitation.” *The New Phytologist* 214: 5–7.

- McPherson, R.A., C.A. Fiebrich, K.C. Crawford, J.R. Kilby, D.L. Grimsley, J.E. Martinez, J.B. Basara, B.G. Illston, D.A. Morris, and K.A. Kloesel. 2007. "Statewide Monitoring of the Mesoscale Environment: A Technical Update on the Oklahoma Mesonet." *Journal of Atmospheric and Oceanic Technology* 24: 301–21.
- Mooney, H., and S. Gulmon. 1979. "Environmental and Evolutionary Constraints on the Photosynthetic Characteristics of Higher Plants." In *Topics in Plant Population Biology* (pp. 316–337). New York: Columbia University Press.
- R Core Team. 2016. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Reich, P.B., Y. Luo, J.B. Bradford, H. Poorter, C.H. Perry, and J. Oleksyn. 2014. "Temperature Drives Global Patterns in Forest Biomass Distribution in Leaves, Stems, and Roots." *Proceedings of the National Academy of Sciences of the United States of America* 111: 13721–6.
- Reichstein, M., M. Bahn, P. Ciais, D. Frank, M.D. Mahecha, S.I. Seneviratne, J. Zscheischler, C. Beer, N. Buchmann, and D.C. Frank. 2013. "Climate Extremes and the Carbon Cycle." *Nature* 500: 287–95.
- Rustad, L., J. Campbell, G. Marion, R. Norby, M. Mitchell, A. Hartley, J. Cornelissen, and J. Gurevitch. 2001. "A Meta-Analysis of the Response of Soil Respiration, Net Nitrogen Mineralization, and Aboveground Plant Growth to Experimental Ecosystem Warming." *Oecologia* 126: 543–62.
- Sala, O.E., L.A. Gherardi, L. Reichmann, E. Jobbagy, and D. Peters. 2012. "Legacies of Precipitation Fluctuations on Primary Production: Theory and Data Synthesis." *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 367: 3135–44.
- Sala, O.E., L. Yahdjian, K. Havstad, and M.R. Aguiar. 2017. *Rangeland Ecosystem Services: Nature's Supply and Humans' Demand. Rangeland Systems: Processes, Management, and Challenges* (pp. 467–89). New York: Springer Series on Environmental Management.
- Schlenker, W., and M.J. Roberts. 2009. "Nonlinear Temperature Effects Indicate Severe Damages to US Crop Yields under Climate Change." *Proceedings of the National Academy of Sciences of the United States of America* 106: 15594–8.
- Schuur, E.A. 2003. "Productivity and Global Climate Revisited: The Sensitivity of Tropical Forest Growth to Precipitation." *Ecology* 84: 1165–70.
- Sherry, R.A., E. Weng, J.A. Arnone, III, D.W. Johnson, D.S. Schimel, P.S. Verburg, L.L. Wallace, and Y. Luo. 2008. "Lagged Effects of Experimental Warming and Doubled Precipitation on Annual and Seasonal Aboveground Biomass Production in a Tallgrass Prairie." *Global Change Biology* 14: 2923–36.
- Sherwood, S., and Q. Fu. 2014. "A drier future?" *Science* 343: 737–9.
- Shi, Z., X. Xu, L. Souza, K. Wilcox, L. Jiang, J. Liang, J. Xia, P. García-Palacios, and Y. Luo. 2016. "Dual Mechanisms Regulate Ecosystem Stability under Decade-Long Warming and Hay Harvest." *Nature communications* 7: 1–6.
- Sinclair, R. 1970. "Convective Heat Transfer from Narrow Leaves." *Australian Journal of Biological Sciences* 23: 309–22.
- Vogel, S. 2009. "Leaves in the Lowest and Highest Winds: Temperature, Force and Shape." *New Phytologist* 183: 13–26.
- Wan, S., Y. Luo, and L. Wallace. 2002. "Changes in Microclimate Induced by Experimental Warming and Clipping in Tallgrass Prairie." *Global Change Biology* 8: 754–68.
- Wang, Y., S.J. Burgess, E.M. de Becker, and S.P. Long. 2020. "Photosynthesis in the Fleeting Shadows: An Overlooked Opportunity for Increasing Crop Productivity?" *The Plant Journal* 101: 874–84.
- Way, D.A., G.G. Katul, S. Manzoni, and G. Vico. 2014. "Increasing Water Use Efficiency along the C3 to C4 Evolutionary Pathway: A Stomatal Optimization Perspective." *Journal of Experimental Botany* 65: 3683–93.
- Wentz, F.J., L. Ricciardulli, K. Hilburn, and C. Mears. 2007. "How Much More Rain Will Global Warming Bring?" *Science* 317: 233–5.
- Wickham, H. 2009. *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag.
- Wilcox, K.R., J.C. Fischer, J.M. Muscha, M.K. Petersen, and A.K. Knapp. 2015. "Contrasting Above- and Belowground Sensitivity of Three Great Plains Grasslands to Altered Rainfall Regimes." *Global Change Biology* 21: 335–44.
- Wilcox, K.R., Z. Shi, L.A. Gherardi, N.P. Lemoine, S.E. Koerner, D. L. Hoover, E. Bork, et al. 2017. "Asymmetric Responses of Primary Productivity to Precipitation Extremes: A Synthesis of Grassland Precipitation Manipulation Experiments." *Global Change Biology* 23: 4376–85.
- Wu, Z., P. Dijkstra, G.W. Koch, J. Peñuelas, and B.A. Hungate. 2011. "Responses of Terrestrial Ecosystems to Temperature and Precipitation Change: A Meta-Analysis of Experimental Manipulation." *Global Change Biology* 17: 927–42.
- Xu, X., S. Niu, R. Sherry, X. Zhou, J. Zhou, and Y. Luo. 2012. "Interannual Variability in Responses of Belowground NPP and NPP Partitioning to Long-Term Warming and Clipping in a Tallgrass Prairie." *Global Change Biology* 18: 1648–56.
- Xu, X., R.A. Sherry, S. Niu, D. Li, and Y. Luo. 2013. "Net Primary Productivity and Rain-Use Efficiency as Affected by Warming, Altered Precipitation, and Clipping in a Mixed-Grass Prairie." *Global Change Biology* 19: 2753–64.
- Xu, X., Y. Luo, Z. Shi, X. Zhou, and D. Li. 2014. "Consistent Proportional Increments in Responses of Belowground Net Primary Productivity to Long-Term Warming and Clipping at Various Soil Depths in a Tallgrass Prairie." *Oecologia* 174: 1045–54.
- Xue, X., Y. Luo, X. Zhou, R. Sherry, and X. Jia. 2011. "Climate Warming Increases Soil Erosion, Carbon and Nitrogen Loss with Biofuel Feedstock Harvest in Tallgrass Prairie." *GCB Bioenergy* 3: 198–207.
- Yachi, S., and M. Loreau. 1999. "Biodiversity and Ecosystem Productivity in a Fluctuating Environment: The Insurance Hypothesis." *Proceedings of the National Academy of Sciences of the United States of America* 96: 1463–8.
- Yamori, W., K. Hikosaka, and D.A. Way. 2014. "Temperature Response of Photosynthesis in C3, C4, and CAM Plants: Temperature Acclimation and Temperature Adaptation." *Photosynthesis Research* 119: 101–17.
- Ye, J.-S., J.-Y. Pei, and C. Fang. 2017. "Under which Climate and Soil Conditions the Plant Productivity–Precipitation Relationship Is Linear or Nonlinear?" *Science of the Total Environment* 616–617: 1174–80.
- Zhou, X., R.A. Sherry, Y. An, L.L. Wallace, and Y. Luo. 2006. "Main and Interactive Effects of Warming, Clipping, and Doubled Precipitation on Soil CO₂ Efflux in a Grassland Ecosystem." *Global Biogeochemical Cycles* 20. <https://doi.org/10.1029/2005GB002526>

Zhu, K., N.R. Chiariello, T. Tobeck, T. Fukami, and C.B. Field. 2016. "Nonlinear, Interacting Responses to Climate Limit Grassland Production under Global Change." *Proceedings of the National Academy of Sciences of the United States of America* 113: 10589–94.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Jung, Chang Gyo, Xia Xu, Zheng Shi, Shuli Niu, Jianyang Xia, Rebecca Sherry, Lifen Jiang, Kai Zhu, Enqing Hou, and Yiqi Luo. 2022. "Warmer and Wetter Climate Promotes Net Primary Production in C₄ Grassland with Additional Enhancement by Hay Harvesting." *Ecosphere* 13(1): e3899. <https://doi.org/10.1002/ecs2.3899>