



On the Sharp Time Decay Rates for the 2D Generalized Quasi-geostrophic Equation and the Boussinesq System

Atanas G. Stefanov¹ · Fazel Hadadifard¹

Received: 27 June 2018 / Accepted: 3 April 2019 / Published online: 10 April 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

We compute the sharp time decay rates of the solutions of the IVP for quasi-geostrophic equation and the Boussinesq model, subject to fractional dissipation. Moreover, we explicitly identify the asymptotic profiles, the kernel of the α stable processes, which are analogues of the Oseen vortices.

Keywords Time decay · Quasi-geostrophic equation · Boussinesq system

Mathematics Subject Classification Primary 35Q35 · 35B40; 76D03 Secondary 76B03 · 76D07

1 Introduction

The initial value problem for the Navier–Stokes equation,

$$\begin{cases} u_t + u \cdot \nabla u - \Delta u = \nabla p, & x \in \mathbf{R}^n, t > 0 \\ u(0, x) := u_0(x), \nabla \cdot u = 0 \end{cases} \quad (1.1)$$

where u is the fluid velocity and p is the pressure, is ubiquitous and much studied model in the modern PDE theory. Basic issues like global well-posedness remain elusively

Communicated by Paul Newton.

Hadadifard is supported in part by a graduate fellowship under NSF-DMS, # 1614734. Stefanov is partially supported by NSF-DMS under # 1614734.

✉ Atanas G. Stefanov
stefanov@ku.edu

Fazel Hadadifard
f.hadadi@ku.edu

¹ Department of Mathematics, University of Kansas, 1460 Jayhawk Boulevard, Lawrence, KS 66045–7523, USA

unresolved in spatial dimensions $n \geq 3$. In the case of two spatial dimensions though, the problem is globally well-posed. This is mostly due to the following representation, which eliminates the pressure term and leads to equivalent *vorticity formulation*,

$$\begin{cases} \omega_t + u \cdot \nabla \omega - \Delta \omega = 0, & x \in \mathbf{R}^2, t > 0 \\ \omega(0, x) := \omega_0(x), \end{cases} \quad (1.2)$$

where the vorticity ω , a scalar¹ quantity, is given by $\omega = \partial_1 u_2 - \partial_2 u_1$. We denote $\nabla^\perp = \begin{pmatrix} -\partial_2 \\ \partial_1 \end{pmatrix}$ for future reference, so that $\omega = \nabla^\perp \cdot \mathbf{u}$. Many generalizations of this model have been considered, in particular to respond to modeling situations where the actual physical dissipation is different than the one provided by the Laplacian, in particular in large-scale atmospheric models and large-scale ocean modeling, see Abidi and Hmidi (2007), Chae (2006) and Jiu et al. (2014). In particular, we consider the following “umbrella” model

$$\begin{cases} \partial_t z + u \cdot \nabla z + |\nabla|^\alpha z = 0, & x \in \mathbf{R}^2, t > 0, \\ u = (\nabla^\perp)^{-\beta} z, \nabla \cdot u = 0. \end{cases} \quad (1.3)$$

where $\alpha > 1$ and $\beta \geq 0$, $(\nabla^\perp)^{-\beta} = \nabla^\perp m_{-\beta-1}(|\nabla|) = \mathbf{m}_{-\beta}(\xi)$, where \mathbf{m}_a is a symbol of order a , see Sect. 2.1 for precise definition.² These types of equations frequently arise in fluid dynamics, and as such, they have been widely studied, especially so in the last 20 years. We refer the reader to the works Abidi and Hmidi (2007), Ben-Artzi (1994), Carpio (1994), Chae (2006), Cordoba and Cordoba (2004), Giga and Kambe (1988), Jiu et al. (2014), Niche and Schonbek (2007), Wu (2012), Yang et al. (2014) and references therein.

A few examples, that we would like to emphasize as model cases, are as follows. The 2D Fractional Navier–Stokes equation arises, if we take $z = \omega$ and $\beta = 1$,

$$\omega_t + u \cdot \nabla \omega + |\nabla|^\alpha \omega = 0. \quad (1.4)$$

If we let $z = \theta$ be the temperature of a flow, $\alpha > 1$ and $\beta = 0$, the resulting equation is the so-called active scalar equation,

$$\theta_t + u \cdot \nabla \theta + |\nabla|^\alpha \theta = 0, \quad (1.5)$$

where $u_1 = -R_2 \theta$, $u_2 = R_1 \theta$, and R_j , $j = 1, 2$ are the Riesz transforms, given by the symbols $m_j(\xi) = i \frac{\xi_j}{|\xi|}$.

¹ Note that in the 3D case, where the vorticity formulation is still available, it is a vector quantity and it does not seem to lead to any improvement over the original formulation

² Note that it is a requirement that $m_{-\beta-1}(|\nabla|)$ is a *radial symbol* of order $-\beta - 1$.

The Boussinesq system, with general dissipations, reads

$$\begin{cases} \partial_t u + u \cdot \nabla u + |\nabla|^{\sigma_1} u = -\nabla p + \theta \mathbf{e}_2, & x \in \mathbb{R}^2, t > 0, \\ \partial_t \theta + u \cdot \nabla \theta + |\nabla|^{\sigma_2} \theta = 0, & x \in \mathbb{R}^2, t > 0, \\ \nabla \cdot u = 0. \end{cases} \quad (1.6)$$

where u is the velocity of the fluid, θ is its temperature, p is the pressure and $\sigma_1, \sigma_2 > 0$ are the dissipation rates for the velocity and the temperature respectively, see Abidi and Hmidi (2007), Brandoles and Schonbek (2012), Chae (2006), Hadadifard and Stefanov (2017), Jiu et al. (2014), Wu (2012) and Yang et al. (2014) for background and various well-posedness results.

We consider the equivalent vorticity formulation, with the usual scalar vorticity variable is given by $\omega = \partial_1 u_2 - \partial_2 u_1$. For the purposes of this work, we will only consider the diagonal case $\sigma_1 = \sigma_2 = \alpha$. That is in vorticity formulation, the system consists of the following coupled equations

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega + |\nabla|^\alpha \omega = \partial_1 \theta, & x \in \mathbb{R}^2, t > 0, \\ \partial_t \theta + u \cdot \nabla \theta + |\nabla|^\alpha \theta = 0, & x \in \mathbb{R}^2, t > 0, \\ u = (\nabla^\perp)^{-1} \omega, \quad \nabla \cdot u = 0. \end{cases} \quad (1.7)$$

1.1 Previous Results

As we have mentioned earlier, a lot of work has been done on the question of well-posedness, regularity of the solutions to these systems. We do not even attempt to overview the results, as this is only tangentially relevant for the current work, but the previously mentioned references contain lots of information about these issues. As the purpose of this paper is to study the long-time behavior of the said models, we discuss some recent works on the topic. Most of the research has been done in the important (and classical) Navier–Stokes case in two- and three-dimensional cases. As the global regularity for this model remains a challenging open problem in 3D, some authors restricted themselves to weak solutions³ or they considered eventual⁴ behavior of strong solutions. In this regard, we would like to reference the following works, Carpio (1994), Gallay and Wayne (2002a,b), Giga and Kambe (1988), Miyakawa and Schonbek (2001), Schonbek (1985, 1991, 1992), Schonbek and Schonbek (2000) and Schonbek and Wiegner (1996).

In Schonbek (1991), the author has exhibited lower time decay bounds for the solutions, which match the upper bounds and are therefore sharp. The approach in Gallay and Wayne (2002a,b), for the same question, uses the method of the so-called scaled variables. This was pioneered in Giga and Kambe (1988) and Carpio (1994), although the idea really took off after the work (Gallay and Wayne 2002a). It showed not only the optimal decay rates for the Navier–Stokes equation (this was actually previously established in Brandoles and Schonbek 2012), but it provided an explicit asymptotic

³ Which may be non-unique.

⁴ That is, past eventual singularity formation.

expansion of the solution, which explains the specific conditions on the initial data in Brandoles and Schonbek (2012), under which there are better decay rates. Recently, Goh and Wayne (2018) have considered the Boussinesq model, with rapid rotation in 3D. They have shown, using the method of scaled variables, convergence to the Oseen vortex and associated leading order asymptotics.

In this paper, we follow this idea, to provide an explicit asymptotic expansion for the two models under consideration—the generalized quasi-geostrophic equation (1.3) and the Boussinesq system (in vorticity formulation) (1.7). Note that we work exclusively in two spatial dimensions. There are several reasons for this—2D is the natural playground for (1.3), while the IVP for the Boussinesq system, the three (and higher)-dimensional case, faces the same difficulties as the Navier–Stokes problem, namely absence of a global regularity theory. Moreover, we explore relatively low levels of dissipation, which in some sense brings the global regularity theory to its limits, and we are still able to analyze the asymptotic behavior. Another interesting feature that we deal with is the fractional dissipation. These have been studied in the recent literature, but there are certain technical (and conceptual!) difficulties associated with them, that we deal with by applying advanced Fourier analysis methods.

1.2 Main Results

The main goal of this work is to establish the sharp time decay rates of (various norms of) the solutions to (1.3) and (1.7). Our results actually provide explicit asymptotic profiles, of which the precise asymptotic rates are a mere corollary.

Since it is clear that the equation for θ in (1.7) is basically⁵ (1.3), it is essential that we start with (1.3). This is the content of our first result, but in order to state it, we shall need to introduce a function $G: \hat{G}(p) = e^{-|p|^\alpha}$, see Sect. 2.4 for proper definitions and properties. This is a variant of the function $e^{-\frac{|x|^2}{2}}$, or the Oseen vortex in the case $\alpha = 2$.

In order to state our results, we shall need to introduce weighted spaces. More specifically, for any $m \geq 0$ we define the Hilbert space $L^2(m)$ as follow

$$L^2(m) = \left\{ f \in L^2 : \|f\|_{L^2(m)} = \left(\int_{\mathbf{R}^2} (1 + |x|^2)^m |f(x)|^2 dx \right)^{\frac{1}{2}} < \infty \right\} \quad (1.8)$$

One can show by means of Hölder's, the embedding $L^2(2) \hookrightarrow L^p(\mathbf{R}^2)$, whenever $1 \leq p \leq 2$. We are now ready for the statement of the main result.

Theorem 1 (Global decay estimates for SQG) *Let $1 < \alpha < 2$, and $\alpha + \beta \leq 3$. Then, assuming that the initial data z_0 is in $L^2(2) \cap L^\infty$, the Cauchy problem (1.3) has a unique, global solution in $L^2(2) \cap L^\infty$. Moreover, for all $\epsilon > 0$, there is a constant*

⁵ Albeit the relation of u with θ is not a direct one, but through the vorticity ω .

$C = C_{\alpha, \beta, \epsilon}$ and for all $p \in [1, 2]$ and $t \geq 0$,

$$\|z(t, \cdot) - \frac{\int_{\mathbf{R}^2} z_0(x) dx}{(1+t)^{\frac{2}{\alpha}}} G\left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}}\right)\|_{L^p} \leq \frac{C_{\alpha, \beta, \epsilon} \|z_0\|_{L^2(2) \cap L^\infty}}{(1+t)^{\frac{3}{\alpha} - \frac{2}{\alpha p} - \epsilon}}. \quad (1.9)$$

Moreover, if $\beta > 1$, we have that (1.9) holds for the full range of indices $1 \leq p < \infty$.

In particular, for generic initial data, i.e., $\int_{\mathbf{R}^2} z_0(x) dx \neq 0$, we have from (1.9), after selecting $\epsilon < \frac{1}{\alpha}$ and for large t , that

$$\|z(t, \cdot)\|_{L^p} \sim (1+t)^{-\frac{2(p-1)}{\alpha p}}, \quad 1 \leq p \leq 2. \quad (1.10)$$

which extends to all $1 \leq p < \infty$, provided $\beta > 1$.

Remark

- Perhaps, we need to explain a bit the conclusion (1.10). The right-hand side of (1.9) is $\frac{C_\epsilon}{(1+t)^{\frac{3}{\alpha} - \frac{2}{\alpha p} - \epsilon}}$, that is, for each $\epsilon > 0$, there is C_ϵ , so that this estimate holds. That is, one gets better statement for large t (which is the goal here), every time ϵ is smaller, but pays with a larger and larger constant, basically our construction here says $\lim_{\epsilon \rightarrow 0^+} C_\epsilon = \infty$. The explicit expression inside the norm in (1.9) has decay rate in L^p , which is $\frac{1}{(1+t)^{\frac{2}{\alpha} - \frac{2}{\alpha p}}}$. Clearly, for small enough $\epsilon > 0$ (namely $\epsilon < \frac{1}{\alpha}$), one has that

$$\frac{C_\epsilon}{(1+t)^{\frac{3}{\alpha} - \frac{2}{\alpha p} - \epsilon}} << \frac{1}{(1+t)^{\frac{2}{\alpha} - \frac{2}{\alpha p}}}.$$

for large t . Thus, fixing $\epsilon < \frac{1}{\alpha}$, say $\epsilon = \frac{1}{2\alpha}$, and applying (1.9) yields

$$\frac{C}{(1+t)^{\frac{2}{\alpha} - \frac{2}{\alpha p}}} - \frac{C_\epsilon}{(1+t)^{\frac{3}{\alpha} - \frac{2}{\alpha p} - \epsilon}} \leq \|z(t)\|_{L^p} \leq \frac{C}{(1+t)^{\frac{2}{\alpha} - \frac{2}{\alpha p}}} + \frac{C_\epsilon}{(1+t)^{\frac{3}{\alpha} - \frac{2}{\alpha p} - \epsilon}}.$$

From this, one gets (1.10).

- Our results extend those in Constantin and Wu (1999), as they provide an upper bound for the time decay, for weak solutions of the SQG.
- In Gallay and Wayne (2002a, b), the authors go one step further in deriving explicitly the next order asymptotic profiles. The analysis required for this step is performed in higher order weighted L^2 space. This cannot be done in this framework, since the function G does not belong to the next order weighted space, namely $L^2(3)$, see Proposition 3. This is in sharp contrast with the case $\alpha = 2$, considered in Gallay and Wayne (2002a, b), where the function is in Schwartz class.
- Related to the previous point, we need to address a problem, where the function G and the heat kernel of the semigroup $e^{\tau \mathcal{L}}$ have limited decay at infinity. Thus, any attempt to use the dynamical system approach in Gallay and Wayne (2002a)

to construct stable manifolds faces serious obstacles. We take a different approach to the problem in that we use *a priori* estimates and estimates on the evolution operator to establish the asymptotic decomposition.

Our next result concerns (1.7).

Theorem 2 (Global decay estimates for Boussinesq) *Let $\alpha \in (1, \frac{3}{2})$. Consider the Cauchy problem for (1.7), with initial data $w_0, \theta_0 \in Y := L^2(2) \cap L^\infty \cap H^1(\mathbf{R}^2)$. Then, the Cauchy problem (1.7) is globally well-posed in Y - that is for every $t > 0$, the solution $(w(t), \theta(t)) \in Y \times Y$.*

Moreover, for every $\delta > 0$, there exists $C = C(\alpha, \delta, \|w_0\|_Y, \|\theta_0\|_Y)$, so that for all $p \in [1, 2]$ and for all $t > 0$,

$$\begin{aligned} \|w(t, \cdot) - \frac{\gamma_2(0)}{(1+t)^{\frac{3}{\alpha}-1}} \partial_1 G \left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}} \right) \\ - \frac{\gamma_1(0)}{(1+t)^{\frac{2}{\alpha}}} G \left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}} \right) \|_{L^p} \leq \frac{C_{\alpha, \delta} \|(w_0, \theta_0)\|_Y}{(1+t)^{\frac{6}{\alpha}-3-\frac{2}{\alpha p}-\delta}}, \end{aligned} \quad (1.11)$$

$$\begin{aligned} \|\theta(t, \cdot) - \frac{\gamma_2(0)}{(1+t)^{\frac{2}{\alpha}}} G \left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}} \right) \|_{L^p} \\ \leq \frac{C_{\alpha, \delta} \|(w_0, \theta_0)\|_Y}{(1+t)^{\frac{5}{\alpha}-2-\frac{2}{\alpha p}-\delta}}, \end{aligned} \quad (1.12)$$

where $\gamma_1(0) = \int_{\mathbf{R}^2} w_0(x) dx$, $\gamma_2(0) = \int_{\mathbf{R}^2} \theta_0(x) dx$. In particular, if $\gamma_2(0) \neq 0$, we have

$$\|w(t, \cdot)\|_{L^p} \sim \frac{1}{(1+t)^{\frac{3}{\alpha}-1-\frac{2}{\alpha p}}}, \|\theta(t, \cdot)\|_{L^p} \sim \frac{1}{(1+t)^{\frac{2}{\alpha}-\frac{2}{\alpha p}}}, \quad (1.13)$$

Remark

- Again, we shall need to explain (1.13). We only discuss the w , the argument for θ is similar. We have the following decay for the two terms inside of (1.11),

$$\begin{aligned} \frac{1}{(1+t)^{\frac{3}{\alpha}-1}} \left\| \partial_1 G \left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}} \right) \right\|_{L^p} &\sim \frac{1}{(1+t)^{\frac{3}{\alpha}-\frac{2}{\alpha p}-1}}, \\ \frac{1}{(1+t)^{\frac{2}{\alpha}}} \left\| G \left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}} \right) \right\|_{L^p} &\sim \frac{1}{(1+t)^{\frac{2}{\alpha}-\frac{2}{\alpha p}}}. \end{aligned}$$

For $\alpha \in (1, \frac{3}{2})$, the slowest rate is $\frac{1}{(1+t)^{\frac{3}{\alpha}-\frac{2}{\alpha p}-1}}$, which is clearly slower than the decay rate on the right-hand side of (1.11). More precisely, for small δ : $\delta = \frac{3}{2\alpha} - 1$, we have that for large t

$$\frac{1}{(1+t)^{\frac{6}{\alpha}-3-\frac{2}{\alpha p}-\delta}} << \frac{1}{(1+t)^{\frac{3}{\alpha}-\frac{2}{\alpha p}-1}}$$

whence (1.13) follows.

- Related to the previous point, for $\alpha \in (\frac{4}{3}, \frac{3}{2})$, the correction term $\frac{\gamma_1(0)}{(1+t)^{\frac{2}{\alpha}}} G\left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}}\right)$ is faster decaying than the error term and we can restate the result as follows

$$\left\| w(t, \cdot) - \frac{\gamma_2(0)}{(1+t)^{\frac{3}{\alpha}-1}} \partial_1 G\left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}}\right) \right\|_{L^p} \leq \frac{C_{\alpha, \delta} \|(w_0, \theta_0)\|_Y}{(1+t)^{\frac{6}{\alpha}-3-\frac{2}{\alpha p}-\delta}},$$

- As in Theorem 1, the results can be extended to provide asymptotic expansions for w, θ in the norms L^p , $p \in (2, \infty)$, with the exact same statement.

The paper is organized as follows. In Sect. 2.1, we introduce some basic Sobolev spaces, weighted L^2 spaces and some relevant estimates that will be useful in the sequel. In Sect. 3.1, we study the operator \mathcal{L} —we establish the basic structure of its spectrum, as well as an explicit form of the semigroup $e^{\tau \mathcal{L}}$. The semigroup is shown to act boundedly on certain weighted L^2 spaces. This is helpful for the study of the nonlinear evolutions problem, but it also helps us identify the spectrum, through the Hille-Yosida theorem, see Sect. 3.4. In Sect. 4, we develop the local and global well-posedness theory for the generalized quasi-geostrophic equation, both in the original variables and then in the scaled variables. This is done via standard energy estimates methods. Even at this level, the optimal decay estimates start to emerge, in the scaled variables context.⁶ Our asymptotic results for the quasi-geostrophic model are in Sect. 6. In it, we use the *a priori* information from Sect. 4, together with new estimates for the Duhamel's operator to derive the precise asymptotic profiles for the solutions. For the Boussinesq system, we provide the necessary local and global well-posedness theory in Sect. 5. Some of these results are basic and could have been recovered from earlier publications. Others provide new *a priori* estimates for the scaled variables system, which are used in Sect. 7. In Sect. 7, we provide the proof of our main result about the precise asymptotic profiles for the Boussinesq evolution.

2 Preliminaries

2.1 Fourier Transform, Function Spaces and Multipliers

The Fourier transform and its inverse are taken in the form

$$\hat{f}(p) = \int_{\mathbf{R}^n} f(x) e^{-ix \cdot p} dx, \quad f(x) = (2\pi)^{-n} \int_{\mathbf{R}^n} \hat{f}(p) e^{ix \cdot p} dp$$

⁶ But at this point, we cannot yet conclude the optimality of these estimates, as we are missing an estimate from below.

Consequently, since $\widehat{-\Delta f}(p) = |p|^2 \widehat{f}(p)$, we define the operators $|\nabla|^a := (-\Delta)^{a/2}$, $a > 0$, via its action on the Fourier side $\widehat{|\nabla|^a f}(p) = |p|^a \widehat{f}(p)$. More generally, the operators $F(|\nabla|)$, for reasonable functions F , are acting as multipliers by $F(|p|)$. We also make use of the following notation—we say that \mathbf{m} is a symbol of order a , $a \in \mathbf{R}$, if it is a smooth, possibly vector-valued function on $\mathbf{R}^n \setminus \{0\}$, satisfying for all multi-indices $\alpha \in \mathbf{N}^n$,

$$|\partial^\alpha \mathbf{m}(\xi)| \leq C_\alpha |\xi|^{a-|\alpha|}.$$

It is actually enough to assume this inequality for a finite set of indices, say $|\alpha| \leq n$. The prototype will be something of the form $m(\xi) = |\xi|^a$, but note that a will be often negative in our applications. We schematically denote a symbol of order a by \mathbf{m}_a .

The L^p spaces are defined by the norm $\|f\|_{L^p} = \left(\int |f(x)|^p dx \right)^{\frac{1}{p}}$, while the weak L^p spaces are

$$L^{p,\infty} = \left\{ f : \|f\|_{L^{p,\infty}} = \sup_{\lambda > 0} \left\{ \lambda |\{x : |f(x)| > \lambda\}|^{\frac{1}{p}} \right\} < \infty \right\}.$$

In this context, recall the Hausdorff–Young inequality which reads as follows: For $p, q, r \in (1, \infty)$ and $1 + \frac{1}{p} = \frac{1}{q} + \frac{1}{r}$

$$\|f * g\|_{L^p} \leq C_{p,q,r} \|f\|_{L^{q,\infty}} \|g\|_{L^r}.$$

For an integer k and $p \in (1, \infty)$, the Sobolev spaces are the closure of the Schwartz functions in the norm $\|f\|_{W^{k,p}} = \|f\|_{L^p} + \sum_{|\alpha| \leq k} \|\partial^\alpha f\|_{L^p}$, while for a non-integer s one takes

$$\|f\|_{W^{s,p}} = \|(1 - \Delta)^{s/2} f\|_{L^p} \sim \|f\|_{L^p} + \||\nabla|^s f\|_{L^p}.$$

The Sobolev embedding theorem states $\|f\|_{L^p(\mathbf{R}^n)} \leq C \||\nabla|^s f\|_{L^q(\mathbf{R}^n)}$, where $1 < q < p < \infty$ and $n(\frac{1}{q} - \frac{1}{p}) = s$, with the usual modification for $p = \infty$, namely $\|f\|_{L^\infty(\mathbf{R}^n)} \leq C_s \|f\|_{W^{s,q}(\mathbf{R}^n)}$, $s > \frac{n}{q}$. In particular, an estimate that will be useful for us, is

$$\|(\nabla^\perp)^{-\beta} f\|_{L^p} \leq C \|f\|_{L^q}, \quad 1 < q < p < \infty, \beta = n \left(\frac{1}{q} - \frac{1}{p} \right) \quad (2.1)$$

This follows from the Mikhlin's criteria for L^p , $1 < p < \infty$ boundedness. Sometimes, we use the following replacement of (2.1), when $p = \infty$ and $\beta < n$,

$$\|(\nabla^\perp)^{-\beta} f\|_{L^\infty} \leq C_\epsilon \left(\|f\|_{L^{\frac{n}{\beta}+\epsilon}} + \|f\|_{L^{\frac{n}{\beta}-\epsilon}} \right). \quad (2.2)$$

We provide a proof for this inequality in “Appendix A.” Note that these estimates hold in a more general setting, when $(\nabla^\perp)^{-\beta}$ is replaced by an arbitrary symbol of order $-\beta$, that is

$$\|m_{-\beta}(\nabla)f\|_{L^\infty} \leq C_\epsilon \left(\|f\|_{L^{\frac{n}{\beta}+\epsilon}} + \|f\|_{L^{\frac{n}{\beta}-\epsilon}} \right). \quad (2.3)$$

Another useful ingredient will be the Gagliardo–Nirenberg interpolation inequality,

$$\||\nabla|^s f\|_{L^p} \leq \||\nabla|^{s_1} f\|_{L^q}^\theta \||\nabla|^{s_2} f\|_{L^r}^{1-\theta},$$

where $s = \theta s_1 + (1 - \theta) s_2$ and $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$.

2.2 The Scaled Variables

We now introduce the scaled variables, for the models under consideration. Basically, the method consists of introducing a new time variable $\tau : e^\tau \sim t$ and the corresponding variables in x are rescaled to accommodate this scaling, by keeping the linear part of the equation autonomous. In this way, an algebraic decay in t will manifest itself as an exponential decay in τ . As is well-known, algebraic decays in time (especially non-integrable ones) are notoriously hard to propagate along nonlinear evolution equations, while any (however small) exponential decay, due to its integrability, is more amenable to this type of analysis. Here are the details.

2.2.1 The Scaled Variables: The SQG Equation

Consider the Eq. (1.3) and use the scaled variables to rewrite the variables in terms of

$$\xi = \frac{x}{(1+t)^{\frac{1}{\alpha}}}, \quad \tau = \ln(1+t). \quad (2.4)$$

We define new functions $Z(\xi, \tau)$ and $U(\xi, \tau)$ correspond to $z(x, t)$ and $u(x, t)$ as follows:

$$z(x, t) = \frac{1}{(1+t)^{1+\frac{\beta-1}{\alpha}}} Z \left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \ln(1+t) \right), \quad (2.5)$$

$$u(x, t) = \frac{1}{(1+t)^{1-\frac{1}{\alpha}}} U \left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \ln(1+t) \right). \quad (2.6)$$

The choices of the parameters are clearly dictated by the stricture of the corresponding equation—the goal is to ensure an autonomous PDE in the new variables. Indeed, a straightforward calculation shows

$$z_t = \frac{Z_\tau}{(1+t)^{2+\frac{\beta-1}{\alpha}}} - \frac{1}{\alpha} \frac{1}{(1+t)^{2+\frac{\beta-1}{\alpha}}} \frac{x}{(1+t)^{\frac{1}{\alpha}}} \cdot \nabla_\xi Z - \frac{1+\frac{\beta-1}{\alpha}}{(1+t)^{2+\frac{\beta-1}{\alpha}}} Z,$$

$$|\nabla|^\alpha z = \frac{1}{(1+t)^{2+\frac{\beta-1}{\alpha}}} |\nabla|^\alpha Z,$$

$$u \cdot \nabla Z = \frac{1}{(1+t)^{2+\frac{\beta-1}{\alpha}}} U \cdot \nabla_{\xi} Z.$$

Hence, $Z(\xi, \tau)$ satisfies the equation

$$Z_{\tau} = \mathcal{L}Z - U \cdot \nabla_{\xi} Z \quad (2.7)$$

where

$$\mathcal{L}Z = -|\nabla|^{\alpha}Z + \frac{1}{\alpha}\xi \cdot \nabla_{\xi}Z + \left(1 + \frac{\beta-1}{\alpha}\right)Z. \quad (2.8)$$

Note that the relation $u = (|\nabla|^{\perp})^{-\beta}z$ transforms into $U = (|\nabla|^{\perp})^{-\beta}Z$. In addition, the property $\nabla \cdot u = 0$ is retained, i.e., $\nabla \cdot U = 0$.

Next, we introduce the scaled variables for the Boussinesq system.

2.2.2 The Scaled Variables: The Boussinesq System

Similar to the SQG case, we use the scaled variables

$$\xi = \frac{x}{(1+t)^{\frac{1}{\alpha}}}, \quad \tau = \ln(1+t).$$

We define new functions $W(\xi, \tau)$, $U(\xi, \tau)$ and $\Theta(\xi, \tau)$, corresponding to $\omega(x, t)$, $u(x, t)$ and $\theta(x, t)$ as follows

$$\begin{aligned} \omega(x, t) &= \frac{1}{(1+t)} W\left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \ln(1+t)\right) \\ u(x, t) &= \frac{1}{(1+t)^{1-\frac{1}{\alpha}}} U\left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \ln(1+t)\right) \\ \theta(x, t) &= \frac{1}{(1+t)^{2-\frac{1}{\alpha}}} \Theta\left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \ln(1+t)\right) \end{aligned}$$

Then, we calculate

$$\begin{aligned} \omega_t &= \frac{W_{\tau}}{(1+t)^2} - \frac{1}{\alpha} \frac{1}{(1+t)^2} \frac{x}{(1+t)^{\frac{1}{\alpha}}} \cdot \nabla_{\xi} W - \frac{1}{(1+t)^2} W, \\ |\nabla|^{\alpha} \omega &= \frac{1}{(1+t)^2} \cdot |\nabla|^{\alpha} W, \quad u \cdot \nabla \omega = \frac{1}{(1+t)^2} U \cdot \nabla W, \quad \partial_1 \theta = \frac{1}{(1+t)^2} \partial_1 \Theta. \end{aligned}$$

For the θ equation similar computation shows that

$$\theta_t = \frac{\Theta_{\tau}}{(1+t)^{3-\frac{1}{\alpha}}} - \frac{1}{\alpha} \frac{1}{(1+t)^{3-\frac{1}{\alpha}}} \frac{x}{(1+t)^{\frac{1}{\alpha}}} \cdot \nabla_{\xi} \Theta - \frac{2 - \frac{1}{\alpha}}{(1+t)^{3-\frac{1}{\alpha}}} \Theta,$$

$$|\nabla|^\alpha \theta = \frac{1}{(1+t)^{3-\frac{1}{\alpha}}} |\nabla|^\alpha \Theta, \quad u \cdot \nabla \theta = \frac{1}{(1+t)^{3-\frac{1}{\alpha}}} U \cdot \nabla \Theta.$$

Therefore $W(\xi, \tau)$ and $\Theta(\xi, \tau)$ satisfy (with the \mathcal{L} defined above in (2.8), but with $\beta = 1$)

$$\begin{cases} W_\tau = \mathcal{L}W - U \cdot \nabla_\xi W + \partial_1 \Theta \\ \Theta_\tau = (\mathcal{L} + 1 - \frac{1}{\alpha})\Theta - (U \cdot \nabla_\xi \Theta) \end{cases} \quad (2.9)$$

Clearly, the relations $\nabla \cdot u = 0$ and $u = (|\nabla|^\perp)^{-1} \omega$ continue to hold for the capital letter variables as well, that is $\nabla \cdot U = 0$ and $U = (|\nabla|^\perp)^{-1} W$. In addition to the above equations, we can define $p(x, t) = \frac{1}{(1+t)^{2-\frac{2}{\alpha}}} P\left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}, \log(1+t)\right)$ and find the following equation for $U(\xi, \tau)$,

$$U_\tau = (\mathcal{L} - \frac{1}{\alpha})U - (U \cdot \nabla_\xi U) - \nabla P + \Theta \cdot e_2 \quad (2.10)$$

2.3 The Fractional Laplacian

First, we record the following kernel representation formula for negative powers of Laplacian. This is nothing, but a fractional integral—for $a \in (0, 2)$,

$$|\nabla|^{-a} f(x) = c_a \int_{\mathbf{R}^2} \frac{f(y)}{|x-y|^{2-a}} dy. \quad (2.11)$$

For positive powers, we have a similar formula—for $a \in (0, 2)$,

$$|\nabla|^a f(x) = C_a p.v. \int_{\mathbf{R}^2} \frac{f(x) - f(y)}{|x-y|^{2+a}} dy.$$

see Cordoba and Cordoba (2004, Proposition 2.1). Next, we have the following result (Cordoba and Cordoba 2004), see also Theorem 2, (Chamorro and Lemarié-Rieusset 2012) for a more refined version.

Lemma 1 *For $p : 1 \leq p < \infty$, $a \in [0, 2]$ and $f \in W^{a,p}(\mathbf{R}^2)$,*

$$\int_{\mathbf{R}^2} |f(x)|^{p-2} f(x) [|\nabla|^a f](x) dx \geq 0. \quad (2.12)$$

If in addition, $p = 2^n$, $n = 1, 2, \dots$, there is the stronger coercivity estimate

$$\int_{\mathbf{R}^2} |f(x)|^{p-2} f(x) [|\nabla|^a f](x) dx \geq \frac{1}{p} \||\nabla|^{\frac{a}{2}} [f^{\frac{p}{2}}]\|_{L^2(\mathbf{R}^2)}^2. \quad (2.13)$$

Finally, for $p \in [1, \infty)$, $a \in (0, 2)$,

$$\int_{\mathbf{R}^2} |f(x)|^{p-2} f(x) [|\nabla|^a f](x) dx \geq \frac{1}{p} \|f\|_{L^{\frac{2p}{2-a}}(\mathbf{R}^2)}^2. \quad (2.14)$$

2.4 The Function G

The function G defined by $\hat{G}(p) = e^{-|p|^\alpha}$, $p \in \mathbf{R}^2$ will be used frequently in the sequel. We list and prove some important properties.

Lemma 2 For any $p \in [2, \infty]$ and $\alpha \in (1, 2)$,

$$(1 + |\xi|^2) G(\xi), (1 + |\xi|^2) \nabla G(\xi) \in L_\xi^p \quad (2.15)$$

In particular, $G, \nabla G \in L^1(\mathbf{R}^2) \cap L^\infty(\mathbf{R}^2)$.

Note For $\alpha \in (1, 2)$, the function G does not belong to $L^2(3)$, due to the lack of smoothness of \hat{G} at zero (or what is equivalent to the lack of decay of G at ∞).

Proof For the L^2 estimate, $\|G\|_{L^2} = \|\hat{G}\|_{L^2} < \infty$. Since $|\cdot|^2 \hat{G}(\cdot)(p) = -\Delta_p \hat{G}(p)$, we obtain by Plancherel's,

$$\||\xi|^2 G(\xi)\|_{L^2} = \|\Delta_p \hat{G}(p)\|_{L^2} = \|\Delta_p e^{-|p|^\alpha}\|_{L^2} = \|(\partial_{\rho\rho} + \frac{1}{\rho} \partial_\rho)(e^{-\rho^\alpha})\|_{L^2(\rho d\rho)}.$$

where in the last identity, we have used that \hat{G} is a radial function. But, $(\partial_{\rho\rho} + \frac{1}{\rho} \partial_\rho)(e^{-\rho^\alpha}) = -\alpha(\alpha-1)\rho^{\alpha-2}e^{-\rho^\alpha} + \alpha^2\rho^{2(\alpha-1)}e^{-\rho^\alpha}$. Therefore, $\||\xi|^2 G(\xi)\|_{L^2}^2 \leq I_1 + I_2$, where $I_1 = \|\rho^{\alpha-2}e^{-\rho^\alpha}\|_{L^2(\rho d\rho)}^2$, $I_2 = \|\rho^{2(\alpha-1)}e^{-\rho^\alpha}\|_{L^2(\rho d\rho)}^2$. We have

$$I_1 \leq \int_0^1 \frac{1}{\rho^{2(2-\alpha)-1}} d\rho + \int_1^\infty \rho^{2(\alpha-2)+1} e^{-2\rho^\alpha} \rho d\rho.$$

Since $2(2-\alpha)-1 < 1$, the first term is bounded. The second term is also bounded by the exponential decay, whence I_1 is bounded. The second term, $I_2 = \|\rho^{2(\alpha-1)}e^{-\rho^\alpha}\|_{L^2(\rho d\rho)}^2$ is also bounded—no singularity at zero and exponential decay at ∞ . This proves the L^2 estimate.

For the L^∞ estimate, we can use the Hausdorff–Young's to bound $\|G\|_{L^\infty} \leq \|\hat{G}\|_{L^1} < \infty$. Similarly,

$$\begin{aligned} \||\xi|^2 G(\xi)\|_{L^\infty} &\leq \|\Delta_p \hat{G}(p)\|_{L^1} \leq \alpha(\alpha-1) \int_0^\infty \rho^{\alpha-2} e^{-\rho^\alpha} \rho d\rho + \alpha^2 \int_0^\infty \rho^{2(\alpha-1)} e^{-\rho^\alpha} \rho d\rho \\ &\leq \alpha(\alpha-1) \int_0^\infty \rho^{\alpha-1} e^{-\rho^\alpha} d\rho + \alpha^2 \int_0^\infty \rho^{2\alpha-1} e^{-\rho^\alpha} d\rho < \infty. \end{aligned}$$

Now the interpolation between L^2 and L^∞ yields $(1 + |\xi|^2) G(\xi) \in L_\xi^p$, $1 \leq p \leq \infty$.

Regarding the claims about ∇G , note that $\widehat{|\cdot|^2 \nabla G}(p) = -\Delta_p[p\hat{G}(p)]$, whence by Plancherel's $\||\xi|^2 \nabla G\|_{L^2} = \|\Delta_p[p e^{-|p|^\alpha}]\|_{L^2} < \infty$. The last conclusion follows easily from an identical argument as the one above, as the central issue was the singularity at zero for $\|\Delta_p e^{-|p|^\alpha}\|_{L^2}$. Now the situation is better as we multiply by p , which actually alleviates the singularity at zero. Similar argument applies to $\||\xi|^2 \nabla G\|_{L^\infty}$, so we omit the details. \square

The following Lemma will be used frequently in the next sections—it is an easy consequence of the Hausdorff–Young's inequality.

Lemma 3 *Let $\alpha > 0$, then for any $t > 0$ and $1 \leq p \leq \infty$,*

$$\|e^{-t|\nabla|^\alpha} f\|_{L^p} \leq C\|f\|_{L^p} \quad (2.16)$$

$$\|e^{-t|\nabla|^\alpha} \nabla f\|_{L^p} \leq C t^{-\frac{1}{\alpha}} \|f\|_{L^p} \quad (2.17)$$

Proof Clearly,

$$e^{-t|\nabla|^\alpha} f = \int G_t(x-y) f(y) dy$$

where $\widehat{G_t}(p) = \widehat{G}(t^{-\frac{1}{\alpha}} p)$. Then $\|e^{-t|\nabla|^\alpha} f\|_{L^p} \leq \|G_t\|_{L^1} \|f\|_{L^p} = C\|f\|_{L^p}$, where $C = \|G\|_{L^1(\mathbb{R}^2)}$.

$$\|e^{-t|\nabla|^\alpha} \nabla f\|_{L^p} = t^{-\frac{1}{\alpha}} \left\| \int \nabla G(t^{-\frac{1}{\alpha}}(\cdot-y)) f(y) dy \right\|_{L^p} \leq C t^{-\frac{1}{\alpha}} \|f\|_{L^p},$$

where $C = \|\nabla G\|_{L^1(\mathbb{R}^2)}$. \square

2.5 Kato–Ponce and Commutator Estimates

The classical by now product rule estimate, usually attributed to Kato–Ponce, can be stated as follows.

Lemma 4 *Let $a \in (0, 1)$ and $1 < p, q, r < \infty$, so that $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$. Then, there exists $C = C_{p,q,r,a}$*

$$\|\nabla|^a [fg]\|_{L^p} \leq C_{p,q,r,a} (\|\nabla|^a f\|_{L^q} \|g\|_{L^r} + \|\nabla|^a g\|_{L^q} \|f\|_{L^r})$$

We also make use of the following Lemma from Hadadifard and Stefanov (2017).

Lemma 5 *Let s_1, s_2 be two reals so that $0 \leq s_1$ and $0 \leq s_2 - s_1 \leq 1$. Let p, q, r be related via the Hölder's $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$, where $2 < q < \infty$, $1 < p, r < \infty$. Finally, let $\nabla \cdot V = 0$. Then for any $a \in [s_2 - s_1, 1]$*

$$\|\nabla|^{-s_1} [|\nabla|^{s_2}, V \cdot \nabla] \varphi\|_{L^p} \leq C \|\nabla|^a V\|_{L^q} \|\nabla|^{s_2 - s_1 + 1 - a} \varphi\|_{L^r} \quad (2.18)$$

In addition, we have the following end-point estimate. For $s_1 > 0, s_2 > 0, s_3 > 0$ and $s_1 < 1, s_3 < 1, s_2 < s_1 + s_3$, there is

$$\||\nabla|^{-s_1} [|\nabla|^{s_2}, |\nabla|^{-s_3} V \cdot \nabla] \varphi\|_{L^2} \leq C \|V\|_{L^\infty} \||\nabla|^{s_2-s_1+1-s_3} \varphi\|_{L^2}. \quad (2.19)$$

2.6 A Variant of the Gronwall's Inequality

We shall need a version of the Gronwall's inequality as follows.

Lemma 6 *Let $\sigma > 0, \mu > 0, \kappa > 0$ and $a \in [0, 1)$. Let A_1, A_2, A_3 be three positive constants so that a function $I : [0, \infty) \rightarrow \mathbf{R}_+$ satisfies $I(\tau) \leq A_1 e^{-\gamma\tau}$, for some real γ and*

$$I(\tau) \leq A_2 e^{-\mu\tau} + A_3 \int_0^\tau \frac{e^{-\sigma(\tau-s)}}{(\min(1, |\tau-s|)^a)} e^{-\kappa s} I(s) ds. \quad (2.20)$$

Then, there exists $C = C_{a,\sigma,\mu,\kappa,\gamma}$, so that

$$I(\tau) \leq C_{a,\sigma,\mu,\kappa,\gamma} (1 + |A_1| + |A_2| + |A_3|) e^{-\mu\tau}.$$

The proof of Lemma 6 is rather elementary, but we provide it for completeness in the “Appendix B.”

3 The Operator \mathcal{L} in $L^2(2)$: Spectral Analysis and Semigroup Estimates

3.1 Spectral Theory for \mathcal{L}

The following result discusses the spectrum of \mathcal{L} .

Proposition 1 *Let \mathcal{L} be as defined in (2.8), then*

(1) The discrete spectrum: *Let $k \in \mathbb{N} \cup \{0\}$ be fixed and $\sigma = (\sigma_1, \sigma_2)$ be such that $|\sigma| = \sigma_1 + \sigma_2 = k$. Then the function $\phi_\sigma(\xi)$ defined by*

$$\phi_\sigma(\xi) = \partial_1^{\sigma_1} \partial_2^{\sigma_2} G, \quad (3.1)$$

is an eigenfunction of \mathcal{L} of the multiplicity greater or equal to $\binom{k+1}{k} = k+1$ related to the eigenvalue $\lambda_k = 1 - \frac{3-\beta+k}{\alpha}$. In fact,

$$\sigma_d(\mathcal{L}) \supseteq \left\{ \lambda_k \in \mathbb{C} : \lambda_k = 1 - \frac{3-\beta+k}{\alpha}; k = 0, 1, 2, \dots \right\}.$$

(2) The continuous spectrum: *Let $\mu \in \mathbb{C}$ be such that $\Re \mu \leq -\frac{1}{\alpha}$ and define, $\psi_\mu \in L^2$ such that*

$$\widehat{\psi_\mu}(p) = |p|^{-\alpha\mu} e^{-|p|^\alpha}. \quad (3.2)$$

Then ψ_μ is an eigenfunction of the operator \mathcal{L} with the corresponding eigenvalue⁷ $\lambda = 1 + \mu - \frac{3-\beta}{\alpha}$. In fact,

$$\sigma_{ess}(\mathcal{L}) \supseteq \left\{ \lambda \in \mathbb{C} : \Re \lambda \leq 1 - \frac{4-\beta}{\alpha} \right\}.$$

Proof Regarding discrete spectrum, we start with a calculation, which will allow us to identify some of the eigenvalues. Let $\phi_0(\xi)$ be a radial function, i.e., $\widehat{\phi_0}(p) = g(|p|)$. Then

$$\begin{aligned} \widehat{\mathcal{L}\phi_0}(p) &= -\widehat{|\nabla|^\alpha \phi_0} + \frac{1}{\alpha} \xi \cdot \widehat{\nabla_\xi \phi_0}(p) + \left(1 + \frac{\beta-1}{\alpha}\right) \widehat{\phi_0}(p) \\ &= -|p|^\alpha \widehat{\phi_0}(p) - \frac{2}{\alpha} \widehat{\phi_0}(p) - \frac{1}{\alpha} \sum_{j=1}^2 p_j \partial_j \widehat{\phi_0}(p) + \left(1 + \frac{\beta-1}{\alpha}\right) \widehat{\phi_0}(p) \\ &= -|p|^\alpha g(|p|) - \frac{2}{\alpha} \widehat{\phi_0}(p) - \frac{1}{\alpha} \sum_{j=1}^2 p_j g'(|p|) \frac{p_j}{|p|} + \left(1 + \frac{\beta-1}{\alpha}\right) \widehat{\phi_0}(p) \\ &= \left(1 + \frac{\beta-3}{\alpha}\right) \widehat{\phi_0}(p) + \left(-|p|^\alpha g(|p|) - \frac{1}{\alpha} |p| g'(|p|)\right) \end{aligned} \quad (3.3)$$

Now if g satisfies,

$$-|p|^\alpha g(|p|) - \frac{1}{\alpha} |p| g'(|p|) = 0 \quad (3.4)$$

then clearly $\lambda = \left(1 - \frac{3-\beta}{\alpha}\right)$ is an eigenvalue for \mathcal{L} . The solution of (3.4), gives the eigenfunction, $\widehat{\phi_0}(p) = e^{-|p|^\alpha}$ or $\phi_0 = G$.

Now, let ϕ_k be an eigenfunction corresponding to the eigenvalue $\lambda_k = \left(1 - \frac{3-\beta+k}{\alpha}\right)$, that is

$$\mathcal{L}\phi_k(\xi) = \left(1 - \frac{3-\beta+k}{\alpha}\right) \phi_k \quad (3.5)$$

Taking a derivative ∂_j in (3.5), we obtain

$$\begin{aligned} \left(1 - \frac{3-\beta+k}{\alpha}\right) \partial_j \phi_k &= \partial_j \mathcal{L}\phi_k(\xi) = -|\nabla|^\alpha \partial_j \phi_k + \frac{1}{\alpha} \partial_j (\xi \cdot \nabla \phi_k) \\ &\quad + \left(1 + \frac{\beta-1}{\alpha}\right) \partial_j \phi_k \\ &= -|\nabla|^\alpha \partial_j \phi_k + \frac{1}{\alpha} \partial_j \phi_k + \frac{1}{\alpha} \xi \cdot \nabla (\partial_j \phi_k) \\ &\quad + \left(1 + \frac{\beta-1}{\alpha}\right) \partial_j \phi_k(\xi) \end{aligned}$$

⁷ Note however that all these eigenvalues are not isolated; hence, they are in the essential spectrum.

$$= \mathcal{L}[\partial_j \phi_k] + \frac{1}{\alpha} \partial_j \phi_k.$$

It follows that

$$\mathcal{L}[\partial_j \phi_k] = \left(1 - \frac{3 - \beta + (k + 1)}{\alpha}\right) \partial_j \phi_k$$

It follows that $\left(1 - \frac{3 - \beta + k + 1}{\alpha}\right)$ is an eigenvalue, corresponding to an eigenfunction $\partial_j \phi_k$. Thus, we have identified a family of eigenvalues and eigenvectors as follows. Fix $k \in \mathbb{N}$, and let (σ_1, σ_2) be so that $\sigma_1 + \sigma_2 = k$. Then, by induction, for the function $\phi_k := \partial_1^{\sigma_1} \partial_2^{\sigma_2} \phi_0$, we have (3.5). Note that what we have proved so far does not guarantee that there is not any more discrete spectrum, but merely an inclusion, as stated.

Regarding essential spectrum, we compute $\widehat{\mathcal{L}\psi_\mu}$. From the calculation (3.3), we have

$$\widehat{\mathcal{L}\psi_\mu}(p) = \left(\mu + 1 + \frac{\beta - 3}{\alpha}\right) \widehat{\psi_\mu}(p),$$

whence ψ_μ is an eigenfunction. Indeed, $\psi_\mu \in L^2(2)$, when $\Re\mu \leq -\frac{1}{\alpha}$. This is easy to see with a computation similar to the ones performed in Lemma 2.

$$\||\xi|^2 \psi_\mu\|_{L^2}^2 = \|\Delta_p \widehat{\psi_\mu}\|_{L^2}^2 = \int_0^\infty |(\partial_{\rho\rho} + \frac{1}{\rho} \partial_\rho) \rho^{-\alpha\mu} e^{-\rho^\alpha}|^2 \rho d\rho.$$

The worst term (when $\alpha > 1$) is exactly $\int_0^1 \rho^{-(3+2\alpha\mu)} d\rho$, which converges for $\Re\mu < -\frac{1}{\alpha}$. \square

3.2 The Semigroup $e^{\tau\mathcal{L}}$

The following proposition yields an explicit formula for the semigroup $e^{\tau\mathcal{L}}$. This is an extension of the formula established in Gallay and Wayne (2002a).

Proposition 2 *The operator \mathcal{L} defines a C_0 semigroup on $L^2(2)(\mathbf{R}^2)$, $e^{\tau\mathcal{L}}$. In fact, we have the following formula for its action*

$$(\widehat{e^{\tau\mathcal{L}} f})(p) = e^{(1 - \frac{3 - \beta}{\alpha})\tau} e^{-a(\tau)|p|^\alpha} \widehat{f}(e^{-\frac{\tau}{\alpha}} p), \quad (3.6)$$

$$(e^{\tau\mathcal{L}} f)(\xi) = \frac{e^{(1 - \frac{1 - \beta}{\alpha})\tau}}{a(\tau)^{\frac{2}{\alpha}}} \int_{\mathbf{R}^2} G\left(\frac{\xi - \eta}{a(\tau)^{\frac{1}{\alpha}}}\right) f(e^{\frac{\tau}{\alpha}} \eta) d\eta, \quad (3.7)$$

where $a(\tau) = 1 - e^{-\tau}$. In particular, for $1 \leq p \leq \infty$,

$$\|e^{\tau \mathcal{L}} f\|_{L^p} \leq C e^{(1 - \frac{1-\beta}{\alpha} - \frac{2}{\alpha p})\tau} \|f\|_{L^p} \quad (3.8)$$

$$\|e^{\tau \mathcal{L}} \nabla f\|_{L^p} \leq C \frac{e^{(1 - \frac{2-\beta}{\alpha} - \frac{2}{\alpha p})\tau}}{a(\tau)^{\frac{1}{\alpha}}} \|f\|_{L^p}. \quad (3.9)$$

Remark Note that $a(\tau) \sim \min(1, \tau)$. This will be used frequently in the sequel.

Proof The generation of the semigroup would follow, once we prove that the function $g : \hat{g}(\tau, p) := e^{(1 - \frac{3-\beta}{\alpha})\tau} e^{-a(\tau)|p|^\alpha} \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}})$ satisfies $\partial_\tau \hat{g}(\tau, p) = \widehat{\mathcal{L}g(\tau, \cdot)}$. Clearly, $\hat{g}(0, p) = \hat{f}(p)$, so $g(0, \xi) = f(\xi)$. Next, we compute $\partial_\tau \hat{g}(\tau, p)$. We have

$$\begin{aligned} \partial_\tau \hat{g}(\tau, p) &= \left[(1 - \frac{3-\beta}{\alpha} - a'(\tau)|p|^\alpha) \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) - \frac{1}{\alpha} e^{-\frac{\tau}{\alpha}} p \cdot \nabla_p \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) \right] \\ &\quad e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha} \\ &= \left(1 + \frac{\beta-3}{\alpha} \right) \hat{g}(\tau, p) + (a(\tau) - 1)|p|^\alpha \hat{g}(\tau, p) \\ &\quad - \frac{1}{\alpha} e^{-\frac{\tau}{\alpha}} p \cdot \nabla_p \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha}, \end{aligned}$$

where we have used the relation $a'(\tau) = 1 - a(\tau)$. Next, by (3.3), we have

$$\widehat{\mathcal{L}g(\tau, \cdot)} = -|p|^\alpha \hat{g}(\tau, p) - \frac{1}{\alpha} \sum_{j=1}^2 p_j \partial_j \hat{g}(\tau, p) + \left(1 + \frac{\beta-3}{\alpha} \right) \hat{g}(\tau, p).$$

But,

$$\begin{aligned} \frac{1}{\alpha} \sum_{j=1}^2 p_j \partial_j \hat{g}(\tau, p) &= \frac{1}{\alpha} \sum_{j=1}^2 p_j \left(-\alpha a(\tau) p_j |p|^{\alpha-2} \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) + e^{-\frac{\tau}{\alpha}} \partial_j \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) \right) \\ &\quad e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha} \\ &= -a(\tau)|p|^\alpha \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha} \\ &\quad + \frac{1}{\alpha} e^{-\frac{\tau}{\alpha}} p \cdot \nabla_p \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha}. \end{aligned}$$

Altogether,

$$\begin{aligned} \widehat{\mathcal{L}g(\tau, \cdot)} &= -|p|^\alpha \hat{g}(\tau, p) + \left(1 + \frac{\beta-3}{\alpha} \right) \hat{g}(\tau, p) + a(\tau)|p|^\alpha \hat{g}(\tau, p) - \\ &\quad - \frac{1}{\alpha} e^{-\frac{\tau}{\alpha}} p \cdot \nabla_p \hat{f}(p \cdot e^{-\frac{\tau}{\alpha}}) e^{\tau(1 - \frac{3-\beta}{\alpha})} e^{-a(\tau)|p|^\alpha}. \end{aligned}$$

An immediate inspection reveals that $\partial_\tau \hat{g}(\tau, p) = \widehat{\mathcal{L}g(\tau, \cdot)}(p)$ and so the semigroup formula (3.6) is established. The formula (3.7) is just a Fourier inversion of (3.6).

Regarding the estimate (3.8), we proceed as follows

$$\|e^{\tau \mathcal{L}} f\|_{L^p} \leq e^{(1-\frac{1-\beta}{\alpha})\tau} \|G_{a(\tau)^{\frac{1}{\alpha}}} \|_{L^1} \|f(e^{\frac{\tau}{\alpha}} \cdot)\|_{L^p} = e^{(1-\frac{1-\beta}{\alpha}-\frac{2}{\alpha p})\tau} \|G\|_{L^1} \|f\|_{L^p}.$$

For (3.9), note that integration by parts yields

$$\begin{aligned} (e^{\tau \mathcal{L}} \partial_j f)(\xi) &= \frac{e^{(1-\frac{1-\beta}{\alpha})\tau}}{a(\tau)^{\frac{2}{\alpha}}} \int_{\mathbf{R}^2} G\left(\frac{\xi - \eta}{a(\tau)^{\frac{1}{\alpha}}}\right) (\partial_j f)(e^{\frac{\tau}{\alpha}} \eta) d\eta \\ &= \frac{e^{(1-\frac{2-\beta}{\alpha})\tau}}{a(\tau)^{\frac{3}{\alpha}}} \int_{\mathbf{R}^2} \partial_j G\left(\frac{\xi - \eta}{a(\tau)^{\frac{1}{\alpha}}}\right) f(e^{\frac{\tau}{\alpha}} \eta) d\eta, \end{aligned}$$

whence

$$\|(e^{\tau \mathcal{L}} \nabla f)(\xi)\|_{L^p} \leq \frac{e^{(1-\frac{2-\beta}{\alpha}-\frac{2}{\alpha p})\tau}}{a(\tau)^{\frac{1}{\alpha}}} \|\nabla G\|_{L^1} \|f\|_{L^p}.$$

□

We need a variant of Proposition A.2 in Gallay and Wayne (2002a), which discusses the commutation of the semigroup with differential operators.

Lemma 7 *We have the following commutation relation for $e^{\tau \mathcal{L}}$*

$$\nabla e^{\tau \mathcal{L}} = e^{\frac{\tau}{\alpha}} e^{\tau \mathcal{L}} \nabla \quad (3.10)$$

Proof Let $u(x, \tau) = e^{\tau \mathcal{L}} f(x)$, then u satisfies the following equation

$$\begin{cases} u_\tau = \mathcal{L}u, \\ u(0, x) = f(x). \end{cases}$$

Clearly, taking a derivative ∂_j in (2.8) yields, for $j = 1, 2$

$$\begin{cases} (\partial_j u)_\tau = \partial_j(\mathcal{L}u) = \mathcal{L}\partial_j u + \frac{1}{\alpha} \partial_j u, \\ \partial_j u(x, 0) = \partial_j f(x), \end{cases}$$

which has the solution $\partial_j u(x, \tau) = e^{\tau[\mathcal{L}+\frac{1}{\alpha}]} \partial_j f(x)$. In other words $\nabla e^{\tau \mathcal{L}} = e^{\frac{\tau}{\alpha}} e^{\tau \mathcal{L}} \nabla$. □

3.3 Semigroup Estimates

We need to address an important question, namely the behavior of the bounded operators $e^{\tau \mathcal{L}}$ on $L^2(2)$. The next Proposition does that. More precisely, we are interested in the decay of the operator norms $\|e^{\tau \mathcal{L}}\|_{L^2(2) \rightarrow L^2(2)}$. Importantly, good decay estimates

only happen, when the functions have mean value zero. The proof of this proposition will be given in “Appendix D.”

Proposition 3 *Let $f \in L^2(2)$, $\hat{f}(0) = 0$ and $\gamma = (\gamma_1, \gamma_2) \in \mathbf{N}^2$, $|\gamma| = 0, 1$ and $0 < \epsilon << 1$. Then there exists $C = C_\epsilon > 0$, such that for any $\tau > 0$,*

$$\|\nabla^\gamma(e^{\tau\mathcal{L}}f)\|_{L^2(2)} \leq C \frac{e^{\left(1-\frac{4-\beta}{\alpha}+\epsilon\right)\tau}}{a(\tau)^{\frac{|\gamma|}{\alpha}}} \|f\|_{L^2(2)}, \quad (3.11)$$

or

$$\|\nabla^\gamma(e^{\tau\mathcal{L}}f)\|_{L^2(2)} \leq C \|f\|_{L^2(2)} \cdot \begin{cases} \tau^{-\frac{|\gamma|}{\alpha}}, & \tau \leq 1 \\ e^{\left(1-\frac{4-\beta}{\alpha}+\epsilon\right)\tau}, & \tau > 1 \end{cases} \quad (3.12)$$

3.4 The Decay Estimates for $e^{\tau\mathcal{L}}$ Give a Description of the Spectrum of \mathcal{L}

In this section, we show that the spectral inclusions in Proposition 1 are actually equalities. We also compute explicitly the Riesz projection \mathcal{P}_0 onto the eigenvalue of \mathcal{L} with the largest real part. In Proposition 1, we have already identified G as being an eigenfunction for \mathcal{L} corresponding to an eigenvalue $\lambda_0 = 1 - \frac{3-\beta}{\alpha}$. On the other hand, applying Proposition 3, for functions with $\hat{f}(0) = 0$ and $\gamma = (0, 0)$, implies

$$\|e^{\tau\mathcal{L}}f\|_{L^2(2)} \leq C_\epsilon e^{\left(1-\frac{4-\beta}{\alpha}+\epsilon\right)\tau} \|f\|_{L^2(2)}. \quad (3.13)$$

Denote the co-dimension one subspace $X_0 = \{f \in L^2(2) : \hat{f}(0) = 0\}$. Clearly, the operator \mathcal{L} acts invariantly on X_0 , since for every $f \in L^2(2) : \int f(\xi)d\xi = 0$, we have $\int_{\mathbf{R}^2} \xi \cdot \nabla f d\xi = 0$, whence $\int \mathcal{L}f(\xi)d\xi = 0$.

Introduce $\mathcal{L}_0 := \mathcal{L}|_{X_0}$, with domain $D(\mathcal{L}_0) = D(\mathcal{L}) \cap X_0 = H^\alpha \cap X_0$. By the Hille-Yosida theorem, this estimate (3.13) implies that the set $\{\lambda : \Re \lambda > \left(1 - \frac{4-\beta}{\alpha}\right)\}$ is in the resolvent set of \mathcal{L}_0 , since the integral representing $(\lambda - \mathcal{L})^{-1}$, namely $\int_0^\infty e^{-\lambda\tau} e^{\tau\mathcal{L}} d\tau$, converges by virtue of (3.13). Combining this with the results from Proposition 1, we conclude that $\sigma(\mathcal{L}) \cap \{\lambda : \Re \lambda > \left(1 - \frac{4-\beta}{\alpha}\right)\}$ is a singleton—the eigenvalue $\lambda_0 = 1 - \frac{3-\beta}{\alpha}$, which is simple, with eigenfunction G . We conclude that

$$\sigma(\mathcal{L}) = \left\{1 - \frac{3-\beta}{\alpha}\right\} \cup \sigma_{ess}(\mathcal{L}); \quad \sigma_{ess}(\mathcal{L}) = \left\{\lambda : \Re \lambda \leq \left(1 - \frac{4-\beta}{\alpha}\right)\right\},$$

Moreover, its Riesz projection \mathcal{P}_0 , a rank one operator, is given by

$$\mathcal{P}_0 f = \left(\int_{\mathbf{R}^2} f(\xi) d\xi \right) G$$

Clearly, such an operator is well-normalized, since $\mathcal{P}_0^2 f = \langle G, 1 \rangle \mathcal{P}_0 f = \hat{G}(0) \mathcal{P}_0 f = \mathcal{P}_0 f$, since $\hat{G}(0) = 1$. The projection $\mathcal{Q}_0 := Id - \mathcal{P}_0$ over the complementary part of the spectrum, satisfies $\mathcal{L}_0 = \mathcal{Q}_0 \mathcal{L} \mathcal{Q}_0$. Also, $\mathcal{Q}_0 : L^2(2) \rightarrow X_0$. Now, (3.13) can be reformulated as

$$\|\nabla^\gamma e^{\tau \mathcal{L}_0} f\|_{L^2(2)} \leq C_\epsilon \frac{e^{\left(1 - \frac{4-\beta}{\alpha} + \epsilon\right)\tau}}{a(\tau)^{\frac{|\gamma|}{\alpha}}} \|f\|_{L^2(2)}. \quad (3.14)$$

for any function f , since $e^{\tau \mathcal{L}_0} f = e^{\tau \mathcal{L}} \mathcal{Q}_0 f$ and the entry $\mathcal{Q}_0 f$ has mean value zero, so (3.13) is applicable. In addition, we can derive estimates for the action of the semigroup $e^{\tau \mathcal{L}}$ on $L^2(2)$, without the mean value zero property $\hat{f}(0) = 0$.

Proposition 4 *Let $f \in L^2(2)$. Then, there exists a constant C , so that*

$$\|\nabla^\gamma (e^{\tau \mathcal{L}} f)\|_{L^2(2)} \leq C \frac{e^{\left(1 - \frac{3-\beta}{\alpha}\right)\tau}}{a(\tau)^{\frac{|\gamma|}{\alpha}}} \|f\|_{L^2(2)}. \quad (3.15)$$

Proof We use the decomposition

$$f = \mathcal{P}_0 f + \mathcal{Q}_0 f = \langle f, 1 \rangle G + [f - \langle f, 1 \rangle G].$$

Thus,

$$e^{\tau \mathcal{L}} f = \langle f, 1 \rangle e^{\tau(1 - \frac{3-\beta}{\alpha})} G + e^{\tau \mathcal{L}_0} [f]$$

It follows that

$$\begin{aligned} \|e^{\tau \mathcal{L}} f\|_{L^2(2)} &\leq C |\langle f, 1 \rangle| e^{\tau(1 - \frac{3-\beta}{\alpha})} \|G\|_{L^2(2)} + C_\epsilon e^{\left(1 - \frac{4-\beta}{\alpha} + \epsilon\right)\tau} \|f\|_{L^2(2)} \\ &\leq C e^{\tau(1 - \frac{3-\beta}{\alpha})} \|f\|_{L^2(2)}, \end{aligned}$$

where we have used (3.14) and $|\langle f, 1 \rangle| \leq C \|f\|_{L^2(2)}$. Similar estimates can be derived, as before, for $|\nabla|^\gamma e^{\tau \mathcal{L}}$, we omit the details. \square

4 Local and Global Well-Posedness of the SQG

The local and global theory of the Cauchy problem for SQG has been well-studied in the literature. Local and global well-posedness holds under very general conditions on initial data. Regardlessly, we will present a few results for our problem (1.3). This is necessary, since we assume a nonstandard relation between u and z , but also because we need precise properties, beyond the scope of the well-posedness. Then, we will turn to properties of the rescaled equation, (2.7). We will do so, both in L^p spaces as

well as in $L^2(2)$ spaces—the reason is that we will use some of our preliminary results as *a priori* estimates in the subsequent Lemmas.

Our first results are about the well-posedness of the standard model (1.3) in L^p spaces.⁸

4.1 Global Well-Posedness and a Priori Estimates in L^p Spaces

Lemma 8 *Suppose that $z_0 \in L^1 \cap L^\infty =: X$. Then, (1.3) is globally well-posed in the space X . Moreover, for every $p \in [1, \infty]$, $t \rightarrow \|z(\cdot, t)\|_{L^p}$ is non-increasing in time.*

Proof We first prove the local existence of the strong solution in the space $C([0, T); X)$, that is, with T to be determined, we are looking for a fixed point of the integral equation

$$z(\xi) = e^{-t|\nabla|^\alpha} z_0 - \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u \cdot z) \, ds. \quad (4.1)$$

According to Lemma (3) $\|e^{-t|\nabla|^\alpha} z_0\|_{L^1 \cap L^\infty} \leq C_0 \|z_0\|_{L^1 \cap L^\infty}$. For any $T > 0$ and $t \in (0, T)$, consider

$$Q(z_1, z_2) := \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u_1 \cdot z_2) \, ds,$$

where u_1 is given by $u_1 = (\nabla^\perp)^{-\beta} z_1$. For $t \in (0, T)$, using (2.17)

$$\begin{aligned} \|Q(z_1(t), z_2(t))\|_{L^1} &= \left\| \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u_1 \cdot z_2) \, ds \right\|_{L^1} \leq C \int_0^t \frac{1}{(t-s)^{\frac{1}{\alpha}}} \|(u_1 \cdot z_2)\|_{L^1} \, ds \\ &\leq C t^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|u_1(s, \cdot)\|_{L^\infty} \sup_{0 \leq s \leq T} \|z_2(s, \cdot)\|_{L^1} \\ &\lesssim T^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} (\|z_1(s, \cdot)\|_{L^{\frac{2}{\beta}+\epsilon}} + \|z_1(s, \cdot)\|_{L^{\frac{2}{\beta}-\epsilon}}) \sup_{0 \leq s \leq T} \|z_2(s, \cdot)\|_{L^1} \\ &\lesssim T^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|z_1\|_X \sup_{0 \leq s \leq T} \|z_2\|_X. \end{aligned}$$

where we have used the Sobolev embedding estimate (2.2). Similarly,

$$\|Q(z_1, z_2)\|_{L^\infty} \leq C T^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|u_1\|_{L^\infty} \sup_{0 \leq s \leq T} \|z_2\|_{L^\infty} \leq C T^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|z_1\|_X \sup_{0 \leq s \leq T} \|z_2\|_X.$$

Finally, following similar path, we also have

$$\|Q(z_1, z_1) - Q(z_2, z_2)\|_X \leq C T^{1-\frac{1}{\alpha}} (\|z_1\|_X + \|z_2\|_X) \|z_1 - z_2\|_X.$$

⁸ The results can be made more precise, in individual L^p spaces, rather than in all L^p spaces. We will not do so here, because our goal is to extend to $L^2(2)$, which is yet smaller space.

Upon introducing $Y_T := \{z : \sup_{0 \leq t \leq T} \|z(t, \cdot)\|_X \leq 2C_0\|z_0\|_X\}$ and taking into account the estimates above, we realize that the mapping (4.1) has a fixed point in the metric space $C([0, T], X)$, for small enough $T = T(\|z_0\|_X)$. In fact, the argument shows that $T \sim \|z_0\|_X^{-\frac{\alpha}{\alpha-1}}$.

For the global existence, we need to show that the $t \rightarrow \|z(t, \cdot)\|_{L^p}$ does not blow up in finite time. In fact, we show that the $t \rightarrow \|z(t, \cdot)\|_{L^p}$ is non-increasing, which will allow us to conclude global existence as well. To that end, we dot product the Eq. (1.3) with $|z|^{p-2}z$, $p \in (1, \infty)$ to get

$$\frac{1}{p} \partial_t \|z\|_{L^p}^p + \int_{\mathbf{R}^2} |\nabla|^\alpha z \cdot |z|^{p-2} z d\xi = 0.$$

By the positivity estimate (2.12), we have $\int_{\mathbf{R}^2} |\nabla|^\alpha z \cdot |z|^{p-2} z d\xi \geq 0$. Therefore, $\partial_t \|z\|_{L^p}^p \leq 0$, and $t \rightarrow \|z(t, \cdot)\|_{L^p}$ is non-increasing in time. For $p = 1$, $p = \infty$ the monotonicity follows from an approximation argument from the cases $1 < p < \infty$. \square

Our next result is about *a priori* estimates in L^p spaces, but this time in the rescaled variable formulation, (2.7). Note that the global existence of the rescaled equation is not in question anymore, due to Lemma 8. However, we show precise decay estimates for the norm of the solution Z . This fairly elementary Lemma already shows the advantage of the rescaled variables approach and its far reaching consequences.

Lemma 9 *Let $Z_0 \in L^1 \cap L^\infty(\mathbf{R}^2)$, $\alpha \in (1, 2)$, $0 \leq \beta < 2$ and $p \in [1, \infty)$. Then the unique global strong solution Z of (2.7) satisfies*

$$\|Z(\tau)\|_{L^p} \leq \|Z_0\|_{L^p} e^{-\tau \left(\frac{2}{p\alpha} - 1 - \frac{\beta-1}{\alpha} \right)}. \quad (4.2)$$

Proof If we dot product (2.7) with $Z|Z|^{p-2}$, we have by the positivity estimate (2.12), $\int_{\mathbf{R}^2} |\nabla|^\alpha Z \cdot |Z|^{p-2} Z d\xi \geq 0$. Furthermore, using the divergent free property of $U(\xi)$

$$\begin{aligned} \frac{1}{p} \frac{d}{d\tau} \|Z\|_{L^p}^p &\leq \frac{1}{\alpha} \int (\xi \cdot \nabla_\xi Z) Z |Z|^{p-2} d\xi - \int (U \cdot \nabla_\xi Z) Z |Z|^{p-2} d\xi \\ &+ \left(1 + \frac{\beta-1}{\alpha} \right) \|Z\|_{L^p}^p = \left(1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right) \|Z\|_{L^p}^p, \end{aligned} \quad (4.3)$$

therefore, we arrive at

$$\frac{1}{p} \frac{d}{d\tau} \|Z\|_{L^p}^p + \left(\frac{2}{\alpha p} - 1 - \frac{\beta-1}{\alpha} \right) \|Z\|_{L^p}^p \leq 0.$$

Now we use the Gronwall's inequality to finish the proof. \square

The Lemma above shows *a priori* bound for $\|Z(\tau, \cdot)\|_{L^p}$, for any $p \in [1, \infty]$, and a decay rate for $p < \frac{2}{\alpha+\beta-1}$, but it is not giving any decay rate for $p \geq \frac{2}{\alpha+\beta-1}$. On the other hand, as we shall see later, the decay rate predicted by Lemma 9 is in fact

optimal for $p = 1$ (but certainly not so, for any other value of p). We can bootstrap the results of Lemma 9 in the next Lemma to find, what *it will turn out to be, the optimal decay rate for any $p \geq 1$* .⁹

Lemma 10 *Let $Z_0 \in L^1 \cap L^\infty(\mathbf{R}^2)$, $1 \leq p \leq \infty$ and $\alpha \in (1, 2)$, $\alpha + \beta \leq 3$. Then, there exists constant $C = C_{p,\alpha,\beta}$, so that the unique global strong solution Z of (2.7) satisfies*

$$\|Z(\tau, \cdot)\|_{L^p} \leq C_{p,\alpha,\beta} \|Z_0\|_{L^1 \cap L^\infty} e^{-(\frac{3-\beta-\alpha}{\alpha})\tau}. \quad (4.4)$$

Proof Recall that the estimate (2.14) is available to us. Taking dot product $|Z|^{p-2}Z$ and taking into account (2.14) which implies $\int_{\mathbf{R}^2} |\nabla|^\alpha Z \cdot |Z|^{p-2}Z d\xi \geq c_{p,\alpha} \|Z\|_{L^{\frac{2p}{2-\alpha}}}^2$.

We further add $C\|Z\|_{L^p}^p$, for some large C , to be determined. We have

$$\frac{1}{p} \frac{d}{d\tau} \|Z\|_{L^p}^p + C\|Z\|_{L^p}^p + c_{p,\alpha} \|Z\|_{L^{\frac{2p}{2-\alpha}}}^2 \leq \left(C + 1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right) \|Z\|_{L^p}^p$$

By Gagliardo–Nirenberg's, with $\gamma = \frac{2p-2}{2p-2+\alpha}$, $\|Z\|_{L^p} \leq \|Z\|_{L^{\frac{2p}{2-\alpha}}}^\gamma \|Z\|_{L^1}^{1-\gamma}$, whence by Young's inequality

$$\begin{aligned} & \frac{1}{p} \frac{d}{d\tau} \|Z\|_{L^p}^p + C\|Z\|_{L^p}^p + c_{p,\alpha} \|Z\|_{L^{\frac{2p}{2-\alpha}}}^p \\ & \leq \left(C + 1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right) \|Z\|_{L^{\frac{2p}{2-\alpha}}}^{p\gamma} \|Z\|_{L^1}^{p(1-\gamma)} \\ & \leq \epsilon_0 \|Z\|_{L^{\frac{2p}{2-\alpha}}}^p + \frac{\left(C + 1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right)^{\frac{1}{1-\gamma}}}{\epsilon_0^{\frac{\gamma}{1-\gamma}}} \|Z\|_{L^1}^p \end{aligned}$$

and $\epsilon_0 > 0$ is a fixed number, say we select it $\epsilon_0 = c_{p,\alpha}$. Then

$$\begin{aligned} & \frac{1}{p} \frac{d}{d\tau} \|Z\|_{L^p}^p + C\|Z\|_{L^p}^p \leq \frac{\left(C + 1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right)^{\frac{1}{1-\gamma}}}{\epsilon_0^{\frac{\gamma}{1-\gamma}}} \|Z\|_{L^1}^p \\ & \leq \frac{\left(C + 1 + \frac{\beta-1}{\alpha} - \frac{2}{\alpha p} \right)^{\frac{1}{1-\gamma}}}{\epsilon_0^{\frac{\gamma}{1-\gamma}}} \|Z_0\|_{L^1}^p e^{-p\tau(\frac{3-\beta-\alpha}{\alpha})}, \end{aligned}$$

where we have used Lemma (9) to estimate $\|Z(\tau, \cdot)\|_{L^1}$. Denoting $\mu := (\frac{3-\beta-\alpha}{\alpha}) \geq 0$, select $C = \mu + 1$. We have

$$I'(\tau) + p(\mu + 1)I(\tau) \leq D\|Z_0\|_{L^1}^p e^{-p\mu\tau},$$

⁹ For generic data.

where $I(\tau) = \|Z(\tau)\|_{L^p}^p$, $D = p^{1+\frac{\gamma}{1-\gamma}} \frac{(\mu+2+\frac{\beta-1}{\alpha}-\frac{2}{\alpha p})^{\frac{1}{1-\gamma}}}{c_\alpha^{\frac{\gamma}{1-\gamma}}}$. Now we use the Gronwall's inequality to derive the estimate

$$I(\tau) \leq e^{-p(\mu+1)\tau} I(0) + \frac{D}{p} \|Z_0\|_{L^1}^p e^{-p\mu\tau}.$$

Taking p^{th} root and simplifying yields the final estimate

$$\|Z(\tau)\|_{L^p} \leq (\|Z_0\|_{L^p} + \left(\frac{D}{p}\right)^{\frac{1}{p}} \|Z_0\|_{L^1}) e^{-\mu\tau} \leq (1 + \left(\frac{D}{p}\right)^{\frac{1}{p}}) \|Z_0\|_{L^1 \cap L^\infty} e^{-\mu\tau}.$$

For the case $p = \infty$, we take limits in the previous identity, for fixed $\tau > 0$, as $p \rightarrow \infty$. Note that by the explicit form of D_p , $\lim_{p \rightarrow \infty} \left(\frac{D}{p}\right)^{\frac{1}{p}} = 1$, so (4.4) holds true in this case with $C = 2$. \square

4.2 Global Solutions and a Priori Estimates in $L^2(2)$

From the previous section, we know that the SQG equation in its standard form, namely (1.3), has global solutions in L^p . Thus, the rescaled equation (2.7) also has unique global (strong) solutions in L^p . We now would like to understand the Cauchy problem in the smaller space $L^2(2)$. In particular, even if the initial data is well-localized, say $Z(0, \cdot) \in L^2(2)$, it is not *a priori* clear why the solution $Z(\tau)$ will stay in $L^2(2)$ for (any) later time $\tau > 0$. In other words, one needs to start with the local well-posedness for (2.7), and then, we shall upgrade it to a global one, by means of *a priori* estimates on $\|Z(\tau)\|_{L^2(2)}$.

Theorem 3 Suppose that $Z_0 \in L^2(2)(\mathbf{R}^2) \cap L^\infty(\mathbf{R}^2) =: X$. Then (2.7) has an unique global strong solution $Z \in C^0([0, \infty]; L^2(2)(\mathbf{R}^2) \cap L^\infty(\mathbf{R}^2))$, with $Z(0) = Z_0$. In addition, there is the *a priori* estimate

$$\|Z(\tau)\|_{L^2(2) \cap L^\infty} \leq C e^{-\tau \left(\frac{3-\alpha-\beta}{\alpha}\right)} \|Z_0\|_{L^2(2) \cap L^\infty}, \quad (4.5)$$

where C is an absolute constant.

Proof We set up a local well-posedness scheme for the integral equation corresponding to (2.7), with initial data $Z(0) = f$, namely

$$Z(\tau) = e^{\tau \mathcal{L}} f - \int_0^\tau e^{(\tau-s)\mathcal{L}} \nabla \cdot (U Z) \, ds, \quad (4.6)$$

where $U = U_Z = (|\nabla|^\perp)^{-\beta} Z$. We have, according to (3.8) and (3.15),

$$\|e^{\tau \mathcal{L}} f\|_{L^2(2)} + \|e^{\tau \mathcal{L}} f\|_{L^\infty} \leq C (e^{(1-\frac{1-\beta}{\alpha})\tau} + e^{(1-\frac{3-\beta}{\alpha})\tau}) \|f\|_{L^2(2) \cap L^\infty}$$

Thus, with $T \leq 1$ to be determined later, set

$$Y_T := \{Z(\tau, \cdot) \in X : \sup_{0 \leq s \leq T} \|Z(s, \cdot)\|_X \leq 2C \left(e^{(1-\frac{1-\beta}{\alpha})} + e^{(1-\frac{3-\beta}{\alpha})} \right) \|f\|_X\},$$

where the bound in Y is selected to be twice the value of the bound above, at $\tau = 1$. For the nonlinear term, we have for each $\tau \in (0, T)$,

$$\begin{aligned} \left\| \int_0^\tau e^{(\tau-s)\mathcal{L}} \nabla \cdot (U_{Z_1} Z_2) \, ds \right\|_{L^\infty} &\leq C \int_0^\tau C \frac{e^{(1-\frac{2-\beta}{\alpha})(\tau-s)}}{a(\tau-s)^{\frac{1}{\alpha}}} \|U_{Z_1}(s) Z_2(s)\|_{L^\infty} \, ds \leq \\ &\leq C \sup_{0 \leq s \leq T} \|U_{Z_1}\|_{L^\infty} \sup_{0 \leq s \leq T} \|Z_2\|_{L^\infty} \int_0^\tau \frac{1}{(\tau-s)^{\frac{1}{\alpha}}} \, ds \leq \\ &\leq CT^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} (\|Z_1\|_{L^{\frac{2}{\beta}+\epsilon}} + \|Z_1\|_{L^{\frac{2}{\beta}-\epsilon}}) \sup_{0 \leq s \leq T} \|Z_2\|_{L^\infty} \\ &\leq CT^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|Z_1\|_X \sup_{0 \leq s \leq T} \|Z_2\|_X, \end{aligned}$$

where we have used (3.9), $e^{(1-\frac{2-\beta}{\alpha})(\tau-s)} \leq 3$, $a(\tau-s) = 1 - e^{-(\tau-s)} \sim (\tau-s)$, for $0 < s < \tau \leq 1$, the Sobolev embedding estimate (2.2) and finally the fact that $X = L^2(2) \cap L^\infty \hookrightarrow L^1 \cap L^\infty$. For the other norm in the definition of X , we have by Lemma 7,

$$\begin{aligned} \left\| \int_0^\tau e^{(\tau-s)\mathcal{L}} \nabla \cdot (U_{Z_1} \cdot Z_2) \, ds \right\|_{L^2(2)} &= \int_0^\tau e^{-\frac{(\tau-s)}{\alpha}} \|\nabla \cdot e^{(\tau-s)\mathcal{L}} (U_{Z_1} \cdot Z_2)\|_{L^2(2)} \, ds \\ &\leq C \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}} e^{(1-\frac{3-\beta}{\alpha})(\tau-s)}}{a(\tau-s)^{\frac{1}{\alpha}}} \|U_{Z_1}(s) \cdot Z_2(s)\|_{L^2(2)} \, ds \leq \\ &\leq C \sup_{0 \leq s \leq T} \|U_{Z_1}(s)\|_{L^\infty} \sup_{0 \leq s \leq T} \|Z_2(s)\|_{L^2(2)} \int_0^\tau \frac{1}{(\tau-s)^{\frac{1}{\alpha}}} \, ds \\ &\leq CT^{1-\frac{1}{\alpha}} \sup_{0 \leq s \leq T} \|Z_1\|_X \sup_{0 \leq s \leq T} \|Z_2\|_{L^2(2)}. \end{aligned}$$

Having these two bilinear estimates allows us to conclude that for sufficiently small T , of the form $T \sim \|f\|_X^{-\frac{\alpha}{\alpha-1}}$ (which should also be taken $T \leq 1$), we have local well-posedness in the space X .

Regarding global existence in $X = L^2(2) \cap L^\infty$, we obviously need *a priori* estimates for the solution to prevent potential blow up. We already have those in L^∞ and in L^2 , by the results of Lemma 10. Thus, it remains to control the norm $J(\tau) := \int_{\mathbf{R}^2} |\xi|^4 |Z(\tau, \xi)|^2 \, d\xi$. To this end, take a dot product of the Eq. (2.7) with

$|\xi|^4 Z$. We have

$$\begin{aligned} \partial_\tau \frac{1}{2} \int |\xi|^4 Z^2 d\xi + \int |\xi|^4 |\nabla|^\alpha Z \cdot Z d\xi = \\ = \frac{1}{\alpha} \int (\xi \cdot \nabla_\xi Z) |\xi|^4 Z d\xi - \int (U \cdot \nabla_\xi Z) |\xi|^4 Z d\xi + (1 + \frac{\beta-1}{\alpha}) \int |\xi|^4 Z^2 d\xi. \end{aligned}$$

We first analyze the terms on the right-hand side. Integration by parts yields

$$\begin{aligned} \frac{1}{\alpha} \int (\xi \cdot \nabla_\xi Z) |\xi|^4 Z d\xi = -\frac{3}{\alpha} \int |\xi|^4 Z^2 d\xi; \quad \int (U \cdot \nabla_\xi Z) |\xi|^4 Z d\xi \\ = -2 \int |\xi|^2 (\xi \cdot U) Z^2 d\xi. \end{aligned}$$

Note that by Young's inequality, we have for all $\epsilon > 0$

$$\begin{aligned} \left| \int |\xi|^2 (\xi \cdot U) Z^2 d\xi \right| &\leq C \int |\xi|^3 \|U\|_{L^\infty} Z^2(\xi) d\xi \\ &\leq \epsilon \int |\xi|^4 Z^2(\xi) d\xi + C\epsilon^{-3} \|U\|_{L^\infty}^4 \|Z\|_{L^2}^2. \end{aligned}$$

By the Sobolev embedding (2.2) and Lemma 10, we have

$$\|U\|_{L^\infty} \leq C \left(\|Z\|_{L^{\frac{2}{\beta}+\epsilon}} + \|Z\|_{L^{\frac{2}{\beta}-\epsilon}} \right) \leq C e^{-(\frac{3-\beta-\alpha}{\alpha})\tau},$$

where $C = C(Z_0)$. So for every $\epsilon > 0$, we have the estimate

$$\left| \int |\xi|^2 (\xi \cdot U) Z^2 d\xi \right| \leq \epsilon \int |\xi|^4 Z^2(\xi) d\xi + C\epsilon^{-3} e^{-6\tau(\frac{3-\beta-\alpha}{\alpha})}.$$

The term $\int |\xi|^4 |\nabla|^\alpha Z \cdot Z d\xi$ will give rise to some harder error terms (involving commutators between the $|\nabla|^{\alpha/2}$ and the weights), which we need to eventually control. It turns out that the most advantageous way to reign in the error terms is to split the weight $|\xi|^4$ between the two entries. More precisely,

$$\begin{aligned} \int |\xi|^4 |\nabla|^\alpha Z \cdot Z d\xi &= \int |\xi|^2 |\nabla|^\alpha Z \cdot |\xi|^2 Z d\xi = \langle |\xi|^2 |\nabla|^{\alpha/2} [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle \\ &= \langle |\nabla|^{\alpha/2} |\xi|^2 [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle - \langle [|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle \\ &= \langle |\xi|^2 [|\nabla|^{\alpha/2} Z], |\nabla|^{\alpha/2} [|\xi|^2 Z] \rangle - \langle [|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle \\ &= \langle |\xi|^2 |\nabla|^{\alpha/2} Z, |\xi|^2 |\nabla|^{\alpha/2} Z \rangle + \langle |\xi|^2 |\nabla|^{\alpha/2} Z, [|\nabla|^{\alpha/2}, |\xi|^2] Z \rangle \\ &\quad - \langle [|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle \\ &= \int |\xi|^4 |\nabla|^{\alpha/2} Z^2 d\xi + \langle |\xi|^2 |\nabla|^{\alpha/2} Z, [|\nabla|^{\alpha/2}, |\xi|^2] Z \rangle \\ &\quad - \langle [|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle. \end{aligned}$$

Denote the error terms $E := \langle |\xi|^2 |\nabla|^{\alpha/2} Z, [|\nabla|^{\alpha/2}, |\xi|^2] Z \rangle - \langle [|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z], |\xi|^2 Z \rangle$. Putting it all together implies

$$\begin{aligned} \frac{1}{2} J'(\tau) + \left(\frac{4-\alpha-\beta}{\alpha} - \epsilon \right) J(\tau) + \int |\xi|^4 |\nabla|^{\frac{\alpha}{2}} Z|^2 d\xi &\leq |E| + C\epsilon^{-3} e^{-6\tau(\frac{3-\beta-\alpha}{\alpha})} \\ &\lesssim \||\xi|^2 |\nabla|^{\alpha/2} Z\|_{L^2} \|[|\nabla|^{\alpha/2}, |\xi|^2] Z\|_{L^2} \\ &\quad + \|[|\nabla|^{\alpha/2}, |\xi|^2] [|\nabla|^{\alpha/2} Z]\|_{L^2} \||\xi|^2 Z\|_{L^2} + \epsilon^{-3} e^{-6\tau(\frac{3-\beta-\alpha}{\alpha})}. \end{aligned} \quad (4.7)$$

□

At this point, it becomes clear that we need to control the commutator expression above. In fact, we have the following Lemma.

Lemma 11 *Let $\alpha \in (1, 2)$. Then, there is $C = C_\alpha$, so that*

$$\|[|\nabla|^{\alpha/2}, |\xi|^2] f\|_{L^2(\mathbf{R}^2)} \leq C \||\xi|^{2-\frac{\alpha}{2}} f\|_{L^2(\mathbf{R}^2)}. \quad (4.8)$$

We postpone the proof of Lemma 11 for the “Appendix,” see Section C. We finish the proof of Theorem 3 based upon it. By Gagliardo–Nirenberg’s inequality

$$\||\xi|^{2-\frac{\alpha}{2}} g\|_{L^2} \leq \||\xi|^2 g\|_{L^2}^{1-\frac{\alpha}{4}} \|g\|_{L^2}^{\frac{\alpha}{4}}.$$

Continuing with our arguments above [see (4.7)], we conclude from Lemma 11 that

$$\begin{aligned} \frac{1}{2} J'(\tau) + \left(\frac{4-\alpha-\beta}{\alpha} - \epsilon \right) J(\tau) + \||\xi|^2 |\nabla|^{\alpha/2} Z\|_{L^2}^2 &\leq \epsilon \||\xi|^2 |\nabla|^{\alpha/2} Z\|_{L^2}^2 \\ &\quad + \epsilon \||\xi|^2 Z\|_{L^2}^2 + C_\epsilon \|Z\|_{L^2}^2 + \epsilon^{-3} e^{-6\tau(\frac{3-\beta-\alpha}{\alpha})} \end{aligned}$$

All in all, for all $\epsilon < 1$, we have by Lemma 10,

$$\begin{aligned} \frac{1}{2} J'(\tau) + \left(\frac{4-\alpha-\beta}{\alpha} - 2\epsilon \right) J(\tau) &\leq C_\epsilon \|Z\|_{L^2}^2 \leq C \|Z_0\|_{L^1 \cap L^\infty}^2 e^{-2\tau(\frac{3-\beta-\alpha}{\alpha})} \\ &\quad + \epsilon^{-3} e^{-6\tau(\frac{3-\beta-\alpha}{\alpha})}. \end{aligned}$$

By Gronwall’s, we finally conclude that

$$J(\tau) \leq J(0) e^{-2\tau(\frac{4-\alpha-\beta}{\alpha} - 2\epsilon)} + C \|Z_0\|_{L^1 \cap L^\infty}^2 e^{-2\tau(\frac{3-\alpha-\beta}{\alpha})}.$$

As a consequence

$$\||\xi|^2 Z(\tau)\|_{L^2} \leq C \|Z_0\|_{L^2(2) \cap L^\infty} e^{-\tau(\frac{3-\alpha-\beta}{\alpha})}.$$

This completes the proof of Theorem 3. □

5 Local and Global Existence of the Solutions to the Boussinesq System

The results of this section closely mirror Sect. 4. Consequently, we omit many of the arguments, when they are virtually the same. There are however a few important distinctions, which we will highlight herein.

5.1 Global Regularity for the Vorticity (ω, θ) Boussinesq System in $L^p(\mathbf{R}^2)$

Our first result is, non-surprisingly, a local existence and uniqueness result in $L^p(\mathbf{R}^2)$. Most of the claims in this Lemma are either well-known or follow classical arguments, but we provide a sketch of the proof for completeness.

Lemma 12 *Suppose that $\omega_0, \theta_0 \in L^p$, $1 \leq p \leq \infty$. Then there exists $T = T(\|(\omega_0, \theta_0)\|_{L^1 \cap L^\infty})$, such that unique strong solutions $\omega, \theta \in C([0, T); L^1 \cap L^\infty)$ exist.*

Moreover, the solutions $\omega(t), \theta(t)$ exist globally. In addition, the function $t \rightarrow \|\theta(t, \cdot)\|_{L^p}$, $1 \leq p \leq \infty$ is non-increasing, $\|\theta(t, \cdot)\|_{L^p} \leq \|\theta_0\|_{L^p}$, $1 < p < \infty$, while

$$\|u(t, \cdot)\|_{L^2} \leq \|u_0\|_{L^2} + t \|\theta_0\|_{L^2}.$$

Proof For the local existence, we work in the space $X = L^1 \cap L^\infty = \cap L^p$. The strong solutions of the system of Eq. (1.7) are solutions of the integral equations

$$\begin{cases} \omega(\xi, t) = e^{-t|\nabla|^\alpha} \omega_0 + \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u \cdot \omega) ds - \int_0^t e^{-(t-s)|\nabla|^\alpha} \partial_1 \theta ds, \\ \theta(\xi, t) = e^{-t|\nabla|^\alpha} \theta_0 + \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u \cdot \theta) ds. \end{cases} \quad (5.1)$$

By (2.16), we have that

$$\|e^{-t|\nabla|^\alpha} \omega_0\|_X + \|e^{-t|\nabla|^\alpha} \theta_0\|_X \leq C(\|\omega_0\|_X + \|\theta_0\|_X)$$

One can now consider the space $Y := \{(\omega, \theta) : \sup_{0 \leq t \leq T} [\|\omega\|_X + \|\theta\|_X] \leq 2C(\|\omega_0\|_X + \|\theta_0\|_X)\}$. For the bilinear forms

$$Q_1(\omega_1, \omega_2) = \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u \cdot \omega_2) ds, \quad Q_2(\omega_1, \theta) = \int_0^t e^{-(t-s)|\nabla|^\alpha} \nabla(u \cdot \theta) ds$$

where $u = (\nabla^\perp)^{-1} \omega_1$, we establish the estimates

$$\begin{aligned} \|Q_1(\omega_1, \omega_2) - Q_1(\tilde{\omega}_1, \tilde{\omega}_2)\|_X &\leq CT^{1-\frac{1}{\alpha}} (\|(\omega_1, \omega_2)\|_X \\ &\quad + \|(\tilde{\omega}_1, \tilde{\omega}_2)\|_X) (\|\omega_1 - \tilde{\omega}_1\|_X + \|\omega_2 - \tilde{\omega}_2\|_X) \\ \|Q_2(\omega_1, \theta) - Q_2(\tilde{\omega}_1, \tilde{\theta})\|_X &\leq CT^{1-\frac{1}{\alpha}} (\|(\omega_1, \theta)\|_X \\ &\quad + \|(\tilde{\omega}_1, \tilde{\theta})\|_X) (\|\omega_1 - \tilde{\omega}_1\|_X + \|\theta - \tilde{\theta}\|_X) \end{aligned}$$

for $j = 1, 2$. This is done in an identical manner as in the proof of Lemma 8. It remains to deal with the integral term $\int_0^t e^{-(t-s)|\nabla|^\alpha} \partial_1 \theta \, ds$, for which we have

$$\begin{aligned} & \left\| \int_0^t e^{-(t-s)|\nabla|^\alpha} \partial_1 (\theta - \tilde{\theta}) \, ds \right\|_{L^1 \cap L^\infty} \leq C \int_0^t \frac{1}{(\tau - s)^{\frac{1}{\alpha}}} \|\theta - \tilde{\theta}\|_{L^1 \cap L^\infty} \, ds \\ & \leq CT^{1-\frac{1}{\alpha}} \sup_{0 < s < T} \|\theta(s) - \tilde{\theta}(s)\|_{L^1 \cap L^\infty}, \end{aligned}$$

for $0 < t < T$. All in all, we can guarantee that with an appropriate choice of T , the nonlinear map given by (5.1) has a fixed point ω, θ in the space X .

Regarding the global well-posedness, we can continue the solution, as long as the norm $t \rightarrow \|\theta(t, \cdot)\|_{L^p}$ stay under control. First, for $1 < p < \infty$, take dot product of the θ equation with $|\theta|^{p-2}\theta$, $p \in (1, \infty)$ and using the fact the positivity estimate (2.12), we obtain

$$\frac{1}{p} \partial_t \|\theta(t, \cdot)\|_{L^p}^p \leq \frac{1}{p} \partial_t \|\theta\|_{L^p}^p + \int_{\mathbf{R}^2} |\theta|^{p-2} \theta \cdot |\nabla|^\alpha \theta \, dx = 0$$

It follows that $t \rightarrow \|\theta(t, \cdot)\|_{L^p}$ is non-increasing in any interval $(0, t)$, whence the solution is global and $\|\theta(t, \cdot)\|_{L^p} \leq \|\theta_0\|_{L^p}$. For $p = 1, p = \infty$, we use approximation arguments to establish the same result.

Finally, we use this information to establish the global well-posedness of the u equation in (1.6). Taking dot product with u , we obtain

$$\begin{aligned} \frac{1}{2} \partial_t \|u(t, \cdot)\|_{L^2}^2 & \leq \frac{1}{2} \partial_t \|u(t, \cdot)\|_{L^2}^2 + \||\nabla|^{\frac{\alpha}{2}} u\|_{L^2}^2 \\ & = \langle u_2, \theta \rangle \leq \|u_2\|_{L^2} \|\theta(t)\|_{L^2} \leq \|u_2(t)\|_{L^2} \|\theta_0\|_{L^2} \end{aligned}$$

It follows that

$$\|u(t, \cdot)\|_{L^2} \leq \|u_0\|_{L^2} + t \|\theta_0\|_{L^2},$$

which provides the necessary bound to conclude global regularity, as stated. \square

The next Lemma provides a global existence and uniqueness result for the (ω, θ) system.

Lemma 13 *Let $\alpha > 1$. Then, assuming $\omega_0 \in L^2, \theta_0 \in H^{\frac{\alpha}{2}}$, the Cauchy problem (1.7) has unique global solutions. In addition, for any $T > 0$, there exists $C = C_{T, \|\omega_0\|_{L^2}, \|\theta_0\|_{H^{\frac{\alpha}{2}}}} > 0$, so that the solutions satisfy*

$$\sup_{0 \leq t \leq T} \|\omega\|_{L^2} + \sup_{0 \leq t \leq T} \||\nabla|^{\frac{\alpha}{2}} \theta\|_{L^2} \leq C. \quad (5.2)$$

Remark The constant C_T obtained in this argument is exponential in T , which is very non-efficient. On the other hand, it is sufficient for our purposes in bootstrapping the solution.

Proof The global regularity for (1.7) is of course very similar to the global regularity established in Lemma 12. For the energy estimates, needed for (5.2), we can dot product the first equation in (1.7) with ω and the second one with $|\nabla|^\alpha \theta$ to get the following energy estimate

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\omega\|_{L^2}^2 + \||\nabla|^{\frac{\alpha}{2}} \theta\|_{L^2}^2 \right) + \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2 + \||\nabla|^\alpha \theta\|_{L^2}^2 \\ & \leq \left| \int \omega \cdot \partial_1 \theta d\xi \right| + \left| \langle [|\nabla|^{\frac{\alpha}{2}}, u \cdot \nabla] \theta, |\nabla|^{\frac{\alpha}{2}} \theta \rangle \right| \\ & := I_1 + I_2. \end{aligned}$$

Then for some $0 < \gamma < 1$,

$$\begin{aligned} I_1 &= \left| \int \omega \cdot \partial_1 \theta d\xi \right| \leq \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2} \|\partial_1 |\nabla|^{-\frac{\alpha}{2}} \theta\|_{L^2} \leq \epsilon \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2 + C_\epsilon \|\partial_1 |\nabla|^{-\frac{\alpha}{2}} \theta\|_{L^2}^2 \\ &\leq \epsilon \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2 + C_\epsilon \||\nabla|^\alpha \theta\|_{L^2}^{2\gamma} \|\theta\|_{L^2}^{2(1-\gamma)} \\ &\leq \epsilon \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2 + \epsilon \||\nabla|^\alpha \theta\|_{L^2}^2 + C_\epsilon \|\theta_0\|_{L^2}^2. \end{aligned}$$

We also have

$$I_2 = \left| \langle [|\nabla|^{\frac{\alpha}{2}}, u \cdot \nabla] \theta, |\nabla|^{\frac{\alpha}{2}} \theta \rangle \right| \leq \||\nabla|^{-\frac{\alpha}{2}} [|\nabla|^{\frac{\alpha}{2}}, u \cdot \nabla] \theta\|_{L^2} \||\nabla|^\alpha \theta\|_{L^2}$$

We can make use of the inequality (2.18) with $a = 1$, $s_1 = s_2 = \frac{\alpha}{2}$, $p = 2$, $q = \frac{8}{4-\alpha}$ and $r = \frac{8}{\alpha}$ to get

$$\begin{aligned} \||\nabla|^{-\frac{\alpha}{2}} [|\nabla|^{\frac{\alpha}{2}}, u \cdot \nabla] \theta(t)\|_{L^2} &\leq C \|\theta\|_{L^{\frac{8}{\alpha}}} \|\nabla u\|_{L^{\frac{8}{4-\alpha}}} \\ &\leq C \|\theta_0\|_{L^{\frac{8}{\alpha}}} \|\omega\|_{L^{\frac{8}{4-\alpha}}} \leq C \|\theta_0\|_{L^{\frac{8}{\alpha}}} \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2} \\ &\leq C \|\theta_0\|_{L^{\frac{8}{\alpha}}} \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^{\frac{1}{2}} \|\omega\|_{L^2}^{\frac{1}{2}}. \end{aligned}$$

where we have used the Sobolev embedding and the Gagliardo–Nirenberg’s inequality. Then,

$$I_2 \leq \epsilon \||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2 + \epsilon \||\nabla|^\alpha \theta\|_{L^2}^2 + C_\epsilon (\|\theta_0\|_{L^{\frac{8}{\alpha}}} \|\omega\|_{L^2}^{\frac{1}{2}})^4.$$

Therefore, for $\epsilon < \frac{1}{2}$, we can hide the terms $\||\nabla|^{\frac{\alpha}{2}} \omega\|_{L^2}^2$ and $\||\nabla|^\alpha \theta\|_{L^2}^2$ and we obtain

$$\frac{d}{dt} \left(\|\omega\|_{L^2}^2 + \||\nabla|^{\frac{\alpha}{2}} \theta\|_{L^2}^2 \right) \leq C \|\theta_0\|_{L^{\frac{8}{\alpha}}}^4 \left(\|\omega\|_{L^2}^2 + \||\nabla|^{\frac{\alpha}{2}} \theta\|_{L^2}^2 \right) + C \|\theta_0\|_{L^2}^2.$$

We use Gronwall’s to conclude (5.2). \square

5.2 Some a Priori Estimates for the Scaled Vorticity Boussinesq Problem (W, Θ) in L^p

We now turn our attention to the scaled vorticity system. By the results of Lemmas 13 and 14, such solutions exist globally, by virtue of the change of variables. Now that we have a global solution, together with the global estimate (5.6), we can actually obtain global *a priori* estimates for Θ in all L^p spaces.

Lemma 14 *Let $p \geq 1$, and $\Theta_0 \in L^1 \cap L^\infty(\mathbf{R}^2) \cap H^\alpha(\mathbf{R}^2)$, $W_0 \in L^2$. Then for any $\tau > 0$, $\Theta \in C^0([0, \tau]; L^p)$, there exists $C = C_{\alpha, p}$ such that*

$$\|\Theta(\tau, \cdot)\|_{L^p} \leq C_{\alpha, p} \|\Theta_0\|_{L^p(\mathbf{R}^2)} e^{(2 - \frac{1}{\alpha} - \frac{2}{\alpha p})\tau}. \quad (5.3)$$

Proof We take a dot product of the Θ equation in (2.9) with $|\Theta|^{p-2}\Theta$, $p \geq 1$. We obtain

$$\frac{1}{p} \partial_\tau \|\Theta\|_{L^p}^p + \int_{\mathbf{R}^2} |\nabla|^\alpha \Theta| \Theta|^{p-2} \Theta d\xi = \left(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}\right) \|\Theta\|_{L^p}^p.$$

Recall however that $\int_{\mathbf{R}^2} |\nabla|^\alpha \Theta| \Theta|^{p-2} \Theta d\xi \geq 0$, by Lemma 1. Thus, integrating this inequality yields (5.3). \square

Lemma 14 provides us with a decay rate for $\Theta(\tau, \cdot)$ for $1 \leq p < \frac{2}{2\alpha-1}$, but clearly an increasing exponential bound for $p \geq \frac{2}{2\alpha-1}$. However, we can use it to get a decay rate for any $p \geq 1$.

Lemma 15 *Let $p \geq 1$, and $\Theta_0 \in L^1 \cap L^\infty(\mathbf{R}^2) \cap H^\alpha(\mathbf{R}^2)$, $W_0 \in L^2$. Then for any $\tau > 0$, $\Theta \in C^0([0, \tau]; L^p)$, there exists $C = C_{\alpha, p}$ such that*

$$\|\Theta(\tau, \cdot)\|_{L^p} \leq C_{\alpha, p} \|\Theta_0\|_{L^p(\mathbf{R}^2)} e^{(2 - \frac{3}{\alpha})\tau}. \quad (5.4)$$

Proof Similar to Lemma 10, we have the following energy estimate

$$\frac{1}{p} \partial_\tau \|\Theta\|_{L^p}^p + c_{p, \alpha} \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^p \leq (2 - \frac{1}{\alpha} - \frac{2}{\alpha p}) \|\Theta\|_{L^p}^p.$$

In other words

$$\begin{aligned} \partial_\tau \|\Theta\|_{L^p}^p + p c_{p, \alpha} \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^p &\leq p(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}) \|\Theta\|_{L^p}^p \\ &\leq \left(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}\right) \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^{\gamma p} \|\Theta\|_{L^1}^{(1-\gamma)p} \\ &\leq \frac{\left[p \left(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}\right)\right]^{\frac{1}{1-\gamma}}}{\epsilon^{\frac{\gamma}{1-\gamma}}} \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^{\gamma p} \|\Theta\|_{L^1}^{(1-\gamma)p} \\ &\leq p \left(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}\right) \|\Theta\|_{L^1}^p + \epsilon \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^p. \end{aligned}$$

Now we use (5.3) with $p = 1$ to get the following energy estimate

$$\partial_\tau \|\Theta\|_{L^p}^p + (pc_{p,\alpha} - \epsilon) \|\Theta\|_{L^{\frac{2p}{2-\alpha}}}^p \leq p(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}) \|\Theta\|_{L^1}^p \leq p(2 - \frac{1}{\alpha} - \frac{2}{\alpha p}) e^{p(2 - \frac{3}{\alpha})\tau}.$$

Finally, we use Gronwall's inequality to finish the proof. \square

We can use above Lemma to find some decay rate for $U(\tau, \cdot)$. We need this to be able to get some bounds for W in higher L^p spaces.

Lemma 16 *Let $U_0 \in L^2(\mathbf{R}^2)$. There exists $C = C_{\alpha,p}$, such that for any $\tau > 0$, $U \in C^0([0, \tau]; L^2)$ and*

$$\|U(\tau, \cdot)\|_{L^2} \leq C_{\alpha,p} \|U_0\|_{L^2(\mathbf{R}^2)} e^{\left(2 - \frac{3}{\alpha}\right)\tau}. \quad (5.5)$$

Proof If we dot product the equation (2.10) with U , we get the following relation

$$\frac{1}{2} \partial_\tau \|U\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}} U\|_{L^2}^2 = \frac{1}{\alpha} \int (\xi \cdot \nabla U) U d\xi + (1 - \frac{1}{\alpha}) \|U\|_{L^2}^2 + \int \Theta \cdot U d\xi.$$

Then

$$\begin{aligned} \partial_\tau \|U\|_{L^2}^2 + 2 \|\nabla|^{\frac{\alpha}{2}} U\|_{L^2}^2 &= 2 \left(1 - \frac{2}{\alpha}\right) \|U\|_{L^2}^2 + \int \Theta \cdot U d\xi \\ &\leq 2 \left(1 - \frac{2}{\alpha}\right) \|U\|_{L^2}^2 + \|\Theta\|_{L^2} \|U\|_{L^2} \\ &\leq 2 \left(1 - \frac{2}{\alpha} + \epsilon\right) \|U\|_{L^2}^2 + C_\epsilon \|\Theta\|_{L^2}^2 \\ &\leq 2 \left(1 - \frac{2}{\alpha} + \epsilon\right) \|U\|_{L^2}^2 + C_\epsilon e^{2(2 - \frac{3}{\alpha})\tau}. \end{aligned}$$

We finish the proof by the Gronwall's inequality. \square

The next lemma provides *a priori* estimates for W and Θ in L^2 spaces, which allows us to conclude global regularity.

Lemma 17 *Let $\alpha \in (1, \frac{3}{2})$, $W_0 \in L^2$. Then the solution W of (2.9), satisfies*

$$\|W(\tau, \cdot)\|_{L^2} + \|\Theta(\tau, \cdot)\|_{L^2} \leq C e^{(2 - \frac{3}{\alpha})\tau}, \quad (5.6)$$

$$\sup_{0 \leq \tau < \infty} \int_0^\tau \left(\|\nabla|^{\frac{\alpha}{2}} W(s)\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}} \Theta(s)\|_{L^2}^2 \right) ds \leq C \quad (5.7)$$

for some $C = C(\|W_0\|_{L^2}, \|\Theta_0\|_{L^2}, \alpha)$, independent on τ .

Proof We dot product the first equation in (2.9) with W , and the second equation with Θ . We also use the trick from Lemma 10—we add the term $A(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2)$, where A is a large constant to be determined. Then

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2 \right) + A(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2) + \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 \\ & \leq \left| \int \partial_1 \Theta W d\xi \right| + \left(A + 1 - \frac{1}{\alpha} \right) \|W\|_{L^2}^2 + \left(A + 2 - \frac{2}{\alpha} \right) \|\Theta\|_{L^2}^2. \end{aligned}$$

But by Gagliardo–Nirenberg (and taking into account that $1 - \frac{\alpha}{2} < \frac{\alpha}{2}$) and Young's inequalities,

$$\begin{aligned} \left| \int \partial_1 \Theta W d\xi \right| & \leq \|\nabla|^{1-\frac{\alpha}{2}} \Theta\|_{L^2} \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2} \leq \epsilon \|\nabla|^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 + \epsilon \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + C_\epsilon \|\Theta\|_{L^2}^2 \\ & \leq \epsilon \|\nabla|^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 + \epsilon \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + C_\epsilon e^{2(2-\frac{3}{\alpha})\tau}. \end{aligned}$$

where we have used the estimate for $\|\Theta\|_{L^2}$ from (5.3), with $p = 2$. We also have

$$\begin{aligned} \left(A + 1 - \frac{1}{\alpha} \right) \|W\|_{L^2}^2 & \leq C \left(A + 1 - \frac{1}{\alpha} \right) \|\nabla U\|_{L^2}^2 \leq C \left(A + 1 - \frac{1}{\alpha} \right) \|U\|_{L^2}^{2\gamma} \|\nabla|^{1+\frac{\alpha}{2}} U\|_{L^2}^{2(1-\gamma)} \\ & \leq C \left(A + 1 - \frac{1}{\alpha} \right) \|U\|_{L^2}^{2\gamma} \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^{2(1-\gamma)} \leq \epsilon \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 \\ & \quad + \frac{[C(A+1-\frac{1}{\alpha})]^{\frac{1}{1-\gamma}}}{\epsilon^{\frac{\gamma}{1-\gamma}}} \|U\|_{L^2}^2 \\ & \leq \epsilon \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + \frac{[C(A+1-\frac{1}{\alpha})]^{\frac{1}{1-\gamma}}}{\epsilon^{\frac{\gamma}{1-\gamma}}} e^{2(2-\frac{3}{\alpha})\tau}. \end{aligned}$$

Considering the estimate for $\|\Theta\|_{L^2}$ from (5.3) (with $p = 2$)

$$\begin{aligned} & \frac{d}{dt} \left(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2 \right) + 2A(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2) + 2(1-2\epsilon) \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 \\ & \quad + 2(1-2\epsilon) \|\nabla|^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 \\ & \leq \frac{2[C(A+1-\frac{1}{\alpha})]^{\frac{1}{1-\gamma}}}{\epsilon^{\frac{\gamma}{1-\gamma}}} e^{2(2-\frac{3}{\alpha})\tau}. \end{aligned}$$

We choose $A = 2(\frac{3}{\alpha} - 2)$ (recall $\alpha < \frac{3}{2}$). Then the last relation has two consequences. First we can drop the term $2(1-2\epsilon) \|\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + 2(1-2\epsilon) \|\nabla|^{\frac{\alpha}{2}} \Theta\|_{L^2}^2$, so

$$\frac{d}{dt} \left(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2 \right) + 4 \left(\frac{3}{\alpha} - 2 \right) \left(\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2 \right) \leq \frac{[C(\frac{5}{\alpha} - 3)]^{\frac{1}{1-\gamma}}}{\epsilon^{\frac{\gamma}{1-\gamma}}} e^{2(2-\frac{3}{\alpha})\tau}.$$

and then use the Gronwall's inequality for the following inequality and get the decay rate (5.6). The second consequence is that we get

$$\int_0^\tau (\|\nabla|^{\frac{\alpha}{2}} W(t)\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}} \Theta(t)\|_{L^2}^2) dt \leq \left(\|W_0\|_{L^2}^2 + \|\Theta_0\|_{L^2}^2 \right) + \frac{C_\epsilon}{2(\frac{3}{\alpha} - 2)}.$$

This implies (5.7). \square

We shall need some *a priori* estimates for $\|W\|_{L^p}$ for some $p > 2$, as these will be necessary in our subsequent considerations. This turns out to be non-trivial. It turns out that it is easier to control $\|W\|_{H^1}$, $\|\Theta\|_{H^1}$ and then use Sobolev embedding to control $\|W\|_{L^p}$, $\|\Theta\|_{L^p}$, $1 < p < \infty$. In this way, we get the control needed, but we end up needing to require smoother H^1 initial data.

Proposition 5 $W_0, \Theta_0 \in H^1$. Then, the global solution satisfies $W, \Theta \in C^0([0, \tau]; H^1(\mathbf{R}^2))$. Moreover,

$$\|W(\tau)\|_{H^1} + \|\Theta(\tau)\|_{H^1} \leq C e^{(2 - \frac{3}{\alpha})\tau}. \quad (5.8)$$

$C = C(\|W_0\|_{H^1}, \|\Theta_0\|_{H^1}, \alpha)$, independent on τ .

Proof Local well-posedness in the space H^1 , for the original (unscaled) equations works as in Lemma 13, so we omit it. Thus, we have local solutions for the scaled system as well. We now need to establish *a priori* estimates to show that these are global.

We differentiate each of the equations in (2.9). Then, we dot product it with¹⁰ ∂W and $\partial \Theta$ respectively. We add the two resulting equations to obtain the following energy inequality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\|\partial W\|_{L^2}^2 + \|\partial \Theta\|_{L^2}^2 \right) + \|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2}^2 \leq \\ & \leq \left| \int \partial_1 \partial \Theta \partial W d\xi \right| + \left(1 - \frac{1}{\alpha} \right) \|\partial W\|_{L^2}^2 + 2 \left(1 - \frac{1}{\alpha} \right) \|\partial \Theta\|_{L^2}^2 + |\langle \partial U \nabla W, \partial W \rangle| \\ & \quad + |\langle \partial U \nabla \Theta, \partial \Theta \rangle|. \end{aligned}$$

By Gagliardo–Nirenbergs' and Young's

$$\|\partial W\|_{L^2}^2 + \|\partial \Theta\|_{L^2}^2 \leq \epsilon (\|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2}^2) + C_\epsilon (\|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2)$$

Next,

$$\begin{aligned} \left| \int \partial_1 \partial \Theta \partial W d\xi \right| & \leq C \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2} \|\nabla|^{2-\frac{\alpha}{2}} W\|_{L^2} \\ & \leq \epsilon (\|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2}^2) + C_\epsilon \|W\|_{L^2}^2, \end{aligned}$$

¹⁰ Here ∂ means either ∂_1 or ∂_2 .

where in the last estimate we have used that $2 - \frac{\alpha}{2} < 1 + \frac{\alpha}{2}$. Finally,

$$\begin{aligned} |\langle \partial U \cdot \nabla W, \partial W \rangle| &= |\langle \nabla \cdot (\partial U W), \partial W \rangle| \leq C \|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2} \|\nabla|^{1-\frac{\alpha}{2}} (\partial U W)\|_{L^2} \\ &\leq \epsilon \|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + C_\epsilon \|\nabla|^{1-\frac{\alpha}{2}} (\partial U W)\|_{L^2}^2 \end{aligned}$$

By product estimates, lemma (4) and Sobolev embedding

$$\begin{aligned} \|\nabla|^{1-\frac{\alpha}{2}} (\partial U W)\|_{L^2} &\leq C(\|\nabla|^{1-\frac{\alpha}{2}} \partial U\|_{L^{\frac{8}{4-\alpha}}} \|W\|_{L^{\frac{8}{\alpha}}} + \|\nabla|^{1-\frac{\alpha}{2}} W\|_{L^{\frac{8}{4-\alpha}}} \|\partial U\|_{L^{\frac{8}{\alpha}}}) \\ &\leq C \|\nabla|^{1-\frac{\alpha}{4}} \partial U\|_{L^2} \|\nabla|^{1-\frac{\alpha}{4}} W\|_{L^2} \leq C \|\nabla|^{1-\frac{\alpha}{4}} W\|_{L^2}^2 \leq \|\nabla|^{1+\frac{\alpha}{2}} W\|_{L^2}^{\frac{2-\frac{\alpha}{2}}{1+\frac{\alpha}{2}}} \|W\|_{L^2}^{\frac{3\alpha}{1+\frac{\alpha}{2}}}, \end{aligned}$$

where we have used $\partial U \sim W$ (in all Sobolev spaces) and Gagliardo–Nirenberg's. This allows us to estimate by Young's

$$|\langle \partial U \cdot \nabla W, \partial W \rangle| \leq 2\epsilon \|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + C_\epsilon \|W\|_{L^2}^{\frac{3\alpha}{\alpha-1}}.$$

Clearly, the appropriate estimate, obtained in the same way holds for

$$|\langle \partial U \nabla \Theta, \partial \Theta \rangle| \leq 2\epsilon \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2}^2 + C_\epsilon \|W\|_{L^2}^{\frac{3\alpha}{\alpha-1}}.$$

All in all, we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|\partial W\|_{L^2}^2 + \|\partial \Theta\|_{L^2}^2 \right) &+ (1 - 6\epsilon)(\|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 + \|\nabla|^{\frac{\alpha}{2}+1} \Theta\|_{L^2}^2) \\ &\leq C_\epsilon (\|W\|_{L^2}^{\frac{3\alpha}{\alpha-1}} + \|W\|_{L^2}^2 + \|\Theta\|_{L^2}^2). \end{aligned}$$

Set $\epsilon = \frac{1}{10}$. For every $A > 0$, there is $c_{\alpha, A}$, so that $\|\nabla|^{\frac{\alpha}{2}+1} W\|_{L^2}^2 \geq A \|\partial W\|_{L^2}^2 - c_{A, \alpha} \|W\|_{L^2}^2$ and similar for Θ , so we end up with

$$\frac{d}{dt} \left(\|\partial W\|_{L^2}^2 + \|\partial \Theta\|_{L^2}^2 \right) + A \left(\|\partial W\|_{L^2}^2 + \|\partial \Theta\|_{L^2}^2 \right) \leq C_{A, \alpha} e^{2(2-\frac{3}{\alpha})\tau}.$$

where we have used the exponential bounds from (5.6). Setting sufficiently large A , namely $A = 2(\frac{3}{\alpha} - 2)$, and applying Gronwall's yields the result. \square

As an immediate corollary, we have control of the L^p norms for W .

Corollary 1 *Let $W_0, \Theta_0 \in H^1$. Then, for all $p \in (2, \infty)$, there is the bound*

$$\|W(\tau, \cdot)\|_{L^p} \leq C(\|W_0\|_{H^1}, \|\Theta_0\|_{H^1}, \alpha, p) e^{(2-\frac{3}{\alpha})\tau}. \quad (5.9)$$

5.3 Global Regularity for the Scaled Vorticity Boussinesq Problem (W, Θ) in $L^2(2) \cap L^\infty(\mathbb{R}^2)$

Let us mention that the results in this section are valid under the assumption that $\alpha \in (1, 2)$. The more restrictive assumption $\alpha \in (1, \frac{3}{2})$ comes up later in Theorem 5, in the more subtle estimates of the evolution of the solution, projected on various pieces of the spectrum.

The next Lemma is a local well-posedness result, which is a companion to Theorem 3.

Lemma 18 *Suppose that $W_0, \Theta_0 \in L^2(2) \cap L^\infty$. Then, there exists time $T = T(\|(W_0, \Theta_0)\|_{L^2(2) \cap L^\infty})$, so that the system of equation (2.9) has unique local solution $W, \Theta \in C^0([0, T]; L^2(2) \cap L^\infty)$ with $W(0) = W_0$ and $\Theta(0) = \Theta_0$.*

Proof We are looking for strong solutions in the space $X = L^2(2) \cap L^\infty$, that is a solutions of the following system of integral equations

$$\begin{aligned} W(\tau) &= e^{\tau \mathcal{L}} W_0 - \int_0^\tau e^{(\tau-s)\mathcal{L}} \nabla(U \cdot W) ds + \int_0^\tau e^{(\tau-s)\mathcal{L}} (\partial_1 \Theta) ds, \\ \Theta(\tau) &= e^{\tau(\mathcal{L}+1-\frac{1}{\alpha})} \Theta_0 - \int_0^\tau e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \nabla(U \cdot \Theta) ds \end{aligned}$$

For the free solutions, according to (3.15) and (3.8),

$$\|e^{\tau \mathcal{L}} W_0\|_{L^2(2) \cap L^\infty} + \|e^{\tau(\mathcal{L}+1-\frac{1}{\alpha})} \Theta_0\|_{L^2(2) \cap L^\infty} \leq C e^\tau (\|W_0\|_{L^2(2) \cap L^\infty} + \|\Theta_0\|_{L^2(2) \cap L^\infty}).$$

For $0 < T < 1$, to be determined, introduce the space

$$Y_T := \{(W, \Theta) : \sup_{0 \leq \tau \leq T} [\|W(\tau, \cdot)\|_X + \|\Theta(\tau, \cdot)\|_X] \leq 2C e(\|W_0\|_{L^2(2) \cap L^\infty} + \|\Theta_0\|_{L^2(2) \cap L^\infty})\}.$$

According to (3.10) and (3.15),

$$\begin{aligned} \left\| \int_0^\tau e^{(\tau-s)\mathcal{L}} \nabla(U \cdot W) ds \right\|_{L^2(2) \cap L^\infty} &\leq \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}} (e^{(1-\frac{2}{\alpha})(\tau-s)} + e^{\tau-s})}{a(\tau-s)^{\frac{1}{\alpha}}} \|U \cdot W\|_{L^2(2) \cap L^\infty} ds \\ &\leq C \sup_{0 \leq \tau \leq T} \|U W\|_{L^2(2) \cap L^\infty} \int_0^\tau \frac{1}{|\tau-s|^{\frac{1}{\alpha}}} ds \leq C T^{1-\frac{1}{\alpha}} \sup_{0 \leq \tau \leq T} \|U\|_{L^\infty} \sup_{0 \leq \tau \leq T} \|W\|_{L^2(2) \cap L^\infty}. \end{aligned}$$

and similarly

$$\left\| \int_0^\tau e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \nabla(U \cdot \Theta) ds \right\|_{L^2(2) \cap L^\infty} \leq C T^{1-\frac{1}{\alpha}} \sup_{0 \leq \tau \leq T} \|U\|_{L^\infty} \sup_{0 \leq \tau \leq T} \|\Theta\|_{L^2(2) \cap L^\infty}.$$

Recalling that $U = (\nabla^\perp)^{-1} W$, we further estimate by (2.2),

$$\|U\|_{L^\infty} \leq C(\|W\|_{L^{2+\epsilon}} + \|W\|_{L^{2-\epsilon}}) \leq C\|W\|_{L^2(2) \cap L^\infty},$$

since $L^2(2) \hookrightarrow L^{2-\epsilon}$ and $L^2(2) \cap L^\infty \hookrightarrow L^1 \cap L^\infty \hookrightarrow L^{2+\epsilon}$. Finally,

$$\left\| \int_0^\tau e^{(\tau-s)\mathcal{L}}(\partial_1 \Theta) ds \right\|_{L^2(2) \cap L^\infty} \leq CT^{1-\frac{1}{\alpha}} \sup_{0 \leq \tau \leq T} \|\Theta\|_{L^2(2) \cap L^\infty}.$$

Clearly, appropriate estimate hold for the differences, whence the integral equations provide a contraction mapping in the space Y_T , provided, $T^{1-\frac{1}{\alpha}} << \frac{1}{2Ce(\|W_0\|_{L^2(2) \cap L^\infty} + \|\Theta_0\|_{L^2(2) \cap L^\infty})}$. \square

Our next result provides a global regularity for the W, Θ system in the space $L^2(2)$.

Lemma 19 *The system of equations (2.9), with $W_0, \Theta_0 \in X = L^2(2) \cap L^\infty$, and also $W_0, \Theta_0 \in H^1(\mathbf{R}^2)$ has an unique global solution, in space X . There exists $C = C(\|W_0\|_X, \|\Theta\|_X)$ such that*

$$\sup_{0 \leq \tau < \infty} (\|W(\tau, \cdot)\|_{L^2(2)} + \|\Theta(\tau, \cdot)\|_{L^2(2)}) \leq C. \quad (5.10)$$

Remark The decay rate by a constant is very inefficient. One could improve the argument below, at a considerable technical price, to obtain better decay estimates. Since the results in Sect. 7 will supersede these anyway, we choose to present the simpler arguments.

Proof The existence of a local solutions are guaranteed by Lemma 18. So, it remains to establish energy estimates, which keep the relevant $L^2(2)$ norms under control. Note that the unweighted portion of the norm has an exponential decay, by (5.3) and (5.6). So, it remains to control the weighted norms.

We run a preliminary argument only on the Θ variable. As usual, this is easier, due to the lack of problematic term $\partial_1 \Theta$, which appears in the equation for W . We dot product the Θ equation in (2.9) with $|\xi|^4 \Theta$. We have

$$\begin{aligned} & \frac{1}{2} \frac{d}{d\tau} \int |\xi|^4 \Theta^2 d\xi + \int |\xi|^4 |\nabla|^\alpha \Theta \cdot \Theta d\xi + \left(\frac{4}{\alpha} - 2 \right) \int |\xi|^4 \Theta^2 d\xi \\ &= - \int (U \cdot \nabla_\xi \Theta) |\xi|^4 \Theta d\xi. \end{aligned}$$

Then

$$- \int (U \cdot \nabla_\xi \Theta) |\xi|^4 \Theta d\xi = 2 \int |\xi|^2 (\xi \cdot U) \Theta^2 d\xi.$$

But

$$\left| \int |\xi|^2 (\xi \cdot U) \Theta^2 d\xi \right| \leq C \int |\xi|^3 \|U\|_{L^\infty} |\Theta|^2 d\xi \leq \epsilon \int |\xi|^4 |\Theta|^2 d\xi + C\epsilon^{-3} \|U\|_{L^\infty}^4 \|\Theta\|_{L^2}^2.$$

Now, according to (2.2), for every $\delta > 0$

$$\begin{aligned} \|U\|_{L^\infty} &\leq C_\delta(\|W\|_{L^{2+\delta}} + \|W\|_{L^{2-\delta}}) \leq C_\delta(e^{(2-\frac{3}{\alpha})\tau} + \|W\|_{L^2}^{\frac{2-2\delta}{2-\delta}} \|W\|_{L^1}^{\frac{\delta}{2-\delta}}) \\ &\leq C_\delta + C_\delta \|W\|_{L^2(2)}^{\frac{\delta}{2-\delta}}. \end{aligned}$$

We also have

$$\begin{aligned} \int |\xi|^4 \Theta |\nabla|^\alpha \Theta d\xi &= \langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} |\nabla|^{\frac{\alpha}{2}} \Theta, |\xi|^2 \Theta \rangle \\ &= \langle |\nabla|^{\frac{\alpha}{2}} [|\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle - \langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle \\ &= \langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \rangle + \langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] \Theta \rangle - \langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle \\ &= \int |\xi|^4 |\nabla|^{\frac{\alpha}{2}} \Theta|^2 d\xi + \langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] \Theta \rangle - \langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle \end{aligned}$$

Now if we define $I(\tau) = \int |\xi|^4 \Theta^2 d\xi$, and put all above together we have the following relation

$$\begin{aligned} \frac{1}{2} I'(\tau) + \left(\frac{4}{\alpha} - 2 - 10\epsilon \right) I(\tau) + \int |\xi|^4 |\nabla|^{\frac{\alpha}{2}} \Theta|^2 d\xi \\ \leq |\langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] \Theta \rangle| + |\langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle| + C_{\delta, \epsilon} \|W(\tau, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}}. \end{aligned}$$

We can use Lemma 11 to get

$$\begin{aligned} |\langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] \Theta \rangle| &\leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2} \| [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] \Theta \|_{L^2} \\ &\leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2} \| |\xi|^{2-\frac{\alpha}{2}} \Theta \|_{L^2} \leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2} \| |\xi|^2 \Theta \|_{L^2}^{1-\frac{\alpha}{4}} \| \Theta \|_{L^2}^{\frac{\alpha}{4}} \\ &\leq \epsilon (\| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^2 + \| |\xi|^2 \Theta \|_{L^2}^2) + C_\epsilon. \end{aligned}$$

For the other term we have

$$\begin{aligned} |\langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 \Theta \rangle| &\leq \| |\xi|^2 \Theta \|_{L^2} \| [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta] \|_{L^2} \\ &\leq \| |\xi|^2 \Theta \|_{L^2} \| |\xi|^{2-\frac{\alpha}{2}} [|\nabla|^{\frac{\alpha}{2}} \Theta] \|_{L^2} \leq \| |\xi|^2 \Theta \|_{L^2} \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^{1-\frac{\alpha}{4}} \| |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^{\frac{\alpha}{4}} \\ &\leq \epsilon (\| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^2 + \| |\xi|^2 \Theta \|_{L^2}^2) + C_\epsilon \| |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^2. \end{aligned}$$

It follows that

$$\begin{aligned} \frac{1}{2} I'(\tau) + \left(\frac{4}{\alpha} - 2 - 20\epsilon \right) I(\tau) + (1 - 5\epsilon) \int |\xi|^4 |\nabla|^{\frac{\alpha}{2}} \Theta|^2 d\xi \\ \leq C_\epsilon + C_{\delta, \epsilon} \|W(\tau, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}} + C_\epsilon \| |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^2. \end{aligned}$$

Choose $\epsilon = \frac{1}{200}$, apply Gronwall's and then (5.7) implies that for every $\delta > 0$, there is C_δ , so that

$$\||\xi|^2 \Theta(\tau, \cdot)\|_{L^2} \leq C_\epsilon + C_\delta e^{-(\frac{4}{\alpha} - 2 - \delta)\tau} + C_\delta \sup_{0 < s < \tau} \|W(s, \cdot)\|_{L^2(2)}^{\frac{2\delta}{2-\delta}}. \quad (5.11)$$

for every $\delta > 0$. In addition, we obtain the L_τ^2 bound

$$\int_0^\tau \||\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta(\tau, \cdot)\|_{L^2}^2 d\tau \leq C + C_\delta \sup_{0 < s < \tau} \|W(s, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}}. \quad (5.12)$$

We are now ready for the bounds for W , which are always harder. If we dot product in (2.9), the first equation with $|\xi|^4 W$, we have the energy equality

$$\begin{aligned} & \frac{1}{2} \frac{d}{d\tau} \int |\xi|^4 W^2 d\xi + \int |\xi|^4 |\nabla|^\alpha W \cdot W d\xi + \left(\frac{3}{\alpha} - 1 \right) \int |\xi|^4 W^2 d\xi \\ &= - \int (U \cdot \nabla_\xi W) |\xi|^4 W d\xi + \int \partial_1 \Theta |\xi|^4 W d\xi \end{aligned}$$

Then $-\int (U \cdot \nabla_\xi W) |\xi|^4 W d\xi = 2 \int |\xi|^2 (\xi \cdot U) W^2 d\xi$. We can bound this term as follows

$$\left| \int |\xi|^2 (\xi \cdot U) W^2 d\xi \right| \leq C \int |\xi|^3 \|U\|_{L^\infty} |W|^2 d\xi \leq \epsilon \int |\xi|^4 |W|^2 d\xi + C\epsilon^{-3} \|U\|_{L^\infty}^4 \|W\|_{L^2}^2.$$

Again, according to (2.1), for every $\delta > 0$

$$\|U\|_{L^\infty} \leq C_\delta (\|W\|_{L^{2+\delta}} + \|W\|_{L^{2-\delta}}) \leq C \left(e^{(2-\frac{3}{\alpha})\tau} + \|W\|_{L^2}^{\frac{2-2\delta}{2-\delta}} \|W\|_{L^1}^{\frac{\delta}{2-\delta}} \right).$$

Taking into account (5.3), (5.9), $L^2(2) \hookrightarrow L^1$ and Young's inequality, allows us to estimate

$$\left| \int |\xi|^2 (\xi \cdot U) W^2 d\xi \right| \leq 2\epsilon \int |\xi|^4 |W|^2 d\xi + C_{\epsilon, \delta} \|W(\tau, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}}.$$

We also have, similar to the Θ variable calculation,

$$\begin{aligned} \int |\xi|^4 W |\nabla|^\alpha W d\xi &= \||\xi|^2 |\nabla|^{\frac{\alpha}{2}} W\|_{L^2}^2 + \langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] W \rangle \\ &\quad - \langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} W], |\xi|^2 W \rangle \end{aligned}$$

Now if we take $J(\tau) = \int |\xi|^4 W^2 d\xi$, and put all above together we have the following relation

$$\begin{aligned} & \frac{1}{2} J'(\tau) + \left(\frac{3}{\alpha} - 1 - 10\epsilon \right) J(\tau) + \int |\xi|^4 |\nabla|^{\frac{\alpha}{2}} W^2 d\xi \\ & \leq |\langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] W \rangle| + |\langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} W], |\xi|^2 W \rangle| \\ & \quad + \left| \int |\xi|^4 (\partial_1 \Theta) W d\xi \right| \\ & \quad + C_\epsilon + C_{\epsilon, \delta} \|W(\tau, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}} = I_1 + I_2 + I_3 + C_\epsilon + C_{\epsilon, \delta} \|W(\tau, \cdot)\|_{L^2(2)}^{\frac{4\delta}{2-\delta}} \end{aligned}$$

We can use Lemma 11 to get

$$\begin{aligned} I_1 &= |\langle |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W, [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] W \rangle| \leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2} \| [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] W \|_{L^2} \\ &\leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2} \| |\xi|^{2-\frac{\alpha}{2}} W \|_{L^2} \leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2} \| |\xi|^2 W \|_{L^2}^{1-\frac{\alpha}{4}} \| W \|_{L^2}^{\frac{\alpha}{4}} \\ &\leq \epsilon (\| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2}^2 + \| |\xi|^2 W \|_{L^2}^2) + C_\epsilon, \end{aligned}$$

where we have used the bounds (5.6) for $\|W\|_{L^2}$. Next, regarding I_2 , we have

$$\begin{aligned} I_2 &= |\langle [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} W], |\xi|^2 W \rangle| \leq \| |\xi|^2 W \|_{L^2} \| [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} W] \|_{L^2} \\ &\leq \| |\xi|^2 W \|_{L^2} \| |\xi|^{2-\frac{\alpha}{2}} |\nabla|^{\frac{\alpha}{2}} W \|_{L^2} \leq \| |\xi|^2 W \|_{L^2} \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2}^{1-\frac{\alpha}{4}} \| |\nabla|^{\frac{\alpha}{2}} W \|_{L^2}^{\frac{\alpha}{4}} \\ &\leq \epsilon (\| |\xi|^2 W \|_{L^2}^2 + \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} W \|_{L^2}^2) + C_\epsilon \| |\nabla|^{\frac{\alpha}{2}} W \|_{L^2}^2. \end{aligned}$$

I_3 is normally a problematic term, but now we have the decay estimates for $\|\Theta\|_{L^2(2)}$, which we have proved in the first part of this Lemma. We have

$$\begin{aligned} I_3 &= \left| \langle \partial_1 \Theta, |\xi|^4 W \rangle \right| \leq \left| \langle |\xi|^2 \partial_1 \Theta, |\xi|^2 W \rangle \right| \leq \left| \langle \partial_1 |\nabla|^{-\frac{\alpha}{2}} |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, |\xi|^2 W \rangle \right| \\ &\quad + \left| \langle [\partial_1 |\nabla|^{-\frac{\alpha}{2}}, |\xi|^2] [|\nabla|^{\frac{\alpha}{2}} \Theta], |\xi|^2 W \rangle \right| := I_{3,1} + I_{3,2}. \end{aligned}$$

$I_{3,1}$ is estimated as follows

$$\begin{aligned} I_{3,1} &= \left| \langle \partial_1 |\nabla|^{-\frac{\alpha}{2}} |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta, |\xi|^2 W \rangle \right| \leq C \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2} \| |\nabla|^{1-\frac{\alpha}{2}} [|\xi|^2 W] \|_{L^2} \\ &\leq \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2} \| |\xi|^2 W \|_{L^2}^{\frac{2\alpha-2}{\alpha}} \| |\nabla|^{\frac{\alpha}{2}} [|\xi|^2 W] \|_{L^2}^{\frac{2-\alpha}{\alpha}} \leq C_\epsilon \| |\xi|^2 |\nabla|^{\frac{\alpha}{2}} \Theta \|_{L^2}^2 + \\ &\quad + \epsilon (\| |\xi|^2 W \|_{L^2}^2 + \| |\nabla|^{\frac{\alpha}{2}} [|\xi|^2 W] \|_{L^2}^2) \end{aligned}$$

We bound the last term, by Lemma 11,

$$\begin{aligned} \|\nabla^{\frac{\alpha}{2}}[|\xi|^2 W]\|_{L^2} &\leq \||\xi|^2 \nabla^{\frac{\alpha}{2}} W\|_{L^2} + \|[\nabla^{\frac{\alpha}{2}}, |\xi|^2] W\|_{L^2} \\ &\leq \||\xi|^2 \nabla^{\frac{\alpha}{2}} W\|_{L^2} + C \||\xi|^{2-\frac{\alpha}{2}} W\|_{L^2} \\ &\leq \||\xi|^2 \nabla^{\frac{\alpha}{2}} W\|_{L^2} + C(\|W\|_{L^2} + \||\xi|^2 W\|_{L^2}). \end{aligned}$$

Collecting terms together yields the following estimate for $I_{3,1}$ and using (5.8),

$$I_{3,1} \leq 2\epsilon(\||\xi|^2 W\|_{L^2}^2 + \||\xi|^2 \nabla^{\frac{\alpha}{2}} W\|_{L^2}^2) + C_\epsilon \||\xi|^2 \nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 + C e^{2(2-\frac{3}{\alpha})\tau}.$$

We can easily bound $I_{3,2}$, provided we know an appropriate estimate for the commutator $[\partial_1 \nabla^{-\frac{\alpha}{2}}, |\xi|^2]$. In fact, this commutator is morally like $[\nabla^{1-\frac{\alpha}{2}}, |\xi|^2]$, which was indeed considered in Lemma 11. However, there does not appear to be an easy way to transfer the estimate (4.8) to it, so we state the relevant estimate here

$$\|\partial_1 \nabla^{-a}, |\xi|^2\|_{L^2} \leq C \||\xi|^{1+a} f\|_{L^2}, \quad a \in (0, 1) \quad (5.13)$$

The proof of (5.13) is postponed to the “Appendix B.”¹¹ Assuming the validity of (5.13), we proceed to bound $I_{3,2}$.

$$\begin{aligned} I_{3,2} &= \left| \langle [\partial_1 \nabla^{-\frac{\alpha}{2}}, |\xi|^2] [\nabla^{\frac{\alpha}{2}} \Theta], |\xi|^2 W \rangle \right| \leq \||\xi|^2 W\|_{L^2} \|[\partial_1 \nabla^{-\frac{\alpha}{2}}, |\xi|^2] [\nabla^{\frac{\alpha}{2}} \Theta]\|_{L^2} \\ &\leq \||\xi|^2 W\|_{L^2} \||\xi|^{1+\frac{\alpha}{2}} \nabla^{\frac{\alpha}{2}} \Theta\|_{L^2} \leq \||\xi|^2 W\|_{L^2} \||\xi|^2 \nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^{\frac{2+\alpha}{4}} \|\nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^{\frac{2-\alpha}{4}} \\ &\leq \epsilon \||\xi|^2 W\|_{L^2}^2 + \|\nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 + C_\epsilon \||\xi|^2 \nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2. \end{aligned}$$

Combining all the estimates, we obtain the following energy inequality

$$\begin{aligned} \frac{1}{2} J'(\tau) &+ \left(\frac{3}{\alpha} - 1 - 20\epsilon \right) J(\tau) + (1 - 5\epsilon) \int |\xi|^4 \|\nabla^{\frac{\alpha}{2}} W\|^2 d\xi \\ &\leq C_\epsilon + C_\delta \|W\|_{L^2(2)}^{\frac{4\delta}{2-\delta}} + C_\epsilon (\||\xi|^2 \nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2 + \|\nabla^{\frac{\alpha}{2}} W\|_{L^2}^2 + \|\nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2) \end{aligned}$$

Applying Gronwall’s and taking into account the L_τ^2 integrability results (5.7) and (5.12), and $\|W\|_{L^2(2)}^2 \leq J(\tau) + C$, we conclude for every $\delta > 0$

$$\begin{aligned} J(\tau) &\leq J(0) e^{-2(\frac{3}{\alpha} - 1 - 20\epsilon)\tau} + C_\epsilon \tau e^{-2(\frac{3}{\alpha} - 1 - 20\epsilon)\tau} + C_\delta \sup_{0 < s < \tau} J(\tau)^{\frac{2\delta}{2-\delta}} + \\ &\quad + C_\epsilon \int_0^\tau (\||\xi|^2 \nabla^{\frac{\alpha}{2}} \Theta(s, \cdot)\|_{L^2}^2 + \|\nabla^{\frac{\alpha}{2}} W(s, \cdot)\|_{L^2}^2 \\ &\quad + \|\nabla^{\frac{\alpha}{2}} \Theta\|_{L^2}^2) ds \leq C_\epsilon + C_\delta \sup_{0 < s < \tau} J(\tau)^{\frac{2\delta}{2-\delta}} \end{aligned}$$

¹¹ In fact, it can be reduced to a similar expression as in the proof of (4.8), so we prove them simultaneously.

Selecting small δ and solving this inequality for $\sup_{0 < s < \tau} J(\tau)$ implies the $\sup_{0 < s < \tau} J(\tau) \leq C$, for all times τ . Inputting this last estimate in (5.11) implies the desired bound for $\|\Theta\|_{L^2(2)}$ as well. \square

6 Global Dynamics of the Solutions of the SQG Model

Theorem 3 already provides pretty good estimate about the behavior of the solutions to the rescaled equation (2.7), in particular the solution Z disperses at ∞ , with the rate of about $e^{-\tau(\frac{3-\alpha-\beta}{\alpha})}$. An obviously question is whether or not this is optimal, that is whether there is a lower bound with the same exponential function, at least for generic data. It turns out that this is indeed the case. In fact, we have a more precise result, namely an asymptotic expansion.

Before we continue with the formal statement of the main result, we need a simple algebraic observation, which is important in the sequel. Recall the generalized Biot-Savart law that we imposed, $u = u_z = (|\nabla|^{\perp})^{-\beta} z$. This naturally transformed into the relation $U = U_z = (|\nabla|^{\perp})^{-\beta} Z$ between the “scaled” velocity U and its vorticity Z . We claim that

$$U_G \cdot \nabla G = 0. \quad (6.1)$$

Indeed, since G is a radial function,¹² say $G(\xi) = \zeta(|\xi|)$, we have that $\nabla G = (\xi_1, \xi_2) \frac{\zeta'(|\xi|)}{|\xi|}$. On the other hand, $U_G = (|\nabla|^{\perp})^{-\beta} G = |\nabla|^{\perp} m_{-\beta-1}(|\nabla|)G$, so $U_G = |\nabla|^{\perp} h(|\xi|)$, where h is a radial function representing $[m_{-\beta-1}(|\nabla|)G]$. That is, $h(|\xi|) = [m_{-\beta-1}(|\nabla|)G](\xi)$. It follows that $U_G = (-\xi_2, \xi_1) \frac{h'(|\xi|)}{|\xi|}$. Thus,

$$U_G \cdot \nabla G = (-\xi_2, \xi_1) \frac{h'(|\xi|)}{|\xi|} \cdot (\xi_1, \xi_2) \frac{\zeta'(|\xi|)}{|\xi|} = 0.$$

We are now ready to state the main theorem of this section.

Theorem 4 *Let $Z_0 \in L^2(2) \cap L^\infty(\mathbf{R}^2)$, $\epsilon > 0$, $\alpha \in (1, 2)$, $\alpha + \beta \leq 3$. Denote $\gamma(0) := \int_{\mathbf{R}^2} Z_0(\xi) d\xi$. Then there exists $C_\epsilon > 0$ such that for any $\tau > 0$,*

$$\|Z(\tau, \cdot) - \gamma(0)e^{-\tau(\frac{3-\alpha-\beta}{\alpha})} G\|_{L^2(2)} \leq C_\epsilon e^{-\tau(\frac{4-\alpha-\beta}{\alpha} - \epsilon)}. \quad (6.2)$$

Assuming in addition that $\beta > 1$, we also have

$$\|\nabla[Z(\tau, \cdot) - \gamma(0)e^{-\tau(\frac{3-\alpha-\beta}{\alpha})} G]\|_{L^2(2)} \leq C_\epsilon e^{-\tau(\frac{4-\alpha-\beta}{\alpha} - \epsilon)}. \quad (6.3)$$

In particular if $\int_{\mathbf{R}^2} Z_0(\xi) d\xi = 0$, then $\|Z\|_{L^2(2)} \leq C_\epsilon e^{-\tau(\frac{4-\alpha-\beta}{\alpha} - \epsilon)}$.

¹² As the Fourier transform of a radial one.

Remark

- We would like to point out that the existence of solution Z (and subsequently $\gamma(\tau)$ and $\tilde{Z}(\tau)$) is not in question anymore, due to the results obtained in Theorem 3. The purpose of this theorem is just to obtain better *a priori* estimates, in the form described in above.
- The requirement $\beta > 1$, imposed so that (6.3) holds is likely only a technical one, but we cannot remove it with our methods.

Proof (Theorem 4) According to the results in Sect. 3.4, $\lambda_0 = -\frac{3-\alpha-\beta}{\alpha} \leq 0$ is an isolated and simple eigenvalue for the operator \mathcal{L} on $L^2(2)$, with eigenfunction G , while the rest of the spectrum is the essential spectrum, which we have identified before, $\sigma_{ess}(\mathcal{L}) = \{\lambda : \Re \lambda \leq -\frac{4-\alpha-\beta}{\alpha}\}$. We have also found the spectral projection $\mathcal{P}_0 f = \langle f, 1 \rangle G$ and $\mathcal{Q}_0 = Id - \mathcal{P}_0$. Thus, we can write

$$Z(\tau, \cdot) = \gamma(\tau)G(\xi) + \tilde{Z}(\tau, \cdot), \quad (6.4)$$

where $\gamma(\tau) = \langle Z(\tau, \cdot), 1 \rangle = \int_{\mathbf{R}^2} Z(\tau, \xi) d\xi$, $\tilde{Z}(\tau) = \mathcal{Q}_0 Z(\tau, \cdot)$. Projecting the equation (2.7), with respect to the spectral decomposition provided by \mathcal{P}_0 and \mathcal{Q}_0 , we obtain an ODE for γ and a PDE for $\tilde{Z}(\tau)$. More precisely,

$$\begin{aligned} \partial_\tau \gamma &= \langle \mathcal{L}Z, 1 \rangle - \langle U \cdot \nabla Z, 1 \rangle = \langle -|\nabla|^\alpha Z + \frac{1}{\alpha} \xi \cdot \nabla_\xi Z + \left(1 + \frac{\beta-1}{\alpha}\right) Z, 1 \rangle - \langle \nabla(U \cdot Z), 1 \rangle \\ &= \frac{\alpha+\beta-3}{\alpha} \gamma(\tau). \end{aligned}$$

Integrating this first-order ODE yields the formula $\gamma(\tau) = \gamma(0) e^{-\tau \frac{3-\alpha-\beta}{\alpha}}$. For the PDE governing $\tilde{Z}(\tau)$, and recalling $\mathcal{L}_0 = \mathcal{L} \mathcal{Q}_0$, we obtain

$$\tilde{Z}_\tau = \mathcal{L}_0 \tilde{Z} - \mathcal{Q}_0 [U \cdot \nabla Z] = \mathcal{L}_0 \tilde{Z} - \mathcal{Q}_0 \left[U \cdot \nabla (\gamma(0) e^{-\tau \frac{3-\alpha-\beta}{\alpha}} G + \tilde{Z}) \right].$$

In its equivalent integral formulation,

$$\begin{aligned} \tilde{Z}(\tau) &= e^{\tau \mathcal{L}_0} \tilde{Z}(0) - \int_0^\tau e^{(\tau-s) \mathcal{L}_0} \mathcal{Q}_0 \left[U \cdot \nabla (\gamma(0) e^{-\tau \frac{3-\alpha-\beta}{\alpha}} G + \tilde{Z}(s)) \right] ds. \end{aligned} \quad (6.5)$$

Note that since $\mathcal{P}_0 \nabla = 0$, we have the commutation relation $\mathcal{Q}_0 \nabla = (Id - \mathcal{P}_0) \nabla = \nabla$, whence one can remove \mathcal{Q}_0 in front of the nonlinearity. By (3.14), we can estimate

$$\begin{aligned} \|\tilde{Z}(\tau)\|_{L^2(2)} &\leq \|e^{\tau \mathcal{L}_0} \tilde{Z}(0)\|_{L^2(2)} \\ &\quad + \int_0^\tau \left\| e^{(\tau-s) \mathcal{L}_0} \left((U_G + U_{\tilde{Z}}) \nabla \cdot (\gamma(0) e^{-s \frac{3-\alpha-\beta}{\alpha}} G) + \tilde{Z}(s) \right) \right\|_{L^2(2)} ds \\ &\leq \|e^{\tau \mathcal{L}_0} \tilde{Z}(0)\|_{L^2(2)} + |\gamma(0)| \int_0^\tau e^{-\frac{(\tau-s)}{\alpha}} e^{-s \frac{3-\alpha-\beta}{\alpha}} \left\| \nabla \cdot e^{(\tau-s) \mathcal{L}_0} (U_{\tilde{Z}} \cdot G) \right\|_{L^2(2)} ds \\ &\quad + \int_0^\tau e^{-\frac{(\tau-s)}{\alpha}} \|\nabla \cdot e^{(\tau-s) \mathcal{L}_0} (U \cdot \tilde{Z})\|_{L^2(2)} ds =: I_1 + I_2 + I_3, \end{aligned}$$

where we have used (6.1). Clearly by (3.11)

$$I_1 \leq C e^{-\tau \left(\frac{4-\beta-\alpha}{\alpha} - \epsilon \right)} \|\tilde{Z}(0)\|_{L^2(2)}$$

Regarding I_2 , we have

$$I_2 \leq |\gamma(0)| \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}} e^{-s \frac{3-\alpha-\beta}{\alpha}} e^{-(\tau-s) \left(\frac{3-\beta-\alpha}{\alpha} \right)} \|U_{\tilde{Z}} \cdot G\|_{L^2(2)}}{a(\tau-s)^{\frac{1}{\alpha}}} ds$$

Now to bound $\|U_{\tilde{Z}} \cdot G\|_{L^2(2)}$ we look at two different cases, namely $0 \leq \beta < 1$ and $1 \leq \beta < 2$. If $0 \leq \beta \leq 1$, then we can use Lemma 2 to get

$$\begin{aligned} \|U_{\tilde{Z}} \cdot G\|_{L^2(2)} &\leq \|U_{\tilde{Z}}\|_{L^{\frac{2}{1-\beta}}} \|(1+|\xi|^2)G\|_{L^{\frac{2}{\beta}}} \\ &\leq C \|U_{\tilde{Z}}\|_{L^{\frac{2}{1-\beta}}} \leq C \|\nabla|^\beta U_{\tilde{Z}}\|_{L^2} \leq C \|\tilde{Z}\|_{L^2} \leq \|\tilde{Z}\|_{L^2(2)}. \end{aligned}$$

If $1 \leq \beta < 2$, then for some $0 < \epsilon \ll 1$ we have

$$\begin{aligned} \|U_{\tilde{Z}} \cdot G\|_{L^2(2)} &\leq \|U_{\tilde{Z}}\|_{L^{\frac{2}{\epsilon}}} \|(1+|\xi|^2)G\|_{L^{\frac{2}{1-\epsilon}}} \\ &\leq C \|U_{\tilde{Z}}\|_{L^{\frac{2}{\epsilon}}} \leq C \|\nabla|^\beta U_{\tilde{Z}}\|_{L^{\frac{2}{\beta+\epsilon}}} \leq C \|\tilde{Z}\|_{L^{\frac{2}{\beta+\epsilon}}} \leq C \|\tilde{Z}\|_{L^2(2)}. \end{aligned}$$

In the last inequality, we used the fact that for $1 < p < 2$, $L^p \hookrightarrow L^2(2)$ and (2.15). Therefore

$$I_2 \leq C \int_0^\tau \frac{e^{-(\tau-s) \left(\frac{4-\beta-\alpha}{\alpha} \right)} e^{-s \frac{3-\alpha-\beta}{\alpha}}}{(\min(1, |\tau-s|)^{\frac{1}{\alpha}})} \|\tilde{Z}(s)\|_{L^2(2)} ds.$$

Finally, we make use of (3.15) to get

$$\begin{aligned} I_3 &\leq \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}} e^{-(\tau-s) \left(\frac{3-\beta-\alpha}{\alpha} \right)} \|U(s)\|_{L^\infty} \|\tilde{Z}(s)\|_{L^2(2)}}{a(\tau-s)^{\frac{1}{\alpha}}} ds \\ &\leq C \int_0^\tau \frac{e^{-(\tau-s) \left(\frac{4-\beta-\alpha}{\alpha} - \epsilon \right)} e^{-s \left(\frac{3-\beta-\alpha}{\alpha} \right)}}{(\min(1, |\tau-s|)^{\frac{1}{\alpha}})} \|\tilde{Z}(s)\|_{L^2(2)} ds, \end{aligned}$$

where we have used that $a(\tau) \sim \min(1, \tau)$, the Sobolev inequality and Theorem 3 to conclude

$$\|U(s)\|_{L^\infty} \leq C \left(\|Z(s)\|_{L^{\frac{2}{\beta}+\epsilon}} + \|Z(s)\|_{L^{\frac{2}{\beta}-\epsilon}} \right) \leq C e^{-s \left(\frac{3-\beta-\alpha}{\alpha} \right)}. \quad (6.6)$$

We are now in a position to use the Gronwall's inequality, more precisely the version displayed in Lemma 6. We apply it with $I(\tau) = \|\tilde{Z}(\tau)\|_{L^2(2)}$, $\mu = \frac{4-\alpha-\beta}{\alpha} - \epsilon$, $\sigma =$

$\frac{4-\alpha-\beta}{\alpha}$, $\kappa = \frac{3-\alpha-\beta}{\alpha}$ and $a = \frac{1}{\alpha} < 1$, for $\epsilon \ll 1$. Recall that by the *a priori* estimates in Theorem 3, we have

$$\|\tilde{Z}(\tau)\|_{L^2(2)} \leq \|Z(\tau)\|_{L^2(2)} + |\gamma(0)|e^{-\tau(\frac{3-\alpha-\beta}{\alpha})} \|G\|_{L^2(2)} \leq Ce^{-\tau(\frac{3-\alpha-\beta}{\alpha})} \leq C,$$

for all $\tau > 0$, since $3 \geq \alpha + \beta$. Thus, all the requirements of Lemma 6 are met and we obtain the bound

$$\|\tilde{Z}(\tau)\|_{L^2(2)} \leq C_\epsilon e^{-\tau(\frac{4-\alpha-\beta}{\alpha}-\epsilon)}. \quad (6.7)$$

Regarding the proof of (6.3), we proceed in a similar fashion. We need to control $\|\partial \tilde{Z}\|_{L^2(2)}$, for large τ , say $\tau \geq 1$. Applying $\partial = \partial_1, \partial_2$ to the integral equation (6.5) and taking $\|\cdot\|_{L^2(2)}$, we obtain

$$\begin{aligned} \|\partial \tilde{Z}(\tau)\|_{L^2(2)} &\lesssim e^{-\tau(\frac{4-\alpha-\beta}{\alpha}-\epsilon)} \|\tilde{Z}(0)\|_{L^2(2)} \\ &+ \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}} e^{-s\frac{3-\alpha-\beta}{\alpha}}}{\min(1, \tau-s)^{\frac{1}{\alpha}}} \|e^{(\tau-s)\mathcal{L}_0} \nabla(U_{\tilde{Z}} \cdot G)\|_{L^2(2)} ds \\ &+ \int_0^\tau \frac{e^{-\frac{(\tau-s)}{\alpha}}}{\min(1, \tau-s)^{\frac{1}{\alpha}}} \|e^{(\tau-s)\mathcal{L}_0} \nabla(U \cdot \tilde{Z})\|_{L^2(2)} ds \lesssim e^{-\tau(\frac{4-\alpha-\beta}{\alpha}-\epsilon)} \\ &+ \int_0^\tau \frac{e^{-s\frac{3-\alpha-\beta}{\alpha}} e^{-(\tau-s)(\frac{5-\alpha-\beta}{\alpha}-\epsilon)}}{\min(1, \tau-s)^{\frac{1}{\alpha}}} \|\nabla[U_{\tilde{Z}}(s)G]\|_{L^2(2)} ds \\ &+ \int_0^\tau \frac{e^{-(\tau-s)(\frac{5-\alpha-\beta}{\alpha}-\epsilon)}}{\min(1, \tau-s)^{\frac{1}{\alpha}}} \|\nabla[U(s)\tilde{Z}(s)]\|_{L^2(2)} ds \end{aligned}$$

We estimate $\|\nabla[U_{\tilde{Z}}(s)G]\|_{L^2(2)} \leq \|\nabla U_{\tilde{Z}}(s)G\|_{L^2(2)} + \|U_{\tilde{Z}}(s)\nabla G\|_{L^2(2)}$. Following the strategy above, for $\beta \leq 1$ and then for $\beta > 1$, we arrive at

$$\|\nabla[U_{\tilde{Z}}(s)G]\|_{L^2(2)} \lesssim \|\tilde{Z}(s)\|_{L^2(2)} + \|\partial \tilde{Z}(s)\|_{L^2(2)} \lesssim e^{-s(\frac{4-\alpha-\beta}{\alpha}-\epsilon)} + \|\partial \tilde{Z}(s)\|_{L^2(2)},$$

where we have used (6.7). For the other term, it is relatively easy to bound $\|\nabla[U(s)\tilde{Z}(s)]\|_{L^2(2)}$, when $\beta > 1$,

$$\begin{aligned} \|\partial[U(s)\tilde{Z}(s)]\|_{L^2(2)} &\lesssim \|\partial U(s)\|_{L^\infty} \|\tilde{Z}(s)\|_{L^2(2)} + \|U(s)\|_{L^\infty} \|\partial \tilde{Z}(s)\|_{L^2(2)} \\ &\lesssim e^{-s(\frac{3-\alpha-\beta}{\alpha})} e^{-s(\frac{4-\alpha-\beta}{\alpha}-\epsilon)} + e^{-s(\frac{3-\beta-\alpha}{\alpha})} \|\partial \tilde{Z}(s)\|_{L^2(2)}. \end{aligned}$$

where we have used (recalling $U \sim |\nabla|^{-\beta} Z$), $\|\partial U(s)\|_{L^\infty} \leq C(\|Z\|_{L^{\frac{2}{\beta-1}+\epsilon}} + \|Z\|_{L^{\frac{2}{\beta-1}-\epsilon}}) \leq Ce^{-s(\frac{3-\alpha-\beta}{\alpha})}$, (6.7), (6.6). Plugging it together yields

$$\|\partial \tilde{Z}(\tau)\|_{L^2(2)} \lesssim e^{-\tau(\frac{4-\alpha-\beta}{\alpha}-\epsilon)} + \int_0^\tau \frac{e^{-(\tau-s)(\frac{5-\alpha-\beta}{\alpha}-\epsilon)} e^{-s(\frac{3-\alpha-\beta}{\alpha})}}{\min(1, \tau-s)^{\frac{1}{\alpha}}} \|\partial \tilde{Z}(s)\|_{L^2(2)}. \quad (6.8)$$

This puts us in a position to use the Gronwall's Lemma 6. Note that in order to do that, we need any *a priori* exponential bound on $\|\partial Z(\tau)\|_{L^2(2)}$, similar to Theorem 3 for $\|Z(\tau)\|_{L^2(2)}$. This is actually easy to achieve, one just has to differentiate the equation and perform very coarse energy estimates.¹³ As a result, Lemma 6 applies and we obtain

$$\|\partial \tilde{Z}(\tau)\|_{L^2(2)} \lesssim e^{-\tau(\frac{4-\alpha-\beta}{\alpha}-\epsilon)}.$$

This is nothing but a reformulation of the statement of (6.3). \square

It is now easy to conclude the main result, Theorem 1. Realizing that $L^2(2) \hookrightarrow L^p$, $1 \leq p \leq 2$, one just needs to translate the L^p estimates for Z , in the language of the original variable z . In fact,

$$\begin{aligned} \|Z(\cdot, \tau) - \gamma(0)e^{-\tau(\frac{3-\alpha-\beta}{\alpha})} G\|_{L^p} &= \left[\int_{\mathbf{R}^2} \left| Z(\xi, \tau) - \gamma(0)e^{-\tau(\frac{3-\alpha-\beta}{\alpha})} G(\xi) \right|^p d\xi \right]^{\frac{1}{p}} \\ &= \left[\int_{\mathbf{R}^2} \left| (1+t)^{1+\frac{\beta-1}{\alpha}} z(x, t) - \frac{\gamma(0)}{(1+t)^{\frac{3-\alpha-\beta}{\alpha}}} G\left(\frac{x}{(1+t)^{\frac{1}{\alpha}}}\right) \right|^p \frac{dx}{(1+t)^{\frac{2}{\alpha}}} \right]^{\frac{1}{p}} \\ &= (1+t)^{1+\frac{\beta-1}{\alpha}-\frac{2}{\alpha p}} \left\| z(t, \cdot) - \frac{\int_{\mathbf{R}^2} z_0(x) dx}{(1+t)^{\frac{2}{\alpha}}} G\left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}}\right) \right\|_{L^p}. \end{aligned}$$

Now if we compare this relation with the result in (6.2), we clearly see that

$$(1+t)^{1+\frac{\beta-1}{\alpha}-\frac{2}{\alpha p}} \left\| z(t, \cdot) - \frac{\int_{\mathbf{R}^2} z_0(x) dx}{(1+t)^{\frac{2}{\alpha}}} G\left(\frac{\cdot}{(1+t)^{\frac{1}{\alpha}}}\right) \right\|_{L^p} \leq \frac{C_\epsilon}{(1+t)^{\frac{4-\alpha-\beta}{\alpha}-\epsilon}},$$

which clearly leads to proof of Theorem 1.

7 Global Dynamics of the Solutions of the Boussinesq Model

In this section, we compute the optimal decay rate in $L^2(2)$ for the solution of the Boussinesq model (2.9). Recall that the relevant operator \mathcal{L} has the form

$$\mathcal{L} = -|\nabla|^\alpha + \frac{1}{\alpha} \xi \cdot \nabla_\xi + 1,$$

¹³ Which will give very inefficient exponential bounds on $\|\partial Z(\tau)\|_{L^2(2)}$, but that is all we need to jump start Lemma 6.

with $\lambda_0(\mathcal{L}) = 1 - \frac{2}{\alpha}$ and $\sigma_{ess}(\mathcal{L}) \subset \{\lambda : \Re \lambda \leq 1 - \frac{3}{\alpha}\}$.

Theorem 5 Suppose $\alpha \in (1, \frac{3}{2})$ and $W_0, \Theta_0 \in Y := L^2(2)(\mathbf{R}^2) \cap L^\infty(\mathbf{R}^2) \cap H^1(\mathbf{R}^2)$. Then for every $\delta > 0$, there exists $C = C_\delta(\|W_0\|_Y, \|\Theta_0\|_Y) > 0$, such that for any $\tau > 0$, the solutions W, Θ for the system of equations (2.9) obey

$$\begin{aligned} & \|W - \gamma_2(0)e^{-(\frac{3}{\alpha}-2)\tau} \partial_1 G - \gamma_1(0)e^{-(\frac{2}{\alpha}-1)\tau} G\|_{L^2(2)} \\ & + \|\Theta - \gamma_2(0)e^{-(\frac{3}{\alpha}-2)\tau} G\|_{L^2(2)} \leq Ce^{-2(\frac{3}{\alpha}-2-\delta)\tau}. \end{aligned} \quad (7.1)$$

where $\gamma_1(0) := \int W_0(\xi) d\xi$, and $\gamma_2(0) := \int \Theta_0(\xi) d\xi$. In particular, if $\widehat{W}_0(0) = \widehat{\Theta}_0(0) = 0$ then

$$\|W\|_{L^2(2)} + \|\Theta\|_{L^2(2)} \leq C_\delta e^{-2(\frac{3}{\alpha}-2-\delta)\tau}. \quad (7.2)$$

Proof Using the spectral decomposition for \mathcal{L} , described in Sect. 3.4, write

$$W(\tau) = \gamma_1(\tau)G(\xi) + \widetilde{W}(\tau) \quad (7.3)$$

$$\Theta(\tau) = \gamma_2(\tau)G(\xi) + \widetilde{\Theta}(\tau) \quad (7.4)$$

where $\gamma_1(\tau) := \langle W(\tau), 1 \rangle$, $\gamma_2(\tau) := \langle \Theta(\tau), 1 \rangle$, $\widetilde{W} = \mathcal{Q}_0 W(\tau, \cdot)$ and $\widetilde{\Theta} = \mathcal{Q}_0 \Theta(\tau, \cdot)$. Then, we derive the equations for γ_1, γ_2 as before - namely

$$\begin{aligned} \partial_\tau \gamma_1 &= \langle W_\tau, 1 \rangle = \langle \mathcal{L}W, 1 \rangle - \langle U \cdot \nabla W, 1 \rangle + \langle \partial_1 \Theta, 1 \rangle \\ &= \langle \mathcal{L}W, 1 \rangle = \langle W, \mathcal{L}^* 1 \rangle = \left(1 - \frac{2}{\alpha}\right) \langle W, 1 \rangle = \left(1 - \frac{2}{\alpha}\right) \gamma_1(\tau) \end{aligned}$$

Similarly, $\partial_\tau \gamma_2 = (2 - \frac{3}{\alpha}) \gamma_2(\tau)$. Solving the ODE's results in the formulas

$$\gamma_1(\tau) = \gamma_1(0)e^{(1-\frac{2}{\alpha})\tau}, \quad \gamma_2(\tau) = \gamma_2(0)e^{(2-\frac{3}{\alpha})\tau}.$$

For the projections over the essential spectrum, we have the following PDE's

$$\begin{aligned} \widetilde{W}_\tau &= \mathcal{L}\widetilde{W} - \mathcal{Q}_0[U \cdot \nabla W - \partial_1 \Theta] = \mathcal{L}\widetilde{W} - \mathcal{Q}_0[U \cdot \nabla(\gamma_1(0) e^{(1-\frac{2}{\alpha})\tau} G + \widetilde{W})] + \\ &\quad + \mathcal{Q}_0[\partial_1(\gamma_2(0) e^{(1-\frac{2}{\alpha})\tau} G + \widetilde{\Theta})], \\ \widetilde{\Theta}_\tau &= (\mathcal{L} + 1 - \frac{1}{\alpha})\widetilde{\Theta} - \mathcal{Q}_0[U \cdot \nabla \Theta] = (\mathcal{L} + 1 - \frac{1}{\alpha})\widetilde{\Theta} - \mathcal{Q}_0[U \cdot \nabla(\gamma_2(0) e^{(2-\frac{3}{\alpha})\tau} G + \widetilde{\Theta})]. \end{aligned}$$

We represent them via the Duhamel's formula

$$\begin{aligned} \widetilde{W}(\tau) &= e^{\tau \mathcal{L}} \widetilde{W}_0 - \int_0^\tau e^{(\tau-s)\mathcal{L}} \mathcal{Q}_0 \left[U \cdot \nabla(\gamma_1(0) e^{(1-\frac{2}{\alpha})s} G + \widetilde{W}(s)) \right] ds \\ &\quad + \int_0^\tau e^{(\tau-s)\mathcal{L}} \mathcal{Q}_0[\partial_1 \Theta(s)] ds, \\ \widetilde{\Theta}(\tau) &= e^{\tau(\mathcal{L}+1-\frac{1}{\alpha})} \widetilde{\Theta}_0 - \int_0^\tau e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{Q}_0 \left[U \cdot \nabla(\gamma_2(0) e^{(2-\frac{3}{\alpha})s} G + \widetilde{\Theta}(s)) \right] ds. \end{aligned}$$

One term deserves a special attention, as it is explicit. Note that $\mathcal{D}_0 \partial_1 = \partial_1$, since $\mathcal{P}_0 \partial_1 = 0$. Also for $\kappa > 0$, since G is an eigenfunction, with eigenvalue $1 - \frac{2}{\alpha}$, we have $e^{\kappa \mathcal{L}} G = e^{(1 - \frac{2}{\alpha})\kappa} G$. By Lemma 7,

$$\begin{aligned} \int_0^\tau e^{(\tau-s)\mathcal{L}} \mathcal{D}_0[\partial_1 \Theta(s)] ds &= \int_0^\tau e^{(\tau-s)\mathcal{L}} [\partial_1 [\gamma_2(0) e^{(2-\frac{3}{\alpha})s} G + \tilde{\Theta}(s)]] ds \\ &= \gamma_2(0) \int_0^\tau e^{(2-\frac{3}{\alpha})s} e^{-\frac{\tau-s}{\alpha}} \partial_1 e^{(\tau-s)\mathcal{L}} [G] ds + \int_0^\tau e^{-\frac{\tau-s}{\alpha}} \partial_1 e^{(\tau-s)\mathcal{L}} \tilde{\Theta}(s) ds \\ &= \gamma_2(0) \partial_1 G \int_0^\tau e^{(2-\frac{3}{\alpha})s} e^{-\frac{\tau-s}{\alpha}} e^{(1-\frac{2}{\alpha})(\tau-s)} ds + \int_0^\tau e^{-\frac{\tau-s}{\alpha}} \partial_1 e^{(\tau-s)\mathcal{L}} \tilde{\Theta}(s) ds \\ &= \gamma_2(0) (e^{(2-\frac{3}{\alpha})\tau} - e^{(1-\frac{3}{\alpha})\tau}) \partial_1 G + \int_0^\tau e^{-\frac{\tau-s}{\alpha}} \partial_1 e^{(\tau-s)\mathcal{L}} \tilde{\Theta}(s) ds. \end{aligned}$$

At this point, it makes more sense to introduce the new variable,

$$W_1(\tau, \xi) := \tilde{W}(\tau, \xi) - \gamma_2(0) (e^{(2-\frac{3}{\alpha})\tau} - e^{(1-\frac{3}{\alpha})\tau}) \partial_1 G =: \tilde{W} - e^{(2-\frac{3}{\alpha})\tau} G_1(\tau, \xi).$$

Note that the decay rate $e^{(2-\frac{3}{\alpha})\tau}$ along the G_1 direction is slower than the decay rate $e^{(1-\frac{2}{\alpha})\tau}$ of the evolution along the G direction. Also, G_1 is basically $\partial_1 G$ multiplied by a bounded function of τ and hence an element of $L^2(2) \cap L^\infty$. For future reference,

$$\|W_1\|_X - C e^{(2-\frac{3}{\alpha})\tau} \leq \|\tilde{W}\|_X \leq \|W_1\|_X + C e^{(2-\frac{3}{\alpha})\tau}. \quad (7.5)$$

for all Banach spaces in consideration herein.

We write the equations for W_1 and $\tilde{\Theta}$ as follows

$$\begin{aligned} W_1(\tau) &= e^{\tau\mathcal{L}} \tilde{W}_0 - \int_0^\tau e^{(\tau-s)\mathcal{L}} \mathcal{D}_0[U \cdot \nabla (\gamma_1(0) e^{(1-\frac{2}{\alpha})s} G + e^{(2-\frac{3}{\alpha})s} G_1 + W_1(s))] ds \\ &\quad + \int_0^\tau e^{-\frac{\tau-s}{\alpha}} \partial_1 e^{(\tau-s)\mathcal{L}} \tilde{\Theta}(s) ds. \\ \tilde{\Theta}(\tau) &= e^{\tau(\mathcal{L}+1-\frac{1}{\alpha})} \tilde{\Theta}_0 - \int_0^\tau e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{D}_0[U \cdot \nabla (\gamma_2(0) e^{(2-\frac{3}{\alpha})s} G + \tilde{\Theta}(s))] ds. \end{aligned}$$

Note that $U = e^{(1-\frac{2}{\alpha})s} U_G + e^{(2-\frac{3}{\alpha})s} U_{G_1} + U_{W_1}$ and $U_G \cdot G = 0$.

We start the estimates for $\tilde{\Theta}$

$$\begin{aligned} \|\tilde{\Theta}\|_{L^2(2)} &\leq C e^{(2-\frac{4}{\alpha}+\delta)\tau} \|\tilde{\Theta}(0)\|_{L^2(2)} \\ &\quad + |\gamma_2(0)| \int_0^\tau e^{(2-\frac{3}{\alpha})s} \|e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{D}_0[U \cdot \nabla G]\|_{L^2(2)} ds + \\ &\quad + \int_0^\tau \|e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{D}_0[U \cdot \nabla \tilde{\Theta}(s)]\|_{L^2(2)} ds =: C e^{(2-\frac{4}{\alpha}+\delta)\tau} + J_1 + J_2 \end{aligned}$$

For all $\delta > 0$ small enough, there is C_δ ,

$$\begin{aligned} J_1 &= \int_0^\tau e^{(2-\frac{3}{\alpha})s} \|e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{Q}_0[U \cdot \nabla G]\|_{L^2(2)} ds \\ &\lesssim \|U_{G_1} G\|_{L^2(2)} \int_0^\tau \frac{e^{(2-\frac{5}{\alpha}+\delta)(\tau-s)} e^{2(2-\frac{3}{\alpha})s}}{(a(\tau-s))^{\frac{1}{\alpha}}} ds \\ &\quad + \int_0^\tau \frac{e^{(2-\frac{5}{\alpha}+\delta)(\tau-s)} e^{(2-\frac{3}{\alpha})s}}{(\min(1, |\tau-s|))^{\frac{1}{\alpha}}} \|U_{W_1}(s, \cdot) \cdot \nabla G\|_{L^2(2)} ds \lesssim e^{2(2-\frac{3}{\alpha})\tau} \\ &\quad + \int_0^\tau \frac{e^{(2-\frac{5}{\alpha}+\delta)(\tau-s)} e^{(2-\frac{3}{\alpha})s}}{(\min(1, |\tau-s|))^{\frac{1}{\alpha}}} (e^{(2-\frac{3}{\alpha})s})^{1-\epsilon} ds \leq C_\delta e^{2(2-\frac{3}{\alpha})\tau}. \end{aligned}$$

where we have used Lemma 2, Gagliardo–Nirenberg's, (5.6), $L^2(2) \hookrightarrow L^1$, (5.10), to estimate

$$\begin{aligned} \|U_{W_1} \nabla G\|_{L^2(2)} &\leq \|U_{W_1}\|_{L^{\frac{2}{\epsilon}}} \|(1+|\xi|^2)|\nabla G|\|_{L^{\frac{2}{1-\epsilon}}} \leq C \|U_{W_1}\|_{L^{\frac{2}{\epsilon}}} \leq C \|W_1\|_{L^{\frac{2}{1+\epsilon}}} \\ &\leq C \|W_1\|_{L^2}^{1-\epsilon} \|W_1\|_{L^1}^\epsilon \leq C (e^{(2-\frac{3}{\alpha})s})^{1-\epsilon}. \end{aligned}$$

Similarly,

$$\begin{aligned} J_2 &= \int_0^\tau \|e^{(\tau-s)(\mathcal{L}+1-\frac{1}{\alpha})} \mathcal{Q}_0[U \cdot \nabla \tilde{\Theta}(s)]\|_{L^2(2)} ds \\ &\leq C \int_0^\tau \frac{e^{(2-\frac{5}{\alpha}+\delta)(\tau-s)}}{(\min(1, |\tau-s|))^{\frac{1}{\alpha}}} \|U(s)\|_{L^\infty} \|\tilde{\Theta}(s)\|_{L^2(2)} ds \end{aligned}$$

Thus, we need a good estimate of $\|U(s)\|_{L^\infty}$. We have by (2.2)

$$\|U(s, \cdot)\|_{L^\infty} \leq C(\|W(s, \cdot)\|_{L^{2+\epsilon}} + \|W(s, \cdot)\|_{L^{2-\epsilon}}).$$

By the *a priori* estimate (5.9), we have a good control of $\|W(s, \cdot)\|_{L^{2+\epsilon}}$, namely $\|W(s, \cdot)\|_{L^{2+\epsilon}} \leq C e^{(2-\frac{3}{\alpha})s}$. For $\|W(s, \cdot)\|_{L^{2-\epsilon}}$, we can control it by (5.10), but this is not efficient for our arguments—we need some, however small, decay in s , which we can then input in the Gronwall's, (2.20). To achieve that, we proceed by Gagliardo–Nirenberg's estimate. Taking into account once again $L^2(2) \hookrightarrow L^1$, and the bounds (5.6),

$$\|W(s, \cdot)\|_{L^{2-\epsilon}} \leq \|W(s, \cdot)\|_{L^2}^{\frac{2-2\epsilon}{2-\epsilon}} \|W(s, \cdot)\|_{L^1}^{\frac{\epsilon}{2-\epsilon}} \leq C (e^{(2-\frac{3}{\alpha})s})^{\frac{2-2\epsilon}{2-\epsilon}}.$$

All in all, for all $\delta > 0$,

$$\|U(s, \cdot)\|_{L^\infty} \leq C_\delta e^{-(\frac{3}{\alpha}-2-\delta)s}. \quad (7.6)$$

This results in the following estimates for J_2

$$J_2 \leq \int_0^\tau \frac{e^{\left(2-\frac{5}{\alpha}+\delta\right)(\tau-s)} e^{-\left(\frac{3}{\alpha}-2-\delta\right)s}}{(\min(1, |\tau-s|)^{\frac{1}{\alpha}})} \|\tilde{\Theta}(s)\|_{L^2(2)} ds$$

Combining all the estimates obtained about¹⁴ $\|\tilde{\Theta}(s)\|_{L^2(2)}$, we have

$$\|\tilde{\Theta}(\tau)\|_{L^2(2)} \leq C e^{-2(\frac{3}{\alpha}-2-\delta)\tau} + \int_0^\tau \frac{e^{\left(2-\frac{5}{\alpha}+\delta\right)(\tau-s)} e^{-\left(\frac{3}{\alpha}-2-\delta\right)s}}{(\min(1, |\tau-s|)^{\frac{1}{\alpha}})} \|\tilde{\Theta}(s)\|_{L^2(2)} ds$$

Applying the Gronwall's, more precisely Lemma 6, we conclude

$$\|\tilde{\Theta}(\tau)\|_{L^2(2)} \leq C_\delta e^{-(\frac{3}{\alpha}-2-\delta)\tau},$$

as stated. For W_1 , we get

$$\begin{aligned} \|W_1\|_{L^2(2)} &\leq C e^{-(\frac{3}{\alpha}-1-\delta)\tau} \|\tilde{W}_0\|_{L^2(2)} \\ &+ \int_0^\tau e^{-\frac{(\tau-s)}{\alpha}} \|\nabla e^{(\tau-s)\mathcal{L}_0} [U \cdot (\gamma_1(0)e^{(1-\frac{2}{\alpha})s}G + e^{(2-\frac{3}{\alpha})s}G_1) + U \cdot W_1]\|_{L^2(2)} ds \\ &+ \int_0^\tau e^{-\frac{(\tau-s)}{\alpha}} \|\partial_1 e^{(\tau-s)\mathcal{L}_0} \tilde{\Theta}(s)\|_{L^2(2)} ds \lesssim e^{(1-\frac{3}{\alpha}+\delta)\tau} \\ &+ \int_0^\tau \frac{e^{(1-\frac{3}{\alpha}+\delta)(\tau-s)} e^{(2-\frac{3}{\alpha})s} \|U(|G| + |G_1|)\|_{L^2(2)}}{(a(\tau-s))^{\frac{1}{\alpha}}} ds \\ &+ \int_0^\tau \frac{e^{(1-\frac{3}{\alpha}+\delta)(\tau-s)} \|U\|_{L^\infty} \|W_1\|_{L^2(2)}}{(a(\tau-s))^{\frac{1}{\alpha}}} ds \\ &+ \int_0^\tau \frac{e^{(1-\frac{4}{\alpha}+\delta)(\tau-s)} \|\tilde{\Theta}(s)\|_{L^2(2)}}{(a(\tau-s))^{\frac{1}{\alpha}}} ds = e^{(1-\frac{3}{\alpha}+\delta)\tau} + I_1 + I_2 + I_3 \end{aligned}$$

For I_1 , we have

$$\begin{aligned} \|U(|G| + |G_1|)\|_{L^2(2)} &\leq \|(e^{(1-\frac{2}{\alpha})s}U_G + e^{(2-\frac{3}{\alpha})s}U_{G_1})(|G| + |G_1|)\|_{L^2(2)} \\ &+ \|U_{W_1}(|G| + |G_1|)\|_{L^2(2)}. \end{aligned}$$

The first term is easily estimated, since $G, G_1 \in L^2(2)$ (whence $U_G, U_{G_1} \in L^\infty$ by Sobolev embedding and Lemma 2)

$$\|(e^{(1-\frac{2}{\alpha})s}U_G + e^{(2-\frac{3}{\alpha})s}U_{G_1})(|G| + |G_1|)\|_{L^2(2)} \leq C e^{(2-\frac{3}{\alpha})s},$$

¹⁴ Note that with our restrictions on α , $(\frac{3}{\alpha}-2) < \frac{4}{\alpha}-2$, so this is the slowest rate on the right-hand sides of $\|\tilde{\Theta}(\tau)\|_{L^2(2)}$.

whence the contribution of these terms is no more than

$$C \int_0^\tau \frac{e^{(1-\frac{3}{\alpha}+\delta)(\tau-s)} e^{2(2-\frac{3}{\alpha})s}}{\min(1, |\tau-s|^{\frac{1}{\alpha}})} ds \leq C e^{2\tau(2-\frac{3}{\alpha})}.$$

For U_{W_1} terms, we can use Lemma 2, the Sobolev inequality and $L^2(2) \hookrightarrow L^{\frac{2}{1+\epsilon}}$ to get

$$\begin{aligned} \|U_{W_1}(s)(|G| + |G_1|)\|_{L^2(2)} &= \|U_{W_1} \cdot (1 + |\xi|^2)(|G| + |G_1|)\|_{L^2} \\ &\leq \|U_{W_1}\|_{L^{\frac{2}{\epsilon}}} \|(1 + |\xi|^2)(|G| + |G_1|)\|_{L^{\frac{2}{1-\epsilon}}} \\ &\leq C \|U_{W_1}\|_{L^{\frac{2}{\epsilon}}} \leq C \|\nabla U_{W_1}\|_{L^{\frac{2}{1+\epsilon}}} \\ &\leq C \|W_1\|_{L^{\frac{2}{1+\epsilon}}} \leq C \|W_1(s)\|_{L^2(2)}. \end{aligned}$$

All together, the contribution of I_1 is estimated by

$$I_1 \leq C e^{-2(\frac{3}{\alpha}-2)\tau} + \int_0^\tau \frac{e^{-(\frac{3}{\alpha}-1-\delta)(\tau-s)} e^{-(\frac{3}{\alpha}-2)s}}{\min(1, |\tau-s|^{\frac{1}{\alpha}})} \|W_1(s)\|_{L^2(2)} ds$$

Regarding I_2 , we first need an appropriate estimate on $\|U\|_{L^\infty}$, which is fortunately already given by (7.6). This then gives the bound for I_2 ,

$$I_2 \leq \int_0^\tau \frac{e^{-(\frac{3}{\alpha}-1-\delta)(\tau-s)} e^{-(\frac{3}{\alpha}-2-\delta)s}}{\min(1, |\tau-s|^{\frac{1}{\alpha}})} \|W_1(s)\|_{L^2(2)} ds$$

Combining all estimates for $\|W_1(\tau)\|_{L^2(2)}$ yields

$$\|W_1(\tau, \cdot)\|_{L^2(2)} \leq C e^{-2(\frac{3}{\alpha}-2)\tau} + \int_0^\tau \frac{e^{-(\frac{3}{\alpha}-1-\delta)(\tau-s)} e^{-(\frac{3}{\alpha}-2-\delta)s}}{\min(1, |\tau-s|^{\frac{1}{\alpha}})} \|W_1(s)\|_{L^2(2)} ds.$$

Applying Lemma 6, with $\mu = 2(\frac{3}{\alpha}-2)$, $\sigma = (\frac{3}{\alpha}-1-\delta)$, $\kappa = (\frac{3}{\alpha}-2-\delta)$ yields

$$\|W_1(\tau, \cdot)\|_{L^2(2)} \leq C e^{-2(\frac{3}{\alpha}-2)\tau}.$$

This is the statement of (7.1) and Theorem 5 is proved in full. \square

Acknowledgements The authors would like to take the opportunity to thank Ryan Goh and Jiahong Wu for stimulating discussions regarding these topics.

Appendix A. Sobolev Embedding at L^∞ : Relation (2.2)

Before we start the proof of (2.2), we recall the following Bernstein inequality. Let g satisfy

$$\text{supp } \hat{g} \subset \{\xi \in R^n : C_1 2^k \leq |\xi| \leq C_2 2^{k+1}\}$$

for some k , and constants $C_1 \leq C_2$. Then for any $\alpha \geq 0$ and $1 \leq p \leq q \leq \infty$,

$$C_1 2^{\alpha k} \|g\|_{L^q(\mathbf{R}^n)} \leq \|\nabla^\alpha g\|_{L^q(\mathbf{R}^n)} \leq C_2 2^{\alpha k + nk(\frac{1}{p} - \frac{1}{q})} \|g\|_{L^p(\mathbf{R}^n)}.$$

Now let $\widehat{P_k f}(\xi) = \widehat{\psi}(2^{-k}\xi) \widehat{f}(\xi)$, where $\widehat{\psi} \in C^\infty$, $\text{supp } \widehat{\psi} \subseteq \{\xi \in R^n : \xi \in (\frac{1}{2}, 2)\}$. Then

$$\|(\nabla^\perp)^{-\beta} f\|_{L^\infty} \leq \sum_{k=0}^{\infty} \|P_k((\nabla^\perp)^{-\beta} f)\|_{L^\infty} + \sum_{k=0}^{\infty} \|P_{-k}((\nabla^\perp)^{-\beta} f)\|_{L^\infty}.$$

We make use of the above Bernstein inequality several times to control each of these terms. Indeed,

$$\begin{aligned} \sum_{k=0}^{\infty} \|P_k((\nabla^\perp)^{-\beta} f)\|_{L^\infty} &\leq \sum_{k=0}^{\infty} 2^{-k\beta} \|P_k f\|_{L^\infty} \leq \sum_{k=0}^{\infty} 2^{-k\beta + nk(\frac{1}{\beta} + \delta)} \|P_k f\|_{L^{\frac{n}{\beta} + \delta}} \\ &\leq \|f\|_{L^{\frac{n}{\beta} + \delta}} \sum_{k=0}^{\infty} 2^{-k\beta(1 - \frac{n}{n + \beta\gamma})} \leq C \|f\|_{L^{\frac{n}{\beta} + \delta}}. \end{aligned}$$

In the same way,

$$\begin{aligned} \sum_{k=0}^{\infty} \|P_{-k}((\nabla^\perp)^{-\beta} f)\|_{L^\infty} &\leq \sum_{k=0}^{\infty} 2^{k\beta} \|P_{-k} f\|_{L^\infty} \leq \sum_{k=0}^{\infty} 2^{k\beta - nk(\frac{1}{\beta} - \delta)} \|P_k f\|_{L^{\frac{n}{\beta} - \delta}} \\ &\leq \|f\|_{L^{\frac{n}{\beta} - \delta}} \sum_{k=0}^{\infty} 2^{k\beta(1 - \frac{n}{n - \beta\gamma})} \leq C \|f\|_{L^{\frac{n}{\beta} - \delta}}. \end{aligned}$$

Appendix B. Generalized Gronwall's Estimate: Lemma 6

The proof of Lemma 6 is straightforward, by a bootstrapping argument. We show that every Lyapunov exponent less than $-\mu$ can be bootstrapped lower. First, relabeling $I(\tau) \rightarrow (1 + |A_1| + |A_2| + |A_3|)^{-1} I(\tau)$, we may assume without loss of generality that $A_1 = A_2 = A_3 = 1$. Next, assume that $\gamma < \mu$ is a Lyapunov exponent, that is $I(\tau) \leq C e^{-\gamma\tau}$. We know by the *a priori* assumed boundedness of $I(\tau)$ there is such an exponent. Applying this in (2.20), we obtain an improved estimate for $I(\tau)$.

Indeed,

$$I(\tau) \leq e^{-\mu\tau} + Ce^{-\sigma\tau} \int_0^\tau \frac{e^{s(\sigma-\kappa-\gamma)}}{(\min(1, |\tau-s|)^a)} ds$$

If $\sigma - \kappa - \gamma \neq 0$, we have for $\tau > 1$,

$$\begin{aligned} \int_0^\tau \frac{e^{s(\sigma-\kappa-\gamma)}}{(\min(1, |\tau-s|)^a)} ds &\leq \int_0^{\tau-1} e^{s(\sigma-\kappa-\gamma)} ds + e^{\tau(\sigma-\kappa-\gamma)} e^{|\sigma-\kappa-\gamma|} \int_{\tau-1}^\tau \frac{1}{|\tau-s|^a} ds \\ &\leq \frac{e^{(\tau-1)(\sigma-\kappa-\gamma)} - 1}{\sigma - \kappa - \gamma} + C_{a,\sigma,\kappa,\gamma} e^{\tau(\sigma-\kappa-\gamma)}. \end{aligned}$$

whence the bound

$$I(\tau) \leq e^{-\mu\tau} + C_{a,\sigma,\kappa,\gamma} e^{-\tau(\kappa+\gamma)}.$$

It follows that $\min(\mu, \gamma + \kappa) > \gamma$ is a new, better Lyapunov exponent than γ .

In general, we can keep $\sigma - \kappa - \gamma$ away from zero (and so the previous argument valid in all cases), if we readjust the γ if necessary.

In practice, starting with $\gamma = 0$, we jump immediately to κ by the previous argument, since $\sigma - \kappa > 0$, by assumption. Since $\kappa < \mu$, we can apply the same argument again with $\gamma = \kappa$. At this point, either $2\kappa > \mu$ and we finish off (by readjusting slightly γ by taking it smaller, like $\gamma = \frac{2\kappa}{3}$, if it happens that, say $|\sigma - 2\kappa| \leq \frac{\kappa}{2}$). If not, that is if $2\kappa < \mu$, take $\gamma = 2\kappa$ to be our new Lyapunov exponent and repeat. Eventually, for some n_0 , $n_0\kappa < \mu \leq (n_0+1)\kappa$ and we will reach a Lyapunov exponent μ .

Appendix C. Commutator Estimates with Weights

In this section, we prove (4.8) and (5.13).

C.1. Proof of (4.8)

Recall, that for $s \in (0, 2)$

$$\begin{aligned} [|\nabla|^s, g]f(x) &= |\nabla|^s(gf) - g|\nabla|^s f \\ &= c_s \int \frac{f(x)g(x) - f(y)g(y)}{|x-y|^{2+s}} dy - g(x)c_s \int \frac{f(x) - f(y)}{|x-y|^{2+s}} dy \\ &= c_s \int \frac{f(y)(g(x) - g(y))}{|x-y|^{2+s}} dy. \end{aligned}$$

Introduce a smooth partition of unity, that is a function $\psi \in C_0^\infty(\mathbf{R})$, $\text{supp } \psi \subset (\frac{1}{2}, 2)$, so that

$$\sum_{k=-\infty}^{\infty} \psi(2^{-k}|\xi|) = 1, \xi \in \mathbf{R}^2, \xi \neq 0.$$

Introduce another C_0^∞ function $\Psi(z) = z^2\psi(z)$, so that we can decompose

$$|\xi|^2 = \sum_{k=-\infty}^{\infty} |\xi|^2 \psi(2^{-k}|\xi|) = \sum_{k=-\infty}^{\infty} 2^{2k} \Psi(2^{-k}|\xi|).$$

We can then write

$$\begin{aligned} F(\xi) &:= [|\nabla|^{\frac{\alpha}{2}}, |\xi|^2] f \sum_k 2^{2k} [|\nabla|^{\frac{\alpha}{2}}, \Psi(2^{-k}\cdot)] f(\xi) \\ &= \sum_k 2^{2k} \int \frac{f(y)(\Psi(2^{-k}\xi) - \Psi(2^{-k}y))}{|\xi - y|^{2+\frac{\alpha}{2}}} dy. \end{aligned}$$

Introducing

$$F_k := \int \frac{|f(y)||\Psi(2^{-k}\xi) - \Psi(2^{-k}y)|}{|\xi - y|^{2+\frac{\alpha}{2}}} dy,$$

we need to control

$$\begin{aligned} \|F\|_{L^2}^2 &= \sum_l \int_{|\xi| \sim 2^l} |F(\xi)|^2 d\xi = \sum_l \int_{|\xi| \sim 2^l} \left| \sum_k 2^{2k} F_k(\xi) \right|^2 d\xi = \\ &= \sum_l \int_{|\xi| \sim 2^l} \left| \sum_{k > l+10} 2^{2k} F_k(\xi) \right|^2 d\xi + \sum_l \int_{|\xi| \sim 2^l} \left| \sum_{k=l-10}^{l+10} 2^{2k} F_k(\xi) \right|^2 d\xi + \\ &\quad + \sum_l \int_{|\xi| \sim 2^l} \left| \sum_{k < l-10} 2^{2k} F_k(\xi) \right|^2 d\xi =: K_1 + K_2 + K_3 \end{aligned}$$

We first consider the cases $k > l+10$. One can estimate easily F_k point-wise. More specifically, since in the denominator of the expression for F_k , we have $|\xi - y| \geq \frac{1}{2}|\xi| \geq 2^{k-3}$,

$$|F_k(\xi)| \leq 2^{-k(2+\frac{\alpha}{2})} \int |f(y)||\Psi(2^{-k}y)| dy \leq C 2^{-k(1+\frac{\alpha}{2})} \|f\|_{L^2(|y| \sim 2^k)},$$

whence

$$\begin{aligned}
 K_1 &\leq \sum_l 2^{2l} \sum_{k_1 > l+10} \sum_{k_2 > l+10} 2^{k_1(1-\frac{\alpha}{2})} \|f\|_{L^2(|y|\sim 2^{k_1})} 2^{k_2(1-\frac{\alpha}{2})} \|f\|_{L^2(|y|\sim 2^{k_2})} \\
 &\leq \sum_{k_1} \sum_{k_2} 2^{2\min(k_1, k_2)} 2^{k_1(1-\frac{\alpha}{2})} \|f\|_{L^2(|y|\sim 2^{k_1})} 2^{k_2(1-\frac{\alpha}{2})} \|f\|_{L^2(|y|\sim 2^{k_2})} \\
 &\leq C \sum_k 2^{k(4-\alpha)} \|f\|_{L^2(|y|\sim 2^k)}^2 \leq C \|\xi\|^{2-\frac{\alpha}{2}} \|f\|^2.
 \end{aligned}$$

where we have used $\sum_{l:l < \min(k_1, k_2) - 10} 2^{2l} \leq C 2^{2\min(k_1, k_2)}$.

For the case $k < l - 10$, we perform similar argument, since

$$|F_k(\xi)| \leq C 2^{-l(2+\frac{\alpha}{2})} 2^k \|f\|_{L^2(|y|\sim 2^k)}.$$

So,

$$\begin{aligned}
 K_3 &\leq C \sum_l 2^{2l} 2^{-l(4+\alpha)} \sum_{k_1 < l-10} \sum_{k_2 < l-10} 2^{3k_1} \|f\|_{L^2(|y|\sim 2^{k_1})} 2^{3k_2} \|f\|_{L^2(|y|\sim 2^{k_2})} \\
 &\leq C \sum_{k_1} \sum_{k_2} 2^{3k_1} \|f\|_{L^2(|y|\sim 2^{k_1})} 2^{3k_2} \|f\|_{L^2(|y|\sim 2^{k_2})} 2^{-(2+\alpha)\max(k_1, k_2)} \\
 &\leq C \sum_k 2^{k(4-\alpha)} \|f\|_{L^2(|y|\sim 2^k)}^2 \leq C \|\xi\|^{2-\frac{\alpha}{2}} \|f\|^2.
 \end{aligned}$$

Finally, for the case $|l - k| \leq 10$, we use

$$|\Psi(2^{-k}\xi) - \Psi(2^{-k}y)| \leq 2^{-k} |\xi - y| |\nabla \Psi(2^{-k}(\xi - y))| \leq C 2^{-k} |\xi - y|,$$

so that

$$|F_k(\xi)| \leq C 2^{-k} \int_{|y|\sim 2^k} \frac{|f(y)|}{|\xi - y|^{1+\frac{\alpha}{2}}} dy = C 2^{-k} |f| \chi_{|y|\sim 2^k} * \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}}.$$

Thus, by Hölder's

$$\begin{aligned}
 K_2 &\leq C \sum_k \int_{|\xi|\sim 2^k} 2^{2k} \left| |f| \chi_{|y|\sim 2^k} * \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}} \right|^2 d\xi \\
 &\leq C \sum_k 2^{2k} \| |f| \chi_{|y|\sim 2^k} * \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}} \|_{L^2(|\xi|\sim 2^k)}^2 \\
 &\leq C \sum_k 2^{k(4-\alpha)} \| |f| \chi_{|y|\sim 2^k} * \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}} \|_{L^{\frac{4}{1+\frac{\alpha}{2}}}(|\xi|\sim 2^k)}^2 \\
 &\leq C \sum_k 2^{k(4-\alpha)} \|f\|_{L^2(|\xi|\sim 2^k)}^2 \leq C \|\xi\|^{2-\frac{\alpha}{2}} \|f\|^2.
 \end{aligned}$$

where we have used the Hausdorff–Young’s inequality

$$\left\| f \chi_{|y| \sim 2^k} * \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}} \right\|_{L^{\frac{4}{\alpha}}} \leq C \left\| \frac{1}{|\cdot|^{1+\frac{\alpha}{2}}} \right\|_{L^{\frac{4}{2-\alpha}}, \infty} \|f\|_{L^2(|\xi| \sim 2^k)} \leq C \|f\|_{L^2(|\xi| \sim 2^k)}.$$

C.2. Proof of (5.13)

For the proof of (5.13), recall the representation formula (2.11). We will reduce to the same expressions as above. With the partition of unity displayed above, write

$$\begin{aligned} [\partial_1 |\nabla|^{-a}, |\xi|^2] f(\xi) &= c_a \sum_{k=-\infty}^{\infty} 2^{2k} [\partial_1 |\nabla|^{-a}, \psi(2^{-k} \cdot)] f \\ &= c_a \sum_{k=-\infty}^{\infty} 2^{2k} \left[\partial_{\xi_1} \int_{\mathbf{R}^2} \frac{\psi(2^{-k} y) f(y)}{|\xi - y|^{2-a}} dy \right. \\ &\quad \left. - \psi(2^{-k} \xi) \partial_{\xi_1} \int_{\mathbf{R}^2} \frac{f(y)}{|\xi - y|^{2-a}} dy \right] \\ &= c_a (a-2) \sum_{k=-\infty}^{\infty} 2^{2k} \int_{\mathbf{R}^2} \frac{\xi_1 - y_1}{|\xi - y|} \frac{(\psi(2^{-k} y) - \psi(2^{-k} \xi)) f(y)}{|\xi - y|^{2-a}} dy \end{aligned}$$

Taking absolute values and estimating yields the bound

$$|[\partial_1 |\nabla|^{-a}, |\xi|^2] f(\xi)| \leq C_a \sum_{k=-\infty}^{\infty} 2^{2k} \int_{\mathbf{R}^2} \frac{|\psi(2^{-k} y) - \psi(2^{-k} \xi)| |f(y)|}{|\xi - y|^{3-a}} dy$$

This is of course exactly the same expression as before for the F_k , with $a := 1 - \frac{\alpha}{2}$. Therefore, we can apply the same estimates to obtain

$$\|[\partial_1 |\nabla|^{-a}, |\xi|^2] f\|_{L^2(\mathbf{R}^2)} \leq C \||\xi|^{1+a} f\|_{L^2}.$$

This establishes (5.13).

Appendix D. Semigroup Estimates: Proposition (3)

Proof We have

$$\begin{aligned} \|\partial^\gamma (e^{\tau \mathcal{L}} f)\|_{L^2(2)}^2 &\leq \int_{\mathbf{R}^2} |\partial^\gamma (e^{\tau \mathcal{L}} f)|^2 d\xi + \int_{\mathbf{R}^2} ||\xi|^2 \partial^\gamma (e^{\tau \mathcal{L}} f)|^2 d\xi \\ &= e^{2(1-\frac{3-\beta}{\alpha})\tau} \left(\int_{\mathbf{R}^2} |p^\gamma [e^{-a(\tau)|p|^\alpha} \widehat{f}(pe^{-\frac{\tau}{\alpha}})]|^2 dp + \int_{\mathbf{R}^2} |\Delta_p [p^\gamma e^{-a(\tau)|p|^\alpha} \widehat{f}(pe^{-\frac{\tau}{\alpha}})]|^2 dp \right) \\ &= e^{2(1-\frac{3-\beta}{\alpha})\tau} \left(\int_{\mathbf{R}^2} |p^\gamma [e^{-a(\tau)|p|^\alpha} \widehat{f}(pe^{-\frac{\tau}{\alpha}})]|^2 dp + \gamma^2 \int_{\mathbf{R}^2} |p^{|\gamma|-1} \nabla_p [e^{-a(\tau)|p|^\alpha} \widehat{f}(pe^{-\frac{\tau}{\alpha}})]|^2 dp \right. \\ &\quad \left. + \int_{\mathbf{R}^2} |p^\gamma \Delta_p [e^{-a(\tau)|p|^\alpha} \widehat{f}(pe^{-\frac{\tau}{\alpha}})]|^2 dp \right). \end{aligned}$$

At this point, it is clear that it is better to divide both sides by $e^{2(1-\frac{3-\beta}{\alpha})\tau}$. Then, we want to control the right-hand side of the following

$$\begin{aligned} \frac{\|\partial^\gamma (e^{\tau \mathcal{L}} f)\|_{L^2(2)}^2}{e^{2(1-\frac{3-\beta}{\alpha})\tau}} &\leq \int_{\mathbf{R}^2} |p^\gamma [e^{-a(\tau)|p|^\alpha} \widehat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp \\ &\quad + \gamma^2 \int_{\mathbf{R}^2} |p|^{|\gamma|-1} \nabla_p [e^{-a(\tau)|p|^\alpha} \widehat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp \\ &\quad + \int_{\mathbf{R}^2} |p^\gamma \Delta_p [e^{-a(\tau)|p|^\alpha} \widehat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp := J_1 + J_2 + J_3. \end{aligned}$$

□

D.1. Estimate for J_1

To control J_1 we divide the argument into two different cases, $\tau \leq 1$ and $\tau > 1$. In the case of $\tau \leq 1$, we have

$$\begin{aligned} J_1 &= \int_{\mathbf{R}^2} |p^\gamma [e^{-a(\tau)|p|^\alpha} \widehat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp = \int_{\mathbf{R}^2} |q|^{2|\gamma|} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\widehat{f}(q)|^2 dq \\ &\leq \int_{\{q: 0 \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} |q|^{2|\gamma|} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\widehat{f}(q)|^2 dq \\ &\quad + \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} |q|^{2|\gamma|} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\widehat{f}(q)|^2 dq \\ &= J_1^1 + J_1^2. \end{aligned}$$

We can estimate, since $\tau \leq 1$,

$$J_1^1 \leq \int_{0 \leq |q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2|\gamma|} |\widehat{f}(q)|^2 dq \leq \frac{1}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2}^2 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2.$$

We treat J_1^2 in a similar manner. Indeed, again for $\tau \leq 1$,

$$\begin{aligned} J_1^2 &\leq \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |q|^{2|\gamma|} |\widehat{f}(q)|^2 dq \\ &\leq \frac{C}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \sum_{j=1}^{\infty} e^{-(j+1)} (j+1)^{\frac{2|\gamma|}{\alpha}} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |\widehat{f}|^2 dq \end{aligned}$$

$$\leq \frac{C}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2}^2 \sum_{j=1}^{\infty} e^{-(j+1)} (j+1)^{\frac{2|\gamma|}{\alpha}} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2$$

After putting together the estimates for J_1^1 and J_1^2 , we get $J_1 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}$.

Regarding the case $\tau > 1$, first note that in this range of τ , $a(\tau) \geq \frac{1}{2}$. Moreover,

$$|\hat{f}(q) - \hat{f}(0)| \leq 2\|\hat{f}\|_{L^\infty}, \quad |\hat{f}(q) - \hat{f}(0)| \leq |q| \|\nabla \hat{f}\|_{L^\infty},$$

whence by interpolation, we conclude that for every $\epsilon > 0$, we have

$$|\hat{f}(q) - \hat{f}(0)| \leq C_\epsilon |q|^{1-\epsilon} \|\nabla^{1-\epsilon} \hat{f}\|_{L^\infty} \leq C_\epsilon |q|^{1-\epsilon} \|f\|_{L^2(2)}, \quad (\text{D.1})$$

where in the last inequality we have used that by Hausdorff–Young's $\|\nabla^{1-\epsilon} \hat{f}\|_{L^\infty} \leq \int_{\mathbf{R}^2} |\xi|^{1-\epsilon} |\hat{f}(\xi)| d\xi \lesssim \|f\|_{L^2(2)}$. Therefore,

$$\begin{aligned} J_1 &= \int_{\mathbf{R}^2} |p^\gamma [e^{-a(\tau)|p|^\alpha} \hat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp = e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \int_{\mathbf{R}^2} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2|\gamma|} |\hat{f}(q)|^2 dq \\ &\leq e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \int_{\mathbf{R}^2} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(|\gamma|+1-\epsilon)} dq \\ &\leq e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(|\gamma|+1-\epsilon)} dq \\ &\quad + e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(|\gamma|+1-\epsilon)} dq = J_1^1 + J_1^2. \end{aligned}$$

Now

$$\begin{aligned} J_1^1 &= e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(|\gamma|+1-\epsilon)} dq \\ &\leq e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} |q|^{2(|\gamma|+1-\epsilon)} dq \leq \\ &\leq C e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \int_0^{\frac{1}{(2a(\tau))^{\frac{1}{\alpha}}}} r^{2(|\gamma|+1-\epsilon)+1} dq \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2}{\alpha}(|\gamma|+2-\epsilon)}} \\ &\leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}. \end{aligned}$$

In a similar way,

$$\begin{aligned}
J_1^2 &= e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq j+1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha}} |q|^{2(|\gamma|+1-\epsilon)} dq \\
&\leq e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq j+1\}} |q|^{2(|\gamma|+1-\epsilon)} dq \\
&\leq e^{\frac{2\tau}{\alpha}(|\gamma|+1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \int_{(\frac{j}{a(\tau)})^{\frac{1}{\alpha}} e^{-\frac{\tau}{\alpha}}}^{(\frac{j+1}{a(\tau)})^{\frac{1}{\alpha}} e^{-\frac{\tau}{\alpha}}} r^{2(|\gamma|+1-\epsilon)+1} dr \leq \\
&\leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{a(\tau)^{\frac{2}{\alpha}(|\gamma|+2-\epsilon)}} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} (j+1)^{2(|\gamma|+2-\epsilon)} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}.
\end{aligned}$$

Therefore for $\tau > 1$ we have $J_1 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}$.

D.2. Estimate for J_2

To control J_2 first note that

$$\nabla e^{-a(\tau)|p|^{\alpha}} = -\alpha a(\tau) p |p|^{\alpha-2} e^{-a(\tau)|p|^{\alpha}}. \quad (\text{D.2})$$

Therefore,

$$\begin{aligned}
J_2 &\leq \alpha^2 |\gamma|^2 a(\tau)^2 \int_{\mathbf{R}^2} |p|^{\gamma|-1} |p|^{\alpha-1} e^{-a(\tau)|p|^{\alpha}} |\widehat{f}(p e^{\frac{-\tau}{\alpha}})|^2 dp \\
&\quad + |\gamma|^2 e^{\frac{-2\tau}{\alpha}} \int_{\mathbf{R}^2} |p|^{\gamma|} e^{-a(\tau)|p|^{\alpha}} \cdot (\nabla \widehat{f})(p e^{\frac{-\tau}{\alpha}})|^2 dp := I_1 + I_2.
\end{aligned}$$

D.2.1. Estimate for I_1

To control the first term I_1 we proceed as follows

$$\begin{aligned}
\frac{I_1}{a(\tau)^2} &\leq \int_{\mathbf{R}^2} e^{-2a(\tau)|p|^{\alpha}} |p|^{2(\alpha+|\gamma|-2)} |\widehat{f}(p \cdot e^{-\frac{\tau}{\alpha}})|^2 dp \\
&= e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{\mathbf{R}^2} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha}} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq \\
&\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq 1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha}} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq \\
&\quad + e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq j+1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha}} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq
\end{aligned}$$

$$= I_1^1 + I_1^2.$$

We can estimate

$$\begin{aligned} I_1^1 &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{|q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2(\alpha+|\gamma|-2)} |\hat{f}(q)|^2 dq \\ &= e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{|q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2(\alpha+|\gamma|-2)} |\hat{f}(q) - \hat{f}(0)|^2 dq \end{aligned}$$

Using the relation (D.1), we obtain

$$\begin{aligned} I_1^1 &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|f\|_{L^2(2)}^2 \int_{|q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2(\alpha+|\gamma|-2)} |q|^{2(1-\epsilon)} dq \\ &= C e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|f\|_{L^2(2)}^2 \int_0^{\frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} r^{2(\alpha+|\gamma|-1-\epsilon)-1} dr = C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{2(1+\frac{|\gamma|}{\alpha}-\epsilon)}}. \end{aligned}$$

therefore, recalling that $a(\tau) \leq 1$, $I_1^1 \leq \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{2(1+\frac{|\gamma|}{\alpha})}}$.

We treat I_1^2 in a similar manner. Again, using (D.1),

$$\begin{aligned} I_1^2 &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{2(\alpha+|\gamma|-2)} |\hat{f}(q) - \hat{f}(0)|^2 dq \\ &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{2(\alpha+|\gamma|-2)} |q|^{2(1-\epsilon)} \|f\|_{L^2(2)}^2 dq \\ &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{2(\alpha+|\gamma|-1-\epsilon)} dq \\ &\leq C \frac{e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|f\|_{L^2(2)}^2}{a(\tau)^{2(1+\frac{|\gamma|}{\alpha})}} \sum_{j=1}^{\infty} e^{-j} j^{\frac{2(\alpha+|\gamma|-1-\epsilon)}{\alpha}} e^{-\frac{2\tau}{\alpha}(\alpha+|\gamma|-1-\epsilon)} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{2(1+\frac{|\gamma|}{\alpha})}}. \end{aligned}$$

After putting together the estimates for I_1^1 and I_1^2 we get $I_1 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}$.

D.2.2. Estimate for I_2

$$\begin{aligned}
I_2 &\leq C e^{-\frac{2\tau}{\alpha}} \int_{\mathbf{R}^2} | |p|^{|\gamma|-1} e^{-a(\tau)|p|^\alpha} (\nabla \hat{f})(p e^{\frac{-\tau}{\alpha}}) |^2 dp \\
&= e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \int_{\mathbf{R}^2} |q|^{2(|\gamma|-1)} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\nabla \hat{f}(q)|^2 dq \\
&\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \sum_{j=0}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} \left(|q|^{2(|\gamma|-1)} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\nabla \hat{f}(q)|^2 dq \right) \\
&= I_2^1 + I_2^2.
\end{aligned}$$

For I_2^1 , we have by Hölder's

$$\begin{aligned}
I_2^1 &\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \int_{0 \leq |q| \leq \frac{e^{\frac{-\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2(|\gamma|-1)} |\nabla \hat{f}(q)|^2 dq \\
&\leq C e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \left(\int_{0 \leq |r| \leq \frac{e^{\frac{-\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} r^{\frac{2(|\gamma|-1)}{1-\epsilon}+1} dr \right)^{1-\epsilon} \\
&= C e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \frac{e^{-\frac{2\tau}{\alpha}(|\gamma|-1)-\frac{2\tau}{\alpha}(1-\epsilon)}}{(2a(\tau))^{\frac{2|\gamma|}{\alpha}-\frac{2\epsilon}{\alpha}}} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2.
\end{aligned}$$

By Sobolev embedding, we have $\|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \leq C \|\nabla \hat{f}\|_{H^{1-\epsilon}(\mathbf{R}^2)}^2 \leq C \|(1-\Delta) \hat{f}\|_{L^2}^2 = C \|f\|_{L^2(2)}^2$. Therefore

$$I_2^1 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2.$$

For I_2^2 , we estimate

$$\begin{aligned}
I_2^2 &\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |q|^{2(|\gamma|-1)} |\nabla \hat{f}|^2 dq \\
&\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \sum_{j=1}^{\infty} e^{-j} \left[\int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |q|^{\frac{2(|\gamma|-1)}{1-\epsilon}} dq \right]^{1-\epsilon}.
\end{aligned}$$

But,

$$\int_{j \leq 2a(\tau)|q e^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |q|^{\frac{2(|\gamma|-1)}{1-\epsilon}} dq \leq C \left(\frac{j}{a(\tau)} \right)^{\frac{2(|\gamma|-1)}{1-\epsilon}+2} e^{-\frac{2\tau}{\alpha}(|\gamma|-1)-\frac{2\tau}{\alpha}},$$

so using again the bound $\|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}} \leq C\|f\|_{L^2(2)}$,

$$\begin{aligned} I_2^2 &\leq C e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \left[\left(\frac{j}{a(\tau)} \right)^{\frac{2(|\gamma|-1)}{1-\epsilon}+2} e^{\frac{-2\tau}{\alpha}(\alpha+|\gamma|-1)-\frac{2\tau}{\alpha}} \right]^{1-\epsilon} \\ &\leq C \frac{e^{-\frac{2\tau(1-\epsilon)}{\alpha}} \|f\|_{L^2(2)}^2}{a(\tau)^{2\frac{|\gamma|}{\alpha}-2\epsilon}} \sum_{j=1}^{\infty} e^{-j} j^{2\alpha+2|\gamma|-2\epsilon} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}. \end{aligned}$$

Hence after putting together the estimates for I_2^1 and I_2^2 we have $I_2 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}}$.

D.3. Estimate for J_3

$$\begin{aligned} J_3 &= \int_{\mathbf{R}^2} |p^{|\gamma|} \Delta_p [e^{-a(\tau)|p|^\alpha} \hat{f}(p e^{-\frac{\tau}{\alpha}})]|^2 dp \leq \int_{\mathbf{R}^2} |p^{|\gamma|} \Delta_p [e^{-a(\tau)|p|^\alpha}] \hat{f}(p e^{-\frac{\tau}{\alpha}})|^2 dp \\ &\quad + 2 \int_{\mathbf{R}^2} |p^{|\gamma|} \nabla e^{-a(\tau)|p|^\alpha} \cdot \nabla (\hat{f}(p e^{-\frac{\tau}{\alpha}}))|^2 dp + \int_{\mathbf{R}^2} |p^{|\gamma|} e^{-a(\tau)|p|^\alpha} \Delta_p (\hat{f}(p e^{-\frac{\tau}{\alpha}}))|^2 dp. \end{aligned}$$

By (D.2) we have,

$$\begin{aligned} \Delta_p [e^{-a(\tau)|p|^\alpha}] &= \sum_{j=1}^2 \partial_j \left(-\alpha a(\tau) p_j |p|^{\alpha-2} e^{-a(\tau)|p|^\alpha} \right) \\ &= -\alpha a(\tau) \sum_{j=1}^2 \left(|p|^{\alpha-2} + (\alpha-2) \frac{p_j^2}{|p|} |p|^{\alpha-3} + p_j |p|^{\alpha-2} (-\alpha a(\tau)) \frac{p_j}{|p|} |p|^{\alpha-1} \right) e^{-a(\tau)|p|^\alpha} \\ &= \left(-\alpha^2 a(\tau) |p|^{\alpha-2} + \alpha^2 a(\tau)^2 |p|^{2(\alpha-1)} \right) e^{-a(\tau)|p|^\alpha}. \end{aligned}$$

Hence, by allowing for a slight abuse of notations by using γ , which is a multi-index instead of $|\gamma|$, its length,

$$\begin{aligned} J_3 &\lesssim a(\tau)^2 \int_{\mathbf{R}^2} | |p|^{\alpha+|\gamma|-2} e^{-a(\tau)|p|^\alpha} \hat{f}(p e^{-\frac{\tau}{\alpha}}) |^2 dp \\ &\quad + a(\tau)^4 \int_{\mathbf{R}^2} | |p|^{2(\alpha-1)+|\gamma|} e^{-a(\tau)|p|^\alpha} \hat{f}(p e^{-\frac{\tau}{\alpha}}) |^2 dp \\ &\quad + a(\tau)^2 e^{-\frac{2\tau}{\alpha}} \int_{\mathbf{R}^2} | |p|^{\alpha+|\gamma|-1} e^{-a(\tau)|p|^\alpha} (\nabla \hat{f})(p e^{-\frac{\tau}{\alpha}}) |^2 dp \\ &\quad + e^{-\frac{4\tau}{\alpha}} \int_{\mathbf{R}^2} | |p|^\gamma e^{-a(\tau)|p|^\alpha} (\Delta \hat{f})(p e^{-\frac{\tau}{\alpha}}) |^2 dp \\ &:= I_3 + I_4 + I_5 + I_6, \end{aligned}$$

D.3.1. Estimate for I_3 and I_4

$$\begin{aligned}
\frac{I_3}{a(\tau)^2} &\leq \int_{\mathbf{R}^2} e^{-2a(\tau)|p|^\alpha} |p|^{2(\alpha+|\gamma|-2)} |\widehat{f}(p \cdot e^{-\frac{\tau}{\alpha}})|^2 dp \\
&= e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{\mathbf{R}^2} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq \\
&\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq \\
&\quad + e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{2(\alpha+|\gamma|-2)} |\widehat{f}(q)|^2 dq \\
&= I_3^1 + I_3^2.
\end{aligned}$$

By comparing I_3 with I_1 it is clear that $I_3^1 = I_1^1$ and $I_3^2 = I_1^2$, and we treat them in the same way. Hence $I_3 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{\frac{2|\gamma|}{\alpha}}}$.

The estimate for I_4 proceeds in an identical manner, but we have a slightly different power of p , so we present it here briefly.

$$\begin{aligned}
\frac{I_4}{a(\tau)^4} &\leq \int_{\mathbf{R}^2} | |p|^{2(\alpha-1)+|\gamma|} e^{-a(\tau)|p|^\alpha} \widehat{f}(p e^{-\frac{\tau}{\alpha}})|^2 dp \\
&= e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \sum_{j=0}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} \left(e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |q|^{4(\alpha-1)+2|\gamma|} |\widehat{f}(q)|^2 dq \right) \\
&:= I_4^2 + I_4^1.
\end{aligned}$$

Denoting by I_4^1 the integral corresponding to $2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1$ and the rest with I_4^2 , we have by (D.1), $|\widehat{f}(q)| = |\widehat{f}(q) - \widehat{f}(0)| \leq C|q|^{1-\epsilon} \|f\|_{L^2(2)}$,

$$\begin{aligned}
I_4^1 &\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} |q|^{4(\alpha-1)+2|\gamma|} |\widehat{f}(q)|^2 dq \\
&\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \|f\|_{L^2(2)}^2 \int_{\{q: 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq 1\}} |q|^{4(\alpha-1)+2|\gamma|+2(1-\epsilon)} dq \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{(a(\tau))^{4+\frac{2|\gamma|}{\alpha}}}.
\end{aligned}$$

For I_4^2 , we have

$$\begin{aligned}
I_4^2 &\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} |q|^{4(\alpha-1)+2|\gamma|} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\widehat{f}(q)|^2 dq \\
&\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \sum_{j=1}^{\infty} e^{-j} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha \leq j+1\}} |q|^{4(\alpha-1)+2|\gamma|} |\widehat{f}(q) - \widehat{f}(0)|^2 dq
\end{aligned}$$

$$\begin{aligned} &\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq j+1\}} |q|^{4(\alpha-1)+2|\gamma|} |q|^{2(1-\epsilon)} dq \\ &\leq e^{\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2)} \|f\|_{L^2(2)}^2 \frac{e^{-\frac{\tau}{\alpha}(4\alpha+2|\gamma|-2\epsilon)}}{(a(\tau))^{4+\frac{2|\gamma|}{\alpha}-\frac{2\epsilon}{\alpha}}} \sum_{j=1}^{\infty} e^{-j} j^{2(2\alpha+|\gamma|-\epsilon)} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{4+\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2. \end{aligned}$$

$$\text{Therefore } I_4 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}}.$$

D.4. Estimate for I_5

$$\begin{aligned} \frac{I_5}{a(\tau)^2} &\leq C e^{-\frac{2\tau}{\alpha}} \int_{\mathbf{R}^2} |p|^{\alpha+|\gamma|-1} e^{-a(\tau)|p|^\alpha} (\nabla \hat{f})(p e^{\frac{-\tau}{\alpha}})^2 dp \\ &= e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{\mathbf{R}^2} |q|^{2(\alpha+|\gamma|-1)} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\nabla \hat{f}(q)|^2 dq \\ &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=0}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq j+1\}} \left(|q|^{2(\alpha+|\gamma|-1)} e^{-2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^\alpha} |\nabla \hat{f}(q)|^2 dq \right) \\ &= I_5^1 + I_5^2. \end{aligned}$$

For I_5^1 , we have by Hölder's

$$\begin{aligned} I_5^1 &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \int_{|q| \leq \frac{e^{\frac{-\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2(\alpha+|\gamma|-1)} |\nabla \hat{f}(q)|^2 dq \\ &= C e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \frac{e^{-\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)-\frac{2\tau}{\alpha}(1-\epsilon)}}{(2a(\tau))^{2+\frac{2|\gamma|}{\alpha}-\frac{2\epsilon}{\alpha}}} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{2+\frac{2|\gamma|}{\alpha}}} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2. \end{aligned}$$

However, by Sobolev embedding, we have as before $\|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \leq C \|f\|_{L^2(2)}^2$.

For I_5^2 , we estimate

$$\begin{aligned} I_5^2 &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{2(\alpha+|\gamma|-1)} |\nabla \hat{f}|^2 dq \\ &\leq e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}}^2 \sum_{j=1}^{\infty} e^{-j} \left[\int_{j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{\frac{2(\alpha+|\gamma|-1)}{1-\epsilon}} dq \right]^{1-\epsilon}. \end{aligned}$$

But,

$$\int_{j \leq 2a(\tau)|q \cdot e^{\frac{\tau}{\alpha}}|^{\alpha} \leq (j+1)} |q|^{\frac{2(\alpha+|\gamma|-1)}{1-\epsilon}} dq \leq C \left(\frac{j}{a(\tau)} \right)^{\frac{2(\alpha+|\gamma|-1)}{1-\epsilon}+2} e^{\frac{-2\tau}{\alpha}(\alpha+|\gamma|-1)-\frac{2\tau}{\alpha}},$$

so using again the bound $\|\nabla \hat{f}\|_{L^{\frac{2}{\epsilon}}} \leq C \|f\|_{L^2(2)}$,

$$\begin{aligned}
I_5^2 &\leq C e^{\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)} \|f\|_{L^2(2)}^2 \sum_{j=1}^{\infty} e^{-j} \left[\left(\frac{j}{a(\tau)} \right)^{\frac{2(\alpha+|\gamma|-1)}{1-\epsilon}+2} e^{-\frac{2\tau}{\alpha}(\alpha+|\gamma|-1)-\frac{2\tau}{\alpha}} \right]^{1-\epsilon} \\
&\leq C \frac{e^{-\frac{2\tau(1-\epsilon)}{\alpha}} \|f\|_{L^2(2)}^2}{a(\tau)^{2+2\frac{|\gamma|}{\alpha}-2\epsilon}} \sum_{j=1}^{\infty} e^{-j} j^{2\alpha+2|\gamma|-2\epsilon} \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{a(\tau)^{2+\frac{2|\gamma|}{\alpha}}}.
\end{aligned}$$

Hence after putting together the estimates for I_5^1 and I_5^2 we have

$$I_5 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)} \|f\|_{L^2(2)}^2}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}}.$$

D.5. Estimate for I_6

In the same way, we can get the estimate for I_6 . Indeed,

$$\begin{aligned}
I_6 &\leq e^{-\frac{4\tau}{\alpha}} \int_{\mathbf{R}^2} \left| |p|^{\gamma} e^{-a(\tau)|p|^\alpha} (\Delta \hat{f})(p e^{-\frac{\tau}{\alpha}}) \right|^2 dp = e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \int_{\mathbf{R}^2} |q|^{2|\gamma|} e^{-2a(\tau)|q \cdot \mathbf{e}^{\frac{\tau}{\alpha}}|^\alpha} |\Delta \hat{f}(q)|^2 dq \\
&\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \left[\int_{\{q: 2a(\tau)|q \cdot \mathbf{e}^{\frac{\tau}{\alpha}}| \leq 1\}} + \sum_{j=1}^{\infty} \int_{\{q: j \leq 2a(\tau)|q \cdot \mathbf{e}^{\frac{\tau}{\alpha}}| \leq j+1\}} \right] \left(|q|^{2|\gamma|} e^{-2a(\tau)|q \cdot \mathbf{e}^{\frac{\tau}{\alpha}}|^\alpha} |\Delta \hat{f}(q)|^2 dq \right) \\
&= I_6^1 + I_6^2.
\end{aligned}$$

For I_6^1 ,

$$\begin{aligned}
I_6^1 &\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \int_{|q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |q|^{2|\gamma|} |\Delta \hat{f}(q)|^2 dq = \frac{e^{\frac{2\tau}{\alpha}(|\gamma|-1)} e^{-\frac{2\tau|\gamma|}{\alpha}}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \int_{|q| \leq \frac{e^{-\frac{\tau}{\alpha}}}{(2a(\tau))^{\frac{1}{\alpha}}}} |\Delta \hat{f}(q)|^2 dq \\
&\leq C \frac{e^{-\frac{2\tau}{\alpha}}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2.
\end{aligned}$$

For I_6^2 , we have

$$\begin{aligned}
I_6^2 &\leq e^{\frac{2\tau}{\alpha}(|\gamma|-1)} \sum_{j=1}^{\infty} e^{-j} \int_{j \leq 2a(\tau)|q \cdot \mathbf{e}^{\frac{\tau}{\alpha}}|^\alpha \leq (j+1)} |q|^{2|\gamma|} |\Delta \hat{f}(q)|^2 dq \\
&\leq \frac{e^{\frac{2\tau}{\alpha}(|\gamma|-1)} e^{-\frac{2\tau|\gamma|}{\alpha}}}{a(\tau)^{\frac{2|\gamma|}{\alpha}}} \sum_{j=1}^{\infty} e^{-j} (j+1)^{\frac{2|\gamma|}{\alpha}} \int |\Delta \hat{f}(q)|^2 dq \leq C \frac{e^{-\frac{2\tau}{\alpha}}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2.
\end{aligned}$$

Therefore,

$$I_6 \leq C \frac{e^{-\frac{2\tau}{\alpha}}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2 \leq C \frac{e^{-\frac{2\tau}{\alpha}(1-\epsilon)}}{(a(\tau))^{\frac{2|\gamma|}{\alpha}}} \|f\|_{L^2(2)}^2.$$

Putting it all together finishes off the proof. \square

References

Abidi, H., Hmidi, T.: On the global well posedness for Boussinesq system. *J. Differ. Equ.* **233**(1), 199–220 (2007)

Ben-Artzi, M.: Global solutions of two-dimensional Navier–Stokes and Euler equations. *Arch. Ration. Mech. Anal.* **128**(4), 329–358 (1994)

Brandoles, L., Schonbek, M.E.: Large time decay and growth for solutions of a viscous Boussinesq system. *Trans. Am. Math. Soc.* **364**(10), 5057–5090 (2012)

Carpio, A.: Asymptotic behavior for the vorticity equations in dimensions two and three. *Commun. Partial Differ. Equ.* **19**(5–6), 827–872 (1994)

Chae, D.: Global regularity for the 2D Boussinesq equations with partial viscosity terms. *Adv. Math.* **203**, 497–513 (2006)

Chamorro, D., Lemarié-Rieusset, P.G.: Quasi-geostrophic equation, nonlinear Bernstein inequalities and α -stable processes. *Rev. Mat. Iberoam.* **28**, 1109–1122 (2012)

Constantin, P., Wu, J.: Behavior of solutions of 2D quasi-geostrophic equations. *SIAM J. Math. Anal.* **30**(5), 937–948 (1999)

Cordoba, A., Cordoba, D.: A maximum principle applied to quasi-geostrophic equations. *Commun. Math. Phys.* **249**(3), 511–528 (2004)

Gallay, T., Wayne, C.E.: Invariant manifolds and long-time asymptotics of the Navier–Stokes and vorticity equations on \mathbf{R}^2 . *Arch. Ration. Mech. Anal.* **163**(3), 209–258 (2002a)

Gallay, T., Wayne, C.E.: Long-time asymptotics of the Navier–Stokes and vorticity equations on \mathbf{R}^3 . *Philos. Trans. R. Soc. Lond.* **360**, 2155–2188 (2002b)

Giga, Y., Kambe, T.: Large time behavior of the vorticity of two-dimensional viscous flow and its application to vortex formation. *Commun. Math. Phys.* **117**(4), 549–568 (1988)

Goh, R., Wayne, E.G.: Vortices in stably-stratified rapidly rotating Boussinesq convection (2018). [arXiv:1802.05369](https://arxiv.org/abs/1802.05369)

Hadadifard, F., Stefanov, A.: On the global regularity of the 2D critical Boussinesq system with $\alpha > \frac{2}{3}$. *Commun. Math. Sci.* **15**(5), 1325–1351 (2017)

Jiu, Q., Miao, C., Wu, J., Zhang, Z.: The two-dimensional incompressible Boussinesq equations with general critical dissipation. *SIAM J. Math. Anal.* **46**(5), 3426–3454 (2014)

Miyakawa, T., Schonbek, M.: On optimal decay rates for weak solutions to the Navier–Stokes equations in \mathbf{R}^n . In: *Proceedings of partial differential equations and applications* (Olomouc, 1999), vol. 126, pp. 443–455 (2001)

Niche, C., Schonbek, M.E.: Decay of weak solutions to the 2D dissipative quasi-geostrophic equation. *Commun. Math. Phys.* **276**(1), 93–115 (2007)

Schonbek, M.E.: L^2 decay for weak solutions of the Navier–Stokes equations. *Arch. Ration. Mech. Anal.* **88**(3), 209–222 (1985)

Schonbek, M.E.: Lower bounds of rates of decay for solutions to the Navier–Stokes equations. *J. Am. Math. Soc.* **4**, 423–449 (1991)

Schonbek, M.E.: Asymptotic behavior of solutions to the three-dimensional Navier–Stokes equations. *Indiana Univ. Math. J.* **41**, 809–823 (1992)

Schonbek, M., Schonbek, T.: On the boundedness and decay of moments of solutions to the Navier–Stokes equations. *Adv. Differ. Equ.* **5**(7–9), 861–898 (2000)

Schonbek, M., Wiegner, M.: On the decay of higher-order norms of the solutions of Navier–Stokes equations. *Proc. R. Soc. Edinburgh Sect. A* **126**(3), 677–685 (1996)

Wu, J.: The 2D incompressible Boussinesq equations. *Peking University Summer School Lecture notes* (2012)

Yang, J.: Large time decay of solutions to the Boussinesq system with fractional dissipation. *J. Math. Anal. Appl.* **453**(1), 607–619 (2017)

Yang, W., Jiu, Q., Wu, J.: Global well-posedness for a class of 2D Boussinesq systems with fractional dissipation. *J. Differ. Equ.* **257**(11), 4188–4213 (2014)