

Research



Cite this article: Tobasco I. 2022 Optimal cooling of an internally heated disc. *Phil. Trans. R. Soc. A* **380**: 20210040.
<https://doi.org/10.1098/rsta.2021.0040>

Received: 25 October 2021

Accepted: 22 January 2022

One contribution of 14 to a theme issue
'Mathematical problems in physical fluid
dynamics (part 1)'.

Subject Areas:

applied mathematics, analysis, differential
equations, fluid mechanics

Keywords:

heat transfer, convective cooling, optimal
design, variational bounds, branching flows

Author for correspondence:

Ian Tobasco

e-mail: itobasco@uic.edu

Dedication: To Charlie Doering, who taught us
how to bound.

Optimal cooling of an internally heated disc

Ian Tobasco

Department of Mathematics, Statistics, and Computer Science,
University of Illinois at Chicago, Chicago, IL 60607, USA

IT, 0000-0003-3618-8036

Motivated by the search for sharp bounds on turbulent heat transfer as well as the design of optimal heat exchangers, we consider incompressible flows that most efficiently cool an internally heated disc. Heat enters via a distributed source, is passively advected and diffused, and exits through the boundary at a fixed temperature. We seek an advecting flow to optimize this exchange. Previous work on energy-constrained cooling with a constant source has conjectured that global optimizers should resemble convection rolls; we prove one-sided bounds on energy-constrained cooling corresponding to, but not resolving, this conjecture. In the case of an enstrophy constraint, our results are more complete: we construct a family of self-similar, tree-like 'branching flows' whose cooling we prove is within a logarithm of globally optimal. These results hold for general space- and time-dependent source-sink distributions that add more heat than they remove. Our main technical tool is a non-local Dirichlet-like variational principle for bounding solutions of the inhomogeneous advection-diffusion equation with a divergence-free velocity.

This article is part of the theme issue 'Mathematical problems in physical fluid dynamics (part 1)'.

1. Introduction

Imagine a fluid in a container that is heated from within, and whose temperature is fixed at its boundary. How should the fluid flow so that the container cools as quickly as possible? This question arises, for instance, in the design of optimal heat exchangers, whose complicated shapes and flows facilitate heat transfer at rates far beyond diffusion [1–10]. More generally, the problem is related to the ongoing search for sharp bounds on turbulent heat transfer in a variety

of settings, including internally heated [11–16] as well as buoyancy-driven convection [17–21], among others. While any bound must explain why the rate of heat transfer associated with a given flow cannot exceed a certain amount, sharp bounds are distinguished in that their values are attained by particular flows—flows that, as a result, are extremal. Finding sharp bounds on convective heat transfer, or at least bounds that are asymptotically sharp in their scaling with respect to physical parameters, has remained a widely open problem in our understanding of turbulence since the pioneering works of Howard & Busse [22,23] and Doering & Constantin [24] on Rayleigh–Bénard convection (for a recent summary of the state of the art, see [25]).

A widely used technique for proving bounds on fluid dynamical quantities is the ‘background method’ introduced by Doering & Constantin [24,26,27], which takes the form of a convex variational problem whose optimal value is the bound. Whether or not this method and its descendants (see e.g. [28,29] or [30] for a recent review) produce sharp bounds on convection remains largely unclear. This has led some to ask whether information beyond the background method might be used, e.g. via the non-quadratic auxiliary functionals of [31–34], the conjecture that global optimizers of certain chaotic evolutions are steady [35], or the maximum principle which is known to improve some bounds [36,37]. Recently, progress with internally heated convection has led to asymptotically sharp bounds on heat transfer between a pair of steady and perfectly balanced sources and sinks [38]. Due to the balanced nature of the source–sink distribution, which puts in just as much heat as it takes away, the overall rate of heat transfer in the bound turns out to be much larger than the usual ‘ultimate scaling’ law; it is nonetheless asymptotically sharp and is saturated by a bulk, convection roll-like flow.

Motivated by these questions, we consider the design of incompressible flows to optimally cool an internally heated disc. We treat a general space- and time-dependent source–sink distribution, under an assumption that more heat is added than is removed. This ensures that some heat must make its way from the bulk to the boundary, and turns out to drive the formation of fine flow microstructure upon optimization (as shown in the upcoming figure 1). Although temperature will be governed in our setup by the usual advection–diffusion equation, we will not impose a momentum equation directly on the velocity, but will instead replace it with a constraint on the mean enstrophy it implies. Optimizing with respect to this constraint leads to a self-similar, tree-like ‘branching flow’ whose cooling we prove is within a logarithm of globally optimal. While the *a priori* bound part of this result can be shown using the background method, we prefer a different approach that highlights the variational structure of the advection–diffusion equation. We also comment on the related energy-constrained problem, for which considerably less is known.

Setting up these problems in detail, we let $D \subset \mathbb{R}^2$ be a disc with radius $R > 0$ and consider a temperature field $T(\mathbf{x}, t)$ that is passively advected and diffused according to a divergence-free velocity field $\mathbf{u}(\mathbf{x}, t)$ we take to be under our control. At the disc’s boundary ∂D we set the temperature to be a constant, say $T = 0$, and let $f(\mathbf{x}, t)$ be a given source–sink distribution. Altogether, T solves the inhomogeneous advection–diffusion equation

$$\begin{cases} \partial_t T + \mathbf{u} \cdot \nabla T = \kappa \Delta T + f & \text{in } D \\ T = 0 & \text{at } \partial D \end{cases} \quad (1.1)$$

where $\kappa > 0$ is the thermal diffusivity. We leave the value of T at the initial time $t = 0$ unspecified, as in the long run it is lost to diffusion. To measure cooling efficiency, we use the mean-square temperature gradient¹

$$\langle |\nabla T|^2 \rangle = \lim_{\tau \rightarrow \infty} \frac{1}{\tau |D|} \int_0^\tau \int_D |\nabla T(\mathbf{x}, t)|^2 \, d\mathbf{x} \, dt,$$

whose value depends on $\mathbf{u}$. The notation $|D|$ stands for the area of the disc. Besides the divergence-free constraint

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } D,$$

¹All limits are understood in the limit superior sense to ensure they are well-defined.

we enforce the no-penetration boundary conditions

$$\mathbf{u} \cdot \hat{\mathbf{n}} = 0 \quad \text{at } \partial D$$

where $\hat{\mathbf{n}}$ is the outwards pointing unit normal.

Given this setup, we minimize $\langle |\nabla T|^2 \rangle$ among all velocities $\mathbf{u}$ with a given value of mean enstrophy $\langle |\nabla \mathbf{u}|^2 \rangle$, or mean kinetic energy $\langle |\mathbf{u}|^2 \rangle$. These ‘enstrophy-’ and ‘energy-constrained optimal cooling problems’ were originally posed in [39] and analysed further in [40]. Actually, those papers focussed on the special case of a constant heat source, e.g. $f = 1$, and were also mostly concerned with the energy-constrained problem. Here, we treat a general source function $f(\mathbf{x}, t)$ which may vary both in space and in time, and may even be negative in some places and at some times allowing for temporary sinks. Before coming to our results, we pause to discuss other possible objective functionals as well as the practical meaning of ‘optimizing over $\mathbf{u}$ ’.

Various minimization problems have been proposed to optimize cooling. In [39], the authors minimize the mean temperature $\langle T \rangle$ subject to an energy-constraint and for a constant positive source. In [40], the same steady minimization is studied alongside a variety of others involving L^p -based quantities, namely, $\langle T^p \rangle$ for $1 \leq p < \infty$ and $\max T$ for $p = \infty$. Multiplying the advection–diffusion equation (1.1) by T and integrating by parts shows that

$$\kappa \langle |\nabla T|^2 \rangle = \langle fT \rangle$$

in general. If f is constant and positive, minimizing $\langle T \rangle$ can be regarded as minimizing $\langle |\nabla T|^2 \rangle$. If f is bounded and uniformly positive, meaning that $C \geq f \geq c$ for some fixed $c, C \in (0, \infty)$, then

$$c \langle T \rangle \leq \kappa \langle |\nabla T|^2 \rangle \leq C \langle T \rangle,$$

so that the minimizations give comparable results. For a general and possibly sign-indefinite f , such a simple relationship need not hold. Our choice to minimize $\langle |\nabla T|^2 \rangle$ as opposed to, say, $\langle |T|^p \rangle$ is partially out of mathematical convenience, but also because the former provides a more direct assessment of long-term diffusive transport—the rate-limiting step in any heat exchange.

Turning to our choice to treat the velocity directly as a control, perhaps it is more reasonable from the viewpoint of applications to think of controlling an applied body force $\mathbf{f}(\mathbf{x}, t)$. One may then seek to optimize cooling subject to a constraint on the power used. Or, one might limit the complexity of the force, e.g. by constraining $\langle |\nabla \mathbf{f}|^2 \rangle$. In any case, the velocity $\mathbf{u}$ will solve the forced, incompressible Navier–Stokes equations

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \frac{1}{\rho} \nabla p = \nu \Delta \mathbf{u} + \frac{1}{\rho} \mathbf{f} \quad \text{in } D, \quad (1.2)$$

where $p(\mathbf{x}, t)$ and $\nu > 0$ are the pressure and kinematic viscosity and $\rho > 0$ is the density. Dotting (1.2) by $\mathbf{u}$, integrating by parts, and remembering that $\nabla \cdot \mathbf{u} = 0$ yields the fundamental ‘balance law’

$$\nu \langle |\nabla \mathbf{u}|^2 \rangle = \frac{1}{\rho} \langle \mathbf{f} \cdot \mathbf{u} \rangle.$$

Thus, a constraint on the mean power $\langle \mathbf{f} \cdot \mathbf{u} \rangle$ can be regarded as a constraint on the mean enstrophy $\langle |\nabla \mathbf{u}|^2 \rangle$. Having found an optimal $\mathbf{u}$, we can then simply read off from (1.2) the corresponding optimal $\mathbf{f}$. This is not to say that every optimal velocity is dynamically stable—in fact, in related problems involving extremal orbits the opposite situation holds [41,42]. Nevertheless, to get started, we set aside the momentum equation henceforth and focus solely on optimizing advection–diffusion.

Switching to non-dimensional variables, let $\kappa = R = 1$. The value of the mean enstrophy $\langle |\nabla \mathbf{u}|^2 \rangle$, or the mean energy $\langle |\mathbf{u}|^2 \rangle$ when we discuss it, will be called the Peclet number Pe as either quantity sets the relative strength of advection to diffusion. (In dimensional variables, $Pe = UR/\kappa$ with U being the imposed velocity scale.) In the advective limit $Pe \rightarrow \infty$, one expects to be able to drive $\langle |\nabla T|^2 \rangle \rightarrow 0$ along a well-chosen sequence of velocities. The question is: at what optimal rate

can this convergence occur? Our main result is a set of upper and lower bounds on $\min(|\nabla T|^2)$ subject to the enstrophy constraint $\langle |\nabla \mathbf{u}|^2 \rangle = Pe^2$ that identify the optimal cooling rate up to a possible logarithm in Pe . Let λ_1 be the first Dirichlet eigenvalue of $-\Delta$ on D , and let $t \wedge s$ denote the minimum of $t, s \in \mathbb{R}$.

Theorem 1.1. *Let $f(\mathbf{x}, t)$ satisfy*

$$\lim_{\tau \rightarrow \infty} \frac{1}{\sqrt{\tau}} \int_0^\tau e^{-\lambda_1((\tau-t) \wedge t)} \|f(\cdot, t)\|_{L^2(D)} dt = 0 \quad \text{and} \quad \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle < \infty$$

and assume $\langle f \rangle > 0$. There exist positive constants $C(f)$, $C'(f)$ and $c(f)$ depending only on f such that

$$C \frac{1}{Pe^{2/3}} \leq \min_{\substack{\mathbf{u}(\mathbf{x}, t) \\ \langle |\nabla \mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} \cdot \hat{\mathbf{n}} = 0 \text{ at } \partial D}} \langle |\nabla T|^2 \rangle \leq C' \frac{\log^{4/3} Pe}{Pe^{2/3}}$$

whenever $Pe \geq c(f)$.

Remark 1.2. The first assumption concerns the possibility that f grows as $t \rightarrow \infty$, and limits that growth such that $\langle |\nabla T|^2 \rangle$ remains finite. Without it, one can make $\langle |\nabla T|^2 \rangle = \infty$ for all $\mathbf{u}$, e.g. with $f = a(t)\varphi_1(\mathbf{x})$ and where $\varphi_1 \in H_0^1(D)$ has $-\Delta\varphi_1 = \lambda_1\varphi_1$. The remaining assumptions on f enter into our bounds on C and C' : we achieve $C \gtrsim \langle f \rangle^2$ and $C' \lesssim \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle$. For the dependence of c , see §2 where we prove the lower bound.

Remark 1.3. The same result holds if the no-penetration boundary conditions $\mathbf{u} \cdot \hat{\mathbf{n}} = 0$ are replaced by the more restrictive no-slip conditions $\mathbf{u} = \mathbf{0}$ at ∂D . This is because we use no-slip velocities for the upper bound. That the lower bound goes through to the no-slip case is clear.

Let us briefly discuss the strategy of our proof. On the one hand, we must explain why no enstrophy-constrained velocity having $\langle |\nabla \mathbf{u}|^2 \rangle = Pe^2$ can lower $\langle |\nabla T|^2 \rangle$ significantly beyond $Pe^{-2/3}$. At the same time, we must construct a sequence of admissible velocities to cool within a logarithm of this bound. The challenge is to find a way of computing heat transfer that at the same time shows how it can be optimized. We follow the approach of our previous articles on maximizing transport across an externally heated fluid layer [43–45], the key difference being the presence of the internal source $f(\mathbf{x}, t)$. We show it is possible to bound $\langle |\nabla T|^2 \rangle$, and in fact to compute it exactly in the steady case, by a pair of variational problems. These problems recall Dirichlet's principle for Poisson's equation, but are non-local due to the mixed character of the operator $\mathbf{u} \cdot \nabla - \Delta$. That there exists a non-local Dirichlet-like principle for the effective diffusivity of a periodic or random fluid flow is well-known in homogenization [46,47]. Our idea is that $\langle |\nabla T|^2 \rangle$ should behave like an effective diffusivity upon optimization. This is very much in the spirit of the general connection between homogenization and optimal design (see, e.g. [48–50]).

The bulk of our work goes into constructing the 'branching flows' behind the upper bound in theorem 1.1. In general, we envision an unsteady, tree-like, multi-scale flow whose features refine from the bulk to the boundary as in the bottom of figure 1. To achieve it, we piece together a family of convection roll-like flows, with the number of rolls in the azimuthal direction θ depending on the radial coordinate r . A similar, albeit steady, branching flow was used in [43,45] to prove nearly sharp bounds on optimal transport through a fluid layer. Later on, a beautiful and fully three-dimensional branching flow was found numerically via gradient ascent [51]. Each of these may be regarded as a more refined version of Busse's 'multi- α ' flows [22], the latter of which do not enforce the full advection–diffusion equation but instead hinge upon various balance laws it implies.

Branching may be anticipated as a mechanism for optimizing heat transfer by asking what it takes to guide Brownian particles from the bulk to the boundary of the disc. Imagine a cloud of particles is released at a radius r and is to be transported outwards across a distance $\sim 1 - r$, say with (non-dimensional) speed $\sim U$. In time τ , the particles diffuse in the θ -direction by a typical distance $\sim \sqrt{\tau}$. In the same time, they advect in the r -direction by a distance $\sim U \cdot \tau$. With the




Figure 1. Divergence-free flow designs for cooling an internally heated unit disc. The top row shows contours of the source functions $f = 1, e^{-4r^2}, e^{-4(1-r)^2}$ and $\sin^2(2\theta)$ from left to right. The middle row shows streamlines of convection roll-like designs adapted to these sources. The bottom row shows their branching flow counterparts. Counterclockwise/clockwise circulations are coloured purple/blue. (Online version in colour.)

goal of not losing too many particles to poorly directed advection, we suggest using streamlines whose azimuthal scale $\ell(r)$ matches that of the Brownian cloud:

$$\ell(r) \sim \sqrt{\frac{1-r}{U}}. \quad (1.3)$$

Two parameters emerge, namely the speed U and the boundary layer width δ_{bl} where the streamlines finally turn around. By the enstrophy constraint,

$$Pe^2 = \langle |\nabla \mathbf{u}|^2 \rangle \sim \int^{1-\delta_{\text{bl}}} \left(\frac{U}{\ell(r)} \right)^2 r \, dr \sim U^3 \log \frac{1}{\delta_{\text{bl}}}.$$

Thinking of an isotropic roll-based boundary layer, we take $\sqrt{\delta_{\text{bl}}/U} \sim \delta_{\text{bl}}$ giving $U \sim 1/\delta_{\text{bl}}$. This determines all parameters via the scalings

$$U \sim \frac{Pe^{2/3}}{\log^{1/3} Pe} \quad \text{and} \quad \delta_{\text{bl}} \sim \frac{\log^{1/3} Pe}{Pe^{2/3}}. \quad (1.4)$$

Even if our thought experiment on how advection can effectively ‘hug’ diffusion suggests a particular way of branching, it is far from a proof of theorem 1.1, or even of its upper bound. To achieve it, we apply our variational bounds to estimate $\langle |\nabla T|^2 \rangle$ on a general roll-based branching flow design. By optimizing over all admissible ‘scale functions’ $\ell(r)$, we discover the very same scalings as in (1.3) and (1.4). In fact, we see no way out of these and the logarithmically corrected

bound they imply: we conjecture that

$$\min_{\substack{\mathbf{u}(\mathbf{x},t) \\ \langle |\nabla \mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} \cdot \hat{\mathbf{n}} = 0 \text{ at } \partial D}} \langle |\nabla T|^2 \rangle = C(f) \cdot \frac{\log^{4/3} Pe}{Pe^{2/3}} + \dots \quad \text{as } Pe \rightarrow \infty,$$

with the dots representing asymptotically negligible terms. This is for two-dimensional cooling; the data in [51] suggest a pure power law scaling for optimal transport across a three-dimensional layer. Maybe there is more room for chaperoning particles in three dimensions versus two? In any case, proving the analogue of our conjecture for a fluid layer would establish a new bound on the longstanding problem of Rayleigh–Bénard convection, as noted in [43]. This bound would state that $Nu \lesssim Ra^{1/2} / \log^2 Ra$ with Nu and Ra being the Nusselt and Rayleigh numbers (the relevant measures of transport and driving there).

Stochastic analysis of optimal cooling is the focus of [40] where the authors estimate, among other things, the mean temperature of steady convection roll-like flows in an internally heated layer with a constant source. Using large deviations-based bounds, they prove that $\min \langle T \rangle \leq c(\delta)(\log Pe / Pe^{1-\delta})$ for any $\delta > 0$, where $\langle |\mathbf{u}|^2 \rangle = Pe^2$. Showing the conjectured optimal scaling

$$\min_{\substack{\mathbf{u}(\mathbf{x},t) \\ \langle |\mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} \cdot \hat{\mathbf{n}} = 0 \text{ at } \partial D}} \langle |\nabla T|^2 \rangle = C(f) \cdot \frac{1}{Pe} + \dots \quad \text{as } Pe \rightarrow \infty$$

put forth for $f = 1$ in [39] remains an open challenge. While the arguments behind theorem 1.1 lead without major modifications to the supposedly sub-optimal lower bound $\min \langle |\nabla T|^2 \rangle \geq C(f)/Pe^2$, they are strong enough to show the conjectured upper bound $\min \langle |\nabla T|^2 \rangle \leq C'(f)/Pe$ and to likewise improve the estimate from [40]. The Pe^{-1} scaling was found in [39] through a matched asymptotic analysis of convection roll-like flows, but without a bound on the error terms. See propositions 2.6 and 5.1 for our lower and upper bounds on energy-constrained cooling.

This paper is organized as follows. We begin in §2 with our variational bounds on cooling. At the end of that section, we choose a background-like test function to prove our lower bounds on the enstrophy- and energy-constrained problems, including the lower bound part of theorem 1.1. The rest of the paper is devoted to upper bounds. First, we discuss the steady enstrophy-constrained problem in §3 to explain our strategy for finding designs. Then, in §4 we construct our branching flows to prove the upper bound part of theorem 1.1. We end in §5 with our upper bound on energy-constrained cooling.

(a) Notation

We write $X \lesssim Y$ if $X \leq cY$ for a fixed numerical constant $c > 0$, i.e. one that is independent of all parameters. We write $X \lesssim_a Y$ if $X \leq cY$ where $c = c(a)$, and $X \ll Y$ if $(X/Y) \rightarrow 0$ in a relevant limit. We abbreviate $X \wedge Y = \min\{X, Y\}$ and $X \vee Y = \max\{X, Y\}$.

We often conflate a point $\mathbf{x}$ with its polar coordinates (r, θ) . The unit vectors $\hat{\mathbf{e}}_r = \mathbf{x}/|\mathbf{x}|$ and $\hat{\mathbf{e}}_\theta = \hat{\mathbf{e}}_r^\perp$, where $(\cdot)^\perp$ is a counterclockwise rotation by $\pi/2$. Given a function $\varphi(\mathbf{x})$, its average on the disc D is

$$\oint_D \varphi = \frac{1}{|D|} \int_D \varphi(\mathbf{x}) \, d\mathbf{x},$$

where $|D|$ is the disc's area. By D_r we mean the concentric disc of radius r . Restricting φ to ∂D_r gives a function of θ , whose average and L^1 -norm are

$$\overline{\varphi}(r) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(r, \theta) \, d\theta \quad \text{and} \quad \|\varphi\|_{L^1_\theta}(r) = \int_0^{2\pi} |\varphi(r, \theta)| \, d\theta.$$

Given a function $\varphi(\mathbf{x}, t)$, we write

$$\langle \varphi \rangle = \limsup_{\tau \rightarrow \infty} \oint_0^\tau \oint_D \varphi = \limsup_{\tau \rightarrow \infty} \frac{1}{\tau} \frac{1}{|D|} \int_0^\tau \int_D \varphi(\mathbf{x}, t) \, d\mathbf{x} \, dt$$

for its (limit superior) space and long-time average.

The Sobolev spaces $L^2(D)$ and $H^1(D)$ are defined as usual, using the norms

$$\|\varphi\|_{L^2(D)} = \sqrt{\int_D |\varphi|^2 dx} \quad \text{and} \quad \|\varphi\|_{H^1(D)} = \sqrt{\int_D |\varphi|^2 + |\nabla \varphi|^2 dx}.$$

We write $H^{-1}(D)$ for the dual of $H_0^1(D)$, the latter indicating H^1 -functions on D with zero trace at ∂D . It will be convenient to normalize their duality bracket $(\cdot, \cdot)$ by $|D|$, and in particular we take

$$(\nabla \cdot \mathbf{m}, \varphi) = - \int_D \mathbf{m} \cdot \nabla \varphi$$

for $\mathbf{m} \in L^2(D; \mathbb{R}^2)$ and $\varphi \in H_0^1(D)$. We use various mixed spaces such as $L^2((0, \infty); H_0^1(D))$ and its local-in-time version $L_{\text{loc}}^2((0, \infty); H_0^1(D))$. For definitions, see a text on partial differential equations (PDEs) such as [52].

2. Variational bounds on cooling

We begin with a general method for bounding $\langle |\nabla T|^2 \rangle$ from above and below. While it is easy to show using integration by parts that

$$\langle |\nabla T|^2 \rangle \lesssim \langle |f|^2 \rangle, \quad (2.1)$$

improving this upper bound and finding a corresponding lower bound is not so simple. Incidentally, it follows from (2.1) and the linearity of the PDE in T that $\langle |\nabla T|^2 \rangle$ does not depend on the exact choice of its initial data, so long as it belongs to $L^2(D)$. To obtain better bounds, we invoke a certain symmetrizing change of variables that couples T to a second ‘temperature’ field arising from its adjoint PDE. The resulting bounds are sharp in the steady case where $\mathbf{u}$ and f do not depend on time.

Define the admissible set

$$\mathcal{A} = \left\{ \theta \in L_{\text{loc}}^2((0, \infty); H_0^1(D)) : \partial_t \theta \in L_{\text{loc}}^2((0, \infty); H^{-1}(D)), \limsup_{\tau \rightarrow \infty} \frac{1}{\sqrt{\tau}} \|\theta(\cdot, \tau)\|_{L^2(D)} < \infty \right\}$$

and let $\Delta^{-1} : H^{-1}(D) \rightarrow H_0^1(D)$ denote the inverse Laplacian with zero Dirichlet boundary conditions. Recall

$$\lambda_1 = \min_{\varphi \in H_0^1(D)} \frac{\int_D |\nabla \varphi|^2}{\int_D |\varphi|^2}.$$

Proposition 2.1. *Let $\mathbf{u}(\mathbf{x}, t)$ be weakly divergence-free and have*

$$\langle |\mathbf{u}|^2 \rangle < \infty$$

and let $f(\mathbf{x}, t)$ satisfy

$$\lim_{\tau \rightarrow \infty} \frac{1}{\sqrt{\tau}} \int_0^\tau e^{-\lambda_1((\tau-t) \wedge t)} \|f(\cdot, t)\|_{L^2(D)} dt = 0.$$

Any weak solution $T(\mathbf{x}, t)$ of

$$\begin{cases} \partial_t T + \mathbf{u} \cdot \nabla T = \Delta T + f & \text{in } D \\ T = 0 & \text{at } \partial D \end{cases}$$

with initial data $T(\cdot, 0) \in L^2(D)$ must obey the bounds

$$\begin{aligned} & \langle 2\xi f - |\nabla \xi|^2 - |\nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)\xi|^2 \rangle \\ & \leq \langle |\nabla T|^2 \rangle \leq \langle |\nabla \eta|^2 + |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla)\eta - f]|^2 \rangle \end{aligned}$$

for all $\xi, \eta \in \mathcal{A}$.

Remark 2.2. The statement that T is a weak solution deserves to be clarified, especially as $\mathbf{u} \cdot \nabla T$ is at first glance only in $L^1(D)$ a.e. in time, and so would appear not to belong to $H^{-1}(D) = (H_0^1(D))'$. (Two dimensions is critical for the relevant Sobolev embedding.) However, $\mathbf{u}$ and ∇T are divergence- and curl-free, and this is enough to compensate for their lack of regularity.

To see why, introduce a stream function $\psi \in H^1(D)$ with $\mathbf{u} = \nabla^\perp \psi$, and note that $\mathbf{u} \cdot \nabla T$ is the Jacobian determinant of the mapping $\mathbf{x} \mapsto (\psi, T)$. Therefore, by the estimate of Coifman, Lions, Meyer and Semmes on Jacobian determinants [53] (see also [54]),

$$\left| \int_D \mathbf{u} \cdot \nabla T \varphi \, d\mathbf{x} \right| \lesssim \|\mathbf{u}\|_{L^2(D)} \|\nabla T\|_{L^2(D)} \|\nabla \varphi\|_{L^2(D)} \quad \forall \varphi \in C_c^\infty(D).$$

Letting $(\cdot, \cdot)$ denote the (normalized) duality bracket between $H^{-1}(D)$ and $H_0^1(D)$, the previous inequality extends to say that

$$|(\mathbf{u} \cdot \nabla T, \varphi)| \lesssim \|\mathbf{u}\|_{L^2(D)} \|\nabla T\|_{L^2(D)} \|\nabla \varphi\|_{L^2(D)} \quad \forall \varphi \in H_0^1(D). \quad (2.2)$$

Thus, $\mathbf{u} \cdot \nabla T \in H^{-1}(D)$ at a.e. time. Clearly, the same holds for ΔT and f .

At this point, the definition of T as a weak solution can be carried out as usual, by requiring that

$$T \in L_{\text{loc}}^2((0, \infty); H_0^1(D)) \quad \text{and} \quad \partial_t T \in L_{\text{loc}}^2((0, \infty); H^{-1}(D)),$$

and enforcing the PDE as an equality on $H^{-1}(D)$. Existence and uniqueness of weak solutions with initial data in $L^2(D)$ follows (see, e.g. [52]).

Remark 2.3. A particular consequence of the previous remark that will be used throughout the proof of proposition 2.1 is that

$$(\mathbf{u} \cdot \nabla T, T) = 0 \quad \text{for a.e. } t,$$

if $\mathbf{u}$ and T are as in the proposition. This follows from a smooth approximation argument and the fact that $T(\cdot, t) \in H_0^1(D)$ a.e. in t . Indeed, if $T(\cdot, t)$ is smooth and compactly supported in D then

$$(\mathbf{u} \cdot \nabla T, T) = \frac{1}{2} \int_D \mathbf{u} \cdot \nabla |T|^2 \, d\mathbf{x} = 0$$

by the divergence theorem. The bilinear form $T \mapsto (\mathbf{u} \cdot \nabla T, T)$ is continuous on $H_0^1(D)$, per (2.2). The claimed identity follows from the density of smooth and compactly supported functions in $H_0^1(D)$.

Remark 2.4. In the case that $\mathbf{u}$ and f are steady, the bounds from the proposition become sharp and are achieved by steady test functions $\eta(\mathbf{x})$ and $\xi(\mathbf{x})$. That is,

$$\begin{aligned} \langle |\nabla T|^2 \rangle &= \min_{\eta \in H_0^1(D)} \int_D |\nabla \eta|^2 + |\nabla \Delta^{-1}[\mathbf{u} \cdot \nabla \eta - f]|^2 \, d\mathbf{x} \\ &= \max_{\xi \in H_0^1(D)} \int_D 2\xi f - |\nabla \xi|^2 - |\nabla \Delta^{-1} \mathbf{u} \cdot \nabla \xi|^2 \, d\mathbf{x}. \end{aligned}$$

The crucial point is that, when $\mathbf{u}$ and f are steady, the above optimizations in η and ξ not only provide bounds on $\langle |\nabla T|^2 \rangle$, but also turn out to be ‘strongly’ dual in that their optimal values are the same (an easy consequence of their Euler–Lagrange equations). See [43] for the full proof of a similar duality arising for steady heat transport through a fluid layer.

Remark 2.5. On the other hand, if f is allowed to depend on time, equality need not hold between our bounds on $\langle |\nabla T|^2 \rangle$. This is because one cannot rule out the possibility that

$$\liminf_{\tau \rightarrow \infty} \int_0^\tau \int_D |\nabla T|^2 < \limsup_{\tau \rightarrow \infty} \int_0^\tau \int_D |\nabla T|^2$$

in general while, as will become clear in the proof below, our bounds actually hold on these limit inferior and limit superior long-time averages (see (2.6) and (2.7)). We do not know if our bounds are sharp when $f(\mathbf{x})$ is steady and $\mathbf{u}(\mathbf{x}, t)$ is unsteady, though we guess the answer is ‘no’.

Proof. Consider the formally adjoint pair of problems

$$\begin{cases} (\pm \partial_t \pm \mathbf{u} \cdot \nabla) T_{\pm} = \Delta T_{\pm} + f & \text{in } D \\ T_{\pm} = 0 & \text{at } \partial D \end{cases}'$$

the + version of which is solved by the given temperature T . As the value of $\langle |\nabla T|^2 \rangle$ does not depend on its initial data, we let $T(\cdot, 0) = 0$ and refer to this version of the temperature as T_+ in the rest of this proof. Fixing an arbitrary time $\tau > 0$ which we will eventually take to ∞ , let T_- be the unique weak solution of the – problem with $T_-(\cdot, \tau) = 0$. (The definition of a ‘weak solution’ is standard; see remark 2.2.) A quick argument using integration by parts and Gronwall’s inequality shows that

$$\begin{aligned} \|T_+(\cdot, \tau)\|_{L^2(D)} \vee \|T_-(\cdot, 0)\|_{L^2(D)} \\ \leq \int_0^\tau e^{-\lambda_1((\tau-t) \wedge t)} \|f(\cdot, t)\|_{L^2(D)} dt \ll \sqrt{\tau} \quad \text{as } \tau \rightarrow \infty \end{aligned}$$

by our hypothesis on f . Briefly, one notes for $Z = \sqrt{\epsilon + \|T_+\|_{L^2(D)}^2}$ that

$$\frac{d}{dt} Z + \lambda_1 Z \leq \lambda_1 \epsilon^{1/2} + \|f\|_{L^2(D)}$$

for all $\epsilon > 0$. Using the integrating factor $e^{-\lambda_1(\tau-t)}$ and taking $\epsilon \rightarrow 0$ yields the + version of the inequality. The – version follows by reversing time.

Having defined $T_{\pm}$, we now change variables to the pair

$$\eta = \frac{1}{2}(T_+ - T_-) \quad \text{and} \quad \xi = \frac{1}{2}(T_+ + T_-)$$

and note they satisfy

$$\begin{cases} (\partial_t + \mathbf{u} \cdot \nabla) \eta = \Delta \xi + f & \text{in } D \\ (\partial_t + \mathbf{u} \cdot \nabla) \xi = \Delta \eta & \text{in } D \\ \eta = \xi = 0 & \text{at } \partial D \end{cases} \quad (2.3)$$

weakly for $t \in (0, \tau)$. It follows from our previous bounds on $T_{\pm}$ that

$$\|\xi(\cdot, 0)\|_{L^2(D)} + \|\xi(\cdot, \tau)\|_{L^2(D)} + \|\eta(\cdot, 0)\|_{L^2(D)} + \|\eta(\cdot, \tau)\|_{L^2(D)} \ll \sqrt{\tau} \quad \text{as } \tau \rightarrow \infty. \quad (2.4)$$

Introduce the notation $\langle \cdot \rangle_\tau = \int_0^\tau \int_D \cdot \, d\mathbf{x} \, dt$ for the truncated space- and time average. Testing the second PDE in (2.3) against ξ , we get that

$$\begin{aligned} \langle \nabla \eta \cdot \nabla \xi \rangle_\tau &= \int_0^\tau \langle -\Delta \eta, \xi \rangle dt = \int_0^\tau \langle -(\partial_t + \mathbf{u} \cdot \nabla) \xi, \xi \rangle dt \\ &= \int_0^\tau \left[-\frac{d}{dt} \frac{1}{2} \int_D |\xi|^2 d\mathbf{x} + \langle \mathbf{u} \cdot \nabla \xi, \xi \rangle \right] dt \\ &= \frac{1}{2\tau} \int_D |\xi(\mathbf{x}, 0)|^2 - |\xi(\mathbf{x}, \tau)|^2 d\mathbf{x} \end{aligned}$$

where again $\langle \cdot, \cdot \rangle$ is the duality bracket normalized by $|D|$. That $\mathbf{u} \cdot \nabla \xi \in H^{-1}(D)$ at a.e. time follows from the assumed mean-square integrability of $\mathbf{u}$ and the statement that it is weakly divergence-free; see remarks 2.2 and 2.3 for the proof that $\langle \mathbf{u} \cdot \nabla \xi, \xi \rangle = 0$. Combined with (2.4), the conclusion is that

$$\langle \nabla \eta \cdot \nabla \xi \rangle_\tau \rightarrow 0 \quad \text{as } \tau \rightarrow \infty.$$

Hence, $T = T_+ = (\eta + \xi)/2$ obeys

$$\begin{aligned} \langle |\nabla T_+|^2 \rangle_\tau &= \langle |\nabla \eta|^2 + |\nabla \xi|^2 \rangle_\tau + o_\tau(1) \\ &= \langle |\nabla \eta|^2 + |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla) \eta - f]|^2 \rangle_\tau + o_\tau(1) \end{aligned} \quad (2.5)$$

with $o_\tau(1)$ denoting a quantity that vanishes as $\tau \rightarrow \infty$. This way of writing the mean-square temperature gradient leads to the upper bound from the claim.

To see why, let $\tilde{\eta} \in \mathcal{A}$ and consider the difference

$$A_\tau := \left\langle |\nabla \tilde{\eta}|^2 + |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla)\tilde{\eta} - f]|^2 \right\rangle_\tau - \left\langle |\nabla \eta|^2 + |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla)\eta - f]|^2 \right\rangle_\tau.$$

We claim it is non-negative up to a term that vanishes as $\tau \rightarrow \infty$. Since $|\cdot|^2$ is convex,

$$\begin{aligned} \frac{1}{2}A_\tau &\geq \left\langle \nabla \eta \cdot \nabla(\tilde{\eta} - \eta) + \nabla \xi \cdot \nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)(\tilde{\eta} - \eta) \right\rangle_\tau \\ &= \int_0^\tau \left[\int_D \nabla \eta \cdot \nabla(\tilde{\eta} - \eta) \, dx - ((\partial_t + \mathbf{u} \cdot \nabla)(\tilde{\eta} - \eta), \xi) \right] dt \\ &= \int_0^\tau (-\Delta \eta + (\partial_t + \mathbf{u} \cdot \nabla)\xi, \tilde{\eta} - \eta) \, dt - \frac{1}{\tau} \left(\int_D (\tilde{\eta} - \eta)\xi \right) \Big|_{t=0}^{t=\tau} \geq -o_\tau(1) \end{aligned}$$

again by (2.4) and the growth condition in our definition of $\mathcal{A}$. Hence,

$$\left\langle |\nabla T_+|^2 \right\rangle_\tau \leq \left\langle |\nabla \tilde{\eta}|^2 + |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla)\tilde{\eta} - f]|^2 \right\rangle_\tau + o_\tau(1) \quad (2.6)$$

for all $\tilde{\eta} \in \mathcal{A}$. This bound is a bit better than the one from the claim (however, see remark 2.5).

Returning to the PDEs in (2.3), we now test the second one against η and integrate by parts, again using the abbreviation $o_\tau(1)$ for terms that limit to zero as $\tau \rightarrow \infty$. The result is that

$$\begin{aligned} \left\langle |\nabla \eta|^2 \right\rangle_\tau &= \int_0^\tau (-\Delta \eta, \eta) \, dt = \int_0^\tau (-(\partial_t + \mathbf{u} \cdot \nabla)\xi, \eta) \, dt \\ &= \int_0^\tau ((\partial_t + \mathbf{u} \cdot \nabla)\eta, \xi) \, dt + o_\tau(1) = \int_0^\tau (\Delta \xi + f, \xi) \, dt + o_\tau(1) \\ &= \left\langle -|\nabla \xi|^2 + f\xi \right\rangle_\tau + o_\tau(1) \end{aligned}$$

where in the second line we applied the first PDE in (2.3). Combined with (2.5), this proves that

$$\left\langle |\nabla T_+|^2 \right\rangle_\tau = \left\langle f\xi \right\rangle_\tau + o_\tau(1)$$

and that

$$\begin{aligned} \left\langle |\nabla T_+|^2 \right\rangle_\tau &= \left\langle 2f\xi - |\nabla \eta|^2 - |\nabla \xi|^2 \right\rangle_\tau + o_\tau(1) \\ &= \left\langle 2f\xi - |\nabla \xi|^2 - |\nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)\xi|^2 \right\rangle_\tau + o_\tau(1). \end{aligned}$$

We are ready to deduce the lower bound from the claim.

Let $\tilde{\xi} \in \mathcal{A}$ and call

$$B_\tau := \left\langle 2f\tilde{\xi} - |\nabla \tilde{\xi}|^2 - |\nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)\tilde{\xi}|^2 \right\rangle_\tau - \left\langle 2f\tilde{\xi} - |\nabla \tilde{\xi}|^2 - |\nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)\tilde{\xi}|^2 \right\rangle_\tau.$$

As before, we use the convexity of $|\cdot|^2$ to write that

$$\begin{aligned} \frac{1}{2}B_\tau &\geq \left\langle -f(\tilde{\xi} - \xi) + \nabla \xi \cdot \nabla(\tilde{\xi} - \xi) + \nabla \eta \cdot \nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)(\tilde{\xi} - \xi) \right\rangle_\tau \\ &= \int_0^\tau \left[\int_D -f(\tilde{\xi} - \xi) + \nabla \xi \cdot \nabla(\tilde{\xi} - \xi) \, dx - ((\partial_t + \mathbf{u} \cdot \nabla)(\tilde{\xi} - \xi), \eta) \right] dt \\ &= \int_0^\tau (-f - \Delta \xi + (\partial_t + \mathbf{u} \cdot \nabla)\eta, \tilde{\xi} - \xi) \, dt - \frac{1}{\tau} \left(\int_D (\tilde{\xi} - \xi)\eta \right) \Big|_{t=0}^{t=\tau} \geq -o_\tau(1). \end{aligned}$$

In the last step, we applied (2.4) to handle the terms at $t=0$ and $t=\tau$. Thus,

$$\left\langle |\nabla T_+|^2 \right\rangle_\tau \geq \left\langle 2f\tilde{\xi} - |\nabla \tilde{\xi}|^2 - |\nabla \Delta^{-1}(\partial_t + \mathbf{u} \cdot \nabla)\tilde{\xi}|^2 \right\rangle_\tau - o_\tau(1) \quad (2.7)$$

for all $\tilde{\xi} \in \mathcal{A}$. Again, the result is a bit better than the bound from the claim. ■

As a first application of our variational bounds on cooling, we obtain the lower bound from theorem 1.1 which applies to all velocities with a specified mean enstrophy.

Proof of the lower bound from theorem 1.1. The lower bound half of proposition 2.1 applied with a steady test function $\xi \in H_0^1(D)$ shows that

$$2\langle f\xi \rangle \leq \langle |\nabla T|^2 \rangle + \langle |\nabla \xi|^2 + |\nabla \Delta^{-1} \nabla \cdot (\mathbf{u}\xi)|^2 \rangle.$$

Scaling $\xi \rightarrow \lambda \xi$ and optimizing over $\lambda \in \mathbb{R}$, we now write that

$$\langle f\xi \rangle^2 \leq \langle |\nabla T|^2 \rangle \langle |\nabla \xi|^2 + |\nabla \Delta^{-1} \nabla \cdot (\mathbf{u}\xi)|^2 \rangle. \quad (2.8)$$

This bound holds for all $\xi \in H_0^1(D)$, and we proceed to make a choice.

Given any $\delta \in (0, 1)$, define

$$\xi_\delta = \chi_\delta(r),$$

where $\chi_\delta(r)$ is a smooth, radial ‘cutoff function’ that goes from zero to one across a small boundary layer. Precisely, $\chi_\delta \in C_c^\infty([0, 1])$ satisfies

$$0 \leq \chi_\delta(r) \leq 1 \quad \forall r \in [0, 1) \quad \text{and} \quad \chi_\delta(r) = 1 \quad \forall r \leq 1 - \delta$$

and has

$$\|\chi'_\delta\|_{L^\infty([0,1])} \lesssim \frac{1}{\delta},$$

with a constant independent of δ . Note that

$$\begin{aligned} |\langle f \rangle| &\leq |\langle f\xi_\delta \rangle| + |\langle f(\xi_\delta - 1) \rangle| \leq |\langle f\xi_\delta \rangle| + \sqrt{\langle |f|^2 \rangle} \sqrt{\langle \mathbb{1}_{r>1-\delta} \rangle} \\ &\leq |\langle f\xi_\delta \rangle| + C'(f)\sqrt{\delta}. \end{aligned}$$

Hence, there exists $\delta_0(f) > 0$ such that

$$\frac{1}{2} |\langle f \rangle| \leq |\langle f\xi_\delta \rangle| \quad \forall \delta \in (0, \delta_0]. \quad (2.9)$$

Similarly,

$$\langle |\nabla \xi_\delta|^2 \rangle = \int_D |\chi'_\delta(r)|^2 \lesssim \int_D \frac{1}{\delta^2} \mathbb{1}_{r>1-\delta} \lesssim \frac{1}{\delta} \quad (2.10)$$

for all δ .

At this point, we have dealt with each of the terms in the bound (2.8) except for the non-local one involving the L^2 -orthogonal projection $\nabla \Delta^{-1} \nabla$. For this, observe that

$$\int_D |\nabla \Delta^{-1} \nabla \cdot \mathbf{v}|^2 \, \mathrm{d}\mathbf{x} = \max_{\theta \in H_0^1(D)} \int_D 2\nabla \theta \cdot \mathbf{v} - |\nabla \theta|^2 \, \mathrm{d}\mathbf{x}$$

for all $\mathbf{v} \in L^2(D)$. So, the estimate

$$\langle |\nabla \Delta^{-1} \nabla \cdot (\mathbf{u}\xi_\delta)|^2 \rangle \lesssim \delta^2 \langle |\nabla \mathbf{u}|^2 \rangle \quad (2.11)$$

follows from its steady version

$$\int_D \nabla \theta \cdot \mathbf{u}\xi_\delta \lesssim \int_D |\nabla \theta|^2 + \delta^2 |\nabla \mathbf{u}|^2 \quad \forall \theta \in H_0^1(D),$$

which we now establish at a.e. time. Since $\mathbf{u}$ is divergence-free and $\xi_\delta = \chi_\delta(r)$ depends only on r ,

$$\begin{aligned} \left| \int_D \nabla \theta \cdot \mathbf{u}\xi_\delta \right| &= \left| \int_D \theta \mathbf{u} \cdot \nabla \xi_\delta \right| \lesssim \int_{1-\delta}^1 |\overline{\theta \mathbf{u} \cdot \hat{\mathbf{e}}_r}| |\chi'_\delta| r \, \mathrm{d}r \\ &\lesssim \int_{1-\delta}^1 |\overline{\theta \mathbf{u} \cdot \hat{\mathbf{e}}_r}| \, \mathrm{d}r. \end{aligned}$$

Note $\bar{\cdot} = \int_0^{2\pi} \cdot d\theta$ gives the θ -average, as defined in §1a. Continuing, we claim that

$$|\overline{\theta \mathbf{u} \cdot \hat{\mathbf{e}}_r}| \lesssim (1-r) \|\nabla \theta\|_{L^2(D)} \|\nabla \mathbf{u}\|_{L^2(D)} \quad \text{for a.e. } r \in (0, 1). \quad (2.12)$$

In fact, (2.12) follows a more or less standard argument involving the Cauchy–Schwarz inequality applied to $(d/dr)\bar{\theta}^2$ and $(d/dr)(\mathbf{u} \cdot \hat{\mathbf{e}}_r)^2$, and then again to $(d/dr)\overline{\theta \mathbf{u} \cdot \hat{\mathbf{e}}_r}$, all of which are derivatives of quantities vanishing at $r = 1$. (For the complete details in an analogous fluid layer setting, see [43, lemma 2.6].) Applying (2.12), we get that

$$\begin{aligned} \int_{1-\delta}^1 |\overline{\theta \mathbf{u} \cdot \hat{\mathbf{e}}_r}| dr &\lesssim \int_{1-\delta}^1 (1-r) dr \cdot \|\nabla \theta\|_{L^2(D)} \|\nabla \mathbf{u}\|_{L^2(D)} \lesssim \delta \|\nabla \theta\|_{L^2(D)} \|\nabla \mathbf{u}\|_{L^2(D)} \\ &\lesssim \int_D |\nabla \theta|^2 + \delta^2 |\nabla \mathbf{u}|^2. \end{aligned}$$

As $\theta \in H_0^1(D)$ was arbitrary, the estimate (2.11) is proved.

Summing up, we have shown via (2.8)–(2.11) with the test function $\xi_\delta = \chi_\delta(r)$ that

$$\langle f \rangle^2 \lesssim \langle |\nabla T|^2 \rangle \left(\frac{1}{\delta} + \delta^2 \langle |\nabla \mathbf{u}|^2 \rangle \right) \quad \forall \delta \in (0, \delta_0(f)].$$

The bound

$$\langle f \rangle^2 \lesssim \langle |\nabla T|^2 \rangle \left(\frac{1}{\delta_0} \vee \langle |\nabla \mathbf{u}|^2 \rangle^{1/3} \right)$$

follows. Indeed, if $\delta_0^{-1} \leq \langle |\nabla \mathbf{u}|^2 \rangle^{1/3}$ we can choose $\delta = \langle |\nabla \mathbf{u}|^2 \rangle^{-1/3}$, otherwise we take $\delta = \delta_0$. This proves the lower bound in theorem 1.1 with $C \gtrsim \langle f \rangle^2$ and $c = \delta_0^{-3}$. ■

A minor modification of the previous proof leads to the (supposedly sub-optimal) lower bound on energy-constrained cooling from the introduction. To achieve it, replace the inequality in (2.11) with the simpler inequality

$$\langle |\nabla \Delta^{-1} \nabla \cdot (\mathbf{u} \xi_\delta)|^2 \rangle \leq \langle |\mathbf{u}|^2 \rangle \quad (2.13)$$

which holds because $\nabla \Delta^{-1} \nabla \cdot$ is an L^2 -orthogonal projection and $|\xi_\delta| \leq 1$. The rest of the proof goes through to yield the following result:

Proposition 2.6. *Let $f(\mathbf{x}, t)$ satisfy*

$$\lim_{\tau \rightarrow \infty} \frac{1}{\sqrt{\tau}} \int_0^\tau e^{-\lambda_1((\tau-t) \wedge t)} \|f(\cdot, t)\|_{L^2(D)} dt = 0 \quad \text{and} \quad \langle |f|^2 \rangle < \infty.$$

There exist positive constants C and $c(f)$, the former numerical and the latter depending only on f , such that

$$\frac{C}{Pe^2} \cdot \langle f \rangle^2 \leq \min_{\substack{\mathbf{u}(\mathbf{x}, t) \\ \langle |\mathbf{u}|^2 \rangle = Pe^2}} \langle |\nabla T|^2 \rangle$$

whenever $Pe \geq c(f)$.

The reader familiar with the background method may wonder whether it also leads to these lower bounds. Indeed it does, the key point being that either (2.11) or (2.13) verifies the relevant spectral constraint, depending on whether one considers enstrophy- or energy-constrained cooling. For more on the connection between our symmetrization-based bounds and those of the background method, see [43, 44].

3. Optimal steady flows for enstrophy-constrained cooling

The rest of this paper is about upper bounds. In this section and the next, we prove the one from theorem 1.1 on enstrophy-constrained cooling. First, to warm up, we study the steady version of

the problem

$$\min_{\substack{\mathbf{u}(\mathbf{x}) \\ f_D |\nabla \mathbf{u}|^2 = Pe^2 \\ \mathbf{u} = \mathbf{0} \text{ at } \partial D}} \langle |\nabla T|^2 \rangle \quad (3.1)$$

for a given source $f(\mathbf{x})$. We return to the unsteady problem in §4. Note we use no-slip boundary conditions from now on. This is compatible with our goal of proving an upper bound on optimal no-penetration flows, as minimizing over no-slip flows can only increase the result.

When $\mathbf{u}$ and f are steady, the upper bound from proposition 2.1 leads with a little effort to the double minimization

$$\min_{\substack{\mathbf{u}(\mathbf{x}), \eta(\mathbf{x}) \\ \mathbf{u} = \mathbf{0}, \eta = 0 \text{ at } \partial D}} \int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 + \frac{1}{Pe^2} \int_D |\nabla \eta|^2 \int_D |\nabla \mathbf{u}|^2, \quad (3.2)$$

whose optimal value is the same as that of the original problem (3.1), and whose optimizers $(\mathbf{u}, \eta)$ yield solutions to it under the rescaling $\mathbf{u} \rightarrow \lambda_{Pe} \mathbf{u}$ with $\lambda_{Pe} = Pe / \sqrt{\langle |\nabla \mathbf{u}|^2 \rangle}$. After explaining this in §3a, we go on in §3b to show how it is possible to achieve

$$\mathbf{u} \cdot \nabla \eta \approx f \quad \text{in } H^{-1}(D)$$

using convection roll-based flows. The estimates given there on the non-local part of (3.2) will come in handy later on when we discuss branching flows.

(a) A change of variables

First, we show that the problems in (3.1) and (3.2) are the same. Applying the upper bound from proposition 2.1—here we use remark 2.4 as it gives the better result in the steady case—we see that

$$\langle |\nabla T|^2 \rangle = \min_{\substack{\eta(\mathbf{x}) \\ \eta = 0 \text{ at } \partial D}} F(\mathbf{u}, \eta), \quad \text{where} \quad F(\mathbf{u}, \eta) = \int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 + \int_D |\nabla \eta|^2.$$

Evidently, minimizing $\langle |\nabla T|^2 \rangle$ is the same as minimizing F . A simple change of variables removes the enstrophy constraint. Consider the substitutions

$$\mathbf{u} = \frac{Pe}{\sqrt{f_D |\nabla \tilde{\mathbf{u}}|^2}} \tilde{\mathbf{u}} \quad \text{and} \quad \eta = \frac{\sqrt{f_D |\nabla \tilde{\mathbf{u}}|^2}}{Pe} \tilde{\eta}, \quad (3.3)$$

where $\tilde{\mathbf{u}}$ is not allowed to be identically zero. Evidently,

$$\int_D |\nabla \mathbf{u}|^2 = Pe^2 \quad \forall \tilde{\mathbf{u}},$$

while

$$F(\mathbf{u}, \eta) = \int_D |\nabla \Delta^{-1}(\tilde{\mathbf{u}} \cdot \nabla \tilde{\eta} - f)|^2 + \frac{1}{Pe^2} \int_D |\nabla \tilde{\mathbf{u}}|^2 \cdot \int_D |\nabla \tilde{\eta}|^2.$$

Thus, the two minimization problems

$$\min_{\substack{\mathbf{u}(\mathbf{x}) \\ f_D |\nabla \mathbf{u}|^2 = Pe^2 \\ \mathbf{u} = \mathbf{0} \text{ at } \partial D}} \langle |\nabla T|^2 \rangle$$

and

$$\min_{\substack{\tilde{\mathbf{u}}(\mathbf{x}), \tilde{\eta}(\mathbf{x}) \\ \tilde{\mathbf{u}} = \mathbf{0}, \tilde{\eta} = 0 \text{ at } \partial D}} \int_D |\nabla \Delta^{-1}(\tilde{\mathbf{u}} \cdot \nabla \tilde{\eta} - f)|^2 + \frac{1}{Pe^2} \int_D |\nabla \tilde{\mathbf{u}}|^2 \cdot \int_D |\nabla \tilde{\eta}|^2$$

are the same. Their optimal values agree, and their optimizers are related via (3.3).

A similar change of variables can be done for the unsteady problem, with the result being a bound rather than an equivalence. Setting

$$\mathbf{u} = \frac{Pe}{\sqrt{|\nabla \tilde{\mathbf{u}}|^2}} \tilde{\mathbf{u}} \quad \text{and} \quad \eta = \frac{\sqrt{|\nabla \tilde{\mathbf{u}}|^2}}{Pe} \tilde{\eta}$$

into the upper bound from proposition 2.1 shows that

$$\langle |\nabla T|^2 \rangle \leq \min_{\substack{\tilde{\eta}(\mathbf{x}, t) \\ \tilde{\eta}=0 \text{ at } \partial D}} \left\langle |\nabla \Delta^{-1}[(\partial_t + \tilde{\mathbf{u}} \cdot \nabla) \tilde{\eta} - f]|^2 + \frac{|\nabla \tilde{\mathbf{u}}|^2}{Pe^2} |\nabla \tilde{\eta}|^2 \right\rangle$$

where $T(\mathbf{x}, t)$ is a temperature field associated with the unsteady velocity $\mathbf{u}(\mathbf{x}, t)$. This is the starting point of §4.

(b) Steady advection

The next step is to see what it takes to drive the non-local terms in these optimizations to zero. Focusing again on a steady source $f(\mathbf{x})$, we ask what it takes for a velocity–test function pair $(\mathbf{u}(\mathbf{x}), \eta(\mathbf{x}))$ to achieve

$$\mathbf{u} \cdot \nabla \eta \approx f \quad \text{in } H^{-1}(D),$$

where the notation means that $\int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2$ is small. Guided by the divergence theorem and our usual assumption that $\nabla \cdot \mathbf{u} = 0$, we see that any successful pair must achieve

$$\int_{\partial V} \mathbf{u} \eta \cdot \hat{\mathbf{n}} \, ds \approx \int_V f \, d\mathbf{x}$$

for $V \subset D$. The following result makes this intuition precise.

Introduce the notation D_r for the disc of radius $r > 0$ centred at the origin of D , and again let

$$\bar{\varphi}(r) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(r, \theta) \, d\theta$$

denote the average over its boundary ∂D_r . As noted in §1a, we allow ourselves to conflate a point $\mathbf{x}$ with its polar coordinates (r, θ) . Any integral over ρ is done with respect to the radial coordinate.

Lemma 3.1. *Let $f \in L^2(D)$ and suppose that $(\mathbf{u}, \eta) \in H^1(D; \mathbb{R}^2) \times H^1(D)$ where $\mathbf{u}$ is divergence-free. Then*

$$\int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 \, d\mathbf{x} = \frac{1}{2\pi} \int_0^1 |\overline{\mathbf{u} \eta \cdot \hat{\mathbf{e}}_r} - F|^2 r \, dr + Q(\mathbf{u} \eta - g \hat{\mathbf{e}}_r),$$

where

$$F(r) = \frac{1}{2\pi r} \int_{D_r} f(\mathbf{x}) \, d\mathbf{x} \quad \text{and} \quad g(r, \theta) = \frac{1}{r} \int_0^r \rho f(\rho, \theta) \, d\rho$$

and

$$Q(v) = \min_{\varphi \in H^1(D)} \int_D \left| -\frac{1}{r} \partial_\theta \varphi + \mathbf{v} \cdot \hat{\mathbf{e}}_r - \overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r} \right|^2 + |\partial_r \varphi + \mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2 \, d\mathbf{x}.$$

Remark 3.2. A particular choice of test function we use often below is

$$\varphi = r \partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r) = r \int_0^\theta \mathbf{v} \cdot \hat{\mathbf{e}}_r(r, \theta') - \overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r}(r) \, d\theta'.$$

It sets the first integral in Q to zero, giving the bound

$$Q(\mathbf{v}) \leq \int_D |\partial_r(r \partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)) + \mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2 \, d\mathbf{x}.$$

Proof. We need the fact that

$$\int_D |\nabla \Delta^{-1} \nabla \cdot \mathbf{v}|^2 = \min_{\substack{\mathbf{m} \in L^2(D) \\ \nabla \cdot \mathbf{m} = 0}} \int_D |\mathbf{m} + \mathbf{v}|^2 = \min_{\varphi \in H^1(D)} \int_D |\nabla^\perp \varphi + \mathbf{v}|^2 \quad (3.4)$$

for any $\mathbf{v} \in L^2(D)$. A quick proof of it goes as follows: let $\zeta \in H_0^1(D)$ satisfy $\Delta \zeta = \nabla \cdot \mathbf{v}$ in D , and note that $\nabla \zeta$ is L^2 -orthogonal to divergence-free $\mathbf{m}$, including $\nabla \zeta - \mathbf{v}$. Hence,

$$\int_D |\mathbf{m} + \mathbf{v}|^2 = \int_D |\mathbf{m} + \mathbf{v} - \nabla \zeta|^2 + \int_D |\nabla \zeta|^2 \geq \int_D |\nabla \zeta|^2 = \int_D |\nabla \Delta^{-1} \nabla \cdot \mathbf{v}|^2$$

and equality holds for $\mathbf{m} = \nabla \zeta - \mathbf{v}$. This proves the first part of (3.4), and the rest of it follows from the usual representation of a divergence-free vector field $\mathbf{m}$ as the perpendicular gradient of a streamfunction φ .

Now using (3.4) and the definition of g , which satisfies $\partial_r(rg) = rf$, write that

$$\int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 = \int_D |\nabla \Delta^{-1} \nabla \cdot (\mathbf{u}\eta - g\hat{\mathbf{e}}_r)|^2 = \min_{\nabla \cdot \mathbf{m} = 0} \int_D |\mathbf{m} + \mathbf{v}|^2$$

where

$$\mathbf{v} = \mathbf{u}\eta - g\hat{\mathbf{e}}_r.$$

In the first step, we used that $\nabla \cdot (g\hat{\mathbf{e}}_r) = f$, which is clear from its expression in polar coordinates. Observe that a vector field $a(r)\hat{\mathbf{e}}_r$ is L^2 -orthogonal to any divergence-free $\mathbf{m}$. Indeed, writing

$$\mathbf{m} = \nabla^\perp \varphi = -\frac{1}{r} \partial_\theta \varphi \hat{\mathbf{e}}_r + \partial_r \varphi \hat{\mathbf{e}}_\theta$$

where φ is 2π -periodic in θ shows that

$$\int_D \mathbf{m} \cdot a(r)\hat{\mathbf{e}}_r = \int_0^1 \left[\int_0^{2\pi} -\frac{1}{r} \partial_\theta \varphi \, d\theta \right] a(r)r \, dr = 0.$$

This prompts the L^2 -orthogonal decomposition

$$\mathbf{v} = \overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r} \hat{\mathbf{e}}_r + \mathbf{w},$$

where $\mathbf{w}$ is the remainder. By orthogonality,

$$\begin{aligned} \min_{\nabla \cdot \mathbf{m} = 0} \int_D |\mathbf{m} + \mathbf{v}|^2 &= \min_{\nabla \cdot \mathbf{m} = 0} \int_D |\mathbf{m} + \mathbf{w}|^2 + \int_D |\overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r} \hat{\mathbf{e}}_r|^2 \\ &= Q(\mathbf{u}\eta - g\hat{\mathbf{e}}_r) + \int_0^1 |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - \bar{g}|^2 r \, dr, \end{aligned}$$

as $\bar{\cdot}$ yields functions of r alone. Since

$$\bar{g}(r) = \frac{1}{2\pi} \int_0^{2\pi} \left[\frac{1}{r} \int_0^r \rho f \, d\rho \right] d\theta = \frac{1}{2\pi r} \int_{D_r} f = F(r),$$

the result is proved. ■

We proceed to construct pairs $(\mathbf{u}, \eta)$ which are not necessarily admissible for the minimization in (3.2), but nevertheless do a good job at achieving $\mathbf{u} \cdot \nabla \eta \approx f$ in $H^{-1}(D)$ for a given $f \in L^2(D)$. We continue to use the functions $F = (1/2\pi r) \int_{D_r} f$ and $g = (1/r) \int_0^r \rho f$ from lemma 3.1. Given $n \in \mathbb{N}$, define the streamfunction

$$\psi(\mathbf{x}) = rg(r, \theta)\Psi(\theta) \quad \text{where} \quad \Psi(\theta) = \frac{\sqrt{2}}{n} \cos(n\theta), \quad (3.5)$$

whose velocity is

$$\mathbf{u} = \nabla^\perp \psi = -(\partial_\theta g \Psi + g \Psi') \hat{\mathbf{e}}_r + rf \Psi \hat{\mathbf{e}}_\theta. \quad (3.6)$$

Likewise, define the test function

$$\eta(\mathbf{x}) = -\Psi'(\theta). \quad (3.7)$$

Lemma 3.3. Let $f \in L^2(D)$. The velocity–test function pair $(\mathbf{u}, \eta)$ in (3.5)–(3.7) satisfies

$$\int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 \, d\mathbf{x} \lesssim \frac{1}{n^2} \int_D |f|^2 + |\nabla f|^2 \, d\mathbf{x}. \quad (3.8)$$

In particular, we have the estimates

$$|\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - F| \lesssim \frac{1}{n} \int_{D_r} |\nabla f| \, d\mathbf{x}, \quad (3.9)$$

$$\|\partial_r(r\partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g))\|_{L_\theta^1} \lesssim \frac{1}{n} \left(\int_{D_r} \frac{|f|}{r} + |\nabla f| \, d\mathbf{x} + r\|f\|_{L_\theta^1} + r^2\|\nabla f\|_{L_\theta^1} \right) \quad (3.10)$$

$$\text{and} \quad \|\mathbf{u}\eta \cdot \hat{\mathbf{e}}_\theta\|_{L_\theta^1} \lesssim \frac{r}{n} \|f\|_{L_\theta^1} \quad (3.11)$$

for a.e. $r > 0$. Furthermore, the streamfunction ψ and velocity $\mathbf{u}$ obey

$$|\psi| \lesssim \frac{1}{n} \int_0^r \rho |f| \, d\rho, \quad |\mathbf{u} \cdot \hat{\mathbf{e}}_r| \lesssim \int_0^r \frac{\rho |f|}{r} + \frac{\rho |\nabla f|}{n} \, d\rho \quad \text{and} \quad |\mathbf{u} \cdot \hat{\mathbf{e}}_\theta| \lesssim \frac{r}{n} |f|, \quad (3.12)$$

and

$$|\nabla \mathbf{u}| \lesssim \int_0^r \frac{n\rho |f|}{r^2} + \frac{n\rho |\nabla f|}{r} + \frac{\rho |\nabla \nabla f|}{n} \, d\rho + |f| + r|\nabla f|, \quad (3.13)$$

and the test function η obeys

$$|\eta| \lesssim 1 \quad \text{and} \quad |\nabla \eta| \lesssim \frac{n}{r} \quad (3.14)$$

for a.e. $\theta \in [0, 2\pi]$ and $r > 0$. The constants implicit in these estimates are independent of all parameters.

Proof. We start at the bottom of the claim and work backwards. The estimates in (3.14) are clear given the formula for η . To prove (3.12) and (3.13), we require the following inequalities involving $g = (1/r) \int_0^r \rho f$:

$$|g| \leq \frac{1}{r} \int_0^r \rho |f| \, d\rho, \quad |\partial_r g| \leq \frac{1}{r^2} \int_0^r \rho |f| \, d\rho + |f| \quad \text{and} \quad |\partial_\theta g| \leq \int_0^r \rho |\nabla f| \, d\rho \quad (3.15)$$

and

$$|\partial_{r\theta} g| \leq \frac{1}{r} \int_0^r \rho |\nabla f| \, d\rho + r|\nabla f| \quad \text{and} \quad |\partial_{\theta\theta} g| \leq r \int_0^r \rho |\nabla \nabla f| \, d\rho + \int_0^r \rho |\nabla f| \, d\rho. \quad (3.16)$$

Now by the definitions of ψ and $\mathbf{u}$,

$$|\psi| \lesssim \frac{r}{n} |g|, \quad |\mathbf{u} \cdot \hat{\mathbf{e}}_r| \lesssim \frac{1}{n} |\partial_\theta g| + |g| \quad \text{and} \quad |\mathbf{u} \cdot \hat{\mathbf{e}}_\theta| \lesssim \frac{r}{n} |f|,$$

and (3.12) is proved. Next, compute the gradient

$$\begin{aligned} \nabla \mathbf{u} &= -\nabla \hat{\mathbf{e}}_r (\partial_\theta g \Psi + g \Psi') - \hat{\mathbf{e}}_r \otimes \nabla (\partial_\theta g \Psi + g \Psi') + \nabla \hat{\mathbf{e}}_\theta (rf \Psi) + \hat{\mathbf{e}}_\theta \otimes \nabla (rf \Psi) \\ &= -\hat{\mathbf{e}}_\theta \otimes \hat{\mathbf{e}}_\theta \frac{1}{r} (\partial_\theta g \Psi + g \Psi') - \hat{\mathbf{e}}_r \otimes \nabla (\partial_\theta g \Psi + g \Psi') - \hat{\mathbf{e}}_r \otimes \hat{\mathbf{e}}_\theta f \Psi + \hat{\mathbf{e}}_\theta \otimes \nabla (rf \Psi), \end{aligned}$$

where

$$\nabla (\partial_\theta g \Psi + g \Psi') = \hat{\mathbf{e}}_r (\partial_{r\theta} g \Psi + \partial_r g \Psi') + \hat{\mathbf{e}}_\theta \frac{1}{r} (\partial_{\theta\theta} g \Psi + 2\partial_\theta g \Psi' + g \Psi'')$$

and

$$\nabla (rf \Psi) = \hat{\mathbf{e}}_r (f + r\partial_r f) \Psi + \hat{\mathbf{e}}_\theta (\partial_\theta f \Psi + f \Psi').$$

Hence,

$$\begin{aligned} |\nabla \mathbf{u}| &\leq \frac{1}{r} |\partial_\theta g \Psi + g \Psi'| + |\partial_{r\theta} g \Psi + \partial_r g \Psi'| + \frac{1}{r} |\partial_{\theta\theta} g \Psi + 2\partial_\theta g \Psi' + g \Psi''| \\ &\quad + |f \Psi| + |(f + r\partial_r f) \Psi| + |\partial_\theta f \Psi + f \Psi'| \\ &\lesssim (|g| + |\partial_\theta g|) \frac{n}{r} + |f| + |\partial_r g| + \left(|\partial_r f| + \frac{1}{r} |\partial_\theta f| + \frac{1}{r} |\partial_{r\theta} g| + \frac{1}{r^2} |\partial_{\theta\theta} g| \right) \frac{r}{n}. \end{aligned}$$

Combined with (3.15) and (3.16), this yields the desired estimates in (3.13).

It remains to show the first half of the claim. We require the inequalities

$$\|g\|_{L_\theta^1} \leq \frac{1}{r} \int_{D_r} |f| \, d\mathbf{x} \quad \text{and} \quad \|\partial_r g\|_{L_\theta^1} \leq \frac{1}{r^2} \int_{D_r} |f| \, d\mathbf{x} + \|f\|_{L_\theta^1} \quad (3.17)$$

and

$$\|\partial_\theta g\|_{L_\theta^1} \leq \int_{D_r} |\nabla f| \, d\mathbf{x} \quad \text{and} \quad \|\partial_{r\theta} g\|_{L_\theta^1} \leq \frac{1}{r} \int_{D_r} |\nabla f| \, d\mathbf{x} + r \|\nabla f\|_{L_\theta^1} \quad (3.18)$$

which follow from (3.15) and (3.16) by integration. Going back to the definitions,

$$\begin{aligned} \mathbf{u}\eta - g\hat{\mathbf{e}}_r &= (\partial_\theta g \Psi + g \Psi') \Psi' \hat{\mathbf{e}}_r - r f \Psi \Psi' \hat{\mathbf{e}}_\theta - g \hat{\mathbf{e}}_r \\ &= (\partial_\theta g \Psi \Psi' + g((\Psi')^2 - 1)) \hat{\mathbf{e}}_r - r f \Psi \Psi' \hat{\mathbf{e}}_\theta. \end{aligned}$$

Averaging the $\hat{\mathbf{e}}_r$ -component in θ and using that $F = \bar{g}$, there follows

$$\begin{aligned} |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - F| &= |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - \bar{g}| \leq \left| \int_0^{2\pi} \partial_\theta g \Psi \Psi' \right| + \left| \int_0^{2\pi} g((\Psi')^2 - 1) \right| \\ &\lesssim \|\partial_\theta g\|_{L_\theta^1} \frac{1}{n} + \left| \int_0^{2\pi} g((\Psi')^2 - 1) \right| \end{aligned}$$

since $|\Psi| \lesssim 1/n$ and $|\Psi'| \lesssim 1$. Introduce the operator ∂_θ^{-1} defined by

$$\partial_\theta^{-1} \varphi(r, \theta) = \int_0^\theta \varphi(r, \theta') - \bar{\varphi}(r) \, d\theta',$$

and observe that $\overline{(\Psi')^2} = 1$. Hence,

$$\begin{aligned} \left| \int_0^{2\pi} g((\Psi')^2 - 1) \right| &= \left| \int_0^{2\pi} \partial_\theta g \partial_\theta^{-1} (\Psi')^2 \right| \\ &\lesssim \|\partial_\theta g\|_{L_\theta^1} \|\partial_\theta^{-1} (\Psi')^2\|_{L_\theta^\infty} \lesssim \|\partial_\theta g\|_{L_\theta^1} \frac{1}{n} \end{aligned} \quad (3.19)$$

and (3.9) is proved.

Continuing with (3.10), write that

$$\begin{aligned} |\partial_r(r \partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g))| &\leq |\partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g)| + |r \partial_r \partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g)| \\ &= |\partial_\theta^{-1}[\partial_\theta g \Psi \Psi' + g((\Psi')^2 - 1)]| + r |\partial_r \partial_\theta^{-1}[\partial_\theta g \Psi \Psi' + g((\Psi')^2 - 1)]|. \end{aligned}$$

Of course,

$$|\partial_\theta^{-1}[\partial_\theta g \Psi \Psi']| \lesssim \|\partial_\theta g\|_{L_\theta^1} \frac{1}{n} \quad \text{and} \quad |\partial_r \partial_\theta^{-1}[g \Psi \Psi']| \lesssim \|\partial_{r\theta} g\|_{L_\theta^1} \frac{1}{n}.$$

Using the operator ∂_θ^{-1} again, we see that

$$|\partial_\theta^{-1}[g((\Psi')^2 - 1)]| \leq \left| \int_0^\theta g((\Psi')^2 - 1) \right| + \left| \int_0^{2\pi} g((\Psi')^2 - 1) \right|$$

and

$$|\partial_r \partial_\theta^{-1}[g((\Psi')^2 - 1)]| \leq \left| \int_0^\theta \partial_r g((\Psi')^2 - 1) \right| + \left| \int_0^{2\pi} \partial_r g((\Psi')^2 - 1) \right|.$$

The last two terms on the right-hand sides above are controlled by the inequality (3.19) and one just like it with $\partial_r g$ in place of g . To deal with the first two terms, we require the inequalities

$$\left\| \int_0^\theta g((\Psi')^2 - 1) \right\|_{L_\theta^1} \lesssim (\|g\|_{L_\theta^1} + \|\partial_\theta g\|_{L_\theta^1}) \frac{1}{n} \quad (3.20)$$

and

$$\left\| \int_0^\theta \partial_r g((\Psi')^2 - 1) \right\|_{L_\theta^1} \lesssim (\|\partial_r g\|_{L_\theta^1} + \|\partial_{r\theta} g\|_{L_\theta^1}) \frac{1}{n}. \quad (3.21)$$

To prove them, divide $[0, 2\pi]$ into the disjoint intervals $I_j = [2\pi j/n, 2\pi(j+1)/n]$ indexed by $j = 0, \dots, n-1$, and note that $(\Psi')^2 - 1$ is $2\pi/n$ -periodic and averages to zero over each I_j . Choosing k such that $\theta \in I_k$, write that

$$\begin{aligned} \left| \int_0^\theta g((\Psi')^2 - 1) \right| &\leq \sum_{j=0}^{k-1} \left| \int_{I_j} \left(g - \int_{I_j} g \right) ((\Psi')^2 - 1) \right| + \left| \int_{2\pi k/n}^\theta g((\Psi')^2 - 1) \right| \\ &\lesssim \sum_{j=0}^{k-1} \int_{I_j} \left| g - \int_{I_j} g \right| + \int_{2\pi k/n}^\theta |g| \lesssim \|\partial_\theta g\|_{L_\theta^1} \frac{1}{n} + \|g\|_{L_\theta^1(I_k)}. \end{aligned}$$

Remembering that $k = k(\theta)$ and integrating this bound over $\theta \in [0, 2\pi]$ yields (3.20). The proof of the inequality from (3.21) is much the same. Altogether, we have shown that

$$\|\partial_\theta^{-1}[\partial_\theta g \Psi \Psi' + g((\Psi')^2 - 1)]\|_{L_\theta^1} \lesssim (\|g\|_{L_\theta^1} + \|\partial_\theta g\|_{L_\theta^1}) \frac{1}{n} \quad (3.22)$$

and

$$\|\partial_r \partial_\theta^{-1}[\partial_\theta g \Psi \Psi' + g((\Psi')^2 - 1)]\|_{L_\theta^1} \lesssim (\|\partial_r g\|_{L_\theta^1} + \|\partial_r \partial_\theta g\|_{L_\theta^1}) \frac{1}{n} \quad (3.23)$$

which when combined with the inequalities in (3.17) and (3.18) lead to the estimate (3.10) in the claim. The estimate (3.11) follows from what we have already proved (namely, (3.12) and (3.14)).

Finally, we prove (3.8). By lemma 3.1 and the remark appearing immediately after,

$$\int_D |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 \lesssim \int_0^1 |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - F|^2 r \, dr + \int_D |\partial_r(r \partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g))|^2 + \int_D |\mathbf{u}\eta \cdot \hat{\mathbf{e}}_\theta|^2.$$

Applying (3.9)–(3.11) along with Jensen's inequality gives the bound

$$\int_0^1 |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - F|^2 r \, dr \lesssim \int_0^1 \left(\frac{r^2}{n^2} \int_{D_r} |\nabla f|^2 \right) r \, dr \lesssim \frac{1}{n^2} \int_D |\nabla f|^2,$$

as well as the bounds

$$\begin{aligned} &\int_D |\partial_r(r \partial_\theta^{-1}(\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r - g))|^2 \\ &\lesssim \int_0^1 \frac{1}{n^2} \left(\int_{D_r} |f|^2 + r^2 \int_{D_r} |\nabla f|^2 + r^2 \|f\|_{L_\theta^2}^2 + r^4 \|\nabla f\|_{L_\theta^2}^2 \right) r \, dr \\ &\lesssim \frac{1}{n^2} \int_D |f|^2 + |\nabla f|^2 \end{aligned}$$

and

$$\int_D |\mathbf{u}\eta \cdot \hat{\mathbf{e}}_\theta|^2 \lesssim \int_0^1 \left(\frac{r^2}{n^2} \|f\|_{L_\theta^2}^2 \right) r \, dr \lesssim \frac{1}{n^2} \int_D |f|^2.$$

The proof is complete. ■

As a quick application of this last result, we show that the velocities in (3.6) have finite enstrophy if f is sufficiently regular. By the pointwise estimate from (3.13) and Jensen's inequality,

$$|\nabla \mathbf{u}|^2 \lesssim \int_0^r \left[\frac{n^2 \rho^2}{r^3} |f|^2 + \frac{n^2 \rho^2}{r} |\nabla f|^2 + \frac{\rho^2 r}{n^2} |\nabla \nabla f|^2 \right] d\rho + |f|^2 + r^2 |\nabla f|^2.$$

Hence,

$$\begin{aligned} & \int_D |\nabla \mathbf{u}|^2 \, d\mathbf{x} \\ & \lesssim \int_0^{2\pi} \int_{0 \leq \rho \leq r \leq 1} \left[\frac{n^2 \rho^2}{r^2} |f(\rho, \theta)|^2 + n^2 \rho^2 |\nabla f(\rho, \theta)|^2 + \frac{\rho^2 r^2}{n^2} |\nabla \nabla f(\rho, \theta)|^2 \right] dr \, d\rho \, d\theta \\ & \quad + \int_D |f|^2 + r^2 |\nabla f|^2 \, d\mathbf{x} \\ & \lesssim \int_D n^2 |f|^2 + n^2 r |\nabla f|^2 + \frac{r}{n^2} |\nabla \nabla f|^2 \, d\mathbf{x}. \end{aligned}$$

4. Unsteady branching flows for enstrophy-constrained cooling

Section 3 considered the steady optimal cooling problem and explained how to find ‘approximate H^{-1} -solutions’ to the corresponding advection equation $\mathbf{u} \cdot \nabla \eta = f$. We now return to the original unsteady setting of theorem 1.1 to prove our upper bound on

$$\min_{\substack{\mathbf{u}(\mathbf{x}, t) \\ \langle |\nabla \mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} = 0 \text{ at } \partial D}} \langle |\nabla T|^2 \rangle$$

for a general source $f(\mathbf{x}, t)$. We do so by constructing a family of well-chosen branching flows $\{\mathbf{u}_{Pe}\}$ whose temperatures $\{T_{Pe}\}$ satisfy

$$\langle |\nabla T_{Pe}|^2 \rangle \lesssim \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle \frac{\log^{4/3} Pe}{Pe^{2/3}} \quad \text{with} \quad \langle |\nabla \mathbf{u}_{Pe}|^2 \rangle = Pe^2.$$

Section 4a starts by defining a general family of convection roll-based branching flows (see the bottom row of figure 1). Section 4b estimates their cooling and §4c optimizes over their free parameters. The upper bound from theorem 1.1 is finally proved at the end of this section.

Picking up where we left off in §3, recall the upper bound

$$\min_{\substack{\mathbf{u}(\mathbf{x}, t) \\ \langle |\nabla \mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} = 0 \text{ at } \partial D}} \langle |\nabla T|^2 \rangle \leq \min_{\substack{\mathbf{u}(\mathbf{x}, t), \eta(\mathbf{x}, t) \\ \mathbf{u} = 0, \eta = 0 \text{ at } \partial D}} \left\langle |\nabla \Delta^{-1}[(\partial_t + \mathbf{u} \cdot \nabla)\eta - f]|^2 + \frac{\langle |\nabla \mathbf{u}|^2 \rangle}{Pe^2} |\nabla \eta|^2 \right\rangle,$$

where on the right-hand side the magnitude of $\mathbf{u}$ is unconstrained. This bound follows from proposition 2.1 and the change of variables $(\mathbf{u}, \eta) \rightarrow (\lambda_{Pe} \mathbf{u}, \lambda_{Pe}^{-1} \eta)$ with $\lambda_{Pe} = Pe / \sqrt{\langle |\nabla \mathbf{u}|^2 \rangle}$, as explained in §3a. A special case occurs for a steady test function $\eta(\mathbf{x})$: the temperature field T associated with $\lambda_{Pe} \mathbf{u}$ obeys

$$\langle |\nabla T|^2 \rangle \leq \langle |\nabla \Delta^{-1}[\mathbf{u} \cdot \nabla \eta - f]|^2 \rangle + \frac{1}{Pe^2} \langle |\nabla \mathbf{u}|^2 \rangle \int_D |\nabla \eta|^2$$

for all $\eta \in H_0^1(D)$. We proceed to define our branching flows.

(a) Branching flows

By a *branching flow* $\mathbf{u}(\mathbf{x}, t)$ and a corresponding (steady) test function $\eta(\mathbf{x})$, we mean a divergence-free velocity field

$$\mathbf{u} = \nabla^\perp \psi \quad \text{with} \quad \psi(\mathbf{x}, t) = \sum_{k=1}^n \chi_k(r) \psi_k(\mathbf{x}, t),$$

and the scalar function

$$\eta(\mathbf{x}) = \sum_{k=1}^n \chi_k(r) \eta_k(\mathbf{x}),$$

where $\{\chi_k\}$, $\{\psi_k\}$ and $\{\eta_k\}$ are as follows. Let

$$F(r, t) = \frac{1}{2\pi r} \int_{D_r} f(\mathbf{x}, t) \, d\mathbf{x} \quad \text{and} \quad g(r, \theta, t) = \int_0^r \rho f(\rho, \theta, t) \, d\rho, \quad (4.1)$$

and set

$$\psi_k = r g \Psi_k(\theta), \quad \eta_k = -\Psi_k'(\theta) \quad \text{and} \quad \Psi_k(\theta) = \sqrt{2} l_k \cos\left(\frac{\theta}{l_k}\right) \quad \text{for } k = 1, \dots, n.$$

The parameters $\{l_k^{-1}\} \subset \mathbb{N}$ and $n \in \mathbb{N}$ are free, and will eventually be optimized when it comes time to prove theorem 1.1. The functions $\{\chi_k\}$ are described in the paragraph after the next.

To help organize the discussion, we always assume that

$$l_1 > l_2 > \dots > l_n. \quad (4.2)$$

We label the largest and smallest scales as

$$l_{\text{bulk}} = l_1 \quad \text{and} \quad l_{\text{bl}} = l_n,$$

noting that they occur in the bulk of the disc and in a boundary layer near $r = 1$, respectively. The individual velocities

$$\mathbf{u}_k = \nabla^\perp \psi_k = -(\partial_\theta g \Psi_k + g \Psi_k') \hat{\mathbf{e}}_r + r f \Psi_k \hat{\mathbf{e}}_\theta$$

are simply unsteady versions of the ones occurring in our prior discussion of roll-like flows, and so are governed by the estimates in lemma 3.3. The individual test functions

$$\eta_k = -\Psi_k'$$

are also like those in the lemma.

The new ingredients are the functions $\{\chi_k\}$ which we use to interpolate between the individual building blocks listed above. We use a family of smooth and compactly supported ‘cutoff functions’ defined via a choice of points $\{r_k\}$, satisfying

$$\frac{1}{2} < r_1 < r_2 < \dots < r_n < 1 \quad (4.3)$$

with

$$r_1 = r_{\text{bulk}} \quad \text{and} \quad r_n = r_{\text{bl}}.$$

These functions can be quite general, but to fix ideas we let

$$\text{supp } \chi_1 \subset (0, r_2), \quad \text{supp } \chi_n \subset (r_{n-1}, 1) \quad (4.4)$$

and

$$\text{supp } \chi_k \subset (r_{k-1}, r_{k+1}) \quad \text{for } k = 2, \dots, n-1.$$

We also let

$$\sum_{k=1}^n (\chi_k(r))^2 = 1 \quad \forall r \in (0, r_{\text{bl}}), \quad (4.5)$$

and take

$$\chi_k \chi_{k'} \neq 0 \iff |k - k'| \leq 1. \quad (4.6)$$

This last condition simplifies the calculation of products such as $\mathbf{u}\eta$.

To specify our cutoff functions further, introduce the lengths

$$\delta_{\text{bulk}} = 1 - r_{\text{bulk}}, \quad \delta_{\text{bl}} = 1 - r_{\text{bl}} \quad \text{and} \quad \delta_k = r_{k+1} - r_k \quad \text{for } k = 1, \dots, n-1$$

and, following the pattern, call

$$\delta_n = \delta_{\text{bl}}.$$

Let

$$|\chi_k| \vee |\chi_{k+1}| \lesssim 1, \quad |\chi'_k| \vee |\chi'_{k+1}| \lesssim \frac{1}{\delta_k} \quad \text{and} \quad |\chi''_k| \vee |\chi''_{k+1}| \lesssim \frac{1}{\delta_k^2} \quad \forall r \in (r_k, r_{k+1}), \quad (4.7)$$

for $k = 1, \dots, n-1$, and let

$$|\chi_n| \lesssim 1, \quad |\chi'_n| \lesssim \frac{1}{\delta_{\text{bl}}} \quad \text{and} \quad |\chi''_n| \lesssim \frac{1}{\delta_{\text{bl}}^2} \quad \forall r \in (r_{\text{bl}}, 1). \quad (4.8)$$

The constants implicit in these hypotheses are independent of all parameters.

The reader looking for specific cutoff functions should consult [43, §5.1]. There, we describe a similar branching flow in a fluid layer, the z -coordinate of which is analogous to r . The exact choice of these functions does not affect the scaling of our bounds in Pe , but does affect their prefactors.

(b) Upper bounds on branching flows

Having defined our branching flows, we proceed to estimate their cooling using the results of §2. We write the estimate in terms of a continuously varying version of the parameters $\{l_k\}$ and $\{r_k\}$, given by

$$\ell(r) = l_k \frac{r_{k+1} - r}{r_{k+1} - r_k} + l_{k+1} \frac{r - r_k}{r_{k+1} - r_k} \quad \text{for } r \in [r_k, r_{k+1}] \quad (4.9)$$

and $k = 1, \dots, n-1$. By construction, $\ell(r_k) = l_k$ for each k . Recall $\delta_k = r_{k+1} - r_k$.

Proposition 4.1. *Let $\{\mathbf{u}\}$ be a family of branching flows as defined in §4a, whose parameters $\{l_k\}_{k=1}^n$ and $\{r_k\}_{k=1}^n$ obey*

$$|l_{k+1} - l_k| \sim l_{k+1} \sim l_k \quad \text{and} \quad \delta_{k+1} \sim \delta_k \quad \text{for } k = 1, \dots, n-1 \quad (4.10)$$

and

$$l_k \lesssim \delta_k \quad \text{for } k = 1, \dots, n \quad (4.11)$$

with fixed numerical prefactors. Define the rescaled velocities

$$\lambda_{Pe} \mathbf{u} \quad \text{with} \quad \lambda_{Pe} = \frac{Pe}{\sqrt{\langle |\nabla \mathbf{u}|^2 \rangle}},$$

and let their temperature fields T_{Pe} solve

$$\begin{cases} \partial_t T_{Pe} + \lambda_{Pe} \mathbf{u} \cdot \nabla T_{Pe} = \Delta T_{Pe} + f & \text{in } D \\ T_{Pe} = 0 & \text{at } \partial D \end{cases}$$

weakly with arbitrary L^2 -initial data. The estimate

$$\langle |\nabla T_{Pe}|^2 \rangle \lesssim C_0(f) \left[l_{\text{bulk}}^2 + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell'(r))^2 dr + \delta_{\text{bl}} + \frac{1}{Pe^2} \left(\frac{1}{l_{\text{bulk}}^2} + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right)^2 \right]$$

holds with

$$C_0(f) = \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle$$

and a numerical prefactor depending only on the ones from (4.10) to (4.11).

By proposition 2.1, we already know that the rescaled velocities $\lambda_{Pe}\mathbf{u}$ achieve

$$\langle |\nabla T_{Pe}|^2 \rangle \leq \langle |\nabla \Delta^{-1}[\mathbf{u} \cdot \nabla \eta - f]|^2 \rangle + \frac{1}{Pe^2} \langle |\nabla \mathbf{u}|^2 \rangle \int_D |\nabla \eta|^2 \quad \forall \eta \in H_0^1(D).$$

To prove proposition 4.1, we shall plug in the test functions η from §4a, and estimate the ensuing mess of terms. We handle the ‘advection term’ involving $\mathbf{u} \cdot \nabla \eta - f$ in §4bi and the ‘enstrophy term’ involving $\langle |\nabla \mathbf{u}|^2 \rangle$ in §4bii.

(i) The advection term

Averaging the result of lemma 3.1 shows that

$$\langle |\nabla \Delta^{-1}(\mathbf{u} \cdot \nabla \eta - f)|^2 \rangle \leq \langle |\overline{\mathbf{u} \cdot \hat{\mathbf{e}}_r \eta} - F|^2 \rangle + \langle Q(\mathbf{u}\eta - g\hat{\mathbf{e}}_r) \rangle,$$

where for the reader’s convenience we repeat the definition of the quadratic form:

$$Q(\mathbf{v}) = \min_{\varphi \in H^1(D)} \int_D \left| -\frac{1}{r} \partial_\theta \varphi + \mathbf{v} \cdot \hat{\mathbf{e}}_r - \overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r} \right|^2 + |\partial_r \varphi + \mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2 \, \mathrm{d}\mathbf{x}. \quad (4.12)$$

We bound the first average in lemma 4.2 and the second one in lemma 4.3.

Lemma 4.2. *Every branching flow–test function pair $(\mathbf{u}, \eta)$ defined in §4a satisfies*

$$\langle |\overline{\mathbf{u} \cdot \hat{\mathbf{e}}_r \eta} - F|^2 \rangle \lesssim l_{\text{bulk}}^2 \langle |\nabla f|^2 \rangle + \delta_{\text{bl}} \langle |f|^2 \rangle.$$

Proof. It suffices to prove the steady analogue of the result at a.e. time, i.e.

$$\int_D |\overline{\mathbf{u} \cdot \hat{\mathbf{e}}_r \eta} - F|^2 \lesssim l_{\text{bulk}}^2 \int_D |\nabla f|^2 + \delta_{\text{bl}} \int_D |f|^2.$$

From the definitions of $\mathbf{u}$ and η and our assumption (4.6),

$$\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r = \sum_{|k-k'| \leq 1} \chi_k \chi_{k'} \mathbf{u}_k \eta_{k'} \cdot \hat{\mathbf{e}}_r$$

so that

$$\left| \overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - \sum_{k=1}^n \chi_k^2 F \right| \leq \sum_{k=1}^n \chi_k^2 |\overline{\mathbf{u}_k \eta_k \cdot \hat{\mathbf{e}}_r} - F| + \sum_{\substack{|k-k'| \leq 1 \\ k \neq k'}} |\chi_k \chi_{k'}| |\overline{\mathbf{u}_k \eta_{k'} \cdot \hat{\mathbf{e}}_r}|.$$

The first term is handled by (3.9), which shows that

$$|\overline{\mathbf{u}_k \eta_k \cdot \hat{\mathbf{e}}_r} - F| \lesssim l_k \int_{D_r} |\nabla f|.$$

To control the second term, write using the L^2 -orthogonality of $\Psi'_k(\theta)$ and $\Psi'_{k'}(\theta)$ that

$$\begin{aligned} |\overline{\mathbf{u}_k \eta_{k'} \cdot \hat{\mathbf{e}}_r}| &= \left| \int_0^{2\pi} \partial_\theta g \Psi_k \Psi'_{k'} + g \Psi'_k \Psi'_{k'} \right| \\ &\leq \left| \int_0^{2\pi} \partial_\theta g \Psi_k \Psi'_{k'} \right| + \left| \int_0^{2\pi} \partial_\theta g \partial_\theta^{-1} (\Psi'_k \Psi'_{k'}) \right| \\ &\lesssim \|\partial_\theta g\|_{L_\theta^1} (l_k \vee l_{k'}) \lesssim (l_k \vee l_{k'}) \int_{D_r} |\nabla f| \end{aligned}$$

where in the last step we used (3.18). Adding up,

$$\left| \overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - \sum_{k=1}^n \chi_k^2 F \right| \lesssim \sum_{|k-k'| \leq 1} |\chi_k \chi_{k'}| (l_k \vee l_{k'}) \int_{D_r} |\nabla f|$$

for $r > 0$.

Continuing, we have by the triangle and Jensen's inequality that

$$\begin{aligned} \int_D |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - F|^2 \, \mathrm{d}\mathbf{x} &\lesssim \int_0^1 |\overline{\mathbf{u}\eta \cdot \hat{\mathbf{e}}_r} - \sum_{k=1}^n \chi_k^2 F|^2 r \, \mathrm{d}r + \int_0^1 \left| 1 - \sum_{k=1}^n \chi_k^2 \right|^2 |F|^2 r \, \mathrm{d}r \\ &\lesssim \sum_{|k-k'|\leq 1} \int_0^1 |\chi_k \chi_{k'}|^2 (l_k \vee l_{k'})^2 r^3 \int_{D_r} |\nabla f|^2 + \int_0^1 \left| 1 - \sum_{k=1}^n \chi_k^2 \right|^2 |F|^2 r \\ &\lesssim l_{\text{bulk}}^2 \int_D |\nabla f|^2 + \delta_{\text{bl}} \int_{r_{\text{bl}}}^1 |F|^2 r. \end{aligned}$$

Recalling the definition of $F = (1/2\pi r) \int_{D_r} f$ from (4.1), we use Jensen's inequality once more to get that

$$\int_{r_{\text{bl}}}^1 |F|^2 r \lesssim \int_{r_{\text{bl}}}^1 \left[\int_{D_r} |f|^2 \right] r \lesssim \int_D |f|^2.$$

The result follows. ■

Lemma 4.3. *Under the assumptions in (4.10), the branching flow—test function pairs $(\mathbf{u}, \eta)$ from §4a satisfy*

$$\langle Q(\mathbf{u}\eta - g\hat{\mathbf{e}}_r) \rangle \lesssim \left(l_{\text{bulk}}^2 + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell'(r))^2 \, \mathrm{d}r + \frac{l_{\text{bl}}^2}{\delta_{\text{bl}}} \right) \langle |f|^2 + |\nabla f|^2 \rangle + \delta_{\text{bl}} \langle |f|^2 \rangle$$

with a constant depending only on those in (4.10).

Proof. Again we argue a.e. in time. Begin with the bound

$$\begin{aligned} Q(\mathbf{u}\eta - g\hat{\mathbf{e}}_r) &\lesssim Q(\mathbf{u}\eta - \sum_{k=1}^n \chi_k^2 g\hat{\mathbf{e}}_r) + \int_D \left| \left(\sum_{k=1}^n \chi_k^2 - 1 \right) g \right|^2 \\ &\lesssim Q(\mathbf{u}\eta - \sum_{k=1}^n \chi_k^2 g\hat{\mathbf{e}}_r) + \delta_{\text{bl}} \int_D |f|^2, \end{aligned}$$

where in the first step we applied the definition (4.12) of the quadratic form Q , and in the second step we used the formula $g = f_0^r \rho f$ from (4.1) along with Jensen's inequality and (4.5). Calling

$$\mathbf{v} = \mathbf{u}\eta - \sum_{k=1}^n \chi_k^2 g\hat{\mathbf{e}}_r$$

and using

$$\varphi = r\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r) = r \int_0^\theta \mathbf{v} \cdot \hat{\mathbf{e}}_r(r, \theta') - \overline{\mathbf{v} \cdot \hat{\mathbf{e}}_r(r)} \, \mathrm{d}\theta'$$

as in remark 3.2, we deduce that

$$\begin{aligned} Q(\mathbf{v}) &\leq \int_D |\partial_r(r\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)) + \mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2 \\ &\lesssim \int_D |\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)|^2 + \int_D r^2 |\partial_r\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)|^2 + \int_D |\mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2. \end{aligned}$$

We estimate these integrals one-by-one.

For the first, note using the definitions from §4a and in particular (4.6) that

$$\begin{aligned}\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r) &= \partial_\theta^{-1}(\mathbf{u}\boldsymbol{\eta} \cdot \hat{\mathbf{e}}_r - \sum_{k=1}^n \chi_k^2 g) = \sum_{|k-k'|\leq 1} \chi_k \chi_{k'} \partial_\theta^{-1}[\mathbf{u}_k \boldsymbol{\eta}_{k'} \cdot \hat{\mathbf{e}}_r - \delta_{kk'} g] \\ &= \sum_{|k-k'|\leq 1} \chi_k \chi_{k'} \partial_\theta^{-1}[\partial_\theta g \Psi_k \Psi'_{k'} + g(\Psi'_k \Psi'_{k'} - \delta_{kk'})]\end{aligned}$$

where $\delta_{kk'}$ is the Kronecker delta function (one if $k = k'$, zero otherwise). Arguing as in the proof of lemma 3.3—see the paragraph leading up to (3.22)—we write that

$$\begin{aligned}\|\partial_\theta^{-1}[\partial_\theta g \Psi_k \Psi'_{k'} + g(\Psi'_k \Psi'_{k'} - \delta_{kk'})]\|_{L_\theta^1} &\lesssim (l_k \vee l_{k'}) (\|g\|_{L_\theta^1} + \|\partial_\theta g\|_{L_\theta^1}) \\ &\lesssim (l_k \vee l_{k'}) \left(\frac{1}{r} \int_{D_r} |f| + \int_{D_r} |\nabla f| \right)\end{aligned}\quad (4.13)$$

where in the last step we used (3.17) and (3.18). Squaring and integrating, there follows

$$\begin{aligned}\int_D |\partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)|^2 &\lesssim \sum_{|k-k'|\leq 1} \int_0^1 |\chi_k \chi_{k'}|^2 (l_k \vee l_{k'})^2 \left(\int_{D_r} |f|^2 + r^2 \int_{D_r} |\nabla f|^2 \right) r \, dr \\ &\lesssim l_{\text{bulk}}^2 \int_D |f|^2 + |\nabla f|^2.\end{aligned}$$

Continuing with the second integral, write using the product rule that

$$\begin{aligned}\partial_r \partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r) &= \sum_{|k-k'|\leq 1} (\chi_k \chi_{k'})' \partial_\theta^{-1}[\partial_\theta g \Psi_k \Psi'_{k'} + g(\Psi'_k \Psi'_{k'} - \delta_{kk'})] \\ &\quad + \sum_{|k-k'|\leq 1} \chi_k \chi_{k'} \partial_r \partial_\theta^{-1}[\partial_\theta g \Psi_k \Psi'_{k'} + g(\Psi'_k \Psi'_{k'} - \delta_{kk'})]\end{aligned}$$

and note the estimate

$$\begin{aligned}\|\partial_r \partial_\theta^{-1}[\partial_\theta g \Psi_k \Psi'_{k'} + g(\Psi'_k \Psi'_{k'} - \delta_{kk'})]\|_{L_\theta^1} &\lesssim (l_k \vee l_{k'}) (\|\partial_r g\|_{L_\theta^1} + \|\partial_{r\theta} g\|_{L_\theta^1}) \\ &\lesssim (l_k \vee l_{k'}) \left(\frac{1}{r^2} \int_{D_r} |f| + \|f\|_{L_\theta^1} + \frac{1}{r} \int_{D_r} |\nabla f| + r \|\nabla f\|_{L_\theta^1} \right)\end{aligned}\quad (4.14)$$

holds in addition to (4.13). Its proof is essentially the same, and again we point the reader to the paragraph leading up to (3.23) for the details. Using both (4.13) and (4.14), we get that

$$\begin{aligned}\int_D r^2 |\partial_r \partial_\theta^{-1}(\mathbf{v} \cdot \hat{\mathbf{e}}_r)|^2 &\lesssim \sum_{|k-k'|\leq 1} \int_0^1 |(\chi_k \chi_{k'})'|^2 (l_k \vee l_{k'})^2 \left(r^2 \int_{D_r} |f|^2 + r^4 \int_{D_r} |\nabla f|^2 \right) r \, dr \\ &\quad + \sum_{|k-k'|\leq 1} \int_0^1 |\chi_k \chi_{k'}|^2 (l_k \vee l_{k'})^2 \left(\int_{D_r} |f|^2 + r^2 \int_{D_r} |\nabla f|^2 + r^2 \|f\|_{L_\theta^2}^2 + r^4 \|\nabla f\|_{L_\theta^2}^2 \right) r \, dr \\ &\lesssim \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell'(r))^2 \, dr + \frac{l_{\text{bl}}^2}{\delta_{\text{bl}}} + l_{\text{bulk}}^2 \right) \int_D |f|^2 + |\nabla f|^2.\end{aligned}$$

Note we used the definition of ℓ and our hypotheses that $|l_{k+1} - l_k| \sim l_{k+1} \sim l_k$ and $\delta_{k+1} \sim \delta_k$ to bring in ℓ' .

Finally, since

$$\begin{aligned}\mathbf{v} \cdot \hat{\mathbf{e}}_\theta &= \mathbf{u} \eta \cdot \hat{\mathbf{e}}_\theta = \sum_{|k-k'| \leq 1} (\chi'_k \psi_k + \chi_k \mathbf{u}_k \cdot \hat{\mathbf{e}}_\theta) \chi_{k'} \eta_{k'} \\ &= - \sum_{|k-k'| \leq 1} \chi'_k \chi_{k'} r g \psi_k \psi'_{k'} + \chi_k \chi_{k'} r f \psi_k \psi'_{k'},\end{aligned}$$

we get by a completely analogous argument that

$$\begin{aligned}\int_D |\mathbf{v} \cdot \hat{\mathbf{e}}_\theta|^2 &\lesssim \sum_{|k-k'| \leq 1} \int_0^1 (|\chi'_k \chi_{k'}|^2 l_k^2 r^2 \|g\|_{L^2_\theta}^2 + |\chi_k \chi_{k'}|^2 l_k^2 r^2 \|f\|_{L^2_\theta}^2) r \, dr \\ &\lesssim \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell'(r))^2 \, dr + \frac{l_{\text{bl}}^2}{\delta_{\text{bl}}} + l_{\text{bulk}}^2 \right) \int_D |f|^2.\end{aligned}$$

In the last step, we applied Jensen's inequality with $g = f_0^r \rho f$. Adding up the estimates and averaging in time proves the result. ■

Combining lemmas 4.2 and 4.3 with our hypothesis that $l_{\text{bl}} \lesssim \delta_{\text{bl}}$ from (4.11) proves the first part of the estimate in proposition 4.1.

(ii) The enstrophy term

We now estimate the gradients of $\mathbf{u}$ and η .

Lemma 4.4. *Under the assumptions in (4.10) and (4.11), the branching flow—test function pairs $(\mathbf{u}, \eta)$ from §4a satisfy*

$$\langle |\nabla \mathbf{u}|^2 \rangle \lesssim \left(\frac{1}{l_{\text{bulk}}^2} + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} \, dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right) \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle$$

and

$$\langle |\nabla \eta|^2 \rangle \lesssim \frac{1}{l_{\text{bulk}}^2} + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} \, dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2}.$$

The constants implicit in these estimates depend only on those in (4.10) and (4.11).

Proof. We start with the formulas

$$\mathbf{u} = \sum_{k=1}^n \chi_k \mathbf{u}_k + \chi'_k \psi_k \hat{\mathbf{e}}_\theta$$

and

$$\nabla \mathbf{u} = \sum_{k=1}^n \chi_k \nabla \mathbf{u}_k + \chi'_k (\mathbf{u}_k \otimes \hat{\mathbf{e}}_r - \hat{\mathbf{e}}_\theta \otimes \mathbf{u}_k^\perp) + \chi''_k \psi_k \hat{\mathbf{e}}_\theta \otimes \hat{\mathbf{e}}_r,$$

which follow from the definitions in §4a and the fact that $\mathbf{u}_k^\perp = -\nabla \psi_k$ as $(\cdot)^\perp$ is a counterclockwise rotation by $\pi/2$. Looking back at the estimates in (3.12) and (3.13), we see that

$$\begin{aligned}|\nabla \mathbf{u}| &\lesssim \sum_{k=1}^n |\chi_k| |\nabla \mathbf{u}_k| + |\chi'_k| |\mathbf{u}_k| + |\chi''_k| |\psi_k| \\ &\lesssim \sum_{k=1}^n |\chi_k| \left(\frac{A_1}{r l_k} + \frac{A_2}{l_k} + r l_k A_3 + |f| + r |\nabla f| \right) \\ &\quad + |\chi'_k| (A_1 + r l_k A_2 + r l_k |f|) + |\chi''_k| r l_k A_1\end{aligned}$$

where

$$A_1 = \frac{1}{r} \int_0^r \rho |f| \, d\rho, \quad A_2 = \frac{1}{r} \int_0^r \rho |\nabla f| \, d\rho \quad \text{and} \quad A_3 = \frac{1}{r} \int_0^r \rho |\nabla \nabla f| \, d\rho.$$

Squaring and integrating, there follows

$$\begin{aligned} \int_D |\nabla \mathbf{u}|^2 &\lesssim \sum_{k=1}^n \int_0^1 |\chi_k|^2 \left(\frac{1}{r^2 l_k^2} \|A_1\|_{L_\theta^2}^2 + \frac{1}{l_k^2} \|A_2\|_{L_\theta^2}^2 + r^2 l_k^2 \|A_3\|_{L_\theta^2}^2 + \|f\|_{L_\theta^2}^2 + r^2 \|\nabla f\|_{L_\theta^2}^2 \right) r \, dr \\ &\quad + \sum_{k=1}^n \int_0^1 |\chi_k'|^2 (\|A_1\|_{L_\theta^2}^2 + r^2 l_k^2 \|A_2\|_{L_\theta^2}^2 + r^2 l_k^2 \|f\|_{L_\theta^2}^2) r \, dr \\ &\quad + \sum_{k=1}^n \int_0^1 |\chi_k''|^2 r^2 l_k^2 \|A_1\|_{L_\theta^2}^2 r \, dr \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

We bound these three sums in turn, using the estimates

$$\|A_1\|_{L_\theta^2}^2 \leq \int_0^r \rho^2 \|f\|_{L_\theta^2}^2 \, d\rho \lesssim \int_{D_r} |f|^2 \, d\mathbf{x}, \quad (4.15)$$

$$\|A_2\|_{L_\theta^2}^2 \leq \int_0^r \rho^2 \|\nabla f\|_{L_\theta^2}^2 \, d\rho \lesssim \int_{D_r} |\nabla f|^2 \, d\mathbf{x} \quad (4.16)$$

and $\|A_3\|_{L_\theta^2}^2 \leq \int_0^r \rho^2 \|\nabla \nabla f\|_{L_\theta^2}^2 \, d\rho \lesssim \int_{D_r} |\nabla \nabla f|^2 \, d\mathbf{x}. \quad (4.17)$

Note these follow from the definitions via Jensen's inequality.

The first sum I_1 involves the cutoff functions $\{\chi_k\}$ directly. Focusing on $r \in (r_{\text{bulk}}, r_{\text{bl}})$ for now, which are always larger than $1/2$ and no larger than 1 , we estimate

$$\begin{aligned} &\sum_{k=1}^n \int_{r_{\text{bulk}}}^{r_{\text{bl}}} |\chi_k|^2 \left(\frac{1}{r^2 l_k^2} \|A_1\|_{L_\theta^2}^2 + \frac{1}{l_k^2} \|A_2\|_{L_\theta^2}^2 + r^2 l_k^2 \|A_3\|_{L_\theta^2}^2 \right) r \, dr \\ &\lesssim \sum_{k=1}^n \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{|\chi_k|^2}{l_k^2} (\|A_1\|_{L_\theta^2}^2 + \|A_2\|_{L_\theta^2}^2 + \|A_3\|_{L_\theta^2}^2) \\ &\lesssim \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} \, dr \cdot \int_D |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \end{aligned}$$

by our definition of ℓ and the hypothesis that $l_{k+1} \sim l_k$. For $r \in (0, r_{\text{bulk}})$, which belong only to the support of χ_1 , we use Fubini's theorem and the first parts of (4.15)–(4.17) to write that

$$\begin{aligned} &\int_0^{r_{\text{bulk}}} |\chi_1|^2 \left(\frac{1}{r^2 l_1^2} \|A_1\|_{L_\theta^2}^2 + \frac{1}{l_1^2} \|A_2\|_{L_\theta^2}^2 + r^2 l_1^2 \|A_3\|_{L_\theta^2}^2 \right) r \, dr \\ &\lesssim \int_{0 \leq \rho \leq r \leq r_{\text{bulk}}} \left[\frac{\rho^2}{r^2 l_{\text{bulk}}^2} \|f(\rho, \cdot)\|_{L_\theta^2}^2 + \frac{\rho^2}{l_{\text{bulk}}^2} \|\nabla f(\rho, \cdot)\|_{L_\theta^2}^2 + \frac{l_{\text{bulk}}^2 \rho^2}{r^2} \|\nabla \nabla f(\rho, \cdot)\|_{L_\theta^2}^2 \right] d\rho \, dr \\ &\lesssim \frac{1}{l_{\text{bulk}}^2} \int_D |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2, \end{aligned}$$

similarly to what we did at the very end of §3. Finally,

$$\begin{aligned} &\int_{r_{\text{bl}}}^1 |\chi_n|^2 \left(\frac{1}{r^2 l_n^2} \|A_1\|_{L_\theta^2}^2 + \frac{1}{l_n^2} \|A_2\|_{L_\theta^2}^2 + r^2 l_n^2 \|A_3\|_{L_\theta^2}^2 \right) r \, dr \\ &\lesssim \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \int_D |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2. \end{aligned}$$

The bound

$$\sum_{k=1}^n \int_0^1 |\chi_k|^2 (\|f\|_{L_\theta^2}^2 + r^2 \|\nabla f\|_{L_\theta^2}^2) r \, dr \lesssim \int_D |f|^2 + |\nabla f|^2$$

is clear. Altogether,

$$I_1 \lesssim \left(\frac{1}{l_{\text{bulk}}^2} + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right) \cdot \int_D |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2.$$

The second sum I_2 involves the derivatives $\{\chi'_k\}$. This time, we only need to handle $r > r_{\text{bulk}}$. Using the second parts of (4.15) and (4.16), we get that

$$\begin{aligned} I_2 &= \sum_{k=1}^n \int_{r_{\text{bulk}}}^1 |\chi'_k|^2 (|A_1|_{L_\theta^2}^2 + r^2 l_k^2 |A_2|_{L_\theta^2}^2 + r^2 l_k^2 |f|_{L_\theta^2}^2) r dr \\ &\lesssim \left(\sum_{k=1}^{n-1} \int_{r_k}^{r_{k+1}} \frac{1}{(\delta_k \wedge \delta_{k+1})^2} + \int_{r_{\text{bl}}}^1 \frac{1}{\delta_{\text{bl}}^2} \right) \cdot \int_D |f|^2 + |\nabla f|^2 \\ &\lesssim \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right) \cdot \int_D |f|^2 + |\nabla f|^2. \end{aligned}$$

The bounds

$$\frac{1}{\delta_k \wedge \delta_{k+1}} \lesssim \frac{1}{\delta_{k+1}} \lesssim \frac{1}{l_{k+1}} \lesssim \frac{1}{\ell(r)} \quad \forall r \in (r_k, r_{k+1})$$

were used to pass between the second and third lines in the estimate above. They hold by our assumptions that $\delta_k \sim \delta_{k+1}$, $l_k \lesssim \delta_k$ and $l_k \sim l_{k+1}$ along with the definition of ℓ .

Finally, we estimate I_3 which involves the second derivatives $\{\chi''_k\}$. Again we use (4.15) to write that

$$\begin{aligned} I_3 &= \sum_{k=1}^n \int_{r_{\text{bulk}}}^1 |\chi''_k|^2 r^2 l_k^2 |A_1|_{L_\theta^2}^2 r dr \\ &\lesssim \left(\sum_{k=1}^{n-1} \int_{r_k}^{r_{k+1}} \frac{(l_k \vee l_{k+1})^2}{(\delta_k \wedge \delta_{k+1})^4} + \int_{r_{\text{bl}}}^1 \frac{l_{\text{bl}}^2}{\delta_{\text{bl}}^4} \right) \cdot \int_D |f|^2 \\ &\lesssim \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right) \cdot \int_D |f|^2. \end{aligned}$$

Adding up the estimates on I_1 , I_2 and I_3 and averaging in time gives the first part of the claim.

A similar, and much simpler, argument proves the desired bound on $\nabla \eta$. In particular, by the definition of η in §4a and the estimates from (3.14), there holds

$$|\nabla \eta| \leq \sum_{k=1}^n |\chi_k| |\nabla \eta_k| + |\chi'_k| |\eta_k| \lesssim \sum_{k=1}^n |\chi_k| \frac{1}{r l_k} + |\chi'_k|.$$

Hence,

$$\int_D |\nabla \eta|^2 \lesssim \sum_{k=1}^n \int_0^1 \left(|\chi_k|^2 \frac{1}{r^2 l_k^2} + |\chi'_k|^2 \right) r dr.$$

We recognize terms like those from the bound on $\nabla \mathbf{u}$, with the difference being that the terms involving f are now replaced by the number one. The same manipulations apply as before. ■

Combining lemma 4.4 with what we proved in §4bi yields the rest of proposition 4.1.

(c) Optimal branching flows

Finally, we optimize over our branching flows to prove the upper bound in theorem 1.1. Recall from proposition 4.1 that the (rescaled) velocities $\lambda_{Pe}\mathbf{u}$ and temperature fields T_{Pe} achieve

$$\left\langle |\nabla T_{Pe}|^2 \right\rangle \lesssim C_0(f) \cdot M(\{l_k\}_{k=1}^n, \{r_k\}_{k=1}^n, n; Pe) \quad \text{and} \quad \left\langle |\nabla(\lambda_{Pe}\mathbf{u})|^2 \right\rangle = Pe^2, \quad (4.18)$$

where

$$C_0 = \left\langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \right\rangle$$

and

$$M = l_{\text{bulk}}^2 + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell'(r))^2 dr + \delta_{\text{bl}} + \frac{1}{Pe^2} \left[\frac{1}{l_{\text{bulk}}^2} + \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{(\ell(r))^2} dr + \frac{\delta_{\text{bl}}}{l_{\text{bl}}^2} \right]^2.$$

We find it useful to work directly with the continuous ‘scale function’ $\ell(r)$ introduced in (4.9), and to wait to enforce the interpolation rule

$$l_k = \ell(r_k) \quad \text{for } k = 1, \dots, n \quad (4.19)$$

until it comes time to select the parameters $\{l_k\}_{k=1}^n$ and $\{r_k\}_{k=1}^n$.

Recall $l_1 = l_{\text{bulk}}$, $l_n = l_{\text{bl}}$, $r_1 = r_{\text{bulk}}$ and $r_n = r_{\text{bl}}$, and consider the one-dimensional variational problem

$$\min_{\substack{\ell(r) \\ \ell(r_{\text{bulk}}) = l_{\text{bulk}} \\ \ell(r_{\text{bl}}) = l_{\text{bl}}}} \int_{r_{\text{bulk}}}^{r_{\text{bl}}} (\ell')^2 dr + \frac{1}{Pe^2} \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{\ell^2} dr \right)^2$$

suggested by minimizing M . Its solution obeys

$$(\ell')^2 \sim \frac{1}{Pe^2} \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{\ell^2} dr \right) \frac{1}{\ell^2}.$$

Setting $\ell(1) = 0$ and integrating yields

$$\ell(r) \sim \frac{1}{Pe^{1/2}} \left(\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{\ell^2} dr \right)^{1/4} \sqrt{1-r}. \quad (4.20)$$

Put another way,

$$\ell(r) = c(Pe) \sqrt{1-r}.$$

The constant $c(Pe)$ and the parameters r_{bulk} , r_{bl} , l_{bulk} and l_{bl} must be determined. For c , note that

$$\int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{1}{\ell^2} dr = \frac{1}{c^2} \int_{r_{\text{bulk}}}^{r_{\text{bl}}} \frac{dr}{1-r} = \frac{1}{c^2} \log \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right)$$

where $\delta_{\text{bulk}} = 1 - r_{\text{bulk}}$ and $\delta_{\text{bl}} = 1 - r_{\text{bl}}$. Setting this into (4.20) and rearranging, we get that

$$c(Pe) \sim \frac{1}{Pe^{1/3}} \log^{1/6} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right).$$

By (4.19),

$$l_{\text{bulk}} = \ell(r_{\text{bulk}}) \sim \frac{\delta_{\text{bulk}}^{1/2}}{Pe^{1/3}} \log^{1/6} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right) \quad \text{and} \quad l_{\text{bl}} = \ell(r_{\text{bl}}) \sim \frac{\delta_{\text{bl}}^{1/2}}{Pe^{1/3}} \log^{1/6} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right).$$

All that remains is to choose δ_{bulk} and δ_{bl} . Note the quantity M from (4.18) satisfies

$$\begin{aligned} M &\lesssim l_{\text{bulk}}^2 + \frac{1}{Pe^2} \frac{1}{l_{\text{bulk}}^4} + \delta_{\text{bl}} + \frac{1}{Pe^2} \frac{\delta_{\text{bl}}^2}{l_{\text{bl}}^4} + \frac{1}{Pe^{2/3}} \log^{4/3} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right) \\ &\lesssim \delta_{\text{bl}} + \frac{1}{Pe^{2/3}} \left[\frac{1}{\delta_{\text{bulk}}^2 \log^{2/3} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right)} + \log^{4/3} \left(\frac{\delta_{\text{bulk}}}{\delta_{\text{bl}}} \right) \right]. \end{aligned}$$

Minimizing with

$$\delta_{\text{bulk}} \sim 1 \quad \text{and} \quad \delta_{\text{bl}} \sim \frac{\log^{1/3} Pe}{Pe^{2/3}}$$

yields the desired bound

$$\langle |\nabla T_{Pe}|^2 \rangle \leq C'(f) \cdot \frac{\log^{4/3} Pe}{Pe^{2/3}}, \quad (4.21)$$

where $C' \lesssim \langle |f|^2 + |\nabla f|^2 + |\nabla \nabla f|^2 \rangle$. Regarding its scaling in Pe , this is the best upper bound achievable by a roll-based branching flow.

To complete the proof of theorem 1.1, we only need verify that our choices for $\{l_k\}_{k=1}^n$, $\{r_k\}_{k=1}^n$ and n are actually admissible in our analysis of branching flows. We do this now.

Proof of the upper bound from theorem 1.1. To be absolutely clear, we fix

$$\ell(r) = \frac{\log^{1/6} Pe}{Pe^{1/3}} \sqrt{1-r}, \quad r \in \left(\frac{1}{2}, 1\right) \quad (4.22)$$

for the remainder of the proof, and let $n \in \mathbb{N}$ satisfy

$$n \leq \log_2 \frac{Pe^{1/3}}{\log^{1/6} Pe} \leq n+1.$$

Implicit in this is the requirement that $2 \leq Pe^{1/3} / \log^{1/6} Pe$. Define the scales $\{l_k\}_{k=1}^n$ by taking

$$l_k = \frac{l_{\text{bulk}}}{2^{k-1}} \quad \text{for } k = 1, \dots, n$$

where $l_{\text{bulk}}^{-1} \in \mathbb{N}$ obeys

$$2 \frac{Pe^{1/3}}{\log^{1/6} Pe} \leq \frac{1}{l_{\text{bulk}}} \leq 4 \frac{Pe^{1/3}}{\log^{1/6} Pe}.$$

For the points $\{r_k\}_{k=1}^n$, enforce the interpolation rule

$$l_k = \ell(r_k),$$

which says here that

$$\frac{l_{\text{bulk}}}{2^{k-1}} = \frac{\log^{1/6} Pe}{Pe^{1/3}} \sqrt{1-r_k} \quad \text{for } k = 1, \dots, n. \quad (4.23)$$

At this point, all available choices have been made and we can go ahead with our proof of the desired bound (4.21). Actually, we already did most of the heavy lifting in the paragraphs above, where we explained how these choices follow from optimizing the result of proposition 4.1. All that remains is to verify the hypotheses from §4a and proposition 4.1.

First, we check the assumptions of §4a. Clearly, l_k decreases with increasing k per (4.2). And as $r \mapsto \ell(r)$ is strictly decreasing, the points r_k increase with increasing k . Taking $k=1$ in (4.23), we get that

$$1 = \frac{1}{l_{\text{bulk}}} \frac{\log^{1/6} Pe}{Pe^{1/3}} \sqrt{1-r_1} \geq 2\sqrt{1-r_1}$$

by our choice of l_{bulk} . So, $r_1 \geq 3/4 > 1/2$ as per (4.3). We have shown that our choices for $\{l_k\}$, $\{r_k\}$ and n constitute a viable branching flow.

Next, we check the hypotheses of proposition 4.1. Its first one (4.10) requires that $|l_{k+1} - l_k| \sim l_{k+1} \sim l_k$ and $\delta_{k+1} \sim \delta_k$ with fixed numerical constants. The former holds by the dyadic nature of l_k . For the latter, introduce the inverse $\ell \mapsto r(\ell)$, which is strictly decreasing, and note that

$\delta_k = r_{k+1} - r_k$ obeys

$$\delta_k = \int_{l_{k+1}}^{l_k} |r'(\ell)| \, d\ell.$$

It suffices to check that $|r'(\ell_k)| \sim |r'(\ell_{k+1})|$. Differentiating (4.22) implicitly and rearranging gives

$$r'(\ell) = -2 \frac{Pe^{1/3}}{\log^{1/6} Pe} \sqrt{1-r}. \quad (4.24)$$

One sees from (4.23) that $1 - r_k \sim 1 - r_{k+1}$. Hence, (4.10) is proved.

Finally, we verify the second hypothesis (4.11) of proposition 4.1, which is that $l_k \lesssim \delta_k$ for each k . Since $\delta_k = r_{k+1} - r_k$ and $l_k \sim |l_{k+1} - l_k|$, we must show that

$$1 \lesssim \left| \frac{r_{k+1} - r_k}{l_{k+1} - l_k} \right|.$$

Referring again to $r(\ell)$, it suffices to check that $1 \lesssim |r'(\ell)|$ for $\ell \in (l_1, l_n)$. Evidently by (4.24),

$$|r'(\ell)| \gtrsim \frac{Pe^{1/3}}{\log^{1/6} Pe} \sqrt{1-r} \geq \frac{Pe^{1/3}}{\log^{1/6} Pe} \sqrt{1-r_n} \sim 1.$$

In the last step we used (4.23) with $k = n$. That $l_{bl} \sim \delta_{bl}$ is clear. Theorem 1.1 is proved. ■

5. Unsteady roll-like flows for energy-constrained cooling

We close by proving the upper bound on energy-constrained cooling from the introduction, by estimating the mean-square temperature gradient of a well-chosen family of convection roll-like flows (see the middle row of figure 1). It is an open challenge to decide whether this bound is sharp in its scaling with respect to Pe , in the advective limit $Pe \rightarrow \infty$. In particular, we note the significant gap between the lower bound in proposition 2.6 and the upper bound achieved below.

Proposition 5.1. *Let $f(\mathbf{x}, t)$ satisfy*

$$\lim_{\tau \rightarrow \infty} \frac{1}{\sqrt{\tau}} \int_0^\tau e^{-\lambda_1((\tau-t) \wedge t)} \|f(\cdot, t)\|_{L^2(D)} \, dt = 0 \quad \text{and} \quad \langle |f|^2 + |\nabla f|^2 \rangle < \infty.$$

There exists a fixed, numerical constant $C' > 0$ such that

$$\min_{\substack{\mathbf{u}(\mathbf{x}, t) \\ \langle |\mathbf{u}|^2 \rangle = Pe^2 \\ \mathbf{u} = \mathbf{0} \text{ at } \partial D}} \langle |\nabla T|^2 \rangle \leq \frac{C'}{Pe} \cdot \langle |f|^2 + |\nabla f|^2 \rangle$$

whenever $Pe \geq 1$. The same bound holds using no-penetration conditions $\mathbf{u} \cdot \hat{\mathbf{n}} = 0$ in place of the no-slip ones $\mathbf{u} = \mathbf{0}$ at ∂D .

Proof. Pure convection roll-like flows occur as the simplest case of our branching flows from §4a, with $n = 1$ and using only $l_1 = l_{\text{bulk}}$ and $\delta_1 = \delta_{bl}$. The rescaled velocities

$$\lambda_{Pe} \mathbf{u} \quad \text{with} \quad \lambda_{Pe} = \frac{Pe}{\sqrt{\langle |\mathbf{u}|^2 \rangle}}$$

generate temperature fields T_{Pe} satisfying

$$\langle |\nabla T_{Pe}|^2 \rangle \lesssim C_0(f) \cdot \left[l_{\text{bulk}}^2 + \delta_{bl} + \frac{1}{Pe^2} \left(\frac{1}{l_{\text{bulk}}^2} + \frac{\delta_{bl}}{l_{\text{bulk}}^2} \right) \right]$$

so long as $l_{\text{bulk}} \lesssim \delta_{bl}$, where now

$$C_0 = \langle |f|^2 + |\nabla f|^2 \rangle.$$

The proof of this is contained in that of proposition 4.1, once one notes that the un-scaled velocities from §4a obey

$$\langle |\mathbf{u}|^2 \rangle \lesssim 1.$$

To see this write

$$\mathbf{u} = \chi_1 \mathbf{u}_1 + \chi_1' \psi_1 \hat{\mathbf{e}}_\theta$$

in the case $n = 1$, and use the definitions to get that

$$|\mathbf{u}| \lesssim 1 + \frac{l_{\text{bulk}}}{\delta_{\text{bl}}} \lesssim 1.$$

(The proof for $n > 1$ is the same, but we leave it to the reader as it is not needed here.) The dependence of C_0 on f comes from lemmas 4.2 and 4.3 and not, in this case, from lemma 4.4. Optimizing gives

$$\delta_{\text{bl}} \sim l_{\text{bulk}} \sim \frac{1}{Pe^{1/2}},$$

and this proves the result. ■

Data accessibility. This article has no additional data.

Authors' contributions. I.T.: conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, resources, validation, visualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing interests. I declare I have no competing interests.

Funding. This work was supported by National Science Foundation Award DMS-2025000.

Acknowledgements. The author thanks Charlie Doering for many inspiring discussions leading to this study, and David Goluskin for noting the relevance of balanced versus unbalanced heating.

References

1. Alam T, Kim M-H. 2018 A comprehensive review on single phase heat transfer enhancement techniques in heat exchanger applications. *Renew. Sustain. Energy Rev.* **81**, 813–839. (doi:10.1016/j.rser.2017.08.060)
2. Alben S. 2017 Improved convection cooling in steady channel flows. *Phys. Rev. Fluids* **2**, 104501. (doi:10.1103/PhysRevFluids.2.104501)
3. Alben S. 2017 Optimal convection cooling flows in general 2D geometries. *J. Fluid Mech.* **814**, 484–509. (doi:10.1017/jfm.2017.35)
4. Catton I. 2011 Conjugate heat transfer within a heterogeneous hierarchical structure. *J. Heat Transfer* **133**, 103001. (doi:10.1115/1.4003576)
5. Feppon F, Allaire G, Bordeu F, Cortial J, Dapogny C. 2019 Shape optimization of a coupled thermal fluid–structure problem in a level set mesh evolution framework. *SeMA J.* **76**, 413–458. (doi:10.1007/s40324-018-00185-4)
6. Feppon F, Allaire G, Dapogny C, Jolivet P. 2020 Topology optimization of thermal fluid–structure systems using body-fitted meshes and parallel computing. *J. Comput. Phys.* **417**, 109574. (doi:10.1016/j.jcp.2020.109574)
7. Feppon F, Allaire G, Dapogny C, Jolivet P. 2021 Body-fitted topology optimization of 2D and 3D fluid-to-fluid heat exchangers. *Comput. Methods Appl. Mech. Eng.* **376**, 113638. (doi:10.1016/j.cma.2020.113638)
8. Geb D, Ge M, Chu J, Catton I. 2013 Measuring transport coefficients in heterogeneous and hierarchical heat transfer devices. *J. Heat Transf.* **135**, 061101. (doi:10.1115/1.4023547)
9. Rastan H, Abdi A, Hamawandi B, Ignatowicz M, Meyer JP, Palm B. 2020 Heat transfer study of enhanced additively manufactured minichannel heat exchangers. *Int. J. Heat Mass Transf.* **161**, 120271. (doi:10.1016/j.ijheatmasstransfer.2020.120271)
10. Weisberg A, Bau HH, Zemel JN. 1992 Analysis of microchannels for integrated cooling. *Int. J. Heat Mass Transf.* **35**, 2465–2474. (doi:10.1016/0017-9310(92)90089-B)
11. Arslan A, Fantuzzi G, Craske J, Wynn A. 2021 Bounds for internally heated convection with fixed boundary heat flux. *J. Fluid Mech.* **922**, R1. (doi:10.1017/jfm.2021.527)
12. Arslan A, Fantuzzi G, Craske J, Wynn A. 2021 Bounds on heat transport for convection driven by internal heating. *J. Fluid Mech.* **919**, A15. (doi:10.1017/jfm.2021.360)
13. Bouillaut V, Lepot S, Aumaître S, Gallet B. 2019 Transition to the ultimate regime in a radiatively driven convection experiment. *J. Fluid Mech.* **861**, R5. (doi:10.1017/jfm.2018.972)

14. Goluskin D. 2015 Internally heated convection beneath a poor conductor. *J. Fluid Mech.* **771**, 36–56. (doi:10.1017/jfm.2015.140)
15. Goluskin D. 2016 *Internally heated convection and Rayleigh-Bénard convection*. Cham, Switzerland: Springer.
16. Lepot S, Aumaitre S, Gallet B. 2018 Radiative heating achieves the ultimate regime of thermal convection. *Proc. Natl Acad. Sci. USA* **115**, 8937–8941. (doi:10.1073/pnas.1806823115)
17. Drivas TD, Nguyen HQ, Nobili C. 2021 Bounds on heat flux for Rayleigh-Bénard convection between Navier-slip fixed-temperature boundaries. Preprint (<https://arxiv.org/abs/2109.13205>)
18. Fantuzzi G. 2018 Bounds for Rayleigh-Bénard convection between free-slip boundaries with an imposed heat flux. *J. Fluid Mech.* **837**, R5. (doi:10.1017/jfm.2017.907)
19. Goluskin D, Doering CR. 2016 Bounds for convection between rough boundaries. *J. Fluid Mech.* **804**, 370–386. (doi:10.1017/jfm.2016.528)
20. Toppaladoddi S, Wells AJ, Doering CR, Wettlaufer JS. 2021 Thermal convection over fractal surfaces. *J. Fluid Mech.* **907**, A12. (doi:10.1017/jfm.2020.826)
21. Whitehead JP, Doering CR. 2011 Ultimate state of two-dimensional Rayleigh-Bénard convection between free-slip fixed-temperature boundaries. *Phys. Rev. Lett.* **106**, 244501. (doi:10.1103/PhysRevLett.106.244501)
22. Busse FH. 1969 On Howard's upper bound for heat transport by turbulent convection. *J. Fluid. Mech.* **37**, 457–477. (doi:10.1017/S0022112069000668)
23. Howard LN. 1963 Heat transport by turbulent convection. *J. Fluid. Mech.* **17**, 405–432. (doi:10.1017/S0022112063001427)
24. Doering CR, Constantin P. 1996 Variational bounds on energy dissipation in incompressible flows. III. Convection. *Phys. Rev. E* **53**, 5957–5981. (doi:10.1103/PhysRevE.53.5957)
25. Doering CR. 2020 Turning up the heat in turbulent thermal convection. *Proc. Natl Acad. Sci. USA* **117**, 9671–9673. (doi:10.1073/pnas.2004239117)
26. Constantin P, Doering CR. 1995 Variational bounds on energy dissipation in incompressible flows. II. Channel flow. *Phys. Rev. E* **51**, 3192–3198. (doi:10.1103/PhysRevE.51.3192)
27. Doering CR, Constantin P. 1994 Variational bounds on energy dissipation in incompressible flows: shear flow. *Phys. Rev. E* **49**, 4087–4099. (doi:10.1103/PhysRevE.49.4087)
28. Kerswell RR. 1998 Unification of variational principles for turbulent shear flows: the background method of Doering-Constantin and the mean-fluctuation formulation of Howard-Busse. *Physica D* **121**, 175–192. (doi:10.1016/S0167-2789(98)00104-3)
29. Plasting SC, Kerswell RR. 2003 Improved upper bound on the energy dissipation rate in plane Couette flow: the full solution to Busse's problem and the Constantin-Doering-Hopf problem with one-dimensional background field. *J. Fluid Mech.* **477**, 363–379. (doi:10.1017/S0022112002003361)
30. Fantuzzi G, Arslan A, Wynn A. 2022 The background method: theory and computations. *Phil. Trans. R. Soc. A* **380**, 20210038. (doi:10.1098/rsta.2021.0038)
31. Chernyshenko SI, Goulart P, Huang D, Papachristodoulou A. 2014 Polynomial sum of squares in fluid dynamics: a review with a look ahead. *Phil. Trans. R. Soc. A* **372**, 20130350. (doi:10.1098/rsta.2013.0350)
32. Olson ML, Goluskin D, Schultz WW, Doering CR. 2021 Heat transport bounds for a truncated model of Rayleigh-Bénard convection via polynomial optimization. *Physica D* **415**, 132748. (doi:10.1016/j.physd.2020.132748)
33. Rosa RMS, Temam RM. 2021 Optimal minimax bounds for time and ensemble averages for the incompressible Navier-Stokes equations. Preprint (<https://arxiv.org/abs/2010.06730>)
34. Tobasco I, Goluskin D, Doering CR. 2018 Optimal bounds and extremal trajectories for time averages in nonlinear dynamical systems. *Phys. Lett. A* **382**, 382–386. (doi:10.1016/j.physleta.2017.12.023)
35. Goluskin D, Fantuzzi G. 2019 Bounds on mean energy in the Kuramoto-Sivashinsky equation computed using semidefinite programming. *Nonlinearity* **32**, 1705–1730. (doi:10.1088/1361-6544/ab018b)
36. Nobili C, Otto F. 2017 Limitations of the background field method applied to Rayleigh-Bénard convection. *J. Math. Phys.* **58**, 093102. (doi:10.1063/1.5002559)
37. Otto F, Seis C. 2011 Rayleigh-Bénard convection: improved bounds on the Nusselt number. *J. Math. Phys.* **52**, 083702. (doi:10.1063/1.3623417)

38. Miquel B, Lepot S, Bouillaut V, Gallet B. 2019 Convection driven by internal heat sources and sinks: heat transport beyond the mixing-length or ‘ultimate’ scaling regime. *Phys. Rev. Fluids* **4**, 121501. (doi:10.1103/PhysRevFluids.4.121501)
39. Marcotte F, Doering CR, Thiffeault J-L, Young WR. 2018 Optimal heat transfer and optimal exit times. *SIAM J. Appl. Math.* **78**, 591–608. (doi:10.1137/17M1150220)
40. Iyer G, Van T-S. 2022 Bounds on the heat transfer rate via passive advection. *SIAM J. Math. Anal.* **54**, 1927–1965. (doi:10.1137/21M1394497)
41. Goluskin D. 2018 Bounding averages rigorously using semidefinite programming: mean moments of the Lorenz system. *J. Nonlinear Sci.* **28**, 621–651. (doi:10.1007/s00332-017-9421-2)
42. Lakshmi MV, Fantuzzi G, Fernández-Caballero JD, Hwang Y, Chernyshenko SI. 2020 Finding extremal periodic orbits with polynomial optimization, with application to a nine-mode model of shear flow. *SIAM J. Appl. Dyn. Syst.* **19**, 763–787. (doi:10.1137/19M1267647)
43. Doering CR, Tobasco I. 2019 On the optimal design of wall-to-wall heat transport. *Commun. Pur. Appl. Math.* **72**, 2385–2448. (doi:10.1002/cpa.21832)
44. Souza AN, Tobasco I, Doering CR. 2020 Wall-to-wall optimal transport in two dimensions. *J. Fluid Mech.* **889**, A34. (doi:10.1017/jfm.2020.42)
45. Tobasco I, Doering CR. 2017 Optimal wall-to-wall transport by incompressible flows. *Phys. Rev. Lett.* **118**, 264502. (doi:10.1103/PhysRevLett.118.264502)
46. Avellaneda M, Majda AJ. 1991 An integral representation and bounds on the effective diffusivity in passive advection by laminar and turbulent flows. *Commun. Math. Phys.* **138**, 339–391. (doi:10.1007/BF02099496)
47. Fannjiang A, Papanicolaou G. 1994 Convection enhanced diffusion for periodic flows. *SIAM J. Appl. Math.* **54**, 333–408. (doi:10.1137/S0036139992236785)
48. Kohn RV, Strang G. 1986 Optimal design and relaxation of variational problems, I. *Commun. Pur. Appl. Math.* **39**, 113–137. (doi:10.1002/cpa.3160390107)
49. Kohn RV, Strang G. 1986 Optimal design and relaxation of variational problems, II. *Commun. Pur. Appl. Math.* **39**, 139–182. (doi:10.1002/cpa.3160390202)
50. Kohn RV, Strang G. 1986 Optimal design and relaxation of variational problems, III. *Commun. Pur. Appl. Math.* **39**, 353–377. (doi:10.1002/cpa.3160390305)
51. Motoki S, Kawahara G, Shimizu M. 2018 Maximal heat transfer between two parallel plates. *J. Fluid Mech.* **851**, R4. (doi:10.1017/jfm.2018.557)
52. Evans C. 2010 *Partial differential equations*, vol. **19**, 2nd edn. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society.
53. Coifman R, Lions P-L, Meyer Y, Semmes S. 1993 Compensated compactness and Hardy spaces. *J. Math. Pures Appl.* **72**, 247–286.
54. Müller S. 1994 Hardy space methods for nonlinear partial differential equations. *Equadiff* **8**, 159–168.