

Measurement of the parity-odd angular distribution of γ rays from polarized neutron capture on ^{35}Cl

N. Fomin^{1,2}, R. Alarcon,³ L. Alonzi,⁴ E. Askanazi,⁴ S. Baeßler,^{4,5} S. Balascuta,^{6,3} L. Barrón-Palos,⁷ A. Barzilov,⁸ D. Blyth,^{3,9} J. D. Bowman,⁵ N. Birge,¹ J. R. Calarco,¹⁰ T. E. Chupp,¹¹ V. Cianciolo,⁵ C. E. Coppola,¹ C. B. Crawford,¹² K. Craycraft,^{1,12} D. Evans,^{4,13} C. Fieseler,¹² E. Frlež,⁴ J. Fry,^{4,13,14} I. Garishvili,^{5,1} M. T. W. Gericke,¹⁵ R. C. Gillis,^{5,13} K. B. Grammer,^{5,1} G. L. Greene,^{1,5} J. Hall,⁴ J. Hamblen,¹⁶ C. Hayes,^{17,1} E. B. Iverson,⁵ M. L. Kabir,^{18,12} S. Kucuker,^{19,1} B. Lauss,²⁰ R. Mahurin,²¹ M. McCrea,^{12,15} M. Maldonado-Velázquez,⁷ Y. Masuda,²² J. Mei,¹³ R. Milburn,¹² P. E. Mueller,⁵ M. Musgrave,^{23,1} H. Nann,^{13,*} I. Novikov,²⁴ D. Parsons,¹⁶ S. I. Penttilä,⁵ D. Počanić,⁴ A. Ramirez-Morales,⁷ M. Root,⁴ A. Salas-Bacci,⁴ S. Santra,²⁵ S. Schröder,^{4,26} E. Scott,¹ P.-N. Seo,^{4,27} E. I. Sharapov,²⁸ F. Simmons,¹² W. M. Snow,¹³ A. Sprow,¹² J. Stewart,¹⁶ E. Tang,^{12,2} Z. Tang,^{13,2} X. Tong,⁵ D. J. Turkoglu,²⁹ R. Whitehead,¹ and W. S. Wilburn²

(NPDGamma Collaboration)

¹University of Tennessee, Knoxville, Tennessee 37996, USA

²Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

³Arizona State University, Tempe, Arizona 85287, USA

⁴University of Virginia, Charlottesville, Virginia 22904, USA

⁵Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

⁶Horia Hulubei National Institute for Physics and Nuclear Engineering, Magurele 077125, Romania

⁷Instituto de Física, Universidad Nacional Autónoma de México, Apartado Postal 20-364, 01000, México

⁸University of Nevada, Las Vegas, Nevada 89154, USA

⁹High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois, 60439, USA

¹⁰University of New Hampshire, Durham, New Hampshire 03824, USA

¹¹University of Michigan, Ann Arbor, Michigan 48109, USA

¹²University of Kentucky, Lexington, Kentucky 40506, USA

¹³Indiana University, Bloomington, Indiana 47405, USA

¹⁴Eastern Kentucky University, Richmond, Kentucky 40475, USA

¹⁵University of Manitoba, Winnipeg, MB, Canada R3T 2N2

¹⁶University of Tennessee, Chattanooga, Tennessee 37403, USA

¹⁷Physics Department, North Carolina State University, Raleigh, North Carolina 27695, USA

¹⁸Mississippi State University, Mississippi State, Mississippi 39759, USA

¹⁹Northwestern University Feinberg School of Medicine, Chicago, Illinois 60611, USA

²⁰Paul Scherrer Institut, CH-5232 Villigen, Switzerland

²¹Middle Tennessee State University, Murfreesboro, Tennessee 37132, USA

²²High Energy Accelerator Research Organization (KEK), Tsukuba-shi, 305-0801, Japan

²³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

²⁴Western Kentucky University, Bowling Green, Kentucky 42101, USA

²⁵Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India

²⁶Saarland University, Institute of Experimental Ophthalmology, Kirrberger Str. 100, Bldg. 22, 66424 Homburg/Saar, Germany

²⁷Triangle Universities Nuclear Lab, Durham, North Carolina 27708, USA

²⁸Joint Institute for Nuclear Research, Dubna 141980, Russia

²⁹National Institute of Standards and Technology, Gaithersburg, Maryland 20899, USA



(Received 1 October 2021; accepted 17 May 2022; published 27 July 2022)

We report a measurement of two energy-weighted γ cascade angular distributions from polarized slow neutron capture on the ^{35}Cl nucleus, one parity-odd correlation proportional to $\vec{s}_n \cdot \vec{k}_\gamma$ and one parity-even correlation proportional to $\vec{s}_n \cdot \vec{k}_n \times \vec{k}_\gamma$. A parity-violating asymmetry can appear in this reaction due to the weak nucleon-nucleon interaction, which mixes opposite parity S- and P-wave levels in the excited compound ^{36}Cl nucleus formed upon slow neutron capture. If parity-violating (PV) and parity-conserving (PC) terms both exist, the

*Deceased

measured differential cross section can be related to them via $\frac{d\sigma}{d\Omega} \propto 1 + A_{\gamma,\text{PV}} \cos \theta + A_{\gamma,\text{PC}} \sin \theta$. The PV and PC asymmetries for energy-weighted γ cascade angular distributions for polarized slow neutron capture on ^{35}Cl averaged over the neutron energies from 2.27–9.53 meV were measured to be $A_{\gamma,\text{PV}} = (-23.9 \pm 0.7) \times 10^{-6}$ and $A_{\gamma,\text{PC}} = (0.1 \pm 0.7) \times 10^{-6}$. These results are consistent with previous experimental results. Systematic errors were quantified and shown to be small compared to the statistical error. These asymmetries in the angular distributions of the γ rays emitted from the capture of polarized neutrons in ^{35}Cl were used to verify the operation and data analysis procedures for the NPDGamma experiment, which measured the parity-odd asymmetry in the angular distribution of γ s from polarized slow neutron capture on protons.

DOI: [10.1103/PhysRevC.106.015504](https://doi.org/10.1103/PhysRevC.106.015504)

I. INTRODUCTION

Parity violation in nuclei in the standard model arises from the weak interaction between nucleons. The parity-odd component of the nucleon-nucleon (NN) weak interaction mixes opposite parity levels. Interference between electromagnetic transitions between these states leads to parity-odd γ asymmetries (from polarized initial states) and circular polarization (from unpolarized initial states). Parity violation at the NN level is poorly understood because the strongly interacting limit of QCD is not solved and the typical size of the parity-odd amplitudes are about 10^{-7} of the dominant strong interaction amplitudes. Several reviews of this subject exist [1–3].

The NPDGamma experiment, which motivated the measurements and results presented in this paper, measured the parity-violating directional asymmetry A_{γ}^{np} in the emission of γ s from polarized neutron capture on liquid parahydrogen, $\frac{d\sigma}{d\Omega} \propto \frac{1}{4\pi}(1 - A_{\gamma}^{np} \cos \theta)$. This reaction isolates the $\Delta I = 1$, $^3S_1 \rightarrow ^3P_1$ component of the weak NN interaction dominated by pion exchange. A_{γ}^{np} can be directly related to the NN weak coupling constant h_{π}^1 in the DDH meson exchange model [4] and to a low-energy constant in pionless effective field theory, $C^{^3S_1 \rightarrow ^3P_1}/C_0$ [5]. The NPDGamma Collaboration reported a result of $A_{\gamma}^{np} = [-3.0 \pm 1.4(\text{stat}) \pm 0.2(\text{sys})] \times 10^{-8}$, which corresponds to a DDH weak πNN coupling of $h_{\pi}^1 = [2.6 \pm 1.2(\text{stat}) \pm 0.2(\text{sys})] \times 10^{-7}$ and a pionless EFT constant of $[-7.4 \pm 3.5(\text{stat}) \pm 0.5(\text{sys})] \times 10^{-11} \text{MeV}^{-1}$ [6].

In the simplest case, that of $\bar{n} + p \rightarrow d + \gamma$, the γ -ray asymmetry expression can be written down in terms of the matrix elements between initial and final states as:

$$A_{\gamma}^{np} \propto \frac{\langle ^3P_1 | \mathbf{E1} | ^3S_1 \rangle}{\langle ^3S_1 | \mathbf{M1} | ^1S_0 \rangle}, \text{ where } \epsilon = \frac{\langle \psi_{\alpha'} | W | \psi_{\alpha} \rangle}{\Delta E} \quad (1)$$

with $\alpha = J, L, S, p$, where p denotes parity. The situation becomes complicated quickly for heavier nuclei as the number of γ -ray transitions grows, making the calculation of the parity-violating asymmetry directly from the strong and weak Hamiltonians virtually impossible.

The parity-odd γ asymmetry then becomes a complicated superposition of asymmetries from both different γ transitions and also from different γ cascade paths on the way to the ground state, each step with its different associated initial state polarization values, which in turn are dependent on the cascade path. Furthermore, the signals from these different γ transition energies are not equally weighted: a γ with twice the energy makes twice the signal size in current mode de-

tection. The γ -ray asymmetry from the decaying compound nucleus as measured by a current-mode γ detector can be written as:

$$A_{\gamma} = \epsilon B_{\gamma}, \quad (2)$$

with

$$B_{\gamma} = \xi \cdot F(J_i, J_f) \times \frac{2Re[\sum_{J_f} \langle J_f^p | \mathbf{E1} | J_i^{p'} \rangle \langle J_i^p | \mathbf{M1} | J_f^p \rangle E_{\gamma,if}^4]}{\sum_{J_f} (|\langle J_f^p | \mathbf{M1} | J_i^{p'} \rangle|^2 + |\langle J_f^p | \mathbf{E1} | J_i^{p'} \rangle|^2) E_{\gamma,if}^4}. \quad (3)$$

B_{γ} describes the γ cascade with transitions between initial and final compound nuclear states with total angular momentum (J_i, J_f) and parity (p, p'), and where $F(J_i, J_f)$ is the angular-momentum coupling factor resulting from the compound state polarization and J_T is the angular momentum of the target nucleus before neutron capture [7]. Finally, ξ is a dilution factor that arises because the current mode γ detector lacks energy resolution and instead sees a superposition of currents from all transitions [7].

In this paper, we present a precise measurement of one parity-odd and one parity-even cascade γ asymmetries in polarized slow neutron capture in ^{35}Cl . The nuclear structure of the ^{35}Cl nucleus is far too complicated to use such an experiment to probe the NN weak interaction amplitudes in a quantitative way. Our motivation to measure parity violation in this nucleus is its usefulness in calibrating the properties of the NPDGamma apparatus. For this purpose it is useful to have a nucleus that possesses a large parity-odd γ asymmetry. ^{35}Cl is already known to possess a very large parity-odd γ asymmetry. Results from previous measurements are summarized in Table I, giving a world average of $A_{\gamma,\text{PV}}^{^{35}\text{Cl}} = (-23.9 \pm 1.36) \times 10^{-6}$. This asymmetry is almost three orders of magnitude larger than in polarized neutron capture in hydrogen. The large parity violation seen in this nucleus is thought to arise from the mixing of the $J^{\pi} = 2^{-}$ p-wave level at

TABLE I. Summary of results for A_{γ} on ^{35}Cl .

Measurement	Result ($\times 10^{-6}$)
Vesna <i>et al.</i> [9]	-27.8 ± 4.9
NPDGamma LANL [10]	-29.1 ± 6.7
ILL [8]	-21.2 ± 1.7




FIG. 1. NPDGamma apparatus, with chlorine target shown inside the detector array (one of the experimental geometries).

+398 eV and a $J^\pi = 2^+$ subthreshold s-wave resonance at -130 eV in the $n + {}^{35}\text{Cl}$ system [8].

In this paper, we describe the early chronology of the NPDGamma apparatus testing, which involved measurements on the ${}^{35}\text{Cl}$ target to test the system as well as validate the calculations of the geometrical factors [11]. The ${}^{35}\text{Cl}$ measurements uncovered a small number of issues and led to some modifications of the experimental setup. While the first (problematic) measurement is described for the purposes of motivating and explaining the experimental improvements, it is not used in the extraction of the PV asymmetry quoted in this work.

II. EXPERIMENTAL SETUP

A. Beam line

The NPDGamma experiment (Fig. 1) was installed on the Fundamental Neutron Physics Beamline (FNBP) of the Spallation Neutron Source (SNS) at Oak Ridge National Laboratory (ORNL). The SNS is a pulsed source operating at 60 Hz, with a liquid Hg target and a supercritical hydrogen moderator. Detailed beam line information is available in Ref. [12]. NPDGamma is installed on the cold, polychromatic beam line 13B, with the center of the detector array located ≈ 17.6 m downstream of the moderator. Two bandwidth choppers were used to select neutrons with wavelength $1.93 < \lambda < 5.6$ Å for the ${}^{35}\text{Cl}$ data for configuration 1 (CONF1) and $2.93 < \lambda < 6.0$ Å for configuration 2 (CONF2).

B. Polarizer

The neutron beam was polarized with a supermirror polarizer (SMP) [13], manufactured by Swiss Neutronics. A compensation magnet was designed and built in order to cancel the fringe field of the SMP and minimize field gradients. A detailed description is available in Ref. [14].

C. RFSR

In order to minimize effects of detector gain drifts and any other time-dependent changes in the experiment, data were first taken and analyzed in units of an eight-step spin sequence ($\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow$), where each step corresponds to a single accelerator pulse. In later data taking, the spin sequence was alternated with its inverse, as is discussed later in the




FIG. 2. (a) Drawing of the first liquid carbon tetrachloride target used in the experiment (CONF1), inside an aluminum container and (b) the improved target made of teflon (CONF2).

paper. In order to reverse the spin of the neutron beam on a pulse-by-pulse basis, we employ an RF spin rotator (RFSR). The advantage of a spin rotator over an adiabatic spin flipper is that it does not change the kinetic energy of the neutrons and leaves the phase space intact. The resonant rotator reverses the spin of the polarized neutron beam by performing NMR as the beam moves through a region with an orthogonal combination of static and RF magnetic fields without a dc field gradient. The neutron spin precesses in the static holding field of $B_0\hat{y}$, and upon entering the RFSR, it will rotate about the effective magnetic field given by:

$$\vec{B}_{\text{eff}} = \left(B_0 - \frac{\Omega}{\mu_n} \hbar \right) \hat{y} + B_1 \hat{z}, \quad (4)$$

where Ω is the resonant frequency. The condition for resonance is met when Ω matches the Larmor frequency ($\omega_0 = \mu_n B_0 / \hbar$). The magnitude of B_1 is inversely proportional to time of flight, allowing us to reverse neutron spin for a range of neutron velocities. More details on the RFSR are available in Ref. [15].

D. Beam monitors

Beam power and stability are measured with a beam monitor that's 15.15 meters downstream of the hydrogen moderator and intercepts the whole area of the beam. This is a multiwire proportional counter (MWPC) with a ${}^3\text{He}$ (filled to 15.1 Torr) and nitrogen (filled to 750 Torr) gas mixture. Details on the construction and performance of the MWPC are available in Ref. [16].

E. Target

The chlorine asymmetry was measured several times, with different chlorine targets. The CONF1 data set was obtained with a target of liquid carbon tetrachloride in a cylindrical aluminum container, as shown in Fig. 2. The inner and outer radii of the aluminum container are 5.71 cm and 6.15 cm, respectively, with a depth of 5.59 mm. The upstream face of the target container is thinner than the downstream face, 0.76 mm compared to 2.67 mm, in order to minimize background from

neutron capture on aluminum. In this configuration, the target vessel was located 4.9 cm downstream of the center of the detector array in the $\hat{z}$ direction. The analysis of the first chlorine results revealed some shortcomings in the experiment design, which will be addressed in the analysis discussion. The CONF2 set of chlorine measurements also used a liquid carbon tetrachloride target, but this time enclosed in a teflon container. Teflon is transparent to slow neutrons and C and F both possess small (n, γ) capture cross sections, minimizing any background in the detected signals. As with the aluminum-cased target, a thin cylinder was used with an outer radius of 8.43 cm and an inner radius of 6.35 cm, where the outer radius refers to the case, and the inner radius is that of the target volume. The front and rear window thicknesses were 0.30 cm and the target volume thickness was 0.56 cm. The target was placed inside the RFSR enclosure, just downstream of the coils.

F. Detector array

The γ rays are detected in an array of 48 CsI current-mode detectors, arranged around the target with an acceptance of $\approx 3\pi$. Each detector consists of two CsI crystals ($15.2 \times 15.2 \times 15.2$ cm³) viewed by a single vacuum photodiode (VPD), whose voltage is read out and converted to current via low-noise solid-state electronics. The detector components and characteristics are described in detail in Ref. [17].

G. Guide field

The apparatus (starting downstream of the polarizer and ending downstream of detector array) is surrounded by four guide coils providing a 9.7 G field to preserve the polarization of the neutrons after the exit of the polarizer. Details are available in Ref. [14].

III. DATA ANALYSIS

A. Overview

Data were taken in two configurations: first with the aluminum-cased target (CONF1), and later with the teflon-cased target (CONF2), after multiple improvements to the experiment. The reasons for the improvements will be discussed below. While raw asymmetries from the first data set will be shown, they will only serve to illustrate the need for experimental modifications. The final physics asymmetries were extracted using data from CONF2.

Detector and monitor data are recorded in units of the previously described eight-step spin sequence, consisting of eight (16.67 ms) neutron pulses. The ninth pulse is used to read out the data. Half of the pulses have neutrons in the spin-up state, and half are spin-down. An asymmetry is calculated for each spin sequence, integrating the voltage over the time-of-flight bins. The accelerator skips proton delivery to the mercury target every 10 s, for diagnostics. Spin sequences with these so-called dropped pulses are eliminated from the analysis. Additionally, the location of the choppers and the experiment are such that the spectrum is not completely clean: there are leakage neutrons at 13–15 Å and 28–30 Å. Spin

sequences with missing leakage neutrons (from previously dropped pulses) are also eliminated from the analysis. Finally, a beam stability cut of 1% is required for all the pulses in a given spin sequence.

B. Background

In addition to the signal from γ rays emitted in the capture of polarized neutrons on ^{35}Cl , there are several sources of background present. The first is the electronic pedestal, which consists of an offset in the ADC as well as an additional pedestal from the solid-state preamplifiers. The electronic pedestal is present when the beam is off and is on the order of a few mV. The container for the second target is made of teflon, making it transparent to cold neutrons, meaning there was no additional background associated with it. In the case of the first chlorine target, the aluminum holder captures a small fraction of the neutron beam, and the γ s from that process are detected along with the chlorine signal. This aluminum background includes both prompt γ s as well as β -delayed γ s. Cold neutrons capture on ^{27}Al , creating an excited state, $^{28}\text{Al}^*$, which decays via a γ cascade (≈ 8 MeV) down to ^{28}Al . This cascade is the prompt γ background. The half-life of ^{28}Al is ≈ 2.2 minutes and it β decays into an excited state of silicon, $^{28}\text{Si}^*$. The radiation from this β decay is the constant background we refer to as β -delayed γ s.

The liquid target is composed of natural chlorine, whose composition is 75.77% ^{35}Cl and 24.23% ^{37}Cl . Their neutron capture cross sections are 43.6 barns and 0.43 barns, respectively, meaning that the contribution from ^{37}Cl is rather small. The β -delayed background from ^{36}Cl is not an issue, as it is stable (half-life of $\approx 3 \times 10^5$ years). However, we expect some β -delayed background from ^{38}Cl , whose half-life is ≈ 37 minutes [18].

We perform a dynamic pedestal subtraction, which removes the electronic pedestal as well as the β -delayed γ signal, leaving only the prompt γ signal from neutron capture on ^{35}Cl . Two regions in the chlorine spectrum are defined, whose average voltages are V1 and V2, as shown in Figs. 3. While the average signal size is different in the two regions, the background we are trying to subtract is the same, assuming that the neutron beam has been on for a long enough period to build up the β -delayed signal. Detailed discussion of the calculation is available in Ref. [19].

C. Geometrical factors

The geometrical factors are average energy weighted functions that are a measure of the emission direction of a photon from the target that deposits energy in a given detector. The calculation of the so-called geometrical factors is required to correct for the position of the detectors relative to the location of neutron capture in the target. The direction of the beam is defined as $+\hat{z}$, upward neutron polarization is the $+\hat{y}$ direction, and $+\hat{x}$ is the beam left direction in order to make the coordinate system right handed (Fig. 4). Spherical coordinates are defined in the usual way with ϕ measured from the $\hat{x}$ axis and θ measured from the $\hat{z}$ axis.




FIG. 3. A section of a typical cold spectrum corresponding to 2.5 neutron pulses is shown. Regions V1 and V2 are indicated corresponding to average voltages in a low and high part of the spectrum. Dotted line represents the constant background that is the sum of the electronic pedestal and β -delayed signals.

In the case of a point source and point detector, the geometrical factors can be written down analytically. The $\hat{x}$ direction, or left-right (L-R), geometrical factor is proportional to the parity allowed asymmetry. It is given by:

$$\begin{aligned} G_{\text{PC}} &= \langle \hat{k}_\gamma \cdot (\vec{\sigma}_n \times \hat{k}_n) \rangle \\ &= \langle \hat{k}_\gamma \cdot (\hat{y} \times \hat{z}) \rangle \\ &= \langle \hat{k}_\gamma \cdot \hat{x} \rangle \\ &= \langle \sin(\theta) \cos(\phi) \rangle. \end{aligned} \quad (5)$$




FIG. 4. Top-down view of the detector array. Beam direction is $+\hat{z}$, beam left is $+\hat{x}$, and $+\hat{y}$ corresponds to the spin-up neutron direction.




FIG. 5. A beam's eye view of a ring of detectors. Detectors i and j are an example of a pair that in the ideal case would have geometrical factors of equal magnitude but opposite sign.

The $\hat{y}$ -direction, or up-down (U-D), geometrical factor is proportional to the parity violating asymmetry. It is given by:

$$G_{\text{PV}} = \langle \hat{k}_\gamma \cdot \vec{\sigma}_n \rangle = \langle \hat{k}_\gamma \cdot \hat{y} \rangle = \langle \sin(\theta) \sin(\phi) \rangle. \quad (6)$$

In the ideal case, a pair of detectors i and j , as shown in Fig. 5 will have the same geometrical factors, but of opposite sign. The geometrical factors account for finite beam, detector dimensions, and neutron scattering in the target vessel and can be computed using MCNPX. Source code modifications are necessary to weight the energy deposition in individual detectors by the initial photon emission direction from neutron capture events. The results of this calculation for G_{PV} are shown in Fig. 6. Detectors with the smallest angle with respect to the vertical, $\hat{y}$ direction, have the largest up-down geometrical factors, as they have the highest sensitivity to an up-down parity violating asymmetry.

The detector response to a Cs source was fit to the MCNPX calculation for the same geometry along with a small rotation, ϕ , in order to account for unequal detector half-crystal efficiencies, giving modified geometrical factors of:

$$\begin{aligned} G'_{\text{PC}} &= \langle \hat{k}_\gamma \cdot \hat{x} \rangle' \\ &= \langle \sin(\theta) \sin(\phi + \delta_\phi) \rangle \\ &= \langle \hat{k}_\gamma \cdot \hat{x} \rangle \cos(\delta_\phi) + \langle \hat{k}_\gamma \cdot \hat{y} \rangle \sin(\delta_\phi) \\ G'_{\text{PC}} &= G_{\text{PC}} \cos(\delta_\phi) + G_{\text{PV}} \sin(\delta_\phi). \end{aligned} \quad (7)$$

and

$$G'_{\text{PV}} = G_{\text{PV}} \cos(\delta_\phi) - G_{\text{PC}} \sin(\delta_\phi). \quad (8)$$

A different set of geometrical factors must be applied whenever the position of the chlorine target is changed relative to the position of the detector, as the acceptance changes. Measurements of the chlorine asymmetry in different geometries were used to obtain the systematic uncertainty associated with




FIG. 6. In the above figure, the *PV* geometrical factors for ideal detectors are shown. Detectors are numbered starting at 0 with the upstream, beam-right detector, and continuing clockwise. For example, detectors 12 and 23 are located in the second ring, top, beam-right and beam-left, respectively.

their determination. Details of the calculation and validation of the geometrical factors is available in Ref. [11].

D. Beam polarization

The asymmetry we are trying to measure is neutron energy independent, but the polarization of the beam is not. Since the beam polarization is a multiplicative correction to the asymmetry, we need to know its energy dependence.

An auxiliary experiment was performed in order to establish the polarization of the beam. A polarized ^3He cell was used as a spin filter, along with the RFSR and a ^3He ion chamber flux monitor to perform a series of transmission measurements. These allow us to extract the polarization of the neutron beam as well as the efficiency of the RFSR without needing to know the polarization of the ^3He cell. This is done by taking advantage of the well-known spin dependence of the capture cross section of cold neutrons in polarized ^3He [20–22].

In order to obtain the polarization and the RFSR efficiency across the whole area of the beam ($12 \times 10 \text{ cm}^2$), the ^3He cell and monitor were moved in a $3 \times 3 \text{ cm}^2$ grid, with nine independent measurements, which were averaged together, weighted by the beam flux in each area. The details of the procedure and analysis are described in Ref. [23]. The average polarization and RFSR efficiency over the area of the beam in the energy range used in the asymmetry measurement were determined to be 0.939 ± 0.004 and 0.974 ± 0.009 , respectively.

E. Asymmetry determination

There is more than one way to extract the physics asymmetry from the detector data. One approach is a pair analysis, where asymmetries are formed for two conjugate detectors, i and j , (at equal and opposite angles relative to the spin of the neutrons, see Fig. 5). If the detector gains are matched to




FIG. 7. Raw geometrical asymmetry determined for each detector pair for the CONF1 data set.

within a few percent, this approach will cancel beam fluctuation effects, as well as any systematic effects common to all detectors. A raw asymmetry is calculated for each eight-step spin sequence ($\uparrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow\uparrow$) using conjugate detector yields, N_i and N_j via the geometrical mean $\sqrt{\alpha}$:

$$A_{ij}^{\text{raw}} = \frac{\sqrt{\alpha} - 1}{\sqrt{\alpha} + 1}, \text{ where } \alpha = \left[\frac{N_i^\uparrow}{N_i^\downarrow} \right] \left[\frac{N_j^\uparrow}{N_j^\downarrow} \right]. \quad (9)$$

The raw chlorine asymmetry is plotted as a function of detector pair number in Fig. 7. Each raw asymmetry consists of contributions from L-R parity-conserving and U-D parity-violating physics asymmetries. The sensitivity to the latter is maximal for detectors whose position is closest to the vertical (most aligned with magnetic field), and decreases as one moves away towards the horizontal, which gives rise to the pattern seen in Fig. 7.

As previously mentioned, many common mode effects will cancel at least to first order in the extraction of pair asymmetries, potentially concealing underlying issues. In the interest of thoroughness and to benchmark the experimental apparatus, we also extract the asymmetries for each individual detector i via:

$$A_i = \frac{N_i^\uparrow - N_i^\downarrow}{N_i^\uparrow + N_i^\downarrow}. \quad (10)$$

In the discussion to follow, we show that detector asymmetries are useful for diagnosing possible problems. However, the error extracted from the RMS width of binned detector asymmetries contains contributions from beam fluctuations, in addition to counting statistics. This contribution is $\approx 15\%$. For this reason, in the final analysis, pair asymmetries were used for the proper propagation of statistical error. The results of the raw detector asymmetry from the first set of chlorine measurements (CONF1) can be seen in Fig. 8. However, there's a visible negative offset to the results, of $\approx 2 \times 10^{-6}$, larger than what is present in the results for detector pairs. This effect was investigated and two causes for it were isolated, to be discussed in the next section.




FIG. 8. Raw asymmetry calculated for each detector. The lines correspond to the calculated geometrical factors, G_{PV}^i , scaled for comparison, to illustrate that the measured raw asymmetries are primarily the result of an PV asymmetry. The grid is to show the highlight the offset in the vertical direction.

IV. SYSTEMATIC EFFECTS

A. Instrumental false asymmetries

Investigations revealed two sources for the offset observed in the CONF1 single-detector raw asymmetries: an exponential transient present in the ADC as well as crosstalk between the ADC channels. The data are recorded in an ADC buffer for the duration of eight accelerator pulses and written to file during the ninth. Connecting a 9V battery as input to one of the ADC channels revealed a transient in the form of an exponential decay at the beginning of each data cycle that we believe corresponds to a discharge of a capacitor in the ADC as can be seen in Fig. 9. Each eight pulses of data had the ($\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow$) spin configuration, leading to a false asymmetry, as the signal is enhanced the most by the transient in the first (spin-up) state. The size of the false asymmetry was comparable in each ring of detectors due to the fact that




FIG. 9. Transient signal measured in a monitor ADC channel, with a 9.48 V battery signal as the input.




FIG. 10. The additive (a) and multiplicative (b) asymmetries measured after the sources of false asymmetries were eliminated. Both instrumental asymmetries are consistent with zero at the 10^{-9} level.

the detector signals were separated into a ring average and a difference from the average for each detector, meaning that 13 signals (ring average and 12 differences) were recorded for each ring (an artifact of the DAQ configuration for a previous iteration of the experiment). This configuration was also responsible for the second source of the offset seen in the asymmetries as a function of detector: the ring averages were read into the same ADC as the spin-dependent information, where nonzero crosstalk between channels was observed. Two hardware changes were implemented to eliminate the above problems. The spin sequence was alternated between $\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow$ (sequence A) and $\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow$ (sequence B) and only pairs of sequences were analyzed, leading to a cancellation of the transient-induced asymmetry. Additionally, the detector signals were kept intact and read out by a reserved ADC in an electrically isolated VME crate. This eliminated all false asymmetries at the 10^{-9} level. Additive and multiplicative asymmetries were measured. The former with beam off, but all other experimental components running. The multiplicative asymmetry was measured in the same conditions, but with a signal from LEDs induced in the CsI crystals, comparable to the size of the beam-on signal. The results from one set of those measurements can be seen in Fig. 10.

B. Selection of data set

The CONF1 data set is not included in our quoted value for the PV chlorine asymmetry in this work. The chlorine measurements were repeated after all of the described instrumentation issues were mitigated and the chlorine target case was also replaced to eliminate the need to subtract the aluminum asymmetry from the prompt γ rays. Finally, the choppers were rephased to eliminate 13–15 Å leakage neutrons, leaving only those from 28–30 Å. Data taken after these changes are referred to as “CONF2”.

C. Leakage neutrons

The choppers allow through 28–30 Å neutrons, which have a lower polarization than the main beam, as a larger fraction of these makes it through the polarizer without a bounce. Additionally, these neutrons are not fully rotated by the RFSR, as the field is not optimized for their energies. This will lower the average beam polarization. The fraction of the signal that comes from the leakage neutrons is 0.2% of the total. Leakage neutrons for pulse X will appear in pulse X + 7. For half of the pulses, the RFSR will not be on, so the polarization of the leakage neutrons will be preserved and be correct. The remaining pulses (with RFSR on) will be rotated nine times, meaning the wrong spin state will come through. Given this situation, and assuming an initial beam polarization of 90% for the long wavelength neutrons, the beam polarization becomes:

$$P'_n = 0.998 \times P_n + 0.9 \times 0.001 + 0.9 \times (-1) \times (.001). \quad (11)$$

With a conservative assumption of 50% uncertainty on the amount of leakage neutrons, this changes P'_n by 0.001, resulting in an uncertainty of 0.1%.

D. Beam depolarization

The neutrons in the beam can be depolarized via incoherent scattering before being captured. This effect was modeled in MCNPX to obtain a depolarization correction. Interactions with the following isotopes were included in the calculation: ^1H , ^6Li , ^{14}N , ^{27}Al , ^{35}Cl , ^{37}Cl , ^{55}Mn , ^{63}Cu , ^{65}Cu , Zn (natural). The calculation shows that 1.6% of the beam is depolarized with a statistical uncertainty of 0.03%.

TABLE II. Summary of systematics with negligible contributions.

Additive Asymmetry	$<1 \times 10^{-9}$
Multiplicative Asymmetry	$<1 \times 10^{-9}$
Stern-Gerlach	8×10^{-11}
γ -ray circ. pol	$<1 \times 10^{-12}$
β decay in flight	$<1 \times 10^{-11}$
Capture on ^6Li	$<1 \times 10^{-11}$
Radiative β -decay	$<1 \times 10^{-12}$
β -delayed Al γ s (internal+external)	$<1 \times 10^{-9}$




FIG. 11. Raw chlorine asymmetries for data set designated CHL3. The line represents the PV geometrical factors scaled by -21.6×10^{-6} . They reproduce the shape of the data well, signaling that the raw asymmetry is due to the large PV component.

E. Systematic effects

Additional physics processes can either result in an up-down asymmetry or a parity conserving left-right asymmetry. The latter, if the detector array is not well aligned, can mix into the up-down asymmetry. They have been previously evaluated (Table II [24]), confirmed, and are negligible for this measurement.

V. RESULTS

The chlorine target was used on several occasions to extract the parity-violating and parity-conserving asymmetries. The CONF2 data sets include running with 36 (intermediate configuration) and 48 (full array) detectors. Figure 11 shows a data set with the full array.

The asymmetry calculated for each detector (or detector pair) contains a PC and PV physics contribution, whose magnitude depends on their respective geometrical factors as shown in Eq. (12). A fit is performed using the 48 (24) detector (pair) asymmetries and the 96 (48) geometrical factors to extract A_{PV} , A_{PC} , and an offset. The last parameter should be

TABLE III. Raw PV and PC asymmetries from CHL1-4, obtained from fits using geometrical factors. Uncertainties shown are statistical only.

Data Set	A_{PV}	A_{PC}
CHL1	$(-21.6 \pm 0.3) \times 10^{-6}$	$(-0.1 \pm 0.3) \times 10^{-6}$
CHL2	$(-21.6 \pm 0.3) \times 10^{-6}$	$(-0.4 \pm 0.3) \times 10^{-6}$
CHL3	$(-20.9 \pm 0.3) \times 10^{-6}$	$(0.1 \pm 0.3) \times 10^{-6}$
CHL4	$(-21.8 \pm 0.2) \times 10^{-6}$	$(0.3 \pm 0.2) \times 10^{-6}$
AVE	$(-21.5 \pm 0.1) \times 10^{-6}$	$(0.1 \pm 0.1) \times 10^{-6}$
Corrected	$(-23.9 \pm 0.1) \times 10^{-6}$	$(0.1 \pm 0.1) \times 10^{-6}$




FIG. 12. Raw PV chlorine asymmetries are shown from measurements done in three different geometrical configurations. The inner error bar is statistical only, whereas the outer error bar is total, once the geometrical factor uncertainty has been added.

consistent with zero, and is used as a diagnostic.

$$A_{\text{raw}} = A_{\text{PV}} \cdot G_{\text{PV}} + A_{\text{PC}} \cdot G_{\text{PC}} + \text{offset}. \quad (12)$$

Four CONF2 data sets (CHL1-4) were taken in multiple geometries (target inside the detector array, center and displaced downstream, as well as target inside the RFSR). Comparing the results from three measurement geometries (Table III) allows for a determination of the uncertainty associated with the geometrical factors. In order for the three results from the different geometrical configurations to be consistent (i.e., χ^2 of one when fit to a constant), a 3% systematic error needed to be assigned to the geometrical factor determination. This is illustrated in Fig. 12.

We also analyzed the parity-conserving left-right asymmetry, $A_{\gamma, \text{PC}}$, with the result of $(0.1 \pm 0.7) \times 10^{-6}$, consistent

with zero. Our result is in agreement with what one would expect on theoretical grounds given the statistical error of our measurement [25].

VI. CONCLUSION

We have performed the most precise measurement of the parity-violating asymmetry in cold neutron capture on ^{35}Cl , yielding $A_{\gamma, \text{PV}} = (-23.9 \pm 0.7) \times 10^{-6}$, with a parity-even asymmetry consistent with zero. We have presented in detail the chronology of testing the experimental design, finding and eliminating sources of false asymmetries, and determining the uncertainty associated with geometrical factors.

ACKNOWLEDGMENTS

We gratefully acknowledge the support of the U.S. Department of Energy Office of Nuclear Physics through Grants DE-AC52-06NA25396, DE-FG02-03ER41258, DE-SC0014622, DE-AC-02-06CH11357, DE-AC05-00OR22725, and DE-SC0008107, the U.S. National Science Foundation through Grants PHY-1306547, PHY-1306942, PHY-1614545, PHY-0855584, PHY-0855610, PHY-1205833, PHY-1506021, PHY-1205393, and PHY-0855694, PAPIIT-UNAM Grants IN110410, IN11193, and IG1011016, CONACYT Grant 080444, the Natural Sciences and Engineering Research Council of Canada (NSERC), and the Canadian Foundation for Innovation (CFI). This research used resources of the Spallation Neutron Source of Oak Ridge National Laboratory, a DOE Office of Science User Facility. J.F., R.C.G., J.M., W.M.S., and Z.T. acknowledge support from the Indiana University Center for Spacetime Symmetries. S. Schröder acknowledges support from the German Academic Exchange Service (DAAD).

- [1] E. G. Adelberger and W. C. Haxton, *Annu. Rev. Nucl. Part. Sci.* **35**, 501 (1985).
- [2] M. R. Schindler and R. P. Springer, *Prog. Part. Nucl. Phys.* **72**, 1 (2013).
- [3] W. C. Haxton and B. R. Holstein, *Prog. Part. Nucl. Phys.* **71**, 185 (2013).
- [4] B. Desplanques, J. F. Donoghue, and B. R. Holstein, *Ann. Phys. (NY)* **124**, 449 (1980).
- [5] H. W. Griesshammer and M. R. Schindler, *Eur. Phys. J. A* **46**, 73 (2010).
- [6] D. Blyth *et al.* (NPDGamma Collaboration), *Phys. Rev. Lett.* **121**, 242002 (2018).
- [7] M. T. Gericke *et al.* (NPDGamma Collaboration), *Phys. Rev. C* **74**, 065503 (2006).
- [8] M. Avenier *et al.*, *Nucl. Phys. A* **436**, 83 (1985).
- [9] V. A. Vesna *et al.*, *JETP Lett.* **36**, 209 (1982).
- [10] G. S. Mitchell *et al.* (NPDGamma Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **521**, 468 (2004).
- [11] K. Grammer, D. Blyth, J. Bowman, N. Fomin, G. Greene, M. Musgrave, E. Tang, and Z. Tang, *Nucl. Instrum. Meth. Phys. Res. A* **903**, 302 (2018).
- [12] N. Fomin *et al.*, *Nucl. Instrum. Meth. Phys. Res. A* **773**, 45 (2015).
- [13] F. Mezei, *Comm. Phys.* **1**, 81 (1976).
- [14] S. Balascuta *et al.*, *Nucl. Instrum. Meth. Phys. Res. A* **671**, 137 (2012).
- [15] P. N. Seo, L. Barron-Palos, J. D. Bowman, T. E. Chupp, C. Crawford, M. Dabaghyan, M. Dawkins, S. J. Freedman, T. Gentile, M. T. Gericke, R. C. Gillis, G. L. Greene, F. W. Hersman, G. L. Jones, M. Kandes, S. Lamoreaux, B. Lauss, M. B. Leuschner, R. Mahurin, M. Mason, J. Mei, G. S. Mitchell, H. Nann, S. A. Page, S. I. Penttila, W. D. Ramsay, A. SalasBacci, S. Santra, M. Sharma, T. B. Smith, W. M. Snow, W. S. Wilburn, and H. Zhu, *Phys. Rev. ST Accel. Beams* **11**, 084701 (2008).
- [16] M. McCrea, Ph.D. thesis, University of Manitoba, 2016.
- [17] M. T. Gericke *et al.*, *Nucl. Instrum. Meth. Phys. Res. A* **540**, 328 (2005).
- [18] G. Audi, F. G. Kondev, M. Wang, W. J. Huang, and S. Naimi, *Chin. Phys. C* **41**, 030001 (2017).
- [19] Z. Tang, Ph.D. thesis, Indiana University, 2014.

- [20] S. Mughabhab, M. Divadeenam, and N. Holden, *Neutron Resonance Parameters and Thermal Cross Sections* (Academic Press, New York, 1981).
- [21] H. H. Barschall and W. Haeblerli, *Polarization Phenomena in Nuclear Reactions: Proceedings of the Third International Symposium* (University of Wisconsin Press, Madison, 1971).
- [22] J. Als-Nielsen and O. Dietrich, [Phys. Rev. **133**, B925 \(1964\)](#).
- [23] M. M. Musgrave *et al.*, [Nucl. Instrum. Meth. Phys. Res. A **895**, 19 \(2018\)](#).
- [24] M. T. Gericke *et al.*, [Phys. Rev. C **83**, 015505 \(2011\)](#).
- [25] M. T. Gericke, J. D. Bowman, and M. B. Johnson, [Phys. Rev. C **78**, 044003 \(2008\)](#).