

ARTICLE

Light-Mediated Aerobic Oxidation of C(sp³)–H Bonds by a Ce(IV) Hexachloride Complex

Yu-Heng Wang, §‡ Qiaomu Yang, § Patrick J. Walsh* and Eric J. Schelter*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x



A photochemical C(sp³)–H oxygenation of alkane and arene substrates catalyzed by [NEt₄]₂[Ce^{IV}Cl₆] under mild conditions (1 atm, 25 °C) is described. Time-course studies reveal that the hydrocarbons are oxidized in a stepwise fashion to afford alcohols, aldehydes, ketones, and carboxylic acids. The catalyst resting state, [Ce^{IV}Cl₆]^{2–}, is observed by UV-visible spectroscopy. On/off light-switching experiments, quantum yield measurements, and the absence of a kinetic isotope effect on parallel C–H/C–D functionalization suggest that ligand-to-metal charge transfer of [NEt₄]₂[Ce^{IV}Cl₆] to generate Cl• is the turnover-limiting step. The involvement of a highly reducing excited-state [NEt₄]₂[Ce^{III}Cl₆]* species as well as photo-excited aldehyde, under black light irradiation appears to facilitate the conversion of primary alcohols and aldehydes to carboxylic acids. Remarkably, this approach is found to be capable of direct activation of light alkanes, including methane and ethane.

Introduction

The oxygenation of organic compounds is a fundamental reaction in biology and chemistry.^{1–3} In biology, cells use oxygen to convert glucose into energy in cellular respiration.⁴ Oxidative reactions, including C–H oxygenations, play a significant role in organic transformations and chemical synthesis.^{5–8} The C–H oxygenation of hydrocarbons has been extensively applied to afford valuable products such as fine chemicals, feedstocks, and pharmaceuticals.^{9–16} Previous approaches to C–H oxygenation, however, typically require excess terminal oxidants (e.g., iodosobenzene, ^tbutyl hydroperoxide),^{15, 16} harsh reaction conditions (e.g., elevated temperature and pressure), and noble-metal catalysts (e.g., Pd, Ir, and Pt) and, therefore, are not environmentally benign (Scheme 1a).^{9–16} The abundance of molecular oxygen in the Earth's atmosphere renders catalytic C–H oxygenation using O₂ as the terminal oxidant particularly attractive.^{11–13, 15–20} Additionally, low-cost metal catalysts are needed for practical applications. The development of new methods utilizing molecular oxygen and earth-abundant catalysts to achieve the C–H oxygenation of hydrocarbons, including light alkanes, is, therefore, a high priority.

Photoredox catalysis provides an opportunity to achieve C(sp³)–H oxygenations under mild and environmentally friendly conditions. Molecular catalyst systems that are low-cost, benign, and use molecular oxygen, have been achieved for light-mediated

C(sp³)–H oxygenations, such as decatungstate (under UVA light), uranyl, acridinium and flavin-based photocatalysts (under blue or purple light), and others (Scheme 1b).^{21–26} Recently, Cl• has been studied extensively and employed in alkylarene oxygenation reactions. Recently, Hu, Jiang and Zeng employed iron catalysts^{27–29}; Wang's group used HCl as a catalyst and Shi and co-workers disclosed cerium photoredox catalysis³⁰ during the final stages of the investigations presented herein. However, the substrate scope of these systems was rather restricted and was generally limited to alkyl aromatics.^{31–38} Notably, Fukuzumi and others reported that stronger alkyl C–H bonds, such as those in cyclohexane, could be photochemically activated by chlorine radical (Cl•) under aerobic conditions. However, the generation of the oxidation products in those studies was not particularly selective.^{39, 40} These constraints inspired us to develop an alternative strategy for light-mediated catalytic C(sp³)–H oxygenations with broader substrate scope and controllable selectivity.



Scheme 1 Overview of the established strategies (a and b) and our strategy c for light-mediated oxygenation of C(sp³)–H bonds.

P. Roy and Diana T. Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States. Email: schelter@sas.upenn.edu; pwalsh@sas.upenn.edu

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

§ These authors contributed equally.

‡ Presently at Department of Chemistry, Tsing Hua University, Hsinchu 30013, Taiwan.

Table 1 Optimization of the reaction conditions using blue LEDs.

| Entry | Variations from standard conditions ^[a] | ¹ H NMR AY (%) ^[b] | | | |
|-------|--|--|-------|--------|-------|
| | | 1a-i | 1a-ii | 1a-iii | total |
| 1 | 18 h, 1 atm air | ≈ 5 | 21 | 26 | 52 |
| 2 | 18 h, 1 atm O ₂ | ≈ 5 | 25 | 28 | 58 |
| 3 | none | ≈ 5 | 24 | 29 | 58 |
| 4 | in the dark | – | – | – | – |
| 5 | in the dark, 80 °C | – | – | – | – |
| 6 | black light ^[c] | – | – | ≈ 5 | ≈ 5 |
| 7 | no [NET ₄] ₂ [Ce ^{IV} Cl ₆] | – | – | – | – |
| 8 | 1 atm N ₂ | – | – | – | – |
| 9 | DMF instead of CH ₃ CN ^[d] | – | – | – | – |
| 10 | DME instead of CH ₃ CN ^[e] | – | – | – | – |
| 11 | 10 mol% [NET ₄] ₂ [Ce ^{IV} Cl ₆] | – | 34 | 40 | 74 |
| 12 | 48 h, 1 atm air | – | 36 | 47 | 83 |

[a] Initial conditions: 0.4 mmol toluene, 0.02 mmol **I**, 4 mL CH₃CN in a 8 mL sample vial, irradiated with a Kessil A160WE blue LED lamp with fan cooling at room temperature (rt, 298 K) under aerobic conditions for 24 h. [b] Yield determined by ¹H NMR spectroscopy with 1,4-dioxane as an internal standard. [c] The reaction was irradiated with four 24 W (100 W equivalent) CFL black light bulbs. [d] DMF = dimethylformamide. [e] DME = dimethoxyethane.

Herein, we report the light-mediated oxygenation of various C(sp³)-H bonds in hydrocarbons catalyzed by [NET₄]₂[Ce^{IV}Cl₆] (**I**) with O₂ as the terminal oxidant (Scheme 1c). The influence of radiation wavelengths on product selectivities and yields was investigated, and the crucial role of light frequency on the catalytic cycle has been revealed.

It was proposed that chlorine radical (Cl•) was responsible for the C(sp³)-H bond cleavage,⁴¹⁻⁴⁹ consistent with recent work from our group, where the generation of Cl• radicals was demonstrated as the rate-limiting step, and the Cl• reactivity could be altered by the addition of suitable additives (alcohols and arenes).⁵⁰ The oxygenation of gaseous alkanes (i.e., CH₄ and C₂H₆) mediated by

[NET₄]₂[Ce^{IV}Cl₆] was studied with the goal of transformation of light alkanes into liquid fuels under mild conditions, in contrast with ceria based catalysts that employ harsh conditions.^{51, 52}

Results and discussion

Our previous studies indicated that the C(sp³)-H bond of toluene was activated by catalytic [NET₄]₂[Ce^{IV}Cl₆] using visible light irradiation under anaerobic conditions.⁵³ This finding led us to question whether photoredox catalyzed oxygenation of C(sp³)-H bonds of hydrocarbons could be achieved by [NET₄]₂[Ce^{IV}Cl₆] under aerobic conditions. Toluene (**1a**) was selected as the model substrate for studying oxygenation in the presence of dioxygen due to its moderate C-H bond strength (bond dissociation energy [BDE] = 88.5 kcal/mol).⁵⁴ The study was initiated with blue light irradiation of a 0.1 M solution of toluene with 5 mol% [NET₄]₂[Ce^{IV}Cl₆] in acetonitrile (CH₃CN) at room temperature (298 K) for 18 h under 1 atm of air. The oxidation products PhCHO (**1a-ii**, BDE of the formyl C-H bond = 89 kcal/mol)⁵⁴ and PhCO₂H (**1a-iii**) were observed in 21% and 26% assay yield, respectively (AY, determined by ¹H NMR integration against an internal standard) (Table 1, entry 1). Benzyl alcohol is presumably an intermediate in this process. However, the benzylic C-H bonds are notably weaker and more reactive in benzyl alcohol (BDE = 80 kcal/mol)⁵⁴ than in toluene, so it is not surprising that benzyl alcohol is only observed in the initial stage of the reaction. The AYs did not change significantly when the reaction was conducted under 1 atm of dioxygen under otherwise identical conditions (entry 2). Increasing the reaction time to 24 h increased the reaction AY to 58% (entry 3). Thus, 24 h was set as the standard reaction time. No reaction was observed in the dark either at room temperature (entry 4) or 80 °C (entry 5), suggesting that product formation is light-mediated.

We next conducted control experiments to verify the importance of the reaction components. The selection of light frequency is critical in light-mediated reactions. We were surprised to find that there was no conversion observed when the reaction was irradiated with black light (λ_{max} = 330 nm) instead of blue light (λ_{max} = 467 nm entry 6). No product was observed in the absence of the cerium catalyst **I** (entry 7) or molecular oxygen (entry 8). The oxidation reaction proved to be sensitive to the type of solvent: no oxidation of toluene was detected in dimethylformamide (DMF) or dimethoxyethane (DME), (entries 9 and 10).⁵⁵ The absence of products was presumably due to the limited solubility of **I** in DME and its relative instability in DMF during the reaction.

We next examined the evolution of toluene oxidation products through a time-course study using ¹H-NMR spectroscopy (Figure 1 and Section II of the Supplemental Information). Over the 24 h course of the oxidation, a low concentration of PhCH₂OH (**1a-i**) was observed in the initial stage of the reaction that was maintained as the reaction proceeded (Figure 1, green trace), indicating that the oxidation of PhCH₂OH is kinetically facile relative to toluene. The production of benzaldehyde (Figure 1, blue trace) and benzoic acid (Figure 1, red trace) occurs through the oxidation of PhCH₂OH. The observation of PhCH₂Cl is attributed to radical coupling of Cl• with the benzylic radical (Figure 1, yellow trace) and is consistent with the proposal that Cl• is the active oxidant. The steady-state concentration of PhCH₂Cl (≈ 10 %) suggests that it can be converted





Fig. 1 Time-course studies of C-H oxygenation of toluene by ^1H NMR spectroscopy: distribution of oxidation products. Reaction conditions: 0.1 M toluene, 0.005 M **I** (5 mol%), 0.5 mL CD_3CN in an NMR tube (0.38 mm thickness), irradiated with a Kessil A160WE blue LED lamp with fan cooling at 298 K under aerobic conditions for 24 h.

into oxidation products PhCHO or PhCO_2H (Figure S17) and that catalyst degradation through chlorine sequestration with the formation of benzyl chloride is reversible. The addition of exogenous $[\text{NEt}_4][\text{Cl}]$ was envisioned to provide a chloride source, which is partially consumed in the formation of benzyl chloride.⁵³ Notably, the addition of $[\text{NEt}_4][\text{Cl}]$ led to a decrease in oxidation products and an increase in side products, such as benzyl chloride (Figure S5). The formation of benzyl hydroperoxide was negligible (< 5%) and only observed in the first hour over the time-course study (Figure S5). The detection of benzyl hydroperoxide indicated that the trapping of benzyl radical by dioxygen is involved in the reaction mechanism.

To probe the reaction mechanism, kinetic isotope effect (KIE) studies on the Ce-catalyzed C-H oxygenation of toluene were conducted (Section IV of the Supplemental Information). A parallel KIE study with PhCH_3 and PhCD_3 revealed a KIE of 1.1. This small KIE indicates that the C-H/C-D cleavage is not involved in the turnover-limiting step (Scheme 2a and Figures S9–10).^{56, 57} A KIE value for an intermolecular competition experiment, in which the reaction was conducted with equimolar amounts of PhCD_3 and PhCH_3 in the same reaction vessel, was found to be 1.0 (Scheme 2b and Figure S11).

These results, together with the on-off experiments and quantum yield measurements (vide infra), support the hypothesis that the photo-induced generation of Cl^\bullet is likely the turnover-limiting step, which is followed by hydrogen atom transfer (HAT) from toluene (Scheme 2c).⁵⁸

Having established the turnover-limiting step, we sought to determine if different light wavelengths would impact the distribution of the oxidation products (i.e., the ratio of PhCO_2H to PhCHO).⁵⁹ The influence of irradiation wavelength on photocatalysis has been explored previously, although heterogeneous catalysts were used in most of these studies.^{60–64} Here, under purple light ($\lambda_{\text{max}} = 390 \text{ nm}$) irradiation, the overall yield of toluene oxygenation products was found to be similar to that with blue light (Table 2, entries 1 vs. 2); however, the selectivity shifted to favor the formation of benzoic acid ($\text{PhCO}_2\text{H} : \text{PhCHO}$ ranged from 1.2 to 1.9




Scheme 2 Kinetic isotope effect experiments: a) two parallel reactions, b) an intermolecular competition, and c) the proposed turnover-limiting step (TLS). Reaction conditions: a) 0.05 mmol PhCH_3 or PhCD_3 , 0.0025 mmol **I**, 0.5 mL CD_3CN in an NMR tube (0.38 mm thickness); b) 0.2 mmol PhCH_3 and PhCD_3 , 0.01 mmol **I**, 2 mL CD_3CN in an 8 mL sample vial. Reactions were irradiated with a Kessil A160WE blue LED lamp with fan cooling at 298 K.

(Table 2, entries 1 vs. 2). This observation implies that the selectivity of C-H oxygenation is affected by the irradiation with different wavelengths (λ_{max}), which provides an opportunity to optimize the reaction using different light sources.⁶⁵ Based on previous studies, a critical step in the catalytic process is the oxidation of Ce(III) back to Ce(IV).^{53, 55} $[\text{NEt}_4]_3[\text{Ce}^{\text{III}}\text{Cl}_6]$ (**II**) is a weak reductant (half-wave potential, $E_{1/2} \approx 0.03 \text{ V}$ vs. Fc^+/Fc), but the photoexcited state of $[\text{Ce}^{\text{III}}\text{Cl}_6]^{3-*}$ (**III**) is a strong reductant ($E_{1/2} \approx -3.45 \text{ V}$ vs. Fc^+/Fc).⁵³ We hypothesized that use of the black light in the catalyst reoxidation step might provide an opportunity to impact catalyst reactivity/selectivity by facilitating the reduction of oxidized intermediates (i.e., peroxides).

We, therefore, set out to examine the impact of a second light source on the oxygenation process.^{66–68} We found that the overall yield of toluene oxygenation was not impacted when blue and black lights were used simultaneously (Table 2, entries 3 vs. 1). The selectivity of products, however, was impacted and favored PhCO_2H ($\text{PhCO}_2\text{H} : \text{PhCHO} = 2.0$). Employing a combination of purple/black lights resulted in an increase in yield (78%) and selectivity ($\text{PhCO}_2\text{H} : \text{PhCHO} = 2.9$) (Table 2, entry 4), revealing that the influence of light source on catalytic oxygenation is considerable and merits further attention.

To further understand how the light source influences the reaction outcome, control experiments were performed on reaction substrates and intermediates using different light sources. Product distributions were analyzed after a 24-h reaction period (Section VIII-X of the Supplemental Information). Initially, *p*-tolunitrile was employed as substrate. Irradiation with a combination of purple and black lights resulted in more carboxylic acid generated (21%) than blue light (trace yield) (Figure S17). When benzyl chloride was used as substrate, it showed the same behaviour (Figure S18).


Table 2 Influence of radiation wavelengths on C–H oxygenation of toluene.

|  | | | | | | |
|---|-----------------------------|---------------------------|-------|-------------------|-------|----------------------------|
| Entry | Light source ^[a] | NMR AY (%) ^[b] | | | | Selectivity (1a-iii/1a-ii) |
| | | 1a-i | 1a-ii | 1a-iii | total | |
| 1 ^[c] | blue LED | ≈ 5 | 24 | 29 | 58 | 1.2 |
| 2 | purple LED ^[d] | ≈ 5 | 19 | 37 | 61 | 1.9 |
| 3 ^[e] | blue LED & black light | ≈ 5 | 16 | 32 | 53 | 2.0 |
| 4 ^[f] | purple LED & black light | ≈ 5 | 19 | 54 | 78 | 2.9 |
| 5 ^[g] | black light | – | – | 84 ^[g] | – | – |

[a] General reaction conditions: 0.4 mmol toluene, 0.02 mmol I, 4 mL CH₃CN, irradiated with *different lights* at 298 K under aerobic conditions for 24 h. [b] Assay yield determined by ¹H NMR spectroscopy with 1,4-dioxane as an internal standard. [c] Entry 1 in Table 1. [d] A Kessil PR160-390 LED lamp (λ_{max} = 390 nm). [e] A Kessil A160WE blue LED lamp (λ_{max} = 467 nm) and four 24 W (100 W equivalent) CFL black light bulbs (λ_{max} = 330 nm). [f] A Kessil PR160-390 LED lamp and four 24 W (100 W equivalent) CFL black light bulbs. [g] [NET₄]₂[Ce^{IV}Cl₆]: 0.0 M; benzaldehyde (**1a-ii**): 0.04 M; toluene: 0.1 M; yield of **1a-iii** was based on 0.04 M benzaldehyde. Benzoic anhydride was included in the benzoic acid yield for convenience.

Control experiments were then performed on PhCH₂OH (**1a-i**), PhCHO (**1a-ii**) and PhC(CH₃)₂OOH. Irradiation of PhCH₂OH under catalytic conditions with purple/black light gave PhCO₂H and PhCHO with a selectivity of 14.5 in a combined AY of 93% (Scheme 3a). Using only blue light led to 83% AY but with decreased selectivity of 3.0. We next examined the oxidation of PhCHO to PhCO₂H. With purple/black or blue light irradiation, both reactions proceeded with similar AY (61–64%, Scheme 3b). The difference in product distributions observed in Scheme 3a and the same AY in Scheme 3b, suggests that there is one reaction pathway producing PhCO₂H (**1a-iii**) from PhCH₂OH (**1a-i**) that bypasses the formation of PhCHO (**1a-ii**).

The light-mediated oxidation of **1a-i** to **1a-iii** has been proposed to proceed via a hydroperoxide intermediate, PhC(CH₃)₂OOH.⁶⁹ As a model peroxide for this intermediate PhC(CH₃)₂OOH was chosen (see Section X of the Supplemental Information). We found that PhC(CH₃)₂OOH was reactive under light irradiation without the cerium catalyst, especially under the purple/black light combination (Figure S23–S25). This indicates that PhC(CH₃)₂OOH can absorb light (hν), causing its decomposition by O–O bond cleavage.⁷⁰ In addition, under irradiation PhC(CH₃)₂OOH is consumed faster in the presence of the cerium(III) catalyst. We hypothesize that the black light photoexcites [NET₄]₃[Ce^{III}Cl₆] species (**II**) to [NET₄]₃[Ce^{III}Cl₆]*, which




Scheme 3 The influence of radiation wavelengths on difference steps in the C–H oxygenation of toluene: a) oxidation of PhCH₂OH to PhCHO and PhCO₂H, b) oxidation of PhCHO to PhCO₂H, and c) a plausible reaction pathway for the oxidation of PhCH₂OH to PhCO₂H via HAT.

reduces the peroxide. In the catalytic oxidation of benzyl alcohol, we propose that the acceleration of the reaction by the purple/black light combination is due to 1) the light absorption by the peroxide (see Supplemental Information, section X) and 2) the generation of the strong photoreductant [NET₄]₃[Ce^{III}Cl₆]* (**III**),⁵³ ultimately generating benzoic acid. At this point, the origin of the different products with purple/black light irradiation is speculated to arise by two possible pathways. One involves reaction of excited state Ce(III) reduction of peroxy intermediates (Scheme 3c). The other involves direct photon absorption by benzylic peroxide intermediates and further reactions. The exact mechanism of the formation of benzoic acid, however, remains under investigation.⁷¹

Aside reaction for benzaldehyde under black light irradiation was also considered (See Supplemental Information, Section IX). The triplet excited state of benzaldehyde (**1a-ii**) was reported to act as a triplet sensitizer capable of C–H activation.^{72–75} A control experiment to probe toluene activation with benzaldehyde as catalyst under black light irradiation was carried out under air (Table 2, entry 5). This control experiment showed that the most significant reaction promoted by benzaldehyde was its oxidation to benzoic acid and benzoic anhydride (84% combined yield after 24 h). This outcome is likely due to either the excited triplet aldehyde reacting with triplet oxygen or singlet aldehyde with excited singlet oxygen via photosensitization.^{76–78} Toluene activation, under these conditions, was not detected.

After investigating the influence of the light sources, we turned to examination of the substrate scope. A variety of hydrocarbons and heteroatom-containing compounds with diverse structures were tested (Scheme 4) under blue light or purple/black light irradiation. In general, better yields and higher selectivity was observed under the dual irradiation conditions. The reaction with para-xylene (**1b**) shows that oxygenation occurred selectively at only one of the methyl groups under the irradiation of blue light. Catalytic oxidation of 2-phenoxy-1-phenylethanol (**1c**), a model for lignin,^{26, 69} generated C–C bond cleavage product phenol as the major component. Benzaldehyde and small amounts of benzoic acid were also



Scheme 4 The substrate scope of the C–H oxygenation mediated by **I** with blue vs. purple/black light irradiation. The yield with blue light (top values) is in blue font while purple/black light is purple. General reaction conditions: 0.4 mmol substrate, 0.02 mmol **I**, 4 mL CH₃CN (or CD₃CN), irradiated with 1) blue and 2) purple/black lights at 298 K under aerobic conditions for 24 h. Assay yield determined by ¹H NMR spectroscopy with 1,4-dioxane as an internal standard. See Sections VIII and XII of the Supplemental Information. [a] Reaction conditions: 1 atm CH₄ (or C₂H₆) and O₂, 0.01 mmol **I**, 4 mL CD₃CN. [b] Turnover numbers (TONs) are based on the concentration of products over the catalyst loading, [product]/[**I**].

observed. Diphenylmethane (**1d**) gave up to 87% yield of benzophenone (**1d-ii**) under purple/black light irradiation. On the other hand, triphenylmethane provided up to 51% yield of the expected alcohol (**1e-i**) under purple/black light irradiation, with further reaction resulting in C–C bond cleavage to afford benzophenone (**1d-ii**). Dihydroanthracene (**1f**) predominantly gave the dual oxidation product 9,10-anthraquinone (**1f-ii**, 83% with purple/black irradiation), which is used in the dye and papermaking industries.

Indane could be oxidized to 1-indanone (**1g-ii**) in 70% yield under purple/black light conditions. The cyclic ether 1,3-dihydroisobenzofuran underwent selective reaction to provide the corresponding lactone **1h-i** in up to 74%, with the remainder of the material further oxidized to the anhydride (**1h-ii**) and its hydrolysis product the diacid (**1h-iii**). Isochromane (**1i**) was chosen to examine the positional selectivity of the oxidation. Interestingly, the selectivity for the benzylic oxidation to the non-benzylic oxidation was ~ 4: 1 under both irradiation conditions. The low chemoselectivity in the isochromane oxidation likely reflects the

unselective nature of Cl•.^{21, 39–41} The sulfur-containing 2-methylthiophene exhibited low conversion to the expected products (**1j-i** and **1j-ii**), perhaps due to oxidation at the sulfur or heteroaromatic system. Substrates containing a nitrogen atom, such as pyrrolidine and piperidine, showed poor conversions (< 5%) under our catalytic system. The coordination of these substrates to the cerium species may result in the deactivation of the cerium catalyst. The light-mediated C–H oxygenation of cyclic alkanes (**1k–1m**), with stronger C–H bonds [BDE (kcal/mol): 95.6 (**1k**), 99.5 (**1l**), 95.7 (**1m**)],^{54, 79, 80} was investigated. The corresponding cyclic ketones were the major products with blue light. The purple/black light combination provided a higher overall yield (i.e., the yield of alcohol + ketone).

The light-mediated activation of light alkanes represents an approach to transforming hydrocarbons, often flared, into valuable chemicals. Under 1 atm total of a mixture of O₂ and methane or ethane, the conversion of methane to formic acid and ethane to ethanol, acetaldehyde, and acetic acid was achieved (Scheme 4, Figure 2 and Section VII of the Supplemental Information for details).





Fig. 2 ^1H NMR spectra of light-mediated C–H oxygenation of (a) CH_4 and (b) C_2H_6 . Reaction conditions: 1 atm CH_4 (or C_2H_6) and O_2 , 0.01 mmol **1**, 4 mL CD_3CN , irradiated with a Kessil A160WE blue LED lamp at 298 K for 12 h. The quantity of oxidation products was determined using 1,4-dioxane ($\delta = 3.60$ ppm) as an internal standard.

In the case of methane, 4 turn-overs were observed (Figure 2). The limited yield is attributed to the low solubility of methane in acetonitrile at low pressures. The selectivity in the ethane oxygenation was also influenced by light sources: ethanol was observed under blue light irradiation while acetaldehyde and acetic acid were observed under all conditions (Scheme 4, **2b**). An overall 20 turn-overs was observed in ethane activation.

Based on the experiments above and literature precedent, a reaction mechanism for the C–H oxygenation is proposed in Scheme 5. Upon irradiation, $[\text{NEt}_4]_2[\text{Ce}^{\text{IV}}\text{Cl}_6]$ undergoes light-induced LMCT and generates Cl^\bullet , which is proposed to be the turnover-limiting step. The Cl^\bullet undergoes HAT with the substrate C–H bond, producing a carbon-centered radical and hydrogen chloride (HCl). The generated hydrocarbon radical is trapped by O_2 to afford the organic superoxide intermediate. Subsequent electron and proton transfer is followed by O–O bond cleavage resulting in oxidized products. It is noteworthy that radical-chain processes are excluded from the proposed mechanism because the quantum yield of the C–H oxygenation ($\Phi = 1.3\%$, Section XI of the Supplemental Information) is smaller than that of the photo-induced generation of Cl^\bullet from $[\text{NEt}_4]_2[\text{Ce}^{\text{IV}}\text{Cl}_6]$ (**1**) ($\Phi = 1.9\%$, section XII of the Supplemental Information). The key findings of this work are: oxygen can work as a trapping reagent for different substrates; oxygen is able to regenerate the cerium catalyst in catalytic conditions; black light is involved in the reduction of oxygenated intermediates, presumably by light absorption of the peroxide intermediates or the excitation of $[\text{NEt}_4]_3[\text{Ce}^{\text{III}}\text{Cl}_6]$ to generate a stronger reductant $[\text{NEt}_4]_3[\text{Ce}^{\text{III}}\text{Cl}_6]^*$. Support for this hypothesis is the change in product distribution in the presence of the dual purple/black light and related control experiments.

Conclusion

In summary, the light-mediated oxygenation of benzylic and alkane $\text{C}(\text{sp}^3)\text{--H}$ bonds is accessible using catalytic $[\text{NEt}_4]_2[\text{Ce}^{\text{IV}}\text{Cl}_6]$ under air. A key finding of this work is that the light impacts two reaction stages driven by the catalyst in both reductive and oxidative quenching steps. The selectivity toward more oxidized products increases under irradiation with a purple/black combination of light, which is



Scheme 5 Proposed catalytic mechanism of C–H oxygenation mediated by $[\text{NEt}_4]_2[\text{Ce}^{\text{IV}}\text{Cl}_6]$.

rationalized by the activation of peroxide species under light irradiation. The excited state $[\text{NEt}_4]_3[\text{Ce}^{\text{III}}\text{Cl}_6]^*$ (**III**) was proposed to play a role in the activation of the peroxide. The oxidation of benzylic C–H bonds proceeds in good yields. The system also shows reactivity toward cycloalkanes and promising hints of reactivity toward challenging light alkanes (i.e., CH_4 and C_2H_6) under mild conditions. The influence of radiation wavelengths on the yield and selectivity of C–H oxygenation in this study is expected to prove strategic in optimizing metal-catalyzed light-mediated $\text{C}(\text{sp}^3)\text{--H}$ oxidation systems.

Author Contributions

YHW and QMY optimized the reaction condition, YHW performed substrates screening and characterization, YHW and QMY performed mechanistic studies and drafted the manuscript. EJS and PWJ supervised all aspects.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Dr. Yusen Qiao for the helpful discussions. The project was supported by Vagelos Institute for Energy Science and Technology (VIEST), including support for Y.H.W.; Q.Y. was supported as part of the Center for Actinide Science and Technology (CAST), an Energy Frontier Research Center (EFRC) funded by the U.S. Department of Energy (DOE), Office of Science, Basic Energy Sciences (BES), under Award Number DE-SC0016568. We also thank the University of Pennsylvania for support of this work. The NSF Major Research Instrumentation Program (award NSF CHE-1827457), the NIH supplements awards 3R01GM118510-03S1 and 3R01GM087605-

06S1, and Vagelos Institute for Energy Science and Technology supported the purchase of the NMRs used in this study.

Notes and references

1. F. Cavani and J. H. Teles, Sustainability in Catalytic Oxidation: An Alternative Approach or a Structural Evolution?, *ChemSusChem*, 2009, **2**, 508-534.
2. I. Hermans, E. S. Spier, U. Neuenschwander, N. Turrà and A. Baiker, Selective Oxidation Catalysis: Opportunities and Challenges, *Top. Catal.*, 2009, **52**, 1162-1174.
3. F. Cavani, Catalytic selective oxidation: The forefront in the challenge for a more sustainable chemical industry, *Catal. Today*, 2010, **157**, 8-15.
4. P. D. Ray and R. C. Fry, in *Systems Biology in Toxicology and Environmental Health*, ed. R. C. Fry, Academic Press, Boston, 2015, DOI: <https://doi.org/10.1016/B978-0-12-801564-3.00002-X>, pp. 11-42.
5. Y. Ishii, S. Sakaguchi and T. Iwahama, Innovation of Hydrocarbon Oxidation with Molecular Oxygen and Related Reactions, *Adv. Synth. Catal.*, 2001, **343**, 393-427.
6. L. Que and W. B. Tolman, Biologically inspired oxidation catalysis, *Nature*, 2008, **455**, 333-340.
7. Z. Guo, B. Liu, Q. Zhang, W. Deng, Y. Wang and Y. Yang, Recent advances in heterogeneous selective oxidation catalysis for sustainable chemistry, *Chem. Soc. Rev.*, 2014, **43**, 3480-3524.
8. S. Valange and J. C. Védrine, General and prospective views on oxidation reactions in heterogeneous catalysis, *Catalysts*, 2018, **8**, 483.
9. L. V. Desai, K. L. Hull and M. S. Sanford, Palladium-Catalyzed Oxygenation of Unactivated sp³ C–H Bonds, *J. Am. Chem. Soc.*, 2004, **126**, 9542-9543.
10. A. E. Shilov and G. B. Shul'pin, Activation of C–H Bonds by Metal Complexes, *Chem. Rev.*, 1997, **97**, 2879-2932.
11. Y.-F. Liang and N. Jiao, Oxygenation via C–H/C–C Bond Activation with Molecular Oxygen, *Acc. Chem. Res.*, 2017, **50**, 1640-1653.
12. D. Munz and T. Strassner, Alkane C–H Functionalization and Oxidation with Molecular Oxygen, *Inorg. Chem.*, 2015, **54**, 5043-5052.
13. L. Boisvert and K. I. Goldberg, Reactions of Late Transition Metal Complexes with Molecular Oxygen, *Acc. Chem. Res.*, 2012, **45**, 899-910.
14. J. A. Labinger, Selective alkane oxidation: hot and cold approaches to a hot problem, *J. Mol. Catal. A: Chem.*, 2004, **220**, 27-35.
15. P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz and L. Ackermann, 3d Transition Metals for C–H Activation, *Chem. Rev.*, 2019, **119**, 2192-2452.
16. J. Das, S. Guin and D. Maiti, Diverse strategies for transition metal catalyzed distal C(sp³)–H functionalizations, *Chem. Sci.*, 2020, **11**, 10887-10909.
17. H. Sterckx, B. Morel and B. U. W. Maes, Catalytic Aerobic Oxidation of C(sp³)–H Bonds, *Angew. Chem. Int. Ed.*, 2019, **58**, 7946-7970.
18. G. Urgoitia, R. SanMartin, M. T. Herrero and E. Domínguez, Recent Advances in Homogeneous Metal-Catalyzed Aerobic C–H Oxidation of Benzylic Compounds, *Catalysts*, 2018, **8**, 640.
19. D. Wang, A. B. Weinstein, P. B. White and S. S. Stahl, Ligand-Promoted Palladium-Catalyzed Aerobic Oxidation Reactions, *Chem. Rev.*, 2018, **118**, 2636-2679.
20. J. Lloret, L. Gómez and M. Costas, in *Alkane CH Activation by Single-Site Metal Catalysis*, Springer, 2012, pp. 143-228.
21. S. Fukuzumi and K. Ohkubo, Selective photocatalytic reactions with organic photocatalysts, *Chem. Sci.*, 2013, **4**, 561-574.
22. X. Zhang, K. Rakesh, L. Ravindar and H.-L. Qin, Visible-light initiated aerobic oxidations: a critical review, *Green Chem.*, 2018, **20**, 4790-4833.
23. G. Laudadio, S. Govaerts, Y. Wang, D. Ravelli, H. F. Koolman, M. Fagnoni, S. W. Djuric and T. Noël, Selective C(sp³)–H aerobic oxidation enabled by decatungstate photocatalysis in flow, *Angew. Chem.*, 2018, **130**, 4142-4146.
24. S. Fukuzumi, Y. M. Lee and W. Nam, Photocatalytic oxygenation reactions using water and dioxygen, *ChemSusChem*, 2019, **12**, 3931-3940.
25. C. Bian, A. K. Singh, L. Niu, H. Yi and A. Lei, Visible-Light-Mediated Oxygenation Reactions using Molecular Oxygen, *Asian J. Org. Chem.*, 2017, **6**, 386-396.
26. P. L. Arnold, J. M. Purkis, R. Rutkauskaitė, D. Kovacs, J. B. Love and J. Austin, Controlled Photocatalytic Hydrocarbon Oxidation by Uranyl Complexes, *ChemCatChem*, 2019, **11**, 3786-3790.
27. S. Li, B. Zhu, R. Lee, B. Qiao and Z. Jiang, Visible light-induced selective aerobic oxidative transposition of vinyl halides using a tetrahalogenoferrate (III) complex catalyst, *Org. Chem. Front.*, 2018, **5**, 380-385.
28. M. Wang, J. Wen, Y. Huang and P. Hu, Selective Degradation of Styrene-Related Plastics Catalyzed by Iron under Visible Light**, *ChemSusChem*, 2021, **14**, 5049-5056.
29. G. Zhang, Z. Zhang and R. Zeng, Photoinduced FeCl₃-Catalyzed Alkyl Aromatics Oxidation toward Degradation of Polystyrene at Room Temperature†, *Chin. J. Chem.*, 2021, **39**, 3225-3230.
30. C.-C. Wang, G.-X. Zhang, Z.-W. Zuo, R. Zeng, D.-D. Zhai, F. Liu and Z.-J. Shi, Photo-induced deep aerobic oxidation of alkyl aromatics, *Sci. China Chem.*, 2021, **64**, 1487-1492.
31. S.-i. Hirashima and A. Itoh, Aerobic visible light-oxidation of aromatic methyl groups to carboxylic acids, *Photochem. Photobiol. Sci.*, 2007, **6**, 521-524.
32. N. Tada, K. Hattori, T. Nobuta, T. Miura and A. Itoh, Facile aerobic photooxidation of methyl group in the aromatic nucleus in the presence of an organocatalyst under VIS irradiation, *Green Chem.*, 2011, **13**, 1669-1671.
33. B. Mühldorf and R. Wolf, Photocatalytic benzylic C–H bond oxidation with a flavin scandium complex, *Chem. Commun.*, 2015, **51**, 8425-8428.
34. K. A. Korvinson, G. N. Hargenrader, J. Stevanovic, Y. Xie, J. Joseph, V. Maslak, C. M. Hadad and K. D. Glusac, Improved Flavin-Based Catalytic Photooxidation of Alcohols through Intersystem Crossing Rate Enhancement, *J. Phys. Chem. A*, 2016, **120**, 7294-7300.
35. P. Dongare, I. MacKenzie, D. Wang, D. A. Nicewicz and T. J. Meyer, Oxidation of alkyl benzenes by a flavin photooxidation catalyst on nanostructured metal-oxide films, *PNAS*, 2017, **114**, 9279-9283.
36. J. Zelenka, E. Svobodová, J. Tarábek, I. Hoskovcová, V. Boguschová, S. Bailly, M. Sikorski, J. Roithová and R. Cibulka, Combining Flavin Photocatalysis and

- Organocatalysis: Metal-Free Aerobic Oxidation of Unactivated Benzylic Substrates, *Org. Lett.*, 2019, **21**, 114-119.
37. A. H. Tolba, F. Vávra, J. Chudoba and R. Cibulka, Tuning Flavin-Based Photocatalytic Systems for Application in the Mild Chemoselective Aerobic Oxidation of Benzylic Substrates, *Eur. J. Org. Chem.*, 2020, **2020**, 1579-1585.
 38. V. Srivastava, P. K. Singh, A. Srivastava and P. P. Singh, Synthetic applications of flavin photocatalysis: a review, *RSC Adv.*, 2021, **11**, 14251-14259.
 39. K. Ohkubo, A. Fujimoto and S. Fukuzumi, Metal-free oxygenation of cyclohexane with oxygen catalyzed by 9-mesityl-10-methylacridinium and hydrogen chloride under visible light irradiation, *Chem. Commun.*, 2011, **47**, 8515-8517.
 40. W. Wu, Z. Fu, X. Wen, Y. Wang, S. Zou, Y. Meng, Y. Liu, S. R. Kirk and D. Yin, Light-triggered oxy-chlorination of cyclohexane by metal chlorides, *Appl. Catal. A: Gen.*, 2014, **469**, 483-489.
 41. L. L. Costanzo, S. Pistarà and G. Condorelli, Photoreduction of cerium(IV) hexachloride in acetonitrile, *Journal of Photochemistry*, 1983, **21**, 45-51.
 42. R. Yuan, S. Fan, H. Zhou, Z. Ding, S. Lin, Z. Li, Z. Zhang, C. Xu, L. Wu, X. Wang and X. Fu, Chlorine-Radical-Mediated Photocatalytic Activation of C-H Bonds with Visible Light, *Angew. Chem. Int. Ed.*, 2013, **52**, 1035-1039.
 43. H.-P. Deng, Q. Zhou and J. Wu, Microtubing-Reactor-Assisted Aliphatic C-H Functionalization with HCl as a Hydrogen-Atom-Transfer Catalyst Precursor in Conjunction with an Organic Photoredox Catalyst, *Angew. Chem. Int. Ed.*, 2018, **57**, 12661-12665.
 44. K. Ohkubo and K. Hirose, Light-Driven C-H Oxygenation of Methane into Methanol and Formic Acid by Molecular Oxygen Using a Perfluorinated Solvent, *Angew. Chem. Int. Ed.*, 2018, **57**, 2126-2129.
 45. S. Rohe, A. O. Morris, T. McCallum and L. Barriault, Hydrogen Atom Transfer Reactions via Photoredox Catalyzed Chlorine Atom Generation, *Angew. Chem. Int. Ed.*, 2018, **57**, 15664-15669.
 46. D. Leifert and A. Studer, The Persistent Radical Effect in Organic Synthesis, *Angew. Chem. Int. Ed.*, 2020, **59**, 74-108.
 47. M. Zidan, A. O. Morris, T. McCallum and L. Barriault, The Alkylation and Reduction of Heteroarenes with Alcohols Using Photoredox Catalyzed Hydrogen Atom Transfer via Chlorine Atom Generation, *Eur. J. Org. Chem.*, 2020, **2020**, 1453-1458.
 48. S. K. Kariofillis and A. G. Doyle, Synthetic and Mechanistic Implications of Chlorine Photoelimination in Nickel/Photoredox C(sp³)-H Cross-Coupling, *Acc. Chem. Res.*, 2021, **54**, 988-1000.
 49. G. S. Yedase, S. Kumar, J. Stahl, B. König and V. R. Yatham, Cerium-photocatalyzed aerobic oxidation of benzylic alcohols to aldehydes and ketones, *Beilstein J. Org. Chem.*, 2021, **17**, 1727-1732.
 50. Q. Yang, Y.-H. Wang, Y. Qiao, M. Gau, P. J. Carroll, P. J. Walsh and E. J. Schelter, Photocatalytic C-H activation and the subtle role of chlorine radical complexation in reactivity, *Science*, 2021, **372**, 847-852.
 51. L. Vivier and D. Duprez, Ceria - based solid catalysts for organic chemistry, *ChemSusChem*, 2010, **3**, 654-678.
 52. T. Montini, M. Melchionna, M. Monai and P. Fornasiero, Fundamentals and catalytic applications of CeO₂-based materials, *Chem. Rev.*, 2016, **116**, 5987-6041.
 53. H. Yin, Y. Jin, J. E. Hertzog, K. C. Mullane, P. J. Carroll, B. C. Manor, J. M. Anna and E. J. Schelter, The Hexachloroacetate(III) Anion: A Potent, Benchtop Stable, and Readily Available Ultraviolet A Photosensitizer for Aryl Chlorides, *J. Am. Chem. Soc.*, 2016, **138**, 16266-16273.
 54. Y.-R. Luo, *Comprehensive handbook of chemical bond energies*, CRC press, 2007.
 55. Y. Qiao, Q. Yang and E. J. Schelter, Photoinduced Miyaura Borylation by a Rare-Earth-Metal Photoreductant: The Hexachloroacetate(III) Anion, *Angew. Chem. Int. Ed.*, 2018, **57**, 10999-11003.
 56. E. M. Simmons and J. F. Hartwig, On the Interpretation of Deuterium Kinetic Isotope Effects in C H Bond Functionalizations by Transition-Metal Complexes, *Angew. Chem. Int. Ed.*, 2012, **51**, 3066-3072.
 57. M. Gómez-Gallego and M. A. Sierra, Kinetic Isotope Effects in the Study of Organometallic Reaction Mechanisms, *Chem. Rev.*, 2011, **111**, 4857-4963.
 58. Photochemical generation of Cl• from via ligand-to-metal charge transfer (LMCT) has been investigated by our group previously, see ref. [36] and [48].
 59. The product ratio of the toluene oxygenation under blue light irradiation is close to 1 (i.e., 1a-iii to 1a-ii, entries 1-3 in Table 1).
 60. A. Enesca and L. Isac, The Influence of Light Irradiation on the Photocatalytic Degradation of Organic Pollutants, *Materials*, 2020, **13**, 2494.
 61. N. Luo, M. Wang, H. Li, J. Zhang, H. Liu and F. Wang, Photocatalytic Oxidation-Hydrogenolysis of Lignin β-O-4 Models via a Dual Light Wavelength Switching Strategy, *ACS Catal.*, 2016, **6**, 7716-7721.
 62. P. Montes-Navajas, M. Serra and H. Garcia, Influence of the irradiation wavelength on the photocatalytic activity of Au-Pt nanoalloys supported on TiO₂ for hydrogen generation from water, *Catalysis Science & Technology*, 2013, **3**, 2252-2258.
 63. C. Gomes Silva, R. Juárez, T. Marino, R. Molinari and H. García, Influence of Excitation Wavelength (UV or Visible Light) on the Photocatalytic Activity of Titania Containing Gold Nanoparticles for the Generation of Hydrogen or Oxygen from Water, *J. Am. Chem. Soc.*, 2011, **133**, 595-602.
 64. G. Li Puma and P. L. Yue, Effect of the Radiation Wavelength on the Rate of Photocatalytic Oxidation of Organic Pollutants, *Ind. Eng. Chem. Res.*, 2002, **41**, 5594-5600.
 65. For the emission spectrum of Kessil A160WE blue LEDs, see: C. C. Le, M. K. Wismer, Z.-C. Shi, R. Zhang, D. V. Conway, G. Li, P. Vachal, I. W. Davies, D. W. C. MacMillan, *ACS Cent. Sci.* 2017, **3**, 647-653. For other Kessil lamp, see website: <https://kessil.com/science/PR160L.php>
 66. I. Ghosh and B. König, Chromoselective Photocatalysis: Controlled Bond Activation through Light-Color Regulation of Redox Potentials, *Angew. Chem. Int. Ed.*, 2016, **55**, 7676-7679.
 67. A. M. Martínez-Gualda, R. Cano, L. Marzo, R. Pérez-Ruiz, J. Luis-Barrera, R. Mas-Ballesté, A. Fraile, V. A. de la Peña O'Shea and J. Alemán, Chromoselective access to Z- or E-allylated amines and heterocycles by a photocatalytic allylation reaction, *Nat. Commun.*, 2019, **10**, 2634.

68. L. Schmermund, S. Reischauer, S. Bierbaumer, C. K. Winkler, A. Diaz-Rodriguez, L. J. Edwards, S. Kara, T. Mielke, J. Cartwright, G. Grogan, B. Pieber and W. Kroutil, Chromoselective Photocatalysis Enables Stereocomplementary Biocatalytic Pathways**, *Angew. Chem. Int. Ed.*, 2021, **60**, 6965-6969.
69. T. S. Symeonidis, I. Tamiolakis, G. S. Armatas and I. N. Lykakis, Green photocatalytic organic transformations by polyoxometalates vs. mesoporous TiO₂ nanoparticles: selective aerobic oxidation of alcohols, *Photochem. Photobiol. Sci.*, 2015, **14**, 563-568.
70. R. Sapach and T. Viraraghavan, An introduction to the use of hydrogen peroxide and ultraviolet radiation: An advanced oxidation process, *Journal of Environmental Science and Health . Part A: Environmental Science and Engineering and Toxicology*, 1997, **32**, 2355-2366.
71. The influence of radiation wavelengths on the O–O cleavage of the alkyl hydroperoxide presumably leads to the different selectivity (see Section X of the Supplemental Information).
72. C. Walling and M. J. Gibian, Hydrogen Abstraction Reactions by the Triplet States of Ketones¹, *J. Am. Chem. Soc.*, 1965, **87**, 3361-3364.
73. J. A. Dantas, J. T. M. Correia, M. W. Paixão and A. G. Corrêa, Photochemistry of Carbonyl Compounds: Application in Metal-Free Reactions, *ChemPhotoChem*, 2019, **3**, 506-520.
74. F. Strieth-Kalthoff and F. Glorius, Triplet Energy Transfer Photocatalysis: Unlocking the Next Level, *Chem*, 2020, **6**, 1888-1903.
75. M. A. Theodoropoulou, N. F. Nikitas and C. G. Kokotos, Aldehydes as powerful initiators for photochemical transformations, *Beilstein J. Org. Chem.*, 2020, **16**, 833-857.
76. Z. E. Hamami, L. Vanoye, P. Fongarland, C. de Bellefon and A. Favre-Reguillon, Metal-free, visible light-promoted aerobic aldehydes oxidation, *J. Flow Chem.*, 2016, **6**, 206-210.
77. C. Grewer and H.-D. Brauer, Mechanism of the Triplet-State Quenching by Molecular Oxygen in Solution, *The Journal of Physical Chemistry*, 1994, **98**, 4230-4235.
78. J. C. Andre, M. Bouchy and M. Niclaude, Initiation of aldehyde photo-oxidation in the liquid phase, *Journal of Photochemistry*, 1976, **5**, 1-12.
79. G. K. Cook and J. M. Mayer, C-H Bond Activation by Metal Oxo Species: Chromyl Chloride Oxidations of Cyclooctane, Isobutane, and Toluene, *J. Am. Chem. Soc.*, 1995, **117**, 7139-7156.
80. X.-S. Xue, P. Ji, B. Zhou and J.-P. Cheng, The Essential Role of Bond Energetics in C–H Activation/Functionalization, *Chem. Rev.*, 2017, **117**, 8622-8648.