

Arctic Rain on Snow Events: Bridging Observations to Understand Environmental and Livelihood Impacts

Mark C. Serreze, Julia Gustafson, Andrew P. Barrett, Matthew L. Druckenmiller,
Shari Fox, Jessica Voveris, Julianne Stroeve, Betsy Sheffield
National Snow and Ice Data Center, Cooperative Institute for Research in Environmental Sciences,
University of Colorado Boulder, Boulder Colorado

Bruce C. Forbes, Sirpa Rasmus, Roza Laptander
Arctic Centre, University of Lapland, Rovaniemi, Finland

Mike Brook, Mike Brubaker, James Temte
Center for Climate and Health, Alaska Pacific University, Anchorage Alaska

Michelle R. McCrystall
University of Manitoba, Winnipeg, Canada

Annett Bartsch
b.geos, Korneuburg, Austria
Austrian Polar Research Institute, Vienna, Austria

Abstract

When rain falls on an existing cover of snow, followed by low temperatures, or falls as freezing rain, it can leave a hard crust. These Arctic rain on snow (ROS) events can profoundly influence the environment and in turn, human livelihoods. Impacts can be immediate (e.g. on human travel, herding, or harvesting) or evolve or accumulate, leading to massive starvation-induced die-offs of reindeer, caribou, and musk oxen, for example. As part of the international Arctic Rain on Snow Study (AROSS) and the HORIZON2020 project CHARTER, we provide here a review and synthesis of Arctic ROS events and their impacts, addressing human-environment relationships, meteorological conditions associated with ROS events, and challenges in their detection. We offer a path to anticipate and mitigate impacts through knowledge co-production. Systematic detection of ROS events, their intensity, and trends across the Arctic region can be approached by combining data from satellite remote sensing, atmospheric reanalyses, and meteorological station records. However, obtaining knowledge and information most germane to impacts, such as the thickness of ice layers, how ice layers form within a snowpack, and antecedent conditions that can amplify impacts, necessitates collaboration and knowledge co-production with community members and Indigenous knowledge-holders.

1. Introduction

There is increasing recognition that Arctic rain on snow (ROS) events, which are projected to become more frequent as the Arctic warms (Rennert et al. 2009, Bintanja and Andry 2017), can have significant and sometimes even catastrophic impacts on the physical and living environments of the North (**Figure 1**). ROS events are generally associated with short-lived (hours to days) warm spells in autumn and winter, linked to extratropical cyclones that generate rain, followed by a rapid drop in temperature (Hansen et al. 2014). The rainwater may freeze on the snow surface or percolate through the snowpack and pool at the ground surface. This depends upon the thickness and density of the snowpack, and the intensity of the precipitation event. After refreezing, ice crusts can variously form on the snow surface, as layers within the snow, or at the base of the snowpack.




Figure 1. Some impacts of Arctic rain on snow events over Arctic lands and ocean.

ROS events can foster slush avalanches when the rain reduces cohesiveness and destabilizes the snow (Hansen et al. 2014). Alternatively, formation of an ice layer can weaken the snow pack through growth of large faceted grains, providing a hard surface upon which a slab avalanche can slide (Jamieson 2006). ROS events influence snow structure, density and conductivity, and hence the ground thermal regime (e.g. Mazurkiewicz et al. 2008, Freudiger et al. 2014, Guan et al. 2016, Romanovsky et al. 2010, Westermann et al. 2011). If enough water pools at the ground surface and freezes, latent heat release alters seasonally frozen ground, increasing the depth of the permafrost active layer and warming underlying permafrost. Severe rain events can remove the insulating ability of the snowpack (Putkonen and Roe 2003, Rennert et al. 2009). Impacts have been documented on vegetation (e.g. stress due to ice encasement) and soil organisms (e.g. through altered temperature and drainage conditions) (Bjerke et al. 2015). ROS events can influence vegetation greenness as detectable with satellite data (Bjerke et al. 2017) as well as CO₂ fluxes (Treharne et al. 2020).

On land, ROS events influence population dynamics of lemmings (Kausrud et al. 2008), voles (Stein et al. 2012), and bird species that seek shelter in the snowpack, including owls, ptarmigan, and grouse (Myserud 2016). Notably, icing can foster fading of lemming cycles, causing crashes of Arctic predators such as snowy owls and Arctic fox, which depend on them (Sokolov et al. 2016). In marine environments, polar bears (*Ursus maritimus*) and ringed seals (*Phoca hispida*) are also impacted – rains

early in the breeding season can melt subnivean lairs and increase cub mortality (Stirling and Smith 2003).

Perhaps the most devastating impact of ROS events, however, is that ice layers, in creating barriers that prohibit foraging (**Figure 2**), can lead to massive die-offs of large herbivores, with concomitant social-economic impacts, especially in regions where humans depend on either wild or semi-domesticated populations of *Rangifer* species (caribou and reindeer) for food, transportation, and other needs. Such impacts can evolve and cascade, presenting threats to shared food systems, cultural vitality, and other issues identified by communities (Egeland et al. 2013, Huntington et al. 2019).




Figure 2. Ice layers within the snowpack associated with two separate ROS events. Photo: Florian Stammer.

Based on analysis of radar backscatter data (Ku-band QuikScat records) from 2000-2009 (Bartsch 2010a, b) and in general agreement with patterns determined from atmospheric reanalysis data (Rennert et al. 2009, Liston and Hiemstra 2011) (see Sections 4.4 and 4.5), ROS events are most frequent over northern Europe and southern Alaska. They occasionally occur over western Siberia, southern Greenland, and parts of Canada but are rare in the cold Canadian Arctic Archipelago (classified as polar desert). Events that do occur here have tended to happen fairly early in the season when it is warmer. It seems likely,

however, that as the Arctic continues to warm, ROS events will become more frequent and spread into more northerly regions.




Figure 3: Potential ROS events per year for November through February of the winters 2000/1–2008/9 based on midwinter daily average backscatter increases of more than 1.5 dB as detected with Ku-band QuikSCAT; no masking applied for lakes (Bartsch 2010a, b, Bartsch et al. 2010, Freund and Bartsch 2020). Also shown are the locations for three ROS case studies: 1- Yamal Peninsula, Russia, 8-10 November 2013; 2- Svalbard, Norway 30-31 January 2012; 3- Banks Island, Canada, 3-8 October 2003.

We provide here a review and synthesis of our current knowledge of Arctic ROS events, emphasizing impacts on reindeer herding livelihoods and wildlife harvesting. We offer a path forward for anticipating and mitigating impacts of these events through knowledge co-production with Indigenous researchers, hunters, harvesters, and other Arctic residents whose knowledge, lived experiences, and detailed

observations of changes in climate and weather are needed for understanding and addressing environmental impacts (e.g. Krupnik and Jolly 2002, ACIA 2005, Gearheard et al. 2010, Pearce et al. 2015) and ensuring that research on the physical aspects of ROS events and anticipated changes are relevant and practical for community use.

The paper starts with an overview of observed impacts, followed by case studies of the meteorology and impacts of three notable events. Attention then turns to data sources and approaches to detect ROS events and their intensity across the Arctic. We then discuss the importance of knowledge co-production for its role in improving event detection from remote sensing and other means, and for targeting research on key aspects of ROS events such as the thickness of ice layers and how they form within a snowpack, as well as on other icing events, including thaw-freeze, or freezing of very wet soil surface/vegetation layers. Our paper represents a contribution to the Arctic Rain on Snow Study (AROSS), a project within the National Science Foundation's Navigating the New Arctic (NNA) initiative seeking to better understand the distribution, severity, and changes in the frequency of ROS and melt-refreeze events in the Arctic and their impacts. It also contributes to CHARTER by comparing the impacts of well-documented events on contrasting social-ecological conditions (e.g. semi-domesticated versus wild ungulates) in Arctic Eurasia and North America.

2. ROS Impacts on Human-Environment Relationships

2.1 Reindeer and Reindeer Herding Communities

Reindeer and caribou are the same species, though reindeer have some morphological differences due to semi-domestication. A mature male reindeer can grow to about 100 kg. Females typically give birth in May and June to a single calf. Reindeer herding is practiced across the circumpolar Arctic, but most activity is in Fennoscandia and parts of Russia. Herding is conducted by families, within herding districts, villages, and state-owned collectives. Reindeer are used for food, clothing, as well as transportation (Forbes and Kumpula 2009). Finland has 54 herding districts and approximately 4,500 reindeer owners, of which an estimated 900 are full-time herders (RHA 2020). Reindeer husbandry is practiced across northern Russia, by Sámi herders on the Kola Peninsula to the Taimyr reindeer herd of northcentral Siberia (the largest in the world of at least a half million animals) to the Chukchi herders in northeastern Siberia. In North America, there are also roughly 10,000 reindeer managed on and around Alaska's Seward Peninsula by approximately 20 herders, who belong to the Kawerak Reindeer Herders Association—a tribal organization that assists in developing of a viable reindeer industry for rural Alaska communities (Kawerak Reindeer Herders Association 2020). There are also many thousands of reindeer throughout Alaska's Nunivak Island, St. Lawrence Island, the Pribilof Islands, and parts of the Aleutian Islands. Reindeer were originally introduced to Alaska in the late 1800s and early 1900s by government officials who worked first with Chukchi and then Sámi herders to introduce reindeer husbandry as a strategy to supplement the food supply of Iñupiaq communities (Andrews 1939).

2.2 ROS Impacts on Reindeer Herding

During spring and summer, reindeer graze on graminoids, herbs and shrubs; in autumn, mushrooms are favored. Once winter sets in, the reindeer diet is largely limited to lichens, typically accessed by digging through the snow. However, in boreal forest regions, arboreal lichen found in old-growth forests becomes important.

The cold season is critical for managed reindeer herds, which consist mainly of pregnant females with high energy demands. Icing for a relatively short period towards the spring melt is part of a typical evolution of the snow cover in Northern Fennoscandia and can be managed by searching for less-affected pastures with softer snow. Herds can also disperse to find more easily available forage, but this necessitates effort and resources to control the animals. If snow is deep or icy, foraging through it in search of lichens consumes considerable energy (Helle 1984). However, especially in the boreal forest zone, if old-growth forests are available, icy snow may actually ease reindeer grazing on arboreal lichen, which often falls in clumps from branches and remains at the surface. Reindeer can also move from tree to tree to reach lichen growing on higher branches.

The situation changes if thick and extensive ice crusts are formed. Impacts depend on many factors, including the timing and intensity of icing events, snowpack characteristics, topography, vegetation, climate, as well as herding culture, traditions, and practices. ROS and other icing events are historically known to cause strong declines in reindeer populations in Fennoscandia (Riseth et al. 2016, Eira et al. 2018). Ice barriers, especially in early winter, can force animals to expand their grazing range, depleting body fat and protein reserves as they travel in search of soft snow. Starvation can occur quickly, or drag on for many months and even longer (Bartsch et al. 2010, Forbes et al. 2016). Spring weight, calf production, calf size, and adult survivability are all lower when reindeer are unable to satisfy their energy needs. Cumulative energy loss can lead to late winter/early spring abortion of calves and low calf survival rates in late spring and early summer (Tveraa et al. 2003, Helle and Kojola 2008). On the Seward Peninsula, weather related stressors, including ROS events, can be one of multiple drivers leading caribou to expand their range onto the peninsula, causing competition with other reindeer for food, often resulting in reindeer being pulled away from their herd (Project Jukebox 2001).

Impacts can be mitigated by supplementary feeding, but this is time consuming and costly (Turunen et al. 2016, Rasmus et al. 2020). Population crashes have sometimes occurred even with supplementary feeding (for example, arboreal lichen supplied by felling old-growth trees, cf. Berg et al. 2011a, b) or when reindeer were moved to alternative pastures (Helle and Jaakkola 2008). Die-offs have also been triggered by difficult conditions of previous seasons, causing reindeer to be in poor condition before the winter. If ROS events occur too frequently, herders may not have enough time to recover the size, demography, and health of their herds (Stammler and Ivanova 2020), illustrating the cumulative impact of ROS events beyond individual episodes or seasons to across annual cycles.

Another challenge is that in a globalizing North, reindeer husbandry shares the same operational space with other land uses, leading to fragmentation, deterioration, and shrinking of pastures (Landauer et al.

submitted). When other land use and infrastructure hinders the mobility and flexibility in pasture use, supplementary winter feeding occurs either in the pasture areas or in enclosures near the settlements, increasing working hours and expenses (Turunen et al. 2016, Turunen and Vuojala-Magga 2014) and ultimately reducing the adaptive capacity of herders to climate change.

2.3 Beyond Reindeer

ROS impacts on human-environment relationships extend far beyond challenges related to reindeer herding. Mass starvation of reindeer, caribou, and musk oxen can lead to trophic cascades as predators and scavengers (e.g. red foxes and corvids) move north to exploit abundant carrion. These predators in turn have a negative impact on medium-sized game species such as ptarmigan and hare, which are also valuable food resources for local people (Sokolov et al. 2016). Turning to the marine environment, Furgal et. al (2002), in documenting details of seal hunting at Arctic Bay, Nunavut, note how unusual weather conditions such as ROS events can affect local ringed seal populations. A local hunter described how a February rain created a thick ice cover over seal breathing holes, leading to fewer seal harvests. Cree hunters in the eastern James Bay region observed that increased autumn and winter precipitation had weakened inland lake ice through changing its composition (e.g. less black ice, more white ice, lack of columnar ice crystals) (Royer et al. 2013). The variation and instability of the ice conditions created hazards for Cree who use frozen rivers and lakes for hunting, fishing, trapping, and travel. Communities in the region reported an increase in ice travel-related accidents among both novice and experienced hunters (Royer et al. 2013).

Rain events during freeze-up and early winter can also lead to break-up of river ice, delaying use of rivers as travel routes via snow machines. While this is caused by a sudden increase in river volume, rather than icing from rain, such events add to the cumulative effects of ROS events on Arctic communities. As one more example, climate extremes in the winter of 2010-2011, including late winter rainfall, decreased sea ice coverage, and high temperatures affected the traditional food system in Iqaluit, Nunavut (Statham et al. 2015). Caribou moving further inland for forage after vegetation iced over, together with the shortened sea ice hunting season, posed a key threat to food security, exacerbated by existing socioeconomic conditions in this coastal community (Statham et al. 2015).

3. Some Notable ROS Events

3.1 Yamal Peninsula, Russia, 8-10 November, 2013

In early November 2013, an ROS event affected most of the southern Yamal Peninsula (Forbes et al. 2016), as well as coastal portions of neighboring regions (Sokolov et al. 2016, Staalesen 2016). Following 24 hours of rain, air temperatures rapidly dropped and remained below freezing throughout autumn and winter. While based on only eight years of data, analysis of radar backscatter data (see Section 4.4) suggests that this region tends to see approximately one ROS event per year (**Figure 3, Region 1**). The salient meteorological feature linked to the event was a low-pressure system moving from the northern North Atlantic into the Yamal Peninsula on November 8. Based on data from the ERA5 atmospheric

reanalysis (Hersbach et al. 2020), anomalously high near-surface air temperatures and positive anomalies in precipitable water (column integrated water vapor) were found over inland Eurasia before the event and then over the Yamal Peninsula (**Figure 4**). While local herders reported a severe icing of pastures following the event, starting on November 10, 2-m temperature anomalies shown by ERA5 are positive even after this date, possibly reflecting limitations of the snow parameterization (Arduini et al. 2019). Ice covered an area of approximately 27,000 km² and completely blocked reindeer from foraging, leading to the death of 61,000 animals between November 2013 and June 2014 (Forbes et al. 2016).




Figure 4. Daily average sea level pressure and precipitable water (top row), and anomalies of daily average 2 m air temperature (bottom row) from ERA5 over the Yamal Peninsula between 7 November and 10 November 2013. Regions with precipitable water less than 10 mm, considered very dry air, are shown in white. Air temperature anomalies are calculated with respect to 1979 to 2020 daily average fields, with a 30-day boxcar averaging filter applied.

Socioeconomic impacts unfolded for years after. Nenets nomads in Arctic Russia were particularly affected (Stammler 2005) – many herders lost most or all of their animals. This left some families stranded on the tundra without viable transportation (Forbes et al. 2016). Additionally, the ground icing made large portions of migration routes inaccessible. In response, nomadic Nenets had to adjust their migration routes and the timing of migration. Nenets were forced to adapt by finding new food sources and breeding techniques in an attempt to rebuild their livelihoods.

During the 5-year period following 2013, several additional significant ROS events occurred on the Yamal Peninsula (Laptander 2020), resulting in a situation where tundra Nenets nomads report not having enough time between events to regenerate the size, demography and health of their herds (Stammler

and Ivanova 2020). By the spring of 2019, some herding communities were still subsisting entirely on fish and trying to rebuild their stock of animals.

3.2 Svalbard, Norway, 30-31 January, 2012

Figure 3 suggests that the Svalbard Archipelago, located between 78-80°N latitude, experiences up to one event per year over the central and northern parts of the largest island of Spitsbergen, and coastal regions of the more northerly Nordaustlandet. A greater number of events are detected on the west and east coasts of Spitsbergen. This reflects the region's proximity to open water and its location along the terminus of the North Atlantic cyclone track, yielding a relatively warm winter climate with respect to latitude (Serreze et al. 2015). During a two-week period in late January and early February 2012, a remarkable event brought above-freezing air temperatures to the entire archipelago and record-breaking precipitation.

Key meteorological aspects are summarized in **Figure 5**. A blocking pattern over northern Fennoscandia fostered the migration of low-pressure systems up into the Arctic and the influx of warm and moist air towards Svalbard (Hansen et al. 2014). The stream of moist air was part of an atmospheric river (Serreze et al. 2015). Many previous studies have documented links between atmospheric rivers and extreme precipitation events, such as along the coasts of California and Greenland. Positive anomalies in precipitable water moved into the area on January 27 and persisted throughout the course of the warm spell. On January 30, record breaking precipitation was recorded at the Ny Ålesund meteorological station, with 98 mm of rain (Serreze et al. 2015). On this day, there was strong high pressure over northern Fennoscandia and Eurasia, and low pressure over Svalbard that originated in the North Atlantic. The low-pressure movement was paired with positive precipitable water anomalies directly over the archipelago, with the largest positive air temperature anomalies located just to the northeast.




Figure 5. Daily average sea level pressure and precipitable water (top row), and anomalies of daily average 2 m air temperature (bottom row) from ERA5 over the North Atlantic and Svalbard Archipelago between 30 January and 2 February 2012. Regions with precipitable water less than 10 mm, considered very dry air, are shown in white. Air temperature anomalies are calculated with respect to 1979 to 2020 daily average fields, with a 30-day boxcar averaging filter applied.

Temperatures then dropped, and Spitsbergen became covered in 10-20 cm of ice (Hansen et al. 2014), covering the low growing tundra vegetation and blocking reindeer from grazing. Despite favorable feeding conditions prior to the ROS event, there was a large starvation induced mortality episode. The number of reindeer carcasses counted in the summer 2012 census was among the highest ever recorded (Hansen et al. 2014).

In Longyearbyen, a slush avalanche destroyed a pedestrian bridge and forced roads to shut down for several days. Ice formation on the Svalbard airport runway forced the airport to shut down on January 29 and 30. The impacts that avalanches and ground ice had on transportation, including transportation by snowmobile, dogsled, and pedestrian transport, resulted in massive income losses for the local tourism industry. After the event, local permafrost experienced warming down to at least 5 m. At several monitoring sites, the ground surface temperature stayed at 0 °C for weeks after the warm spell had passed (Hansen et al. 2014).

3.3 Banks Island, Canada, 3-8 October, 2003

Banks Island, in the western Canadian Arctic Archipelago (Figure 3, Region 3), has a cold and very dry environment classified as polar desert. ROS events from November through February appear to be uncommon. However, in early October 2003, the climatological flow that usually brings dry air from Siberia briefly shifted and warmer, moister air from the Pacific invaded the region (Rennert et al. 2009).

The weather system initially brought approximately six inches of new snow, but from October 3-8 the precipitation fell as drizzly, intermittent rains. This event was the first to be documented with satellite observations (Grenfell and Putkonen 2008).

In the days before the event, the southwesterly flow was attended by a strong low-pressure system centered in the Bering Strait. As the event began, the low tracked along the Alaskan coast towards Banks Island, and was attended by positive anomalies in both air temperature and precipitable water. Slightly positive anomalies of both existed over Banks Island prior to the main ROS event, most notably during October 3-4 (**Figure 6**). On October 7, high pressure moved over Banks Island, bringing colder and drier air. According to reports from hunters in the area at the time, following intermittent rains and falling temperatures, the former 6 inches of new snow was turned into a thick sheet of ice on the northern two thirds of the island (Rennert et al. 2009).




Figure 6. Daily average sea level pressure and precipitable water (top row), and anomalies of daily average 2 m air temperature (bottom row) from ERA5 over the Banks Island between 3 October and 6 October 2003. Regions with precipitable water less than 10 mm, considered very dry air, are shown in white. Air temperature anomalies are calculated with respect to 1979 to 2020 daily average fields, with a 30-day boxcar averaging filter applied.

The surface icing had severe impacts on Banks Island's ungulate populations, especially musk oxen; approximately 20,000 animals died. In addition to the physical barrier created by ice, water that had percolated through the snowpack and pooled at the ground surface caused vegetation to spoil. This forced musk oxen to expand their grazing range. After the event, in mid- to late winter, some animals wandered onto the sea ice in search of food and were left stranded at sea (Rennert et al. 2009).

Significant ungulate mortality events in the western Canadian Arctic date back several decades (Parker et al. 1975, Gunn et al. 1989, Miller and Gunn 2003), but as noted, the October 2003 ROS event was the first that could be documented via satellite data. Rennert et al. (2009) conclude that the severe impacts of the event can be traced to it having occurred early in the cold season so that food sources were blocked for the entire winter. The direct mortality from starvation was most prevalent within the youngest and oldest age groups. Only a few musk oxen were healthy enough to maintain pregnancy, and there were very few calves born the following summer. Rennert et al. (2009) also argues that, as has been observed for reindeer, the lack of new calves coupled with the decline of overall herd health can remove multiple generations of calves from a herd, leading to a population crash years after the ROS event. For this particular event, there were few calves born in summer 2004 with another population decline expected 4-5 years later as the adults aged without replacement (Rennert et al. 2009).

4. ROS Detection and Characteristics

4.1 Overview

A key element of the ongoing Arctic Rain on Snow Study is to provide a pan-Arctic perspective of ROS events, their frequency, intensity and impacts by combining information from meteorological station records, local and Indigenous knowledge holders, satellite remote sensing and atmospheric reanalyses. Surface observations provide ground truth for developing detection algorithms and combining multiple information sources. It is only from local observers that one can obtain critical information bearing on impacts such as how ice layers form, how thick they become and whether they form at the snow surface, within, or at the base of the snowpack. In turn, data from satellite retrievals and atmospheric reanalyses addresses the spatial scale of events, which can range from local to widespread.

4.2 Meteorological Records

An ongoing challenge with station meteorological records is that records span many individual sources. A series of Environment and Climate Change Canada sites are accessible across the Canadian Arctic (Mekis et al. 2018). A number of archives are available through the National Oceanic Atmospheric Administration (NOAA) National Weather Service (NWS). Some records from NOAA/NWS consist of automated surface observation stations and Cooperative Observer Network records, mainly for sites located in Alaska. An additional data source for Alaska and some parts of Canada is snow telemetry (SNOTEL) sites maintained by the Natural Resources Conservation Service (e.g. used for ROS detection in Wilson et al. 2013, Semmens et al. 2013, Pan et al. 2018). Other sources include the Norwegian Meteorological Institute, the Finland Meteorological Institute and the Swedish Meteorological and Hydrological Institute. Some information over the Arctic Ocean is available from the Russian North Pole series of drifting stations (ongoing) and field campaigns. The World Meteorological Organization's Observing Systems Capability Analysis and Review Tool (OSCAR) provides information on data sources across the globe. While station records of air temperature, precipitation and snow height serve only as indicators of ROS, SYNOP (present weather codes) made by observers at the station (codes 20-29 corresponding to precipitation and type) can be highly valuable.

Spatially as well as temporally extensive snow profile observations exist only in rare cases. A so far unique dataset, a collection of more than 500 snow profiles during the period 2010–2017 in Central Spitsbergen, was used to analyze long term patterns directly (Peeters et al. 2019).

4.3 Local Observers

Observations of ROS events from reindeer herders, harvesters and others can greatly augment station records, but more importantly, can provide detailed information on event characteristics key to understanding impacts. Reindeer herders move over wide areas during the year, communicate across herding families on topics related to weather and herd health, and pay close attention to conditions limiting reindeer foraging in winter pastures. Such observations, however, have seldom been systematically collected, one exception being Rasmus et al. (2018), who compiled data on annual icing events from 1948-2016 for the reindeer management area of Finland, based on archival sources (annual management reports of herding districts). Rasmus et al. (2018) focused on the formation of basal ice layers on the pastures. They found that rain and subsequent freezing of the liquid precipitation in the basal layer of the snow cover (or in the whole of a thin snow cover) was reported in 27% of cases. Icing was also found to be regularly attended by mold formation on pastures during early winter, which can have severe negative effects on the condition of reindeer (Kumpula et al. 2000).

Another exception is a collection of 17 interviews with reindeer herders from Alaska’s Seward Peninsula, a collaborative effort between the Reindeer Herders Association (RHA) in Nome, Alaska and the University of Alaska Fairbanks Oral History Department. These interviews, accessible online, discuss herding practices, the generational passing-down of herder knowledge, and how climate and environmental change impact herder livelihoods and the reindeer themselves. As gleaned from these interviews, herders nowadays observe more thawing events interlaced between snowfalls during freeze-up, resulting in a snow pack that is icy and consolidated from top to bottom once the winter cold season sets in, making it more difficult for reindeer to access feed (e.g., Larry Davis, 2001, Tape Number: H2000-102-03; Project Jukebox 2001).

A potentially rich information source is organized observer networks. Generally, observer networks focus on qualitative information such as photos and videos that may be posted on specific observer web-platforms or social media, and first-hand narratives from “on the ground” experience. Some networks are broad in their focus, and the observers that participate are necessarily generalists. However, other observer networks may be highly knowledgeable in specific aspects of their local places and community activities. Observers with the Alaska Arctic Observatory and Knowledge Hub (AAOKH) specialize in observing seasonality, focusing on the snow, ice, weather, coastal processes, and related community travel and hunting activities. The AAOKH observers’ database (<https://eloka-arctic.org/sizonet/>), maintained by the Exchange for Local Observations and Knowledge of the Arctic (ELOKA), contains some detailed observations of winter rain events, often in the context of local travel or hunting activities. For example, an observation from Point Hope in November 2019 described how a rain event led to the break-up of a foot of ice on the Kukpuk River, leaving a wall of ice along the riverbank (Adams 2013a). An observer from Toksook Bay noted in late December 2009 that glare ice,

resulting from a freezing rain event, prevented planes from reaching the village (Adams 2013b). An event that resulted in a change of migration patterns in 2006-07 on Yamal was documented by an anthropologist traveling with reindeer herders (Bartsch et al. 2010). A more extensive event during winter 2013-14 on Yamal similarly triggered substantial reorganization of migration patterns, which were mapped using a participatory approach with reindeer herders, administrators and scientists (Forbes et al. 2016).

Also notable is the Local Environmental Observer (LEO) Network, with observers concentrated in North America and growing membership around the circumpolar North. The LEO Network provides an online platform for sharing information about significant environmental events. Local knowledge experts share observations and engage with topic experts as co-authors on LEO Network posts. Members can also post news articles as another local source for these events. The LEO Network has roots in the Indigenous communities of Alaska, but has grown greatly. As of June 2020, the LEO Network has over 3,500 members from over 50 countries (including all Arctic nations). The LEO editorial process involves validating observations, geocoding, time stamping, sharing them with subject matter experts, organizing them by topic and geography, and publishing them to the online platform (**Figure 7**). Two of the most common event categories are *Weather*, and *Ice and Snow*. Much less common, but still numerous, are events further categorized as *Rain on Snow*, with 47 events in the LEO database as of November 2020. While many of these ROS entries discuss impacts on animals, other impacts are also noted, such as diminished recreational opportunities, public safety issues from icing and infrastructure issues involving power and telephone lines, and flooding.




Figure 7: Schematic information sources and information flow in the LEO network.

4.4 Satellite Retrievals

ROS detection from satellite microwave retrievals (both passive and active) shows considerable promise for providing a pan-Arctic view (e.g. Bartsch et al. 2010, Dolant et al. 2016, 2017, Langlois et al. 2017)

(Table 1). This is because microwave emission or backscatter from the surface is strongly affected by liquid water and snow cover structure, as well as the strength of impact depending on microwave frequency and polarization.

In passive microwave radiometry, the energy detected by the satellite is the measured emission intensity, converted to a brightness temperature that is a function of the physical temperature of the target and the emissivity. The emissivity of the snow cover is determined by its dielectric properties and its surface roughness. When there is liquid water in the snow, emissivity sharply increases because of the much higher dielectric constant of water, leading to a sudden increase in passive microwave brightness temperature. In this case, absorption processes dominate over scattering, and consequently only the topmost snow layers contribute to the measured brightness temperature. The response is larger at higher frequencies because of the change in emission depth associated with melt. The response is also polarization dependent (Anderson 1997). The response of emissivity to liquid water is the basis for snowmelt onset detection from microwave brightness temperatures on land (Frei et al. 2012, Alimasi et al. 2018) as well as on sea ice (e.g. Markus et al. 2009, Wang et al. 2013) and ice sheets (Abdalati and Steffen 1995).

As for active microwave, microwave energy transmitted by a radar is reflected and scattered by the snowpack. The unit of measure is called the backscatter, and depends on the surface area illuminated, and the distance from the receiver to the target. During the transition from dry to wet snow, absorption is the main mechanism, driven by increased dielectric loss from the snowpack, causing the radar backscatter to decrease (Ulaby et al. 1982). Volume scattering occurs under dry snow conditions in the case of short wavelengths.

The strategy of wet snow detection is applicable when measurement intervals are shorter than the ROS events themselves. Given that event duration can be limited to a few hours (Bartsch et al. 2010) it is thus very likely that events are missed. Satellite observations from polar orbits (as is the case for relevant active and passive microwave missions) are usually limited to certain time windows, in the morning and afternoon. An alternative is to focus on ROS impacts on snow structure. The resulting ice layers change the interaction of the microwave radiation with the snowpack. In case of for example active microwave techniques, the backscatter response increases when the snow grain size increases and specifically when crusts are present. Such changes can occur as part of seasonal snow metamorphosis (grain size increases with snow age). These are, however, gradual changes, compared to ROS impacts. The determination of sudden changes can therefore reveal potential ROS events. This has been so far mostly applied to Ku-band radar for monitoring larger areas as in Figure 3 (Bartsch et al. 2010, Bartsch 2010a, b, Wilson et al. 2013, Semmens et al. 2013) although passive microwave data are suitable as well to identify changes (Grenfell and Putkonen 2008, Langlois et al. 2017). C-band radar is also promising (Forbes et al. 2016) and offers potentially high resolution (down to 10m). This is based on the synthetic aperture radar technique. Such detail cannot be achieved with passive microwave to date. What also needs to be considered for this technique is that multiple events lead to stepwise increase of backscatter and the magnitude of change decreases throughout the season. In extreme cases, a maximum level (similar as backscatter levels of ice sheets) is reached, and detection of further events is impeded. The presence of vegetation also leads to volume scattering and increases the backscatter

minimum at the beginning of the snow-covered season, reducing sensitivity to snow structure change (Bartsch et al. 2020).

A challenge with longer wavelengths such as C-band is the sensitivity to the soil beneath the snowpack. Changes in midwinter which are related to the evolution of frozen state dielectric properties (driven by ground temperature) can also lead to high backscatter response similar to snow crusts, and saturate the signal (Bergstedt et al. 2018). Recent fire events can also alter soil wetness and temperature which impacts the snowpack structure for several decades (Bartsch et al. 2020).

In general, retrieval algorithms use threshold values that require tuning with respect to observations and are unlikely to hold everywhere as weather effects and snowpack properties influence the retrievals. Furthermore, the detection of liquid water does not necessarily mean that an ROS event occurred. It can also not be directly inferred if snow is present on the ground. A combination with snow cover datasets based on multi-spectral observations is therefore useful (Pan et al. 2018). A further strategy is limiting analysis to mid-winter when warm spells are less likely. The definition of this period does, however, strongly differ between different studies (Table 1) which complicates comparing results. An alternative is the combination with reanalysis data such as applied by Semmens et al. (2013) and described in Section 4.5. To date, the only study for the entire Arctic land area which is based on satellite records is Bartsch (2010a), derived from Ku-band QuikScat records from 2000-2009 (Figure 3). The algorithm has been initially based on local observations by reindeer herders (Bartsch et al. 2010).

Table 1: Overview of remote sensing and reanalyses studies of ROS events

Source/ Method	Sensor (s) or parameter	Study	Analyses period	Season	Region
Passive microwave: Gradient ratio/ polarization ratio Method: wet snow detection	SSMI	Grenfell and Putkonen 2008	2003/4	Single event	Banks island, Canada
	AMSR-E/2	Semmens et al. 2013	2003-2009	All	Yukon basin
	AMSR-E/2	Dolant et al. 2016	2010/11	All	Nunavik
	SSMI + AMSR-E/2	Langlois et al. 2017	1979-2011	October – May	Canadian Arctic Archipelago
	AMSR-E/2 + Modis	Pan et al. 2018	2003-2016	November – March	Alaska
Active microwave:	Quikscat	Bartsch et al. 2010	2000-2009	Nov – Feb	Russia

Single band backscatter Method: Snow structure change detection		Bartsch 2010a	2000-2009	Nov – Feb	Circumpolar
		Wilson et al. 2013	2000-2009	April-May	Alaska
		Semmens et al. 2013	2003-2009	All	Yukon basin
	ASCAT	Forbes et al. 2016	2013	Single event	South Yamal
Reanalyses - MERRA	Rain and temperature	Liston and Hiemstra et al. 2011	1979–2009	All	Circumpolar
Reanalyses - ERA-40	Rain	Rennert et al. 2009	1957–2002	-	Circumpolar
Reanalyses - seNorge	Rain and snow water equivalent	Pall et al. 2019	1957-2016	at least 3mm of snow water equivalent	Norway

The use of satellite data allows for the determination of spatial patterns. The extent of events is crucial, such as with respect to the migration possibilities of ungulates. Wilson et al. (2013) analyzed event sizes for the first time and found differences with respect to distance from coast for Alaska. Events in proximity to the coast tend to be comparably small and frequent compared to further inland where they are larger and less frequent. Such types of investigations are, however, still lacking across the rest of the Arctic.

4.5 Atmospheric Reanalyses

Atmospheric reanalyses, the most recent example being the European Center for Medium Range Forecasts ERA5 effort, represent retrospective forms of numerical weather predictions models. A great advantage of reanalyses (and operational Numerical Weather Prediction (NWP) systems) is the provision of forecasts of precipitation amount and phase along with information on the vertical temperature and humidity structure. In an operational setting, forecasts of precipitation type and amount are made in a post-processing step using information from NWP systems about the vertical structure of temperature and humidity as well as other variables (Elmore et al. 2015). All of the modern generation of atmospheric reanalyses provide data from 1979 or 1980 onwards (1979 represents the start of the modern satellite era). ERA5 provides data from 1950, and the Japan Meteorological Agency JRA55 provides data from 1958. A national specific dataset used in this context is seNorge, a high-resolution gridded hydrometeorological dataset covering mainland Norway (Pall et al. 2019).

The quality of NWP precipitation forecasts (for both operational systems and reanalysis) depends on both the physics in the model and the amount and quality of observations that can be assimilated to

produce the analysis from which forecasts are generated. It is commonly thought that NWP forecasts in the Arctic are limited by sparse observational data. This is true only to an extent - while the radiosonde network is suboptimal, polar orbiting satellites provide a wealth of data. Uncertainties and biases in reanalysis outputs in the Arctic region have been widely addressed. All reanalyses have biases in Arctic precipitation relative to observations (e.g. Wang et al. 2019, Barrett et al. 2020).

Output from the North American Regional Reanalysis (NARR) has been employed in a number of ROS studies, serving as a basis for evaluation of satellite-based retrievals (Wilson et al. 2013, Semmens et al. 2013). Precipitable water (total column water vapour) from re-analyses data also serves as input for correction of passive microwave observations (Langlois et al. 2017). Re-analyses records are also useful for assessing the reasons behind snow structure changes detected with satellite records. Near-surface air temperature and dew point temperature, visibility, and relative humidity, were used to determine the occurrence of fog by Semmens et al. (2013). The relatively moist and warm conditions forming fog are considered to affect snow and melting. In some years, more than 60% of snow structure change events over the Yukon basin could be attributed to fog and only a maximum of 35% to ROS events.

5. A Path Forward

Knowledge co-production can be viewed as a process through which technical experts and other groups interact with their different ways of viewing and analyzing the world, thereby gaining new knowledge and technologies. While multiple data sources, notably remote sensing and weather and climate models, can be used and combined in ways to improve our ability to detect and forecast ROS events and anticipate future changes, partnering with local observers can convey key information regarding ROS impacts, providing a focus for developing better technologies and providing information useful to stakeholders. For example, while ROS impacts are known to be tied to the formation of ice layers, ice layer formation processes are poorly understood and difficult to model and ground observations are sparse (Kohler and Aanes 2004, Liston and Hiemstra 2011, Vikhamar-Schuler et al. 2013, Rasmus et al., 2014, 2016, Pirazzini et al. 2018). Having detailed surface observations offers a path for developing better process models and for understanding how the development of ice layers ties into the larger-scale meteorological aspects of ROS events.

The approach taken by AROSS revolves around developing a “Data and Knowledge Hub” as a community resource, a key part of which is developing a “data and information package” for every detected ROS event from 1979 onwards, combining physical science and observer components from AAOKH, the LEO Network and other local and Indigenous partners. The database, searchable by event date, location, event duration, area affected, total precipitation as well as whether any local observations are available, is intended to 1) provide a platform for observers and physical scientists to be equal partners in AROSS; 2) promote better understanding of ROS event meteorology, including common and unique features; and 3) as just discussed, provide an avenue for a collaborative and interdisciplinary approach to focus on event characteristics most relevant to impacts.

The physical science component of each data and information package includes event and post-event period gridded atmospheric variables and surface fields, including precipitation, temperature and those grid cells for which ROS is detected. Different fields can be overlain and animated to track the evolution of each event. ROS detection relies on an algorithm that combines information from satellite passive microwave retrievals and output from several different atmospheric reanalysis. Surface observations are used to train, test and validate the algorithm. **Figure 8** provides an example of the information that can be obtained from these data sources, while **Figure 9** provides a schematic of the algorithm. The observation component of each data and information package contains all available information from observers participating in AROSS as text, photographs, videos or from social or mainstream media relevant to the physical expression of the event and any impacts, whether they be immediate, delayed or cumulative. AROSS partners with the international CHARTER effort, focused on developing enhanced active microwave sensing techniques, which allows for better understanding of spatial patterns. Special emphasis is on impact determination via satellite retrievals, paleo-records, local community observations and various documentations of biodiversity. The determination of severity is addressed from multiple viewpoints, based on impacts on reindeer herding, local communities and the natural environment. Ultimately, the goal of AROSS and CHARTER partnership is to develop ROS detection capabilities and integrated outreach and communication products that provide useful information to local communities and herders in the Arctic, as well as to other decision makers interested in the intersection between climate change, ROS events, and their impacts.




Figure 8. Tropospheric air temperature and precipitation amount and phase for the closest MERRA2 (Gelaro et al. 2017) grid point to 67.75° N and 68° E on the Yamal Peninsula for 4 to 11 November 2013 corresponding to the events studied by Forbes et al. (2016). Also shown is precipitation from the ERA5 and MERRA2 reanalyses at the same location, and the ratio of vertical and horizontal

polarization gradient ratios between the 18 GHz and 36 GHz bands from AMSR2, with values <1 indicated liquid water at the surface.




Figure 9. Flow diagram showing potential inputs, and procedure for creating training data and developing the ROS detection algorithm.

6. Conclusions

Despite advances in our understanding of ROS events and their impacts, many gaps in knowledge still exist. These include uncertainties in detecting liquid water on the snowpack from satellite data, detection of ROS events and precipitation amount from atmospheric reanalyses, and the general sparseness of direct surface observations. Another is a lack of understanding of ice layer formation processes.

As the Arctic warms, ROS and winter thaw events are likely to become more frequent and there is growing evidence that changes are already happening (Hansen et al. 2011, Liston and Hiemstra 2011, Ruosteenoja et al. 2016, Vikhamer-Schuler et al. 2016, Luomaranta et al. 2019). Climate models project a shift toward more Arctic rain events (Bintanja and Andry 2017), with total Arctic precipitation expected to increase by 30-60% (Bintanja and Selten, 2014, Bintanja and Andry 2017). Uncertainties abound, however - projected changes in precipitation amount and type range widely between different models using the same emissions scenario, and future emission rates are unknown. In turn, it is possible that a decrease in the duration of the snow cover season could at least in part counterbalance the effects on more rainfall and the frequency of ROS events (Mudryk et al. 2020).

It is clear that to understand Arctic ROS events and their impacts, there needs to be a co-production of knowledge approach, one that bridges scientific disciplines, Indigenous knowledge, local observations and especially ensures Indigenous and local collaboration in the research from the outset. To this end, AROSS and CHARTER have assembled a multi-disciplinary and multi-cultural research team, including atmospheric scientists, harvesters, herders, social scientists, modelers, Indigenous scholars and others. We are working together to develop new understandings of ROS events and their impacts.

Acknowledgements: This study was supported by NSF Grant NNA 1928230. The project has received funding under the European Union's Horizon 2020 Research and Innovation Programme under grant agreement no. 869471.

7. References

Abdalati W and Steffen K 1995 Passive microwave-defined snow melt regions on the Greenland Ice Sheet *Geophys. Res. Lett.* 22 787-790

ACIA 2005 *Arctic Climate Impact Assessment* Cambridge University Press, Cambridge, UK

Adams B, Apangalook L, Apangalook P, John S, Leavitt J, Weyapuk M Jr and other observers 2013a *Local Observations from the Seasonal Ice Zone Observing Network (SIZONet)*. Observation from Guy Omnik, Nov. 1, 2019. Edited by H. Eicken and M. Kaufman. Boulder, Colorado USA: National Snow and Ice Data Center <http://dx.doi.org/10.7265/N5TB14VT>

Adams B, Apangalook L, Apangalook P, John S, Leavitt J, Weyapuk W J and other observers 2013b *Local Observations from the Seasonal Ice Zone Observing Network (SIZONet)*. Observation from Dilool Erickson, Dec. 28, 2009. Edited by H. Eicken and M. Kaufman. Boulder, Colorado USA: National Snow and Ice Data Center <http://dx.doi.org/10.7265/N5TB14VT>

Alimasi, N 2018 A review of passive microwave observations of snow-covered areas over complex Arctic terrain *Bulletin of Glaciological Research* 36 1-13 <https://doi.org/10.5331/bgr.18W01>

Anderson M 1997 Determination of a melt onset date for Arctic sea ice regions using passive microwave data *Ann. Glaciol.* 25 382-387

Andrews C L 1939 *The Eskimo and His Reindeer in Alaska* The Caxon Printers Ltd. Caldwell Idaho 253 pp

Arduini G, Balsamo G, Dutra E, Day J J, Sandu I, Bousssetta S, Haiden T 2019 Impact of a multi-layer snow scheme on near-surface weather forecasts *Journal of Advances in Modeling Earth Systems* 11 4687–4710 <https://doi.org/10.1029/2019MS001725>

Barrett A P, Stroeve J and Serreze M C 2020 Arctic Ocean precipitation from atmospheric reanalyses and comparisons with North Pole Drifting Station records *J. Geophys. Res. Oceans* 125, 1–17

Bartsch A 2010a Spring snowmelt and midwinter thaw and refreeze north of 60°N based on Seawinds QuikScat 2000-2009 *PANGAEA* <https://doi.org/10.1594/PANGAEA.834198>

Bartsch A, Kumpula, T, Forbes B C and Stammeler F 2010 Detection of snow surface thawing and refreezing in the Eurasian Arctic with QuikSCAT: implications for reindeer herding *Ecological Applications* 20 2346-2358.

Bartsch, A 2010b Ten years of SeaWinds on QuikSCAT for snow applications *Remote Sensing* 2 1142-1156, <https://doi.org/10.3390/rs2041142>

Bartsch A, Widhalm B, Leibman M et al. 2020 Feasibility of tundra vegetation height retrieval from Sentinel-1 and Sentinel-2 data *Remote Sensing of Environment* 237 111515 doi: 10.1016/j.rse.2019.111515

Berg A et al. 2011a At this point, the lichens in the trees are their only means of survival: A history of tree cutting for winter reindeer fodder by Sami people in northern Sweden *Environment and History* 17 265-289

Berg A et al. 2011b Cutting of lichen trees: a survival strategy used before the 20th century in northern Sweden *Vegetation History and Archaeobotany* 20 125–133

Bergstedt, H., Zwieback, S, Bartsch, A and Liebman M 2018 Dependence of C-band backscatter on ground temperature, air temperature and snow depth in Arctic permafrost regions *remote Sensing* 10 <https://www.mdpi.com/2072-4292/10/1/142>

Bintanja R and Andry O 2017 Towards a rain-dominated Arctic *Nature Climate Change* 7 263-267 <https://doi.org/10.1038/NCLIMATE3420>

Bintanja R and Selten F M 2014 Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat *Nature* doi:10.1038/nature13259

- Bjerke J W, Tømmervik H, Zielke M. and Jørgensen M 2015 Impacts of snow season on ground-ice accumulation, soil frost and primary productivity in a grassland of sub-Arctic Norway *Env. Res. Lett.* 10 095007
- Bjerke J, Treharne R, Vikhamer-Schuler D et al. 2017 Understanding the drivers of extensive plant damage in boreal and Arctic ecosystems: Insights from field surveys in the aftermath of damage *Science of the Total Environment* 590-600, 1965-1776 <https://doi.org/10.1016/j.scitotenv.2017.05.050>
- Dolant C, Langlois A, Montpetit A, Brucker L, Roy A and Royer A 2016 Development of a rain-on-snow detection algorithm using passive microwave radiometry *Hydrological Processes* 30 3184-3196
- Dolant C, Langlois A, Brucker L, Royer A, Roy A and Montpetit B 2017 Meteorological inventory of rain-on-snow events in the Canadian Arctic Archipelago and satellite detection assessment using passive microwave data *Physical Geography* <https://doi.org/10.1081/02723646.2017.1400339>
- Egeland G M et al. 2013 Back to the future: using traditional food and knowledge to promote a healthy future among Inuit In, *Indigenous Peoples' food systems and well-being: interventions and policies for healthy communities* FAO of the UN Rome 371 pp
- Eira I M G, Oskal A, Hanssen-Bauer I and Mathiesen S D 2018 Snow cover and the loss of traditional indigenous knowledge *Nature Climate Change* 8 928-931 doi: <https://doi.org/10.1038/s41558-018-0319-2>
- Elmore K L, Grams H M, Apps D, and Reeves H D 2015 Verifying forecast precipitation type with mPING *Weather Forecasting* 30 656–667 doi: 10.1175/WAF-D-14-00068.1
- Forbes B C and Kumpula T 2009 The ecological role of reindeer (*rangifer tarandus*) in northern Eurasia *Geography Compass* <https://doi.org/10.1111/j.1749-8198.2009.00250.x>
- Forbes B C, Kumpula T, Meschtyb N et al. 2016 Sea ice, rain-on-snow and tundra reindeer nomadism in Arctic Russia *Biology Letters* 12 doi: 10.1096/rsbl2016.0466
- Frei A, Tedesco M, Lee S, Foster J, Hall D K, Kelly R and Robinson D A 2012 A review of global satellite-derived products *Advances in Space Research* 8, 1007-1029, doi 10.1016/j.asr.2011.12.021
- Freudiger D, Kohn I, Stahl K and Weiler M 2014 Large scale analysis of changing frequencies of rain-on-snow events with flood generation potential *Hydrology and Earth System Sciences* 18 2695–2709.
- Freund K and Bartsch A 2020 Midwinter thaw events over Greenland derived from Seawinds QuikScat 2000-2008 *PANGAEA* [doi:10.1594/PANGAEA.911298](https://doi.org/10.1594/PANGAEA.911298)
- Furgal C M, Innes S and Kovacs K M 2002 Inuit spring hunting techniques and local knowledge of the ringed seal in Arctic Bay (Ikpiarjuk), Nunavut *Polar Research* 21 1-16
- Gearheard S, Pocernich M, Stewart R, Sanguya J and Huntington H P 2010 Linking Inuit knowledge and meteorological station observations to understand changing wind patterns at Clyde River, Nunavut *Climatic Change* 100 267-294 doi: 10.1007/s10584-009-9587-1

- Gelaro R, McCarty Suarez M J et al. 2017 The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2) *J. Climate* 30 5419–5454 doi: 10.1175/JCLI-D-16-0758.1
- Grenfell T C and Putkonen J 2008 A method for the detection of the severe rain-on-snow event on Banks Island, October 2003, using passive microwave remote sensing *Water Resources Res.* 44 W03425 doi:10.1029/2007WR005929
- Guan B, Waliser D E, Ralph F M, Fetzer E J, and Neiman P J 2016 Hydrometeorological characteristics of rain-on-snow events associated with atmospheric rivers *Geophys. Res. Lett.* 43 2964–2973 doi:10.1002/2016GL067978.
- Gunn A, Miller F L and McLean B 1989 Evidence for and possible causes of increased mortality of bull muskoxen during severe winters *Can. J. Zool.* 67 1106–1111.
- Hansen B B, Aanes R, Herfindal I, Kohler J and Sæther B-E 2011 Climate, icing, and wild arctic reindeer: past relationships and future prospects *Ecology* 92 1917–1923 doi:10.1890/11-0095.1
- Hansen B B, Isaksen K, Benestad R E, Kohler J et al. 2014 Warmer and wetter winters: characteristics and implications of an extreme weather event in the High Arctic *Environ. Res. Lett.* 9 114021 doi:10.1088/1748-9326/9/11/114021.
- Helle T 1984 Foraging behaviour of semi-domesticated reindeer (*Rangifer tarandus tarandus*) in relation to snow in Finnish Lapland *Rep. Kevo Subarctic Res. Stat.* 19 35–47
- Helle T and Kojola I 2008 Demographics in an alpine reindeer herd: effects of density and winter weather *Ecography* 31 221–230 doi: 10.1111/j.2008.0906-7590.04912.x
- Helle T and Jaakkola L M 2008 Transition in herd management of semi-domesticated reindeer in northern Finland *Annales Zoologici Fennici* 45 81–101.
- Hersbach, H, Bell B, Berrisford P, et al. The ERA5 global reanalysis *Q J R Meteorol Soc.* 2020; 146: 1999–2049. <https://doi.org/10.1002/qj.3803>
- Huntington H P, Cary M, Apok C et al. 2019 Climate change in context: putting people first in the Arctic *Regional Environmental Change* 19 1217–1223 <https://doi.org/10.1007/s10113-019-01478-9>
- Jamieson B 2006, Formation of refrozen snowpack layers and their role in slab avalanche release, *Rev. Geophys.* 44 RG2001 doi:10.1029/2005RG000176.
- Kausrud K L, Mysterud A, Steen H et al. 2008 Linking climate change to lemming cycles *Nature* 456 93–U3
- Kawerak Reindeer Herders Association, 2020. Reindeer Herders Association. Accessed on 3 December 2020 <https://kawerak.org/natural-resources/reindeer-herders-association/>
- Kohler J and Aanes R 2004 Effect of winter snow on ground-icing on a Svalbard reindeer population: Results of a simple snowpack model *Arctic and Alpine Research* 36 333–341 doi: 10.1657/1523-0430(2004)035[0333: EOWSAG]2.0.CO;2
- Krupnik I and Jolly D eds. 2002. *The Earth is Faster Now: Indigenous Observations of Arctic Environmental Change* Fairbanks, Alaska: Arctic Research Consortium of the United States. xxvii + 356p

- Kumpula J, Parikka, P and Nieminen M 2000 Occurrence of certain microfungi on reindeer pastures in northern Finland during winter 1996–97 *Rangifer* 20, 3–8
- Langlois A, Johnson C A, Montpetit B et al. 2017 Detection of rain-on-snow (ROS) events and ice layer formation using passive microwave radiometry: A context of Peary Caribou habitat in the Canadian Arctic *Rem. Sens. Env.* 189, 84-95
- Landauer M, Rasmus S and Forbes B C 2021 What drives reindeer management in Finland towards social and ecological tipping points? *Regional Environmental Change* (under review)
- Laptander R 2020: *When we got reindeer, we moved to live to the tundra: The spoken and silenced history of the Yamal Nenets* Rovaniemi: Lapin yliopisto. 169 p
- Liston, G E and Hiemstra C A 2011 The changing cryosphere: Pan-Arctic snow trends (1979–2009) *J. Climate* 24 5691–5712
- Luomaranta A, Aalto J and Jylhä K 2019. Snow cover trends in Finland over 1961 - 2014 based on gridded snow depth observations *Int. J. Climatol.* doi:10.1002/joc.6007
- Markus T, Stroeve J C and Miller J 2009 Recent changes in Arctic sea ice melt onset, freezeup, and melt season length *J. Geophys. Res.* 114, C12024, doi:10.1029/2009JC005436
- Mazurkiewicz A B, Callery D G, and McDonnell J J 2008 Assessing the controls of the snow energy balance and water available for runoff in a rain-on-snow environment *J. Hydrol.* 354, 1–14
- Mekis E, Donaldson N, Reid J et al. 2018 An overview of surface-based precipitation observations at Environment and Climate Change Canada *Atmosphere-Ocean* 56 71–95
<https://doi.org/10.1080/07055900.2018.1433627>.
- Miller F L and Gunn A 2003 Catastrophic die-off of Peary caribou on the western Queen Elizabeth Islands, Canadian High Arctic *Arctic* 56 381–390.
- Mudryk L, Santolaria-Otin Krinner M et al. 2020 Historical northern hemisphere snow cover trends and projected changes in the CMIP6 multi-model ensemble *The Cryosphere* 14 <https://doi.org/10.5194/tc-14-2459-2020>
- Mysterud I 2016 Range extensions of some boreal owl species: Comments on snow cover, ice crusts, and climate change *Arctic, Antarctic, and Alpine Research* 48 213-219 doi: 10.1657/AAAR0015-04
- Pall P, Tallaksen L, Stordal F 2019 A Climatology of Rain-on-Snow Events for Norway *J. Climate* 32, 6995-7016 DOI: 10.1175/JCLI-D-18-0529.1
- Pan C G, Kirchner P B, Kimball J S, Kim T and Du Jinyang 2018 Rain-on-snow events in Alaska, their frequency and distribution from satellite observations *Env. Res. Lett.* 13 <https://doi.org/10.1088/1748-9326/aac9d3>

Parker G R, Thomas D C, Broughton E, and Gray D R 1975 Crashes of muskox and Peary caribou populations in 1973 – 74 in the Parry Islands, Arctic Canada. *Canadian Wildlife Service Progress Notes* 56 10 p

Pearce T, Ford J, Willox A C and Smit B 2015 Inuit traditional ecological knowledge (TEK), subsistence hunting and adaptation to climate change in the Canadian Arctic *Arctic* 68 233-245
<http://www.jstor.org/stable/43871322>

Peeters B, Pedersen A O, Loe L E et al. 2019 Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: selection of a climate-cryosphere regime shift *Env. Res. Lett.* 14
<https://doi.org/10.1088/1748-9326/aaefb3>

Pirazzini R, Leppänen L, Picard G et al. 2018 European in-situ snow measurements: Practices and purposes *Sensors* 18 <https://doi.org/10.3390/s18072016>

Project Jukebox 2001 Reindeer Herding: The Past & The Present, University of Alaska Fairbanks Oral History Program, Fairbanks, AK, <http://jukebox.uaf.edu/reindherding/>.

Putkonen J and Roe G 2003 Rain-on-snow events impact soil temperatures and affect ungulate survival *Geophys. Res. Lett.* 30 1188. doi:10.1029/2002GL01632

Rasmus S, Kumpula J and Siitari J 2014 Can a snow structure model estimate snow characteristics relevant for reindeer husbandry? *Rangifer* 34 37-56

Rasmus S, Kivinen S, Bavay M and Heiskanen J 2016 Local and regional variability in snow conditions in northern Finland: a reindeer herding perspective. *Ambio* 45 398-414 doi: 10.1007/s13280-015-0762-5

Rasmus S, Kivinen S, Irannezhad M 2018 Basal ice formation in Northern Finland snow covers during 1948-2016 *Env. Res. Lett.* 13 114009 <https://doi.org/10.1088/1748-9326/aae541>

Rasmus S, Turunen M, Luomaranta et al. 2020 Climate change and reindeer management in Finland: co-analysis of practitioner knowledge and meteorological data for better adaptation *Science of the Total Environment* 710136229 doi: <https://doi.org/10.1016/j.scitotenv.2019.136229>

RHA 2020 Reindeer Herders Association Rovaniemi, Finland

Rennert K J, Roe G R, Putkonen J, and Bitz C M 2009 Soil thermal and ecological impacts of rain on snow events in the circumpolar Arctic *J. Climate* 22 2302-2315 doi: 10.1175/2008JCLI2117.1

Riseth J Å, Tømmervik, H and Bjerke, J W 2016 175 years of adaptation: North Scandinavian Sámi reindeer herding between government policies and winter climate variability (1835–2010) *Journal of Forest Economics* 24 186 – 204 doi: 10.1016/j.jfe.2016.05.002

Romanovsky V E, Smith S L, and Christiansen H H 2010 Permafrost thermal state in the polar northern hemisphere during the International Polar Year 2007–2009: A synthesis *Permafrost and Periglacial Processes* 21 106–116

- Royer M-J S, Herrmann T M, Sonnentag O, Fortier D, Delusca K and Cuciurean R 2013 Linking Cree hunters' and scientific observations of changing inland ice and meteorological conditions in the subarctic eastern James Bay region, Canada *Climatic Change* 119 719-732 doi 10.1007/s10584-013-0773-9
- Ruosteenoja K, Jylhä K and Kämäräinen M 2016 Climate projections for Finland under the RCP forcing scenarios *Geophysica* 51 17-50.
- Semmens K E, Ramage J, Bartch A and Liston G E 2013 Early snowmelt events: detection, distribution and significance in a major sub-arctic watershed *Env. Res. Lett.* 8 <http://iopscience.iop.org/1748-9326/8/1/014020>
- Serreze M C, Crawford A and Barrett A P 2015 Extreme daily precipitation events at Spitsbergen, an Arctic Island *Int. J. Climatol.* doi:10.1002/joc.4308
- Sokolov A A, Sokolova N A, Ims R A, Brucker L and Ehrich D 2016 Emergent rainy winter warm spells may promote boreal predator expansion into the Arctic *Arctic* 69 121-129 <https://www.jstor.org/stable/43871413>
- Staalesen A 2016 On Arctic island, a reindeer tragedy *Independent Barents Observer* 4 May 2016 <http://thebarentsobserver.com/ecology/2016/05/arctic-island-reindeer-tragedy>
- Stammler F 2005 Reindeer Nomads Meet the Market: Culture, Property and Globalisation at the 'End of the Land' Halle Studies in the Anthropology of Eurasia *Lit-Verlag* vol 6 320 pp
- Stammler F, and Ivanova A 2020 From spirits to conspiracy? Nomadic perceptions of climate change, pandemics and disease *Anthropology Today* 36 8-12 <https://doi.org/10.1111/1467-8322.12589>
- Statham S, Ford J, Berrang-Ford L, and Lardeau M-P 2015 Anomalous climatic conditions during winter 2010-2011 and vulnerability of the traditional Inuit food system in Iqaluit, Nunavut *Polar Record* 51 301-317 doi: 10.1017/s0032247414000151
- Stien A, Ims R A, Albon S D 2012 Congruent responses to weather variability in high arctic herbivores *Biology Letters* 8 1002–1005
- Stirling I and Smith T G 2003 Implications of warm temperatures and an unusual rain event for the survival of ringed seals on the coast on southeastern Baffin Island *Arctic* 57, 59–67 <https://doi.org/10.3390/rs8060457>
- Treharne R, Bjerke J W, Tommervik H and Phoenix G K 2020 Extreme event impact on CO2 fluxes across a range of high latitude, shrub dominated ecosystems *Env. Res. Lett.* <https://iopscience.iop.org/article/10.1088/1748-9326/abb0b1/pdf>
- Turunen M and Vuojala-Magga T 2014 Past and present winter feeding of reindeer in Finland: herders adaptive learning of the practices *Arctic* 67 173-188. doi: <http://dx.doi.org/10.14430/arctic4385>

- Turunen M, Rasmus S, Bavay M, Ruosteenoja K and Heiskanen J 2016 Coping with increasingly difficult weather and snow conditions: Reindeer herders' views on climate change impacts and coping strategies. *Climate Risk Management* 11 15–36 doi: 10.1016/j.crm.2016.01.002
- Tveraa T et al. 2003 An examination of a compensatory relationship between food limitation and predation in semi-domestic reindeer *Oecologia* 12 370-376.
- Ulaby F T, Moore, R K and Fung A K 1982 *Microwave Remote Sensing*, Vol. 11 Artech House, Dehram MA
- Vikhamar-Schuler D, Hanssen-Bauer I, Schuler T V, Mathiesen S D and Lehning M 2013 Use of a multi-layer snow model to assess grazing conditions for reindeer *Ann. Glaciol.* 54 doi: 10.3189/2013AoG62A306
- Vikhamar-Schuler D, Isaksen K, Haugen J E et al. 2016 Changes in winter warming events in the Nordic Arctic Region *J. Climate* 29 6223–6244
- Wang C, Graham RM, Wang K, Gerland S. and Granskog M A 2019 Comparison of ERA5 and ERA-Interim near-surface air temperature, snowfall and precipitation over Arctic sea ice: effects on sea ice thermodynamics and evolution *The Cryosphere* 13, 1661–1679, <https://doi.org/10.5194/tc-13-1661-2019>
- Wang L, Derksen C, Brown R, Markus T 2013 Recent changes in pan-Arctic melt onset from satellite passive microwave measurements *Geophys. Res. Lett.* 40, 522-528, <https://doi.org/10.1002/grl.50098>
- Westermann S, Boike J, Langer M, Schuler T V, Etzelmüller B 2011 Modeling the impact of wintertime rain events on the thermal regime of permafrost *Cryosphere* 5 945–59
- Wilson R, Bartsch A, Joly K, Reynolds J H, Orlando A, Loya W M 2013 Frequency, timing, extent, and size of winter thaw-refreeze events in Alaska 2001-2008 detected by remotely sense microwave backscatter data *Polar Biology* 36 419–426 doi 10.1007/s00300-012-1272-6