computational complexity

RANK AND BORDER RANK OF KRONECKER POWERS OF TENSORS AND STRASSEN'S LASER METHOD

AUSTIN CONNER, FULVIO GESMUNDO, JOSEPH M. LANDSBERG, EMANUELE VENTURA

Abstract. We prove that the border rank of the Kronecker square of the little Coppersmith–Winograd tensor $T_{cw,q}$ is the square of its border rank for q > 2 and that the border rank of its Kronecker cube is the cube of its border rank for q > 4. This answers questions raised implicitly by Coppersmith & Winograd (1990, §11) and explicitly by Bläser (2013, Problem 9.8) and rules out the possibility of proving new upper bounds on the exponent of matrix multiplication using the square or cube of a little Coppersmith–Winograd tensor in this range.

In the positive direction, we enlarge the list of explicit tensors potentially useful for Strassen's laser method, introducing a skew-symmetric version of the Coppersmith–Winograd tensor, $T_{skewcw,q}$. For q=2, the Kronecker square of this tensor coincides with the 3×3 determinant polynomial, $\det_3\in\mathbb{C}^9\otimes\mathbb{C}^9$, regarded as a tensor. We show that this tensor could potentially be used to show that the exponent of matrix multiplication is two.

We determine new upper bounds for the (Waring) rank and the (Waring) border rank of det_3 , exhibiting a strict submultiplicative behaviour for $T_{skewcw,2}$ which is promising for the laser method.

We establish general results regarding border ranks of Kronecker powers of tensors, and make a detailed study of Kronecker squares of tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$.

Keywords. Matrix multiplication complexity, Tensor rank, Asymptotic rank, Laser method.

Subject classification. 68Q17; 14L30; 15A69.

1. Introduction

The exponent ω of matrix multiplication is defined as

$$\omega := \inf \left\{ \tau \; \middle| \; \begin{array}{l} \text{two } \mathbf{n} \times \mathbf{n} \text{ matrices may be multiplied} \\ \text{using } O(\mathbf{n}^{\tau}) \text{ arithmetic operations} \end{array} \right\}.$$

1 Page 2 of 40 Conner et al. cc

This is a fundamental constant governing the complexity of the basic operations in linear algebra. It is conjectured that $\omega=2$. There is a classical upper bound $\omega\leq 3$ following from the standard row-by-column multiplication. Starting from the work of Strassen (1969), a great deal of effort has been spent on the research on upper bounds on the exponent, involving methods from combinatorics, probability, and statistical mechanics; we refer to Section 1.4 for a brief history. More recently, Cohn & Umans (2003) uses group-theoretic techniques and in particular the Fourier-transform of finite groups. In this work, we approach the problem via algebraic geometry and representation theory. We obtain both negative and hopeful results.

Our focus will be on Strassen's laser method, see Strassen (1987). This technique was used to achieve Strassen's upper bound of 1988 and essentially all subsequent upper bounds. In order to present the method and our contributions, we adopt the language of tensors.

1.1. Definitions. Let A, B, C be complex vector spaces. A tensor $T \in A \otimes B \otimes C$ has rank one if $T = a \otimes b \otimes c$ for some $a \in A, b \in B, c \in C$. The rank of T, denoted $\mathbf{R}(T)$, is the smallest r such that T is sum of r rank-one tensors. The border rank of T, denoted $\mathbf{R}(T)$, is the smallest r such that T is the limit of a sequence of rank r tensors.

A tensor $T \in A \otimes B \otimes C$ defines a bilinear map $A^* \times B^* \to C$ and a trilinear map $A^* \times B^* \times C^* \to \mathbb{C}$. The matrix multiplication tensor $M_{\mathbf{l},\mathbf{m},\mathbf{n}}$ is the tensor associated to the bilinear map

$$M_{\langle \mathbf{l}, \mathbf{m}, \mathbf{n} \rangle} : \mathrm{Mat}_{\mathbf{l} \times \mathbf{m}} \times \mathrm{Mat}_{\mathbf{m} \times \mathbf{n}} \to \mathrm{Mat}_{\mathbf{l} \times \mathbf{n}}$$

sending a pair of matrices (X,Y) to their product XY. As a trilinear map, the matrix multiplication tensor is $M_{\langle \mathbf{l},\mathbf{m},\mathbf{n}\rangle}(X,Y,Z) = \operatorname{trace}(XYZ)$, where X,Y,Z are matrices of size $\mathbf{l}\times\mathbf{m}$, $\mathbf{m}\times\mathbf{n}$ and $\mathbf{n}\times\mathbf{l}$, respectively. The matrix multiplication tensor has the following important self-reproducing property: $M_{\langle \mathbf{l},\mathbf{m},\mathbf{n}\rangle}\boxtimes M_{\langle \mathbf{l}',\mathbf{m}'\mathbf{n}'\rangle} = M_{\langle \mathbf{l}',\mathbf{m}\mathbf{m}',\mathbf{n}\mathbf{n}'\rangle}$. Write $M_{\langle \mathbf{n}\rangle} := M_{\langle \mathbf{n},\mathbf{n},\mathbf{n}\rangle}$.

The complexity of performing a bilinear map, and in particular the complexity of matrix multiplication, is controlled by the tensor rank of the corresponding tensor. Bini (1980) showed that border rank controls the complexity as well:

$$\omega = \inf \{ \tau : \underline{\mathbf{R}}(M_{\langle \mathbf{n} \rangle}) \in O(\mathbf{n}^{\tau}) \}.$$

Let GL(A) be the general linear group of invertible linear maps $A \to A$ and similarly for B and C. We say that two tensors are *isomorphic* if they are in the same orbit under the natural action of $GL(A) \times GL(B) \times GL(C)$ on $A \otimes B \otimes C$. We will often assume that all tensors involved in the discussion belong to the same space $A \otimes B \otimes C$. This is not restrictive, since we may re-embed the spaces A, B, C into larger spaces whenever it is needed.

Given $T, T' \in A \otimes B \otimes C$, we say that T degenerates to T' whenever $T' \in \overline{GL(A) \times GL(B) \times GL(C) \cdot T}$, the closure of the orbit of T, equivalently in the Euclidean or in the Zariski topology. Border rank is semicontinuous under degeneration: $\mathbf{R}(T') \leq \mathbf{R}(T)$ if T degenerates to T'.

Border rank may be rephrased in terms of degeneration as follows. For a tensor T, one has $\underline{\mathbf{R}}(T) \leq r$ if and only if T is a degeneration of $M_{\langle 1 \rangle}^{\oplus r} = \sum_{i=1}^{r} a_i \otimes b_i \otimes c_i$, where $\{a_i\}$ is a set of linearly independent vectors and similarly for $\{b_i\}$ and $\{c_i\}$. The border subrank of T, denoted $\underline{\mathbf{Q}}(T)$, is the maximum q such that T degenerates to $M_{\langle 1 \rangle}^{\oplus q}$.

For tensors $T \in A \otimes B \otimes C$ and $T' \in A' \otimes B' \otimes C'$, the Kronecker product of T and T' is the tensor $T \boxtimes T' := T \otimes T' \in (A \otimes A') \otimes (B \otimes B') \otimes (C \otimes C')$, regarded as 3-way tensor. Given $T \in A \otimes B \otimes C$, the Kronecker powers of T are $T^{\boxtimes N} \in A^{\otimes N} \otimes B^{\otimes N} \otimes C^{\otimes N}$, defined iteratively. Rank and border rank are submultiplicative under Kronecker product: $\mathbf{R}(T \boxtimes T') \leq \mathbf{R}(T)\mathbf{R}(T')$, $\mathbf{R}(T \boxtimes T') \leq \mathbf{R}(T)\mathbf{R}(T')$, and both inequalities may be strict.

Asymptotic versions of border rank and border subrank, respectively, called asymptotic rank and asymptotic subrank, are defined as follows:

$$\underline{\mathbf{R}}(T) = \lim_{N \to \infty} [\underline{\mathbf{R}}(T^{\boxtimes N})]^{1/N}, \quad \underline{\mathbf{Q}}(T) = \lim_{N \to \infty} [\underline{\mathbf{Q}}(T^{\boxtimes N})]^{1/N}.$$

One has $\omega = \log_2(\mathbf{R}(M_{\langle 2 \rangle}))$; in particular $\omega = 2$ if and only if $\mathbf{R}(M_{\langle \mathbf{n} \rangle}) = \mathbf{n}^2$ for any (and as a consequence all) \mathbf{n} .

1.2. Strassen's laser method and its barriers. The two fundamental ingredients of Strassen's laser method are submultiplicativity of border rank under Kronecker powers and semicontinuity of border rank under degeneration. The laser method relies on an auxiliary tensor T with the property that $\underline{\mathbf{R}}(T)$ is small, and for some large N, $T^{\boxtimes N}$ degenerates to a large matrix multiplication tensor.

Since 1987, only three tensors have been employed in the method and the best upper bounds so far come from the big Coppersmith-Winograd tensor, see Coppersmith & Winograd (1990):

$$T_{CW,q} := \sum_{j=1}^{q} a_0 \otimes b_j \otimes c_j + a_j \otimes b_0 \otimes c_j + a_j \otimes b_j \otimes c_0$$

+ $a_0 \otimes b_0 \otimes c_{q+1} + a_0 \otimes b_{q+1} \otimes c_0 + a_{q+1} \otimes b_0 \otimes c_0 \in (\mathbb{C}^{q+2})^{\otimes 3},$

It was used to prove $\omega < 2.38$ in 1988 and all further improvements to the current best known upper bound $\omega < 2.373$.

Ambainis et al. (2015) gave an explanation for the limited progress since 1988, followed by further explanations in Alman (2019); Alman & Williams

1 Page 4 of 40 Conner *et al.* cc

(2018a,b); Christandl et al. (2021). One major consequence of these results is that $T_{CW,q}$ cannot be used to prove $\omega < 2.3$ using the standard laser method.

A geometric identification of the barrier of Ambainis *et al.* (2015) was given in Christandl *et al.* (2021). Strassen showed $\underline{\mathbf{Q}}(M_{\langle \mathbf{n} \rangle}) \geq \lceil \frac{3}{4}n^2 \rceil$; in Kopparty *et al.* (2020, Theorem 3) equality was proved. This, together with the self-reproducing property of the matrix multiplication tensor, implies that $\underline{\mathbf{Q}}(M_{\langle \mathbf{n} \rangle}) = \mathbf{n}^2$, which is the maximum possible value. A consequence is that no tensor having non-maximal asymptotic subrank can be used to prove $\omega = 2$ via the laser method; Strassen (1991) proved that $\underline{\mathbf{Q}}(T_{CW,q})$ is non-maximal.

The second most effective tensor used for upper bounds via Strassen's laser method is the *small Coppersmith-Winograd tensor*:

$$(1.1) T_{cw,q} := \sum_{j=1}^{q} a_0 \otimes b_j \otimes c_j + a_j \otimes b_0 \otimes c_j + a_j \otimes b_j \otimes c_0 \in (\mathbb{C}^{q+1})^{\otimes 3}.$$

In Coppersmith & Winograd (1990), the laser method was used to (implicitly) prove the following result:

Theorem 1.2 (Coppersmith & Winograd 1990). For all k and q,

(1.3)
$$\omega \leq \log_q(\frac{4}{27}(\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes k}))^{\frac{3}{k}}).$$

One has $\mathbf{R}(T_{cw,q}) = q+2$, which is one more than minimal, see, e.g., Bürgisser et al. (1997, Sec. 15.8). Applying Theorem 1.2 to $T_{cw,8}$ with k=1 one obtains $\omega \leq 2.41$, see Coppersmith & Winograd (1990). Theorem 1.2 implies that if the border rank of the Kronecker square or some higher Kronecker power of $T_{cw,q}$ were strictly submultiplicative, one could get a better bound, and one could even potentially prove $\omega = 2$ using Kronecker powers of $T_{cw,2}$. Indeed, by Bürgisser et al. (1997, Ex. 15.24) one sees that Theorem 1.2 holds replacing $\mathbf{R}(T_{cw,q}^{\boxtimes k})^{\frac{1}{k}}$ with $\mathbf{R}(T_{cw,q})$. In particular, were $\mathbf{R}(T_{cw,2}) = 3$, then Theorem 1.2 would imply $\omega = 2$. This shows that the barriers of Alman & Williams (2018a,b); Ambainis et al. (2015); Christandl et al. (2021) do not apply to $T_{cw,2}$. Previous to our work, the possibility to prove the upper bound $\omega < 2.3$ using the second and third Kronecker power of $T_{cw,q}$ for $3 \leq q \leq 10$ was open, in the sense that the if the state-of-the-art lower bound on $T_{cw,q}^{\boxtimes k}$ were equal to an upper bound, then Theorem 1.2 would have given an improvement. We show that this is not the case.

1.3. Main results. Bläser (2013, Problem 9.8) posed the problem of determining the border rank of $T_{cw,q}^{\boxtimes 2}$. We show:

cc

Theorem 1.4. For all q > 2, $\mathbf{R}(T_{cw,q}^{\boxtimes 2}) = (q+2)^2$; moreover $15 \leq \mathbf{R}(T_{cw,2}^{\boxtimes 2}) \leq 16$.

For all q > 4, $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 3}) = (q+2)^3$; if q = 3, 4 then $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 3}) \ge (q+2)^2(q+1)$; if q = 2, then $\underline{\mathbf{R}}(T_{cw,2}^{\boxtimes 3}) \ge 15 \cdot 3$.

For all q > 4 and all N, $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes N}) \ge (q+1)^{N-3}(q+2)^3$; if q = 3, 4 then $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes N}) \ge (q+2)^2 \cdot (q+1)^{N-2}$; if q = 2 then $\underline{\mathbf{R}}(T_{cw,2}^{\boxtimes 3}) \ge 15 \cdot 3^{N-2}$.

This improves on the previous lower bound from Bläser & Lysikov (2016), which was $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes N}) \geq (q+1)^N + 2^N - 1$ for all q, N.

This result shows that the second and third Kronecker powers of $T_{cw,q}$ cannot give any improvement on the current upper bounds on the exponent. For instance, the lower bound of Bläser & Lysikov (2016) for (q, N) = (3, 3) is $\underline{\mathbf{R}}(T_{cw,3}^{\boxtimes 3}) \geq 71$; if this had been the value of $\underline{\mathbf{R}}(T_{cw,3}^{\boxtimes 3})$ then Theorem 1.2 would give $\omega < 2.15$; however, the lower bound of Theorem 1.4 guarantees $\underline{\mathbf{R}}(T_{cw,3}^{\boxtimes 3}) \geq$ 100, and even if this turns out to be the value of $\underline{\mathbf{R}}(T_{cw,3}^{\boxtimes 3})$, Theorem 1.4 only gives $\omega < 2.46$.

In light of the above-mentioned barriers and Theorem 1.4, one might try to determine better tensors which are not subject to the barriers (similarly to $T_{cw,q}$) and at the same time have strict submultiplicativity of border rank under Kronecker powers.

Inspired by Conner et al. (2019a), we introduce a new family of tensors, which are a skew-symmetric version of the small Coppersmith-Winograd tensors for every even q:

(1.5)
$$T_{skewcw,q} := \sum_{j=1}^{q} a_0 \otimes b_j \otimes c_j + \sum_{j=1}^{q} a_j \otimes b_0 \otimes c_j + \sum_{j=1}^{\frac{q}{2}} (a_{\xi} \otimes b_{\xi + \frac{q}{2}} - a_{\xi + \frac{q}{2}} \otimes b_{\xi}) \otimes c_0 \in (\mathbb{C}^{q+1})^{\otimes 3}.$$

Proposition 2.2 shows Theorem 1.2 holds with $T_{cw,q}$ is replaced by $T_{skewcw,q}$, so in particular $T_{skewcw,2}$ could potentially be used to prove $\omega=2$.

Proposition 3.2 contains more negative news: $\mathbf{R}(T_{skewcw,q}) \geq q + 3$, and in particular $\underline{\mathbf{R}}(T_{skewcw,2}) = 5$. However, we show a strong submultiplicative behaviour for $T_{skewcw,q}$, namely $\mathbf{R}(T_{skewcw,2}^{\boxtimes 2}) \leq 17 < 5^2$. Theorem 1.6 below actually proves a stronger statement. We show in Lemma 2.8 that $T_{skewcw,2}^{\boxtimes 2}$ is isomorphic to the 3×3 determinant polynomial regarded as a tensor and we prove new upper bounds for the symmetric rank (also known as Waring rank, see, e.g., Landsberg (2012, §2.6.6)) and symmetric border rank of the 3×3 determinant polynomial.

1 Page 6 of 40 Conner et al. cc

Theorem 1.6. Let $det_3 \in \mathbb{C}^9 \otimes \mathbb{C}^9 \otimes \mathbb{C}^9$ be the 3×3 determinant polynomial regarded as a symmetric tensor. Then

$$\mathbf{R}_S(\det_3) \le 18$$
, and $\mathbf{R}_S(\det_3) \le 17$.

In Conner *et al.* (2019b), it was shown that $\underline{\mathbf{R}}(\det_3) = 17$ and in particular the second inequality in Theorem 1.6 is an equality.

The proof of Theorem 1.4 is given in Section 3 and the proof of Theorem 1.6 is given in Section 4.

Some of the proofs of this work rely on computer calculations performed by the software Macaulay2 (Grayson & Stillman 2020) and Sage (Sage Developers 2020). The scripts performing these calculations are collected in different appendices in Supplementary Material available at

1.4. Brief history of upper bounds. There was steady progress in the research for upper bounds on ω from 1968 to 1988.

Strassen (1969) gave an algorithm to perform 2×2 matrix multiplication using 7 scalar multiplications (rather than the 8 multiplications of the standard algorithm). This gives the upper bound $\omega < 2.81$. Bini *et al.* (1980), using border rank, showed $\omega < 2.78$.

A major breakthrough due to Schönhage (1981), known as the asymptotic sum inequality, was used to show $\omega < 2.55$ by exploiting the interplay between direct sums and the self-reproducing property of the matrix multiplication tensor. Strassen (1987) introduced the *laser method* and showed $\omega < 2.48$. A refined form of the laser method was used by Coppersmith and Winograd to show $\omega < 2.3755$, see Coppersmith & Winograd (1990).

There was no progress on upper bounds on the exponent until 2010 when, via a further refinement of the method, a series of improvements Stothers (2010), Williams (2012), Le Gall (2014) and Alman & Williams (2021) lowered the upper bound to the current state of the art $\omega < 2.373$.

2. Preliminary results

In this section, we provide some results which will be useful in the rest of the paper.

The following remark shows that submultiplicativity holds asymptotically for most tensors; this appeared implicitly in Strassen (1988, Lemma 3.5) and then explicitly in Christandl *et al.* (2021, Prop. 2.12).

REMARK 2.1. Any $T \in \mathbb{C}^m \otimes \mathbb{C}^m \otimes \mathbb{C}^m$ is a degeneration of $M_{\langle 1,m,m\rangle} \in \mathbb{C}^{m^2} \otimes \mathbb{C}^m \otimes \mathbb{C}^m$, so $T^{\boxtimes 3}$ is a degeneration of $M_{\langle m^2,m^2,m^2\rangle} = M_{\langle 1,m,m\rangle} \boxtimes M_{\langle m,1,m\rangle} \boxtimes M_{\langle m,m,1\rangle}$. In particular $\mathbf{R}(T^{\boxtimes 3}) \leq \mathbf{R}(M_{\langle m^2,m^2,m^2\rangle}) = m^{2\omega}$ and therefore $\mathbf{R}(T) \leq m^{2\omega/3}$. Since $\omega < 2.4$, we have $\mathbf{R}(T) \leq m^{1.6}$.

2.1. $T_{skewcw,q}$ and the laser method. The first result is the analog of Theorem 1.2 for the family $T_{skewcw,q}$:

Proposition 2.2. For all k,

(2.3)
$$\omega \leq \log_q(\frac{4}{27}(\mathbf{R}(T_{skewcw,q}^{\boxtimes k}))^{\frac{3}{k}}).$$

PROOF. Similarly to the case of $T_{cw,q}$, the proof follows immediately from Bürgisser *et al.* (1997, Theorem 15.41), because $T_{skewcw,q}$ has the same "block structure" as $T_{cw,q}$.

In particular, similarly to $T_{cw,q}$, if $\Re(T_{skewcw,2}) = 3$ then $\omega = 2$, and it is potentially possible to improve the current upper bounds on ω using $T_{skewcw,q}$. Therefore, it is important to determine upper bounds on the border rank of the Kronecker powers of $T_{skewcw,q}$, and in particular in the case q = 2.

2.2. Coppersmith–Winograd tensors, symmetries, determinants, and permanents. Let $S^3\mathbb{C}^m$ and $\Lambda^3\mathbb{C}^m$ respectively denote the subspaces of symmetric and skew-symmetric tensors in $\mathbb{C}^m \otimes \mathbb{C}^m \otimes \mathbb{C}^m$. By identifying the three copies of \mathbb{C}^{q+1} in (1.1) and (1.5), we observe that $T_{cw,q}$ is isomorphic to a symmetric tensor and $T_{skewcw,q}$ is isomorphic to a skew-symmetric tensor. Indeed, fixing a basis a_0, \ldots, a_q of \mathbb{C}^{q+1} , the isomorphism $a_j \leftrightarrow b_j \leftrightarrow c_j$ provides

(2.4)
$$T_{cw,q} = a_0(a_1^2 + \dots + a_q^2) \in S^3 \mathbb{C}^{q+1}.$$

Similarly, if q = 2u is even, the isomorphism

$$a_0 \leftrightarrow -b_0 \leftrightarrow c_0$$

$$a_j \leftrightarrow b_j \leftrightarrow -c_{u+j} \quad j = 1, \dots, u$$

$$a_{u+j} \leftrightarrow b_{u+j} \leftrightarrow c_j \quad j = 1, \dots, u$$

provides

$$(2.5) T_{skewcw,q} = a_0 \wedge (a_1 \wedge a_{u+1} + \dots + a_u \wedge a_q) \in \Lambda^3 \mathbb{C}^{q+1}.$$

We introduce some definitions concerning the symmetries of a tensor. The group homomorphism $\Phi: GL(A) \times GL(B) \times GL(C) \rightarrow GL(A \otimes B \otimes C)$

1 Page 8 of 40 Conner et al. cc

defining the natural action on $A \otimes B \otimes C$ has a two dimensional kernel ker $\Phi = \{(\lambda \operatorname{Id}_A, \mu \operatorname{Id}_B, \nu \operatorname{Id}_C) : \lambda \mu \nu = 1\} \simeq (\mathbb{C}^*)^2$.

In particular, the group $(GL(A) \times GL(B) \times GL(C)) / (\mathbb{C}^*)^{\times 2}$ is identified with a subgroup of $GL(A \otimes B \otimes C)$. Given $T \in A \otimes B \otimes C$, the symmetry group of a tensor T is the stabilizer of T in $(GL(A) \times GL(B) \times GL(C)) / (\mathbb{C}^*)^{\times 2}$, that is

(2.6)
$$G_T := \{ g \in (GL(A) \times GL(B) \times GL(C)) / (\mathbb{C}^*)^{\times 2} \mid g \cdot T = T \}.$$

If the three spaces A,B,C are identified, so that $A\otimes B\otimes C\simeq A^{\otimes 3}$, one can consider the action restricted to GL(A) embedded diagonally as $GL^{diag}(A)\subseteq GL(A)^{\times 3}$. In this case, the kernel of the action reduces to the cyclic group $\mathbb{Z}_3=\{\zeta \operatorname{Id}_A: \zeta^3=1\}$ and one can consider a restricted version of the symmetry group

$$G_T^s := G_T \cap GL^{diag}(A) = \{ g \in GL(A)/\mathbb{Z}_3 \mid g \cdot T = T \}.$$

Let \mathfrak{S}_k be the permutation group on k elements.

We record the following observation:

Proposition 2.7. Let $T \in A \otimes B \otimes C$ (resp. $T \in A^{\otimes 3}$). Then

$$G_{T^{\boxtimes N}}\supseteq G_T^{\times N}\rtimes \mathfrak{S}_N\quad (\text{resp. }G_{T^{\boxtimes N}}^s\supseteq G_T^{s\times N}\rtimes \mathfrak{S}_N)$$

where the symmetric group acts by permuting the factors of the direct product.

PROOF. Let $T \in A \otimes B \otimes C$. Every factor G_T in $G_T^{\times N} \rtimes \mathfrak{S}_N$ acts on a single factor of $T^{\boxtimes N}$ and it stabilizes it by definition of G_T . The groups \mathfrak{S}_N permutes the factors of $T^{\boxtimes N}$, which is a Kronecker power, and therefore, it is stabilized.

The statement for $T \in A^{\otimes 3}$ is an immediate consequence.

Consider the action of the symmetric group \mathfrak{S}_3 which permutes the tensor factors. A tensor is symmetric if it is invariant under this action and skew-symmetric if it is skew-invariant. It is easy to observe that Kronecker powers of symmetric tensors are symmetric tensors. Moreover, odd Kronecker powers of skew-symmetric tensors are skew-symmetric and even Kronecker powers of skew-symmetric tensors are symmetric.

We record the expressions of the 3×3 permanent and determinant polynomials as tensors in $\mathbb{C}^9\otimes\mathbb{C}^9\otimes\mathbb{C}^9$. Write $(-1)^{\sigma}$ for the sign of a permutation σ . Then

$$\det_{3} = \frac{1}{6} \sum_{\sigma, \tau \in \mathfrak{S}_{3}} (-1)^{\sigma \tau} a_{\sigma(1)\tau(1)} \otimes b_{\sigma(2)\tau(2)} \otimes c_{\sigma(3)\tau(3)},$$
$$\operatorname{perm}_{3} = \frac{1}{6} \sum_{\sigma, \tau \in \mathfrak{S}_{3}} a_{\sigma(1)\tau(1)} \otimes b_{\sigma(2)\tau(2)} \otimes c_{\sigma(3)\tau(3)}.$$

Lemma 2.8. We have the following isomorphisms of tensors:

$$T_{cw,2}^{\boxtimes 2} \cong \text{perm}_3,$$

 $T_{skewcw,2}^{\boxtimes 2} \cong \text{det}_3.$

From (2.4), we have $T_{cw,2} = a_0(a_1^2 + a_2^2)$. Let $\tilde{a}_1 = (a_1 + \sqrt{-1}a_2)$ and $\tilde{a}_2 = (a_1 - \sqrt{-1}a_2)$, so that $T_{cw,2} = a_0 \tilde{a}_1 \tilde{a}_2$. This shows that after a suitable change of basis $T_{cw,2} = a_0 a_1 a_2$. Its symmetry group is $G_{T_{cw,2}}^s = \mathbb{T}_3^{SL} \rtimes \mathfrak{S}_3$, where \mathbb{T}_3^{SL} denotes the torus of diagonal matrices with determinant one, and \mathfrak{S}_3 acts permuting the three basis elements.

By Proposition 2.7, we deduce that $T_{cw,2}^{\boxtimes 2}$ is a symmetric tensor, with $G_{T_{cw,2}^{\boxtimes 2}}^{s} \supseteq$ $(\mathbb{T}_3^{SL} \rtimes \mathfrak{S}_3)^{\times 2} \rtimes \mathbb{Z}_2$ (and in fact equality holds). This is the stabilizer of the permanent polynomial perm₃. Since the permanent is characterized by its stabilizer (see, e.g., Lemma 2.9 below), we conclude.

The proof for $T_{skewcw,2}$ is similar. From (2.5), we have $T_{skewcw,2} = a_0 \wedge a_1 \wedge a_2$. Therefore $G_{T_{skewcw,2}}^s = SL_3$; indeed $T_{skewcw,2}$ is the unique, up to scale, SL_3 invariant in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$.

By Proposition 2.7, we deduce that $T_{skewcw,2}^{\boxtimes 2}$ is a symmetric tensor, with $G_{T_{skewcw,2}^{\boxtimes 2}}^{Skewcw,2} \supseteq (SL_3)^{\times 2} \rtimes \mathbb{Z}_2$ (and in fact equality holds). This is the stabilizer of the determinant polynomial det₃. Since the determinant is characterized by its stabilizer, we conclude.

The symmetric tensors \det_m and perm_m are characterized by their stabilizers. For the determinant, this fact is classical. For the permanent, the statement, but not the proof, appears in Mulmuley & Sohoni (2008). For completeness, we provide a proof here assuming some familiarity with the representation theory of SL_m and of the symmetric group \mathfrak{S}_m .

LEMMA 2.9. Let $T \in S^m(\mathbb{C}^m \otimes \mathbb{C}^m)$ be a symmetric tensor of order m. If $G_T^s \supseteq (\mathbb{T}^{SL} \rtimes \mathfrak{S}_m)^{\times 2} \rtimes \mathbb{Z}_2$, then $T = \operatorname{perm}_m$, up to scale. If $G_T^s \supseteq (SL_m^{\times 2}) \rtimes \mathbb{Z}_2$, then $T = \det_m$, up to scale.

First consider the case of the determinant. Let $SL_m \times SL_m =$ $SL(E) \times SL(F)$ act on $S^m(E \otimes F)$. This space decomposes as $SL(E) \times SL(F)$ representation as (see, e.g., Landsberg (2012, §6.7.6))

$$S^{m}(E \otimes F) = \bigoplus_{|\pi|=m} S_{\pi}E \otimes S_{\pi}F;$$

this is multiplicity free, with the only trivial module $S_{(1^m)}E\otimes S_{(1^m)}F=\Lambda^m E\otimes$ $\Lambda^m F$. This is the space spanned by \det_m .

In the case of the permanent, note that the decomposition above holds for the action of $(\mathbb{T}^{SL} \rtimes \mathfrak{S}_m)^{\times 2}$ as well. Then, the $\mathbb{T}^{\tilde{SL}(E)} \times \mathbb{T}^{SL(F)}$ -invariant subspace is given by the sum of the weight zero spaces $(S_{\pi}E)_0 \otimes (S_{\pi}F)_0$. By Gay (1976), one has the isomorphism $(S_{\pi}E)_0 \otimes (S_{\pi}F)_0 = [\pi]_E \otimes [\pi]_F$ for the weight zero spaces as $\mathfrak{S}_E \times \mathfrak{S}_F$ -modules. The only trivial representation is the one corresponding to $\pi = (d)$, which is the subspace spanned by perm_m.

Lemma 2.8 guarantees that perm₃ and det₃ are tensors not subject to barriers for the laser method. In particular, either $\mathbf{R}(\det_3) = 9$ or $\mathbf{R}(\mathrm{perm}_3) = 9$ would imply $\omega = 2$.

REMARK 2.10. A similar result holds for higher Kronecker powers. For every k, the even power $T_{skewcw,2}^{\boxtimes 2k}$ is invariant under $SL_3^{\times 2k} \rtimes \mathfrak{S}_{2k}$. There is a unique invariant PasDet_{k,3} for $SL_3^{\times 2k}$ in $S^3((\mathbb{C}^3)^{\otimes 2k})$: it is the generator of the submodules $(\Lambda^3\mathbb{C}^3)^{\otimes 2k}$, known as the Pascal determinant (see, e.g., Landsberg (2012, §8.3)). If any of the Pascal determinants has minimal asymptotic rank, i.e., $\mathfrak{R}(\operatorname{PasDet}_{k,3}) = 3^{2k}$, then $\omega = 2$.

REMARK 2.11. One can regard the 3×3 determinant and permanent as trilinear maps $\mathbb{C}^3 \times \mathbb{C}^3 \times \mathbb{C}^3 \to \mathbb{C}$, where the three copies of \mathbb{C}^3 are the first, second and third column of a 3×3 matrix. From this point of view, the trilinear map given by the determinant is $T_{skewcw,2}$ as a tensor and the one given by the permanent is $T_{cw,2}$ as a tensor. This perspective, combined with the notion of product rank (in the sense of Ilten & Teitler (2016)) provides the upper bounds $\mathbf{R}_S(\text{perm}_3) \leq 16$ and $\mathbf{R}(\text{det}_3) \leq 20$. These bounds already appeared in Derksen (2016); Ilten & Teitler (2016) and are also a consequence of Lemma 2.8.

2.3. Generic tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$. It is a classical fact that a generic tensor in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ has border rank five, see, e.g., Strassen (1983).

Remark 2.12. Computer experiments indicate that for all $T \in \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$, $\mathbf{R}(T^{\boxtimes 2}) \leq 22 < 25$.

The evidence for the remark is obtained as follows. We considered tensors $T \in \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ whose coefficients in a fixed basis were taken independently and uniformly random in [-1,1]. We obtained numerically that $\underline{\mathbf{R}}(T^{\boxtimes 2}) \leq 22$. An instance of this computation is available in Appendix A of Supplementary Material.

PROBLEM 2.13. Prove the claim in Remark 2.12. Even better, give a geometric proof.

Remark 2.12 is not too surprising because $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ is secant defective, in the sense that by a dimension count, one would expect the maximum border

rank of a tensor to be 4, but the actual maximum is 5. This means that for a generic tensor, there is a 8 parameter family of rank 5 decompositions, and it is not surprising that the naïve 64-parameter family of decompositions of the square might have decompositions of lower border rank on the boundary.

3. Koszul flattenings and lower bounds for Kronecker powers

In this section we review Koszul flattenings and prove a result on propagation of Koszul flattening lower bounds under Kronecker products. We will use Koszul flattenings to prove $\underline{\mathbf{R}}(T_{skewcw,a}) \geq q+3$ in Proposition 3.2. Moreover, we prove Theorem 1.4: the proof will follow from Theorem 3.5, Theorem 3.7 and Corollary 3.8.

Fix bases $\{a_i\}$, $\{b_i\}$, $\{c_k\}$ of the vector spaces A, B, C, respectively; fix an integer p. Given a tensor $T = \sum_{ijk} T^{ijk} a_i \otimes b_j \otimes c_k \in A \otimes B \otimes C$, the p-th Koszul flattening of T on the space A is the linear map

$$T_A^{\wedge p}: \Lambda^p A \otimes B^* \to \Lambda^{p+1} A \otimes C$$
$$X \otimes \beta \mapsto \sum_{ijk} T^{ijk} \beta(b_j)(a_i \wedge X) \otimes c_k.$$

Then Landsberg & Ottaviani (2013, Proposition 4.1.1) states

(3.1)
$$\underline{\mathbf{R}}(T) \ge \frac{\operatorname{rank}(T_A^{\wedge p})}{\binom{\dim(A)-1}{p}}.$$

This type of lower bound has a long history. More generally, one considers an embedding of the space $A \otimes B \otimes C$ into a large space of matrices. Then if a rank-one tensor maps to a rank q matrix, a rank r tensor maps to a rank at most rq matrix, so the size rq+1 minors give equations testing for border rank r. In this case, the size of the matrices is $\binom{\mathbf{a}}{p}\mathbf{b}\times\binom{\mathbf{a}}{p+1}\mathbf{c}$ and a rank-one tensor maps to a matrix of rank $\binom{\mathbf{a}-1}{p}$. Here $\mathbf{a}=\mathrm{dim}A,\ \mathbf{b}=\mathrm{dim}B$ and $\mathbf{c}=\mathrm{dim}C$.

In practice, one considers a subspace $A'^* \subseteq A^*$ of dimension 2p+1 and restricts T (considered as a trilinear form) to $A'^* \times B^* \times C^*$ to get an optimal bound, so the denominator $\binom{\dim(A)-1}{p}$ is replaced by $\binom{2p}{p}$ in (3.1). Equivalently, one considers a linear map $\phi: A \to A'$ and the corresponding Koszul flattening map gives a lower bound for $\mathbf{R}(\phi(T))$, which, by linearity, is a lower bound for $\mathbf{R}(T)$.

The case p=1 is a straightening of Strassen's equations Strassen (1983). There are numerous expositions of Koszul flattenings and their generalizations, see, e.g., Landsberg (2012, §7.3), Ballico et al. (2019, §7.2), Derksen & Makam (2018), Landsberg (2017, §2.4), or Efremenko et al. (2018).

We use Koszul flattenings to give the following lower bound on $\underline{\mathbf{R}}(T_{skewcq,q})$:

1 Page 12 of 40 Conner et al. cc

Proposition 3.2. For every even $q \geq 2$, $\underline{\mathbf{R}}(T_{skewcw,q}) \geq q + 3$.

PROOF. Write q = 2u. Fix a space $A' = \langle e_0, e_1, e_2 \rangle$. Define $\phi : A \to A'$ by

$$\phi(a_0) = e_0,$$

 $\phi(a_i) = e_1 \text{ for } i = 1, \dots, u,$
 $\phi(a_s) = e_2 \text{ for } s = u + 1, \dots, q.$

As an element of $\Lambda^3 A \subseteq A \otimes A \otimes A$, we have $T_{skewcw,q} = a_0 \wedge \sum_{i=1}^u a_i \wedge a_{u+i}$ as in (2.5).

We prove that for $T = (\phi \otimes \operatorname{Id}_B \otimes \operatorname{Id}_C)(T_{skewcw,q}) \in A' \otimes B \otimes C$, one obtains $\operatorname{rank}(T_{A'}^{\wedge 1}) = 2(q+2)+1$. This provides the lower bound $\underline{\mathbf{R}}(T) \geq \left\lceil \frac{2(q+2)+1}{2} \right\rceil = q+3$.

We record the images via $T_{A'}^{\wedge 1}$ of a basis of $A' \otimes B^*$. Fix the range of $i = 1, \ldots, u$:

$$T_{A'}^{\wedge 1}(e_{0} \otimes \beta_{0}) = (e_{0} \wedge e_{1}) \otimes \sum_{i=1}^{u} c_{u+i} - (e_{0} \wedge e_{2}) \otimes \sum_{i=1}^{u} c_{i},$$

$$T_{A'}^{\wedge 1}(e_{0} \otimes \beta_{i}) = (e_{0} \wedge e_{2}) \otimes c_{0},$$

$$T_{A'}^{\wedge 1}(e_{0} \otimes \beta_{u+i}) = (e_{0} \wedge e_{1}) \otimes c_{0},$$

$$T_{A'}^{\wedge 1}(e_{1} \otimes \beta_{0}) = (e_{1} \wedge e_{2}) \otimes \sum_{i=1}^{u} c_{u+i},$$

$$T_{A'}^{\wedge 1}(e_{1} \otimes \beta_{i}) = (e_{0} \wedge e_{1}) \otimes c_{u+i} + e_{1} \wedge e_{2} \otimes c_{0},$$

$$T_{A'}^{\wedge 1}(e_{1} \otimes \beta_{u+i}) = e_{0} \wedge e_{1} \otimes c_{i},$$

$$T_{A'}^{\wedge 1}(e_{2} \otimes \beta_{0}) = (e_{1} \wedge e_{2}) \otimes \sum_{i=1}^{u} c_{i},$$

$$T_{A'}^{\wedge 1}(e_{2} \otimes \beta_{i}) = e_{0} \wedge e_{2} \otimes c_{u+i},$$

$$T_{A'}^{\wedge 1}(e_{2} \otimes \beta_{u+i}) = (e_{0} \wedge e_{2}) \otimes c_{i} - e_{1} \wedge e_{2} \otimes c_{0}.$$

Notice that the image of $\sum_{i=1}^{u} (e_1 \otimes \beta_i) - \sum_{i=1}^{u} (e_2 \otimes \beta_{u+i}) - e_0 \otimes \beta_0$ is (up to scale) $e_1 \wedge e_2 \otimes e_0$.

From the contributions above, we deduce that the image of $T_{A'}^{\wedge 1}$ contains the three subspaces

$$\langle e_0 \wedge e_1, e_0 \wedge e_2, e_1 \wedge e_2 \rangle \otimes \langle c_0 \rangle, \langle e_1 \wedge e_2 \rangle \otimes \langle \sum_{i=1}^u c_i, \sum_{i=1}^u c_{u+i} \rangle, \langle e_0 \wedge e_1, e_0 \wedge e_2 \rangle \otimes \langle c_1, \dots, c_q \rangle.$$

These subspaces are in direct sum, therefore we conclude

$$rank(T_{A'}^{\wedge 1}) \ge 3 + 2 + 2q = 2q + 5.$$

3.1. Propagation of lower bounds under Kronecker products. Christandl et al. (2019, 2018), it was shown that generalized flattening lower bounds are multiplicative under the unflattened tensor product. The same result does not hold for Kronecker products. However, we provide a partial multiplicativity result for Koszul flattenings lower bounds.

A tensor $T \in A \otimes B \otimes C$, with dim $B = \dim C$ is 1_A -generic if $T(A^*) \subseteq B \otimes C$ contains a full rank element.

PROPOSITION 3.3. Let $T_1 \in A_1 \otimes B_1 \otimes C_1$ with dim $B_1 = \dim C_1$ be a tensor. Let A' be a quotient of A_1 with dim A' = 2p + 1 and suppose T_1 has a Koszul flattening lower bound for border rank $\underline{\mathbf{R}}(T) \geq r$ given by $T_{1A'}^{\wedge p}$. Let $T_2 \in$ $A_2 \otimes B_2 \otimes C_2$, with dim $B_2 = \dim C_2 = \mathbf{b}_2$ be 1_{A_2} -generic. Then

(3.4)
$$\underline{\mathbf{R}}(T_1 \boxtimes T_2) \ge \left\lceil \frac{\operatorname{rank}(T_1^{\wedge p}) \cdot \mathbf{b}_2}{\binom{2p}{p}} \right\rceil.$$

In particular, if $\frac{\operatorname{rank}(T_1^{\Lambda_p})}{\binom{2p}{2}} \in \mathbb{Z}$, then $\underline{\mathbf{R}}(T_1 \boxtimes T_2) \geq r\mathbf{b}_2$.

Identify T_1 with its image in $A' \otimes B_1 \otimes C_1$. The lower bound for T_1 Proof. is

$$\underline{\mathbf{R}}(T_1) \ge \left\lceil \frac{\operatorname{rank}(T_1 \stackrel{\wedge p}{A'})}{\binom{2p}{p}} \right\rceil.$$

Let $\alpha \in A_2^*$ be such that $T(\alpha) \in B_2 \otimes C_2$ has full rank \mathbf{b}_2 , which exists by 1_{A_2} -genericity. Define $\psi:A'\otimes A_2\to A'$ by $\psi=\mathrm{Id}_{A'}\otimes \alpha$ and set $\Psi:=$ $\psi \otimes \operatorname{Id}_{B_1 \otimes C_1 \otimes B_2 \otimes C_2}$. Then $(\Psi(T_1 \boxtimes T_2)^{\wedge p}_{A'})$ provides the desired lower bound.

Indeed, the linear map $(\Psi(T_1 \boxtimes T_2)_{A'}^{\wedge p})$ coincides with $T_1_{A'}^{\wedge p} \boxtimes T_1(\alpha)$. matrix rank is multiplicative under Kronecker product, we conclude.

3.2. A lower bound for the Kronecker square of $T_{cw,q}$. In this section, we give a proof of the first statement in Theorem 1.4.

The statement for q=2, can be checked explicitly. The lower bound $\underline{\mathbf{R}}(T_{cw,2}^{\boxtimes 2}) \geq$ 15 follows from the p=2 Koszul flattening lower bound. The upper bound is immediate by submultiplicativity. In Conner et al. (2020b), the lower bound was improved showing that $\underline{\mathbf{R}}(T_{cw,2}^{\boxtimes 2}) = 16$.

THEOREM 3.5. Let $q \geq 3$. Then $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 2}) = (q+2)^2$.

Recall the expression of $T_{cw,q}$ from (1.1). When q=3, the result is true by a direct calculation using the p=2 Koszul flattening with a sufficiently generic restriction $A \to \mathbb{C}^5$.

1 Page 14 of 40 Conner et al. cc

Assume q > 3. Write $a_{ij} = a_i \otimes a_j \in A^{\otimes 2}$ and similarly for $B^{\otimes 2}$ and $C^{\otimes 2}$. Let $A' = \langle e_0, e_1, e_2 \rangle$ and define the linear map $\phi_2 : A^{\otimes 2} \to A'$ by

$$\phi_2(a_{00}) = \phi_2(a_{01}) = \phi_2(a_{10}) = e_0 + e_1,$$

$$\phi_2(a_{11}) = e_0,$$

$$\phi_2(a_{02}) = \phi_2(a_{20}) = e_1 + e_2$$

$$\phi_2(a_{33}) = \phi_2(a_{21}) = e_2$$

$$\phi_2(a_{0i}) = \phi_2(a_{i0}) = e_1 \quad \text{for } i = 3, \dots, q$$

$$\phi_2(a_{ij}) = 0 \quad \text{for all other pairs } (i, j).$$

Write $T_q := \phi_2(T_{cw,q}^{\boxtimes 2}) \in A' \otimes B^{\otimes 2} \otimes C^{\otimes 2}$. Consider the p = 1 Koszul flattening $(T_q)_{A'}^{\wedge 1} : A' \otimes B^{\otimes 2^*} \to \Lambda^2 A' \otimes C^{\otimes 2}$.

We are going to prove that $\operatorname{rank}((T_q)_{A'}^{\wedge 1}) = 2(q+2)^2$. This provides the lower bound $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 2}) \geq (q+2)^2$ and equality follows because the upper bound is immediate by submultiplicativity.

We proceed by induction on q. When q=4 one does a direct computation with the p=1 Koszul flattening, which is left to the reader, and which provides the base of the induction.

Write $W_j = a_0 \otimes b_j \otimes c_j + a_j \otimes b_0 \otimes c_j + a_j \otimes b_j \otimes c_0$. Then $T_{cw,q} = \sum_{j=1}^q W_j$, so that $T_{cw,q}^{\boxtimes 2} = \sum_{ij} W_i \boxtimes W_j$.

If
$$q \geq 4$$
, write $T_{cw,q} = T_{cw,q-1} + W_q$, so $T_{cw,q}^{\boxtimes 2} = T_{cw,q-1}^{\boxtimes 2} + T_{cw,q-1} \boxtimes W_q + W_q \boxtimes T_{cw,q-1} + W_q \boxtimes W_q$. Let $S_q = \phi_2((T_{cw,q-1} \boxtimes W_q + W_q \boxtimes T_{cw,q-1} + W_q \boxtimes W_q))$.

Write $U_1 = A' \otimes \langle \beta_{ij} : i, j = 0, \dots, q - 1 \rangle$ and $U_2 = A' \otimes \langle \beta_{qi}, \beta_{iq} : i = 0, \dots, q \rangle$ so that $U_1 \oplus U_2 = A' \otimes B^{\otimes 2*}$. Similarly, define $V_1 = \Lambda^2 A' \otimes \langle c_{ij} : i, j = 0, \dots, q - 1 \rangle$ and $V_2 = \Lambda^2 A' \otimes \langle c_{qi}, c_{iq} : i = 0, \dots, q \rangle$, so that $V_1 \oplus V_2 = \Lambda^2 A' \otimes C^{\otimes 2}$. Observe that $(T_{q-1})_A^{\wedge 1}$ is identically 0 on U_2 and its image is contained in V_1 . Moreover, the image of U_1 under $(S_q)_{A'}^{\wedge 1}$ is contained in V_1 . Representing the Koszul flattening in blocks, we have

$$(T_{q-1})_{A'}^{\wedge 1} = \begin{bmatrix} M_{11} & 0 \\ 0 & 0 \end{bmatrix} \qquad (S_q)_{A'}^{\wedge 1} = \begin{bmatrix} N_{11} & N_{12} \\ 0 & N_{22} \end{bmatrix}$$

therefore $rank((T_q)_{A'}^{\land 1}) \ge rank(M_{11} + N_{11}) + rank(N_{22}).$

First, we prove that $\operatorname{rank}(M_{11}+N_{11}) \geq \operatorname{rank}(M_{11}) = 2(q+1)^2$. This follows by a degeneration argument.

Consider the degeneration given by the linear maps $(g_{\varepsilon}, h_{\varepsilon}) \in GL(B^{\otimes 2}) \times GL(C^{\otimes 2})$ with

$$g_{\varepsilon}: b_{iq} \mapsto \varepsilon b_{iq} \qquad h_{\varepsilon}: c_{iq} \mapsto \varepsilon c_{iq} b_{qi} \mapsto \varepsilon b_{qi} \qquad c_{qi} \mapsto \varepsilon c_{qi} b_{ij} \mapsto b_{ij} \text{ if } i, j \neq q \qquad c_{ij} \mapsto c_{ij} \text{ if } i, j \neq q$$

Let $T_{q,\varepsilon} = (g_{\varepsilon}, h_{\varepsilon}) \cdot T_q$. We have $T_{q,\varepsilon} = T_{q-1} + S_{q,\varepsilon}$ where $S_{q,\varepsilon} = (g_{\varepsilon}, h_{\varepsilon}) \cdot S_q$. In particular $\lim_{\varepsilon\to 0} S_{q,\varepsilon} = 0$. Moreover, the degeneration preserves the spaces U_1, U_2, V_1, V_2 , and therefore, the Koszul flattening of $T_{q,\varepsilon}$ has the same block structure as the one of T_q with

$$(S_{q,\varepsilon})_{A'}^{\wedge 1} = \left[\begin{array}{cc} N_{11}(\varepsilon) & N_{12}(\varepsilon) \\ 0 & N_{22}(\varepsilon) \end{array} \right].$$

Since $\lim_{\varepsilon\to 0} S_{q,\varepsilon} = 0$, we have $\lim_{\varepsilon\to 0} N_{11}(\varepsilon) \to 0$. The value of rank $(M_{11} +$ $N_{11}(\varepsilon)$) is constant for (generic) $\varepsilon \neq 0$, and by semicontinuity we obtain

$$\operatorname{rank}(M_{11}) = \operatorname{rank}(\lim_{\varepsilon \to 0} (M_{11} + N_{11}(\varepsilon)) \le \operatorname{rank}(M_{11} + N_{11}).$$

By the induction hypothesis rank $(M_{11}) = 2(q+1)^2$, threfore rank $(M_{11}+N_{11}) \ge$ $2(q+1)^2$.

We show that $rank(N_{22}) = 2(2q + 3)$. The following equalities are modulo V_1 . Moreover, each equality is modulo the tensors resulting from the previous ones. They are all straightforward applications of the Koszul flattening map, which in these cases, can always be performed on some copy of $W_i \boxtimes W_j$.

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{qj}) \equiv e_1 \wedge e_0 \otimes c_{qj} \quad \text{for } j = 3, \dots, q$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{jq}) \equiv e_1 \wedge e_0 \otimes c_{jq} \quad \text{for } j = 3, \dots, q$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{3q}) \equiv e_0 \wedge e_1 \otimes c_{0q}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{q3}) \equiv e_0 \wedge e_1 \otimes c_{q0}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{q1}) \equiv e_0 \wedge e_1 \otimes c_{q1}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{1q}) \equiv e_0 \wedge e_1 \otimes c_{1q}$$

Further passing modulo $\langle e_0 \wedge e_1 \rangle \otimes C$, we obtain

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{0q}) \equiv e_0 \wedge e_2 \otimes c_{2q}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{q0}) \equiv e_0 \wedge e_2 \otimes c_{q2}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{q2}) \equiv e_0 \wedge e_2 \otimes c_{0q}$$

$$(S_q)_{A'}^{\wedge 1}(e_0 \otimes \beta_{2q}) \equiv e_0 \wedge e_2 \otimes c_{q0}$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{20}) \equiv e_1 \wedge e_2 \otimes c_{0q}$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{02}) \equiv e_1 \wedge e_2 \otimes c_{q0}$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{02}) \equiv e_1 \wedge e_2 \otimes c_{2q}$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{q0}) \equiv e_1 \wedge e_2 \otimes c_{2q}$$

$$(S_q)_{A'}^{\wedge 1}(e_1 \otimes \beta_{0q}) \equiv e_1 \wedge e_2 \otimes c_{2q}$$

1 Page 16 of 40 Conner et al. cc

and modulo the above,

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{qj}) \equiv e_2 \wedge (e_0 + e_1) \otimes c_{qj} \quad \text{for } j = 3, \dots, q$$

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{jq}) \equiv e_2 \wedge (e_0 + e_1) \otimes c_{jq} \quad \text{for } j = 3, \dots, q$$

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{q1}) \equiv e_2 \wedge (e_0 + e_1) \otimes c_{q1}$$

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{1q}) \equiv e_2 \wedge (e_0 + e_1) \otimes c_{1q}.$$

Finally passing modulo $\langle e_1 \wedge e_2 \rangle$, we have

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{q0}) \equiv e_2 \wedge e_0 \otimes c_{q1}$$

$$(S_q)_{A'}^{\wedge 1}(e_2 \otimes \beta_{0q}) \equiv e_2 \wedge e_0 \otimes c_{1q}.$$

All the tensors listed above are linearly independent. Adding all the contributions together, we obtain

$$\operatorname{rank}((S_q)_{A'}^{\wedge 1}) = [2(q-3)+1] + 4 + 8 + 2 + [2(q-3)+1] + 4 = 2(2q+3)$$

as desired, and since $2(q+3)^2 = 2(q+1)^2 + 2(2q+3)$, this concludes the proof.

We will provide a second proof of Theorem 3.5, which will generalize to the proof of Theorem 3.7. More precisely, we will give a representation-theoretic argument to compute the rank of the Koszul flattening map considered in the proof above. The same representation-theoretic technique will apply for the third Kronecker power.

3.3. A short detour on computing ranks of equivariant maps. We briefly explain how to exploit Schur's Lemma (see, e.g., Fulton & Harris (1991, §1.2)) to compute the rank of an equivariant linear map. This is a standard technique, used extensively, e.g., in Gesmundo *et al.* (2017); Landsberg & Ottaviani (2015) and will reduce the proof of Theorem 3.5 and Theorem 3.7 to the computation of the ranks of specific linear maps in small dimension.

Let G be a reductive group. In the proof of Theorem 3.5 and Theorem 3.7, G will be the product of symmetric groups. Let Λ_G be the set of irreducible representations of G. For $\lambda \in \Lambda_G$, let W_{λ} denote the corresponding irreducible module.

Suppose U, V are two representations of G. Write $U = \bigoplus_{\lambda \in \Lambda_G} W_{\lambda}^{\oplus m_{\lambda}}$, $V = \bigoplus_{\lambda \in \Lambda_G} W_{\lambda}^{\oplus \ell_{\lambda}}$, where m_{λ} is the multiplicity of W_{λ} in U and ℓ_{λ} is the multiplicity of W_{λ} in V. The direct summand corresponding to λ is called the *isotypic component* of type λ .

Let $f: U \to V$ be a G-equivariant map. By Schur's Lemma, f decomposes as $f = \bigoplus f_{\lambda}$, where $f_{\lambda} : W_{\lambda}^{\bigoplus m_{\lambda}} \to W_{\lambda}^{\bigoplus \ell_{\lambda}}$ are G-equivariant. Consider multiplicity spaces M_{λ}, L_{λ} with $\dim M_{\lambda} = m_{\lambda}$ and $\dim L_{\lambda} = \ell_{\lambda}$ so that $W_{\lambda}^{\oplus m_{\lambda}} \simeq M_{\lambda} \otimes W_{\lambda}$ as a G-module, where G acts trivially on M_{λ} and similarly $W_{\lambda}^{\oplus \ell_{\lambda}} \simeq L_{\lambda} \otimes W_{\lambda}$.

By Schur's Lemma, the map $f_{\lambda}: M_{\lambda} \otimes W_{\lambda} \to L_{\lambda} \otimes W_{\lambda}$ decomposes as $f_{\lambda} =$ $\phi_{\lambda} \otimes \operatorname{Id}_{[\lambda]}$, where $\phi_{\lambda} : M_{\lambda} \to L_{\lambda}$. Thus $\operatorname{rank}(f)$ can be expressed in terms of $\operatorname{rank}(\phi_{\lambda})$ and the dimension of the multiplicity spaces W_{λ} for $\lambda \in \Lambda_G$:

$$\operatorname{rank}(f) = \sum_{\lambda} \operatorname{rank}(\phi_{\lambda}) \cdot \dim W_{\lambda}.$$

The ranks $\operatorname{rank}(\phi_{\lambda})$ can be computed via restrictions of f. For every λ , fix a nonzero vector $w_{\lambda} \in W_{\lambda}$, so that $M_{\lambda} \otimes \langle w_{\lambda} \rangle$ is a subspace of U. Here and in what follows, for a subset $X \subset V$, $\langle X \rangle$ denotes the span of X. Then the rank of the restriction of f to $M_{\lambda} \otimes \langle w_{\lambda} \rangle$ coincides with the rank of ϕ_{λ} .

The second proof of Theorem 3.5 and proof of Theorem 3.7 will follow the algorithm described above, exploiting the symmetries of $T_{cw,q}$. Consider the action of the symmetry group \mathfrak{S}_q on $A \otimes B \otimes C$ defined by permuting the basis elements with indices $\{1,\ldots,q\}$. More precisely, a permutation $\sigma\in\mathfrak{S}_q$ induces the linear map defined by $\sigma(a_i) = a_{\sigma(i)}$ for $i = 1, \ldots, q$ and $\sigma(a_0) = a_0$. The group \mathfrak{S}_q acts on B, C similarly, and the simultaneous action on the three factors defines an \mathfrak{S}_q -action on $A \otimes B \otimes C$. The tensor $T_{cw,q}$ is invariant under this action.

3.4. Second Proof of Theorem 3.5. We use the method explained in Section 3.3 to give a representation-theoretic proof of Theorem 3.5.

PROOF (Proof of Theorem 3.5). As before, the case q=3 can be verified explicitly. For $q \geq 4$, we apply the p = 1 Koszul flattening map to the same restriction of $T_{cw,q}^{\boxtimes 2}$ as the first proof, although to be consistent with the code on the website, we use the less appealing swap of the roles of a_2 and a_3 in the projection ϕ_2 .

The tensor $T_{cw,q}$ is invariant under the action of \mathfrak{S}_q acting on the indices $\{1,\ldots,q\}$ of the basis elements of \mathbb{C}^{q+1} . Therefore, $T_{cw,q}^{\boxtimes 2}$ is invariant under the action of $\mathfrak{S}_q \times \mathfrak{S}_q$ on $A^{\otimes 2} \otimes B^{\otimes 2} \otimes C^{\otimes 2}$. Let $\Gamma := \mathfrak{S}_{q-3} \times \mathfrak{S}_{q-3}$ where \mathfrak{S}_{q-3} is the permutation group on $\{4,\ldots,q\}$; $T_{cw,q}^{\boxtimes 2}$ is invariant under the action of Γ.

Moreover, the projection ϕ_2 is invariant under the action of Γ .

In general, the map $A \otimes B \otimes C \to \text{Hom}(B^* \otimes \Lambda^p A, C \otimes \Lambda^{p+1} A)$ is equivariant for the action of $GL(A) \times GL(B) \times GL(C)$. Using this fact, and the invariancy with respect to Γ described above, we deduce $(\phi_2(T_{cw,q}^{\boxtimes 2}))_{A'}^{\wedge 1}$ is Γ -equivariant.

We now apply the method described in Section 3.3 to compute rank $((T_q)_{A'}^{\wedge 1})$.

1 Page 18 of 40 Conner et al.

Let [triv] denote the trivial \mathfrak{S}_{q-3} -representation and let V denote the standard representation, that is the Specht module associated to the partition (q-4,1) of q-3. We have $\dim[\text{triv}]=1$ and $\dim V=q-4$. When q=4 only the trivial representation appears.

The spaces B, C are isomorphic as \mathfrak{S}_{q-3} -modules and they decompose as $B = C = [\operatorname{triv}]^{\oplus 5} \oplus V$. After fixing a 5-dimensional multiplicity space \mathbb{C}^5 for the trivial isotypic component, we write $B^* = C = \mathbb{C}^5 \otimes [\operatorname{triv}] \oplus V$. To distinguish the two \mathfrak{S}_{q-3} -actions, we write $B^{*\otimes 2} = ([\operatorname{triv}]_L^{\oplus 5} \oplus V_L) \otimes ([\operatorname{triv}]_R^{\oplus 5} \oplus V_R)$ and similarly for $C^{\otimes 2}$

Thus,

$$B^{*\otimes 2} = C^{\otimes 2} = \mathbb{C}^{5\otimes 2} \otimes ([\operatorname{triv}]_L \otimes [\operatorname{triv}]_R) \oplus$$

$$\mathbb{C}^5 \otimes ([\operatorname{triv}]_L \otimes V_R) \quad \oplus$$

$$\mathbb{C}^5 \otimes (V_L \otimes [\operatorname{triv}]_R) \quad \oplus$$

$$(V_L \otimes V_R).$$

Write W_1, \ldots, W_4 for the four irreducible representations in the decomposition above and let M_1, \ldots, M_4 be the four corresponding multiplicity spaces.

Recall from Fulton (1997) that a basis of V is given by standard Young tableaux of shape (q-4,1) (with entries in $4,\ldots,q$ for consistency with the action of \mathfrak{S}_{q-3}); let w_{std} be the vector corresponding to the standard tableau having $4,6,\ldots,q$ in the first row and 5 in the second row. We refer to Fulton (1997, §7) for the straightening laws of the tableaux. Let w_{triv} be a generator of the trivial representation [triv]. Writing $\mathbb{C}^{q+1} = \langle e_0,\ldots,e_q \rangle$, we explicitly have $w_{std} = e_5 - e_4$ and the multiplicity space 5-dimensional multiplicity space of the trivial representation is $\langle e_0,\ldots,e_3,\sum_{q=1}^q e_j \rangle$.

For each of the four isotypic components in the decomposition above, we fix a vector $w_i \in W_i$ and explicitly realize the subspaces $M_i \otimes \langle w_i \rangle$ of $B^{*\otimes 2}$ as follows:

W_i	w_i	$\dim M_i$	$M_i \otimes \langle w_i angle$
$[\operatorname{triv}]_L \otimes [\operatorname{triv}]_R$	$w_{ m triv} \otimes w_{ m triv}$	25	$\langle \beta_{ij} : i, j = 0, \dots, 3 \rangle \oplus$ $\langle \sum_{j=4}^{q} \beta_{ij} : i = 0, \dots, 3 \rangle \oplus$ $\langle \sum_{i=4}^{q} \beta_{ij} : j = 0, \dots, 3 \rangle \oplus$ $\langle \sum_{i,j=4}^{q} \beta_{ij} \rangle$
$[\mathrm{triv}]_L \otimes V_R$	$w_{ m triv}\otimes w_{std}$	5	$\langle \beta_{i5} - \beta_{i4} : i = 0, \dots, 3 \rangle \oplus \\ \langle \sum_{i=4}^{q} (\beta_{i5} - \beta_{i4}) \rangle$
$V_L \otimes [\mathrm{triv}]_R$	$w_{std} \otimes w_{ ext{triv}}$	5	$\langle \beta_{5j} - \beta_{4j} : j = 0,, 3 \rangle \oplus \\ \langle \sum_{j=4}^{q} (\beta_{5j} - \beta_{4j}) \rangle$
$V_L \otimes V_R$	$w_{std} \otimes w_{std}$	1	$\langle \beta_{55} - \beta_{45} - \beta_{54} + \beta_{44} \rangle$.

The subspaces in $C^{\otimes 2}$ are realized similarly.

Since $(T_{cw,q}^{\boxtimes 2})_{A'}^{\wedge 1}$ is Γ -equivariant, by Schur's Lemma, it has the isotypic decomposition $(T_{cw,q}^{\boxtimes 2})_{A'}^{\wedge 1} = f_1 \oplus f_2 \oplus f_3 \oplus f_4$, where

$$(3.6) f_i: A' \otimes (M_i \otimes W_i) \to \Lambda^2 A' \otimes (M_i \otimes W_i).$$

As explained in Section 3.3, it suffices to compute the ranks of the four restrictions $\Phi_i: A' \otimes M_i \otimes \langle w_i \rangle \to \Lambda^2 A' \otimes M_i \otimes \langle w_i \rangle$ to the multiplicities spaces.

The four matrices representing Φ_1, \ldots, Φ_4 are computed by a routine which exploits their structure. The script to compute the matrices and their ranks is available in Appendix D of Supplementary Material. The method to compute the matrices is explained in Section 6.

The script provides an expression for the entries of the matrices Φ_i which are univariate polynomials in q up to a global univariate polynomial factor. The expressions are valid for $q \geq 5$. The rank of the Koszul flattening in the cases q=3 and q=4 is computed directly.

We determine a lower bound on rank(Φ_i) by computing a matrix $P_i \cdot \Phi_i \cdot Q_i$, where P_i is a rectangular matrix whose entries are rational functions of q (well defined for $q \geq 5$) and Q_i is a rectangular matrix whose entries are constant. The resulting matrix $P_i \cdot \Phi_i \cdot Q_i$ is a square matrix, upper triangular with ± 1 on the diagonal, so that the size of $P_i\Phi_iQ_i$ gives a lower bound on rank (Φ_i) .

We summarize the results of the script in the following table.

W_i	$\dim W_i$	$\dim M_i$	$\operatorname{rank}(\Phi_i)$	contribution to total rank
$[\mathrm{triv}]_L \otimes [\mathrm{triv}]_R$	1	25	72	72
$[\mathrm{triv}]_L \otimes V_R$	q-4	5	12	12(q-4)
$V_L \otimes [\mathrm{triv}]_R$	q-4	5	12	12(q-4)
$V_L \otimes V_R$	$(q-4)^2$	1	2	$2(q-4)^2$

Adding the total contributions, we obtain

$$\operatorname{rank}(T_{A'}^{\wedge 1}) = 2 \cdot (q-4)^2 + 12 \cdot (q-4) + 12 \cdot (q-4) + 72 \cdot 1 = 2(q+2)^2.$$

This concludes the proof of Theorem 3.5.

3.5. A lower bound for the Kronecker cube of $T_{cw,q}$. In this section, we use the method explained in Section 3.3 and illustrated in Section 3.4 to prove the second part of Theorem 1.4.

1 Page 20 of 40 Conner et al. cc

THEOREM 3.7. Let $q \geq 5$. Then $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 3}) = (q+2)^3$.

PROOF. We will give a lower bound on $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 3})$ by computing its Koszul flattening for p=2. Write $a_{ijk}=a_i\otimes a_j\otimes a_k\in A^{\otimes 3}$ and similarly for $B^{\otimes 3}$ and $C^{\otimes 3}$. Let $\{\alpha_{ijk}\}\subseteq A^{*\otimes 3}$ be the dual basis to $\{a_{ijk}\}\subseteq A^{\otimes 3}$. Let $A'=\langle e_0,\ldots,e_4\rangle$ be a 5-dimensional space and let $\{e^0,\ldots,e^4\}$ be the dual basis of $\{e_0,\ldots,e_4\}$ and define $\phi_3:A^{\otimes 3}\to A'$ to be the linear map whose transpose $\phi_3^T:A'^*\to A^{*\otimes 3}$ is given by

$$\phi_3^T(e^0) = \alpha_{000}$$

$$\phi_3^T(e^1) = \sum_{i=1}^q (\alpha_{i00} + \alpha_{0i0} + \alpha_{00i})$$

$$\phi_3^T(e^2) = \alpha_{001} + \alpha_{010} + \alpha_{012} + \alpha_{102} + \alpha_{110} + \alpha_{121} + \alpha_{200} + \alpha_{211}$$

$$\phi_3^T(e^3) = \alpha_{022} + \alpha_{030} + \alpha_{031} + \alpha_{100} + \alpha_{103} - \alpha_{120} + \alpha_{210} + \alpha_{212} + \alpha_{300}$$

$$\phi_3^T(e^4) = \alpha_{002} + \alpha_{004} + \alpha_{011} + \alpha_{014} + \alpha_{020} + \alpha_{023} + \alpha_{032} + \alpha_{040} + \alpha_{100} + \alpha_{122} + \alpha_{220} + \alpha_{303}.$$

Let $T_q = \phi_3(T_{cw,q}^{\boxtimes 3}) \in A' \otimes B^{\otimes 3} \otimes C^{\otimes 3}$ and consider the Koszul flattening

$$(T_q)^{\wedge 2}_{A'}: \Lambda^2 A' \otimes B^{* \otimes 3} \to \Lambda^3 A' \otimes C^{\otimes 3}$$

We will show rank $((T_q)_{A'}^{\wedge 2}) = 6(q+2)^3$, which implies $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes 3}) \geq (q+2)^3$.

We employ the same method as in Section 3.4 in the case of $T_{cw,q}^{\boxtimes 2}$. The Koszul flattening is equivariant for the action of $\Gamma = \mathfrak{S}_{q-4}^{\times 3}$ where \mathfrak{S}_{q-4} acts on $\{5,\ldots,q\}$. In particular, \mathbb{C}^{q+1} splits under the action of \mathfrak{S}_{q-4} into a 6-dimensional subspace of invariants $\mathbb{C}^6 \otimes [\text{triv}] = \langle e_0,\ldots,e_4,e_5+\cdots+e_q \rangle$ and a copy of the standard representation $V = \langle e_i - e_5 : i = 6,\ldots,q \rangle$, with $\dim V = q - 5$.

Hence, the spaces $B^{\otimes 3}$ and $C^{\otimes 3}$ split into the direct sum of 8 isotypic components for the action of Γ as follows (we use indices 1, 2, 3 to denote the trivial or the standard representation on the first, second or third factor):

$$B^{*\otimes 3} \simeq C^{\otimes 3} = (\mathbb{C}^{6})^{\otimes 3} \otimes ([\operatorname{triv}]_{1} \otimes [\operatorname{triv}]_{2} \otimes [\operatorname{triv}]_{3}) \oplus$$

$$(\mathbb{C}^{6})^{\otimes 2} \otimes \Big[([\operatorname{triv}]_{1} \otimes [\operatorname{triv}]_{2} \otimes V_{3}) \oplus$$

$$([\operatorname{triv}]_{1} \otimes V_{2} \otimes [\operatorname{triv}]_{3}) \oplus$$

$$(V_{1} \otimes [\operatorname{triv}]_{2} \otimes [\operatorname{triv}]_{3}) \Big] \oplus$$

$$(\mathbb{C}^{6}) \otimes \Big[([\operatorname{triv}]_{1} \otimes V_{2} \otimes V_{3}) \oplus$$

$$(V_{1} \otimes V_{2} \otimes [\operatorname{triv}]_{3}) \oplus$$

$$(V_{1} \otimes [\operatorname{triv}]_{2} \otimes V_{3}) \Big] \oplus$$

$$V_{1} \otimes V_{2} \otimes V_{3}$$

Similarly to the square case, for each of the eight isotypic components, we consider $w_i \in W_i$ where W_i is the corresponding irreducible and we compute the rank of the restriction $\Psi_i: \Lambda^2 A' \otimes M_i \otimes \langle w_i \rangle \to \Lambda^3 A' \otimes M_i \otimes \langle w_i \rangle$ of the Koszul flattening.

The matrices representing the maps Ψ_i are computed exploiting the structure of the tensors involved, following the method described in Section 6. The expression computed by the script is valid for $q \geq 6$. The case q = 5 is computed explicitly. Their ranks are computed by reducing Ψ_i to a triangular matrix as in the previous case.

The ranks of the restrictions are recorded in the following table:

W_i	\dimW_i	$\dim M_i$	${\rm rank}(\Psi_i)$	total contribution
$[\mathrm{triv}]_1 \otimes [\mathrm{triv}]_2 \otimes [\mathrm{triv}]_3$	1	$6^3 = 216$	2058	2058
$[\text{triv}]_1 \otimes [\text{triv}]_2 \otimes V_3$ (and permutations)	(q-5) (three times)	$6^2 = 36$ (three times)	294 (three times)	$3 \cdot 294(q-5)$
$[ext{triv}]_1 \otimes V_2 \otimes V_3$ (and permutations)	$(q-5)^2$ (three times)	6 (three times)	42 (three times)	$3\cdot 42(q-5)^2$
$V_1 \otimes V_2 \otimes V_3$	$(q-5)^3$	1	6	$6(q-5)^3$

Adding all the contributions together, we obtain

$$\operatorname{rank}(T_{A'}^{\wedge 2}) = 6(q-5)^3 + 3 \cdot 42(q-5)^2 + 3 \cdot 294(q-5) + 2058 \cdot 1 = 6 \cdot (q+2)^3.$$

This concludes the proof of Theorem 3.7.

The third part of Theorem 1.4 is a consequence of Proposition 3.3 and Theorem 3.7 for the case q > 5 and Proposition 3.3 and Theorem 3.5 in the case q=4. We record it explicitly in the following Corollary

COROLLARY 3.8. For all q > 4 and all N, $\underline{\mathbf{R}}(T_{cw,a}^{\boxtimes N}) \ge (q+1)^{N-3}(q+2)^3$, and $\mathbf{R}(T_{cw,4}^{\boxtimes N}) \ge 36 \times 5^{N-2}.$

PROOF. If q > 4, let $T_1 = T_{cw,q}^{\boxtimes 3}$ and $T_2 = T_{cw,q}^{\boxtimes N-3}$. Since T_2 is 1_A -generic, the lower bound $\underline{\mathbf{R}}(T_{cw,q}^{\boxtimes N}) \geq (q+1)^{N-3}(q+2)^3$ follows by Proposition 3.3.

If q=4, let $T_1=T_{cw,q}^{\boxtimes 2}$ and $T_2=T_{cw,q}^{\boxtimes N-2}$. Again, since T_2 is 1_A -generic, the lower bound $\underline{\mathbf{R}}(T_{cw,4}^{\boxtimes N})\geq (4+2)^2\times 5^{N-2}=36\times 5^{N-2}$ follows by Proposition 3.3.

This concludes the proof of Theorem 1.4.

1 Page 22 of 40 Conner et al.

4. Upper bounds for Waring rank and border Waring rank of det₃

In this section, we give a proof of Theorem 1.6. We briefly recall the definition of Waring rank and border Waring rank. A symmetric tensor $T \in S^d \mathbb{C}^m \subseteq \mathbb{C}^{m \otimes d}$ has Waring rank one if $T = a^{\otimes d}$ for some $a \in \mathbb{C}^m$. The Waring rank of T, denoted $\mathbf{R}_S(T)$, is the smallest r such that T is sum of r tensors of Waring rank one. The border Waring rank of T, denoted $\mathbf{R}_S(T)$, is the smallest r such that T is limit of a sequence of tensors of Waring rank r. If T is regarded as a homogeneous polynomial of degree d, then $a \in \mathbb{C}^m$ can be regarded as a linear form and $a^{\otimes d}$ coincides with the d-th power of a: in this setting, the Waring rank is the minimum number of summands in an expression of T as sum of powers of linear forms.

4.1. Waring rank of det₃. Theorem 1.6 will be a consequence of Theorem 4.1 and Theorem 4.2 below.

THEOREM 4.1. The Waring rank of det₃ is at most 18: $\mathbf{R}_S(\det_3) \leq 18$.

PROOF. We give the rank 18 decomposition for det₃ explicitly, as a collection of 18 linear forms on $\mathbb{C}^9 = \mathbb{C}^3 \otimes \mathbb{C}^3$ whose cubes add up to det₃. The linear forms are given in coordinates recorded in the matrices below: the 3×3 matrix (ζ_{ij}) represents the linear form $\sum_{ij} \zeta_{ij} x_{ij}$. This presentation highlights some of the symmetries of the decomposition. Let $\vartheta = \exp(2\pi i/6)$ and let $\overline{\vartheta}$ be its inverse. The tensor det₃ = $T_{skewcw,2}^{\boxtimes 2} = \det(x_{ij}) \in S^3(\mathbb{C}^3 \otimes \mathbb{C}^3)$ satisfies

$$\det_3 = \sum_{i=1}^{18} L_i^3$$

where L_1, \ldots, L_{18} are the 18 linear forms given by the following coordinates:

$$\begin{split} L_1 &= \begin{pmatrix} -\vartheta & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{0}{\vartheta} \\ 0 & 0 & \frac{1}{\vartheta} \end{pmatrix} \qquad L_2 = \begin{pmatrix} -\overline{\vartheta} & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & \vartheta \end{pmatrix} \qquad L_3 = \begin{pmatrix} -\overline{\vartheta} & 0 & 0 \\ 0 & \frac{1}{3} \overline{\vartheta} & 0 \\ 0 & 0 & \frac{1}{\vartheta} \end{pmatrix} \\ L_4 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -\overline{\vartheta} \\ 0 & -\frac{1}{3} \vartheta & 0 \end{pmatrix} \qquad L_5 = \begin{pmatrix} \overline{\vartheta} & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -\frac{1}{3} \vartheta & 0 \end{pmatrix} \qquad L_6 = \begin{pmatrix} \vartheta & 0 & 0 \\ 0 & 0 & -\overline{\vartheta} \\ 0 & -\frac{1}{3} \vartheta & 0 \end{pmatrix} \\ L_7 &= \begin{pmatrix} 0 & \frac{1}{3} \overline{\vartheta} & 0 \\ -\vartheta & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \qquad L_8 &= \begin{pmatrix} 0 & \frac{1}{3} \overline{\vartheta} & 0 \\ -\overline{\vartheta} & 0 & 0 \\ 0 & 0 & -\overline{\vartheta} \end{pmatrix} \qquad L_9 &= \begin{pmatrix} 0 & \frac{1}{3} \vartheta & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ L_{10} &= \begin{pmatrix} 0 & -\frac{1}{3} \vartheta & 0 \\ 0 & 0 & \overline{\vartheta} \\ -1 & 0 & 0 \end{pmatrix} \qquad L_{11} &= \begin{pmatrix} 0 & -\frac{1}{3} \overline{\vartheta} & 0 \\ 0 & 0 & \vartheta \\ -1 & 0 & 0 \end{pmatrix} \qquad L_{12} &= \begin{pmatrix} 0 & \frac{1}{3} \vartheta & 0 \\ 0 & 0 & -1 \\ -1 & 0 & 0 \end{pmatrix} \\ L_{13} &= \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \end{pmatrix} \qquad L_{14} &= \begin{pmatrix} 0 & 0 & 1 \\ \overline{\vartheta} & 0 & 0 \\ 0 & \frac{1}{3} \vartheta & 0 \end{pmatrix} \qquad L_{15} &= \begin{pmatrix} 0 & 0 & 1 \\ \vartheta & 0 & 0 \\ 0 & \frac{1}{3} \overline{\vartheta} & 0 \end{pmatrix} \\ L_{16} &= \begin{pmatrix} 0 & 0 & \overline{\vartheta} \\ 0 & -\frac{1}{3} \overline{\vartheta} & 0 \\ 0 & 0 & 0 \end{pmatrix} \qquad L_{17} &= \begin{pmatrix} 0 & 0 & \overline{\vartheta} \\ 0 & -\frac{1}{3} \overline{\vartheta} & 0 \\ 0 & -\frac{1}{3} \overline{\vartheta} & 0 \end{pmatrix} \qquad L_{18} &= \begin{pmatrix} 0 & 0 & \vartheta \\ 0 & -\frac{1}{3} \overline{\vartheta} & 0 \\ 0 & -\frac{1}{3} \overline{\vartheta} & 0 \end{pmatrix} \end{split}$$

The equality can be verified by hand. A Macaulay file performing the calculation is available in Appendix B of Supplementary Material. The Waring decomposition of Theorem 4.1 was generalized in Johns & Teitler (2020) giving an upper bound for the Waring rank of the determinant polynomial \det_m . \square

4.2. Waring border rank of det₃. The statement for the border rank is given by the following result. As in the previous proof, the border rank upper bound is proved explicitly giving linear forms, depending on a parameter t, whose cubes provide a border rank expression for det₃. The algebraic numbers involved are more complicated than in the previous case.

The result was achieved by numerical methods, which allowed us to sparsify the decomposition and ultimately determine the value of the coefficients. A detailed explanation of the method is given in Section 4.3.

THEOREM 4.2. The border Waring rank of det₃ is at most 17: $\mathbf{R}_{S}(\det_{3}) \leq 17$.

The 17 linear forms providing a border rank decomposition of det₃ are described below. Consider

$$\begin{split} L_1(t) &= \begin{pmatrix} z_1 & 0 & 0 \\ 0 & z_2 t & 0 \\ -1 & 0 & 0 \end{pmatrix} & L_2(t) &= \begin{pmatrix} z_3 & 0 & 0 \\ z_4 & 0 & z_5 t \\ z_6 & 0 & 0 \end{pmatrix} & L_3(t) &= \begin{pmatrix} -z_{36} & z_7 t & 0 \\ -z_{38} & 0 & -z_{39} t \\ 0 & 0 & t \end{pmatrix} \\ L_4(t) &= \begin{pmatrix} 0 & 0 & t \\ -z_{34} & 0 & 0 \\ 0 & z_8 t & -z_{35} t \end{pmatrix} & L_5(t) &= \begin{pmatrix} 0 & -z_{19} t & -z_{20} t \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} & L_6(t) &= \begin{pmatrix} -z_{22} & z_9 t & 0 \\ -z_{23} & 0 & -z_{24} t \\ -z_{25} & 0 & 0 \end{pmatrix} \\ L_7(t) &= \begin{pmatrix} z_{10} & z_{11} t & 0 \\ z_{12} & 0 & z_{13} t \\ z_{14} & 0 & 0 \end{pmatrix} & L_8(t) &= \begin{pmatrix} z_{15} & -t & 0 \\ z_{16} & 0 & z_{17} t \\ z_{18} & 0 & 0 \end{pmatrix} & L_9(t) &= \begin{pmatrix} 0 & z_{19} t & z_{20} t \\ 0 & z_{21} t & 0 \\ 1 & 0 & 0 \end{pmatrix} \\ L_{10}(t) &= \begin{pmatrix} -z_{41} & 0 & 0 \\ 0 & 0 & 0 \\ -z_{44} & 0 & 0 \end{pmatrix} & L_{11}(t) &= \begin{pmatrix} z_{22} & 0 & 0 \\ z_{23} & 0 & z_{24} t \\ z_{25} & 0 & 0 \end{pmatrix} & L_{12}(t) &= \begin{pmatrix} -z_{31} & z_{26} t & 0 \\ 0 & z_{27} t & 0 \\ 0 & 0 & 0 & t \end{pmatrix} \\ L_{13}(t) &= \begin{pmatrix} z_{28} & z_{29} t & 0 \\ z_{30} & 0 & -t \\ 0 & t & 0 \end{pmatrix} & L_{14}(t) &= \begin{pmatrix} z_{31} & z_{32} t & 0 \\ 0 & z_{33} t & -t \end{pmatrix} & L_{15}(t) &= \begin{pmatrix} 0 & 0 & -t \\ z_{34} & 0 & 0 \\ 0 & 0 & z_{35} t \end{pmatrix} \\ L_{16}(t) &= \begin{pmatrix} z_{36} & z_{37} t & 0 \\ z_{38} & 0 & z_{39} t \\ 0 & z_{40} t & -t \end{pmatrix} & L_{17}(t) &= \begin{pmatrix} z_{41} & z_{42} t & 0 \\ 0 & z_{44} & 0 & 0 \\ 0 & z_{44} & 0 & 0 \end{pmatrix} \end{aligned}$$

The coefficients z_1, \ldots, z_{44} are algebraic numbers described as follows. Let y_* be a real root of the polynomial

$$x^{27} - 2x^{26} + 17x^{25} - 29x^{24} + 81x^{23} + 52x^{22} - 726x^{21} + 3451x^{20} - 10901x^{19} + 25738x^{18} \\ - 50663x^{17} + 72133x^{16} - 72973x^{15} + 10444x^{14} + 138860x^{13} - 308611x^{12} + 427344x^{11} \\ - 267416x^{10} - 196096x^9 + 762736x^8 - 1236736x^7 + 1092352x^6 - 537600x^5 - 42240x^4 \\ + 684032x^3 - 1136640x^2 + 1146880x - 520192.$$

For i = 1, ..., 44, we consider algebraic numbers y_i in the field extension $\mathbb{Q}[y_*]$, described as a polynomial of degree (at most) 26 in y_* with rational coefficients. Notice that all the y_i 's are real. The expressions of the y_1, \ldots, y_{44} in terms of 1 Page 24 of 40 Conner et al. cc

 y_* are provided in the file yy_exps in Appendix C of Supplementary Material. Let z_j be the unique real cubic root of y_j .

We are going to prove that, with this choice of coefficients z_i ,

(4.3)
$$t^{2} \det_{3} + O(t^{3}) = \sum_{i=1}^{17} L_{i}(t)^{3}.$$

The condition $t^2 \det_3 + O(t^3) = \sum_{i=1}^{17} L_i(t)^3$ is equivalent to the fact that the degree 0 and the degree 1 components of $\sum_{i=1}^{17} L_i(t)^3$ vanish and that the degree 2 component equals \det_3 . Given the sparse structure of the $L_i(t)$, this reduces to a system of 54 cubic equations in the 44 unknowns z_1, \ldots, z_{44} . Our goal is to show that the algebraic numbers described above are a solution of this system.

We show that the z_i 's satisfy each equation as follows. After evaluating the equations at the z_i 's, there are two possible cases

- 1. all monomials appearing in the equation are elements of $\mathbb{Q}[y_*]$; we say that this is an equation of type 1; there are 14 such equations;
- 2. at least one monomial appearing in the equation is not an element of $\mathbb{Q}[y_*]$; we say that this is an equation of type 2; there are 40 such equations.

For equations of type 1, we provide expressions of each monomial in terms of y_* . To verify that each expression is indeed equal to the corresponding monomial, it suffices to compare the cube of the given expression and the expression obtained by evaluating the monomial at the y_j 's. Finally, the equation can be verified in $\mathbb{Q}[y_*]$. This is performed by the file checkingType1eqns.m2.

For equations of type 2, let u be one of the monomials which do not belong to $\mathbb{Q}[y_*]$. We claim that it is possible to choose the monomial in such a way that $\mathbb{Q}[u^3] = \mathbb{Q}[y_*]$. For each equation, we choose one of the monomials and we verify the claim as follows. The element u^3 has an expression in terms of y_* which equals the chosen monomial evaluated at the y_i 's. Let M_u be the 27×27 matrix with rational entries such that

$$(1, u^3, \dots, u^{3 \cdot 26}) = (1, y_*, \dots, y_*^{26}) \cdot M_u;$$

 M_u can be computed directly by considering the expressions of the powers of u^3 in terms of y_* . Then $\mathbb{Q}[u^3] = \mathbb{Q}[y_*]$ if and only if M_u is full rank.

In particular y_* has an expression in terms of u^3 , which can be computed inverting the matrix M_u . A consequence of this is that $\mathbb{Q}[u] = \mathbb{Q}[y_*, u]$.

At this point, we observe that $\mathbb{Q}[u]$ contains the other monomials occurring in the equation as well. To see this, we proceed as in the case of equations of type 1. For each monomial occurring in the equation, we provide an expression in

terms of u (in fact, to speed up the calculation, we provide an expression in terms of u and y_* , which is equivalent to an expression in u because $\mathbb{Q}[u^3] =$ $\mathbb{Q}[y_*]$ and y_* has a unique expression in terms of u^3); we compare the cube of this expression (appropriately reduced modulo the minimal polynomial of y_* and the relation between u^3 and u_*) with the expression obtained by evaluating the monomial at the y_i 's (expressed in terms of y_*). This shows that all monomials occurring in the expression belong to $\mathbb{Q}[u]$, and verifies that the given expressions are indeed equal to the corresponding monomials. Finally, the equation is verified in $\mathbb{Q}[u]$ as in the case of type 1. This is performed by the file checkingType2eqns.m2.

4.3. Discussion of how the decomposition was obtained. were accomplished by finding solutions of polynomial equations by nonlinear optimization. In each case, this was accomplished using a variant of Newton's method applied to the mapping of variable values to corresponding polynomial values. The result of this procedure in each case is limited precision machine floating point numbers.

First, we attempted to solve the equations describing a Waring rank 17 decomposition of det₃ with nonlinear optimization, namely, $\det_3 = \sum_{i=1}^{17} (w_i')^{\otimes 3}$, where $w_i' \in \mathbb{C}^{3\times 3}$. Instead of finding a solution to working precision, we obtained a sequence of local refinements to an approximate solution where the distance between det₃ and its approximation is slowly converging to zero, and some of the parameter values are exploding to infinity. Numerically, these are Waring decompositions of polynomials very close to det_3 .

Next, this approximate solution needed to be upgraded to a solution to equation (4.3).

We found a choice of parameters in the neighborhood of a solution, and then applied local optimization to solve to working precision. We used the following method: Consider the linear mapping $M: \mathbb{C}^{17} \to S^3(\mathbb{C}^{3\times 3}), M(e_i) = (w_i')^{\otimes 3}$, and let $M = U\Sigma V^*$ be its singular value decomposition (with respect to the standard inner products for the natural coordinate systems). We observed that the singular values seemed to be naturally partitioned by order of magnitude. We estimated this magnitude factor as $t_0 \approx 10^{-3}$, and wrote Σ' as Σ where we multiplied each singular value by $(t/t_0)^k$, with k chosen to agree with this observed partitioning, so that the constants remaining were reasonably sized. Finally, we let $M' = U\Sigma'V^*$, which has entries in $\mathbb{C}[[t]]$. Thus M' is a representation of the map M with a parameter t.

Next, for each i, we optimized to find a best fit to the equation $(a_i + tb_i +$ $t^2c_i)^{\otimes 3}=M'(e_i)$, which is defined by polynomial equations in the entries of a_i, b_i and c_i . The a_i, b_i and c_i we constructed in this way proved to be a good initial guess to optimize equation (4.3), and we immediately saw quadratic convergence to a solution to machine precision. At this point, we greedily sparsified the solution by speculatively zero-ing values and reoptimizing, rolling

1 Page 26 of 40 Conner et al. cc

back one step in case of failure. After sparsification, it turned out the c_i were not needed. The resulting matrices are those given in the proof.

To compute the minimal polynomials and other integer relationships between quantities, we used Lenstra–Lenstra–Lovász integer lattice basis reduction Lenstra et al. (1982). As an example, let $\zeta \in \mathbb{R}$ be approximately an algebraic number of degree k. Let N be a large number inversely proportional to the error of ζ . Consider the integer lattice with basis $\{e_i + \lfloor N\zeta^i \rfloor e_{k+1}\} \subset \mathbb{Z}^{k+2}$, for $0 \le i \le k$. Then elements of this lattice are of the form $v_0e_0 + \cdots + v_ke_k + Ee_{k+1}$, where $E \approx Np(\zeta)$, $p = v_0 + v_1x + \cdots x_kx^k$. Polynomials p for which ζ is an approximate root are distinguished by the property of having relatively small Euclidean norm in this lattice. Computing a small norm vector in an integer lattice is accomplished by LLL reduction of a known basis.

For example, the fact that the number field of degree 27 obtained by adjoining any z_{α}^3 to \mathbb{Q} contains all the rest was determined via LLL reduction, looking for expressions of z_{α}^3 as a polynomial in z_{β}^3 for some fixed β . These expressions of z_{α}^3 in a common number field can be checked to have the correct minimal polynomial, and thus agree with our initial description of the z_{α} . LLL reduction was also used to find the expressions of values as polynomials in the primitive root of the various number fields.

After refining the known value of the parameters to 10,000 bits of precision using Newton's method, LLL reduction was successful in identifying the minimal polynomials. The degrees were simply guessed, and the results checked by evaluating the computed polynomials in the parameters to higher precision.

REMARK 4.4. With the minimal polynomial information, it is possible to check that equation (4.3) is satisfied to any desired precision by the parameters.

5. Tight tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$

Following an analysis started in Conner *et al.* (2020a), we consider Kronecker squares of tight tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$. We compute their symmetry groups and numerically provide bounds to their tensor rank and border rank, highlighting the submultiplicativity properties.

We refer to Bläser (2013); Bürgisser et al. (1997); Conner et al. (2020a); Strassen (1994) for an exposition of the role of tightness in Strassen's work and in the laser method. In Lemma 5.1, we explicitly show that $T_{cw,q}$ and $T_{CW,q}$ are tight tensors. This fact was known and appears implicitly in Bläser (2013); Christandl et al. (2021) and other related works: However, we are not aware of a reference where the proof is given in its entirety.

5.1. Tight tensors. Recall the map $\Phi: GL(A) \times GL(B) \times GL(C) \rightarrow$ $GL(A \otimes B \otimes C)$ from Section 2.2 defining the action of $GL(A) \times GL(B) \times GL(C)$ on $A \otimes B \otimes C$. Its differential $d\Phi$ defines a map at the level of Lie algebras, mapping $\mathfrak{gl}(A) \oplus \mathfrak{gl}(B) \oplus \mathfrak{gl}(C)$ to a subalgebra of $\mathfrak{gl}(A \otimes B \otimes C)$. This subalgebrais isomorphic to $(\mathfrak{gl}(A) \oplus \mathfrak{gl}(B) \oplus \mathfrak{gl}(C))/\mathbb{C}^2$ where $\mathbb{C}^2 \simeq \ker d\Phi =$ $\{(\lambda_A \mathrm{Id}_A, \lambda_B \lambda \mathrm{Id}_B, \lambda_C \lambda \mathrm{Id}_C) : \lambda_A + \lambda_B + \lambda_C = 0\}$ is the Lie algebra of the 2dimensional kernel of Φ . Write $\mathfrak{g}_T \subseteq \mathfrak{gl}(A) \oplus \mathfrak{gl}(B) \oplus \mathfrak{gl}(C)$ for the annihilator of T under this action.

A tensor $T \in A \otimes B \otimes C$ is tight if $\mathfrak{g}_T/\mathbb{C}^2$ contains a regular semisimple element. Given a basis $\{a_i : i = 1, ..., \dim A\}$ of A and similarly for B and C, write T_{ijk} for the coordinates of a tensor T in the induced basis $\{a_i \otimes b_j \otimes c_k\}$ of $A \otimes B \otimes C$. The support of a tensor $T \in A \otimes B \otimes C$ is

$$supp(T) = \{(i, j, k) : T_{ijk} \neq 0\}.$$

Tightness can be defined combinatorially with respect to a basis, see, e.g., Conner et al. (2020a, Def. 1.3). Explicitly, T is tight if and only if there exist bases of A, B, C and injective functions $s_A : \{1, \ldots, \dim A\} \to \mathbb{Z}, s_B :$ $\{1,\ldots,\dim B\}\to\mathbb{Z},\ s_C:\{1,\ldots,\dim C\}\to\mathbb{Z}\ \text{such that}$

$$s_A(i) + s_B(j) + s_C(k) = 0$$
 for every $(i, j, k) \in \text{supp}(T)$.

The following result was "known to the experts" but since we do not have a reference for it, we provide its proof.

LEMMA 5.1. The tensors $T_{cw,q}$ and $T_{CW,q}$ are tight.

Proof. Write q = 2u or q = 2u + 1 depending on the parity of q. Consider the change of basis

$$a_0 \mapsto a_0$$

$$a_j \mapsto \frac{\sqrt{2}}{2}(a_j + a_{u+j}) \quad \text{for } j = 1, \dots, u$$

$$a_{u+j} \mapsto \frac{\sqrt{-2}}{2}(a_j - a_{u+j}) \quad \text{for } j = 1, \dots, u$$

$$a_q \mapsto a_q \quad \text{(if } q \text{ is odd)}$$

$$a_{q+1} \mapsto a_{q+1}$$

and similarly on B and C.

After this change of basis, regarding $T_{cw,q}$ and $T_{CW,q}$ as symmetric tensors in S^3A , we have

$$T_{cw,2u} = a_0 \left(\sum_{j=1}^u a_j a_{u+j} \right),$$

$$T_{CW,2u} = a_0 \left(\sum_{j=1}^u a_j a_{u+j} \right) + a_0^2 a_{q+1},$$

1 Page 28 of 40 Conner et al. cc

or

$$T_{cw,2u+1} = a_0 \left(\sum_{j=1}^u a_j a_{u+j} + a_q^2 \right),$$

$$T_{CW,2u+1} = a_0 \left(\sum_{j=1}^u a_j a_{u+j} + a_q^2 \right) + a_0^2 a_{q+1},$$

depending on the parity of q.

Define $s = s_A = s_B = s_C$ by

$$s(0) = 2,$$

 $s(j) = 2 + j$ for $j = 1, ..., u,$
 $s(u + j) = -j - 4$ for $j = 1, ..., u,$
 $s(q) = -1$ if q is odd,
 $s(q + 1) = -4.$

It is easy to verify that s(i) + s(j) + s(k) = 0 if $(i, j, k) \in \text{supp}(T_{CW,q})$. Moreover, since $\text{supp}(T_{cw,q}) \subseteq \text{supp}(T_{CW,q})$, the same holds for $(i, j, k) \in \text{supp}(T_{cw,q})$. This concludes the proof.

The combinatorial characterization of tightness makes it clear that this property only depends on the support of a tensor in a given basis; we say that a support S is tight if every tensor having support S is tight.

A tensor $T \in A \otimes B \otimes C$ is *concise* if the induced linear maps $T_A : A^* \to B \otimes C$, $T_B : B^* \to A \otimes C$, $T_C : C^* \to A \otimes B$ are injective. We say that a concise tensor $T \in \mathbb{C}^m \otimes \mathbb{C}^m \otimes \mathbb{C}^m$ has *minimal rank* (resp. *minimal border rank*) if $\mathbf{R}(T) = m$ (resp. $\mathbf{R}(T) = m$).

Given concise tensors $T_1 \in A_1 \otimes B_1 \otimes C_1$ and $T_2 \in A_2 \otimes B_2 \otimes C_2$, Conner *et al.* (2020a, Theorem 4.1) guarantees that

$$\mathfrak{g}_{T_1\boxtimes T_2}\supseteq\mathfrak{g}_{T_1}\otimes \operatorname{Id}_{A_2\otimes B_2\otimes C_2}+\operatorname{Id}_{A_1\otimes B_1\otimes C_1}\otimes\mathfrak{g}_{T_2};$$

moreover if $\mathfrak{g}_{T_1} = 0$ and $\mathfrak{g}_{T_2} = 0$ then equality holds $\mathfrak{g}_{T_1 \boxtimes T_2} = 0$.

The strict containment in (5.2) occurs, for instance, in the case of the matrix multiplication tensor. In Conner *et al.* (2020a), we posed the problem of characterizing tensors $T \in A \otimes B \otimes C$ such that $\mathfrak{g}_T \otimes \operatorname{Id}_{A \otimes B \otimes C} + \operatorname{Id}_{A \otimes B \otimes C} \otimes \mathfrak{g}_T$ is strictly contained in $\mathfrak{g}_{T^{\boxtimes 2}} \subset \mathfrak{gl}(A^{\boxtimes 2}) + \mathfrak{gl}(B^{\boxtimes 2}) + \mathfrak{gl}(C^{\boxtimes 2})$.

Proposition 5.4 provides several additional examples of tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ for which this containment is strict.

5.2. Tight supports in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$. From Conner *et al.* (2020a, Proposition 2.14), one obtains an exhaustive list of unextendable tight supports for tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$, up to the action of $\mathbb{Z}_2 \times \mathfrak{S}_3$, where \mathfrak{S}_3 acts permuting the factors and \mathbb{Z}_2 acts by reversing the order of the basis elements. In fact,

tightness is invariant under the action of the full \mathfrak{S}_3 acting by permutation on the basis vectors. This additional simplification, pointed out by J. Hauenstein, provides the following list of 9 unextendable tight supports up to the action of $((\mathfrak{S}_3)^{\times 3}) \rtimes \mathfrak{S}_3$.

```
T_1 = \{(1,1,3), (1,2,2), (2,1,2), (3,3,1)\};
T_2 = \{(1, 1, 3), (1, 3, 2), (2, 3, 1), (3, 2, 2)\}:
\mathcal{T}_3 = \{(1,1,3), (1,2,2), (1,3,1), (2,1,2), (3,2,1)\};
T_4 = \{(1,1,3), (1,2,2), (2,1,2), (2,3,1), (3,2,1)\};
T_5 = \{(1,1,3), (1,2,2), (2,3,1), (3,1,2), (3,2,1)\};
\mathcal{T}_6 = \{(1,1,3), (1,3,2), (2,2,2), (3,1,2), (3,3,1)\};
\mathcal{T}_7 = \{(1,1,3), (1,2,2), (1,3,1), (2,1,2), (2,2,1), (3,1,1)\};
\mathcal{T}_8 = \{(1,1,3), (1,3,2), (2,2,2), (2,3,1), (3,1,2), (3,2,1)\};
\mathcal{T}_9 = \{(1,2,3), (1,3,2), (2,1,3), (2,2,2), (2,3,1), (3,1,2), (3,2,1)\};
```

Supports S_2 and S_3 of Conner *et al.* (2020a) are equivalent to support $S_1 = T_1$; supports S_8 and S_{10} are equivalent to support $S_6 = T_4$.

The following result characterizes tight tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ up to isomorphism.

Proposition 5.3. Let $T \in \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ be a tight tensor with unextendable tight support in some basis. Then, up to permuting the three factors, T is isomorphic to exactly one of the following.

```
T_1 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_2 \otimes c_2 + a_2 \otimes b_1 \otimes c_2 + a_3 \otimes b_3 \otimes c_1
                   T_2 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_3 \otimes c_2 + a_2 \otimes b_3 \otimes c_1 + a_3 \otimes b_2 \otimes c_2
                   T_3 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_2 \otimes c_2 + a_1 \otimes b_3 \otimes c_1 + a_2 \otimes b_1 \otimes c_2 + a_3 \otimes b_2 \otimes c_1
                   T_4 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_2 \otimes c_2 + a_2 \otimes b_1 \otimes c_2 + a_2 \otimes b_3 \otimes c_1 + a_3 \otimes b_2 \otimes c_1
                   T_5 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_2 \otimes c_2 + a_2 \otimes b_3 \otimes c_1 + a_3 \otimes b_1 \otimes c_2 + a_3 \otimes b_2 \otimes c_1
                   T_6 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_3 \otimes c_2 + a_2 \otimes b_2 \otimes c_2 + a_3 \otimes b_1 \otimes c_2 + a_3 \otimes b_3 \otimes c_1
                   T_7 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_2 \otimes c_2 + a_1 \otimes b_3 \otimes c_1 + a_2 \otimes b_1 \otimes c_2 + a_2 \otimes b_2 \otimes c_1 + a_3 \otimes b_4 \otimes c_4 + a_4 \otimes b_4 \otimes c_5 + a_5 \otimes c_5 \otimes c_6 \otimes c_6
                   T_8 := a_1 \otimes b_1 \otimes c_3 + a_1 \otimes b_3 \otimes c_2 + a_2 \otimes b_2 \otimes c_2 + a_2 \otimes b_3 \otimes c_1 + a_3 \otimes b_1 \otimes c_2 + a_3 \otimes b_3 \otimes c_4 + a_3 \otimes b_4 \otimes c_5 + a_4 \otimes b_5 \otimes c_5 + a_5 \otimes b_5 \otimes c_5 \otimes c_5 + a_5 \otimes b_5 \otimes c_5 \otimes c_5 + a_5 \otimes b_5 \otimes c_5 \otimes c_5 \otimes c_5 + a_5 \otimes b_5 \otimes c_5 \otimes c_5
T_{9,\mu} := a_1 \otimes b_2 \otimes c_3 + a_1 \otimes b_3 \otimes c_2 + a_2 \otimes b_1 \otimes c_3 + a_2 \otimes b_2 \otimes c_2 + a_2 \otimes b_3 \otimes c_1 + a_3
```

The result of Conner et al. (2020a, Proposition 2.14) and the discussion above shows that T is, up to permutation of the factors, equivalent to a tensor with support \mathcal{T}_i for some $i = 1, \dots, 9$.

 $\otimes b_1 \otimes c_2 + \mu \cdot a_3 \otimes b_2 \otimes c_1$ for some $\mu \in \mathbb{C} \setminus \{0\}$.

For i = 1, ..., 8, it is straightforward to verify that all tensors with support \mathcal{T}_i are isomorphic, via the change of bases given by three diagonal matrices.

The case of \mathcal{T}_9 is slightly more involved but essentially the same argument shows that a tensor T with support \mathcal{T}_9 is isomorphic to $T_{9,\mu}$, for some μ .

Finally, we have to show that any two of the tensors in the statement are not isomorphic. For tensors having distinct supports, this is a consequence of Proposition 5.4 below: indeed, if T, T' are two of the tensors above, Proposition 5.4 shows that either $\dim \mathfrak{g}_T \neq \dim \mathfrak{g}_{T'}$ or $\dim \mathfrak{g}_{T^{\boxtimes 2}} \neq \dim \mathfrak{g}_{T'^{\boxtimes 2}}$.

As for the tensors with support T_9 , we proceed as follows. Let $T = T_{9,\mu}$ and $T' = T_{9,\mu'}$ with $\mu \neq \mu'$. We show that T is not isomorphic to T'. Suppose by contradiction that there is a triple of 3×3 matrices $g = (g_A, g_B, g_C) \in GL_3 \times GL_3 \times GL_3$ with g(T) = T'. One sees that in each case, g_A, g_B, g_C have to be diagonal matrices, and an explicit calculation shows that there is no triple of diagonal matrices such that g(T) = T'.

We point out that T_7 is isomorphic to the Coppersmith-Winograd tensor $T_{CW,1}$, as well as to the structure tensor of the algebra $\mathbb{C}[x]/(x^3)$.

The tensors $T_{cw,2}$ and $T_{skewcw,2}$ are degenerations of $T_{9,\mu}$, respectively, for $\mu = 1$ and $\mu = -1$. In particular, they do not have an unextendable tight support in some basis.

PROPOSITION 5.4. For $i=1,\ldots,9$, the following table records $\dim \mathfrak{g}_{T_i}$ and $\dim \mathfrak{g}_{T^{\boxtimes 2}}$.

T	$\dim \mathfrak{g}_T$	$\dim \mathfrak{g}_{T^{\boxtimes 2}}$
T_1	5	22
T_2	3	9
T_3	5	13
T_4	4	9
T_5	3	7
T_6	2	5
T_7	6	28
T_8	1	2
$T_{9,-1}$	5	10
$T_{9,\mu}$ (for $\mu \neq 0,-1$)	1	2

In summary

 $\dim \mathfrak{g}_{T^{\boxtimes 2}} > 2 \dim \mathfrak{g}_T$

for tight tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ with unextendable tight supports $\mathcal{T}_1, \ldots, \mathcal{T}_7$.

PROOF. For T_1, \ldots, T_8 and for the $T_{9,-1}$, the proof follows by a direct calculation. The first part of the Macaulay2 file symmetryTightSupports.m2 in Appendix E of Supplementary Material computes the dimension of the symmetry algebras of interest in these cases.

The second part of the file deals with the case $T_{9,\mu}$ when $\mu \neq -1$. By tightness, $\dim \mathfrak{g}_{T_{9,\mu}} \geq 1$.

Consider the linear map $\omega_{T_{9,\mu}}: \mathfrak{gl}(A)+\mathfrak{gl}(B)+\mathfrak{gl}(C) \to A\otimes B\otimes C$ defined by $(X,Y,Z)\mapsto (X,Y,Z).T_{9,\mu}$. Then $\mathfrak{g}_{T_{9,\mu}}=[\ker(\omega_{T_{9,\mu}})]/\mathbb{C}^2$, where \mathbb{C}^2 corresponds to $\ker d\Phi$.

The second part of the file symmetryTightSupports.m2 computes a matrix representation of $\omega_{T_{9,\mu}}$, depending on a parameter μ (t in the file). Let F_{μ} be this 27×27 matrix representation. Then, it suffices to select a 24×24 submatrix whose determinant is a nonzero univariate polynomial in μ . If μ is a value for which $\dim \mathfrak{g}_{T_{9,\mu}} > 1$, then μ has to be a root of this univariate polynomial.

In the example computed in the file, we select a 24×24 submatrix whose determinant is $(\mu+1)^6\mu$, showing that the only possible values of μ for which $\dim \mathfrak{g}_{T_{9,\mu}} > 1$ are $\mu = 0$ or $\mu = -1$. The case $\mu = -1$ was considered separately. The case $\mu = 0$ does not correspond to a unextendable support, so it is not of interest. We point out that however, $\operatorname{rank}(\omega_{T_{9,0}}) = 24$, namely $\dim \mathfrak{g}_{T_{9,0}} = 1$.

For $T_{9,\mu}^{\boxtimes 2}$, we follow essentially the same argument. By tightness, and (5.2), we obtain $\dim \mathfrak{g}_{T_{9,\mu}^{\boxtimes 2}} \geq 2$. The third part of symmetryTightSupports.m2 computes a matrix representation of the map $\omega_{T_{9,\mu}^{\boxtimes 2}}$, depending on a parameter μ : this is a 729×243 matrix of rank at most 239.

In the example computed in the file, we select a 239 \times 239 submatrix whose determinant is the univariate polynomial $\mu^{8}(\mu+1)^{12}$. As before, we conclude.

We also provide the values of the border rank of the tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ having unextendable tight support and numerical evidence for the values of border rank of their Kronecker square. They are recorded in the following table. The values of the border rank for the T_i 's are straightforward to verify. The lower bounds for the Kronecker squares are obtained via Koszul flattenings. In the cases labeled by N/A the upper bounds coincide with the multiplicative upper bound; in the other cases, the upper bound is obtained via numerical methods, and the last column of the table records the ℓ_2 distance (in the given basis) between the tensor obtained via the numerical approximation and the Kronecker square. The numerical approximations are recorded in supplementary files in Appendix F of Supplementary Material.

1 Page 32 of 40 Conner et al.

T	$\mathbf{R}(T)$	$\underline{\mathbf{R}}(T^{\boxtimes 2})$	ℓ_2 error for upper bound in $T^{\boxtimes 2}$ decomposition
T_1	3	9	N/A
T_2	4	[11, 14]	0.000155951
T_3	4	[11, 14]	0.00517612
T_4	4	14	0.0144842
T_5	4	[11, 15]	0.0237172
T_6	4	[11, 15]	0.00951205
T_7	3	9	N/A
T_8	4	[14, 16]	N/A
$T_{9,-1}$	5	[16, 19]	0.0231353
$T_{9,\mu}$ (for $\mu \neq 0, -1$)	4	[15, 16]	N/A

6. A method to compute flattenings of structured tensors

In this section, we explain how to compute the matrices Φ_1, \ldots, Φ_4 in Section 3.4 and the matrices Ψ_1, \ldots, Ψ_8 in Section 3.5.

The matrices Φ_1, \ldots, Φ_4 and Ψ_1, \ldots, Ψ_8 arise via a series of tensor contractions of highly structured tensors. In this section, we introduce the notion of *box* parametrized sequence of tensors. Lemma 6.2 below shows that contraction of box parametrized tensors gives rise to box parametrized tensors; in addition, the expression of the tensors resulting from the contraction is particularly easy to control.

We will then show that the tensors in Section 3.4 and Section 3.5 which give rise to the matrices Φ_1, \ldots, Φ_4 and Ψ_1, \ldots, Ψ_8 are box parametrized. This allows us to track down the entries of the final matrices as functions of the dimension q.

The full calculation of the matrices is left to the scripts available in Appendix D of Supplementary Material.

The point of view is partially inspired by the interpretation of tensors in communication models, where a tensor on k factors is regarded as a function from $\underbrace{\mathbb{N} \times \cdots \times \mathbb{N}}_{k} \to \mathbb{C}$ with finite support sending a k-tuple of integers to the cor-

responding coefficient of the tensor. Explicitly, for every $j=1,\ldots,k$ fix a basis $\{v_i^{(j)}\}$ on the j-th factor: given a finite support $\Sigma\subseteq\mathbb{N}^{\times k}$, the tensor $T=\sum_{(i_1,\ldots,i_k)\in\Sigma}t_{i_1,\ldots,i_k}v_{i_1}^{(1)}\otimes\cdots\otimes v_{i_k}^{(k)}$ corresponds to the function defined by $T(i_1,\ldots,i_k)=t_{i_1,\ldots,i_k}$. We do not explicitly write the dimensions of the factors.

Let $\mathcal{T}=\{T_q:q\in\mathbb{N}\}$ be a sequence of tensors of order k. We say that \mathcal{T} is

basic box-parametrized if, for every q

$$T_q = p(q) \sum_{(i_1, \dots, i_k) \in \Sigma_q} v_{i_1}^{(1)} \otimes \dots \otimes v_{i_k}^{(k)}$$

where p(q) is a univariate polynomial in q and the support Σ_q is defined by conditions $\eta_i q + \vartheta_i \leq i_i \leq H_i q + \Theta_i$ for $\eta_i, H_i \in \{0,1\}$ and $\vartheta_i, \Theta_i \in \mathbb{Z}_{>0}$, and any number (not depending on q) of equalities $i_i = i_{i'}$ among indices. Without loss of generality, assume that the inequalities are sharp for every j, in the sense that for every i_j satisfying the j-th inequality, the basis element $v_{i_j}^{(j)}$ does appear in T_q . We often say that \mathcal{T} is basic box-parametrized for $q \geq q_0$ for some q_0 , in the sense that the sequence has the desired structure for $q \geq q_0$.

Example 6.1. The sequence $T_q = v_0^{(1)} \otimes \sum_{i=1}^q v_i^{(2)} \otimes v_i^{(3)}$ is basic box-parametrized for $q \geq 1$, with support Σ_q defined by the conditions

$$0 \le i_1 \le 0$$
, $1 \le i_2 \le q$, $1 \le i_2 \le q$, $i_2 = i_3$.

 \Diamond

We define a contraction operation between the j_1 -th and the j_2 -th factor of \mathcal{T} , obtained by summing over the corresponding indices: in other words, the contraction is the image of T via the trace map $\sum u_i^{(j_1)} \otimes u_i^{(j_2)}$ applied to the j_1 -th and j_2 -th factors, where $\{u_i^{(j)}\}$ is the dual basis to the fixed basis $\{v_i^{(j)}\}$ on the j-th factor.

LEMMA 6.2. Let \mathcal{T} , \mathcal{T}' be basic box-parametrized tensors for $q \geq q_0$ and $q \geq q'_0$ respectively. Then

- $\circ \mathcal{T} \otimes \mathcal{T}'$ is basic box-parametrized for $q \geq \max\{q_0, q_0'\}$;
- \circ the contraction of \mathcal{T} on factors j_1 and j_2 is basic box-parametrized for $q \ge \max\{|\vartheta_{j_1} - \vartheta_{j_2}|, |\Theta_{j_1} - \Theta_{j_2}|, q_0\};$ moreover, if the univariate coefficient p(q) of T is a polynomial of degree e, then the coefficient of the tensor resulting from the contraction has degree at most e + 1.

Proof. The first statement is immediate.

For the second statement, without loss of generality assume $j_1 = 1$ and $j_2 = 2$. First observe that if \mathcal{T} is basic box-parametrized, then summing over the first index, or equivalently applying the linear map $\sum_i u_i^{(1)}$, generates a basic boxparametrized tensor; the coefficient of this tensor has the same degree as the coefficient of \mathcal{T} unless the first index i_1 is not related by equality to any other index, and $\eta_1 = 0$ and $H_1 = 1$; in the latter case, the degree of the coefficient is increased by one.

1 Page 34 of 40 Conner et al. cc

Now, contraction of \mathcal{T} on factors 1 and 2 is equivalent to first imposing the equality $i_1=i_2$ on the support Φ_q and then summing up on the first and second index. Imposing the equality $i_1=i_2$ effects the inequalities of i_1 and i_2 as follows:

$$\max\{\eta_1 q + \vartheta_1, \eta_2 q + \vartheta_2\} \le i_1 = i_2 \le \min\{H_1 q + \Theta_1, H_2 q + \Theta_2\}.$$

Each of the two bounds can be replaced by one of the two linear functions (uniformly in q) whenever $q \geq \{|\vartheta_1 - \vartheta_2|, |\Theta_1 - \Theta_2|\}$. This, together with the previous observation, concludes the proof.

Given two sequences of tensors $\mathcal{T}^{(1)}$, $\mathcal{T}^{(2)}$ of order k, we define their sum as $\mathcal{T}_1 + \mathcal{T}_2 = \{T_q^{(1)} + T_q^{(2)} : q \in \mathbb{N}\}$. We say that a sequence \mathcal{T} is box parametrized (for $q \geq q_0$) if \mathcal{T} is a finite sum of basic box-parametrized sequences of tensors (for $q \geq q_0$). Observe that a sequence of tensors with constant dimensions is box parametrized if and only if its coefficients are univariate polynomials in q.

We will show that the maps Φ_1, \ldots, Φ_4 in the proof of Theorem 3.5 in Section 3.4 and the maps Ψ_1, \ldots, Ψ_8 in the proof of Theorem 3.7 in Section 3.5 are box parametrized.

The scripts in Appendix D perform the contraction of box parametrized tensors according to Lemma 6.2, keeping track of the univariate polynomial coefficients and of the lower bound q_0 for which the expressions are valid. The final result is that the maps Φ_1, \ldots, Φ_4 are box parametrized for $q \geq 5$ and the maps Ψ_1, \ldots, Ψ_8 are box parametrized for $q \geq 6$.

In the following, we show that the tensors involved in the various contractions are box parametrized. Lemma 6.2 guarantees that the results of the contractions are box parametrized as well.

First, notice that $T_{cw,q}$ is box parametrized for $q \geq 1$, as it is the sum of three tensors as the ones described in Example 6.1. By Lemma 6.2, we deduce that $T_{cw,q}^{\otimes 2}$ (regarded as a tensor of order 6) and $T_{cw,q}^{\otimes 3}$ (regarded as a tensor of order 9) are box parametrized. In all three cases, the polynomials defining the coefficients have degree 0.

6.1. Restriction. We show that the two restriction maps $\phi_2: A^{\otimes 2} \to \mathbb{C}^3$ and $\phi_3: A^{\otimes 3} \to \mathbb{C}^5$ are box parametrized as tensors of order 3 and 4 respectively.

Write $\phi_2 = X_0 \otimes e_0 + X_1 \otimes e_1 + X_2 \otimes e_2$, where $\mathbb{C}^3 = \langle e_0, e_1, e_2 \rangle$ and $X_0, X_1, X_2 \in A^{\otimes 2^*}$. It suffices to show that X_0, X_1, X_2 are box parametrized, regarded as tensors of order two. Using a basis dual to the basis of $A^{\otimes 2}$, we have

$$X_0 = \alpha_0 \otimes \alpha_1 + \alpha_1 \otimes \alpha_0 + \alpha_1 \otimes \alpha_1$$

$$X_1 = \alpha_0 \otimes \sum_{i=1}^{q} \alpha_i + \sum_{i=1}^{q} \alpha_i \otimes \alpha_0$$

$$X_2 = \alpha_0 \otimes \alpha_2 + \alpha_2 \otimes \alpha_0 + \alpha_2 \otimes \alpha_1 + \alpha_3 \otimes \alpha_3.$$

This shows that X_0, X_1, X_2 are box parametrized.

Similarly, write $\phi_3 = Y_0 \otimes e_0 + \cdots + Y_4 \otimes e_4$, where $\mathbb{C}^5 = \langle e_0, \dots, e_4 \rangle$ and $Y_0, \ldots, Y_4 \in A^{\otimes 3^*}$. Directly from the definition in Section 3.5, it is immediate that Y_0, \ldots, Y_4 are box parametrized and therefore ϕ_3 is box parametrized as well.

Applying Lemma 6.2, we deduce that the two sequences $\phi_2(T_{cw,a}^{\otimes 2})$ and $\phi_3(T_{cw,a}^{\otimes 3})$ are box parametrized.

6.2. Koszul maps. The Koszul differentials on \mathbb{C}^3 and \mathbb{C}^5 used in the definition of the Koszul flattenings are the skew-symmetric projections $\mathbb{C}^3 \otimes \mathbb{C}^3 \to$ $\Lambda^2\mathbb{C}^3$ and $\Lambda^2\mathbb{C}^5\otimes\mathbb{C}^5\to\Lambda^3\mathbb{C}^5$. They are both fixed size, therefore they are box parametrized.

By Lemma 6.2, we deduce that the resulting Koszul flattenings $(\phi_2(T_{cw,q}^{\boxtimes 2}))^{\wedge 1}$ and $(\phi_3(T_{cw,q}^{\boxtimes 3}))^{\wedge 2}$ are box parametrized, regarded as tensors of order 6 and 8 respectively.

6.3. Diagonalizing maps. Recall that the maps Φ_1, \ldots, Φ_4 in the proof of Theorem 3.5 and the maps Ψ_1, \ldots, Ψ_8 in the proof of Theorem 3.7 are the restrictions of $(\phi_2(T_{cw,q}^{\boxtimes 2}))^{\wedge 1}$ and $(\phi_3(T_{cw,q}^{\boxtimes 3}))^{\wedge 2}$ to the multiplicity spaces of the isotypic components for the action of \mathfrak{S}_{q-3} and \mathfrak{S}_{q-5} .

We analyze the square case in detail. For the square case, let \mathcal{M} be the matrix of change of basis on \mathbb{C}^q from the basis $\{e_1,\ldots,e_q\}$ to the basis $\{e_1, e_2, e_3, \sum_{4}^{q} e_i, e_5 - e_4, \dots, e_q - e_{q-1}\}$. Explicitly

$$\mathcal{M} = \begin{bmatrix} \text{Id}_3 & & & & \\ & 1 & 1 & \cdots & 1 \\ & -1 & 1 & & \\ & & \ddots & \ddots & \\ & & & -1 & 1 \end{bmatrix}.$$

In particular, \mathcal{M} diagonalizes the action of \mathfrak{S}_{q-3} and therefore the change of basis defined by $\mathrm{Id}_{\mathbb{C}^3} \boxtimes \mathcal{M}^{\boxtimes 2}$ on $\mathbb{C}^3 \otimes B^{\otimes 2}$ brings the matrix representing $(\phi_2(T_{cw,q}^{\boxtimes 2}))^{\wedge 1}$ into a block diagonal matrix, whose diagonal blocks are matrices representing the maps $f_i: \mathbb{C}^3 \otimes (M_i \otimes W_i) \to \Lambda^2 \mathbb{C}^3 \otimes (M_i \otimes W_i)$ from (3.6); denote the diagonal blocks by $f_1^{\mathcal{M}}, \ldots, f_4^{\mathcal{M}}$.

Because of our choice of basis, the multiplicity subspaces $\mathbb{C}^3 \otimes \langle w_i \rangle \otimes M_i$ and $\Lambda^2 \mathbb{C}^3 \otimes \langle w_i \rangle \otimes M_i$ described in Section 3.4 are spanned by basis vectors, so that the matrices representing Φ_1, \ldots, Φ_4 are given by submatrices of $f_1^{\mathcal{M}}, \ldots, f_4^{\mathcal{M}}$. More precisely, setting π_{inv} , π_{std} to be the matrices of the two coordinate 1 Page 36 of 40 Conner et al. cc

projections of \mathbb{C}^q onto $\langle e_1, \ldots, e_4 \rangle$ and $\langle e_5 \rangle$, we have

$$\Phi_{1} = (\operatorname{Id}_{\Lambda^{2}\mathbb{C}^{3}} \boxtimes \pi_{inv} \boxtimes \pi_{inv}) \circ f_{1}^{\mathcal{M}} \circ (\operatorname{Id}_{\mathbb{C}^{3}} \boxtimes \pi_{inv} \boxtimes \pi_{inv})^{T},
\Phi_{2} = (\operatorname{Id}_{\Lambda^{2}\mathbb{C}^{3}} \boxtimes \pi_{inv} \boxtimes \pi_{std}) \circ f_{2}^{\mathcal{M}} \circ (\operatorname{Id}_{\mathbb{C}^{3}} \boxtimes \pi_{inv} \boxtimes \pi_{std})^{T},
\Phi_{3} = (\operatorname{Id}_{\Lambda^{2}\mathbb{C}^{3}} \boxtimes \pi_{std} \boxtimes \pi_{inv}) \circ f_{3}^{\mathcal{M}} \circ (\operatorname{Id}_{\mathbb{C}^{3}} \boxtimes \pi_{std} \boxtimes \pi_{inv})^{T},
\Phi_{4} = (\operatorname{Id}_{\Lambda^{2}\mathbb{C}^{3}} \boxtimes \pi_{std} \boxtimes \pi_{std}) \circ f_{4}^{\mathcal{M}} \circ (\operatorname{Id}_{\mathbb{C}^{3}} \boxtimes \pi_{std} \boxtimes \pi_{std})^{T}.$$

Since the composition can be performed on the single factors, by Lemma 6.2 it suffices to show that the four matrices $\mathcal{M}^{-1} \circ \pi_{inv}^T$, $\mathcal{M}^{-1} \circ \pi_{std}^T$, $\pi_{inv} \circ \mathcal{M}$ and $\pi_{std} \circ \mathcal{M}$ are box parametrized.

From the structure of \mathcal{M} , it is clear that $\pi_{inv} \circ \mathcal{M}$ and $\pi_{std} \circ \mathcal{M}$ are box parametrized. The computation of \mathcal{M}^{-1} is straightforward, and it is easy to see that $\mathcal{M}^{-1} \circ \pi_{inv}^T$, $\mathcal{M}^{-1} \circ \pi_{std}^T$ are box parametrized.

This shows that Φ_1, \ldots, Φ_4 are box parametrized. The script available in Appendix D computes the box parametrized representation of Φ_1, \ldots, Φ_4 starting from the box parametrized version of T_{cw} , the restriction map ϕ_2 , the Koszul differential and the four matrices $\mathcal{M}^{-1} \circ \pi_{inv}^T$, $\mathcal{M}^{-1} \circ \pi_{std}^T$, $\pi_{inv} \circ \mathcal{M}$ and $\pi_{std} \circ \mathcal{M}$.

The cube case is similar. Now, restriction space \mathbb{C}^3 is a \mathbb{C}^5 , the top left block in the matrix \mathcal{M} is a 5×5 identity block, the result of the conjugation by \mathcal{M} is block diagonal with 8 blocks, corresponding to the eight isotypic components. The coordinate projections π_{inv} and π_{std} are onto $\langle e_1, \ldots, e_6 \rangle$ and $\langle e_7 \rangle$. The script computes the box parametrized representation of the matrices Ψ_1, \ldots, Ψ_8 .

Acknowledgements

Landsberg supported by NSF grant AF-1814254. Gesmundo acknowledges support from VILLUM FONDEN via the QMATH Centre of Excellence (Grant no. 10059). We thank the anonymous referees for their careful reading of the paper and useful suggestions.

Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is

not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Funding

Open Access funding enabled and organized by Projekt DEAL

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- J. Alman (2019). Limits on the Universal Method for Matrix Multiplication. In 34th Comp. Compl. Conf. (CCC 2019). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik.
- J. Alman & V. V. Williams (2018a). Further Limitations of the Known Approaches for Matrix Multiplication. In 9th Innov. Th. Comp. Science Conf., ITCS 2018, January 11–14, 2018, Cambridge, MA, USA, 25:1–25:15.
- J. Alman & V. V. Williams (2018b). Limits on all known (and some unknown) approaches to matrix multiplication. In 2018 IEEE 59th Ann. Symp. Found. Comp. Sc. (FOCS), 580-591.
- J. Alman & V. V. Williams (2021). A refined laser method and faster matrix multiplication. In *Proc.* 2021 ACM-SIAM Symp. Disc. Alq. (SODA), 522–539. SIAM.
- A. Ambainis, Y. Filmus & F. Le Gall (2015). Fast matrix multiplication: limitations of the Coppersmith–Winograd method. In *Proc. of the 47th ACM Symp. Th. Comp.*, 585–593. ACM.
- E. Ballico, A. Bernardi, M. Christandl & F. Gesmundo (2019). On the partially symmetric rank of tensor products of W-states and other symmetric tensors. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **30**, 93–124.
- D. Bini (1980). Relations between exact and approximate bilinear algorithms. Applications. *Calcolo* 17(1), 87–97.
- D. Bini, G. Lotti & F. Romani (1980). Approximate solutions for the bilinear form computational problem. SIAM J. Comput. 9(4), 692–697.
- M. Bläser (2013). Fast Matrix Multiplication. Theory of Computing, Graduate Surveys 5, 1–60.
- M. Bläser & V. Lysikov (2016). On degeneration of tensors and algebras. In 41st International Symposium on Mathematical Foundations of Computer Science, volume 58 of LIPIcs. Leibniz Int. Proc. Inform., Art. No. 19, 11. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern.

1 Page 38 of 40 Conner et al.

- P. BÜRGISSER, M. CLAUSEN & M. A. SHOKROLLAHI (1997). Algebraic complexity theory, volume 315 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin. ISBN 3-540-60582-7.
- M. CHRISTANDL, F. GESMUNDO & A. K. JENSEN (2019). Border rank is not multiplicative under the tensor product. SIAM J. Appl. Alq. Geom. 3, 231–255.
- M. Christandl, A. K. Jensen & J. Zuiddam (2018). Tensor rank is not multiplicative under the tensor product. *Lin. Alg. Appl.* **543**, 125–139.
- M. Christandl, P. Vrana & J. Zuiddam (2021). Barriers for Fast Matrix Multiplication from Irreversibility. *Theory of Computing* 17(2), 1–32.
- H. Cohn & C. Umans (2003). A group theoretic approach to fast matrix multiplication. In *Proc. of the 44th Symp. Found. Comp. Science*, number 2, 438–449.
- A. Conner, F. Gesmundo, J. M. Landsberg & E. Ventura (2019a). Tensors with maximal symmetries. arXiv:1909.09518.
- A. Conner, F. Gesmundo, J. M. Landsberg, E. Ventura & Y. Wang (2021) Towards a geometric approach to Strassen's asymptotic rank conjecture. Collectanea Mathematica $\bf 72(1)$, 63–86
- A. Conner, A. Harper & J. M. Landsberg (2019b). New lower bounds for matrix multiplication and the 3×3 determinant. arXiv:1911.07981.
- A. Conner, H. Huang & J. M. Landsberg (2020b). Bad and good news for Strassen's laser method: Border rank of the 3×3 permanent and strict submultiplicativity. arXiv:2009.11391.
- D. COPPERSMITH & S. WINOGRAD (1990). Matrix multiplication via arithmetic progressions. J. Symb. Comput. 9(3), 251–280.
- H. Derksen (2016). On the nuclear norm and the singular value decomposition of tensors. Found. Comp. Math. 16(3), 779–811.
- H. Derksen & V. Makam (2018). On non-commutative rank and tensor rank. Linear Multilinear Algebra **66**(6), 1069–1084.
- K. Efremenko, A. Garg, R. Oliveira & A. Wigderson (2018). Barriers for rank methods in arithmetic complexity. In 9th Innovations in Theoretical Computer Science, volume 94 of LIPIcs. Leibniz Int. Proc. Inform., Art. No. 1, 19. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern.
- W. Fulton (1997). Young tableaux. With applications to representation theory and geometry, volume 35 of London Mathematical Society Student Texts. Cambridge University Press, Cambridge. ISBN 0-521-56144-2; 0-521-56724-6, x+260.
- W. Fulton & J. Harris (1991). Representation theory: a first course, volume 129 of Graduate Texts in Mathematics. Springer-Verlag, New York.
- D. A. GAY (1976). Characters of the Weyl group of SU(n) on zero weight spaces and centralizers of permutation representations. Rocky Mountain J. Math. $\mathbf{6}(3)$, 449–455.

- F. GESMUNDO, C. IKENMEYER & G. PANOVA (2017). Geometric complexity theory and matrix powering. *Diff. Geom. Appl.* **55**, 106–127.
- D. R. Grayson & M. E. Stillman (2020). Macaulay 2, a software system for research in algebraic geometry (Version 1.16. Available at http://www.math.uiuc.edu/Macaulay2/.
- N. ILTEN & Z. TEITLER (2016). Product ranks of the 3×3 determinant and permanent. Canad. Math. Bull. **59**(2), 311–319.
- G. JOHNS & Z. TEITLER Z (2020). An improved upper bound for the Waring rank of the determinant. arXiv:2004.06158
- S. KOPPARTY, G. MOSHKOVITZ & J. ZUIDDAM (2020). Geometric Rank of Tensors and Subrank of Matrix Multiplication. In 35th Comp. Compl. Conf. (CCC 2020), volume 169 of LIPIcs. Leibniz Int. Proc. Inform., 35:1–35:21. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, Dagstuhl, Germany.
- J. M. Landsberg (2012). Tensors: Geometry and Applications, volume 128 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI.
- J. M. LANDSBERG (2017). Geometry and complexity theory, volume 169 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge.
- J. M. LANDSBERG & G. OTTAVIANI (2013). Equations for secant varieties of Veronese and other varieties. *Ann. Mat. Pura Appl.* (4) **192**(4), 569–606. ISSN 0373-3114.
- J. M. LANDSBERG & G. OTTAVIANI (2015). New lower bounds for the border rank of matrix multiplication. *Th. of Comp.* **11**(11), 285–298.
- F. Le Gall (2014). Powers of tensors and Fast Matrix Multiplication. In Proc.~39th Int. Symp. Symb. Alg. Comp., 296–303. ACM.
- A. K. Lenstra, H. W. Lenstra Jr & L. Lovász (1982). Factoring Polynomials with Rational Coefficients. *Math. Ann* **261**(4), 515-534.
- K. D. Mulmuley & M. Sohoni (2008). Geometric Complexity Theory. II. Towards explicit obstructions for embeddings among class varieties. $SIAM\ J.\ Comput.\ 38(3),\ 1175-1206.$ ISSN 0097-5397.
- SAGE DEVELOPERS (2020). SageMath, the Sage Mathematics Software System (Version 9.0). Available at https://www.sagemath.org.
- A. SCHÖNHAGE (1981). Partial and total matrix multiplication. SIAM J. Comp. 10(3), 434–455.
- A. Stothers (2010). On the Complexity of Matrix Multiplication. Ph.D. thesis, U. Edinburgh.
- V. Strassen (1969). Gaussian elimination is not optimal. *Numerische mathematik* **13**(4), 354–356.
- V. STRASSEN (1983). Rank and optimal computation of generic tensors. Lin. Alg. Appl. 52/53, 645-685. ISSN 0024-3795.

1 Page 40 of 40 Conner et al.

- V. Strassen (1987). Relative bilinear complexity and matrix multiplication. J. Reine Angew. Math. 375/376, 406–443.
- V. STRASSEN (1988). The asymptotic spectrum of tensors. *J. Reine Angew. Math.* **384**, 102–152.
- V. STRASSEN (1991). Degeneration and complexity of bilinear maps: some asymptotic spectra. J. Reine Angew. Math. 413, 127–180. ISSN 0075-4102. URL https://doi.org/10.1515/crll.1991.413.127.
- V. STRASSEN (1994). Algebra and complexity. In First European Congress of Mathematics Paris, July 6–10, 1992, 429–446. Springer.
- V. V. WILLIAMS (2012). Multiplying matrices faster than Coppersmith–Winograd. In *Proc. 44th ACM Symp. Th. Comp. STOC'12*, 887–898. ACM.

Manuscript received June 17, 2020

Austin Conner

Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

(current) Department of Mathematics, Harvard University, Cambridge, MA 02138, USA

connerad@math.tamu.edu

Joseph M. Landsberg Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA jml@math.tamu.edu

Fulvio Gesmundo

Dept. Math. Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen O., Denmark
(current) Max Planck Institute for Mathematics in the Sciences, Inselstrasse 22, 04103 Leipzig, Germany gesmundo@mis.mpg.de

EMANUELE VENTURA

Mathematisches Institut, Universität Bern, Sidlerstrasse 5, 3012, Bern, Switzerland

(current) Università di Torino, Dipartimento di Matematica, via Carlo Alberto 10, 10123 Torino, Italy emanuele ventura@unito.it