

Optics Letters

Hyperuniform scalar random fields for lensless, multispectral imaging systems: erratum

YUYAO CHEN,¹  WESLEY A. BRITTON,² AND LUCA DAL NEGRO^{1,2,3,*}

¹Department of Electrical and Computer Engineering and Photonics Center, Boston University, 8 Saint Mary's Street, Boston, Massachusetts 02215, USA

²Division of Materials Science and Engineering, Boston University, 15 Saint Mary's Street, Brookline, Massachusetts 02246, USA

³Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215, USA

*Corresponding author: dalnegro@bu.edu

Received 23 March 2022; posted 24 March 2022; published 4 April 2022

We present an erratum to our Letter [*Opt. Lett.* **46**, 5360 (2021)]. This erratum refers to Fig. 3, where a previous version was wrongly uploaded during the final resubmission of the paper. This correction has no influence on the text, the results, and the conclusions of the original Letter. © 2022 Optica Publishing Group

<https://doi.org/10.1364/OL.458682>

There was an unintended error in Fig. 3 that did not appear in its final version in the original Letter [1]. The correct version

of Fig. 3 is now shown here in Fig. 1. We note that this figure is now consistent with the published text of the paper and that neither the results nor the conclusions of the original Letter are affected by this correction.

Funding. Army Research Laboratory (W911NF-12-2-0023); National Science Foundation (ECCS-2015700).

Disclosures. The authors declare no conflicts of interest.

REFERENCE

1. Y. Chen, W. A. Britton, and L. Dal Negro, *Opt. Lett.* **46**, 5360 (2021).




Fig. 1. (a) Azimuthally averaged and (b) radially averaged MTFs for the different lensless PSFs indicated in the legend. (c) Isotropic parameter γ computed for different lensless PSFs considering 50 realizations of uniformly random initial conditions. (d) Sparsity of the CH-based PSF with respect to the model parameter ϵ in the CH equation.