

The IGRINS YSO Survey I. Stellar parameters of pre-main sequence stars in Taurus-Auriga

RICARDO LÓPEZ-VALDIVIA,¹ KIMBERLY R. SOKAL,¹ GREGORY N. MACE,¹ BENJAMIN T. KIDDER,¹ MARYAM HUSSAINI,¹ LARISSA NOFI,^{2,3} L. PRATO,² CHRISTOPHER M. JOHNS-KRULL,⁴ HEERYOUNG OH,^{1,5} JAE-JOON LEE,⁵ CHAN PARK,⁵ JAE SOK OH,⁵ ADAM KRAUS,¹ KYLE F. KAPLAN,⁶ JOE LLAMA,² ANDREW W. MANN,⁷ HWIHYUN KIM,⁸ MICHAEL A. GULLY-SANTIAGO,¹ HYE-IN LEE,⁵ SOOJONG PAK,⁹ NARAE HWANG,⁵ AND DANIEL T. JAFFE¹

¹The University of Texas at Austin, Department of Astronomy, 2515 Speedway, Stop C1400, Austin, TX 78712-1205

²Lowell Observatory, 1400 W. Mars Hill Rd, Flagstaff, AZ 86001, USA

³Institute for Astronomy, University of Hawaii Manoa, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

⁴Physics & Astronomy Dept., Rice University, 6100 Main St., Houston, TX 77005

⁵Korea Astronomy and Space Science Institute, 776 Daedeok-daero, Yuseong-gu, Daejeon 34055, Korea

⁶SOFIA Science Center - USRA, NASA Ames Research Center, Moffett Field, CA 94035, USA

⁷Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

⁸Gemini Observatory/NSF's NOIRLab, Casilla 603, La Serena, Chile

⁹School of Space Research, Kyung Hee University, 1732 Deogyong-daero, Giheung-gu, Yongin-si, Gyeonggi-do 17104, Korea

Submitted to ApJ

ABSTRACT

We present fundamental parameters for 110 canonical K- & M-type ($1.3\text{--}0.13 M_{\odot}$) Taurus-Auriga young stellar objects (YSOs). The analysis produces a simultaneous determination of effective temperature (T_{eff}), surface gravity ($\log g$), magnetic field strength (B), and projected rotational velocity ($v \sin i$). Our method employed synthetic spectra and high-resolution ($R \sim 45,000$) near-infrared spectra taken with the Immersion GRating INfrared Spectrometer (IGRINS) to fit specific K-band spectral regions most sensitive to those parameters. The use of these high-resolution spectra reduces the influence of distance uncertainties, reddening, and non-photospheric continuum emission on the parameter determinations. The median total (fit + systematic) uncertainties were 170 K, 0.28 dex, 0.60 kG, 2.5 km s^{-1} for T_{eff} , $\log g$, B, and $v \sin i$, respectively. We determined B for 41 Taurus YSOs (upper limits for the remainder) and find systematic offsets (lower T_{eff} , higher $\log g$ and $v \sin i$) in parameters when B is measurable but not considered in the fit. The average $\log g$ for the Class II and Class III objects differs by 0.23 ± 0.05 dex, which is consistent with Class III objects being the more evolved members of the star-forming region. However, the dispersion in $\log g$ is greater than the uncertainties, which highlights how the YSO classification correlates with age ($\log g$), yet there are exceptionally young (lower $\log g$) Class III YSOs and relatively old (higher $\log g$) Class II YSOs with unexplained evolutionary histories. The spectra from this work are provided in an online repository along with TW Hydrae Association (TWA) comparison objects and the model grid used in our analysis.

Keywords: infrared: stars — stars: pre-main sequence — stars: fundamental parameters

1. INTRODUCTION

Young stellar objects (YSOs) are stars at an early stage of evolution, and the disks surrounding them provide the raw material out of which planets can form. An empirically determined lifetime for protoplanetary disks is key to differentiating theories of planet formation and

migration (e.g., Baruteau et al. 2014). At a population level, observations of disk fractions in star forming regions and moving groups, with ages determined by comparing stellar properties to theoretical isochrones, have established a characteristic timescale of 2-3 Myr for YSO disk dissipation (e.g. Haisch et al. 2001; Hernández et al. 2007; Mamajek 2009; Ribas et al. 2014). These studies also make it clear that there is heterogeneity in the YSO population, with very young clusters having disk-

less members and a small number of systems with thick disks remaining at 8-10 Myr (e.g. TW Hydra, Sokal et al. 2018, and references therein). There are some indications of differences as a function of mass (Ribas et al. 2015; Galli et al. 2015), but mass-dependence does not explain many of the exceptional cases.

The combination of ALMA imaging of the outer disks of YSOs (e.g., ALMA Partnership et al. 2015; Barenfeld et al. 2017; Huang et al. 2018; Long et al. 2019; Pérez et al. 2020) and infrared interferometry and spectroscopy of inner disks (e.g., Muzerolle et al. 2003; Varga et al. 2017; Boccaletti et al. 2020) is providing new details on disk structure, kinematics, and composition. Both the puzzle of lifetime variation and the new details about the disks themselves drive a need for more precise information about the stellar hosts. Effective temperature (T_{eff}) and surface gravity ($\log g$), as proxies for YSO masses and ages, respectively (see Figure 9 of Sokal et al. 2018), permit comparisons of disk classifications with evolutionary models (Baraffe et al. 2015; Simon et al. 2019). Since angular momentum transport is very important both to the disks and to the evolution of stellar rotation (e.g., Attridge & Herbst 1992; Herbst et al. 2007; Bouvier et al. 2014), measures of the rotation are also needed. Magnetic interactions between a star and disk are crucial for accretion onto the star and material flow through the disk (e.g., Camenzind 1990; Koenigl 1991; López-Santiago et al. 2016). We therefore also need estimates of stellar magnetic field strength.

However, measurements of YSO parameters are influenced by interstellar reddening, continuum veiling, and stellar spots. The impacts of interstellar reddening are reduced at longer wavelengths, making the infrared more amenable to YSO studies than visible-light observations. The presence of strong magnetic fields (B) is associated with cool spots (e.g., Crockett et al. 2012; Gully-Santiago et al. 2017), Zeeman-split atomic spectral lines (e.g., Johns-Krull et al. 1999; Doppmann et al. 2003; Yang et al. 2005; Yang & Johns-Krull 2011; Lavail et al. 2017; Sokal et al. 2018; Lavail et al. 2019; Sokal et al. 2020) and changes in absorption line profiles. Veiling (r) is a non-stellar continuum emission that reduces the depth of the photospheric lines in YSO spectra (Joy 1949). It has been explained as the consequence of different physical processes including an active chromosphere (e.g., Calvet et al. 1984; Basri & Bertout 1989; Hartigan et al. 1989; Hartmann & Kenyon 1990; Valenti et al. 1993; Cieza et al. 2005), emission produced by accretion onto the star (e.g., Kenyon & Hartmann 1987; Johns-Krull & Valenti 2001; Gahm et al. 2008), excess emission from dust at the sublimation radius of the inner disk wall (e.g., Natta et al. 2001; Muzerolle et al. 2003) and the

emission from warm gas located inside the dust sublimation radius of the disk (e.g., Fischer et al. 2011). These variable processes can result in YSO parameters that differ significantly between literature references because the YSOs were observed at different times, in different physical states, and with instruments observing different wavelengths.

We have employed the Immersion GRating Infrared Spectrometer (IGRINS; Yuk et al. 2010; Park et al. 2014; Mace et al. 2016) to survey the Taurus-Auriga star-forming region and determine YSO properties while minimizing the effects listed above. With its broad spectral grasp (1.45-2.5 μm) at high resolution ($R \sim 45,000$), IGRINS simultaneously observes numerous spectral features of YSOs. Additionally, we can average over variability and derive typical parameters for each target by combining multiple observation epochs. In 2014, IGRINS was commissioned on the 2.7 m Harlan J. Smith Telescope at McDonald Observatory. Observations initially focused on the Taurus-Auriga complex (often recognized as Taurus) because it is a close ($d = 140$ pc, Kenyon et al. 1994; Galli et al. 2018) and young (about 1-5 Myr, Kraus & Hillenbrand 2009; Gennaro et al. 2012) star-forming region that provides observations of YSOs at various stages of evolution. The population of Taurus is at an age coincident with disk dissipation timescales and includes both Class II and Class III YSOs.

The IGRINS YSO Survey is uniquely suited to the determination of physical parameters because:

- The 1.45–2.5 μm spectral region includes both photospheric and disk contributions to the YSO spectrum.
- The fixed spectral format of IGRINS provides similar spectral products for each object at each epoch.
- Multiple-epoch observations provide a means to characterize and average over variability.
- The spectral resolution of $\sim 7 \text{ km s}^{-1}$ is smaller than the typical rotational velocity of young stars and can reliably resolve line-splitting by magnetic fields (B) $\gtrsim 1.0 \text{ kG}$.

In this paper, we present the simultaneous determination of T_{eff} , $\log g$, B , and projected rotational velocity ($v \sin i$) of 119 K- & M-type YSOs located in Taurus, of which 110 are reliable. In Section 2 we describe the sample and the IGRINS observations. Section 3 details the parameter determination and sources of uncertainty, while in Section 4 we discuss the results and provide analysis. In Section 5 we provide a summary and the conclusions of this work.

2. OBSERVATIONS AND SAMPLE

IGRINS employs a silicon immersion grating as the primary disperser (Jaffe et al. 1998; Wang et al. 2010; Gully-Santiago et al. 2010) and volume phase holographic gratings to cross-disperse the H- and K-band echellograms onto Teledyne Hawaii-2RG arrays. This setup provides a compact design with high sensitivity and a significant single-exposure spectral grasp (Yuk et al. 2010; Park et al. 2014). IGRINS has a fixed spectral format and no moving optics, so the science products remain unchanged no matter where it is installed. IGRINS has increased its scientific value by traveling between McDonald Observatory, the Lowell Discovery Telescope (LDT), and the Gemini South telescope (Mace et al. 2018).

Flat-field calibration frames were taken at the start of each night and used for bad pixel masking and 2D aperture definitions. All the YSOs in our sample were observed by nodding the targets along the slit in patterns made up of AB or BA pairs. An A0V telluric standard star was observed at a similar airmass within a period of two hours before or after the science target. Flexure between the target and A0V observations was sub-pixel (Mace et al. 2016). The airmass during the IGRINS observations ranged from 1.00 to 2.17. We used the IGRINS exposure time calculator¹ to estimate the exposure time of each object. The total exposure time varied from minutes to hours to achieve a signal-to-noise ratio (SNR) > 100.

All IGRINS data were reduced using the IGRINS pipeline (Lee, Gullikson, & Kaplan 2017)² which produces a telluric corrected spectrum with the wavelength solution derived from OH night sky emission lines at shorter wavelengths and telluric absorption lines at wavelengths greater than 2.2 μm . Telluric correction was performed by dividing the target spectrum by an A0V spectral standard and multiplying by a standard Vega model. We corrected for any sub-pixel shifts between the target and A0V observations (due to instrument flexure) by aligning the spectra at a strong telluric absorption feature. Finally, the wavelength solution of the target was corrected for the barycenter velocity derived using ZBARYCORR (Wright & Eastman 2014) along with the Julian date at the midpoint of observation and the telescope site. The telluric correction uncertainties were propagated into the telluric-corrected spectra by

combining the observed uncertainties of the target and standard spectra in quadrature.

To construct the YSO sample used in this work, we looked into the reduced science spectral archive of IGRINS for targets classified as Taurus members by Luhman et al. (2010), with a spectral type (SpT) between K0 and M5 (Luhman et al. 2017), and a minimum average SNR of 50 in the K-band. The latter condition resulted in more than 500 single visit observations of 139 Taurus YSOs observed with IGRINS on the McDonald 2.7 m telescope and LDT between 2014 and 2017. This sample contains many of the canonical Taurus-Auriga members with K-band magnitudes < 10 mag and according to the classifications made by Kenyon & Hartmann (1995), Luhman et al. (2010), Rebull et al. (2010), Esplin et al. (2014), and Kraus et al. (2017), there are 9 Class I, 91 Class II and 39 Class III YSOs. Finally, we visually checked the K-band spectrum of these 139 YSOs, and we excluded objects with shallow (< 2%) lines due to high veiling or $v \sin i$ values. After this cleaning step, our final sample (see Table 1), contained 119 YSOs (84 Class II and 35 Class III). Figure 1 depicts the spectral type (Luhman et al. 2017), 2MASS K-band magnitude (Cutri et al. 2003), and the number-of-epoch distributions of the 119 YSOs that define our sample.

By compiling the survey sample from the IGRINS spectral archive, we identify some biases in the sample selection. First, for the same effective temperature ($T_{\text{eff}} \lesssim 4000$ K), younger stars are brighter because of their larger radii (Hayashi 1961). These young and bright stars were easier to observe at high spectral resolution. The full census of Taurus YSOs is > 400 members (Luhman 2018). Our survey includes only ~45% (116 out of 258) of the Taurus YSOs between spectral types K0 and M5, since we have not observed the faintest or more widely distributed members (Kraus et al. 2017). Lastly, binary stars are the specific focus of some IGRINS YSO observing programs. The combined flux of the two binary components makes them brighter, which results in higher SNRs, but their analysis is also complicated by line blending in the composite spectrum. In this work we have identified known binaries (or multiples) in the literature and have compared the single and multiple samples. A more rigorous identification, analysis, and characterization of binarity will be the topic of a future study.

The final combined spectrum of each target was made up of between 1 and 10 epochs, with a median value of 4 epochs. If more than ten epochs were available, we chose the best ten based on a combination of airmass, SNR, and the quality of the telluric correction. We shifted each epoch to the wavelength rest-frame by

¹ <https://wikis.utexas.edu/display/IGRINS/SNR+Estimates+and+Guidelines>

² <https://github.com/igrins/plp/tree/v2.1-alpha.3>




Figure 1. Luhman et al. (2017) spectral types (left panel), 2MASS K-band magnitudes (middle panel), and the total number of combined epochs (right panel) for our sample of 119 YSOs. More than 70% of the stars in our sample have spectral type K9 or later. Our sample also has a mean K magnitude of 8.4 ± 0.9 mag and a median value of 4 combined epochs per target.

determining the radial velocity shift of Na lines at 2.206 and 2.209 μm . Then, we produced the combined spectrum by taking a weighted average of all epochs, where the weight corresponded to the SNR at each data point. The standard deviation of the mean gave the final uncertainties per data point.

In 2018, we observed members of the TW Hydrae Association (TWA; Kastner et al. 1997) while IGRINS visited the 8.1 m Gemini South telescope (Mace et al. 2018). We reduced the TWA objects in the same fashion as the Taurus sample, but all TWA observations were single epoch and final combined spectra were not produced. The collection of IGRINS spectra is available from the Harvard Dataverse (López-Valdivia 2021a,b)³.

3. STELLAR PARAMETER DETERMINATION

To determine T_{eff} , $\log g$, B , and $v \sin i$ we used a Markov chain Monte Carlo (MCMC) algorithm comparing observations with synthetic spectra, which we describe here in detail.

3.1. Theoretical grid

We computed a four-dimensional (T_{eff} , $\log g$, B , $v \sin i$) grid of synthetic spectra using the MOOGSTOKES code (Deen 2013). MOOGSTOKES is a customization of the one-dimensional LTE radiative transfer code MOOG (Snedden 1973) that synthesizes the emergent spectrum of a star taking into account the Zeeman splitting produced by the presence of a photospheric magnetic field.

It assumes a uniform and purely radial magnetic field and uniform T_{eff} and $\log g$, producing a disk-averaged spectrum broadened to the user-specified spectral resolution and $v \sin i$ values.

To synthesize a spectrum with MOOGSTOKES, the program required a model atmosphere and an atomic/molecular line list. The T_{eff} , $\log g$, and metallicity of the resulting spectrum are defined by the model atmosphere, while B , $v \sin i$, and spectral resolution are user-selected quantities. In this work, we used the MARCS atmospheric models (Gustafsson et al. 2008) with solar metallicity (suitable for YSOs; Padgett 1996; Santos et al. 2008; D’Orazi et al. 2011) and the Vienna Atomic Line Database (VALD3; Ryabchikova et al. 2015). Often, the atomic data coming from databases are not accurate enough to reproduce the Solar spectrum at high resolution, requiring modifications to some of the absorption line oscillator strengths and van der Waals constants. In this work we used the astrophysical-inferred modifications to the VALD atomic data presented by Flores et al. (2019). Low-mass stars have micro-turbulence values between 0 and 2 km s^{-1} (e.g., Gray 2005; Reid & Hawley 2005; Bean et al. 2006), thus for this study we used a micro-turbulence of 1 km s^{-1} . Finally, we selected B values up to 4 kG, $v \sin i$ between 2 and 50 km s^{-1} , and matched the IGRINS spectral resolution ($R \sim 45,000$).

Our grid of synthetic spectra covers the parameter space as follows: from 3000 to 5000 K in T_{eff} (steps of 100 K up to 4000 K, and 250 K above 4000 K), from 3.0 to 5.0 dex in $\log g$ (steps of 0.5 dex), from 0 to 4 kG

³ https://dataverse.harvard.edu/dataverse/igrins_ysos

in B (steps of 0.5 kG), and values from 2 to 50 km s⁻¹ in $v \sin i$ (steps of 2 km s⁻¹). The grid steps⁴ are well suited to our study as they are enough to see the effects of these parameters on the spectra. Smaller step sizes in the model grid would drastically increase the computation time without significantly altering the synthetic spectra. Instead, the grid was linearly interpolated at values between grid points. The grid of synthetic spectra does not cover the entire IGRINS spectrum, but only the spectral regions described in Section 3.2. The grid of synthetic spectra is also available from Harvard Data-verse (López-Valdivia 2021c).

3.2. Spectral Regions

We employed four spectral regions in the K-band to determine the stellar parameters in this work. These regions are in four different K-band IGRINS orders. Each region is sensitive to changes in different stellar parameters, as shown by their use previously in K and M stars (e.g., Rajpurohit et al. 2018; Sokal et al. 2018; Flores et al. 2019).

Sokal et al. (2018) used three spectral intervals in the K-band around Na ($\sim 2.210 \mu\text{m}$), Ti ($\sim 2.222 \mu\text{m}$), and CO ($\sim 2.295 \mu\text{m}$) features to determine T_{eff} , $\log g$, and B of the star TW Hydra. Recently, Flores et al. (2019) included the Ca I lines around $2.264 \mu\text{m}$ in the determination of the stellar parameters of the young stars BP Tau and V347 Aur with iSHELL (Rayner et al. 2016) infrared spectra at a similar spectral resolution to IGRINS.

All spectral lines depend, to some extent, on each of the stellar parameters. Isolating the contribution to the line profile of each parameter is only as accurate as the models and line lists. To reduce degeneracies, it was important to use a collection of spectral lines that are primarily sensitive to variations of just one stellar parameter.

To assess the sensitivity of the selected spectral regions to the stellar parameters, we took the difference between two synthetic spectra varied by 300 K in T_{eff} , 0.5 dex in $\log g$ or 1.0 kG in B. Figure 2 depicts the dependency of each spectral region on the stellar parameters. We label the four intervals as follows:

Na region (2.2045–2.2105 μm): The Na doublet ($\lambda \sim 2.2062$ and $2.2090 \mu\text{m}$) is the dominant feature of this region, but two Sc I lines (~ 2.2058 and $2.2071 \mu\text{m}$) and a Si I line ($\sim 2.2068 \mu\text{m}$) are also present. The Na lines show sensitivity to the magnetic field, while the Sc I and

Si I are more sensitive to T_{eff} and respond in opposite ways as T_{eff} increases.

Ti region (2.2205–2.2346 μm): This region has three Ti I lines at ~ 2.2217 , $\sim 2.2239 \mu\text{m}$, and ~ 2.2315 that are mainly sensitive to B. There are also two Fe I lines mainly sensitive to changes in T_{eff} . The Ti lines present greater Zeeman splitting when the B-field increases.

Ca region (2.2606–2.2664 μm): This region includes three Ca I lines at ~ 2.2614 , 2.2631 , and $2.2657 \mu\text{m}$ and one Fe I at $\sim 2.2626 \mu\text{m}$. These lines together help us to determine T_{eff} and $\log g$ because their dependency on B is small.

CO region (2.2986–2.3036 μm): This region contains six CO lines sensitive to $\log g$, and a weak Sc I line ($\sim 2.2993 \mu\text{m}$) sensitive to B. As the $\log g$ increases the CO line depths decrease. This CO region is smaller than others have used, which provides better continuum flattening.

The combined use of the four regions located in the K-band allows for the accurate simultaneous determination of T_{eff} , $\log g$, B, and $v \sin i$ of our YSOs sample.

3.3. Determining Stellar Parameters

To determine stellar parameters we continuum normalized each spectral region through an interactive python script. First, the wavelength range of the spectral region is equally divided into n bins. Then, we computed the median (μ_{flux}) and the standard deviation (σ_{flux}) of the flux in each bin; we then excluded points out of $\mu_{\text{flux}} \pm 0.5\sigma_{\text{flux}}$, and we recomputed the median flux of the bin (μ_{clip}). Finally, we fit a polynomial of order k to the μ_{clip} of the n bins. The values of n and k vary between objects and spectral regions and could be interactively modified to obtain a better normalization. We typically used between 10 and 20 bins and between 1 and 4 for the polynomial order.

Then, we carried out a MCMC analysis, as implemented in the code *emcee* (Foreman-Mackey et al. 2013), using the four K-band spectral regions (Na, Ti, Ca, and CO). We compared observed and synthetic spectra by allowing T_{eff} , $\log g$, B, $v \sin i$, and K-band veiling (r_K) to vary along with small continuum ($< 3\%$) and wavelength ($< 0.6 \text{\AA}$) offsets. In each MCMC trial we linearly interpolated within the four-dimensional (T_{eff} , $\log g$, B, $v \sin i$) synthetic spectral grid described in Section 3.1, to obtain the corresponding spectrum with the sampled set of parameters. The interpolated synthetic spectrum was then artificially veiled and re-normalized for each region following this equation:

$$F_v = \frac{F + r_K}{1 + r_K} \quad (1)$$

⁴ The grid steps of T_{eff} and $\log g$ are defined by the model atmosphere, while we selected the steps of B and $v \sin i$.




Figure 2. Dependence on stellar parameters for the selected spectral regions. We used a synthetic spectrum with $T_{\text{eff}} = 3600$ K, $\log g = 4.0$ dex, $B = 2.0$ kG, and $v \sin i = 10$ km s $^{-1}$ (solid black line) as our reference. We show the difference caused by a change of 300 K in T_{eff} (green dashed line), of 0.5 dex in $\log g$ (blue dash-dotted line), or 1.0 kG in B (red dotted line). In each comparison, we fixed two of the three parameters to the reference value and varied the third to a higher value. Varying parameters to lower values results in similar behavior in the opposite directions. A similar comparison can be seen in Sokal et al. (2018).

where F_v , F , and r_K are the veiled flux, the synthetic flux, and the K-band veiling value (Basri & Batalha 1990). A single, best-fit veiling value was determined for all the K-band wavelength regions at once. Note that the r_K values determined with our MCMC analysis are not strictly a veiling value, but a combination of the true veiling with secondary systematic offsets between the model and target continua.

We then evaluated each MCMC with a likelihood function based on the sum of the χ^2 statistics divided by the number of pixels in each region. Finally, from the posterior probability distributions of the MCMC, we took the 50th percentile as the most likely value for the stellar parameters.

In Figure 3, we show the posterior distributions of the MCMC analysis for the star IW Tau as an illustrative example. Figure 3 also shows the dependence between stellar parameters like $\log g$ and r_K . A good fit to the observation can be achieved with a slightly lower $\log g$, or a higher r_K because both $\log g$ or veiling varies the CO line depths. The degeneracy between these parameters adds to the uncertainty in each quantity separately.

3.4. Uncertainties

The 16th and 84th percentiles of the posterior probability distributions of the stellar parameters represent the formal uncertainties of our fitting method. Most of the uncertainties were found to be nearly symmetric, and we report the larger of the two percentiles as the corresponding symmetric uncertainty (σ_{fit}) for each parameter. The median fit uncertainties are 152 K in T_{eff} , 0.25 dex in $\log g$, 0.54 kG in B, 1.9 km s⁻¹ in $v \sin i$, and 0.10 in r_K .

Additionally, we quantified the systematic uncertainties (σ_{sys}) by comparing our stellar parameters to previously published values. Stars with precise physical parameters provided by interferometric observations help mitigate model dependency by calibrating relationships between T_{eff} and stellar radius. The works of Mann et al. (2013, 2015) and Newton et al. (2015) used 20+ stars with interferometric measurements to calibrate their model-independent relationships with ~ 150 K precision. To quantify the σ_{sys} for T_{eff} and $\log g$, we determined the stellar parameters of seven field M stars with metallicity close to the solar value ($[\text{Fe}/\text{H}] = 0.0 \pm 0.10$ dex) from Mann et al. (2013).

We used the same fitting routines as for the YSO sample, but assuming $r_K = 0.0$ since these stars are on the main sequence and diskless. In Table 2, we compared our T_{eff} and $\log g$ values with those obtained by Mann et al. (2013). They determined T_{eff} , $[\text{Fe}/\text{H}]$, mass (M) and radius (R), which we used in the following equation

to compute their $\log g$ values:

$$\log g_{\text{cal}} = \log(M/M_{\odot}) - 2 \log(R/R_{\odot}) + 4.437 \quad (2)$$

We found an average difference in $\log g$ (ours - Mann) of -0.04 dex and a standard deviation of the residuals of 0.13 dex. For T_{eff} we found a mean difference of -69 K and a standard deviation of the residuals of 75 K. Therefore, we assigned 0.13 dex and 75 K as the systematic uncertainties for $\log g$ and T_{eff} , respectively.

To obtain the systematic uncertainty on $v \sin i$, we compared measurements for 52 YSOs in common with Nofi et al. (2021). In that work, they determined the $v \sin i$ of pre-main-sequence stars using the same single-epoch, uncombined IGRINS K-band spectra that we have used. Their method used the cross-correlation technique outlined in Hartmann et al. (1986) and Soderblom et al. (1989) along with synthetic spectra. In brief, a stellar spectrum with rotationally broadened lines was cross-correlated against an unbroadened spectrum. Then the width of the cross-correlation function is used to measure the rotational broadening of the spectrum. Generally, our $v \sin i$ values were in good agreement with those of Nofi et al. (2021). We found a mean difference between the $v \sin i$ values (ours - Nofi) of 0.2 km s⁻¹ and a standard deviation of the residuals of 1.7 km s⁻¹. We assigned this 1.7 km s⁻¹ as the $v \sin i$ systematic uncertainty.

Over 85% of our $v \sin i$ values are ≤ 30 km s⁻¹, with the median value being 16.6 km s⁻¹. In Figure 4 we compare $v \sin i$ measurements for 60 YSOs in common with Nguyen et al. (2012). They used high-resolution ($R=60,000$) optical (4800–9300 Å) spectra to determine $v \sin i$ for pre-main-sequence stars in the Chamaeleon I and Taurus-Auriga star-forming regions. Their method compared each echelle order of the target spectrum with slow-rotating template stars, which were artificially broadened to different values of $v \sin i$ to find the best fit. Finally, they removed 2.7σ outliers around the mean value and computed a weighted average over all the remaining echelle orders (for more details see also Nguyen et al. 2009). We found our $v \sin i$ values in agreement with those of Nguyen et al. (2012), with an mean difference of 0.08 km s⁻¹, and a error on the mean of 0.3 km s⁻¹.

Finally, we collected 12 determinations (for 11 YSOs) of B-field strength from the literature (Johns-Krull 2007; Sokal et al. 2020; Flores et al. 2020) to assess the systematic uncertainty of our B values (see Table 3). We found a mean difference in the residuals (ours - literature) of -0.27 kG and a standard deviation of 0.26 kG. We assigned 0.26 kG as our systematic uncertainty in the B-field measurements.

IW Tau




Figure 3. Posterior distributions and degeneracies (after removing the burn-in phase) of T_{eff} , $\log g$, B , $v \sin i$, and r_K determined for IW Tau. The median value and the 1σ uncertainties are printed above each distribution.

Table 1. Compiled information for the 119 stars in our Taurus YSO sample, including T_{eff} , $\log g$, B , and $v \sin i$. Columns 1-8 provide target information from the literature, while Columns 9-16 present information and parameters from this work. The Ref column identifies the source of the class and binary (objects with companions closer than 2 arcseconds) classification. The SpT and the K magnitude comes from Luhman et al. (2017) and Cutri et al. (2003), respectively. Uncertainties are the 1σ total (fit + systematic) uncertainties. The N column is the number of epochs combined for each object. The SNR_K column is the median SNR of the combined spectrum for the spectral regions used in this work. The flag column equals 0, 1, 2 or 3 indicates good, acceptable, outside/edge of grid, or poor determinations. The full version of this table is available in the online version of the paper.

2MASS ID	Alt.	Name	K (mag)	SpT	Class	Ref	Binary	Ref	N	SNR _K	flag	T_{eff} (K)	$\log g$ (dex)	B (kG)	$v \sin i$ (km s ⁻¹)	r_K
J041141760+2806096	[BCG93]	1	9.9	M5.0	II	1,2,3,4	?	—	2	172	0	3281±197	3.88±0.40	<1.09	9.9±2.9	1.10±0.21
J05075496+2500156	[BCG93]	12	10.4	M3.7	II	2,4	N	16	2	64	3	—	—	—	—	—
J04182909+2826191	[SS94]	V410 Anon 25	9.9	M3.5	III	2,4,5	?	—	1	87	1	3619±328	4.05±0.59	<2.32	30.7±7.4	0.39±0.20
J04174965+2829362	[SS94]	V410 X-ray 1	9.1	M3.7	II	2,3,4	N	16	2	182	0	3315±129	3.37±0.25	1.22±0.33	9.7±2.0	0.28±0.06
J04183444+2830302	[SS94]	V410 X-ray 2	9.2	M0.0	II	2,4	?	—	2	182	0	4231±143	4.53±0.27	1.84±0.49	10.9±2.9	0.44±0.08
J04345542+2428531	AA	Tau	8.0	M0.6	II	1,2,3,4	N	16	10	356	0	3751±171	3.87±0.26	2.16±0.41	12.5±2.3	1.56±0.12
J04191583+2906269	BP	Tau	7.7	M0.5	II	1,2,3,4	N	16	7	353	0	3719±131	4.01±0.25	2.19±0.37	9.9±2.3	1.36±0.11
J04335200+2250301	CI	Tau	7.8	K5.5	II	2,3,4	N	16	10	931	0	3951±94	3.77±0.17	1.95±0.31	12.5±1.9	2.34±0.09
J04265440+2606510	CoKu FV	Tau c	8.9	M2.5	II	1,2,3,4	Y	24	6	302	0	3456±136	3.53±0.23	<2.01	38.6±2.4	0.33±0.06
J04355349+2254089	CoKu HP	Tau G3	8.8	M0.6	III	1,2,3,4,5	Y	16	6	239	0	3760±164	3.84±0.26	<2.65	43.3±2.9	0.25±0.07
J04354093+2411087	CoKu	Tau 3	8.4	M0.5	II	1,2,3,4	N	16	5	249	0	3665±154	3.85±0.26	2.33±0.34	8.2±2.1	0.77±0.08
J04411681+2840000	CoKu	Tau 4	8.7	M1.1	II	1,2,4	Y	13	6	254	0	3706±121	3.70±0.21	<1.90	26.3±2.2	0.22±0.04
J04144786+2648110	CX	Tau	8.8	M2.5	II	1,2,3,4	N	16	2	152	0	3520±180	3.61±0.30	<1.39	20.5±2.5	0.28±0.08
J04173372+2820468	CY	Tau	8.6	M2.3	II	1,2,3,4	N	16	2	203	0	3445±155	3.73±0.28	<1.31	10.8±2.3	0.81±0.13
J04183158+2816585	CZ	Tau	9.4	M4.2	II	1,2,3,4	Y	18	4	220	1	3385±211	3.83±0.35	<1.70	31.4±3.1	0.64±0.12
J04183112+2816290	DD	Tau	7.9	M3.5	II	1,2,3,4	Y	6	4	251	0	3250±207	3.47±0.37	<1.48	16.9±3.1	1.82±0.26
J04215563+2755060	DE	Tau	7.8	M2.3	II	1,2,3,4	N	16	4	251	0	3463±158	3.45±0.27	<1.01	9.4±2.2	1.11±0.11
J04270280+2542223	DF	Tau	6.7	M2.0	II	1,2,3,4	Y	24	4	293	3	—	—	—	—	—
J04270469+2606163	DG	Tau	7.0	K7.0	II	1,2,3,4	N	16	5	553	2	3969±216	3.20±0.28	<1.78	26.9±4.3	4.82±0.94
J04294155+2632582	DH	Tau	8.2	M2.3	II	1,2,3,4	N	16	10	399	0	3477±125	3.89±0.22	2.21±0.32	8.4±2.1	1.24±0.10

References—(1)Kenyon & Hartmann (1995); (2)Luhman et al. (2010); (3)Rebull et al. (2010); (4)Espino et al. (2014); (5)Kraus et al. (2017); (6)Bouvier et al. (1992); (7)Correia et al. (2006); (8)Duchêne et al. (1999); (9)Dyck et al. (1982); (10)Ghez et al. (1993); (11)Haas et al. (1990); (12)Herbig & Bell (1988); (13)Ireland & Kraus (2008); (14)Konopacki et al. (2007); (15)Kraus et al. (2006); (16)Kraus et al. (2011); (17)Leinert et al. (1991); (18)Leinert et al. (1993); (19)Mathieu et al. (1991); (20)Mathieu et al. (1996); (21)Richichi et al. (1999); (22)Ruiz-Rodríguez et al. (2016); (23)Schaefer et al. (2014); (24)Simon et al. (1992); (25)Walter et al. (1988); (26)Weintraub (1989).




Figure 4. Comparison of $v \sin i$ values between [Nguyen et al. \(2012\)](#) and this work for 60 YSOs in common. The dashed line shows the one-to-one relation. The mean difference between both studies is $0.08 \pm 0.3 \text{ km s}^{-1}$, and we found a correlation coefficient of 0.97. Our error bars are the fit uncertainties.

Table 2. Effective temperature and surface gravity determined for our calibration sample as well as those determinations of [Mann et al. \(2013\)](#). The fit errors are reported on our T_{eff} and $\log g$ determinations.

Star	Mann et al. (2013)		this work	
	T_{eff}	$\log g$	T_{eff}	$\log g$
	(K)	(dex)	(K)	(dex)
GJ 338A	3953 ± 41	4.71 ± 0.05	3813 ± 56	4.68 ± 0.09
GJ 338B	3926 ± 37	4.72 ± 0.04	3785 ± 54	4.69 ± 0.08
GJ 436	3520 ± 66	4.77 ± 0.06	3587 ± 59	4.90 ± 0.11
GJ 570A	4588 ± 58	4.57 ± 0.07	4536 ± 50	4.36 ± 0.11
GJ 702B	4475 ± 33	4.66 ± 0.04	4391 ± 38	4.50 ± 0.14
GJ 809	3744 ± 27	4.72 ± 0.04	3604 ± 63	4.59 ± 0.09
GJ 887	3695 ± 35	4.78 ± 0.05	3699 ± 67	4.95 ± 0.08

The systematic uncertainties were added in quadrature to the fit errors to compute the total error of each parameter (σ_{tot}), which we report in Table 1.

4. RESULTS AND ANALYSIS

We plotted the spectral regions used in our analysis for all the observed and the best-fit spectra, as in Figure 5, to categorize the quality of our parameter determinations. Through a ranking of the χ^2 statistics and visual

Table 3. Literature and this work magnetic field strength determinations for 11 YSOs. The reference column are JF07 = [Johns-Krull \(2007\)](#), F19 = [Flores et al. \(2019\)](#), and S20 = [Sokal et al. \(2020\)](#). We report the fit uncertainties on our B values.

Star	B literature	Ref.	B this work
	(kG)		(kG)
AA Tau	2.78 ± 0.28	JK07	2.16 ± 0.32
BP Tau	2.17 ± 0.22	JK07	2.19 ± 0.27
BP Tau	2.50 ± 0.16	F19	2.19 ± 0.27
CI Tau	2.15 ± 0.15	S20	1.95 ± 0.16
CY Tau	1.16 ± 0.12	JK07	1.31 ± 0.37
DE Tau	1.12 ± 0.11	JK07	1.01 ± 0.24
DH Tau	2.68 ± 0.27	JK07	2.21 ± 0.18
DK Tau	2.64 ± 0.26	JK07	2.55 ± 0.58
DN Tau	2.00 ± 0.20	JK07	1.54 ± 0.26
GI Tau	2.73 ± 0.27	JK07	1.92 ± 0.36
GK Tau	2.28 ± 0.23	JK07	2.05 ± 0.87
GM Aur	2.22 ± 0.22	JK07	2.11 ± 0.22

inspection, we marked each set of parameters in Table 1 with a numerical flag equal to 0, 1, or 3 if they are good, acceptable, or poor determinations, respectively. We also computed the lower and upper limits of each parameter using its total uncertainty, to look for values outside or at the edge of our grid. We find that six lower limits of $\log g$ meet this criterion. We flagged these parameters with a flag equal to 2 and have retained these measurements in our analysis since they are otherwise classified as ‘good’ determinations. We found that our parameter determination method needs a minimum median SNR $\gtrsim 80$ to produce reliable results. A median SNR $\gtrsim 150$ is generally required for good parameter determinations. In total, we identified 80, 24, 9, and 6 stars whose parameters are good, acceptable, poor, and outside/edge of grid, respectively. We exclude poor determinations in further analysis, and we will refer to the 110 remaining YSO measurements as our Taurus sample.

4.1. Magnetic fields

While Zeeman line splitting effects are most significant in the Ti region, the Na region is also sensitive to variations in the B field (see Figure 2). However, our ability to measure Zeeman line separations is primarily a function of the line widths, which are dominated by $v \sin i$. [Hussaini et al. \(2020\)](#) measured the line splitting of the Na and Ti spectral regions in IGRINS data and the MOOGSTOKES spectra employed in this work. That work found that the retrieval of B is limited to




Figure 5. The four IGRINS K-band spectral regions of the star IW Tau (solid black line). We include the best-fit synthetic spectrum (red dashed line) with $T_{\text{eff}} = 3663$ K, $\log g = 4.07$ dex, $B = 2.28$ kG, $v \sin i = 8.3$ km s $^{-1}$, and $r_K = 0.10$. In the bottom panels, we present the residuals between the observed and the synthetic spectra. The level of agreement between the best-fit-parameter synthetic spectrum and the IW Tau observed spectrum was considered a good determination.

$\sim v \sin i / 10$ in the Ti region and $\sim v \sin i / 5$ in the Na region. In other words, an object with $v \sin i = 15$ km s $^{-1}$ must have $B > 1.5$ kG for Zeeman splitting to be measurable in the Ti region and > 3 kG to be measurable in the Na region. Additionally, the IGRINS spectral resolution limits the minimum measurable B field from line splitting to ~ 0.7 kG. For this work, we conservatively considered a detection limit of $B \geq (v \sin i / 8)$ kG, and no lower than 1 kG. Values of B field lower than 1 kG can not be distinguished from a non-detection at the spectral resolution of IGRINS when simultaneously fitting for other model parameters. We found that 41 of our B field determinations meet the detection threshold.

The remaining upper-limit measurements are provided in Table 1.

4.2. Impact of B on the stellar parameters

To evaluate the impact of B on T_{eff} , $\log g$, $v \sin i$, and r_K , we fixed B to 0 kG and redetermined the stellar parameters for the 41 YSOs with B-field detections in Table 1. In Figure 6 we compare the stellar parameters determined in our primary analysis with those obtained by disabling the B parameter. We further consider the low B (objects with $B \leq 2$ kG) and the high B ($B > 2$ kG) results in this comparison.

We found that all the T_{eff} values determined without considering the B-field effects are systematically colder than those found considering the B-field. For low B (18 objects), we found a mean difference of -42 ± 4 K, while for high B (23 objects), this difference increases slightly to -73 ± 9 K. Both values are well within the median total (fit+systemic) T_{eff} error, which is ~ 170 K. Similarly, r_K is lower when B is not taken into account, with a mean difference of -0.14 ± 0.02 for both the low and high B bins. Contrary to T_{eff} and r_K , most of the $\log g$ values are higher by 0.08 ± 0.01 dex and 0.03 ± 0.01 dex for the low and high B bins, respectively, when not considering B-fields in the fit. These $\log g$ differences are within the $\log g$ median total (fit+systemic) error of 0.28 dex. It is likely that the degeneracy between $\log g$ and r_K discussed in Section 3.3 is why $\log g$ is overestimated and r_K is underestimated when B is not included in the parameter determinations.

The most prominent effect of ignoring the B-field strength when determining parameters is seen in the $v \sin i$ values. All $v \sin i$ determinations are higher without B-field included in the fit, and the difference is greater for larger values of B . We found for the low-B bin a mean difference of 1.6 ± 0.2 km s $^{-1}$. For the high-B bin, the difference increases to 3.8 ± 0.3 km s $^{-1}$. This trend is explained by excess rotational broadening of the synthetic spectral lines to try and replicate the B-field induced Zeeman line-splitting. The absence of B-field considerations by Nguyen et al. (2012) and Nofi et al. (2021) is likely responsible for some of the differences between those works and ours, which adds to the systematic uncertainties.

The previous tests suggest that it is possible to obtain suitable stellar parameters, even when excluding the B-field from the determination. However, the parameters will have systematic offsets, with magnetically active YSO parameters being the most impacted.

4.3. Effective Temperature

Kenyon & Hartmann (1995) presented a conversion between spectral type and T_{eff} for young stars, based on




Figure 6. Stellar parameters determined with and without taking into account the effects of B-field strength. The dashed line is the one-to-one relation, while the dotted lines demarcate the median fit error intervals namely: ± 150 K, ± 0.25 dex, ± 1.9 km s $^{-1}$, and ± 0.10 for T_{eff} , $\log g$, $v \sin i$, and r_K , respectively. The data are color-coded by the determined strength of B.

the work of [Schmidt-Kaler \(1982\)](#) and [Straižys \(1992\)](#). Shortly after, [Luhman \(1999\)](#) developed a new temperature scale for young M stars between the giant and dwarf scales. However, improvements made in the last two decades in the modeling of cool star atmospheres, and the development of new instruments, have resulted in several studies that have determined effective temperatures (or spectral type) either for field (e.g., [Casagrande et al. 2008](#); [Rajpurohit et al. 2013](#); [Rajpurohit et al. 2018](#); [López-Valdivia et al. 2019](#)) or for young (e.g., [Luhman et al. 2003a](#); [Luhman et al. 2003b](#); [Herczeg & Hillenbrand 2014](#); [Cottaar et al. 2014](#); [Yao et al. 2018](#)) K & M stars.

In Figure 7 we compare the T_{eff} values we determined to spectral types from [Luhman et al. \(2017\)](#), along with three different temperature scales of pre-main-sequence stars ([Luhman et al. 2003b](#); [Herczeg & Hillenbrand 2014](#); [Pecaut & Mamajek 2013](#)). To analyze how multiplicity plays a role in determining T_{eff} , we divided our sample into single (51) and binary (44) stars based on classifications from the literature. These identifications are reported in Table 1. We consider a star a binary if it has a companion closer than 2 arcseconds because the resultant IGRINS spectrum will contain some amount of flux from both components.

The amount of contamination from a binary companion is not easily quantified. An equal brightness bi-




Figure 7. Effective temperature as a function of [Luhman et al. \(2017\)](#) spectral type for our Taurus-Auriga known single and binary (multiple) YSOs. The dashed, dotted-dashed, and dotted lines are the temperature scales of [Herczeg & Hillenbrand \(2014\)](#), [Luhman et al. \(2003b\)](#) and [Pecaut & Mamajek \(2013\)](#), respectively. Typically the error on the spectral type is about 1 subclass. There is good agreement between our temperatures for M0–M4 single and K6–M5 binary stars with the published temperature scales. For single K stars, our T_{eff} values are cooler than the published temperature scales. The binary stars HQ Tau, T Tau, and RW Aur present the largest discrepancies. Our error bars are the total (fit + systematic) T_{eff} uncertainties.

nary with no velocity offsets will have composite line profiles, often producing poor fits to the data. If the binary has equal brightness and radial velocity offsets (a double-lined spectroscopic binary), then line blending in the combined spectrum will often result in poor parameter determinations. Generally, equal brightness binaries have been excluded from our analysis by visual inspection in the initial sample selection and at the parameter-fit quality check. Binaries with disproportionate fluxes will have the same issues as the equal flux binaries, but at lower levels. Stellar parameters obtained for the known binary stars should be used with caution since secondary contamination is possible, although the spectral fit is dominated by the primary star’s contribution. The binary identification here allows us to look for systematic differences in the data.

In general our temperatures follow the scales of [Pecaut & Mamajek \(2013\)](#), [Herczeg & Hillenbrand \(2014\)](#) and

[Luhman et al. \(2003b\)](#) for both the single and binary samples. Early K-type binaries (T Tau, RW Aur, and HQ Tau) show large discrepancies which is possibly due to wrong SpT classifications, as [Flores et al. \(2020\)](#) also found a cooler temperature for T Tau ($T_{\text{eff}}=3976\pm90$ K), cool spots or the binary nature of these YSOs biasing measurements to cooler temperatures. A more detailed multi-epoch and multi-wavelength study should be carried out at higher angular resolution to understand the large variation in temperatures for these early-K binaries.

A temperature scale for the single stars was computed using the mean value and the standard deviation within ± 0.5 spectral type sub-classes bins. This IGRINS temperature scale, reported in Table 4, represents an alternative to previous published pre-main-sequence stars temperature scales, with the advantage of being obtained through a simultaneous determination of various

Table 4. Temperature scales of Luhman et al. (2003b, L03), Pecaut & Mamajek (2013, PM13), Herczeg & Hillenbrand (2014, HH14) and that determined here for single YSOs. The last column correspond to the number of objects with which we computed the mean T_{eff} value and the standard deviation within a ± 0.5 spectral type subclasses. If just one object was included in the spectral type bin, we report the value of T_{eff} (and its error) found in our analysis.

SpT	L03	PM13	HH14	This work	N
	(K)	(K)	(K)	(K)	
K0	–	5030	4870	4606 ± 233	1
K2	–	4760	4710	4450 ± 123	1
K5	–	4140	4210	4055 ± 78	2
K7	–	3970	4020	3905 ± 63	2
M0	–	3770	3900	3769 ± 111	5
M1	3705	3630	3720	3696 ± 56	9
M2	3560	3490	3560	3535 ± 79	9
M3	3415	3360	3410	3496 ± 26	5
M4	3270	3160	3190	3319 ± 58	3
M5	3125	2880	2980	3269 ± 48	2

atmospheric parameters and the inclusion of the magnetic field effects.

4.4. Surface gravity, stellar ages and masses

The log g – T_{eff} plane, also known as the spectroscopic Hertzsprung-Russell diagram (sHR; Langer & Kudritzki 2014), helps estimate stellar properties such as age and mass because log g and T_{eff} are proxies for those properties (See Figure 9 of Sokal et al. 2018). According to the evolutionary models of Baraffe et al. (2015), over the mass range of 0.05 – $1.4 M_{\odot}$, the log g changes by about 0.5 dex between 1 to 10 Myr. Therefore, two populations of different ages should occupy different parts of the sHR.

With this in mind, we observed members of the TW Hydrae Association (TWA; Kastner et al. 1997). TWA is a nearby (~ 60 pc; Zuckerman & Song 2004; Gaia Collaboration et al. 2018) and young (~ 7 – 10 Myr; Ducourant et al. 2014; Herczeg & Hillenbrand 2015; Sokal et al. 2018) group of stars, that serves as an evolved counterpart to our Taurus sample. We analyzed and determined the stellar parameters (Table 5) for all 19 TWA objects in the same fashion as the Taurus sample. In Figure 8 we assemble the sHR of the Taurus and TWA YSOs, and compare them to the 1, ~ 5 , and 10 Myr isochrones of Baraffe et al. (2015).

As expected, the Taurus and TWA YSOs populate different parts of the sHR, exhibiting an offset in log g .

Most of the Taurus sample is located between the 1 and 5 Myr isochrones while the TWA members are located below the 5 Myr isochrone. We found a mean log g (and error on the mean) of 3.87 ± 0.03 dex and 4.22 ± 0.03 dex, for Taurus and TWA, respectively.

We applied to the log g distributions of Taurus and TWA a Kolmogorov-Smirnov (KS) test⁵. The KS test explores the null hypothesis that two samples are drawn from the same distribution. If the probability value ($\mathcal{P}(\text{KS})$) obtained from this test is higher than a certain threshold, often set at 1%, 5%, or 10%, we cannot reject the null hypothesis. We found a $\mathcal{P}(\text{KS}) \sim 10^{-8}$ that the log g distributions of Taurus and TWA come from the same parent distribution, validating that precise log g values can provide meaningful age proxies for populations of stars.

For the mean T_{eff} and log g values, and the set of isochrones from Baraffe et al. (2015), we can make a rough estimate of the age of Taurus and TWA. Our calculations lead to a Taurus mean age of ~ 2.5 Myr and a TWA mean age of ~ 10 Myr. However, the dispersion within these groups is larger than the differences between them. The parameters of the least evolved TWA members are more similar to the more evolved members of Taurus than to the most evolved members of TWA.

4.5. Class II vs. Class III YSOs

Based on the shape of their spectral energy distributions, Adams, Lada, & Shu (1987) divided YSOs into three different morphological classes (I, II, and III) using the spectral index $n = d \log(\lambda F_{\lambda}) / d \log \lambda$, and suggested that such classes represent an evolutionary sequence in the formation of low-mass stars. Since then, the sequence has been extended to Class 0 YSOs embedded in an infalling cold dust envelope (Andre et al. 1993), flat spectrum YSOs (Greene et al. 1994) between Class I and II, and post-Class III transition disk objects without the cold dust corresponding to flux from 5 – $20 \mu\text{m}$ (Lada et al. 2006). If the classification of YSOs by their SED slope, as suggested, represents an evolutionary scheme, the populations of YSOs would differ in terms of stellar properties, such as log g , given that stars undergo gravitational contraction while they evolve towards the main-sequence (Herbig 1962).

To test this, we collected YSO classifications from the literature for 84 Class II and 35 Class III (78 and 32 with good or acceptable parameters, respectively) objects in our sample (Kenyon & Hartmann 1995; Luhman et al. 2010; Rebull et al. 2010; Esplin et al. 2014; Kraus et al.

⁵ We have used the KS_2samp function, which is part of the python package Scipy.stats

Table 5. Stellar parameters and basic information for 19 TWA members. The quality flag is the same as in Table 1. K magnitudes come from 2MASS (Cutri et al. 2003) while the SpT references are indicated. The errors on the stellar parameters are the total (fit + systematic) uncertainties.

2MASS ID	Name	SpT	Ref.	K	flag	T_{eff}	log g	B	$v \sin i$	r_K
				(mag)		(K)	(dex)	(kG)	(km s ⁻¹)	
J11015191-3442170	TWA 1	M0.5	1	7.3	0	3783±108	4.35±0.18	2.75±0.30	8.4±2.0	0.53±0.05
J11091380-3001398	TWA 2	M2.2	1	6.7	0	3558± 92	4.07±0.16	<1.38	15.9±1.9	0.06±0.02
J11102788-3731520	TWA 3A	M4.1	1	6.8	1	3285± 75	4.00±0.13	<0.97	11.9±1.7	0.29±0.01
–	TWA 3B	M4.0	1	–	1	3355± 75	4.20±0.13	<1.78	15.8±1.7	0.16±0.01
J11220530-2446393	TWA 4	K6.0	1	5.6	0	4257±103	4.51±0.21	<1.10	10.2±2.1	0.31±0.04
J10423011-3340162	TWA 7	M3.2	1	6.9	0	3328± 80	4.27±0.13	2.19±0.26	7.4±1.7	0.10±0.01
J11324124-2651559	TWA 8A	M2.9	1	7.4	0	3398± 75	4.30±0.13	2.86±0.26	7.5±1.7	0.13±0.01
J11482422-3728491	TWA 9A	K6.0	1	7.8	0	4043± 94	4.32±0.17	2.29±0.30	12.0±2.0	0.17±0.03
J11482373-3728485	TWA 9B	M3.4	1	9.2	0	3351± 75	4.30±0.14	<1.30	10.9±1.7	0.09±0.01
J12350424-4136385	TWA 10	M3.0	2	8.2	1	3358± 75	4.23±0.13	2.54±0.26	9.5±1.7	0.10±0.01
J12360055-3952156	TWA 11B	M2.5	3	8.3	0	3532±115	4.22±0.18	2.23±0.34	14.8±2.1	0.02±0.02
12354893-3950245,	TWA 11C	M4.5	2	8.9	1	3391±122	4.45±0.22	<1.98	22.8±2.3	0.36±0.04
J11210549-3845163	TWA 12	M2.75	2	8.1	0	3534±104	4.19±0.17	2.67±0.32	19.7±2.0	0.02±0.02
J11211723-3446454	TWA 13A	M1.1	1	7.5	0	3638± 96	4.14±0.16	2.12±0.29	14.2±1.9	0.04±0.02
J11211745-3446497	TWA 13B	M1.0	1	7.5	0	3673± 93	4.13±0.16	1.71±0.30	13.7±1.9	0.04±0.02
J12345629-4538075	TWA 16	M3.0	2	8.1	0	3445± 91	4.15±0.16	1.85±0.28	12.4±1.9	0.07±0.02
J12072738-3247002	TWA 23	M3.5	1	7.8	1	3405±101	4.15±0.18	<1.42	19.1±1.9	0.16±0.03
J12153072-3948426	TWA 25	M0.5	1	7.3	0	3707± 99	4.14±0.17	2.21±0.29	15.0±1.9	0.06±0.03
J10120908-3124451	TWA 39	M4.0	4	8.0	1	3316± 88	4.06±0.18	<1.34	18.0±2.0	0.28±0.03

References— (1) Herczeg & Hillenbrand (2014); (2) Luhman et al. (2017); (3) Webb et al. (1999); (4) Riedel et al. (2014)

2017). We used a KS test to quantify if the stellar parameters determined for the Class II and Class III stars are statistically different.

We found a $\mathcal{P}(\text{KS})$ of 32%, 9%, and 6% that the Class II and Class III objects come from the same parent distributions of T_{eff} , $v \sin i$, and B respectively.⁶ This probability dropped to values lower than 1% for log g and r_K . These probability values indicate that the Class II and Class III YSOs of our sample are statistically indistinguishable in terms of T_{eff} , marginally indistinguishable in terms of $v \sin i$ and B, but different in terms of r_K and log g.

The result of the KS tests for T_{eff} and r_K are not surprising. The distribution of temperatures (SpT) reflects the initial mass function of Taurus. Also, Class II YSOs host disks and are actively accreting, which correlate with high levels of veiling, while Class III objects have lost their disks and show little or no veiling. The veiling values we measure for Class III objects is closer to zero than what we find for Class II YSOs, consistent with the expected behavior. For $v \sin i$, the $\mathcal{P}(\text{KS})$ of 9% that we found is in agreement with the findings of Nguyen et al. (2009), who found that low-mass accre-

tors and non-accretors in Taurus have a $\mathcal{P}(\text{KS}) = 10\%$ of coming from the same $v \sin i$ parent distribution.

Since log g can be used as a proxy for age (Yao et al. 2018), the different distributions for Class II and Class III YSOs support the idea of an evolutionary sequence (see Figure 9). We found a mean log g of 3.83 ± 0.03 and 4.06 ± 0.04 dex for Class II and Class III Taurus objects, respectively. The most interesting sources in each log g distribution are those in the wings, where evolutionary differences between the lowest log g Class III YSOs and the highest log g Class II YSOs are still not well understood.

Finally, the spatial distribution for the IGRINS Taurus sample (Figure 10) does not show significant clustering of Class II or Class III objects, nor a trend with log g, suggesting that there is not a star formation gradient across the cloud. Yet, this is a 2-dimensional view of Taurus, and it will require the entire population of Taurus YSOs (>400; Luhman 2018) studied in the context of 3-dimensional substructures (Krolkowski et al. 2021) before we can fully understand the spatial distribution in relation to the parameters we have determined.

4.6. Important Considerations

There are many ways to perform the parameter determination that we present here. As outlined above, there are a variety of models to choose from and possi-

⁶ We considered just the 41 B determinations in this test, not the limits.




Figure 8. Spectroscopic Hertzsprung-Russell diagram. The solid and empty circles represent our determinations for Class II and Class III Taurus YSOs, while the empty squares are our determinations for the TWA members. These different YSO samples present an offset in $\log g$, which suggests an age ladder with the Taurus YSOs younger than their TWA counterparts. We also include the Baraffe et al. (2015) isochrones of 1 (dashed line), ~ 5 (dash-dotted line) and 10 Myr (dotted line). The error bar represents our median fit uncertainties.

ble approaches. We have shown that magnetic fields are a critical component of the parameter determination, and we allowed many of the YSO parameters to remain variables in the fit rather than fixing them to assumed values. During our tests, we found that $\log g$ was the most sensitive parameter to variations in our approach. Additionally, we fit for all parameters simultaneously rather than taking an iterative approach to the fitting. While deciding on the best method for parameter determination, by minimizing fit residuals and validating the results against the literature, we have identified some important considerations when performing YSO parameter determinations.

Use of consistent T_{eff} estimators: IGRINS provides us with both the H- and K-band spectrum of our sources, but we have only employed the K-band in our analysis. There are temperature sensitive spectral features in the H-band (the OH region of López-Valdivia et al. 2019) that we initially included in our analysis. The result of including this H-band region was an offset in $\log g$ towards higher values, inconsistent with the evolution-

ary stage of the Taurus star-forming region. Using both the H- and K-band spectra meant that $\log g$ came primarily from the CO region and T_{eff} came from the OH region, which are about $0.7 \mu\text{m}$ apart. Furthermore, Gully-Santiago et al. (2017) found that T_{eff} varies as a function of wavelength when a single temperature component was fit to the IGRINS spectrum of the spotted YSO LkCa 4. The temperature in the CO region of LkCa 4 is ~ 700 K cooler than the value determined in the OH region (see Figure 3 of Gully-Santiago et al. 2017). In this work, differences between the H- and K-band temperatures were compensated for by the remaining stellar parameters, which primarily impacted the $\log g$ determination. While the broad spectral grasp of IGRINS permits the analysis of additional features, it also probes different components of the YSO photosphere, which can produce conflicting measurements. We have mitigated these effects by fitting only the K-band regions of the Taurus YSOs.

Degeneracy between r_K and $\log g$: Another consideration was the degeneracy between $\log g$ and r_K . As seen




Figure 9. Probability densities of $\log g$ for 78 Class II (dashed) and 32 Class III (solid) Taurus YSOs. We also included the $\log g$ probability density of the TWA YSOs as a comparison. We computed a KS probability between the Class II and Class III of $\sim 3 \times 10^{-3}$. If we compare the $\log g$ distributions of the Taurus and TWA YSOs, we also found low KS probabilities of $\sim 1.5 \times 10^{-9}$ and $\sim 7 \times 10^{-5}$, for Class II and Class III, respectively. These low probability values show that Taurus Class II and Class III YSOs, as well as the TWA objects, are different in terms of their $\log g$.




Figure 10. The spatial distribution of the Taurus sample color-coded by $\log g$. The 78 Class II (circles) and 32 Class III (crosses) YSOs are distributed similarly across the sky. There is no significant clustering of Class II or Class III objects in the sample we have studied.

in the diagonal distribution of r_K and $\log g$ in the sub-panel of Figure 3, higher values of r_K results in lower $\log g$ values, and vice versa. This degeneracy results at a fixed T_{eff} because r_K and $\log g$ impact the CO lines in nearly in the same way, and the CO lines are the most $\log g$ sensitive lines in our analysis. This degeneracy increases the uncertainties for each parameter and we concluded that $\log g$ and r_K need to be determined simultaneously and from the same spectrum. The use of fixed r_K or $\log g$ values from the literature will disproportionately impact the other parameter.

The atomic data: Our method relies on the comparison of observed and synthetic spectra, and the line list and the model atmosphere selected is a crucial component of the result. We initially determined the stellar parameters using the unmodified VALD line list expecting that the use of different spectral regions would average the inaccuracies of the atomic data. However, the resultant stellar parameters were offset from previous determinations in the literature. We then modified the oscillator strengths and the van der Waals constants of our spectral regions to the values of Flores et al. (2019), which improved the stellar parameter determinations and the consistency with the literature. We used the Flores et al. (2019) modifications because they were obtained from spectra with a spectral resolution close to IGRINS and for the same spectral regions we have employed.

Comparison with the literature: The bright YSOs in this work are canonical members of the Taurus-Auriga star forming region and have numerous published parameters. Many of the prior works in the literature (e.g., Johns-Krull 2007; Pecaute & Mamajek 2013; Herczeg & Hillenbrand 2014) fixed parameters that we have left as variables in the fits. Prior studies also analyzed shorter wavelength regions of the YSO spectra, which are dominated by the hotter photosphere. Additionally, the variable nature of YSOs means that observations made at different epochs likely have different physical properties. In comparing our results to the literature, we find that the observed spectra and methods of Flores et al. (2019) provide the most direct comparisons and consistency.

5. SUMMARY AND CONCLUSIONS

We employed four different spectral regions of the infrared K-band and an iterative MCMC approach to determine T_{eff} , $\log g$, B , and $v \sin i$ for a sample of canonical YSOs in the Taurus-Auriga star-forming region. Our analyses used a solar metallicity synthetic spectral grid of MARCS atmosphere models and the spectral syn-

thesis code MOOGSTOKES. Our observational data were obtained with the Immersion GRating INfrared Spectrometer (IGRINS).

We were able to reliably determine T_{eff} , $\log g$, and $v \sin i$ for 110 YSOs. Our ability to determine B is dependent on the $v \sin i$ and limited by the spectral resolution of IGRINS. We established that B values >1 kG or $(v \sin i/8)$ kG are required to provide meaningful B -field detections, and that $\text{SNR} > 150$ is desirable for overall parameter fitting. Measurable B -fields were found for 41 of the YSOs, and upper limits were identified for the remaining objects.

The T_{eff} we derived agrees with previous temperature scales for spectral type M YSOs, but it is systematically cooler for K stars. This could be a result of the models used in this analysis compared to others or the wavelength regions studied, which are more sensitive to the cool spotted regions of hotter YSOs. Our results show that the Class II and Class III objects are statistically similar in terms of T_{eff} , marginally similar in $v \sin i$ and B , and different in $\log g$ and r_K . The differences found in the distributions of $\log g$ and r_K correspond to the expected behavior if Class III are more evolved than Class II objects. We found mean $\log g$ values of 3.83 ± 0.03 , 4.06 ± 0.04 , and 4.22 ± 0.03 dex for the Taurus Class II and Class III objects and 19 TWA members, respectively. These differences are statistically significant and affirms the age gradient between these young star populations.

We thank the anonymous referees for suggestions that have improved our methodology and scientific results, as well as for inspiring the summary of important considerations in Section 4.6. This work used the Immersion Grating Infrared Spectrograph (IGRINS) that was de-

veloped under a collaboration between the University of Texas at Austin and the Korea Astronomy and Space Science Institute (KASI) with the financial support of the US National Science Foundation under grant AST-1229522 and AST-1702267, of the University of Texas at Austin, and of the Korean GMT Project of KASI. These results made use of the Lowell Discovery Telescope. Lowell is a private, non-profit institution dedicated to astrophysical research and public appreciation of astronomy and operates the LDT in partnership with Boston University, the University of Maryland, the University of Toledo, Northern Arizona University and Yale University. This paper includes data taken at The McDonald Observatory of The University of Texas at Austin. Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina), and Ministério da Ciência, Tecnologia e Inovação (Brazil). This material is based upon work supported by the National Science Foundation under Grant No. AST-1908892 to G. Mace.

Software: IGRINS pipeline package (Lee et al. 2017), MOOGSTOKES (Deen 2013), MARCS models (Gustafsson et al. 2008), ZBARYCORR (Wright & Eastman 2014), astroquery (Ginsburg et al. 2019), emcee (Foreman-Mackey et al. 2013), matplotlib (Hunter 2007), numpy (Harris et al. 2020), pandas (Wes McKinney 2010), astropy (Astropy Collaboration et al. 2013, 2018), scipy (Virtanen et al. 2020)

REFERENCES

- Adams, F. C., Lada, C. J., & Shu, F. H. 1987, *ApJ*, 312, 788
- ALMA Partnership, Brogan, C. L., Pérez, L. M., et al. 2015, *ApJL*, 808, L3
- Andre, P., Ward-Thompson, D., & Barsony, M. 1993, *ApJ*, 406, 122
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- Attridge, J. M., & Herbst, W. 1992, *ApJL*, 398, L61
- Baraffe, I., Homeier, D., Allard, F., & Chabrier, G. 2015, *A&A*, 577, A42
- Barenfeld, S. A., Carpenter, J. M., Sargent, A. I., Isella, A., & Ricci, L. 2017, *ApJ*, 851, 85
- Baruteau, C., Crida, A., Paardekooper, S. J., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 667
- Basri, G., & Batalha, C. 1990, *ApJ*, 363, 654
- Basri, G., & Bertout, C. 1989, *ApJ*, 341, 340
- Bean, J. L., Sneden, C., Hauschildt, P. H., Johns-Krull, C. M., & Benedict, G. F. 2006, *ApJ*, 652, 1604
- Boccaletti, A., Di Folco, E., Pantin, E., et al. 2020, *A&A*, 637, L5

- Bouvier, J., Matt, S. P., Mohanty, S., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 433
- Bouvier, J., Tessier, E., & Cabrit, S. 1992, *A&A*, 261, 451
- Calvet, N., Basri, G., & Kuhi, L. V. 1984, *ApJ*, 277, 725
- Camenzind, M. 1990, *Reviews in Modern Astronomy*, 3, 234
- Casagrande, L., Flynn, C., & Bessell, M. 2008, *MNRAS*, 389, 585
- Cieza, L. A., Kessler-Silacci, J. E., Jaffe, D. T., Harvey, P. M., & Evans, Neal J., I. 2005, *ApJ*, 635, 422
- Correia, S., Zinnecker, H., Ratzka, T., & Sterzik, M. F. 2006, *A&A*, 459, 909
- Cottaar, M., Covey, K. R., Meyer, M. R., et al. 2014, *The Astrophysical Journal*, 794, 125
- Crockett, C. J., Mahmud, N. I., Prato, L., et al. 2012, *ApJ*, 761, 164
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *VizieR Online Data Catalog*, II/246
- Deen, C. P. 2013, *AJ*, 146, 51
- Doppmann, G. W., Jaffe, D. T., & White, R. J. 2003, *AJ*, 126, 3043
- D’Orazi, V., Biazzo, K., & Randich, S. 2011, *A&A*, 526, A103
- Duchêne, G., Monin, J. L., Bouvier, J., & Ménard, F. 1999, *A&A*, 351, 954
- Ducourant, C., Teixeira, R., Galli, P. A. B., et al. 2014, *A&A*, 563, A121
- Dyck, H. M., Simon, T., & Zuckerman, B. 1982, *ApJL*, 255, L103
- Esplin, T. L., Luhman, K. L., & Mamajek, E. E. 2014, *ApJ*, 784, 126
- Fischer, W., Edwards, S., Hillenbrand, L., & Kwan, J. 2011, *ApJ*, 730, 73
- Flores, C., Connelley, M. S., Reipurth, B., & Boogert, A. 2019, *ApJ*, 882, 75
- Flores, C., Reipurth, B., & Connelley, M. S. 2020, *arXiv e-prints*, arXiv:2006.10139
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Gahm, G. F., Walter, F. M., Stempels, H. C., Petrov, P. P., & Herczeg, G. J. 2008, *A&A*, 482, L35
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1
- Galli, P. A. B., Bertout, C., Teixeira, R., & Ducourant, C. 2015, *A&A*, 580, A26
- Galli, P. A. B., Loinard, L., Ortiz-Léon, G. N., et al. 2018, *ApJ*, 859, 33
- Gennaro, M., Prada Moroni, P. G., & Tognelli, E. 2012, *MNRAS*, 420, 986
- Ghez, A. M., Neugebauer, G., & Matthews, K. 1993, *AJ*, 106, 2005
- Ginsburg, A., Sipőcz, B. M., Brasseur, C. E., et al. 2019, *AJ*, 157, 98
- Gray, D. F. 2005, *The Observation and Analysis of Stellar Photospheres* (Cambridge University Press), doi:10.1017/CBO9781316036570
- Greene, T. P., Wilking, B. A., Andre, P., Young, E. T., & Lada, C. J. 1994, *ApJ*, 434, 614
- Gully-Santiago, M., Wang, W., Deen, C., et al. 2010, in *Modern Technologies in Space- and Ground-based Telescopes and Instrumentation*, ed. E. Atad-Ettinger & D. Lemke, Vol. 7739, International Society for Optics and Photonics (SPIE), 1353 – 1359. <https://doi.org/10.1117/12.857568>
- Gully-Santiago, M. A., Herczeg, G. J., Czekala, I., et al. 2017, *ApJ*, 836, 200
- Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951
- Haas, M., Leinert, C., & Zinnecker, H. 1990, *A&A*, 230, L1
- Haisch, Karl E., J., Lada, E. A., & Lada, C. J. 2001, *ApJL*, 553, L153
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357–362
- Hartigan, P., Hartmann, L., Kenyon, S., Hewett, R., & Stauffer, J. 1989, *ApJS*, 70, 899
- Hartmann, L., Hewett, R., Stahler, S., & Mathieu, R. D. 1986, *ApJ*, 309, 275
- Hartmann, L. W., & Kenyon, S. J. 1990, *ApJ*, 349, 190
- Hayashi, C. 1961, *PASJ*, 13, 450
- Herbig, G. H. 1962, *Advances in Astronomy and Astrophysics*, 1, 47
- Herbig, G. H., & Bell, K. R. 1988, in *Third Catalog of Emission-Line Stars of the Orion Population : 3 : 1988*
- Herbst, W., Eislöffel, J., Mundt, R., & Scholz, A. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 297
- Herczeg, G. J., & Hillenbrand, L. A. 2014, *The Astrophysical Journal*, 786, 97
- . 2015, *ApJ*, 808, 23
- Hernández, J., Calvet, N., Briceño, C., et al. 2007, *ApJ*, 671, 1784
- Huang, J., Andrews, S. M., Cleeves, L. I., et al. 2018, *ApJ*, 852, 122
- Hunter, J. D. 2007, *Computing in Science Engineering*, 9, 90
- Hussaini, M., Mace, G. N., López-Valdivia, R., Honaker, E. J., & Han, E. 2020, *Research Notes of the American Astronomical Society*, 4, 241
- Ireland, M. J., & Kraus, A. L. 2008, *ApJL*, 678, L59

- Jaffe, D. T., Keller, L. D., & Ershov, O. A. 1998, in *Infrared Astronomical Instrumentation*, ed. A. M. Fowler, Vol. 3354, International Society for Optics and Photonics (SPIE), 201 – 212. <https://doi.org/10.1117/12.317263>
- Johns-Krull, C. M. 2007, *ApJ*, 664, 975
- Johns-Krull, C. M., & Valenti, J. A. 2001, *ApJ*, 561, 1060
- Johns-Krull, C. M., Valenti, J. A., & Koresko, C. 1999, *ApJ*, 516, 900
- Joy, A. H. 1949, *ApJ*, 110, 424
- Kastner, J. H., Zuckerman, B., Weintraub, D. A., & Forveille, T. 1997, *Science*, 277, 67
- Kenyon, S. J., Gomez, M., Marzke, R. O., & Hartmann, L. 1994, *AJ*, 108, 251
- Kenyon, S. J., & Hartmann, L. 1987, *ApJ*, 323, 714
- . 1995, *ApJS*, 101, 117
- Koenigl, A. 1991, *ApJL*, 370, L39
- Konopacky, Q. M., Ghez, A. M., Rice, E. L., & Duchêne, G. 2007, *ApJ*, 663, 394
- Kraus, A. L., Herczeg, G. J., Rizzuto, A. C., et al. 2017, *ApJ*, 838, 150
- Kraus, A. L., & Hillenbrand, L. A. 2009, *ApJ*, 704, 531
- Kraus, A. L., Ireland, M. J., Martinache, F., & Hillenbrand, L. A. 2011, *ApJ*, 731, 8
- Kraus, A. L., White, R. J., & Hillenbrand, L. A. 2006, *ApJ*, 649, 306
- Krolikowski, D. M., Kraus, A. L., & Rizzuto, A. C. 2021, *arXiv e-prints*, arXiv:2105.13370
- Lada, C. J., Muench, A. A., Luhman, K. L., et al. 2006, *AJ*, 131, 1574
- Langer, N., & Kudritzki, R. P. 2014, *A&A*, 564, A52
- Lavail, A., Kochukhov, O., & Hussain, G. A. J. 2019, *A&A*, 630, A99
- Lavail, A., Kochukhov, O., Hussain, G. A. J., et al. 2017, *A&A*, 608, A77
- Lee, J.-J., Gullikson, K., & Kaplan, K. 2017, *igrins/plp* 2.2.0. Zenodo, doi:10.5281/zenodo.845059. <https://doi.org/10.5281/zenodo.845059>
- Leinert, C., Haas, M., Mundt, R., Richichi, A., & Zinnecker, H. 1991, *A&A*, 250, 407
- Leinert, C., Zinnecker, H., Weitzel, N., et al. 1993, *A&A*, 278, 129
- Long, F., Herczeg, G. J., Harsono, D., et al. 2019, *ApJ*, 882, 49
- López-Santiago, J., Crespo-Chacón, I., Flaccomio, E., et al. 2016, *A&A*, 590, A7
- López-Valdivia, R. 2021a, *Reduced IGRINS Spectra: Taurus-Auriga*, vDRAFT VERSION, Harvard Dataverse, doi:10.7910/DVN/F9N2E0. <https://doi.org/10.7910/DVN/F9N2E0>
- . 2021b, *Reduced IGRINS spectra: TW Hydrae Association*, vDRAFT VERSION, Harvard Dataverse, doi:10.7910/DVN/BHSMT6. <https://doi.org/10.7910/DVN/BHSMT6>
- . 2021c, *Codes and tools for IGRINS data: Synthetic grid*, vDRAFT VERSION, Harvard Dataverse, doi:10.7910/DVN/ALOPIS. <https://doi.org/10.7910/DVN/ALOPIS>
- López-Valdivia, R., Mace, G. N., Sokal, K. R., et al. 2019, *ApJ*, 879, 105
- Luhman, K. L. 1999, *The Astrophysical Journal*, 525, 466
- . 2018, *AJ*, 156, 271
- Luhman, K. L., Allen, P. R., Espaillat, C., Hartmann, L., & Calvet, N. 2010, *The Astrophysical Journal Supplement Series*, 186, 111
- Luhman, K. L., Briceño, C., Stauffer, J. R., et al. 2003a, *The Astrophysical Journal*, 590, 348
- Luhman, K. L., Mamajek, E. E., Shukla, S. J., & Loutrel, N. P. 2017, *AJ*, 153, 46
- Luhman, K. L., Stauffer, J. R., Muench, A. A., et al. 2003b, *The Astrophysical Journal*, 593, 1093
- Mace, G., Kim, H., Jaffe, D. T., et al. 2016, in *Proc. SPIE, Vol. 9908, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 99080C
- Mace, G., Sokal, K., Lee, J.-J., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10702, *Proc. SPIE*, 107020Q
- Mamajek, E. E. 2009, in *American Institute of Physics Conference Series*, Vol. 1158, *Exoplanets and Disks: Their Formation and Diversity*, ed. T. Usuda, M. Tamura, & M. Ishii, 3–10
- Mann, A. W., Feiden, G. A., Gaidos, E., Boyajian, T., & von Braun, K. 2015, *ApJ*, 804, 64
- Mann, A. W., Gaidos, E., & Ansdell, M. 2013, *ApJ*, 779, 188
- Mathieu, R. D., Adams, F. C., & Latham, D. W. 1991, *AJ*, 101, 2184
- Mathieu, R. D., Martin, E. L., & Magazzu, A. 1996, in *American Astronomical Society Meeting Abstracts*, Vol. 188, *American Astronomical Society Meeting Abstracts #188*, 60.05
- Muzerolle, J., Calvet, N., Hartmann, L., & D’Alessio, P. 2003, *ApJL*, 597, L149
- Natta, A., Prusti, T., Neri, R., et al. 2001, *A&A*, 371, 186
- Newton, E. R., Charbonneau, D., Irwin, J., & Mann, A. W. 2015, *ApJ*, 800, 85
- Nguyen, D. C., Brandeker, A., van Kerkwijk, M. H., & Jayawardhana, R. 2012, *ApJ*, 745, 119
- Nguyen, D. C., Jayawardhana, R., van Kerkwijk, M. H., et al. 2009, *ApJ*, 695, 1648

- Nofi, L. A., Johns-Krull, C. M., López-Valdivia, R., et al. 2021, arXiv e-prints, arXiv:2103.01246
- Padgett, D. L. 1996, *ApJ*, 471, 847
- Park, C., Jaffe, D. T., Yuk, I.-S., et al. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9147, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 1
- Pecaut, M. J., & Mamajek, E. E. 2013, *ApJS*, 208, 9
- Pérez, S., Casassus, S., Hales, A., et al. 2020, *ApJL*, 889, L24
- Rajpurohit, A. S., Allard, F., Rajpurohit, S., et al. 2018, *A&A*, 620, A180
- Rajpurohit, A. S., Reylé, C., Allard, F., et al. 2013, *A&A*, 556, A15
- Rayner, J., Tokunaga, A., Jaffe, D., et al. 2016, in *Ground-based and Airborne Instrumentation for Astronomy VI*, ed. C. J. Evans, L. Simard, & H. Takami, Vol. 9908, International Society for Optics and Photonics (SPIE), 2401 – 2417. <https://doi.org/10.1117/12.2232064>
- Rebull, L. M., Padgett, D. L., McCabe, C.-E., et al. 2010, *ApJS*, 186, 259
- Reid, I. N., & Hawley, S. L. 2005, *New light on dark stars : red dwarfs, low-mass stars, brown dwarfs* (Springer, Berlin, Heidelberg), doi:10.1007/3-540-27610-6
- Ribas, Á., Bouy, H., & Merín, B. 2015, *A&A*, 576, A52
- Ribas, Á., Merín, B., Bouy, H., & Maud, L. T. 2014, *A&A*, 561, A54
- Richichi, A., Ragland, S., Calamai, G., et al. 1999, *A&A*, 350, 491
- Riedel, A. R., Finch, C. T., Henry, T. J., et al. 2014, *AJ*, 147, 85
- Ruíz-Rodríguez, D., Ireland, M., Cieza, L., & Kraus, A. 2016, *MNRAS*, 463, 3829
- Ryabchikova, T., Piskunov, N., Kurucz, R. L., et al. 2015, *PhyS*, 90, 054005
- Santos, N. C., Melo, C., James, D. J., et al. 2008, *A&A*, 480, 889
- Schaefer, G. H., Prato, L., Simon, M., & Patience, J. 2014, *AJ*, 147, 157
- Schmidt-Kaler, T. 1982, in *Landolt-Börnstein - Group VI Astronomy and Astrophysics*, ed. K. Schaifers & H. H. Voigt, Vol. 2B (Springer-Verlag Berlin Heidelberg). https://materials.springer.com/lb/docs/sm_lbs_978-3-540-31378-6_1
- Simon, M., Chen, W. P., Howell, R. R., Benson, J. A., & Slowik, D. 1992, *ApJ*, 384, 212
- Simon, M., Guilloteau, S., Beck, T. L., et al. 2019, *ApJ*, 884, 42
- Snedden, C. A. 1973, PhD thesis, THE UNIVERSITY OF TEXAS AT AUSTIN.
- Soderblom, D. R., Pendleton, J., & Pallavicini, R. 1989, *AJ*, 97, 539
- Sokal, K. R., Deen, C. P., Mace, G. N., et al. 2018, *ApJ*, 853, 120
- Sokal, K. R., Johns-Krull, C. M., Mace, G. N., et al. 2020, *ApJ*, 888, 116
- Straižys, V. 1992, *Multicolor stellar photometry* (Pachart Publishing House)
- Valenti, J. A., Basri, G., & Johns, C. M. 1993, *AJ*, 106, 2024
- Varga, J., Gabányi, K. É., Ábrahám, P., et al. 2017, *A&A*, 604, A84
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261
- Walter, F. M., Brown, A., Mathieu, R. D., Myers, P. C., & Vrba, F. J. 1988, *AJ*, 96, 297
- Wang, W., Gully-Santiago, M., Deen, C., Mar, D. J., & Jaffe, D. T. 2010, in *Modern Technologies in Space- and Ground-based Telescopes and Instrumentation*, ed. E. Atad-Etchedgui & D. Lemke, Vol. 7739, International Society for Optics and Photonics (SPIE), 1575 – 1583. <https://doi.org/10.1117/12.857164>
- Webb, R. A., Zuckerman, B., Platais, I., et al. 1999, *ApJL*, 512, L63
- Weintraub, D. A. 1989, PhD thesis, California Univ., Los Angeles.
- Wes McKinney. 2010, in *Proceedings of the 9th Python in Science Conference*, ed. Stéfan van der Walt & Jarrod Millman, 56 – 61
- Wright, J. T., & Eastman, J. D. 2014, *PASP*, 126, 838
- Yang, H., & Johns-Krull, C. M. 2011, *ApJ*, 729, 83
- Yang, H., Johns-Krull, C. M., & Valenti, J. A. 2005, *ApJ*, 635, 466
- Yao, Y., Meyer, M. R., Covey, K. R., Tan, J. C., & Da Rio, N. 2018, *ApJ*, 869, 72
- Yuk, I.-S., Jaffe, D. T., Barnes, S., et al. 2010, in *Proc. SPIE*, Vol. 7735, *Ground-based and Airborne Instrumentation for Astronomy III*, 77351M
- Zuckerman, B., & Song, I. 2004, *ARA&A*, 42, 685