

The Chow t -structure on the ∞ -category of motivic spectra

By TOM BACHMANN, HANA JIA KONG, GUOZHEN WANG, and ZHOULI XU

Abstract

We define the Chow t -structure on the ∞ -category of motivic spectra $\mathcal{SH}(k)$ over an arbitrary base field k . We identify the heart of this t -structure $\mathcal{SH}(k)^{c\heartsuit}$ when the exponential characteristic of k is inverted. Restricting to the cellular subcategory, we identify the Chow heart $\mathcal{SH}(k)^{c\heartsuit, \text{cell}}$ as the category of even graded $\text{MU}_2\text{-MU}$ -comodules. Furthermore, we show that the ∞ -category of modules over the Chow truncated sphere spectrum $\mathbb{1}_{c=0}$ is algebraic.

Our results generalize the ones in Gheorghe–Wang–Xu in three aspects: to integral results; to all base fields other than just $\mathbb{C}$; to the entire ∞ -category of motivic spectra $\mathcal{SH}(k)$, rather than a subcategory containing only certain cellular objects.

We also discuss a strategy for computing motivic stable homotopy groups of $(p$ -completed) spheres over an arbitrary base field k using the Postnikov–Whitehead tower associated to the Chow t -structure and the motivic Adams spectral sequences over k .

Contents

1. Introduction	708
1.1. Overview	708
1.2. Main result	709
1.3. Reconstruction theorems	710
1.4. Proof strategy of Theorem 1.5	713
1.5. Proof strategy for Theorem 1.1	713
1.6. Towards computing motivic stable homotopy groups of spheres	714
1.7. Organization	717
1.8. Conventions and notation	718
1.9. Acknowledgements	719
2. Elementary properties	719

Keywords: motivic homotopy theory, t -structure, homotopy groups of spheres, Adams spectral sequence

AMS Classification: Primary: 14F42, 18G80, 55Q45.

© 2022 Department of Mathematics, Princeton University.

2.1.	First properties	720
2.2.	Compatibility with filtered colimits	721
2.3.	Interaction with tensor products	721
2.4.	Right completeness	722
2.5.	Interaction with η -periodization	723
2.6.	Interaction with base change	723
2.7.	Cellular Chow t -structure	724
3.	Relationship to algebraic cobordism	726
3.1.	Chow degrees	726
3.2.	MGL-homology in a more general t -structure	727
3.3.	t -structure as a direct colimit	728
3.4.	MGL-completion and Adams–Novikov spectral sequence	730
3.5.	First consequences	731
4.	Reconstruction theorems	732
4.1.	Comonadic descent	733
4.2.	Modules over $\mathrm{MGL}_{c=0}$	736
4.3.	The Chow heart	742
4.4.	W -cellular objects	743
5.	Towards computations of motivic stable stems	749
5.1.	Algebraicity of the motivic Adams spectral sequence in the Chow heart	750
5.2.	The strategy	753
5.3.	Continuous Adams spectral sequences	755
5.4.	Over the complex numbers	757
5.5.	Over the real numbers	758
5.6.	Over finite fields	761
Appendix A.	A cell structure for MGL	765
Appendix B.	A vanishing result for algebraic cobordism	767
Appendix C.	Motivic cohomology	767
References		768

1. Introduction

1.1. *Overview.* Motivic stable homotopy theory, introduced by Voevodsky and Morel, is a subject that has successfully applied abstract homotopy theory to solve problems in number theory and algebraic geometry.

Recent work by Gheorghe–Wang–Xu [GWX21] has reversed this information flow — using motivic stable homotopy theory, the computation of classical stable homotopy groups of spheres, a fundamental problem in topology, has been extended into a much larger range [IWX20a], [IWX20b].

Gheorghe–Wang–Xu [GWX21] defined and studied a subcategory of (p -completed) cellular objects of the ∞ -category of motivic spectra over the

complex numbers $\mathcal{SH}(\mathbb{C})$, and they identify its heart as the abelian category of p -complete, even graded comodules over the Hopf algebroid $\mathrm{MU}_{2*}\mathrm{MU}$, which is equivalent to the abelian category quasi-coherent sheaves on the moduli stack of formal groups over $\mathbb{Z}_p$ -algebras, and is central to chromatic homotopy theory.

A natural question to ask is whether we can generalize this work

- to the entire motivic stable homotopy category, rather than just a subcategory consisting of certain cellular objects over the p -completed sphere spectrum;
- to an integral rather than p -complete result;
- and to other base fields than just $\mathbb{C}$.

In this paper, we achieve all three goals by defining a t -structure, which we call the *Chow t -structure*, on the motivic stable homotopy category $\mathcal{SH}(k)$ over any base field k . Its non-negative part $\mathcal{SH}(k)_{c \geq 0}$ is generated under colimits and extensions by Thom spectra $\mathrm{Th}(\xi)$ associated to K -theory points $\xi \in K(X)$ (equivalently formal differences of vector bundles $\xi = [V_1] - [V_2]$) on smooth and proper schemes X (see [Definition 2.1](#)). We denote the truncation functors by $E \mapsto E_{c=0}, E_{c \geq 0}$, and so on.

One of our key theorems is that the motivic bigraded homotopy groups of a homotopy object with respect to the Chow t -structure can be expressed by Ext groups over $\mathrm{MU}_{2*}\mathrm{MU}$ of its MGL homology ([Theorem 1.1](#)). Using the theory of comonadic descent, we give a concrete algebraic description of the Chow heart $\mathcal{SH}(k)^{c\heartsuit}$ ([Theorem 1.5](#)). We show that the ∞ -category of modules over $\mathbb{1}_{c=0}$ (where $\mathbb{1}$ denotes the motivic sphere spectrum) is algebraic ([Theorem 1.8](#)). Restricting to the subcategory of cellular modules over $\mathbb{1}_{c=0}$, we get an equivalence with Hovey's stable category $\mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU})$ ([Theorem 1.9](#)).

This equivalence between stable ∞ -categories results in an isomorphism between the motivic Adams spectral sequence and the algebraic Novikov spectral sequence, that is, the Adams spectral sequence in the stable category $\mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU})$. Since the latter spectral sequence is purely algebraic, its computation is much more accessible and can be done by computer programs. Therefore, this isomorphism allows us to fully compute the motivic Adams spectral sequence for all spectra in $\mathcal{SH}(k)^{c\heartsuit}$ within any reasonable range. We can then adapt the same methodology as in [\[GWX21\]](#), [\[IWX20a\]](#), [\[IWX20b\]](#); using naturality of the motivic Adams spectral sequences, we can obtain new information on differentials and extensions from spectra in the Chow heart $\mathcal{SH}(k)^{c\heartsuit}$ to the sphere spectrum, and therefore compute motivic stable stems over any base field k in a reasonable range.

1.2. Main result. From now on, we denote by e the exponential characteristic of the field k . We prove the following key theorem about the Chow t -structure:

THEOREM 1.1 (see [Theorem 3.14](#)). *Let $E \in \mathcal{SH}(k)[1/e]$. Then there is a canonical isomorphism*

$$\pi_{2w-s,w}E_{c=i} \cong \mathrm{Ext}_{\mathrm{MU}_{2*}\mathrm{MU}}^{s,2w}(\mathrm{MU}_{2*}, \mathrm{MGL}_{2*+i,*}E).$$

Here, on the right-hand side, s is the homological degree, and $2w$ is the internal degree.

Over $k = \mathbb{C}$ for E the sphere spectrum $\mathbb{1}$, [Theorem 1.1](#) tells us

$$\pi_{2w-s,w}\mathbb{1}_{c=0} \cong \mathrm{Ext}_{\mathrm{MU}_{2*}\mathrm{MU}}^{s,2w}(\mathrm{MU}_{2*}, \mathrm{MU}_{2*})$$

as an integral statement. This is in contrast to Isaksen's result [[Isa19](#), Prop. 6.2.5]

$$\pi_{2w-s,w}\mathbb{1}_p^\wedge/\tau \cong \mathrm{Ext}_{\mathrm{BP}_{2*}\mathrm{BP}}^{s,2w}(\mathrm{BP}_{2*}, \mathrm{BP}_{2*})$$

at each prime p , where $\tau : \Sigma^{0,-1}\mathbb{1}_p^\wedge \rightarrow \mathbb{1}_p^\wedge$ is the endomorphism of [[Ghe18](#), [Isa19](#)].

We have the following easy corollary of [Theorem 1.1](#), reproving [[Ghe18](#), Th. 3.13].

COROLLARY 1.2. *Over $\mathbb{C}$, both $\mathbb{1}_{c=0}$ and $\mathbb{1}_p^\wedge/\tau$ are $\mathcal{E}_\infty$ -rings.*

Remark 1.3. Another easy corollary of [Theorem 1.1](#) is that smashing with MGL detects Chow- ∞ -connectivity ([Corollary 3.17](#)).

1.3. *Reconstruction theorems.* It is shown in [[GWX21](#), Cor. 1.2] that the category $\mathbb{1}_p^\wedge/\tau\text{-Mod}$ of cellular modules over $\mathbb{1}_p^\wedge/\tau$ is purely algebraic:¹ the heart of the t -structure defined in [[GWX21](#)] is equivalent to $\mathrm{MU}_{2*}\mathrm{MU}_p^\wedge\text{-CoMod}$; the entire module category can be identified with Hovey's *stable category of comodules* $\mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU}_p^\wedge)$. (See [[Hov04](#)] for a definition of this category.) Other versions of this result over $\mathbb{C}$ can be found in [[Kra18](#)], [[Pst18](#)], [[GIKR21](#)]. We upgrade this to an integral result over arbitrary fields, using $\mathbb{1}_{c=0}$ as a replacement for $\mathbb{1}_p^\wedge/\tau$ (which makes sense by the discussion after [Theorem 1.1](#)).

1.3.1. *Description of the Chow heart.* The category $\mathcal{SH}(k)^{c\heartsuit}[1/e]$ can be described as a category of presheaves over the category of pure MGL-motives, together with some extra data.

Definition 1.4. The category of pure MGL-motives, denoted by $\mathrm{PM}_{\mathrm{MGL}}(k)$, is the smallest idempotent complete additive subcategory of $\mathrm{MGL}_{c=0}\text{-Mod}$ containing the object $X\{i\} := (\Sigma^{2i,i}X_+ \wedge \mathrm{MGL})_{c=0}$ for each $i \in \mathbb{Z}$ and smooth proper variety X .

¹It is known that over the complex numbers, the p -completion and the $\mathrm{H}\mathbb{Z}/p$ -nilpotent completion for the sphere are equivalent.

When it is clear from the context, we abuse notation and denote $X\{0\}$ by X .

It turns out that $\mathrm{PM}_{\mathrm{MGL}}(k)$ is an additive ordinary 1-category, and the mapping set $[X, Y\{*\}]_{\mathrm{PM}_{\mathrm{MGL}}(k)}$ is equivalent to $\mathrm{MGL}^{2*,*}(Y \times X)$. This definition of pure MGL-motives coincides with the definition in [NZ06] by taking the idempotent completion of the category of MGL-correspondences.

The category $\mathrm{PM}_{\mathrm{MGL}}(k)$ is naturally enriched in $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules. There is thus a classical notion of enriched presheaves on $\mathrm{PM}_{\mathrm{MGL}}(k)$ (see, e.g., [Rie14, §3.5]).

THEOREM 1.5 (see Section 4.3 and Remark 4.17). *The functor sending $F \in \mathcal{SH}(k)^{c\heartsuit}[1/e]$ to the presheaf on $\mathrm{PM}_{\mathrm{MGL}}(k)$ given by $F_*(X) = [\Sigma^{2*,*}X_+, F \wedge \mathrm{MGL}]$ induces an equivalence of categories between $\mathcal{SH}(k)^{c\heartsuit}[1/e]$ and the category of enriched presheaves on $\mathrm{PM}_{\mathrm{MGL}}(k)$ (with values in $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules).*

Remark 1.6. Explicitly, the structure of enriched presheaves on $\mathrm{PM}_{\mathrm{MGL}}(k)$ requires the following compatibility: for every graded MGL-correspondence $\alpha : X \rightarrow Y$, the following diagram commutes:

$$\begin{array}{ccc} F(Y)_* & \xrightarrow{\alpha^*} & F(X)_* \\ \Delta_{F,Y} \downarrow & & \Delta_{F,X} \downarrow \\ \mathrm{MU}_{2*}\mathrm{MU} \otimes_{\mathrm{MU}_{2*}} F(Y)_* & \xrightarrow{\Delta(\alpha)^*} & \mathrm{MU}_{2*}\mathrm{MU} \otimes_{\mathrm{MU}_{2*}} F(X)_* \end{array}$$

Here $\Delta(\alpha)$ is the effect of comultiplication. We give the full explicit description in Section 4.3.

Remark 1.7. P. Sechin is studying a category of “Landweber-equivariant Grothendieck motives” over a field k [Sec20]. One may show that this category embeds fully faithfully into $\mathcal{SH}(k)^{c\heartsuit}$.

1.3.2. *Algebraicity of $\mathbb{1}_{c=0}\text{-Mod}[1/e]$.* By [Bac21, Lemma 29],

$$\mathbb{1}_{c=0}\text{-Mod}[1/e]^{c\heartsuit} \simeq \mathcal{SH}(k)[1/e]^{c\heartsuit}.$$

The Chow heart $\mathcal{SH}(k)[1/e]^{c\heartsuit}$ is an abelian category. In general, for a compactly generated abelian category A viewed as the heart of its derived category, we consider the category $\mathcal{Hov}(A) := \mathrm{Ind}(\mathrm{Thick}(A^\omega))$, where A^ω denotes the subcategory of the compact objects (see Section 4.1.5). We prove that the module category $\mathbb{1}_{c=0}\text{-Mod}[1/e]$ is algebraic.

THEOREM 1.8 (see Proposition 4.3). *There is a symmetric monoidal equivalence*

$$\mathbb{1}_{c=0}\text{-Mod}[1/e] \simeq \mathcal{Hov}(\mathbb{1}_{c=0}\text{-Mod}[1/e]^{c\heartsuit}).$$

1.3.3. *Results in the cellular case.* In the cellular case, the results take the following simpler form.

THEOREM 1.9 (see [Corollary 4.29](#)).

(1) *There is an equivalence*

$$\mathcal{SH}(k)[1/e]^{\text{cell}, c\heartsuit} \simeq \text{MU}_{2*}\text{MU-CoMod}[1/e].$$

(2) *The cellular subcategory is equivalent to Hovey's stable category of comodules*

$$\mathbb{1}_{c=0}\text{-Mod}^{\text{cell}}[1/e] \simeq \mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})[1/e].$$

Remark 1.10. It comes as a surprise that [Theorem 1.9](#) does not depend on the base field k . In fact, let l/k be a field extension. The base change functor $\mathcal{SH}(k)^{\text{cell}} \rightarrow \mathcal{SH}(l)^{\text{cell}}$ is right- t -exact (for the Chow t -structures) and induces

$$\mathcal{SH}(k)[1/e]^{\text{cell}, c\heartsuit} \simeq \text{MU}_{2*}\text{MU-CoMod}[1/e] \simeq \mathcal{SH}(l)[1/e]^{\text{cell}, c\heartsuit}.$$

This does *not* imply that base change is conservative on cellular spectra that are Chow bounded below, since the Chow t -structure is not left complete (e.g., all η -periodic spectra are Chow ∞ -connective, by [Proposition 2.17](#)).

In more general cases involving field extensions, we have the following results.

Definition 1.11 (see [Definition 4.19](#)). Let W be a set of smooth proper schemes over k that contains $\text{Spec}(k)$ and is closed under finite products. Define the W -cellular category, denoted by $\mathcal{SH}(k)^{\text{wcell}}$ to be the subcategory of $\mathcal{SH}(k)$ generated under taking colimits and desuspensions by objects of the form $\text{Th}(\xi)$ for $\xi \in K(X)$ and $X \in W$.

We also define the Chow t -structure on the W -cellular category.

THEOREM 1.12 (see [Corollary 4.27](#) and [4.28](#)). *Let G and W be as in one of the following situations:*

- (1) *Let l/k be a finite Galois extension with Galois group G , and let W be $\{\text{Spec}(l') \mid l' \text{ is a subextension of } l/k\}$.*
- (2) *Let W be $\{\text{Spec}(l) \mid l/k \text{ is a finite separable extension}\}$, and let $G = \text{Gal}(k)$ be the absolute Galois group.*

We have

$$\begin{aligned} \mathcal{SH}(k)[1/e]^{\text{wcell}, c\heartsuit} &\simeq \underline{\text{MU}_{2*}}\text{MU-CoMod}[1/e], \\ \mathbb{1}_{c=0}\text{-Mod}[1/e]^{\text{wcell}} &\simeq \mathcal{H}\text{ov}(\underline{\text{MU}_{2*}}\text{MU})[1/e], \\ \text{MGL}_{c=0}\text{-Mod}[1/e]^{\text{wcell}, c\heartsuit} &\simeq \underline{\text{MU}_{2*}}\text{-Mod}[1/e]. \end{aligned}$$

Here $\underline{\mathrm{MU}}_{2*}\mathrm{MU}$ and $\underline{\mathrm{MU}}_{2*}$ denote the corresponding constant Mackey functors, and $\mathrm{-CoMod}$, Hov and $\mathrm{-Mod}$ are performed relative to the category of G -Mackey functors.

Remark 1.13. In [BHS20], Burklund, Hahn and Senger prove a similar result in the special case when the field extension is $\mathbb{C}/\mathbb{R}$ with the Galois group C_2 .

1.4. *Proof strategy of Theorem 1.5.* We obtain the description of the Chow heart using the theory of comonadic descent; to show it applies, we use results that follow from Theorem 1.1. In the discussion below we omit $[1/e]$ from the notation. More precisely, for a motivic commutative ring spectrum $A \in \mathrm{CAlg}(\mathcal{SH}(k))$, there is an induced Chow t -structure on the category $A\text{-Mod}$. (Just let the non-negative part be generated by free A -modules of the form $A \wedge \mathrm{Th}(\xi)$, for X smooth proper and $\xi \in K(X)$.) We focus on the induced t -structures on $\mathbb{1}_{c=0}\text{-Mod}$ and also $\mathrm{MGL}_{c=0}\text{-Mod}$. The free-forgetful adjunction

$$\mathbb{1}_{c=0}\text{-Mod} \rightleftarrows \mathrm{MGL}_{c=0}\text{-Mod}$$

defines a comonad C over $\mathrm{MGL}_{c=0}\text{-Mod}$. We show that when restricted to subcategories of suitably bounded objects, the adjunction is comonadic. Therefore $\mathbb{1}_{c=0}\text{-Mod}^{c^\heartsuit}$ is equivalent to the category of $C^{c^\heartsuit}$ -comodules in $\mathrm{MGL}_{c=0}\text{-Mod}^{c^\heartsuit}$, where $C^{c^\heartsuit}$ denotes the restriction of C .

Using spectral Morita theory, we can identify the category $\mathrm{MGL}_{c=0}\text{-Mod}$ as a presheaf category. Under this equivalence, we give an explicit description of the comonad $C^{c^\heartsuit}$. In the cellular case this turns out to be exactly the comonad describing $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules over MU_{2*} -modules. This description together with the equivalences $\mathcal{SH}(k)^{c^\heartsuit} \simeq \mathbb{1}_{c=0}\text{-Mod}^{c^\heartsuit} \simeq C^{c^\heartsuit}\text{-CoMod}$ results in Theorem 1.5.

Remark 1.14 (work of Bondarko). The Chow t -structure on $\mathrm{MGL}_{c=0}\text{-Mod}$ is adjacent to a weight structure in the sense of Bondarko [Bon10]. Bondarko's work then yields an explicit description of $\mathrm{MGL}_{c=0}\text{-Mod}^{c^\heartsuit} \simeq \mathrm{MGL}\text{-Mod}^{c^\heartsuit}$ (see [Bon10, Th. 4.4.2(4)]) and a similar description of all of $\mathrm{MGL}_{c=0}\text{-Mod}$ is easily obtained. This proof is equivalent to what we summarized above as “spectral Morita theory.”

Remark 1.15. The proofs show that the scalar extension functor $\mathrm{MGL}\text{-Mod} \rightarrow \mathrm{MGL}_{c=0}\text{-Mod}$ identifies with Bondarko's *weight complex functor*.

1.5. *Proof strategy for Theorem 1.1.* Before explaining some further applications of Theorem 1.1, let us sketch its proof. Our approach uses another t -structure on $\mathcal{SH}(k)$, the homotopy t -structure. Its non-negative part $\mathcal{SH}(k)_{\geq 0}$ is generated under colimits and extensions by $\{\Sigma_+^\infty X \wedge \mathbb{G}_m^{\wedge n} \mid n \in \mathbb{Z}, X \in \mathrm{Sm}_k\}$ (see Section 3.3). For all $d \geq 0$, the intersections

$$I^d := \Sigma^d \mathcal{SH}(k)_{\geq 0} \cap \mathcal{SH}(k)_{c \geq 0}$$

define a sequence of further t -structures. We write $\tau_{=0}^d$ for the 0-th truncation functor with respect to I^d . It turns out that these t -structures form a direct system with the Chow t -structure as the colimit (Lemma 3.13).

One key property of these t -structures is the following vanishing result.

PROPOSITION 1.16 (see Proposition 3.12). *Let $E \in \mathcal{SH}(k)[1/e]$.*

- (1) $\pi_{*,*}\tau_{=0}^d E$ is concentrated in Chow degrees ≤ 0 .
- (2) $\mathrm{MGL}_{2*,*}\tau_{=0}^d E$ equals $\mathrm{MGL}_{2*,*}E$ and vanishes for other bidegrees.

As a consequence of Proposition 1.16, the Adams–Novikov spectral sequence for $\tau_{=0}^d E$ collapses and converges to $\pi_{*,*}\tau_{=0}^d E$. In other words, we have canonical isomorphisms

$$\pi_{2w-s,w}(\tau_{=0}^d E) \cong \mathrm{Ext}_{\mathrm{MU}_*\mathrm{MU}}^{s,2w}(\mathrm{MU}_*, \mathrm{MGL}_{2*,*}E).$$

Since we can approximate $E_{c=0}$ as a colimit by $\tau_{=0}^d E$, the same result holds for the Chow t -structure. This is exactly the statement of Theorem 1.1.

1.6. *Towards computing motivic stable homotopy groups of spheres.* We now turn to computational applications of Theorems 1.1 and 1.9.

Computations of motivic stable homotopy groups of spheres is currently a very popular subject. Most results are in the following two categories, using mainly two different computational tools.

The first type of results gives a description for certain bidegrees of motivic stable stems over an arbitrary base field k , in terms of number theoretical invariants of the base field k . Two major results in this direction are

- Morel’s work [Mor06] on the 0-line, i.e., $\bigoplus_s \pi_{s,s}\mathbb{1}$, in terms of Milnor K-theory;
- Röndigs–Spitzweck–Østvær’s more recent work [RSØ19] on the 1-line, i.e., $\bigoplus_s \pi_{s+1,s}\mathbb{1}$, in terms of hermitian and Milnor K -groups of the base field k whose characteristic is not 2.

These computations use the slice spectral sequence and the results are for the integral sphere $\mathbb{1}$.

The second type of results determine the motivic stable stems over a specific field k in a reasonable but much larger range of bidegrees. The computational tool is the motivic Adams spectral sequence. These results provide p -primary information one prime at a time, where p is different from the exponential characteristic of the base field k . Most notably,

- over $\mathbb{C}$, Isaksen–Wang–Xu [IWX20a] have computed $\bigoplus_w \pi_{s,w}\mathbb{1}_2^\wedge$ for $s \leq 90$;
- over $\mathbb{R}$, Belmont–Isaksen [BI20] have computed $\bigoplus_w \pi_{s,w}\mathbb{1}_2^\wedge$ for $s - w \leq 11$;
- over finite fields, Wilson–Østvær [WØ17] have computed $\pi_{s,0}\mathbb{1}_2^\wedge$ for $s \leq 18$.

Our goal is to pursue the second direction using the motivic Adams spectral sequence, and to use the Chow t -structure to do computations over an arbitrary

field k , so the p -primary computations can be achieved in a large and reasonable range.

The computation of the E_2 -page of the motivic Adams spectral sequence depends on understanding the motivic homology of a point and the motivic Steenrod algebra action on it. The mod p homology of a point is computed in terms of Milnor K -theory by work of Voevodsky [Voe03a], [Voe11] when the base field contains a primitive p -th root of unity and has characteristic coprime to p . The motivic Steenrod algebra is computed by work of Voevodsky [Voe03b] and Hoyois–Kelly–Østvær [HKØ17]. With the knowledge of the structure of the motivic Steenrod algebra, the motivic Adams E_2 -page can be fully determined by a purely homological algebra computation.

The next step, which is also the hardest part, is to determine the motivic Adams differentials. According to the Mahowald Uncertainty Principle [Goe07], [Xu17], any Adams type spectral sequence that converges to the stable homotopy groups of spheres has infinitely many non-zero differentials after the E_2 -page, and any method that is used to compute these non-zero differentials leaves infinitely many unsolved by that method. The spirit of the Mahowald Uncertainty Principle is that we need to combine all known methods together to push forward these computations.

Nevertheless, classically and over the base field $\mathbb{C}$, Gheorghe–Wang–Xu [GWX21] has developed a new method to compute the motivic Adams differentials, which is so far the most efficient one. The most crucial part of this method is to obtain the following isomorphism of two spectral sequences, using an equivalence of two stable ∞ -categories.

THEOREM 1.17 (Theorem 1.3 of [GWX21]). *For each prime p , there is an isomorphism of spectral sequences, between the $\mathbb{H}\mathbb{Z}/p$ -based motivic Adams spectral sequence for $\mathbb{1}_p^\wedge/\tau$ and the algebraic Novikov spectral sequence for BP_{2*} .*

Since the algebraic Novikov spectral sequence is a purely algebraic spectral sequence, and can be computed in a large range by an automated computer program [IWX20a], [IWX20b], Theorem 1.17 allows us to obtain non-zero Adams differentials for free.

Using Theorems 1.8 and 1.9, we generalize Theorem 1.17 to the following Theorem 1.18 that works for any spectrum F in $\mathcal{SH}(k)^{c\heartsuit}$ and over an arbitrary base field k . This allows us to obtain non-zero Adams differentials for various spectra for free in a similar fashion to the case over $\mathbb{C}$ but over an arbitrary base field k .

In the discussion below, we again implicitly invert the exponential characteristic e of the base field throughout to ease notation.

Let $\mathbb{H}\mathbb{Z}/p$ denote the motivic Eilenberg–MacLane spectrum. Under the equivalence in Theorem 1.9,

$$\mathbb{1}_{c=0}\text{-}\mathcal{Mod}^{\mathrm{cell}} \simeq \mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU}),$$

the spectrum $\mathbb{H}\mathbb{Z}/p \wedge \mathbb{1}_{c=0}$ corresponds to the $\mathrm{MU}_{2*}\mathrm{MU}$ -comodule

$$H = \mathrm{MU}_{2*}\mathrm{MU}/(p, a_1, a_2, \dots).$$

The equivalence in [Theorem 1.5](#) gives an equivalence between the $\mathbb{H}\mathbb{Z}/p$ -based motivic Adams spectral sequences and the H -based Adams type spectral sequences in the category of stable $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules.

THEOREM 1.18 (see [Theorem 5.3](#) and [5.4](#)). *Let $F \in \mathcal{SH}(k)^{c\heartsuit}$. Let $M = \mathrm{MGL}_{2*,*}F$ be the associated $\mathrm{MU}_{2*}\mathrm{MU}$ -comodule. Then the trigraded motivic Adams spectral sequence for F based on $\mathbb{H}\mathbb{Z}/p$ is isomorphic (with all higher and multiplicative structure) to the trigraded algebraic Novikov spectral sequence based on H for M .*

See [Definition 5.2](#) for the algebraic Novikov spectral sequence for an object M in $\mathcal{H}\mathrm{ov}(\mathrm{MU}_{2*}\mathrm{MU})$ based on a commutative monoid H .

Now we discuss how to compute the motivic homotopy groups of the p -completion of a general motivic spectrum X with the property that $X \simeq X_{c \geq 0}$ using [Theorem 1.18](#). Note that the motivic sphere spectrum is such an example. Consider its Postnikov–Whitehead tower with respect to the Chow t -structure:

$$\begin{array}{ccc} & \downarrow & \\ & X_{c \geq 2} & \longrightarrow X_{c=2} \\ & \downarrow & \\ & X_{c \geq 1} & \longrightarrow X_{c=1} \\ & \downarrow & \\ X & \xlongequal{\quad} X_{c \geq 0} & \longrightarrow X_{c=0}. \end{array}$$

For every term $X_{c=n}$ in the Postnikov–Whitehead tower, consider the motivic Adams spectral sequences based on $\mathbb{H}\mathbb{Z}/p$ and the induced maps among them. This gives us the following tower of motivic Adams spectral sequences:

$$\begin{array}{ccc} & \downarrow & \\ & \mathbf{motASS}(X_{c \geq 2}) \rightarrow \mathbf{motASS}(X_{c=2}) = \mathbf{algNSS}(\mathrm{MGL}_{2*+2,*}X) & \\ & \downarrow & \\ & \mathbf{motASS}(X_{c \geq 1}) \rightarrow \mathbf{motASS}(X_{c=1}) = \mathbf{algNSS}(\mathrm{MGL}_{2*+1,*}X) & \\ & \downarrow & \\ \mathbf{motASS}(X) = \mathbf{motASS}(X_{c \geq 0}) \rightarrow \mathbf{motASS}(X_{c=0}) = \mathbf{algNSS}(\mathrm{MGL}_{2*,*}X). & & \end{array}$$

For every comodule $\mathrm{MGL}_{2*+n,*}X$, its algebraic Novikov spectral sequence can be computed in a large range by a computer program. This gives many non-zero differentials in the motivic Adams spectral sequence for $X_{c=n}$. We may then

pull back these Adams differentials to the motivic Adams spectral sequences for $X_{c \geq n}$ and push them forward to the motivic Adams spectral sequences for X .

The strategy for computing motivic Adams differentials for X can be summarized in the following steps. See [Section 5.2](#) for more details.

- (1) Compute $\text{Ext}_A^{*,*,*}(\mathbb{H}\mathbb{Z}/p_{*,*}, \mathbb{H}\mathbb{Z}/p_{*,*}X)$ over the k -motivic Steenrod algebra by a computer program. These Ext-groups consist of the E_2 -page of the motivic Adams spectral sequence for X based on $\mathbb{H}\mathbb{Z}/p$.
- (2) Compute by a computer program the algebraic Novikov spectral sequence based on $\text{BP}_{2*}\text{BP}/I$ for the BP_{2*}BP -comodules $\text{BPGL}_{2*+n,*}X$ for every n in a reasonable range.
- (3) Identify the k -motivic Adams spectral sequence based on $\mathbb{H}\mathbb{Z}/p$ for each $X_{c=n}$, with the algebraic Novikov spectral sequence based on $\text{BP}_{2*}\text{BP}/I$ for $\text{BPGL}_{2*+n,*}X$ for $n \geq 0$. This computation includes an identification of the abutments with the multiplicative (and higher) structures.
- (4) Compute the mod p motivic homology of $X_{c \geq n}$ using the universal coefficient spectral sequence (see Propositions 7.7 and 7.10 of [\[DI05\]](#))

$$\bigoplus_{k \geq n} \text{Tor}_{*,*}^{\text{BP}_{2*}}(\text{BPGL}_{2*+k,*}X, \mathbb{Z}/p) \implies \mathbb{H}\mathbb{Z}/p_{*,*}X_{c \geq n}.$$

- (5) Compute by a computer program the E_2 -pages of the $\mathbb{H}\mathbb{Z}/p$ -based motivic Adams spectral sequence for $X_{c \geq n}$, using the computation of $\mathbb{H}\mathbb{Z}/p_{*,*}X_{c \geq n}$ in step (4).
- (6) Pull back motivic Adams differentials for $X_{c=n}$ to motivic Adams differentials for $X_{c \geq n}$, and then push forward to motivic Adams differentials for X .
- (7) Deduce additional Adams differentials for X with a variety of ad hoc arguments. The most important methods are Toda bracket shuffles and comparison to known results in the $\mathbb{C}$ -motivic Adams spectral sequence.

The inputs of our strategy are $\mathbb{H}\mathbb{Z}/p_{*,*}X$ and $\text{MGL}_{*,*}X$ (or $\text{BPGL}_{*,*}X$). In the case of the sphere spectrum $X = \mathbb{1}$, it is more plausible to know $\pi_{*,*}\text{BPGL}_p^\wedge$ (over some fields) rather than $\text{BPGL}_{*,*}$. In this case, we modify our strategy by using pro-objects $(\mathbb{1}/p^n)_n$ as a substitute. See [Section 5.3](#) for more details.

As an example, we can show that over finite fields, our method gives proofs of previously undetermined Adams differentials in [\[WØ17\]](#). See [Section 5.6](#) for more details.

1.7. Organization. The rest of this article is organized as follows. In [Section 2](#), we define the Chow t -structure and discuss some of its basic properties. In [Section 3](#), we discuss the connection between the Chow t -structure and the homotopy t -structure. In particular, we prove one of our main theorems, [Theorem 1.1](#) as [Theorem 3.14](#). In [Section 4](#), we use the theory of comonadic descent, the Barr–Beck–Lurie theorem and spectral Morita theory to give a

concrete description of the Chow heart and to prove the algebraicity of the stable ∞ -category $\mathbb{1}_{c=0}\text{-Mod}[1/e]$. In particular, we prove [Theorems 1.5, 1.8, 1.9 and 1.12](#). In [Section 5](#), we prove [Theorem 1.18](#) and discuss applications toward computing motivic stable homotopy groups of spheres. In [Appendices A, B and C](#), we recall well-known results regarding a cell structure of MGL, a vanishing result of MGL, and motivic cohomology of a point.

1.8. *Conventions and notation.* Following, e.g., [NS17, Def. 2.1], we call a symmetric monoidal ∞ -category $\mathcal{C}$ *presentably symmetric monoidal* if $\mathcal{C}$ is presentable and the tensor product preserves colimits in each variable separately.

Given a morphism of schemes $f : X \rightarrow Y$, we denote by $f^* : \mathcal{SH}(Y) \rightarrow \mathcal{SH}(X)$ the induced base change functor and by $f_* : \mathcal{SH}(X) \rightarrow \mathcal{SH}(Y)$ its right adjoint. If f is smooth, then f^* has a left adjoint, which we denote by $f_\#$.

Here is a table of further notation that we employ.

$\underline{\text{map}}(-, -)$	internal mapping object	
Spc	∞ -category of spaces	
$\mathcal{SH}$	∞ -category of spectra	
Ab	1-category of abelian groups	
$\mathcal{P}_\Sigma, \mathcal{P}_{\mathcal{SH}}, \mathcal{P}_{\text{Ab}}$	product-preserving presheaves	Section 4.2.2
$\text{CB}^\bullet(A)$	cobar construction	[MNN17, Constr. 2.7]
$A\text{-Mod}$	modules over monoid A	[Lur17, Def. 4.2.1.13]
$C\text{-CoMod}$	comodules over comonad C	[Lur17, Def. 4.2.1.13]
$\mathcal{Hov}(C)$	stable category of comodules	Section 4.1.5
$\text{Alg}(-), \text{CAlg}(-)$	(commutative) algebra objects	
MU	complex cobordism spectrum	
$\text{Map}(-, -)$	mapping space in an ∞ -category	
$\underline{\text{map}}(-, -)$	mapping spectrum in a stable ∞ -category	
$[-, -]$	homotopy classes of maps, i.e. $\pi_0 \text{Map}(-, -)$	
$\mathcal{O}, \mathcal{O}_X$	trivial line bundle	
$n\mathcal{O}, \mathcal{O}^n$	trivial vector bundle of rank n	
Sm_S	category of smooth, finite type S -schemes	
$\text{Spc}(S)$	∞ -category of motivic spaces over S	[BH21, §2.2]
$\mathcal{SH}(S)$	∞ -category of motivic spectra over S	[BH21, §4.1]
$S^{p,q}$	bigraded sphere $S^{p-q} \wedge \mathbb{G}_m^{\wedge q}$	
$\Sigma^{p,q}$	functor of suspension by $S^{p,q}$	
$\text{Th}(\xi) = \text{Th}_S(\xi)$	Thom spectrum of $\xi \in K(X), X \in \text{Sm}_S$	[BH21, §16.2]
MGL	algebraic cobordism motivic spectrum	[BH21, §16.2]
$E_a^\wedge$	a -completion of a (motivic) spectrum	
$E_{\text{MGL}}^\wedge$	MGL-nilpotent completion	[Man21, §7.1]
$L_{\text{MGL}}E$	MGL-localization	[Man21, §2.4]
$\mathcal{SH}(S)_{\geq 0, \leq 0}$	homotopy t -structure	[BH21, §B]

$\pi_{p,q}E$	bigraded homotopy groups	
$\underline{\pi}_{p,q}E$	bigraded homotopy sheaves	
$\mathcal{SH}(S)_{c \geq 0, c \leq 0}$	Chow t -structure	Definition 2.1
$\mathcal{SH}(S)^{\text{pure}}$		Definition 2.10
$\mathcal{SH}(S)^{\text{lisse}}$		Definition 2.13
$\mathcal{SH}(S)^{\text{pure}, \text{ Tate} \geq d}$		Appendix A
$c(p, q)$	Chow degree	Definition 3.1

1.9. *Acknowledgements.* We thank Marc Levine and the University of Duisburg-Essen for the conference SPP 1786 Jahrestagung in 2019, where part of this collaboration was started. We thank Mark Behrens, Robert Burklund, Lars Hesselholt, Dan Isaksen, Fangzhou Jin, Haynes Miller, Markus Spitzweck, Glen Wilson, and Paul Arne Østvær for conversations regarding this project. We thank three anonymous referees for their careful read-through of the manuscript and helpful suggestions and comments. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1926686. The third author was partially supported by grant NSFC-11801082 and the Shanghai Rising-Star Program under agreement No. 20QA1401600. The fourth author was partially supported by the National Science Foundation under Grant No. DMS-2105462.

2. Elementary properties

Let S be a scheme.

Definition 2.1. Denote by $\mathcal{SH}(S)_{c \geq 0}$ the subcategory generated under colimits and extensions by motivic Thom spectra $\text{Th}(\xi)$ for X smooth and proper over S and $\xi \in K(X)$ arbitrary. This is the non-negative part of a t -structure on $\mathcal{SH}(S)$ [[Lur17](#), Prop. 1.4.4.11(2)] called the *Chow t -structure*. We denote the non-positive part by $\mathcal{SH}(S)_{c \leq 0}$, the heart by $\mathcal{SH}(S)^{c\heartsuit}$, and we write

$$E \mapsto E_{c \geq 0}, E_{c \leq 0}, E_{c=0}$$

for the truncation functors. We also put $\mathcal{SH}(S)_{c \geq n} = \Sigma^n \mathcal{SH}(S)_{c \geq 0}$ and define $\mathcal{SH}(S)_{c \leq n}, E_{c \geq n}$, etc. similarly.

Example 2.2. By definition, for any $n \in \mathbb{Z}$, the bigraded sphere $S^{2n,n} \simeq \text{Th}(n\mathcal{O})$ is in $\mathcal{SH}(S)_{c \geq 0}$.

Remark 2.3. For any ring spectrum $A \in \text{CAlg}(\mathcal{SH}(S))$, we can define a Chow t -structure on the category of A -modules. Its non-negative part is the subcategory generated under colimits and extensions by the free A -modules $A \wedge \text{Th}(\xi)$ for X smooth and proper over S and $\xi \in K(X)$.

2.1. *First properties.*

LEMMA 2.4. *Let $\mathcal{C}$ be a stable presentable category, $\mathcal{S} \subset \mathcal{C}$ a set of objects, and denote by $\mathcal{C}_{\geq 0} \subset \mathcal{C}$ the subcategory generated under colimits and extensions by $\mathcal{S}$. Then $E \in \mathcal{C}_{\leq 0}$ if and only if $[\Sigma^i S, E] = 0$ for all $S \in \mathcal{S}$ and $i > 0$.*

Proof. By definition, $E \in \mathcal{C}_{\leq 0}$ if and only if $\text{Map}(\Sigma X, E) = 0$ for all $X \in \mathcal{C}_{\geq 0}$ (see, e.g., [Lur17, Rem. 1.2.1.3]). Necessity is thus clear. To prove sufficiency, let $E \in \mathcal{C}$ satisfy the stated condition, and write $\mathcal{C}_E \subset \mathcal{C}$ for the subcategory of all $F \in \mathcal{C}$ such that $\text{Map}(\Sigma F, E) = 0$. Then $\mathcal{S} \subset \mathcal{C}_E$ by assumption, and $\mathcal{C}_E$ is closed under colimits and extensions. Thus $\mathcal{C}_{\geq 0} \subset \mathcal{C}_E$, which concludes the proof. $\square$

Thus given $E \in \mathcal{SH}(S)$, we have $E \in \mathcal{SH}(S)_{c \leq 0}$ if and only if $[\Sigma^i \text{Th}(\xi), E] = 0$ for all $i > 0$ (and ξ a K -theory point on a smooth proper scheme X over S). We will use this characterization without further comment from now on.

The category $\mathcal{SH}(S)$ is symmetric monoidal, so it carries a notion of strong duals (see, e.g., [DP83]). The following is a straightforward generalization of [Rio05, Th. 2.2].

LEMMA 2.5. *For $f : X \rightarrow S$ smooth and proper and $\xi \in K(X)$, the spectrum $\text{Th}_S(\xi) \in \mathcal{SH}(S)$ is strongly dualizable with dual $\text{Th}_S(-T_X - \xi)$, where T_X denote the tangent bundle. More generally, if $E \in \mathcal{SH}(X)$ is strongly dualizable with dual DE , then $f_{\#}(E)$ is strongly dualizable with dual $f_{\#}(\text{Th}(-T_X) \wedge DE)$.*

Proof. Since the Thom spectrum functor turns sums into smash products (essentially by construction, see, e.g., [BH21, §16.2]), $\text{Th}(\xi) \in \mathcal{SH}(X)$ is strongly dualizable (in fact invertible) with dual $\text{Th}(-\xi)$. By definition, we have $\text{Th}_S = f_{\#} \circ \text{Th}$; it thus suffices to prove the second statement. This follows from the natural equivalences

$$\begin{aligned} \underline{\text{map}}(f_{\#}(E), -) &\simeq f_* \underline{\text{map}}(E, f^*(-)) \simeq f_*(DE \wedge f^*(-)) \\ &\simeq f_{\#}(\text{Th}(-T_X) \wedge DE \wedge f^*(-)) \simeq f_{\#}(\text{Th}(-T_X) \wedge DE) \wedge -; \end{aligned}$$

here we used the relationship between $f^!$ and f^* and f_* [Hoy17, Th. 6.18(1,2)] and the projection formula [Hoy17, Th. 1.1(7)]. $\square$

Remark 2.6.

- (1) Note that the spectra $\text{Th}_S(\xi)$ are compact, being strongly dualizable in a category with compact unit (see, e.g., [Hoy17, Prop. 6.4(3)] for the latter).
- (2) The dualizability of the generators is important for many of the properties of the Chow t -structure.

2.2. Compatibility with filtered colimits.

PROPOSITION 2.7. *The Chow t -structure is compatible with filtered colimits: $\mathcal{SH}(S)_{c \leq 0}$ is closed under filtered colimits (and so are $\mathcal{SH}(S)_{c \geq 0}$ and $\mathcal{SH}(S)_{c=0}$).*

Proof. The first claim is immediate from Remark 2.6(1) (stating that $\mathrm{Th}(\xi)$ is compact) and Lemma 2.4 (stating that $E \in \mathcal{SH}(S)_{c \leq 0}$ if and only if $\mathrm{Map}(\Sigma \mathrm{Th}(\xi), E) \simeq 0$ for certain X, ξ). By definition $\mathcal{SH}(S)_{c \geq 0}$ is even closed under all colimits, and finally $\mathcal{SH}(S)_{c=0}$ is closed under filtered colimits, being the intersection of $\mathcal{SH}(S)_{c \leq 0}$ and $\mathcal{SH}(S)_{c \geq 0}$. $\square$

COROLLARY 2.8. *The functor $E \mapsto E_{c \leq 0} : \mathcal{SH}(S) \rightarrow \mathcal{SH}(S)$ preserves filtered colimits.*

Proof. The functor factors as $\mathcal{SH}(S) \rightarrow \mathcal{SH}(S)_{c \leq 0} \hookrightarrow \mathcal{SH}(S)$; here the first functor is a left adjoint by definition, and the second one preserves filtered colimits by Proposition 2.7. $\square$

2.3. Interaction with tensor products.

PROPOSITION 2.9. *The Chow t -structure is compatible with the symmetric monoidal structure; i.e., the non-negative part of the Chow t -structure is closed under taking smash products:*

$$\mathcal{SH}(S)_{c \geq 0} \wedge \mathcal{SH}(S)_{c \geq 0} \subset \mathcal{SH}(S)_{c \geq 0}.$$

Proof. Since $\mathcal{SH}(S)$ is presentably symmetric monoidal, it suffices to show that the generators of $\mathcal{SH}(S)_{c \geq 0}$ are closed under binary smash products. This is clear since $\mathrm{Th}(\xi) \wedge \mathrm{Th}(\zeta) \simeq \mathrm{Th}(\xi \times \zeta)$, and smooth proper S -schemes are closed under fiber products over S . $\square$

Definition 2.10. Denote by $\mathcal{SH}(S)^{\mathrm{pure}} \subset \mathcal{SH}(S)$ the smallest subcategory that is closed under filtered colimits and extensions and contains $\mathrm{Th}(\xi)$ for any K -theory point ξ on a smooth proper scheme X over S .

Note that by Definition 2.1, the subcategory $\mathcal{SH}(S)_{c \geq 0}$ is defined to be closed under arbitrary colimits. Thus $\mathcal{SH}(S)^{\mathrm{pure}} \subset \mathcal{SH}(S)_{c \geq 0}$. By Proposition 2.9, we have that $\mathcal{SH}(S)^{\mathrm{pure}} \wedge \mathcal{SH}(S)_{c \geq 0} \subset \mathcal{SH}(S)_{c \geq 0}$. Interestingly, the same holds for the non-negative part of the Chow t -structure, essentially by duality (see Lemma 2.5).

PROPOSITION 2.11. *We have*

$$\mathcal{SH}(S)^{\mathrm{pure}} \wedge \mathcal{SH}(S)_{c \leq 0} \subset \mathcal{SH}(S)_{c \leq 0}.$$

Proof. Let $\mathcal{C} \subset \mathcal{SH}(S)$ denote the subcategory of all spectra Z such that

$$Z \wedge \mathcal{SH}(S)_{c \leq 0} \subset \mathcal{SH}(S)_{c \leq 0}.$$

Since $\mathcal{SH}(S)_{c \leq 0}$ is closed under filtered colimits (by [Proposition 2.7](#)) and extensions (as is the non-positive part of any t -structure), so is the category $\mathcal{C}$. Hence to show that $\mathcal{SH}(S)^{\text{pure}} \subset \mathcal{C}$, it suffices to show that $\text{Th}(\zeta) \in \mathcal{C}$ for any K -theory point ζ on a smooth proper scheme X .

Let E be a spectrum in $\mathcal{SH}(S)_{c \leq 0}$, and let ξ be a K -theory point on a smooth proper scheme Y . For any $i > 0$, we find by duality (see [Remark 2.6\(2\)](#)) that

$$\begin{aligned} [\Sigma^i \text{Th}(\xi), E \wedge \text{Th}(\zeta)] &\simeq [\Sigma^i \text{Th}(\xi) \wedge \text{Th}(-T_X - \zeta), E] \\ &\simeq [\Sigma^i \text{Th}(\xi \times (-T_X - \zeta)), E] = 0. \end{aligned}$$

This concludes the proof. $\square$

COROLLARY 2.12. *For $X \in \mathcal{SH}(S)^{\text{pure}}$, $Y \in \mathcal{SH}(S)$, we have*

$$Y_{c \leq 0} \wedge X \simeq (Y \wedge X)_{c \leq 0}, \quad Y_{c \geq 0} \wedge X \simeq (Y \wedge X)_{c \geq 0}$$

and

$$Y_{c=0} \wedge X \simeq (Y \wedge X)_{c=0}.$$

In particular, (taking $Y = \mathbb{1}$) we have

$$X_{c \leq 0} \simeq X \wedge \mathbb{1}_{c \leq 0}.$$

Proof. For any $E \in \mathcal{SH}(S)$, there is a unique up to equivalence cofiber sequence

$$E' \rightarrow E \rightarrow E'',$$

where the first term is in $\mathcal{SH}(S)_{c \geq 0}$ and the last term is in $\mathcal{SH}(S)_{c < 0}$ [[BBD82](#), Prop. 1.3.3(ii)]. Therefore, any stable endofunctor of $\mathcal{SH}(S)$ preserving the non-negative and non-positive parts of a t -structure must commute with truncation. This holds for $- \wedge X$ by [Propositions 2.11](#) and [2.9](#). $\square$

2.4. Right completeness.

Definition 2.13. Write $\mathcal{SH}(S)^{\text{lisse}}$ for the stable presentable subcategory generated by $\mathcal{SH}(S)^{\text{pure}}$.

Thus $\mathcal{SH}(S)_{c \geq 0} \subset \mathcal{SH}(S)^{\text{lisse}}$, and so it defines a t -structure on $\mathcal{SH}(S)^{\text{lisse}}$. This is just the restriction of the Chow t -structure.

PROPOSITION 2.14. *The Chow t -structure on $\mathcal{SH}(S)^{\text{lisse}}$ is right complete; i.e., we have*

$$\cap_n \mathcal{SH}(S)_{c \leq n}^{\text{lisse}} = 0.$$

The above definition of right completeness coincides with other possible ones, by (the dual of) [[Lur17](#), Prop. 1.2.1.19] and [Proposition 2.7](#).

Proof. This is a formal consequence of the fact that $\mathcal{SH}(S)^{\text{lisse}}$ is generated by $\mathcal{SH}(S)_{c \geq 0}$. $\square$

Remark 2.15. The Chow t -structure is typically not left complete; i.e., $\cap_n \mathcal{SH}(S)_{c \geq n}^{\text{lisse}} \neq 0$. See [Proposition 2.17](#) below.

2.5. Interaction with η -periodization.

Definition 2.16. A spectrum $E \in \mathcal{SH}(S)^{\text{lisse}}$ is *Chow- ∞ -connective* if $E \in \mathcal{SH}(S)_{c \geq n}$ for all n .

PROPOSITION 2.17. *If $E \in \mathcal{SH}(S)^{\text{lisse}}$ is η -periodic, then E is Chow- ∞ -connective.*

Proof. Right completeness and compatibility with filtered colimits ([Proposition 2.7](#)) imply that

$$E \simeq \operatorname{colim}_{n \rightarrow -\infty} E_{c \geq n};$$

indeed the cofiber of the natural map is in $\mathcal{SH}(S)_{c \leq n}$ for every n . Since

$$\mathbb{G}_m^{\wedge -k} \simeq \Sigma^k \operatorname{Th}(-k\mathcal{O}) \in \mathcal{SH}(S)_{c \geq k},$$

we have $E_{c \geq n} \wedge \mathbb{G}_m^{\wedge -k} \in \mathcal{SH}(S)_{c \geq n+k}$ and hence

$$E \simeq E[1/\eta] \simeq \operatorname{colim}_{n \rightarrow -\infty} E_{c \geq n}[1/\eta] \simeq \operatorname{colim}_{n \rightarrow -\infty} \operatorname{colim}_{k \rightarrow \infty} E_{c \geq n} \wedge \mathbb{G}_m^{\wedge -k} \in \mathcal{SH}(S)_{c \geq \infty}.$$

This was to be shown. $\square$

Example 2.18. We work entirely in $\mathcal{SH}(S)^{\text{lisse}}$. For $E \in \mathcal{SH}(S)^{\text{lisse}}$, the fiber of the η -completion map $E \rightarrow E_\eta^\wedge$ is η -periodic, and hence Chow- ∞ -connective. Thus η -completion induces an equivalence on Chow homotopy objects.

Remark 2.19. Let k be a field, and let e be the exponential characteristic of k . Then $\mathcal{SH}(k)[1/e] \subset \mathcal{SH}(k)^{\text{lisse}}$ (combine [\[EK20, Cor. 2.1.7\]](#) and [\[LYZR19, Prop. B.1\]](#)). Hence the above results also apply to $\mathcal{SH}(k)[1/e]$.

2.6. Interaction with base change. For future reference, we record the following facts about the interaction between the Chow t -structure and base change. Recall the following standard definition.

Definition 2.20. Let $\mathcal{C}$ and $\mathcal{D}$ be two categories with t -structures. We say a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *right- t -exact* (respectively *left- t -exact*) if F preserves the non-negative (respectively non-positive) parts of t -structures.

PROPOSITION 2.21. *Let $f : S' \rightarrow S$ be a morphism of schemes.*

- (1) *The functor $f^* : \mathcal{SH}(S) \rightarrow \mathcal{SH}(S')$ is right- t -exact and $f_* : \mathcal{SH}(S') \rightarrow \mathcal{SH}(S)$ is left- t -exact.*
- (2) *If f is smooth and proper, then f^* is also left- t -exact and its left adjoint $f_\# : \mathcal{SH}(S') \rightarrow \mathcal{SH}(S)$ is right- t -exact.*

Proof. The right adjoint of a right- t -exact functor is left- t -exact and vice versa (see, e.g., [\[Bac18b, Lemma 5\]](#)). It thus suffices to show the right- t -exactness claims.

(1) For any $X \in \mathrm{Sm}_S$ and $\xi \in K(X)$, we have $f^*\mathrm{Th}(\xi) \simeq \mathrm{Th}(f^*\xi)$. It follows that f^* sends the generators of $\mathcal{SH}(S)_{c \geq 0}$ into $\mathcal{SH}(S')_{c \geq 0}$. The claim follows since f^* also preserves colimits and extensions.

(2) For $X \in \mathrm{Sm}_{S'}$ and $\xi \in K(X)$, we have $f_\#(\mathrm{Th}(\xi)) = \mathrm{Th}_S(\xi)$. Since f is proper, if X is proper over S' , then X is proper over S ; thus $f_\#$ preserves the generators of $\mathcal{SH}(-)_{c \geq 0}$. We conclude as in (1). $\square$

Remark 2.22. If f is not smooth and proper, then f^* is not in general left- t -exact. Indeed if this was the case for all f smooth, then $f_\#$ would be right- t -exact for all such f , and we would find that $\Sigma^{2n,n}\Sigma^\infty X_+ \in \mathcal{SH}(S)_{c \geq 0}$ for all $n \in \mathbb{Z}$ and all $X \in \mathrm{Sm}_S$. This would imply that $\mathcal{SH}(S)_{c \geq 0} = \mathcal{SH}(S)$, which is false in general (see, e.g., [Proposition 3.6](#)).

Example 2.23. We give an explicit example showing that f^* fails to be left- t -exact when f is not smooth and proper.

Let k be an infinite perfect field, and let U be an open subscheme of $\mathbb{A}_k^1$ such that $\emptyset \neq U \subsetneq \mathbb{A}_k^1$. Then

$$\tilde{\Sigma}U \simeq \mathbb{A}_k^1/U \simeq \bigvee_{x \in \mathbb{A}_k^1 \setminus U} S^{2,1} \wedge x_+,$$

where by $\tilde{\Sigma}$ we mean the unreduced suspension. It follows from [Proposition 3.6](#) that the spheres $S^{2,1} \wedge x_+$ are not in $\mathcal{SH}(S)_{c \geq 1}$, and consequently $\Sigma^\infty \tilde{\Sigma}U \notin \mathcal{SH}(k)_{c \geq 1}$. Since k is infinite, U has a rational point and so $\Sigma_+^\infty U \simeq \mathbb{1} \vee \Sigma^{-1}\Sigma^\infty \tilde{\Sigma}U$. Writing $f_U : U \rightarrow \mathrm{Spec}(k)$ for the structure morphism, we deduce that $f_{U\#}(\mathbb{1}) \simeq \Sigma_+^\infty U \notin \mathcal{SH}(k)_{c \geq 0}$. In particular, f_U^* is not left- t -exact.

Example 2.24. We now improve the construction of the previous example to show that even the base change along field extensions is not exact. Suppose that $0 \notin U$. Then $\Sigma_+^\infty U$ admits $\Sigma^{-1}S^{2,1}$ as a retract. In fact, we find that the pro-object

$$\text{“}\lim_{\emptyset \neq U} \text{”} \Sigma_+^\infty U$$

also admits $S^{2,1}$ as a retract. Writing

$$f : \mathrm{Spec}(k(t)) = \lim_{\emptyset \neq U \subset \mathbb{A}^1 \setminus 0} U \rightarrow \mathrm{Spec}(k)$$

we find that by continuity (see, e.g., [\[Hoy15, Lemma A.7\]](#))

$$[\Sigma^{-2,-1}\mathbb{1}, f^*\Sigma^{-1}(\mathbb{1}_{c \leq 0})]_{\mathcal{SH}(k(t))} \simeq \mathrm{colim}_U [\Sigma^{-2,-1}U_+, \Sigma^{-1}(\mathbb{1}_{c \leq 0})]_{\mathcal{SH}(k)} \neq 0,$$

since the latter group admits $[\mathbb{1}, \mathbb{1}_{c \leq 0}]_{\mathcal{SH}(k)}$ as a summand (which is non-zero by [Theorem 3.14](#)). It follows that f^* is not t -exact.

2.7. Cellular Chow t -structure. We denote by $\mathcal{SH}(S)^{\mathrm{cell}} \subset \mathcal{SH}(S)$ the subcategory of *cellular spectra*, i.e., the localizing subcategory generated by the bigraded spheres $S^{p,q}$, $p, q \in \mathbb{Z}$.

Definition 2.25. Denote by $\mathcal{SH}(S)_{c \geq 0}^{\text{cell}}$ the subcategory generated under colimits and extensions by $S^{2n,n}$ for $n \in \mathbb{Z}$. This is the non-negative part of a t -structure on $\mathcal{SH}(S)^{\text{cell}}$ [Lur17, Prop. 1.4.4.11(2)] called the *cellular Chow t -structure*. We denote the non-positive part by $\mathcal{SH}(S)_{c \leq 0}^{\text{cell}}$.

Remark 2.26. By definition we have

$$\mathcal{SH}(S)_{c \geq 0}^{\text{cell}} \subset \mathcal{SH}(S)_{c \geq 0} \cap \mathcal{SH}(S)^{\text{cell}},$$

but we do not know if the reverse inclusion holds, in general. This would hold if cellularization is t -exact (rather than just left t -exact, which it always is). Lemma 4.22 implies that this is true over fields.

Lemma 2.4 characterizes $\mathcal{SH}(S)_{c \leq 0}^{\text{cell}}$ in terms the vanishing of certain homotopy groups.

We now have enough notation in place to justify the multiplicative aspects of $\mathbb{1}_{c=0}$ and $\mathbb{1}_p^\wedge/\tau$ over $\mathbb{C}$ more fully. Specifically, we make the following claims:

- (1) The truncation functor $\mathcal{SH}(k)_{c \geq 0} \rightarrow \mathcal{SH}(k)_{c=0}$ is symmetric monoidal.
- (2) Its right adjoint $\mathcal{SH}(k)_{c=0} \hookrightarrow \mathcal{SH}(k)_{c \geq 0}$ is lax symmetric monoidal.
- (3) The p -completion functor $\mathcal{SH}(k) \rightarrow \mathcal{SH}(k)_p^\wedge$ is symmetric monoidal.
- (4) Its right adjoint $\mathcal{SH}(k)_p^\wedge \hookrightarrow \mathcal{SH}(k)$ is lax symmetric monoidal.
- (5) The canonical inclusion $\mathcal{SH}(k)^{\text{cell}} \hookrightarrow \mathcal{SH}(k)$ is symmetric monoidal.
- (6) Its right adjoint (cellularization) $\mathcal{SH}(k) \rightarrow \mathcal{SH}(k)^{\text{cell}}$ is lax symmetric monoidal.

The right adjoint of any symmetric monoidal functor is lax symmetric monoidal, whence statements (2), (4), and (6) follow from (1), (3) and (5), respectively, and (5) is obvious. (1) and (3) hold because the functors are localizations compatible with the tensor product (i.e., the tensor product preserves weak equivalences in either variable), which for (1) follows from Proposition 2.9 and for (3) follows from the fact that the tensor product preserves cofibers in each variable.

Proof of Corollary 1.2. Recall that over $\mathbb{C}$, $\tau : \Sigma^{0,-1} \mathbb{1}_p^\wedge \rightarrow \mathbb{1}_p^\wedge$ is the endomorphism of [Ghe18], [Isa19]. Since $S^{0,-1} = \Sigma^2 \text{Th}(-\mathcal{O}) \in \Sigma \mathcal{SH}(k)_{c > 0}$, we find that the canonical map $\mathbb{1}_p^\wedge \rightarrow (\mathbb{1}_{c=0})_p^\wedge$ annihilates $\tau : \Sigma^{0,-1} \mathbb{1}_p^\wedge \rightarrow \mathbb{1}_p^\wedge$. There is thus an induced map

$$\mathbb{1}_p^\wedge/\tau \rightarrow (\mathbb{1}_{c=0})_p^\wedge,$$

which by Theorem 1.1 and [Isa19, Prop. 6.2.5] induces an isomorphism on $\pi_{*,*}$. We deduce that $\mathbb{1}_p^\wedge/\tau$ is the p -complete cellularization of $\mathbb{1}_{c=0}$. The above discussion implies that $\mathbb{1}_{c=0}$ and thus $\mathbb{1}_p^\wedge/\tau$ are both $\mathcal{E}_\infty$ -rings, reproving [Ghe18, Th. 3.13]. $\square$

Most results established above for the Chow t -structure on $\mathcal{SH}(S)$ also hold for the cellular version.

Example 2.27. Let k be a field of characteristic $\neq p$ with $k^\times/p \simeq \{1\}$, and containing a primitive p -th root of unity. Then by [Voe11], we have $\pi_{*,*}(\mathrm{HZ}/p) \simeq K_*^M(k)/p[\tau] \simeq \mathbb{F}_p[\tau]$. This implies that $\mathrm{HZ}/(p, \tau) \in \mathcal{SH}(k)^{\mathrm{cell}, c^\heartsuit}$. Indeed we have $\mathrm{MGL} \in \mathcal{SH}(k)_{c \geq 0}^{\mathrm{cell}}$ (Theorem A.1). By the Hopkins–Morel isomorphism [Hoy15], we have $\mathrm{HZ} \in \mathcal{SH}(k)_{c \geq 0}^{\mathrm{cell}}$. Therefore $\mathrm{HZ}/(p, \tau) \in \mathcal{SH}(k)_{c \geq 0}^{\mathrm{cell}}$. On the other hand, $\mathrm{HZ}/(p, \tau) \in \mathcal{SH}(k)_{c \leq 0}^{\mathrm{cell}}$ just means that $\pi_{2m+i,m}(\mathrm{HZ}/(p, \tau))$ vanishes for $m \in \mathbb{Z}, i > 0$, which is clear since $K_*^M(k)/p \simeq \mathbb{F}_p$.

Remark 2.28. Continuing Example 2.27, write

$$P^i : \mathrm{HZ}/p \rightarrow \Sigma^{2i(p-1), i(p-1)} \mathrm{HZ}/p$$

for the reduced power operation. Since P^i commutes with τ (note that if $p = 2$, then $\sqrt{-1} \in k$ since $k^\times/2 \simeq \{1\}$), it defines

$$P^i/\tau : \mathrm{HZ}/(p, \tau) \rightarrow \Sigma^{2i(p-1), i(p-1)} \mathrm{HZ}/(p, \tau).$$

Noting that $\Sigma^{2n,n} \mathrm{HZ}/(p, \tau) \in \mathcal{SH}(k)^{\mathrm{cell}, c^\heartsuit}$, which is an additive 1-category, we deduce that the operation P^i/τ is $\mathbb{Z}/p$ -linear, i.e., it comes from a map in the $\mathbb{A}^1$ -derived category.

3. Relationship to algebraic cobordism

In this section, we show that the Chow t -structure is the colimit of a directed systems of t -structures, in the sense of Lemma 3.13. The non-negative parts and the non-positive parts of these t -structures can be partially characterized by taking MGL-homology (Lemma 3.12). As a result, the homotopy groups of Chow homotopy objects can be computed by the motivic Adams–Novikov spectral sequence, which collapse at the E_2 -page (Theorem 3.14). In fact, we prove these results for a more general collection of t -structures, which include Chow t -structure as a special case.

Throughout this section, we let $S = \mathrm{Spec}(k)$ be the spectrum of a field of exponential characteristic e , and we implicitly invert e throughout. Thus we write $\mathcal{SH}(k)$ for $\mathcal{SH}(k)[1/e]$, and so on. The main reason for this is that we need to use a vanishing result for algebraic cobordism (Theorem B.1), which is proved by relating algebraic cobordism to higher Chow groups, and this relationship is currently only known away from the characteristic [Hoy15]. Another simplification also occurs, which is that Remark 2.19 applies.

3.1. Chow degrees.

Definition 3.1. Given a bidegree $(p, q) \in \mathbb{Z} \times \mathbb{Z}$, we call $c(p, q) = p - 2q$ the *associated Chow degree*. Given a bigraded abelian group $M_{*,*}$, we say that $M_{*,*}$ is concentrated in Chow degrees $\geq d$ (respectively $\leq d, = d$) if $M_{p,q} = 0$ for $c(p, q) < d$ (respectively $> d, \neq d$).

Example 3.2. By [Example 2.2](#), spheres of non-negative Chow degree are in $\mathcal{SH}_{c \geq 0}(k)$. Therefore, if $E \in \mathcal{SH}(k)_{c \leq 0}$, then $\pi_{*,*}E$ is concentrated in Chow degrees ≤ 0 .

Example 3.3. Let $E \in \mathcal{SH}(k)$ with $\pi_{*,*}E$ concentrated in Chow degrees ≤ 0 . Consider the cofiber sequence

$$\mathbb{G}_m^{\wedge n} \wedge E \xrightarrow{\eta^n} E \rightarrow E/\eta^n.$$

The homotopy groups $\pi_{*,*}(\mathbb{G}_m^{\wedge n} \wedge E) \simeq \pi_{*-n, *-n}E$ are concentrated in Chow degree $\leq -n$. Hence we have $\pi_{p,q}(E/\eta^n) \simeq \pi_{p,q}(E)$ for any fixed (p, q) and n sufficiently large. As a result, $E \rightarrow E/\eta^n$ induces an isomorphism on $\pi_{*,*}$.

3.2. MGL-homology in a more general t -structure. In anticipation of our treatment of cellular objects later, we generalize our setting slightly. Thus we fix

$$I \subset \mathcal{SH}(k)_{c \geq 0}.$$

We assume throughout that I is closed under colimits and extensions and generated by a set of compact object; thus, in particular, I defines a t -structure. We denote its non-negative and non-positive parts by

$$\mathcal{SH}(k)_{I \geq 0} := I \text{ and } \mathcal{SH}(k)_{I \leq 0},$$

and we denote the truncations by $\tau_{I \geq n}, \tau_{I \leq n}$ and $\tau_{I=n}$. We also assume that $\mathbb{1} \in I$ and $\Sigma^{2,1}I = I$.

Example 3.4. Taking $I = \mathcal{SH}(k)_{c \geq 0}$, we get $\tau_{I \geq 0} = (-)_{c \geq 0}$, and so on. This is the maximal choice.

Example 3.5. Taking I to be the subcategory generated under colimits and extensions by the $S^{2n,n}$, we obtain the smallest possible choice.

PROPOSITION 3.6. *Let $E \in \mathcal{SH}(k)_{I \geq 0}$ (resp. $E \in \mathcal{SH}(k)_{I \leq 0}$, $E \in \mathcal{SH}(k)_{I=0}$). Then*

- (1) $\text{MGL} \wedge E \in \mathcal{SH}(k)_{I \geq 0}$ (resp. $\text{MGL} \wedge E \in \mathcal{SH}(k)_{I \leq 0}$, $\text{MGL} \wedge E \in \mathcal{SH}(k)_{I=0}$),
and
- (2) $\text{MGL}_{*,*}E$ is concentrated in Chow degrees ≥ 0 (resp. ≤ 0 , $= 0$).

Proof. (1) In the notation of [Section A](#), we have $\text{MGL} \in \mathcal{SH}(k)^{\text{pure Tate} \geq 0}$ (see [Theorem A.1](#)). Thus (1) is immediate from (the proofs of) [Propositions 2.9](#) and [2.11](#).

(2) The ≤ 0 part of (2) follows from (1) and the fact that $S^{2n,n} \in I$. It hence remains to prove the ≥ 0 part of (2). It is sufficient to show the statement is true for the maximal choice of I ([Example 3.4](#)).

Let $\mathcal{C} \subset \mathcal{SH}(k)$ denote the subcategory of all spectra E such that $\text{MGL}_{*,*}E$ is concentrated in Chow degrees ≥ 0 . It suffices to show that $\mathcal{SH}(k)_{c \geq 0} \subset \mathcal{C}$.

The category $\mathcal{C}$ is closed under small sums, cofibers, and extensions, and hence to show that $\mathcal{SH}(k)_{c \geq 0} \subset \mathcal{C}$ it is enough to show that $\mathrm{Th}(\xi) \in \mathcal{C}$ for every K -theory point ξ on a smooth proper variety X . We thus need to show that $\mathrm{MGL}_{2n+i,n} \mathrm{Th}(\xi) = 0$ for $i < 0$. Using the Thom isomorphism and duality we rewrite this as

$$[S^{2n+i,n}, \mathrm{MGL} \wedge \mathrm{Th}(\xi)] \cong [\Sigma^{2n+i,n} \mathrm{Th}(-T_X - \xi), \mathrm{MGL}] \cong \mathrm{MGL}^{2s-i,s} X,$$

where $s = \dim X + \mathrm{rank}(\xi) - n$. Thus the result follows from [Theorem B.1](#). $\square$

COROLLARY 3.7. *For $E \in \mathcal{SH}(k)$ arbitrary, we have*

$$\mathrm{MGL}_{p,q} \tau_{I=i} E = \begin{cases} 0 & \text{if } c(p, q) \neq i, \\ \mathrm{MGL}_{p,q}(E) & \text{if } c(p, q) = i. \end{cases}$$

Proof. By replacing E with $\Sigma^i E$, we reduce to the case that $i = 0$.

The Chow degree non-zero part of the statement is immediate from [Proposition 3.6](#).

For the Chow degree zero part, we consider two cofiber sequences

$$\tau_{I \geq 1} E \rightarrow \tau_{I \geq 0} E \rightarrow \tau_{I=0} E, \text{ and } \tau_{I \geq 0} E \rightarrow E \rightarrow \tau_{I \leq -1} E.$$

They yield associated long exact sequences

$$\begin{aligned} \cdots \rightarrow \mathrm{MGL}_{2*,*} \tau_{I \geq 1} E &\rightarrow \mathrm{MGL}_{2*,*} \tau_{I \geq 0} E \rightarrow \mathrm{MGL}_{2*,*} \tau_{I=0} E \\ &\rightarrow \mathrm{MGL}_{2*-1,*} \tau_{I \geq 1} E \rightarrow \cdots, \\ \cdots \rightarrow \mathrm{MGL}_{2*+1,*} \tau_{I \leq -1} E &\rightarrow \mathrm{MGL}_{2*,*} \tau_{I \geq 0} E \rightarrow \mathrm{MGL}_{2*,*} E \\ &\rightarrow \mathrm{MGL}_{2*,*} \tau_{I \leq -1} E \rightarrow \cdots. \end{aligned}$$

By [Proposition 3.6](#), in each long exact sequence, the first and the last term in the above part vanish. Therefore, we have equivalences

$$\mathrm{MGL}_{2*,*} \tau_{I=0} E \cong \mathrm{MGL}_{2*,*} \tau_{I \geq 0} E \cong \mathrm{MGL}_{2*,*} E. \quad \square$$

3.3. t -structure as a direct colimit. We now bring the homotopy t -structure into play.

Definition 3.8. Let $\mathcal{SH}(k)_{\geq 0}$ denote the subcategory of $\mathcal{SH}(k)$ generated under colimits and extensions by $\Sigma_+^\infty X \wedge \mathbb{G}_m^{\wedge n}$ for $X \in \mathrm{Sm}_k$ and $n \in \mathbb{Z}$. This defines the non-negative part of the homotopy t -structure.

Since we are working over fields, the subcategory $\mathcal{SH}(k)_{\geq 0}$ is characterised by a vanishing of homotopy sheaves [[Hoy15](#), Th. 2.3]:

$$(3.3.1) \quad \begin{aligned} \mathcal{SH}(k)_{\geq 0} &= \{E \in \mathcal{SH}(k) \mid \underline{\pi}_{i+n,n}(E) = 0 \text{ for } i < 0, n \in \mathbb{Z}\}, \\ \mathcal{SH}(k)_{\leq 0} &= \{E \in \mathcal{SH}(k) \mid \underline{\pi}_{i+n,n}(E) = 0 \text{ for } i > 0, n \in \mathbb{Z}\}. \end{aligned}$$

Recall that the notation $\mathcal{SH}(k)_{\geq d}$ denotes the subcategory $\Sigma^d \mathcal{SH}(k)_{\geq 0}$.

THEOREM-DEFINITION 3.9. *Let I^d be the category $\mathcal{SH}(k)_{\geq d} \cap I$. This is the non-negative part of a t -structure on $\mathcal{SH}(k)$.*

Proof. We need to show that I^d is closed under colimits and extensions and is presentable [Lur17, Prop. 1.4.4.11(1)]. This is true for $\mathcal{SH}(k)_{\geq d}$ and I by [Lur17, Prop. 1.4.4.11(2)]; the case of I^d follows since subcategories with these properties are closed under limits (and so, in particular, intersections; use that limits of presentable categories along left adjoint functors are presentable [Lur09, Prop. 5.5.3.13]). $\square$

Example 3.10. Let I be $\mathcal{SH}(k)_{c \geq 0}$. For a smooth proper variety X and a K -theory point ξ on X of rank $\geq d$, we have $\mathrm{Th}(\xi) \in I^d$. Indeed it suffices to check that $\mathrm{Th}(\xi) \in \mathcal{SH}(k)_{\geq d}$, which follows from [BH21, Lemma 13.1].

Example 3.11. Let $E \in I$ be compact. Then $E \in I^d$ for d sufficiently small. This follows from the facts that (1) the subcategory of bounded below spectra is thick, (2) the subcategory of compact spectra is the thick subcategory generated by spectra of the form $\Sigma^{n,n}\Sigma_+^\infty X$ for $n \in \mathbb{Z}$ and $X \in \mathrm{Sm}_k$ [Nee01, Lemma 4.4.5] [DI05, Th. 9.2] and (3) these generators are bounded below, by definition.

We use the notations $\tau_{\leq n}^d$, $\tau_{\geq n}^d$, $\tau_{=n}^d$ to denote the truncation functors associated to the t -structure with non-negative part I^d .

The following is a key technical result.

PROPOSITION 3.12. *Let $E \in I^d$. Then we have the following:*

- (1) $\tau_{\leq 0}^d E \in I^d$,
- (2) $\mathrm{MGL}_{2*,*}\tau_{\leq 0}^d E \cong \mathrm{MGL}_{2*,*}E$.

Suppose now that $E \in I^{d+1}$. Then we have additionally the following:

- (3) $\pi_{*,*}\tau_{\leq 0}^d E$ is concentrated in Chow degrees ≤ 0 , and
- (4) $\mathrm{MGL}_{*,*}\tau_{\leq 0}^d E$ is concentrated in Chow degree 0.

Proof. (1) This is immediate from the cofiber sequence $\tau_{\geq 1}^d E \rightarrow E \rightarrow \tau_{\leq 0}^d E$, since $E \in I^d$ and $\tau_{\geq 1}^d E \in \Sigma I^d \subset I^d$.

(2) Consider the long exact sequence

$$\begin{aligned} \cdots \rightarrow \mathrm{MGL}_{2*,*}\tau_{\geq 1}^d E \rightarrow \mathrm{MGL}_{2*,*}E \rightarrow \mathrm{MGL}_{2*,*}\tau_{\leq 0}^d E \\ \rightarrow \mathrm{MGL}_{2*-1,*}\tau_{\geq 1}^d E \rightarrow \cdots \end{aligned}$$

It suffices to show that $\mathrm{MGL}_{2*+i,*}\tau_{\geq 1}^d E = 0$ for $i = 0, -1$. Since $\tau_{\geq 1}^d E \in \Sigma I^d \subset \mathcal{SH}(k)_{c \geq 1}$, the result follows from Proposition 3.6.

(3) We need to show that $\pi_{2n+i,n}(\tau_{\leq 0}^d E) = 0$ for $i > 0$. This vanishing arises for two slightly different reasons, depending on if (a) $n + i > d$ or (b) $n + i \leq d$.

(a) When $n + i > d$, we have $S^{2n+i,n} = \Sigma^{n+i}\mathbb{G}_m^{\wedge n} \in \Sigma\mathcal{SH}(k)_{\geq d}$. Therefore we have $S^{2n+i,n} \in \Sigma I^d$. Since $\tau_{\leq 0}^d E$ is in the ≤ 0 part of the t -structure corresponding to I^d , the homotopy groups $\pi_{2n+i,i}(\tau_{\leq 0}^d E)$ vanishes by definition.

(b) When $n+i \leq d$, by [equation \(3.3.1\)](#), if $F \in \mathcal{SH}(k)_{\geq d+1}$, then $\pi_{2n+i,n}(F) = 0$. Therefore it suffices to show that $\tau_{\leq 0}^d E \in \mathcal{SH}(k)_{\geq d+1}$. Consider the cofiber sequence in (1). Since $\tau_{\geq 1}^d E \in \Sigma I^d \subset \mathcal{SH}(k)_{\geq d+1}$ and $E \in I^{d+1} \subset \mathcal{SH}(k)_{\geq d+1}$, the result follows.

(4) By (1) we have $\tau_{\leq 0}^d E \in \mathcal{SH}(k)_{c \geq 0}$. Hence $\mathrm{MGL}_{*,*}\tau_{\leq 0}^d E$ is concentrated in Chow degrees ≥ 0 , by [Proposition 3.6\(2\)](#). It remains to show that $\mathrm{MGL}_{*,*}\tau_{\leq 0}^d E$ is concentrated in Chow degrees ≤ 0 .

Let $\mathcal{C} \subset \mathcal{SH}(k)$ be the subcategory consisting all spectra M such that $\pi_{*,*}(M \wedge \tau_{\leq 0}^d E)$ is concentrated in Chow degrees ≤ 0 . It suffices to show that $\mathrm{MGL} \in \mathcal{C}$. Note that $\mathcal{C}$ is closed under filtered colimits, extensions, wedge sums, and $\Sigma^{2n,n}$ for any $n \in \mathbb{Z}$. By (3), $\mathbb{1} \in \mathcal{C}$, and hence (in the notation of [Section A](#)) $\mathcal{SH}(k)^{\mathrm{pure Tate} \geq 0} \subset \mathcal{C}$. The result thus follows from [Theorem A.1](#). $\square$

PROPOSITION 3.13. *Let $E \in \mathcal{SH}(k)$. There are directed systems*

$$\tau_{\geq 0}^n E \rightarrow \tau_{\geq 0}^{n-1} E \rightarrow \cdots \rightarrow \tau_{I \geq 0} E$$

and

$$\tau_{\leq 0}^n E \rightarrow \tau_{\leq 0}^{n-1} E \rightarrow \cdots \rightarrow \tau_{I \leq 0} E,$$

which are in fact colimit diagrams.

Proof. Since $I^n \subset I^{n-1} \subset \cdots \subset \mathcal{SH}(k)_{I \geq 0}$, the directed systems exist.

To see that the second one is a colimit diagram, it suffices to show that $\mathrm{colim}_n \tau_{\leq 0}^n E \in \mathcal{SH}(k)_{I \leq 0}$. (Indeed the fiber of $E \rightarrow \mathrm{colim}_n \tau_{\leq 0}^n E$ is $\mathrm{colim}_n \tau_{> 0}^n E \in \mathcal{SH}(k)_{I > 0}$, so we conclude by [\[BBD82, Prop. 1.3.3\(ii\)\]](#).) This follows from [Example 3.11](#), which shows that every generator of $\mathcal{SH}(k)_{I \geq 0} = I$ is in I^n for n sufficiently small.

Taking the fiber of the constant colimit diagram E mapping to the second one yields the first one (up to a shift), which is thus also a colimit diagram. This concludes the proof. $\square$

3.4. *MGL-completion and Adams–Novikov spectral sequence.* We come to the main result of this section.

Recall from [\[DI10, §7\]](#) or [\[HKO11\]](#) the usual MGL based motivic Adams–Novikov spectral sequence

$$\mathrm{Ext}_{\mathrm{MGL}_{*,*}\mathrm{MGL}}^{*,*,*}(\mathrm{MGL}_{*,*}, \mathrm{MGL}_{*,*}X) \implies \pi_{*,*}X_{\mathrm{MGL}}^{\wedge}.$$

Recall that $(\mathrm{MGL}_{2*,*}, \mathrm{MGL}_{2*,*}\mathrm{MGL})$ is a Hopf algebroid canonically isomorphic to $(\mathrm{MU}_{2*}, \mathrm{MU}_{2*}\mathrm{MU})$. (Combine [\[NSØ09, Cor. 6.7, Lemma 6.4\]](#) and [\[Spi20, Cor. 6.7\]](#); recall our convention that we implicitly invert e .) In particular, for

$E \in \mathcal{SH}(k)$, the graded abelian group $\mathrm{MGL}_{2*,*}E$ is canonically a comodule over $\mathrm{MU}_{2*}\mathrm{MU}$.

THEOREM 3.14. *Let $E \in \mathcal{SH}(k)_{I \geq 0}$. Then*

- (1) *the canonical map $\tau_{I \leq 0}E \rightarrow \tau_{I \leq 0}(E)_{\mathrm{MGL}}^{\wedge}$ to the MGL-nilpotent completion induces an isomorphism on $\pi_{*,*}$, and*
- (2) $\pi_{2w-s,w}\tau_{I \leq 0}E \cong \mathrm{Ext}_{\mathrm{MU}_*\mathrm{MU}}^{s,2w}(\mathrm{MU}_*, \mathrm{MGL}_{2*,*}E)$.

Proof. We first prove part (2) of the theorem. We have isomorphisms

$$\begin{aligned} \tau_{I \leq 0}E &\stackrel{(*)}{\simeq} \tau_{I \leq 0}\tau_{I \geq 0}E \stackrel{L.3.13}{\simeq} \tau_{I \leq 0} \operatorname{colim}_d \tau_{\geq 0}^d E \\ &\stackrel{C.2.8}{\simeq} \operatorname{colim}_d \tau_{I \leq 0} \tau_{\geq 0}^d E \stackrel{L.3.13}{\simeq} \operatorname{colim}_{n,d} \tau_{\leq 0}^n \tau_{\geq 0}^d E, \end{aligned}$$

where $(*)$ holds because $E \in \mathcal{SH}(k)_{I \geq 0}$. The third equivalence uses [Corollary 2.8](#), which is stated for the largest I . By a similar proof we can show it also works for other choices of I . Since the right-hand side of the isomorphism in part (2) is compatible with filtered colimits in E [[Hov04](#), Lemma 3.2.2(b)], we may replace E by $\tau_{\geq 0}^d E$ and so we may assume that $E \in I^d$. Let $n < d$. We shall show that $\pi_{2w-s,w}\tau_{\leq 0}^n E \simeq \mathrm{Ext}_{\mathrm{MU}_*\mathrm{MU}}^{s,2w}(\mathrm{MGL}_{2*,*}E)$. Taking the (constant) colimit as $n \rightarrow -\infty$ will yield the result.

By [Lemma 3.12](#)(2) and (4), we find that $\mathrm{MGL}_{2*+i,*}\tau_{\leq 0}^n E = \mathrm{MGL}_{2*,*}E$ for $i = 0$, and otherwise it vanishes. Hence the Adams–Novikov spectral sequence for $\tau_{\leq 0}^n E$ again collapses, and it suffices to show that this spectral sequence converges to $\pi_{*,*}\tau_{\leq 0}^n E$ or, in other words, that $\tau_{\leq 0}^n E$ is MGL-nilpotent complete (on homotopy groups). By [Lemma 3.12](#)(1), $\tau_{\leq 0}^n(E)$ is connective in the homotopy t -structure; hence by [[Man21](#), §5.1 and Th. 7.3.5] we have

$$\tau_{\leq 0}^n(E)_{\mathrm{MGL}}^{\wedge} \simeq L_{\mathrm{MGL}}\tau_{\leq 0}^n(E) \simeq \tau_{\leq 0}^n(E)_{\eta}^{\wedge}.$$

It is thus sufficient to show that $\tau_{\leq 0}^n(E) \rightarrow \tau_{\leq 0}^n(E)_{\eta}^{\wedge}$ induces an equivalence on $\pi_{*,*}$. This follows from [Lemma 3.12](#)(3) and [Example 3.3](#).

This concludes the proof of part (2).

For part (1), [Corollary 3.7](#) implies that the Adams–Novikov spectral sequence for $\tau_{I \leq 0}E$ collapses and

$$\pi_{2w-s,w}\tau_{I \leq 0}(E)_{\mathrm{MGL}}^{\wedge} \cong \mathrm{Ext}_{\mathrm{MU}_*\mathrm{MU}}^{s,2w}(\mathrm{MU}_*, \mathrm{MGL}_{2*,*}E).$$

Hence part (1) follows from part (2). $\square$

3.5. *First consequences.* [Theorem 3.14](#) is the central result of this paper. We establish here some of its immediate consequences, many of which will be amplified in [Section 4](#).

LEMMA 3.15. *Let $E \in \mathcal{SH}(k)$. If $\pi_{i,0}(E \wedge \mathrm{Th}(\xi)) = 0$ for all $i \in \mathbb{Z}$ and K -theory points ξ on smooth proper varieties X , then $E \simeq 0$.*

Proof. By duality, the assumption is equivalent to $[\Sigma^i \mathrm{Th}(\xi), E] = 0$ for all $i \in \mathbb{Z}$ and K -theory points ξ on smooth proper varieties X . Taking $\xi = n\mathcal{O}$ shows that $[\Sigma^{i,j} \Sigma_+^\infty X, E] = 0$ for all $i, j \in \mathbb{Z}$ and all X smooth and proper. This implies that $E = 0$ by [Remark 2.19](#). $\square$

Remark 3.16. Let $E \in \mathcal{SH}(k)^{c\heartsuit}$. By [Theorem 3.14](#), the map $\alpha : E \rightarrow E_{\mathrm{MGL}}^\wedge$ induces an isomorphism on $\pi_{*,*}$. Let X be smooth projective and $\xi \in K(X)$. By [Corollary 2.12](#) we have $E \wedge \mathrm{Th}(\xi) \in \mathcal{SH}(k)^{c\heartsuit}$, and hence $E \wedge \mathrm{Th}(\xi) \rightarrow (E \wedge \mathrm{Th}(\xi))_{\mathrm{MGL}}^\wedge$ again induces an equivalence on $\pi_{*,*}$. But tensoring with the strongly dualizable object $\mathrm{Th}(\xi)$ preserves limits, so $(E \wedge \mathrm{Th}(\xi))_{\mathrm{MGL}}^\wedge \simeq E_{\mathrm{MGL}}^\wedge \wedge \mathrm{Th}(\xi)$. [Lemma 3.15](#) now implies that α is an equivalence. This is a special case of [Proposition 4.3](#) in the next section.

We deduce the following strengthening of [Proposition 2.17](#).

COROLLARY 3.17. *Let $E \in \mathcal{SH}(k)$. Then E is Chow- ∞ -connective if and only if $E \wedge \mathrm{MGL} \simeq 0$.*

Proof. By right completeness ([Proposition 2.14](#)), $E \in \mathcal{SH}(k)_{c \geq n}$ if and only if $E_{c=i} = 0$ for $i < n$. (Indeed necessity is clear, and for sufficiency note that $E_{c < n} \in \cap_m \mathcal{SH}(k)_{c < m} = 0$.) Consequently, E is Chow- ∞ -connective if and only if $E_{c=i} = 0$ for all i .

Hence we find

$$\begin{aligned}
& E \text{ is Chow-}\infty\text{-connective} \\
& \iff E_{c=i} = 0 \text{ for all } i \\
& \xLeftrightarrow{L.3.15} \pi_{*,*}(E_{c=i} \wedge \mathrm{Th}(\xi)) = 0 \text{ for all } i, X \text{ smooth proper and } \xi \in K(X) \\
& \xLeftrightarrow{C.2.12} \pi_{*,*}(E \wedge \mathrm{Th}(\xi))_{c=i} = 0 \text{ for all } i \\
& \xLeftrightarrow{T.3.14+T.A.1} \mathrm{MGL}_{*,*}(E \wedge \mathrm{Th}(\xi))_{c=i} = 0 \text{ for all } i \\
& \xLeftrightarrow{C.3.7} \mathrm{MGL}_{2*+i,*}(E \wedge \mathrm{Th}(\xi)) = 0 \text{ for all } i \\
& \iff \pi_{*,*}(\mathrm{MGL} \wedge E \wedge \mathrm{Th}(\xi)) = 0 \\
& \xLeftrightarrow{L.3.15} \mathrm{MGL} \wedge E = 0
\end{aligned}$$

$\square$

4. Reconstruction theorems

We keep the conventions from the last section: $S = \mathrm{Spec}(k)$ is the spectrum of a field of exponential characteristic e , which is implicitly inverted throughout.

In this section we will amplify the results of the last section by deducing fairly explicit descriptions of the categories $\mathcal{SH}(k)^{c\heartsuit}$ and $\mathbb{1}_{c=0}\text{-Mod}$. We begin in [Section 4.1](#) by describing these categories in terms of a certain comonad C on $\mathrm{MGL}_{c=0}\text{-Mod}$. Then in [Section 4.2](#) we explain how work of Bondarko supplies

a very explicit description of the category $\mathrm{MGL}_{c=0}\text{-Mod}$, and we also identify the comonad C . Finally in [Section 4.3](#), we use this to provide an explicit description of $\mathcal{SH}(k)^{c\heartsuit}$.

4.1. Comonadic descent.

4.1.1. Suppose we have a presentably symmetric monoidal category $\mathcal{D}$ and $A \in \mathrm{CAlg}(\mathcal{D})$. We obtain a free-forgetful adjunction

$$F : \mathcal{D} \rightleftarrows A\text{-Mod} : U$$

and the endofunctor

$$C := FU = \otimes A : A\text{-Mod} \rightarrow A\text{-Mod},$$

which acquires the structure of a comonad [[Lur17](#), Prop. 4.7.3.3].

We denote by $C\text{-CoMod}$ the category of comodules under C (see [[Lur17](#), Def. 4.2.1.13]), and hence obtain a factorization [[Lur17](#), §4.7.4]

$$\mathcal{D} \rightleftarrows C\text{-CoMod} \rightleftarrows A\text{-Mod},$$

where $C\text{-CoMod} \rightarrow A\text{-Mod}$ is the forgetful functor that we denote by H (with right adjoint the cofree comodule functor) and $\mathcal{D} \rightarrow C\text{-CoMod}$ sends X to $X \otimes A$ with its canonical comodule structure.

4.1.2. *Cobar resolution.* We can form the cobar resolution [[MNN17](#), Constr. 2.7]

$$\mathbb{1}_{\mathcal{D}} \rightarrow \mathrm{CB}^{\bullet}(A) := \left(A \rightrightarrows A \otimes A \overset{\rightarrow}{\rightrightarrows} \dots \right);$$

this as a coaugmented cosimplicial object in $\mathrm{CAlg}(\mathcal{D})$. Taking module categories, we obtain a coaugmented cosimplicial category

$$(4.1.1) \quad \mathcal{D} \rightarrow \mathrm{CB}^{\bullet}(A)\text{-Mod} := \left(A\text{-Mod} \rightrightarrows (A \otimes A)\text{-Mod} \overset{\rightarrow}{\rightrightarrows} \dots \right).$$

It follows from [[Lur17](#), Th. 4.7.6.2] that

$$\lim_{\Delta} \mathrm{CB}^{\bullet}(A)\text{-Mod} \simeq C\text{-CoMod}.$$

Let Δ_s be the subcategory of Δ with the same objects, but where the morphisms are given by injective order preserving maps between nonempty linearly ordered sets. We will often be interested in the restriction of diagram (4.1.1) to the coinital subcategory $\Delta_s \hookrightarrow \Delta$ [[Lur09](#), Lemma 6.5.3.7]; i.e., view this as a coaugmented *semi*-cosimplicial object. The limit of any cosimplicial object is the same as the limit of its associated semi-cosimplicial object. Therefore we also have

$$\lim_{\Delta_s} \mathrm{CB}^{\bullet}(A)\text{-Mod} \simeq C\text{-CoMod}.$$

4.1.3. *Monoidal structure.* The category $C\text{-CoMod}$ is in fact symmetric monoidal, and the left adjoint functors above are symmetric monoidal. Indeed (4.1.1) is a diagram of symmetric monoidal categories and symmetric monoidal functors (since it comes from a diagram of commutative rings and commutative ring maps). Since limits of symmetric monoidal categories are computed on the underlying categories [Lur17, Cor. 3.2.2.5], it follows that the limit $C\text{-CoMod}$ is symmetric monoidal, as desired.

4.1.4. *t-structure.* Suppose that $\mathcal{D}$ carries a t -structure and $\otimes A : \mathcal{D} \rightarrow \mathcal{D}$ is t -exact. We can give each of the categories $A^{\otimes n}\text{-Mod}$ the t -structure detected by the forgetful functor to $\mathcal{D}$ (see, e.g., [Bac21, Lemma 29]).

When viewed as a semi-cosimplicial category, the maps in (4.1.1) become t -exact functors.

For $-\infty \leq m \leq n \leq +\infty$, we denote by $A^{\otimes p}\text{-Mod}_{[m,n]}$ the subcategory of objects bounded in the t -structure. Then $\text{CB}^\bullet(A)\text{-Mod}_{[m,n]}$ is a full subdiagram of (4.1.1). Since limits preserve fully faithful functors, we find that

$$C\text{-CoMod}_{[m,n]} := \lim_{\Delta_s} \text{CB}^\bullet(A)\text{-Mod}_{[m,n]}$$

is a full subcategory of $C\text{-CoMod}$; in fact, $C\text{-CoMod}_{[m,n]}$ is equivalent to $H^{-1}(A\text{-Mod}_{[m,n]})$, where $H : C\text{-CoMod} \rightarrow A\text{-Mod}$ is the forgetful functor.

Let $C_{[m,n]} : A\text{-Mod}_{[m,n]} \rightarrow A\text{-Mod}_{[m,n]}$ denote the restriction of C , and $C^\heartsuit = C_{[0,0]}$. We note that by construction,

$$(4.1.2) \quad C\text{-CoMod}_{[m,n]} \simeq C_{[m,n]}\text{-CoMod}.$$

One checks immediately that

$$C\text{-CoMod}_{\geq 0} := C\text{-CoMod}_{[0,\infty]} \quad \text{and} \quad C\text{-CoMod}_{\leq 0} := C\text{-CoMod}_{[-\infty,0]}$$

define a t -structure on $C\text{-CoMod}$.

4.1.5. *Compact generation.* We consider the t -structure on $C\text{-CoMod}$ defined in Section 4.1.4.

Suppose that $C\text{-CoMod}^\heartsuit$ is compactly generated. Let $C\text{-CoMod}^{\heartsuit\omega}$ denote the category of its compact objects.

We consider the category

$$\mathcal{H}\text{ov}(C) := \text{Ind}(\text{Thick}(C\text{-CoMod}^{\heartsuit\omega})),$$

where $\text{Thick}(C\text{-CoMod}^{\heartsuit\omega})$ denotes the thick subcategory of $C\text{-CoMod}$ generated by $C\text{-CoMod}^{\heartsuit\omega}$, and Ind denotes the category obtained by freely adding filtered colimits (see [Lur09, 5.3.5.1]). We obtain an adjunction [Lur09, Props. 5.3.5.10 and 5.3.5.13]

$$\mathcal{H}\text{ov}(C) \rightleftarrows C\text{-CoMod}.$$

Remark 4.1. The assumption that $C\text{-CoMod}^\heartsuit$ is compactly generated need not imply that $C\text{-CoMod}$ is compactly generated. Moreover if $E \in C\text{-CoMod}^\heartsuit$ is compact, it need not be the case that $E \in C\text{-CoMod}$ is compact.

Remark 4.2. If C is the comonad describing comodules over some Hopf algebroid Ψ , then under mild assumptions $\mathcal{H}ov(C)$ coincides with Hovey's category $\mathcal{H}ov(\Psi)$ [Hov04]. See [BH18, §3] for a treatment in the language of ∞ -categories. In this situation we will use the notations $\mathcal{H}ov(C)$ and $\mathcal{H}ov(\Psi)$ interchangeably.

4.1.6. *Modules over $\mathbb{1}_{c=0}$.* We apply the above discussion with the category $\mathcal{D} = \mathbb{1}_{c=0}\text{-Mod}$ and $A = \text{MGL} \wedge \mathbb{1}_{c=0}$ (which is equivalent to $\text{MGL}_{c=0}$ by Corollary 2.12). We obtain

$$\mathbb{1}_{c=0}\text{-Mod} \rightleftarrows C\text{-CoMod} \rightleftarrows \text{MGL}_{c=0}\text{-Mod}.$$

We consider the t -structure on $\mathbb{1}_{c=0}\text{-Mod}$ induced by the Chow t -structure (Remark 2.3).

PROPOSITION 4.3.

- (1) *The free functor $\bar{F} : \mathbb{1}_{c=0}\text{-Mod} \rightarrow C\text{-CoMod}$ is t -exact and symmetric monoidal.*
- (2) *For $-\infty \leq m \leq n < \infty$, the restriction $\mathbb{1}_{c=0}\text{-Mod}_{[m,n]} \rightarrow C\text{-CoMod}_{[m,n]}$ is an equivalence. In particular,*

$$\mathbb{1}_{c=0}\text{-Mod}^{c^\heartsuit} \simeq C\text{-CoMod}^{c^\heartsuit}.$$

- (3) *The functor $\bar{F}$ induces a symmetric monoidal equivalence $\mathbb{1}_{c=0}\text{-Mod} \simeq \mathcal{H}ov(C)$.*

Proof. (1) This is clear by construction.

(2) We shall apply the Barr-Beck-Lurie theorem [Lur17, Th. 4.7.3.5]. It hence suffices to show that (a) $F : \mathbb{1}_{c=0}\text{-Mod}_{[m,n]} \rightarrow \text{MGL}_{c=0}\text{-Mod}_{[m,n]}$ is conservative, and (b) F -split totalizations exist in $\mathbb{1}_{c=0}\text{-Mod}_{[m,n]}$ and are preserved by F .

(a) For $E \in \mathbb{1}_{c=0}\text{-Mod}$, we have $E \wedge_{\mathbb{1}_{c=0}} \text{MGL}_{c=0} \simeq E \wedge \text{MGL}$, and this is zero if and only if E has vanishing Chow homotopy objects (Corollary 3.17). Thus if E is additionally Chow bounded from the left, then $E \simeq 0$ by right completeness.

(b) Since our categories are presentable, totalizations (and in fact all small limits) exist [Lur09, Cor. 5.5.2.4]. The truncation functors $(-)_{c \geq m}$ commute with limits, and the subcategories of n -coconnective objects are preserved by limits. It follows that limits in $\mathbb{1}_{c=0}\text{-Mod}_{[m,n]}$ are computed by first computing in $\mathbb{1}_{c=0}\text{-Mod}$ and then applying $(-)_{c \geq m}$, and similarly for $\text{MGL}_{c=0}\text{-Mod}_{[m,n]}$. Being t -exact, F commutes with $(-)_{c \geq m}$, and hence it is

enough to show that $F : \mathbb{1}_{c=0}\text{-Mod}_{[-\infty, n]} \rightarrow \text{MGL}_{c=0}\text{-Mod}_{[-\infty, n]}$ preserves totalizations. Since the forgetful functors are conservative and preserves limits, it is enough to show that $\wedge \text{MGL}$ preserves totalizations in $\mathcal{SH}(k)_{c \leq n}$, i.e., $\text{MGL} \wedge \lim_{n \in \Delta} E^n \simeq \lim_{n \in \Delta} \text{MGL} \wedge E^n$. We are dealing with Chow-bounded above objects, so it suffices to show that the induced maps on Chow homotopy objects are isomorphisms (Proposition 2.14). This follows from t -exactness of smashing with MGL (Corollary 2.12) and [Lur17, Prop. 1.2.4.5(5)], i.e., the fact that totalizations of bounded above spectra behave like finite limits on homotopy objects. More precisely, for $E^\bullet \in \mathcal{SH}(k)_{c \leq n}$ and $i \in \mathbb{Z}$, there exists $N \gg 0$ such that

$$\begin{aligned} (\text{MGL} \wedge \lim_{n \in \Delta} E^n)_{c=i} &\simeq \text{MGL} \wedge (\lim_{n \in \Delta} E^n)_{c=i} \\ &\simeq \text{MGL} \wedge (\text{Tot}^N E^\bullet)_{c=i} \\ &\simeq (\text{MGL} \wedge \text{Tot}^N E^\bullet)_{c=i} \\ &\simeq (\text{Tot}^N (\text{MGL} \wedge E^\bullet))_{c=i} \simeq (\lim_{n \in \Delta} \text{MGL} \wedge E^n)_{c=i}; \end{aligned}$$

here Tot^N refers to the N -th partial totalization, i.e., the limit over $\Delta_{\leq N}$.

(3) Let $\mathcal{G} \subset \mathbb{1}_{c=0}\text{-Mod}$ the class of objects of the form $\text{Th}(\xi) \wedge \mathbb{1}_{c=0} \simeq \text{Th}(\xi)_{c=0}$ (see Corollary 2.12). They form a compact generating family, and $\mathcal{G} \subset \mathbb{1}_{c=0}\text{-Mod}^{c^\heartsuit}$. There is an induced symmetric monoidal functor

$$\text{Ind}(\text{Thick}(\mathcal{G})) \rightarrow \text{Ind}(\text{Thick}(\bar{F}\mathcal{G})),$$

which is an equivalence by (2). The left-hand side is $\mathbb{1}_{c=0}\text{-Mod}$. The right-hand side is $\mathcal{Hov}(C)$, by definition (and (2)). This is the desired result. $\square$

COROLLARY 4.4. *There are canonical symmetric monoidal equivalences*

$$\mathcal{SH}(k)^{c^\heartsuit} \simeq \mathbb{1}_{c=0}\text{-Mod}^{c^\heartsuit} \simeq C^{c^\heartsuit}\text{-CoMod},$$

where $C^{c^\heartsuit}$ is the comonad on $\text{MGL}_{c=0}\text{-Mod}^{c^\heartsuit}$ obtained by restricting C .

Proof. The functor $\mathcal{SH}(k) \rightarrow \mathbb{1}_{c=0}\text{-Mod}$ induces an equivalence of the hearts, e.g., by [Bac21, Lemma 29]. The second equivalence follows from Proposition 4.3(2) via equation (4.1.2). $\square$

4.2. Modules over $\text{MGL}_{c=0}$.

4.2.1. *Pure MGL-motives.* For a smooth proper variety X and $i \in \mathbb{Z}$, denote by $X\{i\} \in \text{MGL}_{c=0}\text{-Mod}$ the object $(\Sigma^{2i,i} X_+ \wedge \text{MGL})_{c=0} \simeq \Sigma^{2i,i} (X_+)_{c=0} \wedge \text{MGL}$. By the Thom isomorphism, these are generators of $\text{MGL}_{c=0}\text{-Mod}$ (as a localizing subcategory). Write $\text{PM}_{\text{MGL}}(k) \subset \text{MGL}_{c=0}\text{-Mod}$ for the smallest idempotent complete additive subcategory containing the objects $X\{i\}$. For now we view this as a spectrally enriched category. By duality and adjunction,

we have

$$\begin{aligned} \mathrm{Map}_{\mathrm{PM}_{\mathrm{MGL}}(k)}(X\{i\}, Y\{j\}) &\simeq \mathrm{Map}_{\mathrm{PM}_{\mathrm{MGL}}(k)}(\mathrm{MGL}_{c=0}, (X \times Y)\{j - d_X - i\}) \\ &\simeq \mathrm{Map}_{\mathcal{SH}(k)}(S^{2(i+d_X-j), (i+d_X-j)}, ((X \times Y)_+)_{c=0} \wedge \mathrm{MGL}). \end{aligned}$$

[Proposition 3.6](#) thus implies that

$$\pi_* \mathrm{Map}_{\mathrm{PM}_{\mathrm{MGL}}(k)}(X\{i\}, Y\{j\}) \simeq \mathrm{MGL}_{*+2(i+d_X-j), (i+d_X-j)}(((X \times Y)_+)_{c=0})$$

is concentrated in degree 0. In other words, our spectrally enriched category $\mathrm{PM}_{\mathrm{MGL}}(k)$ is just an additive ordinary 1-category. The above computation together with [Corollary 3.7](#) shows that

$$\begin{aligned} (4.2.1) \quad \mathrm{Hom}_{\mathrm{PM}_{\mathrm{MGL}}(k)}(X\{i\}, Y\{j\}) &\simeq \mathrm{MGL}_{2(i+d_X-j), (i+d_X-j)}(X \times Y) \\ &\simeq \mathrm{MGL}^{2(d_Y+j-i), (d_Y+j-i)}(X \times Y). \end{aligned}$$

Remark 4.5. For future use, we point out the following generalization. If $B \in \mathcal{SH}(k)$ is any oriented ring spectrum, we can form a category PM_B of *pure B -motives*. It is the idempotent complete additive 1-category generated by objects $X\{i\}_B$ and morphisms

$$\mathrm{Hom}_{\mathrm{PM}_B}(X\{i\}_B, Y\{j\}_B) = [\Sigma^{2i, i} \Sigma_+^\infty X \wedge B, \Sigma^{2j, j} \Sigma_+^\infty Y \wedge B]_{B\text{-Mod}}.$$

This construction enjoys the following properties:

- (1) If M is in $B\text{-Mod}$, then the functor $M_*(X) := [\Sigma^{2*, *}\Sigma_+^\infty X \wedge B, M]_{B\text{-Mod}}$ defines a linear presheaf on PM_B .
- (2) If $u : A \rightarrow B$ is a morphism of oriented ring spectra and F is any A -module, then $B \otimes_A F$ is a B -module and we have a canonical map $u_F : F \rightarrow B \otimes_A F$. Given $\alpha : X \rightarrow Y \in \mathrm{PM}_A$ and $s \in F_*$, we have $u_{F*}(\alpha^* s) = u_*(\alpha)^*(u_{F*} s)$.

Remark 4.6. Given an oriented cohomology theory A^* on smooth proper k -varieties, one can define a 1-category of pure A -motives [[NZ06](#), §6]. It is generated as an idempotent complete additive 1-category by objects $X\{i\}$ with sets of maps

$$[X\{i\}, Y\{j\}] = A^{j+d_Y-i}(X \times Y)$$

and composition given by convolution. Taking $A^* = B^{2*, *}$ for some oriented ring spectrum B , we recover the category PM_B . (In particular, taking $A^* = \mathrm{MGL}^{2*, *}$ we recover $\mathrm{PM}_{\mathrm{MGL}}(k)$.) Given the above description, the only part of this assertion that we have not proved yet is that composition in PM_A is given by convolution; this is a purely formal consequence of the rigidity of the generators.

4.2.2. *Spectral Morita theory.* By [Remark 2.19](#) and the Thom isomorphism, $\mathrm{PM}_{\mathrm{MGL}}(k)$ compactly generates $\mathrm{MGL}_{c=0}\text{-Mod}$. Since we have an explicit description of $\mathrm{PM}_{\mathrm{MGL}}(k)$ as a spectrally enriched category, we should be able to recover $\mathrm{MGL}_{c=0}\text{-Mod}$ by a variant of Morita theory, such as [\[SS03\]](#). For an ∞ -category $\mathcal{D}$ with finite coproducts, we use the notation

$$\begin{aligned}\mathcal{P}_{\Sigma}(\mathcal{D}) &= \mathrm{Fun}^{\times}(\mathcal{D}^{\mathrm{op}}, \mathcal{S}\mathrm{pc}), & \mathcal{P}_{\mathcal{SH}}(\mathcal{D}) &= \mathrm{Fun}^{\times}(\mathcal{D}^{\mathrm{op}}, \mathcal{SH}), \\ \mathcal{P}_{\mathrm{Ab}}(\mathcal{D}) &= \mathrm{Fun}^{\times}(\mathcal{D}^{\mathrm{op}}, \mathrm{Ab}),\end{aligned}$$

where $\mathrm{Fun}^{\times}$ denotes the category of product-preserving functors. Provided that $\mathcal{D}$ is additive, there are equivalences [\[Lur18, Rem. C.1.5.9\]](#)

$$\mathcal{P}_{\mathcal{SH}}(\mathcal{D})_{\geq 0} \simeq \mathcal{P}_{\Sigma}(\mathcal{D}) \quad \text{and} \quad \mathcal{P}_{\mathcal{SH}}(\mathcal{D})^{\heartsuit} \simeq \mathcal{P}_{\mathrm{Ab}}(\mathcal{D}).$$

Note that $\mathcal{P}_{\mathcal{SH}}(\mathcal{D})$ has a natural (pointwise) t -structure.

LEMMA 4.7. *Let $\mathcal{D}$ be a small semi-additive ∞ -category. The full subcategory $\mathcal{P}_{\mathcal{SH}}(\mathcal{D})_{\geq 0}$ consisting of functors $F : \mathcal{D}^{\mathrm{op}} \rightarrow \mathcal{SH}_{\geq 0} \subset \mathcal{SH}$ is generated under colimits and extension by the image of the canonical functor $\mathcal{D} \rightarrow \mathcal{P}_{\mathcal{SH}}(\mathcal{D})$. In particular, $\mathcal{P}_{\mathcal{SH}}(\mathcal{D})_{\geq 0}$ is the non-negative part of a t -structure. Its non-positive part consists of the functors $\mathcal{D}^{\mathrm{op}} \rightarrow \mathcal{SH}_{\leq 0} \subset \mathcal{SH}$.*

Proof. For $d \in \mathcal{D}$, write $R_d \in \mathcal{P}_{\mathcal{SH}}(\mathcal{D})$ for the “representable functor.” We have

$$(*) \quad \mathrm{map}(R_d, R_e) \simeq \mathrm{Map}(d, e)^{gp} \in \mathcal{SH}_{\geq 0},$$

where the superscript gp denotes group completion of the additive $\mathcal{E}_{\infty}$ -monoid structure. Consider the t -structure on $\mathcal{P}_{\mathcal{SH}}(\mathcal{D})$ generated by the objects R_d for $d \in \mathcal{D}$. We have an adjunction $\mathcal{SH} \rightleftarrows \mathcal{P}_{\mathcal{SH}}(\mathcal{D}) : ev_d$. The left adjoint is right- t -exact, since it sends the generator $1 \in \mathcal{SH}$ to R_d , whence the right adjoint ev_d is left- t -exact. By $(*)$ it is also right- t -exact. Thus the functors ev_d for $d \in \mathcal{D}$ form a conservative t -exact collection, whence the non-negative and non-positive parts of the t -structure are as claimed. $\square$

PROPOSITION 4.8. *We have a canonical t -exact, symmetric monoidal equivalence*

$$\mathrm{MGL}_{c=0}\text{-Mod} \simeq \mathcal{P}_{\mathcal{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k)).$$

Proof. This is standard. To be more precise, the symmetric monoidal category $\mathcal{P}_{\mathcal{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k))$ can be obtained as the stabilization of a localization of $\mathcal{P}(\mathrm{PM}_{\mathrm{MGL}}(k))$ (see, e.g., [\[Aok20, Rem. 2.10\]](#)). The universal properties of presheaves, day convolution, localization and stabilization thus imply that there exists a unique cocontinuous symmetric monoidal functor

$$\mathcal{P}_{\mathcal{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k)) \rightarrow \mathrm{MGL}_{c=0}\text{-Mod}$$

extending the inclusion $\mathrm{PM}_{\mathrm{MGL}}(k) \rightarrow \mathrm{MGL}_{c=0}\text{-Mod}$. Since it induces equivalences on mapping spectra between compact generators, it is an equivalence. By

[Lemma 4.7](#), this equivalence identifies the non-negative parts of the t -structures, and hence is t -exact. $\square$

Given $E \in \mathrm{MGL}_{c=0}\text{-Mod}$, denote by $\pi_i^c(E) \in \mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))$ the presheaf with $\pi_i^c(E)(X) = \pi_i \mathrm{Map}_{\mathrm{MGL}_{c=0}\text{-Mod}}(X, E)$, $\forall X \in \mathrm{PM}_{\mathrm{MGL}}(k)$.

COROLLARY 4.9.

- (1) The functor $\pi_0^c : \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit} \rightarrow \mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))$ is an equivalence.
- (2) For $E \in \mathrm{MGL}_{c=0}\text{-Mod}$, we have $E \in \mathrm{MGL}_{c=0}\text{-Mod}_{c \geq 0}$ (respectively $E \in \mathrm{MGL}_{c=0}\text{-Mod}_{c \leq 0}$) if and only if $\pi_i^c(E) = 0$ for all $i < 0$ (respectively $i > 0$).
- (3) The Chow t -structure on $\mathrm{MGL}_{c=0}\text{-Mod}$ is non-degenerate [BBD82, p. 32].

Proof. By [Proposition 4.8](#), all statements translate into assertions about $\mathcal{P}_{\mathrm{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k))$, which are easily verified. $\square$

Remark 4.10. An object $F \in \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit} \simeq \mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))$ consists of the following data:

- for every smooth proper variety X , we have a graded abelian group $F(X)_* = \pi_0^c(F)(X\{\ast\})$;
- for every graded MGL-correspondence $\alpha : X \rightarrow Y$ (i.e., $\alpha \in \mathrm{MGL}^{2*,*}(X \times Y)$), we have a homomorphism $\alpha^* : F(Y)_* \rightarrow F(X)_*$

subject to the conditions that

- for composable MGL-correspondences α, β , we have $\alpha^* \beta^* = (\beta \alpha)^*$,
- $\mathrm{id}^* = \mathrm{id}$ and $0^* = 0$, as well as
- for parallel MGL-correspondences α, β , we have $\alpha^* + \beta^* = (\alpha + \beta)^*$.

For example, since $\mathrm{MGL}_{2*,*} \cong \mathrm{MU}_{2*,*}$, each $F(X)_*$ is an $\mathrm{MU}_{2*,*}$ -module, and all the α^* are automatically $\mathrm{MU}_{2*,*}$ -module maps.

Remark 4.11. The category $\mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))$ has enough projective objects, namely the representable presheaves. When viewed as objects of the category $\mathcal{P}_{\mathrm{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k))$, they have mapping spectra concentrated in degree 0 (see the proof of [Lemma 4.7](#)). Using [Lur17, Prop. 1.3.3.7] (or spectra Morita theory) this implies $\mathcal{P}_{\mathrm{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k)) \simeq D(\mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k)))$.

To summarize, we have

$$\begin{array}{ccc} \mathrm{PM}_{\mathrm{MGL}}(k) \hookrightarrow \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit} & \xrightarrow{\simeq} & \mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k)) \\ \downarrow & & \downarrow \\ \mathrm{MGL}_{c=0}\text{-Mod} & \xrightarrow[\simeq]{} \mathcal{P}_{\mathrm{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k)) & \xleftarrow[\simeq]{} D(\mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))) \end{array}$$

where the downwards arrows are the inclusions of the hearts.

4.2.3. *Identification of the monad.* Denote by

$$\mathrm{Fun}_0^L(\mathrm{MGL}_{c=0}\text{-Mod}, \mathrm{MGL}_{c=0}\text{-Mod}) \subset \mathrm{Fun}(\mathrm{MGL}_{c=0}\text{-Mod}, \mathrm{MGL}_{c=0}\text{-Mod})$$

the subcategory of those functors F that preserve colimits and such that

$$F(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}) \subset \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}.$$

LEMMA 4.12. *The restriction*

$$\begin{aligned} \mathrm{Fun}_0^L(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}, \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}) \\ \rightarrow \mathrm{Fun}^L(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}, \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}) \end{aligned}$$

is an equivalence.

Proof. By [Sos19, Lemma 3.2] we have

$$\mathrm{Fun}^L(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}, \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}) \simeq \mathrm{Fun}^\oplus(\mathrm{PM}_{\mathrm{MGL}}(k), \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}),$$

where $\mathrm{Fun}^\oplus$ denotes the category of biproduct preserving functors. Therefore, we deduce that

$$\mathrm{Fun}_0^L(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}, \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}) \simeq \mathrm{Fun}^\oplus(\mathrm{PM}_{\mathrm{MGL}}(k), \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}).$$

This latter category identifies with $\mathrm{Fun}^L(\mathcal{P}_\Sigma(\mathrm{PM}_{\mathrm{MGL}}(k)), \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit})$ by [Lur09, Prop. 5.5.8.15] and [BH21, Lemma 2.8]. Finally we have

$$\begin{aligned} \mathrm{Fun}^L(\mathcal{P}_\Sigma(\mathrm{PM}_{\mathrm{MGL}}(k)), \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}) \\ \simeq \mathrm{Fun}^L(\mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k)), \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}), \end{aligned}$$

since $\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}$ is a 1-category and $\mathrm{PM}_{\mathrm{MGL}}(k)$ is additive. This concludes the proof. $\square$

COROLLARY 4.13. *Restriction induces an equivalence*

$$\begin{aligned} \{ \text{cocontinuous comonads on } \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od} \text{ preserving the heart} \} \\ \simeq \{ \text{cocontinuous comonads on } \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit} \}. \end{aligned}$$

Proof. Since comonads on $\mathcal{D}$ are by definition given by $\mathrm{Alg}(\mathrm{Fun}(\mathcal{D}, \mathcal{D})^{\mathrm{op}})$, this follows from the fact that the restriction equivalence of Lemma 4.12 is compatible with the composition monoidal structures. $\square$

Under the above equivalence, the comonad C corresponds to its restriction to the heart, which we denote by $C^{c\heartsuit}$ or also by C when it is clear in the context. We describe this restriction.

By equation (4.2.1), $[X\{i\}, Y\{j\}] \simeq [X\{0\}, Y\{j-i\}]$. Therefore we can view $[X\{*\}, Y\{*\}]$ as a single graded group by taking the first $*$ to be 0. Observe that for smooth proper varieties X, Y , the mapping set, $[X\{0\}, Y\{*\}]_{\mathrm{PM}_{\mathrm{MGL}}(k)}$, is an $\mathrm{MU}_{2*}\mathrm{MU}$ -comodule; indeed we have seen that up to some shift in degrees

it identifies with $\mathrm{MGL}^{2*,*}(X \times Y)$. In other words, for any graded MGL-correspondence $\alpha : X \rightarrow Y \in [X\{0\}, Y\{*\}]_{\mathrm{PM}_{\mathrm{MGL}}(k)}$, we obtain

$$\begin{aligned} \Delta(\alpha) &= \sum_i p_i \otimes \alpha_i \\ &\in \mathrm{MU}_{2*}\mathrm{MU} \otimes_{\mathrm{MU}_{2*}} \mathrm{MGL}_{2*,*}(X \times Y) \\ &\simeq \mathrm{MU}_{2*}\mathrm{MU} \otimes_{\mathrm{MU}_{2*}} [X\{0\}, Y\{*\}]_{\mathrm{PM}_{\mathrm{MGL}}(k)}. \end{aligned}$$

Recall our description of $\mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit}$ in [Remark 4.10](#). Let $F \in \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit}$; thus F is a kind of presheaf on smooth proper varieties together with some extra data, namely an action by MGL-correspondences. We wish to describe $CF \in \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit}$; again this is a presheaf with an action by MGL-correspondences.

Remark 4.14. We have $CF = \mathrm{MGL}_{c=0} \wedge_{\mathbb{1}_{c=0}} F$, which has *two* structures as an $\mathrm{MGL}_{c=0}$ -module. Since the underlying spectra are the same, the right module structure has the same value on sections as the left module structure when viewed as an object of $\mathcal{P}_{\mathcal{SH}}(\mathrm{PM}_{\mathrm{MGL}}(k))$, however the action by graded MGL-correspondences differs. The “correct” action is on the left, and given a correspondence $\alpha : X \rightarrow Y$, we denote it by $\alpha_L^* : CF(Y) \rightarrow CF(X)$.

PROPOSITION 4.15.

- (1) Given $F \in \mathrm{MGL}_{c=0}\text{-Mod}^{c\heartsuit} \simeq \mathcal{P}_{\mathrm{Ab}}(\mathrm{PM}_{\mathrm{MGL}}(k))$, the object CF is given on sections by

$$(CF)(X)_* = \mathrm{MU}_{2*}\mathrm{MU} \otimes_{\mathrm{MU}_{2*}} F(X)_*.$$

Given an MGL-correspondence $\alpha : X \rightarrow Y$, the action $\alpha_L^* : CF(Y)_* \rightarrow CF(X)_*$ is given by $\Delta(\alpha)^*$. In other words, for $s \in F(Y)$ and $p \in \mathrm{MU}_{2*}\mathrm{MU}$, we have

$$\alpha_L^*(p \otimes s) = \sum_i p p_i \otimes \alpha_i^*(s)$$

in the notation for $\Delta(\alpha)$ of above.

- (2) The counit map $CF \rightarrow F$ is given on sections by $p \otimes s \mapsto \epsilon(p)s$, where ϵ is the counit of the Hopf algebroid $(\mathrm{MU}_{2*}, \mathrm{MU}_{2*}\mathrm{MU})$.
 (3) The comultiplication map $CF \rightarrow C^2F$ is given on sections by $p \otimes s \mapsto \Delta(p) \otimes s$.

Proof. (1) Given $F \in \mathrm{MGL}_{c=0}\text{-Mod}$, we have

$$CF = \mathrm{MGL}_{c=0} \wedge_{\mathbb{1}_{c=0}} F \simeq (\mathrm{MGL}_{c=0} \wedge_{\mathbb{1}_{c=0}} \mathrm{MGL}_{c=0}) \wedge_{\mathrm{MGL}_{c=0}} F.$$

Since also $\mathrm{MGL}_{c=0} \simeq \mathrm{MGL} \wedge \mathbb{1}_{c=0}$ ([Corollary 2.12](#)) we get

$$\mathrm{MGL}_{c=0} \wedge_{\mathbb{1}_{c=0}} \mathrm{MGL}_{c=0} \simeq (\mathrm{MGL} \wedge \mathrm{MGL}) \wedge \mathbb{1}_{c=0} \simeq \mathrm{MGL}_{c=0}[a_1, a_2, \dots],$$

where $\mathrm{MU}_{2*}\mathrm{MU} = \mathrm{MU}_*[a_1, a_2, \dots]$ (use [\[NSØ09, Lemma 6.4\]](#)). The description of the sections of CF follows.

To obtain the description of α_L^* , we apply [Remark 4.5](#) with $A = \text{MGL}_{c=0}$, $M = CF$, $B = \text{MGL}_{c=0} \wedge_{\mathbb{1}_{c=0}} \text{MGL}_{c=0}$ and u the map inserting a unit on the right. Part (1) of the remark implies that α_L^* is MU_{2*}MU -linear. Thus to determine $\alpha_L^*(p \otimes s)$, we may assume that $p = 1$. We find that $B \wedge_A F \simeq CF$, $u_{F*}(s) = 1 \otimes s$ and $u_*(\alpha) = \Delta(\alpha)$. Part (2) of the remark thus asserts the claimed formula $\alpha_L^*(1 \otimes s) = \Delta(\alpha)^*(1 \otimes s)$.

(2) The counit map is the module action

$$CF = \text{MGL} \wedge F \rightarrow F,$$

which under $\text{MGL} \wedge F \simeq (\text{MGL} \wedge \text{MGL}) \wedge_{\text{MGL}} F$ corresponds to the multiplication map $\text{MGL} \wedge \text{MGL} \rightarrow \text{MGL}$ followed by the old module action on F . The result follows since the multiplication in MGL corresponds to the counit in the associated Hopf algebroid.

(3) By definition, the comultiplication $\text{MGL} \wedge \text{MGL} \rightarrow \text{MGL} \wedge \text{MGL} \wedge \text{MGL}$ is by inserting a unit in the middle. The proof is similar to (2). $\square$

Example 4.16. In the special case when $\alpha \in \text{MU}_{2*} = [\mathbb{1}\{0\}, \mathbb{1}\{*\}]_{\text{PM}_{\text{MGL}}(k)}$, we get $\Delta(\alpha) = \eta_L(\alpha) \otimes \text{id}$, where η_L is the left unit of the Hopf algebroid $(\text{MU}_*, \text{MU}_*\text{MU})$. Therefore the MU_{2*} -module structure on $(CF)(X)_*$ is indeed the left MU_{2*} -module structure on $\text{MU}_{2*}\text{MU} \otimes_{\text{MU}_{2*}} F(X)_*$.

4.3. *The Chow heart.* Using [Corollary 4.4](#) and [Proposition 4.15](#), we can now describe the category $\mathcal{SH}(k)^{c^\heartsuit}$ explicitly, by spelling out what modules under the comonad $C^{c^\heartsuit}$ mean in terms of [Remark 4.10](#). Namely, we have $\mathcal{SH}(k)^{c^\heartsuit} \simeq C^{c^\heartsuit}\text{-CoMod}$, where we view $C^{c^\heartsuit}$ as a comonad on $\mathcal{P}_{\text{Ab}}(\text{PM}_{\text{MGL}}(k))$. An object $F \in C^{c^\heartsuit}\text{-CoMod}$ is given by the following data:

- for every smooth proper variety X , we have a graded MU_{2*}MU -comodule $F(X)_* := [\Sigma^{2*,*}X_+, F \wedge \text{MGL}]$, and
- for every graded MGL -correspondence $\alpha : X \rightarrow Y$, we have an MU_{2*} -linear map $\alpha^* : F(Y)_* \rightarrow F(X)_*$

subject to the following conditions:

- $(\alpha\beta)^* = \beta^*\alpha^*$, $(\alpha + \beta)^* = \alpha^* + \beta^*$, $\text{id}^* = \text{id}$, $0^* = 0$, and
- for every graded MGL -correspondence $\alpha : X \rightarrow Y$, the following diagram commutes:

$$\begin{array}{ccc} F(Y)_* & \xrightarrow{\alpha^*} & F(X)_* \\ \Delta_{F,Y} \downarrow & & \Delta_{F,X} \downarrow \\ \text{MU}_{2*}\text{MU} \otimes_{\text{MU}_{2*}} F(Y)_* & \xrightarrow{\Delta(\alpha)^*} & \text{MU}_{2*}\text{MU} \otimes_{\text{MU}_{2*}} F(X)_* \end{array}$$

A morphism from F to G consists of an MU_{2*}MU -comodule map $F(X)_* \rightarrow G(X)_*$ for every smooth proper variety X , compatible with pullback along graded MGL -correspondences in the evident way.

We thank an anonymous referee for pointing out the following.

Remark 4.17. As we have observed above, the mapping sets in $\mathrm{PM}_{\mathrm{MGL}}(k)$ are naturally $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules. In fact, this makes $\mathrm{PM}_{\mathrm{MGL}}(k)$ into a category *enriched* in $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules. The compatibility condition displayed above precisely means that $\mathcal{SH}(k)^{c\heartsuit}[1/e]$ is equivalent to the category of *enriched* presheaves on $\mathrm{PM}_{\mathrm{MGL}}(k)$ (see, e.g., [Rie14, §3.5]).

We can also prove the following.

PROPOSITION 4.18. *There is a canonical equivalence*

$$\mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}[1/e]_b \simeq D^b(\mathcal{SH}(k)^{c\heartsuit}).$$

Proof. We implicitly invert e throughout. We know by Remark 4.11 that $\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od} \simeq D(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit})$. Consider the comonadic adjunction

$$F : \mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}_b \rightleftarrows \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}_b \simeq D^b(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}) : R.$$

The functor F is t -exact and conservative by Proposition 4.3, and R is t -exact since $\mathrm{MGL}_{c=0} \in \mathcal{SH}(k)_{c \geq 0}$. The category $\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}$ has enough injectives, being a presheaf category. Since F is a conservative exact left adjoint, the category $\mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}$ also has enough injectives, namely those of the form RI for $I \in \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}$ injective. To conclude, by the dual of [Lur17, Prop. 1.3.3.7], it suffices to prove that if $X \in \mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit}$, then $[X, \Sigma^i RI]_{\mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}_b} = 0$ for $i > 0$. But by adjunction, this is the same as $[FX, \Sigma^i I]_{D^b(\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}^{c\heartsuit})}$, which vanishes since F is t -exact and I is injective. $\square$

4.4. W -cellular objects.

Definition 4.19. Let W be a set of smooth proper schemes over k that contains $\mathrm{Spec}(k)$ and is closed under finite products. Define the W -cellular category, denoted by $\mathcal{SH}(k)^{\mathrm{wcell}}$, to be the subcategory of $\mathcal{SH}(k)$ generated under taking colimits and desuspensions by objects of the form $\mathrm{Th}(\xi)$ for $\xi \in K(X)$ and $X \in W$.

Remark 4.20. When W contains only $\mathrm{Spec}(k)$, this specializes to the cellular subcategory in Section 2.7. Since this is the smallest choice, all cellular objects are W -cellular for any choice of W .

Fix W . In the discussion below, cellular means W -cellular.

4.4.1. *The W -cellular Chow t -structure.* Given a stable presentable category $\mathcal{D}$ under $\mathcal{SH}(k)$, we denote by $\mathcal{D}^{\mathrm{wcell}}$ the subcategory generated under colimits and desuspensions by the images of $\mathcal{SH}(k)^{\mathrm{wcell}}$. Similarly, we write $\mathcal{D}_{\geq 0}^{\mathrm{wcell}}$ for the subcategory generated under colimits and extensions by images of the objects of the form $\mathrm{Th}(\xi)$ for $\xi \in K(X)$ and $X \in W$. This defines a

t -structure on $\mathcal{D}^{\text{wcell}}$, whose truncation we denote by $\tau_{c \leq 0}^{\text{wcell}}, \tau_{c \geq 0}^{\text{wcell}}, \tau_{c=0}^{\text{wcell}}$. We call it cellular Chow t -structure. The inclusion $\mathcal{D}^{\text{wcell}} \hookrightarrow \mathcal{D}$ admits a right adjoint, called the cellularization functor $\mathcal{D} \rightarrow \mathcal{D}^{\text{wcell}}$. In the following discussion, we take $\mathcal{D}$ to be $\mathcal{SH}(k)$ or $\mathcal{SH}(k) \xrightarrow{\wedge^A} A\text{-Mod}$ for some $A \in \text{CAlg}(\mathcal{SH}(k))$.

We have the following results in the cellular case, analogous to [Lemma 3.15](#).

LEMMA 4.21. *Let $E \in \mathcal{SH}(k)^{\text{wcell}}$. If $\pi_{i,0}(E \wedge \text{Th}(\xi)) = 0$ for all $i \in \mathbb{Z}$ and K -theory points ξ on smooth proper varieties $X \in W$, then $E \simeq 0$.*

Proof. The proof is similar to that of [Lemma 3.15](#). $\square$

Let $E \mapsto E^{\text{wcell}}$ denote the cellularization of $E \in \mathcal{SH}(k)$. A key fact is as follows.

LEMMA 4.22. *The cellularization functor $\mathcal{SH}(k) \rightarrow \mathcal{SH}(k)^{\text{wcell}}$ is t -exact for the Chow t -structures. Equivalently, for $E \in \mathcal{SH}(k)_{c \geq 0}$, we have $E^{\text{wcell}} \in \mathcal{SH}(k)_{c \geq 0}^{\text{wcell}}$.*

Proof. The equivalence follows from the fact that $\mathcal{SH}(k)_{c \geq 0}^{\text{wcell}} \subset \mathcal{SH}(k)_{c \geq 0}$, whence cellularization is always left- t -exact, being right adjoint to a right- t -exact functor.

Now let $E \in \mathcal{SH}(k)_{c \geq 0}$. To prove that $E^{\text{wcell}} \in \mathcal{SH}(k)_{c \geq 0}^{\text{wcell}}$ it is enough to show that $\tau_{c < 0}^{\text{wcell}} E^{\text{wcell}} \simeq 0$. Similar to the proof of [Proposition 2.14](#), the cellular Chow t -structure is right complete. Therefore we only need to show that $\tau_{c=i}^{\text{wcell}} E^{\text{wcell}} \simeq 0$ for all $i < 0$.

By [Lemma 4.21](#), it suffice to show $\pi_{s,0}(\tau_{c=i}^{\text{wcell}} E^{\text{wcell}} \wedge \text{Th}(\xi)) = 0$ for all $s \in \mathbb{Z}$ and K -theory points ξ on smooth proper varieties $X \in W$. [Theorem 3.14](#) applied to $I = \mathcal{SH}(k)_{c \geq 0}^{\text{wcell}}$ shows that

$$\pi_{*,*}(\tau_{c=i}^{\text{wcell}} E^{\text{wcell}} \wedge \text{Th}(\xi)) \simeq \text{Ext}_{\text{MU}_{2*}\text{MU}}^{*,*}(\text{MU}_{2*}, \text{MGL}_{2*+i,*} E^{\text{wcell}} \wedge \text{Th}(\xi)).$$

Here we also use the cellular version of [Proposition 2.11](#) that gives $(\tau_{c=i}^{\text{wcell}} E^{\text{wcell}}) \wedge \text{Th}(\xi) \simeq \tau_{c=i}^{\text{wcell}}(E^{\text{wcell}} \wedge \text{Th}(\xi))$. Since MGL is cellular, we have

$$\text{MGL}_{2*+i,*} E^{\text{wcell}} \wedge \text{Th}(\xi) \simeq \text{MGL}_{2*+i,*} E \wedge \text{Th}(\xi),$$

which vanishes by [Proposition 3.6\(2\)](#). Therefore when $i \leq 0$, the bigraded homotopy groups of $\tau_{c=i}^{\text{wcell}} E^{\text{wcell}} \wedge \text{Th}(\xi)$ all vanish. The result follows. $\square$

LEMMA 4.23. *Let $A \in \text{CAlg}(\mathcal{SH}(k)_{c \geq 0})$, and consider the following commutative diagram of left adjoint functors:*

$$\begin{array}{ccccc} \mathcal{SH}(k) & \xrightarrow{\wedge^A \text{wcell}} & A^{\text{wcell}}\text{-Mod} & \xrightarrow{\wedge_{A^{\text{wcell}}} A} & A\text{-Mod} \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{SH}(k)^{\text{wcell}} & \xrightarrow{\wedge^A \text{wcell}} & A^{\text{wcell}}\text{-Mod}^{\text{wcell}} & \xrightarrow[a]{\wedge_{A^{\text{wcell}}} A} & A\text{-Mod}^{\text{wcell}} \end{array}$$

All of the right adjoints are t -exact, all the horizontal right adjoints are conservative, and the bottom right horizontal functor a is an equivalence.

Proof. If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor of presentable stable ∞ -categories such that $\mathcal{D}$ is generated under colimits by the essential image of F , and F admits a right adjoint G , then G is conservative. Indeed given $\alpha : X \rightarrow Y \in \mathcal{D}$ such that $G\alpha$ is an equivalence, the class of objects $T \in \mathcal{D}$ such that $\text{Map}(T, \alpha)$ is an equivalence is closed under colimits and contains the essential image of F , hence is all of $\mathcal{D}$, and so α is an equivalence. Conservativity of the horizontal right adjoints follows from this observation, taking F to be a horizontal left adjoint. This also means that a is an equivalence if and only if it is fully faithful, which we may test on the compact generators $\Sigma^{p,q} A^{\text{wcell}} \wedge \text{Th}(\xi)$ for K -theory points ξ on smooth proper varieties $X \in W$. In other words we need to show that $\pi_{*,*}(A^{\text{wcell}} \wedge \text{Th}(\xi)) \simeq \pi_{*,*}(A \wedge \text{Th}(\xi))$. This holds by definition (and dualizability of $\text{Th}(\xi)$).

We prove the t -exactness of the right adjoints. Since all the left adjoints are right t -exact, the right adjoints are left t -exact. Therefore it suffices to show that all the right adjoints are right t -exact.

First we prove the statement for the top horizontal arrows. We denote the functors by notation as in the following diagram:

$$F : \mathcal{SH}(k) \xrightleftharpoons[U']{F'} A^{\text{wcell}}\text{-Mod} \xrightleftharpoons[R]{L} A\text{-Mod} : U,$$

where F and U are the composites. By assumption, $A \in \mathcal{SH}(k)_{c \geq 0}$. Thus for any $E \in \mathcal{SH}(k)_{c \geq 0}$, $UFE \simeq A \wedge E \in \mathcal{SH}(k)_{c \geq 0}$. Since the non-negative part of $A\text{-Mod}$ is generated by $F(\mathcal{SH}(k)_{c \geq 0})$ and U preserves colimits (its left adjoint preserving compact generators), U is right- t -exact. By [Lemma 4.22](#),

$$A^{\text{wcell}} \in \mathcal{SH}(k)_{c \geq 0}^{\text{wcell}} \subset \mathcal{SH}(k)_{c \geq 0}.$$

We may thus apply the same argument to $A^{\text{wcell}}\text{-Mod}$ and deduce that the U' is t -exact. Now we show the right t -exactness of R . Given $E \in A\text{-Mod}_{\geq 0}$, by t -exactness of U and U' , we have $0 \simeq U(\tau_{<0}E) \simeq (U'R(E))_{<0} \simeq U'\tau_{<0}R(E)$. Using conservativity of U' , we conclude that $\tau_{<0}R(E) \simeq 0$ and equivalently $R(E) \in A^{\text{wcell}}\text{-Mod}_{\geq 0}$.

The same argument proves the t -exactness statements for the bottom left arrow (and the bottom right arrows are t -exact, being equivalences).

It remains to check that the vertical right adjoints (cellularizations) are t -exact. This is true for the left-hand vertical right adjoint by [Lemma 4.22](#). Exactness of the other two vertical right adjoints follows formally, since a functor is t -exact if and only if its composite with a t -exact conservative functor is. $\square$

4.4.2. *W-cellular reconstruction.*

COROLLARY 4.24. *Let $-\infty \leq m \leq n \leq \infty$ and $A \in \mathbf{CAlg}(\mathcal{SH}(k)_{c \geq 0})$. The composite functor*

$$A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} \hookrightarrow A\text{-}\mathcal{M}\mathrm{od} \xrightarrow{\tau_{[m,n]}} A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}$$

is fully faithful.

Proof. Since cellularization preserves the $\geq m$ part by Lemma 4.23, we have an adjunction

$$A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]}^{\mathrm{wcell}} \rightleftarrows A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]} : (-)^{\mathrm{wcell}}.$$

Consider the following diagram of adjunction pairs:

$$\begin{array}{ccc} A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]}^{\mathrm{wcell}} & \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{(-)^{\mathrm{wcell}}} \end{array} & A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]} \\ \tau_{\leq n}^{\mathrm{wcell}} \downarrow \uparrow i & & \tau_{\leq n} \downarrow \uparrow \\ A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} & & A\text{-}\mathcal{M}\mathrm{od}_{[m,n]} \end{array}$$

We denote by $L : A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]}^{\mathrm{wcell}} \rightarrow A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}$ the composite of the left adjoints and R its right adjoint. Note that the vertical adjunctions are in fact localizations, namely annihilating the $> n$ parts. It follows that there is an induced adjunction of the localizations

$$\tau_{\leq n} : A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} \rightleftarrows A\text{-}\mathcal{M}\mathrm{od}_{[m,n]} : (-)^{\mathrm{wcell}},$$

where left adjoint is the composite $L \circ i$, and thus it is equivalent to the functor we are required to show is fully faithful. It thus suffices to show that $RL(E) \simeq E$ for any $E \in A\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} \subset A\text{-}\mathcal{M}\mathrm{od}_{[m,\infty]}^{\mathrm{wcell}}$. Since cellularization is t -exact by Lemma 4.23, we have

$$RL(E) \simeq (\tau_{\leq n} E)^{\mathrm{wcell}} \simeq \tau_{\leq n}^{\mathrm{wcell}}(E^{\mathrm{wcell}}) \simeq \tau_{\leq n}^{\mathrm{wcell}}(E) \simeq E.$$

This concludes the proof. $\square$

COROLLARY 4.25. *Let $-\infty \leq m \leq n < \infty$.*

(1) *The following diagram is a pullback square:*

$$\begin{array}{ccc} \mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} & \longrightarrow & \mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]} \\ \downarrow & & \downarrow \\ \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}} & \longrightarrow & \mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]}. \end{array}$$

(2) *The comonad C on $\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}$ restricts to a comonad $C_{[m,n]}^{\mathrm{wcell}}$ on $\mathrm{MGL}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}}$ and*

$$C_{[m,n]}^{\mathrm{wcell}}\text{-CoMod} \simeq \mathbb{1}_{c=0}\text{-}\mathcal{M}\mathrm{od}_{[m,n]}^{\mathrm{wcell}}.$$

Proof. (1) In other words, given $E \in \mathbb{L}_{c=0}\text{-Mod}_{[m,n]}$, E is in (the essential image of the fully faithful inclusion of) $\mathbb{L}_{c=0}\text{-Mod}_{[m,n]}^{\text{wcell}}$ if and only if $E \wedge \text{MGL}$ is in (the essential image of) $\text{MGL}_{c=0}\text{-Mod}_{[m,n]}^{\text{wcell}}$. Since necessity is clear, we show sufficiency. Since $\wedge \text{MGL}$ is conservative on $\mathbb{L}_{c=0}\text{-Mod}_{[m,n]}$ as we proved in Proposition 4.3, it is enough to show that $E^{\text{wcell}} \wedge \text{MGL} \simeq E \wedge \text{MGL}$. Since MGL is cellular, we have $E^{\text{wcell}} \wedge \text{MGL} \simeq (E \wedge \text{MGL})^{\text{wcell}}$, and $E \wedge \text{MGL}$ is cellular in $\text{MGL}_{c=0}\text{-Mod}_{[m,n]}$ by assumption, so the result follows.

(2) C restricts as claimed since $\text{MGL}_{c=0} \wedge \text{MGL} \in \text{MGL}_{c=0}\text{-Mod}^{\text{wcell}, c\heartsuit}$. Now by construction, under the equivalence $C_{[m,n]}\text{-CoMod} \simeq \mathbb{L}_{c=0}\text{-Mod}_{[m,n]}$, $C_{[m,n]}^{\text{wcell}}\text{-CoMod}$ is equivalent to the full subcategory of $\mathbb{L}_{c=0}\text{-Mod}_{[m,n]}$ spanned by those objects E such that $E \wedge \text{MGL}$ is cellular in $\text{MGL}_{c=0}\text{-Mod}_{[m,n]}$; this is $\mathbb{L}_{c=0}\text{-Mod}_{[m,n]}^{\text{wcell}}$ by (1). $\square$

Arguing as in the proof of Proposition 4.8, we have the following cellular analogue.

PROPOSITION 4.26. *We have a canonical t -exact, symmetric monoidal equivalence*

$$\text{MGL}_{c=0}\text{-Mod}^{\text{wcell}} \simeq \mathcal{P}_{\mathcal{SH}}(\text{PM}_{\text{MGL}}^{\text{wcell}}(k)),$$

where $\text{PM}_{\text{MGL}}^{\text{wcell}}(k) \subset \text{PM}_{\text{MGL}}(k)$ denote the full subcategory spanned by $X\{i\}$ for $X \in W$ and $i \in \mathbb{Z}$.

Proof. The proof is similar to that of Proposition 4.8 by replacing the whole categories with the cellular categories. $\square$

4.4.3. *Field extensions.* We discuss the reconstruction theorems in the cellular cases for different choices of W .

First, set W to be $\{\text{Spec}(l) \mid l/k \text{ is a finite separable extension}\}$. Let $G = \text{Gal}(k)$ be the absolute Galois group. Recall that the stable category of genuine G -spectra $\mathcal{SH}(BG)$ [BH21, Exam. 9.12] admits a t -structure with heart the category of G -Mackey functors [BH21, Prop. 9.11]

$$\mathcal{P}_{\text{Ab}}(\text{Span}(\text{Fin}_G)) =: \text{Mack}_G.$$

Here Fin_G denotes the category of finite discrete G -sets. The canonical symmetric monoidal cocontinuous functor $\mathcal{SH} \rightarrow \mathcal{SH}(BG)$ induces the “constant Mackey functor”

$$\text{Ab} \rightarrow \text{Mack}_G, A \mapsto \underline{A}.$$

This is symmetric monoidal and so preserves rings, Hopf algebroids, etc. In particular, from the usual Hopf algebroid $(\text{MU}_{2*}, \text{MU}_{2*}\text{MU})$ we obtain the constant Hopf algebroid in (graded) Mackey functors $(\underline{\text{MU}}_{2*}, \underline{\text{MU}}_{2*}\text{MU})$.

COROLLARY 4.27. *Set W to be $\{\mathrm{Spec}(l) \mid l/k \text{ is a finite separable extension}\}$, and let $G = \mathrm{Gal}(k)$ be the absolute Galois group. We have*

$$\begin{aligned}\mathcal{SH}(k)^{\mathrm{wcell}, c^\heartsuit} &\simeq \underline{\mathrm{MU}}_{2*} \mathrm{MU}\text{-}\mathrm{CoMod}, \\ \mathbb{1}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}} &\simeq \mathcal{H}\mathrm{ov}(\underline{\mathrm{MU}}_{2*} \mathrm{MU}), \\ \mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}, c^\heartsuit} &\simeq \underline{\mathrm{MU}}_{2*}\text{-}\mathrm{Mod}.\end{aligned}$$

Here $\text{-}\mathrm{CoMod}$, $\mathcal{H}\mathrm{ov}$ and $\text{-}\mathrm{Mod}$ are performed relative to Mack_G .

Proof. Recall the cocontinuous symmetric monoidal functor $c : \mathcal{SH}(BG) \rightarrow \mathcal{SH}(k)$ [BH21, Prop. 10.6]. For $X \in \mathrm{Fin}_G$, it satisfies $c(\Sigma_+^\infty X) \simeq \Sigma_+^\infty cX$ for some cX finite étale over k , and hence $c(\mathcal{SH}(BG)_{\geq 0}) \subset \mathcal{SH}(k)_{c \geq 0}$. There is thus an induced functor

$$c : \mathrm{Mack}_G \simeq \mathcal{SH}(BG)^\heartsuit \rightarrow \mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}, c^\heartsuit}.$$

Let $\mathrm{Mack}_G^\mathbb{Z}$ denote the category of $\mathbb{Z}$ -graded Mackey functors. The invertible object $\Sigma^{2,1} \mathrm{MGL}_{c=0} \in \mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}, c^\heartsuit}$ determines a symmetric monoidal functor $c^\mathbb{Z} : \mathrm{Mack}_G^\mathbb{Z} \rightarrow \mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}, c^\heartsuit}$, which admits a right adjoint $c_\mathbb{Z}^*$.

We claim $c_\mathbb{Z}^*(\mathrm{MGL}_{c=0}) \simeq \underline{\mathrm{MU}}_{2*}$ and $\mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}^{\mathrm{wcell}, c^\heartsuit} \simeq \underline{\mathrm{MU}}_{2*}\text{-}\mathrm{Mod}$.

Since both sides are of the form $\mathcal{P}_{\mathrm{Ab}}$ and the functor is surjective on strongly dualizable generators, the second claim follows from the first. Let X be the finite G -set G/H for some subgroup $H \subset G$. Using Section 4.2, we compute that

$$\begin{aligned}c_\mathbb{Z}^*(\mathrm{MGL}_{c=0})(X)_i &\simeq \mathrm{Map}_{\mathrm{MGL}_{c=0}\text{-}\mathrm{Mod}}(c(X)(i), \mathrm{MGL}_{c=0}) \\ &\simeq \mathrm{MGL}_{2i, i}(\mathrm{Spec}(l^H)) \simeq \mathrm{MU}_{2i}.\end{aligned}$$

This is indeed the formula expected for the sections of a constant Mackey functor. In other words, $c_\mathbb{Z}^*(\mathrm{MGL}_{c=0})(-)_*$ and $\underline{\mathrm{MU}}_{2*} \in \mathrm{Mack}_G = \mathcal{P}_{\mathrm{Ab}}(\mathrm{Span}(\mathrm{Fin}_G))$ have canonically identical sections. What we need to check is that given $\alpha : X \rightarrow Y \in \mathrm{Span}(\mathrm{Fin}_G)$, the pullback maps α^* of the two presheaves agree. Any map in $\mathrm{Span}(\mathrm{Fin}_G)$ is a composite of a “usual” map $f : X \rightarrow Y \in \mathrm{Fin}_G$ and a “reverse” map $g^t : Y \rightarrow Z \in \mathrm{Span}(\mathrm{Fin}_G)$ corresponding to $g : Z \rightarrow Y \in \mathrm{Fin}_G$. It suffices to separately compare pullbacks along f (restriction maps) and pullbacks along g^t (transfers). Our formula for $c_\mathbb{Z}^*(\mathrm{MGL}_{c=0})(X)_*$ is actually functorial in $X \in \mathrm{Fin}_G$, i.e., the restriction maps are as expected. It remains to compare the transfers. We employ the following trick. The category Fin_G can be made into a site, where the coverings are those maps of finite G -sets that are surjective (ignoring the G -actions); this is often called the *étale topology*. Denote by $\mathcal{Shv}_{\mathrm{Ab}}(\mathrm{Span}(\mathrm{Fin}_G)) \subset \mathcal{P}_{\mathrm{Ab}}(\mathrm{Span}(\mathrm{Fin}_G))$ the full subcategory of those Mackey functors whose underlying presheaves on Fin_G are étale sheaves, and by $\mathcal{Shv}_{\mathrm{Ab}}(\mathrm{Fin}_G) \subset \mathcal{P}_{\mathrm{Ab}}(\mathrm{Fin}_G)$ the category of étale sheaves. Note that the underlying presheaf $\underline{\mathrm{MU}}_{2*} \in \mathcal{P}_{\mathrm{Ab}}(\mathrm{Fin}_G)$ is a sheaf in

the étale topology. By [BH21, Prop. C.11, Cor. C.13], the forgetful functor $\mathrm{Shv}_{\mathrm{Ab}}(\mathrm{Span}(\mathrm{Fin}_G)) \rightarrow \mathrm{Shv}_{\mathrm{Ab}}(\mathrm{Fin}_G)$ is an equivalence. In other words, the transfers are uniquely determined by the restriction maps.

We have thus established the third equivalence. The other two follow formally from Propositions 4.25 and 4.15, i.e., the identification of $\mathcal{SH}(k)^{\mathrm{wcell}, c\heartsuit}$ with $C^{\mathrm{wcell}}\text{-CoMod}$, and the explicit description of C , translated into the category $\mathrm{MGL}_{c=0}\text{-Mod}^{\mathrm{wcell}, c\heartsuit} \simeq \underline{\mathrm{MU}}_{2*}\text{-Mod}$ (and similarly for $\mathbb{1}_{c=0}\text{-Mod}^{\mathrm{wcell}}$). $\square$

We have similar results for another two choices of W :

COROLLARY 4.28. *Let l/k be a finite Galois extension with Galois group G . Let W be $\{\mathrm{Spec}(l') \mid l' \text{ is a subextension of } l/k\}$. We have*

$$\begin{aligned} \mathcal{SH}(k)^{\mathrm{wcell}, c\heartsuit} &\simeq \underline{\mathrm{MU}}_{2*}\mathrm{MU}\text{-CoMod}, \\ \mathbb{1}_{c=0}\text{-Mod}^{\mathrm{wcell}} &\simeq \mathcal{H}\mathrm{ov}(\underline{\mathrm{MU}}_{2*}\mathrm{MU}), \\ \mathrm{MGL}_{c=0}\text{-Mod}^{\mathrm{wcell}, c\heartsuit} &\simeq \underline{\mathrm{MU}}_{2*}\text{-Mod}. \end{aligned}$$

Again, everything is relative to Mack_G , but this time for the finite group G .

Proof. Similarly, by [BH21, Prop. 10.6], we have a cocontinuous symmetric monoidal functor $c_{l/k} : \mathcal{SH}(BG) \rightarrow \mathcal{SH}(k)$. The results follow from the same arguments as in the proof of Corollary 4.27. $\square$

As mentioned in Remark 4.20, we can also obtain results for the ordinary cellular category by taking W to be $\{\mathrm{Spec}(k)\}$.

COROLLARY 4.29. *We have canonical, symmetric monoidal, t -exact equivalences*

$$\begin{aligned} \mathcal{SH}(k)^{\mathrm{cell}, c\heartsuit} &\simeq \mathrm{MU}_{2*}\mathrm{MU}\text{-CoMod}, \\ \mathbb{1}_{c=0}\text{-Mod}^{\mathrm{cell}} &\simeq \mathcal{H}\mathrm{ov}(\mathrm{MU}_{2*}\mathrm{MU}), \\ \mathrm{MGL}_{c=0}\text{-Mod}^{\mathrm{cell}, c\heartsuit} &\simeq \mathrm{MU}_{2*}\text{-Mod}. \end{aligned}$$

Proof. It follows from Corollary 4.28 by taking the trivial extension. $\square$

5. Towards computations of motivic stable stems

In this section, we discuss a strategy for computing differentials in the motivic Adams spectral sequence based on $\mathrm{H}\mathbb{Z}/p$ for a general motivic spectrum X and, in particular, consequentially computing motivic stable stems. This strategy has worked very well for $\mathbb{1}$ over $\mathbb{C}$ in the work [GWX21], [IWX20a], [IWX20b]. Since the main structural theorems in this paper work for any field k , whose exponential characteristic is different from the prime p , we can now generalize the strategy over $\mathbb{C}$ to the field k . We will first discuss the general strategy over k and then take a closer look at the cases $k = \mathbb{C}, \mathbb{R}$, and finite fields.

5.1. *Algebraicity of the motivic Adams spectral sequence in the Chow heart.* The crucial point of the strategy for computing Adams differentials over $\mathbb{C}$ is the following theorem [GWX21, Th. 1.3], describing the algebraicity of the motivic Adams spectral sequence of $\mathbb{1}_p^\wedge/\tau$.

THEOREM 5.1 (Gheorghe–Wang–Xu). *For each prime p , there is an isomorphism of spectral sequences, between the HZ/p -based motivic Adams spectral sequence for $\mathbb{1}_p^\wedge/\tau$, and the algebraic Novikov spectral sequence for BP_{2*} .*

Since the algebraic Novikov spectral sequence for BP_{2*} can be completely computed by a computer program [IWX20b] within any reasonable range, [Theorem 5.1](#) provides a source of nontrivial differentials in the motivic Adams spectral sequence for $\mathbb{1}_p^\wedge/\tau$. We can then pull back these differentials along the map of spectral sequences induced by the unit map $\mathbb{1}_p^\wedge \rightarrow \mathbb{1}_p^\wedge/\tau$:

$$\mathbf{motASS}(\mathbb{1}_p^\wedge) \longrightarrow \mathbf{motASS}(\mathbb{1}_p^\wedge/\tau),$$

and push forward these differentials along the map of spectral sequences induced by the quotient map $\mathbb{1}_p^\wedge/\tau \rightarrow \Sigma^{1,-1}\mathbb{1}_p^\wedge$:

$$\mathbf{motASS}(\mathbb{1}_p^\wedge/\tau) \longrightarrow \mathbf{motASS}(\Sigma^{1,-1}\mathbb{1}_p^\wedge).$$

As a consequence, we obtain differentials in the motivic Adams spectral sequence for $\mathbb{1}_p^\wedge$.

Over an arbitrary field k , the substitute for $\mathbb{1}_p^\wedge/\tau$ over $\mathbb{C}$ consists of the objects $\mathbb{1}_{c=n}$ for $n \geq 0$. They all live in a suspension of the Chow heart. [Theorem 5.4](#) below is a generalization of [Theorem 5.1](#).

In general, let $F \in \mathcal{SH}(k)^{c\heartsuit}$. We can attempt to compute $\pi_{*,*}F$ by running the HZ/p -based motivic Adams spectral sequence. This is the trigraded spectral sequence obtained from the graded cosimplicial spectrum

$$\mathrm{map}(S^{0,*}, F \wedge \mathrm{CB}^\bullet(\mathrm{HZ}/p)) \simeq \mathrm{map}(S^{0,*}, F \wedge_{\mathbb{1}_{c=0}} \mathrm{CB}_{\mathbb{1}_{c=0}}^\bullet(\mathbb{1}_{c=0} \wedge \mathrm{HZ}/p)).$$

The Hopkins–Morel isomorphism (with the exponential characteristic e implicitly inverted) [[Hoy15](#)]

$$\mathrm{HZ} \simeq \mathrm{MGL}/(a_1, a_2, \dots)$$

implies that

$$\mathbb{1}_{c=0} \wedge \mathrm{HZ}/p \simeq \mathbb{1}_{c=0} \wedge \mathrm{MGL}/(p, a_1, a_2, \dots)$$

is cellular in $\mathbb{1}_{c=0}\text{-Mod}$, where $p \neq e$. Under the equivalence (see [Corollary 4.29](#))

$$\mathbb{1}_{c=0}\text{-Mod}^{\mathrm{cell}} \simeq \mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU}),$$

$\mathbb{1}_{c=0} \wedge \mathrm{HZ}/p$ thus corresponds to $\mathrm{MU}_{2*}\mathrm{MU}/(p, a_1, a_2, \dots)$.

We give a more general definition of the algebraic Novikov spectral sequence in the category of $\mathcal{Hov}(\mathrm{MU}_{2*}\mathrm{MU})$.

Definition 5.2. Let X be an object in $\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})$, and let H be a commutative monoid in $\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})$. Define the *algebraic Novikov spectral sequence* based on H for X to be the spectral sequence associated to the cosimplicial object $X \wedge \text{CB}^\bullet(H)$ in $\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})$, of the form

$$\begin{aligned} E_1^{s,i,t} &= \pi_s \text{map}_{\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})}(\Sigma^t \text{MU}_{2*}, X \wedge \text{CB}^i(H)) \\ &\implies \pi_{s-i} \text{map}_{\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})}(\Sigma^t \text{MU}_{2*}, X), \end{aligned}$$

with differentials $d_r : E_r^{s,i,t} \rightarrow E_r^{s+r-1,i+r,t}$. Here $\Sigma^t \text{MU}_{2*}$ denote MU_{2*} with the internal degree of every element shifted positively by t . A similar definition of the algebraic Novikov spectral sequence applies to the category of $\mathcal{H}\text{ov}(\text{BP}_{2*}\text{BP})$.

THEOREM 5.3. *Let $F \in \mathcal{SH}(k)^{\text{cell}, c^\heartsuit}$. Write M for the MU_{2*}MU -comodule associated to F under the equivalence in [Corollary 4.29](#), and write H for the comodule $\text{MU}_{2*}\text{MU}/(p, a_1, a_2, \dots)$. The trigraded motivic Adams spectral sequence for F based on $\mathbb{H}\mathbb{Z}/p$ is equivalent (with all higher and multiplicative structure) to the algebraic Novikov spectral sequence based on H for M .*

Proof. We have canonical equivalences of graded cosimplicial spectra:

$$\begin{aligned} &\text{map}(\mathbb{1}, F \wedge \text{CB}^\bullet(\mathbb{H}\mathbb{Z}/p)) \\ &\simeq \text{map}(\mathbb{1}, F \wedge_{\mathbb{1}_{c=0}} \text{CB}^\bullet_{\mathbb{1}_{c=0}}(\mathbb{1}_{c=0} \wedge \mathbb{H}\mathbb{Z}/p)) \\ &\simeq \text{map}_{\mathbb{1}_{c=0}\text{-Mod}}(\mathbb{1}_{c=0}, F \wedge_{\mathbb{1}_{c=0}} \text{CB}^\bullet_{\mathbb{1}_{c=0}}(\mathbb{1}_{c=0} \wedge \mathbb{H}\mathbb{Z}/p)) \\ &\simeq \text{map}_{\mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})}(\text{MU}_{2*}, M \wedge \text{CB}^\bullet(H)); \end{aligned}$$

here the first equivalence follows from the fact that F is a module over $\mathbb{1}_{c=0}$, the second is by adjunction, and the third is by [Corollary 4.29](#). $\square$

More generally we have the following.

THEOREM 5.4. *Let $F \in \mathcal{SH}(k)^{c^\heartsuit}$. Write M for the MU_{2*}MU -comodule associated to F^{cell} under the equivalence in [Corollary 4.29](#), and write H for the comodule $\text{MU}_{2*}\text{MU}/(p, a_1, a_2, \dots)$. The same equivalence of spectral sequences holds.*

Proof. Indeed for general $F \in \mathcal{SH}(k)^{c^\heartsuit}$, its $\mathbb{H}\mathbb{Z}/p$ -based motivic Adams spectral sequence coincides with that for F^{cell} . $\square$

The category of p -local MU_{2*}MU -comodules and the category of BP_{2*}BP -comodules are equivalent as abelian categories (cf. [Mor85, Prop. 1.2.3]). The comodule $\text{MU}_{2*}\text{MU}/(p, a_1, a_2, \dots)$ is p -local, and it corresponds to

$$\text{BP}_{2*} \otimes_{\text{MU}_{2*}} \text{MU}_{2*}\text{MU}/(p, a_1, a_2, \dots) \cong \text{BP}_{2*}\text{BP}/I$$

as a $\mathrm{BP}_2\mathrm{BP}$ -comodule, where $I = (p, v_1, v_2, \dots)$ is the augmentation ideal of BP_2 . Therefore, the algebraic Novikov spectral sequence of an $\mathrm{MU}_2\mathrm{MU}$ -comodule X based on $H = \mathrm{MU}_2\mathrm{MU}/(p, a_1, a_2, \dots)$, is equivalent to the algebraic Novikov spectral sequence of its p -localization $X_{(p)}$ based on H , and is equivalent to the algebraic Novikov spectral sequence of its corresponding $\mathrm{BP}_2\mathrm{BP}$ -comodule based on $\mathrm{BP}_2\mathrm{BP}/I$.

COROLLARY 5.5. *Let $F \in \mathcal{SH}(k)^{c\vee}$. Then the trigraded motivic Adams spectral sequence for F based on HZ/p is equivalent (with all higher and multiplicative structure) to the trigraded algebraic Novikov spectral sequence for $\mathrm{BPGL}_{2*,*}F$ based on $\mathrm{BP}_2\mathrm{BP}/I$.*

Proof. Immediate from [Theorem 5.4](#) and the above discussion. $\square$

The original definition of the algebraic Novikov spectral sequence ([\[Mil75\]](#), [\[Nov67\]](#)) is defined at each prime p in the category of $\mathrm{BP}_2\mathrm{BP}$ -comodules, and only for BP_2 . It converges to the E_2 -page of the Adams-Novikov spectral sequence $\mathrm{Ext}_{\mathrm{BP}_2\mathrm{BP}}^{*,*}(\mathrm{BP}_2, \mathrm{BP}_2)$ for the (p -completed) sphere spectrum. It is defined by filtering the cobar complex using the powers of the augmentation ideal $I = (p, v_1, v_2, \dots)$.

In the case of BP_2 , our [Definition 5.2](#) of the algebraic Novikov spectral sequence based on $\mathrm{BP}_2\mathrm{BP}/I$ is equivalent to the original one. This is proved in [\[GWX21, §9\]](#).

A natural question is the following.

Question 5.6. For a $\mathrm{BP}_2\mathrm{BP}$ -comodule Y , we have the following two definitions of the algebraic Novikov spectral sequence:

- the one using the filtration of the powers of the augmentation ideal I , and
- the one in [Definition 5.2](#) based on $\mathrm{BP}_2\mathrm{BP}/I$.

When are they equivalent?

We give a sufficient condition for such an equivalence in [Proposition 5.8](#).

We first recall the following definition in homological algebra (see, e.g., [\[Seg01\]](#)).

Definition 5.7. A BP_2 -module Y has *Levin index* 1 if the following maps that are induced by $I^n \rightarrow I^{n-1}$ are zero for all $n \geq 1$:

$$\mathrm{Ext}_{\mathrm{BP}_2}^{*,*}(I^{n-1}Y, \mathbb{F}_p) \rightarrow \mathrm{Ext}_{\mathrm{BP}_2}^{*,*}(I^nY, \mathbb{F}_p),$$

where $I = (p, v_1, v_2, \dots)$.

It is proved in [\[Seg01\]](#) and [\[GWX21, Lemma 9.7\]](#) that BP_2 itself has Levin index 1.

PROPOSITION 5.8. *Let Y be a $\mathrm{BP}_2\mathrm{BP}$ -comodule whose underlying BP_2 -module has Levin index 1. Then the two definitions of the algebraic Novikov spectral sequence for Y in Question 5.6 are equivalent.*

Proof. The proof is essentially identical to the proof of Theorem 8.3 of [GWX21], where the technical condition that BP_2 has Levin index 1 ([GWX21, Lemma 9.7]) is replaced by the general condition that Y has Levin index 1. $\square$

Remark 5.9. In general, without the condition that Y has Levin index 1, the two definitions of the algebraic Novikov spectral sequences are not equivalent. In fact, for $F \in \mathcal{SH}(k)^{c\heartsuit}$, suppose that $\mathrm{BPGL}_{2*,*}F$ has Levin index at least 2. Then the algebraic Novikov spectral sequence for $\mathrm{BPGL}_{2*,*}F$ defined by powers of I is isomorphic to a modified version of the motivic Adams spectral sequence in the sense of [BHHM08, §3].

5.2. *The strategy.* For a motivic spectrum X with the property that $X \simeq X_{c \geq 0}$, we consider its Postnikov–Whitehead tower with respect to the Chow t -structure:

$$\begin{array}{ccc} & \downarrow & \\ & X_{c \geq 2} & \longrightarrow X_{c=2} \\ & \downarrow & \\ & X_{c \geq 1} & \longrightarrow X_{c=1} \\ & \downarrow & \\ X & \xlongequal{\quad} X_{c \geq 0} & \longrightarrow X_{c=0}. \end{array}$$

Remark 5.10.

- By Corollary 3.7 and for degree reasons, if we take $\mathrm{MGL}_{*,*}$ of each term of this Postnikov–Whitehead tower, the corresponding Chow spectral sequence computing $\mathrm{MGL}_{*,*}X$ collapses at its E_1 -page.
- If we take $\pi_{*,*}$ of each term of this Postnikov–Whitehead tower, by Theorem 3.14, we have

$$\pi_{*,*}X_{c=n} \simeq \mathrm{Ext}_{\mathrm{MU}_{2*}\mathrm{MU}}^{*,*}(\mathrm{MU}_{2*}, \mathrm{MGL}_{2*+n,*}X).$$

The E_1 -page of the corresponding tower spectral sequence computing $\pi_{*,*}X$ is isomorphic to the E_2 -page of the motivic Adams–Novikov spectral sequence. Over $\mathbb{C}$, for the case $X = \mathbb{1}_p^\wedge$, it was proved in [Isa19] that after a re-grading, these two spectral sequences are isomorphic. We suspect that this is an isomorphism of spectral sequences after a re-grading over a general field.

Now consider the motivic Adams spectral sequences based on $\mathbb{H}\mathbb{Z}/p$ for every term in the Postnikov–Whitehead tower with respect to the Chow t -structure, and the induced maps among them. This gives us the following tower of motivic Adams spectral sequences:

$$\begin{array}{c}
 \downarrow \\
 \mathbf{motASS}(X_{c \geq 2}) \rightarrow \mathbf{motASS}(X_{c=2}) = \mathbf{algNSS}(\mathbf{BPGL}_{2*+2,*}X) \\
 \downarrow \\
 \mathbf{motASS}(X_{c \geq 1}) \rightarrow \mathbf{motASS}(X_{c=1}) = \mathbf{algNSS}(\mathbf{BPGL}_{2*+1,*}X) \\
 \downarrow \\
 \mathbf{motASS}(X) = \mathbf{motASS}(X_{c \geq 0}) \rightarrow \mathbf{motASS}(X_{c=0}) = \mathbf{algNSS}(\mathbf{BPGL}_{2*,*}X).
 \end{array}$$

We now have a strategy for computing differentials in the motivic Adams spectral sequence for X based on $\mathbb{H}\mathbb{Z}/p$. The input of our computations is $\mathbf{BPGL}_{*,*}X$ and $\mathbb{H}\mathbb{Z}/p_{*,*}X$ over the base field k .

For every $\mathbf{BP}_*\mathbf{BP}$ -comodule $\mathbf{BPGL}_{2*+n,*}X$, its algebraic Novikov spectral sequence can be computed in a large range by a computer program. This gives many non-zero differentials in the motivic Adams spectral sequence for $X_{c=n}$. We may then pull back these Adams differentials to the motivic Adams spectral sequences for $X_{c \geq n}$, and push forward them to the motivic Adams spectral sequences for X .

Here are detailed steps of this strategy:

- (1) Compute $\mathrm{Ext}_A^{*,*,*}(\mathbb{H}\mathbb{Z}/p_{*,*}, \mathbb{H}\mathbb{Z}/p_{*,*}X)$ over the k -motivic Steenrod algebra by a computer program. These Ext-groups consist of the E_2 -page of the motivic Adams spectral sequence for X based on $\mathbb{H}\mathbb{Z}/p$.
 - Over $\mathbb{C}$ for the spectrum $\mathbb{1}_2^\wedge$, this was computed in a large range in [IWX20a], [IWX20b].
 - Over $\mathbb{R}$ for the spectrum $\mathbb{1}_2^\wedge$, this was computed in a range in [DI17], [BI20] using the ρ -Bockstein spectral sequence and computations over $\mathbb{C}$. This ρ -Bockstein computation can also be automated.
 - Over finite fields for the spectrum $\mathbb{1}_2^\wedge$, this was computed in [WØ17] by a long exact sequence with the other two terms the $\mathbb{C}$ -motivic Ext. The boundary homomorphism corresponds to the Steenrod operation action on powers of τ .
- (2) Compute by a computer program the algebraic Novikov spectral sequence based on $\mathbf{BP}_{2*}\mathbf{BP}/I$ for the $\mathbf{BP}_{2*}\mathbf{BP}$ -comodules $\mathbf{BPGL}_{2*+n,*}X$ for every n in a reasonable range. The structures of these comodules $\mathbf{BPGL}_{2*+n,*}X$ depend on each field k for $n \geq 1$. For $n = 0$, $\mathbf{BPGL}_{2*+n,*} = \mathbf{BP}_{2*}$ after p -completion for all fields k . The case for $\mathbf{BP}_{2*}$ was computed in a large range in [IWX20a], [IWX20b]. This computation includes all differentials

and the multiplicative structure of the cohomology of the Hopf algebroid $(\mathrm{BP}_{2*}, \mathrm{BP}_{2*}\mathrm{BP})$.

- (3) Identify the k -motivic Adams spectral sequence based on $\mathrm{H}\mathbb{Z}/p$ for each $X_{c=n}$, with the algebraic Novikov spectral sequence based on $\mathrm{BP}_{2*}\mathrm{BP}/I$ for $\mathrm{BPGL}_{2*+n,*}X$, for $n \geq 0$. This computation includes an identification of the abutments with the multiplicative (and higher) structures.
- (4) Compute the mod p motivic homology of $X_{c \geq n}$ using the universal coefficient spectral sequence (see Propositions 7.7 and 7.10 of [DI05]):

$$\bigoplus_{k \geq n} \mathrm{Tor}_{*,*}^{\mathrm{BP}_{2*}}(\mathrm{BPGL}_{2*+k,*}X, \mathbb{Z}/p) \implies \pi_{*,*}X_{c \geq n} \wedge \mathrm{H}\mathbb{Z}/p.$$

Here we use the fact that $\mathbb{1}_{c=0} \wedge \mathrm{BPGL}/(p, v_1, \dots)$ is a cellular module over $\mathbb{1}_{c=0} \wedge \mathrm{BPGL}$ and the equivalences

$$\begin{aligned} X_{c \geq n} \wedge \mathrm{H}\mathbb{Z}/p &\simeq X_{c \geq n} \wedge \mathrm{BPGL}/(p, v_1, \dots) \\ &\simeq (X_{c \geq n} \wedge \mathrm{BPGL}) \wedge_{(\mathbb{1}_{c=0} \wedge \mathrm{BPGL})} (\mathbb{1}_{c=0} \wedge \mathrm{BPGL}/(p, v_1, \dots)), \end{aligned}$$

and the isomorphisms

$$\begin{aligned} \pi_{*,*}\mathbb{1}_{c=0} \wedge \mathrm{BPGL} &\cong \mathrm{BP}_{2*}, \\ \pi_{*,*}X_{c \geq n} \wedge \mathrm{BPGL} &\cong \bigoplus_{k \geq n} \mathrm{BPGL}_{2*+k,*}X, \\ \pi_{*,*}\mathbb{1}_{c=0} \wedge \mathrm{BPGL}/(p, v_1, \dots) &\cong \mathbb{Z}/p. \end{aligned}$$

- (5) Compute by a computer program the E_2 -pages of the motivic Adams spectral sequence for $X_{c \geq n}$ based on $\mathrm{H}\mathbb{Z}/p$, using the computation of $\pi_{*,*}X_{c \geq n} \wedge \mathrm{H}\mathbb{Z}/p$ in step (4).
- (6) Pull back motivic Adams differentials for $X_{c=n}$ to motivic Adams differentials for $X_{c \geq n}$, and then push forward to motivic Adams differentials for X .
- (7) Deduce additional Adams differentials for X with a variety of ad hoc arguments. The most important methods are Toda bracket shuffles and comparison to known results in the $\mathbb{C}$ -motivic Adams spectral sequence.

Most of the computations in steps (1)–(5) can be automated. In the rest of this section, we will elaborate on step (6) of this strategy, in the cases of $X = \mathbb{1}$ and $k = \mathbb{C}$, $\mathbb{R}$ and finite fields.

5.3. Continuous Adams spectral sequences. For the sphere spectrum $\mathbb{1}$, the inputs of our strategy are $\mathrm{H}\mathbb{Z}/p_{*,*}$ and $\mathrm{BPGL}_{*,*}$. In practice, it is more plausible to know $\pi_{*,*}(\mathrm{BPGL}_p^\wedge)$ rather than $\mathrm{BPGL}_{*,*}$. In order to take the p -completion into account, we consider pro-objects as a substitute.

Definition 5.11. Let $(X_i)_{i \in I}$ be a pro-object in $\mathcal{SH}(k)$. We call the cosimplicial object

$$\Delta \ni n \mapsto \lim_i [X_i \wedge (\mathbb{H}\mathbb{Z}/p)^{\wedge n}]$$

the *continuous $\mathbb{H}\mathbb{Z}/p$ -resolution* of $(X_i)_{i \in I}$, and the associated spectral sequence the *continuous Adams spectral sequence* of $(X_i)_{i \in I}$.

Remark 5.12. Given $X \in \mathcal{SH}(k)$, write

$$F_p(X) = (\cdots \rightarrow X \xrightarrow{p} X \xrightarrow{p} X) \in \text{Pro}(\mathcal{SH}(k)),$$

and $(X/p^n)_n \in \text{Pro}(\mathcal{SH}(k))$ for the pro- p -completion. Then there is a fiber sequence

$$F_p(X) \rightarrow X \rightarrow (X/p^n)_n,$$

where in the middle we mean the constant pro-object. Thus if $Y \in \mathcal{SH}(k)$ such that $p : Y \rightarrow Y$ is equivalent to the trivial map, then $F_p(X) \wedge Y \simeq 0$ in $\text{Pro}(\mathcal{SH}(k))$ and so $X \wedge Y \simeq (X/p^n \wedge Y)_n$ in $\text{Pro}(\mathcal{SH}(k))$. In particular, the continuous $\mathbb{H}\mathbb{Z}/p$ -resolution of the pro- p -completion $(X/p^n)_n$ is just the usual $\mathbb{H}\mathbb{Z}/p$ -resolution of X .

We now consider the following tower of pro-objects:

$$\cdots \rightarrow ((\mathbb{1}/p^n)_{c \geq i})_n \rightarrow \cdots \rightarrow ((\mathbb{1}/p^n)_{c \geq 1})_n \rightarrow ((\mathbb{1}/p^n)_{c \geq 0})_n \simeq (\mathbb{1}/p^n)_n.$$

The layers are given by

$$((\mathbb{1}/p^n)_{c=i})_n \in \text{Pro}(\mathcal{SH}(k)^{c\heartsuit}).$$

Their continuous Adams spectral sequences are related in the way outlined above to the continuous Adams spectral sequence for $((\mathbb{1}/p^n)_{c \geq 0})_n$, which is just the pro- p -completion of $\mathbb{1}$, and so by [Remark 5.12](#) we recover information about the usual motivic Adams spectral sequence. It remains to identify the continuous Adams spectral sequence of the layers; we can achieve this under some simplifying assumptions.

PROPOSITION 5.13. *Suppose that $\pi_{*,*}\text{BPGL}_p^\wedge$ has bounded p -torsion. Put*

$$B_*^i = \pi_{2*+i,*}\text{BPGL}_p^\wedge,$$

viewed as a BP_{2}BP -comodule. Then $(\pi_{*,*}(\mathbb{1}/p^n)_{c=i})_n$ corresponds to the pro- p -completion of $B_*^i \in \mathcal{H}\text{ov}(\text{MU}_{2*}\text{MU})$. In particular, the continuous Adams spectral sequence of $((\mathbb{1}/p^n)_{c=i})_n$ is isomorphic to the algebraic Novikov spectral sequence of B_*^i .*

Proof. We may replace BP by MU throughout. We know that $(\mathbb{1}/p^n)_{c=i} \in \mathcal{SH}(k)^{\text{cell}, c\heartsuit}$ corresponds to the MU_{2*}MU -comodule $X_n^i = \pi_{2*+i,*}(\text{MGL}/p^n)$.

Let K_n^i be the kernel of multiplication by p^n on B_*^i . Then we have a short exact sequence

$$0 \rightarrow (B^i/p^n)_n \rightarrow (X_n^i)_n \rightarrow (K_n^{i-1})_n \rightarrow 0$$

of pro-objects in $\mathrm{MU}_{2*}\mathrm{MU}$ -comodules. The bonding maps in $(K_n^{i-1})_n$ are given by multiplication by p , and since the p -torsion is bounded, sufficiently long composites of bonding maps are zero. It follows that $(K_n^{i-1})_n \simeq 0$ as a pro-object, and so $(B^i/p^n)_n \simeq (X_n^i)_n$ as pro-objects. Let

$$Y_n^i = \mathrm{cof}(B^i \xrightarrow{p^n} B^i) \in \mathcal{H}\mathrm{ov}(\mathrm{MU}_{2*}\mathrm{MU});$$

in other words, $(Y_n^i)_n$ is the pro- p -completion of B^i in the derived category of comodules. We have a cofiber sequence

$$\Sigma(K_n^i)_n \rightarrow (Y_n^i)_n \rightarrow (B^i/p^n)_n,$$

and so arguing as before and combining with the previous result, we find that

$$(Y_n^i)_n \simeq (B^i/p^n)_n \simeq (X_n^i)_n.$$

The last sentence follows from [Remark 5.12](#). $\square$

5.4. Over the complex numbers. Over the complex numbers, this strategy has been proved to be very successful. In [\[IWX20a\]](#), [\[IWX20b\]](#), using this method, the knowledge of both $\mathbb{C}$ -motivic and classical stable homotopy groups of spheres at the prime 2 was extended from 60 to 90. It is also used in [\[GWX21, §10\]](#) to provide one-line proofs for many historically hard Adams differentials.

In fact, over $\mathbb{C}$, we have $\pi_{*,*}\mathrm{BPGL}_p^\wedge = (\mathrm{BP}_{2*})_p^\wedge[\tau] = \mathbb{Z}_p^\wedge[\tau, v_1, v_2, \dots]$, with $|\tau| = (0, -1)$, $|v_i| = (2p^i - 2, p^i - 1)$. In other words, we have

$$\pi_{2*+n,*}\mathrm{BPGL}_p^\wedge \cong \begin{cases} \Sigma^{0,-m}(\mathrm{BP}_{2*})_p^\wedge & \text{if } n = 2m \text{ and } m \geq 0, \\ 0 & \text{otherwise,} \end{cases}$$

where $\Sigma^{0,-m}(\mathrm{BP}_{2*})_p^\wedge$ means every element in $(\mathrm{BP}_{2*})_p^\wedge$ is shifted by bidegree $(0, -m)$.

This corresponds to the fact that

$$\pi_{*,*}(\mathbb{1}_p^\wedge)_{c=n} \cong \begin{cases} \pi_{*,*}\Sigma^{0,-m}\mathbb{1}_p^\wedge/\tau & \text{if } n = 2m \text{ and } m \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

By Proposition 7.2 of [\[GWX21\]](#), we have $\mathrm{MGL}_{*,*}\mathbb{1}_p^\wedge \cong \pi_{*,*}\mathrm{MGL}_p^\wedge$ over $\mathbb{C}$. By a similar proof to that of Proposition 7.2 of [\[GWX21\]](#), we have $\mathrm{BPGL}_{*,*}\mathbb{1}_p^\wedge \cong \pi_{*,*}\mathrm{BPGL}_p^\wedge$. Then the tower of motivic Adams spectral sequences based on $\mathrm{H}\mathbb{Z}/p$ becomes

$$\begin{array}{c}
\downarrow \\
\mathbf{motASS}((\mathbb{1}_p^\wedge)_{c \geq 4}) \rightarrow \mathbf{motASS}((\mathbb{1}_p^\wedge)_{c=4}) \xlongequal{\quad} \mathbf{algNSS}(\Sigma^{0,-2}(\mathbf{BP}_{2*})_p^\wedge) \\
\downarrow \\
\mathbf{motASS}((\mathbb{1}_p^\wedge)_{c \geq 2}) \rightarrow \mathbf{motASS}(\mathbb{1}_p^\wedge)_{c=2} \xlongequal{\quad} \mathbf{algNSS}(\Sigma^{0,-1}(\mathbf{BP}_{2*})_p^\wedge) \\
\downarrow \\
\mathbf{motASS}(\mathbb{1}_p^\wedge) \xlongequal{\quad} \mathbf{motASS}(\mathbb{1}_p^\wedge) \longrightarrow \mathbf{motASS}((\mathbb{1}_p^\wedge)_{c=0}) = \mathbf{algNSS}((\mathbf{BP}_{2*})_p^\wedge)
\end{array}$$

Here we may view $\Sigma^{m,n}(\mathbf{BP}_{2*})_p^\wedge$ as an element in the derived category of p -completed $\mathbf{BP}_{2*}\mathbf{BP}$ -comodules: It is a cochain complex concentrated in cohomological degree $2n - m$, with the comodule $\Sigma^{2n}(\mathbf{BP}_{2*})_p^\wedge$ in that cohomological degree. Under the equivalence of the categories in [Corollary 4.29](#), $\Sigma^{m,n}(\mathbf{BP}_{2*})_p^\wedge$ corresponds to $\Sigma^{m,n}\mathbb{1}_p^\wedge/\tau$.

In this case, all horizontal arrows are identical, up to a shift of degrees. So the information above the bottom row is not of much use. This is indeed the case in the work of [\[IWX20a\]](#), [\[IWX20b\]](#), where only the naturality of the motivic Adams spectral sequences of the unit map $\mathbb{1}_p^\wedge \rightarrow \mathbb{1}_p^\wedge/\tau$ was used.

5.5. *Over the real numbers.* Over the real numbers, at the prime 2, recall that we have

$$\pi_{*,*}\mathbf{HZ}/2 = \mathbb{Z}/2[\rho, \tau],$$

with $|\rho| = (-1, -1)$, $|\tau| = (0, -1)$. The class ρ is a map

$$\rho : \mathbb{1} \rightarrow \mathbb{G}_m = \Sigma^{1,1}\mathbb{1},$$

that corresponds to $-1 \in \mathbb{R}^\times$ (see [\[Bac18a\]](#) for example), and it can be further viewed as a map

$$\rho : \Sigma^{-1,-1}\mathbb{1}_2^\wedge \rightarrow \mathbb{1}_2^\wedge.$$

We denote by $\mathbb{1}_2^\wedge/\rho$ the cofiber of ρ .

By [Corollary 1.9](#) of [\[BS20\]](#), there is an equivalence between the category of cellular spectra over $\mathbb{C}$ and the category of cellular modules over $\mathbb{1}_2^\wedge/\rho$ over $\mathbb{R}$, under which the $\mathbb{C}$ -motivic sphere $\mathbb{1}_2^\wedge$ corresponds to $\mathbb{1}_2^\wedge/\rho$ over $\mathbb{R}$.

The class τ then corresponds to a map

$$\tau : \Sigma^{0,-1}\mathbb{1}_2^\wedge/\rho \rightarrow \mathbb{1}_2^\wedge/\rho.$$

We denote by $\mathbb{1}_2^\wedge/(\rho, \tau)$ the cofiber of τ .

The $\mathbb{R}$ -motivic $\pi_{*,*}\mathbf{BPGL}_2^\wedge$ was computed by Hu–Kriz [\[HK01\]](#) and Hill [\[Hil11\]](#).

PROPOSITION 5.14. *We have*

$$\pi_{*,*}\mathrm{BPGL}_2^\wedge \cong \mathbb{Z}_2^\wedge \left[\begin{array}{cccccc} \rho, & & & & & \\ v_0, & \tau^2 v_0, & \tau^4 v_0, & \tau^6 v_0, & \tau^8 v_0, & \dots \\ v_1, & & \tau^4 v_1, & & \tau^8 v_1, & \dots \\ v_2, & & & & \tau^8 v_2, & \dots \\ \dots & & & & & \end{array} \right] / \left[\begin{array}{l} v_0 = 2 \\ \rho v_0 = 0 \\ \rho^3 v_1 = 0 \\ \rho^7 v_2 = 0 \\ \dots \end{array} \right]$$

and the generators satisfy the further relations

$$\tau^{2^{i+1} \cdot j} v_i \cdot \tau^{2^{k+1} \cdot l} v_k = \tau^{2^{i+1}(j+2^{k-i}l)} v_i v_k$$

when $i \leq k$, as if the class τ were an element in this ring.

Here the bidegrees of the generators are

$$|\tau| = (0, -1), \quad |\rho| = (-1, -1), \quad |v_n| = (2^{n+1} - 2, 2^n - 1).$$

For the Chow degrees, we have τ and ρ in degrees 2 and 1, and we have all v_n 's in degree 0. Therefore, we can read off the Chow degree n part of the $\mathbb{R}$ -motivic $\pi_{*,*}\mathrm{BPGL}_2^\wedge$. The first few are the following:

- **Chow = 0:** $\mathbb{Z}_2^\wedge[v_0, v_1, \dots] = (\mathrm{BP}_{2*})_2^\wedge$.
- **Chow = 1:** $\rho \cdot \mathbb{Z}_2^\wedge[v_0, v_1, \dots] / \rho v_0 = \Sigma^{-1, -1} \mathrm{BP}_{2*} / 2$.
- **Chow = 2:** $\rho^2 \cdot \mathbb{Z}_2^\wedge[v_0, v_1, \dots] / \rho v_0 = \Sigma^{-2, -2} \mathrm{BP}_{2*} / 2$.
- **Chow = 3:** $\rho^3 \cdot \mathbb{Z}_2^\wedge[v_0, v_1, \dots] / (\rho v_0, \rho^3 v_1) = \Sigma^{-3, -3} \mathrm{BP}_{2*} / (2, v_1)$.
- **Chow = 4:** $\rho^4 (\mathrm{BP}_{2*})_2^\wedge \oplus \tau^2 v_0 (\mathrm{BP}_{2*})_2^\wedge = \Sigma^{-4, -4} \mathrm{BP}_{2*} / (2, v_1) \oplus \Sigma^{0, -2} (\mathrm{BP}_{2*})_2^\wedge$.
- **Chow = 5:** $\rho^5 (\mathrm{BP}_{2*})_2^\wedge = \Sigma^{-5, -5} \mathrm{BP}_{2*} / (2, v_1)$.
- **Chow = 6:** $\rho^6 (\mathrm{BP}_{2*})_2^\wedge = \Sigma^{-6, -6} \mathrm{BP}_{2*} / (2, v_1)$.
- **Chow = 7:** $\rho^7 (\mathrm{BP}_{2*})_2^\wedge = \Sigma^{-7, -7} \mathrm{BP}_{2*} / (2, v_1, v_2)$.
- **Chow = 8:** $\rho^8 (\mathrm{BP}_{2*})_2^\wedge \oplus \frac{\tau^4 v_0 (\mathrm{BP}_{2*})_2^\wedge + \tau^4 v_1 (\mathrm{BP}_{2*})_2^\wedge}{\tau^4 v_0 \cdot v_1 - \tau^4 v_1 \cdot v_0}$
 $= \Sigma^{-8, -8} \mathrm{BP}_{2*} / (2, v_1, v_2) \oplus (\Sigma^{0, -4} (\mathrm{BP}_{2*})_2^\wedge + \Sigma^{2, -3} \mathrm{BP}_{2*} / 2)$.

On the spectrum level, this corresponds to the fact that

$$\pi_{*,*}(\mathbb{1}_2^\wedge)_{c=0} \cong \pi_{*,*} \mathbb{1}_2^\wedge / (\rho, \tau).$$

In the category of $\mathrm{BP}_{2*}\mathrm{BP}$ -comodules, we have short exact sequences

$$0 \longrightarrow \mathrm{BP}_{2*} / I_n \xrightarrow{v_{n+1}} \mathrm{BP}_{2*} / I_n \longrightarrow \mathrm{BP}_{2*} / I_{n+1} \longrightarrow 0,$$

where $I_n = (2, v_1, \dots, v_n)$. This implies that we could realize the comodule BP_{2*}/I_n inductively by the spectrum $\mathbb{1}_2^\wedge/(\rho, \tau, 2, v_1, \dots, v_n)$. These spectra are all $\mathcal{E}_\infty$.

In higher Chow degrees, these comodules are more complicated. However, [Corollary 4.29](#) tells us they correspond to module spectra over $\mathbb{1}_2^\wedge/(\rho, \tau)$. By [Proposition 5.14](#), these comodules can be filtered such that the subquotients are shifts of BP_{2*}/I_n . Then by naturality, we can use the motivic Adams spectral sequences for shifts of the spectra $\mathbb{1}_2^\wedge/(\rho, \tau, 2, v_1, \dots, v_n)$ to study the one for $(\mathbb{1}_2^\wedge)_{c=n}$.

For example in Chow degree 8, by [Corollary 4.29](#), the second summand of the comodule, denoted by M_8 , corresponds to a module spectrum over $\mathbb{1}_2^\wedge/(\rho, \tau)$, denoted by X_8 . Then the short exact sequence of comodules

$$0 \longrightarrow \Sigma^{0,-4}(\mathrm{BP}_{2*})_2^\wedge \longrightarrow M_8 \longrightarrow \Sigma^{2,-3}\mathrm{BP}_{2*}/2 \longrightarrow 0$$

gives us a cofiber sequence of module spectra over $\mathbb{1}_2^\wedge/(\rho, \tau)$

$$\Sigma^{0,-4}\mathbb{1}_2^\wedge/(\rho, \tau) \longrightarrow X_8 \longrightarrow \Sigma^{2,-3}\mathbb{1}_2^\wedge/(\rho, \tau, 2).$$

This cofiber sequence gives us maps of motivic Adams spectral sequences:

$$\mathbf{motASS}(\Sigma^{0,-4}\mathbb{1}_2^\wedge/(\rho, \tau)) \rightarrow \mathbf{motASS}(X_8) \rightarrow \mathbf{motASS}(\Sigma^{2,-3}\mathbb{1}_2^\wedge/(\rho, \tau, 2)).$$

We anticipate that this would allow us to compute motivic Adams differentials for X_8 from the ones for $\Sigma^{0,-4}\mathbb{1}_2^\wedge/(\rho, \tau)$ and $\Sigma^{2,-3}\mathbb{1}_2^\wedge/(\rho, \tau, 2)$, which are purely algebraic. This gives us a tower of motivic Adams spectral sequences:

$$\begin{array}{c} \downarrow \\ \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c \geq 3}) \rightarrow \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c=3}) \xlongequal{\text{algNSS}} (\Sigma^{-3,-3}\mathrm{BP}_{2*}/(2, v_1)) \\ \downarrow \\ \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c \geq 2}) \rightarrow \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c=2}) \xlongequal{\text{algNSS}} (\Sigma^{-2,-2}\mathrm{BP}_{2*}/2) \\ \downarrow \\ \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c \geq 1}) \rightarrow \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c=1}) \xlongequal{\text{algNSS}} (\Sigma^{-1,-1}\mathrm{BP}_{2*}/2) \\ \downarrow \\ \mathbf{motASS}(\mathbb{1}_2^\wedge) \xlongequal{\text{algNSS}} \mathbf{motASS}(\mathbb{1}_2^\wedge) \longrightarrow \mathbf{motASS}((\mathbb{1}_2^\wedge)_{c=0}) \xlongequal{\text{algNSS}} (\mathrm{BP}_{2*})_2^\wedge. \end{array}$$

Comparing to the case over $\mathbb{C}$, it is clear that over $\mathbb{R}$ different rows contain different algebraic information of motivic Adams differentials. We anticipate that this allows us to obtain motivic Adams differentials for $\mathbb{1}_2^\wedge$ from the algebraic Novikov spectral sequences for various $\mathrm{BP}_{2*}\mathrm{BP}$ -comodules.

5.6. *Over finite fields.* Next, we discuss the cases of finite fields $\mathbb{F}_q$, where q is a power of an odd prime. We are going to concentrate on the computations at the prime 2.

Recall from [HKØ17] and Theorem 8.2 of [Kyl19] (see also [WØ17] and Appendix C) that we have

$$\bullet \pi_{*,*}\mathrm{HZ}/2 = \begin{cases} \mathbb{Z}/2[\tau, u]/u^2, & Sq^1\tau = 0 \quad \text{if } q \equiv 1 \pmod{4}, \\ \mathbb{Z}/2[\tau, \rho]/\rho^2, & Sq^1\tau = \rho \quad \text{if } q \equiv 3 \pmod{4}, \end{cases}$$

where $|\tau| = (0, -1)$, $|u| = |\rho| = (-1, -1)$;

$$\bullet \pi_{*,*}\mathrm{HZ}_2^\wedge = \begin{cases} \mathbb{Z}_2^\wedge & \text{if } (*, *) = (0, 0), \\ (\mathbb{Z}/(q^w - 1))_2^\wedge & \text{if } (*, *) = (-1, -w) \text{ and } w \geq 1, \\ 0 & \text{otherwise;} \end{cases}$$

$$\bullet \pi_{*,*}\mathrm{BPGL}_2^\wedge = \pi_{*,*}\mathrm{HZ}_2^\wedge \otimes \mathrm{BP}_{2*}.$$

We can read off the Chow degree n part of $\pi_{*,*}\mathrm{BPGL}_2^\wedge$. In particular, it is concentrated in Chow degrees 0 and $2w - 1$ for $w \geq 1$. In positive Chow degrees, we have

$$\mathrm{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=2w} = 0,$$

$$\mathrm{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=2w-1} = \Sigma^{-1, -w}\mathrm{BP}_{2*}/2^{\nu_2(q^w-1)},$$

where $\nu_2(-)$ is the 2-adic valuation function.

Remark 5.15. We remark that in positive chow degrees, most of these BP_{2*} -modules do not have Levin index 1. For example in Chow degree 1, it is $\Sigma^{-1, -1}\mathrm{BP}_{2*}/2^{\nu_2(q-1)}$. In particular, when $q \equiv 5 \pmod{8}$, we have $\Sigma^{-1, -1}\mathrm{BP}_{2*}/4$, and when $q \equiv 1 \pmod{8}$, we have $\Sigma^{-1, -1}\mathrm{BP}_{2*}/8 \cdot 2^m$ for $m \geq 0$. Both of them do not have Levin index 1. In fact, one can show that $\mathrm{BP}_{2*}/2^{\nu_2(q^w-1)}$ has Levin index 1 if and only if $\nu_2(q^w - 1) = 1$.

For every spectrum $(\mathbb{1}_2^\wedge)_{c=n}$, by a change-of-ring isomorphism and the description of $\mathrm{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=n}$, the universal coefficient spectral sequence collapses for degree reasons (see Propositions 7.7 and 7.10 of [DI05] and step (4) of our strategy in Section 5.2):

$$\mathrm{Tor}_{*,*}^{(\mathrm{BP}_{2*})_2^\wedge}(\mathrm{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=n}, \mathbb{Z}/2) \implies \mathrm{HZ}/2_{*,*}(\mathbb{1}_2^\wedge)_{c=n}.$$

We have that for $w \geq 1$,

$$\mathrm{HZ}/2_{*,*}(\mathbb{1}_2^\wedge)_{c=2w} = 0,$$

$$\mathrm{HZ}/2_{*,*}(\mathbb{1}_2^\wedge)_{c=2w-1} = \begin{cases} \mathbb{Z}/2\{\tau^w\} \oplus \mathbb{Z}/2\{u\tau^{w-1}\} & \text{if } q \equiv 1 \pmod{4}, \\ \mathbb{Z}/2\{\tau^w\} \oplus \mathbb{Z}/2\{\rho\tau^{w-1}\} & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

Here τ^w , $u\tau^{w-1}$, $\rho\tau^{w-1}$ denote the generators of the groups $\mathbb{Z}/2$, in bidegrees $(0, -w)$, $(-1, -w)$, $(-1, -w)$. Note that

$$\mathrm{HZ}/2_{*,*}(\mathbb{1}_2^\wedge)_{c=0} = \mathrm{HZ}/2\{1\},$$

generated by 1 in bidegree $(0, 0)$. By [Corollary 4.29](#), these spectra $(\mathbb{1}_2^\wedge)_{c=n}$ are cellular modules over $(\mathbb{1}_2^\wedge)_{c=0}$. Therefore, we learn that for $w \geq 1$,

$$\begin{aligned}\pi_{*,*}(\mathbb{1}_2^\wedge)_{c=2w} &= 0, \\ \pi_{*,*}(\mathbb{1}_2^\wedge)_{c=2w-1} &= \pi_{*,*}\Sigma^{-1,-w}(\mathbb{1}_2^\wedge)_{c=0}/2^{\nu_2(q^w-1)}.\end{aligned}$$

In particular, the spectrum $(\mathbb{1}_2^\wedge)_{c=2w-1}$ has the same homotopy groups as a 2-cell complex over $(\mathbb{1}_2^\wedge)_{c=0}$, with cells in bidegrees $(-1, -w)$ and $(0, -w)$, and attaching map $2^{\nu_2(q^w-1)}$, or equivalently $q^w - 1$.

We emphasize that the spectrum $(\mathbb{1}_2^\wedge)_{c=0}$ is very well understood independent of the base field (whose characteristics is not 2, which includes these finite fields $\mathbb{F}_q$). By [Corollaries 4.29](#) and [5.5](#), the motivic Adams spectral sequence of $(\mathbb{1}_2^\wedge)_{c=0}$ is isomorphic to the algebraic Novikov spectral sequence for BP_{2*} , which is again independent of the base field. Over $\mathbb{C}$, this spectral sequence is isomorphic to the motivic Adams spectral sequence of $\mathbb{1}_2^\wedge/\tau$, and it is well understood in a large range of dimensions [[IWX20a](#)], [[IWX20b](#)]. For the spectrum $(\mathbb{1}_2^\wedge)_{c=0}/2^{\nu_2(q^w-1)}$, its motivic Adams spectral sequence is isomorphic to the algebraic Novikov spectral sequence for $\mathrm{BP}_{2*}/2^{\nu_2(q^w-1)}$, which can also be understood through known computations over $\mathbb{C}$.

As an application of the Postnikov–Whitehead tower of the Chow t -structure, we compute all Adams differentials on powers of τ , reproving [Corollary 7.12](#) of [[WØ17](#)]. By Leibniz’s rule, we only need to compute all non-zero differentials on classes of the form τ^{2^n} .

PROPOSITION 5.16. *Let $m(n) = \nu_2(q^{2^n} - 1)$, where $\nu_2(-)$ is the 2-adic valuation function.*

(1) *When $q \equiv 1 \pmod{4}$, all non-zero Adams differentials on the classes τ^{2^n} are*

$$d_{m(n)}(\tau^{2^n}) = h_0^{m(n)} u \tau^{2^n-1} \text{ for } n \geq 0.$$

(2) *When $q \equiv 3 \pmod{4}$, all non-zero Adams differentials on the classes τ^{2^n} are*

$$d_{m(n)}(\tau^{2^n}) = h_0^{m(n)} \rho \tau^{2^n-1} \text{ for } n \geq 1.$$

Proof. Suppose that $q \equiv 1 \pmod{4}$. For degree reasons, the only non-zero Adams differential on τ^{2^n} has the form

$$d_m(\tau^{2^n}) = h_0^m u \tau^{2^n-1}$$

for some integer m , depending on n .

Let $w = 2^n$. From the above discussion, we have

$$\mathrm{HZ}/2_{*,*}(\mathbb{1}_2^\wedge)_{c=2^{n+1}-1} = \mathbb{Z}/2\{\tau^{2^n}\} \oplus \mathbb{Z}/2\{u\tau^{2^n-1}\}.$$

So the differentials of interest are present in the motivic Adams spectral sequence for $(\mathbb{1}_2^\wedge)_{c=2^{n+1}-1}$, and the ones for $\mathbb{1}_2^\wedge$ follows from naturality of the motivic Adams spectral sequences and the following zigzag in the Postnikov–Whitehead tower:

$$\begin{array}{ccccc} \text{motASS} & & \text{motASS} & & \text{algNSS} \\ ((\mathbb{1}_2^\wedge)_{c \geq 2^{n+1}-1}) & \longrightarrow & ((\mathbb{1}_2^\wedge)_{c=2^{n+1}-1}) & \xlongequal{\quad} & (\text{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=2^{n+1}-1}) \\ \downarrow & & & & \\ \text{motASS}(\mathbb{1}_2^\wedge). & & & & \end{array}$$

We have

$$\text{BPGL}_{*,*}(\mathbb{1}_2^\wedge)_{c=2^{n+1}-1} = \Sigma^{-1,-2^n} \text{BP}_{2*} / 2^{\nu_2(q^{2^n}-1)}.$$

Therefore, in the algebraic Novikov spectral sequence, we have

$$d_{m(n)}(\tau^{2^n}) = h_0^{m(n)} u \tau^{2^n-1},$$

where $m(n) = \nu_2(q^{2^n}-1)$. This completes the proof for the cases $q \equiv 1 \pmod{4}$.

For the cases $q \equiv 3 \pmod{4}$, note that τ^2 is an indecomposable element on the motivic Adams E_2 -page. The rest of the proof works in the exact same way. $\square$

In [WØ17], Wilson–Østvær computed $\pi_{n,0} \mathbb{1}_2^\wedge$ for $0 \leq n \leq 20$ when $q \equiv 1 \pmod{4}$ and for $0 \leq n \leq 18$ when $q \equiv 3 \pmod{4}$. The difference of ranges is due to the computation of an Adams d_2 -differential from the 20-stem to the 19-stem in the cases $q \equiv 1 \pmod{4}$, while the answer for an Adams d_2 -differential is unknown in the cases $q \equiv 3 \pmod{4}$. This d_2 -differential is also the hardest one in the computation up to the 20-stem. In fact, Wilson–Østvær proved

PROPOSITION 5.17 ([WØ17, Prop. 7.16, Rem. 7.19]). *Over $\mathbb{F}_q$, where q is a power of an odd prime, in the motivic Adams spectral sequence for $\mathbb{1}_2^\wedge$, we have*

- (1) $d_2(\tau g) = u h_0^2 g$, when $q \equiv 5 \pmod{8}$;
- (2) $d_2(\tau g) = 0$, when $q \equiv 1 \pmod{8}$;
- (3) $d_2(\tau^2 g) = \rho \tau h_0^2 g$ or 0, when $q \equiv 3 \pmod{4}$.

As an application of Proposition 5.16 and the Chow t -structure method, we give much simpler proofs of the Adams d_2 -differentials in the cases (1) and (2) in Proposition 5.17. We also compute the unknown Adams d_2 -differential in the cases $q \equiv 3 \pmod{4}$ in Proposition 5.17. The intuitions of our proofs are explained in Remarks 5.18 and 5.20.

Proof of Proposition 5.17 (1) and (2). In both cases $q \equiv 1, 5 \pmod{8}$, as discussed in [WØ17], for degree reasons, the only possibilities are

$$d_2(\tau g) = u h_0^2 g \text{ or } 0.$$

On the motivic Adams E_2 -page, we have

$$\begin{aligned}\tau g \cdot d_0 &= \tau e_0^2 \neq 0, \\ uh_0^2 g \cdot d_0 &= uh_0^2 e_0^2 \neq 0.\end{aligned}$$

See [WØ17, Prop. 7.1] for the multiplicative structure of the Adams E_2 -page.

By Proposition 5.16(1), for the case $n = 0$, we have

$$d_2(\tau) = \begin{cases} 0 & \text{if } q \equiv 1 \pmod{8}, \\ uh_0^2 & \text{if } q \equiv 5 \pmod{8}. \end{cases}$$

By Leibniz's rule, we have $d_2(e_0^2) = 0$. Thus,

$$d_2(\tau \cdot e_0^2) = d_2(\tau) \cdot e_0^2 = \begin{cases} 0 & \text{if } q \equiv 1 \pmod{8}, \\ uh_0^2 e_0^2 & \text{if } q \equiv 5 \pmod{8}. \end{cases}$$

The element d_0 is a permanent cycle for degree reasons. So we have $d_2(\tau g \cdot d_0) = d_2(\tau g) \cdot d_0$ and

$$d_2(\tau g) = \begin{cases} 0 & \text{if } q \equiv 1 \pmod{8}, \\ uh_0^2 g & \text{if } q \equiv 5 \pmod{8}. \end{cases} \quad \square$$

Remark 5.18. There are two reasons why it is harder to come up with direct proofs of Proposition 5.17(1) and (2).

- The element τg is indecomposable on the motivic Adams E_2 -page.
- On the E_2 -pages, the element τg is detected by $(\mathbb{1}_2^\wedge)_{c=0}$, while its potential target $uh_0^2 g$ is detected by $(\mathbb{1}_2^\wedge)_{c=1}$. They live in different Chow degrees.

After multiplication by d_0 , both $\tau g \cdot d_0 = \tau e_0^2$ and $uh_0^2 g \cdot d_0 = uh_0^2 e_0^2$ are detected by $(\mathbb{1}_2^\wedge)_{c=1}$ on the motivic Adams E_2 -pages, so this d_2 -differential can be obtained through the algebraic Novikov spectral sequence, something purely algebraic.

PROPOSITION 5.19. *In the motivic Adams spectral sequence for $\mathbb{1}_2^\wedge$ over $\mathbb{F}_q$, where q is a power of an odd prime, and $q \equiv 3 \pmod{4}$, we have*

$$d_2(\tau^2 g) = 0.$$

Proof. For degree reasons, the only possibilities are

$$d_2(\tau^2 g) = \rho \tau h_0^2 g \text{ or } 0.$$

On the motivic Adams E_2 -page, we have

$$\begin{aligned}\tau^2 g \cdot d_0 &= \tau^2 e_0^2 \neq 0, \\ \rho \tau h_0^2 g \cdot d_0 &= \rho \tau h_0^2 e_0^2 \neq 0.\end{aligned}$$

See [WØ17, Prop. 7.1] for the multiplicative structure of the Adams E_2 -page.

By Proposition 5.16(2) for the case $n = 1$, we have

$$d_2(\tau^2) = 0.$$

(In fact, we have $d_3(\tau^2) = \rho\tau h_0^3$ when $q \equiv 3 \pmod{8}$, and τ^2 supports non-zero d_4 or higher when $q \equiv 7 \pmod{8}$.)

By Leibniz's rule, we have

$$d_2(\tau^2 g) \cdot d_0 = d_2(\tau^2 g \cdot d_0) = d_2(\tau^2 \cdot e_0^2) = d_2(\tau^2) \cdot e_0^2 = 0.$$

The element d_0 is a permanent cycle for degree reasons. Since $\rho\tau h_0^2 g \cdot d_0 \neq 0$, we have

$$d_2(\tau^2 g) = 0. \quad \square$$

Remark 5.20. Similarly to [Remark 5.18](#), on the E_2 -pages, the element $\tau^2 g$ is detected by $(\mathbb{1}_2^\wedge)_{c=1}$, while its potential target $\rho\tau h_0^2 g$ is detected by $(\mathbb{1}_2^\wedge)_{c=3}$. They live in different Chow degrees. After multiplication by d_0 , they are both detected by $(\mathbb{1}_2^\wedge)_{c=3}$, and the d_2 -differential can be obtained through the algebraic Novikov spectral sequence.

Remark 5.21. We note that by Leibniz's rule, $d_2(\tau^2 g) = 0$ in all cases for q .

Appendix A. A cell structure for MGL

We record the fact that the Thom spectrum MGL admits a very well-behaved cell structure. We suspect that this is well known, but we do not know a reference.

Write $\mathcal{SH}(S)^{\text{pure Tate} \geq d}$ for the subcategory generated under filtered colimits, extensions and sums by $\text{Th}_S(\mathcal{O}^n) \simeq S^{2n,n}$ with $n \geq d$. The result we need is as follows.

THEOREM A.1. *We have $\text{MGL} \in \mathcal{SH}(S)^{\text{pure Tate} \geq 0}$. The same is true for the motivic Thom spectra MSL, MSp .*

We need a few preparations for the proof.

LEMMA A.2. *Let $X \in \text{Sm}_S$, $U \subset X$ open with smooth complement $Z \subset X$. Let V be a vector bundle on X . Then there is a cofibration sequence*

$$\text{Th}(V|_U) \rightarrow \text{Th}(V) \rightarrow \text{Th}(V|_Z \oplus N_{Z/X}) \in \mathcal{Spc}(S)_*.$$

Proof. We have

$$\text{Th}(V)/\text{Th}(V|_U) = \frac{V/V \setminus 0}{V|_U/V|_U \setminus 0} \simeq \frac{V}{V \setminus 0 \cup V|_U} = V/V \setminus Z,$$

and that latter space is equivalent to $\text{Th}(N_{Z/V})$ by purity. The short exact sequence (see, e.g., [\[Sta18, Tag 0690\]](#))

$$0 \rightarrow N_{Z/X} \rightarrow N_{Z/V} \rightarrow N_{X/V} \simeq V \rightarrow 0$$

implies the desired result by [\[Hoy17, \(3.26\)\]](#). $\square$

- Denote by $\mathcal{Spc}(S)_*^{\text{puretate} \geq d}$ the smallest full subcategory of $\mathcal{Spc}(S)_*$ that
- (1) contains $*$ and $S^{2n,n}$ for $n \geq d$,
 - (2) is closed under filtered colimits, and
 - (3) if $A \rightarrow B \rightarrow C$ is a *cofiber* sequence with $A, C \in \mathcal{Spc}(S)_*^{\text{puretate} \geq d}$, then $B \in \mathcal{Spc}(S)_*^{\text{puretate} \geq d}$.

THEOREM A.3 (Wendt). *Let S be the spectrum of a field, G a split reductive group, $X = G/P$ a homogeneous space and V a vector bundle on X of rank d . Then $\text{Th}(V) \in \mathcal{Spc}(S)_*^{\text{puretate} \geq d}$.*

Proof. We use the ideas of [Wen10, Prop. 2.2]. The Bruhat decomposition provides a filtration

$$Z_0 = \emptyset \subset Z_1 \subset \cdots \subset Z_n = X$$

by closed subschemes, such that $Z_i \setminus Z_{i-1}$ is a finite disjoint union of affine spaces (see [Wen10, Proof of Prop. 3.7]). Via Lemma A.2, we obtain cofibration sequences

$$\text{Th}(V|_{X \setminus Z_i}) \rightarrow \text{Th}(V|_{X \setminus Z_{i-1}}) \rightarrow \text{Th}(N_{(Z_i \setminus Z_{i-1})/(X \setminus Z_{i-1})} \oplus V|_{Z_i \setminus Z_{i-1}}).$$

For $i = n$, we have $X \setminus Z_i = \emptyset$ and hence the left-hand Thom space is contractible; by induction on i it thus suffices to show that if E is a vector bundle on $Z_i \setminus Z_{i-1}$, then $\text{Th}(E) \in \mathcal{Spc}(S)_*^{\text{puretate} \geq d}$. But $Z_i \setminus Z_{i-1}$ is a disjoint union of affine spaces, so $N \oplus V|_{Z_i \setminus Z_{i-1}}$ is trivial by Quillen–Suslin [Qui76], and the result follows. $\square$

Remark A.4. The assumption that S is the spectrum of a field was used twice in the above proof:

- (1) to know that there is a Bruhat decomposition of G/P into affine space cells, and
- (2) to know that vector bundles on affine spaces are trivial.

For usual symplectic and special linear Grassmannians, (1) holds over $\mathbb{Z}$ (and hence any base). Result (2) will hold over any base S such that the Bass–Quillen conjecture holds (i.e., any vector bundle on $\mathbb{A}_S^n$ is extended from S) and such that all vector bundles on S are trivial, e.g., $S = \text{Spec}(\mathbb{Z})$ (by Lindel’s theorem — see, e.g., [AHW17, Th. 5.2.1]). In fact, in order to establish Theorem A.1 we only need the stable analog. Since Thom spectra of vector bundles only depend on their classes in homotopy K -theory [BH21, Rem. 16.11], this holds over any base.

Proof of Theorem A.1. Since the spectra MGL , MSL , MSp are stable under base change, we may assume that $S = \text{Spec}(\mathbb{Z})$. Each of these spectra is obtained as

$$\text{colim}_i \text{Th}(V_i \ominus \mathcal{O}^{rk(V_i)})$$

for suitable Grassmannians (usual, special linear or symplectic) X_i and vector bundles V_i on them. We have

$$\begin{aligned} \mathrm{Th}(V_i \ominus \mathcal{O}^{rk(V_i)}) &\simeq \Sigma^\infty \mathrm{Th}(V_i) \wedge S^{-2rk(V_i), -rk(V_i)} \\ &\in \Sigma^\infty \mathcal{S}_{\mathrm{pc}}(S)_*^{\mathrm{pure} \geq rk(V_i)} \wedge S^{-2rk(V_i), -rk(V_i)} \\ &\subset \mathcal{SH}(S)^{\mathrm{pure} \geq 0}, \end{aligned}$$

here we have used [Remark A.4](#) for the first containment. This concludes the proof. $\square$

Appendix B. A vanishing result for algebraic cobordism

We are confident that the following result is well known, but we could not locate a reference.

THEOREM B.1. *Let X be essentially smooth over a semi-local principal ideal domain, and let $S \subset \mathbb{N}$ denote the set of positive residue characteristics of X . Then for $i > 0$, we have*

$$\mathrm{MGL}^{2*+i,*}(X)[1/S] = 0.$$

Proof. The slice tower for $\mathrm{MGL}[1/S]$ of the form

$$\mathrm{MGL}[1/S] = f_0(\mathrm{MGL})[1/S] \leftarrow f_1(\mathrm{MGL})[1/S] \leftarrow \cdots$$

has increasing connectivity and subquotients given by $\Sigma^{2*,*}\mathrm{HZ}[1/S] \otimes L_{2*}$ (where L_{2t} is the degree $2t$ part of the Lazard ring). This follows by combining [\[Spi10, proof of Th. 4.7\]](#), [\[Spi18, Th. 11.3\]](#) and [\[SS18, Prop. 3.7\]](#). Induction up to tower thus reduces to proving the same result for motivic cohomology. By construction,

$$H^{p,q}(X, \mathbb{Z}) = \mathrm{CH}^q(X, 2q - p)$$

and hence it is enough to show that $\mathrm{CH}^*(X, -i) = 0$ for $i > 0$. This follows from the fact that $\mathrm{CH}^q(X, -i) = H_{-i}(z^q(X))$, where $z^q(X)$ is a complex concentrated in non-negative degrees [\[Lev01, middle of p.3\]](#). (This is where we use that the base is semi-local.) $\square$

Appendix C. Motivic cohomology

The motivic cohomology of a point with finite coefficients can be computed as a consequence of the Beilinson–Lichtenbaum and Bloch–Kato conjectures. These were established by Voevodsky, Rost and others [\[Voe03a\]](#), [\[Voe11\]](#). For a textbook treatment, see [\[HW19\]](#). We collect some of their results here.

THEOREM C.1. *Let k be a field of exponential characteristic e and $n > 0$ coprime to e .*

- (1) The cycle class map induces an equivalence $L_{\text{ét}}\mathbb{Z}/n(d) \simeq L_{\text{ét}}\mu_n^{\otimes d}$, as complexes of sheaves on Sm_k .
- (2) The induced map $\mathbb{Z}/n(d) \rightarrow L_{\text{ét}}\mu_n^{\otimes d}$ induces an equivalence $\mathbb{Z}/n(d) \simeq \tau_{\geq -d}L_{\text{ét}}\mu_n^{\otimes d}$, the truncation as complexes of Nisnevich sheaves on Sm_k .
- (3) There is a canonical isomorphism $H_{\text{ét}}^m(k, \mu_n^{\otimes m}) \simeq K_m^M(k)/n$.

Proof.

- (1) This is well known; see, e.g., [MVW06, Th. 10.3].
- (2) The case $n = \ell$ is [HW19, Th. C]. The general case follows from this since $\mathbb{Z}/n$ is a sum of iterated extensions of $\mathbb{Z}/\ell$ (for various ℓ).
- (3) By (2), we have $H_{\text{ét}}^m(k, \mu_n^{\otimes m}) \simeq H^{m,m}(k, \mathbb{Z}/m)$. This is the same as $H^{m,m}(k, \mathbb{Z})/n$ by [MVW06, Th. 3.6], which is $K_m^M(k)/n$ by [MVW06, Th. 5.1]. $\square$

COROLLARY C.2. *Using the assumptions above,*

- (1) *We have*

$$\pi_{p,q}(\mathrm{H}\mathbb{Z}/n) \simeq H_{\text{ét}}^{-p}(k, \mu_n^{\otimes -q}) \text{ for } 0 \geq p \geq q, \text{ and } \simeq 0 \text{ else.}$$

- (2) *Suppose that k contains a primitive n -th root of unity. Then*

$$\pi_{*,*}(\mathrm{H}\mathbb{Z}/n) \simeq H_{\text{ét}}^*(k, \mathbb{Z}/n)[\tau] \simeq K_*^M(k)/n[\tau].$$

Here $H_{\text{ét}}^*(k, \mathbb{Z}/n) \simeq K_*^M(k)/n$ is placed in bidegree $(-*, -*)$, and $|\tau| = (0, -1)$.

Proof. In (2) we have $\mu_n^{\otimes r} \simeq \mathbb{Z}/n$ for all r . The element τ corresponds to

$$1 \in H_{\text{ét}}^0(k, \mu_n^{\otimes 1}) \simeq H_{\text{ét}}^0(k, \mathbb{Z}/n) \simeq \mathbb{Z}/n.$$

Both (1) and (2) are now a direct consequences of Theorem C.1. $\square$

References

- [Aok20] K. AOKI, The weight complex functor is symmetric monoidal, *Adv. Math.* **368** (2020), 107145, 10. MR 4082495. Zbl 1440.18027. <https://doi.org/10.1016/j.aim.2020.107145>.
- [AHW17] A. ASOK, M. HOYOIS, and M. WENDT, Affine representability results in $\mathbb{A}^1$ -homotopy theory, I: vector bundles, *Duke Math. J.* **166** no. 10 (2017), 1923–1953. MR 3679884. Zbl 1401.14118. <https://doi.org/10.1215/00127094-0000014X>.
- [Bac18a] T. BACHMANN, Motivic and real étale stable homotopy theory, *Compos. Math.* **154** no. 5 (2018), 883–917. MR 3781990. Zbl 06855294. <https://doi.org/10.1112/S0010437X17007710>.
- [Bac18b] T. BACHMANN, On the conservativity of the functor assigning to a motivic spectrum its motive, *Duke Math. J.* **167** no. 8 (2018), 1525–1571. MR 3807316. Zbl 1390.14064. <https://doi.org/10.1215/00127094-2018-0002>.

- [Bac21] T. BACHMANN, Motivic Tambara functors, *Math. Z.* **297** no. 3-4 (2021), 1825–1852. [MR 4229625](#). [Zbl 07333904](#). <https://doi.org/10.1007/s00209-020-02581-x>.
- [BH21] T. BACHMANN and M. HOYOIS, *Norms in Motivic Homotopy Theory*, *Astérisque* **425**, Soc. Math. France, Paris, 2021. [MR 4288071](#). [Zbl 07403459](#). <https://doi.org/10.24033/ast>.
- [BH18] T. BARTHEL and D. HEARD, Algebraic chromatic homotopy theory for BP_*BP -comodules, *Proc. Lond. Math. Soc.* (3) **117** no. 6 (2018), 1135–1180. [MR 3893176](#). [Zbl 1412.55005](#). <https://doi.org/10.1112/plms.12158>.
- [BHHM08] M. BEHRENS, M. HILL, M. J. HOPKINS, and M. MAHOWALD, On the existence of a v_2^{32} -self map on $M(1, 4)$ at the prime 2, *Homology Homotopy Appl.* **10** no. 3 (2008), 45–84. [MR 2475617](#). [Zbl 1162.55010](#). <https://doi.org/10.4310/HHA.2008.v10.n3.a4>.
- [BS20] M. BEHRENS and J. SHAH, C_2 -equivariant stable homotopy from real motivic stable homotopy, *Ann. K-Theory* **5** no. 3 (2020), 411–464. [MR 4132743](#). [Zbl 07237238](#). <https://doi.org/10.2140/akt.2020.5.411>.
- [BBD82] A. A. BEĬLINSON, J. BERNSTEIN, and P. DELIGNE, Faisceaux pervers, in *Analysis and Topology on Singular Spaces, I* (Luminy, 1981), *Astérisque* **100**, Soc. Math. France, Paris, 1982, pp. 5–171. [MR 0751966](#). [Zbl 0536.14011](#).
- [BI20] E. BELMONT and D. C. ISAKSEN, $\mathbb{R}$ -motivic stable stems, 2020. [arXiv 2001.03606](#).
- [Bon10] M. V. BONDARKO, Weight structures vs. t -structures; weight filtrations, spectral sequences, and complexes (for motives and in general), *J. K-Theory* **6** no. 3 (2010), 387–504. [MR 2746283](#). [Zbl 1303.18019](#). <https://doi.org/10.1017/is010012005jkt083>.
- [BHS20] R. BURKLUND, J. HAHN, and A. SENGER, Galois reconstruction of Artin–Tate $\mathbb{R}$ -motivic spectra, 2020. [arXiv 2010.10325](#).
- [DP83] A. DOLD and D. PUPPE, Duality, trace and transfer, in *Topology* (Moscow, 1979), *Trudy Mat. Inst. Steklov.* **154**, 1983, pp. 81–97. [MR 0733829](#). [Zbl 0556.55006](#).
- [DI05] D. DUGGER and D. C. ISAKSEN, Motivic cell structures, *Algebr. Geom. Topol.* **5** (2005), 615–652. [MR 2153114](#). [Zbl 1086.55013](#). <https://doi.org/10.2140/agt.2005.5.615>.
- [DI10] D. DUGGER and D. C. ISAKSEN, The motivic Adams spectral sequence, *Geom. Topol.* **14** no. 2 (2010), 967–1014. [MR 2629898](#). [Zbl 1206.14041](#). <https://doi.org/10.2140/gt.2010.14.967>.
- [DI17] D. DUGGER and D. C. ISAKSEN, Low-dimensional Milnor-Witt stems over $\mathbb{R}$, *Ann. K-Theory* **2** no. 2 (2017), 175–210. [MR 3590344](#). [Zbl 1400.14064](#). <https://doi.org/10.2140/akt.2017.2.175>.
- [EK20] E. ELMANTO and A. A. KHAN, Perfection in motivic homotopy theory, *Proc. Lond. Math. Soc.* (3) **120** no. 1 (2020), 28–38. [MR 3999675](#). [Zbl 1440.14123](#). <https://doi.org/10.1112/plms.12280>.

- [Ghe18] B. GHEORGHE, The motivic cofiber of τ , *Doc. Math.* **23** (2018), 1077–1127. [MR 3874951](#). [Zbl 1407.55007](#). <https://doi.org/10.4153/cjm-1971-110-9>.
- [GIKR21] B. GHEORGHE, D. C. ISAKSEN, A. KRAUSE, and N. RICKA, $\mathbb{C}$ -motivic modular forms, *J. Eur. Math. Soc. (JEMS)* (2021), Published online first: 2021-11-09. <https://doi.org/10.4171/JEMS/1171>.
- [GWX21] B. GHEORGHE, G. WANG, and Z. XU, The special fiber of the motivic deformation of the stable homotopy category is algebraic, *Acta Math.* **226** no. 2 (2021), 319–407. [MR 4281382](#). [Zbl 07378147](#). <https://doi.org/10.4310/acta.2021.v226.n2.a2>.
- [Goe07] P. GOERSS, The Adams–Novikov spectral sequence and the homotopy groups of spheres, 2007, available on author’s website.
- [HW19] C. HAESEMEYER and C. A. WEIBEL, *The Norm Residue Theorem in Motivic Cohomology*, *Ann. of Math. Stud.* **200**, Princeton Univ. Press, Princeton, NJ, 2019. [MR 3931681](#). [Zbl 1433.14001](#). <https://doi.org/10.1515/9780691189635>.
- [Hil11] M. A. HILL, Ext and the motivic Steenrod algebra over $\mathbb{R}$, *J. Pure Appl. Algebra* **215** no. 5 (2011), 715–727. [MR 2747214](#). [Zbl 1222.55014](#). <https://doi.org/10.1016/j.jpaa.2010.06.017>.
- [Hov04] M. HOVEY, Homotopy theory of comodules over a Hopf algebroid, in *Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-Theory*, *Contemp. Math.* **346**, Amer. Math. Soc., Providence, RI, 2004, pp. 261–304. [MR 2066503](#). [Zbl 1067.18012](#). <https://doi.org/10.1090/conm/346/06291>.
- [Hoy15] M. HOYOIS, From algebraic cobordism to motivic cohomology, *J. Reine Angew. Math.* **702** (2015), 173–226. [MR 3341470](#). [Zbl 1382.14006](#). <https://doi.org/10.1515/crelle-2013-0038>.
- [Hoy17] M. HOYOIS, The six operations in equivariant motivic homotopy theory, *Adv. Math.* **305** (2017), 197–279. [MR 3570135](#). [Zbl 1400.14065](#). <https://doi.org/10.1016/j.aim.2016.09.031>.
- [HKØ17] M. HOYOIS, S. KELLY, and P. A. ØSTVÆR, The motivic Steenrod algebra in positive characteristic, *J. Eur. Math. Soc. (JEMS)* **19** no. 12 (2017), 3813–3849. [MR 3730515](#). [Zbl 1386.14087](#). <https://doi.org/10.4171/JEMS/754>.
- [HK01] P. HU and I. KRIZ, Real-oriented homotopy theory and an analogue of the Adams–Novikov spectral sequence, *Topology* **40** no. 2 (2001), 317–399. [MR 1808224](#). [Zbl 0967.55010](#). [https://doi.org/10.1016/S0040-9383\(99\)00065-8](https://doi.org/10.1016/S0040-9383(99)00065-8).
- [HKO11] P. HU, I. KRIZ, and K. ORMSBY, Remarks on motivic homotopy theory over algebraically closed fields, *J. K-Theory* **7** no. 1 (2011), 55–89. [MR 2774158](#). [Zbl 1248.14026](#). <https://doi.org/10.1017/is010001012jkt098>.
- [Isa19] D. C. ISAKSEN, *Stable Stems*, **262** Mem. Amer. Math. Soc. no. 1269, Amer. Math. Soc., 2019. [MR 4046815](#). [Zbl 1454.55001](#). <https://doi.org/10.1090/memo/1269>.

- [IWX20a] D. C. ISAKSEN, G. WANG, and Z. XU, More stable stems, 2020. [arXiv 2001.04511](#).
- [IWX20b] D. C. ISAKSEN, G. WANG, and Z. XU, Stable homotopy groups of spheres, *Proc. Natl. Acad. Sci. USA* **117** no. 40 (2020), 24757–24763. [MR 4250190](#). <https://doi.org/10.1073/pnas.2012335117>.
- [Kra18] A. KRAUSE, Periodicity in motivic homotopy theory and over BP_*BP , 2018, Bonn University thesis. Available at <https://hdl.handle.net/20.500.11811/7592>.
- [Kyl19] J. I. KYLLING, Algebraic cobordism of number fields, 2019. [arXiv 1901.04031](#).
- [Lev01] M. LEVINE, Techniques of localization in the theory of algebraic cycles, *J. Algebraic Geom.* **10** no. 2 (2001), 299–363. [MR 1811558](#). [Zbl 1077.14509](#).
- [LYZR19] M. LEVINE, Y. YANG, G. ZHAO, and J. RIOU, Algebraic elliptic cohomology theory and flops I, *Math. Ann.* **375** no. 3-4 (2019), 1823–1855. [MR 4023393](#). [Zbl 1433.14015](#). <https://doi.org/10.1007/s00208-019-01880-x>.
- [Lur09] J. LURIE, *Higher Topos Theory*, *Ann. of Math. Stud.* **170**, Princeton Univ. Press, Princeton, NJ, 2009. [MR 2522659](#). [Zbl 1175.18001](#). <https://doi.org/10.1515/9781400830558>.
- [Lur17] J. LURIE, Higher algebra, September 2017, available on author’s webpage.
- [Lur18] J. LURIE, Spectral algebraic geometry, February 2018, available on author’s webpage.
- [Man21] L. MANTOVANI, Localizations and completions of stable ∞ -categories, 2021. [arXiv 2105.02285](#).
- [MNN17] A. MATHEW, N. NAUMANN, and J. NOEL, Nilpotence and descent in equivariant stable homotopy theory, *Adv. Math.* **305** (2017), 994–1084. [MR 3570153](#). [Zbl 1420.55024](#). <https://doi.org/10.1016/j.aim.2016.09.027>.
- [MVW06] C. MAZZA, V. VOEVODSKY, and C. WEIBEL, *Lecture Notes on Motivic Cohomology*, *Clay Mathematics Monographs* **2**, Amer. Math. Soc., Providence, RI, 2006. [MR 2242284](#). [Zbl 1115.14010](#).
- [Mil75] H. R. MILLER, *Some Algebraic Aspects of the Adams–Novikov Spectral Sequence*, ProQuest LLC, Ann Arbor, MI, 1975, Thesis (Ph.D.)–Princeton University. [MR 2625232](#).
- [Mor85] J. MORAVA, Noetherian localisations of categories of cobordism comodules, *Ann. of Math.* (2) **121** no. 1 (1985), 1–39. [MR 0782555](#). [Zbl 0572.55005](#). <https://doi.org/10.2307/1971192>.
- [Mor06] F. MOREL, $\mathbb{A}^1$ -algebraic topology, in *International Congress of Mathematicians. Vol. II*, Eur. Math. Soc., Zürich, 2006, pp. 1035–1059. [MR 2275634](#). [Zbl 1097.14014](#).
- [NSØ09] N. NAUMANN, M. SPITZWECK, and P. A. ØSTVÆR, Motivic Landweber exactness, *Doc. Math.* **14** (2009), 551–593. [MR 2565902](#). [Zbl 1230.55005](#).

- [Nee01] A. NEEMAN, *Triangulated Categories*, *Ann. of Math. Stud.* **148**, Princeton Univ. Press, Princeton, NJ, 2001. MR 1812507. Zbl 0974.18008. <https://doi.org/10.1515/9781400837212>.
- [NZ06] A. NENASHEV and K. ZAINOULLINE, Oriented cohomology and motivic decompositions of relative cellular spaces, *J. Pure Appl. Algebra* **205** no. 2 (2006), 323–340. MR 2203620. Zbl 1101.14025. <https://doi.org/10.1016/j.jpaa.2005.06.021>.
- [NS17] T. NIKOLAUS and S. SAGAVE, Presentably symmetric monoidal ∞ -categories are represented by symmetric monoidal model categories, *Algebr. Geom. Topol.* **17** no. 5 (2017), 3189–3212. MR 3704256. Zbl 1380.55018. <https://doi.org/10.2140/agt.2017.17.3189>.
- [Nov67] S. P. NOVIKOV, Methods of algebraic topology from the point of view of cobordism theory, *Izv. Akad. Nauk SSSR Ser. Mat.* **31** no. 4 (1967), 855–951. MR 0221509. Zbl 0169.54503. <https://doi.org/10.1070/IM1967v001n04ABEH000591>.
- [Pst18] P. PSTRĄGOWSKI, Synthetic spectra and the cellular motivic category, 2018. [arXiv 1803.01804](https://arxiv.org/abs/1803.01804).
- [Qui76] D. QUILLEN, Projective modules over polynomial rings, *Invent. Math.* **36** (1976), 167–171. MR 0427303. Zbl 0337.13011. <https://doi.org/10.1007/BF01390008>.
- [Rie14] E. RIEHL, *Categorical Homotopy Theory*, *New Math. Monogr.* **24**, Cambridge Univ. Press, Cambridge, 2014. MR 3221774. Zbl 1317.18001. <https://doi.org/10.1017/CBO9781107261457>.
- [Rio05] J. RIOU, Dualité de Spanier-Whitehead en géométrie algébrique, *C. R. Math. Acad. Sci. Paris* **340** no. 6 (2005), 431–436. MR 2135324. Zbl 1068.14021. <https://doi.org/10.1016/j.crma.2005.02.002>.
- [RSØ19] O. RÖNDIGS, M. SPITZWECK, and P. A. ØSTVÆR, The first stable homotopy groups of motivic spheres, *Ann. of Math. (2)* **189** no. 1 (2019), 1–74. MR 3898173. Zbl 1406.14018. <https://doi.org/10.4007/annals.2019.189.1.1>.
- [SS18] J. SCHMIDT and F. STRUNK, Stable $\mathbb{A}^1$ -connectivity over Dedekind schemes, *Ann. K-Theory* **3** no. 2 (2018), 331–367. MR 3781430. Zbl 1423.14158. <https://doi.org/10.2140/akt.2018.3.331>.
- [SS03] S. SCHWEDE and B. SHIPLEY, Stable model categories are categories of modules, *Topology* **42** no. 1 (2003), 103–153. MR 1928647. Zbl 1013.55005. [https://doi.org/10.1016/S0040-9383\(02\)00006-X](https://doi.org/10.1016/S0040-9383(02)00006-X).
- [Sec20] P. SECHIN, Landweber-equivariant Grothendieck motives: a toy example of non-oriented phenomena, 2020, Seminar on $\mathbb{A}^1$ -topology, motives and K-theory. Available at <https://indico.eimi.ru/event/127/>.
- [Şeg01] L. M. ŞEGA, Homological properties of powers of the maximal ideal of a local ring, *J. Algebra* **241** no. 2 (2001), 827–858. MR 1843329. Zbl 0998.13005. <https://doi.org/10.1006/jabr.2001.8786>.
- [Sos19] V. SOSNILO, Theorem of the heart in negative K-theory for weight structures, *Doc. Math.* **24** (2019), 2137–2158. MR 4033821. Zbl 1441.18008.

- [Spi10] M. SPITZWECK, Relations between slices and quotients of the algebraic cobordism spectrum, *Homology Homotopy Appl.* **12** no. 2 (2010), 335–351. MR 2771593. Zbl 1209.14019. <https://doi.org/10.4310/HHA.2010.v12.n2.a11>.
- [Spi18] M. SPITZWECK, A commutative $\mathbb{P}^1$ -spectrum representing motivic cohomology over Dedekind domains, *Mém. Soc. Math. Fr. (N.S.)* no. 157 (2018), 110. MR 3865569. Zbl 1408.14081. <https://doi.org/10.24033/msmf.465>.
- [Spi20] M. SPITZWECK, Algebraic cobordism in mixed characteristic, *Homology Homotopy Appl.* **22** no. 2 (2020), 91–103. MR 4093171. Zbl 1440.14126. <https://doi.org/10.4310/hha.2020.v22.n2.a5>.
- [Sta18] THE STACKS PROJECT AUTHORS, Stacks project, 2018. Available at <https://stacks.math.columbia.edu>.
- [Voe03a] V. VOEVODSKY, Motivic cohomology with $\mathbf{Z}/2$ -coefficients, *Publ. Math. Inst. Hautes Études Sci.* no. 98 (2003), 59–104. MR 2031199. Zbl 1057.14028. <https://doi.org/10.1007/s10240-003-0010-6>.
- [Voe03b] V. VOEVODSKY, Reduced power operations in motivic cohomology, *Publ. Math. Inst. Hautes Études Sci.* no. 98 (2003), 1–57. MR 2031198. Zbl 1057.14027. <https://doi.org/10.1007/s10240-003-0009-z>.
- [Voe11] V. VOEVODSKY, On motivic cohomology with $\mathbf{Z}/l$ -coefficients, *Ann. of Math.* (2) **174** no. 1 (2011), 401–438. MR 2811603. Zbl 1236.14026. <https://doi.org/10.4007/annals.2011.174.1.11>.
- [Wen10] M. WENDT, More examples of motivic cell structures, 2010. arXiv 1012.0454.
- [WØ17] G. M. WILSON and P. A. ØSTVÆR, Two-complete stable motivic stems over finite fields, *Algebr. Geom. Topol.* **17** no. 2 (2017), 1059–1104. MR 3623682. Zbl 1361.14020. <https://doi.org/10.2140/agt.2017.17.1059>.
- [Xu17] Z. XU, Conference talk “Computing stable homotopy groups of spheres” at Homotopy Theory: Tools and Applications, 2017, University of Illinois at Urbana-Champaign.

(Received: January 26, 2021)

(Revised: August 31, 2021)

MATHEMATISCHES INSTITUT, LMU MUNICH, MUNICH, GERMANY

E-mail: tom.bachmann@zoho.com

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ, USA

E-mail: hanajk@ias.edu

SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, FUDAN UNIVERSITY,
SHANGHAI, CHINA

E-mail: wangguozhen@fudan.edu.cn

UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093, USA

E-mail: xuzhouli@ucsd.edu