DOI: 10.1111/gcb.16033

#### REVIEW



## Socio-ecological drivers of multiple zoonotic hazards in highly urbanized cities

Matthew A. Combs<sup>1</sup> | Pallavi A. Kache<sup>1</sup> | Meredith C. VanAcker<sup>1</sup> | Nichar Gregory<sup>1</sup> | Laura D. Plimpton<sup>1</sup> | Danielle M. Tufts<sup>1,2</sup> | Maria P. Fernandez<sup>1,3</sup> Maria A. Diuk-Wasser<sup>1</sup>

#### Correspondence

Maria A. Diuk-Wasser, Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, NY 10027, USA.

## Email: mad2256@columbia.edu

#### **Funding information**

National Center for Emerging and Zoonotic Infectious Diseases, Grant/ Award Number: U01CK000509-01: National Science Foundation Dynamics of Integrated Socio-Ecological Systems (CNH2) Program, Grant/Award Number: 1924061

[Correction added on 9-February-2022, after first online publication: The copyright line was changed.]

#### **Abstract**

The ongoing COVID-19 pandemic is a stark reminder of the devastating consequences of pathogen spillover from wildlife to human hosts, particularly in densely populated urban centers. Prevention of future zoonotic disease is contingent on informed surveillance for known and novel threats across diverse human-wildlife interfaces. Cities are a key venue for potential spillover events because of the presence of zoonotic pathogens transmitted by hosts and vectors living in close proximity to dense human settlements. Effectively identifying and managing zoonotic hazards requires understanding the socioecological processes driving hazard distribution and pathogen prevalence in dynamic and heterogeneous urban landscapes. Despite increasing awareness of the human health impacts of zoonotic hazards, the integration of an eco-epidemiological perspective into public health management plans remains limited. Here we discuss how landscape patterns, abiotic conditions, and biotic interactions influence zoonotic hazards across highly urbanized cities (HUCs) in temperate climates to promote their efficient and effective management by a multi-sectoral coalition of public health stakeholders. We describe how to interpret both direct and indirect ecological processes, incorporate spatial scale, and evaluate networks of connectivity specific to different zoonotic hazards to promote biologically-informed and targeted decision-making. Using New York City, USA as a case study, we identify major zoonotic threats, apply knowledge of relevant ecological factors, and highlight opportunities and challenges for research and intervention. We aim to broaden the toolbox of urban public health stakeholders by providing ecologicallyinformed, practical guidance for the evaluation and management of zoonotic hazards.

#### KEYWORDS

hazard, landscape, management, mesomammal, mosquito, rodent, spillover, tick, urban, zoonosis

### 1 | INTRODUCTION

Zoonotic diseases, caused by pathogens transmitted between human and wildlife populations, most often emerge at human-wildlife interfaces where there is an increased likelihood of direct or indirect contact between people and infectious hosts and/or vectors (Hassell et al., 2017; Soulsbury & White, 2015). Urban landscapes are increasingly recognized as habitat for wildlife across both green

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Global Change Biology published by John Wiley & Sons Ltd.

<sup>&</sup>lt;sup>1</sup>Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, New York, USA

<sup>&</sup>lt;sup>2</sup>Infectious Diseases and Microbiology Department, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

<sup>&</sup>lt;sup>3</sup>Paul G. Allen School for Global Animal Health, Washington State University, Pullman, Washington, USA

spaces (e.g., vegetated or natural landcover) and gray spaces (e.g., built structures, impervious surface, and associated infrastructure) leading to increased interactions with human populations (Deplazes et al., 2004; Hansford et al., 2017; Mckinney, 2008; Rothenburger et al., 2017). Globally, cities are growing both in their number of inhabitants-with over 68% of the world's populations expected to be urban-dwelling by 2080 (United Nations, 2018), and in their geographic footprint—with urban land cover steadily increasing (Seto et al., 2012). The process of urbanization contributes to biotic homogenization (McKinney, 2006), invasive species introductions (Blair, 1996; Shochat et al., 2010), and results in a wealth of exploitable resources, which promotes the increased richness and abundance of zoonotic host species and their associated pathogens (Gibb et al., 2020). The COVID-19 pandemic highlighted the risk of zoonotic disease emergence and the unique vulnerability of cities to emerging zoonotic pathogens, emphasizing concerns regarding effective prevention and response to spillover events (Alirol et al., 2011; Bradley & Altizer, 2007; Mackenstedt et al., 2015; Neiderud, 2015). However, there is a persistent need to understand how urbanization alters ecological processes that underlie human zoonotic risk and spillover potential in cities (Karesh et al., 2012) (Figure 1).

Recent discussions of urban zoonoses have focused on factors influencing spillover events (Hassell et al., 2017; Plowright et al., 2017, 2021), pathogen-landscape interactions (Eisenberg et al., 2007; Lambin et al., 2010), and the impacts of urbanization on animal health (Bengis et al., 2004; Bradley & Altizer, 2007; Mackenstedt et al., 2015). The emphasis of these reviews is on the combined influence of environmental change, wildlife biology, and human risk

factors (including exposure and vulnerability, see Table 1 for glossary of terms) to broadly illustrate the drivers of zoonotic disease emergence. However, a detailed assessment of the ecological drivers of zoonotic hazards is needed to improve pre-spillover surveillance, research, and management of hazards in cities. Such efforts require collaboration between public health practitioners, wildlife biologists, urban planners, community leaders and others invested in the health of local human communities (hereafter "public health stakeholders") that often go beyond the scope of a single entity. Multi-sectoral partnerships have shown promise for tackling chronic physical and mental health issues and enabling action on climate change adaptation and mitigation (Ramaswami et al., 2016). Similar efforts are essential to prevent urban zoonotic disease, and begin with a thorough understanding of the foundational biology and socio-ecological interactions driving the presence and prevalence of urban zoonotic hazards.

While drivers of urban zoonoses have been mostly studied in tropical regions where health impacts are greatest (Gottdenker et al., 2014; White & Razgour, 2020), the emergence and persistence of these diseases has been overlooked in highly urbanized cities (HUCs) in temperate climates, which also support a diverse suite of zoonotic hazards. These often include human commensal rodents and mesomammals (Feng & Himsworth, 2014; Plumer et al., 2014; Tufts et al., 2021), *Ixodidae* ticks (Adalsteinsson et al., 2018; Mancini et al., 2014; Steere, 1994), *Aedes* spp., and *Culex* spp. mosquitoes (Calhoun et al., 2007; Goodman et al., 2018; Muir & Kay, 1998; Shragai & Harrington, 2019), all of which can harbor and transmit zoonotic pathogens. In our definition, HUCs are those with more than 500 k




FIGURE 1 Socio-ecological drivers influencing zoonotic hazards in HUCs. (a) Individual drivers illustrated across a simplified urban landscape and color coded based on whether they are predominantly related to landscape (blue), abiotic (purple), or biotic (green) factors. (b) Thematic flow chart illustrating how socio-ecological processes drive the intensity of zoonotic hazard, a foundational component of zoonotic risk, which ultimately impacts the likelihood of zoonotic spillover events [Colour figure can be viewed at wileyonlinelibrary.com]

TABLE 1 Glossary of terms

Term	Definition		
Zoonotic disease	Any disease caused by a pathogen transmitted between non-human animals and humans		
Hazard	Potential source of harm (e.g., infected host or vector, shed pathogens) expected to contribute to zoonotic disease, at varying intensities over space and time		
Exposure	Likelihood of human contact with hazards from interaction with the environment		
Vulnerability	Human or societal condition altering the likelihood of harm, given exposure		
Risk	The likelihood of adverse outcomes caused by a hazard, given exposure and vulnerability		
Spillover	Event in which zoonotic pathogen enters the human population		
Reservoir Host	A species competent to harbor a particular pathogen, such that it sustains the pathogen in the environment and serves as a source of human or vector infection		
Vector	An organism capable of transmitting a pathogen between non-human animals and humans		
Pathogen	An infectious agent (bacteria, virus, endoparasite, or microorganism) capable of causing disease		
Eco-epidemiology	The study of human disease that incorporates information from human populations and societies as well as environmental and biological factors		
Ecosystem Disservice	Any consequence of interacting ecological factors or agents that lead to negative outcomes for human and economic well-being		
Dilution Effect	Theory that increasing biodiversity results in decreased human disease risk due to increased abundance of lower-competence hosts		

residents, citywide sanitation infrastructure, life expectancy above 75 years, as well as a prolonged and pervasive historical conversion of earthen land cover to impervious surfaces. Here, we focus on the ecological conditions and emergent processes accompanying urbanization that drive differences in hazard intensity across HUCs in temperate climates through (1) the distribution and abundance of hosts and vectors and (2) patterns of pathogen prevalence across hosts, vectors, and the environment.

Each HUC supports multiple zoonotic hosts and vectors that experience the heterogeneity of cities differently based on their biological traits and habitat requirements, creating divergent outcomes for the pathogens they harbor and transmit across space, within or among species. Understanding and managing these hazards requires acknowledging several key concepts highlighted throughout this perspective article. (1) There is an inherent ecological tradeoff between green or gray infrastructure and the presence of zoonotic hazards (Diuk-Wasser et al., 2021; Jennings et al., 2019; Lõhmus & Balbus, 2015; Soulsbury & White, 2015; Taguchi et al., 2020). While urban green spaces provide ecosystem services, including climate mitigation and physical and mental health benefits (Gómez-Baggethun & Barton, 2013; Gregory McPherson, 1992; Luederitz et al., 2015), these same spaces produce ecosystem disservices when they increase human exposure to pathogens and act as venues for zoonotic spillover (Shackleton et al., 2016; Vanwambeke et al., 2019). Efforts to increase urban biodiversity or human well-being through urban greening (Fuller et al., 2007) may also alter disease risk, although

biodiversity-disease relationships remain widely debated (Ogden & Tsao, 2009; Randolph & Dobson, 2012; Rohr et al., 2020; Salkeld et al., 2013). Similarly, while gray spaces provide human communities and economies room to grow, these spaces support zoonotic hazards that we experience with more direct spatial and temporal overlap. (2) Urban zoonotic hazards are a component of socio-ecological systems; environmental conditions in cities (including the distribution and prevalence of hazards) are co-produced by the interaction of biophysical and social processes (McGinlay et al., 2016). A city's greening or graying process depends on its history, socio-cultural values, and socio-economic drivers of land use (Des Roches et al., 2020; Pickett et al., 2001). Thus, effective and equitable prevention of urban zoonoses is facilitated by an integrated socio-ecological framework for understanding disease emergence. (3) Plans to study and manage hazards should be species-specific. Because of differences in their spatial scale of movement, each host and vector species experiences a unique pattern of landscape connectivity, which depends on traits such as migration propensity, dispersal distance, resource needs, physiological constraints, and response to disturbance and fluctuating population densities (Jetz et al., 2004; Peters et al., 2019; Tucker et al., 2014). Research and management strategies should reflect the scale at which sampling is most informative and management efforts are most effective.

In this article, we first describe how characteristics of urban socio-ecological landscapes, in tandem with their unique abiotic conditions and biotic interactions, determine zoonotic hazards distribution and prevalence. Then using New York City (NYC), USA as a case study, we apply these concepts to evaluate different zoonotic hazards and spillover prevention strategies that incorporate the city's ecological context. Throughout, we discuss implications for pathogen surveillance, research, and mitigation to improve understanding and outcomes of urban zoonotic emergence and persistence (Box 1).

#### 2 | LANDSCAPE CHARACTERISTICS

The intensity of zoonotic hazards exhibits strong spatial heterogeneity across HUCs. Effectively evaluating and managing these hazards requires an understanding of how landscape patterns influence

# BOX 1 Priority areas for strategic surveillance, research, and mitigation of urban zoonotic hazards

#### (1) Surveillance:

- a. Identify baseline distribution and abundance of zoonotic hosts and vectors across diverse urban sites (i.e., land use, wildlife community, vegetation characteristics, history, socioeconomic status); balance financial limitations with long-term active samplings of key sites and passive sampling through community reporting.
- b. Leverage molecular approaches to identify variable pathogen prevalence in hosts, vectors, and environment.
- Monitor changes in hazard intensity linked to dynamic urban landscape (i.e., greening or degreening of neighborhoods, development).

#### (2) Research:

- a. Establish multi-sectoral collaborations with diverse public health stakeholders (municipal departments, epidemiologists, community leaders, disease ecologists, social scientists, non-profit groups) to study influence of socio-ecological landscape attributes, abiotic conditions, biotic interactions across local multihazard suite.
- b. Create and maintain public data repositories on host, vector, and pathogen distribution, and fine-scale spatial environmental data.

#### (3) Prevention:

- a. Develop ecologically-informed management strategies by mathematical modeling of habitat, pathogen dynamics, and host/vector connectivity; prioritize interventions on prevalence hotspots in conjunction with societal risk factors.
- b. Implement management strategies at spatial scale relevant to species-specific biology and dispersal.

populations of hosts, vectors, and associated pathogens as they interact with one another and the environment. In particular, it is important to address the influence of landscape composition, configuration, and connectivity on specific hazards, and identify the spatial scale at which eco-epidemiological processes take place. But understanding current landscape patterns and their influence on zoonotic hazards necessitates knowledge of their historical and ongoing socio-ecological drivers.

Urban development is driven by natural ecological processes and biophysical features such as topography, hydrology, and native vegetation, in combination with pervasive social forces (Des Roches et al., 2020; Keeler et al., 2019). Indeed, widespread land cover conversion and patterns of fragmentation among HUCs are largely driven by societal values informed by culture, economics, and local history. During early stages of temperate HUC growth, the adoption of Eurocentric ideals of nature and aesthetics promoted gridded urban forms and open lawns, which greatly contributed to their biotic homogenization (Ignatieva & Stewart, 2009; Loughran, 2020; Shackleton & Gwedla, 2021). During post-industrial growth, systematic racism and classism was institutionalized in many cities through land use policy, development projects, and lending decisions (Schell et al., 2020), which was driven in large part by capitalist systems that prioritized profit and exploited the a-spatiality of marginalized communities (Bledsoe & Wright, 2019). These legacies drive current patterns of habitat suitability for hazards through unequal distribution of green space as well as investment in infrastructure and services across racial- and class-based lines (Schell et al., 2020; Venter et al., 2020). For example, brown rat habitat and elevated pathogen prevalence among adult mosquitos are consistently correlated with low-income areas (Dowling et al., 2013; Harrigan et al., 2010; Johnson et al., 2016; LaDeau et al., 2013; Masi et al., 2010; Rothenburger et al., 2017). Social factors, such as income, education, and employment, also affect people's interactions with the environment through their impact on personal exposure, vulnerability, and coping capacity, but these risk factors are more fully described elsewhere (Hosseini et al., 2017; Solar & Irwin, 2014). We focus here on how public health stakeholders can account for socio-ecologically complex landscape attributes such as composition, configuration, connectivity, and scale in their study and management of zoonotic hazards (Frank et al., 2017; Ostrom, 2018).

Landscape composition describes the identity of patch types and their relative geographic coverage (Ostfeld et al., 2005). The land cover attributes used to define composition and identify suitable niche space may differ for each reservoir host or vector species. For instance, variation in the vegetative understory in patches of urban green space impacts the presence of both mice and ticks (Adalsteinsson et al., 2018), altering local Lyme disease (LD) hazard. Thus, satellite data describing canopy cover may not be as informative as vegetative surveys that directly sample variation in relevant tick and host habitat. Similarly, urban gray spaces differ dramatically in their intensity and type of human use and maintenance, and these characteristics can change between and within neighborhoods. For instance, increasing abandonment rates within neighborhoods

correlates to increases in the abundance of rodent hosts and fitness of mosquito vectors (Katz et al., 2020; Peterson et al., 2020a). Such responses impact public health outcomes for spatially adjacent communities (Gulachenski et al., 2016), with homeless populations often experiencing the highest exposure rates to hosts and vectors (Leibler et al., 2018). Identifying zoonotic host habitat in gray spaces may be possible through evaluation of easily recorded, physical attributes like building age or land use; however, ephemeral qualities should also be considered. For example, the likelihood of standing water or solid waste buildup may influence mosquito-borne pathogens and leptospirosis transmission (Murray et al., 2020) by increasing mosquito and commensal rodent presence (Krystosik et al., 2020), respectively. Further, in evaluating when and where to perform surveillance or interventions it is important to consider temporal changes due to seasonality, maintenance frequency, or longer trends of de-urbanization and re-greening (Eskew & Olival, 2018), as well as links between broad-scale habitat attributes and fine-scale patterns of landscape composition.

Landscape configuration reflects the spatial arrangement of patches and their proximity or isolation to one another. In HUCs, patch configuration depend on land use and zoning policies, as well as historical contingencies and natural topography or hydrology that drive local decision making. The degree of landscape fragmentation is metric of configuration that is sometimes used as a proxy for biological community composition and increased potential for zoonotic hazards (Allan et al., 2003; Brock et al., 2019; Diuk-Wasser et al., 2021; Zolnik et al., 2015). Increased landscape fragmentation is associated with increased edges between green and gray spaces that influence hazard distribution and creates increased opportunities for human-wildlife interactions (Barding & Nelson, 2008), For example, in gray spaces, adult mosquitos fly through the landscape (Morlan & Hayes, 1958; Muir & Kay, 1998) and rodents migrate within and between city blocks (Byers et al., 2019), putting residents at risk simply by living within host and/or vector dispersal distance of habitat supporting hazards. While evidence for the impact of green space edges in forests, parks, or yards on vector abundance or pathogen prevalence is mixed (Finch et al., 2014; Hansford et al., 2017; Horobik et al., 2006), assessing the ecology of spillover events at the transition zone between different land covers should be a research priority.

Landscape connectivity results from the integration of composition and configuration and describes both the network of habitable patches available (structural connectivity) and the realized movement of organisms and gene flow among those patches (functional connectivity) (Brooks, 2003). Incorporating connectivity into urban zoonoses mitigation is important because networks of connected habitat permit the flow of pathogens as hosts and vectors move through cities among suitable patches (Ostfeld et al., 2005). For example, taking a species-specific view, habitat connectivity for white-tailed deer based on green space coverage strongly predicts the density and infection prevalence of ticks, and thus LD risk in New York City (Vanacker et al., 2019). In HUCs, linear features like railways, greenbelts, and riverbanks provide movement corridors

allowing for rapid dissemination of pathogens across landscapes, suggesting zoonotic disease emergence can occur rapidly via a small number of linked habitat patches. Behaviorally-flexible species like red foxes often use urban infrastructure in unintended ways, for instance by dispersing along roadways (Kimmig et al., 2020), effectively introducing parasitic infections and occasionally the rabies virus to new areas (Mackenstedt et al., 2015; Plumer et al., 2014; Smith et al., 2003). Given the idiosyncratic ways in which different wildlife hosts and vectors navigate urban landscapes, a key research priority should be to understand movement behavior and connectivity for local zoonotic threats through observations, tracking, and spatial modeling (Deplazes et al., 2004; Hemming-Schroeder et al., 2018; Heylen et al., 2019; Richardson et al., 2017).

Networks of habitat connectivity may result in either spatial concentration or spread of zoonotic hosts and their pathogens. This outcome depends on species-specific responses to the distribution of landscape features and should be incorporated into models of zoonotic risk and the development of hazard management programs. Movement barriers reduce connectivity by limiting species movement through physiological limitations (e.g., inability to cross waterways, impervious surface) or behavioral avoidance (e.g., road traffic, noise) (Clark et al., 2010; Fusco et al., 2021; Munshi-South, 2012). Typically, habitat isolation reduces the invasion potential of pathogens, but if pathogens generate immune-dependent responses in host populations, patch isolation can increase future susceptibility. For example, fruit bats forming high-density groups in urban Australia experienced large outbreaks of Hendra virus when decreased connectivity among patches reduced local population immunity (Plowright et al., 2011). Connectivity networks also influence the formation of, and dispersal from, pathogen hotspots, where local hazard intensity is significantly higher than baseline levels, increasing spillover potential (Paull et al., 2012). Prevalence of Leptospira interrogans and Bartonella tribocorum in urban rats is driven in part by reduced movement among city blocks, leading to high contact rates among high-density groups (Byers et al., 2020). This suggests that zoonotic hazards can persist without widespread movement when infections are endemic and contact rate is high among hosts.

The spatial scale at which the composition, configuration, and connectivity of landscapes is assessed is central to understanding and managing zoonotic hazards (Fischhoff et al., 2019; McGarigal et al., 2016; Richardson et al., 2016). The spatial scale is composed of the scope (i.e., spatial bounds) and grain (i.e., resolution of observed detail) and it must match the resource needs and movement patterns of the particular host(s) or vector(s) (Cushman et al., 2016). For instance, larval mosquito habitat requires only small amounts of standing water that may be perceptible only when composition is characterized at very fine spatial grain, while large mammals or mesomammal host habitat requires larger vegetated patches that are observable at coarser spatial grain. In HUCs, the relevant scale for effective management may differ from the scale at which species sampling or treatments are most practical, creating additional economic and policy hurdles for stakeholders. Brown rats, for example, are often managed across individual properties despite ample

evidence that block-level or neighborhood-level management is required to limit host and pathogen presence (Byers et al., 2019, 2020; Combs et al., 2019). Dispersal distance is a key characteristic influencing the potential for human-wildlife interactions, thus spatial grain of analyses must be smaller than this distance to accurately encompass movement dynamics into management approaches (Cushman et al., 2016). A wealth of high-resolution spatial data describing human population, wildlife, and the environment is often available for HUCs, creating tremendous opportunities to assess the links between urban form, biodiversity, and zoonotic disease emergence at appropriate spatial scales.

#### 3 | ABIOTIC CONDITIONS

Landscape changes driven by urbanization significantly alter abiotic conditions, including temperature, light, noise, heavy metal pollution, and hydrology, each of which have downstream effects on wildlife hosts, arthropod vectors, and their associated pathogens (Alberti, 2005). The widespread presence of impervious surfaces (e.g., roads, buildings, parking lots) is a defining characteristic of HUCs. This graying of urban landscapes drastically alters abiotic conditions and presents ecological tradeoffs regarding the presence and intensity of zoonotic hazards, and their impacts on human health. While impervious surfaces may act as a buffer against pathogen transmission when hosts and vectors are restricted to green space, these land cover types support the densification of human settlements, which can increase hazard presence and opportunities for zoonotic spillover by providing high-quality habitat for a suite of urban-adapted commensal rodents and mosquito vectors (Baiwa. 2018; Feng & Himsworth, 2014).

High impervious surface coverage also leads to increased ambient temperatures in urban landscapes, known as the "heat island" effect (Rizwan et al., 2008). Elevated temperature in cities has a strong direct effect on populations of small ectothermic arthropods (LaDeau et al., 2015; Youngsteadt et al., 2017), including observed increases in Aedes spp. mosquitos (de Azevedo et al., 2018) and increasing overwintering survival of Phlebotomus sand flies, the vector for leishmaniasis (Trajer et al., 2014). Heat island effects may have a strong impact in temperate HUCs specifically, by increasing the number of days above threshold activity temperatures for vectors like Ixodes ticks, though this mechanism requires further study (Duffy & Campbell, 1994; Gray, 2008). Underground infrastructure in cities also provides a geothermally stable environment that supports arthropod and rodent populations year-round (Bajomi et al., 2013; Byrne & Nichols, 1999; Channon et al., 2006). Green infrastructure (e.g., parks, green roofs, plantings) offers climate mitigation for humans while also providing thermal refugia for vectors by buffering against extreme temperatures (Venter et al., 2020), presenting a tradeoff between ecosystem services and disservices. Cities should seek to better characterize the thermal landscape at scales relevant to hosts and vectors of concern, as well as integrate future warming conditions

due to climate change and expected consequences on species activity and range shifts (Gray, 2008; Ryan et al., 2018).

Impervious surfaces can also drastically influence hydrology by reducing drainage and restructuring watersheds (Pickett et al., 2001; Shuster et al., 2005). At broad spatial scales, this can increase the risk of flooding events, which is correlated with Leptospira spp. infection in humans (Naing et al., 2019) and contaminated drinking water, though this phenomenon is more likely in cities lacking robust sanitation systems (Ko et al., 1999; Lau et al., 2010; Rydin et al., 2012). Impervious surfaces also cause environmentally acquired pathogens to concentrate downstream along urban watersheds, increasing hazard exposure (Mallin et al., 2000; VanWormer et al., 2016). At fine spatial scales, impervious surfaces increase opportunities for standing water, which provides breeding habitat for urban mosquitoes and has been correlated with increased Leptospira spp. infection in brown rats due to environmental persistence of bacteria shed in urine of infected hosts (Murray et al., 2020).

Urban environments can alter urban wildlife behavior and hostparasite interactions through light and noise pollution (Francis et al., 2011; Francis et al., 2015; Singh et al., 2014). Migrating birds are attracted to regions with more artificial light at night, which may result in increased deposition of bird-fed ticks and associated pathogens around urban areas (Brinkerhoff et al., 2011; Ogden et al., 2008). Light and noise have variable effects on arthropod activity in nontemperate cities (McMahon et al., 2017; Pacheco-Tucuch et al., 2012) and alter gene expression in Culex pipiens (Honnen et al., 2016), though downstream consequences of these genomic effects for zoonoses are currently unknown. Furthermore, artificial light at night has been linked to increased West Nile Virus (WNV) exposure in Florida (Kernbach et al., 2021), a pattern that may hold true in temperate HUCs as well. These forms of pollution are pervasive in HUCs and should be recognized for their potential impacts on zoonotic hazards.

Zoonotic systems in HUCs may be influenced by heavy metal pollution, which has historically tainted cities and may have persistent physiological and immune effects, despite ongoing cleanup efforts (Perugini et al., 2011; Rodríguez Martín et al., 2015; Swaileh & Sansur, 2006). Heavy metal contamination is spatially aggregated in post-industrial sites, often positively correlated with neighborhood poverty (Aelion et al., 2013), and can have varied downstream influences on reservoir host immunity and pathogen dynamics (Sánchez et al., 2020). For example, urban pigeons with higher lead concentrations suffered higher intensities of blood pathogens, but those with increased zinc concentrations experienced protective effects against Chlamydiaceae infection, a family of potentially human zoonotic bacteria (Gasparini et al., 2014). Lead exposure and cadmium exposure have been found to increase susceptibility of brown rats to bacterial challenge, suggesting that heavy metal pollution could increase their capacity to host zoonotic agents (Cook et al., 1975). In contrast, a recent study of white-footed mice on polluted sites showed no evidence of reduced immunocompetence (Biser et al., 2004). Further work is needed to elucidate how pollution-mediated

variation in host susceptibility impacts zoonotic disease risk across urban landscapes.

#### 4 | BIOTIC INTERACTIONS

Species interactions in wildlife communities and their reactions to abiotic factors and socio-ecological landscape features structure the presence of zoonotic hazards in HUCs, their pathogen burden, and human-wildlife interactions that may result in spillover events (Plowright et al., 2017). Identifying and managing zoonotic hazards is thus contingent on understanding (i) how species traits influence their distribution, (ii) how intra- and inter-species interactions, including indirect effects of community diversity, influence pathogen prevalence, and (iii) how resource availability modifies these processes. With this knowledge, public health stakeholders can better implement both direct interventions (e.g., species removal, pathogen-targeted vaccinations) and indirect management solutions (e.g., habitat modification, policy changes).

For reservoir hosts in HUCs, behavioral traits, particularly flexibility to resource shifts and human disturbance, allow colonization and population persistence in habitable patches of either green and/ or gray space (Lowry et al., 2012). Generalist species often dominate urban wildlife communities, and typically exhibit a dependence on humans for resources (i.e., human commensal ecology) and high spatial overlap with urban residents (Pickett et al., 2011). Intense urbanization can also filter for individuals with increased boldness and reduced stress responses, possibly increasing opportunities for human-wildlife interaction (Atwell et al., 2012; Carrete & Tella, 2017: Lowry et al., 2012). Additionally, behavioral and phenotypically plastic traits interact with landscape heterogeneity to determine species-specific dispersal success and home ranges (Baguette et al., 2013). Generally, host movement is restricted in cities compared to non-urban habitats due to patchy distribution of suitable habitat (Tucker et al., 2018); however, those species capable of surviving within and traversing the urban matrix can play a major role in the presence and prevalence of zoonotic pathogens across the landscape (Firth et al., 2014; Vanacker et al., 2019). By modeling both habitat suitability and dispersal pathways of hosts in HUCs, public health stakeholders can better understand the ecological determinants of their distribution and incorporate these findings into targeted management solutions that reflect the spatial scale relevant to the target species.

Arthropod vectors usually require bloodmeals, providing direct links between the distribution of habitat, hosts, and the patchy distribution of mosquitos and ticks across green and gray space (Bajwa, 2018). For example, *Culex* mosquitoes are attracted to breeding habitats high in organic matter and bacteria (Burkett-Cadena & Mullen, 2007), and preferentially feed on avian hosts (Bernard et al., 2001). This limits them to avian habitats, which may consist of relatively small patches of green space compared to mammal habitat and is often interspersed across human-populated areas, creating opportunities for host shifts from avian to human populations (Kilpatrick

et al., 2006). In contrast, *Ae. aegypti* and *Ae. albopictus*, important vectors of arboviruses, exploit small artificial container habitats in cities (e.g., rainwater in tires, tree holes, etc.) (Hawley, 1988), and blood feed largely on mammals (Faraji et al., 2014; Valerio et al., 2008), including humans (Rose et al., 2020). As a result, these mosquitoes are typically distributed across a broader range of green and gray spaces (Pless et al., 2021). Ticks are an increasingly important vectors of bacterial, viral, and parasitic pathogens in cities globally (Dautel & Kahl, 1999; Hansford et al., 2017; Heylen et al., 2019; LaDeau et al., 2015; Lydecker et al., 2019; Steere, 1994; Vanacker et al., 2019). Pathogen prevalence in urban ticks has been linked to green space structure, including understory composition (Adalsteinsson et al., 2018), and its influence on the movement and community dynamics of hosts (see discussion of dilution effect below).

Though available niche space for zoonotic hazards in urban landscapes may remain unoccupied when landscape barriers and species traits limit successful colonization, urbanization often facilitates introductions (Reed et al., 2020). HUCs are often global and regional hubs of interconnectivity through trade and travel, providing opportunities for the movement of hosts, vectors, and pathogens into and between cities (Padayachee et al., 2017; Reed et al., 2020). Ports of entry in HUCs are particularly vulnerable to invasion given the daily flow of people, goods, and biota moving through them. Pathogen prevalence and diversity in urban rats is often higher around ports, suggesting periodic introductions (Rothenburger et al., 2017). In some scenarios, ecological priority effects may buffer the threat of new invasions if niches are filled such that increased competition limits the successful establishment of additional species or populations (Fraser et al., 2014). However, high propagule pressure (a composite measure of the number of non-native individuals dispersing into a region), continued anthropogenic disturbance, and increased resource availability may allow invasion despite high occupancy and competition with established populations. Additionally, elevated frequency and intensity of human-wildlife interactions due to high human density in HUCs may allow spillover to occur rapidly upon arrival such that population establishment is not always necessary for introduced species to act as important zoonotic hazards. Prioritizing targeted screening for introductions where zoonotic potential is high (e.g., where high propagule pressure and high human density overlap; Little et al., 2017) as well as protocols for rapid identification of potential threats, can provide crucial and cost-effective strategies for limiting zoonotic hazards in HUCs.

The role of biodiversity and abundance of hosts in habitat patches has become an increasingly important and debated factor in the study of zoonotic hazards (Randolph & Dobson, 2012; Wood & Lafferty, 2013). The dilution effect hypothesis posits that increased vertebrate diversity begets an increasing proportion of low-competence hosts that reduce overall environmental prevalence of specific pathogens and thus the hazard or risk to humans (Keesing et al., 2010; LoGiudice et al., 2003). Direct population regulation of zoonotic hosts or vectors through competition or predation is also possible through increases in diversity (e.g., predator reduction of vector mosquito abundance; Dambach, 2020), though this differs

from canonical dilution effects based on indirect interactions. The dilution effect has spurred a strong public health impetus for managing urban environments to maximize biodiversity. However, several arguments suggest these effects are not experienced in many temperate HUCs.

First, we must note that the diversity-risk relationship is often depicted as nonlinear, whereby risk increases in low diversity systems and only decreases after a diversity threshold is met (Diuk-Wasser et al., 2021; Kilpatrick et al., 2017; Rohr et al., 2020; Wood & Lafferty, 2013). Across most small and isolated habitat patches in HUCs, host diversity often remains too low to reach a dilution threshold (Diuk-Wasser et al., 2021). Second, while the classic dilution hypothesis predicts higher diversity systems results in an abundance of "dilution" hosts, a recent meta-analysis revealed that the increased diversity in cities is composed of a large proportion of zoonotic hosts, resulting in pathogen amplification rather than dilution (Gibb et al., 2020). Indeed, species experiencing global abundance increases by adapting to human-dominated landscapes pose the greatest risk of viral spillover (Johnson et al., 2020). Furthermore, while re-greening through de-urbanization in HUCs may create natural experiments by increasing urban diversity, recent work indicates hosts in these environments harbor increased pathogen loads and infection prevalence (Peterson et al., 2020). Third, transmission mode (i.e., frequency dependent vs. density dependent) can dramatically impact infection rates, suggesting changes in diversity in HUCs will lead to different pathogen-specific outcomes depending on underlying biology (Dobson, 2004; Faust et al., 2017). Finally, diversity increases in HUCs are associated with greater beta diversity through increased number of unique niche types rather than alpha diversity from species sharing common habitats (Mckinney, 2008). Given trends of reduced movement and concentrated resources in cities (Tucker et al., 2018), this suggests many urban species do not co-occur or interact with the same suite of vectors and pathogens, which is an assumption of the dilution effect. Thus, increases in biodiversity, either through natural dispersal or human-assisted colonization into these areas, may be more likely to increase human disease risk by increasing abundance of competent zoonotic hosts and facilitating greater diversity of pathogenic hazards.

Practical guidance for public health stakeholders in HUCs regarding biodiversity will require a local perspective and context, as outcomes for human disease may be idiosyncratic (Salkeld et al., 2013). Rather than blanket promotion of biodiversity as a buffer against infectious disease (Randolph & Dobson, 2012), each HUC should seek to understand the unique suite of species and ecological processes driving hazards locally and then balance the value of ecosystem services provided by increasing biodiversity against costs of species-specific zoonotic risks.

Urban areas often exhibit increased availability of anthropogenic food resources, either from household and commercial trash or intentionally provided food subsidies, with clear consequences for zoonotic hazards (Altizer et al., 2018; Becker et al., 2018). These ample and clumped resource distributions directly influence zoonotic hazards by increasing host density (Becker & Hall, 2014; Murray et al.,

2016), affecting host immune response and thus pathogen susceptibility (Murray et al., 2016), increasing host aggregation, and altering host movement patterns (either increasing dispersal to, or decreasing dispersal from, high-resource sites) (Becker et al., 2018). Highresource density can drive local pathogen load and prevalence in hosts indirectly by altering behavior and interspecific and intraspecific interactions among species (Becker et al., 2018; Moyers et al., 2018). For example, raccoons feeding at clumped resources exhibited higher endoparasite prevalence and parasite diversity (Wright & Gompper, 2005). The dynamics of pathogen-host interactions and resultant zoonotic hazards will depend on specific host immune responses and changes in dispersal (Becker & Hall, 2014; Becker et al., 2018), suggesting a need for further research across local contexts in HUCs. Ultimately, human activity drives the distribution of clumped anthropogenic resources in HUCs, suggesting policy decisions and cultural attitudes around trash containment and wildlife feeding remain important levers in the toolbox of public health stakeholders. Ideally, waste and wildlife managers can develop plans to reduce access to supplemental anthropogenic resources because it increases the likelihood of spatial overlap among hosts and humans, but effective economical waste management remains a major challenge for cities globally.

## 5 | CASE STUDY: NEW YORK CITY, NEW YORK, UNITED STATES

New York City (NYC) supports the largest human population (8.4 M) and highest density (28k people/mi<sup>2</sup>) of any city in the United States. NYC has also been a hub for international travel and trade for centuries, serving as an entry point for the invasion of zoonotic reservoirs, such as brown rats (Armitage, 1993), and pathogenic agents such as West Nile Virus (Sejvar, 2003). The five boroughs of NYC exhibit strong socioeconomic disparities and differences in land use history that drive differential hazard presence (Figure 2). NYC supports a spectrum of green and gray space connectivity, from the large, connected forest patches of Staten Island and Bronx to the intensely developed urban core of Manhattan (Figure 2a). Furthermore, a wealth of research by academic institutions and municipal agencies actively document the distribution and pathogenicity of several zoonotic hazards across NYC (Figure 2c-f) (Bajwa, 2018; Johnson et al., 2016; Little et al., 2017; Vanacker et al., 2019; Walsh, 2014). These attributes make NYC a useful case study to understand how urban ecoepidemiological processes influence zoonotic disease emergence and the roles of diverse stakeholders in surveillance, research, and identification of relevant ecological drivers.

#### 5.1 | Rodent-borne zoonoses

Brown rats and house mice host a diverse community of zoonotic pathogens (Table 2). While city agencies track disease outbreaks, genetic and observational studies by academics provide detailed




FIGURE 2 Eco-epidemiology of zoonotic hazards in New York City described by: distribution of green and gray land cover (a), median household income (b), density of *Aedes albopictus* (c) and *Culex pipiens* (d) mosquitos interpolated from citywide sampling points, *Ixodes scapularis* density and infection prevalence with *Borrelia burgdorferi* (e), and predicted abundance of *Rattus norvegicus* (f). Land cover data sourced from https://opendata.cityofnewyork.us/. Income data mapped by zip code, sourced from the American Community Survey. *Aedes* and *Culex* density adapted from Bajwa (2018) reproduced with permission from the American Mosquito Control Association. *Ixodes* and *Borrelia* data provided by authors of Vanacker et al. (2019). *Rattus* data adapted from Walsh (2014) [Colour figure can be viewed at wileyonlinelibrary.com]

characterization of pathogen and ectoparasite threats (Firth et al., 2014; Frye et al., 2015; Williams et al., 2018). Of most concern in NYC is *Leptospira interrogans*, which is shed through rodent urine and environmentally transmitted, with 58 local cases of Leptospirosis reported between 2006 and 2021 (Bassett, 2017; Chokshi, 2021).

Modeling efforts led by city agencies and various academic groups help identify drivers of rodent habitat and exposure (Childs et al., 1998; Johnson et al., 2016; Walsh, 2014), informed by public datasets of approximately 1.8 million city inspections and 23 million reported complaints between 2010 and 2020, highlighting the role of government and community engagement. Lower socioeconomic wealth is consistently identified as driving rat abundance, as are the characteristics of abiotic structures including building age, vacancy rates, and sewer and subway infrastructure (Johnson et al., 2016; Walsh, 2014). NYC recently began neighborhood-based management approaches (i.e., vs property-level treatments) to match the spatial scale of rodent activity (Mayor, 2017). Studies outside of NYC clearly identify biotic factors such as anthropogenic food and vegetated areas as key sources of rodent resources and habitat (Feng & Himsworth, 2014; Traweger et al., 2006; van Adrichem et al., 2013),

though rats in HUCs do not require green space and regularly live at high densities in gray space environments.

#### 5.2 | Aedes-borne zoonoses

Surveys led by city and state agencies and academics help track the diverse community of *Aedes* spp. mosquitos and their associated pathogens in the NYC area (Table 2), where *Ae.* albopictus is the most epidemiologically-significant vector (Bajwa, 2018; Little et al., 2017; McMillan et al., 2020). While most *Aedes*-borne zoonoses in NYC are imported, laboratory studies reveal *Ae. albopictus* from NY State are competent for Zika virus transmission (Chouin-Carneiro et al., 2016) and in a European HUC a researchers from non-profit, for profit, and government researchers reported local chikungunya transmission (Grandadam et al., 2011), suggesting potential spillover and transmission risks in NYC.

Surveys reveal widespread Aedes spp. distribution across Staten Island with localized hotpots in other NYC boroughs with both green and gray spaces. While standing water availability provides necessary larval habitat, modeling efforts suggest Ae. albopictus

TABLE 2 Major zoonotic hazards of New York City

	Host or vector	Zoonotic pathogens	Socio-ecological drivers in NYC
Rodents	Rattus norvegicus Mus musculus	Bartonella spp., Leptospira interrogans, Clostridium difficile, Seoul Hantavirus, others Shigella spp., Salmonella spp., Clostridium difficile, and Leptospira interrogans, others	L: High-poverty areas linked to increased host abundance and pathogen prevalence; neighborhood-scale treatments match spatial scale of rodent activity      A: Built structures and underground infrastructure constitutes major habitat. Older buildings and higher vacancy rates increase abundance      B: Human food waste is major and abundant resource. Reducing trash limits local carrying capacity
Aedes spp. mosquitos	Aedes albopictus Ae. canadensis, Ae. vexans, Ae. trivittatus	Zika virus, dengue virus, chikungunya viruses Eastern equine encephalitis virus, Jamestown Canyon virus, La Crosse virus	L: Aedes found across all urbanization levels; abundance increases in low-to medium-intensity development A: Standing water availability limits larval habitat (natural and artificial containers). Macroclimate variation influences population survival, microclimate variation influences individual behavior and survival B: Vegetation offers sugar meals, resting habitat, and habitat for animal host
Culex spp. mosquitos	Culex pipiens f. pipiens Culex pipiens f. molestus	West Nile Virus (WNV), St. Louis encephalitis	L: WNV incidence higher in high- poverty areas. Neighborhood- scale management targets known disease hotspots (e.g., northern Queens). Stronger association with green space habitat than Aedes spp., especially for aboveground inhabiting C.p. pipiens; C.p. molestus preferentially inhabits underground habitats A: Temperature influences vectorial capacity. Macroclimate variation influences population survival, microclimate variation influences individual behavior and survival B: Canopy and vegetation offer habitat for avian hosts, resting habitat, and useful targets for preventative insecticide treatments
Mesomammals	Felis catus Procyon lotor	Bartonella spp. and Toxoplasma gondii Baylisascaris procyonis, rabies lyssavirus, T. gondii	L: Use green-space habitat but survive across gradient of connectivity.  Cats survive in smaller patches in mix of green and gray space.  High poverty linked to higher cat density and Toxoplasma gondii contamination  A: Mesomammals are robust to normal range of climate variation  B: Clumped food resources from intentional feeding or human food waste increase local density, increase pathogen transmission within/between co-feeding species

TABLE 2 (Continued)

	Host or vector	Zoonotic pathogens	Socio-ecological drivers in NYC
Ticks	Ixodes scapularis Dermacentor variabilis Amblyomma americanum	Borrelia burgdorferi, Babesia microti, Anaplasma phagocytophilum, Powassan virus Rickettsia rickettsii Ehrlichia chaffeensis, Ehrlichiia ewingii	L: Ticks restricted to green land cover.  Ixodes more limited to forest with leaf litter, Amblyomma survives on mowed lawns. Increasing green space connectivity increases  Borrelia prevalence and tick density. Broad scale required for hazard assessment  A: Temperature influences tick activity. Microclimate variation influences fine-scale distribution during questing and resting.  B: Influence of host community on vector and pathogen. Deer required for tick reproduction; small mammal reservoir diversity may influence pathogen prevalence.

Note: Hazards are separated by taxa groups and include relevant host or vector species, zoonotic pathogens of importance, and ecological drivers of hazard intensity broken up by type (L: landscape characteristics, A: abiotic conditions, B: biotic interactions).

abundance correlates with increasing low- and medium-intensity urban development (Kache et al., 2020) as well as seasonal meteorological conditions (Little et al., 2017), highlighting both landscape and abiotic drivers of this hazard. Critical biotic factors include presence of vegetation for sugar meals and resting habitat (Fikrig et al., 2020; Samson et al., 2013), as well as distributions of mammalian hosts that are themselves structured by the socio-ecological landscape (Faraji et al., 2014; Goodman et al., 2018).

Mesomammal-borne zoonoses

et al., 2006; Loss et al., 2009).

#### 5.3 | Culex-borne zoonoses

In NYC, the most important and prevalent *Culex* spp. mosquito-borne pathogen is West Nile Virus (WNV), which is vectored locally by the *Culex pipiens* complex. City agencies lead vector surveillance and track human cases. Across the five boroughs, 381 WNV cases have been reported between 2000 and 2019, with the highest case counts in Queens where WNV was first detected in 1999 (NYC Department of Health and Mental Hygiene, 2019).

Both *Culex pipiens* f. pipiens and *Culex pipiens* f. molestus are locally abundant (Bajwa, 2018) but exhibit diverse activity patterns, physiology, and spatial distribution across subterranean and aboveground environments (Vinogradova, 2000), allowing them to inhabit a wide range of conditions across NYC's green and gray spaces. Academic studies continue to reveal cryptic biological variation in *Culex* species, which helps inform municipal surveillance, identification, and management efforts (Kilpatrick et al., 2010).

Despite few studies describing ecological drivers of *Culex*-borne hazards in NYC, others have identified a correlation between WNV cases and urbanization and poverty (Andreadis et al., 2004; Poh et al., 2020), though the causal mechanisms remain poorly characterized. Temperature is the key abiotic driver of *Culex* spp. populations and vectoral capacity (Ciota et al., 2014; Reisen, 2013). In NYC,

NYC supports a community of native and introduced mesomammals (WildlifeNYC, 2021). While few mesomammal-borne zoonoses are reported in NYC, local studies of raccoons and feral cats conducted by city agencies and academics have identified a diverse group of zoonotic pathogens and vectors from hosts and their environment (Table 2; Bassett, 2018; Rainwater et al., 2017; Tufts et al., 2021; Tyungu et al., 2020). These hosts are under-surveyed and associated zoonotic spillover events are underreported, but recent but unreviewed estimates suggest NYC supports tens of thousands of free roaming cats, and Central Park alone supports around 500 raccoons (Slavinski et al., 2012).

passerine birds are the major WNV reservoir and pathogen prev-

alence may be linked to host-specific competency and preference

by vectors (Bernard et al., 2001; Kilpatrick et al., 2006; Kramer &

Bernard, 2001; Nasci et al., 2002). The role of increasing avian di-

versity for WNV hazard is debated; it appears negatively correlated

at broad scales (Allan et al., 2009), but found to be uncorrelated for

passerine birds and at fine spatial scales in Chicago, USA (Ezenwa

Supplemental feeding by local residents appears to be the main biotic driver of mesomammal populations, causing dense and clumped populations that may allow increased transmission within and between co-feeding species (Bozek et al., 2007; Rainwater et al., 2017). They are robust to most climactic conditions, suggesting little role for abiotic variation. Their capacity for long-distance movements enables mesomammals to survive across a gradient of green connectivity, from networks of forested patches to small and isolated parks or gardens surrounded by the gray spaces. Socio-ecological landscape characteristics are useful predictors of mesomammal

zoonotic hazard, given studies of *Toxocara cati*, a cat-borne parasite, reveal higher levels of egg contamination in soils of lower-income NYC neighborhoods (Tyungu et al., 2020).

#### 5.5 | Tick-borne zoonoses

NYC supports four major Ixodid tick species including the black-legged tick (Ixodes scapularis), the American dog tick (Dermacentor variabilis), the Ione star tick (Amblyomma americanum), and the recently introduced Asian longhorned tick (Haemaphysalis longicornis) (Barbot, 2020; Tufts et al., 2021). Their distributions, densities, and pathogen communities are surveyed by city and state agencies as well as academic researchers through yearly observational sampling (e.g., tick dragging, carbon dioxide traps, microscopy) and molecular approaches (e.g., qPCR). LD remains the most common tick-borne disease in NYC with 812 cases between 2010 and 2019, though human babesiosis is becoming increasingly prevalent (average 62 cases/year; Barbot, 2020).

Sampling indicates tick populations are largely restricted to urban green spaces and adjacent neighborhoods in Staten Island and Bronx in NYC. Modeling efforts further emphasize the critical role of biotic factors for urban tick-borne hazards, revealing positive associations between NYC canopy cover connectivity and both I. scapularis density and prevalence of Borrelia burgdorferi, the etiological agent of LD (Vanacker et al., 2019). Evidence for dilution effects in HUCs like NYC remains mixed (Salkeld et al., 2013) and an area of active research. Neighborhood greenness and biodiversity are correlated with economic wealth in HUCs and linked to racist development policies (i.e., the luxury effect) (Schell et al., 2020), illustrating strong links between tick-borne hazards socio-ecological landscape drivers. Community science reporting further enables research of human behavior and demographics on tick-borne hazard exposure (Bron et al., 2020). Abiotic factors ultimately govern survival of individual ticks via species-specific tolerances for humidity and temperature in HUCs (Diuk-Wasser et al., 2021), where microclimate conditions are often linked to fragmentation and land use patterns (Tuff et al., 2016).

#### 6 | CONCLUSION

The COVID-19 pandemic put a spotlight on the potential devastating human health impacts of zoonotic pathogen emergence, reinforcing the importance of identifying local hazards and understanding their ecological relationships with diverse environments to prevent future spillover events. The persistent risk of known and novel zoonotic diseases in close proximity to high-density human settlements in HUCs should be a call to action to reassess the ecological factors that influence both the distribution of hazards and their varying pathogen prevalence to improve active assessment and mitigation of potential human zoonoses. In this article, we have provided an assessment of how factors relating to the socio-ecological landscape,

abiotic conditions, and biotic interactions influence multiple zoonotic hazards in HUCs and provided examples of these how these patterns and processes play out in NYC.

Each city has its unique suite of hazards and socio-ecological landscape patterns to consider, as well as its own practical limitations in available budget, labor, and political support for wildlife management and spillover mitigation. Effective zoonotic spillover prevention requires key collaborations among public health stakeholders, a strategy suggested by a chorus of zoonoses research groups (Eskew & Olival, 2018; Gottdenker et al., 2014; Hansen et al., 2017; Hassell et al., 2017; Magle et al., 2019; Plowright et al., 2021; Vanwambeke et al., 2019). Municipal departments of health and city biologists can work with disease ecologists and modelers to improve surveillance networks and better understand local host/vector distributions, connectivity networks, and interactions with heterogeneous landscapes. Social scientists and community leaders provide necessary context to socio-ecological frameworks, help prioritize programs toward high-risk communities, and promote local engagement for community-based surveillance and reporting efforts. Urban planners, non-profit public health advocates, and policy specialists will also be key partners for developing feasible mitigation goals and securing political support and funding. Establishing these multi-sectoral partnerships and promoting a common understanding of the ecological drivers of local zoonotic hazards are crucial steps toward pre-spillover prevention in HUCs.

We recommend several steps for surveillance that can inform city-specific hazard assessment and response plans. First, it is critical to establish baseline information on the geographic distribution of hazards through field surveys, community reporting, and local expertise. Multi-city information networks can bolster these efforts by promoting methodological standardization and enabling comparative research that informs local efforts (Magle et al., 2019). Given that exhaustive surveys are often unfeasible, sub-sampling areas of interest and leveraging predictive models of habitat distribution can provide valuable information on hazard intensity across complex urban landscapes. Second, this information must be paired with molecular approaches to identify variable pathogen prevalence and eco-epidemiological modeling approaches to predict how they might change over time (Fountain-Jones et al., 2018; Lloyd-Smith et al., 2009). Incorporating community science in aspects of surveillance (e.g., reporting wildlife sightings and encounters with vectors) would also supplement efforts where resources are limited.

While this article does not focus on variable zoonotic exposure (i.e., the likelihood of human-pathogen contact from human interaction with the environment) or risk factors (i.e., the likelihood of adverse effects, given exposure and vulnerability) across HUCs, we recommend management plans also assess these metrics to focus efforts and limited resources on the most at-risk communities. Together, data on hazards, exposure and vulnerability allows for a comprehensive assessment of risk and the creation of quantitative risk maps, which are key tools to facilitate public education of local hazards and develop actionable and targeted management plans (Beard et al., 2018; Morandeira et al., 2019; Ostfeld et al., 2005;

vonHedemann et al., 2015). Finally, management plans should seek to implement integrated pest management approaches when possible, to reduce local densities and sustainably limit hazard intensity but may often require rapid and/or cost-effective measures to directly reduce hazards, especially when faced with urgent needs (Witmer, 2007).

Reducing zoonotic risk across HUCs is challenging for many reasons. Urban landscapes have dynamic demographic, environmental, cultural, and political conditions, suggesting risk maps along with public health programs should be frequently updated to reflect changes over time. Additionally, the basic biology of many hosts, vectors, and pathogens remains understudied, particularly in the context of hyper-local and novel conditions presented within each city, indicating a continued need for collaborative research. Given the continued growth in the number, density, and geographic footprint of HUCs globally, ensuring prevention and preparedness through biologically informed strategies to surveille, research, and manage zoonoses is key to reducing the burden of these preventable human diseases

#### **ACKNOWLEDGEMENTS**

We thank Waheed Bajwa and Sally Slavinski and others in the New York City Department of Health and Mental Hygiene and the New York City Department of Parks and Recreation for useful discussions and collaboration. We also thank the Journal of the American Mosquito Control Association for use of figures. This publication was supported in part by Cooperative Agreement Number U01CK000509-01, funded by the Centers for Disease Control and Prevention and the National Science Foundation's Dynamics of Integrated Socio-Environmental Systems (CNH2) program (Award #1924061). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention, the Department of Health and Human Services or the National Science Foundation.

#### DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

## ORCID

Matthew A. Combs https://orcid.org/0000-0003-1928-2485

Pallavi A. Kache https://orcid.org/0000-0002-8104-9599

Meredith C. VanAcker https://orcid.org/0000-0002-5690-9139

Nichar Gregory https://orcid.org/0000-0002-0872-9950

Laura D. Plimpton https://orcid.org/0000-0002-2632-9272

Danielle M. Tufts https://orcid.org/0000-0002-8084-7556

Maria P. Fernandez https://orcid.org/0000-0001-8645-2267

Maria A. Diuk-Wasser https://orcid.org/0000-0001-8809-3050

### REFERENCES

Adalsteinsson, S. A., Shriver, W. G., Hojgaard, A., Bowman, J. L., Brisson, D., D'Amico, V., & Buler, J. J. (2018). Multiflora rose invasion amplifies prevalence of Lyme disease pathogen, but not necessarily

- Lyme disease risk. *Parasites and Vectors*, 11(1), 1–10. https://doi. org/10.1186/s13071-018-2623-0
- Aelion, C. M., Davis, H. T., Lawson, A. B., Cai, B., & McDermott, S. (2013). Associations between soil lead concentrations and populations by race/ethnicity and income-to-poverty ratio in urban and rural areas. *Environmental Geochemistry and Health*, 35(1), 1–12. https://doi.org/10.1007/s10653-012-9472-0
- Alberti, M. (2005). The effects of urban patterns on ecosystem function. International Regional Science Review, 28(2), 168–192. https://doi.org/10.1177/0160017605275160
- Alirol, E., Getaz, L., Stoll, B., Chappuis, F., & Loutan, L. (2011). Urbanisation and infectious diseases in a globalised world. The Lancet Infectious Diseases, 11(2), 131–141. https://doi.org/10.1016/ 51473-3099(10)70223-1
- Allan, B. F., Keesing, F., & Ostfeld, R. S. (2003). Effect of forest fragmentation on lyme disease risk. *Conservation Biology*, 17(1), 267–272. https://doi.org/10.1046/j.1523-1739.2003.01260.x
- Allan, B. F., Langerhans, R. B., Ryberg, W. A., Landesman, W. J., Griffin, N. W., Katz, R. S., Oberle, B. J., Schutzenhofer, M. R., Smyth, K. N., de St. Maurice, A., Clark, L., Crooks, K. R., Hernandez, D. E., McLean, R. G., Ostfeld, R. S., & Chase, J. M. (2009). Ecological correlates of risk and incidence of West Nile virus in the United States. *Oecologia*, 158(4), 699–708. https://doi.org/10.1007/s00442-008-1169-9
- Altizer, S., Becker, D. J., Epstein, J. H., Forbes, K. M., Gillespie, T. R., Hall, R. J., Hawley, D. M., Hernandez, S. M., Martin, L. B., Plowright, R. K., Satterfield, D. A., & Streicker, D. G. (2018). Food for contagion: Synthesis and future directions for studying host-parasite responses to resource shifts in anthropogenic environments. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1745). https://doi.org/10.1098/rstb.2017.0102
- Andreadis, T. G., Anderson, J. F., Vossbrinck, C. R., & Main, A. J. (2004). Epidemiology of West Nile Virus in Connecticut: A five-year analysis of mosquito data 1999–2003. *Vector Borne Zoonotic Diseases*, 4(4), 360–378. https://doi.org/10.1089/vbz.2004.4.360
- Armitage, P. L. (1993). Commensal rats in the new world. The Biologist, 40(4), 174-178.
- Atwell, J. W., Cardoso, G. C., Whittaker, D. J., Campbell-Nelson, S., Robertson, K. W., & Ketterson, E. D. (2012). Boldness behavior and stress physiology in a novel urban environment suggest rapid correlated evolutionary adaptation. *Behavioral Ecology*, 23(5), 960–969. https://doi.org/10.1093/beheco/ars059
- Baguette, M., Blanchet, S., Legrand, D., Stevens, V. M., & Turlure, C. (2013). Individual dispersal, landscape connectivity and ecological networks. *Biological Reviews*, 88(2), 310–326. https://doi.org/10.1111/brv.12000
- Bajomi, D., Kiss, Z., & Nagai, Y. (2013). 40 Years of rat control in Budapest: The importance of sewer treatments for effective rat control in urban areas. *International Pest Control*, 55(1), 1–5.
- Bajwa, W. I. (2018). A taxonomic checklist and relative abundance of the mosquitoes of New York City. *Journal of the American Mosquito Control Association*, 34(2), 138–142. https://doi.org/10.2987/ 18-6738.1
- Barbot, O. (2020). 2020 Health advisory # 22: tick-borne disease advisory.
- Barding, E. E., & Nelson, T. A. (2008). Raccoons use habitat edges in northern Illinois. *American Midland Naturalist*, 159(2), 394–402.
- Bassett, M. T. (2017). Leptospirosis cluster in the concourse area of the bronx (Vol. 3, pp. 1–2). New York City Department of Health and Mental Hygiene. https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/leptospirosis-in-the-bronx.pdf
- Barbot, O. (2018). Canine distemper virus identified in raccoons from central park. *Veterinary Alert*, 7, 1–2. https://www1.nyc.gov/assets/doh/downloads/pdf/zoo/18vet07.pdf
- Beard, R., Wentz, E., & Scotch, M. (2018). A systematic review of spatial decision support systems in public health informatics supporting the identification of high risk areas for zoonotic disease outbreaks.

- International Journal of Health Geographics, 17(1), 1–19. https://doi.org/10.1186/s12942-018-0157-5
- Becker, D. J., & Hall, R. J. (2014). Too much of a good thing: Resource provisioning alters infectious disease dynamics in wildlife. *Biology Letters*, 10(7). https://doi.org/10.1098/rsbl.2014.0309
- Becker, D. J., Snedden, C. E., Altizer, S., & Hall, R. J. (2018). Host dispersal responses to resource supplementation determine pathogen spread in wildlife metapopulations. *American Naturalist*, 192(4), 503–517. https://doi.org/10.1086/699477
- Becker, D. J., Streicker, D. G., & Altizer, S. (2018). Using host species traits to understand the consequences of resource provisioning for host-parasite interactions. *Journal of Animal Ecology*, 87(2), 511–525. https://doi.org/10.1111/1365-2656.12765
- Bengis, R. G., Leighton, F. A., Fischer, J. R., Artois, M., Mörner, T., & Tate, C. M. (2004). The role of wildlife in emerging and re-emerging zoonoses. OIE Revue Scientifique Et Technique, 23(2), 497–511. https:// doi.org/10.20506/rst.23.2.1498
- Bernard, K. A., Maffei, J. G., Jones, S. A., Kauffman, E. B., Ebel, G. D., Dupuis, A. P., Ngo, K. A., Nicholas, D. C., Young, D. M., Shi, P.-Y., Kulasekera, V. L., Eidson, M., White, D. J., Stone, W. B., & Kramer, L. D. (2001). West nile virus infection in birds and mosquitoes, New York State, 2000. Emerging Infectious Diseases, 7(4), 679–685. https://doi.org/10.3201/eid0704.017415
- Biser, J. A., Vogel, L. A., Berger, J., Hjelle, B., & Loew, S. S. (2004). Effects of heavy metals on immunocompetence of white-footed mice (*Peromyscus leucopus*). *Journal of Wildlife Diseases*, 40(2), 173–184. https://doi.org/10.7589/0090-3558-40.2.173
- Blair, R. B. (1996). Land use and avian species diversity along an urban gradient. *Ecological Applications*, 6(2), 506–519. https://doi. org/10.2307/2269387
- Bledsoe, A., & Wright, W. J. (2019). The anti-blackness of global capital. Environment and Planning D: Society and Space, 37(1), 8–26. https://doi.org/10.1177/0263775818805102
- Bozek, C. K., Prange, S., & Gehrt, S. D. (2007). The influence of anthropogenic resources on multi-scale habitat selection by raccoons. *Urban Ecosystems*, 10(4), 413-425. https://doi.org/10.1007/s1125 2-007-0033-8
- Bradley, C. A., & Altizer, S. (2007). Urbanization and the ecology of wildlife diseases. *Trends in Ecology and Evolution*, 22(2), 95–102. https://doi.org/10.1016/j.tree.2006.11.001
- Brinkerhoff, R. J., Folsom-O'Keefe, C. M., Tsao, K., & Diuk-Wasser, M. A. (2011). Do birds affect Lyme disease risk? Range expansion of the vector-borne pathogen Borrelia burgdorferi. Frontiers in Ecology and the Environment, 9(2), 103–110. https://doi.org/10.1890/090062
- Brock, P. M., Fornace, K. M., Grigg, M. J., Anstey, N. M., William, T., Cox, J., Drakeley, C. J., Ferguson, H. M., & Kao, R. R. (2019). Predictive analysis across spatial scales links zoonotic malaria to deforestation. *Proceedings of the Royal Society B: Biological Sciences*, 286(1894), 20182351. https://doi.org/10.1098/rspb.2018.2351
- Bron, G. M., Fernandez, M. D. P., Larson, S. R., Maus, A., Gustafson, D., Tsao, J. I., Diuk-Wasser, M. A., Bartholomay, L. C., & Paskewitz, S. M. (2020). Context matters: Contrasting behavioral and residential risk factors for Lyme disease between high-incidence states in the Northeastern and Midwestern United States. *Ticks and Tick-Borne Diseases*, 11(6), 101515. https://doi.org/10.1016/j. ttbdis.2020.101515
- Brooks, C. P. (2003). A scalar analysis of landscape connectivity. *Oikos*, 102(2), 433-440.
- Burkett-Cadena, N. D., & Mullen, G. R. (2007). Field comparison of bermuda-hay infusion to infusions of emergent aquatic vegetation for collecting female mosquitoes. *Journal of the American Mosquito Control Association*, 23(2), 117–123.
- Burkhalter, K. L., Campbell, S. R., Daniels, T. J., Marfin, A. A., Kerst, A. J., Aspen, S. E., Brookes, K., Falco, R. C., Ludwig, G. V., Nasci, R. S., Lanciotti, R. S., Gottfried, K. L., Kramer, L. D., Komar, N., & Moore, C. G. (2002). Detection of West Nile virus-infected mosquitoes

- and seropositive juvenile birds in the vicinity of virus-positive dead birds. American Journal of Tropical Medicine and Hygiene, 67(5), 492–496. https://doi.org/10.4269/ajtmh.2002.67.492
- Byers, K. A., Booker, T. R., Combs, M., Himsworth, C. G., Munshi-South, J., Patrick, D. M., & Whitlock, M. C. (2020). Using genetic relatedness to understand heterogeneous distributions of urban ratassociated pathogens. *Evolutionary Applications*.
- Byers, K. A., Lee, M. J., Patrick, D. M., & Himsworth, C. G. (2019). Rats about town: A systematic review of rat movement in urban ecosystems. *Frontiers in Ecology and Evolution*, 7, 1–12. https://doi.org/10.3389/fevo.2019.00013
- Byrne, K., & Nichols, R. A. (1999). *Culex pipiens* in London Underground tunnels: Differentiation between surface and subterranean populations. *Heredity*, 82(1), 7–15. https://doi.org/10.1038/sj.hdy. 6884120
- Calhoun, L. M., Avery, M., Jones, L. A., Gunarto, K., King, R., Roberts, J., & Fox, M. (2007). Combined sewage overflows (CSO) are major urban breeding sites for Culex quinquefasciatus in Atlanta, Georgia. American Journal of Tropical Medicine and Hygiene, 77(3), 478–484. https://doi.org/10.4269/ajtmh.2007.77.478
- Carrete, M., & Tella, J. L. (2017). Behavioral correlations associated with fear of humans differ between rural and urban burrowing owls. Frontiers in Ecology and Evolution, 5, 1–9. https://doi.org/10.3389/fevo.2017.00054
- Channon, D., Channon, E., Roberts, T., & Haines, R. (2006). Hotspots: Are some areas of sewer network prone to re-infestation by rats (*Rattus norvegicus*) year after year? *Epidemiology and Infection*, 134(1), 41–48. https://doi.org/10.1017/S0950268805004607
- Childs, J. E., McLafferty, S. L., Sadek, R., Miller, G. L., Khan, A. S., DuPree, E. R., Advani, R., & Glass, G. E. (1998). Epidemiology of rodent bites and prediction of rat infestation in New York City. *American Journal of Epidemiology*, 148(1), 78–87. https://doi.org/10.1093/oxfordjournals.aje.a009563
- Chokshi, D. A. (2021). Increase in leptospirosis cases in New York City (p. 35). New York City Department of Health and Mental Hygiene.
- Chouin-Carneiro, T., Vega-Rua, A., Vazeille, M., Yebakima, A., Girod, R., Goindin, D., Dupont-Rouzeyrol, M., Lourenço-de-Oliveira, R., & Failloux, A.-B. (2016). Differential Susceptibilities of Aedes aegypti and Aedes albopictus from the Americas to Zika Virus. PLoS Neglected Tropical Diseases, 10(3), 1-11. https://doi.org/10.1371/journal.pntd.0004543
- Ciota, A. T., Matacchiero, A. C., Kilpatrick, A. M., & Kramer, L. D. (2014). The effect of temperature on life history traits of *Culex* mosquitoes. *Journal of Medical Entomology*, 51(1), 55–62. https://doi.org/10.1603/ME13003
- Clark, R. W., Brown, W. S., Stechert, R., & Zamudio, K. R. (2010). Roads, interrupted dispersal, and genetic diversity in timber rattlesnakes. *Conservation Biology*, 24(4), 1059–1069. https://doi.org/10.1111/j.1523-1739.2009.01439.x
- Combs, M., Byers, K. A., Himsworth, C. G., & Munshi-South, J. (2019). Harnessing population genetics for pest management: Theory and application for urban rats. *Human Wildlife Interactions*, 13(2), 250–263. https://doi.org/10.26077/0bcx-v339
- Cook, J. A., Hoffmann, E. O., & Luzio, N. R. D. (1975). Influence of lead and cadmium on the susceptibility of rats to bacterial challenge. Proceedings of the Society for Experimental Biology and Medicine, 150(3), 741–747. https://doi.org/10.3181/00379727-150-39117
- Cushman, S. A., McRae, B. H., & McGarigal, K. (2016). Basics of landscape ecology: An introduction to landscapes and poulation processes for landscape genetics. In N. Balkenhol, S. A. Cushman, A. T. Storfer, & L. P. Waits (Eds.), Landscape genetics: concepts, methods, application (pp. 11–34). Wiley Blackwell.
- Dambach, P. (2020). The use of aquatic predators for larval control of mosquito disease vectors: Opportunities and limitations. *Biological Control*, 150, 104357. https://doi.org/10.1016/j.biocontrol.2020. 104357

- Dautel, H. & Kahl, O. (1999). Ticks (Acari: Ixodoidea) and their medical importance in the urban environment. Proceedings of the Third International Conference on Urban Pests: 19–22 July 1999, (January), 73–82.
- de Azevedo, T. S., Bourke, B. P., Piovezan, R., & Sallum, M. A. M. (2018). The influence of urban heat Islands and socioeconomic factors on the spatial distribution of *Aedes aegypti* larval habitats. *Geospatial Health*, 13(1), 179–187. https://doi.org/10.4081/gh.2018.623
- Deplazes, P., Hegglin, D., Gloor, S., & Romig, T. (2004). Wilderness in the city: The urbanization of Echinococcus multilocularis. *Trends in Parasitology*, 20(2), 77–84. https://doi.org/10.1016/j. pt.2003.11.011
- Des Roches, S., Brans, K. I., Lambert, M. R., Rivkin, L. R., Savage, A. M., Schell, C. J., & Alberti, M. (2020). Socio-eco-evolutionary dynamics in cities. *Evolutionary Applications*, 14(1), 248–267. https://doi.org/10.1111/eva.13065
- Diuk-Wasser, M. A., VanAcker, M. C., & Fernandez, M. P. (2021). Impact of land use changes and habitat fragmentation on the ecoepidemiology of tick-borne diseases. *Journal of Medical Entomology*, 58(4), 1546–1564. https://doi.org/10.1093/jme/tjaa209
- Dobson, A. (2004). Population dynamics of pathogens with multiple host species. American Naturalist, 164(S5), S64-S78. https://doi. org/10.1086/424681
- Dowling, Z., Ladeau, S. L., Armbruster, P., Biehler, D., & Leisnham, P. T. (2013). Socioeconomic status affects mosquito (Diptera: Culicidae) larval habitat type availability and infestation level. *Journal of Medical Entomology*, 50(4), 764–772. https://doi.org/10.1603/me12250
- Duffy, D. C., & Campbell, S. R. (1994). Ambient air temperature as a predictor of activity of adult Ixodes scapularis (Acari: Ixodidae). *Journal of Medical Entomology*, 31(1), 178–180. https://doi.org/10.1093/jmedent/31.1.178
- Eisenberg, J. N. S., Desai, M. A., Levy, K., Bates, S. J., Liang, S., Naumoff, K., & Scott, J. C. (2007). Environmental determinants of infectious disease: A framework for tracking causal links and guiding public health research. Environmental Health Perspectives, 115(8), 1216–1223. https://doi.org/10.1289/ehp.9806
- Eskew, E. A., & Olival, K. J. (2018). De-urbanization and zoonotic disease risk. *EcoHealth*, 15(4), 707–712. https://doi.org/10.1007/s10393-018-1359-9
- Ezenwa, V. O., Godsey, M. S., King, R. J., & Guptill, S. C. (2006). Avian diversity and West Nile virus: Testing associations between biodiversity and infectious disease risk. *Proceedings of the Royal Society* B: Biological Sciences, 273(1582), 109–117. https://doi.org/10.1098/ rspb.2005.3284
- Faraji, A., Egizi, A., Fonseca, D. M., Unlu, I., Crepeau, T., Healy, S. P., & Gaugler, R. (2014). Comparative host feeding patterns of the asian tiger mosquito, *Aedes albopictus*, in urban and suburban northeastern USA and implications for disease transmission. *PLoS Neglected Tropical Diseases*, 8(8). https://doi.org/10.1371/journ al.pntd.0003037
- Faust, C. L., Dobson, A. P., Gottdenker, N., Bloomfield, L. S. P., McCallum,
  H. I., Gillespie, T. R., Diuk-Wasser, M., & Plowright, R. K. (2017).
  Null expectations for disease dynamics in shrinking habitat:
  Dilution or amplification? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1722). https://doi.org/10.1098/rstb.2016.0173
- Feng, A. Y. T., & Himsworth, C. G. (2014). The secret life of the city rat: A review of the ecology of urban Norway and black rats (*Rattus norvegicus* and *Rattus rattus*). *Urban Ecosystems*, 17(1), 149–162. https://doi.org/10.1007/s11252-013-0305-4
- Fikrig, K., Peck, S., Deckerman, P., Dang, S., Fleur, K. S., Goldsmith, H., & Harrington, L. C. (2020). Sugar feeding patterns of New York Aedes albopictus mosquitoes are affected by saturation deficit, flowers, and host seeking. PLoS Neglected Tropical Diseases, 14(10), 1–27. https://doi.org/10.1371/journal.pntd.000824s4

- Finch, C., Al-Damluji, M. S., Krause, P. J., Niccolai, L., Steeves, T., O'Keefe, C. F., & Diuk-Wasser, M. A. (2014). Integrated assessment of behavioral and environmental risk factors for lyme disease infection on Block Island, Rhode Island. PLoS One, 9(1). https://doi.org/10.1371/journal.pone.0084758
- Firth, C., Bhat, M., Firth, M. A., Williams, S. H., Frye, M. J., Simmonds, P., Conte, J. M., Ng, J., Garcia, J., Bhuva, N. P., Lee, B., Che, X., Quan, P.-L., & Lipkin, W. I. (2014). Detection of zoonotic pathogens and characterization of novel viruses carried by commensal *Rattus norvegicus* in New York City. MBio, 5(5), 1–16. https://doi.org/10.1128/mBio.01933-14
- Fischhoff, I. R., Bowden, S. E., Keesing, F., & Ostfeld, R. S. (2019). Systematic review and meta-analysis of tick-borne disease risk factors in residential yards, neighborhoods, and beyond. BMC Infectious Diseases, 19(1), 1-11. https://doi.org/10.1186/s1287 9-019-4484-3
- Fountain-Jones, N. M., Pearse, W. D., Escobar, L. E., Alba-Casals, A., Carver, S., Davies, T. J., Kraberger, S., Papeş, M., Vandegrift, K., Worsley-Tonks, K., & Craft, M. E. (2018). Towards an ecophylogenetic framework for infectious disease ecology. *Biological Reviews*, 93(2), 950–970. https://doi.org/10.1111/brv.12380
- Francis, C. D., Ortega, C. P., & Cruz, A. (2011). Noise pollution filters bird communities based on vocal frequency. *PLoS One*, *6*(11). https://doi.org/10.1371/journal.pone.0027052
- Francis, M. J., Spooner, P. G., & Matthews, A. (2015). The influence of urban encroachment on squirrel gliders (*Petaurus norfolcensis*): Effects of road density, light and noise pollution. *Wildlife Research*, 42(4), 324–333. https://doi.org/10.1071/WR14182
- Frank, B., Delano, D., & Caniglia, B. S. (2017). Urban Systems: A Socio-Ecological System Perspective. *Sociology International Journal*, 1(1). https://doi.org/10.15406/sij.2017.01.00001
- Fraser, C. I., Banks, S. C., & Waters, J. M. (2014). Priority effects can lead to underestimation of dispersal and invasion potential. *Biological Invasions*, 17(1), 1–8. https://doi.org/10.1007/s10530-014-0714-1
- Frye, M. J., Firth, C., Bhat, M., Firth, M. A., Che, X., Lee, D., Williams, S. H., & Lipkin, W. I. (2015). Preliminary survey of ectoparasites and associated pathogens from Norway rats in New York City. *Journal of Medical Entomology*, 52(2), 253–259. https://doi.org/10.1093/jme/tjv014
- Fuller, R. A., Irvine, K. N., Devine-Wright, P., Warren, P. H., & Gaston, K. J. (2007). Psychological benefits of greenspace increase with biodiversity. *Biology Letters*, 3(4), 390–394. https://doi.org/10.1098/rsbl.2007.0149
- Fusco, N., Carlen, E., & Munshi-South, J. (2021). Urban landscape genetics: Are biologists keeping up with the pace of urbanization? *Current Landscape Ecology Reports*, 1–11. https://doi.org/10.1007/s40823-021-00062-3
- Gasparini, J., Jacquin, L., Laroucau, K., Vorimore, F., Aubry, E., Castrec-Rouëlle, M., & Frantz, A. (2014). Relationships between metals exposure and epidemiological parameters of two pathogens in urban pigeons. *Bulletin of Environmental Contamination and Toxicology*, 92(2), 208–212. https://doi.org/10.1007/s00128-013-1172-7
- Gibb, R., Redding, D. W., Chin, K. Q., Donnelly, C. A., Blackburn, T. M., Newbold, T., & Jones, K. E. (2020). Zoonotic host diversity increases in human-dominated ecosystems. *Nature*, 584. https://doi. org/10.1038/s41586-020-2562-8
- Gómez-Baggethun, E., & Barton, D. N. (2013). Classifying and valuing ecosystem services for urban planning. *Ecological Economics*, 86, 235–245. https://doi.org/10.1016/j.ecolecon.2012.08.019
- Goodman, H., Egizi, A., Fonseca, D. M., Leisnham, P. T., & Ladeau, S. L. (2018). Primary blood-hosts of mosquitoes are influenced by social and ecological conditions in a complex urban landscape. Parasites and Vectors, 11(1), 1-10. https://doi.org/10.1186/s1307 1-018-2779-7
- Gottdenker, N. L., Streicker, D. G., Faust, C. L., & Carroll, C. R. (2014). Anthropogenic land use change and infectious diseases: A review of

- the evidence. *EcoHealth*, 11(4), 619-632. https://doi.org/10.1007/s10393-014-0941-z
- Grandadam, M., Caro, V., Plumet, S., Thiberge, J.-M., Souarès, Y., Failloux, A.-B., Tolou, H. J., Budelot, M., Cosserat, D., Leparc-Goffart, I., & Desprès, P. (2011). Chikungunya virus, Southeastern France. *Emerging Infectious Diseases*, 17(5), 910–913. https://doi.org/10.3201/eid1705.101873
- Gray, J. S. (2008). Ixodes ricinus seasonal activity: Implications of global warming indicated by revisiting tick and weather data. *International Journal of Medical Microbiology*, 298(suppl 1), 19–24. https://doi.org/10.1016/j.ijmm.2007.09.005
- Gregory McPherson, E. (1992). Accounting for benefits and costs of urban greenspace. Landscape and Urban Planning, 22(1), 41–51. https://doi.org/10.1016/0169-2046(92)90006-L
- Gulachenski, A., Ghersi, B. M., Lesen, A. E., & Blum, M. J. (2016). Abandonment, ecological assembly and public health risks in counter-urbanizing cities. Sustainability (Switzerland), 8(5), 1–26. https://doi.org/10.3390/su8050491
- Hansen, A., Xiang, J., Liu, Q., Tong, M. X., Sun, Y., Liu, X., Chen, K., Cameron, S., Hanson-Easey, S., Han, G.-S., Weinstein, P., Williams, C., & Bi, P. (2017). Experts' perceptions on China's capacity to manage emerging and re-emerging zoonotic diseases in an era of climate change. Zoonoses and Public Health, 64(7), 527–536. https://doi.org/10.1111/zph.12335
- Hansford, K. M., Fonville, M., Gillingham, E. L., Coipan, E. C., Pietzsch, M. E., Krawczyk, A. I., Vaux, A. G. C., Cull, B., Sprong, H., & Medlock, J. M. (2017). Ticks and Borrelia in urban and peri-urban green space habitats in a city in southern England. Ticks and Tick-Borne Diseases, 8(3), 353–361. https://doi.org/10.1016/j. ttbdis.2016.12.009
- Harrigan, R. J., Thomassen, H. A., Buermann, W., Cummings, R. F., Kahn, M. E., & Smith, T. B. (2010). Economic conditions predict prevalence of West Nile virus. *PLoS One*, 5(11), 1–8. https://doi.org/10.1371/journal.pone.0015437
- Hassell, J. M., Begon, M., Ward, M. J., & Fèvre, E. M. (2017). Urbanization and disease emergence: Dynamics at the wildlife-livestock-human interface. *Trends in Ecology and Evolution*, 32(1), 55–67. https://doi. org/10.1016/j.tree.2016.09.012
- Hawley, W. A. (1988). The biology of Aedes albopictus. Journal of the American Mosquito Control Association, 1–39.
- Hemming-Schroeder, E., Lo, E., Salazar, C., Puente, S., & Yan, G. (2018). Landscape genetics: A toolbox for studying vector-borne diseases. Frontiers in Ecology and Evolution, 6, 1–11. https://doi.org/10.3389/fevo.2018.00021
- Heylen, D., Lasters, R., Adriaensen, F., Fonville, M., Sprong, H., & Matthysen, E. (2019). Ticks and tick-borne diseases in the city: Role of landscape connectivity and green space characteristics in a metropolitan area. *Science of the Total Environment*, 670, 941–949. https://doi.org/10.1016/j.scitotenv.2019.03.235
- Honnen, A. C., Johnston, P. R., & Monaghan, M. T. (2016). Sex-specific gene expression in the mosquito *Culex pipiens* f. molestus in response to artificial light at night. *BMC Genomics*, 17(1), 1–10. https://doi.org/10.1186/s12864-015-2336-0
- Horobik, V., Keesing, F., & Ostfeld, R. S. (2006). Abundance and *Borrelia burgdorferi*-infection prevalence of nymphal *Ixodes scapularis* ticks along forest-field edges. *EcoHealth*, 3(4), 262–268. https://doi.org/10.1007/s10393-006-0065-1
- Hosseini, P. R., Mills, J. N., Prieur-Richard, A.-H., Ezenwa, V. O., Bailly, X., Rizzoli, A., Suzán, G., Vittecoq, M., García-Peña, G. E., Daszak, P., Guégan, J.-F., & Roche, B. (2017). Does the impact of biodiversity differ between emerging and endemic pathogens? The need to separate the concepts of hazard and risk. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1722). https://doi.org/10.1098/rstb.2016.0129
- Ignatieva, M. E., & Stewart, G. H. (2009). Landscape design in former colonial cities. In M. J. Mcdonnell, A. K. Hahs, & J. H. Breuste (Eds.),

- Ecology of cities and towns: a comparative approach (pp. 399–421). Cambridge University Press.
- Jennings, V., Browning, M. H. E. M., & Rigolon, A. (2019). Urban green spaces—Public health and sustainabilty in the United States.
- Jetz, W., Carbone, C., Fulford, J., & Brown, J. H. (2004). The scaling of animal space use. Science, 306, 266. https://doi.org/10.1126/scien ce.1102138
- Johnson, C. K., Hitchens, P. L., Pandit, P. S., Rushmore, J., Evans, T. S., Young, C. C. W., & Doyle, M. M. (2020). Global shifts in mammalian population trends reveal key predictors of virus spillover risk. *Proceedings of the Royal Society B: Biological Sciences*, 287(1924). https://doi.org/10.1098/rspb.2019.2736
- Johnson, S., Bragdon, C., & Olson, C. (2016). Characteristics of the built environment and the presence of the Norway rat in New York City: Results from a neighborhood rat surveillance program, 2008–2010. *Journal of Environmental Health*, 78(10), 22–29.
- Kache, P. A., Eastwood, G., Collins-Palmer, K., Katz, M., Falco, R. C., Bajwa, W. I., Armstrong, P. M., Andreadis, T. G., & Diuk-Wasser, M. A. (2020). Environmental determinants of Aedes albopictus abundance at a northern limit of its range in the United States. American Journal of Tropical Medicine and Hygiene, 102(2), 436–447. https://doi.org/10.4269/ajtmh.19-0244
- Karesh, W. B., Dobson, A., Lloyd-Smith, J. O., Lubroth, J., Dixon, M. A., Bennett, M., Aldrich, S., Harrington, T., Formenty, P., Loh, E. H., Machalaba, C. C., Thomas, M. J., & Heymann, D. L. (2012). Ecology of zoonoses: Natural and unnatural histories. *The Lancet*, 380(9857), 1936–1945. https://doi.org/10.1016/S0140-6736(12)61678-X
- Katz, G., Leisnham, P. T., & Ladeau, S. L. (2020). Aedes albopictus body size differs across neighborhoods with varying infrastructural abandonment. *Journal of Medical Entomology*, 57(2), 615–619. https://doi.org/10.1093/jme/tjz170
- Keeler, B. L., Hamel, P., McPhearson, T., Hamann, M. H., Donahue, M. L., Meza Prado, K. A., Arkema, K. K., Bratman, G. N., Brauman, K. A., Finlay, J. C., Guerry, A. D., Hobbie, S. E., Johnson, J. A., MacDonald, G. K., McDonald, R. I., Neverisky, N., & Wood, S. A. (2019). Socialecological and technological factors moderate the value of urban nature. *Nature Sustainability*, 2(1), 29–38. https://doi.org/10.1038/ s41893-018-0202-1
- Keesing, F., Belden, L. K., Daszak, P., Dobson, A., Harvell, C. D., Holt, R. D., Hudson, P., Jolles, A., Jones, K. E., Mitchell, C. E., Myers, S. S., Bogich, T., & Ostfeld, R. S. (2010). Impacts of biodiversity on the emergence and transmission of infectious diseases. *Nature*, 468(7324), 647–652. https://doi.org/10.1038/nature09575
- Kernbach, M. E., Martin, L. B., Unnasch, T. R., Hall, R. J., Jiang, R. H. Y., & Francis, C. D. (2021). Light pollution affects West Nile virus exposure risk across Florida. Proceedings of the Royal Society B: Biological Sciences, 288(1947), https://doi.org/10.1098/rspb.2021.0253
- Kilpatrick, A. M., Dobson, A. D. M., Levi, T., Salkeld, D. J., Swei, A., Ginsberg, H. S., Kjemtrup, A., Padgett, K. A., Jensen, P. M., Fish, D., Ogden, N. H., & Diuk-Wasser, M. A. (2017). Lyme disease ecology in a changing world: Consensus, uncertainty and critical gaps for improving control. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1722), 20160117. https://doi.org/10.1098/rstb.2016.0117
- Kilpatrick, A. M., Fonseca, D. M., Ebel, G. D., Reddy, M. R., & Kramer, L. D. (2010). Spatial and temporal variation in vector competence of Culex pipiens and Cx. restuans mosquitoes for West Nile virus. American Journal of Tropical Medicine and Hygiene, 83(3), 607–613. https://doi.org/10.4269/ajtmh.2010.10-0005
- Kilpatrick, A. M., Kramer, L. D., Jones, M. J., Marra, P. P., & Daszak, P. (2006). West Nile virus epidemics in North America are driven by shifts in mosquito feeding behavior. *PLoS Biology*, 4(4), 606–610. https://doi.org/10.1371/journal.pbio.0040082
- Kimmig, S. E., Beninde, J., Brandt, M., Schleimer, A., Kramer-Schadt, S., Hofer, H., Börner, K., Schulze, C., Wittstatt, U., Heddergott, M., Halczok, T., Staubach, C., & Frantz, A. C. (2020). Beyond the

- landscape: Resistance modelling infers physical and behavioural gene flow barriers to a mobile carnivore across a metropolitan area. *Molecular Ecology*, 29(3), 466–484. https://doi.org/10.1111/mec.15345
- Ko, A. I., Galvão Reis, M., Ribeiro Dourado, C. M., Johnson, W. D., & Riley, L. W. (1999). Urban epidemic of severe leptospirosis in Brazil. *Lancet*, 354(9181), 820–825. https://doi.org/10.1016/S0140-6736(99)80012-9
- Kramer, L. D., & Bernard, K. A. (2001). West nile virus in birds and mammals. Annals of the New York Academy of Sciences, 951(1), 84–93. https://doi.org/10.1111/j.1749-6632.2001.tb02687.x
- Krystosik, A., Njoroge, G., Odhiambo, L., Forsyth, J. E., Mutuku, F., & LaBeaud, A. D. (2020). Solid wastes provide breeding sites, burrows, and food for biological disease vectors, and urban zoonotic reservoirs: A call to action for solutions-based research. Frontiers in Public Health, 7, 1-17. https://doi.org/10.3389/fpubh.2019.00405
- LaDeau, S. L., Allan, B. F., Leisnham, P. T., & Levy, M. Z. (2015). The ecological foundations of transmission potential and vector-borne disease in urban landscapes. Functional Ecology, 29(7), 889–901. https://doi.org/10.1111/1365-2435.12487
- LaDeau, S. L., Leisnham, P. T., Biehler, D., & Bodner, D. (2013). Higher mosquito production in low-income neighborhoods of baltimore and washington, DC: Understanding ecological drivers and mosquito-borne disease risk in temperate cities. *International Journal of Environmental Research and Public Health*, 10(4), 1505– 1526. https://doi.org/10.3390/ijerph10041505
- Lambin, E. F., Tran, A., Vanwambeke, S. O., Linard, C., & Soti, V. (2010). Pathogenic landscapes: Interactions between land, people, disease vectors, and their animal hosts. *International Journal of Health Geographics*, 9(5), 54. https://doi.org/10.1186/1476-072X-9-54
- Lau, C. L., Smythe, L. D., Craig, S. B., & Weinstein, P. (2010). Climate change, flooding, urbanisation and leptospirosis: Fuelling the fire? Transactions of the Royal Society of Tropical Medicine and Hygiene, 104(10), 631–638. https://doi.org/10.1016/j.trstmh.2010.07.002
- Leibler, J. H., Robb, K., Joh, E., Gaeta, J. M., & Rosenbaum, M. (2018). Self-reported animal and ectoparasite exposure among urban homeless people. *Journal of Health Care for the Poor and Underserved*, 29(2), 664–675. https://doi.org/10.1353/hpu.2018.0050
- Little, E., Bajwa, W., & Shaman, J. (2017). Local environmental and meteorological conditions influencing the invasive mosquito Ae. albopictus and arbovirus transmission risk in New York City. PLoS Neglected Tropical Diseases, 11(8), 1–19. https://doi.org/10.1371/journal.pntd.0005828
- Lloyd-Smith, J. O., George, D., Pepin, K. M., Pitzer, V. E., Pulliam, J. R. C., Dobson, A. P., Hudson, P. J., & Grenfell, B. T. (2009). Epidemic dynamics at the human-animal interface. *Science*, 326(5958), 1362–1367. https://doi.org/10.1126/science.1177345
- LoGiudice, K., Ostfeld, R. S., Schmidt, K. A., & Keesing, F. (2003). The ecology of infectious disease: Effects of host diversity and community composition on Lyme disease risk. *Proceedings of the National Academy of Sciences*, 100(2), 567–571. https://doi.org/10.1073/pnas.0233733100
- Lõhmus, M., & Balbus, J. (2015). Making green infrastructure healthier infrastructure. *Infection Ecology & Epidemiology*, *5*(1), 30082. https://doi.org/10.3402/iee.v5.30082
- Loss, S. R., Hamer, G. L., Walker, E. D., Ruiz, M. O., Goldberg, T. L., Kitron, U. D., & Brawn, J. D. (2009). Avian host community structure and prevalence of West Nile virus in Chicago, Illinois. *Oecologia*, 159(2), 415–424. https://doi.org/10.1007/s00442-008-1224-6
- Loughran, K. (2020). Urban parks and urban problems: An historical perspective on green space development as a cultural fix. *Urban Studies*, 57(11), 2321–2338. https://doi.org/10.1177/0042098018 763555
- Lowry, H., Lill, A., & Wong, B. B. M. (2012). Behavioural responses of wildlife to urban environments. *Biological Reviews*, 88(3), 537–549. https://doi.org/10.1111/brv.12012

- Luederitz, C., Brink, E., Gralla, F., Hermelingmeier, V., Meyer, M., Niven, L., Panzer, L., Partelow, S., Rau, A.-L., Sasaki, R., Abson, D. J., Lang, D. J., Wamsler, C., & von Wehrden, H. (2015). A review of urban ecosystem services: Six key challenges for future research. *Ecosystem Services*, 14, 98-112. https://doi.org/10.1016/j. ecoser.2015.05.001
- Lydecker, H. W., Hochuli, D. F., & Banks, P. B. (2019). Peri-urban black rats host a rich assembly of ticks and healthier rats have more ticks. *Ticks and Tick-Borne Diseases*, 10(4), 749–753. https://doi.org/10.1016/j.ttbdis.2019.03.007
- Mackenstedt, U., Jenkins, D., & Romig, T. (2015). The role of wildlife in the transmission of parasitic zoonoses in peri-urban and urban areas. *International Journal for Parasitology: Parasites and Wildlife*, 4(1), 71–79. https://doi.org/10.1016/j.ijppaw.2015.01.006
- Magle, S. B., Fidino, M., Lehrer, E. W., Gallo, T., Mulligan, M. P., Ríos, M. J., Ahlers, A. A., Angstmann, J., Belaire, A., Dugelby, B., Gramza, A., Hartley, L., MacDougall, B., Ryan, T., Salsbury, C., Sander, H., Schell, C., Simon, K., St Onge, S., & Drake, D. (2019). Advancing urban wild-life research through a multi-city collaboration. Frontiers in Ecology and the Environment, 17(4), 232–239. https://doi.org/10.1002/fee.2030
- Mallin, M. A., Williams, K. E., Esham, E. C., & Lowe, R. P. (2000). Effect of human development on bacteriological water quality in coastal watersheds. *Ecological Applications*, 10(4), 1047–1056.
- Mancini, F., Di Luca, M., Toma, L., Vescio, F., Bianchi, R., Khoury, C., Marini, L., Rezza, G., & Ciervo, A. (2014). Prevalence of tick-borne pathogens in an urban park in Rome, Italy. Annals of Agricultural and Environmental Medicine, 21(4), 723–727. https://doi.org/10.5604/12321966.1129922
- Masi, E., Pino, F. A., Santos, M. D. G. S., Genehr, L., Albuquerque, J. O. M., Bancher, A. M., & Alves, J. C. M. (2010). Socioeconomic and environmental risk factors for urban rodent infestation in Sao Paulo, Brazil. *Journal of Pest Science*, 83(3), 231–241. https://doi.org/10.1007/s10340-010-0290-9
- Mayor, O. (2017). De Blasio Administration Announces \$32 Million Neighborhood Rat Reduction Plan. https://www1.nyc.gov/offic e-of-the-mayor/news/472-17/de-blasio-administration-32-million-neighborhood-rat-reduction-plan#/0
- McGarigal, K., Wan, H. Y., Zeller, K. A., Timm, B. C., & Cushman, S. A. (2016). Multi-scale habitat selection modeling: a review and outlook. *Landscape Ecology*, 31(6), 1161–1175. https://doi.org/10.1007/s10980-016-0374-x
- McGinlay, J., Gowing, D. J. G., & Budds, J. (2016). Conserving socioecological landscapes: An analysis of traditional and responsive management practices for floodplain meadows in England. *Environmental Science and Policy*, 66, 234–241. https://doi. org/10.1016/j.envsci.2016.07.008
- McKinney, M. L. (2006). Urbanization as a major cause of biotic homogenization. *Biological Conservation*, 127(3), 247–260. https://doi.org/10.1016/j.biocon.2005.09.005
- Mckinney, M. L. (2008). Effects of urbanization on species richness: A review of plants and animals. *Urban Ecosystems*, 11(2), 161–176. https://doi.org/10.1007/s11252-007-0045-4
- McMahon, T. A., Rohr, J. R., & Bernal, X. E. (2017). Light and noise pollution interact to disrupt interspecific interactions. *Ecology*, *98*(5), 1290–1299. https://doi.org/10.1002/ecy.1770
- McMillan, J. R., Armstrong, P. M., & Andreadis, T. G. (2020). Patterns of mosquito and arbovirus community composition and ecological indexes of arboviral risk in the Northeast United States. PLoS Neglected Tropical Diseases, 14(2), 1–21. https://doi.org/10.1371/ journal.pntd.0008066
- Morandeira, N. S., Castesana, P. S., Cardo, M. V., Salomone, V. N., Vadell, M. V., & Rubio, A. (2019). An interdisciplinary approach to assess human health risk in an urban environment: A case study in temperate Argentina. *Heliyon*, *5*(10), e02555. https://doi.org/10.1016/j. heliyon.2019.e02555

- Morlan, H. B., & Hayes, R. O. (1958). Urban dispersal and activity of *Aedes aegypti. Mosquito News*, 18(2), 137–144.
- Moyers, S. C., Adelman, J. S., Farine, D. R., Thomason, C. A., & Hawley, D. M. (2018). Feeder density enhances house finch disease transmission in experimental epidemics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1745). https://doi.org/10.1098/rstb.2017.0090
- Muir, L. E., & Kay, B. H. (1998). Aedes aegypti survival and dispersal estimated by mark-release-recapture in northern Australia. American Journal of Tropical Medicine and Hygiene, 58(3), 277–282. https://doi.org/10.4269/ajtmh.1998.58.277
- Munshi-South, J. (2012). Urban landscape genetics: Canopy cover predicts gene flow between white-footed mouse (*Peromyscus leucopus*) populations in New York City. *Molecular Ecology*, 21(6), 1360–1378. https://doi.org/10.1111/j.1365-294X.2012.05476.x
- Murray, M. H., Becker, D. J., Hall, R. J., & Hernandez, S. M. (2016). Wildlife health and supplemental feeding: A review and management recommendations. *Biological Conservation*, 204, 163–174. https://doi. org/10.1016/j.biocon.2016.10.034
- Murray, M. H., Fidino, M., Fyffe, R., Byers, K. A., Pettengill, J. B., Sondgeroth, K. S., Killion, H., Magle, S. B., Rios, M. J., Ortinau, N., & Santymire, R. M. (2020). City sanitation and socioeconomics predict rat zoonotic infection across diverse neighbourhoods. *Zoonoses and Public Health*, 67(6), 673–683. https://doi.org/10.1111/zph.12748
- Naing, C., Reid, S. A., Aye, S. N., Htet, N. H., & Ambu, S. (2019). Risk factors for human leptospirosis following flooding: A meta-analysis of observational studies. *PLoS One*, 14(5), 1–15. https://doi. org/10.1371/journal.pone.0217643
- Neiderud, C. J. (2015). How urbanization affects the epidemiology of emerging infectious diseases. African Journal of Disability, 5(1). https://doi.org/10.3402/iee.v5.27060
- NYC Department of Health and Mental Hygiene. (2019). Communicable disease surveillance data.
- Ogden, N. H., Lindsay, L. R., Hanincová, K., Barker, I. K., Bigras-Poulin, M., Charron, D. F., & Thompson, R. A. (2008). Role of migratory birds in introduction and range expansion of *Ixodes scapularis* ticks and of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in Canada. *Applied and Environmental Microbiology*, 74(6), 1780–1790. https:// doi.org/10.1128/AEM.01982-07
- Ogden, N. H., & Tsao, J. I. (2009). Biodiversity and Lyme disease: Dilution or amplification? *Epidemics*, 1(3), 196–206. https://doi.org/10.1016/j.epidem.2009.06.002
- Ostfeld, R. S., Glass, G. E., & Keesing, F. (2005). Spatial epidemiology: An emerging (or re-emerging) discipline. *Trends in Ecology and Evolution*, 20(6), 328–336. https://doi.org/10.1016/j.tree.2005.03.009
- Ostrom, E. (2018). A general framework for analyzing sustainability of social-ecological systems. *Science*, 362, 763.
- Pacheco-Tucuch, F. S., Ramirez-Sierra, M. J., Gourbière, S., & Dumonteil, E. (2012). Public street lights increase house infestation by the chagas disease vector *Triatoma dimidiata*. *PLoS One*, *7*(4), 3–9. https://doi.org/10.1371/journal.pone.0036207
- Padayachee, A. L., Irlich, U. M., Faulkner, K. T., Gaertner, M., Procheş, Ş., Wilson, J. R. U., & Rouget, M. (2017). How do invasive species travel to and through urban environments? *Biological Invasions*, 19(12), 3557–3570. https://doi.org/10.1007/s10530-017-1596-9
- Paull, S. H., Song, S., McClure, K. M., Sackett, L. C., Kilpatrick, A. M., & Johnson, P. T. J. (2012). From superspreaders to disease hotspots: Linking transmission across hosts and space. Frontiers in Ecology and the Environment, 10(2), 75–82. https://doi.org/10.1890/110111
- Perugini, M., Manera, M., Grotta, L., Abete, M. C., Tarasco, R., & Amorena, M. (2011). Heavy metal (Hg, Cr, Cd, and Pb) contamination in urban areas and wildlife reserves: Honeybees as bioindicators. *Biological Trace Element Research*, 140(2), 170–176. https://doi.org/10.1007/s12011-010-8688-z
- Peters, W., Hebblewhite, M., Mysterud, A., Eacker, D., Hewison, A. J. M., Linnell, J. D. C., Focardi, S., Urbano, F., De Groeve, J., Gehr, B.,

- Heurich, M., Jarnemo, A., Kjellander, P., Kröschel, M., Morellet, N., Pedrotti, L., Reinecke, H., Sandfort, R., Sönnichsen, L., ... Cagnacci, F. (2019). Large herbivore migration plasticity along environmental gradients in Europe: Life-history traits modulate forage effects. *Oikos*, 128(3), 416–429. https://doi.org/10.1111/oik.05588
- Peterson, A. C., Ghersi, B. M., Campanella, R., Riegel, C., Lewis, J. A., & Blum, M. J. (2020). Rodent assemblage structure reflects socioecological mosaics of counter-urbanization across post-Hurricane Katrina New Orleans. *Landscape and Urban Planning*, 195, 103710. https://doi.org/10.1016/j.landurbplan.2019.103710
- Peterson, A. C., Ghersi, B. M., Riegel, C., Wunder, E. A., Childs, J. E., & Blum, M. J. (2020). Amplification of pathogenic *Leptospira* infection with greater abundance and co-occurrence of rodent hosts across a counter-urbanizing landscape. *Molecular Ecology*. https://doi.org/10.1111/mec.15710
- Pickett, S., Cadenasso, M. L., Grove, J. M., Boone, C. G., Groffman, P. M., Irwin, E., Kaushal, S. S., Marshall, V., McGrath, B. P., Nilon, C. H., Pouyat, R. V., Szlavecz, K., Troy, A., & Warren, P. (2011). Urban ecological systems: Scientific foundations and a decade of progress. *Journal of Environmental Management*, 92(3), 331–362. https://doi.org/10.1016/j.jenvman.2010.08.022
- Pickett, S. T. A., Cadenasso, M. L., Grove, J. M., Nilon, C. H., Pouyat, R. V., Zipperer, W. C., & Costanza, R. (2001). Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas. *Annual Review of Ecology and Systematics*, 32(1), 127–157. https://doi.org/10.1146/annurev.ecolsys.32.081501.114012
- Pless, E., Saarman, N. P., Powell, J. R., Caccone, A., & Amatulli, G. (2021). A machine-learning approach to map landscape connectivity in Aedes aegypti with genetic and environmental data. *Proceedings of the National Academy of Sciences of the United States of America*, 118(9), 1–8. https://doi.org/10.1073/pnas.2003201118
- Plowright, R. K., Foley, P., Field, H. E., Dobson, A. P., Foley, J. E., Eby, P., & Daszak, P. (2011). Urban habituation, ecological connectivity and epidemic dampening: The emergence of hendra virus from flying foxes (Pteropus spp.). Proceedings of the Royal Society B: Biological Sciences, 278(1725), 3703–3712. https://doi.org/10.1098/rspb. 2011.0522
- Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., & Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews Microbiology*, 15(8), 502–510. https://doi.org/10.1038/nrmicro.2017.45
- Plowright, R. K., Reaser, J. K., Locke, H., Woodley, S. J., Patz, J. A., Becker, D. J., & Hudson, P. J. (2021). Land use-induced spillover: A call to action to safeguard environmental, animal, and human health. *The Lancet Planetary Health*, *5*(4), e237–e245. https://doi.org/10.1016/S2542-5196(21)00031-0
- Plumer, L., Davison, J., Saarma, U., & Cameron, E. Z. (2014). Rapid urbanization of red foxes in estonia: Distribution, behaviour, attacks on domestic animals, and health-risks related to zoonotic diseases. *PLoS One*, 9(12), 1–15. https://doi.org/10.1371/journal.pone.0115124
- Poh, K. C., Medeiros, M. C. I., & Hamer, G. L. (2020). Landscape and demographic determinants of Culex infection with West Nile virus during the 2012 epidemic in Dallas County, TX. Spatial and Spatio-Temporal Epidemiology, 33, 100336. https://doi.org/10.1016/j.sste.2020.100336
- Rainwater, K. L., Marchese, K., Slavinski, S., Humberg, L. A., Dubovi, E. J., Jarvis, J. A., & Calle, P. P. (2017). Health survey of free-ranging raccoons (*Procyon lotor*) in central park, New York, New York, USA: Implications for human and domestic animal health. *Journal of Wildlife Diseases*, 53(2), 272–284. https://doi.org/10.7589/2016-05-096
- Ramaswami, A., Russell, A. G., Culligan, P. J., Rahul Sharma, K., & Kumar, E. (2016). Meta-principles for developing smart, sustainable, and healthy cities. *Science*, *352*(6288), 940–943. https://doi.org/10.1126/science.aaf7160

- Randolph, S. E., & Dobson, A. D. M. (2012). Pangloss revisited: A critique of the dilution effect and the biodiversity-buffers-disease paradigm. *Parasitology*, 139(7), 847–863. https://doi.org/10.1017/s0031182012000200
- Reed, E. M. X., Serr, M. E., Maurer, A. S., & Burford Reiskind, M. O. (2020). Gridlock and beltways: The genetic context of urban invasions. *Oecologia*, 192(3), 615–628. https://doi.org/10.1007/s00442-020-04614-v
- Reisen, W. K. (2013). Ecology of West Nile virus in North America. Viruses, 5(9), 2079-2105. https://doi.org/10.3390/v5092079
- Richardson, J. L., Brady, S. P., Wang, I. J., & Spear, S. F. (2016). Navigating the pitfalls and promise of landscape genetics. *Molecular Ecology*, 25(4), 849–863. https://doi.org/10.1111/mec.13527
- Richardson, J. L., Burak, M. K., Hernandez, C., Shirvell, J. M., Mariani, C., Carvalho-Pereira, T. S. A., Pertile, A. C., Panti-May, J. A., Pedra, G. G., Serrano, S., Taylor, J., Carvalho, M., Rodrigues, G., Costa, F., Childs, J. E., Ko, A. I., & Caccone, A. (2017). Using fine scale spatial genetics of Norway rats to improve control efforts and reduce leptospirosis risk in urban slum environments. Evolutionary Applications, 10(4), 323-337. https://doi.org/10.1111/eva.12449
- Rizwan, A. M., Dennis, L. Y. C., & Liu, C. (2008). A review on the generation, determination and mitigation of Urban Heat Island. *Journal of Environmental Sciences*, 20(1), 120–128. https://doi.org/10.1016/S1001-0742(08)60019-4
- Rodríguez Martín, J. A., De Arana, C., Ramos-Miras, J. J., Gil, C., & Boluda, R. (2015). Impact of 70 years urban growth associated with heavy metal pollution. *Environmental Pollution*, 196, 156–163. https://doi. org/10.1016/j.envpol.2014.10.014
- Rohr, J. R., Civitello, D. J., Halliday, F. W., Hudson, P. J., Lafferty, K. D., Wood, C. L., & Mordecai, E. A. (2020). Towards common ground in the biodiversity-disease debate. *Nature Ecology and Evolution*, 4(1), 24–33. https://doi.org/10.1038/s41559-019-1060-6
- Rose, N. H., Sylla, M., Badolo, A., Lutomiah, J., Ayala, D., Aribodor, O. B., Ibe, N., Akorli, J., Otoo, S., Mutebi, J.-P., Kriete, A. L., Ewing, E. G., Sang, R., Gloria-Soria, A., Powell, J. R., Baker, R. E., White, B. J., Crawford, J. E., & McBride, C. S. (2020). Climate and Urbanization Drive Mosquito Preference for Humans. *Current Biology*, 1-10. https://doi.org/10.1016/j.cub.2020.06.092
- Rothenburger, J. L., Himsworth, C. H., Nemeth, N. M., Pearl, D. L., & Jardine, C. M. (2017). Environmental factors and zoonotic pathogen ecology in urban exploiter species. *EcoHealth*, *14*(3), 630–641. https://doi.org/10.1007/s10393-017-1258-5
- Ryan, S. J., Carlson, C. J., Mordecai, E. A., & Johnson, L. R. (2018). Global expansion and redistribution of Aedes-borne virus transmission risk with climate change. *PLoS Neglected Tropical Diseases*, 13(3), 1–20. https://doi.org/10.1371/journal.pntd.0007213
- Rydin, Y., Bleahu, A., Davies, M., Davila, J. D., Friel, S., di Grandis, G., & Wilson, J. (2012). Shaping cities for health: The complexities of planning urban environments in the 21st century. *The Lancet*, *379*, 2079–2108. https://doi.org/10.1016/S0140-6736(12)60435-8.Shaping
- Salkeld, D. J., Padgett, K. A., & Jones, J. H. (2013). A meta-analysis suggesting that the relationship between biodiversity and risk of zoonotic pathogen transmission is idiosyncratic. *Ecology Letters*, 16(5), 679–686. https://doi.org/10.1111/ele.12101
- Samson, D. M., Qualls, W. A., Roque, D., Naranjo, D. P., Alimi, T., Arheart, K. L., Müller, G. C., Beier, J. C., & Xue, R.-D. (2013). Resting and energy reserves of Aedes albopictus collected in common land-scaping vegetation in St. Augustine, Florida. Journal of the American Mosquito Control Association, 29(3), 231–236. https://doi.org/10.2987/13-6347R.1
- Sánchez, C. A., Altizer, S., & Hall, R. J. (2020). Landscape-level toxicant exposure mediates infection impacts on wildlife populations: Toxicants and infection in wildlife. *Biology Letters*, 16(11). https://doi.org/10.1098/rsbl.2020.0559rsbl20200559
- Schell, C. J., Dyson, K., Fuentes, T. L., Des Roches, S., Harris, N. C., Miller, D. S., Woelfle-Erskine, C. A., & Lambert, M. R. (2020). The

- ecological and evolutionary consequences of systemic racism in urban environments. *Science*, *369*(6509). https://doi.org/10.1126/SCIENCE.AAY4497
- Sejvar, J. J. (2003). West Nile virus: An historical overview. *Ochsner Journal*, 5(3), 6-10.
- Seto, K. C., Güneralp, B., & Hutyra, L. R. (2012). Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences of the United States of America*, 109(40), 16083–16088. https://doi.org/10.1073/pnas.1211658109
- Shackleton, C. M., & Gwedla, N. (2021). The legacy effects of colonial and apartheid imprints on urban greening in South Africa: Spaces, species, and suitability. *Frontiers in Ecology and Evolution*, 8. https://doi.org/10.3389/fevo.2020.579813
- Shackleton, C. M., Ruwanza, S., Sinasson Sanni, G. K., Bennett, S., De Lacy, P., Modipa, R., Mtati, N., Sachikonye, M., & Thondhlana, G. (2016). Unpacking Pandora's box: Understanding and categorising ecosystem disservices for environmental management and human wellbeing. *Ecosystems*, 19(4), 587–600. https://doi.org/10.1007/s10021-015-9952-z
- Shochat, E., Lerman, S. B., Anderies, J. M., Warren, P. S., Faeth, S. H., & Nilon, C. H. (2010). Invasion, competition, and biodiversity loss in urban ecosystems. *BioScience*, 60(3), 199–208. https://doi. org/10.1525/bio.2010.60.3.6
- Shragai, T., & Harrington, L. C. (2019). Aedes albopictus (Diptera: Culicidae) on an invasive edge: Abundance, spatial distribution, and habitat usage of larvae and pupae across urban and socioeconomic environmental gradients. Journal of Medical Entomology, 56(2), 472–482. https://doi.org/10.1093/jme/tjy209
- Shuster, W. D., Bonta, J., Thurston, H., Warnemuende, E., & Smith, D. R. (2005). Impacts of impervious surface on watershed hydrology: A review. *Urban Water Journal*, 2(4), 263–275. https://doi.org/10.1080/15730620500386529
- Singh, S., Tandia, N., Kumar, N., Paliwal, R., & Panicker, A. (2014). Environmental pollution and wildlife health. *Pharma Science Monitor*, 5(2), 169–175.
- Slavinski, S., Humberg, L., Lowney, M., Simon, R., Calvanese, N., Bregman, B., Kass, D., & Oleszko, W. (2012). Trap-vaccinate- release program to control raccoon rabies, New York, USA. Emerging Infectious Diseases, 18(7), 1170-1172. https://doi.org/10.3201/ eid1807.111485
- Smith, G. C., Gangadharan, B., Taylor, Z., Laurenson, M. K., Bradshaw, H., Hide, G., & Craig, P. S. (2003). Prevalence of zoonotic important parasites in the red fox (Vulpes vulpes) in Great Britain. Veterinary Parasitology, 118(1–2), 133–142. https://doi.org/10.1016/j.vetpar.2003.09.017
- Solar, O., & Irwin, A. (2014). A conceptual framework for action on the social determinants of health: Discussion paper for the Commission on Social Determinants of Health (DISCUSSION PAPER).
- Soulsbury, C. D., & White, P. C. L. (2015). Human-wildlife interactions in urban areas: A review of conflicts, benefits and opportunities. *Wildlife Research*, 42(7), 541–553. https://doi.org/10.1071/WR14229
- Steere, A. C. (1994). Lyme disease: A growing threat to urban populations. Proceedings of the National Academy of Sciences of the United States of America, 91(7), 2378–2383. https://doi.org/10.1073/pnas.91.7.2378
- Swaileh, K. M., & Sansur, R. (2006). Monitoring urban heavy metal pollution using the house sparrow (Passer domesticus). *Journal of Environmental Monitoring*, 8(1), 209–213. https://doi.org/10.1039/b510635d
- Taguchi, V., Weiss, P., Gulliver, J., Klein, M., Hozalski, R., Baker, L., Finlay, J., Keeler, B., & Nieber, J. (2020). It is not easy being green: Recognizing unintended consequences of green stormwater infrastructure. Water, 12(2). https://doi.org/10.3390/w12020522

- Trajer, A., Mlinarik, L., Juhasz, P., & Bede-Fazekas, A. (2014). The combined impact of urban heat island, thermal bridge effect of buildings and future climate change on the potential overwintering of Phlebotomus species in a Central European metropolis. *Applied Ecology and Environmental Research*, 12(4), 887–908. https://doi.org/10.15666/aeer/1204\_887908
- Traweger, D., Travnitzky, R., Moser, C., Walzer, C., & Bernatzky, G. (2006). Habitat preferences and distribution of the brown rat (*Rattus norvegicus Berk.*) in the city of Salzburg (Austria): implications for an urban rat management. *Journal of Pest Science*, 79(3), 113–125. https://doi.org/10.1007/s10340-006-0123-z
- Tucker, M. A., Böhning-Gaese, K., Fagan, W. F., Fryxell, J. M., Van Moorter, B., Alberts, S. C., Ali, A. H., Allen, A. M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar, B., Belant, J. L., Bertassoni, A., Beyer, D., Bidner, L., van Beest, F. M., Blake, S., Blaum, N., ... Mueller, T. (2018). Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science*, 359, 466-469. https://doi.org/10.1126/science.aam9712
- Tucker, M. A., Ord, T. J., & Rogers, T. L. (2014). Evolutionary predictors of mammalian home range size: Body mass, diet and the environment. Global Ecology and Biogeography, 23(10), 1105–1114. https://doi. org/10.1111/geb.12194
- Tuff, K. T., Tuff, T., & Davies, K. F. (2016). A framework for integrating thermal biology into fragmentation research. *Ecology Letters*, 19(4), 361–374. https://doi.org/10.1111/ele.12579
- Tufts, D. M., Goodman, L. B., Benedict, M. C., Davis, A. D., VanAcker, M. C., & Diuk-Wasser, M. (2021). Association of the invasive Haemaphysalis longicornis tick with vertebrate hosts, other native tick vectors, and tick-borne pathogens in New York City, USA. *International Journal for Parasitology*, 51(2-3), 149–157. https://doi.org/10.1016/j.ijpara.2020.08.008
- Tyungu, D. L., McCormick, D., Lau, C. L., Chang, M., Murphy, J. R., Hotez, P. J., & Pollack, H. (2020). Toxocara species environmental contamination of public spaces in New York city. PLoS Neglected Tropical Diseases, 14(5), 1–13. https://doi.org/10.1371/journal.pntd.0008249
- United Nations. (2018). World urbanization prospects: The 2018 revision. United Nations Department of Economic and Social Affairs. https://doi.org/10.4054/demres.2005.12.9
- Valerio, L., Marini, F., Bongiorno, G., Facchinelli, L., Pombi, M., Caputo, B., & della Torre, A. (2008). Blood-feeding preferences of Aedes albopictus (Diptera: Culicidae) in urban and rural settings within the province of Rome, Italy. Parassitologia, 50(1-2), 103-104.
- van Adrichem, M. H. C., Buijs, J. A., Goedhart, P. W., & Verboom, J. (2013). Factors influencing the density of the brown rat (*Rattus norvegicus*) in and around houses in Amsterdam. *Lutra Journal of the Dutch Mammal Society*, 56(2), 77–91.
- VanAcker, M. C., Little, E. A. H., Molaei, G., Bajwa, W. I., & Diuk-Wasser, M. A. (2019). Enhancement of risk for lyme disease by landscape connectivity, New York, New York, USA. *Emerging Infectious Diseases*, 25(6), 1136-1143. https://doi.org/10.3201/eid2506.181741
- Vanwambeke, S. O., Linard, C., & Gilbert, M. (2019). Emerging challenges of infectious diseases as a feature of land systems. Current Opinion in Environmental Sustainability, 38, 31–36. https://doi.org/10.1016/j. cosust.2019.05.005
- VanWormer, E., Carpenter, T. E., Singh, P., Shapiro, K., Wallender, W. W., Conrad, P. A., Largier, J. L., Maneta, M. P., & Mazet, J. A. K. (2016). Coastal development and precipitation drive pathogen flow from land to sea: Evidence from a Toxoplasma gondii and felid host system. Scientific Reports, 6, 1–9. https://doi.org/10.1038/srep29252
- Venter, Z. S., Krog, N. H., & Barton, D. N. (2020). Linking green infrastructure to urban heat and human health risk mitigation in Oslo,

- Norway. Science of the Total Environment, 709, 136193, https://doi.org/10.1016/j.scitotenv.2019.136193
- Venter, Z. S., Shackleton, C. M., Van Staden, F., Selomane, O., & Masterson, V. A. (2020). Green Apartheid: Urban green infrastructure remains unequally distributed across income and race geographies in South Africa. *Landscape and Urban Planning*, 203(0349), 103889. https://doi.org/10.1016/j.landurbplan.2020.103889.
- Vinogradova, E. B. (2000). Culex pipiens pipiens mosquitoes: taxonomy, distribution, ecology, physiology, genetics, applied importance and control. Issue 2. Pensoft Publishers.
- vonHedemann, N., Butterworth, M. K., Robbins, P., Landau, K., & Morin, C. W. (2015). Visualizations of mosquito risk: A political ecology approach to understanding the territorialization of hazard control. *Landscape and Urban Planning*, 142, 159–169. https://doi.org/10.1016/j.landurbplan.2015.03.001
- Walsh, M. G. (2014). Rat sightings in New York City are associated with neighborhood sociodemographics, housing characteristics, and proximity to open public space. *PeerJ*, 2, e533. https://doi.org/10.7717/peeri.533
- White, R. J., & Razgour, O. (2020). Emerging zoonotic diseases originating in mammals: A systematic review of effects of anthropogenic land-use change. *Mammal Review*, 50(4), 336–352. https://doi.org/10.1111/mam.12201
- WildlifeNYC. (2021). Animals in NYC. Retrieved from https://www1.nyc. gov/site/wildlifenyc/animals/animals.page
- Williams, S. H., Che, X., Paulick, A., Guo, C., Lee, B., Muller, D., Uhlemann, A.-C., Lowy, F. D., Corrigan, R. M., & Lipkin, W. I. (2018). New York City house mice (*Mus musculus*) as potential reservoirs for pathogenic bacteria and antimicrobial resistance determinants. *MBio*, 9(2), 1–17. https://doi.org/10.1128/mBio.00624-18
- Witmer, G. W. (2007). The ecology of vertebrate pests and integrated pest management (IPM), (August).
- Wood, C. L., & Lafferty, K. D. (2013). Biodiversity and disease: A synthesis of ecological perspectives on Lyme disease transmission. *Trends in Ecology and Evolution*, 28(4), 239–247. https://doi.org/10.1016/j.tree.2012.10.011
- Wright, A. N., & Gompper, M. E. (2005). Altered parasite assemblages in raccoons in response to manipulated resource availability. *Oecologia*, 144(1), 148–156. https://doi.org/10.1007/s0044 2-005-0018-3
- Youngsteadt, E., Ernst, A. F., Dunn, R. R., & Frank, S. D. (2017). Responses of arthropod populations to warming depend on latitude: evidence from urban heat islands. *Global Change Biology*, 23(4), 1436–1447. https://doi.org/10.1111/gcb.13550
- Zolnik, C. P., Falco, R. C., Kolokotronis, S. O., & Daniels, T. J. (2015). No observed effect of landscape fragmentation on pathogen infection prevalence in blacklegged ticks (*Ixodes scapularis*) in the Northeastern United States. *PLoS One*, 10(10), 1–12. https://doi.org/10.1371/journal.pone.0139473

How to cite this article: Combs, M. A., Kache, P. A., VanAcker, M. C., Gregory, N., Plimpton, L. D., Tufts, D. M., Fernandez, M. P., & Diuk-Wasser, M. A. (2022). Socio-ecological drivers of multiple zoonotic hazards in highly urbanized cities. *Global Change Biology*, 28, 1705–1724. <a href="https://doi.org/10.1111/gcb.16033">https://doi.org/10.1111/gcb.16033</a>