

Reconstructed evolutionary patterns for crocodile-line archosaurs demonstrate impact of failure to log-transform body size data

Roger B. J. Benson¹✉, Pedro Godoy^{2,3}, Mario Bronzati⁴, Richard J. Butler⁵ & William Gearty⁶

ARISING FROM M.T. Stockdale and M.J. Benton. *Communications Biology* <https://doi.org/10.1038/s42003-020-01561-5> (2021)

Pseudosuchia includes crocodylians, plus all extinct species more closely related to them than to birds. They appeared around 250 million years ago and have a rich fossil history, with extinct diversity exceeding that of their living members^{1–3}. Recently, Stockdale & Benton⁴ presented analyses of a new dataset of body size estimates spanning the entire evolutionary history of Pseudosuchia. They quantified patterns of average body size, body size disparity through time and rates of evolution along phylogenetic lineages. Their results suggest that pseudosuchians exhibited considerable variation in rates of body size evolution, for which they provided various group-specific explanations and asserted the importance of climatic drivers. This differs from two recent studies that analysed a substantial portion of pseudosuchian body size evolution and proposed that adaptation to aquatic life, a biological innovation of some subgroups, was the main driver of body size evolution, with patterns of disparity also being influenced by size-dependent extinction risk^{5,6}. Here we show that the analytical results of Stockdale & Benton⁴ are strongly influenced by a methodological error in their body size index. Specifically, that they chose not to log-transform measurement data prior to analyses.

Stockdale & Benton⁴ recorded 21 measurements across 280 species. Most of their measurements (17) were cranial and four were from the limb skeleton. They submitted these measurements to a principal components analysis (PCA) and iterative missing data estimation procedure⁷, then used principal component 1 (PC1) scores as a size index. This method was intended to overcome the problem of using skull length on its own as a size proxy, which may be biased by variation in relative head size and snout length⁴. Indeed, Stockdale & Benton⁴ suggested that this bias might explain key differences between their findings and the results of the previous studies^{5,6}. Nevertheless, their PC1 size index is highly correlated to skull length (Fig. 1 and see below; regardless of data treatment), and one previous study did also address the biasing effects of snout length variation by excluding the snout from skull length measurements⁵. Therefore, we did not expect such strongly different results on that basis alone. Instead, we argue that key differences result mainly from the fact that previous studies used log-transformed size indices^{5,6}, whereas Stockdale & Benton⁴ did not.

Measurements of real-world objects are taken in additive (or absolute) units such as millimetres (mm). This presents a well-understood statistical problem because biological variation is often multiplicative (=relative, or proportional), such that variance increases with increasing scale (heteroskedasticity; e.g. refs. ^{8,9}). For example, a hypothetical group of rodents has species-mean body masses ranging from 100 to 200 grams, whereas hypothetical artiodactyls range from 100 to 200 kg. Both groups exhibit identical (two-fold) relative variation, but this is not reflected when additive units are analysed because the absolute difference between the largest and smallest artiodactyls is 1000 times that in rodents. Therefore, size variation among large-bodied species is substantially over-weighted unless measurements are log-transformed (see refs. ^{8,9}). To ignore this, either by analysing or simulating trait data in a non-logged context, is to model a version of evolution in which it is as easy for a mouse population to evolve a body size increase of 10 kg as it is for an elephant population.

When using log units instead of additive units, identical proportional increases are indexed by identical numerical increases, entirely solving the problem (e.g. our rodents span from 2.0 to 2.3 log₁₀-grams, cf. 5.0–5.3 log₁₀-grams in artiodactyls). Therefore, evolutionary rate studies have routinely used log-transformed measurements for more than 70 years^{8–10}. This is especially important when measurements span across orders of magnitude⁹, as with pseudosuchians which range from the largest species, *Sarcosuchus imperator* (skull length = 1650 mm) to the smallest, *Knoetschkesuchus guimarotae* (33 mm). This represents an estimated 125,000-fold variation in body mass, approximating that body mass scales with the cube of linear dimensions (or 15,000-fold, conservatively reducing the skull of *Sarcosuchus* by 50% to account for its proportionally long snout). Although we illustrate the nature of this problem using extremes, it cannot be addressed just by excluding small-and large-bodied taxa from analyses, because variance increases as a continuous function of scale.

We replicated the analyses of Stockdale & Benton⁴, using log₁₀-transformed measurements. This resulted in a modified version of their PC1 size index that scales linearly with, and is strongly correlated to, log-transformed skull length (Fig. 1b; $p < 0.001$;

¹Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK. ²Departamento de Zoologia, Universidade Federal do Paraná, 81531-980 Curitiba, Paraná, Brazil. ³Department of Anatomical Sciences, Stony Brook University, Stony Brook, NY, USA. ⁴Department of Biology, University of Sao Paulo, 13840-091 Ribeirão Preto, SP, Brazil. ⁵School of Geography, Earth & Environmental Sciences, University of Birmingham, Birmingham B15 2TT, UK. ⁶School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA. ✉email: roger.benson@earth.ox.ac.uk




Fig. 1 Effects of log-transformation on the PC1 size index of Stockdale & Benton (2021). **a** Original (untransformed) version of the PC1 size index shows a curved relationship with log-transformed skull length (Spearman's $\rho = 0.98$, $p < 0.00001$, $N = 202$; using rank-based correlation due to the curved nature of the relationship). **b** Log-transformed version of the PC1 size index shows a linear relationship with log-transformed skull length (see text for correlation test results). **c** Evolutionary rates estimated from the original (untransformed) version of the PC1 size index are not independent of size. **d** Evolutionary rates estimated from the log-transformed version of the PC1 size index are independent of size.

$R^2 = 0.85$; $N = 202$; Pearson's product-moment). In contrast, the non-logged version⁴ shows a curved relationship (Fig. 1a) that upweights relative variation among large-bodied species—~80% of the variation in their index (y-axis) represents less than 25% of variation in relative (log-transformed) size. Due to space limitations, we focus only on macroevolutionary model comparisons and variation in evolutionary rates mapped to phylogeny. These analyses are central to the conclusions of Stockdale & Benton⁴ because they document variation in the tempo and mode of evolution among groups of different ages. Therefore, they provide process-based explanations that underpin the interpretation of their other analytical outputs, such as patterns of variation in disparity and average rates through time.

Re-analysis of rate variation among phylogenetic lineages¹¹ using the non-logged PC1 size index returns similar results to those of Stockdale & Benton⁴ (Fig. 2a). Evolutionary rates returned by this analysis are measured in additive (or absolute) length units per million years and high rates occur in two contexts: (1) On the lineages leading to some large-bodied species such as the notosuchian *Razanandrongobe*, the phytosaur *Angistorhinus*, and the early crocodylomorph *Carnufex*; and (2) In some clades or grades of generally large-bodied species, including various Triassic pseudosuchians, Tethysuchia, Thalattosuchia and some eusuchians. However, these instances of high evolutionary rates are artefacts resulting from an underlying correlation of high rates with large body sizes (Fig. 1c; $R^2 = 0.21$; $p < 0.001$; $N = 281$ species; phylogenetic least squares regression [PGLS]; $R^2 = 0.46$; $p < 0.001$; $N = 559$ species and nodes; ordinary

least squares regression [OLS]). This is expected when data are not log-transformed because using non-logged measurements inflates the amount of evolutionary change inferred to have occurred among large-bodied species.

Analysis of our log-transformed version of the PC1 size index documents rate variation in relative length units per million years and is uncorrelated or very weakly correlated with variation in absolute size (Fig. 1d; $R^2 = 0.03$; $p = 0.004$; $N = 281$ species; PGLS; $R^2 = 0.008$; $p = 0.19$; $N = 559$; OLS). This rate variation shows a very different pattern to that of Stockdale & Benton⁴ (Fig. 2b), rejecting the occurrence of high evolutionary rates in large-bodied groups and species, including those listed above. This removes the need for hypotheses such as early evolutionary radiation of Triassic pseudosuchians, island endemism in *Razanandrongobe*, and viviparity in thalattosuchians⁴. These processes may well be important drivers of phenotypic evolution and species diversification in those taxa, but they did not result in above-background rates of body size evolution. Instead, we find a much more even distribution of high rates through time and among large- and small-bodied lineages, including high rates involved in the attainment of small body size in groups such as atoposaurids and shartegosuchids.

Nevertheless, biological interpretation of these patterns should be avoided because we find no support for the variable-rate model compared to a uniform-rate Brownian motion (BM) model (marginal likelihood = -477.2 compared to -470.7 for BM). Therefore, the variable rates model may be over-parameterised and should not be interpreted closely, overturning the




Fig. 2 Phylogenetic patterns of rate variation inferred from original and log-transformed data showing highly different results. Rate variation mapped to phylogeny using colours based on **a** the original (untransformed) version of the PC1 size index, compared to those from **b** the log-transformed version of the PC1 size index. Yellow circles at the tips of the tree are scaled according to species body size.

conclusions of Stockdale & Benton⁴, including their time series of rate variation, which is based on this model. We also find strong support for a constrained, Ornstein-Uhlenbeck (OU), model compared to BM (marginal likelihood_{OU} = -463.5), consistent with previous studies^{5,6}, but differing from Stockdale & Benton⁴. The constraint parameter of the OU model (α) is estimated as 0.016. This corresponds to a phylogenetic half life¹² of 43 million years, which is short compared to the study duration of ~250 million years. Pseudosuchian body size evolution is therefore highly distinguishable from Brownian (diffusive) evolution, consistent with the importance of functional and ecological limits to body size evolution at large phylogenetic scales^{5,6}. A speciation (kappa) model is best supported (marginal likelihood_{kappa} = -454.0), but we disagree with the interpretation⁴ that this provides evidence of punctuational evolution, given that only a small proportion of species that ever lived are actually sampled in the fossil record.

Our analyses of log-transformed data reject key aspects of the conclusions of Stockdale & Benton⁴. More broadly, they demonstrate that the decision not to log-transform measurements introduces substantial errors to inferences of variation in the rate of evolution, and this should be accounted for in future studies.

Methods

We used the measurements provided by Stockdale & Benton⁴ to reproduce their published analyses of rate variation, also analysing a version in which the input data were log-transformed prior to analysis. Principal component analysis (PCA) with iterative missing data imputation was carried out in PAST version 3.17. We used the scores of the first principal component (PC1) as a body size index. Rate variation was evaluated on the time-scaled phylogeny of Stockdale & Benton⁴, using the 'varRates' model of BayesTraits version 3.0.2¹¹ and running our mcmc analyses for 2 million generations, with 50% of these discarded as burn-in (compared to 2 million generations with 10,000 discarded as burn-in by Stockdale & Benton⁴). Results were summarised using tools available at www.evolution.reading.ac.uk/VarRatesWebPP. We used mean scalar values as an

estimate of evolutionary rates following Stockdale & Benton⁴, but our scripts allow the use of alternative rate summary metrics (Supplementary data).

Phylogenetic rate variation was visualised by painting rate colours to phylogenetic branches using functions from the R package ape version 5.0¹³, in R version 4.0.3¹⁴. We also used functions from ape to extract estimated body sizes at internal nodes of the phylogeny and to graphically compare body size to evolutionary rates (Fig. 2). We statistically tested the correlation of evolutionary rates to body size using ordinary least squares regression of node- and tip-rates to body size, and also using phylogenetic generalised least squares regression (PGLS) to compare summed root-to-tip rates to the PC1 body size index, following e.g. ref. 15. This was implemented using custom code (Supplementary data) and the pglsl function of the R package caper version 1.0.1¹⁶.

Finally, to evaluate support for a variable rates model compared to alternatives, we conducted a model comparison analysis in BayesTraits¹¹. This was based on stepping-stone sampling, running 10 stones each for 100,000 generations following 1,000,000 generations of burn-in, comparing varRates to speciation (kappa), Ornstein-Uhlenbeck and Brownian motion models.

Data availability

The datasets generated and analysed during the current study are available in the Figshare repository <https://doi.org/10.6084/m9.figshare.15147306>. This includes measurements from the supplementary data of Stockdale & Benton⁴, as well as other data required to replicate our analyses.

Code availability

The analytical scripts used in the current study are available in the Figshare repository¹⁷.

Received: 12 March 2021; Accepted: 25 January 2022;

Published online: 25 February 2022

References

1. Bronzati, M., Montefeltro, F. C. & Langer, M. C. Diversification events and the effects of mass extinctions on Crocodyliformes evolutionary history. *R. Soc. Open Sci.* **2**, 140385 (2015).
2. Mannion, P. D. et al. Climate constrains the evolutionary history and biodiversity of crocodylians. *Nat. Commun.* **6**, 8438 (2015).

3. Stubbs, T. L., Pierce, S. E., Rayfield, E. J. & Anderson, P. S. Morphological and biomechanical disparity of crocodile-line archosaurs following the end-Triassic extinction. *Proc. R. Soc. Lond. Ser. B* **280**, 20131940 (2013).
4. Stockdale, M. T. & Benton, M. J. Environmental drivers of body size evolution in crocodile-line archosaurs. *Commun. Biol.* **4**, 38 (2021).
5. Godoy, P. L., Benson, R. B., Bronzati, M. & Butler, R. J. The multi-peak adaptive landscape of crocodylomorph body size evolution. *BMC Evol. Biol.* **19**, 167 (2019).
6. Gearty, W. & Payne, J. L. Physiological constraints on body size distributions in Crocodyliformes. *Evolution* **74**, 245–255 (2020).
7. Hammer, Ø., Harper, D. A. & Ryan, P. D. PAST: Paleontological statistics software package for education and data analysis. *Palaeontol. Electron.* **4**, 4–9 (2001).
8. Gingerich, P. D. Arithmetic or geometric normality of biological variation: an Empirical Test of Theory. *J. Theor. Biol.* **204**, 201–221 (2000).
9. Hunt, G. & Carrano, M. T. Models and methods for analysing phenotypic evolution in lineages and clades. *Paleontol. Soc. Pap.* **16**, 245–269 (2010).
10. Haldane, B. S. Suggestions as to quantitative measurement of rates of evolution. *Evolution* **3**, 51–56 (1949).
11. Meade, A. & Pagel, M. *BayesTraits V3. 02: a computer package for analyses of trait evolution* <http://www.evolution.rdg.ac.uk/BayesTraits.html> (2019).
12. Hansen, T. F. Stabilizing selection and the comparative analysis of adaptation. *Evolution* **51**, 1341–1351 (1997).
13. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).
14. R Core Team. *R: A language and environment for statistical computing*. (R Foundation for Statistical Computing, Vienna, Austria, 2020). <https://www.R-project.org/>.
15. Avaria-Llautureo, J., Hernández, C. E., Rodríguez-Serrano, E. & Venditti, C. The decoupled nature of basal metabolic rate and body temperature in endotherm evolution. *Nature* **572**, 651–654 (2019).
16. Orme, D. et al. *caper: Comparative Analyses of Phylogenetics and Evolution in R. R package version 1.0.1*. <https://CRAN.R-project.org/package=caper> (2018).
17. Benson, R. B. J. et al. *Reconstructed evolutionary patterns for crocodile-line archosaurs demonstrate impact of failure to log-transform body size*. <https://doi.org/10.6084/m9.figshare.15147306> (2021).

Acknowledgements

We thank Max Stockdale and Mike Benton for sharing data required to replicate their analyses, and for discussion. We thank Chris Venditti for writing with statistical advice on our preprint. We thank three referees for reviewing our manuscript.

Author contributions

R.B.J.B. designed the analyses, performed analysis and drafted the manuscript. P.G. performed analyses. R.B.J.B., P.G., M.B., R.J.B. and W.G. developed the concepts and contributed to the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Roger B. J. Benson.

Peer review information *Communications Biology* thanks the anonymous reviewers for their contribution to the peer review of this work. Primary Handling Editor: Caitlin Karniski.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com