

# Direct and indirect effects of altered temperature regimes and phenological mismatches on insect populations

Mariana Abarca<sup>1</sup> and Ryan Spahn<sup>2</sup>



Climate change is transforming ecosystems by altering species ranges, the composition of communities, and trophic interactions. Here, we synthesize recent reviews and subsequent developments to provide an overview of insect ecological and evolutionary responses to altered temperature regimes. We discuss both direct responses to thermal stress and indirect responses arising from phenological mismatches, altered host quality, and changes in natural enemy activity. Altered temperature regimes can increase exposure to both cold and heat stress and result in phenological and morphological mismatches with adjacent trophic levels. Host plant quality varies in a heterogeneous way in response to altered temperatures with both increases and decreases observed. Density-dependent effects, spatial heterogeneity, and rapid evolutionary change provide some resilience to these threats.

## Addresses

<sup>1</sup> Department of Biological Sciences, Smith College, Northampton, MA, United States

<sup>2</sup> Department of Biological Sciences, George Washington University, DC, 20052, United States

Corresponding author: Abarca, Mariana ([mabarca@smith.edu](mailto:mabarca@smith.edu))

**Current Opinion in Insect Science** 2021, **47**:67–74

This review comes from a themed issue on **Global change biology**

Edited by **Shannon M Murphy** and **Gina M Wimp**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 11th May 2021

<https://doi.org/10.1016/j.cois.2021.04.008>

2214-5745/© 2021 Elsevier Inc. All rights reserved.

## Introduction

The profound environmental transformation resulting from global climate change is a major threat to biodiversity. Altered weather and precipitation patterns as well as increased frequencies of extreme events are important drivers of insect population declines [1,2,3,4,5]. Altered temperature regimes can affect insects both directly (e.g. by exceeding their physiological thermal limits) and indirectly, through phenological alterations [6,7,8] and via adjacent trophic level effects [9,10]. These effects are not always negative, however. Warmer temperatures can

benefit some insect populations by allowing them to grow faster and complete more generations in a growing season, depending on local initial conditions and population-specific thermal sensitivities [11]. Insects possess characteristics that can allow them to survive and adapt to rapid environmental change, including high mobility, remarkable phenotypic plasticity, and short generation times [12,13]. However, extreme temperatures can push insects beyond their adaptive limits, disrupting interactions and food webs, and ultimately leading to outbreak-breakdown cycles that affect ecosystem functioning and resilience [4,14].

Over the last several decades, empirical studies documenting the physiological, ecological, and evolutionary effects of altered temperature regimes on insects have proliferated, but major gaps remain. Assessing the magnitude and pervasiveness of the threat that climate change poses for insects is complicated by a paucity of long-term monitoring data and by the cyclic dynamics of insect populations that often include recurrent outbreaks and declines caused by a variety of well-known intrinsic and extrinsic factors [15]. Additionally, attributing population responses to causative pathways of direct and indirect effects is challenging to do in practice. Nonetheless, we attempt to separate them conceptually in this review to highlight emerging patterns and to encourage future studies that explicitly address these important questions. Here, we synthesize 29 recent reviews and subsequent developments to provide an overview of insect ecological and evolutionary responses to direct and indirect effects of altered temperature regimes. More specifically, we discuss 1) direct responses to thermal stress, and 2) indirect responses arising from phenological mismatches, decreased resource quality, and changes in arthropod natural enemy activity.

## Direct effects of thermal stress

Warmer temperatures can result in increased exposure of insects to both heat and cold stress (Table 1). Heat stress impairs insect neural performance, muscular control, immune function, and can result in coma and eventually death [16,17]. In addition, exposure to heat stress can result in morphological changes that may compromise dispersal capabilities [18]. As a result, weather events that expose insects to higher temperatures and lower humidity (e.g. summer droughts) often result in increased mortality [19] and may trigger population declines [20,21]. Multiple

**Table 1**

**Recent studies documenting direct effects of altered temperature regimes on insects, bold numbers correspond to review articles and italics to studies in which the focal species is not a model organism or agricultural pest.**

Direct effects	
Exposure to:	Alterations in:
Cold stress [ <b>4*4</b> ,30,31]	Development time [19,78]
Heat stress [ <b>4*4</b> , <i>17*17</i> ••]	Growth [78,79]
Depletion of energy reserves [25]	Mortality [ <b>4*4</b> , <i>17*17</i> ••,26,27]
Population declines [14,20]	Diapause timing [ <b>19</b> 19,2525,55*55•]
Population outbreaks [20]	Voltinism [13*]
Loss of endosymbiont protection against parasitoids [19]	Caterpillar capacity to induce plant defenses [78]
	Immune [16,1919]
	Phenology
	[ <b>3*3</b> •,18,45,5050,55*55•,62]
	Wing length [18,79]
	Body size [18,80]
	Metabolic rate [66]
	Expression of polyphenisms [81]

factors affect heat tolerance, including sex, life stage, color, body size, presence of other environmental stressors, resource deprivation, parent's thermal environment [17\*\*], caste [22\*\*], and the presence of endosymbionts. Bacteria such as *Serratia symbiotica* and *Rickettsia* can protect their hosts against heat stress [19]. In addition, insects use a diverse array of behavioral and physiological mechanisms to mitigate the harmful effects of heat stress. Many of these mechanisms are energetically costly and these costs can restrict the evolution of thermal tolerance [17\*\*]. Consistently, upper thermal limits are more conserved among species and life stages than lower thermal limits [17\*\*,23,24\*], suggesting greater negative impacts on populations when upper limits are exceeded.

Warmer temperatures during autumn and winter can have stressful effects despite falling within insects' physiological thermal limits. Warmer winters can result in depletion of energy reserves of overwintering insects [25], as well as increased mortality of both overwintering larvae [26] and diapausing pupae [27]. Counterintuitively, milder winters can result in increased exposure to cold stress. Snow often provides insulation from extreme cold, so milder winters in which this protective layer melts can increase exposure to inclement temperatures [25]. Additionally, warmer temperatures during winter and spring can trigger premature diapause termination and advance phenology [25,28]. Phenological advances in spring can expose vulnerable life stages to frosts [20].

Cold stress affects multiple metabolic pathways, and can cause mechanical damage and cellular dehydration, loss of neuromuscular function, and imbalance in ion and

water homeostasis [29,30]. These metabolic changes can result in increased incidence of deformities, premature pupation, decreased mobility, longevity, and fecundity, and male biased sex ratios of parasitoids [9\*]. Thus, events such as late spring frosts can trigger dramatic population declines for a multitude of reasons [20].

Insects possess a diversity of adaptations to cope with cold stress [12,25,29], and traits associated with cold hardiness exhibit variable genetic architecture, heritability, and a strong genetic basis for plasticity [25]. In addition, cold hardiness [25] and lower thermal limits vary within and among species and life stages [31,32]. Notably, there is some evidence suggesting that cold hardiness can evolve independently among life stages in *Drosophila melanogaster* [33] and the butterfly *Bicyclus aninana* [34]. Biotic interactions can also affect cold hardiness, as when hymenopteran parasitoids physiologically manipulate their hosts to avoid freezing by lowering their supercooling points [9\*]. Taken together, these variable patterns indicate high potential for cold hardiness to evolve [25]. Despite this potential, studies documenting cold hardiness evolution in response to recent climate change are lacking [25]. By contrast, there are multiple studies showing recent local adaptation in diapause induction and termination [25].

Terrestrial insects can thermoregulate behaviorally and implement microenvironmental modifications to buffer stressful temperatures. Some of the most striking examples come from social insects, as they can build complex insulating structures (e.g. termite nests) and engage in collaborative thermoregulation. Bees can regurgitate gut fluids that cool the nest as they evaporate, and they also use the bodies of workers as insulation to protect the young from heat stress [22\*\*]. While sociality allows for sophisticated thermoregulation strategies, it also entails trade-offs, for example, cuticular hydrocarbons play a role in thermoregulation, but selection to enhance this role can be restricted due to their key function in colony communication [22\*\*]. Other thermoregulation strategies simply consist in moving across microhabitats. Vulnerability to climate change can be buffered or magnified by the small microhabitats terrestrial insects can exploit, as surface temperatures can differ greatly (>30°C) from air temperatures, and among features of the same tree trunk, rock or set of leaves (~10°C) [35]. By contrast, microhabitat variation is more restricted in aquatic environments, increasing insect vulnerability to climate change [36]. While still water bodies exhibit thermal stratification, water's high-specific heat capacity buffers temperature changes, particularly in rapidly moving streams, resulting in a more homogeneous thermal environment [36]. In high-elevation streams, the effect of warmer temperatures on insects is further complicated by changes in oxygen concentration, water flow, ultraviolet radiation and salinity [37]. Warming of streams stimulates

metabolic demand for oxygen [37] and can lead to declines in insect abundance and altered community composition [38].

At the macro scale, vulnerability to warming can be estimated by calculating thermal safety margins, which are the difference between mean habitat temperature and the insect's maximum thermal limits [39]. Tropical species were believed to be particularly vulnerable to warming due to their smaller safety margins, as they historically experienced ambient temperatures close to their physiological thermal limits [39]. However, a new analysis suggests that temperate species might face levels of threat from climate change similar to those of their tropical counterparts [40]. This re-analysis accounted for latitudinal differences in activity periods by calculating thermal limits using mean temperature during the months insects are active, instead of annual mean temperature [40]. There are, however, geographic areas of particular interest because of the accelerated rates of change they are experiencing. High elevation streams, where aquatic insects are the dominant fauna [37], and the arctic, where snow cover is the main regulator of phenology [41] are relatively understudied areas that are experiencing rapid changes. In the Arctic, warming has resulted in an increase of populations of herbivores and a decrease of detritivores [41].

In addition to a geographic bias, taxonomic and methodological biases also affect our understanding of insect responses to thermal stress. These responses have been studied mostly in laboratory settings [24<sup>\*</sup>] and with a strong focus on agricultural pests [11,42,43], model organisms [17<sup>\*\*</sup>,29], and, in the case of social insects, bees and ants [22<sup>\*\*</sup>]. Most of the work on genetic variation in heat resistance, selection experiments, and evolutionary change have been performed in *Drosophila* [17<sup>\*\*</sup>,24<sup>\*</sup>]. This bias is problematic because insects are a heterogeneous group and vary in key traits that mediate their ability to detect and respond to warming. Recent studies focusing on species outside these popular study groups are highlighted in [Tables 1 and 2](#).

### Indirect effects — phenological mismatches

Temperature, rainfall, and photoperiod are the main drivers of phenology [12,44,45]. Thus, shifts in temperature regimes that alter the historical correlation of thermal and photoperiodic conditions can disrupt insect seasonal activities (e.g. breeding) [46]. Warmer temperatures during the growing season typically accelerate ectotherm development; however, not all species respond at the same rate, which can result in consumers missing resource availability [47]. In temperate areas, warming during winter can result in delayed phenology, because some species require exposure to a certain period of low temperatures before they can resume development in the spring [12,48,49<sup>\*\*</sup>]. More commonly, however, warmer

springs result in phenological advances and accelerated development [25]. Differences in sensitivity and responsiveness to these environmental changes can result in phenological mismatches and disrupt historical patterns of trophic interactions [47,50].

Phenological mismatches arise when consumers and resources occur asynchronously. Assessing synchrony at the community level involves evaluating the overlap of the occurrence distribution of the relevant phenophase of consumers and producers [7<sup>\*</sup>]. For herbivores, synchrony involves fitting their larval stage within the period of foliage availability [51]; for pollinators, synchronizing foraging with flower occurrence; for parasitoids, co-existence between ovipositing adults and susceptible life stages of the hosts. The shapes of these occurrence distributions vary according to both heat accumulation and life history traits. The degree of overlap in the distributions of interactants, as well as the abundance of resources can define the severity and outcomes of mismatches [7<sup>\*</sup>]. For example, when resources are abundant, mild mismatches, in which occurrence curves partially overlap, are not likely to have negative effects on consumers [7<sup>\*</sup>]. Evaluating the role of recent climate change in triggering or exacerbating phenological mismatches requires establishing a pre-climate change baseline, which is the degree of synchrony a given interaction had before recent climate change (~1980) [52]. This baseline is important because cases of adaptive asynchrony are commonly due to factors other than climate change, including trade-offs such as fertility vs offspring survival, or bet-hedging strategies [53,54]. For example, larval starvation due to asynchrony with host plants is common in the bay Checkerspot (*Euphydryas editha*), because of a trade-off between fertility and offspring survival. Females that feed for longer periods are more fecund, but their offspring are less likely to occur within the period of host plant availability [53].

The likelihood of a phenological mismatch, as well as its outcome for producers and consumers, varies according to interaction type, consumer diet breadth, and the environmental cues triggering phenology of each trophic level. Mutualisms, where both interactants experience selection for synchrony, are expected to be more robust to mismatches than antagonisms, where interactants experience opposite pressures, as synchrony is favorable for consumers and unfavorable for producers. Parasitoids constitute a special case of antagonism because they often follow phenological cues similar to those of their hosts and even directly from their host, by engaging in endogenous physiological synchronization [6,55<sup>\*</sup>]. Despite this intimacy, a long-term field study showed that different weather components explain variation in the phenology of aphids and their parasitoids [49<sup>\*\*</sup>].

Table 2

Recent studies documenting indirect effects of altered temperature regimes on insects, bold numbers correspond to review articles and italics to studies in which the focal species is not a model organism or agricultural pest

## Indirect effects

## Phenological mismatches

[66,77,88,99,4545,51,5252,5353,5657,5959,6060,6161,8282,8383]

*Altered performance of parasitoids:*

Survival [44,99,19,68]

Development [44,99,68]

Flight [44,99]

Host-finding efficiency [44,99]

Oviposition behavior [44]

Parasitism rate [1919,8282]

Host preference [99,1919]

Sex allocation [99,1919]

Mating [99]

Fecundity [99]

*Altered performance of predators:*

Consumption rate [44]

Growth [44]

Behavior [44,7272,2323]

*Altered resource quality:*

Foliage quality [1010,1313,66]

Concentration of plant defenses [44,78]

Emission of volatile compounds [44,99,7171]

Plant/floral morphology [44,88]

Plant physiology [44]

Plant biomass [78]

Trichome density [78]

Host quality for parasitoids [1919]

*Altered relationships with endosymbionts*

[1919,7575,76,77]

Studies documenting the consequences of phenological mismatches between plants and pollinators have focused primarily on the effects on plant fitness. Thus, there are multiple field empirical studies documenting detrimental effects of phenological mismatches on plant reproduction [28,56], but a dearth of studies documenting the effects on pollinator populations or individual fitness [8]. A simulation using empirical interaction network data found that pollinators are expected to experience severe population declines due to mismatches [57]. Consistently, an analysis of the same historical data set identified phenological mismatches between pollinators and flower resources as a factor contributing to the extirpation of pollinators and degradation of plant-pollinator interaction networks [58], with higher vulnerability of specialists than generalists [8,58]. An analysis of over 2000 species of European pollinators documented not only phenological advances, but also flight period reductions in the last 60 years; as a result, flight period overlap of pollinators decreased, limiting redundancy and complementarity of pollinator assemblages, which could result in decreases in pollinator function [59]. An experimental study on two species of solitary bees (*Osmia*) showed that males advanced their spring phenology in response to warmer temperatures, but females did not, and that male bees emerged on or before the onset of host flowering and females after flowering [60]. A recent study reported a pollinator behavior that, if widespread, could serve to mitigate asynchrony with floral resources. Pashalidou *et al.* [61] found that when facing pollen scarcity, workers of the bumblebee *Bombus terrestris* damaged plants, and this damage accelerated flower production. To our knowledge this is the only study documenting such a response, so it remains unclear how ubiquitous this behavior is for pollinators.

There are more studies directly documenting adverse effects of phenological mismatches on consumers in plant-herbivore than in plant-pollinator interactions. Mismatches among herbivores and their host plants can have adverse effects on the consumers and trigger evolutionary responses. The best documented example of recent evolutionary change in an insect due to the pressures of phenological mismatches is the case of the winter moth (*Operophtera brumata*), whose larvae hatch in spring and feed on expanding oak foliage. Warmer temperatures have led to earlier hatching and have precipitated caterpillar population declines; after a decade of this selective pressure, the *O. brumata* population responded by delaying its hatching time, decreasing asynchrony with foliage availability [7,62]. However, not all phenological mismatches have negative consequences on insect populations, as mechanisms such as density-dependent compensation can offset negative effects on population size [49]. In this context, density-dependent compensation occurs when asynchrony-triggered mortality is compensated by a subsequent increase in fecundity of the survivors due to reduced competition, resulting in a stable population size.

Environmental heterogeneity is another factor that may buffer consumers from phenological mismatches [28]. Caterpillar peak abundance has been found to vary across habitat types within a region [63] and across altitudinal gradients [64]. In addition, intraspecific and interspecific sensitivity to phenological cues can vary across both space and time [45]. A study of butterfly communities in a Mediterranean ecosystem where seasonality is driven by temperature and rainfall, showed that early spring species were more sensitive to temperature than late species, and that butterfly phenology responded more

strongly to variation in temperature than to rainfall [45]. In tropical ecosystems, rainfall mediates plant-herbivore relationships [65], so altered precipitation patterns could trigger asynchrony between consumers and producers; however, this possibility remains understudied as research on phenological mismatches has been strongly biased towards temperate systems [48].

### Indirect effects – host quality and natural enemies

Altered temperature regimes can also affect insects indirectly by altering the quality of the resources they utilize and/or the performance of their natural enemies (Table 2). These alterations are important because resource quality influences numerous life history traits including lifespan, fecundity, immune response, metabolic and growth rates, and tolerance to thermal stress [10,26,43,66]. Warming can affect plant traits relevant for pollinators such as plant height, floral diameter, volatile emissions, and the nutritional quality of pollen and nectar [8]. In addition, changes in floral morphology, in conjunction with decreasing body size of insects (a common response to temperature increases), can create morphological mismatches between plants and pollinators, particularly for specialist pollinators [8]. Miller-Struttman *et al.* [67] found that two alpine bumblebee species have evolved shorter tongue lengths over the past 40 years, but the flowers they pollinate have not become shallower, suggesting that generalist pollinators will be more successful as this environment continues to change. Similarly, higher temperatures can affect foliage quality for herbivores, as temperature affects plant growth, primary production, and inducibility of plant defenses (Table 2); however, the effects of warming on foliage quality are not uniform, as both increases and decreases in nutrient concentration have been reported [4<sup>•</sup>,10,13<sup>•</sup>]. Moreover, many aspects of plant quality are more heavily influenced by atmospheric CO<sub>2</sub> concentration as well as drought, which are simultaneously increasing due to climate change [10,13<sup>•</sup>,21]. Altered foliage quality in combination with the direct effects of thermal stress on herbivores often results in increased feeding rates, particularly in agricultural systems [13<sup>•</sup>].

Agricultural systems face the additional challenge that parasitoids, important biocontrol agents, are particularly susceptible to altered temperature regimes. Parasitoids and other natural enemies frequently exhibit lower thermal tolerances than their hosts [9<sup>•</sup>,68] and have to cope with the cumulative effects of warming on lower trophic levels (Table 2). For example, water-stressed plants hosted fewer and smaller aphids, which in turn exhibited reduced parasitism rates [69,70]. Many parasitoids rely on chemical cues produced by plants to locate hosts, but the integrity of these signals can be compromised by altered temperature regimes, resulting in failed parasitoid recruitment and decreased rates of parasitism [9<sup>•</sup>].

Similarly, temperature can influence volatile emissions of predators, which in turn can affect predator-prey interactions. Ladybeetle larvae emit chemicals which inform oviposition decisions of conspecific females, and Sentis *et al.* [71] found that increased temperature altered ladybeetle infochemical emission and perception. The alterations in infochemicals resulted in decreased oviposition at higher temperatures, which should reduce local predation pressure on aphids.

Altered temperature regimes can also influence the behavior of natural enemies. Heat waves and cold snaps can have negative consequences for parasitoid flying, feeding, and host finding, resulting in decreased performance [9<sup>•</sup>]. Likewise, exposure to thermal stress can alter predator growth and consumption rates [4<sup>•</sup>]. Some predators can mitigate the effects of suboptimal temperatures by changing their habitat domain or foraging behavior. Sit-and-wait ambush spiders have been shown to alter their position in the grass canopy and adjust the time of day during which they forage in response to increased temperature, subsequently influencing predation on grasshopper prey [72,73]. High temperatures can also decrease foraging success of insectivorous birds by drying out soil, which makes it more difficult for birds to probe for invertebrate prey [74].

Finally, endosymbionts are well-known for their ability to mediate interactions between hosts and parasitoids by conferring resistance to hosts through multiple pathways, including enhanced encapsulation response and improved thermal tolerance [19]. However, endosymbionts themselves are sensitive to changes in temperature so the benefits they provide to hosts are also subject to change [19]. For example, endosymbionts that typically protect their aphid hosts against parasitoids (e.g. *Hamiltonella defensa*) fail to do so under heat stress [75]. Moreover, high temperatures can result in loss of host endosymbionts altogether, as evidenced by reduced vertical transmission of *Wolbachia* [76] and *Spiroplasma* [77]. Additional field studies, particularly in natural systems, are necessary to further our understanding of temperature effects on tri-trophic dynamics.

### Conclusions

Altered temperature regimes affect insects through multiple mechanisms, both directly impacting their metabolism and phenology, and indirectly through their effects on adjacent trophic levels. Direct and indirect pressures exerted by thermal stress occur simultaneously and their distinction in practice is complicated; however, a conceptual separation of these effects can aid in study design and interpretation. Understanding the responses of insect populations to altered temperature regimes requires the integration of results from laboratory and field experiments, as well as long-term monitoring programs. Thanks to a proliferation of studies on thermal tolerances and insect responses at a variety of organizational levels, this

integration is increasingly possible. However, some challenges remain given the rich diversity of insect taxa and variation in their traits. More field studies, particularly in natural systems, are necessary to further our understanding of temperature effects on insects that manifest through indirect effects on adjacent trophic levels. Similarly, there is a remarkable imbalance in the predicted effects of phenological mismatches on pollinators and the dearth of field studies documenting their individual or population-level responses. The assessment of these mismatches has been biased, focusing mainly on the effects on plants, probably due to the challenges involved in measuring pollinator mortality due to starvation. Phenological mismatches may very well be an underappreciated mechanism of insect decline, overlooked due to the logistical constraints associated with their assessment.

Altered temperature regimes pose an important challenge for ectotherms and it is particularly worrisome that insects appear to exhibit stronger potential for evolutionary change in response to cold stress than to heat stress. However, spatial heterogeneity, density-dependent effects, and short-term evolutionary change in seasonal traits can provide some resilience to the negative effects of climate change.

### Funding sources

This work was funded by Georgetown University and George Washington University.

### Conflict of interest statement

Nothing declared.

### Acknowledgements

We thank the Lill (George Washington University) and Weiss (Georgetown University) labs as well as two anonymous reviewers for helpful comments on an early version of this manuscript.

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Forister ML, Pelton EM, Black SH: **Declines in insect abundance and diversity: we know enough to act now.** *Conserv Sci Pract* 2019, **1**:1-8.
  2. Sánchez-Bayo F, Wyckhuys KAG: **Worldwide decline of the entomofauna: a review of its drivers.** *Biol Conserv* 2019, **232**:8-27.
  3. Eggleton P: **The state of the world's insects.** *Annu Rev Environ Resour* 2020, **45**:61-82.  
Comprehensive review of insects including their taxonomy, ecological and economical importance, evidence for recent declines and the main causes of these declines.
  4. Harvey JA, Heinen R, Gols R, Thakur MP: **Climate change-mediated temperature extremes and insects: from outbreaks to breakdowns.** *Glob Change Biol* 2020, **26**:6685-6701.  
Comprehensive review integrating effects of thermal extremes on individuals (insect physiology, development, reproduction and behavior), trophic interactions (plant-insect interactions, third and fourth trophic levels), and community level processes.
  5. Wagner DL: **Insect declines in the Anthropocene.** *Annu Rev Entomol* 2020, **65**:457-480.  
Comprehensive review of the different threats insects face and how they have resulted in population declines. The consequences of population declines of abundant species for vertebrate consumers and ecosystem services are also discussed.
  6. Damien M, Tougeron K: **Prey-predator phenological mismatch under climate change.** *Curr Opin Insect Sci* 2019, **35**:60-68.
  7. Visser ME, Gienapp P: **Evolutionary and demographic consequences of phenological mismatches.** *Nat Ecol Evol* 2019, **3**:879-885.  
Clear and thorough explanation of a theoretical framework to study the demographic and evolutionary effects of phenological mismatches. This review includes examples, definitions, and future directions.
  8. Gérard M, Vanderplanck M, Wood T, Michez D: **Global warming and plant-pollinator mismatches.** *Emerg Top life Sci* 2020, **4**:77-86.
  9. Chidawanyika F, Mudavanhu P, Nyamukondiwa C: **Global climate change as a driver of bottom-up and top-down factors in agricultural landscapes and the fate of host-parasitoid interactions.** *Front Ecol Evol* 2019, **7**:1-13.  
A review of how tri-trophic interactions in agricultural systems may be altered by climate change, including the bottom-up and top-down mechanisms driving these changes.
  10. Jactel H, Koricheva J, Castagneyrol B: **Responses of forest insect pests to climate change: not so simple.** *Curr Opin Insect Sci* 2019, **35**:103-108.
  11. Lehmann P, Ammunét T, Barton M, Battisti A, Eigenbrode SD, Jepsen JU, Kalinkat G, Neuvonen S, Niemelä P, Terblanche JS *et al.*: **Complex responses of global insect pests to climate warming.** *Front Ecol Environ* 2020, **18**:141-150.
  12. Tauber MJ, Tauber CA, Masaki S: *Seasonal Adaptations of Insects.* Oxford University Press; 1986.
  13. Hamann E, Blevins C, Franks SJ, Jameel MI, Anderson JT: **Climate change alters plant-herbivore interactions.** *New Phytol* 2021, **229**:1894-1918 <http://dx.doi.org/10.1111/nph.17036>.  
Review and phylogenetically corrected meta-analysis of herbivore responses at the individual and population levels to climate change manipulations in the field and under controlled conditions.
  14. Patterson TA, Grundel R, Dzurisin JDK, Knutson RL, Hellmann JJ: **Evidence of an extreme weather-induced phenological mismatch and a local extirpation of the endangered Karner blue butterfly.** *Conserv Sci Pract* 2020, **2**:1-13.
  15. Barbosa P, Letourneau DK, Agrawal AA (Eds): *Insect Outbreaks Revisited.* J. Wiley-Blackwell; 2012.
  16. Shamakhi L, Zibae A, Karimi-Malati A, Hoda H: **Effect of thermal stress on the immune responses of *Chilo suppressalis* walker (Lepidoptera: Crambidae) to *Beauveria bassiana*.** *J Therm Biol* 2019, **84**:136-145.
  17. González-Tokman D, Córdoba-Aguilar A, Dáttilo W, Lira-Noriega A, Sánchez-Guillén RA, Villalobos F: **Insect responses to heat: physiological mechanisms, evolution and ecological implications in a warming world.** *Biol Rev* 2020, **3**:802-821.  
Comprehensive review of the physiological mechanisms that allow insects to cope with heat stress: from the neuronal pathways that mediate the detection and response to heat; metabolic and stress responses to heat, thermoregulation; and how hormones that regulate development and behaviour respond to heat. Include a list of heat shock proteins and cognates, highlight gaps and outline future research.
  18. Jourdan J, Baranov V, Wagner R, Plath M, Haase P: **Elevated temperatures translate into reduced dispersal abilities in a natural population of an aquatic insect.** *J Anim Ecol* 2019, **88**:1498-1509.
  19. Thierry M, Hrček J, Lewis OT: **Mechanisms structuring host-parasitoid networks in a global warming context: a review.** *Ecol Entomol* 2019, **44**:581-592.
  20. Marquis RJ, Lill JT, Forkner RE, Le Corff J, Landosky JM, Whitfield JB: **Declines and resilience of communities of leaf chewing insects on Missouri oaks following spring frost and summer drought.** *Front Ecol Evol* 2019, **7**:1-11.

21. Gely C, Laurance SGW, Stork NE: **How do herbivorous insects respond to drought stress in trees?** *Biol Rev* 2020, **95**:434-448.
22. Perez R, Aron S: **Adaptations to thermal stress in social insects: recent advances and future directions.** *Biol Rev* 2020, **95**:1535-1553.
- Thermal tolerance and adaptation: behavioural (passive and active nest thermoregulation), morphological (body size, hair and leg length) physiological and molecular (Cuticular hydrocarbons, cryoprotectants, heat shock proteins and antioxidants) in social insects.
23. Addo-bediako A, Chown SL, Gaston KJ: **Thermal tolerance, climatic variability and latitude.** *Proc R Soc B Biol Sci* 2000, **267**:739-745.
24. Kellermann V, van Heerwaarden B: **Terrestrial insects and climate change: adaptive responses in key traits.** *Physiol Entomol* 2019, **44**:99-115.
- Review of traits that mediate responses to climate change: upper thermal limits, desiccation resistance, phenology, plasticity. Includes a discussion of how these traits mediate responses to climate change and review studies documenting their evolutionary responses.
25. Marshall KE, Gotthard K, Williams CM: **Evolutionary impacts of winter climate change on insects.** *Curr Opin Insect Sci* 2020, **41**:54-62 <http://dx.doi.org/10.1016/j.cois.2020.06.003>.
26. Abarca M, Larsen EA, Ries L: **Heatwaves and novel host consumption increase overwinter mortality of an imperiled wetland butterfly.** *Front Ecol Evol* 2019, **7**:1-13.
27. Klockmann M, Fischer K: **Strong reduction in diapause survival under warm and humid overwintering conditions in a temperate-zone butterfly.** *Popul Ecol* 2019, **61**:150-159.
28. Inouye DW: **Effects of climate change on alpine plants and their pollinators.** *Ann N Y Acad Sci* 2020, **1469**:26-37.
29. Lubawy J, Urbański A, Colinet H, Pflüger HJ, Marciniak P: **Role of the insect neuroendocrine system in the response to cold stress.** *Front Physiol* 2020, **11**:1-11.
30. Lubawy J, Daburon V, Chowanski S, Stocinska M, Colinet H: **Thermal stress causes DNA damage and mortality in a tropical insect.** *J Exp Biol* 2019, **22**:1-9.
31. Amiresmaeili N, Romeis J, Collatz J: **Cold tolerance of the drosophila pupal parasitoid *Trichopria drosophilae*.** *J Insect Physiol* 2020, **125**:104087.
32. Kingsolver JG, Buckley LB: **Ontogenetic variation in thermal sensitivity shapes insect ecological responses to climate change.** *Curr Opin Insect Sci* 2020, **41**:17-24.
33. Freda PJ, Ali ZM, Heter N, Ragland GJ, Morgan TJ: **Stage-specific genotype-by-environment interactions for cold and heat hardiness in *Drosophila melanogaster*.** *Heredity (Edinb)* 2019, **123**:479-491.
34. Dierks A, Kölzow N, Franke K, Fischer K: **Does selection on increased cold tolerance in the adult stage confer resistance throughout development?** *J Evol Biol* 2012, **25**:1650-1657.
35. Pincebourde S, Woods HA: **There is plenty of room at the bottom: microclimates drive insect vulnerability to climate change.** *Curr Opin Insect Sci* 2020, **41**:63-70.
36. Shah AA, Dillon ME, Hotaling S, Woods HA: **High elevation insect communities face shifting ecological and evolutionary landscapes.** *Curr Opin Insect Sci* 2020, **41**:1-6.
37. Birrell JH, Shah AA, Hotaling S, Giersch JJ, Williamson CE, Jacobsen D, Woods HA: **Insects in high-elevation streams: Life in extreme environments imperiled by climate change.** *Glob Change Biol* 2020, **26**:6667-6684.
38. Baranov V, Jourdan J, Pilotto F, Wagner R, Haase P: **Complex and nonlinear climate-driven changes in freshwater insect communities over 42 years.** *Conserv Biol* 2020, **34**:1241-1251.
39. Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC, Martin PR: **Impacts of climate warming on terrestrial ectotherms across latitude.** *Proc Natl Acad Sci U S A* 2008, **105**:6668-6672.
40. Johansson F, Orizaola G, Nilsson-örtman V: **Temperate insects with narrow seasonal activity periods can be as vulnerable to climate change as tropical insect species.** *Sci Rep* 2020, **10**:1-8.
41. Hoyer TT: **Arthropods and climate change — Arctic challenges and opportunities.** *Curr Opin Insect Sci* 2020, **41**:40-45 <http://dx.doi.org/10.1016/j.phrs.2020.104743>.
42. Gu LL, Li MZ, Wang GR, Liu XD: **Multigenerational heat acclimation increases thermal tolerance and expression levels of Hsp70 and Hsp90 in the rice leaf folder larvae.** *J Therm Biol* 2019, **81**:103-109.
43. Mutamiswa R, Machekano H, Nyamukondiwa C, Chidawanyika F: **Host plant-related responses on the thermal fitness of *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae).** *Arthropod Plant Interact* 2020, **14**:463-471.
44. Franklin KA: **Light and temperature signal crosstalk in plant development.** *Curr Opin Plant Biol* 2009, **12**:63-68.
45. Gutiérrez D, Wilson RJ: **Intra- and interspecific variation in the responses of insect phenology to climate.** *J Anim Ecol* 2021, **90**:248-259 <http://dx.doi.org/10.1111/1365-2656.13348>.
46. Tsai HY, Rubenstein DR, Fan YM, Yuan TN, Chen BF, Tang Y, Chen IC, Shen SF: **Locally-adapted reproductive photoperiodism determines population vulnerability to climate change in burying beetles.** *Nat Commun* 2020, **11**:1-12.
47. Visser ME, Both C: **Shifts in phenology due to global climate change: the need for a yardstick.** *Proc Biol Sci* 2005, **272**:2561-2569.
48. Renner SS, Zohner CM: **Climate change and phenological mismatch in trophic interactions among plants, insects, and vertebrates.** *Annu Rev Ecol Syst* 2018, **49** <http://dx.doi.org/10.1146/annurev-ecolsys-110617-062535>.
49. Senior VL, Evans LC, Leather SR, Oliver TH, Evans KL: **Phenological responses in a sycamore-aphid-parasitoid system and consequences for aphid population dynamics: a 20 year case study.** *Glob Change Biol* 2020, **26**:2814-2828.
- A 20-year study documenting phenology of a tri-trophic system including a plant, two aphid species (one specialist one generalist) and their parasitoids. Main findings include identifying the individual and combined roles of direct effects of weather, phenological mismatches and parasitoid attack on aphid population growth.
50. Chmura HE, Kharouba HM, Ashander J, Ehlman SM, Rivest EB, Yang LH: **The mechanisms of phenology: the patterns and processes of phenological shifts.** *Ecol Monogr* 2019, **89**.
51. Singer MC, Parmesan C: **Misunderstood mismatch.** *EcoEvoRxiv* 2020 <http://dx.doi.org/10.32942/osf.io/rxmct>. Preprint.
52. Kharouba HM, Wolkovich EM: **Disconnects between ecological theory and data in phenological mismatch research.** *Nat Clim Change* 2020, **10**:406-415.
53. Singer MC, Parmesan C: **Phenological asynchrony between herbivorous insects and their hosts: signal of climate change or pre-existing adaptive strategy?** *Philos Trans R Soc Lond B Biol Sci* 2010, **365**:3161-3176.
54. Johansson J, Jonzén N: **Game theory sheds new light on ecological responses to current climate change when phenology is historically mismatched.** *Ecol Lett* 2012, **15**:881-888.
55. Tougeron K, Brodeur J, Le Lann C, van Baaren J: **How climate change affects the seasonal ecology of insect parasitoids.** *Ecol Entomol* 2020, **45**:167-181.
- Review of parasitoid traits that face selective pressures from climate change and the expected outcomes of this selection. Discussion of the consequences of climate change for insect ecology and expected scenarios both supported by empirical evidence and plausible but not supported yet.
56. Kudo G, Cooper EJ: **When spring ephemerals fail to meet pollinators: mechanism of phenological mismatch and its impact on plant reproduction.** *Proc R Soc B Biol Sci* 2019, **286**.
57. Memmott J, Craze PG, Waser NM, Price MV: **Global warming and the disruption of plant-pollinator interactions.** *Ecol Lett* 2007, **10**:710-717.

58. Burkle LA, Marlin JC, Knight TM: **Plant-pollinator interactions over 120 years: loss of species, co-occurrence, and function.** *Science (80-)* 2013, **339**:1611-1616.
59. Duchenne F, Thébaud E, Michez D, Elias M, Drake M, Persson M, Piot JS, Pollet M, Vanormelingen P, Fontaine C: **Phenological shifts alter the seasonal structure of pollinator assemblages in Europe.** *Nat Ecol Evol* 2020, **4**:115-121.
60. Kehrberger S, Holzschuh A: **Warmer temperatures advance flowering in a spring plant more strongly than emergence of two solitary spring bee species.** *PLoS One* 2019, **14**:1-15.
61. Pashalidou FG, Lambert H, Peybernes T, Mescher MC, de Moraes CM: **Bumble bees damage plant leaves and accelerate flower production when pollen is scarce.** *Science (80-)* 2020, **368**:881-884.
62. Van Asch M, Salis L, Holleman LJM, Van Lith B, Visser ME: **Evolutionary response of the egg hatching date of a herbivorous insect under climate change.** *Nat Clim Change* 2013, **3**:244-248.
63. Persche ME, Pidgeon AM: **Habitat resilience for songbirds: the role of topographic position in a mixed deciduous forest.** *For Ecol Manage* 2020, **472**:118238.
64. Shutt JD, Burgess MD, Phillimore AB: **A spatial perspective on the phenological distribution of the spring woodland caterpillar peak.** *Am Nat* 2019, **194**:E109-E121.
65. Andrade JF, Alvarado F, Carlos Santos J, Santos BA: **Rainfall reduction increases insect herbivory in tropical herb communities.** *J Veg Sci* 2020, **31**:487-496.
66. Hill SJ, Silcocks SC, Andrew NR: **Impacts of temperature on metabolic rates of adult *Extatosoma tiaratum* reared on different host plant species.** *Physiol Entomol* 2020, **45**:7-15.
67. Miller-Struttman NE, Geib JC, Franklin JD, Kevan PG, Holdo RM, Ebert-May D, Lyn AM, Kettenbach JA, Elizabeth H, Galen C: **Functional mismatch in a bumble bee pollination mutualism under climate change.** *Science (80-)* 2015, **349**:75-78.
68. Moore ME, Kester KM, Kingsolver JG: **Rearing temperature and parasitoid load determine host and parasitoid performance in *Manduca sexta* and *Cotesia congregata*.** *Ecol Entomol* 2020, **45**:79-89.
69. Ahmed SS, Liu D, Simon JC: **Impact of water-deficit stress on tritrophic interactions in a wheat-aphid-parasitoid system.** *PLoS One* 2017, **12**:1-17.
70. Nguyen LTH, Monticelli LS, Desneux N, Metay-Merrien C, Amiens-Desneux E, Lavoit AV: **Bottom-up effect of water stress on the aphid parasitoid *aphidius ervi*.** *Entomol Gen* 2018, **38**:15-27.
71. Sentis A, Ramon-Portugal F, Brodeur J, Hemptinne JL: **The smell of change: warming affects species interactions mediated by chemical information.** *Glob Change Biol* 2015, **21**:3586-3594.
72. Barton BT, Schmitz OJ: **Experimental warming transforms multiple predator effects in a grassland food web.** *Ecol Lett* 2009, **12**:1317-1325.
73. Barton BT, Schmitz OJ: **Opposite effects of daytime and nighttime warming on top-down control of plant diversity.** *Ecology* 2018, **99**:13-20.
74. Bretagnolle V, Terraube J: **Predator-prey interactions and climate change.** In *Effects of Climate Change on Birds*. Edited by Dunn PO, Møller AP. Oxford University Press; 2019:199-220.
75. Bensadia F, Boudreault S, Guay JF, Michaud D, Cloutier C: **Aphid clonal resistance to a parasitoid fails under heat stress.** *J Insect Physiol* 2006, **52**:146-157.
76. Hurst GDD, Johnson AP, Schulenburg JHGVD, Fuyama Y: **Male-killing *Wolbachia* in *Drosophila*: a temperature-sensitive trait with a threshold bacterial density.** *Genetics* 2000, **156**:699-709.
77. Anbutsu H, Goto S, Fukatsu T: **High and low temperatures differently affect infection density and vertical transmission of male-killing *Spiroplasma* symbionts in *Drosophila* hosts.** *Appl Environ Microbiol* 2008, **74**:6053-6059.
78. Paudel S, Lin PA, Hoover K, Felton GW, Rajotte EG: **Asymmetric responses to climate change: temperature differentially alters herbivore salivary elicitor and host plant responses to herbivory.** *J Chem Ecol* 2020, **46**:891-905.
79. Carnicer J, Stefanescu C, Vives-Inglá M, López C, Cortizas S, Wheat C, Vila R, Llusà J, Peñuelas J: **Phenotypic biomarkers of climatic impacts on declining insect populations: a key role for decadal drought, thermal buffering and amplification effects and host plant dynamics.** *J Anim Ecol* 2019, **88**:376-391.
80. Schneider D, Ramos AG, Córdoba-Aguilar A: **Multigenerational experimental simulation of climate change on an economically important insect pest.** *Ecol Evol* 2020, **23**:12893-12909 <http://dx.doi.org/10.1002/ece3.6847>.
81. Richard G, Le Trionnaire G, Danchin E, Sentis A: **Epigenetics and insect polyphenism: mechanisms and climate change impacts.** *Curr Opin Insect Sci* 2019, **35**:138-145.
82. Frank SD, Just MG: **Can cities activate sleeper species and predict future forest pests? A case study of scale insects.** *Insects* 2020, **11**.
83. Ren P, Néron V, Rossi S, Liang E, Bouchard M, Deslauriers A: **Warming counteracts defoliation-induced mismatch by increasing herbivore-plant phenological synchrony.** *Glob Change Biol* 2020, **26**:2072-2080.