

Neutron–Mirror-Neutron Oscillation and Neutron Star Cooling

Itzhak Goldman¹,[✉] Rabindra N. Mohapatra,² Shmuel Nussinov,³ and Yongchao Zhang⁴

¹*Afeke College and Tel Aviv University, 6195001 Tel Aviv, Israel*

²*Maryland Center for Fundamental Physics and Department of Physics, University of Maryland, College Park, Maryland 20742, USA*

³*Tel Aviv University, 6195001 Tel Aviv, Israel*

⁴*School of Physics, Southeast University, Nanjing 211189, China*



(Received 23 March 2022; accepted 21 July 2022; published 5 August 2022)

It was pointed out in a recent paper that the observed cooling rate of old, cold neutron stars (NS) can provide an upper limit on the transition rate of neutron to mirror neutron ($n - n'$). This limit is so stringent that it would preclude any discovery of $n \rightarrow n'$ oscillation in the current round of terrestrial searches for the process. Motivated by this crucially important conclusion, we critically analyze this suggestion and note an interesting new effect present in nearly exact mirror models for $n \rightarrow n'$ oscillation, which significantly affects this bound. The new element is the β decay $n' \rightarrow p' + e' + \bar{\nu}_e$, which creates a cloud of mirror particles n' , p' , e' , and D' inside the NS core. The e' can “rob” the energy generated by the $n \rightarrow n'$ transition via $e - e'$ scattering enabled by the presence of a (minute) millicharge in mirror particles. This energy is emitted as unobserved mirror photons via fast mirror bremsstrahlung leading to a relaxation of this upper limit.

DOI: [10.1103/PhysRevLett.129.061103](https://doi.org/10.1103/PhysRevLett.129.061103)

Introduction.—Neutron stars (NSs) and their origin from supernovae have played an important role in constraining physics beyond the standard model (BSM) [1]. One class of BSM scenarios which can lead to new effects in NSs are the mirror models, which consist of a mirror sector coexisting with the standard model (SM) and which contains a parity symmetric duplicate of the particles and forces of the SM [2]. When the mirror parity is nearly exact, all particles in the two sectors including the neutron and mirror neutron are nearly degenerate. This raises the possibility of neutrons oscillating to mirror neutrons ($n \rightarrow n'$) [3] if the sum of ordinary (B) and mirror (B') baryon numbers is conserved. This phenomenon has been proposed as a solution to the neutron lifetime anomaly [4]. There are a number of experiments already carried out or planned to search for this $n \rightarrow n'$ oscillation [5]. It is therefore important to know if there are any constraints on the $n - n'$ mixing parameter $\epsilon_{nn'}$ from astrophysical settings. Since NSs are extremely rich in neutrons, they are a perfect laboratory for testing implications of $n \rightarrow n'$ oscillation.

The transition of an ordinary neutron n to a mirror neutron n' is followed by a migration of the latter toward the NS center under gravity. The hole left will then be filled by another neutron at the Fermi level, and in the process

energy is liberated [6]. If the process is fast enough, it would lead to a fully mixed star. The resulting mass loss of a NS will not only lead to changes in the orbital period of a binary pulsar [7], but also affect the luminosity of a single NS [6,8]. The observational constraints on the rate of the binary periods for several binary pulsars were shown to lead to upper bounds on $\epsilon_{nn'}$ of 10^{-13} eV [9]. On the other hand, taking the coldest NS, i.e., PSR J2144-3933 [10], it was argued in Refs. [6,8] that one gets $\epsilon_{nn'} \leq 10^{-17}$ eV. Both the bounds are valid for $n - n'$ mass difference up to 15 MeV [9]. This luminosity limit is particularly important, since currently planned terrestrial experiments are sensitive to $\epsilon_{nn'}$ at the level of 10^{-17} eV [5]. Note that in terrestrial searches for $n - n'$ oscillation, to maintain coherent buildup of the mirror neutron wave function along the neutron beam, and allow for such sensitive measurements, one must require a remarkably precise degeneracy between the neutron and its mirror partner of $\delta_{nn'}/m_n \leq 10^{-26}$ with $\delta_{nn'} \equiv |m_{n'} - m_n|$.

In this Letter, we critically analyze the luminosity bound, by following the evolution of the n' generated in $n \rightarrow n'$ transition a bit longer. We observe that in almost exact mirror models, the mirror neutrons generated inside the NS β decay producing mirror fermions e' , p' , and $\bar{\nu}_e$ leading eventually to a cloud of e' and deuterons D' . These mirror particles then provide a competing cooling channel via the emission of mirror photons γ' , and reduce the photonic signal claimed in Ref. [8] considerably, relaxing the upper bounds on $\epsilon_{nn'}$. For a relatively wide acceptable range of interactions between the ordinary and mirror sectors, mediated by the millicharge of mirror particles [11], the

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

nucleons and electrons of the visible sector in this core region of the NS can transfer their energy to the mirror particles. The latter then emit this energy via mirror photons γ' , which do not interact with the ordinary nucleons and electrons and can freely escape. The philosophy of this Letter is similar to that in Ref. [12]. The millicharge on mirror particles arises if γ and γ' have kinetic mixing.

$n \rightarrow n'$ Transition.—Initially, shortly after its birth, a NS is relatively hot and cools down via volume emission of neutrino pairs. At the time of observation, the star may be still cooling off or, if some other sources of energy exist, it may have settled into a thermal steady state, with the thermal energy emitted as electromagnetic radiation often as a black body radiation [13,14]. Let us apply this scenario to the pulsar PSR J2144-3933. In a steady state, the NS black body luminosity is given by the Stefan-Boltzmann formula $\mathcal{L}_{\text{NS}} = 4\pi\sigma_{\text{SB}}R^2T_s^4$, where σ_{SB} is the Stefan-Boltzmann constant, R is the radius of the NS, and its external surface temperature T_s is maintained by the constant internal energy source. If we have observational limits on the luminosity, this implies upper bounds on the rate of internal heat production. It is important to note that there is a ~ 100 meter thick nuclear “thermal blanket” just under the surface [15]. It causes the internal temperature, which is almost uniform over the NS, to drop dramatically by a factor of ~ 100 as we move out from the inside across the blanket toward the surface. The estimated upper bound on surface temperature $T_s \sim 42000$ K of the coldest pulsar PSR J2144-3933 would then correspond to the internal temperature $T_{\text{int}} \simeq 0.35$ keV, which would play an important role in obtaining upper bounds on any heat generating mechanism.

If the $n \rightarrow n'$ processes were the only source of heat supply, then in a steady state the overall $n - n'$ transition rate would be given by $d\mathcal{N}_{n'}/dt = \mathcal{L}_{\text{NS}}/\Delta E$, where $\Delta E \sim 30$ MeV is the energy initially gained by ordinary nucleons in each $n \rightarrow n'$ transition. For PSR J2144-3933, taking $R = 11$ km, the rate of generating new mirror neutrons turns out to be

$$\frac{d\mathcal{N}_{n'}}{dt} \sim 0.45 \times 10^{32} \left(\frac{T_s}{42000 \text{ K}} \right)^4 \text{ sec}^{-1}. \quad (1)$$

During its long lifetime of 330 million years, about $\mathcal{N}_{n'} \sim 10^{48}$ neutrons would have converted into mirror neutrons. This comprises a tiny $\mathcal{N}_{n'}/\mathcal{N}_n \sim 10^{-9}$ fraction of the total neutron number $\mathcal{N}_n \sim 2 \times 10^{57}$ in the star, with no change of the gravity fields and of the local density profile of the ordinary NS. Some pulsars have temperatures up to 100 times higher yielding $d\mathcal{N}_{n'}/dt \sim 10^{40} \text{ sec}^{-1}$, and were also used to bound high $\epsilon_{nn'}$ values [13,14].

Neighboring neutrons rush into the “hole” formed by $n \rightarrow n'$ transition, and the work done in the process is ~ 30 MeV on average and becomes the kinetic energy of these nucleons. The nucleons collide with neighboring neutrons with density $n_N \sim 10^{39} \text{ cm}^{-3}$, and very quickly

settle into the spatially and temporally fixed internal temperature $T_{\text{int}} (\sim 0.35 \text{ keV})$. It should be noted that only the $f = kT/E_F$ fraction of nucleons and electrons in the high energy tail of the degenerate Fermi-Dirac energy distribution are not Pauli blocked and can be excited (or de-excited) to higher (or lower) empty energy states, reducing the specific heat and the heat content Q^* of the NS by a factor of f . It is then given by

$$Q^* = \mathcal{N}_n f^2 E_F. \quad (2)$$

Upon using $kT \simeq 0.35 \text{ keV}$ for PSR J2144-3933 and $E_F = 30 \text{ MeV}$, we find $Q^* \sim 10^{52} \text{ keV}$, with only the $f \sim 10^{-5}$ fraction of these end point “active” electrons partaking in electron scattering or any other dynamic processes, which will play an important role in the following calculations.

n' Decay and the $e' - D'$ fluid.—In connection with the extreme degeneracy of n and n' , there are three extra light neutrinos and the mirror photon in exact mirror models. To bring about consistency between three extra neutrinos and an extra photon contributing to the energy density in the big bang nucleosynthesis (BBN) epoch of the universe with the Planck data [16], we require that there be asymmetric inflation implemented [17]. This will remove the BBN problem by lowering the reheat temperature in the mirror sector by a factor of 3, thus diluting the impact of the extra mirror neutrinos and the mirror photon on BBN.

The β decay $n' \rightarrow p' + e' + \bar{\nu}'_e$ of n' proceeds in the same manner as n , and will have the same rate of $\sim (800 \text{ sec})^{-1}$ as n decay in vacuum, so long as the Fermi energy of the electron is much smaller than the Q value of 0.7 MeV of the β decay [18]. The p' s, like the n' s, are gravitationally bound to the NS, and local mirror charge neutrality forces the number densities $n_{e',p'}(r)$ of e' and p' to be the same at all $r < R$, i.e., $n_{e'}(r) = n_{p'}(r)$. The mirror neutrons and mirror protons slow down and form mirror deuterons D' , since the process $p' + n' \rightarrow D' + \gamma'$ is faster than the inverse beta decay $e' + p' \rightarrow n' + \nu'$. All the p' s are “eaten up” to form D' , and the number of e' s that will remain is only half the number of n' produced. The resulting γ' s escape taking away part of the energy released in $n \rightarrow n'$ transition but it does not drain the energy generated by neutrons falling from the Fermi surface, which is drained away via $e - e'$ scattering. Charge neutrality requires that $n_{e'}(r) = n_{D'}(r)$, with $n_{D'}(r)$ the number density of D' . The new processes we consider are depicted schematically in Fig. 1.

The e' and D' constitute a fluid that is supported against the gravity of the ordinary NS by degenerate pressure, which is dominated by that of the e' . The mass density of the fluid is dominated by the D' . The corresponding hydrostatic equation is

$$\frac{\partial}{\partial r} P_{e'}(r) = -\rho(r)g(r), \quad (3)$$




FIG. 1. A schematic depiction of what happens after the $n \rightarrow n'$ transition takes place in a NS. In the right panel we enlarge the “mirror star” region in the left panel.

where $\rho(r) = n_{e'}(r)m_{D'}$ is the mass density of the D' , and the e' pressure for a given Fermi momentum p_F is

$$P_{e'} = \frac{8\pi}{3m_e\hbar^3} \int_0^{p_F} dp \frac{p^4}{\sqrt{1 + (p/m_{e'}c)^2}}. \quad (4)$$

For the small radii considered, the gravitational acceleration can be approximated by

$$g(r) = \frac{G_N M(r)}{r^2} = \frac{4\pi}{3} G_N \rho_0 r, \quad (5)$$

where G_N is the Newtonian constant of gravitation. For $r < 2$ km the density $\rho_0 \simeq 10^{15}$ gr cm $^{-3}$ in the center of the NS is almost a constant. The general relativistic modifications of the hydrostatic equation are very small, at the level of 10^{-3} . We can solve the hydrostatic equation (3) analytically and get

$$n_{e'}(r) = \frac{8\pi}{3m_{e'}^3 c^3 \hbar^3} \left[\left(\sqrt{X_F^2(0) + 1} - \frac{r^2}{2r_0^2} \right)^2 - 1 \right]^{3/2}, \quad (6)$$

where $r_0 = (3m_{e'}c^2/4\pi G_N \rho_0 m_{D'})^{1/2} \simeq 0.296$ km, and $X_F(0) = p_F(0)/m_{e'}c$. Then the number of the e' up to the radius r is

$$\mathcal{N}_{e'}(r) = \int_0^r 4\pi n_{e'}(x) x^2 dx. \quad (7)$$

The fluid is confined inside a sphere with radius R_c so that $n_{e'}(R_c) = 0$. Once $X_F(0)$ is given, R_c , $n_{e'}(r)$, and the

total number $\mathcal{N}_{e'} = \mathcal{N}_{D'}$ are determined by pure numbers and fundamental constants. This resembles the case of the Chandrasekhar mass. The dimensionless constant $X_F(0)$ is determined by $\mathcal{N}_{e'}(R_c) = 5 \times 10^{47}$ so that $\mathcal{N}_{e'} = \mathcal{N}_{D'}$ is half of the total n' generated. We obtain $X_F(0) \simeq 8.9$, implying $E_F(0) \simeq 4$ MeV and $R_c \simeq 1.18$ km. More details can be found in the Supplemental Material [19].

Energy drain from the visible sector to the mirror fluid.—In deriving the strict bound by using the electromagnetic luminosity $\mathcal{L} = dW/dt$ of the NS, a key point is that the rate of $n \rightarrow n'$ transition is constant and independent of any thermal or other variations (except for stopping when the mixed star forms, which happens after many Hubble times for the small values of $\epsilon_{nn'}$ considered). The $\sim 50\%$ of the heat generated which resides in the SM component is then radiated via a fixed black body luminosity [8]. Having all the mirror particles segregated in a “core region” (the orange region in Fig. 1) comprising $\sim 0.1\%$ of the star volume would have seemed to minimize their ability to intercept and impede ordinary heat emission and photon radiation from the mirror free, large outer region. This, in turn, would have suggested only minor luminosity reduction and no relaxing of the bounds on $\epsilon_{nn'}$. However, a more careful scrutiny shows that this simplistic argument is misleading.

The energy emission from the core will be dominated by the radiation of mirror photons, while the heat is continuously transferred from the normal sector to the mirror sector by scatterings of the normal and mirror electrons in the core region. For sufficiently large millicharge ϵ , the heat emission rate from the mirror particles may overtake the

normal emission rate from the external surface by an appreciable factor. The ordinary photonic energy may then account only for a small part of the energy generated inside the star. Furthermore, the cumulative effect of this over most of the star's history will reduce its heat content and push the internal and external surface temperatures to zero, quenching the photonic emission and destroying the steady state model envisioned.

Thanks to the mutual mirror electromagnetic scattering of the mirror particles inside the core region and attendant emission of the fast escaping mirror photons, the time required for their cooling off and equilibrating at a temperature T' is very short on a typical thermal timescale of $t_{\text{thermal}} = W^*/(dW/dt)$, where $W^* = Q^*$ is the total heat content of the star. Using Eq. (2) we find $t_{\text{thermal}} \sim 3 \times 10^{15}$ sec, which happens to be close to the age of the star.

Since the emission of heat from the mirror sector is much faster than heat transfer between the sectors, any amount of heat in the mirror sector will be emitted rather than go back to the normal sector, which also implies that $T' \leq T$. To avoid a detailed discussion at the particle scattering level, we first view the core region as a black body for the *mirror* photons with temperature T' , as indeed it absorbs any such photon falling on it. The surface of area $4\pi R_c^2$ of the inner “core region” serves effectively as an additional boundary, through which the heat in the normal component of the surrounding star can be emitted. The mirror electrons in the core will then radiate their heat content to the outside with the rate of black body luminosity: $\mathcal{L}' = 4\pi\sigma_{\text{SB}}R_c^2T'^4$. Relative to the internal core region surface $4\pi R_c^2$, the stellar surface is larger—by roughly a factor of 100. However, the thermal blanket makes the internal temperature about one hundredfold bigger than the surface temperature. Thanks to the possibility that $T'^4 \geq 10^8 T_s^4$, even if we keep $T' < T$ to make $e \rightarrow e'$ energy transfers more than the reverse transfer, we can still, in principle, have the rate of mirror photon emission almost 6 orders of magnitude bigger than that of the ordinary photons, so long as $R_c \geq 1$ km.

However, to verify that this indeed happens, we need to check how many $e - e'$ collisions occur per second (which we denote by $\dot{N}_{\text{col}}$) between the $\mathcal{N}_{e'}(r < R_c) \sim 10^{38} R_c^3 \text{ cm}^{-3}$ electrons in the core region and the $\mathcal{N}_{e'} \sim 5 \times 10^{47}$ mirror electrons. If the total energy transferred per second via these collisions from the ordinary to mirror electrons much exceeds the stellar luminosity, namely the inequality

$$\dot{N}_{\text{col}} \Delta E \sim \dot{N}_{\text{col}} \Delta T \gg \mathcal{L}_{\text{NS}} \sim 2 \times 10^{36} \text{ keV sec}^{-1} \quad (8)$$

holds, then the mirror luminosity dominates and the scenario envisioned in deriving the strict upper bounds on $\epsilon_{nn'}$ becomes inoperative. On the other hand, if the inequality in Eq (8) is (strongly) reversed, then the scenario

above involving the β decay of the mirror neutron will be irrelevant.

For the average energy transfer of $\Delta T \sim 0.35$ keV, Eq. (8) becomes $\dot{N}_{\text{col}} \geq 10^{37} \text{ sec}^{-1}$. Each electron and also each mirror electron move with the speed of light c . Then we can express $\dot{N}_{\text{cal}}$ with energy transfer of ~ 0.35 keV in a manner that is symmetric between the ordinary and mirror sectors:

$$\dot{N}_{\text{cal}} = \frac{c f f' \mathcal{N}_e(r < R_c) \mathcal{N}_{e'} \sigma_{ee'}}{(4\pi/3) R_c^3}, \quad (9)$$

where $f' = kT'/E'_F \sim 10^{-4}$ is the fraction of the “active” mirror electrons. For $R_c = 1.2$ km, the condition $\dot{N}_c \gg 10^{37} \text{ sec}^{-1}$ translates into the following requirement on the $e - e'$ scattering cross section:

$$\sigma_{ee'} \simeq \epsilon^2 \sigma_{ee} \gg 10^{-50} \text{ cm}^2, \quad (10)$$

where σ_{ee} is the standard Rutherford scattering cross section of electrons in the same kinematic configuration. For the formula for $\sigma_{ee'}$, see the Supplemental Material [19]. Including only the Feynman diagram for the t -channel photon exchange, the cross section $\sigma_{ee'}$ is calculated by having the relativistic e and e' with energies $E_F \simeq 10E'_F \simeq 35$ MeV collide at random relative direction in the laboratory frame and transferring an energy of $T \sim 0.35$ keV between them. Using a plasmon mass as the cutoff, we estimate this cross section to be $\sigma_{ee'} \simeq 4\pi\alpha^2\epsilon^2/E_F T \simeq 10^{-23}\epsilon^2 \text{ cm}^2$, which leaves us with the rather weak, easy to satisfy requirement

$$\epsilon^2 \gg 10^{-27}. \quad (11)$$

The strongest upper bounds $\epsilon \leq 10^{-12}$ [20] do not apply here, as in mirror models the dark matter (DM) is made of neutral objects such as the $p' - e'$ composite mirror hydrogen, deuteron, or helium. On the other hand, $\epsilon \leq 10^{-9}$ required for cosmological constraint consistent with BBN limits is more directly applicable here [21] whereas a weaker limit of $\epsilon \leq 10^{-7}$ comes from the consistency of asymmetric inflation [22]. This still leaves 9 orders of magnitude margin for satisfying Eq. (11).

We also note that even though the photonic cooling of ultracold NS (UCNS) is not a reliable way to set bounds on the $n \rightarrow n'$ transition rate for near exact mirror models and slow $n \rightarrow n'$ transition, there are situations when it works: e.g., we could have (i) a near exact mirror symmetry but the millicharge of the mirror fermions $\epsilon \leq 10^{-13}$ or (ii) an asymmetric mirror model with $m_{p'} \geq m_{n'}$ where n' is the DM of the universe, so that β decay of the mirror neutron is forbidden. It can also work in other dark baryon contexts, such as those suggested in connection with the neutron lifetime anomaly.

An advantage of the heating up argument as compared with the orbital period stability method [9] is that in the former one can use all pulsars, whereas the latter case requires binary pulsars [7,9]. Unfortunately, unlike the misquote in Ref. [8], the spinning period changes of single pulsars—which, as part of the ambitious nanogravity project, is determined in many cases with stunning accuracy—cannot be used, as it is affected by relatively large and incalculable changes due to magnetic braking, etc. This is the reason why binary pulsars were used in Refs. [7,9].

Conclusion.—To summarize our main result, the photonic luminosities of UCNSs do not necessarily imply robust bounds on $\epsilon_{nn'}$. In particular, they do NOT exclude discovery via terrestrial measurements of the tiny $\epsilon_{nn'} \sim \mathcal{O}(10^{-17})$ eV. This happens due to the beta decay of n' following $n \rightarrow n'$ transition and the subsequent deuteron formation. Our main assumption, the existence of a millicharge ϵ , is definitely allowed and possibly even favored within mirror models. In this scenario, under the joint effect of the weight of the mirror deuterons and the Fermi energy of the mirror electrons, the mirror deuterons and electrons form a configuration resembling that of a “mini white dwarf” inside the NS. A remarkable feature of this configuration is its universality stemming from, and in analogy with, the features of NSs and actual white dwarfs. Within this structure, heat is transferred relatively fast (on characteristic thermal timescales of the NS) from the heat reservoir in the normal matter of the NS to the mirror sector, and is radiated via mirror photons.

The work of R. N. M. is supported by the US National Science Foundation Grant No. PHY-1914631. Y. Z. is supported by the National Natural Science Foundation of China under Grant No. 12175039, the 2021 Jiangsu Shuangchuang (Mass Innovation and Entrepreneurship) Talent Program No. JSSCBS20210144, and the “Fundamental Research Funds for the Central Universities.”

[1] G. G. Raffelt, *Stars as Laboratories for Fundamental Physics* (Chicago University Press, Chicago, 1996).
 [2] T. D. Lee and C. N. Yang, *Phys. Rev.* **104**, 254 (1956); I. Yu. Kobzarev, L. B. Okun, and I. Ya. Pomeranchuk, *Yad. Fiz.* **3**, 1154 (1966); S. I. Blinnikov and M. Yu. Khlopov, *Sov. Astron.* **27**, 371 (1983); R. Foot, H. Lew, and R. R. Volkas, *Phys. Lett. B* **272**, 67 (1991); H. M. Hodges, *Phys. Rev. D* **47**, 456 (1993); R. Foot and R. R. Volkas, *Phys. Rev. D* **52**, 6595 (1995); Z. G. Berezhiani and R. N. Mohapatra, *Phys. Rev. D* **52**, 6607 (1995); Z. Berezhiani, D. Comelli, and F. L. Villante, *Phys. Lett. B* **503**, 362 (2001); Z. Berezhiani, P. Ciarcelluti, D. Comelli, and F. L. Villante, *Int. J. Mod. Phys. D* **14**, 107 (2005); A. Y. Ignatiev and R. R. Volkas, *Phys. Rev. D* **68**, 023518 (2003); R. Foot, *Int. J. Mod. Phys. A* **29**, 1430013 (2014); F. Sandin and P. Ciarcelluti, *Astropart. Phys.* **32**, 278 (2009).

[3] Z. Berezhiani and L. Bento, *Phys. Rev. Lett.* **96**, 081801 (2006).
 [4] Z. Berezhiani, *Lett. High Energy Phys.* **2**, 118 (2019); *Eur. Phys. J. C* **79**, 484 (2019).
 [5] N. J. Ayres, Z. Berezhiani, R. Biondi, G. Bison, K. Bodek, V. Bondar, P. J. Chiu, M. Daum, R. T. Dinani, C. B. Doorenbos *et al.*, *Symmetry* **14**, 503 (2022); G. Ban *et al.*, *Phys. Rev. Lett.* **99**, 161603 (2007); Y. N. Pokotilovski, *Phys. Lett. B* **639**, 214 (2006); A. P. Serebrov, E. B. Aleksandrov, N. A. Dovator, S. P. Dmitriev, A. K. Fomin, P. Geltenbort, A. G. Kharitonov, I. A. Krasnoschekova, M. S. Lasakov, A. N. Murashkin *et al.*, *Phys. Lett. B* **663**, 181 (2008); C. Abel *et al.* (nEDM Collaboration), *Phys. Lett. B* **812**, 135993 (2021); P. Mohanmurthy, A. R. Young, J. A. Winger, and G. Zsigmond, [arXiv:2201.04191](https://arxiv.org/abs/2201.04191); L. J. Broussard, J. L. Barrow, L. DeBeer-Schmitt, T. Dennis, M. R. Fitzsimmons, M. J. Frost, C. E. Gilbert, F. M. Gonzalez, L. Heilbronn, E. B. Iverson *et al.*, [arXiv:2111.05543](https://arxiv.org/abs/2111.05543); Y. Kamyshev, J. Ternullo, L. Varriano, and Z. Berezhiani, *Symmetry* **14**, 230 (2022); Z. Berezhiani, R. Biondi, P. Geltenbort, I. A. Krasnoschekova, and V. E. Varlamov, *Eur. Phys. J. C* **78**, 717 (2018).
 [6] Z. Berezhiani, R. Biondi, M. Mannarelli, and F. Tonelli, *Eur. Phys. J. C* **81**, 1036 (2021); M. Mannarelli, Z. Berezhiani, R. Biondi, and F. Tonelli, *Proceedings of the Nordita ESS Workshop* (2018).
 [7] I. Goldman and S. Nussinov, *J. High Energy Phys.* **08** (2010) 091.
 [8] D. McKeen, M. Pospelov, and N. Raj, *Phys. Rev. Lett.* **127**, 061805 (2021).
 [9] I. Goldman, R. N. Mohapatra, and S. Nussinov, *Phys. Rev. D* **100**, 123021 (2019).
 [10] S. Guillot, G. G. Pavlov, C. Reyes, A. Reisenegger, L. Rodriguez, B. Rangelov, and O. Kargaltsev, *Astrophys. J.* **874**, 175 (2019).
 [11] Bob Holdom, *Phys. Lett.* **166B**, 196 (1986).
 [12] Zurab Berezhiani, *Int. J. Mod. Phys. A* **19**, 3775 (2004).
 [13] D. G. Yakovlev and C. J. Pethick, *Annu. Rev. Astron. Astrophys.* **42**, 169 (2004).
 [14] A. Y. Potekhin, D. A. Zyuzin, D. G. Yakovlev, M. V. Beznogov, and Yu. A. Shibarov, *Mon. Not. R. Astron. Soc.* **496**, 5052 (2020).
 [15] M. V. Beznogov, A. Y. Potekhin, and D. G. Yakovlev, *Phys. Rep.* **919**, 1 (2021).
 [16] N. Aghanim *et al.* (Planck Collaboration), *Astron. Astrophys.* **641**, A1 (2020).
 [17] Z. G. Berezhiani, A. D. Dolgov, and R. N. Mohapatra, *Phys. Lett. B* **375**, 26 (1996).
 [18] As in the normal sector, there might be Zeeman effect in the mirror sector which lead to corrections to the n' mass thus affecting the mirror β decay. However, such effect is much less important than the $n - n'$ mass splitting $\Delta M \sim \mathcal{O}(\text{MeV})$ under gravity in the star, and can be safely neglected.
 [19] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.061103> for the supplementary material has two parts: First part presents a derivation and plot of the profile of mirror electron as a function of radial distance from the center of the neutron

- star; Second part gives the formula for the $e - e'$ scattering due to millicharge of e'^* .
- [20] H. Vogel and J. Redondo, *J. Cosmol. Astropart. Phys.* **02** (2014) 029.
- [21] Z. Berezhiani and A. Lapedi, *Phys. Lett. B* **681**, 276 (2009).
- [22] K. S. Babu and R. N. Mohapatra, *Symmetry* **14**, 731 (2022).