

# Phase separation of Dri1 contributes to heterochromatin formation in *Schizosaccharomyces pombe*

Hyoju Ban<sup>1\*</sup>, Wenqi Sun<sup>2,3\*</sup>, Yong Chen<sup>2§</sup>, Fei Li<sup>1§</sup>

## **Abstract**

The RNA binding protein Dri1 facilitates heterochromatin assembly via the RNAi pathway and histone deacetylases (HDAC). Dri1 contains an intrinsically disordered region (IDR) and three zinc fingers at its C-terminus, which are important for its role in heterochromatin silencing. Both IDR and zinc fingers have been implicated in mediating liquid-liquid phase separation (LLPS). In this study, we investigated the phase separation properties of Dri1. We observed that Dri1 undergoes phase separation *in vitro*. Dri1 also exhibits liquid-like behavior *in vivo*. Combined with our previous findings, our data support a model in which the phase-separated condensates formed by Dri1 may help recruit RNAi components and HDAC to mediate heterochromatin assembly.

<sup>&</sup>lt;sup>1</sup>Department of Biology, New York University, New York, NY 10003, USA.

<sup>&</sup>lt;sup>2</sup>Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai, 200031, China.

<sup>&</sup>lt;sup>3</sup>University of Chinese Academy of Sciences, Beijing 100049, China.

<sup>§</sup>To whom correspondence should be addressed: yongchen@sibcb.ac.cn; fl43@nyu.edu

<sup>\*</sup>These authors contributed equally.




Figure 1. Dri1 exhibits liquid-like phase-separated behavior in vitro and in vivo.

A. Fluorescence microscopy images of liquid-like droplets formed by Dri1-C-GFP at 150 mM NaCl. B. Time-lapse fluorescence microscopy of droplet-like structures formed by Dri1-C-GFP at 150 mM NaCl. Note: the spherical droplets in the solution can flow and fuse. C. Fluorescence microscopy images of liquid-like droplets form by Dri1-C-GFP in the presence of 10% Dextran or Ficoll at 150 mM NaCl. D. Top: Lowly overexpressed GFP-Dri1 induces liquid-like condensates *in vivo*. Time-lapse fluorescence microscopy analysis of a typical live cell expressing GFP-Dri1. Lower: Time-lapse fluorescence microscopy of a typical live cell expressing GFP-Dri1-ΔC. Cells with lowly overexpressed GFP-Dri1-ΔC displays either diffuse signal or 1-2 GFP foci. Unlike the full-length Dri1, these GFP foci are immobile.

## **Description**

Liquid-liquid phase separation (LLPS) of proteins has been implicated in a variety of biological processes, such as formation of membrane less organelles and transcriptional regulation (Hnisz *et al.* 2017; Shin and Brangwynne 2017). LLPS can drive compartmentalization that concentrates biochemical reactions within cells. Heterochromatin is a transcriptionally silent chromatin domain in eukaryotes that is essential for genome stability and gene expression regulation. HP1, a conserved heterochromatin protein, in *Drosophila* and human cells has been shown to exhibit liquid-like phase-separated behavior (Larson *et al.* 2017; Strom *et al.* 2017). But the role of LLPS in heterochromatin formation remains to be elucidated. The fission yeast *Schizosaccharomyces pombe* has emerged as an excellent model system for the study of heterochromatin

#### 4/19/2022 - Open Access

formation (He et al. 2014). We recently showed that a RNA-binding protein Dri1 plays an important role in heterochromatin assembly in fission yeast (Ban et al. 2021). Dri1 physically associates with heterochromatic transcripts, and is required for the recruitment of the RNA-induced transcriptional silencing (RITS) complex. Furthermore, Dri1 contributes to the association of the Sir2 histone deacetylase with heterochromatin (Ban et al. 2021). In addition to the RNA recognition motif (RRM), Dri1 contains an intrinsically disordered region (IDR) at its C-terminus, which is shared by many proteins undergoing LLPS (Shin and Brangwynne 2017). The C-terminus of Dri1 also contains three zinc fingers. Deletion of the disorder domain or zinc fingers results in defects in heterochromatin silencing, indicating that they are important for the role of Dri1 in heterochromatin silencing (Ban et al. 2021). Both IDR and zinc fingers have been shown to be able to drive protein phase separation (Schwartz et al. 2013; Shin and Brangwynne 2017; Maharana et al. 2018). However, whether Dri1 has the liquidliquid phase separation property has not been demonstrated. To determine whether Dri1 can undergo LLPS in vitro, we fused the different fragments of Dri1 with GFP and test their liquid-droplet formation. Full-length Dri1 fused with GFP (Dri1-GFP) formed solid-like precipitation under physiological conditions (150 mM NaCl, pH7.0-8.0). On the contrary, the C-terminus of Dri1 including three zinc fingers and the IDR with GFP (Dri1-C-GFP) displayed liquid-like phase separation under the same condition. Brightly fluorescent, micrometer-sized droplets are visible for Dri1-C-GFP by fluorescence microscopy (Fig. 1A). In addition, our time-lapse fluorescence microscopy analysis revealed that the spherical droplets in the solution can flow and fuse (Fig. 1B). Thus, Dri1-C-GFP can undergo LLPS in vitro. To further investigate how intracellular crowding may affect LLPS property of Dri1, we examined Dri1-C-GFP in the presence of crowding agents, inert macromolecules used to study crowding, including 10% Dextran or Ficoll. We found that Dri1-C-GFP also underwent LLPS in crowders tested, and the number of droplets is substantially increased (Fig. 1C), suggesting that the crowding agents promote LLPS of Dri1 in vitro. We also tested whether Dri1 has liquid-like phase-separated behavior in vivo. We have previously shown that the signal of GFP-Dri1 expressed at its endogenous level in fission yeast is very weak (Ban et al. 2021). We therefore constructed GFP-Dri1 under the thiamine-repressible *nmt1* promoter, and lowly overexpressed GFP-Dri1 by supplementing with 0.05 mM thiamine for 18 hours. We previously showed that lowly overexpressed GFP-Dri1 formed distinct foci mostly in the nucleus (Ban et al. 2021). Careful examination of the GFP-Dri1 signal revealed that these foci are usually spherical, a typical shape formed for proteins having liquid-like state (Fig. 1D). Our time-lapse microscopy analysis further revealed that the GFP-Dri1 foci were very dynamic in movement, and can frequently fuse together (Fig. 1D), suggesting that Dri1 has liquid-liquid phase separation properties in vivo. Furthermore, we deleted the C-terminus of Dri1, which contains the IDR region and three zinc fingers (Dri1- $\Delta C$ ). We found that lowly overexpression of GFP-Dri1- $\Delta C$  results in few immobile foci or dispersed GFP signal, indicating that the liquid phase separation property is lost in the mutant (Fig. 1D). It has been shown that overexpression of proteins with LLPS property impairs cell growth (Bolognesi et al. 2016). In agreement with this, we also showed that that strong overexpression of Dri1 results in severe growth defects (Ban et al. 2021). Our data thus demonstrate that C-terminus of Dri1 displays liquid-like property, which contributes to its role in heterochromatin silencing. We propose that the phase separated condensates formed by Dri1 may help recruit RNAi components and HDAC to mediate heterochromatin assembly.

## Methods

## Media and DNA constructs

Standard media for fission yeast growth were used (Moreno *et al.* 1991). Full-length Dri1 and C-terminus-deleted Dri1 (Dri1- $\Delta$ C) were cloned into the pREP1 vector containing in-frame GFP under the *nmt1* promoter. The constructs were lowly overexpressed by supplementing the medium with 0.05 mM thiamine for 18 hours.

#### Protein expression and purification

Dri1 fragments were cloned into a modified pET28b vector with a 6xHis-SUMO protein fused at the N terminus and a GFP protein fused at the C terminus, and were expressed in Rosetta cells. Dri1 fragments were purified by using Ni-NTA agarose beads (Qiagen). Ulp1 protease was subsequently added to remove the N-terminal 6xHis-SUMO tag, and Dri1-GFP proteins were collected after ULP1 on-beads digestion. The proteins were further purified by gel-filtration chromatography on Hiload Superdex 200 column (GE Healthcare).

#### In vitro phase separation assays

For the droplet formation, proteins were diluted to the final concentration (30 mM) with the phase separation buffer (25mM Tris-HCl pH 8.0,150 mM NaCl, 0.2mM DTT) in 0.2 ml PCR tubes. In experiments using Dextran-70 or Ficoll 400 as crowding agents, crowders were added to reach the final concentrations (10%) in the phase separation buffer. Fluorescent images were acquired on a Zeiss LSM 710 microscope with a 63x objective.

### Microscopic analysis



#### 4/19/2022 - Open Access

Microscopic analysis was performed as described (Dong *et al.* 2016). Briefly, fluorescence signal was captured by DeltaVision system (Applied Precision, Issaquah, WA). Images were taken as z-stacks of 0.2-μm increments with an oil immersion objective (×100).

## Reagents

All the *in vivo* experiments were performed using the haploid strains of *S. pombe* (FL 255  $h^-$  *leu1-32 ura4-D18 ade6-216 his3-D1*).

#### References

Ban H, Sun W, Chen YH, Chen Y, Li F. 2021. Dri1 mediates heterochromatin assembly via RNAi and histone deacetylation. Genetics 218: . PubMed ID: <u>33693625</u>

Bolognesi B, Lorenzo Gotor N, Dhar R, Cirillo D, Baldrighi M, Tartaglia GG, Lehner B. 2016. A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity upon Increased Protein Expression. Cell Rep 16: 222-231. PubMed ID: 27320918

Dong Q, Yin FX, Gao F, Shen Y, Zhang F, Li Y, He H, Gonzalez M, Yang J, Zhang S, Su M, Chen YH, Li F. 2016. Ccp1 Homodimer Mediates Chromatin Integrity by Antagonizing CENP-A Loading. Mol Cell 64: 79-91. PubMed ID: <u>27666591</u>

He H, Gonzalez M, Zhang F, Li F. 2014. DNA replication components as regulators of epigenetic inheritance--lesson from fission yeast centromere. Protein Cell 5: 411-9. PubMed ID: <u>24691906</u>

Hnisz D, Shrinivas K, Young RA, Chakraborty AK, Sharp PA. 2017. A Phase Separation Model for Transcriptional Control. Cell 169: 13-23. PubMed ID: <u>28340338</u>

Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA, Redding S, Narlikar GJ. 2017. Liquid droplet formation by HP1 $\alpha$  suggests a role for phase separation in heterochromatin. Nature 547: 236-240. PubMed ID: 28636604

Maharana S, Wang J, Papadopoulos DK, Richter D, Pozniakovsky A, Poser I, Bickle M, Rizk S, Guillén-Boixet J, Franzmann TM, Jahnel M, Marrone L, Chang YT, Sterneckert J, Tomancak P, Hyman AA, Alberti S. 2018. RNA buffers the phase separation behavior of prion-like RNA binding proteins. Science 360: 918-921. PubMed ID: 29650702

Moreno S, Klar A, Nurse P. 1991. Molecular genetic analysis of fission yeast Schizosaccharomyces pombe. Methods Enzymol 194: 795-823. PubMed ID: <u>2005825</u>

Schwartz JC, Wang X, Podell ER, Cech TR. 2013. RNA seeds higher-order assembly of FUS protein. Cell Rep 5: 918-25. PubMed ID: 24268778

Shin Y, Brangwynne CP. 2017. Liquid phase condensation in cell physiology and disease. Science 357: PubMed ID: 28935776

Strom AR, Emelyanov AV, Mir M, Fyodorov DV, Darzacq X, Karpen GH. 2017. Phase separation drives heterochromatin domain formation. Nature 547: 241-245. PubMed ID: <u>28636597</u>

Funding: This project was supported by NIH grant R35GM134920-01 and NSF grant MCB-1934628.

**Author Contributions:** Hyoju Ban: investigation, methodology, formal analysis. Wenqi Sun: investigation, methodology, formal analysis. Yong Chen: conceptualization, methodology, funding acquisition, supervision, writing - review editing. Fei Li: conceptualization, methodology, funding acquisition, supervision, writing - original draft.

Reviewed By: Qingfei Zheng

History: Received February 1, 2022 Revision Received April 15, 2022 Accepted April 19, 2022 Published April 19, 2022

**Copyright:** © 2022 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Ban, H; Sun, W; Chen, Y; Li, F (2022), Phase separation of Dri1 contributes to heterochromatin formation in *Schizosaccharomyces pombe*. microPublication Biology. <u>10.17912/micropub.biology.000559</u>