

Large time behavior, bi-Hamiltonian structure and kinetic formulation for complex Burgers equation

Yu Gao^{*1}, Yuan Gao^{†2}, and Jian-Guo Liu^{‡3}

¹Department of Mathematics, The University of Hong Kong, Pokfulam, Hong Kong.

²Department of Mathematics, Duke University, Durham, NC 27708, USA.

³Department of Mathematics and Department of Physics, Duke University, Durham, NC 27708, USA.

Abstract

We prove the existence and uniqueness of positive analytical solutions with positive initial data to the mean field equation (the Dyson equation) of the Dyson Brownian motion through the complex Burgers equation with a force term on the upper half complex plane. These solutions converge to a steady state given by Wigner's semicircle law. A unique global weak solution with nonnegative initial data to the Dyson equation is obtained and some explicit solutions are given by Wigner's semicircle laws. We also construct a bi-Hamiltonian structure for the system of the real and imaginary components of the complex Burgers equation (coupled Burgers system). We establish a kinetic formulation for the coupled Burgers system and prove the existence and uniqueness of entropy solutions. The coupled Burgers system in Lagrangian variable naturally leads to two interacting particle systems, Fermi-Pasta-Ulam-Tsingou model with nearest-neighbor interactions, and Calogero-Moser model. These two particle systems yield the same Lagrangian dynamics in the continuum limit.

1 Introduction

Complex Burgers equation arises, although in different ways, from many different fields such as fluid mechanics, random surface minimizing problem and Burgers turbulence in quantum chromodynamics, which always unveils some mechanisms of singularity formations. We only list several examples here. [31] use complex Burgers equation to construct a family of singular solution to zero-gravity water wave system. [11] use the complex Burgers equation to study the limit shape and singularity formations of random surface models. For other applications of complex Burgers equation such as singularity tracking in the evolution of the complex system and the large-N limit of induced quantum chromodynamics we refer to [9] and the references therein.

In this paper, we study the complex Burgers equation with a force term $\gamma^2 z$ on the upper half complex plane $\mathbb{C}_+ := \{z : \Im(z) > 0\}$

$$\partial_t g + g \partial_z g = \gamma^2 z, \quad z \in \mathbb{C}_+, t > 0. \quad (1.1)$$

Here, $\gamma \geq 0$ is a constant. We use $\Re(z)$ and $\Im(z)$ to stand for the real and imaginary parts of a complex number z respectively.

Take the trace of a solution $g(z, t)$ to (1.1) on the real line and there are two real functions $u(x, t)$ and $\rho(x, t)$ such that

$$g(x, t) + \gamma x = u(x, t) + i\pi\rho(x, t), \quad x \in \mathbb{R}, t > 0, \quad (1.2)$$

*gaoyu90@hku.hk

†yuangao@math.duke.edu

‡jliu@phy.duke.edu

where π is the circumference ratio. If $g(z, t)$ is a \mathbb{C}_+ -holomorphic function, then we have the following relation between u and ρ

$$u(x, t) = (\pi H\rho)(x, t), \quad (1.3)$$

where $H\rho$ stands for the Hilbert transform of ρ given by

$$(H\rho)(x, t) = \frac{1}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{\rho(y, t)}{x - y} dy, \quad x \in \mathbb{R}.$$

1 Take (1.2) into (1.1) and we obtain the following nonlocal partial differential equation for ρ

$$\partial_t \rho + \partial_x [\rho(u - \gamma x)] = 0, \quad u(x, t) = (\pi H\rho)(x, t), \quad x \in \mathbb{R}, \quad t > 0. \quad (1.4)$$

2 The equation for u can be obtained from (1.4) by the Hilbert transform (see (2.6)). We refer to
3 (1.4) as the Dyson equation which is a mean field equation for the Dyson Brownian motion as
4 described below.

The $N \times N$ complex Hermitian matrices form a N^2 dimensional linear vector space over field \mathbb{R} . Consider a Hermitian matrices valued Ornstein-Uhlenbeck (OU) process $A(t) = (A_{jk}(t))_{N \times N}$ given by

$$\begin{cases} dA_{jj}(t) = \frac{1}{\sqrt{N}} dB_{jj}(t) - \gamma A_{jj}(t) dt, \quad j = 1, \dots, N, \\ d\Re A_{jk}(t) = \frac{1}{\sqrt{2N}} dB_{jk}^R(t) - \gamma \Re A_{jk}(t) dt, \quad j < k, \\ d\Im A_{jk}(t) = \frac{1}{\sqrt{2N}} dB_{jk}^I(t) - \gamma \Im A_{jk}(t) dt, \quad j < k, \end{cases} \quad (1.5)$$

with $A(0) = 0$. Here $B_{jj}(t)$ ($1 \leq j \leq N$), $B_{jk}^R(t)$, $B_{jk}^I(t)$ ($1 \leq j < k \leq N$), are N^2 independent standard Brownian motions in \mathbb{R} . The eigenvalues $\lambda_1(t) \leq \dots \leq \lambda_N(t)$ of $A(t)$ form some real stochastic processes. By applying Ito's formula to $\lambda_j(t) (= \lambda_j(A(t)))$, one can show that $\lambda_j(t)$ evolve by ([7, 8, 29])

$$d\lambda_j(t) = \frac{1}{\sqrt{N}} dB_j(t) + \frac{1}{N} \sum_{k \neq j} \frac{dt}{\lambda_j(t) - \lambda_k(t)} - \gamma \lambda_j(t) dt, \quad 1 \leq j \leq N. \quad (1.6)$$

This evolution of eigenvalues are referred to as the Dyson Brownian motion. One can refer to [8, 29] for more details about random matrices and the Dyson Brownian motion. It is well known that the effects of harmonic trap term $-\gamma \lambda_j(t) dt$ in the OU process (1.6) can be reformulated into the case $\gamma = 0$, i.e. (1.6) without the trap term, by a space-time rescaling. We describe this space-time rescaling for complex Burgers equation below. Let g be a solution to (1.1) and set

$$\tilde{g}(w, \tau) \sqrt{1 + 2\gamma\tau} = g(z, t) + \gamma z, \quad z = \frac{w}{\sqrt{1 + 2\gamma\tau}}, \quad t = \frac{1}{2\gamma} \log(1 + 2\gamma\tau). \quad (1.7)$$

Then, \tilde{g} is a solution to the Complex Burgers equation without the force term:

$$\partial_\tau \tilde{g} + \tilde{g} \partial_w \tilde{g} = 0. \quad (1.8)$$

5 Note that $\tilde{g}(\cdot, \tau)$ is a \mathbb{C}_+ -holomorphic ($\overline{\mathbb{C}_+}$ -holomorphic) solution to (1.8) if and only if $g(\cdot, t)$ is a
6 \mathbb{C}_+ -holomorphic ($\overline{\mathbb{C}_+}$ -holomorphic) solution to (1.1).

The mean field limit of the Dyson Brownian motion (1.6) yields the Dyson equation (1.4) ([25, 5, 2]), and (1.4) is a gradient flow in the probability measure space with Wasserstein distance with respect to a free energy functional given by [1, Chapter 11]

$$\begin{aligned} E(\rho(\cdot, t)) &= \frac{\gamma}{2} \int_{\mathbb{R}} x^2 \rho(x, t) dx - \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \log|x - y| \rho(x, t) \rho(y, t) dx dy \\ &=: E_h(\rho(\cdot, t)) + E_i(\rho(\cdot, t)). \end{aligned} \quad (1.9)$$

Here E_h is a harmonic trap energy and E_i is an interaction energy. Then, the Dyson equation (1.4) is recast to

$$\partial_t \rho - \partial_x \left[\rho \partial_x \left(\frac{\delta E}{\delta \rho} \right) \right] = 0, \quad \frac{\delta E}{\delta \rho} = \frac{\gamma}{2} x^2 - \int_{\mathbb{R}} \log |x - y| \rho(y, t) dy. \quad (1.10)$$

1 With initial data $\rho_0 > 0$ and $\rho_0 \in L^2(\mathbb{R}) \cap C^{0,\delta}(\mathbb{R})$, Castro and Córdoba [4] proved global
 2 existence and uniqueness of real analytical solutions for $t > 0$ to the case $\gamma = 0$ of (1.4). This
 3 instantaneous analytical property is suggested by the gradient flow structure (1.10). However,
 4 if there is $x_0 \in \mathbb{R}$ such that $\rho(x_0) = 0$, then the solution ρ will blow up in $H^s(\mathbb{R})$, $s > \frac{3}{2}$ at
 5 finite time [4]. Thanks to the transformation in (1.7), these two results hold also for $\gamma > 0$; see
 6 Theorem 2.1 and Remark 2.1. Moreover, we prove the existence and uniqueness of global weak
 7 solution $\rho \in L^\infty(0, T; H^{\frac{1}{2}}(\mathbb{R}) \cap L^1_+(\mathbb{R}))$ to (1.4) in Theorem 2.2. The global regularity or finite
 8 time blow-up in the space $H^s(\mathbb{R})$, $s \in (\frac{1}{2}, \frac{3}{2}]$ remain open.

The steady state for the Dyson equation is given by Wigner's semicircle law

$$\mu_1(dx) = \rho_1(x) dx := \frac{\sqrt{(4 - x^2)_+}}{2\pi} dx, \quad (1.11)$$

9 which has a compact support. Hence the solution ρ is not absolutely continuous with respect to
 10 the steady state and the relative entropy method can not be directly applied here. There are two
 11 methods to prove the convergence of solution ρ to its steady state. (i) For strictly positive initial
 12 data $\rho_0(x) > 0$, following the idea of [25] we prove the pointwise convergence as t goes to infinity
 13 using analytical method; see Appendix A. (ii) Notice the free energy $E(\rho)$ given by (1.9) for the
 14 Dyson equation consists a harmonic trap energy E_h and an interaction energy E_i . Since E_i is
 15 convex along generalized Wasserstein geodesics and E_h is γ -convex along Wasserstein geodesics,
 16 the standard gradient flow theory yields W_2 -contraction and hence the exponential convergence
 17 to the steady state in Wasserstein distance (see Remark 2.4 and Carrillo et. al. [3]).

Consider the complex Burgers equation (1.1) with $\gamma = 0$. If $g(x, t)$ given by (1.2) is no longer
 a trace of a \mathbb{C}_+ -holomorphic function, then the relation between u and ρ in (1.3) does not hold.
 We need to treat u and ρ independently. Take (1.2) into (1.1) and we obtain the following system
 on the real line

$$\begin{cases} \rho_t + (\rho u)_x = 0, & x \in \mathbb{R}, t > 0, \\ \partial_t u + u \partial_x u - \pi^2 \rho \partial_x \rho = 0. \end{cases} \quad (1.12)$$

Unfortunately, for the Cauchy problem, the above system is ill-posed as described below. We
 introduce the following system of conservation law with general constant $\alpha \in \mathbb{R}$

$$\begin{cases} \partial_t \rho + \partial_x (\rho u) = 0, & x \in \mathbb{R}, t > 0, \\ \partial_t u + \partial_x \left(\frac{u^2 + \alpha \rho^2}{2} \right) = 0. \end{cases} \quad (1.13)$$

18 Due to the relation between System (1.12) ((1.13) with $\alpha = -\pi^2$) and the complex Burgers equation
 19 (1.1), we call System (1.13) as the coupled Burgers system in this paper. System (1.13) can be
 20 rewritten as the following quasi-linear system

$$\frac{\partial}{\partial t} \begin{pmatrix} \rho \\ u \end{pmatrix} + A(\rho, u) \frac{\partial}{\partial x} \begin{pmatrix} \rho \\ u \end{pmatrix} = 0, \quad A(\rho, u) = \begin{pmatrix} u & \rho \\ \alpha \rho & u \end{pmatrix}. \quad (1.14)$$

21 The eigenvalues of A are given by $u \pm \sqrt{\alpha} \rho$, where $\sqrt{\alpha} = \sqrt{-1} \sqrt{|\alpha|} = i \sqrt{|\alpha|}$ for $\alpha < 0$. When
 22 $\alpha > 0$, this system is a hyperbolic system of conservation laws. When $\alpha < 0$ and $\rho \neq 0$, A has
 23 two imaginary eigenvalues and System (1.13) is elliptic and ill-posedness. For $\alpha \neq 0$, we set the
 24 eigenvalues as

$$f_+ := u + \sqrt{\alpha} \rho, \quad f_- := u - \sqrt{\alpha} \rho. \quad (1.15)$$

A linear transformation from the coupled Burgers system (1.13) shows that the eigenvalues satisfy the following decoupled Burgers equations

$$\partial_t f_+ + f_+ \partial_x f_+ = 0, \quad x \in \mathbb{R}, t > 0, \quad (1.16)$$

$$\partial_t f_- + f_- \partial_x f_- = 0, \quad x \in \mathbb{R}, t > 0. \quad (1.17)$$

1 When $\alpha < 0$, (1.17) is just the conjugate of equation (1.16). When $\alpha = -\pi^2$, (1.16) is exactly the
 2 complex Burgers equation (1.1) ($\gamma = 0$) on the real line.

For $\alpha > 1$, notice that f_{\pm} are Riemann invariants of the following system of isentropic gas dynamics:

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0, & x \in \mathbb{R}, t > 0, \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_x p = 0, \end{cases} \quad (1.18)$$

where the pressure p is given by

$$p(x, t) = \frac{\alpha}{3} \rho^3(x, t). \quad (1.19)$$

3 Formally, system (1.18) is a nonlinear transformation of the coupled Burgers system (1.13) and it
 4 expresses in physics the conservation of mass and the conservation of momentum, i.e. $m := \rho u$,
 5 for an isentropic gas system. In the quasi-linear form, we have

$$\frac{\partial}{\partial t} \begin{pmatrix} \rho \\ m \end{pmatrix} + B(\rho, m) \frac{\partial}{\partial x} \begin{pmatrix} \rho \\ m \end{pmatrix} = 0, \quad B(\rho, m) = \begin{pmatrix} 0 & 1 \\ -\frac{m^2}{\rho^2} + \alpha \rho^2 & \frac{2m}{\rho} \end{pmatrix}. \quad (1.20)$$

The functions $f_{\pm} = u \pm \sqrt{\alpha} \rho$ are also the eigenvalues of B . Notice that classical solutions of the coupled Burgers system (1.13) are also classical solutions to (1.18). However, when shock appears, shock speed for the coupled Burgers system (1.13) and (1.18) are different. For smooth solutions of System (1.18), the following conservation of energy holds

$$\partial_t E + \partial_x [u(E + p)] = 0, \quad (1.21)$$

where the total energy density is given by

$$E(x, t) = \frac{1}{2} \rho u^2 + \frac{p}{2} = \frac{1}{2} \rho u^2 + \frac{\alpha}{6} \rho^3. \quad (1.22)$$

6 Although there is no bi-Hamiltonian structure for Burgers equation, we use the decoupled
 7 Burgers equations (1.16) and (1.17) to construct a bi-Hamiltonian structure for the coupled Burgers
 8 system (1.13) (see Theorem 3.1). Moreover, we obtain infinite many conserved quantities for the
 9 coupled Burgers system (1.13). Bi-Hamiltonian structures for System (1.18) and p-system (which
 10 is the gas dynamics in Lagrangian coordinates; see (1.23) below) are also obtained. To discover
 11 a bi-Hamiltonian structure or a Lax pair for an integrable system is very important. Indeed,
 12 according to the fundamental theorem of Magri [18], any bi-Hamiltonian system associated with a
 13 nondegenerate Hamiltonian pair induces a hierarchy of commuting Hamiltonian flows and, provided
 14 enough of these Hamiltonians are functionally independent, is therefore completely integrable. For
 15 general discussions about Hamiltonian structures for systems of hyperbolic conservation laws, one
 16 can refer to [22].

17 When $\alpha > 0$, we establish a kinetic formulation for the coupled Burgers system (1.13). Using
 18 the kinetic formulation, we define a class of entropy pairs to the coupled Burgers system (1.13).
 19 Notice that our definition of entropies corresponds to the counter part (in the sense as explained in
 20 Remark 4.2) of entropies used in [15] for System (1.18). In [15], Lions, Perthame and Tadmor proved
 21 the existence of global entropy solutions to (1.18) and the uniqueness is unknown. In contrast,
 22 we prove the existence and uniqueness of entropy solutions to the coupled Burgers system (1.13)
 23 (see Section 4.2). Moreover, we show that an entropy solution to the coupled Burgers system
 24 (1.13) corresponds to an entropy solution to the decoupled Burgers equations (1.16) and (1.17)
 25 (see Proposition 4.3). For more details on relations of entropy solutions and weak solutions to
 26 kinetic equations, one can refer to [24].

We also derive the Lagrangian dynamics (see (5.11)) for the coupled Burgers system (1.13), which resembles the gas dynamics in Lagrangian variables, or p-system [27]

$$\begin{cases} \partial_t \tau - \partial_\xi V = 0, \\ \partial_t V + \partial_\xi p = 0, \end{cases} \quad (1.23)$$

where $\tau(\xi, t) = 1/\rho(X(\xi, t), t) = X_\xi(\xi, t)$ stands for the specific volume and ξ is the Lagrangian labels. $X(\xi, t)$ is the flow map according to velocity field $u(X(\xi, t), t)$ (see (5.5)). V is the velocity in Lagrangian variable $V(\xi, t) := u(X(\xi, t), t)$ and $p(\tau) = \alpha/(3\tau^3)$ is the pressure given by (1.19) (see more details in Section 5.1). The Lagrangian dynamics of the coupled Burgers system (1.13) naturally leads to a spring-mass system (Fermi-Pasta-Ulam-Tsingou model) such that each mass evolves by the elastic force between adjacent mass that are reciprocal proportion to the cubic of distances between the mass and the adjacent masses (see (5.15)). Instead of the nearest-neighbor interaction, if the mass interacts with all the other masses with the same manner, we obtain the Calogero-Moser model with different coefficients. As it is known, the Calogero-Moser model is an integrable systems with a Lax-pair; see [21]. An interesting fact is that the continuum limit of the Calogero-Moser model gives the same Lagrangian dynamics of the coupled Burgers system (1.13); see [19].

The rest of this paper is organized as follows. In Section 2, we prove the global existence and uniqueness of real analytical solutions to complex Burgers equation (1.1) and the Dyson equation (1.4) ($\gamma \geq 0$) with strictly positive initial datum $\rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$, $s > 1/2$. We also obtain the pointwise convergence to the steady state for analytical solutions. Some explicit solutions are constructed by using Wigner's semicircle law, which converge to the steady state exponentially when $\gamma > 0$. The same explicit solution is given in Appendix B by the Stieltjes transform of Wigner's semicircle law μ_1 . Moreover, we prove the global existence of weak solutions in $H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$ for nonnegative initial data. In Section 3, we construct bi-Hamiltonian structures for the coupled Burgers system (1.13), isentropic gas system (1.18) and p-system (1.23). In Section 4, we establish kinetic formulation for the coupled Burgers system (1.13) with $\alpha > 0$. The existence and uniqueness of entropy solutions to (1.13) are also proved. In Section 5, we study the Lagrangian dynamics for the coupled Burgers system (1.13) and explore the connection between the Lagrangian dynamics system and a Fermi-Pasta-Ulam-Tsingou model with nearest-neighbor interactions. In Appendix A, we give the proof of Theorem 2.1.

2 Complex Burgers equation and the Dyson Brownian motion

Recall the Dyson Brownian motion (1.6). The eigenvalues λ_j given by (1.6) evolve by Brownian motion, combined with a deterministic repulsion force that repels nearby eigenvalues from each other with a strength inversely proportional to the separation. Notice that System (1.6) can also be rewritten as

$$d\lambda_j(t) = \frac{1}{\sqrt{N}} dB_j(t) - \partial_{\lambda_j} \Phi(\lambda_1(t), \dots, \lambda_N(t)), \quad 1 \leq j \leq N, \quad (2.1)$$

with potential function given by

$$\Phi(\lambda_1(t), \dots, \lambda_N(t)) := \frac{\gamma}{2} \sum_{j=1}^N \lambda_j^2(t) - \frac{1}{2N} \sum_{j=1}^N \sum_{k \neq j} \log |\lambda_j(t) - \lambda_k(t)|. \quad (2.2)$$

It can be proved that the eigenvalues almost surely not collide with each other (see [25, 16, 14]) and the solutions to System (1.6) exist globally. Hence, the empirical measure

$$\mu^N(t) := \frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j(t)} \quad (2.3)$$

is well defined for $t \in [0, \infty)$. One can prove that $\mu^N(t)$ converges to some probability measure satisfying the Dyson equation (1.4) ([25, 5, 2]).

1 Next, we derive the complex Burgers equation (1.1) from the Dyson equation (1.4). For $f, g \in$
 2 $L^p(\mathbb{R})$ ($p > 1$), the Hilbert transform has the following properties (see e.g. [23]):

$$H(Hf) = -f, \quad \partial_x(Hf) = H\partial_x f,$$

3 and

$$H(fHg + gHf) = HfHg - fg.$$

4 Applying the Hilbert transform to the Dyson equation (1.4) yields

$$\partial_t(H\rho) + \pi H\rho H\partial_x\rho - \pi\rho\partial_x\rho - \gamma\partial_x H(\rho x) = 0.$$

Moreover, for any function $g : \mathbb{R} \rightarrow \mathbb{R}$, we have

$$\begin{aligned} H(xg(x)) &= \frac{1}{\pi} \text{p. v.} \int_{\mathbb{R}} \frac{yg(y)}{x-y} dy = \frac{1}{\pi} \text{p. v.} \int_{\mathbb{R}} \frac{(y-x)g(y)}{x-y} dy + \frac{1}{\pi} \text{p. v.} \int_{\mathbb{R}} \frac{xg(y)}{x-y} dy \\ &= xHg(x) - \frac{1}{\pi} \int_{\mathbb{R}} g(x) dx, \end{aligned} \quad (2.4)$$

which implies

$$H(\rho x) = -\frac{\|\rho(t)\|_{L^1}}{\pi} + \frac{1}{\pi} ux. \quad (2.5)$$

Combining the above two equations, we have

$$\partial_t u + u\partial_x u - \pi^2 \rho\partial_x\rho - \gamma\partial_x(ux) = 0. \quad (2.6)$$

5 Set

$$f = u - i\pi\rho, \quad u = \pi H\rho.$$

6 Hence, f gives the trace of an analytic function in the upper half plane. Combining (1.4) and (2.6)
 7 yields

$$\partial_t f + f\partial_x f - \gamma\partial_x(fx) = 0, \quad x \in \mathbb{R}, \quad t > 0.$$

This corresponds to the following complex equation in \mathbb{C}_+ :

$$\partial_t f + f\partial_z f - \gamma\partial_z(fz) = \partial_t f + f\partial_z f - \gamma z\partial_z f - \gamma f = 0, \quad t > 0. \quad (2.7)$$

8 By the linear transformation $g(z, t) = f(z, t) - \gamma z$, we have

$$\partial_t g + g\partial_z g - \gamma^2 z = \partial_t f + (f - \gamma z)(\partial_z f - \gamma) - \gamma^2 z = \partial_t f + f\partial_z f - \gamma z\partial_z f - \gamma f = 0,$$

9 which is the Burgers equation with force term $\gamma^2 z$ (1.1). Moreover, from the above computation
 10 we see that the Dyson equation (1.4) with $\gamma = 0$ is equivalent to the coupled Burgers system (1.13)
 11 with $\alpha = -\pi^2$ and $u = \pi H\rho$.

12 2.1 Analytical solutions to the Dyson equation (1.4), convergence to 13 steady state and finite time blow up

14 In this subsection, we prove the existence and uniqueness of positive analytical solutions to the
 15 Dyson equation (1.4) with $\gamma > 0$ and initial datum $0 < \rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ ($s > 1/2$) by
 16 proving the well-posedness results for complex Burgers equation (1.1). We also show the pointwise
 17 convergence to the steady state for analytical solutions.

Let $\rho_0(x) > 0$ and $\rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ with $s > 1/2$ be the initial datum for the Dyson equation (1.4). The initial datum ρ_0 can be extended to a \mathbb{C}_+ -holomorphic function by Hilbert transform (also called Stieltjes transform, Borel transform or Markov function) for positive measures:

$$f_0(z) := \frac{1}{\pi} \int_{\mathbb{R}} \frac{\rho_0(s)}{z-s} ds, \quad z = x + iy \in \mathbb{C}_+. \quad (2.8)$$

Let

$$g_0(z) := f_0(z) - \gamma z, \quad z = x + iy \in \mathbb{C}_+. \quad (2.9)$$

Then, g_0 is a \mathbb{C}_+ -holomorphic function. Consider the following Cauchy problem of the Burgers equation with force term $\gamma^2 z$ in \mathbb{C}_+ :

$$\begin{cases} [\partial_t g + g \partial_z g](z, t) = \gamma^2 z, \quad z = x + iy \in \mathbb{C}_+, \\ g(z, 0) = g_0(z). \end{cases} \quad (2.10)$$

First let us list some simple estimates for the Dyson equation (1.4).

Fact 1 (L^1 -conservation law): $\|\rho(t)\|_{L^1(\mathbb{R})} = \|\rho_0\|_{L^1(\mathbb{R})}$.

Fact 2 (Second moment estimate): Multiplying (1.4) by x^2 and taking integral yield

$$\frac{d}{dt} \int_{\mathbb{R}} x^2 \rho(x, t) dx = 2\pi \int_{\mathbb{R}} x \rho H \rho dx - 2\gamma \int_{\mathbb{R}} x^2 \rho(x, t) dx$$

Notice from (2.5), we have

$$\int_{\mathbb{R}} x \rho H \rho dx = \frac{1}{2\pi} \|\rho\|_{L^1}^2,$$

hence

$$\frac{d}{dt} \int_{\mathbb{R}} x^2 \rho(x, t) dx = \|\rho_0\|_{L^1}^2 - 2\gamma \int_{\mathbb{R}} x^2 \rho(x, t) dx,$$

which implies

$$m_2(t) = \frac{\|\rho_0\|_{L^1}^2}{2\gamma} - \frac{\|\rho_0\|_{L^1}^2 - 2\gamma m_2(0)}{2\gamma} e^{-2\gamma t}, \quad \forall t > 0.$$

Fact 3 (L^2 estimate): Multiplying (1.4) by ρ and integration by parts show that

$$\frac{d}{dt} \int_{\mathbb{R}} \rho^2 dx + 2 \int_{\mathbb{R}} \int_{\mathbb{R}} \rho(x, t) \frac{|\rho(x, t) - \rho(y, t)|^2}{|x - y|^2} dx dy = \gamma \int_{\mathbb{R}} \rho^2 dx;$$

see more details in the proof of Theorem 2.2.

Fact 4 ($\dot{H}^{\frac{1}{2}}$ estimate):

$$\frac{d}{dt} \|(-\Delta)^{1/4} \rho\|_{L^2}^2 + \pi \int_{\mathbb{R}} (\partial_x H \rho)^2 \rho dx + \pi \int_{\mathbb{R}} \rho (\partial_x \rho)^2 dx = 2\gamma \|(-\Delta)^{1/4} \rho\|_{L^2}^2;$$

see more details in the proof of Theorem 2.2.

Fact 5 (Entropy estimate): Taking the time derivative to $\int_{\mathbb{R}} \rho \log \rho dx$ and integration by parts show that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} \rho \log \rho dx &= \int_{\mathbb{R}} \partial_t \rho (\log \rho + 1) dx = \int_{\mathbb{R}} -(\rho H \rho + \gamma x \rho)_x (\log \rho + 1) dx \\ &= \int_{\mathbb{R}} (H \rho - \gamma x) \rho_x dx = -\|(-\Delta)^{1/4} \rho\|_{L^2}^2 \end{aligned}$$

Fact 6 (Energy dissipation): Since the Dyson equation is a W_2 -gradient flow with respect to the energy (1.9), we have the following energy dissipation property

$$\begin{aligned} \frac{d}{dt} E(\rho) &= \int_{\mathbb{R}} \frac{\delta E}{\delta \rho} \cdot \partial_t \rho dx = - \int_{\mathbb{R}} \rho \left| \partial_x \left(\frac{\delta E}{\delta \rho} \right) \right|^2 dx \\ &= - \int_{\mathbb{R}} \rho(x, t) |\gamma x - \pi H \rho(x, t)|^2 dx. \end{aligned}$$

¹ Now we have the following theorem:

1 **Theorem 2.1.** Let $\gamma \geq 0$ and $0 < \rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ with $s > 1/2$. Then,

2 (i) The complex Burgers equation (2.10) has a unique $\overline{\mathbb{C}_+}$ -holomorphic solution $g(\cdot, t)$ for $t \in$
3 $(0, \infty)$, and $\frac{\partial^k}{\partial t^k} g(\cdot, t)$ is an analytical function of z on $\overline{\mathbb{C}_+}$ for any positive integer k and $t > 0$.

4 (ii) For any $t > 0$, the trace of $f(z, t) = g(z, t) + \gamma z$ on the real line gives a positive analytical
5 solution $\rho(x, t) > 0$ to the Dyson equation (1.4) with $\rho(x, 0) = \rho_0(x)$ and $\frac{\partial^k}{\partial t^k} \rho(x, t)$ is an analytical
6 function of $x \in \mathbb{R}$ for any positive integer k . The following estimates hold:

(a) The total mass $\|\rho(t)\|_{L^1}$ is conserved:

$$\|\rho(t)\|_{L^1} = \|\rho_0\|_{L^1}. \quad (2.11)$$

(b) If $x^2 \rho_0 \in L^1(\mathbb{R})$, then the second moment $m_2(t) := \int_{\mathbb{R}} x^2 \rho(x, t) dx$ satisfies

$$m_2(t) = \begin{cases} \frac{\|\rho_0\|_{L^1}^2}{2\gamma} - \frac{\|\rho_0\|_{L^1}^2 - 2\gamma m_2(0)}{2\gamma} e^{-2\gamma t}, & \gamma > 0, \\ m_2(0) + \|\rho_0\|_{L^1}^2 t, & \gamma = 0. \end{cases} \quad (2.12)$$

(c) The following energy dissipation holds:

$$\frac{d}{dt} E(\rho) = - \int_{\mathbb{R}} \rho(x, t) |\gamma x - \pi H \rho(x, t)|^2 dx, \quad (2.13)$$

7 with E defined by (1.9).

(d) If $\rho_0 \log \rho_0 \in L^1(\mathbb{R})$, then the entropy $\theta(t) := \int_{\mathbb{R}} \rho(x, t) \log \rho(x, t) dx$ satisfies

$$\theta(t) \leq \gamma \|\rho_0\|_{L^1} t + \theta(0). \quad (2.14)$$

8 (iii) For $\gamma > 0$, $g(z, t)$ converges to the steady state:

$$\lim_{t \rightarrow \infty} g(z, t) = -\sqrt{\gamma^2 z^2 - 2\gamma}, \quad \forall z \in \mathbb{C}_+,$$

and $\rho(x, t)$ converges to the steady state given by semicircle law:

$$\lim_{t \rightarrow \infty} \rho(x, t) = \rho_{\infty}(x) := \frac{\sqrt{(2\gamma - \gamma^2 x^2)_+}}{\pi}, \quad \forall x \in \mathbb{R}. \quad (2.15)$$

9 (iv) For $\gamma = 0$, the solution $g(z, t)$ and $\rho(x, t)$ converge to steady state after scaling in the
10 following sense:

$$e^t g\left(e^t z, \frac{e^{2t}-1}{2}\right) - z \rightarrow -\sqrt{z^2 - 2} \quad \text{as } t \rightarrow \infty.$$

11 and

$$e^t \rho\left(e^t x, \frac{e^{2t}-1}{2}\right) \rightarrow \frac{\sqrt{(2-x^2)_+}}{\pi} \quad \text{as } t \rightarrow \infty.$$

We remark that part (i) of Theorem 2.1 is derived directly by combining the solutions given by [4] and the space-time rescaling (1.7) as described below. Consider the following complex Burgers equation

$$\begin{cases} [\partial_{\tau} \tilde{g} + \tilde{g} \partial_w \tilde{g}](w, \tau) = 0, & w \in \mathbb{C}_+, \\ \tilde{g}(w, 0) = g_0(w) + \gamma w, \end{cases} \quad (2.16)$$

12 where g_0 is defined by (2.9). Castro and Córdoba [4] proved global existence and uniqueness
13 of \mathbb{C}_+ -holomorphic solution \tilde{g} to (2.16) by the method of characteristics. For $t > 0$, $\tilde{g}(\cdot, t)$ is
14 $\overline{\mathbb{C}_+}$ -holomorphic. Hence, from (1.7) we obtain a \mathbb{C}_+ -holomorphic solution g to (2.10) with initial
15 datum g_0 and for $t > 0$, $g(\cdot, t)$ is $\overline{\mathbb{C}_+}$ -holomorphic. This proves part (i) of Theorem 2.1. For part
16 (ii), let

$$f(z, t) := g(z, t) + \gamma z, \quad z \in \overline{\mathbb{C}_+}, \quad t > 0.$$

1 Then, f is a to \mathbb{C}_+ -holomorphic solution to (2.7) with initial datum f_0 given by (2.8) and for $t > 0$,
 2 $g(\cdot, t)$ is $\overline{\mathbb{C}_+}$ -holomorphic. Consider the trace of f on the real line and define

$$f(x, t) := u(x, t) - i\pi\rho(x, t), \quad x \in \mathbb{R}, \quad t > 0.$$

3 Then, we have $u = \pi H\rho$ and $\rho(x, t)$ is an analytical solution to the Dyson equation (1.4) with
 4 initial datum ρ_0 . This proves part (ii) of Theorem 2.1.

5 Since, the acceleration of characteristics for complex Burgers (2.10) is not zero, which is different
 6 with (2.16). This also brings some detailed information of solutions. Therefore, for completeness
 7 and to unveil those information, we provide a direct proof for Theorem 2.1 in Appendix A.

8 **Remark 2.1** (Finite time blow up). Note that condition $\rho_0 > 0$ is essential to Theorem 2.1.
 9 Castro and Córdoba [4, Theorem 4.4, Remark 4.5] proved that if $\rho_0 \geq 0$ and $\rho_0 \in H^2(\mathbb{R})$, then
 10 there exists a unique local solution $\rho \in C([0, T]; H^2(\mathbb{R})) \cap C^1([0, T]; H^1(\mathbb{R}))$ to (1.4) with $\gamma = 0$.
 11 Moreover, if $\rho_0(x_0) = 0 = \inf_{x \in \mathbb{R}} \rho_0(x)$ for some point $x_0 \in \mathbb{R}$, the solution blows up in finite time
 12 (see [4, Theorem 4.8, Remark 4.9]). Precisely, along the trajectories of characteristics $X(x_0, t)$
 13 starting from x_0 , we have

$$X(x_0, t) = H\rho_0(x_0)t + x_0,$$

14 and

$$\partial_x H\rho(X(x_0, t), t) \rightarrow -\infty \quad \text{as } t \rightarrow t^* := -\frac{1}{\partial_x H\rho_0(x_0)}.$$

15 Due to (1.7), there exists a unique local solution $\tilde{\rho}$ to (1.4) for $\gamma > 0$ given by

$$\tilde{\rho}(y, \tau) = e^{\gamma\tau} \rho\left(e^{\gamma\tau} y, \frac{e^{2\gamma\tau} - 1}{2\gamma}\right), \quad y \in \mathbb{R}, \quad \tau > 0.$$

16 Moreover, we have

$$\partial_y H\tilde{\rho}(y, \tau) = e^{2\gamma\tau} \partial_x H\rho\left(e^{\gamma\tau} y, \frac{e^{2\gamma\tau} - 1}{2\gamma}\right).$$

17 Let

$$t = \frac{e^{2\gamma\tau} - 1}{2\gamma}, \quad y = e^{-\gamma\tau} X(x_0, t) = e^{-\gamma\tau} \left[\frac{H\rho_0(x_0)(e^{2\gamma\tau} - 1)}{2\gamma} + x_0 \right],$$

18 and

$$\tau^* = \frac{1}{2\gamma} \log(1 + 2\gamma t^*).$$

Then, we have

$$\begin{aligned} \lim_{\tau \rightarrow \tau^*} \partial_y H\tilde{\rho}(y, \tau) &= \lim_{\tau \rightarrow \tau^*} e^{2\gamma\tau} \partial_x H\rho\left(e^{\gamma\tau} y, \frac{e^{2\gamma\tau} - 1}{2\gamma}\right) \\ &= e^{2\gamma\tau^*} \lim_{t \rightarrow t^*} \partial_x H\rho(X(x_0, t), t) = -\infty. \end{aligned}$$

19 Hence, the solution to (1.4) with $\gamma > 0$ also blows up in finite time.

20 2.2 Explicit solutions to the Dyson equation (1.4) from semicircle law 21 and exponential convergence to the steady state for $\gamma > 0$

22 In this subsection, we give some explicit solutions to the Dyson equation (1.4) by using Wigner's
 23 semicircle law (1.11). When $\gamma > 0$ the explicit solutions converge exponentially to steady state
 24 given by (2.15).

25 2.2.1 An explicit solution to the Dyson equation (1.4) with $\gamma = 0$

26 For $\gamma = 0$, notice that $\sqrt{N}A(t)/\sqrt{t}$ is a Wigner matrix (Hermitian matrix with i.i.d entries which
 27 have mean zero and variance one), where $A(t)$ is defined by (1.5) with $A(0) = 0$. Let $\{\lambda_j(t)\}_{j=1}^N$ be
 28 the eigenvalues of matrix $A(t)$. Hence, as N goes to infinity, the empirical measure $\frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j(t)/\sqrt{t}}$
 29 almost surely converges to Wigner's semicircle law $\mu_1(x)$ given by (1.11) weakly in probability
 30 measure space (see [30] or [29, Theorem 2.4.2]). On the other hand, the empirical measure $\mu^N(t) =$
 31 $\frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j(t)}(x)$ almost surely converges to a measure solution $\rho(x, t)$ of the Dyson equation (1.4)
 32 with $\gamma = 0$ [25]. We can obtain the relation between $\rho(x, t)$ and $\mu_1(x)$ by the following lemma.

1 **Lemma 2.1.** For any constant $a > 0$, if we have the following narrow convergences in probability
 2 measure space $\mathcal{P}(\mathbb{R})$:

$$\tilde{\nu}^N(x) := \frac{1}{N} \sum_{j=1}^N \delta_{x_j/a}(x) \rightarrow \tilde{\nu}(x) \quad \text{and} \quad \nu^N(x) := \frac{1}{N} \sum_{j=1}^N \delta_{x_j}(x) \rightarrow \nu(x)$$

for two probability measures $\tilde{\nu}, \nu$, then we have

$$\nu(x) = \frac{1}{a} \tilde{\nu}\left(\frac{x}{a}\right). \quad (2.17)$$

Proof. For any test function $\varphi \in C_b(\mathbb{R})$, we have

$$\begin{aligned} \int_{\mathbb{R}} \varphi(x) d\tilde{\nu}(x) &= \lim_{N \rightarrow \infty} \int_{\mathbb{R}} \varphi(x) d\tilde{\nu}^N(x) = \frac{1}{N} \sum_{j=1}^N \varphi(x_j/a) \\ &= \lim_{N \rightarrow \infty} \int_{\mathbb{R}} \varphi(y/a) d\nu^N(y) = a \lim_{N \rightarrow \infty} \int_{\mathbb{R}} \varphi(x) d\nu^N(ax) \\ &= \int_{\mathbb{R}} \varphi(y/a) d\nu(y) = a \int_{\mathbb{R}} \varphi(x) d\nu(ax). \end{aligned}$$

3 Hence, $a\nu(ax) = \tilde{\nu}(x)$, which implies (2.17). □

4

From Lemma 2.1, we choose ρ as the rescaling of ρ_1 defined in (1.11)

$$\rho(x, t) = \frac{1}{\sqrt{t}} \rho_1\left(\frac{x}{\sqrt{t}}\right) = \frac{\sqrt{(4t - x^2)_+}}{2\pi t}, \quad (2.18)$$

where $\rho(x, t)$ is the limit of the empirical measure $\frac{1}{N} \sum_{j=1}^N \delta_{\lambda_j(t)}(x)$ for $\gamma = 0$. This implies $\rho(x, t)$ is a kind of self-similar rarefaction wave solution of the Dyson equation (1.4) with $\gamma = 0$. Next, we calculate $u(x, t)$ using the Hilbert transform of $\pi\rho(x, t)$ and then verify the obtained (ρ, u) satisfies (1.4) ($\gamma = 0$). For $x \in \mathbb{R} \setminus [-2\sqrt{t}, 2\sqrt{t}]$, by changing of variable with $y = 2\sqrt{t} \sin \theta$, we have

$$\begin{aligned} (\pi H\rho)(x, t) &= \frac{1}{2t\pi} \int_{-2\sqrt{t}}^{2\sqrt{t}} \frac{\sqrt{4t - y^2}}{x - y} dy \\ &= \frac{1}{2t\pi} \int_{-\pi/2}^{\pi/2} (x + 2\sqrt{t} \sin \theta) d\theta + \frac{4t - x^2}{2t\pi} \int_{-\pi/2}^{\pi/2} \frac{1}{x - 2\sqrt{t} \sin \theta} d\theta \\ &= \frac{x}{2t} + \frac{4t - x^2}{2t\pi} \int_{-\pi/2}^{\pi/2} \frac{1}{x - 2\sqrt{t} \sin \theta} d\theta \\ &= \frac{x}{2t} + \frac{4t - x^2}{2t\pi} \frac{2}{\sqrt{x^2 - 4t}} \left[\arctan\left(\frac{x - 2\sqrt{t}}{\sqrt{x^2 - 4t}}\right) + \arctan\left(\frac{x + 2\sqrt{t}}{\sqrt{x^2 - 4t}}\right) \right]. \end{aligned} \quad (2.19)$$

Using the fact

$$\arctan x + \arctan y = \begin{cases} \frac{\pi}{2} & \text{for } x \cdot y = 1, x, y > 0, \\ -\frac{\pi}{2} & \text{for } x \cdot y = 1, x, y < 0, \end{cases}$$

we obtain

$$(\pi H\rho)(x, t) = \begin{cases} \frac{x}{2t} + \frac{\sqrt{x^2 - 4t}}{2t}, & x < -2\sqrt{t}, \\ \frac{x}{2t} - \frac{\sqrt{x^2 - 4t}}{2t}, & x > 2\sqrt{t}. \end{cases} \quad (2.20)$$

For $x \in [-2\sqrt{t}, 2\sqrt{t}]$, we have

$$\begin{aligned} (\pi H\rho)(x, t) &= \frac{1}{2t\pi} \text{p. v.} \int_{-2\sqrt{t}}^{2\sqrt{t}} \frac{\sqrt{4-y^2}}{x-y} dy \\ &= \frac{1}{2t\pi} \lim_{\epsilon \rightarrow 0} \left(\int_{-2\sqrt{t}}^{x-\epsilon} + \int_{x+\epsilon}^{2\sqrt{t}} \right) \frac{\sqrt{4-y^2}}{x-y} dy. \end{aligned}$$

Then, using similar calculation as (2.19) we have $(\pi H\rho)(x, t) = \frac{x}{2t}$, $x \in [-2\sqrt{t}, 2\sqrt{t}]$. Therefore we have

$$u(x, t) = (\pi H\rho)(x, t) = \begin{cases} \frac{x + \sqrt{x^2 - 4t}}{2t}, & x < -2\sqrt{t}, \\ \frac{x}{2t}, & x \in [-2\sqrt{t}, 2\sqrt{t}], \\ \frac{x - \sqrt{x^2 - 4t}}{2t}, & x > 2\sqrt{t}, \end{cases} \quad (2.21)$$

1 and (ρ, u) satisfies (1.4) ($\gamma = 0$) with initial datum

$$\rho(x, 0) = \delta(0), \quad u(x, 0) = (\pi H\rho)(x, 0) = \text{p.v.} \frac{1}{x}. \quad (2.22)$$

2 Notice that the above self-similar solution (ρ, u) corresponds to the self-similar solution to complex
3 Burgers equation given in [20, Section 1.2].

4 In Appendix B we will give the same explicit solution by the Stieltjes transform of Wigner's
5 semicircle law μ_1 (see (B.8)).

6 **Remark 2.2** (Connection with Barenblatt solutions to porous media equation). Consider the
7 following one dimensional porous media equation:

$$\partial_t h = \frac{\pi^2}{3} \partial_{xx}(h^3), \quad h|_{t=0} = \delta(0).$$

8 It has a self-similar solution called Barenblatt solution (see [26, Page 104]) given by

$$h(x, t) = \frac{\sqrt{(4\sqrt{t} - x^2)_+}}{2\pi\sqrt{t}} = \frac{1}{t^{1/4}} \cdot \frac{\sqrt{(4 - (\frac{x}{t^{1/4}})^2)_+}}{2\pi}.$$

9 Notice that

$$\rho(x, t) = h(x, t^2) = \frac{\sqrt{(4t - x^2)_+}}{2\pi t}$$

10 is exactly the explicit solution (2.18) to the Dyson equation (1.4) with $\gamma = 0$.

11 2.2.2 An explicit solution to the Dyson equation (1.4) with $\gamma > 0$ and exponential 12 convergence to the steady state

13 When $\gamma > 0$, we first show that (2.18) with $t = \frac{1}{2\gamma}$ gives a steady state of (1.4) with $\gamma > 0$.
14 Actually, we have

$$\rho\left(x, \frac{1}{2\gamma}\right) = \frac{\sqrt{(2\gamma - \gamma^2 x^2)_+}}{\pi},$$

and

$$u\left(x, \frac{1}{2\gamma}\right) = \pi H\rho\left(x, \frac{1}{2\gamma}\right) = \begin{cases} \gamma x + \sqrt{\gamma^2 x^2 - 2\gamma}, & x < -\sqrt{2}, \\ \gamma x, & x \in [-\sqrt{2}, \sqrt{2}], \\ \gamma x - \sqrt{\gamma^2 x^2 - 2\gamma}, & x > \sqrt{2}. \end{cases}$$

Define

$$\rho_\infty(x) := \rho\left(x, \frac{1}{2\gamma}\right), \quad u_\infty(x) = u\left(x, \frac{1}{2\gamma}\right), \quad (2.23)$$

1 and then

$$\rho_\infty(u_\infty - \gamma x) \equiv 0,$$

2 which implies that ρ_∞ is a steady state of the Dyson equation (1.4) when $\gamma > 0$. Due to the
3 convexity of the energy E in (1.9), the steady state is the minimizer and it is unique (see Remark
4 2.4).

Next, we construct an explicit solution which converges to ρ_∞ exponentially. Let $\sigma(t)$ be an unknown function and $\sigma(0) = \sigma_0 > 0$ and assume solution $\rho(x, t)$ to (1.4) with $\gamma > 0$ has the following form

$$\rho(x, t) = \frac{\sqrt{(2\sigma(t) - x^2)_+}}{\pi\sigma(t)}. \quad (2.24)$$

Correspondingly, we have

$$u(x, t) = \pi H \rho(x, t) = \begin{cases} \frac{x + \sqrt{x^2 - 2\sigma(t)}}{\sigma(t)}, & x < -\sqrt{2\sigma(t)}, \\ \frac{x}{\sigma(t)}, & |x| \leq \sqrt{2\sigma(t)}, \\ \frac{x - \sqrt{x^2 - 2\sigma(t)}}{\sigma(t)}, & x > \sqrt{2\sigma(t)}. \end{cases}$$

5 Obviously, (ρ, u) satisfies (1.4) when $|x| > \sqrt{2\sigma(t)}$. Next, we consider the case $|x| \leq \sqrt{2\sigma(t)}$ to
6 obtain a proper ordinary differential equation for $\sigma(t)$ such that (ρ, u) is a solution of (1.4). Direct
7 calculations show that

$$\partial_t \rho = -\frac{\sqrt{2\sigma - x^2}}{\pi\sigma^2} \dot{\sigma} + \frac{\dot{\sigma}}{\pi\sigma\sqrt{2\sigma - x^2}}, \quad \partial_x \rho = -\frac{x}{\pi\sigma\sqrt{2\sigma - x^2}},$$

8 and

$$\rho + x\partial_x \rho = 2\sigma \left(\frac{\sqrt{2\sigma - x^2}}{\pi\sigma^2} - \frac{1}{\pi\sigma\sqrt{2\sigma - x^2}} \right).$$

Take the above equalities into (1.4) and we obtain

$$\begin{aligned} \partial_t \rho + \partial_x [\rho(u - \gamma x)] &= \partial_t \rho + \left(\frac{1}{\sigma} - \gamma \right) (\rho + x\partial_x \rho) \\ &= (-\dot{\sigma} + 2 - 2\gamma\sigma) \left(\frac{\sqrt{2\sigma - x^2}}{\pi\sigma^2} - \frac{1}{\pi\sigma\sqrt{2\sigma - x^2}} \right) = 0, \quad |x| \leq \sqrt{2\sigma(t)}. \end{aligned}$$

9 Hence, we have

$$\dot{\sigma}(t) = 2 - 2\gamma\sigma, \quad \sigma(0) = \sigma_0 > 0,$$

10 which implies

$$\sigma(t) = \frac{1}{\gamma} - \frac{1 - \gamma\sigma_0}{\gamma} e^{-2\gamma t} > 0.$$

Hence, for any $\sigma_0 > 0$, an explicit solution to (1.4) is given by

$$\rho(x, t) = \frac{\sqrt{(2\gamma[1 - (1 - \gamma\sigma_0)e^{-2\gamma t}] - \gamma^2 x^2)_+}}{\pi[1 - (1 - \gamma\sigma_0)e^{-2\gamma t}]}. \quad (2.25)$$

11 This solution tends to ρ_∞ (defined by (2.23)) exponentially as $t \rightarrow \infty$.

2.3 Global weak solutions of the Dyson equation (1.4)

In Theorem 2.1, we proved global existence and uniqueness of a positive analytical solution to (1.4) with a strictly positive initial datum $\rho_0 > 0$ and $\rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ with $s > 1/2$. If $\rho_0 \geq 0$ and $\rho_0(x_0) = 0$ for some $x_0 \in \mathbb{R}$, the solution to (1.4) blows up in finite time (see Remark 2.1) in the sense that $\partial_x H\rho$ goes to $-\infty$. Consequently, there is also a finite time blow up in the space $H^s(\mathbb{R})$ for $s > 3/2$. Next, we show global existence of weak solution in $\dot{H}^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$. Note that we have interpolation inequality

$$\|\rho\|_{L^2} \leq 3\|\rho\|_{L^1}^{1/2}\|\rho\|_{\dot{H}^{1/2}}^{1/2}.$$

Hence $\rho \in \dot{H}^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$ is equivalent to $\rho \in H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$. Let us define the weak solutions:

Definition 2.1. For $T > 0$, $\rho_0 \in H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$ and $\rho_0 \geq 0$, a nonnegative function $\rho \in L^\infty(0, T; H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})) \cap W^{1, \infty}(0, T; H^{-m}(\mathbb{R}))$ for some $m > 0$ is said to be a weak solution of the Dyson equation (1.4) if

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}} \partial_t \phi(x, t) \rho(x, t) \, dx \, dt + \int_{\mathbb{R}} \phi(x, 0) \rho_0(x) \, dx \\ &= -\frac{1}{2} \int_0^T \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\partial_x \phi(x, t) - \partial_x \phi(y, t)}{x - y} \rho(x, t) \rho(y, t) \, dx \, dy \, dt \\ & \quad + \gamma \int_0^T \int_{\mathbb{R}} x \partial_x \phi(x, t) \rho(x, t) \, dx \, dt, \end{aligned} \quad (2.26)$$

holds for any test function $\phi \in C_c^\infty(\mathbb{R} \times [0, T))$.

Theorem 2.2. Assume $0 \leq \rho_0 \in H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})$ and $m_2(0) := \int_{\mathbb{R}} x^2 \rho_0(x) \, dx < \infty$. Then, there exists a unique global nonnegative weak solution to the Dyson equation (1.4) satisfying

$$\rho \in L^\infty(0, T; H^{1/2}(\mathbb{R}) \cap L^1(\mathbb{R})) \cap W^{1, \infty}(0, T; H^{-3}(\mathbb{R}))$$

for any time $T > 0$. Moreover, we have the following estimates

(a)

$$\|\rho(t)\|_{H^{1/2}} \leq e^{\gamma t} \|\rho_0\|_{H^{1/2}}, \quad t > 0, \quad (2.27)$$

(b) The mass $\|\rho(t)\|_{L^1}$ is conserved:

$$\|\rho(t)\|_{L^1} = \|\rho_0\|_{L^1}. \quad (2.28)$$

(c) For a.e. $t > 0$, the second moment $m_2(t) := \int_{\mathbb{R}} x^2 \rho(x, t) \, dx$ satisfies

$$m_2(t) \leq \begin{cases} \frac{\|\rho_0\|_{L^1}^2}{2\gamma} - \frac{\|\rho_0\|_{L^1}^2 - 2\gamma m_2(0)}{2\gamma} e^{-2\gamma t}, & \gamma > 0, \\ m_2(0) + \|\rho_0\|_{L^1}^2 t, & \gamma = 0. \end{cases} \quad (2.29)$$

(d) The following energy dissipation holds:

$$E(\rho(\cdot, t)) + \int_0^t \int_{\mathbb{R}} \rho(x, s) |\gamma x - \pi H\rho(x, s)|^2 \, dx \, ds \leq E(\rho_0) \quad \text{for any } t > 0, \quad (2.30)$$

with E defined by (1.9).

(e) If $\rho_0 \log \rho_0 \in L^1(\mathbb{R})$, then the entropy $\theta(t) := \int_{\mathbb{R}} \rho(x, t) \log \rho(x, t) \, dx$ satisfies

$$\theta(t) \leq \gamma \|\rho_0\|_{L^1} t + \theta(0), \quad t > 0. \quad (2.31)$$

1 *Proof.* Let $\varphi_\epsilon > 0$ ($\epsilon > 0$) be the standard Friedrichs mollifier. Set

$$\rho_0^\epsilon = \rho_0 * \varphi_\epsilon.$$

Then, for nontrivial initial datum ρ_0 , we have $\rho_0^\epsilon(x) > 0$ for $x \in \mathbb{R}$ and $\rho_0^\epsilon \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ ($s > 1/2$). Moreover, from Young's inequality for convolution, we have

$$\|\rho_0^\epsilon\|_{L^2} \leq \|\rho_0\|_{L^2}, \quad \|\rho_0^\epsilon\|_{\dot{H}^{1/2}} \leq \|\rho_0\|_{\dot{H}^{1/2}}, \quad \|\rho_0^\epsilon\|_{L^1} = \|\rho_0\|_{L^1}. \quad (2.32)$$

By Theorem 2.1, we have a global positive analytical solution ρ^ϵ to (1.4) with initial data ρ_0^ϵ :

$$\partial_t \rho^\epsilon + \partial_x [\rho^\epsilon (\pi H \rho^\epsilon - \gamma x)] = 0. \quad (2.33)$$

2 **Step 1.** Uniform estimates for ρ^ϵ .

First, multiplying (2.33) by ρ^ϵ and integration by parts show that

$$\frac{d}{dt} \int_{\mathbb{R}} \frac{(\rho^\epsilon)^2}{2} dx + \frac{\pi}{2} \int_{\mathbb{R}} (\rho^\epsilon)^2 \partial_x H \rho^\epsilon dx - \frac{\gamma}{2} \int_{\mathbb{R}} (\rho^\epsilon)^2 dx = 0.$$

Since the second term on the left hand side is

$$\begin{aligned} \frac{\pi}{2} \int_{\mathbb{R}} (\rho^\epsilon)^2 \partial_x H \rho^\epsilon dx &= \int_{\mathbb{R}} \int_{\mathbb{R}} (\rho^\epsilon(x, t))^2 \frac{\rho^\epsilon(x, t) - \rho^\epsilon(y, t)}{|x - y|^2} dy dx \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} \rho^\epsilon(x) \frac{|\rho^\epsilon(x, t) - \rho^\epsilon(y, t)|^2}{|x - y|^2} dx dy, \end{aligned}$$

3 we obtain

$$\frac{d}{dt} \int_{\mathbb{R}} (\rho^\epsilon)^2 dx + 2 \int_{\mathbb{R}} \int_{\mathbb{R}} \rho^\epsilon(x, t) \frac{|\rho^\epsilon(x, t) - \rho^\epsilon(y, t)|^2}{|x - y|^2} dx dy = \gamma \int_{\mathbb{R}} (\rho^\epsilon)^2 dx. \quad (2.34)$$

Grönwall's inequality and (2.32) imply

$$\|\rho^\epsilon(t)\|_{L^2}^2 \leq e^{\gamma t} \|\rho_0^\epsilon\|_{L^2}^2 \leq e^{\gamma t} \|\rho_0\|_{L^2}^2, \quad t > 0. \quad (2.35)$$

Second, multiplying (2.33) by $H \rho_x^\epsilon$ gives the following estimate:

$$\frac{1}{2} \frac{d}{dt} \|(-\Delta)^{1/4} \rho^\epsilon\|_{L^2}^2 + \pi \int_{\mathbb{R}^2} (\partial_x H \rho^\epsilon)^2 \rho^\epsilon dx + \pi \int_{\mathbb{R}} \partial_x \rho^\epsilon \partial_x H \rho^\epsilon H \rho^\epsilon dx - \gamma \int_{\mathbb{R}} \partial_x H \rho^\epsilon \partial_x (x \rho^\epsilon) = 0. \quad (2.36)$$

On the one hand, we have

$$\pi \int_{\mathbb{R}} \partial_x \rho^\epsilon \partial_x H \rho^\epsilon H \rho^\epsilon dx = -\pi \int_{\mathbb{R}} H (\partial_x \rho^\epsilon \partial_x H \rho^\epsilon) \rho^\epsilon dx = -\frac{\pi}{2} \int_{\mathbb{R}} [(\partial_x H \rho^\epsilon)^2 - (\partial_x \rho^\epsilon)^2] \rho^\epsilon dx. \quad (2.37)$$

On the other hand, we estimate the last term in (2.36) as below. Due to (2.4), we derive

$$\begin{aligned} \gamma \int_{\mathbb{R}} \partial_x H \rho^\epsilon \partial_x (x \rho^\epsilon) &= -\gamma \int_{\mathbb{R}} \partial_x \rho^\epsilon \partial_x H (x \rho^\epsilon) = -\gamma \int_{\mathbb{R}} \partial_x \rho^\epsilon \partial_x (x H \rho^\epsilon) dx \\ &= -\gamma \int_{\mathbb{R}} \partial_x \rho^\epsilon H \rho^\epsilon dx - \gamma \int_{\mathbb{R}} x \partial_x \rho^\epsilon \partial_x H \rho^\epsilon dx \\ &= \gamma \|(-\Delta)^{1/4} \rho^\epsilon\|_{L^2}^2 - \gamma \int_{\mathbb{R}} x \partial_x \rho^\epsilon \partial_x H \rho^\epsilon dx. \end{aligned}$$

4 Use (2.4) again and we have

$$-\gamma \int_{\mathbb{R}} x \partial_x \rho^\epsilon \partial_x H \rho^\epsilon dx = \gamma \int_{\mathbb{R}} H (x \partial_x \rho^\epsilon) \partial_x \rho^\epsilon dx = \gamma \int_{\mathbb{R}} x \partial_x H \rho^\epsilon \partial_x \rho^\epsilon dx.$$

This implies $\gamma \int_{\mathbb{R}} x \partial_x \rho^\epsilon \partial_x H \rho^\epsilon dx = 0$ and hence

$$\gamma \int_{\mathbb{R}} \partial_x H \rho^\epsilon \partial_x (x \rho^\epsilon) = \gamma \|(-\Delta)^{1/4} \rho^\epsilon\|_{L^2}^2. \quad (2.38)$$

1 Combining (2.36), (2.37) and (2.38) shows

$$\frac{d}{dt} \|(-\Delta)^{1/4} \rho^\epsilon\|_{L^2}^2 + \pi \int_{\mathbb{R}} (\partial_x H \rho^\epsilon)^2 \rho^\epsilon dx + \pi \int_{\mathbb{R}} \rho^\epsilon (\partial_x \rho^\epsilon)^2 dx = 2\gamma \|(-\Delta)^{1/4} \rho^\epsilon\|_{L^2}^2. \quad (2.39)$$

Grönwall's inequality and (2.32) imply

$$\|\rho^\epsilon(t)\|_{H^{1/2}}^2 \leq e^{2\gamma t} \|\rho_0\|_{H^{1/2}}^2, \quad t > 0. \quad (2.40)$$

Inequalities (2.35) and (2.40) yield

$$\|\rho^\epsilon(t)\|_{H^{1/2}}^2 \leq e^{2\gamma t} \|\rho_0\|_{H^{1/2}}^2, \quad t > 0. \quad (2.41)$$

2 and hence we have

$$\rho^\epsilon \in L^\infty(0, T; H^{1/2}(\mathbb{R})) \quad \text{for any } T > 0.$$

Third, for time regularity, the following estimate holds for any $\phi \in C_c^\infty(\mathbb{R})$

$$\begin{aligned} & \int_{\mathbb{R}} \phi(x) \partial_t \rho^\epsilon(x, t) dx \\ &= -\frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\partial_x \phi(x) - \partial_x \phi(y)}{x - y} \rho^\epsilon(x, t) \rho^\epsilon(y, t) dx dy + \gamma \int_{\mathbb{R}} x \phi_x(x, t) \rho^\epsilon(x, t) dx \\ &\leq C(\|\partial_{xx} \phi\|_{L^\infty} + (m_2(\rho^\epsilon) + \|\rho^\epsilon\|_{L^1}) \|\partial_x \phi\|_{L^\infty}) \leq C\|\phi\|_{H^3}, \end{aligned}$$

and hence

$$\|\partial_t \rho^\epsilon\|_{L^\infty(0, \infty; H^{-3}(\mathbb{R}))} \leq C, \quad \partial_t \rho^\epsilon \in L^\infty(0, \infty; H^{-3}(\mathbb{R})). \quad (2.42)$$

3 **Step 2.** Take limits for ρ^ϵ as ϵ goes to 0.

4 First, from uniform estimates (2.41) and (2.42) in Step 1, there exist $\rho \in L^\infty(0, T; H^{1/2}(\mathbb{R})) \cap$
5 $W^{1, \infty}(0, T; H^{-3}(\mathbb{R}))$ and a subsequence of $\{\rho^\epsilon\}_{\epsilon > 0}$ (still denoted as $\{\rho^\epsilon\}_{\epsilon > 0}$) such that

$$\rho^\epsilon \xrightarrow{*} \rho \quad \text{in } L^\infty(0, T; H^{1/2}(\mathbb{R})) \quad \text{as } \epsilon \rightarrow 0,$$

6 and

$$\partial_t \rho^\epsilon \xrightarrow{*} \partial_t \rho \quad \text{in } L^\infty(0, T; H^{-3}(\mathbb{R})) \quad \text{as } \epsilon \rightarrow 0.$$

7 Hence, we have (2.27).

8 Second, from (2.41) and (2.42), by Lions-Aubin Lemma, we also know

$$\rho^\epsilon \rightarrow \rho \quad \text{in } L^\infty(0, T; L_{loc}^2(\mathbb{R})) \quad \text{as } \epsilon \rightarrow 0,$$

9 and consequently

$$\rho^\epsilon \rightarrow \rho \quad \text{in } L^\infty(0, T; L_{loc}^1(\mathbb{R})) \quad \text{as } \epsilon \rightarrow 0. \quad (2.43)$$

10 Due to $\|\rho^\epsilon(t)\|_{L^1} \equiv \|\rho_0\|_{L^1}$, we have

$$\|\rho(t)\|_{L^1} = \lim_{R \rightarrow +\infty} \int_{-R}^R \rho(x, t) dx = \lim_{R \rightarrow +\infty} \lim_{\epsilon \rightarrow 0} \int_{-R}^R \rho^\epsilon(x, t) dx \equiv \|\rho_0\|_{L^1},$$

11 where the last step we used the uniform bound of second momentum for ρ^ϵ (2.12). Hence,

$$\rho \in L^\infty(0, T; L^1(\mathbb{R})),$$

and (2.28) holds. For any test function $\phi \in C_c^\infty(\mathbb{R} \times [0, T])$, by (2.32) we have

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}} \partial_t \phi(x, t) \rho^\epsilon(x, t) \, dx \, dt + \int_{\mathbb{R}} \phi(x, 0) \rho_0^\epsilon(x) \, dx \\ &= -\frac{1}{2} \int_0^T \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\partial_x \phi(x, t) - \partial_x \phi(y, t)}{x - y} \rho^\epsilon(x, t) \rho^\epsilon(y, t) \, dx \, dy \, dt \\ & \quad + \gamma \int_0^T \int_{\mathbb{R}} x \partial_x \phi(x, t) \rho^\epsilon(x, t) \, dx \, dt, \end{aligned} \quad (2.44)$$

1 By the strong convergence of ρ^ϵ in (2.43), we can take the limit as $\epsilon \rightarrow 0$ in (2.44) and conclude
2 that ρ satisfies (2.26). Hence, ρ is a global weak solution to (1.4).

3 **Step 3.** Consequent estimates for $H\rho$. First, from (2.41) and

$$\|(-\Delta)^{\frac{1}{4}} \rho^\epsilon\|_{L^2}^2 = \int_{\mathbb{R}} (H\rho^\epsilon) H \partial_x (H\rho^\epsilon) \, dx = \|(-\Delta)^{\frac{1}{4}} (H\rho^\epsilon)\|_{L^2}^2, \quad (2.45)$$

4 we have uniform estimates

$$\|H\rho^\epsilon\|_{L^\infty(0, T; H^{\frac{1}{2}}(\mathbb{R}))} \leq C \text{ for any } T > 0. \quad (2.46)$$

Second, from the equation for u^ϵ (2.6) with $u^\epsilon = \pi H\rho^\epsilon$, we have for any $\phi \in C_c^\infty(\mathbb{R})$

$$\begin{aligned} & \int_{\mathbb{R}} \phi(x) \partial_t u^\epsilon(x, t) \, dx = \int_{\mathbb{R}} \partial_x \phi(x) \left[\frac{(u^\epsilon)^2}{2} - \frac{\pi^2}{2} (\rho^\epsilon)^2 - \gamma x u^\epsilon \right] \, dx \\ & \leq C \|\partial_x \phi\|_{L^\infty} \|\rho^\epsilon\|_{L^2}^2 + \int_{\mathbb{R}} \gamma H(x \partial_x \phi) \rho^\epsilon \, dx = C \|\partial_x \phi\|_{L^\infty} \|\rho^\epsilon\|_{L^2}^2 + \int_{\mathbb{R}} \gamma H(\partial_x \phi) x \rho^\epsilon \, dx \\ & \leq C [\|\partial_x \phi\|_{L^\infty} \|\rho^\epsilon\|_{L^2}^2 + (m_2(\rho^\epsilon)^{\frac{1}{2}} \|\rho^\epsilon\|_{L^2}^{\frac{1}{2}}) \|H \partial_x \phi\|_{L^4}] \leq C (\|\partial_x \phi\|_{L^\infty} + \|\partial_x \phi\|_{L^4}), \end{aligned}$$

and hence

$$\|\partial_t u^\epsilon\|_{L^\infty(0, \infty; H^{-2}(\mathbb{R}))} \leq C, \quad \partial_t u^\epsilon \in L^\infty(0, \infty; H^{-2}(\mathbb{R})). \quad (2.47)$$

Similar to ρ^ϵ , combining (2.46), (2.47) and Lions-Aubin Lemma, we also know for $u = \pi H\rho \in L^\infty(0, T; H^{1/2}(\mathbb{R})) \cap W^{1, \infty}(0, T; H^{-2}(\mathbb{R}))$,

$$u^\epsilon \xrightarrow{*} u \text{ in } L^\infty(0, T; H^{1/2}(\mathbb{R})) \text{ as } \epsilon \rightarrow 0, \quad (2.48)$$

$$\partial_t u^\epsilon \xrightarrow{*} \partial_t u \text{ in } L^\infty(0, T; H^{-2}(\mathbb{R})) \text{ as } \epsilon \rightarrow 0, \quad (2.49)$$

$$u^\epsilon \rightarrow u \text{ in } L^\infty(0, T; L_{loc}^2(\mathbb{R})) \text{ as } \epsilon \rightarrow 0. \quad (2.50)$$

5 Consequently, we have for a.e. $t \in [0, T]$

$$H\rho^\epsilon(\cdot, t) \rightarrow H\rho(\cdot, t) \text{ for a.e. } x \in \mathbb{R} \text{ as } \epsilon \rightarrow 0. \quad (2.51)$$

6 **Step 4.** The uniqueness of weak solutions is a direct result of the contraction property of
7 Wasserstein distance as stated in (2.58).

8 **Step 5.** We prove properties (2.29), (2.30), and (2.31) below.

Due to (2.12), we have

$$m_2^\epsilon(t) = \begin{cases} \frac{\|\rho_0\|_{L^1}^2}{2\gamma} - \frac{\|\rho_0\|_{L^1}^2 - 2\gamma m_2^\epsilon(0)}{2\gamma} e^{-2\gamma t}, & \gamma > 0, \\ m_2^\epsilon(0) + \|\rho_0\|_{L^1}^2 t, & \gamma = 0, \end{cases} \quad (2.52)$$

9 where

$$m_2^\epsilon(t) := \int_{\mathbb{R}} x^2 \rho^\epsilon(x, t) \, dx.$$

1 Due to strong convergence of ρ^ϵ to ρ in $L^\infty(0, T; L^1_{loc}(\mathbb{R}))$, for a.e. $t \in (0, T)$ we have

$$\rho^\epsilon(\cdot, t) \rightarrow \rho(\cdot, t) \text{ for a.e. } x \in \mathbb{R} \text{ as } \epsilon \rightarrow 0. \quad (2.53)$$

2 To take the limit in (2.52), first notice $m_2^\epsilon(0) \rightarrow m_2(0)$ by Young's convolution inequality. Second,
3 by Levi's lemma and Fatou's lemma, we have

$$\begin{aligned} m_2(t) &= \lim_{N \rightarrow +\infty} \int_{\mathbb{R}} (x^2)_N \rho \, dx \leq \lim_{N \rightarrow +\infty} \liminf_{\epsilon \rightarrow 0} \int_{\mathbb{R}} (x^2)_N \rho^\epsilon \, dx \\ &\leq \lim_{N \rightarrow +\infty} \liminf_{\epsilon \rightarrow 0} \int_{\mathbb{R}} x^2 \rho^\epsilon \, dx \leq \liminf_{\epsilon \rightarrow 0} m_2^\epsilon(t), \end{aligned} \quad (2.54)$$

4 where $(x^2)_N$ means the cutoff $(x^2)_N = \min\{x^2, N\}$. Hence, we obtain (2.29).

5 For the energy dissipation (2.30), we prove it by taking limit in (2.13), Levi's lemma and Fatou's
6 Lemma. First by pointwise convergence of ρ^ϵ in (2.53), pointwise convergence of $H\rho^\epsilon$ in (2.51) and
7 Fatou's lemma, we have

$$\int_0^T \int_{\mathbb{R}} \rho |\gamma x - \pi H \rho(x, t)|^2 \, dx \leq \liminf_{\epsilon \rightarrow 0} \int_0^T \int_{\mathbb{R}} \rho^\epsilon |\gamma x - \pi H \rho^\epsilon(x, t)|^2 \, dx. \quad (2.55)$$

8 Second, there exists a constant c such that $K(x, y) := \frac{1}{2}\gamma(x^2 + y^2) + \log \frac{1}{|x-y|} + c \geq 0$, so we rewrite
9 the energy as

$$E(\rho) = \frac{1}{2} \int_{\mathbb{R}^2} K(x, y) \rho(x) \rho(y) \, dx \, dy - \frac{c}{2}. \quad (2.56)$$

10 Denote the cutoff of K as $K_N(x, y) := \min\{K(x, y), N\}$ such that $0 \leq K_N(x, y) \leq K(x, y)$, which
11 increasingly converges to $K(x, y)$ for a.e. $(x, y) \in \mathbb{R}^2$. Then by Levi's lemma and Fatou's Lemma,
12 we obtain

$$\begin{aligned} E(\rho) + \frac{c}{2} &= \lim_{N \rightarrow +\infty} \frac{1}{2} \int_{\mathbb{R}^2} K_N(x, y) \rho(x) \rho(y) \, dx \, dy \\ &\leq \lim_{N \rightarrow +\infty} \liminf_{\epsilon \rightarrow 0} \frac{1}{2} \int_{\mathbb{R}^2} K_N(x, y) \rho^\epsilon(x) \rho^\epsilon(y) \, dx \, dy \\ &\leq \lim_{N \rightarrow +\infty} \liminf_{\epsilon \rightarrow 0} \frac{1}{2} \int_{\mathbb{R}^2} K(x, y) \rho^\epsilon(x) \rho^\epsilon(y) \, dx \, dy \\ &\leq \liminf_{\epsilon \rightarrow 0} E(\rho^\epsilon) + \frac{c}{2}. \end{aligned} \quad (2.57)$$

13 The entropy inequality (2.31) can be obtained by (2.14) and the weak lower semi-continuity of
14 the entropy [10].

15

□

16 **Remark 2.3.** We shall remark that the global existence of weak solutions for the following non-
17 conservative equation remains open:

$$\partial_t \rho - u \partial_x \rho = 0, \quad u = H\rho.$$

18 We refer to [6, 28] for in depth study of this equation with or without a viscous term.

Remark 2.4 (Exponential convergence to the steady state). Carrillo et. al. [3] proved the
existence and uniqueness of probability solutions by using gradient flow structure in Wasserstein
distance. Notice the free energy $E(\rho)$ given by (1.9) for the Dyson equation consists a harmonic trap
energy E_h and an interaction energy E_i . E_i is convex (or displacement convex) along generalized
Wasserstein geodesics and E_h is γ -convex along Wasserstein geodesics as explained below. Assume
 $\rho_0, \rho_1 \in \mathcal{P}_{AC}(\mathbb{R})$ and $T : \rho_0 \, dx \rightarrow \rho_1 \, dy$ is W_2 -optimal transport (Bernier's map). Then $\rho_t :=$
 $[tI + (1-t)T]_{\#} \rho_0$ is a Wasserstein geodesics (or displacement interpolation between ρ_0 and ρ_1).
From the definition of push forward (see [1, Section 5.2]),

$$\begin{aligned} E_h(\rho_t) &= \gamma \int_{\mathbb{R}} \frac{x^2}{2} \rho_t(dx) = \gamma \int_{\mathbb{R}} \frac{[tx + (1-t)T(x)]^2}{2} \rho_0(dx) \\ &= \gamma \int_{\mathbb{R}} \frac{tx^2 - t(1-t)(x - T(x))^2 + (1-t)T^2(x)}{2} \rho_0(dx) \\ &= tE_h(\rho_0) + (1-t)E_h(\rho_1) - \gamma \frac{t(1-t)}{2} W_2^2(\rho_0, \rho_1). \end{aligned}$$

Therefore $E_h(\rho)$ is γ -geodesically convex (see [1, Definition 2.4.3]). For the geodesical convexity of the interaction energy $E_i(\rho)$, due to the singularity in logarithmic function, it relies heavily on monotonicity of optimal map. We illustrate the idea for $\rho_0, \rho_1 \in \mathcal{P}_{AC}(\mathbb{R}), \rho_0 > 0$, which ensures the optimal map T is strictly increasing.

$$\begin{aligned}
E_i(\rho_t) &= \int_{\mathbb{R}^2} -\log(|x-y|)\rho_t(dx)\rho_t(dy) \\
&= \int_{\mathbb{R}^2} -\log(|t(x-y) + (1-t)(T(x)-T(y))|)\rho_0(x)\rho_0(y) dx dy \\
&\leq t \int_{\mathbb{R}^2} -\log|x-y|\rho_0(x)\rho_0(y) dx dy + (1-t) \int_{\mathbb{R}^2} -\log|T(x)-T(y)|\rho_0(x)\rho_0(y) dx dy \\
&= t \int_{\mathbb{R}^2} -\log|x-y|\rho_0(x)\rho_0(y) dx dy + (1-t) \int_{\mathbb{R}^2} -\log|x-y|\rho_1(x)\rho_1(y) dx dy \\
&= tE_i(\rho_0) + (1-t)E_i(\rho_1),
\end{aligned}$$

where we used the convexity of logarithmic function in the first inequality and strict increase of T in the third equality. However, without the strictly increasing property, we refer to [3, Proposition 2.7], where Carrillo et. al. proved the generalized geodesic convex of $E_i(\rho)$ using the essential monotonicity property (excluding a null set) of the optimal transport maps between absolutely continuous probability measures in one dimension. The standard gradient flow theory [1, Theorem 11.2.1] yields the exponential convergence to the steady state in W_2 distance; see also [3, 2]. More precisely, if ρ and $\tilde{\rho}$ are two probability measure solutions for initial date ρ_0 and $\tilde{\rho}_0$ separately, then we have

$$W_2(\rho(t), \tilde{\rho}(t)) \leq e^{-\gamma t} W_2(\rho_0, \tilde{\rho}_0). \quad (2.58)$$

1 This implies the uniqueness of probability measure solutions and exponential convergence to the
2 steady state.

3 When $\gamma > 0$, we also remark that γ -convexity of E implies the uniqueness of the steady state
4 (minimizer). Indeed, if μ and ν are two distinct minimizers, consider $\mu_{1/2} := [\frac{1}{2}I + \frac{1}{2}\tilde{T}]_{\#}\mu$, where
5 \tilde{T} is Bernier's map between μ and ν . Then, we have

$$E(\mu_{1/2}) \leq \frac{1}{2}[E(\mu) + E(\nu)] - \frac{\gamma}{8}W_2^2(\mu, \nu) < \frac{1}{2}[E(\mu) + E(\nu)],$$

6 which is a contradiction with that μ and ν are distinct.

7 **3 Bi-Hamiltonian structures**

8 In this section, we construct a bi-Hamiltonian structure for the coupled Burgers system (1.13)
9 by using the decoupled Burgers equations (1.16) and (1.17). First, we present infinite many
10 conserved quantities for the coupled Burgers system (1.13). Recall (1.15). Because $\int_{\mathbb{R}} f_{\pm}^k(x, t) dx$
11 are conserved quantities of the decoupled Burgers equations (1.16) and (1.17), we have the following
12 proposition.

Proposition 3.1. *Let (ρ, u) be a classical solution to the coupled Burgers system (1.13). Then, quantities*

$$\lambda_1 \int_{\mathbb{R}} (u + \sqrt{\alpha\rho})^{k_1} dx + \lambda_2 \int_{\mathbb{R}} (u - \sqrt{\alpha\rho})^{k_2} dx \quad (3.1)$$

13 *are conserved for any constants $\lambda_1, \lambda_2 \in \mathbb{C}$ and any positive integers $k_1, k_2 \in \mathbb{N}_+$.*

14 **Remark 3.1.** Notice that when $\alpha < 0$, we have $f_- = \bar{f}_+$. When $\lambda_1 = \bar{\lambda}_2$ and $k_1 = k_2 = k$ in
15 (3.1), we have

$$\overline{\lambda_1(u + \sqrt{\alpha\rho})^k} = \lambda_2(u - \sqrt{\alpha\rho})^k.$$

16 In this case, (3.1) gives real conserved quantities.

Next, we consider the case for $k_1 = k_2 = 3$ in (3.1) and derive a bi-Hamiltonian structure for the coupled Burgers system (1.13). Define the following functionals of $f_{\pm}(= u \pm \sqrt{\alpha}\rho)$:

$$H_1^f(f_+, f_-) := \int_{\mathbb{R}} \frac{f_+^3 + f_-^3}{12} dx, \quad H_2^f(f_+, f_-) := \int_{\mathbb{R}} \frac{f_+^3 - f_-^3}{12\sqrt{\alpha}} dx. \quad (3.2)$$

Due to Remark 3.1, we know that both H_1^f and H_2^f are real conserved quantities. Moreover, the decoupled Burgers equations (1.16) and (1.17) can be rewritten as

$$\begin{cases} \partial_t f_+ + 2\partial_x \left(\frac{\delta H_1^f}{\delta f_+} \right) = 0, \\ \partial_t f_- + 2\partial_x \left(\frac{\delta H_1^f}{\delta f_-} \right) = 0, \end{cases} \quad \text{and} \quad \begin{cases} \partial_t f_+ + 2\sqrt{\alpha}\partial_x \left(\frac{\delta H_2^f}{\delta f_+} \right) = 0, \\ \partial_t f_- - 2\sqrt{\alpha}\partial_x \left(\frac{\delta H_2^f}{\delta f_-} \right) = 0. \end{cases} \quad (3.3)$$

Define

$$H_1^u(\rho, u) := \int_{\mathbb{R}} \left(\frac{1}{6}u^3 + \frac{\alpha}{2}\rho^2 u \right) dx, \quad H_2^u(\rho, u) := \int_{\mathbb{R}} \left(\frac{1}{2}\rho u^2 + \frac{\alpha}{6}\rho^3 \right) dx. \quad (3.4)$$

Then, direct calculations show that

$$H_j^u(\rho, u) = H_j^f(f_+, f_-), \quad j = 1, 2, \quad (3.5)$$

1 and we have the following theorem:

2 **Theorem 3.1.** *For $\alpha \neq 0$, the coupled Burgers system (1.13) has a bi-Hamiltonian structure:*

$$\frac{\partial}{\partial t} \begin{pmatrix} \rho \\ u \end{pmatrix} = J \begin{pmatrix} \frac{\delta H_1^u}{\delta \rho} \\ \frac{\delta H_1^u}{\delta u} \end{pmatrix} = K \begin{pmatrix} \frac{\delta H_2^u}{\delta \rho} \\ \frac{\delta H_2^u}{\delta u} \end{pmatrix}, \quad (3.6)$$

where J and K are anti-symmetric operators given by

$$J := \begin{pmatrix} -\frac{1}{\alpha}\partial_x & 0 \\ 0 & -\partial_x \end{pmatrix}, \quad K := \begin{pmatrix} 0 & -\partial_x \\ -\partial_x & 0 \end{pmatrix}. \quad (3.7)$$

Proof. Due to $f_{\pm} = u \pm \sqrt{\alpha}\rho$, we have $\rho = \frac{1}{2\sqrt{\alpha}}(f_+ - f_-)$ and $u = \frac{1}{2}(f_+ + f_-)$. From (3.3), we obtain

$$\begin{cases} \partial_t \rho + \frac{1}{\sqrt{\alpha}}\partial_x \left(\frac{\delta H_1^f}{\delta f_+} - \frac{\delta H_1^f}{\delta f_-} \right) = 0, \\ \partial_t u + \partial_x \left(\frac{\delta H_1^f}{\delta f_+} + \frac{\delta H_1^f}{\delta f_-} \right) = 0, \end{cases} \quad \text{and} \quad \begin{cases} \partial_t \rho + \partial_x \left(\frac{\delta H_2^f}{\delta f_+} + \frac{\delta H_2^f}{\delta f_-} \right) = 0, \\ \partial_t u + \sqrt{\alpha}\partial_x \left(\frac{\delta H_2^f}{\delta f_+} - \frac{\delta H_2^f}{\delta f_-} \right) = 0. \end{cases} \quad (3.8)$$

Due to (3.5), we have the following relations:

$$\frac{\delta H_j^u}{\delta \rho} = \sqrt{\alpha} \left(\frac{\delta H_j^f}{\delta f_+} - \frac{\delta H_j^f}{\delta f_-} \right), \quad \frac{\delta H_j^u}{\delta u} = \frac{\delta H_j^f}{\delta f_+} + \frac{\delta H_j^f}{\delta f_-}, \quad j = 1, 2. \quad (3.9)$$

Put (3.9) into (3.8) and we obtain

$$\begin{cases} \partial_t \rho + \frac{1}{\alpha}\partial_x \left(\frac{\delta H_1^u}{\delta \rho} \right) = 0, \\ \partial_t u + \partial_x \left(\frac{\delta H_1^u}{\delta u} \right) = 0, \end{cases} \quad \text{and} \quad \begin{cases} \partial_t \rho + \partial_x \left(\frac{\delta H_2^u}{\delta \rho} \right) = 0, \\ \partial_t u + \partial_x \left(\frac{\delta H_2^u}{\delta \rho} \right) = 0, \end{cases}$$

3 which is (3.6). □

1 From Theorem 3.1, we can directly obtain a bi-Hamiltonian structure for System (1.18), as
 2 shown in the following corollary:

3 **Corollary 3.1.** For $\alpha \neq 0$, the isentropic gas dynamics (1.18) can be rewritten as the following
 4 bi-Hamiltonian structure:

$$\frac{\partial}{\partial t} \begin{pmatrix} \rho \\ m \end{pmatrix} = \tilde{J} \begin{pmatrix} \frac{\delta H_1^m}{\delta \rho} \\ \frac{\delta H_1^m}{\delta m} \end{pmatrix} = \tilde{K} \begin{pmatrix} \frac{\delta H_2^m}{\delta \rho} \\ \frac{\delta H_2^m}{\delta m} \end{pmatrix}, \quad (3.10)$$

where \tilde{J} and \tilde{K} are anti-symmetric operators given by

$$\tilde{J} = \begin{pmatrix} -\frac{1}{\alpha} \partial_x & -\frac{1}{\alpha} \partial_x u \\ -\frac{1}{\alpha} u \partial_x & -\frac{1}{\alpha} u \partial_x u - \rho \partial_x \rho \end{pmatrix}, \quad \tilde{K} = \begin{pmatrix} 0 & -\partial_x \rho \\ -\rho \partial_x & -u \partial_x \rho - \rho \partial_x u \end{pmatrix}, \quad (3.11)$$

and the Hamiltonians are given by

$$H_1^m(\rho, m) = \int_{\mathbb{R}} \left(\frac{m^3}{6\rho^3} + \frac{\alpha}{2} m \rho \right) dx, \quad H_2^m(\rho, m) = \int_{\mathbb{R}} \left(\frac{m^2}{2\rho} + \frac{\alpha}{6} \rho^3 \right) dx. \quad (3.12)$$

5 *Proof.* Due to $m = \rho u$, we have

$$H_j^m(\rho, m) := H_j^u(\rho, u), \quad j = 1, 2.$$

6 Moreover, we have the following relations:

$$\frac{\delta H_j^u}{\delta \rho} = \frac{\delta H_j^m}{\delta \rho} + u \frac{\delta H_j^m}{\delta m}, \quad \frac{\delta H_j^u}{\delta u} = \rho \frac{\delta H_j^m}{\delta m},$$

7 or equivalently

$$\begin{pmatrix} \frac{\delta H_j^u}{\delta \rho} \\ \frac{\delta H_j^u}{\delta u} \end{pmatrix} = \begin{pmatrix} 1 & u \\ 0 & \rho \end{pmatrix} \begin{pmatrix} \frac{\delta H_j^m}{\delta \rho} \\ \frac{\delta H_j^m}{\delta m} \end{pmatrix}, \quad j = 1, 2. \quad (3.13)$$

Combining (3.6) and (3.13), we obtain

$$\begin{aligned} \frac{\partial}{\partial t} \begin{pmatrix} \rho \\ m \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ u & \rho \end{pmatrix} \begin{pmatrix} \partial_t \rho \\ \partial_t u \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ u & \rho \end{pmatrix} J \begin{pmatrix} 1 & u \\ 0 & \rho \end{pmatrix} \begin{pmatrix} \frac{\delta H_1^m}{\delta \rho} \\ \frac{\delta H_1^m}{\delta m} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ u & \rho \end{pmatrix} K \begin{pmatrix} 1 & u \\ 0 & \rho \end{pmatrix} \begin{pmatrix} \frac{\delta H_2^m}{\delta \rho} \\ \frac{\delta H_2^m}{\delta m} \end{pmatrix}. \end{aligned} \quad (3.14)$$

Hence, we have

$$\tilde{J} = \begin{pmatrix} 1 & 0 \\ u & \rho \end{pmatrix} J \begin{pmatrix} 1 & u \\ 0 & \rho \end{pmatrix} = \begin{pmatrix} -\frac{1}{\alpha} \partial_x & -\frac{1}{\alpha} \partial_x u \\ -\frac{1}{\alpha} u \partial_x & -\frac{1}{\alpha} u \partial_x u - \rho \partial_x \rho \end{pmatrix},$$

and

$$\tilde{K} = \begin{pmatrix} 1 & 0 \\ u & \rho \end{pmatrix} K \begin{pmatrix} 1 & u \\ 0 & \rho \end{pmatrix} = \begin{pmatrix} 0 & -\partial_x \rho \\ -\rho \partial_x & -u \partial_x \rho - \rho \partial_x u \end{pmatrix}.$$

8 Hence, we obtain a bi-Hamiltonian structure for System (1.18). \square

Notice that H_2^m is nothing but the total energy of System (1.18), which is given by

$$H_2^m(\rho, m) = \int_{\mathbb{R}} E(x, t) dx = \int_{\mathbb{R}} \left(\frac{1}{2} \rho u^2 + \frac{\alpha}{6} \rho^3 \right) dx = \int_{\mathbb{R}} \left(\frac{m^2}{2\rho} + \frac{\alpha \rho^3}{6} \right) dx. \quad (3.15)$$

9 where $E(x, t)$ is defined by (1.22).

1 **Remark 3.2** (A bi-Hamiltonian structure for p-system (1.23)). Set

$$\eta(\xi, t) := \frac{1}{\tau(\xi, t)}, \quad \xi \in (0, 1), \quad t > 0.$$

Then, the p-system (1.23) becomes the following system for (η, V) :

$$\begin{cases} \partial_\xi \eta = -\eta^2 \partial_\xi V, \\ \partial_t V = -\alpha \eta^2 \partial_\xi \eta. \end{cases} \quad (3.16)$$

We have the following bi-Hamiltonian structure for System (3.16):

$$\frac{\partial}{\partial t} \begin{pmatrix} \eta \\ V \end{pmatrix} = \begin{pmatrix} -\frac{3}{4\alpha} \eta \partial_\xi \eta & -\frac{1}{4\alpha} \eta \partial_\xi V + \frac{3}{2\alpha} V \partial_\xi \eta - \frac{1}{\alpha} \partial_\xi \eta V \\ -\frac{1}{4\alpha} V \partial_\xi \eta + \frac{3}{2\alpha} \eta \partial_\xi V - \frac{1}{\alpha} \eta V \partial_\xi & \frac{1}{4\alpha} V \partial_\xi V + \frac{1}{\alpha} \eta \partial_\xi \eta \end{pmatrix} \begin{pmatrix} \frac{\delta H_1^\eta}{\delta \eta} \\ \frac{\delta H_1^\eta}{\delta V} \end{pmatrix},$$

and

$$\frac{\partial}{\partial t} \begin{pmatrix} \eta \\ V \end{pmatrix} = \begin{pmatrix} 0 & -\eta^2 \partial_\xi \\ -\partial_\xi \eta^2 & 0 \end{pmatrix} \begin{pmatrix} \frac{\delta H_2^\eta}{\delta \eta} \\ \frac{\delta H_2^\eta}{\delta V} \end{pmatrix},$$

where

$$H_1^\eta(\eta, V) = \int_{\mathbb{R}} \left(\frac{V^3}{6\eta} + \alpha \frac{\eta V}{2} \right) d\xi, \quad H_2^\eta(\eta, V) = \int_{\mathbb{R}} \left(\frac{V^2}{2} + \alpha \frac{\eta^2}{6} \right) d\xi.$$

2 4 Kinetic formulations and entropy solutions for the coupled Burgers system (1.13) with $\alpha > 0$

3
4 In this section, we study the kinetic formulation for the coupled Burgers system (1.13) with $\alpha > 0$.
5 In contrast, Lions, Perthame and Tadmor [15] studied System (1.18) and they used the kinetic
6 formulation to obtain global entropy solutions without uniqueness. Here, we show the existence
7 and uniqueness of global entropy solutions for (1.13).

8 4.1 Kinetic formulations

Kinetic formulation is a method which use the distribution function $\kappa(v, x, t)$ at time t in the phase plane for velocity v and the position x to study the continuum equation for $u(x, t)$ (and $\rho(x, t)$). At fixed continuum variable (x, t) , u and ρ are some v -moments of κ . In the local thermal equilibrium the distribution function $\kappa(v, x, t)$ can be described by v -equilibrium distribution $\chi(v; u, \rho)$ with parameters u and ρ , i.e. $\kappa(v, x, t) = \chi(v; u(x, t), \rho(x, t))$. In kinetic theory, the v -equilibrium distribution is also known as Maxwellian. Following the idea of the celebrated work by Lions, Perthame and Tadmor [15], we use the combinations of Heaviside function,

$$H(v) = \begin{cases} 1, & v \geq 0, \\ 0, & v < 0, \end{cases}$$

to construct the equilibrium distribution. Let (ρ, u) be a solution to the coupled Burgers system (1.13) with $\alpha > 0$. Recall (1.15)

$$f_\pm = u \pm \sqrt{\alpha} \rho.$$

9 Then, f_\pm are solutions to the decoupled Burgers equations (1.16) and (1.17). We use the following
10 v -equilibrium distributions

$$\begin{aligned} \chi_+(v; \rho, u) &:= H(v) - H(v - f_+), & \chi_-(v; \rho, u) &:= H(v) - H(v - f_-), \\ \chi(v; \rho, u) &:= \frac{1}{2\sqrt{\alpha}}(\chi_+ - \chi_-) = \frac{1}{2\sqrt{\alpha}}[H(v - f_-) - H(v - f_+)], \end{aligned} \quad (4.1)$$

1 and

$$\hat{\chi}(v; \rho, u) := \frac{1}{2}(\chi_+ + \chi_-) = \frac{1}{2}[2H(v) - H(v - f_-) - H(v - f_+)].$$

For any nonnegative integer k , direct calculations show that the following k -moments equality holds

$$\int_{\mathbb{R}} v^k \chi_{\pm}(v; \rho, u) dv = \frac{f_{\pm}^{k+1}}{k+1}. \quad (4.2)$$

Hence, the conserved quantities given by (3.1) correspond to the integration (w.r.t. x variable) of the following kinetic formulations:

$$\lambda_1 \int_{\mathbb{R}} v^{k_1} \chi_+(v; \rho, u) dv + \lambda_2 \int_{\mathbb{R}} v^{k_2} \chi_-(v; \rho, u) dv, \quad \lambda_i \in \mathbb{C}, \quad k_i \in \mathbb{N}_+, \quad i = 1, 2. \quad (4.3)$$

Choosing $\lambda_1 = \lambda_2 = \frac{1}{4}$ and $k_1 = k_2 = 2$, we obtain the Hamiltonian H_1^u and choosing $\lambda_1 = -\lambda_2 = \frac{1}{4\sqrt{\alpha}}$ and $k_1 = k_2 = 2$, we obtain the Hamiltonian H_2^u given by (3.4). More precisely, we have

$$H_1^u(\rho, u) = \frac{1}{2} \int_{\mathbb{R}} v^2 \hat{\chi}(v; \rho, u) dv, \quad H_2^u(\rho, u) = \frac{1}{2} \int_{\mathbb{R}} v^2 \chi(v; \rho, u) dv. \quad (4.4)$$

2 By (4.2), the decoupled Burgers equations (1.16) and (1.17) have the following kinetic formulations:

$$\int_{\mathbb{R}} (\partial_t \chi_{\pm} + v \partial_x \chi_{\pm}) dv = \partial_t f_{\pm} + f_{\pm} \partial_x f_{\pm} = 0.$$

3 Besides, we also have

$$u = \int_{\mathbb{R}} \hat{\chi}(v; \rho, u) dv, \quad \rho = \int_{\mathbb{R}} \chi(v; \rho, u) dv.$$

4 Hence, the coupled Burgers system (1.13) has the following kinetic formulation:

$$\int_{\mathbb{R}} \begin{pmatrix} \partial_t \chi + v \partial_x \chi \\ \partial_t \hat{\chi} + v \partial_x \hat{\chi} \end{pmatrix} dv = \begin{pmatrix} \partial_t \rho + \partial_x(\rho u) \\ \partial_t u + \partial_x \left(\frac{u^2 + \alpha \rho^2}{2} \right) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (4.5)$$

5 Moreover, direct calculations show that

$$\begin{pmatrix} \rho u \\ E \end{pmatrix} = \int_{\mathbb{R}} \begin{pmatrix} v \\ \frac{v^2}{2} \end{pmatrix} \chi(v; \rho, u) dv, \quad (4.6)$$

where E is the total energy given by (1.22). Comparing with (4.5), we have the following kinetic formulation for the isentropic gas system (1.18):

$$\int_{\mathbb{R}} \begin{pmatrix} 1 \\ v \end{pmatrix} (\partial_t \chi + v \partial_x \chi) dv = \begin{pmatrix} \partial_t \rho + \partial_x(\rho u) \\ \partial_t(\rho u) + \partial_x(\rho u^2 + p) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

6 4.2 Existence and uniqueness of entropy solutions for the coupled Burgers system (1.13)

7

The notion of the entropy-entropy-flux pair refers to the pair of regular functions (η, q) defined on the space of the states (ρ, u) for which every classical solution (ρ, u) of the coupled Burgers system (1.13), also satisfies

$$\partial_t \eta(\rho, u) + \partial_x q(\rho, u) = 0. \quad (4.7)$$

8 Combining the coupled Burgers system (1.13) and (4.7) gives

$$(\partial_\rho q - u \partial_\rho \eta - \alpha \rho \partial_u \eta) \partial_x \rho + (\partial_u q - \rho \partial_\rho \eta - u \partial_u \eta) \partial_x u = 0,$$

which holds for any smooth solutions (ρ, u) . The entropy pair (η, q) can be found by solving the following Euler-Poisson-Darboux equations. First, given $\psi(u), g(u)$, we solve $\eta(\rho, u)$ satisfying

$$\begin{cases} \partial_{\rho\rho}\eta - \alpha\partial_{uu}\eta = 0, \\ \eta(0, u) = \psi(u), \quad \partial_{\rho}\eta(0, u) = g(u). \end{cases} \quad (4.8)$$

Then we solve the entropy flux q by

$$\partial_u q = \rho\partial_{\rho}\eta + u\partial_u\eta, \quad \partial_{\rho}q = u\partial_{\rho}\eta + \alpha\rho\partial_u\eta. \quad (4.9)$$

1 From (4.8), we know $\partial_{u\rho}q = \partial_{\rho u}q$ so (4.9) is solvable. We have the following results:

2 **Proposition 4.1.** *For two given functions $\psi, g \in C^2(\mathbb{R})$, let $(\eta(0, u), \partial_{\rho}\eta(0, u)) = (\psi(u), g(u))$ be*
 3 *the initial datum for (4.8). Then:*

(i) *The solution $\eta(\rho, u)$ to (4.8) can be recast in a kinetic representation:*

$$\eta(\rho, u) = \int_{\mathbb{R}} \psi'(v)\hat{\chi}(v; \rho, u) dv + \int_{\mathbb{R}} g(v)\chi(v; \rho, u) dv \quad (4.10)$$

(ii) *When $\psi = 0$ and $\rho \geq 0$, we have kinetic representations*

$$\eta_g(\rho, u) := \int_{\mathbb{R}} g(v)\chi(v; \rho, u) dv, \quad q_g(\rho, u) := \int_{\mathbb{R}} vg(v)\chi(v; \rho, u) dv. \quad (4.11)$$

4 *Moreover, η_g is convex with respect to (ρ, m) if and only if $g(v)$ is convex, where $m = \rho u$.*

(iii) *When $g = 0$, we have kinetic representations*

$$\eta_{\psi}(\rho, u) := \int_{\mathbb{R}} \psi'(v)\hat{\chi}(v; \rho, u) dv \quad (4.12)$$

$$q_{\psi}(\rho, u) := \int_{\mathbb{R}} v\psi'(v)\hat{\chi}(v; \rho, u) dv = \frac{\phi(u + \sqrt{\alpha}\rho) + \phi(u - \sqrt{\alpha}\rho)}{2}, \quad (4.13)$$

5 *where $\phi'(v) = v\psi'(v)$ for $v \in \mathbb{R}$. Moreover, η_{ψ} is convex with respect to (ρ, u) if and only if ψ is*
 6 *a convex function.*

Proof. (i) By the d'Alembert's formula, we have

$$\eta(\rho, u) = \frac{\psi(f_+) + \psi(f_-)}{2} + \frac{1}{2\sqrt{\alpha}} \int_{f_-}^{f_+} g(v) dv, \quad (4.14)$$

7 where $f_{\pm}(\rho, u) = u \pm \sqrt{\alpha}\rho$. Formula (4.10) is exactly the kinetic formulation for the formula (4.14).

8 (ii) First we verify (4.11) satisfies (4.9). We have

$$\partial_u \eta_g = g(u + \sqrt{\alpha}\rho) - g(u - \sqrt{\alpha}\rho), \quad \partial_{\rho} \eta_g = \sqrt{\alpha} \left(g(u + \sqrt{\alpha}\rho) + g(u - \sqrt{\alpha}\rho) \right),$$

and

$$\begin{aligned} \partial_u q_g &= (u + \sqrt{\alpha}\rho)g(u + \sqrt{\alpha}\rho) - (u - \sqrt{\alpha}\rho)g(u - \sqrt{\alpha}\rho) \\ &= u\partial_u \eta_g + \rho\partial_{\rho} \eta_g. \end{aligned}$$

9 Similarly, we also have $\partial_{\rho} q_{\psi} = u\partial_{\rho} \eta_{\psi} + \alpha\rho\partial_u \eta_{\psi}$.

Second, we check the convexity condition for η_g in terms of (ρ, m) , where $m = \rho u$. By changing of variables $v = u \pm \xi\rho$, we have

$$\begin{aligned} \eta_g(\rho, u) &= \int_{\mathbb{R}} g(v)\chi(v; \rho, u) dv = \frac{1}{2\sqrt{\alpha}} \int_{f_-}^{f_+} g(v) dv \\ &= \frac{1}{2\sqrt{\alpha}} \int_{-\sqrt{\alpha}}^{\sqrt{\alpha}} \rho g\left(\frac{m}{\rho} + \xi\rho\right) d\xi. \end{aligned} \quad (4.15)$$

1 Taking derivatives of (4.15), we can obtain

$$\partial_{\rho\rho}\eta_g = \frac{1}{2\sqrt{\alpha}} \int_{-\sqrt{\alpha}}^{\sqrt{\alpha}} \rho g'' \left(\frac{m}{\rho} + \xi\rho \right) \left(-\frac{m}{\rho^2} + \xi \right)^2 d\xi + \frac{1}{\sqrt{\alpha}} \int_0^{\sqrt{\alpha}} \xi \left[g'(u + \xi\rho) - g'(u - \xi\rho) \right] d\xi.$$

2 When g is convex, g' is increasing and $g'' > 0$. Hence $\partial_{\rho\rho}\eta_g \geq 0$. Moreover, we have

$$\partial_{\rho m}\eta_g = \frac{1}{2\sqrt{\alpha}} \int_{-\sqrt{\alpha}}^{\sqrt{\alpha}} g'' \left(\frac{m}{\rho} + \xi\rho \right) \left(-\frac{m}{\rho^2} + \xi \right) d\xi,$$

3 and

$$\partial_{mm}\eta_g = \frac{1}{2\sqrt{\alpha}} \int_{-\sqrt{\alpha}}^{\sqrt{\alpha}} \frac{1}{\rho} g'' \left(\frac{m}{\rho} + \xi\rho \right) d\xi \geq 0.$$

4 By Hölder's inequality, we can obtain

$$\partial_{\rho\rho}\eta_g \cdot \partial_{mm}\eta_g - (\partial_{\rho m}\eta_g)^2 \geq 0.$$

5 Hence, $\eta_g(\rho, u)$ is convex about (ρ, m) . When g is not convex, we have $g'' < 0$ in some interval.
6 This implies $\partial_{mm}\eta_g < 0$. Hence, η_g is not convex. This proves that η_g is convex if and only if g is
7 convex.

(iii) First, we verify equalities in (4.9) hold for (η_ψ, q_ψ) . For $f_\pm = u \pm \sqrt{\alpha}\rho$, we have

$$\begin{aligned} \partial_u q_\psi &= \frac{f_+ \psi'(f_+) + f_- \psi'(f_-)}{2} = u \frac{\psi'(f_+) + \psi'(f_-)}{2} + \rho \frac{\sqrt{\alpha} \psi'(f_+) - \sqrt{\alpha} \psi'(f_-)}{2} \\ &= u \partial_u \eta_\psi + \rho \partial_\rho \eta_\psi. \end{aligned}$$

8 Similarly, $\partial_\rho q_\psi = u \partial_\rho \eta_\psi + \alpha \rho \partial_u \eta_\psi$. Hence, equalities in (4.9) hold for (η_ψ, q_ψ) . This proves that
9 q_ψ is the corresponding entropy flux of η_ψ .

Second, we check the convexity condition for η_ψ in terms of (ρ, u) . Notice that

$$\eta_\psi(\rho, u) = \int_{\mathbb{R}} \psi'(v) \hat{\chi}(v; \rho, u) dv = \frac{\psi(f_+) + \psi(f_-)}{2}, \quad (4.16)$$

10 where $f_\pm = u \pm \sqrt{\alpha}\rho$. Taking derivative of (4.16), we can obtain

$$\partial_{\rho u} \eta_\psi = \frac{\sqrt{\alpha}(\psi''(f_+) - \psi''(f_-))}{2}, \quad \partial_{\rho\rho} \eta_\psi = \frac{\alpha(\psi''(f_+) + \psi''(f_-))}{2},$$

11 and

$$\partial_{uu} \eta_\psi = \frac{\psi''(f_+) + \psi''(f_-)}{2}.$$

When ψ is convex, we have

$$\partial_{uu} \eta_\psi \geq 0, \partial_{uu} \eta_\psi \geq 0, \quad \text{and} \quad \partial_{\rho\rho} \eta_\psi \partial_{uu} \eta_\psi \geq (\partial_{\rho u} \eta_\psi)^2,$$

12 which means η_ψ is convex with respect to (ρ, u) . Conversely, if η_ψ is convex with respect to (ρ, u) ,
13 $\psi(u) = \eta_\psi(0, u)$ is convex. □

15 In [15], Lions, Perthame and Tadmor studied the kinetic formulation of the isentropic gas
16 system (1.18). The convex entropies they used to define solutions corresponds to $\eta_g(\rho, u)$ given
17 by (4.10) for convex functions g . The corresponding entropy flux are given by (4.11). Recall their
18 definition of the entropy solutions to the isentropic gas system (1.18) (see [15, Definition 2]).

Definition 4.1. A couple (ρ, m) is called an entropy solution of (1.18) if it satisfies

$$\partial_t \eta_g(\rho, u) + \partial_x q_g(\rho, u) \leq 0, \quad (4.17)$$

19 in distribution sense for all convex entropies η_g given by (4.10) with convex g .

An important example for is taking $g(v) = \frac{v^2}{2}$ in (4.15). Direct calculations show that the entropy has the following kinetic formulation

$$\eta_g = \frac{1}{2}\rho u^2 + \frac{\alpha}{6}\rho^3 =: E$$

1 and the entropic flux is

$$q_g = \int_{\mathbb{R}} v g(v) \chi(v; \rho, u) dv = \frac{1}{4\sqrt{\alpha}} \int_{f_-}^{f_+} v^3 dv = \frac{1}{16\sqrt{\alpha}} (f_+^4 - f_-^4) = u(E + p).$$

Then (4.17) in Definition 4.1 becomes

$$\partial_t E + \partial_x [(E + p)u] \leq 0 \quad (4.18)$$

2 in the distributional sense. Notice that $g(v) = \frac{v^2}{2}$ is convex and hence E is convex with respect to
3 (ρ, m) .

4 **Remark 4.1.** Note that global existence of entropy solutions to System (1.18) was proved [15]. It
5 is shown in [15] that (ρ, m) is a weak entropy solution with respect to the family $\{\eta_g\}$, if and only
6 if the kinetic function $\chi(v; \rho, u)$ given by (4.1) is a weak solution of the kinetic equation

$$\partial_t \chi + v \partial_x \chi = -\partial_{vv} \mu,$$

7 for some finite Radon measure $\mu \in \mathcal{M}_+$. Hence, the entropy inequality (4.17) has a kinetic formu-
8 lation:

$$\partial_t \eta_g(\rho, u) + \partial_x q_g(\rho, u) = \int_{\mathbb{R}} g(v) (\partial_t \chi + v \partial_x \chi) dv = - \int_{\mathbb{R}} g''(v) d\mu \leq 0$$

9 in the distributional sense for all $g \in C_0^2(\mathbb{R})$ and $g'' \geq 0$ on the support of μ .

10 4.2.1 Existence and uniqueness of entropy solutions of (1.13)

11 Next, to obtain the uniqueness of entropy solutions, we consider the entropy solutions of the
12 coupled Burgers system (1.13). We have the following proposition for entropy pairs (η, q) :

Proposition 4.2. *Let $\psi_1, \psi_2 \in C^2(\mathbb{R})$ be two convex functions. Define*

$$\begin{aligned} \eta(\rho, u) &:= k_1 \int_{\mathbb{R}} \psi_1'(v) \chi_+(v; \rho, u) dv + k_2 \int_{\mathbb{R}} \psi_2'(v) \chi_-(v; \rho, u) dv \\ &= k_1 \psi_1(u + \sqrt{\alpha}\rho) + k_2 \psi_2(u - \sqrt{\alpha}\rho), \end{aligned} \quad (4.19)$$

and

$$\begin{aligned} q(\rho, u) &:= k_1 \int_{\mathbb{R}} v \psi_1'(v) \chi_+(v; \rho, u) dv + k_2 \int_{\mathbb{R}} v \psi_2'(v) \chi_-(v; \rho, u) dv \\ &= k_1 \phi_1(u + \sqrt{\alpha}\rho) + k_2 \phi_2(u - \sqrt{\alpha}\rho), \end{aligned} \quad (4.20)$$

13 where k_1 and k_2 are two nonnegative real numbers and ϕ satisfies $\phi_i'(v) = v \psi_i'(v)$ for $i = 1, 2$
14 and $v \in \mathbb{R}$. Then, $\eta(\rho, u)$ are convex entropies with respect to (ρ, u) . Moreover, $q(\rho, u)$ is the
15 corresponding entropy flux of $\eta(\rho, u)$.

16 *Proof.* The proof is similar to Proposition 4.1 and we omit it. \square

17 **Remark 4.2.** When $k_1 = k_2 = \frac{1}{2}$ and $\psi_1 = \psi_2 = \psi$, the entropy η defined by (4.19) is equivalent
18 to η_ψ given in (4.10). Recall Definition 4.1. For System (1.18), the entropy is defined by η_g which
19 is one part of (4.10). If we use the counter part η_ψ in (4.10) to define entropy class and entropy
20 solutions of the coupled Burgers system (1.13), we can also obtain global existence of solutions.
21 This can not ensure the uniqueness of entropy solutions. However, if we use the entropies given
22 by (4.19), which can be viewed as a class of entropies modifying η_ψ , to define entropy solutions of
23 the coupled Burgers system (1.13), we can obtain the stability (hence uniqueness) of solutions (see
24 Theorem 4.1).

1 We give the definition of entropy solutions of the coupled Burgers system (1.13).

Definition 4.2. A couple (ρ, u) is called an entropy solution of the coupled Burgers system (1.13) if $\rho \geq 0$ and it satisfies

$$\partial_t \eta(\rho, u) + \partial_x q(\rho, u) \leq 0, \quad (4.21)$$

2 in distribution sense for any convex entropies (η, q) given by (4.19), (4.20).

3 Next, we present an important result about the equivalent relations between entropy solutions
4 of the coupled Burgers system (1.13) and solutions of the decoupled Burgers equations (1.16) and
5 (1.17).

6 **Proposition 4.3.** If (ρ, u) is an entropy solution to the coupled Burgers system (1.13), then $f_{\pm} =$
7 $u \pm \sqrt{\alpha} \rho$ are entropy solutions to the decoupled Burgers equations (1.16) and (1.17). Conversely, if
8 f_{\pm} such that $f_+ \geq f_-$ are entropy solutions to the decoupled Burgers equations (1.16) and (1.17),
9 then $(\rho, u) = \left(\frac{f_+ + f_-}{2}, \frac{f_+ - f_-}{2\sqrt{\alpha}} \right)$ is an entropy solution to the coupled Burgers system (1.13).

Proof. **Step 1.** Assume (ρ, u) is an entropy solution to the coupled Burgers system (1.13). Hence,
the inequality (4.21) holds for any η given by (4.19). For any convex function ψ , let $k_1 = 1$, $k_2 = 0$
and $\psi_1 = \psi$ in (4.19). At this time, the inequality (4.21) gives

$$\partial_t \psi(f_+) + \partial_x \phi(f_+) \leq 0, \quad (4.22)$$

where $\phi'(v) = v\psi'(v)$ and $f_+ = u + \sqrt{\alpha}\rho$. Similarly, when $k_1 = 0$, $k_2 = 1$ and $\psi_2 = \psi$, we can
obtain

$$\partial_t \psi(f_-) + \partial_x \phi(f_-) \leq 0 \quad (4.23)$$

10 in distribution sense. Inequalities (4.22) and (4.23) are exactly the entropy inequalities for the
11 decoupled Burgers equations (1.16) and (1.17). Hence, f_{\pm} are entropy solutions to (1.16) and
12 (1.17).

13 **Step 2.** Let f_{\pm} be an entropy solution of the decoupled Burgers equations (1.16) and (1.17).
14 Due to $f_+ \geq f_-$, we have $\rho = \frac{f_+ - f_-}{2\sqrt{\alpha}} \geq 0$. Moreover, inequality (4.22) holds for any entropy
15 pair (ψ_1, ϕ_1) with $\phi_1'(v) = v\psi_1'(v)$, and inequality (4.23) holds for any entropy pair (ψ_2, ϕ_2) with
16 $\phi_2'(v) = v\psi_2'(v)$. The linear combination of (4.22) and (4.23) with nonnegative coefficients k_1 and
17 k_2 generates the inequality (4.21). Hence, (ρ, u) is an entropy solution to the coupled Burgers
18 system (1.13). □

19
20 Due to the well-posedness of the scalar conservation law (Burgers equation), we have the fol-
21 lowing well-posedness result for the coupled Burgers system (1.13)

22 **Theorem 4.1.** Let $\rho_0(x)$ and $u_0(x)$ be two bounded measurable functions satisfying $\rho_0 \geq 0$. Then

23 (i) There exist a unique entropy solution $(\rho(x, t), u(x, t))$ to the coupled Burgers system (1.13)
24 such that $\rho \geq 0$ and $(\rho, u)|_{t=0} = (\rho_0, u_0)$.

(ii) Let $(\tilde{\rho}, \tilde{u})$ be another entropy solution of the coupled Burgers system (1.13) subject to initial
datum $(\tilde{\rho}_0(x), \tilde{u}_0(x))$ with $\tilde{\rho}_0 \geq 0$. If $u_0 - \tilde{u}_0, \rho_0 - \tilde{\rho}_0 \in L^1(\mathbb{R})$, then $u(\cdot, t) - \tilde{u}(\cdot, t), \rho(\cdot, t) - \tilde{\rho}(\cdot, t) \in$
 $L^1(\mathbb{R})$ and

$$\sqrt{\alpha} \|\rho(\cdot, t) - \tilde{\rho}(\cdot, t)\|_{L^1} + \|u(\cdot, t) - \tilde{u}(\cdot, t)\|_{L^1} \leq 2(\sqrt{\alpha} \|\rho_0 - \tilde{\rho}_0\|_{L^1} + \|u_0 - \tilde{u}_0\|_{L^1}). \quad (4.24)$$

Proof. (i) Consider the decoupled Burgers equations (1.16) and (1.17) with initial datum $f_{\pm}(x, 0) :=$
 $u_0(x) \pm \sqrt{\alpha}\rho_0(x)$. Then, there is a unique entropy solutions $f_{\pm}(x, t)$ to (1.16) and (1.17) respec-
tively. Due to $\rho_0 \geq 0$, we have $f_+(x, 0) \geq f_-(x, 0)$. Hence, from [27, Proposition 2.3.6], we have
 $f_+(x, t) \geq f_-(x, t)$ for any $t > 0$ and $x \in \mathbb{R}$. By Proposition 4.3, there is a unique solution to the
coupled Burgers system (1.13) given by

$$u(x, t) = \frac{f_+(x, t) + f_-(x, t)}{2}, \quad \rho(x, t) = \frac{f_+(x, t) - f_-(x, t)}{2\sqrt{\alpha}}. \quad (4.25)$$

1 Moreover, we have $\rho \geq 0$.

(ii) Let $f_{\pm} := u \pm \sqrt{\alpha}\rho$ and $\tilde{f}_{\pm} := \tilde{u} \pm \sqrt{\alpha}\tilde{\rho}$. Then, f_{\pm} and \tilde{f}_{\pm} are entropy solutions to the decoupled Burgers equations (1.16) and (1.17). By the stability results for scalar conservation law (see [27, Proposition 2.3.6]), we have

$$\begin{aligned} & \sqrt{\alpha}\|\rho(\cdot, t) - \tilde{\rho}(\cdot, t)\|_{L^1} + \|u(\cdot, t) - \tilde{u}(\cdot, t)\|_{L^1} \\ &= \left\| \frac{f_+(\cdot, t) - f_-(\cdot, t)}{2} - \frac{\tilde{f}_+(\cdot, t) - \tilde{f}_-(\cdot, t)}{2} \right\|_{L^1} \\ & \quad + \left\| \frac{f_+(\cdot, t) + f_-(\cdot, t)}{2} - \frac{\tilde{f}_+(\cdot, t) + \tilde{f}_-(\cdot, t)}{2} \right\|_{L^1} \\ &\leq \|f_+(\cdot, 0) - \tilde{f}_+(\cdot, 0)\|_{L^1} + \|f_-(\cdot, 0) - \tilde{f}_-(\cdot, 0)\|_{L^1} \\ &\leq 2(\sqrt{\alpha}\|\rho_0 - \tilde{\rho}_0\|_{L^1} + \|u_0 - \tilde{u}_0\|_{L^1}). \end{aligned}$$

2

□

3 **Remark 4.3.** We remark that f_+ and f_- are entropy solutions to the decoupled Burgers equations
4 (1.16) and (1.17) respectively if and only if there are two positive Radon measures $\mu_+, \mu_- \in \mathcal{M}_+(\mathbb{R})$
5 such that the kinetic functions $\chi_{\pm}(v; \rho, u)$ given by (4.1) are weak solution of the kinetic equations
6 [24]

$$\partial_t \chi_{\pm} + v \partial_x \chi_{\pm} = \partial_v \mu_{\pm}.$$

7 Actually, for an entropy pair (ψ, ϕ) , one has

$$\partial_t \psi(f_{\pm}) + \partial_x \phi(f_{\pm}) = \int_{\mathbb{R}} \psi'(v) (\partial_t \chi_{\pm} + v \partial_x \chi_{\pm}) dv = - \int_{\mathbb{R}} \psi''(v) \mu_{\pm} dv \leq 0$$

8 in distribution sense. For more detailed discussions of using these kinetic density functions to study
9 the coupled Burgers system (1.13) for (ρ, u) , one can refer to [24].

10 **Remark 4.4** (Difference between Definition 4.1 and Definition 4.2). From (4.4), the Hamiltonian
11 H_1^u corresponds to (4.19) for $\psi_1(v) = \psi_2(v) = v^3$ and $k_1 = k_2 = 1/12$. At this time ψ is not convex
12 and hence from Proposition 4.1, we know that H_1^u is not convex with respect to (ρ, u) . When
13 $\rho \geq 0$, we also have H_2^u is convex with respect to (ρ, u) . However, we have

$$H_2^u(\rho, u) = \frac{1}{4} \int_{\mathbb{R}} v^2 \chi_+(v; \rho, u) dv - \frac{1}{4} \int_{\mathbb{R}} v^2 \chi_-(v; \rho, u) dv,$$

14 which is not a proper entropy as in Definition 4.2.

15 Similarly, one can show that H_1^m is not a proper entropy as in Definition 4.1, while H_2^m is a
16 convex entropy for system (1.18).

17 To end this subsection, we give the kinetic formulation for the well known Lax entropy [13].
18 Let the solution of the wave equation (4.8) have the form $\eta(k; \rho, u) = e^{ku} \sigma(k; \rho)$ for some constant
19 parameter $k \neq 0$. Then, equation (4.8) becomes the ODE

$$\sigma_{\rho\rho}(\rho) = \alpha k^2 \sigma(\rho), \quad \sigma(0) = 0, \quad \sigma'(0) = 1.$$

20 Hence, we have

$$\sigma(\rho) = \frac{e^{\sqrt{\alpha}k\rho} - e^{-\sqrt{\alpha}k\rho}}{2\sqrt{\alpha}k}, \quad \eta(k; \rho, u) = \frac{e^{kf_+} - e^{kf_-}}{2\sqrt{\alpha}k}.$$

This yields a family of Lax entropy pairs:

$$\eta(k; \rho, u) = \frac{e^{kf_+} - e^{kf_-}}{2\sqrt{\alpha}k}, \quad q(k; \rho, u) = \frac{(kf_+ - 1)e^{kf_+} - (kf_- - 1)e^{kf_-}}{2\sqrt{\alpha}k^2}. \quad (4.26)$$

21 Note that both η and q are real functions. When $g(v) = e^{kv}$ in (4.15), $\eta_g(\rho, m)$ recovers the Lax
22 entropy given in (4.26).

5 Lagrangian dynamics for (1.13) and its relation with Calogero-Moser model

In this section, we derive the Lagrangian dynamics for the coupled Burgers system (1.13), which recovers the dynamics (1.23) for gas. Moreover, we present a nonlinear spring-mass system (Fermi-Pasta-Ulam-Tsingou model) with nearest-neighbor interactions and its continuum limit yields the Lagrangian dynamics of the coupled Burgers system (1.13).

5.1 Lagrangian dynamics for the coupled Burgers system (1.13)

Consider an initial datum for the coupled Burgers system (1.13):

$$u(x, 0) = u_0(x), \quad \rho(x, 0) = \rho_0(x), \quad x \in \mathbb{R}. \quad (5.1)$$

Assume that initial density function $\rho_0 : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\rho_0(x) > 0$ and the total mass $\|\rho_0\|_{L^1} = 1$. Define the initial cumulative mass distribution function Z_0 :

$$Z_0(x) := \int_{-\infty}^x \rho_0(y) dy \quad \text{for } x \in \mathbb{R}. \quad (5.2)$$

Then, function $Z_0 : \mathbb{R} \rightarrow (0, 1)$ is strictly increasing. Hence, there is an inverse function $X_0 : (0, 1) \rightarrow \mathbb{R}$ such that

$$Z_0(X_0(\xi)) = \xi, \quad X_0(Z_0(x)) = x \quad \text{for } x \in \mathbb{R}, \quad \xi \in (0, 1). \quad (5.3)$$

Moreover, we have

$$Z_0(0) = X_0(0) = 0, \quad \text{and} \quad \frac{1}{X_0'(\xi)} = Z_0'(x) = \rho_0(x) \quad \text{for } \xi = Z_0(x). \quad (5.4)$$

Here, x is the Eulerian coordinates and we take ξ as the Lagrangian coordinates.

Give an Eulerian velocity field $u : \mathbb{R} \times [0, \infty) \rightarrow \mathbb{R}$. Define the flow map $X(\xi, t)$ satisfying

$$\begin{cases} \dot{X}(\xi, t) = u(X(\xi, t), t), & \xi \in (0, 1), \quad t > 0, \\ X(\xi, 0) = X_0(\xi). \end{cases} \quad (5.5)$$

Here, $\dot{X}(\xi, t)$ denotes $\partial_t X(\xi, t)$. Hence, we have $\partial_\xi \dot{X} = \partial_x u \partial_\xi X$ and thus

$$\partial_\xi X(\xi, t) = X_0'(\xi) e^{\int_0^t \partial_x u(X(\xi, s), s) ds} > 0, \quad \xi \in (0, 1). \quad (5.6)$$

Define the density function in Lagrangian coordinates at time t as:

$$\rho(X(\xi, t), t) := \frac{1}{\partial_\xi X(\xi, t)}, \quad (5.7)$$

Hence,

$$\rho(x, t) dx = d\xi \quad \text{and} \quad \partial_t \rho + \partial_x(\rho u) = 0, \quad \rho(x, 0) = \rho_0(x), \quad (5.8)$$

which is the first equation in the coupled Burgers system (1.13). We also have local mass conservation law:

$$\int_{X(\xi_1, t)}^{X(\xi_2, t)} \rho(x, t) dx = \xi_1 - \xi_2 = \int_{X_0(\xi_2)}^{X_0(\xi_1)} \rho_0(x) dx \quad \text{for any } \xi_i \in (0, 1), \quad i = 1, 2.$$

By (5.7), we obtain

$$\partial_x \rho(X(\xi, t), t) = \frac{1}{\partial_\xi X(\xi, t)} \partial_\xi \left(\frac{1}{\partial_\xi X(\xi, t)} \right) = -\frac{\partial_{\xi\xi} X(\xi, t)}{(\partial_\xi X)^3(\xi, t)},$$

which gives

$$-(\alpha\rho\partial_x\rho)(X(\xi, t), t) = \alpha\frac{\partial_{\xi\xi}X(\xi, t)}{(\partial_\xi X)^4(\xi, t)}. \quad (5.9)$$

Set

$$V(\xi, t) := u(X(\xi, t), t). \quad (5.10)$$

Combining (5.9), the coupled Burgers system (1.13) is recast to the Lagrangian dynamics:

$$\begin{cases} \dot{X}(\xi, t) = V(\xi, t), & \xi \in (0, 1), t > 0, \\ \dot{V}(\xi, t) = \alpha\frac{\partial_{\xi\xi}X(\xi, t)}{(\partial_\xi X)^4(\xi, t)} = -\frac{\alpha}{3}\partial_\xi\left(\frac{1}{(\partial_\xi X)^3(\xi, t)}\right), \end{cases} \quad (5.11)$$

subject to initial datum

$$\begin{cases} X(\xi, 0) = X_0(\xi), & \xi \in (0, 1), \\ V(\xi, 0) = u_0(\xi). \end{cases} \quad (5.12)$$

¹ Here, u_0 is given by (5.1) and $X_0(\xi)$ is given by (5.3). Taking derivative of the first equation in ² (5.11) with respect to ξ , we can recover the dynamics (1.23) for gas with $\tau(\xi, t) := X_\xi(\xi, t)$.

Next, we briefly show least action principle for the Lagrangian dynamics (5.11). Corresponding to the total energy $H_2^m(\rho, m)$ given by (3.12), we use Legendre transformation to obtain the Lagrangian functional as

$$\mathcal{L}(\rho, u) = \int_{\mathbb{R}} m\frac{\delta H_2^m}{\delta m} dx - H_2^m(\rho, m) = \int_{\mathbb{R}} \left(\frac{1}{2}\rho u^2 - \frac{\alpha}{6}\rho^3\right) dx$$

³ The momentum m is recovered by taking the variation of \mathcal{L} with respect to u :

$$m = \frac{\delta\mathcal{L}}{\delta u} = \rho u.$$

The action is defined by

$$\mathcal{A}(X) = \frac{1}{2}\int_0^1\int_{\mathbb{R}} \left(\rho u^2 - \frac{\alpha}{3}\rho^3\right) dx dt = \frac{1}{2}\int_0^1\int_0^1 \left(\dot{X}^2(\xi, t) - \frac{\alpha}{3(\partial_\xi X)^2(\xi, t)}\right) d\xi dt. \quad (5.13)$$

Next, consider two increasing functions for $\xi \in [0, 1]$: $X(\xi, 0) = X_0(\xi)$ and $X(\xi, 1) = X_1(\xi)$. We formally show that the coupled Burgers system (1.13) corresponds to a critical path of the action $\mathcal{A}(X)$ in some manifold connecting X_0 and X_1 for $t \in [0, 1]$. For any $Y \in C_c^\infty((0, 1) \times (0, 1))$, we have

$$\begin{aligned} \int_0^1\int_0^1 \frac{\delta\mathcal{A}}{\delta X} \cdot Y d\xi dt &= \lim_{\epsilon \rightarrow 0} \frac{\mathcal{A}(X + \epsilon Y) - \mathcal{A}(X)}{\epsilon} \\ &= \frac{1}{2}\int_0^1\int_0^1 \left(2\dot{X}\dot{Y} + \frac{2\alpha}{3(\partial_\xi X)^3}\partial_\xi Y\right) d\xi dt = \int_0^1\int_0^1 \left[-\ddot{X} - \partial_\xi\left(\frac{\alpha}{3(\partial_\xi X)^3}\right)\right] Y d\xi dt. \end{aligned}$$

This gives

$$\frac{\delta\mathcal{A}}{\delta X} = -\ddot{X} - \partial_\xi\left(\frac{\alpha}{3(\partial_\xi X)^3}\right).$$

Take $\frac{\delta\mathcal{A}}{\delta X} = 0$, and we have

$$\ddot{X} - \alpha\frac{\partial_{\xi\xi}X}{(\partial_\xi X)^4} = 0, \quad (5.14)$$

⁴ which corresponds to the Lagrangian dynamics (5.11).

5.2 A spring-mass system with nearest-neighbor interactions

In this subsection, we present a local interaction model for N masses and show that the Lagrangian dynamics system (5.11) is exactly the continuum limit equation of this model. For N ordered masses $x_1(t) < \dots < x_N(t)$, each mass is evolved by a force generated by interactions between nearest neighbors and the model is described by

$$\begin{cases} \dot{x}_j(t) = v_j(t), & 1 \leq j \leq N, \\ \dot{v}_j(t) = \frac{\alpha}{3N^2} \left[\frac{1}{(x_{j+1}(t) - x_j(t))^3} + \frac{1}{(x_{j-1}(t) - x_j(t))^3} \right]. \end{cases} \quad (5.15)$$

Here we assume

$$x_0 = x_{N+1} = +\infty, \quad \text{and} \quad \frac{1}{(x_0(t) - x_1(t))^3} = \frac{1}{(x_{N+1}(t) - x_N(t))^3} = 0. \quad (5.16)$$

The masses accelerated by an repulsive force if $\alpha < 0$. While $\alpha > 0$, the masses attract each other. System (5.15) is a Hamiltonian system corresponding to the Hamiltonian functional:

$$H(x, p) = \frac{N}{2} \sum_{j=1}^N p_j^2 - \frac{\alpha}{12N^3} \sum_{j=1}^N \sum_{k=j\pm 1}^N \frac{1}{(x_j - x_k)^2}. \quad (5.17)$$

Momentum p_i equals to mass $1/N$ times velocity v_i which means $v_i = Np_i$. Hence, (5.15) equals to

$$\begin{cases} \dot{x}_j(t) = \partial_{p_j} H, & 1 \leq j \leq N, \\ \dot{p}_j(t) = -\partial_{x_j} H. \end{cases}$$

Model (5.15) describes local interactions between masses and their nearest-neighbors, which is a special case of the Fermi-Pasta-Ulam-Tsingou lattice system. We compare (5.15) with another Fermi-Pasta-Ulam-Tsingou lattice system, Toda lattice, given by the system of ordinary differential equations

$$\frac{d^2 q_j}{dt^2} = e^{q_{j+1} - q_j} - e^{q_j - q_{j-1}}, \quad j \in \mathbb{Z}. \quad (5.18)$$

Note that Toda lattice is an integrable system. We do not know whether System (5.15) is an integrable system or not. However, if each mass interacts with all the other masses with the same manner, we can obtain an integrable global interaction model, the Calogero-Moser model (see Remark 5.1).

Next, we formally derive the continuum limit of the local interaction mass system. To do this, we assume the masses initially distribute uniformly and $x_j(t) = X(\xi, t)$, $x_{j+1}(t) = X(\xi + 1/N, t)$ and $x_{j-1}(t) = X(\xi - 1/N, t)$ for some $\xi \in (0, 1)$ and $2 \leq j \leq N - 1$. Using Taylor expansion, we have

$$x_{j+1}(t) - x_j(t) = \partial_\xi X(\xi, t)N^{-1} + \frac{1}{2}\partial_{\xi\xi} X(\xi, t)N^{-2} + \frac{1}{6}\partial_{\xi\xi\xi} X(\xi, t)N^{-3} + O((N^{-4}),$$

and

$$x_{j-1}(t) - x_j(t) = -\partial_\xi X(\xi, t)N^{-1} + \frac{1}{2}\partial_{\xi\xi} X(\xi, t)N^{-2} - \frac{1}{6}\partial_{\xi\xi\xi} X(\xi, t)N^{-3} + O((N^{-4}).$$

Hence, we can obtain

$$\begin{aligned}
& \frac{\alpha}{3N^2} \left[\frac{1}{(x_{j+1}(t) - x_j(t))^3} + \frac{1}{(x_{j-1}(t) - x_j(t))^3} \right] \\
&= \frac{\alpha}{3N^2} \frac{(x_{j-1} + x_{j+1} - 2x_j) \left[(x_{j-1} - x_j)^2 + (x_{j+1} - x_j)^2 - (x_{j-1} - x_j)(x_{j+1} - x_j) \right]}{(x_{j+1} - x_j)^3 (x_{j-1} - x_j)^3} \\
&= \frac{\alpha}{3N^2} \frac{\left[\partial_{\xi\xi} X(\xi, t) N^{-2} + O(N^{-4}) \right] \cdot \left[3(\partial_{\xi} X)^2(\xi, t) N^{-2} + O(N^{-3}) \right]}{(\partial_{\xi} X)^6(\xi, t) N^{-6} + O(N^{-7})} \\
&= \frac{\alpha}{3N^2} \frac{N^{-4} \left[\partial_{\xi\xi} X(\xi, t) + O(N^{-2}) \right] \cdot \left[3(\partial_{\xi} X)^2(\xi, t) + O(N^{-1}) \right]}{N^{-6} (\partial_{\xi} X)^6(\xi, t) + O(N^{-1})} \\
&= \alpha \frac{\partial_{\xi\xi} X(\xi, t) + O(N^{-1})}{(\partial_{\xi} X)^4(\xi, t) + O(N^{-1})}.
\end{aligned}$$

1 Let $N \rightarrow \infty$ and we obtain

$$\lim_{N \rightarrow \infty} \frac{\alpha}{3N^2} \left[\frac{1}{(x_{j+1}(t) - x_j(t))^3} + \frac{1}{(x_{j-1}(t) - x_j(t))^3} \right] = \alpha \frac{\partial_{\xi\xi} X(\xi, t)}{(\partial_{\xi} X)^4(\xi, t)}.$$

2 This gives the continuum coupled Burgers system in Lagrangian coordinate (5.11).

Remark 5.1. If each mass interact with all the other masses with the same manner (the force between each pair of two masses are reciprocal proportion to the cubic of distance between them), we can obtain an integrable global interaction model, the Calogero-Moser model [21]:

$$\begin{cases} \dot{x}_j(t) = v_j(t), \\ \dot{v}_j(t) = \frac{4\alpha}{N^2 \pi^2} \sum_{k=1, k \neq j}^N \frac{1}{(x_j(t) - x_k(t))^3}, \quad 1 \leq j \leq N. \end{cases} \quad (5.19)$$

The coefficients of (5.19) are different from the coefficients in (5.15). System (5.19) is also a Hamiltonian system and the rescaled ($p_j = v_j/N$) Hamiltonian is given by

$$\tilde{H}(x, q) = \frac{N}{2} \sum_{j=1}^N p_j^2 + \frac{\alpha}{2N^3 \pi^2} \sum_{j=1}^N \sum_{k \neq j}^N \frac{1}{(x_j - x_k)^2}. \quad (5.20)$$

3 By using the Euler-MacLaurin asymptotic expansion for the Riemann integral of functions, Menon
4 [19] showed that System (5.11) is the $N \rightarrow \infty$ limit of the Calogero-Moser system corresponding
5 to the rescaled Hamiltonian (5.20). As shown by [19, Eqs. (5.13),(5.26)], the Hamiltonian \tilde{H}
6 corresponds to the total energy H_2^m (see (3.15)) of System (1.18).

7 Acknowledgements

8 We are grateful to Govind Menon for some helpful discussions. We would like to thank the support
9 by the National Science Foundation under grants DMS 1514826 and 1812573 (JGL).

10 References

- 11 [1] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows: in metric spaces and in the space of*
12 *probability measures*. Springer Science & Business Media, 2008.
- 13 [2] R. J. Berman and M. Önnheim. Propagation of chaos for a class of first order models with
14 singular mean field interactions. *SIAM J. Math. Anal.*, 51(1):159–196, 2019.
- 15 [3] J. A. Carrillo, L. C.F. Ferreira, and J. C. Precioso. A mass-transportation approach to a one
16 dimensional fluid mechanics model with nonlocal velocity. *Adv. Math.*, 231(1):306–327, 2012.

- 1 [4] A. Castro and D. Córdoba. Global existence, singularities and ill-posedness for a nonlocal
2 flux. *Adv. Math.*, 219(6):1916–1936, 2008.
- 3 [5] E. Cépa and D. Lépingle. Diffusing particles with electrostatic repulsion. *Probab. Theory
4 Related Fields*, 107(4):429–449, 1997.
- 5 [6] A. Córdoba, D. Córdoba, and M. A. Fontelos. Formation of singularities for a transport
6 equation with nonlocal velocity. *Ann. of Math.*, pages 1377–1389, 2005.
- 7 [7] F.J. Dyson. A Brownian-motion model for the eigenvalues of a random matrix. *J. Math.
8 Phys.*, 3(6):1191–1198, 1962.
- 9 [8] L. Erdos and H.-T. Yau. Dynamical approach to random matrix theory. *Courant Lecture
10 Notes in Mathematics*, 28, 2017.
- 11 [9] D. J. Gross and A. Matytsin. Some properties of large-N two-dimensional Yang-Mills theory.
12 *Nuclear Physics B*, 437(3):541–584, 1995.
- 13 [10] R. Jordan, D. Kinderlehrer, and F. Otto. The variational formulation of the Fokker–Planck
14 equation. *SIAM J. Math. Anal.*, 29(1):1–17, 1998.
- 15 [11] R. Kenyon and A. Okounkov. Limit shapes and the complex Burgers equation. *Acta Math.*,
16 199(2):263–302, 2007.
- 17 [12] S. G. Krantz and H. R. Parks. *The implicit function theorem: history, theory, and applications*.
18 Springer Science & Business Media, 2012.
- 19 [13] Peter D Lax. Hyperbolic systems of conservation laws ii. *Comm. Pure Appl. Math.*, 10(4):537–
20 566, 1957.
- 21 [14] Lei Li, Jian-Guo Liu, and Pu Yu. On the mean field limit for brownian particles with coulomb
22 interaction in 3d. *Journal of Mathematical Physics*, 60(11):111501, 2019.
- 23 [15] P. L Lions, B Perthame, and E Tadmor. Kinetic formulation of the isentropic gas dynamics
24 and p-systems. *Comm. Math. Phys.*, 163(2):415–431, 1994.
- 25 [16] J.-G. Liu and R. Yang. Propagation of chaos for large Brownian particle system with Coulomb
26 interaction. *Research in the Mathematical Sciences*, 3(1):40, 2016.
- 27 [17] Jian-Guo Liu and Robert Pego. On generating functions of hausdorff moment sequences.
28 *Transactions of the American Mathematical Society*, 368(12):8499–8518, 2016.
- 29 [18] F. Magri. A simple model of the integrable Hamiltonian equation. *J. math. phys*, 19(5):1156–
30 1162, 1978.
- 31 [19] G. Menon. The complex Burgers equation, the HCIZ integral and the Calogero-Moser system.
32 *preprint*, <http://www.dam.brown.edu/people/menon/talks/cmsa.pdf>.
- 33 [20] G. Menon. Lesser known miracles of Burgers equation. *Acta Math. Sci.*, 32(1):281–294, 2012.
- 34 [21] J. Moser. Three integrable Hamiltonian systems connected with isospectral deformations. In
35 *Surveys in Applied Mathematics*, pages 235–258. Elsevier, 1976.
- 36 [22] P. J. Olver and Y. Nutku. Hamiltonian structures for systems of hyperbolic conservation laws.
37 *J. math. phys*, 29(7):1610–1619, 1988.
- 38 [23] J. N. Pandey. *The Hilbert Transform of Schwartz Distributions and Applications*. Wiley, New
39 York, 1996.
- 40 [24] B. Perthame and A. E. Tzavaras. Kinetic formulation for systems of two conservation laws
41 and elastodynamics. *Arch. Ration. Mech. Anal.*, 155(1):1–48, 2000.
- 42 [25] L. Rogers and Z. Shi. Interacting Brownian particles and the wigner law. *Probab. Theory
43 Related Fields*, 95(4):555–570, 1993.

- 1 [26] S. Salsa. *Partial differential equations in action: from modelling to theory*, volume 99.
2 Springer, 2016.
- 3 [27] D. Serre. Systems of conservation laws 1. *Systems of Conservation Laws*, 255(7):286, 1999.
- 4 [28] L. Silvestre and V. Vicol. On a transport equation with nonlocal drift. *Transactions of the*
5 *American Mathematical Society*, 368(9):6159–6188, 2015.
- 6 [29] T. Tao. *Topics in Random Matrix Theory*. American Mathematical Society, 2012.
- 7 [30] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. *Annals*
8 *Math.*, 62:548–564, 1955.
- 9 [31] N. M. Zubarev and EA Karabut. Exact local solutions for the formation of singularities on
10 the free surface of an ideal fluid. *JETP Letters*, 107(7):412–417, 2018.

11 A Proof of Theorem 2.1

Consider the initial datum given by (2.8). Direct calculation shows that

$$\begin{aligned} f_0(z) &= \frac{1}{\pi} \int_{\mathbb{R}} \frac{\rho_0(s)}{z-s} ds = \frac{1}{\pi} \int_{\mathbb{R}} \frac{x-s}{y^2+(x-s)^2} \rho_0(s) ds - \frac{i}{\pi} \int_{\mathbb{R}} \frac{y}{y^2+(x-s)^2} \rho_0(s) ds \\ &=: R\rho_0(x, y) - iP\rho_0(x, y), \end{aligned}$$

where $P\rho_0(x, y)$ and $R\rho_0(x, y)$ are given by the convolution of ρ_0 with the Poisson kernel and the conjugate Poisson kernel given by

$$P_y(x) := \frac{1}{\pi} \frac{y}{y^2+x^2} \quad \text{and} \quad R_y(x) := \frac{1}{\pi} \frac{x}{y^2+x^2}. \quad (\text{A.1})$$

12 Furthermore, we have

$$\lim_{y \rightarrow 0^+} [R\rho_0(x, y) - iP\rho_0(x, y)] = H\rho_0(x) - i\rho_0(x) \text{ for a.e. } x \in \mathbb{R}.$$

13 Recall that the following properties of Poisson kernel:

14 (i) If $h \in L^2(\mathbb{R})$, then

$$Rh(x, y) = PHh(x, y) \text{ on } \mathbb{R}_+^2.$$

15 (ii) If $h \in L^\infty(\mathbb{R})$ and is vanishing at infinity, then

$$\lim_{y \rightarrow +\infty} Ph(x, y) = 0, \quad x \in \mathbb{R},$$

16 and

$$\lim_{x \rightarrow \pm\infty} Ph(x, y) = 0, \quad y \geq 0.$$

17 (iii) If $h \in L^\infty(\mathbb{R})$, then $Ph(x, y)$ is a bounded function on \mathbb{R}_+^2 .

18 Next, we prove the existence and uniqueness of \mathbb{C}_+ -holomorphic solutions to (2.10) by the
19 characteristics method. Consider the characteristics given by

$$\frac{d}{dt} Z(w, t) = g(Z(w, t), t), \quad Z(w, 0) = w \in \mathbb{C}_+. \quad (\text{A.2})$$

20 Then,

$$\frac{d^2}{dt^2} Z(w, t) = \frac{d}{dt} g(Z(w, t), t) = [\partial_t g + g \partial_z g](Z(w, t), t) = \gamma^2 Z(w, t),$$

21 with initial date

$$Z(w, 0) = w, \quad \left. \frac{d}{dt} Z(w, t) \right|_{t=0} = g_0(w), \quad w \in \mathbb{C}_+.$$

Equation (A.2) gives the following complex trajectories:

$$Z(w, t) = \begin{cases} w \cosh \gamma t + \frac{1}{\gamma} g_0(w) \sinh \gamma t, & \gamma > 0, \\ g_0(w)t + w = f_0(w)t + w, & \gamma = 0. \end{cases} \quad (\text{A.3})$$

1 Here, we only treat the case for $\gamma > 0$ where the convergence to the steady state for analytical
2 solutions happens. For the well-posedness results of the case $\gamma = 0$, one can refer to [4]. Let

$$Z(w, t) = Z_1(x, y, t) + iZ_2(x, y, t), \quad w = x + iy \in \mathbb{C}_+,$$

and we have

$$Z_1(x, y, t) = x \cosh \gamma t + \frac{\pi}{\gamma} R\rho_0(x, y) \sinh \gamma t - x \sinh \gamma t = xe^{-\gamma t} + \frac{\pi}{\gamma} R\rho_0(x, y) \sinh \gamma t, \quad (\text{A.4})$$

$$Z_2(x, y, t) = y \cosh \gamma t - \frac{\pi}{\gamma} P\rho_0(x, y) \sinh \gamma t - y \sinh \gamma t = ye^{-\gamma t} - \frac{\pi}{\gamma} P\rho_0(x, y) \sinh \gamma t. \quad (\text{A.5})$$

3 Because the initial data $g_0(w)$ in (2.10) is a \mathbb{C}_+ -holomorphic function, $Z(w, t)$ given by (A.3) is
4 \mathbb{C}_+ -holomorphic of w for any $t \geq 0$. Next, we present a lemma to show that for any fixed time
5 $t > 0$ the backward characteristics of (A.3) are well defined on the set $\overline{\mathbb{C}_+}$. We have:

6 **Lemma A.1.** *Let $0 < \rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ with $s > 1/2$. For fixed $t_0 > 0$ and fixed $Z = Z_1 + iZ_2 \in$
7 $\overline{\mathbb{C}_+}$, there exists a unique $w = x + iy \in \mathbb{C}_+$ such that (A.4) and (A.5) hold.*

8 *Proof.* Given $t_0 > 0$, denote

$$a := e^{-\gamma t_0}, \quad b := \frac{\pi}{\gamma} \sinh \gamma t_0.$$

Then (A.4) and (A.5) become

$$Z_1 = ax + bR\rho_0(x, y), \quad Z_2 = ay - bP\rho_0(x, y).$$

9 **Step 1.** In this step, we prove that for any x , there exists a unique $y > 0$ satisfies (A.5) for
10 $Z_2 \geq 0$ and $t_0 > 0$.

Because $P\rho_0(x, y) > 0$ is a bounded function on \mathbb{R}_+^2 , by the property of Poisson kernel we have

$$\lim_{y \rightarrow +\infty} Z_2(x, y, t_0) = +\infty, \quad \lim_{y \rightarrow 0^+} Z_2(x, y, t_0) = -b\rho_0(x) < 0. \quad (\text{A.6})$$

11 Hence, for any fixed $Z_2 \geq 0$, there exists a point $y > 0$ depending on x such that

$$Z_2 = ay - bP\rho_0(x, y).$$

Now we prove that y is unique. Suppose that there exist $y_1 > y_2$ such that

$$\begin{aligned} Z_2 &= ay_1 - bP\rho_0(x, y_1), \\ Z_2 &= ay_2 - bP\rho_0(x, y_2), \end{aligned}$$

12 which implies

$$y_1, y_2 > Z_2/a \quad \text{and} \quad \frac{P\rho_0(x, y_1)}{y_1 - Z_2/a} = \frac{P\rho_0(x, y_2)}{y_2 - Z_2/a}.$$

13 Because function

$$h(y) = \frac{y}{y - Z_2/a} \cdot \frac{1}{y^2 + (x - s)^2}$$

14 is a decreasing function for $y > Z_2/a$, we obtain a contradiction.

Now we denote by $y_{Z_2}(x)$ the solution of (A.5) with fixed $Z_2 \geq 0$, $t_0 > 0$ and $x \in \mathbb{R}$. Hence,
we obtain

$$ay_{Z_2}(x) - Z_2 = bP\rho_0(x, y_{Z_2}(x)). \quad (\text{A.7})$$

Step 2. In this step, we prove there exists a unique x satisfies (A.4) for fixed Z_1, Z_2 and t_0 . Taking derivative of (A.7) with respect to x gives

$$\frac{d}{dx}y_{Z_2}(x) = \frac{\partial_x P\rho_0(x, y_{Z_2}(x))}{a/b - \partial_y P\rho_0(x, y_{Z_2}(x))}. \quad (\text{A.8})$$

Since $\rho_0 \in H^s(\mathbb{R}) \cap L^1(\mathbb{R})$ ($s > 1/2$), it follows that $H\rho_0 \in L^\infty(\mathbb{R})$ and therefore $R\rho_0 = PH\rho_0$ is a bounded function over \mathbb{R}_+^2 . Furthermore,

$$\lim_{x \rightarrow \pm\infty} [ax + bR\rho_0(x, y_{Z_2}(x))] = \pm\infty. \quad (\text{A.9})$$

1 Hence, for any $Z_1 \in \mathbb{R}$, we can find a $x \in \mathbb{R}$ such that

$$Z_1 = ax + bR\rho_0(x, y_{Z_2}(x)).$$

2 To prove the uniqueness, we only have to prove the following function

$$q(x) = ax + bR\rho_0(x, y_{Z_2}(x)),$$

is an increasing function. By using (A.8) and the Cauchy-Riemann equations

$$\partial_x R\rho_0 = -\partial_y P\rho_0, \quad \partial_x P\rho_0 = \partial_y R\rho_0, \quad (\text{A.10})$$

3 and taking derivative of $q(x)$ gives

$$\frac{d}{dx}q(x) = \frac{b(a/b + \partial_x R\rho_0)^2 + (\partial_x P\rho_0)^2}{a/b + \partial_x R\rho_0}(x, y_{Z_2}(x)).$$

To prove the increasing of $q(x)$, it is sufficient to prove

$$a/b + \partial_x R\rho_0(x, y) > 0 \quad (\text{A.11})$$

4 for any $(x, y) \in \mathbb{R}_+^2$ satisfying $ay - bP\rho_0(x, y) \geq 0$ and $y > 0$. Suppose that

$$a/b + \partial_x R\rho_0(x_0, y_0) \leq 0$$

for some point $(x_0, y_0) \in \mathbb{R}_+^2$ with $ay_0 - bP\rho_0(x_0, y_0) \geq 0$. Then, we have

$$\begin{aligned} -a/b \geq \partial_x R\rho_0(x_0, y_0) &= \frac{1}{\pi} \int_{\mathbb{R}} \frac{y_0^2 - (x_0 - s)^2}{[y_0^2 + (x_0 - s)^2]^2} \rho_0(s) ds > \frac{1}{\pi} \int_{\mathbb{R}} \frac{-y_0^2 - (x_0 - s)^2}{[y_0^2 + (x_0 - s)^2]^2} \rho_0(s) ds \\ &= \frac{1}{\pi} \int_{\mathbb{R}} \frac{-1}{y_0^2 + (x_0 - s)^2} \rho_0(s) ds = -\frac{P\rho_0(x_0, y_0)}{y_0}, \end{aligned}$$

5 which implies a contradiction:

$$ay_0 - bP\rho_0(x_0, y_0) < 0.$$

6

□

7 From the above lemma, we know that the backward characteristics are well defined. More
8 importantly, for any $Z \in \overline{\mathbb{C}_+}$ the initial point w must be interior point in \mathbb{C}_+ . For any $t \geq 0$, we
9 denote the backward characteristics as:

$$Z^{-1}(\cdot, t) : \overline{\mathbb{C}_+} \rightarrow \mathbb{C}_+.$$

10 From the uniqueness in Lemma A.1, $Z^{-1}(\cdot, t)$ is an 1 – 1 map.

11 *Proof of Theorem 2.1.* For simplicity, we only consider the case $\gamma = 1$. The proof for arbitrary
12 $\gamma > 0$ is similar.

13 **Step 1.** Proof of (i). From Lemma A.1, we have $\overline{\mathbb{C}_+} \subset \{Z(w, t) : w \in \mathbb{C}_+\}$ and $Z^{-1}(\cdot, t)$ is
14 well defined on $\overline{\mathbb{C}_+}$ for any fixed time $t > 0$. Denote the preimage of $Z(\cdot, t)$ as:

$$Z^{-1}(\overline{\mathbb{C}_+}, t) := \{w \in \mathbb{C}_+; Z(w, t) \in \overline{\mathbb{C}_+}\}.$$

1 Denote

$$a(t) := e^{-t}, \quad b(t) := \pi \sinh t.$$

For $(x, y) \in \mathbb{R}_+^2$ and $Z_2(x, y, t) \geq 0$, by the Cauchy-Riemann equation (A.10), we have

$$\begin{aligned} |Z_w(w, t)| &= \left| \frac{\partial(Z_1, Z_2)}{\partial(x, y)} \right| (x, y) = \begin{vmatrix} \partial_x Z_1 & \partial_y Z_1 \\ \partial_x Z_2 & \partial_y Z_2 \end{vmatrix} = \begin{vmatrix} a(t) + b(t)\partial_x R\rho_0 & b(t)\partial_y R\rho_0 \\ -b\partial_x P\rho_0 & a(t) - b(t)\partial_y P\rho_0 \end{vmatrix} \\ &= \left[a(t) + b(t)\partial_x R\rho_0 \right]^2 + \left[b(t)\partial_x P\rho_0 \right]^2 \Big|_{(x, y)} > 0. \end{aligned} \quad (\text{A.12})$$

2 Due to (A.6) and (A.9), we obtain

$$|Z(w, t)| \rightarrow +\infty \quad \text{as} \quad |w| \rightarrow +\infty.$$

3 which means $Z(\cdot, t)$ is proper [12, Definition 6.2.2]. By the Hadamard's global inverse function
4 theorem [12, Theorem 6.2.8], there exists a inverse function $Z^{-1}(\cdot, t)$ such that

$$Z^{-1}(\cdot, t) : \overline{\mathbb{C}_+} \rightarrow Z^{-1}(\overline{\mathbb{C}_+}, t)$$

5 is a bijection. We also know Z^{-1} is $\overline{\mathbb{C}_+}$ -holomorphic since Z is \mathbb{C}_+ -holomorphic. Moreover, for
6 any $z \in \overline{\mathbb{C}_+}$, there exists $w = Z^{-1}(z, t) \in \mathbb{C}_+$. Due to $z = Z(Z^{-1}(z, t), t) \in \overline{\mathbb{C}_+}$ and $|Z_w(w, t)| \neq 0$
7 (by (A.12)), we have

$$\partial_t Z^{-1}(z, t) = -\frac{\partial_t Z(w, t)}{\partial_w Z(w, t)}, \quad w = Z^{-1}(z, t).$$

Because of (A.3), we know $\frac{\partial^k}{\partial t^k} Z(w, t)$ is \mathbb{C}_+ -holomorphic for any positive integer k . Hence,
 $\frac{\partial^k}{\partial t^k} Z^{-1}(z, t)$ is $\overline{\mathbb{C}_+}$ -holomorphic for any positive integer k . From (A.3), we have

$$z = Z^{-1}(z, t) \cosh t + g_0(Z^{-1}(z, t)) \sinh t, \quad z \in \overline{\mathbb{C}_+}. \quad (\text{A.13})$$

8 By (A.2), we obtain

$$g(Z(w, t), t) = \frac{d}{dt} Z(w, t) = w \sinh t + g_0(w) \cosh t.$$

Hence,

$$g(z, t) = Z^{-1}(z, t) \sinh t + g_0(Z^{-1}(z, t)) \cosh t, \quad (\text{A.14})$$

9 which is a $\overline{\mathbb{C}_+}$ -holomorphic solution to the complex Burgers equation (2.10) satisfying $g(z, 0) =$
10 $g_0(z)$. Moreover, due to the time regularity for $Z^{-1}(z, t)$, we know that $\frac{\partial^k}{\partial t^k} g(z, t)$ is $\overline{\mathbb{C}_+}$ -holomorphic
11 for any positive integer k and $t > 0$.

Step 2. Proof of (ii). A $\overline{\mathbb{C}_+}$ -holomorphic solution to (2.7) is given by

$$f(z, t) := g(z, t) + z, \quad z \in \overline{\mathbb{C}_+}, \quad t > 0, \quad (\text{A.15})$$

with initial datum $f_0(z) = \pi R\rho_0(x, y) - i\pi P\rho_0(x, y)$, $z = x + iy \in \mathbb{C}_+$. Combining (A.13) and
(A.14), we obtain

$$f(z, t) = f_0(Z^{-1}(z, t))e^t \quad \text{and} \quad z = e^{-t}Z^{-1}(z, t) + f_0(Z^{-1}(z, t)) \sinh t, \quad z \in \overline{\mathbb{C}_+}. \quad (\text{A.16})$$

12 Consider the trace of $f(z, t)$ on the real line and define:

$$f(x, t) =: u(x, t) - i\pi\rho(x, t).$$

13 Due to Lemma A.1, for any $x \in \mathbb{R}$, we have $Z^{-1}(x, t) =: a_x + ib_x \in \mathbb{C}_+$ with some positive real
14 number $b_x > 0$. From (A.16), we have

$$f(x, t) = f_0(a_x + ib_x)e^t = \pi R\rho_0(a_x, b_x)e^t - i\pi P\rho_0(a_x, b_x)e^t$$

Therefore,

$$\rho(x, t) = P\rho_0(a_x, b_x)e^t > 0, \quad x \in \mathbb{R}. \quad (\text{A.17})$$

- 1 Hence, $\rho(x, t)$ is a positive analytical solution to the Dyson equation (1.4). Moreover, by the
 2 uniqueness of solutions to the characteristics equation (2.10) we know analytical solutions to the
 3 Dyson equation (1.4) is unique.

The energy estimate (1.21) follows from

$$\begin{aligned} \frac{d}{dt}E(\rho) &= \int_{\mathbb{R}} \frac{\delta E}{\delta \rho} \cdot \partial_t \rho \, dx = - \int_{\mathbb{R}} \rho \left| \partial_x \left(\frac{\delta E}{\delta \rho} \right) \right|^2 \, dx \\ &= - \int_{\mathbb{R}} \rho(x, t) |\gamma x - \pi H \rho(x, t)|^2 \, dx. \end{aligned}$$

- 4 For (2.12), direct calculations show that $\|\rho(t)\|_{L^1(\mathbb{R})} = \|\rho_0\|_{L^1(\mathbb{R})}$. Multiplying (1.4) by x^2 and
 5 taking integral yield

$$\frac{d}{dt} \int_{\mathbb{R}} x^2 \rho(x, t) \, dx = 2\pi \int_{\mathbb{R}} x \rho H \rho \, dx - 2 \int_{\mathbb{R}} x^2 \rho(x, t) \, dx = \|\rho_0\|_{L^1}^2 - 2 \int_{\mathbb{R}} x^2 \rho(x, t) \, dx,$$

which implies (2.12). Inequality (2.14) follows from Grönwall's inequality gives and the following estimate

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} \rho \log \rho \, dx &= \int_{\mathbb{R}} \partial_t \rho (\log \rho + 1) \, dx = \int_{\mathbb{R}} -(\rho H \rho + \gamma x \rho)_x (\log \rho + 1) \, dx \\ &= \int_{\mathbb{R}} (H \rho - \gamma x) \rho_x \, dx = -\|(-\Delta)^{1/4} \rho\|_{L^2}^2 + \gamma \|\rho_0\|_{L^1}. \end{aligned}$$

- 6 **Step 3.** We prove (iii) following the idea of [25]. Recall formula (A.16). For fixed $z \in \overline{\mathbb{C}_+}$,
 7 denote

$$z_r(t) + iz_i(t) := e^{-t} Z^{-1}(z, t).$$

- 8 Next, we prove that $z_r(t) + iz_i(t)$ converges to a point $w = z_r^* + iz_i^* \in \mathbb{C}_+$ as $t \rightarrow \infty$. To this end,
 9 we first prove $|z_r(t)|$ and $z_i(t)$ are all bounded from above and below uniformly in time t .

- 10 Because

$$f_0(Z^{-1}(z, t)) = \pi R \rho_0(e^t z_r(t), e^t z_i(t)) - i\pi P \rho_0(e^t z_r(t), e^t z_i(t)),$$

by (A.16), we have

$$z = z_r(t) + \pi R \rho_0(e^t z_r(t), e^t z_i(t)) \sinh t + i [z_i(t) - \pi P \rho_0(e^t z_r(t), e^t z_i(t)) \sinh t]. \quad (\text{A.18})$$

- 11 Due to $\pi P \rho_0(e^t z_r(t), e^t z_i(t)) \sinh t \geq 0$, we have

$$z_i(t) \geq \Im(z) > 0.$$

Moreover, we have

$$\begin{aligned} \Im(z) &= z_i(t) - \pi P \rho_0(e^t z_r(t), e^t z_i(t)) \sinh t \\ &= z_i(t) - \int_{\mathbb{R}} \frac{e^t z_i(t)}{e^{2t} z_i^2(t) + (e^t z_r(t) - s)^2} \rho_0(s) \, ds \sinh t \\ &\geq z_i(t) - \int_{\mathbb{R}} \frac{e^{2t} z_i(t)}{2e^{2t} z_i^2(t) + 2(e^t z_r(t) - s)^2} \rho_0(s) \, ds \\ &\geq z_i(t) - \frac{1}{z_i(t)}, \end{aligned}$$

- 12 which implies

$$z_i(t) \leq \Im(z) + 1.$$

- 13 Hence, $z_i(t)$ is bounded as

$$0 < \Im(z) \leq z_i(t) \leq \Im(z) + 1$$

1 Next, we prove

$$\sup_{t \geq 0} |z_r(t)| < +\infty.$$

2 We prove this by a contradiction argument. If there exists $t_n \rightarrow \infty$ such that $z_r(t_n) \rightarrow \infty$, then
 3 by the dominated convergence theorem we have

$$\pi R \rho_0(e^{t_n} z_r(t_n), e^{t_n} z_i(t_n)) \sinh t_n = \int_{\mathbb{R}} \frac{e^{t_n} z_r(t_n) - s}{e^{2t_n} z_i^2(t_n) + (e^{t_n} z_r(t_n) - s)^2} dx \sinh t_n \rightarrow 0, \quad n \rightarrow \infty.$$

4 By (A.18), we obtain a contradiction that

$$\Re(z) = z_r(t_n) + \pi R \rho_0(e^{t_n} z_r(t_n), e^{t_n} z_i(t_n)) \sinh t_n \rightarrow \infty.$$

5 Since $|z_r(t)|$ and $z_i(t)$ are bounded, there exist $t_n \rightarrow \infty$ and two constant $z_r^*, z_i^* > 0$ such that

$$z_r(t_n) \rightarrow z_r^*, \quad z_i(t_n) \rightarrow z_i^*, \quad n \rightarrow \infty.$$

6 Let $w := z_r^* + iz_i^*$. For any $s \in \mathbb{R}$, we have

$$\frac{e^{t_n} z_r(t_n) - s}{e^{2t_n} z_i^2(t_n) + (e^{t_n} z_r(t_n) - s)^2} \sinh t_n \rightarrow \frac{z_r^*}{2(z_i^*)^2 + 2(z_r^*)^2}, \quad n \rightarrow \infty.$$

Then, by the dominated convergence theorem we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \pi R \rho_0(e^t z_r(t_n), e^t z_i(t_n)) \sinh t_n \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}} \frac{e^{t_n} z_r(t_n) - s}{e^{2t_n} z_i^2(t_n) + (e^{t_n} z_r(t_n) - s)^2} dx \sinh t_n \\ &= \frac{z_r^*}{2(z_i^*)^2 + 2(z_r^*)^2}. \end{aligned}$$

Similarly, we have

$$\lim_{n \rightarrow \infty} \pi P \rho_0(e^t z_r(t_n), e^t z_i(t_n)) \sinh t = \frac{z_i^*}{2(z_i^*)^2 + 2(z_r^*)^2}.$$

7 Hence, from (A.18) we obtain

$$z = w + \frac{1}{2} \frac{z_r^* - iz_i^*}{(z_i^*)^2 + (z_r^*)^2} = w + \frac{1}{2w}.$$

8 Similar to the calculation of (B.5), we know that the above equation has a unique solution in \mathbb{C}_+ :

$$w = \frac{1}{z - \sqrt{z^2 - 2}}.$$

9 Hence, we have

$$e^{-t} Z^{-1}(z, t) = z_r(t) + iz_i(t) \rightarrow w = \frac{1}{z - \sqrt{z^2 - 2}}, \quad t \rightarrow \infty.$$

By (A.16) and using the dominated convergence theorem again, we have

$$\begin{aligned} f(z, t) &= f_0(Z^{-1}(z, t)) e^t \\ &= \int_{\mathbb{R}} \frac{e^{2t} z_r(t) - s}{e^{2t} z_i^2(t) + [e^t z_r(t) - s]^2} \rho_0(s) ds - i \int_{\mathbb{R}} \frac{e^{2t} z_i(t)}{e^{2t} z_i^2(t) + [e^t z_r(t) - s]^2} \rho_0(s) ds \\ &\rightarrow \frac{z_r^* - iz_i^*}{(z_i^*)^2 + (z_r^*)^2} = \frac{1}{w} = z - \sqrt{z^2 - 2}. \end{aligned}$$

The trace of $z - \sqrt{z^2 - 2}$ on the real line is

$$f_\infty(x) := \pi H \rho_\infty(x) - i\pi \rho_\infty(x) = \begin{cases} x + \sqrt{x^2 - 2}, & x < -\sqrt{2}, \\ x - i\sqrt{2 - x^2}, & x \in [-\sqrt{2}, \sqrt{2}], \\ x - \sqrt{x^2 - 2}, & x > \sqrt{2}, \end{cases}$$

1 Hence,

$$\rho(x, t) \rightarrow \rho_\infty(x) = \frac{\sqrt{(2 - x^2)_+}}{\pi},$$

2 which proves part (iii) in Theorem 2.1.

3 **Step 4.** We prove (iv). From (1.7), if $g(z, t)$ and $\rho(x, t)$ are analytical solutions to (2.16) and
4 (1.4) with $\gamma = 0$, then

$$\tilde{g}(z, t) := e^t g\left(e^t z, \frac{e^{2t} - 1}{2}\right) - z, \quad z \in \overline{\mathbb{C}_+}, \quad t > 0,$$

5 is a $\overline{\mathbb{C}_+}$ -holomorphic solution to (2.10), and

$$\tilde{\rho}(x, t) := e^t \rho\left(e^t x, \frac{e^{2t} - 1}{2}\right), \quad x \in \mathbb{R}, \quad t > 0,$$

6 gives an analytical solution to (1.4) for $\gamma = 1$. By part (iii), we obtain part (iv). □

7

8 B An explicit solution to (1.4) with $\gamma = 0$

9 In this section, by the Stieltjes transform of Wigner's semicircle law μ_1 in (1.11), we recover an
10 explicit solution, which is same as the explicit solution of the Dyson equation (1.4) constructed in
11 (2.18) and (2.21) (see (B.8)).

12 First, we begin by taking Stieltjes transformation. Let $f_1(z)$ be the Stieltjes transform of the
13 Wigner's semicircle law μ_1 given by (1.11):

$$f_1(z) = \int_{-2}^2 \frac{1}{z - y} \mu_1(dy), \quad z \in \mathbb{C} \setminus [-2, 2].$$

Let $y = 2 \cos \theta$ for $\theta \in [-\pi, 0]$, $\alpha = -\theta$ and we have

$$\begin{aligned} f_1(z) &= \frac{1}{2\pi} \int_{-2}^2 \frac{\sqrt{4 - y^2}}{z - y} dy = \frac{1}{\pi} \int_{-\pi}^0 \frac{2 \sin^2 \theta}{z - 2 \cos \theta} d\theta = \frac{1}{\pi} \int_0^\pi \frac{2 \sin^2 \alpha}{z - 2 \cos \alpha} d\alpha \\ &= \frac{1}{\pi} \int_{-\pi}^\pi \frac{\sin^2 \theta}{z - 2 \cos \theta} d\theta. \end{aligned}$$

14 Let $\zeta = e^{i\theta}$ and we obtain

$$f_1(z) = \frac{1}{4\pi i} \oint_{|\zeta|=1} \frac{(\zeta^2 - 1)^2}{\zeta^2(\zeta^2 + 1 - z\zeta)} d\zeta.$$

15 Set

$$h(\zeta) := \frac{(\zeta^2 - 1)^2}{\zeta^2(\zeta^2 + 1 - z\zeta)}.$$

16 Function $h(\zeta)$ has three poles: $\zeta_0 = 0$, $\zeta_1 = \frac{z + \sqrt{z^2 - 4}}{2}$, and $\zeta_2 = \frac{z - \sqrt{z^2 - 4}}{2}$. Next, we choose the
17 branch cut of $\sqrt{z^2 - 4}$. Due to

$$\sqrt{z^2 - 4} = |z^2 - 4|^{1/2} e^{\frac{i}{2}[\arg(z-2) + \arg(z+2)]},$$

we see that -2 and 2 are branch points. We take the branch cut along the interval $[-2, 2]$ and we set $\arg(z-2) = \pi$ and $\arg(z+2) = 0$ for z on the upside of the branch cut. In this case, on the upside of $[-2, 2]$ we have $\sqrt{z^2-4} = i\sqrt{4-x^2}$ while on the downside of $[-2, 2]$, $\sqrt{z^2-4} = -i\sqrt{4-x^2}$. Moreover, the square root of z^2-4 has a positive imaginary part when $z \in \mathbb{C}_+$ and it has a negative imaginary part when $z \in \mathbb{C}_- := \{z : \Im(z) < 0\}$. Hence, for the imaginary part, we have

$$|\Im(z - \sqrt{z^2-4})| < |\Im(z + \sqrt{z^2-4})| \quad \text{for } z \in \mathbb{C} \setminus [-2, 2]. \quad (\text{B.1})$$

1 which implies

$$|\Im(\zeta_2)| < |\Im(\zeta_1)| \quad \text{for } z \in \mathbb{C} \setminus [-2, 2].$$

2 Due to $\zeta_1\zeta_2 = 1$, we have

$$|\zeta_2| < 1 \text{ and } |\zeta_1| > 1 \quad \text{for } z \in \mathbb{C} \setminus [-2, 2].$$

3 And we obtain for $z \in \mathbb{C} \setminus [-2, 2]$,

$$\text{Res } h(\zeta_0) = \lim_{\zeta \rightarrow \zeta_0} \frac{d}{d\zeta} [(\zeta - \zeta_0)^2 h(\zeta)] = z, \quad \text{Res } h(\zeta_2) = -\sqrt{z^2-4}$$

Hence, by the Residue theorem,

$$f_1(z) = \frac{z - \sqrt{z^2-4}}{2}, \quad \mathbb{C} \setminus [-2, 2]. \quad (\text{B.2})$$

Second, we show $f_1(-z)$ is a Herglotz (Pick) function, which is analytical on $\mathbb{C} \setminus [-2, 2]$ and $\Im(z)\Im(f_1(-z)) > 0$ for $\Im(z) \neq 0$ and we show the decay order of $\Re(f_1)$ and $\Im(f_1)$ as $\Re(z)$ and $\Im(z)$ tends to infinity. Direct calculations give

$$2\Re(\sqrt{z})\Im(\sqrt{z}) = \Im(z), \quad \Re(\sqrt{z})^2 = \frac{|z| + \Re(z)}{2}, \quad \text{and} \quad \Im(\sqrt{z})^2 = \frac{|z| - \Re(z)}{2}. \quad (\text{B.3})$$

4 For $z = x + iy$, we obtain

$$\Im(z^2-4) = 2xy, \quad \Re(\sqrt{z^2-4})^2 = \frac{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} + x^2 - y^2 - 4}{2},$$

5 and

$$\Im(\sqrt{z^2-4})^2 = \frac{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} - (x^2 - y^2 - 4)}{2}.$$

Recall that in our settings of branch cut, the square root of z^2-4 has positive imaginary part when $z \in \mathbb{C}_+$ and it has negative imaginary part when $z \in \mathbb{C}_-$. Due to (B.3), we know that the sign of $\Re(\sqrt{z^2-4})$ is the same as $\Im(\sqrt{z^2-4})$ when $xy > 0$ and they have different signs if $xy < 0$. By elementary calculations, we have

$$\Re(f_1(z)) = \begin{cases} \frac{\sqrt{2}x - \sqrt{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} + (x^2-y^2-4)}}{2\sqrt{2}}, & x > 0, \\ \frac{\sqrt{2}x + \sqrt{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} + (x^2-y^2-4)}}{2\sqrt{2}}, & x < 0, \end{cases} \quad (\text{B.4})$$

and

$$\Im(f_1(z)) = \begin{cases} \frac{\sqrt{2}y - \sqrt{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} - (x^2-y^2-4)}}{2\sqrt{2}} < 0, & y > 0, \\ \frac{\sqrt{2}y + \sqrt{\sqrt{(x^2-y^2-4)^2 + 4x^2y^2} - (x^2-y^2-4)}}{2\sqrt{2}} > 0, & y < 0. \end{cases} \quad (\text{B.5})$$

6 From the sign in (B.5), $\Im(z) > 0$ implies $\Im(-z) < 0$ and $\Im(f(-z)) > 0$. Therefore we have
7 $\Im(z) \cdot \Im(f_1(-z)) \geq 0$ and thus $f_1(-z)$ is a Herglotz function. Moreover, for fixed $y \in \mathbb{R}$ in (B.4),

1 dividing $\Re(f_1(z))$ by x shows that $\Re(f_1(z))$ decays in the order $O(|x|^{-1})$ as $|x| \rightarrow \infty$. Similarly,
 2 $\Im(f_1(z))$ decays in the order $O(|y|^{-1})$ as $|y| \rightarrow \infty$ for fixed $x \in \mathbb{R}$.

3 Third, we use f_1 to recover the explicit solution to the Dyson equation (1.4) with $\gamma = 0$ given
 4 by (2.18) and (2.21). Define

$$f_t(z) = \frac{1}{\sqrt{t}} f_1\left(\frac{z}{\sqrt{t}}\right), \quad z \in \mathbb{C} \setminus [-2\sqrt{t}, 2\sqrt{t}].$$

5 Then, direct checking shows that $f_t(z)$ is a self-similar solution to complex Burgers equation (1.1).

6 Finally, we try to obtain the traces of f_1 on the upper and lower half planes respectively. In
 7 the above settings of branch cut, we have

$$\arg(z - 2) = \pi = \arg(z + 2), \quad z \in (-\infty, -2),$$

8 which implies

$$\sqrt{z^2 - 4} = \sqrt{x^2 - 4}e^{i\pi} = -\sqrt{x^2 - 4}, \quad z = x \in (-\infty, -2).$$

9 Similarly, we have

$$\sqrt{z^2 - 4} = \sqrt{x^2 - 4}e^{i2\pi} = \sqrt{x^2 - 4}, \quad z = x \in (2, +\infty).$$

Hence, the trace of function $f_1(z)$ defined by (B.2) from the upper half plane \mathbb{C}_+ is given by

$$f_1(x+) = \begin{cases} \frac{x + \sqrt{x^2 - 4}}{2}, & x < -2, \\ \frac{x - i\sqrt{4 - x^2}}{2}, & x \in [-2, 2], \\ \frac{x - \sqrt{x^2 - 4}}{2}, & x > 2. \end{cases} \quad (\text{B.6})$$

The trace of function $f_1(z)$ given by (B.2) from the lower half plane \mathbb{C}_- is

$$f_1(x-) = \begin{cases} \frac{x + \sqrt{x^2 - 4}}{2}, & x < -2, \\ \frac{x + i\sqrt{4 - x^2}}{2}, & x \in [-2, 2], \\ \frac{x - \sqrt{x^2 - 4}}{2}, & x > 2. \end{cases} \quad (\text{B.7})$$

10 Direct computations show that $\frac{1}{\sqrt{t}}f_1\left(\frac{x}{\sqrt{t}}\pm\right)$ are solutions to complex Burgers equation on the real
 11 line \mathbb{R} .

Recall Section 2. If ρ is a solution to the Dyson equation (1.4), then $g = \pi H\rho - i\pi\rho - x$ is a solution to the complex Burgers equation (1.1) on the real line and $f = \pi H\rho - i\pi\rho$ gives a trace of an analytical function on the upper half plane. Hence, we use the trace $f_1(x+)$ (given by (B.6)) to define

$$f(x, t) = \frac{1}{\sqrt{t}} f_1\left(\frac{x}{\sqrt{t}}\right) = \begin{cases} \frac{x}{2t} + \frac{\sqrt{x^2 - 4t}}{2t}, & x < -2\sqrt{t}, \\ \frac{x}{2t} - i\frac{\sqrt{4t - x^2}}{2t}, & x \in [-2\sqrt{t}, 2\sqrt{t}], \\ \frac{x}{2t} - \frac{\sqrt{x^2 - 4t}}{2t}, & x > 2\sqrt{t}, \end{cases}$$

and

$$u(x, t) = \begin{cases} \frac{x + \sqrt{x^2 - 4t}}{2t}, & x < -2\sqrt{t}, \\ \frac{x}{2t}, & x \in [-2\sqrt{t}, 2\sqrt{t}], \\ \frac{x - \sqrt{x^2 - 4t}}{2t}, & x > 2\sqrt{t}, \end{cases} \quad \rho(x, t) = \frac{\sqrt{(4t - x^2)_+}}{2\pi t}. \quad (\text{B.8})$$

1 To the end of this section, we provide another method to prove $f_1(-z)$ is a Herglotz (Pick)
 2 analytic on $(-\infty, -2) \cup (2, +\infty)$. Recall $\mu_1(dy) = \frac{1}{2\pi} \sqrt{(4-y^2)_+} dy$. Then changing of variable
 3 $y = t - 2$ gives that

$$f_1(-z) = \int_0^4 \frac{1}{-z+2-t} \frac{1}{2\pi} \sqrt{(t(4-t))_+} dt. \quad (\text{B.9})$$

4 Define the measure $d\mu_*(t) := \frac{1}{2\pi} \sqrt{(t(4-t))_+} dt$ and recast $f_1(-z)$ as

$$f_1(-z) = \int_0^4 \frac{1}{-z+2-t} d\mu_*(t). \quad (\text{B.10})$$

5 Changing variable $-z+2 = \frac{1}{w}$ gives

$$f_1(-z) = \int_0^4 \frac{1}{\frac{1}{w}-t} d\mu_*(t) = \int_0^4 \frac{w}{1-wt} d\mu_*(t) =: wF(w). \quad (\text{B.11})$$

6 Here $F(w) = \int_0^4 \frac{1}{1-wt} d\mu_*(t)$ is a Pick function analytic on $(-\infty, \frac{1}{4})$. $F(w)$ is also the generating
 7 function of a completely monotone sequence $\{A_n(2, 2)\}_{n \geq 0}$ [17, Lemma 3], where $A_n(2, 2)$ is the
 8 general Fuss-Catalan numbers (also called Raney numbers) with index $(2, 2)$. Therefore from [17,
 9 Corollary 1 (iii)], $F_1(w) := wF(w)$ is a Pick function analytic on $w \in (-\infty, \frac{1}{4})$. From the relation
 10 $-z+2 = \frac{1}{w}$, we know $w(z) = \frac{1}{2-w}$ is a pick function mapping $(-\infty, -2) \cup (2, +\infty)$ to $(-\infty, \frac{1}{4})$.
 11 Therefore the composition $f_1(-z) = F_1 \circ w(z)$ is a Pick function analytic on $z \in (-\infty, -2) \cup (2, +\infty)$.