



The relationship between PM_{2.5} and anticyclonic wave activity during summer over the United States

Ye Wang^{1,2}, Natalie Mahowald², Peter Hess³, Wenxiu Sun^{3,a}, and Gang Chen⁴

¹College of General Aviation and Flight, Nanjing University of Aeronautics and Astronautics, Nanjing, Jiangsu, China

²Department of Earth and Atmospheric Science, Cornell University, Ithaca, NY, USA

³Department of Biological and Environmental Engineering, Cornell University, Ithaca, NY, USA

⁴Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA, USA

^anow at: BloomSky Inc., Burlingame, CA, USA

Correspondence: Ye Wang (wangye08@nuaa.edu.cn)

Received: 9 September 2021 – Discussion started: 12 October 2021

Revised: 6 May 2022 – Accepted: 7 May 2022 – Published: 13 June 2022

Abstract. To better understand the role of atmospheric dynamics in modulating surface concentrations of fine particulate matter (PM_{2.5}), we relate the anticyclonic wave activity (AWA) metric and PM_{2.5} data from the Interagency Monitoring of Protected Visual Environment (IMPROVE) data for the period of 1988–2014 over the US. The observational results are compared with hindcast simulations over the past 2 decades using the National Center for Atmospheric Research–Community Earth System Model (NCAR CESM). We find that PM_{2.5} is positively correlated (up to $R = 0.65$) with AWA changes close to the observing sites using regression analysis. The composite AWA for high-aerosol days (all daily PM_{2.5} above the 90th percentile) shows a similarly strong correlation between PM_{2.5} and AWA. The most prominent correlation occurs in the Midwestern US. Furthermore, the higher quantiles of PM_{2.5} levels are more sensitive to the changes in AWA. For example, we find that the averaged sensitivity of the 90th-percentile PM_{2.5} to changes in AWA is approximately 3 times as strong as the sensitivity of 10th-percentile PM_{2.5} at one site (Arendtsville, Pennsylvania; 39.92° N, 77.31° W). The higher values of the 90th percentile compared to the 50th percentile in quantile regression slopes are most prominent over the northeastern US. In addition, future changes in US PM_{2.5} based only on changes in climate are estimated to increase PM_{2.5} concentrations due to increased AWA in summer over areas where PM_{2.5} variations are dominated by meteorological changes, especially over the western US. Changes between current and future climates in AWA can explain up to 75 % of PM_{2.5} variability using a linear regression model. Our analysis indicates that higher PM_{2.5} concentrations occur when a positive AWA anomaly is prominent, which could be critical for understanding how pollutants respond to changing atmospheric circulation as well as for developing robust pollution projections.

1 Introduction

Particulate matter less than 2.5 µm in diameter (PM_{2.5}) poses a considerable air quality concern due to its impacts on human health (Liu et al., 2020). PM_{2.5} has been linked to the increased possibility of mortality (Krewski et al., 2009). Continuing exposure to PM_{2.5} can exacerbate existing cardio-

vascular and respiratory problems and cause lung damage (Bernard et al., 2001). It can also alter the body's defense system against foreign materials and even lead to premature death (Kappos et al., 2004). Furthermore, PM_{2.5} could contribute to the degradation of visibility (Hand et al., 2011; Ashley et al., 2015) and the alteration of the hydrological cycle through changing rainfall formation mechanisms in

clouds (Rosenfeld et al., 2008). Once deposited onto snow PM_{2.5} can cause an increase in the melting of snow and ice (Painter et al., 2007) or modify land or ocean biogeochemistry (Mahowald et al., 2011). It also influences Earth's energy balance directly by scattering solar incoming radiation back into space (Charlson et al., 1992) or indirectly by altering cloud albedo and lifetime (Albrecht, 1989; Arimoto, 2001). Aerosols can also alter the large-scale atmospheric circulation and lead to historical changes to wintertime mid-latitude winter extremes over northern Eurasia (Wang et al., 2020). Thus, understanding and predicting potential changes in PM_{2.5} is crucial to both human health concerns and current environmental issues.

Climate change and meteorological factors may have substantial impacts on surface concentrations of PM_{2.5}. A number of studies have demonstrated the influences of weather and climate change on PM_{2.5} variability (Brasseur et al., 2006; Liao et al., 2006; Dawson et al., 2007; Jacob and Winner, 2009; Thishan Dharshana et al., 2010; Tai et al., 2012; Liu et al., 2017; Chen et al., 2018; Wang et al., 2019; Wang and Zhang, 2020), showing connections between specific meteorological conditions and PM_{2.5} concentration responses. Daily variation in temperature, rainfall, moisture, and circulation can explain up to 50 % of the variability in PM_{2.5} (Tai et al., 2010). Meanwhile, Aw and Kleeman (2003) calculated a reduction in PM_{2.5} concentrations caused by rising temperature over southern California in a modeling study. Furthermore, Tai et al. (2012) identified a cyclone passage with the associated cold front as a mechanism for temperature being strongly correlated with interannual variability of PM_{2.5} in the Midwestern US. Similarly, increased atmospheric stagnation may also have the potential to aggravate PM_{2.5} air quality in the future climate (Liao et al., 2006; Leibensperger et al., 2008). Mickley et al. (2004) also proposed an increased severity of summertime PM episodes due to a warmer future climate in the northeastern and Midwestern US. In addition, a significantly negative correlation caused by the longer lifespan of gas-phase-produced sulfate was found between aerosol sulfate and cloud cover compared to aqueous-phase-produced sulfate (Koch et al., 2003). These studies suggest that changes in meteorology linked to climate modification may cause variations in PM_{2.5} levels and exposure risks. However, questions remain regarding whether and how PM_{2.5} concentrations directly related to atmospheric general circulation. Understanding the relationship between meteorology and PM_{2.5} levels will be critical to the understanding of pollutant response to a changing circulation due to climate change as well as the development of robust pollution projections.

Many studies have focused on relating wave activity with extreme weather and climate events. Weather extremes are strongly influenced by the natural variability of the atmosphere at synoptic, intra-seasonal and interannual timescales. For example, persistent high-pressure blocking systems can cause extreme cold winter temperatures in Europe (Wool-

ings et al., 2008) or summer heat waves (Coumou et al., 2015). Midlatitude weather extremes can be influenced by the major modes of climate variability such as Arctic Oscillation (Michel and Rivière, 2011) or El Niño–Southern Oscillation (ENSO) (Ryoo et al., 2013). Furthermore, blocking anticyclones can decay by releasing accumulated wave activity as a stationary Rossby wave train (Takaya and Nakamura, 2001). Meanwhile, blocking highs are connected with high-amplitude, quasi-stationary anticyclonic anomalies that result in protracted and unusual weather events (Nakamura et al., 1997). In addition, temperature extremes are more likely to connect with wave events through the large-amplitude troughs and ridges (Pfahl and Wernli, 2012; Chen et al., 2015; Martineau et al., 2017). Shen and Mickley (2017) identify the association between warm tropical Atlantic sea surface temperatures and enhanced subsidence, reduced precipitation and increased temperatures through stationary wave propagation in the eastern US. However, no previous studies have investigated how wave activity can lead to changes in PM_{2.5} concentrations, although studies have shown a robust correlation between surface ozone concentration and a measure of wave activity over the US using a linear regression model (Sun et al., 2019).

Here we propose the use of anticyclonic wave activity (AWA), the anticyclonic part of local finite-amplitude wave activity (LWA) with quantile regressions, to be an effective method for diagnosing the tendency and sensitivities in the transport of PM_{2.5} concentrations, using a similar approach to a previous study focused on ozone (Sun et al., 2019). LWA assesses longitude-by-longitude anomalies of the finite-amplitude Rossby wave activity deviating from the circle of equivalent latitude using the meridional displacement of potential vorticity (PV) (Huang and Nakamura, 2016). As an example, the prominent Northern Hemisphere blocking episode that occurred in late October 2012 was well explained using LWA (Huang and Nakamura, 2016). Furthermore, recent modeling studies have illuminated the value of quantile regression in examining the influences of global changes on local air quality. For example, Porter et al. (2015) employ quantile regression to diagnose the meteorological sensitivities of higher ozone and PM_{2.5} levels by using observed and reanalysis meteorological data in the US over the past decade.

While meteorology sets the stage for the occurrence of dangerous pollutant levels in the present and future climates, the connection between the meteorological events and extreme pollution events is still not sufficiently understood (Dawson et al., 2014). In the present study, we apply a univariate linear regression model to analyze the daily PM_{2.5}–AWA relationship from observations and simulations in the US, using a similar methodology as that used to examine LWA and ozone relationships (Sun et al., 2019). We also use quantile regression to calculate PM_{2.5} sensitivities to AWA across quantiles from 10 % to 90 %. In addition, the coefficients of the slope for the linear regression model between

PM_{2.5} and AWA in the current climate are evaluated to see how much they can be used to predict future PM_{2.5} changes. Such an exploration of PM_{2.5} from atmospheric motion can provide new insights into understanding the mechanism of PM_{2.5} changes as well as the sensitivity of higher PM_{2.5} concentrations to climate change.

2 Methodology

In order to quantify the relationship between AWA and surface PM_{2.5} concentrations during the summer months, five different combinations of datasets, one observational one and four model outputs, are used in this study. Each of the five cases uses surface PM_{2.5} concentrations and geopotential heights (for calculating a modified AWA, as explained below). Each modeling dataset uses 20 years for the analysis, while the observations are analyzed for the years between 1988 and 2014. Both PM_{2.5} and AWA time series are detrended and deseasonalized by eliminating the 21 d smoothed seasonal cycle from the original data. We use the resulting stationary residuals to focus on the synoptic-scale variability, minimizing aliasing from regular seasonal variations or long-term tendencies. This study is a followup study to Sun et al. (2019), using the same approach used there to analyze ozone, and here we focus on analyzing PM_{2.5}. The details of the methods are discussed in more detail in that paper and summarized here.

2.1 Observational data

We use PM_{2.5} measurements from the Interagency Monitoring of Protected Visual Environment (IMPROVE) monitoring sites, which are located in National Parks and Class I Wilderness areas across the US (<http://vista.cira.colostate.edu/improve/>, last access: 1 September 2021). PM_{2.5} concentrations are reported every 3 d as mass per air volume at local temperature and pressure. Hand et al. (2011, 2012) have described the details regarding IMPROVE site locations, sampling, analysis approach and detailed information of network operations. All stations with at least 1000 valid PM_{2.5} values between 1988 and 2014 are collected in this study, totaling 150 stations for summer (June, July, August; JJA hereafter). For the PM_{2.5} observations (called Obs case), the full names, states, latitudes and longitudes as well as short names are listed in Table S1 in the Supplement, and locations are shown in Fig. 1. We chose three representative stations in different parts of the country to investigate the relation between AWA and PM_{2.5} in detail. The IMPROVE station names are AREN1 (Arendtsville, Pennsylvania; 39.92° N, 77.31° W; in the northeast), SIPS1 (Sipsey Wilderness, Alabama; 34.34° N, 87.34° W; in the southeast) and LAVO1 (Lassen Volcanic National Park, California; 40.54° N, 121.58° W; in the west), which are shown with the red dots in Fig. 1. They match the sites PSU106 (40.72° N, 77.93° W; in the northeast), SND152 (34.29° N, 85.97° W;

in the southeast) and LAV410 (40.54° N, 121.58° W; in the west), respectively, which have differing impacts of meteorological persistence on the distribution and extremes of ozone in Sun et al. (2017) to allow a comparison between the ozone and PM_{2.5} response to AWA. Long-range transport from Asia and meteorology are dominant drivers of pollutants at LAVO1, where anthropogenic influence is at a minimum as it is a clean-air site in California (VanCuren and Gustin, 2015).

In order to compare these in situ observations to the large-scale meteorology, the 500 hPa geopotential height (m) from the European Reanalysis Interim version is used (ERA-Interim) for the time period of January 1991 to December 2010 (Dee et al., 2011). The ERA-Interim is a global reanalysis of recorded meteorological data over the past 3.5 decades and was undertaken by the European Centre for Medium-Range Weather Forecasts. This gridded dataset is created at approximately 0.7° spatial resolution with 37 vertical levels.

2.2 Model output

We compare the IMPROVE data with output from the Community Earth System Model (CESM) as simulated for the Chemistry-Climate Model Initiative (CCMI) (Eyring et al., 2013). As a state-of-the-art Earth system modeling framework coordinated by the National Center for Atmospheric Research (NCAR), the CESM employed here is configured to fully couple the Community Atmosphere Model version 4 (CAM4) (Neale et al., 2010), the Community Land model version 4.0 (CLM4.0) (Oleson et al., 2010), the Parallel Ocean Program version 2 (POP2) (Smith et al., 2010) and the Los Alamos sea ice model (CICE version 4) (Hunke and Lipscomb, 2008). All simulations are performed under current land cover conditions. The top of the simulated atmosphere reaches 40 km with a horizontal resolution of 2.5° longitude by 1.9° latitude. The model has been widely used for modeling the Earth's past, present and future climate states (Neale et al., 2010; Hurrell et al., 2013).

To simulate the atmospheric chemistry, we include the CAM4-Chem module, which has been widely studied with regard to its representation of atmospheric chemistry in the atmosphere (Aghedo et al., 2011; Lamarque et al., 2011a, b, 2012; Lamarque and Solomon, 2010). CAM4-Chem employs the Bulk Aerosol Model (BAM) with externally mixed aerosols considering black carbon, organic carbon, sulfate, sea spray and desert dust, which simulates coarse-mode aerosols in four bins for the latter two. Details of this implementation are discussed in Lamarque et al. (2012). CAM4 uses the Zhang–McFarlane deep-convection scheme (Zhang and McFarlane, 1995), the Hack shallow-cumulus scheme (Hack et al., 2006), Holtslag and Boville's (1993) planetary boundary layer process, and the parameterization of cloud microphysics and macrophysics by Rasch and Kristjánsson (1998) and Zhang et al. (2003). Additionally, the convective momentum transport is utilized to parameterize deep




Figure 1. Geographical location of IMPROVE sites for PM_{2.5}. Red dots are three representative sites.

convection (Richter and Rasch, 2008). These two major revisions caused improvements in such aspects as the Madden–Julian oscillation and ENSO (Neale et al., 2008). The improved trend and magnitude of surface PM_{2.5} using this free-running model have been evaluated elsewhere (Tilmes et al., 2016).

The chemical emissions and forcing details for each of the model simulations are listed in Table 1. The simulation using specified dynamics (REFC1SD) for current levels of PM_{2.5} from 1991 to 2010 is driven by analyzed meteorological data from Modern-Era Retrospective Analysis for Research and Applications (MERRA) (see Tilmes et al., 2016). This simulation follows the conventions of the CCM1 (Eyring et al., 2013). For the AWA analysis for this case, we use the 500 hPa geopotential height from MERRA, which should be very similar to that from ERA-Interim, since they use largely the same observations. To compare the relationship between AWA and PM_{2.5} concentrations in online simulations, three simulations forced by trace gas projections and an interactively coupled ocean are employed. The GCM2000 and GCM2100 simulations are 25-year runs branched from the CCM1 reference simulations in the year 2000 and the year 2100, respectively. Simulations over the first 5 years are discarded as spin-up, and results from the latter 20 years are discussed here (2006–2025 for GCM2000 and 2106–2125 for GCM2100). Note that while the CO₂ concentrations in the GCM2000 and GCM2100 simulations are kept at the year 2000 and 2100 level, respectively, the concentrations of all other greenhouse gases including methane remain constant at the year 2000. In particular, the emissions and prescribed chemical species for longer-lived substances follow

the protocol defined by CCM1 hindcast simulations for the year 2000 (Eyring et al., 2013), which are repeated for all the simulated model years. Another future run (REFC2) is forced by future climate combined with future emissions following the REFC2 CCM1 modeling protocol. In this run, greenhouse gas forcing and emissions follow the RCP6 scenario. The relationship between ozone and AWA has been examined in the GCM2000, GCM2100 and REFC2 simulations in Sun et al. (2019). Characteristics of the REFC1SD simulation are given in Phalitnonkiat et al. (2018). Note that our REFC2 set-up covers volcanic eruptions in the past, but possible volcanic eruptions in the future are not included (Eyring et al., 2013).

2.3 AWA calculation

To calculate AWA, we adopt the procedures in Chen et al. (2015) and Huang and Nakamura (2016). A dynamical quantity, q (here we use Z_{500} , geopotential height at 500 hPa), approximately decreases with latitude in the Northern Hemisphere. For a given value of $q = Q$, we introduce an equivalent latitude $\phi_e(Q)$ as

$$\phi_e(Q) = \arcsin \left[1 - \frac{S(Q)}{2\pi a^2} \right]. \quad (1)$$

Here, $S(Q)$ is the area bounded by the Q contour towards the North Pole and a denotes Earth's radius. Defining an eddy term as $\hat{q} \equiv q - Q$ and separating the southward and northward displacements in the Q contour, we calculate the cyclonic (southern), anticyclonic (northern) and total LWA at

Table 1. The model case names used in this study. Designated model descriptions for the four attribution cases. Online means that the model calculates the meteorology prognostically online. Case names are based on names described previously (Eyring et al., 2013; Sun et al., 2019).

Simulation (years)	Model case name	GHG ¹ forcing	SST ² and sea ice	Emissions	Meteorological driver
REFC1SD (1991–2010)	f.e11.TSREFC1SD.f19.f19.ccmi23.001	CMIP5 ³	HadISST2 ⁴	Anthropogenic and biomass burning emission: MACCity ⁵ Biogenic emissions: MEGAN2 ⁶	MERRA ⁷
GCM2000 (2006–2025)	b.e11.TSREFC2.femis2000.y2000.f19.f19.ccmi23.001	CO ₂ = 369 ppm	Online ⁸	Anthropogenic and biomass burning from AR5 ⁹ Biogenic emissions: monthly values from MEGAN2 for 2000	Online
GCM2100 (2106–2125)	b.e11.TSREFC2.femis2000.y2100.f19.f19.ccmi23.001	CO ₂ = 669 ppm	Online ⁸	Same as GCM2000	Online ⁸
REFC2 (2080–2099)	b.e11.TSREFC2.f19.g16.ccmi23.001.cam.h7	A1B scenario	Online	A1B scenario	Online

¹ Greenhouse gas. ² Sea surface temperature. ³ Coupled Model Intercomparison Project. ⁴ Hadley Centre Sea Ice and Sea Surface Temperature dataset (Titchner and Rayner, 2014).

⁵ Granier et al. (2011). ⁶ Guenther et al. (2012). ⁷ Modern-Era Retrospective Analysis for Research and Applications (Rienecker et al., 2011). ⁸ Tilmes et al. (2016). ⁹ Assessment report 5 (Eyring et al., 2013).

the longitude λ and latitude ϕ_e by

$$A_C(\lambda, \phi_e) = \frac{a}{\cos \phi_e} \int_{\hat{q} \leq 0, \phi \leq \phi_e(Q), \lambda = \text{const}} \hat{q} \cos \phi d\phi, \quad (2)$$

$$A_A(\lambda, \phi_e) = \frac{a}{\cos \phi_e} \int_{\hat{q} \geq 0, \phi \geq \phi_e(Q), \lambda = \text{const}} \hat{q} \cos \phi d\phi, \quad (3)$$

$$A_T(\lambda, \phi_e) = A_C - A_A. \quad (4)$$

Studies on finite-amplitude wave activity (FAWA) have identified the link between the pattern of atmospheric circulation and large-scale wave dynamics (Nakamura and Solomon, 2011; Methven, 2013; Chen and Plumb, 2014; Lu et al., 2015). LWA adds the longitude dimension to the zonally average quantity FAWA and is calculated from the meridional displacement of quasi-geostrophic PV from zonal symmetry (Nakamura and Zhu, 2010). LWA helps differentiate longitudinally isolated events and describe extreme weather events at the local scales (Huang and Nakamura, 2016; Chen et al., 2015). Chen et al. (2015) used local finite-amplitude wave activity based on the 500 hPa geopotential height for characterizing midlatitude weather events. The total wave activity is composed of the cyclonic wave activity residing to the south of the equivalent latitude and the anticyclonic wave activity to the north (see Fig. 1 in Sun et al., 2019). In this study, we focus on AWA to characterize its connection with changes in PM_{2.5} concentrations. Over the US in summer LWA is dominated by its anticyclonic component (Sun et al., 2019). Sun et al. (2019) also used AWA to characterize ozone variability.

2.4 Quantile regression

Quantile regression is used to estimate the slopes for several conditional quantile functions (Koenker and Bassett, 1978).

It characterizes the connection between a range of predictor variables and specified percentiles (or quantiles) of the response variable. For example, Porter et al. (2015) analyzed the sensitivities of ozone and PM_{2.5} concentrations for response quantiles ranging from 2 % to 98 %. The parameters of quantile regression models evaluate the change in a specific quantile of the response variable caused by a one-unit change in the predictor variable. This permits us to measure how some percentiles of the PM_{2.5} may be more influenced by AWA than others, and this is indicated by changes in the regression coefficient. In order to illustrate the sensitivity of the PM_{2.5} concentration at different quantiles, we apply linear quantile regression for percentiles from the 10th to the 90th one at the AREN1 site. And then we compare the 90th-percentile quantile regression coefficient with the 50th-percentile quantile regression coefficient at each station.

2.5 The univariate linear regression model

To help explore and measure the likely relationship between AWA and PM_{2.5} levels, we use the univariate linear regression model, similar to a previous study focused on ozone (Sun et al., 2019). Here the slope of PM_{2.5} with respect to wave activity ($S_{i0,j0}(i,j)$) on the daily timescale is used to show the linear association between changes (in time) of the normalized PM_{2.5} at a point (i_0, j_0) and the normalized wave activity at another point. We use the projection of PM_{2.5} onto AWA to reveal how closely the AWA anomaly field resembles the spatial pattern that enhances PM_{2.5} on the daily timescale during the summer. The projection of AWA ($p_{i0,j0}$) at all points in the domain onto $S_{i0,j0}$ is defined ac-

cording to the following equation:

$$p_{i0,j0} = \text{AWA} \cdot S_{i0,j0} = \sum_j \sum_i \text{AWA}(i,j) \times S_{i0,j0}(i,j). \quad (5)$$

The similarity between the AWA spatial pattern and the PM_{2.5}–AWA regression coefficients' spatial structure is estimated by the projection value. The interannual change in PM_{2.5} due to changes in AWA is predicted based on a linear regression model as in the following equation (see Sun et al., 2019, for more discussion):

$$\text{PM}_{2.5} = \beta p + \alpha. \quad (6)$$

The change in PM_{2.5} (denoted by $\Delta\text{PM}_{2.5}$) in the future due to the change in AWA is calculated using the following equation, which is the climatological difference in a future climate and the present climate ($\text{AWA}_f - \text{AWA}_p$); project it onto S , and multiply it by the slope β :

$$\Delta\text{PM}_{2.5} = \beta [(\overline{\text{AWA}_f} - \overline{\text{AWA}_p}) \cdot S]. \quad (7)$$

Here, S is calculated from the values for the present climate. The projected value can measure the similarity between the AWA change and the PM_{2.5}'s trend with AWA by compressing the information of the AWA field into a single variable. This variable incorporates the non-local effect of AWA on PM_{2.5}'s variability.

2.6 The composite methodology

We use a composite methodology which is based around the most polluted (> 90th percentile) daily PM_{2.5} and the corresponding anomalies at every station. Composite 500 hPa geopotential height and AWA for daily values of PM_{2.5} larger than the 90th percentile are produced by separately averaging all daily anomaly values of the corresponding 500 hPa geopotential height and AWA. The composite methodology can average out much of the variability. Composite Madden–Julian Oscillation cycles of precipitation and ozone for each phase are examined by averaging together all daily anomaly values for the given quantity separately (Sun et al., 2014). In addition, the meteorological conditions conducive to a high-ozone event are investigated by compositing about the first day of each high-ozone event in the northeastern region (Sun et al., 2017).

3 Results and discussion

The monthly mean PM_{2.5} surface concentrations with standard deviations for different scenarios at three representative sites are shown in Fig. 2. PM_{2.5} concentrations are largest during summer at the three sites. The climatological average for PM_{2.5} is greater at the eastern than at the western sites. A statistically significant correlation ($r > 0.80$, $p < 0.01$; Fig. 2) for current PM_{2.5} is found between observations and simulations of monthly mean climatological averages (REFC1SD and GCM2000) at three representative sites.

The highest correlation coefficients between model and observations (0.93) are seen at SIPS1, perhaps due to the large seasonal variation in PM_{2.5} concentrations (Fig. 2b). The future PM_{2.5} concentrations are increased in GCM2100 under current emissions compared with current climate PM_{2.5} simulations. There is a strong decrease in climatological mean for future PM_{2.5} at AREN1 under future emissions and meteorology (REFC2), while the climatological average for future PM_{2.5} has no significant change under future emissions at SIPS1 and LAVO1. Such differences in the monthly mean averages for PM_{2.5} suggest that emission changes are more important than climate changes at AREN1, but it is not clear which is more important at SIPS1 or LAVO1.

Focusing on the spatial distribution, the highest PM_{2.5} concentrations over the 20-year average in summer occur in the south–central US (Fig. 3a; green lines; GCM2000 case). The 20-year averaged AWA on summer days exhibits a maximum over the southwestern US (Fig. 3a; shaded). The difference between two current climate simulations (REFC1SD minus GCM2000) for summertime AWA is shown in Fig. 3b. The reduced AWA in the reanalysis forced simulation is found across most of the US, with the largest reduction in the southwestern US as previously shown (Sun et al., 2019). In contrast, the AWA in summer is higher over the northeastern US in the forced simulations. The corresponding changes in summertime PM_{2.5} concentration caused by a combination of different emissions and possibly changes in AWA are similar to changes in AWA, although the reduction is largely over the south–central US. The difference between two future scenarios (GCM2100; REFC2) and the current climate scenario (GCM2000) has a similar pattern (illustrated in Fig. 3c and d), which shows a large increase in AWA in the southwestern US, but there is a difference in the amplitude of these changes (contrast Fig. 3c with Fig. 3d) (Sun et al., 2019). There is an increase in PM_{2.5} concentration for the future scenario with current emissions (GCM2100), while there is a decrease in PM_{2.5} concentrations when future emissions are used (REFC2), showing the importance of future potential decreases in emissions.

3.1 Relationship between PM_{2.5} concentrations and AWA at specific stations

For the three observation sites highlighted here (AREN1, SIPS1 and LAVO1), the PM_{2.5} concentrations are positively correlated with AWA in the areas close to the sites where presumably at least some of emissions of PM_{2.5} are located (Fig. 4). The relationships between daily PM_{2.5} concentrations and AWA using CESM simulations presented here offer a test of the consistency between observational and model relationships in characterizing the response of PM_{2.5} to AWA. The highest regression coefficient occurs in the observational (Obs) and the reanalysis-driven simulated cases (REFC1SD), as opposed to the case-coupled meteorology (GCM2000) (Fig. 4a, b and c and Fig. 4d, e and f in




Figure 2. Climatological monthly mean average with standard deviation for PM_{2.5} used in this study at three sites (AREN1 **a**, SIPS1 **b** and LAVO1 **c**) for five scenarios used in this study. Red (r) represents the correlation coefficient between observation (Obs) and simulation for current from the REFC1SD simulation. Blue r represents the correlation coefficient between observation (Obs) and simulation for current from GCM2000 simulation. The p values are included.




Figure 3. Wave activity (AWA: shaded using the legend in 10^8 m^2) and PM_{2.5} concentrations (green contour lines in $\mu\text{g m}^{-3}$) for **(a)** the current climate (GCM2000, 2006–2025 summer days' average); **(b)** reanalysis-driven case (REFC1SD, 1991–2010 summer days' average) minus the current climate online case (GCM2000, 2006–2025 summer days' average); **(c)** future climate with current emission (GCM2100, 2106–2125 summer days' average) minus the current climate (GCM2000, 2006–2025 summer days' average); **(d)** future climate with future emission (REFC2, 2080–2099 summer days' average) minus current climate (GCM2000, 2006–2025 summer days' average). Three black dots are representative stations (AREN1, SIPS1 and LAVO1).

contrast to Fig. 4g, h and i). The highest spatial regression coefficients for sites AREN1 and SIPS1 are located southward of the sites, while they are located to the northwest at LAVO1. Overall the model simulates similar spatial patterns to the observations for the case of the reanalysis-driven simu-

lations (REFC1SD) but does less well for the coupled model simulations (GCM2000). Of course, model predictions are not perfect and include uncertain emissions and boundary conditions as well as errors in model physical and chemical processes, which may be driving these inconsistencies be-

tween modeled and observed relationships. In addition, Sun et al. (2019) show considerable variation in the AWA pattern between different ensemble simulations, suggesting that even on the timescale of 20 years there is considerable internal variability between model ensemble members.

To obtain a more in-depth understanding of the physical mechanisms behind the relationship between PM_{2.5} concentration and AWA, we consider the composite AWA for high-PM_{2.5} days at AREN1, SIPSI and LAVO1 (Fig. 5). The pattern for the composite AWA corresponding to daily PM_{2.5} above the 90th quantile is most similar to regression coefficients between daily AWA and PM_{2.5} by comparing different quantiles (Fig. S1 in the Supplement). The composite AWA calculated by averaging together all AWA corresponding to daily PM_{2.5} above the 90th quantile shows a similar strong connection between PM_{2.5} and AWA as that seen for the average (Fig. 4 vs. Fig. 5). The composite AWAs are extremely pronounced over the areas where PM_{2.5} data originate. The geopotential composites for the highest pollution days generally showed spatial distributions which are similar to the regression coefficient distributions (Fig. 5 vs. Fig. 4) at the three representative sites. The areas with the largest values of the composite AWA are located southward of the AREN1 and SIPSI sites. But at LAVO1 the maximum is located to the northwest for the observational and reanalysis-driven cases (Obs and REFC1SD) and eastward for the coupled model case (GCM2000). Overall, the composite AWA for PM_{2.5} also shows that the daily PM_{2.5} above its 90th quantile connects strongly with AWA during summer. Note that there is a spatial displacement between the maximum of geopotential height and the maximum of AWA, since wave activity fundamentally measures the waviness of atmospheric general circulation rather than the magnitude.

3.2 Relationship of AWA and PM_{2.5} regionally

Next we consider how the local relationship between PM_{2.5} and AWA changes in space. To simplify the visualization of the spatial variability in the local relationship, we use the result from the previous section that the maximum regression coefficients between PM_{2.5} and AWA are usually close to the site where PM_{2.5} is measured (Fig. 5). The highest composite AWA anywhere in the domain and the highest regression coefficient with AWA are shown at each grid point in Fig. 6. If we look at the relationship between the PM_{2.5} concentration and the wave activity at each location, it can be seen that PM_{2.5} concentrations are positively correlated with AWA throughout the US but with varying strengths (Fig. 6). A roughly similar spatial distribution is obtained with either the composite AWA for high PM_{2.5} (Fig. 6a, c, e, g and i) or the regression coefficients between the PM_{2.5} and AWA (Fig. 6b, d, f, h and j) for the different observational or model combinations, showing the consistency of the approaches. The pattern of the distribution is consistent for observational (Fig. 6a and b) and simulated (Fig. 6c and d) data

when forced with reanalysis winds, with the largest values in the upper Midwest into the mid-Atlantic states. The climate model simulations tend to have a stronger correlation in the western/mountain regions, than seen in the reanalysis winds (Fig. 6c vs. Fig. 6e), and there are some hints of this in the observations in Arizona, for example (Fig. 6a). The correlations and composites tend to become stronger in the western/mountain regions in the future model simulations (Fig. 6g and i vs. Fig. 6e).

The consistency in the composite AWA under high-PM_{2.5} conditions and the correlations suggests that either of these metrics can be useful tools to identify PM_{2.5} and AWA relationships. The anticyclonic condition is usually characterized by, low-level divergence, subsiding air, light wind, no rainfall and high surface pressure. Taken together, the results above demonstrate the positive connection of PM_{2.5} concentrations with anticyclonic conditions everywhere across the US, which is likely accounted for by arid weather and sinking inversions. This is consistent with the PM_{2.5} sensitivity study of Tai et al. (2010). They illustrated strong linkage between high PM_{2.5} concentrations, high 850 hPa geopotential height and stagnation, which is characterized by an anticyclonic condition. The significant association of PM_{2.5} with stagnation is also demonstrated by Cheng et al. (2007) in their examination of four Canadian cities. Furthermore, similar results were reported that greatly strengthening ozone is associated with increased stagnation (Wu et al., 2008; Sun et al., 2014).

The composite AWA is for PM_{2.5} that is larger than the 90th quantile, while the regression coefficient is for all PM_{2.5}. The composite AWA and regression coefficient have similar spatial distributions suggesting the positive connection between daily PM_{2.5} and AWA is mainly produced by high PM_{2.5} concentration above its 90th quantile. Overall, the relationship between the behavior of AWA and extreme PM_{2.5} concentration is generally consistent with the existing meteorological studies (Woolings et al., 2008; Coumou et al., 2015; Michel and Rivière, 2011; Ryoo et al., 2013).

3.3 The sensitivity of quantiles in PM_{2.5} concentrations to AWA

To examine the sensitivity of different levels of PM_{2.5} concentrations to AWA, we fit the linear regression and quantile regression for AWA and daily PM_{2.5} for summers between 1988 to 2014 from IMPROVE monitoring sites for different percentiles (10th to 90th percentiles) using an “impact region” of AWA at the AREN1 site. Here the averaged AWA over the impact region is defined as an elliptic area bounded by the maximum and minimum longitude and latitude of the maximum composite AWA for PM_{2.5} larger than the 90th percentile minus the 0.05 contour line (blue elliptic circle in Fig. 5a). One can clearly see that the higher percentiles of PM_{2.5} are more sensitive to the change in the averaged AWA over the impact region; e.g., the 90th percentile




Figure 4. Composite 500 hPa geopotential height anomaly (contours; positive values are represented by green lines and negative values by magenta lines) and regression coefficients (shaded) between daily AWA and PM_{2.5} at sites (denoted by the black dots) (a, d, g) AREN1, (b, e, h) SIPS1 and (c, f, i) LAVO1 in the study domain for daily JJA time series of current climates. The top row shows results using IMPROVE PM_{2.5} and reanalysis AWA, the middle row uses the reanalysis-driven simulated PM_{2.5} (REFC1SD) and reanalysis AWA, and the bottom row uses current-climate-simulated PM_{2.5} and AWA (GCM2000). Stippling indicates the regions that are statistically significant at the 5 % confidence level. Unit: $10^{-8} \mu\text{g m}^{-3} \text{m}^2$ for regression coefficients.

of PM_{2.5} is approximately 3 times more sensitive to the averaged AWA over the impact region when compared with the 10th percentile of PM_{2.5} (Fig. 7a). The correlation coefficient of 0.36 between JJA deseasonalized PM_{2.5} and the impact region's average AWA implies that the vast majority of all variability is being driven by factors other than AWA at the AREN1 site. It must be noted that this lack of overall correlation implies other drivers of PM_{2.5} variability at sites like this.

In order to examine whether the finding that a high quantile of PM_{2.5} is more sensitive to the AWA than the low quantile of PM_{2.5} applies to the other sites, we calculate the difference of the 90th-percentile quantile regression coefficient (slope) from the 50th-percentile quantile regression coefficient at the 5 % significance level across all sites (90th-percentile quantile regression coefficient (slope) minus the 50th-percentile quantile regression coefficient, shown in Fig. 7b). Out of the 150 sites, 145 sites show that 90th-percentile PM_{2.5} increases more than the 50th percentile of PM_{2.5} with the enhancement of the AWA. In the northeast region (north and east of New York state with New York state included), this relationship is the most pronounced. This difference in response between the highest and median PM_{2.5} values indicates the different sensitivities within various per-

centiles of the PM_{2.5} levels. These results are to some extent consistent with those from Porter et al. (2015), which addressed the greater sensitivities to mean daily temperature at the highest concentration percentiles in predicting summertime PM_{2.5}, but with PM_{2.5} sensitivities to temperature peaks entirely in the east due to the regionality of PM_{2.5} speciation.

3.4 Projected PM_{2.5} concentrations due to changes in future AWA

The strong association between PM_{2.5} concentrations and AWA in the current climate prompts us to investigate the extent to which we can utilize a linear regression model to predict changes in PM_{2.5} concentrations from AWA change in a future climate. Employing daily present-day summertime concentrations of PM_{2.5} and AWA for the current climate from the coupled model simulation (GCM2000, 2006–2025) and Eqs. (5)–(6), we derive how much of PM_{2.5}'s interannual variance can be explained by the projection of JJA AWA anomalies onto the daily PM_{2.5}–AWA regression coefficient pattern. The coefficient of determination (R^2) of the linear regression model using simulated PM_{2.5} and AWA for the present climate varies from 0 to 0.75 depending on the grid box (Fig. 8). This means that the projected value (using only AWA changes) captures up to 75 % of the interannual




Figure 5. Composite 500 hPa geopotential height anomaly (contours; positive values are represented by green lines and negative values by magenta lines) and (shaded) corresponding AWA for PM_{2.5} larger than the 90th quantile at sites (denoted by the black dots) (a, d, g) AREN1, (b, e, h) SIPS1 and (c, f, i) LAVO1. The top row shows results using IMPROVE PM_{2.5} and reanalysis AWA, the middle row uses the reanalysis-driven simulated PM_{2.5} (REFC1SD) and reanalysis AWA, and the bottom row uses current-climate-simulated PM_{2.5} and AWA (GCM2000). Stippling indicates the regions that are statistically significant at the 5 % confidence level. Unit: meters for 500 hPa geopotential height and 10^8 m^2 for AWA. The blue outlined area in (a) is the impact region, which is defined as the region of the maximum regression coefficient minus 0.05.

variability in PM_{2.5} over the Great Plains and the west. Wise and Comrie (2005) similarly determined R^2 values of 0.1–0.5 for associations of PM with atmospheric variables across sites in the southwest, and here we see a comparable relationship across the southwest, although these studies use different methodologies as well as considering different time periods. Because of the high correlation coefficients (75 %) this suggests that the regression results reveal the broad population instead of a small number of influential outliers (Cook, 1979). The R^2 measures the part of variance of PM_{2.5} that can be explained by the linear regression model (Kutner et al., 2004).

Next we explore how much of the future change in PM_{2.5} concentrations can be predicted just on the basis of changes in AWA. Using PM_{2.5}–AWA relationships determined from current coupled model output (GCM2000), future PM_{2.5} changes can be estimated by using the linear relationship fitted with the current data and projected change in AWA in the future (as shown in Eqs. 5–7). Here we assume that the linear relationship between the predictors and PM_{2.5} does

not change very much in the future compared to the present to extrapolate the current linear relationship between PM_{2.5} and AWA to the future. Future climate change is simulated to cause an increase in PM_{2.5} concentrations over most of the US if there are no changes in emissions (GCM2100; Fig. 9a). Using the CAM4-Chem fitted slope pattern and regression coefficient, enhanced PM_{2.5} is found over most of the western US and a small area in the northeast (Fig. 9b). The prediction suggests that the significant increase in future PM_{2.5} resulted from AWA changes arising over the western US, which are up to $0.92 \mu\text{g m}^{-3}$. The Great Plains, the south-central part, the Midwest and the southwest show a small decrease in PM_{2.5} in the future, where a negative value of the projection value occurs when projecting the positive anomaly onto it. The projection of PM_{2.5} change in most parts of the south is less reliable because of the low R^2 in this region in the interannual variability metrics (Fig. 8).

In order to investigate how much change in PM_{2.5} can be caused by the change in AWA, the fraction of predicted JJA PM_{2.5} change in total JJA PM_{2.5} change from the simula-




Figure 6. The maximum of the composite AWA distribution for PM_{2.5} larger than the 90th quantile (shaded) (a, c, e, g, i) and the centers of the spatial regression coefficient distribution between PM_{2.5} and AWA (b, d, f, h, j): observations (Obs, first row), current climate from the reanalysis-driven simulation (REFC1SD, second row), current climate from the coupled model simulation (GCM2000, third row), future climate with current emission (GCM2100, fourth row) and future climate with future emission (REFC2, bottom row). At each grid point, the highest composite AWAs anywhere in the domain based on the PM_{2.5} larger than the 90th quantile and the highest regression coefficient between AWA and PM_{2.5} are shown. In (a) and (b), the three representative sites are denoted by the black dots. In (a) and (b) the different shapes (circles or triangles) indicate whether the number of values for every grid that are statistically significant (at the 5% confidence level) is more than 30% or not. The different colors indicate different highest composite AWA and regression coefficients as indicated in the legend. In (c) through (j) the number of values for every grid that are statistically significant at the 5% confidence level is shown (in black contours).




Figure 7. (a) Scatterplot for site AREN1 JJA deseasonalized $\text{PM}_{2.5}$ and the impact region's average AWA with linear and quantile regression results. (b) The subtraction of the 50th-percentile quantile regression slope from the 90th-percentile quantile regression slope between $\text{PM}_{2.5}$ and the impact region's average AWA across all 150 sites in the US (at the 5 % significance level).




Figure 8. $\text{PM}_{2.5}$'s interannual variance explained (R^2) by the linear regression model using the AWA projection value as the explanatory variable with modeled results. Stippling indicates where R^2 is significant (at the 5 % significance level) in model grids.

tion is calculated in Fig. 10. Overall, the maximum values are found in the western US. We infer that AWA can be generally utilized in $\text{PM}_{2.5}$ predictions during the summer in the US, especially over the western US. The impact of AWA change alone on summer $\text{PM}_{2.5}$ concentrations is likely to be quite significant (above 50 %) in the western US, suggesting that AWA is a suitable tool for air quality predictions for most regions where meteorology dominates. These results are somewhat consistent with Tagaris et al. (2007), who showed that the Midwest was modeled to have larger daily average $\text{PM}_{2.5}$ levels in the future, but our signal is in the west. In addition, studies suggest increases in $\text{PM}_{2.5}$ over polluted regions in the future climate caused by intensified stagnation (Liao et al., 2006; Jacob and Winner, 2009), which is consistent with the anticyclonic condition seen with AWA in this study.

Future changes in 500 hPa JJA geopotential height anomaly between the future simulation (GCM2100) relative to the current simulation (GCM2000) (shown in Fig. S2) can account for the $\text{PM}_{2.5}$ variability resulting from changes in AWA. The 500 hPa geopotential height increases everywhere by approximately 25–55 m over the entire US. This strengthened geopotential height can be explained by mid-latitude to high-latitude warming in the future climate. The increased geopotential height at higher latitudes is consistent with other model projections (Vavrus et al., 2017). The increase in 500 hPa height, which shows a distinct anticyclonic pattern centering over the western US and the adjacent ocean, is consistent with changes in a suite of atmospheric variables related to changes in $\text{PM}_{2.5}$ concentrations (Dawson et al., 2007; Tai et al., 2010; Porter et al., 2015).

Increased AWA over most parts of US in the future climate are projected to increase $\text{PM}_{2.5}$ levels over western regions where meteorology dominates fluctuations in $\text{PM}_{2.5}$. This is consistent with some studies reporting increased $\text{PM}_{2.5}$ concentrations due to more common and extended stagnation periods across northern midlatitudes in the future climate if anthropogenic emissions remain constant (Mickley et al., 2004; Liao et al., 2006; Leibensperger et al., 2008; Jacob and Winner, 2009). Similarly in simulations, a mean increase of $0.24 \mu\text{g m}^{-3}$ in summer average $\text{PM}_{2.5}$ levels with the largest growth of $0.93 \mu\text{g m}^{-3}$ in the Midwest has been shown (Tai et al., 2010). An increase in $\text{PM}_{2.5}$ despite globally increasing precipitation is also obtained by using coupled chemistry–climate models, revealing a decreased precipitation on a large scale across polluted regions and seasons (Fang et al., 2011; Kloster et al., 2010). In contrast, Avise et al. (2009) found that changes in meteorology tend to reduce summertime $\text{PM}_{2.5}$ concentrations (approximately $-1 \mu\text{g m}^{-3}$) in most regions with the maximum reductions over the southeastern




Figure 9. (a) Simulated change in JJA PM_{2.5} between simulated future climate (future climate with current emission, GCM2100, 2106–2125 mean) and current climate (current climate from the coupled model simulation, GCM2000, 2006–2025 mean). (b) Predicted JJA PM_{2.5} change using the linear regression model fitted with simulated current PM_{2.5} (GCM2000, 2006–2025 mean). Stippling indicates where R^2 is significant (at the 5 % significance level) at model grids. Unit: $\mu\text{g m}^{-3}$ for PM_{2.5}.




Figure 10. The fraction of predicted JJA PM_{2.5} change using simulated data fitted (current climate from the coupled model simulation, GCM2000, 2006–2025 mean; Fig. 9b divided by Fig. 9a) model accounts for the total JJA PM_{2.5} change from simulations. The fraction that is less than zero is regarded as zero. Stippling indicates where R^2 is significant (at the 5 % significance level) at model grids.

US caused by intensified wet deposition. Furthermore, some models suggest there could be a regional increase in summertime PM_{2.5} over the eastern US due to lower precipitation (Racherla and Adams, 2006), while Tagaris et al. (2007) found that climate change, alone, with no emissions increase or controls affects the US PM_{2.5} concentrations slightly. The discrepancy between studies mentioned above and the results given here is most likely attributable to differences in model formulation. Although earlier studies predict important changes in PM_{2.5} levels in a warming climate throughout the US, there is no consistency across studies (Jacob and Winner, 2009). The unpredictable sensitivity of PM_{2.5} levels to climate change could be explained by the complication of the reliance of different PM_{2.5} elements on climatic variables and the uncertainties in regional boundary layer ventilation

and precipitation as well as the diversity of PM_{2.5} components (Racherla and Adams, 2006; Pye et al., 2009).

It should be noted that the change in surface PM_{2.5} predicted by the future change in AWA using the univariate linear regression models is different from the simulated future change in PM_{2.5} from the constant-emissions run, which simulate the most significant change in the eastern US (Fig. 9a), consistent with the other model projections (Tai et al., 2012). This discrepancy mostly results from the distribution of projected AWA change (Fig. 3). Figure 9b only includes the change caused by the change in AWA. These studies did not account for the change in PM_{2.5} separately due to stagnation, temperature or other meteorological conditions, which could also play a significant part in the PM_{2.5} changes. In addition, the predicting capability of the linear models is limited, and the model only looks at linear relationships between PM_{2.5} levels and AWA. Also, it only looks at the mean of PM_{2.5} levels.

PM_{2.5} generally consists of multiple different aerosols each with different sources and variability; for example, the most important in the US are sulfate, organic matter, elemental carbon, nitrate, ammonium and desert dust. The different PM_{2.5} components respond to meteorological variables differently. The sulfate fraction of PM_{2.5} is predicted to be higher due to faster SO₂ oxidation under a warmer climate, while the nitrate and organic fraction is predicted to be lower due to volatility (Dawson et al., 2007; Kleeman, 2008; Tai et al., 2010). Increased temperatures can lead to higher biogenic emissions of PM_{2.5} precursors including agricultural ammonia, soil NO_x and volatile organic compounds (Pinder et al., 2004; Bertram et al., 2005; Guenther et al., 2006; Riddick et al., 2016). Aqueous-phase sulfate and ammonium nitrate production increase with higher relative humidity (Liao et al., 2006; Dawson et al., 2007). Wildfires are an important source of black and organic carbon and they can increase

or decrease depending on the local changes in climate and land use (Park et al., 2007; Spracklen et al., 2009; Kloster et al., 2012). Future exploration of the different components of aerosols and how each responds to climate could provide more information about the effect on each type, but for these simulations, only PM_{2.5} was output and thus is not available for this study.

4 Conclusions

We employed a univariate linear regression model to determine the correlation of PM_{2.5} levels and AWA on synoptic scales over the US. This analysis demonstrates that PM_{2.5} is positively linked to the local anticyclonic finite-amplitude wave activity over the past 2 decades during JJA, and the high PM_{2.5} concentrations are more sensitive to the AWA than those low ones. The relationship between AWA and PM_{2.5} in model-simulated and observational data agrees in its general pattern and amplitude. These results provide insights into the drivers behind high-PM_{2.5} pollution episodes in the observed record, emphasizing the significance of atmospheric circulation to the pollutant accumulation.

We found that AWA is positively correlated with PM_{2.5} at every available station in the summer using regression analysis in this study. The most prominent relationship between PM_{2.5} and AWA occurs in the Midwestern US for the current climate, while it moves westward in the future climates. The composite AWA for PM_{2.5} larger than its 90th percentile can also demonstrate the positive relationship between PM_{2.5} and AWA. Climate change in the future is likely to cause a response in regional PM_{2.5}. The sensitivities of PM_{2.5} levels to changes in AWA are, generally, more robust for higher percentiles through quantile regression, which is most prominent in the northeastern US. It means that changes to AWA are likely to influence the extent of PM_{2.5} extremes more strongly than they influence moderate PM_{2.5} levels. This study presents new perspectives to explore both the observed and simulated PM_{2.5} responses to climate change. Furthermore, the contrast of observed sensitivities to those simulated by CESM could determine essential model biases relating to the prediction of future PM_{2.5}, potentially offering perceptions into the fundamental mechanistic reasons behind those biases.

The coefficient of determination of the linear regression model using simulated PM_{2.5} and AWA for the present climate is up to 75 % over the Great Plains and the west, which shows that the daily variation in AWA can project up to 75 % of interannual PM_{2.5} variability across the US. These effects suggest that AWA could have significant impacts on the PM_{2.5} levels. Significant regional variation is found in these results, indicating that while the positive association between PM_{2.5} and AWA is generally consistent, the extent to which AWA influences PM_{2.5} is local.

Data availability. PM_{2.5} data are available at <http://views.cira.colostate.edu/fed/QueryWizard/> (Hand et al., 2011). The ERA-Interim reanalysis data sets (<https://www.ecmwf.int/en/newsletter/147/news/era5-reanalysis-production>, Hersbach and Dee, 2016) were retrieved from ECMWF's Meteorological Archival and Retrieval System (MARS). See <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-interim> (ECMWF, 2021) for further details. The CCMI output data (Tilmes et al., 2016) can be downloaded from the Centre for Environmental Data Analysis (<http://data.ceda.ac.uk/badc/wcrp-ccmi/data/CCMI-1/output/>).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-22-7575-2022-supplement>.

Author contributions. YW analyzed the observational data and visualized the model output. YW and NM wrote the paper. PH and WS provided critical feedback and helped shape the research and analysis. GC provided the expertise on local finite-amplitude wave activity. All authors reviewed the paper.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We acknowledge high-performance computing support from Cheyenne (<https://doi.org/10.5065/D6RX99HX>) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation. We also thank Simone Tilmes.

Financial support. This research has been supported by the Nanjing University of Aeronautics and Astronautics (grant no. NS2019037) and the National Science Foundation (grant no. NSF AGS-1608775).

Review statement. This paper was edited by Jianzhong Ma and reviewed by Yuan Wang and one anonymous referee.

References

- Aghedo, A. M., Bowman, K. W., Worden, H. M., Kulawik, S. S., Shindell, D. T., Lamarque, J. F., Faluvegi, G., Parrington, M., Jones, D. B. A., and Rast, S.: The vertical distribution of ozone instantaneous radiative forcing from satellite and chemistry climate models, *J. Geophys. Res.*, 116, D01305, <https://doi.org/10.1029/2010jd014243>, 2011.

- Albrecht, B.: Aerosols, cloud microphysics and fractional cloudiness, *Science*, 245, 1227–1230, 1989.
- Arimoto, R.: Eolian dust and climate: Relationships to sources, tropospheric chemistry, transport and deposition, *Earth Sci. Rev.*, 54, 29–42, 2001.
- Ashley, W. S., Strader, S., Dziubla, D. C., and Haberlie, A.: Driving blind: Weather related vision hazards and fatal motor vehicle crashes, *B. Am. Meteorol. Soc.*, 96, 755–778, <https://doi.org/10.1175/BAMS-D-14-00026.1>, 2015.
- Avise, J., Chen, J., Lamb, B., Wiedinmyer, C., Guenther, A., Salathé, E., and Mass, C.: Attribution of projected changes in summertime US ozone and PM_{2.5} concentrations to global changes, *Atmos. Chem. Phys.*, 9, 1111–1124, <https://doi.org/10.5194/acp-9-1111-2009>, 2009.
- Aw, J. and Kleeman, M. J.: Evaluating the first-order effect of intraannual temperature variability on urban air pollution, *J. Geophys. Res.-Atmos.*, 108, 4365, <https://doi.org/10.1029/2002jd002688>, 2003.
- Bernard, S. M., Samet, J. M., Grambsch, A., Ebi, K. L., and Romieu, I.: The potential impact of climate variability and change on air pollution-related health effects in the United States, *Environ. Health Perspect.*, 109, 199–209, 2001.
- Bertram, T. H., Heckel, A., Richter, A., Burrows, J. P., and Cohen, R. C.: Satellite measurements of daily variations in soil NO_x emissions, *Geophys. Res. Lett.*, 32, L24812, <https://doi.org/10.1029/2005gl024640>, 2005.
- Brasseur, G. P., Schultz, M., Granier, C., Saunio, M., Diehl, T., Botzet, M., Roeckner, E., and Walters, S.: Impact of climate change on the future chemical composition of the global troposphere, *J. Climate*, 19, 3932–3951, 2006.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley Jr., J. A., Hansen, J. E., and Hofmann, D. J.: Climate forcing by anthropogenic aerosols, *Science*, 255, 423–430, 1992.
- Cheng, C. S., Campbell, M., Li, Q., Li, G., Auld, H., Day, N., Pengelly, D., Gingrich, S., and Yap, D.: A synoptic climatological approach to assess climatic impact on air quality in south-central Canada. Part II: future estimates, *Water Air Soil Pollut.*, 182, 117–130, 2007.
- Chen, G. and Plumb, A.: Effective isentropic diffusivity of tropospheric transport, *J. Atmos. Sci.*, 71, 3499–3520, 2014.
- Chen, G., Lu, J., Burrows, D. A., and Leung, L. R.: Local finite-amplitude wave activity as an objective diagnostic of midlatitude extreme weather, *Geophys. Res. Lett.*, 42, 10952–10960, <https://doi.org/10.1002/2015GL066959>, 2015.
- Chen, Z., Xie, X., Cai, J., Chen, D., Gao, B., He, B., Cheng, N., and Xu, B.: Understanding meteorological influences on PM_{2.5} concentrations across China: a temporal and spatial perspective, *Atmos. Chem. Phys.*, 18, 5343–5358, <https://doi.org/10.5194/acp-18-5343-2018>, 2018.
- Cook, R.: Influential Observations in Linear Regression, *J. Am. Stat. Assoc.*, 74, 169–174, <https://doi.org/10.2307/2286747>, 1979.
- Coumou, D., Lehmann, J., and Beckmann, J.: The weakening summer circulation in the Northern Hemisphere mid-latitudes, *Science*, 348, 324–327, 2015.
- Dawson, J. P., Adams, P. J., and Pandis, S. N.: Sensitivity of PM_{2.5} to climate in the Eastern US: a modeling case study, *Atmos. Chem. Phys.*, 7, 4295–4309, <https://doi.org/10.5194/acp-7-4295-2007>, 2007.
- Dawson, J. P., Bloomer, B. J., Winner, D. A., and Weaver, C. P.: Understanding the meteorological drivers of U.S. particulate matter concentrations in a changing climate, *B. Am. Meteorol. Soc.*, 95, 521–532, <https://doi.org/10.1175/BAMS-D-12-00181.1>, 2014.
- Dee, D., Uppala, S., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hersbach, H., Holm, E., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A., MongeSanz, B., Morcrette, J. J., Park, B. K., Peubey, C., Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitars, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteorol. Soc.*, 137, 553–597, 2011.
- ECMWF: ECMWF Reanalysis – Interim (ERA-Interim), European Centre for Medium-Range Weather Forecasts [data set], <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-interim>, last access: 1 September 2021.
- Eyring, V., Lamarque, J. F., Hess, P., Arfeuille, F., Bowman, K., Chipperfield, M. P., Duncan, B., Fiore, A., Gettelman, A., Giorgetta, M. A., Granier, C., Hegglin, M., Kinnison, D., Kunze, M., Langematz, U., Luo, B., Martin, R., Matthes, K., Newman, P. A., Peter, T., Robock, A., Ryerson, T., Saiz-Lopez, A., Salawitch, R., Schultz, M., Shepherd, T. G., Shindell, D., Staehelin, J., Tegtmeier, S., Thomason, L., Tilmes, S., Vernier, J. P., Waugh, D. W., and Young, P. J.: Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) community simulations in support of upcoming ozone and climate assessments, *SPARC Newsletter*, 40, 48–66, 2013.
- Fang, Y., Fiore, A. M., Horowitz, L. W., Gnanadesikan, A., Held, I., Chen, G., Vecchi, G., and Levy, H. I.: The impacts of changing transport and precipitation on pollutant distributions in a future climate, *J. Geophys. Res.*, 116, D18303, <https://doi.org/10.1029/2011JD015642>, 2011.
- Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier Van Der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J. F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J. C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., Van Aardenne, J., Van Der Werf, G. R., and Van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Clim. Change*, 109, 163–190, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- Hack, J. J., Caron, J. M., Yeager, S. G., Oleson, K. W., Holland, M. M., Truesdale, J. E., and Rasch, P. J.: Simulation of the Global Hydrological Cycle in the CCSM Community Atmosphere Model Version 3 (CAM3): Mean Features, *J. Climate*, 19, 2199–2221, <https://doi.org/10.1175/jcli3755.1>, 2006.

- Hand, J., Copeland, S., Day, D., Dillner, A., Indresand, H., Malm, W., McDade, C., Moore, C., Pitchford, M. L., Schichtel, B., and Watson, J.: IMPROVE (Interagency Monitoring of Protected Visual Environments): Spatial and seasonal patterns and temporal variability of haze and its constituents in the United States: Report V June 2011, Coop. Inst. for Res. in the Atmos., Fort Collins, Colo., <http://vista.cira.colostate.edu/Improve/spatial-and-seasonal-patterns-and-temporal-variability-of-haze-and-its-constituents-in-the-united-states-report-v-june-2011> (last access: 1 September 2021), 2011 (data available at: <http://views.cira.colostate.edu/fed/QueryWizard/>, last access: 1 September 2021).
- Hand, J., Schichtel, B., Pitchford, M., Malm, W., and Frank, N.: Seasonal composition of remote and urban fine particulate matter in the United States, *J. Geophys. Res.*, 117, D05209, <https://doi.org/10.1029/2011JD017122>, 2012.
- Hersbach, H. and Dee, D.: ERA5 reanalysis is in production, ECMWF Newsletter, 147, <https://www.ecmwf.int/en/newsletter/147/news/era5-reanalysis-production> (last access: 1 September 2021), 2016.
- Holtzlag, A. A. M. and Boville, B. A.: Local versus non-local boundary-layer diffusion in a global climate model, *J. Climate*, 6, 1825–1842, [https://doi.org/10.1175/1520-0442\(1993\)006<1825:lvnbl>2.0.co;2](https://doi.org/10.1175/1520-0442(1993)006<1825:lvnbl>2.0.co;2), 1993.
- Huang, C. S. and Nakamura, N.: Local finite-amplitude wave activity as a diagnostic of anomalous weather events, *J. Atmos. Sci.*, 73, 211–229, 2016.
- Hunke, E. C. and Lipscomb, W. H.: CICE: The Los Alamos Sea Ice Model, documentation and software, version 4.0, Tech. Rep., LA-CC-06-012, Los Alamos Natl. Lab., Los Alamos, N. M., 2008 (https://csdms.colorado.edu/w/images/CICE_documentation_and_software_user's_manual.pdf, last access: 1 September 2021), 2008.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A framework for collaborative research, *B. Am. Meteorol. Soc.*, 94, 1339–1360, 2013.
- Jacob, D. J. and Winner, D. A.: Effect of climate change on air quality, *Atmos. Environ.*, 43, 51–63, 2009.
- Kappos, A. D., Bruckmann, P., Eikmann, T., Englert, N., Heinrich, U., Hoppe, P., Koch, E., Krause, G. H. M., Kreyling, W. G., Rauchfuss, K., Rombout, P., Schulz-Klemp, V., Thiel, W. R., and Wichmann, H. E.: Health effects of particles in ambient air, *Int. J. Hyg. Environ. Health*, 207, 399–407, 2004.
- Kleeman, M. J.: A preliminary assessment of the sensitivity of air quality in California to global change, *Clim. Change*, 87, S273–S292, <https://doi.org/10.1007/S10584-007-9351-3>, 2008.
- Kloster, S., Dentener, F., Feichter, J., Raes, F., Lohmann, U., Roeckner, E., and Fischer-Bruns, I.: A GCM study of future climate response to aerosol pollution reductions, *Clim. Dynam.*, 34, 1177–1194, 2010.
- Kloster, S., Mahowald, N. M., Randerson, J. T., and Lawrence, P. J.: The impacts of climate, land use, and demography on fires during the 21st century simulated by CLM-CN, *Biogeosciences*, 9, 509–525, <https://doi.org/10.5194/bg-9-509-2012>, 2012.
- Koch, D., Park, J., and Del Genio, A.: Clouds and sulfate are anti-correlated: A new diagnostic for global sulfur models, *J. Geophys. Res.*, 108, 4781, <https://doi.org/10.1029/2003JD003621>, 2003.
- Koenker, R. and Bassett, G.: Regression Quantiles, *Econometrica*, 46, 33–50, 1978.
- Krewski, D., Jerrett, M., Burnett, R. T., Ma, R., Hughes, E., Shi, Y., Turner, M. C., Pope, C. A., Thurston, G., Calle, E. E., and Thun, M. J.: Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality, HEI Research Report, 140, 5–114, <https://pubmed.ncbi.nlm.nih.gov/19627030> (last access: 1 September 2021), 2009.
- Kutner, M. H., Nachtsheim, C. J., Neter, J., and Li, W.: Applied linear statistical models, McGraw-Hill/Irwin, New York, NY, USA, ISBN: 9780073108742, 2004.
- Lamarque, J. F. and Solomon, S.: Impact of changes in climate and halocarbons on recent lower stratosphere ozone and temperature trends, *J. Climate*, 23, 2599–2611, <https://doi.org/10.1175/2010jcli3179.1>, 2010.
- Lamarque, J. F., Kyle, G., Meinshausen, M., Riahi, K., Smith, S., van Vuuren, D., Conley, A., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the Representative Concentration Pathways, *Clim. Change*, 109, 191–212, <https://doi.org/10.1007/s10584-011-0155-0>, 2011a.
- Lamarque, J. F., McConnell, J. R., Shindell, D. T., Orlando, J. J., and Tyndall, G. S.: Understanding the drivers for the 20th century change of hydrogen peroxide in Antarctic ice-cores, *Geophys. Res. Lett.*, 38, L04810, <https://doi.org/10.1029/2010gl045992>, 2011b.
- Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, <https://doi.org/10.5194/gmd-5-369-2012>, 2012.
- Leibensperger, E. M., Mickley, L. J., and Jacob, D. J.: Sensitivity of US air quality to mid-latitude cyclone frequency and implications of 1980–2006 climate change, *Atmos. Chem. Phys.*, 8, 7075–7086, <https://doi.org/10.5194/acp-8-7075-2008>, 2008.
- Liao, H., Chen, W. T., and Seinfeld, J. H.: Role of climate change in global predictions of future tropospheric ozone and aerosols, *J. Geophys. Res.*, 111, D12304, <https://doi.org/10.1029/2005JD006852>, 2006.
- Liu, J., Zheng, Y., Geng, G., Hong, C., Li, M., Li, X., Liu, F., Tong, D., Wu, R., Zheng, B., He, K., and Zhang, Q.: Decadal changes in anthropogenic source contribution of PM_{2.5} pollution and related health impacts in China, 1990–2015, *Atmos. Chem. Phys.*, 20, 7783–7799, <https://doi.org/10.5194/acp-20-7783-2020>, 2020.
- Liu, Y., Zhao, N., Vanos, J., and Cao, G.: Effects of synoptic weather on ground-level PM_{2.5} concentrations in the United States, *Atmos. Environ.*, 148, 297–305, 2017.
- Lu, J., Chen, G., Leung, L. R., Burrows, D. A., Yang, Q., Sakaguchi, K., and Hagos, S.: Toward the dynamical convergence on the Jet-stream in aquaplanet AGCMs, *J. Climate*, 28, 6763–6782, 2015.
- Mahowald, N., Ward, D., Kloster, S., Flanner, M. G., Heald, C. L., Heavens, N. G., Hess, P. G., Lamarque, J. F., and Chuang, P.

- Y.: Aerosol impacts on climate and biogeochemistry, *Ann. Rev. Environ. Resour.*, 36, 45–74, 2011.
- Martineau, P., Chen, G., and Burrows, D. A.: Wave events: climatology, trends, and relationship to Northern Hemisphere winter blocking and weather extremes, *J. Climate*, 30, 5675–5697, 2017.
- Methven, J.: Wave activity for large-amplitude disturbances described by the primitive equations on the sphere, *J. Atmos. Sci.*, 70, 1616–1630, 2013.
- Michel, C. and Rivière, G.: The link between Rossby wave breakings and weather regime transitions, *J. Atmos. Sci.*, 68, 1730–1748, 2011.
- Mickley, L. J., Jacob, D. J., Field, B. D., and Rind, D.: Effects of future climate change on regional air pollution episodes in the United States, *Geophys. Res. Lett.*, 31, L24103, <https://doi.org/10.1029/2004gl021216>, 2004.
- Nakamura, H., Nakamura, M., and Anderson, J.: The role of high- and low-frequency dynamics in blocking formation, *Mon. weather rev.*, 125, 2074–2093, 1997.
- Nakamura, N. and Zhu, D.: Finite-amplitude wave activity and diffusive flux of potential vorticity in eddy-mean flow interaction, *J. Atmos. Sci.*, 67, 2701–2716, 2010.
- Nakamura, N. and Solomon, A.: Finite-amplitude wave activity and mean flow adjustments in the atmospheric general circulation. Part II: Analysis in the isentropic coordinate, *J. Atmos. Sci.*, 68, 2783–2799, 2011.
- Neale, R. B., Richter, J. H., and Jochum, M.: The impact of convection on ENSO: From a delayed oscillator to a series of events, *J. Climate*, 21, 5904–5924, 2008.
- Neale, R. B., Richter, J. H., Conley, A. J., Park, S., Lauritzen, P. H., Gettelman, A., Williamson, D. L., Rasch, P. J., Vavrus, S. J., Taylor, M. A., Collins, W. D., Zhang, M., and Lin, S.-J.: Description of the NCAR Community Atmosphere Model (CAM 4.0), NCAR Tech. Note, TN-485, 212 pp., Natl. Cent. for Atmos. Res., Boulder, Colo, http://www.cesm.ucar.edu/models/cesm1.0/cam/docs/description/cam4_desc.pdf (last access: 1 September 2021), 2010.
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddes, J., Heald, C. L., Hoffman, F., Lamarque, J., Mahowald, N., Niu, G., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang, Z., Zeng, X., Zeng, X., and Decker, M.: Technical Description of version 4.0 of the Community Land Model (CLM), NCAR Technical Note NCAR/TN-478+STR, <https://doi.org/10.5065/D6FB50WZ>, 2010.
- Painter, T. H., Barrett, A. P., Landry, C. C., Neff, J. C., Cassidy, M. P., Lawrence, C. R., McBride, K. E., and Farmer, G. L.: Impact of disturbed desert soils on duration of mountain snow cover, *Geophys. Res. Lett.*, 34, L12502, <https://doi.org/10.1029/2007GL030284>, 2007.
- Park, R. J., Jacob, D. J., and Logan, J. A.: Fire and biofuel contributions to annual mean aerosol mass concentrations in the United States, *Atmos. Environ.*, 41, 7389–7400, 2007.
- Pfahl, S. and Wernli, H.: Quantifying the relevance of atmospheric blocking for co-located temperature extremes in the Northern Hemisphere on (sub-)daily time scales, *Geophys. Res. Lett.*, 39, L12807, <https://doi.org/10.1029/2012GL052261>, 2012.
- Phalithonkiat, P., Hess, P. G. M., Grigoriu, M. D., Samorodnitsky, G., Sun, W., Beaudry, E., Tilmes, S., Deushi, M., Josse, B., Plummer, D., and Sudo, K.: Extremal dependence between temperature and ozone over the continental US, *Atmos. Chem. Phys.*, 18, 11927–11948, <https://doi.org/10.5194/acp-18-11927-2018>, 2018.
- Pinder, R. W., Pekney, N. J., Davidson, C. I., and Adams, P. J.: A process-based model of ammonia emissions from dairy cows: Improved temporal and spatial resolution, *Atmos. Environ.*, 38, 1357–1365, 2004.
- Porter, W. C., Heald, C. L., Cooley, D., and Russell, B.: Investigating the observed sensitivities of air-quality extremes to meteorological drivers via quantile regression, *Atmos. Chem. Phys.*, 15, 10349–10366, <https://doi.org/10.5194/acp-15-10349-2015>, 2015.
- Pye, H. O. T., Liao, H., Wu, S., Mickley, L. J., Jacob, D. J., Henze, D. K., and Seinfeld, J. H.: Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, *J. Geophys. Res.*, 114, D01205, <https://doi.org/10.1029/2008JD010701>, 2009.
- Racherla, P. N. and Adams, P. J.: Sensitivity of global tropospheric ozone and fine particulate matter concentrations to climate change, *J. Geophys. Res.*, 111, D24103, <https://doi.org/10.1029/2005JD006939>, 2006.
- Rasch, P. and Kristjánsson, J.: A comparison of the CCM3 model climate using diagnosed and predicted condensate parameterizations, *J. Climate*, 11, 1587–1614, 1998.
- Richter, J. H. and Rasch, P. J.: Effects of convective momentum transport on the atmospheric circulation in the Community Atmosphere Model, version 3, *J. Climate*, 21, 1487–1499, 2008.
- Riddick, S., Ward, D., Hess, P., Mahowald, N., Massad, R., and Holland, E.: Estimate of changes in agricultural terrestrial nitrogen pathways and ammonia emissions from 1850 to present in the Community Earth System Model, *Biogeosciences*, 13, 3397–3426, <https://doi.org/10.5194/bg-13-3397-2016>, 2016.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G. K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, <https://doi.org/10.1175/JCLI-D-11-00015.1>, 2011.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissel, A., and Andreae, M. O.: Flood or drought: How do aerosols affect precipitation? *Science*, 321, 1309–1313, 2008.
- Ryoo, J. M., Kaspi, Y., Waugh, D. W., Kiladis, G. N., Waliser, D. E., Fetzner, E. J., and Kim, J.: Impact of Rossby wave breaking on U. S. West Coast winter precipitation during ENSO events, *J. Climate*, 26, 6360–6382, <https://doi.org/10.1175/JCLI-D-12-00297.1>, 2013.
- Shen, L. and Mickley, L. J.: Seasonal prediction of US summertime ozone using statistical analysis of large scale climate patterns, *P. Natl. Acad. Sci. USA*, 114, 2491–2496, 2017.
- Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P., Hecht, M., Jayne, S., Jochum, M., Large, W., Lindsay, K., Maltrud, M.,

- Norton, N., Peacock, S., Vertenstein, M., and Yeager, S.: The Parallel Ocean Program (POP) reference manual, Ocean component of the Community Climate System Model (CCSM), LANL Tech. Report, LAUR-10-01853, 141 pp., <https://www.cesm.ucar.edu/models/cesm2/ocean/doc/sci/POPRefManual.pdf> (last access: 1 September 2021), 2010.
- Spracklen, D. V., Mickley, L. J., Logan, J. A., Hudman, R. C., Yevich, R., Flannigan, M. D., and Westerling, A. L.: Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous aerosol concentrations in the western United States, *J. Geophys. Res.-Atmos.*, 114, D20301, <https://doi.org/10.1029/2008jd010966>, 2009.
- Sun, W., Hess, P., and Tian, B.: The response of the equatorial tropospheric ozone to the Madden-Julian Oscillation in TES satellite observations and CAM-chem model simulation, *Atmos. Chem. Phys.*, 14, 11775–11790, <https://doi.org/10.5194/acp-14-11775-2014>, 2014.
- Sun, W., Hess, P., and Liu, C.: The impact of meteorological persistence on the distribution and extremes of ozone, *Geophys. Res. Lett.*, 44, 1545–1553, 2017.
- Sun, W., Hess, P., Chen, G., and Tilmes, S.: How waviness in the circulation changes surface ozone: a viewpoint using local finite-amplitude wave activity, *Atmos. Chem. Phys.*, 19, 12917–12933, <https://doi.org/10.5194/acp-19-12917-2019>, 2019.
- Tagaris, E., Manomaiphiboon, K., Liao, K. J., Leung, L. R., Woo, J. H., He, S., Amar, P., and Russell, A. G.: Impacts of global climate change and emissions on regional ozone and fine particulate matter concentrations over the United States, *J. Geophys. Res.*, 112, D14312, <https://doi.org/10.1029/2006jd008262>, 2007.
- Tai, A. P. K., Mickley, L. J., and Jacob, D. J.: Correlations between fine particulate matter (PM_{2.5}) and meteorological variables in the United States: Implications for the sensitivity of PM_{2.5} to climate change, *Atmos. Environ.*, 44, 3976–3984, 2010.
- Tai, A. P. K., Mickley, L. J., Jacob, D. J., Leibensperger, E. M., Zhang, L., Fisher, J. A., and Pye, H. O. T.: Meteorological modes of variability for fine particulate matter (PM_{2.5}) air quality in the United States: implications for PM_{2.5} sensitivity to climate change, *Atmos. Chem. Phys.*, 12, 3131–3145, <https://doi.org/10.5194/acp-12-3131-2012>, 2012.
- Takaya, K. and Nakamura, H.: A formulation of a phase-independent wave-activity flux for stationary and migratory quasigeostrophic eddies on a zonally varying basic flow, *J. Atmos. Sci.*, 58, 608–627, 2001.
- Thishan Dharshana, K. G., Kravtsov, S., and Kahl, J. D. W.: Relationship between synoptic weather disturbances and particulate matter air pollution over the United States, *J. Geophys. Res.-Atmos.*, 115, D24219, <https://doi.org/10.1029/2010jd014852>, 2010.
- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Marsh, D., Garcia, R. R., Smith, A. K., Neely, R. R., Conley, A., Vitt, F., Val Martin, M., Tanimoto, H., Simpson, I., Blake, D. R., and Blake, N.: Representation of the Community Earth System Model (CESM1) CAM4-chem within the Chemistry-Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 9, 1853–1890, <https://doi.org/10.5194/gmd-9-1853-2016>, 2016 (data available at: <http://data.ceda.ac.uk/badc/wcrp-ccmi/data/CCMI-1/output/>, last access: 1 September 2021).
- Titchner, H. A. and Rayner, N. A.: The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2:1. sea ice concentrations, *J. Geophys. Res.-Atmos.*, 119, 2864–2889, <https://doi.org/10.1002/2013JD020316>, 2014.
- VanCuren, R. and Gustin, M. S.: Identification of sources contributing to PM_{2.5} and ozone at elevated sites in the western U.S. by receptor analysis: Lassen Volcanic National Park, California, and at Great Basin National Park, Nevada, *Sci. Total Environ.*, 530, 505–518, <https://doi.org/10.1016/j.scitotenv.2015.03.091>, 2015.
- Vavrus, S. J., Wang, F., Martin, J. E., Francis, J. A., Peings, Y., and Cattiaux, J.: Changes in North American atmospheric circulation and extreme weather: Influence of Arctic amplification and Northern Hemisphere snow cover, *J. Climate*, 30, 4317–4333, 2017.
- Wang, X. and Zhang, R.: Effects of atmospheric circulations on the interannual variation in PM_{2.5} concentrations over the Beijing–Tianjin–Hebei region in 2013–2018, *Atmos. Chem. Phys.*, 20, 7667–7682, <https://doi.org/10.5194/acp-20-7667-2020>, 2020.
- Wang, Y., Le, T., Chen, G., Yung, Y. L., and Jiang, J. H.: Reduced European aerosol emissions suppress winter extremes over northern Eurasia, *Nat. Clim. Chang.*, 10, 225–230, <https://doi.org/10.1038/s41558-020-0693-4>, 2020.
- Wang, Z., Chen, G., Gu, Y., Zhao, B., and Liou, K.: Large-scale meteorological control on the spatial pattern of winter-time PM_{2.5} pollution over China, *Atmos. Sci. Lett.*, 20, e938, <https://doi.org/10.1002/asl.938>, 2019.
- Wise, E. K. and Comrie, A. C.: Meteorologically adjusted urban air quality trends in the Southwestern United States, *Atmos. Environ.*, 39, 2969–2980, <https://doi.org/10.1016/j.atmosenv.2005.01.024>, 2005.
- Woolings, T. J., Hoskins, B. J., Blackburn, M., and Berrisford, P.: A new Rossby wave breaking interpretation of the North Atlantic Oscillation, *J. Atmos. Sci.*, 65, 609–626, <https://doi.org/10.1175/2007JAS2347.1>, 2008.
- Wu, S., Mickley, L. J., Leibensperger, E. M., Jacob, D. J., Rind, D., and Streets, D. G.: Effects of 2000–2050 global change on ozone air quality in the United States, *J. Geophys. Res.*, 113, D06302, <https://doi.org/10.1029/2007JD008917>, 2008.
- Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Centre general circulation model, *Atmos. Ocean*, 33, 407–446, 1995.
- Zhang, M., Lin, W., Bretherton, C. S., Hack, J. J., and Rasch, P. J.: A modified formulation of fractional stratiform condensation rate in the NCAR Community Atmospheric Model (CAM2), *J. Geophys. Res.*, 108, 4035, <https://doi.org/10.1029/2002JD002523>, 2003.