

Anisotropic active Brownian particle with a fluctuating propulsion forceJean-Luc Thiffeault ^{*}*Department of Mathematics, University of Wisconsin–Madison, Madison, Wisconsin 53706, USA*Jiajia Guo *Department of Mathematics, University of Michigan, Ann Arbor, Michigan 48109, USA*

(Received 23 February 2021; accepted 23 June 2022; published 15 July 2022)

The active Brownian particle (ABP) model describes a swimmer, synthetic or living, whose direction of swimming is a Brownian motion. The swimming is due to a propulsion force, and the fluctuations are typically thermal in origin. We present a two-dimensional model where the fluctuations arise from nonthermal noise in a propelling force acting at a single point, such as that due to a flagellum. We take the overdamped limit and find several modifications to the traditional ABP model. Since the fluctuating force causes a fluctuating torque, the diffusion tensor describing the process has a coupling between translational and rotational degrees of freedom. An anisotropic particle also exhibits a mass-dependent noise-induced drift, which does not disappear in the overdamped limit. We show that these effects have measurable consequences for the long-time diffusivity of active particles, in particular adding a contribution that is independent of where the force acts.

DOI: [10.1103/PhysRevE.106.L012603](https://doi.org/10.1103/PhysRevE.106.L012603)

Modeling swimming microorganisms is a challenge, since biological entities resist a simple, uniform description. Nevertheless we need models to explain physical observations and develop intuition, and the hope is that the models capture some essential aspect of an organism's behavior. For microswimmers, most modeling efforts impose some randomness to the motion. The simplest approach is to use a fixed propulsion speed, together with a random reorientation mechanism. The random reorientation comes in two main flavors: a run-and-tumble process where the organism makes large excursions and changes its orientation sporadically, [1–7], and a Brownian process where the direction of swimming gradually varies [8–14]. Both of these models have their place, but in this Letter we focus primarily on the latter, Brownian approach.

In this Letter we present a simple model of random microorganism motion where the swimmer is propelled by a fluctuating force acting at a point (Fig. 1). The randomness is built into the force as a covariance matrix, and is not due to interactions with the medium (though such interactions could be included as well). Our goal is to derive effective equations of motion for this simple configuration, which is meant to represent an organism with a single flagellum. The resulting equations have some points of commonality with the well-known active Brownian particle model (ABP), but differ in crucial ways. In particular, there is an inherent coupling between translational and rotation diffusivities. In addition, there is a noise-induced drift that is present regardless of which stochastic interpretation (Itô or Stratonovich) is used.

The stochastic equations (SDEs) for the two-dimensional (2D) ABP model are [8–14]

$$\dot{\mathbf{x}} = (U + \sqrt{2D_{\parallel}} \dot{w}_{\parallel}) \mathbf{p}_{\parallel} + \sqrt{2D_{\perp}} \mathbf{p}_{\perp} \dot{w}_{\perp}, \quad (1a)$$

$$\dot{\phi} = \Omega + \sqrt{2D_r} \dot{w}_r. \quad (1b)$$

The swimmer is moving at constant speed U in the direction $\mathbf{p}_{\parallel}(\phi)$ and rotating at constant angular speed Ω . The translational noises $\sqrt{2D_{\parallel}} \dot{w}_{\parallel}$ and $\sqrt{2D_{\perp}} \dot{w}_{\perp}$ are respectively along $(\mathbf{p}_{\parallel})$ and perpendicular $(\mathbf{p}_{\perp})$ to the direction of swimming, and the rotational noise $\sqrt{2D_r} \dot{w}_r$ affects the swimming direction. The $w_i(t)$ are independent standard Wiener processes. Equation (1) has been very successful in modeling the swimming and collective behavior of many microorganisms [15–21]. The noises are often taken to be due to thermal fluctuations, in which case they satisfy the Einstein-Smoluchowski relations $D_i = (\beta \sigma_i)^{-1}$, for $i \in \{\parallel, \perp, r\}$, where σ_i are the components of the diagonal grand resistance tensor and β is the inverse temperature $1/k_B T$.

In this Letter we derive a modified ABP model by assuming that the noise is due to a fluctuating propulsion force acting at a single point on the particle, rather than a thermal bath. We will find several new effects: a new noise-induced drift term, as well as a diffusion matrix that couples the rotational and angular degrees of freedom.

A particle subjected to a fluctuating force $\mathbf{f}(\phi, t) = (F_{\parallel} + \sqrt{2E_{\parallel}} \dot{w}_{\parallel}) \mathbf{p}_{\parallel} + (F_{\perp} + \sqrt{2E_{\perp}} \dot{w}_{\perp}) \mathbf{p}_{\perp}$ acting at the point $\ell \mathbf{p}_{\parallel}$ with respect to the center of reaction [22] obeys the Langevin equations

$$m \dot{\mathbf{u}} = -\mathbb{K} \cdot \mathbf{u} + \mathbf{f}, \quad I \dot{\omega} = -\sigma_r \omega + \tau, \quad (2)$$

where m is the mass, I the moment of inertia, $\mathbf{u}$ the velocity, ω the angular velocity, and $\mathbb{K} = \mathbb{Q} \cdot \text{diag}(\sigma_{\parallel}, \sigma_{\perp}) \cdot \mathbb{Q}^T$ the

*jeanluc@math.wisc.edu




FIG. 1. A 2D particle with orientation ϕ subject to a time-dependent force $\mathbf{f}$ acting at a point $\ell \mathbf{p}_{\parallel}(\phi)$ with respect to its center of reaction.

resistance matrix, with $\mathbb{Q}(\phi)$ a 2×2 rotation matrix. The force exerts a torque $\tau(t) = \ell (F_{\perp} + \sqrt{2E_{\perp}} \dot{w}_{\perp})$ [23].

A brief note on the validity of Eq. (2) is in order. We follow many authors such as [24,25] and use a linear damping law in Eq. (2), which as first pointed out by Lorentz [26] is strictly only valid in the limit where the fluid density is less than the particle density [27–32]. The theory could be extended to allow for a memory kernel, the so-called Basset-Boussinesq integral term [33,34], but then the process is non-Markovian and we cannot recover a simple Fokker-Planck equation as detailed below. Nevertheless, we expect that this memory effect is unlikely to *decrease* correlations, and so the effects presented here might be modified but would not disappear.

We rewrite the system (2) in the standard form

$$\frac{d\hat{\mathbf{x}}}{dt} = \hat{\mathbf{u}}, \quad \frac{d\hat{\mathbf{u}}}{dt} = \hat{\mathbb{B}} \cdot (\hat{\mathbf{U}} - \hat{\mathbf{u}}) + \hat{\Sigma} \cdot \dot{\mathbf{w}}, \quad (3)$$

where $\hat{\mathbf{x}} = (\mathbf{x}, \phi)$, $\hat{\mathbf{u}} = (\mathbf{u}, \omega)$, $\dot{\mathbf{w}} = (\dot{w}_{\parallel}, \dot{w}_{\perp})$, $\hat{\mathbb{B}} = \text{diag}(\mathbb{K}/m, \sigma_r/I)$, $\hat{\mathbf{U}} = (\mathbf{U}, \Omega) = (\mathbb{K}^{-1} \cdot \mathbf{F}, \ell F_{\perp}/\sigma_r)$, and

$$\hat{\Sigma} = \begin{pmatrix} (\sqrt{2E_{\parallel}}/m) \mathbf{p}_{\parallel} & (\sqrt{2E_{\perp}}/m) \mathbf{p}_{\perp} \\ 0 & \sqrt{2E_{\perp}} \ell / I \end{pmatrix}. \quad (4)$$

The third components of hat-wearing vectors and matrices pertain to angular quantities.

Typically, in the overdamped limit (small mass, or large drag) the term $d\hat{\mathbf{u}}/dt$ in (3) is neglected, resulting in the equation

$$\frac{d\hat{\mathbf{x}}}{dt} = \hat{\mathbf{U}} + \hat{\mathbb{B}}^{-1} \cdot \hat{\Sigma} \cdot \dot{\mathbf{w}}. \quad (5)$$

This recovers something close to the standard ABP model (1), except that here there are only two rather than three independent noises: the rotational noise is correlated to the translational noise, since the former is caused by the torque of the latter. We will see the consequences of this correlation below.

But first note that taking the overdamped limit in this way is suspicious. The underdamped equations (3) have the same form independent of the interpretation given to the stochastic product (i.e., Itô or Stratonovich), even though the noise appears multiplicative at first glance. However, the noise coupling matrix $\hat{\mathbb{B}}^{-1} \cdot \hat{\Sigma}$ in Eq. (5) leads to a nonvanishing drift term when the stochastic product is interpreted in the

Stratonovich sense [35]. This suggests that Eq. (5) has a uniquely defined noise-induced drift term [36–38], but the naive way of passing from (3) to (5) does not tell us what form it should take.

A more systematic approach is required to find the missing noise-induced drift term Eq. (5). Instead working with SDEs, we take the overdamped limit of the Fokker-Planck equation for the probability density $p(\hat{\mathbf{x}}, \hat{\mathbf{u}}, t)$ corresponding to Eq. (3) (see [36,39–41] for an SDE approach):

$$\varepsilon^2 \partial_t p + \varepsilon \nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbf{u}} p) + \varepsilon \nabla_{\hat{\mathbf{u}}} \cdot (\hat{\mathbb{B}} \cdot \hat{\mathbf{U}} p) = \mathcal{L} p, \quad (6)$$

where ε is a formal expansion parameter, with $\varepsilon \rightarrow 0$ the overdamped limit, and

$$\mathcal{L} p := \nabla_{\hat{\mathbf{u}}} \cdot (\hat{\mathbb{B}} \cdot \hat{\mathbf{u}} p) + \nabla_{\hat{\mathbf{u}}} \otimes \nabla_{\hat{\mathbf{u}}} : (\hat{\mathbb{E}} p) \quad (7)$$

with $\hat{\mathbb{E}} := \frac{1}{2} \hat{\Sigma} \cdot \hat{\Sigma}^{\top}$ [42]. The parameter ε expresses the long-time and large-scale rescalings of t and $\hat{\mathbf{x}}$ for which the $\hat{\mathbf{u}}$ degrees of freedom equilibrate.

Now we proceed order by order with an expansion $p = p_0 + \varepsilon p_1 + \dots$. At leading order we have $\mathcal{L} p_0 = 0$, with solution $p_0 = P(\hat{\mathbf{x}}, t) \varphi(\hat{\mathbf{x}}, \hat{\mathbf{u}})$, where P is yet to be determined and $\varphi(\hat{\mathbf{x}}, \hat{\mathbf{u}})$ is the invariant density for an Ornstein-Uhlenbeck process [43]:

$$\varphi = (2\pi)^{-3} (\det \hat{\mathbb{A}})^{-1/2} \exp\left(-\frac{1}{2} \hat{\mathbf{u}} \cdot \hat{\mathbb{A}}^{-1} \cdot \hat{\mathbf{u}}\right). \quad (8)$$

Here the symmetric positive-definite matrix $\hat{\mathbb{A}}(\hat{\mathbf{x}})$ is the unique solution to the continuous-time Lyapunov equation [44]

$$\hat{\mathbb{B}} \cdot \hat{\mathbb{A}} + \hat{\mathbb{A}} \cdot \hat{\mathbb{B}}^{\top} = 2\hat{\mathbb{E}}, \quad (9)$$

where in our case $\hat{\mathbb{B}} = \hat{\mathbb{B}}^{\top}$. When $\hat{\mathbb{B}}$ commutes with $\hat{\mathbb{E}}$, as occurs for thermal fluctuations, the solution to (9) is $\hat{\mathbb{A}} = \hat{\mathbb{E}} \cdot \hat{\mathbb{B}}^{-1}$; this is not the case here, and we find instead

$$\hat{\mathbb{A}} = \hat{\mathbb{Q}} \cdot \begin{pmatrix} \frac{E_{\parallel}}{m\sigma_{\parallel}} & 0 & 0 \\ 0 & \frac{E_{\perp}}{m\sigma_{\perp}} & \frac{2E_{\perp}\ell}{m\sigma_r + I\sigma_{\perp}} \\ 0 & \frac{2E_{\perp}\ell}{m\sigma_r + I\sigma_{\perp}} & \frac{E_{\perp}\ell^2}{I\sigma_r} \end{pmatrix} \cdot \hat{\mathbb{Q}}^{\top} \quad (10)$$

where $\hat{\mathbb{Q}}(\phi) = \text{diag}(\mathbb{Q}, 1)$ is a 3×3 rotation matrix about the third axis.

At the next order in ε , we have $\mathcal{L} p_1 = \nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbf{u}} \varphi P) - \hat{\mathbf{u}} \cdot \hat{\mathbb{A}}^{-1} \cdot \hat{\mathbb{B}} \cdot \hat{\mathbf{U}} \varphi P$. The solution can be written in two pieces, $p_1 = p_1^{(1)} + p_1^{(2)}$, with $p_1^{(1)} = (\nabla_{\hat{\mathbf{x}}} P - \hat{\mathbf{U}} \cdot \hat{\mathbb{B}}^{\top} \cdot \hat{\mathbb{A}}^{-1} P) \cdot \hat{\chi}^{(1)}$ and $p_1^{(2)} = -\frac{1}{2} P \nabla_{\hat{\mathbf{x}}} \hat{\mathbb{A}}^{-1} : \hat{\chi}^{(2)}$, where $\hat{\chi}^{(1)}$ and $\hat{\chi}^{(2)}$ satisfy

$$\mathcal{L} \hat{\chi}^{(1)} = \hat{\mathbf{u}} \varphi, \quad \mathcal{L} \hat{\chi}^{(2)} = \hat{\mathbf{u}} \hat{\mathbf{u}} \varphi. \quad (11)$$

It is easy to solve for $\hat{\chi}^{(1)} = -\hat{\mathbb{A}} \cdot \hat{\mathbb{B}}^{-\top} \cdot \hat{\mathbb{A}}^{-1} \cdot \hat{\mathbf{u}} \varphi$; $\hat{\chi}^{(2)}$ is harder to solve for in general. However, we shall not need its precise expression in our derivation.

At the next and final order in ε we get from Eq. (6) $\mathcal{L} p_2 = \nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbf{u}} p_1) + \nabla_{\hat{\mathbf{u}}} \cdot (\hat{\mathbb{B}} \cdot \hat{\mathbf{U}} p_1) + \partial_t p_0$, to which we need only apply a solvability condition by integrating over $\hat{\mathbf{u}}$ (denoted by angle brackets):

$$\partial_t P = -\nabla_{\hat{\mathbf{x}}} \cdot \langle \hat{\mathbf{u}} p_1 \rangle. \quad (12)$$

To evaluate the average $\langle \hat{\mathbf{u}} p_1 \rangle$, first note that the adjoint to $\mathcal{L}$ is

$$\mathcal{L}^* g = -\hat{\mathbf{u}} \cdot \hat{\mathbb{B}}^{\top} \cdot \nabla_{\hat{\mathbf{u}}} g + \hat{\mathbb{E}} : \nabla_{\hat{\mathbf{u}}} \otimes \nabla_{\hat{\mathbf{u}}} g, \quad (13)$$

which satisfies $\langle g\mathcal{L}f \rangle = \langle (\mathcal{L}^*g)f \rangle$ for functions f and g vanishing as $|\hat{\mathbf{u}}| \rightarrow \infty$. Multiplying the $\hat{\chi}^{(1)}$ equation in (11) by $\hat{\mathbf{u}}$, we have

$$\langle \hat{\mathbf{u}} \mathcal{L} \hat{\chi}^{(1)} \rangle = \langle \hat{\mathbf{u}} \hat{\mathbf{u}} \varphi \rangle = \hat{\mathbb{A}}. \quad (14)$$

But then using the adjoint property in (14) gives

$$\langle (\mathcal{L}^* \hat{\mathbf{u}}) \hat{\chi}^{(1)} \rangle = \langle (-\hat{\mathbb{B}} \cdot \hat{\mathbf{u}}) \hat{\chi}^{(1)} \rangle = -\hat{\mathbb{B}} \cdot \langle \hat{\mathbf{u}} \hat{\chi}^{(1)} \rangle$$

from which we obtain $\langle \hat{\mathbf{u}} \hat{\chi}^{(1)} \rangle = -\hat{\mathbb{B}}^{-1} \cdot \hat{\mathbb{A}}$. We can play a similar trick with the $\hat{\chi}^{(2)}$ equation to obtain $\langle \hat{\mathbf{u}} \hat{\chi}^{(2)} \rangle = -\hat{\mathbb{B}}^{-1} \cdot \langle \hat{\mathbf{u}} \hat{\mathbf{u}} \hat{\mathbf{u}} \varphi \rangle$, where the fourth moment for the Gaussian φ is easily obtained. We have thus evaluated the required average $\langle \hat{\mathbf{u}} \hat{\chi}^{(2)} \rangle$ without needing to solve for $\hat{\chi}^{(2)}$.

After a lengthy but straightforward calculation we find $\langle \hat{\mathbf{u}} p_1 \rangle = \hat{\mathbf{U}} P - \nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbb{A}} P) \cdot \hat{\mathbb{B}}^{-\top}$, which we insert back into (12) to finally obtain

$$\partial_t P + \nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbf{U}} P) = \nabla_{\hat{\mathbf{x}}} \cdot (\nabla_{\hat{\mathbf{x}}} \cdot (\hat{\mathbb{A}} P) \cdot \hat{\mathbb{B}}^{-\top}). \quad (15)$$

We rewrite (15) in a more convenient form and obtain the first main result of this Letter:

$$\partial_t P + \nabla_{\mathbf{x}} \cdot ((\mathbf{U} + \mathbf{V})P) + \partial_{\phi}(\Omega P) = \nabla_{\hat{\mathbf{x}}} \otimes \nabla_{\hat{\mathbf{x}}} : (\hat{\mathbb{D}} P), \quad (16)$$

where the noise-induced drift [37,38,41,45–48] is

$$\mathbf{V} = \frac{2\ell E_{\perp}(\sigma_{\parallel}^{-1} - \sigma_{\perp}^{-1})}{\sigma_{\tau}(1 + I\sigma_{\perp}/m\sigma_{\tau})} \mathbf{p}_{\parallel} \quad (17)$$

and the translational-rotational grand diffusion tensor is

$$\hat{\mathbb{D}} = \hat{\mathbb{Q}} \cdot \begin{pmatrix} D_{\parallel} & 0 & 0 \\ 0 & D_{\perp} & \sqrt{D_{\perp} D_r} \\ 0 & \sqrt{D_{\perp} D_r} & D_r \end{pmatrix} \cdot \hat{\mathbb{Q}}^{\top} \quad (18)$$

with $D_{\parallel} = E_{\parallel}/\sigma_{\parallel}^2$, $D_{\perp} = E_{\perp}/\sigma_{\perp}^2$, and $D_r = E_{\perp}\ell^2/\sigma_{\tau}^2$. The diffusion tensor couples translational and rotational noises. Our result is closely related to [47], but here the induced drift is due to angular dependence rather than spatial inhomogeneity.

To go back and compare to the overdamped result Eq. (5) obtained by simply neglecting the particle mass, the Fokker-Planck equation (16) implies the SDE

$$\frac{d}{dt} \begin{pmatrix} \mathbf{x} \\ \phi \end{pmatrix} = \begin{pmatrix} \mathbf{U} + \mathbf{V} \\ \Omega \end{pmatrix} + \sqrt{2\hat{\mathbb{D}}} \cdot \dot{\mathbf{w}}, \quad (19)$$

where $\sqrt{2\hat{\mathbb{D}}} = \hat{\mathbb{B}}^{-1} \cdot \hat{\Sigma}$. Note the additional drift $\mathbf{V}$. The drift $\mathbf{V}$ implies that the particle appears to swim at a constant speed as in the ABP model (1) for long times, even for $\mathbf{U} = 0$. The drift $\mathbf{V}$ is only present when the fluctuating force exerts a torque; it is an inertial effect that vanishes for isotropic particles ($\sigma_{\parallel} = \sigma_{\perp}$). It does *not* vanish for zero mass, since it involves the ratio I/m .

It is natural to form Péclet numbers based on the advective time $a/|\mathbf{V}|$ and diffusive times $a^2/D_{\perp}$ and $1/D_r$, with a the particle size:

$$\text{Pe}_{\perp} = \frac{|\mathbf{V}|a}{D_{\perp}} = \frac{2a\ell\sigma_{\perp}^2}{\sigma_{\tau}} \frac{|\sigma_{\parallel}^{-1} - \sigma_{\perp}^{-1}|}{1 + I\sigma_{\perp}/m\sigma_{\tau}} \sim \frac{\ell}{a},$$

$$\text{Pe}_r = \frac{|\mathbf{V}|}{D_r a} = \frac{2\sigma_{\tau}}{a\ell} \frac{|\sigma_{\parallel}^{-1} - \sigma_{\perp}^{-1}|}{1 + I\sigma_{\perp}/m\sigma_{\tau}} \sim \frac{a}{\ell}.$$

$\text{Pe}_{\perp}$ is not large, but also not necessarily small. Pe_r is a dimensionless correlation length that diverges as $\ell \rightarrow 0$, since the rotational diffusivity then vanishes.

We can compute the long-time effective diffusivity of the active particle. Here there are two new effects: the noise-induced drift $\mathbf{V}$ and the coupling terms $\sqrt{D_{\perp} D_r}$ in the grand diffusion tensor $\hat{\mathbb{D}}$. Recall that $\hat{\mathbf{x}} = (\mathbf{x}, \phi)$, so $\hat{x}_3 = \phi$. The overdamped Fokker-Planck equation (16) for $P(\hat{\mathbf{x}}, t)$ is

$$\begin{aligned} & \partial_t P + W_i \partial_{x_i} P + \Omega \partial_{\phi} P \\ & = \partial_{x_i} \partial_{x_j} (D_{ij} P) + 2\partial_{x_i} \partial_{\phi} (\hat{D}_{i3} P) + \partial_{\phi}^2 (D_r P), \end{aligned} \quad (20)$$

where $\mathbf{W} = \mathbf{U} + \mathbf{V} = W \mathbf{p}_{\parallel}$ is the total drift, and indices are summed over 1,2. To find the effective diffusivity, we rescale (20) to focus on large scales $\delta^{-1} \sim \ell^{-1}$ and long times δ^{-2} , with δ a small parameter. We let $\partial_t \rightarrow \partial_t + \delta^2 \partial_T$, and $\partial_{\mathbf{x}} \rightarrow \partial_{\mathbf{x}} + \delta \partial_X$ and expand $P = \mathcal{P}(X, T) + \delta P_1(\phi; X, T) + \delta^2 P_2(\phi; X, T) + \dots$, where we anticipated the functional dependencies to abridge the derivation. (Our approach is equivalent to that of [49], which averages over angles, or [4], which expands P in harmonics.) At order δ^1 we have $D_r \partial_{\phi}^2 P_1 - \Omega \partial_{\phi} P_1 = W_i \partial_{x_i} \mathcal{P} - 2\partial_{x_i} \partial_{\phi} (\hat{D}_{i3} \mathcal{P})$, with a simple solution linear in $\cos \phi$ and $\sin \phi$. At order δ^2 we have the solvability condition

$$\begin{aligned} \partial_T \mathcal{P} &= \langle W_i (W_j - 2\partial_{\phi} \hat{D}_{j3}) / D_r + D_{ij} \rangle \partial_{x_i} \partial_{x_j} \mathcal{P} \\ &=: D_{\text{eff}} \nabla_X^2 \mathcal{P}, \end{aligned} \quad (21)$$

where angle brackets are repurposed for angular averaging, and the effective diffusivity is

$$D_{\text{eff}} = \frac{1}{2}(D_{\parallel} + D_{\perp}) + \tilde{D}, \quad (22a)$$

$$\tilde{D} := \frac{W D_r}{2(D_r^2 + \Omega^2)} \left(W + \frac{2E_{\perp}\ell}{\sigma_{\perp}\sigma_{\tau}} \right). \quad (22b)$$

Equation (21) displays the expected long-time isotropy of the probability density. Compare $\tilde{D}$ to $U^2/2D_r$ for the ABP model (1) [8,50–54].

The new diffusivity $\tilde{D}$ combines contributions from the swimming U , from the noise-induced drift $\mathbf{V}$, and from the coupling terms in $\hat{\mathbb{D}}$. From here we set $\mathbf{U} = \Omega = D_{\parallel} = 0$ to highlight the new effects: the particle is “shaking its hips” but would be a nonswimmer if not for the noise-induced drift; see also [55] for a deterministic version. (The swimmer is a “treadmill” or reciprocal swimmer that does not strictly swim, but only diffuses [56–58].) In that case after using (17) Eq. (22b) becomes

$$\tilde{D}_0 = \frac{2D_{\perp}(1 + I\sigma_{\parallel}/m\sigma_{\tau})}{(1 + I\sigma_{\perp}/m\sigma_{\tau})^2} \frac{\sigma_{\perp}}{\sigma_{\parallel}} \left(\frac{\sigma_{\perp}}{\sigma_{\parallel}} - 1 \right). \quad (23)$$

The form (23) for $\tilde{D}_0$ has two striking features. First, it is negative for particles with $\sigma_{\perp} < \sigma_{\parallel}$, so that it hinders diffusion. In fact, the combination $\tilde{D}_0 + \frac{1}{2}D_{\perp}$ attains a minimum of zero for $\sigma_{\perp} = \sigma_{\parallel}/(2 + I\sigma_{\parallel}/m\sigma_{\tau})$. A particle satisfying this relation can only diffuse through $D_{\parallel}$ and thermal noise.

The second striking feature of (23) is that it is independent of ℓ . This is a paradox: for $\ell = 0$, we have $\mathbf{V} = 0$ and $\hat{D}_{i3} = 0$, so none of the effects mentioned here occur. The resolution




FIG. 2. The mean-squared displacement (variance) $\text{Var} x$, averaged over 5000 oblate nonswimming ($\mathbf{U} = \mathbf{\Omega} = 0$) active particles for various values of ℓ . The upper dotted line is $4 \times \frac{1}{2}(D_{\parallel} + D_{\perp})t$, and the bottom dashed line is $4D_{\text{eff}}t$. As ℓ becomes smaller, there is a longer transient before the behavior begins to follow Eq. (21). This transient diverges as $\ell \rightarrow 0$. Parameter values are $m = I = 0.05$, $\sigma_{\parallel} = 2$, $E_{\perp} = \sigma_{\perp} = \sigma_r = 1$, $E_{\parallel} = 0$.

is that there is a transient of duration $D_r^{-1} = \sigma_r^2/E_{\perp}\ell^2 \sim \delta^{-2}$ before the long-time form (21) applies, and this transient becomes infinite as $\ell \rightarrow 0$. This transient can be seen in the simulations of the full inertial equations (2) in Fig. 2,

It is important to note that the ratio $\tilde{D}_0/D_{\perp}$ is rarely negligible: all the dimensionless ratios appearing on the right of

Eq. (26) are typically of order one. The transient timescale D_r^{-1} can be estimated by $a^2/D_{\perp}$, where a is the particle size; if D_r^{-1} is very long, then $D_{\perp}$ was likely negligible to begin with. The modifications discussed in this paper are thus likely to be relevant in many applications.

So why have these types of corrections not been observed? Many authors simulate the ABP model directly, since the inertial equations (2) are expensive to solve due the small step size required, in which case the new effects are ruled out. Particle anisotropy is also seldom considered. Experimentally, diffusivities are measured directly from the distributions of displacements, and so any connection between the rotational and translational diffusivities is typically lost. One approach might be to obtain the covariance matrix $\hat{\mathbf{A}}$ directly, by measuring the correlations between translational and rotational velocities. A nonzero correlation would indicate a coupling as predicted here.

In future work we will generalize the derivation to arbitrary three-dimensional active particles [25,59], with the fluctuating force not necessarily applied on an axis of symmetry. There are several other possible extensions, such as the inclusion of multiple forces and torques acting on the body. The consequences to swim pressure [60,61], run-and-tumble dynamics [1,4], non-Newtonian swimming [62], velocity-dependent friction [63], and particle interactions [53,64] also remain to be investigated.

The authors thank Saverio Spagnolie, Hongfei Chen, Scott Hottovy, Christina Kurzthaler, Eric Lauga, Cesare Nardini, John Wettlaufer, and Ehud Yariv for helpful comments and discussions.

-
- [1] G. Subramanian and D. L. Koch, Critical bacterial concentration for the onset of collective swimming, *J. Fluid Mech.* **632**, 359 (2009).
 - [2] R. W. Nash, R. Adhikari, J. Tailleur, and M. E. Cates, Run-and-Tumble Particles with Hydrodynamics: Sedimentation, Trapping, and Upstream Swimming, *Phys. Rev. Lett.* **104**, 258101 (2010).
 - [3] K. Martens, L. Angelani, R. D. Leonardo, and L. Bocquet, Probability distributions for the run-and-tumble bacterial dynamics: An analogy to the Lorentz model, *Eur. Phys. J. E* **35**, 84 (2012).
 - [4] M. E. Cates and J. Tailleur, When are active Brownian particles and run-and-tumble particles equivalent? Consequences for motility-induced phase separation, *Europhys. Lett.* **101**, 20010 (2013).
 - [5] J. Elgeti and G. Gompper, Run-and-tumble dynamics of self-propelled particles in confinement, *Europhys. Lett.* **109**, 58003 (2015).
 - [6] B. Ezhilan, R. Alonso-Matilla, and D. Saintillan, On the distribution and swim pressure of run-and-tumble particles in confinement, *J. Fluid Mech.* **781**, R4 (2015).
 - [7] M. Lee, K. Szuttor, and C. Holm, A computational model for bacterial run-and-tumble motion, *J. Chem. Phys.* **150**, 174111 (2019).
 - [8] F. Peruani and L. G. Morelli, Self-Propelled Particles with Fluctuating Speed and Direction of Motion in Two Dimensions, *Phys. Rev. Lett.* **99**, 010602 (2007).
 - [9] S. van Teeffelen and H. Löwen, Dynamics of a Brownian circle swimmer, *Phys. Rev. E* **78**, 020101(R) (2008).
 - [10] A. Baskaran and M. C. Marchetti, Hydrodynamics of self-propelled hard rods, *Phys. Rev. E* **77**, 011920 (2008).
 - [11] P. Romanczuk and L. Schimansky-Geier, Brownian Motion with Active Fluctuations, *Phys. Rev. Lett.* **106**, 230601 (2011).
 - [12] P. Romanczuk, M. Bär, W. Ebeling, B. Lindner, and L. Schimansky-Geier, Active Brownian particles, *Eur. Phys. J.: Spec. Top.* **202**, 1 (2012).
 - [13] C. Kurzthaler, S. Leitmann, and T. Franosch, Intermediate scattering function of an anisotropic active Brownian particle, *Sci. Rep.* **6**, 36702 (2016).
 - [14] C. Kurzthaler and T. Franosch, Intermediate scattering function of an anisotropic Brownian circle swimmer, *Soft Matter* **13**, 6396 (2017).
 - [15] B.-q. Ai, Q.-y. Chen, Y.-f. He, F.-g. Li, and W.-r. Zheng, Rectification and diffusion of self-propelled particles in a two-dimensional corrugated channel, *Phys. Rev. E* **88**, 062129 (2013).

- [16] A. P. Solon, Y. Fily, A. Baskaran, M. E. Cates, Y. Kafri, M. Kardar, and J. Tailleur, Pressure is not a state function for generic active fluids, *Nat. Phys.* **11**, 673 (2015).
- [17] A. Zöttl and H. Stark, Emergent behavior in active colloids, *J. Phys.: Condens. Matter* **28**, 253001 (2016).
- [18] C. G. Wagner, M. F. Hagan, and A. Baskaran, Steady-state distributions of ideal active Brownian particles under confinement and forcing, *J. Stat. Mech.* (2017) 043203.
- [19] G. S. Redner, M. F. Hagan, and A. Baskaran, Structure and Dynamics of a Phase-Separating Active Colloidal Fluid, *Phys. Rev. Lett.* **110**, 055701 (2013).
- [20] J. Stenhammar, D. Marenduzzo, R. Allen, and M. E. Cates, Phase behaviour of active Brownian particles: The role of dimensionality, *Soft Matter* **10**, 1489 (2014).
- [21] H. Chen and J.-L. Thiffeault, Shape matters: a Brownian microswimmer in a channel, *J. Fluid Mech.* **916**, 38 (2021).
- [22] J. Happel and H. Brenner, *Low Reynolds number hydrodynamics* (Martinus Nijhoff, The Hague, 1983).
- [23] We assume for simplicity that the center of mass coincides with the center of reaction.
- [24] A. J. Majda and P. R. Kramer, Stochastic mode reduction for particle-based simulation methods for complex microfluid systems, *SIAM J. Appl. Math.* **64**, 401 (2004).
- [25] S. Delong, F. Balboa Usabiaga, and A. Donev, Brownian dynamics of confined rigid bodies, *J. Chem. Phys.* **143**, 144107 (2015).
- [26] H. A. Lorentz, *Lessen over Theoretische Natuurkunde. VoL V. Kinetische Problemen* (E. J. Brill, Leiden, 2011).
- [27] E. H. Hauge and A. Martin-Löf, Fluctuating hydrodynamics and Brownian motion, *J. Stat. Phys.* **7**, 259 (1973).
- [28] E. J. Hinch, Application of the Langevin equation to fluid suspensions, *J. Fluid Mech.* **72**, 499 (1975).
- [29] D. Dürr, S. Goldstein, and J. L. Lebowitz, A mechanical model of Brownian motion, *Commun. Math. Phys.* **78**, 507 (1981).
- [30] J.-N. Roux, Brownian particles at different times scales: A new derivation of the Smoluchowski equation, *Physica A* **188**, 526 (1992).
- [31] L. Bocquet and J. Piasecki, Microscopic derivation of non-Markovian thermalization of a Brownian particle, *J. Stat. Phys.* **87**, 1005 (1997).
- [32] A. Donev and E. Vanden-Eijnden, Dynamic density functional theory with hydrodynamic interactions and fluctuations, *J. Chem. Phys.* **140**, 234115 (2014).
- [33] A. B. Basset, *Treatise on Hydrodynamics* (Deighton Bell, London, 1888), Vol. 2, Chap. 22, pp. 285–297.
- [34] J. Boussinesq, *Théorie Analytique de la Chaleur* (L'École Polytechnique, Paris, 1903), Vol 2, p. 224.
- [35] B. Øksendal, *Stochastic Differential Equations*, 6th ed. (Springer, Berlin, 2003).
- [36] R. Kupferman, G. A. Pavliotis, and A. M. Stuart, Itô versus Stratonovich white-noise limits for systems with inertia and colored multiplicative noise, *Phys. Rev. E* **70**, 036120 (2004).
- [37] A. W. C. Lau and T. C. Lubensky, State-dependent diffusion: Thermodynamic consistency and its path integral formulation, *Phys. Rev. E* **76**, 011123 (2007).
- [38] O. Farago, Noise-induced drift in two-dimensional anisotropic systems, *Phys. Rev. E* **96**, 042141 (2017).
- [39] S. Bo and A. Celani, White-noise limit of nonwhite nonequilibrium processes, *Phys. Rev. E* **88**, 062150 (2013).
- [40] G. A. Pavliotis, *Stochastic Processes and Applications* (Springer, Berlin, 2014).
- [41] S. Hottovy, A. McDaniel, G. Volpe, and J. Wehr, The Smoluchowski-Kramers limit of stochastic differential equations with arbitrary state-dependent friction, *Commun. Math. Phys.* **336**, 1259 (2015).
- [42] The tensor $\mathbb{E}$ has zero determinant, indicative of a degenerate parabolic problem since there are fewer noises than equations. In practice this is inconsequential, since we can add a bit of thermal noise to remove the degeneracy.
- [43] H. Risken, *The Fokker-Planck Equation: Methods of Solution and Applications*, 2nd ed. (Springer, Berlin, 1996).
- [44] The solution of this matrix problem is implemented as `LyapunovSolve` in *Mathematica*, `sylvester` in *Matlab*, and `scipy.linalg.solve_continuous_lyapunov` in *Python*.
- [45] P. S. Grassia, E. J. Hinch, and L. C. Nitsche, Computer simulations of Brownian motion of complex systems, *J. Fluid Mech.* **282**, 373 (1995).
- [46] S. Hottovy, G. Volpe, and J. Wehr, Noise-induced drift in stochastic differential equations with arbitrary friction and diffusion in the Smoluchowski-Kramers limit, *J. Stat. Phys.* **146**, 762 (2012).
- [47] S. Hottovy, G. Volpe, and J. Wehr, Thermophoresis of Brownian particles driven by coloured noise, *Europhys. Lett.* **99**, 60002 (2012).
- [48] G. Volpe and J. Wehr, Effective drifts in dynamical systems with multiplicative noise: a review of recent progress, *Rep. Prog. Phys.* **79**, 053901 (2016).
- [49] R. N. Zia and J. F. Brady, Single-particle motion in colloids: Force-induced diffusion, *J. Fluid Mech.* **658**, 188 (2010).
- [50] J. R. Howse, R. A. L. Jones, A. J. Ryan, T. Gough, R. Vafabakhsh, and R. Golestanian, Self-Motile Colloidal Particles: From Directed Propulsion to Random Walk, *Phys. Rev. Lett.* **99**, 048102 (2007).
- [51] B. Lindner and E. M. Nicola, Diffusion in different models of active Brownian motion, *Eur. Phys. J.: Spec. Top.* **157**, 43 (2008).
- [52] R. Golestanian, Anomalous Diffusion of Symmetric and Asymmetric Active Colloids, *Phys. Rev. Lett.* **102**, 188305 (2009).
- [53] E. Fodor, C. Nardini, M. E. Cates, J. Tailleur, P. Visco, and F. van Wijland, How Far from Equilibrium Is Active Matter? *Phys. Rev. Lett.* **117**, 038103 (2016).
- [54] L. Caprini and U. M. B. Marconi, Inertial self-propelled particles, *J. Chem. Phys.* **154**, 024902 (2021).
- [55] J.-L. Thiffeault, Moving forward by shaking sideways, *Symmetry* **14**, 620 (2022).
- [56] D. G. Crowdy and Y. Or, Two-dimensional point singularity model of a low-Reynolds-number swimmer near a wall, *Phys. Rev. E* **81**, 036313 (2010).
- [57] E. Lauga, Enhanced Diffusion by Reciprocal Swimming, *Phys. Rev. Lett.* **106**, 178101 (2011).
- [58] K. Obuse and J.-L. Thiffeault, A low-Reynolds-number treadmilling swimmer near a semi-infinite wall, in *IMA Volume on Natural Locomotion in Fluids and on Surfaces: Swimming, Flying, and Sliding*, The IMA Volumes in Mathematics and its Applications, edited by S. Childress, A. Hosoi, W. W. Schultz, and J. Wang (Springer, New York, 2012), pp. 197–206.

- [59] I. M. Ilie, W. J. Briels, and W. K. den Otter, An elementary singularity-free rotational Brownian dynamics algorithm for anisotropic particles, *J. Chem. Phys.* **142**, 114103 (2015).
- [60] S. C. Takatori, W. Yan, and J. F. Brady, Swim Pressure: Stress Generation in Active Matter, *Phys. Rev. Lett.* **113**, 028103 (2014).
- [61] S. C. Takatori and J. F. Brady, Swim stress, motion, and deformation of active matter: effect of an external field, *Soft Matter* **10**, 9433 (2014).
- [62] C. Datt and G. J. Elfring, Active Particles in Viscosity Gradients, *Phys. Rev. Lett.* **123**, 158006 (2019).
- [63] U. Erdmann, W. Ebeling, L. Schimansky-Geier, and F. Schweitzer, Brownian particles far from equilibrium, *Eur. Phys. J. B* **15**, 105 (2000).
- [64] N. K. Marath and J. S. Wettlaufer, Hydrodynamic interactions and the diffusivity of spheroidal particles, *J. Chem. Phys.* **151**, 024107 (2019).