

RECEIVED: November 8, 2021

REVISED: March 1, 2022

ACCEPTED: March 14, 2022

PUBLISHED: April 13, 2022

Open F-branes

Machiko Hatsuda^{a,b} and Warren Siegel^{c,1}

^a*Department of Radiological Technology, Faculty of Health Science, Juntendo University, Yushima, Bunkyo-ku, Tokyo 113-0034, Japan*

^b*KEK Theory Center, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan*

^c*C.N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, NY, U.S.A.*

E-mail: mhatsuda@juntendo.ac.jp, siegel@insti.physics.sunysb.edu

ABSTRACT: We include in F-theory, through open Type I F-theory branes (F-branes), string theories with $N = 1$ supersymmetry, both Type I and heterotic. Type I branes are distinguished from Type II by worldvolume parity projection. The same open Type I branes describe both open Type I superstrings and closed heterotic upon different sectionings from F-branes to worldsheets, while closed Type I superstrings arise from closed Type I branes. (Type II superstrings come from closed Type II branes, as described previously.)

F-theory manifests the exceptional-group U-duality symmetry, with all massless bosonic fields in a single gauge coset. This coset branches to the usual bosonic supergravity fields upon sectioning. We examine in detail the simple case of $D = 3$ F-theory: parity projection reduces the Type II coset $SL(5)/SO(3,2)$ to the Type I coset $SO(3,3)/SO(2,1)^2 = SL(4)/SO(2,2)$.

KEYWORDS: F-Theory, P-Branes, String Duality, Supersymmetry and Duality

ARXIV EPRINT: [2110.13010](https://arxiv.org/abs/2110.13010)

¹<http://insti.physics.sunysb.edu/~siegel/plan.html>.

Contents

1	Type II F-theory	1
1.1	Couplings and massless fields	1
1.2	Sectioning	2
2	Type I F-theory	2
2.1	Geometry	2
2.2	Parity	4
2.3	Symmetries	5
3	M-theory	7
4	T-theory	8
4.1	Type II	8
4.2	3D	10
4.3	Type I	12
5	3D F-theory	13
5.1	Type II	13
5.2	Type I and heterotic	15
5.3	Stringy supergravities	16
6	Conclusions	18
A	Cosets	19
B	$F \rightarrow T$ and M	19

1 Type II F-theory

1.1 Couplings and massless fields

In T-duality, (a component of) the spacetime metric is transformed into its inverse (up to B -field dependence). This implies the usual $R \rightarrow 1/R$ transformation, as the metric of the compactified dimensions can be expressed in terms of “angles” as $ds^2 = R^2 d\theta^2$. More properly, the gravitational constant (in Einstein frame, or α' in string frame) can always be absorbed into the metric, reappearing through its vacuum value. Thus R and α' appear in the combination R^2/α' , so actually $R^2/\alpha' \rightarrow \alpha'/R^2$ (and R and α' always appear in this combination in $\text{AdS}_5 \times S^5$).

In S-duality (in Type IIB), the dilaton appears as part of a metric of scalars, which transforms into its inverse in the same way. (Only the coset is different.) Now it’s the string coupling g that appears as the vacuum value. Thus α' and g , and their dualities, appear in

a completely analogous way in the field theory action, but differently in the first-quantized string action. Also, in compactifying 11D supergravity to 10D Type IIA, the dilaton, and thus g , arises from a component of the metric associated with the compactification radius R .

In $D < 10$ (after field-theory duality transformations), all the compactification scalars merge into a larger coset space (U-duality). The purpose of F-theory is to exhibit this (spontaneously broken) symmetry before compactification. The metric, scalars, and other massless bosonic fields then appear as parts of a higher-dimensional gauge field [1, 2] and their field theoretical formulation is the exceptional field theory [3–7]. As these symmetries relate the various superstring theories, F-theory is meant to unify them perturbatively.

1.2 Sectioning

The limit $R \rightarrow \infty$ (or 0) makes winding modes (or momenta) infinitely massive, for each dimension of toroidal compactification. The Virasoro ($\mathcal{S}$) section condition is the remnant of the $L_0 = \bar{L}_0$ constraint that enforces the decompactification limit in any combination of R 's going to 0 or ∞ . This constraint appears already in T-theory, the manifestly T-duality covariant formulation of string theory [8]. Solving the constraint chooses the vacuum, producing either Type IIA or B superstring theory via spontaneous breakdown. There is also an intermediate Virasoro condition, linear in both zero and nonzero modes, that acts as a first-order form of the selfduality condition on the field strength of X .

The generalization [1, 2] appears in F-theory, as a vector in the “spatial” directions of the worldvolume [9]. Thus string theories related by S-duality (different vacua for Type IIB) are also chosen. F-theory also has a Gauss ($\mathcal{U}$) section condition [9] that mutually constrains spacetime and the worldvolume. It's the zero-mode part of Gauss's law for the gauge field X .

2 Type I F-theory

2.1 Geometry


As for strings, we'll classify branes as Types I and II, corresponding to the number of supersymmetries:

- (I) Like Type I strings, the states of Type I branes, both open and closed, are constrained to be (graded) symmetric under σ_1 reflection (“parity”). In particular, this is true for open branes because of boundary conditions. These open branes describe both heterotic and open Type I strings, while the closed branes describe closed Type I strings.
- (II) Type II branes are only closed, and have no parity projection. As described in previous papers, these describe Type IIA and B strings (as well as M-theory) [9–17].

For simplicity, we consider closed branes to have the topology (for fixed τ) of a torus, and open branes to have the topology of a torus of 1 less dimension $\times$ a line segment. (The line segment σ_1 is sometimes described as the simple orbifold S^1/Z_2 .) Thus we discuss branes with only 0 (closed) or 2 (open) boundaries. “Internal symmetry” degrees of freedom live on the boundaries; in general these contribute to the worldvolume energy-momentum tensor (generalized Virasoro operators).

Our description of heterotic strings in terms of open branes will follow that of Horáva and Witten [18], with a few important differences:

- (1) Our branes are fundamental, not solitons. This should allow perturbative S-matrix calculations in the higher dimensions.
- (2) Upon double dimensional reduction, the Lorentz symmetry of spacetime was found tied to that of the worldvolume (as applied to Θ); for us the 2 are the same before compactification. This is related to the fact that STU symmetries are manifest without compactification, by construction.
- (3) Instead of orbifolding spacetime (S^1/Z_2), which in turn orbifolds the embedded worldvolume, we orbifold only the worldvolume. This has several advantages:
 - (a) The F-branes can end on space-filling branes, not just 9-branes. So ordinary Type I superstrings can be obtained, not just Type IA (I'). Thus both $E_8 \times E_8$ and $SO(32)$ are allowed. (But this does not make obvious why Type I $E_8 \times E_8$ is not allowed.)
 - (b) We derive Type I and heterotic strings by worldvolume compactification, not (explicitly) spacetime (although the latter may be implied by constraints). However, since the worldvolume (σ) derivatives represent the same group as the spacetime (X) derivatives, their covariantizations both have vielbeins that are elements of the coset space containing the bosonic spacetime fields [9]. Thus the effects of both worldvolume reduction and the parity symmetry that projects Type II to Type I F-branes can be seen directly on the bosonic fields, without the need to examine spacetime compactification.
 - (c) The worldvolume of the closed F-brane (“donut”) can’t be compactified along its long(est) circle (without double compactification), unlike the open F-brane (“cylinder”). So open F-branes can yield either Type I open or heterotic (closed), while closed F-branes can give only Type I closed:



Internal symmetry regions are in red. We have illustrated only 2 of the worldvolume dimensions, σ_1 and σ_2 (and not τ nor the other σ 's).

For the open F-brane, σ_1 is the coordinate for the line segment, and σ_2 is one of the toroidal coordinates. For the closed F-brane, σ_1 is for the major circumference, and σ_2 is for the minor one, because closed branes can form by joining ends of open ones.

Heterotic and open Type I strings follow from open branes by 2 different choices of section: if σ_2 is killed by sectioning (along with the rest of the toroidal coordinates), we get an open string, and each boundary is compressed to a point. On the other hand, if σ_1 is eliminated (along with the extra toroidal coordinates), we get a closed string, and the 2 boundaries are compressed into the bulk of the string, together.

Gravity arising from open F-branes might appear unusual, but a similar situation occurs in string theory, where the open-closed string 2-point function implies that closed string fields (in particular, the metric) already are contained in the open string field (the metric from classically massive spin-2 fields) [19].

To treat $SO(32)$ vs. $E_8 \times E_8$ by sectioning, we can choose the boundaries of open F-branes to live in 496 extra dimensions. The (boundary) background metric and 2-form for this space can be chosen to give the current algebras for these 2 groups with appropriate level number, in a way that manifests these symmetries, but spontaneously breaks $SO(496)$ [20].

2.2 Parity

For (super)strings, Type II is reduced to Type I by constraining *states* (or background fields, not just the action) to be invariant under worldsheet parity, $\sigma \rightarrow -\sigma$. This is sometimes called a type of orbifolding, but is more a type of projection, like GSO (which is for the worldsheet symmetry of 2π rotation, which affects only fermions). In T-theory, this is accompanied by switching the independent X 's for left and right modes, to preserve their (anti)selfduality condition. The selfdual and anti-selfdual currents are $\partial_\tau X^N \hat{\eta}_{NM} \pm \partial_\sigma X^N \eta_{NM}$ where $\hat{\eta}_{MN}$ and η_{MN} are the double Minkowski metric and the $O(D,D)$ invariant metric respectively. The selfduality condition is that the anti-selfdual current is 0. It relates the dual coordinate X_m and the usual coordinate X^m in the co/contravariant basis $M = ({}_m, {}^m)$. The selfduality condition is preserved under the worldsheet parity transformation $\sigma \rightarrow -\sigma$ together with interchanging the left and right coordinates in the left/right basis.

In F-theory the target space coordinate X and the worldvolume coordinates σ are different representations of the exceptional symmetry groups, and currents are field strengths $\partial_\sigma X$. A naive parity transformation would violate the selfduality constraint on the X field strengths. (For spacetime dimension $D > 1$, worldvolume dimension d is always even.) So instead we need a rotation as an even number of reflections. For consistency with Type I and heterotic strings, this should change the sign of σ_1 but not τ nor σ_2 , so it's most symmetry-preserving to change signs for *all* of σ *except* τ and σ_2 . This rotation then implies the same rotation on X .

For example, for $D = 3$ ($d = 6$), where in the Lagrangian formalism we have σ_m and X^{mn} (antisymmetric), $\mathcal{P}$ (arity) is represented on the indices m, n by the sign changes

$$\mathcal{P} : \quad (0, 2) \rightarrow +, \quad (1, 3, 4, 5) \rightarrow - \quad (2.1)$$

as a rotation in the Lagrangian symmetry $L = \text{SO}(3, 3)$. This breaks that symmetry to $\text{SO}(2, 2)$.

2.3 Symmetries

This parity (as well as boundary conditions), since it distinguishes only σ_2 (as well as τ), effectively reduces symmetries of the Hamiltonian by 1 worldvolume dimension, $\Delta d = 1$. Generally σ , including τ , is a bispinor: $\sigma^{(\alpha\beta)}$, $\sigma_{\dot{\alpha}\dot{\beta}}$, or $\sigma^{[\alpha\beta]}$ [13]:

H	SO	SL	H'(SU)	
Real	1	2	3	$X^{(\alpha\beta)}$ SO
Complex	0		4	$X^{\alpha\dot{\beta}}$ SU
Pseudo-real	7	6	5	$X^{[\alpha\beta]}$ USp
	$\sigma^{(\alpha\beta)}$	$\sigma_{\dot{\alpha}\dot{\beta}}$	$\sigma^{[\alpha\beta]}$	
	$Y^{(\alpha'\beta')}$	$Y^{\alpha'\dot{\beta}'}$	$Y^{[\alpha'\beta']}$	


Since fermions are representation of the tangent space symmetry $H \subset G$, bosonic coordinates are bispinors of H in the superalgebra. Numbers are spacetime dimensions of corresponding string theories. The primed objects are for extended supersymmetry for that dimension, with additional D' -dimensional spacetime coordinates Y . The (spacetime) spinor size increases by a factor of 2 from one oval to the next. In the oval for 3 and 4 X 's are real and complex representations for $D=3$ and 4 theories respectively. As the exceptional group G -symmetry in the Hamiltonian formalism is enlarged to F -symmetry in the Lagrangian formalism, its tangent space symmetry H is also enlarged to L -symmetry. Going from the Lagrangian to the Hamiltonian formalism breaks L -symmetry (always a GL group) to H by picking the matrix coefficient of τ in the matrix σ to be a group metric for $\mathcal{G} = \text{SO}, \text{SL}$, or Sp . (See also appendix A.) Doing the same for σ_2 yields 2 of the same type of group metric, reducing the H group $\mathcal{G}(2n)$ to $H_I = \mathcal{G}(n)^2 = \mathcal{G}_+ \times \mathcal{G}_-$, where $\mathcal{G}(n)$ is the usual covering group of $\text{SO}(D-1, 1)$, and $\pm$ will eventually have the usual string identifications of left and right.

For Type II theories F -theory, which has the most symmetry, is broken to either M -theory (which has $(D+1)$ -dimensional $N = 1$ supergravity as its low-energy limit) or T -theory (with manifest T -duality) by solving either an $\mathcal{S}$ or $\mathcal{U}$ sectioning condition (reducing both X and σ), or S -theory (the standard formulation of superstring theory) by solving both.

$$\begin{array}{ccc}
 & \text{FII} & \\
 \begin{array}{c} \textcolor{blue}{S} \\ \swarrow \end{array} & & \begin{array}{c} \searrow \textcolor{blue}{U} \\ \text{TII} \end{array} \\
 \text{M} & & \\
 \begin{array}{c} \textcolor{blue}{U} \\ \swarrow \end{array} & & \begin{array}{c} \searrow \textcolor{blue}{S} \\ \text{SIIB} \end{array} \\
 & \text{SIIA} &
 \end{array} \quad (2.2)$$

Since only $\mathcal{U}$ sectioning forces reduction of σ , the brane pictures appearing above (cylinder and donut) apply to both F-theory and M-theory, while the string pictures (worldsheet only) apply to both T-theory and S-theory.

F-symmetry has the Dynkin diagram [14]:



$D+D' = 10$ is the maximal case, which has 12 nodes. The black node is dropped unless $D' = 0$. Remove node M or T for M or T-theory, both for S-theory. Remove $\mathcal{H}$ for the Hamiltonian subgroup of any of these Lagrangian groups. Y, X, λ (parameters for X 's gauge invariance), σ , and $\mathcal{H}-\sigma$ are the nodes for their representations. The subgroup L of the F/L coset is a Wick rotation of the maximal compact subgroup of F. (It also corresponds to the covering group of the Lorentz group $SO(D-1,1)$, but with twice the argument, due to doubled spinor indices on supersymmetry. Embedding the $SO(D,D)$ of Type II T-theory in E_{D+1} was also considered in [21], but there E_{D+1} was considered as a symmetry of the superstring supergravity of $D+1$, not D , dimensions.)

The effect of going from the Lagrangian formalism to the Hamiltonian is to eliminate τ , and thus reduce the Lagrangian symmetry F to the Hamiltonian symmetry G, the symmetry of the background fields, which removes the last node on the right. Projecting from Type II to Type I F-branes then distinguishes σ_2 (parity changes signs for all worldvolume coordinates except τ and σ_2), knocking off the next node on the right, which breaks the symmetry down to $G_{FI} = SO(D,D)$. Since the heterotic string contracts all but τ and σ_2 , it maintains that symmetry. But Type I strings contract σ_2 instead of σ_1 , killing the next node on the right, further breaking the symmetry to $G_I = GL(D)$. Note that these reductions in σ (d) drop nodes from the right, whereas reduction in X (D) drops nodes from the left.

The resulting string of cosets, from consecutively eliminating τ, σ_1, σ_2 from the symmetry, is

$$\frac{F}{L} \rightarrow \frac{G}{H} \rightarrow \frac{G_{FI}}{H_{FI}} = \frac{SO(D,D)}{SO(D-1,1)^2} \rightarrow \frac{G_I}{H_I} = \frac{GL(D)}{SO(D-1,1)} \quad (2.3)$$

(For each of these cosets A/B , B is a Wick rotation of the maximal compact subgroup of the split group A.)

It might seem strange that G_{FI}/H_{FI} is also G/H for Type II T-theory (both A and B), until one remembers that the complete bosonic sector of $N = 1$ stringy supergravity is the same as the NS-NS sector of $N = 2$, as easily seen from the heterotic construction: vector $\times$ vector, no symmetrization. (If $D \neq 3, 4, 6, 10$, there would also be scalars $\times$ vector, since supersymmetry requires vector multiplet $\times$ vector.) The coset G_I/H_I is then the coset of the NS-NS sector of Type I T-theory, namely just gravity + physical scalar (as the determinant of the metric).

In the string frame for T-theory, the “dilaton” is an unphysical, compensating scalar from ghost $\times$ ghost. In this reduction, applying parity to G/H restricts it to not just

$G_{\text{FI}}/H_{\text{FI}}$, but also the dilaton, while not eliminating spacetime coordinates. Then, applying sectioning reduces the spacetime coordinates, but for Type I simply breaks that coset into the metric (the last coset) + the 2-form (but doesn't break the coset for heterotic).

$$\frac{G}{H} \rightarrow GL(1) \frac{SO(D,D)}{SO(D-1,1)^2} = \text{dilaton} \oplus \frac{GL(D)}{SO(D-1,1)} \oplus \text{2-form} \quad (2.4)$$

Although the fields and number of coordinates are the same for Type I and heterotic T-theories, the symmetries are different because different coordinates have been eliminated in the sectioning.

Cosets of bosonic fields:

- (i) G/H : type II F-theory
- (ii) $SO(D,D)/SO(D-1,1)^2$: type I F-theory, heterotic supergravity, NS-NS Type II T-theory
- (iii) $GL(D)/SO(D-1,1)$: NS-NS Type I T-theory

For example:

$$D = 3: \quad \frac{SL(6)}{SO(3,3)} \rightarrow \frac{SL(5)}{SO(3,2)} \rightarrow \frac{SL(4)}{SO(2,2)} = \frac{SO(3,3)}{SO(2,1)^2} \rightarrow \frac{SL(3)}{SO(2,1)} \quad (2.5)$$

$$D = 4: \quad \frac{SO(6,6)}{SO(6,C)} \rightarrow \frac{SO(5,5)}{SO(5,C)} \rightarrow \frac{SO(4,4)}{SO(4,C)} = \frac{SO(4,4)}{SL(2,C)^2} = \frac{SO(4,4)}{SO(3,1)^2} \rightarrow \frac{SL(4)}{SO(3,1)} \quad (2.6)$$

Couplings can be studied through these reductions, as vacuum values of compactification scalars. Since the same coset fields appear as metrics for both the worldvolume and spacetime, these scalars measure distances in both.

3 M-theory

Although we will concentrate on T-theory because it incorporates the greater number of S-theories, we briefly divert to M-theory to see how the original Horava-Witten treatment fits in. M-theory is best described [16] by a worldvolume theory where X^M is a vector gauge field, so it carries the same index as σ_M . (We can also include a dual X , related by a duality constraint, useful for describing a background 6-form field.) Note that our arguments for M-theory and its relation to T-theory can be applied all the way to the critical dimension. (Because of the reliance on exceptional groups, detailed discussions for F-theory need to be specialized for each D , and in this paper we concentrate on $D = 3$ for simplicity.)

The bosonic fields are then the metric G_{MN} and 3-form A_{MNP} , where the indices are for $GL(D+1)$. (We ignore the 6-form, which can be relevant in higher D : see the counting arguments in appendix B.) Parity projection is easy to apply, as fixing σ_2 fixes the corresponding index on the fields. (Background fields don't carry the τ index.) Separating $M = (2, m)$ for the $GL(D)$ index m , we're left with fields with even numbers of $GL(D)$ indices:

$$G_{mn}, \quad G_{22} = \varphi, \quad A_{2mn} = B_{mn} \quad (3.1)$$

We can call this theory M'-theory. It has the same σ 's and X 's as M-theory, but parity projection has reduced the number of fields, and consequently reduced the coset space from $GL(D+1)/SO(D,1)$ to $GL(D)/SO(D-1,1)$. Thus, although still a $(D+1)$ -dimensional theory, it has the symmetries of D-dimensional $N = 1$ supergravity. This is analogous to the situation described by Horava and Witten, although our worldvolume is larger, and X -parity projection is a consequence of σ -parity projection.

However, according to our picture, M'-theory can also be obtained by first applying parity projection, and then $\mathcal{S}$ sectioning:

$$\begin{array}{ccc}
 & \text{FII} & \\
 \mathcal{P} \swarrow & & \searrow \mathcal{S} \\
 \text{FI} & & \text{M} \\
 \searrow & & \swarrow \\
 & \text{M}' &
 \end{array} \tag{3.2}$$

We next reduce to S-theory by applying $\mathcal{U}$ sectioning, the single constraint $\partial^M P_M = 0$. To obtain a D-dimensional theory, this requires

$$\partial^M P_M = 0 \Rightarrow \partial^M = \partial^2, \quad P_M = P_m \tag{3.3}$$

Thus the worldvolume is reduced to the worldsheet, while spacetime is reduced to D dimensions, while the above massless background fields are retained. According to our picture for parity preserving (τ, σ_2) while contracting all the rest, this is (the supergravity sector of) the heterotic string.

Other routes to heterotic S-theory will be described below. ($N = 1$ supersymmetric S-theories follow from different orderings of application of $\mathcal{P}, \mathcal{S}, \mathcal{U}$.)

4 T-theory

4.1 Type II

Before deriving Type I gravity (NS-NS sector) by $\mathcal{P}$ projection from Type II in T-theory, we first review Type II [8, 22, 23]. Working in the Hamiltonian formalism, we express the curved-space currents in terms of the flat-space currents as

$$\triangleright_A = E_A^M \mathring{\triangleright}_M, \quad \mathring{E}_A^M = \frac{1}{\sqrt{2}} \begin{pmatrix} \delta_a^m & -\eta_{am} \\ \delta_{\bar{a}}^m & \eta_{\bar{a}m} \end{pmatrix} \tag{4.1}$$

where the flat index $A = (a, \bar{a})$ is in a left/right basis, while the curved index $M = (m, {}^m)$ is in a co/contra-variant basis. ($\mathring{E}$ is the matrix that converts between the 2 bases. The flat currents can be taken as $\mathring{\triangleright}_M = (P_m, X'^m)$ in a basis where X is not doubled.) The $SO(D,D)$ metric in these 2 bases is

$$\eta_{AB} = \begin{pmatrix} \eta_{ab} & 0 \\ 0 & -\eta_{\bar{a}\bar{b}} \end{pmatrix}, \quad \eta_{MN} = \begin{pmatrix} 0 & \delta_m^n \\ \delta_n^m & 0 \end{pmatrix} \tag{4.2}$$

Since the concept of left and right will not survive in F-theory, we'll generally work instead in a co/contra-variant basis also for flat indices, $A = ({}_a, {}^a)$. Then the metric is the same for both flat and curved indices, and the flat vielbein is simply the identity, but the local $\text{SO}(D-1,1)^2$ transformations are more complicated (not block diagonal).

The orthogonality condition on E is

$$\eta^{AB} E_B{}^N \eta_{NM} = E_M{}^A \quad (4.3)$$

in terms of its inverse. In a gauge that breaks $\text{SO}(D-1,1)^2$ to the diagonal $\text{SO}(D-1,1)$, a convenient form for the solution to orthogonality for the section $\partial^m = 0$, where e and B have the usual gauge transformations, is

$$E_A{}^M = \hat{E}_A{}^N B_N{}^M = \begin{pmatrix} e_a{}^n & 0 \\ 0 & e_n{}^a \end{pmatrix} \begin{pmatrix} \delta_n^m & B_{nm} \\ 0 & \delta_m^n \end{pmatrix} = \begin{pmatrix} e_a{}^m & e_a{}^n B_{nm} \\ 0 & e_m{}^a \end{pmatrix} \quad (4.4)$$

or for $\partial_m = 0$ (with the T-dual gauge transformations),

$$E_A{}^M = \begin{pmatrix} e_a{}^n & 0 \\ 0 & e_n{}^a \end{pmatrix} \begin{pmatrix} \delta_n^m & 0 \\ B^{nm} & \delta_m^n \end{pmatrix} = \begin{pmatrix} e_a{}^m & 0 \\ -B^{mn} e_n{}^a & e_m{}^a \end{pmatrix} \quad (4.5)$$

To treat more general sections, or describe both at the same time, we'll sometimes write this as

$$E_A{}^M = \begin{pmatrix} e_a{}^n & 0 \\ 0 & e_n{}^a \end{pmatrix} \begin{pmatrix} \delta_n^m & B_{nm} \\ B^{nm} & \delta_m^n \end{pmatrix} = \begin{pmatrix} e_a{}^m & e_a{}^n B_{nm} \\ -B^{mn} e_n{}^a & e_m{}^a \end{pmatrix}, \quad B^{mn} B_{np} = 0 \quad (4.6)$$

(maintaining orthogonality).

In the absence of fermions, one can also work directly in terms of the metric

$$M^{MN} = \hat{\eta}^{AB} E_A{}^M E_B{}^N, \quad \hat{\eta}_{AB} = \begin{pmatrix} \eta_{ab} & 0 \\ 0 & \eta_{\bar{a}\bar{b}} \end{pmatrix} \quad (4.7)$$

This produces the usual

$$M^{MN} = \begin{pmatrix} G^{mn} & G^{mp} B_{pn} \\ -B_{mp} G^{pn} & G_{mn} - B_{mp} G^{pq} B_{qn} \end{pmatrix} \quad (4.8)$$

or (as follows from T-duality $\partial^m \leftrightarrow \partial_m$)

$$M^{MN} = \begin{pmatrix} G^{mn} - B^{mp} G_{pq} B^{qn} & -B^{mp} G_{pn} \\ G_{mp} B^{pn} & G_{mn} \end{pmatrix} \quad (4.9)$$

and in the general case

$$M^{MN} = \begin{pmatrix} G^{mn} - B^{mp} G_{pq} B^{qn} & G^{mp} B_{pn} - B^{mp} G_{pn} \\ G_{mp} B^{pn} - B_{mp} G^{pn} & G_{mn} - B_{mp} G^{pq} B_{qn} \end{pmatrix} \quad (4.10)$$

The gauge transformations are

$$\delta E_A{}^M = \lambda^N \partial_N E_A{}^M - E_A{}^N (\partial_N \lambda^M - \partial^M \lambda_N) \quad (4.11)$$

with indices raised and lowered with η . The non-transport term is an orthogonal transformation, and so preserves orthogonality.

We then identify the scalar density dilaton as transforming so $\int \Phi^2$ is invariant,

$$\delta \Phi = \lambda^M \partial_M \Phi + \frac{1}{2} \Phi \partial_M \lambda^M \quad (4.12)$$

After sectioning, the density Φ relates to a scalar φ as

$$\Phi^2 = \varphi^2 \sqrt{-G} \quad (4.13)$$

Besides the NS-NS sector, described by the coset $\text{SO}(D,D)/\text{SO}(D-1,1)^2$, and the ghost-ghost sector described by the dilaton, the bosons also include the R-R sector, described by an $\text{SO}(D,D)$ Weyl spinor [21, 24–29]. It has its own gauge transformation and field strength

$$\delta R = \gamma^M \partial_M \Lambda, \quad F = \gamma^M \partial_M R \quad (4.14)$$

where the $\text{SO}(D,D)$ γ -matrices satisfy

$$\{\gamma^M, \gamma^N\} = 2\eta^{MN} \quad (4.15)$$

(since $(\gamma^M \partial_M)^2 = \square = 0$ is the section constraint). Its coordinate transformation can then be written gauge covariantly (a simple way to derive the noncovariant form) as

$$\delta R = \frac{1}{2} (\gamma^M \lambda_M) F = (\delta_0 + \delta_1) R \quad (4.16)$$

where

$$\delta_0 R = \lambda^M \partial_M R + \frac{1}{4} \gamma^{[M} \gamma^{N]} (\partial_M \lambda_N) R + \frac{1}{2} (\partial_M \lambda^M) R, \quad \delta_1 R = -\frac{1}{2} \gamma^M \partial_M (\gamma^N \lambda_N R) \quad (4.17)$$

with a usual tensor density transformation δ_0 plus a gauge transformation piece δ_1 . Since the R-R fields are weight $\frac{1}{2}$, their (quadratic) action is already weight 1 without coupling to the dilaton Φ (also weight $\frac{1}{2}$).

This spinor gauge field resembles the *coordinates* X of 4D F-theory [10]. On sectioning, the resulting formalism for bispinors resembles that of the so-called “Dirac-Kähler” formalism [30–33].

4.2 3D

For the case $D = 3$, we can use “spinor notation”, and replace A (and M) of $\text{SO}(3,3)$ ($A = 1-6$) with $[AB]$ of $\text{SL}(4)$ ($A = 1-4$), to avoid introducing too many new kinds of indices. In this notation γ -matrices, the flat-space metric, and the Levi-Civita tensor of $\text{SO}(3,3)$ are all expressed in terms of Kronecker δ ’s and the ε -tensor of $\text{SL}(4)$:

$$\gamma_{CD}^{[AB]} = \delta_C^A \delta_D^B, \quad \gamma^{[AB]CD} = \varepsilon^{ABCD} \quad (4.18a)$$

$$\eta^{[AB][CD]} = \varepsilon^{ABCD}, \quad \varepsilon^{[AB][CD][EF][GH]} = \varepsilon^{A[C[E[G^H]F]D]B} \quad (4.18b)$$

Then for the coset of NS-NS fields it's useful to work with the 4-representation $\mathcal{E}$ of $SL(4)$ instead of the 6-representation E :

$$E_{AB}{}^{MN} = \mathcal{E}_{[A}{}^M \mathcal{E}_{B]}{}^N \quad (4.19)$$

especially for later embedding in the $SL(5)$ of F-theory.

The coordinate transformations are then

$$\delta \mathcal{E}_A{}^M = \frac{1}{2} \lambda^{NP} \partial_{NP} \mathcal{E}_A{}^M - \mathcal{E}_A{}^N \left(\partial_{NP} \lambda^{MP} - \frac{1}{4} \delta_N^M \partial_{PQ} \lambda^{PQ} \right) \quad (4.20a)$$

$$\delta \Phi = \frac{1}{2} \lambda^{MN} \partial_{MN} \Phi + \frac{1}{4} \Phi \partial_{MN} \lambda^{MN} \quad (4.20b)$$

$$\delta R_M = \frac{1}{2} \lambda^{NP} \partial_{NP} R_M + (\partial_{MP} \lambda^{NP}) R_N \quad (4.20c)$$

The non-transport term on $\mathcal{E}$ is an $SL(4)$ transformation. (Note that for R we have $\partial_{MP} \lambda^{NP} = (\partial_{MP} \lambda^{NP} - \frac{1}{4} \delta_M^N \partial_{PQ} \lambda^{PQ}) + \frac{1}{4} \delta_M^N \partial_{PQ} \lambda^{PQ}$.) We also have the R-R gauge transformations

$$\delta R_M = \partial_{MN} \Lambda^N \quad (4.21)$$

We can again derive the R coordinate transformations from the gauge covariant version,

$$\delta R_M = \frac{1}{2} \lambda^{NP} F_{MNP}, \quad F_{MNP} = \frac{1}{2} \partial_{[MN} R_{P]} \quad (4.22)$$

or similar expressions with $F^M = \frac{1}{6} \varepsilon^{MNPQ} F_{NPQ}$.

It's convenient to decompose indices as

$$V_{[AB]} = (V_{1a}, V_{[ab]}) \equiv (V_a, \varepsilon_{abc} V^c) \rightarrow (V_a, V^a) \quad (4.23)$$

and similarly for curved indices, where $A = (1, a)$, $a = (2, 3, 4)$. This gives the desired form of the $SO(3,3)$ metric

$$V^2 = \frac{1}{4} \varepsilon^{ABCD} V_{AB} V_{CD} \rightarrow 2V^a V_a \quad (4.24)$$

This allows us to straightforwardly identify components of $\mathcal{E}$ with those of E , and we find

$$\mathcal{E}_A{}^M = \begin{matrix} & 1 & m \\ a & \begin{pmatrix} e^{1/2} & e^{1/2} \frac{1}{2} \varepsilon^{mnp} B_{np} \\ -e^{-1/2} e_a{}^m \frac{1}{2} \varepsilon_{mnp} B^{np} & e^{-1/2} e_a{}^m \end{pmatrix} \end{matrix} \quad (4.25a)$$

$$= \begin{pmatrix} e^{1/2} & 0 \\ 0 & e^{-1/2} e_a{}^n \end{pmatrix} \begin{pmatrix} 1 & \frac{1}{2} \varepsilon^{mpq} B_{pq} \\ -\frac{1}{2} \varepsilon_{npq} B^{pq} & \delta_n^m \end{pmatrix} \quad (4.25b)$$

with a gauge condition again breaking $SO(2,1)^2$ to $SO(2,1)$.

Under this decomposition, the R-R gauge transformations branch into

$$\delta R_m = \partial_m \Lambda + \varepsilon_{mnp} \partial^n \Lambda^p, \quad \delta R = \partial_m \Lambda^m \quad (4.26)$$

Depending on how we section to S-theory (which X 's survive, x^m or x_m), this identifies these fields as either

$$\partial_m : \quad R_m = 1\text{-form}, \quad R = 3\text{-form (IIA)} \quad (4.27a)$$

$$\partial^m : \quad R_m = 2\text{-form}, \quad R = 0\text{-form (IIB)} \quad (4.27b)$$

This illustrates that both theories arise from different choices of section of the same theory (and the fact that T-duality relates them).

To avoid redefining the NS-NS fields (and switching up and down indices), one can also start with different chirality spinors, and section the same way:

$$R_M \quad \Rightarrow \quad R_m = 1\text{-form}, \quad R = 3\text{-form (IIA)} \quad (4.28a)$$

$$R^M \quad \Rightarrow \quad R^m = 2\text{-form}, \quad R = 0\text{-form (IIB)} \quad (4.28b)$$

(with corresponding changes in the gauge transformations from R_M to R^M). But F-theory doesn't prefer this.

A third alternative is to introduce both R_M and R^M , and relate them by duality. Since this is enforced via field equations, it requires introduction of the bosonic coset: in a flat background, the condition would break $\text{SO}(D,D)$ to $\text{SO}(D-1,1)^2$ (just as selfduality of vectors in 4D $N = 8$ supergravity requires the bosons to avoid breaking E_7 to $\text{SU}(8)$ and $\text{GL}(4)$ to $\text{SO}(3,1)$).

Although $\text{SO}(2,1)$ spinors aren't chiral, $\text{SO}(3,3)$ are, so in this way there can be 2 sections of Type II T-theory to S-theory.

4.3 Type I

We now impose also parity projection on both $G = \text{SO}(D,D)$ and $H = \text{SO}(D-1,1)_L \times \text{SO}(D-1,1)_R$, where the parity matrix $\mathcal{P}$ in both bases is

$$\mathcal{P}_A{}^B = \begin{pmatrix} 0 & \delta_a^{\bar{b}} \\ \delta_{\bar{a}}^b & 0 \end{pmatrix}, \quad \mathcal{P}_M{}^N = \begin{pmatrix} \delta_m^n & 0 \\ 0 & -\delta_n^m \end{pmatrix} \quad (4.29)$$

(clearly switching L and R, and changing $\sigma \rightarrow -\sigma$), and parity projection on E and an element of the H gauge group Λ is

$$\mathcal{P}_A{}^B E_B{}^N \mathcal{P}_N{}^M = E_A{}^M, \quad \mathcal{P}_A{}^C \Lambda_C{}^D \mathcal{P}_D{}^B = \Lambda_A{}^B \quad (4.30)$$

(It's actually easier to solve $\mathcal{P}$ projection before orthogonality.) The combination of these 2 conditions, besides identifying the L and R parts of H, also constrains

$$E_A{}^M = \frac{1}{\sqrt{2}} \begin{pmatrix} e_a{}^m & -e_{ma} \\ e_a{}^m & e_{ma} \end{pmatrix} \quad (4.31)$$

in the mixed basis or

$$E_A{}^M = \begin{pmatrix} e_a{}^m & 0 \\ 0 & e_m{}^a \end{pmatrix} \quad (4.32)$$

in the co/contra-variant basis for both. So the coset is reduced to $GL(D)/SO(D-1,1)$ of ordinary gravity.

Then

$$\eta M \eta = M^{-1}, \quad \mathcal{P} M \mathcal{P} = M \Rightarrow M^{MN} = \begin{pmatrix} G^{mn} & 0 \\ 0 & G_{mn} \end{pmatrix} \quad (4.33)$$

Parity projection on R in components (or the Ramond-Neveu-Schwarz formalism) is not so transparent: on field strengths, one simply antisymmetrizes on left and right spinor indices. (NS-NS fields are symmetrized in vector indices, R-R are antisymmetrized in spinor indices, according to graded symmetrization.) On the other hand the physical, transverse R-R degrees of freedom are given by antisymmetrizing spinor indices of the little group. However, since F-theory unifies NS-NS and R-R fields into a single gauge field, the procedure there is straightforward, as we'll see below, and the decomposition of F Type I to T Type I will illustrate the R-R projection. The reason for the difference is that branes, unlike the string worldsheet, can carry R-R charges, so background fields must include the gauge fields explicitly. (In [23] extra currents for R-R charges were introduced into T-theory. These currents are automatically incorporated in F-theory, as part of the same exceptional symmetry representation that includes the currents for the NS-NS “charges”: all bosonic currents are “unified”.)

Note that in $D = 3$, the R-R field strength in Type I is given by antisymmetrizing 2-valued spinor indices of $SO(2,1)$, and thus a scalar, which is the same as a 3-form, the field strength of a 2-form. On the other hand, the physical R-R degrees of freedom are given by antisymmetrizing 1-valued spinor indices of $SO(1)$, which gives nothing, because a 2-form has none.

5 3D F-theory

5.1 Type II

For F-theory we again look at the simple case of $D = 3$, which is actually simpler than the general T-theory case.

The solution to the generalized orthogonality condition [9]

$$E_{AB}{}^{MN} = \mathcal{E}_{[A}{}^M \mathcal{E}_{B]}{}^N \quad (5.1)$$

is simply the statement that the (inverse) worldvolume vielbein $\mathcal{E}$ is just a different $SL(5)/SO(3,2)$ representation of (the bosonic part of) the spacetime vielbein E ($\partial^M = \partial/\partial\sigma_M$):

$$\triangleright_{AB} = \frac{1}{2} E_{AB}{}^{MN} \triangleright_{MN}, \quad \mathcal{D}^A = \mathcal{E}_M{}^A \partial^M \quad (5.2)$$

The gauge transformation is

$$\delta \mathcal{E}_A{}^M = \frac{1}{2} \lambda^{NP} \partial_{NP} \mathcal{E}_A{}^M - \mathcal{E}_A{}^N \left(\partial_{NP} \lambda^{MP} - \frac{1}{5} \delta_N^M \partial_{PQ} \lambda^{PQ} \right) \quad (5.3)$$

The non-transport term is $SL(5)$.

The goal is to reduce Type II F-gravity to Type I and heterotic T-gravities. The procedure we described was:

- (1) First parity project Type II F-theory with respect to all σ but σ_2 to Type I F-theory.
- (2) ($\mathcal{U}$ -)Section to preserve σ_1 as the surviving σ for Type I T-theory and σ_2 for heterotic.

However, the simplest way to find these theories is to instead

- (a) First apply σ_2 -preserving sectioning to Type II F-gravity, producing the known Type II T-gravity (NS-NS and R-R sectors).
- (b) Then parity project by reflecting all but σ_2 labels for heterotic or σ_1 labels for Type I. (This is a reversal of the labeling of σ_1 and σ_2 for the Type I case.)

$$\begin{array}{ccc}
 & \text{FII} & \\
 \textcolor{blue}{\mathcal{P}} \swarrow & & \searrow \textcolor{blue}{\mathcal{U}} \\
 \text{FI} & & \text{TII} \\
 \searrow & & \swarrow \\
 & \text{TI, TH} &
 \end{array} \tag{5.4}$$

In this subsection, we apply the first step, finding Type II T-gravity. For $D = 3$, this means separating the indices as $A \rightarrow (2, a)$ ($a = 1, 3, 4, 5$), thus separating $\mathcal{E}$ into NS-NS, R-R, and dilaton fields, as the $\text{SL}(4)/\text{SO}(2,2)$ fields described above.

We then drop all $X^{[MN]}$'s with M or $N = 2$. Looking at the F-gravity coordinate transformation, for the moment we can ignore the transport term $\lambda \cdot \partial$ and density term $\partial \cdot \lambda$. We then see that dropping ∂_{2m} implies that fixing the Lorentz gauge $\mathcal{E}_2^m = 0$ is preserved by the coordinate transformation, while breaking $\text{SO}(3,2) \rightarrow \text{SO}(2,2)$. We next identify the dilaton Φ as the only scalar (density, with the appropriate weight), $\mathcal{E}_a^m \in \text{SL}(4)/\text{SO}(2,2)$ in the defining representation, and the R-R field R_m as a spinor, all transforming as expected for Type II T-gravity: as above,

$$\delta \mathcal{E}_a^m = \frac{1}{2} \lambda^{np} \partial_{np} \mathcal{E}_a^m - \mathcal{E}_a^n \left(\partial_{np} \lambda^{mp} - \frac{1}{4} \delta_n^m \partial_{pq} \lambda^{pq} \right) \tag{5.5a}$$

$$\delta \Phi = \frac{1}{2} \lambda^{mn} \partial_{mn} \Phi + \frac{1}{4} \Phi \partial_{mn} \lambda^{mn} \tag{5.5b}$$

$$\delta R_m = \frac{1}{2} \lambda^{np} \partial_{np} R_m + (\partial_{mp} \lambda^{np}) R_n + \partial_{mn} \lambda^{n2} \tag{5.5c}$$

The result is

$$\mathcal{E}_A^M = \begin{array}{cc} & \begin{array}{cc} 2 & m \end{array} \\ \begin{array}{c} a \end{array} & \begin{pmatrix} \Phi^{4/5} & 0 \\ \Phi^{-1/5} \mathcal{E}_a^m R_m & \Phi^{-1/5} \mathcal{E}_a^m \end{pmatrix} \end{array} = \begin{pmatrix} \Phi^{4/5} & 0 \\ 0 & \Phi^{-1/5} \mathcal{E}_a^n \end{pmatrix} \begin{pmatrix} 1 & 0 \\ R_n & \delta_n^m \end{pmatrix} \tag{5.6}$$

Then

$$E_{AB}^{MN} = \mathcal{E}_{[A}^M \mathcal{E}_{B]}^N = \begin{array}{cc} & \begin{array}{cc} 2m & mn \end{array} \\ \begin{array}{c} ab \end{array} & \begin{pmatrix} \Phi^{3/5} \mathcal{E}_a^m & 0 \\ \Phi^{-2/5} E_{ab}^{mn} R_n & \Phi^{-2/5} E_{ab}^{mn} \end{pmatrix} \end{array} \tag{5.7}$$

where $E_{ab}^{mn} = \mathcal{E}_{[a}^m \mathcal{E}_{b]}^n \in \text{SO}(3,3)/\text{SO}(2,1)^2$ in its defining representation. $\mathcal{E}$ is more convenient for algebra, but E relates more to general D . In particular, a similar decomposition was found in [23], where currents coupling to R-R charges were added to 10D T-theory.

This E_{ab}^{mn} appears as the same type of coset in Type I F-theory, but X and σ are bigger.

5.2 Type I and heterotic

To be slightly more general (for later applications), we examine the coset $GL(N)/SO(N)$ (without regard to signature), and consider a parity transformation that changes an arbitrary number of signs n ,

$$\mathcal{P} = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \quad (5.8)$$

Then

$$\mathcal{P} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathcal{P} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow b = c = 0 \quad (5.9)$$

simply forces the matrix to be block diagonal. (This is also clear from the index structure.) Thus

$$\frac{GL(N)}{SO(N)} \rightarrow \frac{GL(N-n)GL(n)}{SO(N-n)SO(n)} \quad (5.10)$$

In the present case, this means

$$\frac{SL(5)}{SO(3,2)} \rightarrow \frac{SL(4)}{SO(2,2)} GL(1) \quad (5.11)$$

Applying this restriction to the above Type II expressions,

$$\mathcal{E}_A^M = \begin{matrix} & 2 & m \\ a & \begin{pmatrix} \Phi^{4/5} & 0 \\ 0 & \Phi^{-1/5} \mathcal{E}_a^m \end{pmatrix} \end{matrix} \quad (5.12)$$

and

$$E_{AB}^{MN} = \mathcal{E}_{[A}^M \mathcal{E}_{B]}^N = \begin{matrix} & 2m & mn \\ ab & \begin{pmatrix} \Phi^{3/5} \mathcal{E}_a^m & 0 \\ 0 & \Phi^{-2/5} E_{ab}^{mn} \end{pmatrix} \end{matrix} \quad (5.13)$$

Since the heterotic coset is the same as Type II T-theory (NS-NS sector), identifying σ_2 as the surviving σ produces a T-theory action identical to that for the NS-NS sector of Type II [8], as expected, since NS-NS is the entire bosonic sector of heterotic string supergravity (neglecting the Yang-Mills supermultiplets).

To compare Type I T-theory to heterotic, we perform the same decomposition as for Type II T-theory, distinguishing the “1” label from the remaining. Combining the above results for F II $\rightarrow$ T II and T II $\rightarrow$ ordinary fields, (now $m = 3, 4, 5$)

$$\mathcal{E}_A^M = \begin{matrix} & 2 & 1 & m \\ & \Phi & 0 & 0 \\ 1 & \begin{pmatrix} e^{1/2}(R_1 + \frac{1}{2}\varepsilon^{mnp}B_{mn}R_p) & e^{1/2} & e^{1/2}\frac{1}{2}\varepsilon^{mnp}B_{np} \\ e^{-1/2}e_a^m(R_m - \frac{1}{2}\varepsilon_{mnp}B^{np}R_1) & -e^{-1/2}e_a^m\frac{1}{2}\varepsilon_{mnp}B^{np} & e^{-1/2}e_a^m \end{pmatrix} \end{matrix} \Phi^{-1/5} \quad (5.14)$$

with (R_1, R_m) as either odd (1 and 3) forms for IIA ($B^{mn} = 0$) or even (0 and 2) for IIB ($B_{mn} = 0$) upon reduction to S-theory:

$$R_1 + \frac{1}{2}\varepsilon^{mnp}B_{mn}R_p = \begin{cases} \varepsilon^{mnp}\left(\frac{1}{6}C_{mnp} + \frac{1}{2}B_{mn}A_p\right) & \text{(IIA)} \\ \tilde{\varphi} & \text{(IIB)} \end{cases} \quad (5.15a)$$

$$R_m - \frac{1}{2}\varepsilon_{mnp}B^{np}R_1 = \begin{cases} A_m & \text{(IIA)} \\ \frac{1}{2}\varepsilon_{mnp}(\tilde{B}^{np} - B^{np}\tilde{\varphi}) & \text{(IIB)} \end{cases} \quad (5.15b)$$

We then drop all components of $\mathcal{E}_A^M$ with an odd number of A and $M = 2$ for heterotic or 1 for Type I. Thus both $N = 1$ theories have e_a^m and Φ , but heterotic has B_{mn} while Type I has $\tilde{B}^{mn}$. As we already saw in Type II T-theory, coordinate transformations required that B (NS-NS) couples to the dilaton Φ , while $\tilde{B}$ (R-R) doesn't. (But the dilaton doesn't appear as an independent field in F-theory or M-theory, as is the case in 11D supergravity.)

Parity projection is with respect to both indices and coordinates: this picks out IIB rather than IIA, which we have distinguished in the 3D case with respect to coordinate dependence.

The fact that Type II supergravity can be projected in 2 different ways to get either Type I or heterotic supergravity is not new. What is new is that Type II F-theory can be projected to get Type I F-theory, which has 2 different sections that describe either Type I or heterotic string theory.

5.3 Stringy supergravities

Besides being simpler than their 10D analogs, in 3D (1) $N = 2$ is nonmaximal, so the supermultiplet is reducible, and (2) auxiliary fields and superspace formulations are known.

The Type IIA theory is known from M-theory (4D $N=1$ supergravity with a 3-form gauge auxiliary) and T-theory [34], F-theory [35], and our rederivation above: $N = 2$ supergravity coupled to a vector-tensor multiplet. The dimension-0 field content is the graviton and differential forms,

$$e_a^m, \quad \varphi, \quad A_m, \quad B_{mn}, \quad C_{mnp} \quad (5.16)$$

(The fermions are the usual gravitini and spin- $\frac{1}{2}$. The dimension-1 non-gauge auxiliary fields are a vector and scalars.) This separates into fields for pure supergravity

$$e_a^m, \quad C_{mnp} \quad (5.17)$$

and the matter multiplet

$$\varphi, \quad A_m, \quad B_{mn} \quad (5.18)$$

The reduction of $N = 2$ supergravity to $N = 1$ is unique (drop C), but the vector-tensor multiplet breaks into an $N = 1$ vector multiplet

$$A_m \quad (5.19)$$

and an $N = 1$ tensor multiplet

$$\varphi, \quad B_{mn} \quad (5.20)$$

By the usual construction of heterotic supergravity as vector $\times$ vector multiplet without symmetrization, its content is

$$e_a{}^m, \quad \varphi, \quad B_{mn} \quad (5.21)$$

using the tensor multiplet.

On the other hand (chirality) Type IIB, as described above, uses for the $N = 2$ matter a variant of the scalar multiplet: the total dimension-0 field content

$$e_a{}^m, \quad \varphi, \quad \tilde{\varphi}, \quad B_{mn}, \quad \tilde{B}_{mn} \quad (5.22)$$

divides up into that for $N = 1$ supergravity

$$e_a{}^m \quad (5.23)$$

and for matter

$$\varphi, \quad \tilde{\varphi}, \quad B_{mn}, \quad \tilde{B}_{mn} \quad (5.24)$$

That latter branches into 2 $N = 1$ tensor multiplets


$$\varphi, \quad B_{mn}; \tilde{\varphi}, \quad \tilde{B}_{mn} \quad (5.25)$$

As we saw above, Type I then chooses 1 of these multiplets, using the same scalar (dilaton) as heterotic, but a different 2-form.

In summary, we have from Type II theories to Type I theories in the $D = 3$ case:

	st	wv	Type II coset+	Type II fields	Type I $\mathcal{P}(\text{arity})$	Type I coset+	Type I fields
F	X^{MN}	σ_M, τ		$\mathcal{E}_A{}^M$	$\sigma_2 \rightarrow \sigma_2$		$\mathcal{E}_a{}^m, \Phi$
	$M = 1-5$	$M = 1-5$	$\frac{\text{SL}(5)}{\text{SO}(3,2)}$	$A, M = 1-5$	$\sigma_m \rightarrow -\sigma_m$	$\frac{\text{SL}(4)}{\text{SO}(2,2)}$	$a, m = 1, 3, 4, 5$
	10	$5 + 1$		14	$m = 1, 3, 4, 5$	+1	$9 + 1$
M	X^M	σ_M, τ		$e_A{}^M, A_{MNP}$	$\sigma_2 \rightarrow \sigma_2$		$e_a{}^m, \varphi, B_{mn}$
	$M = 1-4$	$M = 1-4$	$\frac{\text{GL}(4)}{\text{SO}(3,1)}$	$A, M = 1-4$	$\sigma_m \rightarrow -\sigma_m$	$\frac{\text{GL}(3)}{\text{SO}(2,1)}$	$a, m = 1, 3, 4$
	4	$4 + 1$	+4	$10 + 4$	$m = 1, 3, 4$	+1 + 3	$6 + 1 + 3$
T	X^{MN}	σ, τ		$E_{AB}{}^{MN}, \Phi, R_M$			$e_a{}^m, \varphi, R_m$
	$M = 1-4$		$\frac{O(3,3)}{\text{SO}(2,1)^2}$	$A, M = 1-4$	$\sigma \rightarrow -\sigma$	$\frac{\text{GL}(3)}{\text{SO}(2,1)}$	$a, m = 1, 2, 3$
	6	$1 + 1$	+1 + 4	$9 + 1 + 4$		+1 + 3	$6 + 1 + 3$

All the different theories can now be represented as



For clarity, we haven't split some theories, and therefore didn't indicate that (1) $M \rightarrow \text{SIIA}$ only, (2) $M' \rightarrow \text{SH}$ only, and (3) only $\text{SIIB} \rightarrow \text{SI/SH}$. We can also just list the theories by which combinations of $\mathcal{P}, \mathcal{S}, \mathcal{U}$ have been applied:

$$\begin{array}{ll}
 \text{---} & \text{FII} \\
 \mathcal{P} & \text{FI} \\
 \mathcal{S} & \text{M} \\
 \mathcal{U} & \text{TII} \\
 \mathcal{P}, \mathcal{S} & \text{M}' \\
 \mathcal{P}, \mathcal{U} & \text{TI/TH} \\
 \mathcal{S}, \mathcal{U} & \text{SII(A/B)} \\
 \mathcal{P}, \mathcal{S}, \mathcal{U} & \text{SI/SH}
 \end{array} \tag{5.27}$$

6 Conclusions

There are a number of topics for further study:

- (1) We have assumed a restricted topology. For σ_1 and σ_2 , we considered only cylinders (open branes) and donuts (closed branes), which are the only orientable surfaces with vanishing Euler number. This implies some kind of constraint on curvature. We also assumed tori with respect to the remaining worldvolume dimensions, but did not make an explicit application.
- (2) There are some results that will probably require quantization and consideration of anomalies: e.g., opposite chirality of Θ and internal degrees of freedom, and choice of Yang-Mills symmetry groups. (Previously we have quantized only indirectly by first reducing to S-theory.)
- (3) In particular, what precludes the existence of $E_8 \times E_8$ Type I superstrings, since the corresponding open F-brane is apparently the origin of the corresponding heterotic superstring?
- (4) Alternatively, might the Yang-Mills symmetry groups follow as solutions of the quantum corrected background field equations for the metric of the 496 scalars? Quantum F-theory isn't yet understood (except on sectioning to S-theory), but maybe some semiclassical arguments would suffice. Alternatively, the usual determination by anomalies would require only the massless sector of F-theory.
- (5) What are symmetry relations or dualities between Type I and Type II branes? Can symmetry breaking be attributed to super-Higgs? What's the relation to the boundary currents?
- (6) What are symmetry relations or dualities between open and closed Type I branes? In particular, how can supergravity come from not only closed branes (Type I superstrings), but also open (heterotic superstrings)?

In future papers we plan to address these questions, as well as give details on

- (a) the worldvolume formulation,
- (b) the F-gravity action,
- (c) the $D = 4$ case, and
- (d) seeing if the reduction of the symmetry groups from exceptional groups to $SO(D,D)$ allows a simple treatment of the general case.


Acknowledgments

We thank Dharmesh Jain for help with the bibliography, and his style type `hephys`. WS is supported by NSF grant PHY-1915093.

A Cosets

The cosets F/L for the Lagrangians and G/H for the Hamiltonians are [14]:

	D	d	F	G	L	H	
	0	2	$GL(2)$	$GL(1)$	$GL(1, \mathbf{C})$	I	
	1	3	$GL(3)$	$GL(2)$	$GL(2)$	$SO(1,1)$	
	2	4	$SL(4)SL(2)$	$SL(3)SL(2)$	$GL(2)^2$	$GL(2)$	
	3	6	$SL(6)$	$SL(5)$	$GL(4)$	$Sp(4)$	(A.1)
	4	12	$SO(6,6)$	$SO(5,5)$	$GL(4, \mathbf{C})$	$Sp(4, \mathbf{C})$	
	5	56	$E_{7(7)}$	$E_{6(6)}$	$U^*(8)$	$USp(4,4)$	
	6	?	?	$E_{7(7)}$	$U^*(8)^2$	$SU^*(8)$	
	7	?	?	$E_{8(8)}$	$U^*(16)$	$SO^*(16)$	



(Reduction from Lagrangian to Hamiltonian is indicated by τ , spontaneous symmetry breaking by the vacuum $|0\rangle$.) The question marks are for infinite-dimensional extended Lie algebras, as indicated by the Dynkin diagram in the main text.

B $F \rightarrow T$ and M

We present here a counting comparison of the fields in Type II F-theory, from G/H , and T-theory, from $SO(D,D)/SO(D-1,1)^2 = D^2$ (NS-NS) + $SO(D,D)$ Weyl spinor = 2^{D-1} (R-R). The discrepancy is attributed to the dilaton (compensating) multiplet, which comes from the ghost-ghost sector. We also include M-theory, as vielbein $GL(D+1)/SO(D,1) +$

3-form + 6-form in 1 extra dimension.

D	F	T	M
0	$1 - 0 = 1$	$0 + 0 = 0$	$1 + 0 = 1$
1	$4 - 1 = 3$	$1 + 1 = 2$	$3 + 0 = 3$
2	$11 - 4 = 7$	$4 + 2 = 6$	$6 + 1 = 7$
3	$24 - 10 = 14$	$9 + 4 = 13$	$10 + 4 = 14$
4	$45 - 20 = 25$	$16 + 8 = 24$	$15 + 10 = 25$
5	$78 - 36 = 42$	$25 + 16 = 41$	$21 + 20 + 1 = 42$
6	$133 - 63 = 70$	$36 + 32 = 68$	$28 + 35 + 7 = 70$
7	$248 - 120 = 128$	$49 + 64 = 113$	$36 + 56 + 28 = 120$

(B.1)

Note that the M-theory counting agrees with F except for $D = 7$ (lacking an $SO(7,1)$ vector), and T lacks only the dilaton for $D < 6$ ($D = 6$ needs an extra scalar, $D = 7$ an extra $SO(7,7)$ vector).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Coimbra, C. Strickland-Constable and D. Waldram, $E_{d(d)} \times \mathbb{R}^+$ generalised geometry, connections and M-theory, *JHEP* **02** (2014) 054 [[arXiv:1112.3989](https://arxiv.org/abs/1112.3989)] [[INSPIRE](#)].
- [2] D.S. Berman, M. Cederwall, A. Kleinschmidt and D.C. Thompson, The gauge structure of generalised diffeomorphisms, *JHEP* **01** (2013) 064 [[arXiv:1208.5884](https://arxiv.org/abs/1208.5884)] [[INSPIRE](#)].
- [3] O. Hohm and H. Samtleben, Exceptional form of $D = 11$ supergravity, *Phys. Rev. Lett.* **111** (2013) 231601 [[arXiv:1308.1673](https://arxiv.org/abs/1308.1673)] [[INSPIRE](#)].
- [4] O. Hohm and H. Samtleben, Exceptional field theory. I. $E_{6(6)}$ covariant form of M-theory and type IIB, *Phys. Rev. D* **89** (2014) 066016 [[arXiv:1312.0614](https://arxiv.org/abs/1312.0614)] [[INSPIRE](#)].
- [5] O. Hohm and H. Samtleben, Exceptional field theory. II. $E_{7(7)}$, *Phys. Rev. D* **89** (2014) 066017 [[arXiv:1312.4542](https://arxiv.org/abs/1312.4542)] [[INSPIRE](#)].
- [6] O. Hohm and H. Samtleben, Exceptional field theory. III. $E_{8(8)}$, *Phys. Rev. D* **90** (2014) 066002 [[arXiv:1406.3348](https://arxiv.org/abs/1406.3348)] [[INSPIRE](#)].
- [7] D.S. Berman and C.D.A. Blair, The geometry, branes and applications of exceptional field theory, *Int. J. Mod. Phys. A* **35** (2020) 2030014 [[arXiv:2006.09777](https://arxiv.org/abs/2006.09777)] [[INSPIRE](#)].
- [8] W. Siegel, Superspace duality in low-energy superstrings, *Phys. Rev. D* **48** (1993) 2826 [[hep-th/9305073](https://arxiv.org/abs/hep-th/9305073)] [[INSPIRE](#)].
- [9] W.D. Linch, III and W. Siegel, F-theory from fundamental five-branes, *JHEP* **02** (2021) 047 [[arXiv:1502.00510](https://arxiv.org/abs/1502.00510)] [[INSPIRE](#)].
- [10] W.D. Linch and W. Siegel, F-theory with worldvolume sectioning, *JHEP* **04** (2021) 022 [[arXiv:1503.00940](https://arxiv.org/abs/1503.00940)] [[INSPIRE](#)].
- [11] W.D. Linch and W. Siegel, Critical super F-theories, [arXiv:1507.01669](https://arxiv.org/abs/1507.01669) [[INSPIRE](#)].
- [12] C.-Y. Ju and W. Siegel, Gauging unbroken symmetries in F-theory, *Phys. Rev. D* **94** (2016) 106004 [[arXiv:1607.03017](https://arxiv.org/abs/1607.03017)] [[INSPIRE](#)].

- [13] W.D. Linch and W. Siegel, *F-brane dynamics*, [arXiv:1610.01620](#) [INSPIRE].
- [14] W. Siegel and D. Wang, *Enlarged exceptional symmetries of first-quantized F-theory*, [arXiv:1806.02423](#) [INSPIRE].
- [15] W. Siegel and D. Wang, *F-theory superspace backgrounds*, [arXiv:1910.01710](#) [INSPIRE].
- [16] W. Siegel and D. Wang, *M-theory from F-theory*, [arXiv:2010.09564](#) [INSPIRE].
- [17] M. Hatsuda and W. Siegel, *Perturbative F-theory 10-brane and M-theory 5-brane*, *JHEP* **11** (2021) 201 [[arXiv:2107.10568](#)] [INSPIRE].
- [18] P. Hořava and E. Witten, *Heterotic and type-I string dynamics from eleven-dimensions*, *Nucl. Phys. B* **460** (1996) 506 [[hep-th/9510209](#)] [INSPIRE].
- [19] W. Siegel, *Hidden gravity in open string field theory*, *Phys. Rev. D* **49** (1994) 4144 [[hep-th/9312117](#)] [INSPIRE].
- [20] S.J. Gates, Jr. and W. Siegel, *Leftons, rightons, nonlinear σ models, and superstrings*, *Phys. Lett. B* **206** (1988) 631 [INSPIRE].
- [21] D.C. Thompson, *Duality invariance: from M-theory to double field theory*, *JHEP* **08** (2011) 125 [[arXiv:1106.4036](#)] [INSPIRE].
- [22] M. Hatsuda, K. Kamimura and W. Siegel, *Superspace with manifest T-duality from type-II superstring*, *JHEP* **06** (2014) 039 [[arXiv:1403.3887](#)] [INSPIRE].
- [23] M. Hatsuda, K. Kamimura and W. Siegel, *Ramond-Ramond gauge fields in superspace with manifest T-duality*, *JHEP* **02** (2015) 134 [[arXiv:1411.2206](#)] [INSPIRE].
- [24] D. Brace, B. Morariu and B. Zumino, *T duality and Ramond-Ramond backgrounds in the matrix model*, *Nucl. Phys. B* **549** (1999) 181 [[hep-th/9811213](#)] [INSPIRE].
- [25] M. Fukuma, T. Oota and H. Tanaka, *Comments on T dualities of Ramond-Ramond potentials on tori*, *Prog. Theor. Phys.* **103** (2000) 425 [[hep-th/9907132](#)] [INSPIRE].
- [26] S.F. Hassan, *SO(d, d) transformations of Ramond-Ramond fields and space-time spinors*, *Nucl. Phys. B* **583** (2000) 431 [[hep-th/9912236](#)] [INSPIRE].
- [27] A. Rocén and P. West, *E_{11} , generalised space-time and IIA string theory: the $R \otimes R$ sector*, in *Strings, gauge fields, and the geometry behind: the legacy of Maximilian Kreuzer*, A. Rebhan, L. Katzarkov, J. Knapp, R. Rashkov and E. Scheidegger eds., *World Scientific*, Singapore (2010), p. 403 [[arXiv:1012.2744](#)] [INSPIRE].
- [28] O. Hohm, S.K. Kwak and B. Zwiebach, *Unification of type II strings and T-duality*, *Phys. Rev. Lett.* **107** (2011) 171603 [[arXiv:1106.5452](#)] [INSPIRE].
- [29] O. Hohm, S.K. Kwak and B. Zwiebach, *Double field theory of type II strings*, *JHEP* **09** (2011) 013 [[arXiv:1107.0008](#)] [INSPIRE].
- [30] D. Iwanenko and L. Landau, *Zur Theorie des magnetischen Elektrons. I* (in German), *Z. Phys.* **48** (1928) 340.
- [31] C. Lanczos, *Die tensoranalytischen Beziehungen der Diracschen Gleichung* (in German), *Z. Phys.* **57** (1929) 447 [[physics/0508002](#)] [INSPIRE].
- [32] A.W. Conway, *Quaternion treatment of the relativistic wave equation*, *Proc. Roy. Soc. Lond. A* **162** (1937) 145.
- [33] E. Kähler, *Der innere Differentialkalkül* (in German), *Rend. Mat. Appl.* **21** (1962) 425.
- [34] M. Poláček and W. Siegel, *T-duality off shell in 3D type II superspace*, *JHEP* **06** (2014) 107 [[arXiv:1403.6904](#)] [INSPIRE].
- [35] W.D. Linch and W. Siegel, *F-theory superspace*, [arXiv:1501.02761](#) [INSPIRE].