

# Cyanobacterial blooms alter benthic community structure and parasite prevalence among invertebrates in Florida Bay, USA

Elizabeth Duermit-Moreau<sup>1,\*</sup>, Jamie Bojko<sup>2,3,4</sup>, Donald C. Behringer<sup>1,2</sup>

<sup>1</sup>Fisheries and Aquatic Sciences, University of Florida, Gainesville, FL 32611, USA
 <sup>2</sup>Emerging Pathogens Institute, University of Florida, Gainesville, FL 32611, USA
 <sup>3</sup>School of Health and Life Sciences, Teesside University, Darlington DL1 1HG, UK
 <sup>4</sup>National Horizons Centre, Teesside University, Middlesbrough TS1 3BX, UK

ABSTRACT: Many marine habitats are at risk due to increasing frequency, intensity, and persistence of harmful algal blooms. Repeated cyanobacterial harmful algal blooms (cyanoHABs) in Florida Bay, USA, kill sponges, resulting in reduced filtration and loss of shelter for benthic species. The loss of these key ecosystem functions can impact disease dynamics if fewer pathogens are filtered from the water column (dilution), if shelter loss increases host density in remaining shelters and a directly transmitted disease is present (host regulation), or if shelter loss changes species distributions and foraging patterns (trophic exposure). We show persistent impacts to hard-bottom communities relative to non-impacted communities 2 yr after a significant cyanoHAB. We compared benthic structure, invertebrate epibenthic/infaunal community composition, and parasitism among macroinvertebrates, stone crab Menippe mercenaria, and Caribbean spiny lobster *Panulirus arqus*. On sites degraded by cyanoHABs, we found more, smaller sponges, indicating regrowth. Despite this evidence of recovery, epibenthic/infaunal invertebrate communities were distinct and more diverse on unimpacted sites. Additionally, there were fewer, smaller bivalves on impacted sites. The bivalve Tucetona pectinata, prey for stone crabs, was nearly absent on impacted sites, resulting in decreased prevalence of the apicomplexan gregarine Nematopsis sp., which is trophically transmitted from T. pectinata to M. mercenaria. Panulirus argus virus 1 also appears to be affected by cyanoHABs, as it was absent on impacted sites but present in 26.5% of spiny lobster on unimpacted sites. Impacts remain evident 2 yr after significant cyanoHABs, which does not bode well for these areas considering the frequent reoccurrence of blooms.

KEY WORDS: Habitat degradation  $\cdot$  Harmful algal bloom  $\cdot$  HAB  $\cdot$  Florida Bay  $\cdot$  Benthic invertebrate community

-Resale or republication not permitted without written consent of the publisher

# 1. INTRODUCTION

Habitat degradation due to anthropogenic and natural disturbance is a key issue affecting marine environments worldwide (McCauley et al. 2015) and undermining the ecosystem services they provide (Costanza et al. 1997, Smale et al. 2019). In areas with increased nutrient load (i.e. eutrophica-

tion) and certain climate change impacts, such as ocean acidification and warming waters, the frequency, intensity, and persistence of harmful algal blooms (HABs) is on the rise (Fu et al. 2012). These blooms, both toxic and nontoxic, can directly and indirectly harm wildlife (Fire & van Dolah 2012, Capper et al. 2013, Gravinese et al. 2018), humans (Carmichael 2001, Ibelings et al. 2014), and ecosys-

tems (Sunda et al. 2006, Paerl & Huisman 2008). Additionally, HABs that impact foundational species can lead to cascading ecosystem consequences (Butler et al. 1995).

There are also many ways that environmental change can interact with disease to affect host populations (Lafferty et al. 2004, Behringer & Duermit-Moreau 2021). Susceptibility to parasitism and disease can vary ontogenetically (Behringer 2012), by sex (Zuk & McKean 1996), habitat (Altizer et al. 2011), and with a myriad of other factors that are influenced by environmental change (Nunn et al. 2003, Luis et al. 2013). Specifically, environmental stressors that are linked to habitat degradation, such as increased water temperature (Karvonen et al. 2010), pollution (Khan 1990), malnutrition (Beck & Levander 2000), and eutrophication (Lafferty 1997), have also been linked to an increased susceptibility to parasites in a variety of habitats (Holmes 1996). Changing environmental conditions can also directly impact parasites and disease-causing agents, leading to changes in virulence (Braid et al. 2005), abundance/ activity (Groner et al. 2014), new emergence/outbreak (Travers et al. 2009), and range shifts (Burge et al. 2014). These interactive effects alter the hostpathogen-environment triad in ways that can result in net positive or net negative outcomes for any given host species.

Florida Bay, USA, is home to an expansive limestone hard-bottom marine habitat, covering approximately 30% of the shallow water environment (Bertelsen et al. 2009). These locations are designated as 'Essential Fish Habitat' for several commercially fished teleosts and invertebrates (NOAA 2021). This limestone hard-bottom habitat is dominated by sponges, which provide structural complexity to an otherwise low-relief habitat (Behringer & Butler 2006). Over 60 species of sponges found in this habitat are essential in driving benthic–pelagic coupling of nutrients by filter-feeding (Valentine & Butler 2019) and as shelter for numerous species (Westinga & Hoetjes 1981).

In addition to being important habitat features, sponges are also extremely efficient filter-feeders responsible for linking benthic macrofauna with nutrients in the water column by filtering microbes and assimilating dissolved organic carbon and particulate organic carbon (Reiswig 1971, Valentine & Butler 2019). They effectively remove pathogenic bacteria from the water column (Maldonado et al. 2010) and may gain significant nutrition from filtering viruses (Hadas et al. 2006). This filtration may significantly lower disease-causing agents in areas with dense sponge populations.

These ecosystem functions are severely impaired after periodic cyanobacterial blooms, which result in sponge die-offs (Butler et al. 1995, Peterson et al. 2006). The non-toxic but mucilaginous Synechococcus spp. cause these cyanobacteria harmful algal blooms (hereafter referred to as cyanoHABs), reduce light penetration, and appear to smother sponges (Puls 2015), resulting in extreme habitat degradation (Butler et al. 1995). Such die-offs have occurred periodically since 1990, with varying intensity (Cannizzaro et al. 2019). The most recent bloom resulting in documented habitat degradation occurred between 2016 and 2017 (Cannizzaro et al. 2019). A bloom in 2007 degraded sponge communities over 500 km<sup>2</sup>, resulting in the loss of over 90% of sponges at severely affected sites (M. Butler & D. Behringer unpubl. data). These die-offs reduce filtration capacity that can lead to further decreases in water quality and increases in turbidity (Peterson et al. 2006), resulting in a negative feedback loop that further degrades the environment and nearby seagrass beds (Glibert et al. 2009, Hall et al. 2016).

There are 3 main pathways in which degradation of the hard-bottom habitat in Florida Bay can directly and drastically increase exposure to harmful parasites. First, in the 'dilution' pathway (Keesing & Ostfeld 2021) the diminished filtration of bacteria and viruses by sponges can lead to greater proliferation of these harmful organisms (Hadas et al. 2006, Maldonado et al. 2010). Second, in the 'susceptible host regulation' pathway (Keesing et al. 2006), the loss of sponge shelters can artificially increase population density in remaining shelters, thereby increasing the potential for direct transmission of diseases (Hughes et al. 2002). Finally, in the 'trophic' pathway (Wood et al. 2010), prey availability may decrease with shelter loss (Butler et al. 2016), which could impact foraging habits (Sigler et al. 2009) or result in consumption of suboptimal prey (Amélineau et al. 2019) and thereby expose individuals to new parasites.

Hard-bottom degradation from cyanoHABs may also decrease exposure to parasites through these pathways. If degraded habitat is less attractive to hosts, decreasing host density, the susceptible host regulation pathway would lead to a decrease in directly transmitted parasites. The trophic pathway may lead to a decrease in trophically transmitted parasites that are present in healthy hard-bottom habitat if intermediate hosts are negatively impacted by shelter loss or other effects of cyanoHABs. In these scenarios, parasite prevalence may be an indication of ecosystem health (Hechinger & Lafferty 2005, Johnson et al. 2016). Restoration projects are increas-

Table 1. Known parasites in dominant decapod crustaceans in Florida Bay hard-bottom habitat, namely Florida stone crab
Menippe mercenaria and Caribbean spiny lobster Panulirus argus. Transmission and exposure pathways influence the
expected outcomes of degradation by cyanobacterial harmful algal blooms on parasite prevalence in these host species

Host	Parasite	Туре	Transmission	Impact	Exposure pathway	Expected outcome of degradation to parasites
Menippe mercenaria	Nematopsis sp.	Apicomplexan gregarine	Trophic	Unknown	Trophic	<ul> <li>(1) ↑ if bivalve density increases</li> <li>(2) ↓ if bivalve density decreases</li> </ul>
	Hemato- dinium sp.	Dinoflagellate	Environmental	Mortality	Dilution	(1) ↑ if sponges filter pathogen
Panulirus	PaV1	DNA virus	Direct	Mortality	Dilution	(1) ↑ if sponges filter pathogen
argus					Susceptible host regulation	<ul> <li>(1) ↑ if host density increases due to shelter limitation</li> <li>(2) ↓ if host density decreases due to habitat quality</li> </ul>
	herrnkindi host	Susceptible host regulation	<ul> <li>(1) ↑ if host density increases due to shelter limitation</li> <li>(2) ↓ if host density decreases due to habitat quality</li> </ul>			

ingly using prevalence of parasites with complex life cycles as a measure of success (Huspeni & Lafferty 2004, Moore et al. 2020).

These exposure pathways have the potential to impact a wide range of hard-bottom dwellers, including the Florida stone crab *Menippe mercenaria* and the Caribbean spiny lobster *Panulirus argus*. As 2 dominant benthic predators and scavengers that compete for sponge shelter (Behringer & Hart 2017), they could be heavily affected by habitat degradation and loss of sponge habitat. These decapods are known hosts of viral and protistan parasites with distinct transmission and exposure pathways that are expected to be impacted in different ways by habitat degradation (Table 1).

We took advantage of a natural experiment to examine how habitat degradation caused by cyanoHABs impacts benthic community structure in Florida Bay, and how these changes may affect parasite diversity and prevalence in Florida stone crabs and Caribbean spiny lobsters. We first established the differences in the benthic coverage and structure between sites impacted by the periodic cyanoHABs and unimpacted sites to determine whether there are habitat differences that would support changes in parasitism due to dilution and/or susceptible host regulation pathways. Next, we examined epibenthic and infaunal invertebrate communities on these sites, to see whether and how they were impacted by the blooms to test the trophic exposure pathway. Finally, we screened M. mercenaria and P. argus and several prey species for parasites to determine

whether exposure pathways and expected outcomes of cyanoHABs and degradation align. Additionally, we screened sympatric decapods of similar size in search of potential sinks and sources of parasites.

#### 2. MATERIALS AND METHODS

#### 2.1. Habitat

Five hard-bottom sites were surveyed in the Florida Keys (3 'unimpacted' and 2 'impacted') in July and September 2019 (Fig. 1). Sites were chosen based on known history of cyanoHABs and sponge mortality, as well as sufficiently high stone crab density to allow for collections. At each site, benthic coverage and habitat were characterized to quantify any differences between impacted and unimpacted habitats. Within each site, 4 non-overlapping  $2 \text{ m} \times 25 \text{ m}$  belt transects were used to record the number and size (diameter and height) of structures (sponges, solution holes [small pits in karst limestone], coral heads, and rocks). All structures larger than 10 cm × 5 cm were identified and counted along the belt transect, and the measurements from the first 10 on each transect were recorded. Benthic substrate coverage was assessed along the same transect using the point intercept method along each 25 m transect (Alvarez-Filip et al. 2011). The percent cover of benthic substrate (sponge, seagrass, coral, soft coral, red macroalgae, green macroalgae, brown macroalgae, or sand) was estimated every 25 cm along the transect for a total of 100 estimates per transect.




Fig. 1. Sampling sites (n = 5) in Florida Bay, USA: from south to north, the 3 unimpacted sites are Shands, Russel, and Craig Keys and the 2 impacted sites are Lignumvitae Key 1 (LV1) and Lignumvitae Key 2 (LV2). Hatching shows extent of the 2 most recent major cyanobacterial harmful algal blooms that occurred in 2013–2014 and 2016–2017. Inset map shows the state of Florida, with arrow indicating the location of Florida Bay

# 2.2. Benthic community

We then sampled each site for epibenthic and infaunal organisms, to assess differences in prey availability and overall invertebrate communities among the sites and between hard-bottom and seagrass within sites. Sampling was conducted using a Venturi suction sampler (Orth & van Montfrans 1987) with a 1 mm² mesh sieve. At each site, 0.25 m² plots (n = 5 within hard-bottom and 5 within seagrass) were sampled for 5 min or until all sediment had been removed, whichever came first. Samples were then frozen at  $-20^{\circ}$ C for later identification of all invertebrates to the lowest possible taxonomic level. We weighed the first 5 individuals of each species in a sample individually, and the remaining were counted and weighed collectively by species.

# 2.3. Parasite communities

At each site, we collected stone crabs (n = 20–25) and spiny lobsters (n = 15–20) for parasite profile analysis. Additionally, we collected 3 mollusks (n = 5 each of bittersweet comb *Tucetona pectinata*, American star snail *Lithopoma americanum*, and tulip snail *Fasciolaria tulipa*) and 2 sympatric crabs (n  $\le 10$  each of blotched swimming crab *Achelous spinimanus* and West Indian spider crab *Maguimithrax spinosissimus*) for parasitology. The mollusks repre-

sent potential intermediate hosts of trophically transmitted Nematopsis sp. and the sympatric crabs represent potential alternative hosts of *Hematodinium* sp. The mollusk species we collected were presumed to be prey items because the shells from at least 1 of each species were found in or around a stone crab den. Several live T. pectinata were collected directly from stone crab dens and crushed shells of this species were frequently observed around their dens. Crustaceans were collected by hand, as were mollusks, although some additional bivalves were collected ad hoc via suction sampling. Each animal was measured and sexed, and the shelter type and any conspecifics sharing shelter were noted for crustaceans.

All individuals were euthanized and necropsied within 24 h of capture. For molecular diagnostics, skeletal muscle, hepatopancreas, gill, and antennal gland were biopsied and fixed in 1 ml of 99% ethanol.

Skeletal muscle, hepatopancreas, gonad, gill, heart, midgut, antennal gland, and epithelial tissue were biopsied for histological processing and were submerged in Davidson's saltwater fixative (Hopwood 1969) for 24–48 h, then rinsed in tap water and transferred to 70% ethanol. Fixed samples were processed for hematoxylin and alcoholic eosin (H&E) staining using standard methods (Feldman & Wolfe 2014). Disease screening was conducted using an Olympus BH-2 light microscope. Screening was conducted by viewing one entire section of each organ at various magnifications (4–100×).

All M. mercenaria, A. spinimanus, and M. spinosissimus were additionally screened for Hematodinium sp. infection via conventional PCR (Bojko et al. 2018), and all P. argus were screened for Panulirus argus virus 1 (PaV1) and Ameson herrnkindi microsporidia infections using conventional PCR (Baker et al. 1994, Montgomery-Fullerton et al. 2007, Ovcharenko et al. 2010, Moss et al. 2012). DNA was extracted from gill, hemolymph, and muscle tissue for Hematodinium sp., PaV1, and A. herrnkindi, respectively, using Qiagen DNeasy blood and tissue extraction kits according to manufacturer's protocols. PCR reactions consisted of 1.25 U of Taq polymerase, 2.5 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP, 1 µmol of each primer, and 3 µl of DNA template (10–100 ng  $\mu$ l<sup>-1</sup>) in a 50  $\mu$ l reaction volume (Table A1 in the Appendix). Amplicons were visualized on a 2% agarose gel (120 V, 60 min) and appropriately sized bands were extracted from the

gel, purified, and sent for forward sequencing (Eurofins genomic sequencing services; www.eurofins.com). Sequence data were compared with other isolates in GenBank using NCBI BLASTn.

## 2.4. Statistical analyses

To assess any differences between shelter availability, which would affect the susceptible host regulation exposure pathway, at impacted and unimpacted sites number and size of (1) all shelters, (2) all sponge shelters, (3) loggerhead sponge shelters, and (4) solution holes encountered on belt transects were compared. Loggerhead sponges Spheciospongia vesparium were singled out because of their importance as habitat for spiny lobster and stone crab. We used Kruskal-Wallis rank sum *H*-tests (hereafter KW tests) with Bonferroni correction to account for data with non-homogeneous variances that did not meet the assumptions of parametric statistics. We approximated sponge biomass density to assess filtration and dilution capabilities by multiplying the average sponge area (diameter × height) per transect by percent of sponge coverage for that transect and compared impacted to unimpacted sites using a KW test. To assess habitat quality, which could also impact parasitology through the susceptible host regulation pathway, we used KW tests to compare the percent coverage of each bottom type from the transects (abiotic, macroalgae, corals, seagrass, and sponges) between impacted and unimpacted sites.

To characterize differences in invertebrate communities, average mass and abundance of invertebrates per sample and biomass and abundance of phyla (with Mollusca split into Bivalvia, Gastropoda, and Polyplacophora, and Arthropoda split into Decapoda and Isopoda) were compared between site types using KW tests, only including species found on both impacted and unimpacted sites to remove bias in species size. To determine if there were differences in epibenthic and infaunal invertebrate community composition, and thus prey availability, at impacted and unimpacted sites, a Bray-Curtis distance matrix (Bray & Curtis 1957) was calculated in the 'vegan' package in R (Oksanen et al. 2019) using the counts of species found in each sample. A principal coordinates analysis (PCoA) was used to visualize the similarity in overall species composition found among samples and permutational multivariate analysis of variance (PERMANOVA) with 1000 permutations to compare impacted and unimpacted sites. Biodiversity was compared between site types using KW tests

on the Shannon diversity index (Shannon 1948), species richness, and species evenness (Simpson 1949). To determine which benthic species are characteristic of impacted and unimpacted sites, respectively, an indicator species analysis from the 'labdsv' package (Dufrene & Legendre 1997, Roberts 2019) was applied to the species abundance by sample matrix. This analysis accounts for invertebrate site fidelity and relative abundance to identify species that preferentially occur at impacted and unimpacted sites, respectively.

To test for differences between the prevalence of parasites or diseases observed at impacted and unimpacted sites, chi-squared tests with Yates continuity correction were used in instances where expected values were <5. For significant chi-squared results with multiple comparisons (e.g. 5 sites), pairwise chi-squared tests with Bonferroni correction were used. All data were analyzed in RStudio (Version 1.0.153, R version 4.1.2).

## 3. RESULTS

#### 3.1. Habitat

Most shelters encountered were sponges (mean ±  $SD = 13.1 \pm 8.0$  per 25 m transect), although solution holes  $(0.69 \pm 0.8)$ , rocks  $(0.63 \pm 1.8)$ , and coral heads  $(0.5 \pm 0.8)$  were also found along transects. Unimpacted sites had significantly fewer overall shelters than impacted sites, and fewer sponge shelters in particular (Fig. 2a-c; shelters: H = 4.5, p = 0.03; sponges: H = 5.7, p = 0.02); however, there were no differences between the number of loggerhead sponge shelters (H = 1.6, p = 0.20) or solution holes (H = 0.02, p = 0.89). The mean size of all shelters, sponge shelters, and loggerhead sponge shelters at unimpacted sites was larger than at impacted sites (Fig. 2b-d; shelters: H = 50.5, p < 0.001; sponges: H =49.2, p < 0.001; loggerhead sponges: H = 49.2, p < 0.001). Sponge biomass density did not significantly differ between impacted (18.0  $\pm$  29.1 cm<sup>2</sup>) and unimpacted sites (11.0  $\pm$  8.2 cm<sup>2</sup>; H = 0.18, p = 0.67).

At unimpacted sites, benthic coverage was dominated by red macroalgae *Laurencia* spp. (35.7  $\pm$  33.9%), sand (34.3  $\pm$  16.9%), and various green macroalgae (e.g. *Halimeda* spp., *Udotea* spp., and *Penicillus* spp.; 23.2  $\pm$  16.6%). The impacted site benthos was dominated by sand (60.5  $\pm$  18.0%), turtle grass *Thalassia testudinum* (19.9  $\pm$  27.6%), and green macroalgae (15.2  $\pm$  16.7%). Other coverage encountered along transects included the sponges *Ircinia* 




Fig. 2. Number of (a) shelters, (b) sponge shelters, and (c) loggerhead sponge shelters per transect and the diameter in cm of (d) shelters, (e) sponges, and (f) loggerhead sponges at impacted and unimpacted sites surveyed. The horizontal lines indicate the means, the box dimensions include the 25<sup>th</sup> and 75<sup>th</sup> percentiles, and the whiskers extend to the minimum and maximum, with outliers shown as closed circles. Asterisks indicate shelters at a given site that are significantly greater in number or size based on Kruskal-Wallis H-tests

strobilina and *I. campana*, golfball coral *Favia fragum*, the sea fan *Gorgonia ventalina*, and the brown macroalgae *Sargassum* spp. Pairwise comparisons of benthic coverage revealed greater macroalgal coverage on unimpacted sites (H = 7.7, p = 0.005), greater abiotic coverage on impacted sites (H = 5.7, p = 0.017), and no differences in corals (H = 0.11, p = 0.74), seagrass (H = 0.44, p = 0.51), or sponge benthic coverage (H = 0.02, p = 0.88; Fig. 3).

# 3.2. Benthic community

We identified 1734 individual invertebrates from 102 taxa (Table 2) belonging to the phyla Mollusca, Arthropoda, Echinodermata, Annelida, and Sipunculida. Neither the number of individuals (H=0.22, p=0.64) nor total biomass per sample (H=0.24, p=0.63) differed between unimpacted ( $n=40.3\pm34.1$ ; biomass =  $7.03\pm6.40$  g) and impacted ( $n=34.2\pm27.2$ ; biomass =  $5.92\pm4.40$  g) sites. There were more sipunculids in samples from impacted ( $n=7.1\pm6.9$ ) vs. unimpacted sites ( $n=3.3\pm3.9$ ; H=5.56, df = 1, p=0.02). All other phyla and major taxonomic groups showed no differences between the number of individuals per sample at impacted vs. unimpacted

sites. When comparing just species found at both impacted and unimpacted sites, there were differences in the mass of the major taxonomic groups at each site type; individual bivalves ( $H=13.3,\ p<0.001$ ) were on average  $5.5\times$  heavier on unimpacted ( $0.93\pm1.27$  g) than impacted sites ( $0.17\pm0.17$  g), and chitons ( $H=4.5,\ p=0.03$ ) were on average  $2.5\times$  heavier on unimpacted ( $0.02\pm0.01$  g) than impacted ( $0.009\pm0.006$  g) sites. No other phyla or major taxonomic groups were significantly different in mass between impacted and unimpacted sites.

The centroids and dispersion of epibenthic and infaunal communities differed between impacted and unimpacted sites (PERMANOVA  $R^2$  = 0.078, p = 0.001) and between seagrass and hard-bottom (PERMANOVA  $R^2$  = 0.037, p = 0.03), but there was no interaction between site type and bottom type (PERMANOVA  $R^2$  = 0.018, p = 0.65; Fig. 4a). No differences in community structure metrics were identified at seagrass vs. hard-bottom habitats (diversity: H = 0.71, p = 0.40; richness: H = 0.23, p = 0.63; evenness: H = 3.0, p = 0.08); therefore, these bottom types were combined for the remaining analyses.

Communities at unimpacted sites had higher species diversity (H = 9.4, p = 0.002; Fig. 4b), richness (H = 4.3, p = 0.04; Fig. 4c), and evenness (H = 4.3)

5.0, p = 0.03; Fig. 4d). At unimpacted sites, *Tucetona pectinata* and Atlantic oyster drill *Urosalpinx cinerea* were identified as indicator species (Dufrene & Legendre 1997), whereas indicator groups

on impacted sites included sipunculids, flyspeck cerith *Cerithium muscarum*, stocky cerith *C. litteratum*, and Morton's egg cockle *Laevicardium mortoni*.




Fig. 3. Benthic coverage was assessed using the point-intercept methods on 25 m transects (n = 4) per site. (a) Percent cover by coverage type, colored by site type. Boxplot limits as in Fig. 2. Asterisks indicate significance based on Kruskal-Wallis *H*tests. (b) Total percent coverage by site type




Fig. 4. Epibenthic and infaunal invertebrate communities were assessed through suction sampling and species identification. Using (a) principal coordinates analysis (PCoA), which calculates the similarity between entire suction samples, we show that invertebrate communities differ between impacted and unimpacted hardbottom. Invertebrate communities on healthy hardbottom have higher (b) Shannon Diversity index, (c) species richness, and (d) species evenness. Boxplot limits as in Fig. 2.

Asterisks indicate significance based on Kruskal-Wallis H-tests

Table 2. Individual invertebrates identified from suction samples collected at impacted and unimpacted sites, listed to lowest taxon identified. Indicator values from indicator species analysis, calculated based on site fidelity and relative abundance, are listed. Significant indicator species are denoted with asterisks ( $^*p < 0.05$ ;  $^{**}p < 0.01$ ;  $^{***}p < 0.001$ )

Taxon	Impacted	Unimpacted	Taxon	Impacted	Unimpacte
Annelida			Echinodermata		
Amphinomidae	0.00	0.12	Amphiuridae	0.20	0.41
Diopatra cuprea	0.00	0.04	Holothuroidea	0.00	0.08
Nereididae	0.21	0.45	Ophiuridae	0.01	0.20
Polychaeta	0.11	0.11	Oreaster reticulatus	0.00	0.04
Polynoidae	0.00	0.04	Mollusca		
Sabellidae	0.04	0.09	Acanthochitona pygmaea	0.03	0.02
Syllidae	0.08	0.50	Acanthopleura pygmaea	0.00	0.02
Terebellidae	0.03	0.34*	Americardia media	0.03	0.13
Arthropoda			Anadara sp.	0.00	0.02
Achelous floridanus	0.00	0.12	Bivalvia	0.00	0.04
Achelous gibbesii	0.00	0.04	Bostrycapulus aculeatus	0.00	0.04
Achelous gibbesii Achelous spinicarpus	0.00	0.04	Bulla striata	0.07	0.12
	0.00	0.04	Cerithium eburneum	0.26	0.24
Achelous spinimanus Alpheus christofferseni	0.00	0.04	Cerithium litteratum	0.26	0.24
Alpheus floridanus	0.13	0.01	Cerithium nitteratum Cerithium muscarum	0.59*	0.00
1					
Alpheus packardii	0.26	0.22	Cerithium sp.	0.30*	0.00
Alpheus sp.	0.02	0.09	Cerodrillia thea	0.03	0.03
Alpheus verrilli	0.00	0.04	Chama congregata	0.00	0.04
Anomura	0.05	0.00	Chione cancellata	0.00	0.15
Anthuridae	0.01	0.09	Columbella mercatoria	0.00	0.12
Brachycarpus biunguiculatus	0.00	0.04	Conasprella jaspidea	0.15	0.06
Calappa flammea	0.05	0.00	Conus sp.	0.00	0.04
Callinectes sp.	0.05	0.00	Costoanachis avara	0.00	0.08
Caridea	0.06	0.02	Cylindrobulla beauii	0.00	0.08
Coryrhynchus sidneyi	0.00	0.04	Drilliidae	0.00	0.04
Cymothoidae	0.00	0.04	Eoacmaea pustula	0.00	0.08
Decapoda	0.05	0.00	Favartia cellulosa	0.02	0.05
Ebalia sp.	0.03	0.02	Fulvia laevigata	0.18	0.04
Epialtus elongatus	0.01	0.12	Gastropoda	0.00	0.08
Erichsonella floridana	0.03	0.02	Ischnochiton papillosus	0.00	0.08
Euryplax nitida	0.05	0.00	Laevicardium mortoni	0.53***	0.00
Grapsidae	0.00	0.04	Laevicardium pictum	0.05	0.00
Latreutes fuscorum	0.00	0.12	Limaria pellucida	0.00	0.08
<i>Leptochela</i> sp.	0.00	0.04	Lithopoma phoebium	0.00	0.15
Lysmata rathbunae	0.00	0.04	Lottia antillarum	0.00	0.04
Neogonodactylus bredini	0.00	0.04	Mitrella dichroa	0.00	0.04
Omalacantha bicornuta	0.01	0.20	Modulus modulus	0.16	0.21
Paguristes sp.	0.20	0.10	Natica tedbayeri	0.02	0.02
Pagurus sp.	0.04	0.07	Niveria quadripunctata	0.05	0.00
Palaemonidae	0.00	0.04	Olivella sp.	0.00	0.08
Panopeus herbstii	0.00	0.08	Olivella watermani	0.00	0.12
Paracerceis caudata	0.02	0.05	Parvilucina crenella	0.02	0.25
Penaeidae	0.03	0.03	Phrontis vibex	0.02	0.02
Penaeus aztecus	0.05	0.00	Polyplacophora	0.05	0.10
Penaeus setiferus	0.00	0.08	Prunum apicinum	0.05	0.13
<i>Periclemenes</i> sp.	0.08	0.05	Tegula fasciata	0.01	0.16
Pitho lherminieri	0.25	0.26	Tellina sp.	0.08	0.20
Portunus sayi	0.05	0.00	Tucetona pectinata	0.00	0.44***
Precessa sp.	0.00	0.04	Urosalpinx cinerea	0.02	0.50**
Rimapenaus constrictus	0.00	0.04	Vermicularia spirata	0.10	0.00
Stenopus sp.	0.00	0.08	Vokesimurex rubidus	0.00	0.04
Tyche emarginata	0.00	0.04	Sipunculida	0.67***	0.16
Xanthidae	0.17	0.16		5.07	5.10

# 3.3. Parasite communities


There was no difference in size between lobsters collected at unimpacted (mean  $\pm$  SD carapace length [CL] 47.9  $\pm$  14.7 mm; n = 49) versus at impacted sites (52.4  $\pm$  17.4 mm CL; n = 22; H = 1.1, p = 0.31). There was no evidence of *Ameson herrnkindi* infection via PCR. One lobster exhibited gross signs of PaV1 infection during necropsy, and 18.3% of *Panulirus argus* were positive for PaV1 via PCR. Significantly more lobsters from unimpacted sites (26.5%) were infected with PaV1 than those collected from impacted sites (0%;  $\chi^2$  = 5.5, df = 1, p = 0.02).


Menippe mercenaria (n = 65) measured 68.1  $\pm$  20.7 mm in carapace width (CW) at unimpacted sites and 69.0  $\pm$  13.2 mm CW (n = 44) at impacted sites, which did not differ significantly (H = 0.0002, p = 0.99). M. mercenaria were collected from solution holes at unimpacted (56.9%) and impacted (95.5%) sites; however, more crabs were collected from living shelters, such as sponges and coral heads, on unimpacted (26.2%) vs. impacted sites (4.5%). There was no evidence of Hematodinium sp. infection via histopathological screening, and all PCR assays were negative for Hematodinium sp.

Histopathological screening identified the trophont stage of a trophically transmitted apicomplexan gregarine, Nematopsis sp., in the gut of 20.2% of M. mercenaria (Fig. 5a). No significant difference in the percentage of infected crabs between unimpacted (23.1%) and impacted sites  $(15.9\%; \chi^2 = 0.45, df = 1,$ p = 0.50) was observed; however, there was a site effect ( $\chi^2 = 12.9$ , df = 4, p = 0.01), and 1 unimpacted site (Russel Key) had significantly lower prevalence (4.0%) than the other 2 unimpacted sites: Craig Key  $(31.6\%; \chi^2 = 4.2, df = 1, p = 0.05)$  and Shands Key  $(38.1\%; \chi^2 = 6.4, df = 1, p = 0.005)$ . When site was removed from the comparison, unimpacted sites (35.0%) had significantly higher prevalence of the gregarine infection relative to impacted sites (11.4%;  $\chi^2 = 5.4$ , df = 1, p = 0.02; Fig. 5c).

Putative prey species collected for parasite screening included Lithopoma americanum (n = 2), T. pectinata (n = 12), and Fasciolaria tulipa (n = 4). Through histopathological analysis, the oocyst stage of a Nematopsis sp., was observed in the gills of T. pectinata (Fig. 5b), as well as a trematode in the digestive gland. Prevalence of Nematopsis sp. was 91.7%; however, no T. pectinata were found and necropsied at impacted sites (Table 3).

Maguimithrax spinosissimus (n = 7) (CW =  $41.5 \pm 20.7$  mm) and Achelous spinimanus (n = 4) (CW =  $52.7 \pm 12.0$  mm) were also collected. Histopathologi-








Fig. 5. Nematopsis sp. gregarine apicomplexan (a) trophont stage infecting stone crab Menippe mercenaria hindgut tissue and (b) oocyte stage infecting Tucetona pectinata primary gill filament. (c) Prevalence of Nematopsis sp. infecting M. mercenaria on impacted and unimpacted sites.

Boxplot limits as in Fig. 2

cal analysis revealed a trematode encysted in the primary gill filament of 1 M. spinosissimus (Fig. 6a) and a systemic infection of Hematodinium sp. in the

Table 3. Percent prevalence of identified parasites in *Menippe mercenaria*, *Panulirus argus*, *Tucetona pectinata*, and *Maguimithrax spinosissimus*. For size measurements, carapace width is given for *M. mercenaria* and *M. spinosissimus*; carapace length is given for *P. argus*, and shell width is given for *T. pectinata*. Prevalence of *Hematodinium* sp., *Panulirus argus* virus 1 (PaV1), and *Ameson herrnkindi* detected via conventional PCR. Prevalence of *Nematopsis* sp. detected via histopathology. nd: data were not available; na: not applicable

Site type	Sex distribution (M:F)	n	Size (mm) (mean ± SD)	Hematodinium sp.	Nematopsis sp.	PaV1	Ameson herrnkindi
Menippe mercenaria							
Unimpacted	35:30	65	$68.1 \pm 20.7$	0	35.0 <sup>a</sup> , (23.1)	na	na
Impacted	18:26	44	$69.0 \pm 13.2$	0	11.4	na	na
Panulirus argus							
Unimpacted	23:26	49	$47.9 \pm 14.7$	na	na	26.5	0
Impacted	11:11	22	$52.4 \pm 17.4$	na	na	0	0
Tucetona pectinata							
Unimpacted	nd	12	$17.9 \pm 4.3$	na	91.7	na	na
Impacted	nd	0	na	na	0	na	na
Maguimithrax spinosi	ssimus						
Unimpacted	5:0	5	$41.2 \pm 25.3$	20	0	na	na
Impacted	2:0	2	$42.4 \pm 1.9$	0	0	na	na

<sup>&</sup>lt;sup>a</sup>Nematopsis sp. prevalence given without crabs collected at site Russel Key (prevalence with Russel Key in parentheses)


hepatopancreas, muscle, and gill of another M. spinosissimus (Fig. 6b). The Hematodinium internal transcribed spacer (ITS) sequence obtained via PCR from this M. spinosissimus in the Florida Keys (GenBank accession no. OM913495) showed 100% similarity to H. perezi isolated from Callinectes sapidus in Texas, USA (accession KX244634, e-value =  $1 \times 10^{-162}$ ) and environmental water samples in Maryland (accession KF727429, e-value =  $7 \times 10^{-162}$ ) (Li et al. 2010, Pagenkopp Lohan et al. 2012, 2013). This is the first report of H. perezi in M. spinosissimus.

# 4. DISCUSSION

In this study, we documented for the first time significantly different epibenthic and infaunal invertebrate communities at unimpacted hard-bottom sites and those impacted by repeated cyanoHABs in the Florida Keys. The difference is particularly notable in the relatively greater mass of bivalves at unimpacted sites compared to impacted sites, and the low abundance of the bivalve Tucetona pectinata at impacted sites. Furthermore, T. pectinata appears to be an intermediate host of the apicomplexan gregarine Nematopsis sp., which is more prevalent in the final host, Menippe mercenaria, at unimpacted sites vs. impacted sites, possibly because of the near absence of the intermediate host at impacted sites. This relationship supports the trophic exposure pathway and expected outcome of decreased bivalve density. We also document support for the susceptible host regulation exposure pathway of PaV1 via decreased prevalence on impacted sites. Observation of these disease- and community-based differences persisting 2 yr after a major cyanoHAB indicates that the longevity of the impacts these blooms have can be far-reaching and pervasive in multiple ecological systems. While sponge communities appear to be rebounding with a greater abundance of smaller sponges on sites affected by the blooms, the persistent reoccurrence of the blooms threatens the long-term resilience of sponge communities.

# 4.1. Ecological resilience of invertebrate communities in the Florida Keys

After an intense cyanoHAB in Florida Bay in 1991, Butler et al. (1995) found that more than 40% of loggerhead sponges and 80% of other sponge species had perished, whereas 95% remained healthy on the periphery of the bloom. Our impacted sites were located within the heavily affected area of subsequent blooms from 2013–2014 and 2016–2017 (Cannizzaro et al. 2019, K. Hubbard pers. comm.) These were sampled in 2019, allowing our study to take place just 2 yr after an extensive sponge mortality event. While a time-series or data from immediately before and after the bloom for comparison would be the best way to affirm that sponges are rebounding






Fig. 6. Parasites (arrows) of *Maguimithrax spinosissimus*: (a) trematode metacercaria encysted in primary gill filament and (b) *Hematodinium perezi* infecting hepatopancreas

between blooms, our data show that more numerous, but smaller, sponges are emerging at degraded sites and represent a good indication of resilience. The nearshore environment of the middle Florida Keys is similar among these sites in terms of depth, water temperature, and habitat (E. Duermit-Moreau unpubl. data). In taking advantage of a natural experiment, we have been able to gather a rich dataset in a natural environment; however, the lack of sampling prior to the blooms may limit our conclusions.

Further evidence of resilience to cyanoHABs is apparent in the overlapping portions of epibenthic and infaunal invertebrate communities at impacted and unimpacted hard-bottom sites. When sponge communities are restored, or environmental conditions allow their regrowth, associated fauna can also return (Butler et al. 2016); however, the overall epibenthic and infaunal communities were distinct and indicative of wider impacts of the cyanoHABs. In

particular, the differences in bivalve communities could indicate a direct impact of the blooms on bivalves. Toxin-producing algal blooms affect bivalve larval development (Rolton et al. 2014), clearance rates (Leverone et al. 2007), cellular immunity (Lassudrie et al. 2020), and mortality (Leverone et al. 2006, Griffith et al. 2019). Unlike those HABs, these cyanoHABs do not produce toxins, so we believe the key to this impact is that most bivalves, including T. pectinata, are filter feeders, as are sponges. Sponge filtration and water pumping is inhibited by the sticky mucilage produced by Synechococcus sp. during high bloom densities (Phlips et al. 1989, Puls 2015) and this mucilage can have a similar effect on bivalves, resulting in direct mortality or inhibiting growth and reproduction and should be examined further.

# 4.2. Invertebrate biomass and trophic relationships at healthy and degraded sites

Decreased mass of bivalves can have important implications for the health of an ecosystem. In some habitats, such as oyster reefs, they are the dominant filter feeders, capable of increasing water clarity (Beck et al. 2011). They constitute an important prey resource for many vertebrate and crustacean species. *M. mercenaria* are voracious consumers of bivalves (Rindone & Eggleston 2011, Duermit et al. 2015), and we observed extensive evidence of this at unimpacted sites. We specifically observed crushed shells in and around occupied dens and live individuals within dens at sites where this prey item was present.

This trophic link between M. mercenaria and T. pectinata is likely the transmission pathway for Nematopsis sp. (Apicomplexa) found here in the gut and gill of these species, respectively. N. ostrearum and N. prytherchi have been previously reported in M. mercenaria (Sprague 1949, Sprague & Orr 1955); however, no gregarines have been reported in T. pectinata. Most apicomplexan gregarines parasitize just a single invertebrate host and have an environmental (free-living or spore) stage, but those in the genus Nematopsis have complex lifecycles. Their oocytes can infect bivalves, which act as intermediate hosts, and their trophonts infect decapod crustaceans, the definitive hosts (Sprague & Orr 1955). Molecular identification of this parasite in the oocyte stage has not yet proved successful (Silva et al. 2019), limiting our identification technique. We used a dichotomous key (Clopton 2002) and prior description of the parasite in M. mercenaria (Sprague & Orr

1955) to identify the apicomplexan gregarine to genus level for each species in which it was present. There is little evidence in the literature, or in our histological examination, of any pathological effects of *Nematopsis* sp. for either bivalve or stone crab, and there is no indication of zoonosis, so we are not concerned about emergent diseases at this time.

# 4.3. Parasite community structure across healthy and degraded sites

With the evidence of the trophic exposure pathway and suppressed bivalve density at impacted sites, there is potential for the gregarine to be used as an indicator of ecosystem health to further understand stone crab foraging. Stone crabs sampled from impacted habitats have altered stable isotope signatures, indicating a change in the base of the food web and a broadening of their diet, with some individuals feeding at lower trophic levels (D. Pharo & D. Behringer unpubl.). The decrease in size and abundance of bivalves, a preferred prey item, at impacted sites may be the cause of these trophic changes, requiring stone crabs to feed on less preferred prey (such as annelids and small gastropods). Due to the high prevalence of gregarines in T. pectinata (91.7%), gregarines in the gut of stone crab could be an indication of crabs that are feeding on this species of bivalve and are therefore foraging in healthier hardbottom habitat.

Spatial variation in parasite pressure can be due to the environmental requirements of the parasite (Owens 1983), but for parasites with complex life cycles, spatial variation can depend on the presence of one or more hosts (Jokela & Lively 1995). This pattern has led to increased work in the field of environmental parasitology (Lafferty 1997, Gagne et al. 2022), which argues that thorough knowledge of parasites within food webs can be used as indicators for anthropogenic impacts (Gilbert & Avenant-Oldewage 2021, Pravdová et al. 2021), habitat degradation (Sitko & Heneberg 2020), and the success of habitat restoration or conservation efforts (Moore et al. 2020, Braicovich et al. 2021). Moore et al. (2020) found that parasite abundance and richness mirrored that of free-living species following oyster reef restoration and that both measurements resembled those collected for healthy reefs after 16 mo.

A second association noted between parasitism and habitat in our study included PaV1, a member of the *Mininucleoviridae* (Subramaniam et al. 2020), which was absent in lobsters collected from impacted

sites, relative to a 26.5% prevalence of the virus in lobsters from healthy sites. PaV1 can cause nearly 100% mortality in juveniles over 2-8 wk of infection (Shields & Behringer 2004). In the interim, it is directly transmitted between conspecifics, which are gregarious, sharing shelters in rocks, coral heads, and large sponges (Behringer et al. 2018). A loss of large sponges at impacted sites, resulting in shelter limitation, has been shown to regulate susceptible hosts via increased density of lobsters in the few remaining shelters and thus possibly greater infection transmission (Butler et al. 2015). Despite this increase in density, the prevalence of PaV1 does not appear to increase at impacted sites because healthy lobsters are able to detect and avoid diseased conspecifics (Behringer et al. 2006, Butler et al. 2015). We found fewer lobsters overall at impacted sites; however, they were present at similar densities within dens to those found on unimpacted sites. This observation differs from previous studies that found lower densities of lobsters at degraded sites (Butler et al. 2015).

Contrary to previous studies, we found a relatively high prevalence of PaV1-infected lobsters at unimpacted sites. PaV1 can vary in space and time (Davies et al. 2020) so this snapshot of disease prevalence could simply indicate an irregular fluctuation. The prevalence of PaV1 at unimpacted sites could also be indicative of susceptible host regulation via a more attractive settlement habitat for postlarval and early benthic juvenile lobsters, the latter of which are most susceptible to PaV1 (Moss et al. 2012). Unimpacted sites in our study were dominated by macroalgae, with the red macroalgae Laurencia spp. being particularly common, whereas impacted sites had less macroalgal cover and more barren sand and rock. The presence of Laurencia is an important settlement cue for competent postlarvae, so this could be attracting more potentially infected and susceptible individuals to unimpacted sites.

In contrast to the trophic and susceptible host regulation exposure pathways, we did not find support for the dilution pathway in this system. We saw similar sponge biomass density among the sites, suggesting that filtration of pathogenic bacteria and viruses may be similar, thus not diluted on unimpacted sites. Another way that environmental change is predicted to affect host–pathogen dynamics is by altering host susceptibility (Harvell et al. 2002, 2004). For example, warming water temperatures have been shown to increase prevalence and severity of epizootic shell disease in American lobster *Homarus americanus* (Tlusty et al. 2007, Groner et al. 2018). While we did

not explore this prediction in the present study, it could play a role in this system.

We also screened for *Hematodinium* sp. in *M. mer*cenaria, Maquimithrax spinosissimus, and Achelous spinimanus, which is the cause of various decapod diseases and a significant mortality driver for many crustacean species globally (Stentiford & Shields 2005, Stentiford et al. 2012). Hematodinium sp. can negatively impact crab populations and fisheries and is of great interest to fishers and fisheries managers (Small 2012). Despite finding no molecular or histological evidence of Hematodinium sp. infection in M. mercenaria, we did find molecular evidence of an H. perezi infection in 1 M. spinosissimus individual, which establishes the presence of this pathogen in the nearshore hard-bottom habitat of the Florida Keys that is shared with M. mercenaria. The isolate found here is very similar to that found in stone crab in Georgia, USA (Sheppard et al. 2003). This pathogen can survive as a free-living stage in the environment outside a host (Pitula et al. 2012). It is also capable of being transferred between host species (Li et al. 2021), so it is important to continue to monitor stone crabs for infection using molecular techniques that allow for identification of specific isolates (Bojko et al. 2018).

# 4.4. Conclusions

We have documented several important ecological changes associated with the periodic cyanoHABs occurring in Florida Bay. Invertebrate communities differ between impacted and unimpacted hardbottom sites, and the loss of bivalves at impacted sites could be triggering bottom-up trophic changes, particularly for stone crabs. The apicomplexan gregarine Nematopsis sp. is present in stone crabs and their prey, T. pectinata, and could be used as an indicator of ecosystem health because it is less prevalent in stone crabs found at impacted sites. Evidence of resilience in the hard-bottom ecosystem is apparent, due to the presence of many small sponges at impacted sites, indicating regrowth since the major bloom 2 yr prior. *Hematodinium* sp. was not detected from stone crabs at either site, despite being identified in another crab species. A high prevalence of PaV1 in lobsters from unimpacted sites could be an indication of better settlement habitat for postlarval spiny lobsters. Future studies should examine the direct impacts of these cyanoHABs on bivalves and whether sponge restoration could help mitigate the impacts reported here.

Acknowledgements. We thank Kate Rose, Natalie Stephens, Doug Marcinek, Caleb Branam, Hannah Powless, Dalton Goss, Martin S. J. Rogers, Allie Marshbanks, and Sarah Garavaglia for assistance in the field and lab. Special thanks to Lucas Jennings for species identification expertise and guidance and to Martin S. J. Rogers for assistance with mapping. We appreciate Florida Bay Interagency Science Center for the use of their field station. This work was supported by an Aylesworth Foundation Scholarship and FL Skin Diver Association Scholarship to E.D.M.

## LITERATURE CITED

- Altizer S, Bartel R, Han BA (2011) Animal migration and infectious disease risk. Science 331:296–302
- Alvarez-Filip L, Gill JA, Dulvy NK (2011) Complex reef architecture supports more small-bodied fishes and longer food chains on Caribbean reefs. Ecosphere 2:118
- Amélineau F, Grémillet D, Harding AMA, Walkusz W, Choquet R, Fort J (2019) Arctic climate change and pollution impact little auk foraging and fitness across a decade. Sci Rep 9:1014
- Baker MD, Vossbrinck CR, Maddox JV, Undeen AH (1994)
  Phylogenetic relationships among Vairimorpha and
  Nosema species (Microspora) based on ribosomal RNA
  sequence data. J Invertebr Pathol 64:100–106
- Beck MA, Levander OA (2000) Host nutritional status and its effect on a viral pathogen. J Infect Dis 182:S93–S96
- Beck MW, Brumbaugh RD, Airoldi L, Carranza A and others (2011) Oyster reefs at risk and recommendations for conservation, restoration, and management. BioScience 61: 107–116
- Behringer DC (2012) Diseases of wild and cultured juvenile crustaceans: insights from below the minimum landing size. J Invertebr Pathol 110:225–233
- Behringer DC, Butler MJ (2006) Stable isotope analysis of production and trophic relationships in a tropical marine hard-bottom community. Oecologia 148:334–341
- Behringer DC, Duermit-Moreau E (2021) Crustaceans, One Health and the changing ocean. J Invertebr Pathol 186: 107500
- Behringer DC, Hart JE (2017) Competition with stone crabs drives juvenile spiny lobster abundance and distribution. Oecologia 184:205–218
- Behringer DC, Butler MJ, Shields JD (2006) Avoidance of disease by social lobsters. Nature 441:421
- Behringer DC, Karvonen A, Bojko J (2018) Parasite avoidance behaviours in aquatic environments. Philos Trans R Soc B 373:20170202
- Bertelsen RD, Butler MJ IV, Herrnkind WF, Hunt JH (2009)
  Regional characterisation of hard-bottom nursery habitat
  for juvenile Caribbean spiny lobster (*Panulirus argus*)
  using rapid assessment techniques. NZ J Mar Freshw
  Res 43:299–312
- Braicovich PE, Irigoitia MM, Bovcon ND, Timi JT (2021) Parasites of *Percophis brasiliensis* (Percophidae) benefited from fishery regulations: indicators of success for marine protected areas? Aquat Conserv 31:139–152
- RS, Friedman CS (2005) Health and survival of red abalone, *Haliotis rufescens*, under varying temperature,

- food supply, and exposure to the agent of withering syndrome. J Invertebr Pathol 89:219–231
- Bray JR, Curtis JT (1957) An ordination of the upland forest communities of Southern Wisconsin. Ecol Monogr 27: 325–349
- Burge CA, Eakin CM, Friedman CS, Froelich B and others (2014) Climate change influences on marine infectious diseases: implications for management and society. Annu Rev Mar Sci 6:249–277
- Butler J, Stanley JA, Butler MJ IV (2016) Underwater soundscapes in near-shore tropical habitats and the effects of environmental degradation and habitat restoration. J Exp Mar Biol Ecol 479:89–96
- Butler MJ IV, Hunt JH, Herrnkind WF, Childress MJ and others (1995) Cascading disturbances in Florida Bay, USA: cyanobacteria blooms, sponge mortality, and implications for juvenile spiny lobsters *Panulirus argus*. Mar Ecol Prog Ser 129:119–125
- Butler MJ IV, Behringer DC Jr, Dolan TW III, Moss J, Shields JD (2015) Behavioral immunity suppresses an epizootic in Caribbean spiny lobsters. PLOS ONE 10:e0126374
- \*Cannizzaro JP, Barnes BB, Hu C, Corcoran AA and others (2019) Remote detection of cyanobacteria blooms in an optically shallow subtropical lagoonal estuary using MODIS data. Remote Sens Environ 231:111227
- \*Capper A, Flewelling LJ, Arthur K (2013) Dietary exposure to harmful algal bloom (HAB) toxins in the endangered manatee (*Trichechus manatus latirostris*) and green sea turtle (*Chelonia mydas*) in Florida, USA. Harmful Algae 28:1–9
- Carmichael WW (2001) Health effects of toxin-producing cyanobacteria: 'The cyanoHABs'. Hum Ecol Risk Assess 7:1393–1407
  - Clopton RE (2002) The gregarines: a generic level review. In: Lee JJ, Leedale G, Patterson D, Bradbury PC (eds) Illustrated guide to the Protozoa, 2<sup>nd</sup> edn. Society of Protozoologists, Lawrence, KS, p 205–288
- Costanza R, D'Arge R, de Groot R, Farber S and others (1997) The value of the world's ecosystem services and natural capital. Nature 387:253–260
- Davies CE, Briones-Fourzán P, Barradas-Ortiz C, Negrete-Soto F, Moo-Cocom G, Lozano-Álvarez E (2020) Do ecological characteristics drive the prevalence of *Panulirus argus* virus 1 (PaV1) in juvenile Caribbean spiny lobsters in a tropical reef lagoon? PLOS ONE 15:e0229827
- Duermit E, Kingsley-Smith PR, Wilber DH (2015) The consequences of claw removal on stone crabs *Menippe* spp. and the ecological and fishery implications. N Am J Fish Manag 35:895–905
- Dufrene M, Legendre P (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. Ecol Monogr 67:345–366
  - Feldman AT, Wolfe D (2014) Tissue processing and hematoxylin and eosin staining. In: Day CE (ed) Histopathology. Humana Press, New York, NY, p 31–44
  - Fire SE, van Dolah FM (2012) Marine biotoxins: emergence of harmful algal blooms as health threats to marine wildlife. In: Aguirre AA, Ostfeld RS, Daszak P (eds) New directions in conservation medicine: applied cases of ecological health. Oxford University Press, New York, NY, p 374–383
- Fu FX, Tatters AO, Hutchins DA (2012) Global change and the future of harmful algal blooms in the ocean. Mar Ecol Prog Ser 470:207–233
- Gagne RB, Crooks KR, Craft ME, Chiu ES and others (2022)

- Parasites as conservation tools. Conserv Biol 36:e13719
  Gilbert BM, Avenant-Oldewage A (2021) Monogeneans as bioindicators: a meta-analysis of effect size of contaminant exposure toward Monogenea (Platyhelminthes). Ecol Indic 130:108062
  - Glibert PM, Heil CA, Rudnick DT, Madden CJ, Beach WP, Boyer JN, Kelly SP (2009) Florida Bay: water quality status and trends, historic and emerging algal bloom problems. Contrib Mar Sci 38:5–17
- Gravinese PM, Kronstadt SM, Clemente T, Cole C and others (2018) The effects of red tide (*Karenia brevis*) on reflex impairment and mortality of sublegal Florida stone crabs, *Menippe mercenaria*. Mar Environ Res 137: 145–148
- Griffith AW, Shumway SE, Gobler CJ (2019) Differential mortality of North Atlantic bivalve molluscs during harmful algal blooms caused by the dinoflagellate, Cochlodinium (a.k.a. Margalefidinium) polykrikoides. Estuaries Coasts 42:190–203
- Groner ML, Gettinby G, Stormoen M, Revie CW, Cox R (2014) Modelling the impact of temperature-induced life history plasticity and mate limitation on the epidemic potential of a marine ectoparasite. PLOS ONE 9:e88465
- Groner ML, Shields JD, Landers DF, Swenarton J, Hoenig JM (2018) Rising temperatures, molting phenology, and epizootic shell disease in the American lobster. Am Nat 192:E163–E177
- \*Hadas E, Marie D, Shpigel M, Ilan M (2006) Virus predation by sponges is a new nutrient-flow pathway in coral reef food webs. Limnol Oceanogr 51:1548–1550
- Hall MO, Furman BT, Merello M, Durako MJ (2016) Recurrence of *Thalassia testudinum* seagrass die-off in Florida Bay, USA: initial observations. Mar Ecol Prog Ser 560: 243–249
- Harvell CD, Mitchell CE, Ward JR, Altizer S, Dobson AP, Ostfeld RS, Samuel MD (2002) Climate warming and disease risks for terrestrial and marine biota. Science 296: 2158–2163
- Harvell D, Aronson R, Baron N, Connell J and others (2004)
  The rising tide of ocean diseases: unsolved problems and research priorities. Front Ecol Environ 2:375–382
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. Proc R Soc B 272:1059–1066
- Holmes JC (1996) Parasites as threats to biodiversity in shrinking ecosystems. Biodivers Conserv 5:975–983
- Hopwood D (1969) Fixatives and fixation: a review. Histochem J 1:323–360
- \*Hughes WOH, Eilenberg J, Boomsma JJ (2002) Trade-offs in group living: transmission and disease resistance in leafcutting ants. Proc R Soc B 269:1811–1819
- Huspeni TC, Lafferty KD (2004) Using larval trematodes that parasitize snails to evaluate a saltmarsh restoration project. Ecol Appl 14:795–804
- Ibelings BW, Backer LC, Kardinaal WEA, Chorus I (2014) Current approaches to cyanotoxin risk assessment and risk management around the globe. Harmful Algae 40:63–74
- Johnson PTJ, Wood CL, Joseph MB, Preston DL, Haas SE, Springer YP (2016) Habitat heterogeneity drives the host-diversity-begets-parasite-diversity relationship: evidence from experimental and field studies. Ecol Lett 19: 752-761
- Jokela J, Lively CM (1995) Spatial variation in infection by digenetic trematodes in a population of freshwater snails (Potamopyrgus antipodarum). Oecologia 103:509–517

- Karvonen A, Rintamäki P, Jokela J, Valtonen ET (2010) Increasing water temperature and disease risks in aquatic systems: Climate change increases the risk of some, but not all, diseases. Int J Parasitol 40:1483–1488
- Keesing F, Ostfeld RS (2021) Dilution effects in disease ecology. Ecol Lett 24:2490–2505
- Keesing F, Holt RD, Ostfeld RS (2006) Effects of species diversity on disease risk. Ecol Lett 9:485–498
- Khan RA (1990) Parasitism in marine fish after chronic exposure to petroleum hydrocarbons in the laboratory and to the Exxon Valdez oil spill. Bull Environ Contam Toxicol 44:759–763
- Lafferty KD (1997) Environmental parasitology: What can parasites tell us about human impacts on the environment? Parasitol Today 13:251–255
- Lafferty KD, Porter JW, Ford SE (2004) Are diseases increasing in the ocean? Annu Rev Ecol Evol Syst 35:31–54
- Lassudrie M, Hégaret H, Wikfors GH, da Silva PM (2020) Effects of marine harmful algal blooms on bivalve cellular immunity and infectious diseases: a review. Dev Comp Immunol 108:103660
- Leverone JR, Blake NJ, Pierce RH, Shumway SE (2006) Effects of the dinoflagellate *Karenia brevis* on larval development in three species of bivalve mollusc from Florida. Toxicon 48:75–84
- Leverone JR, Shumway SE, Blake NJ (2007) Comparative effects of the toxic dinoflagellate *Karenia brevis* on clearance rates in juveniles of four bivalve molluscs from Florida, USA. Toxicon 49:634–645
- Li C, Shields JD, Miller TL, Small HJ, Pagenkopp KM, Reece KS (2010) Detection and quantification of the free-living stage of the parasitic dinoflagellate *Hematodinium* sp. in laboratory and environmental samples. Harmful Algae 9: 515–521
- Li M, Huang Q, Lv X, Song S, Li C (2021) The parasitic dinoflagellate *Hematodinium* infects multiple crustaceans in the polyculture systems of Shandong Province, China. J Invertebr Pathol 178:107523
- Luis AD, Hayman DTS, O'Shea TJ, Cryan PM and others (2013) A comparison of bats and rodents as reservoirs of zoonotic viruses: Are bats special? Proc R Soc B 280: 20122753
- Maldonado M, Zhang X, Cao X, Xue L, Cao H, Zhang W (2010) Selective feeding by sponges on pathogenic microbes: a reassessment of potential for abatement of microbial pollution. Mar Ecol Prog Ser 403:75–89
- McCauley DJ, Pinsky ML, Palumbi SR, Estes JA, Joyce FH, Warner RR (2015) Marine defaunation: animal loss in the global ocean. Science 347:1255641
- Montgomery-Fullerton MM, Cooper RA, Kauffman KM, Shields JD, Ratzlaff RE (2007) Detection of *Panulirus argus* Virus 1 in Caribbean spiny lobsters. Dis Aquat Org 76:1–6
- Moore CS, Gittman RK, Puckett BJ, Wellman EH, Blakeslee AMH (2020) If you build it, they will come: Restoration positively influences free-living and parasite diversity in a restored tidal marsh. Food Webs 25:e00167
- Moss J, Butler MJ IV, Behringer DC, Shields JD (2012) Genetic diversity of the Caribbean spiny lobster virus, Panulirus argus virus 1 (PaV1), and the discovery of PaV1 in lobster postlarvae. Aquat Biol 14:223–232
  - NOAA (National Oceanic and Atmospheric Administration) (2021) Essential fish habitat mapper. https://www.fisheries.noaa.gov/resource/map/essential-fish-habitat-mapper (accessed 12 October 2021)

- Nunn CL, Altizer S, Jones KE, Sechrest W (2003) Comparative tests of parasite species richness in primates. Am Nat 162:597–614
  - Oksanen AJ, Blanchet FG, Friendly M, Kindt R and others (2019) Vegan: community ecology package. R package version 2.5-7. https://CRAN.R-project.org/package=vegan
- Orth R, van Montfrans J (1987) Utilization of a seagrass meadow and tidal marsh creek by blue crabs *Callinectes sapidus*. I. Seasonal and annual variations in abundance with emphasis on post-settlement juveniles. Mar Ecol Prog Ser 41:283–294
- Covcharenko MO, Bacela K, Wilkinson T, Ironside JE, Rigaud T, Wattier RA (2010) Cucumispora dikerogammari n. gen. (Fungi: Microsporidia) infecting the invasive amphipod Dikerogammarus villosus: a potential emerging disease in European rivers. Parasitology 137:191–204
- Owens L (1983) Bopyrid parasite *Epipenaeon ingens* Nobili as a biological marker for the banana prawn, *Penaeus merguiensis* de Man. Mar Freshw Res 34:477–481
- Paerl HW, Huisman J (2008) Climate: Blooms like it hot. Science 320:57–58
- Pagenkopp Lohan KM, Reece KS, Miller TL, Wheeler KN, Small HJ, Shields JD (2012) The role of alternate hosts in the ecology and life history of *Hematodinium* sp., a parasitic dinoflagellate of the blue crab (*Callinectes sapidus*). J Parasitol 98:73–84
- Pagenkopp Lohan KM, Small HJ, Shields JD, Place AR, Reece KS (2013) Conservation in the first internal transcribed spacer (ITS1) region of *Hematodinium perezi* (genotype III) from *Callinectes sapidus*. Dis Aquat Org 103:65–75
- Peterson BJ, Chester CM, Jochem FJ, Fourqurean JW (2006)
  Potential role of sponge communities in controlling
  phytoplankton blooms in Florida Bay. Mar Ecol Prog Ser
  328:93–103
- Phlips EJ, Zeman C, Hansen P (1989) Growth, photosynthesis, nitrogen fixation and carbohydrate production by a unicellular cyanobacterium, *Synechococcus* sp. (Cyanophyta). J Appl Phycol 1:137–145
- Pitula JS, Dyson WD, Bakht HB, Njoku I, Chen F (2012)
  Temporal distribution of genetically homogenous 'free-living' *Hematodinium* sp. in a Delmarva coastal ecosystem. Aquat Biosyst 8:16
- Pravdová M, Kolářová J, Grabicová K, Mikl L and others (2021) Associations between pharmaceutical contaminants, parasite load and health status in brown trout exposed to sewage effluent in a small stream. Ecohydrol Hydrobiol 21:233–243
  - Puls D (2015) Effects of simulated cyanobacteria blooms and sediment resuspension on several sponge species from Florida Bay. MSc thesis, University of Florida, Gainesville, FL
- Reiswig HM (1971) Particle feeding in natural populations of three marine demosponges. Biol Bull (Woods Hole) 141: 568–591
- Rindone RR, Eggleston DB (2011) Predator–prey dynamics between recently established stone crabs (*Menippe* spp.) and oyster prey (*Crassostrea virginica*). J Exp Mar Biol Ecol 407:216–225
  - Roberts DW (2019) Labdsv: ordination and multivariate analysis for ecology. R package version 1.3-1. https:// CRAN.R-project.org/package=labdsv
- Rolton A, Vignier J, Soudant P, Shumway SE, Bricelj VM, Volety AK (2014) Effects of the red tide dinoflagellate, Karenia brevis, on early development of the eastern oys-

- ter Crassostrea virginica and northern quahog Mercenaria mercenaria. Aquat Toxicol 155:199–206
- Shannon CE (1948) A mathematical theory of communication. Bell Syst Tech J 27:379–423
  - Sheppard M, Walker A, Frischer ME, Lee RF (2003) Histopathology and prevalence of the parasitic dinoflagellate, *Hematodinium* sp, in crabs (*Callinectes sapidus*, *Callinectes similis, Neopanope sayi, Libinia emarginata*, *Menippe mercenaria*) from a Georgia estuary. J Shellfish Res 22:873–880
- Shields JD, Behringer DC Jr (2004) A new pathogenic virus in the Caribbean spiny lobster *Panulirus argus* from the Florida Keys. Dis Aquat Org 59:109–118
- Sigler MF, Tollit DJ, Vollenweider JJ, Thedinga JF and others (2009) Steller sea lion foraging response to seasonal changes in prey availability. Mar Ecol Prog Ser 388: 243–261
- Silva TJ, Soares EC, Casal G, Rocha S and others (2019)
  Ultrastructure of phagocytes and oocysts of *Nematopsis*sp. (Apicomplexa, Porosporidae) infecting *Crassostrea*rhizophorae in Northeastern Brazil. Rev Bras Parasitol
  Vet 28:97–104
- Simpson EH (1949) Measurement of diversity. Nature 163:
- Sitko J, Heneberg P (2020) Systemic collapse of a host–parasite trematode network associated with wetland birds in Europe. Parasitol Res 119:935–945
- Smale DA, Wernberg T, Oliver ECJ, Thomsen M and others (2019) Marine heatwaves threaten global biodiversity and the provision of ecosystem services. Nat Clim Change 9:306–312
- Small HJ (2012) Advances in our understanding of the global diversity and distribution of *Hematodinium* spp.—significant pathogens of commercially exploited crustaceans. J Invertebr Pathol 110:234–246
  - Sprague V (1949) Species of *Nematopsis* in *Ostrea virginica*. J Parasitol 35:243–256
- Sprague V, Orr PE Jr (1955) Nematopsis ostrearum and N.

- prytherchi (Eugregarinina: Porosporidae) with special reference to the host–parasite relations. J Parasitol 41:  $89{\text -}104$
- Stentiford GD, Shields JD (2005) A review of the parasitic dinoflagellates *Hematodinium* species and *Hematodinium*-like infections in marine crustaceans. Dis Aquat Org 66:47–70
- Stentiford GD, Neil DM, Peeler EJ, Shields JD and others (2012) Disease will limit future food supply from the global crustacean fishery and aquaculture sectors.

  J Invertebr Pathol 110:141–157
- Subramaniam K, Behringer DC, Bojko J, Yutin N and others (2020) A new family of DNA viruses causing disease in crustaceans from diverse aquatic biomes. MBio 11: e02938-19
- Sunda WG, Graneli E, Gobler CJ (2006) Positive feedback and the development and persistence of ecosystem disruptive algal blooms. J Phycol 42:963–974
- Tlusty MF, Smolowitz RM, Halvorson HO, DeVito SE (2007) Host susceptibility hypothesis for shell disease in American lobsters. J Aquat Anim Health 19:215–225
- Travers MAS, Basuyaux O, le Goïc N, Huchette S, Nicolas JL, Koken M, Paillard C (2009) Influence of temperature and spawning effort on *Haliotis tuberculata* mortalities caused by *Vibrio harveyi*: an example of emerging vibriosis linked to global warming. Glob Change Biol 15:1365–1376
- Valentine MM, Butler MJ IV (2019) Sponges structure water-column characteristics in shallow tropical coastal ecosystems. Mar Ecol Prog Ser 608:133–147
- Westinga E, Hoetjes PC (1981) The intrasponge fauna of Spheciospongia vesparia (Porifera, Demospongiae) at Curação and Bonaire. Mar Biol 62:139–150
- Wood CL, Lafferty KD, Micheli F (2010) Fishing out marine parasites? Impacts of fishing on rates of parasitism in the ocean. Ecol Lett 13:761–775
- Zuk M, McKean KA (1996) Sex differences in parasite infections: patterns and processes. Int J Parasitol 26: 1009-1024

# **Appendix**

Table A1. Forward and reverse primer sequences used for the amplification of pathogen groups via PCR from genomic template, extracted from host and pathogen tissues. Each PCR run included an initial 5 min denaturation step and a 5 min final extension step, according to the first and final temperatures, respectively, noted in the thermocycler (Tc) settings. The amplification stage consisted of 35 cycles of all 3 temperatures in the Tc settings, with each temperature being held for 1 min or as detailed below. Dir.: direction

Infection	n ——		— Primer — — — — — — — — — — — — — — — — — — —	Tc settings	Amplicon	Reference
	Dir.	Name	Sequence (5'–3')	(°C)	size (bp)	
PaV1						
	Fwd	45aF	TTC CAG CCC AGG TAC GTA TC	94-63-72	499	Montgomery-Fullerton
	Rev	543aR	AAC AGA TTT TCC AGC AGC GT	(45s-45s-1 mir	1)	et al. (2007), Moss et al. (2012)
Ameson	n herrn	kindi				
	Fwd	V1F	CAC CAG GTT GAT TCT GCC TGA C	94-52-72	1100	Baker et al. (1994),
	Rev	MC3R	GAT AAC GAC GGG CGG TGT GTA CA	A		Ovcharenko et al. (2010)
Hemato	diniun	n sp.				,
1 <sup>st</sup>	Fwd	2009ITS1F	AAC CTG CGG AAG GAT CAT TC	94-60-72	500	Bojko et al. (2018)
round	Rev	2009ITS1&2R	TAG CCT TGC CTG ACT CAT G			, ,
$2^{\rm nd}$	Fwd	2009ITS1F	AAC CTG CGG AAG GAT CAT TC	94-60-72	350	
round	Rev	2009ITS1R	CCG AGC CGA GGC ATT CAT CGC T			