

STUDIES

Spatial variation in bidirectional pollinator-mediated interactions between two co-flowering species in serpentine plant communities

Amber Stanley^{1,2,†}, Carlos Martel^{1,3,†} and Gerardo Arceo-Gómez^{1,*}

¹Department of Biological Sciences, East Tennessee State University, Johnson City, TN 37614, USA, ²Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA, ³Instituto de Ciencias Ómicas y Biotecnología Aplicada, Pontificia Universidad Católica del Perú, San Miguel 15088, Lima, Peru

*Corresponding author's e-mail address: gomezg@etsu.edu

†These authors share first authorship.

Populations & Communities. Chief Editor: Jean Burns

Associate Editor: Karina Boege

Abstract

Pollinator-mediated competition and facilitation are two important mechanisms mediating co-flowering community assembly. Experimental studies, however, have mostly focused on evaluating outcomes for a single interacting partner at a single location. Studies that evaluate spatial variation in the bidirectional effects between co-flowering species are necessary if we aim to advance our understanding of the processes that mediate species coexistence in diverse co-flowering communities. Here, we examine geographic variation (i.e. at landscape level) in bidirectional pollinator-mediated effects between co-flowering *Mimulus guttatus* and *Delphinium uliginosum*. We evaluated effects on pollen transfer dynamics (conspecific and heterospecific pollen deposition) and plant reproductive success. We found evidence of asymmetrical effects (one species is disrupted and the other one is facilitated) but the effects were highly dependent on geographical location. Furthermore, effects on pollen transfer dynamics did not always translate to effects on overall plant reproductive success (i.e. pollen tube growth) highlighting the importance of evaluating effects at multiple stages of the pollination process. Overall, our results provide evidence of a spatial mosaic of pollinator-mediated interactions between co-flowering species and suggest that community assembly processes could result from competition and facilitation acting simultaneously. Our study highlights the importance of experimental studies that evaluate the prevalence of competitive and facilitative interactions in the field, and that expand across a wide geographical context, in order to more fully understand the mechanisms that shape plant communities in nature.

Keywords: Competition; *Delphinium uliginosum*; facilitation; heterospecific pollen; *Mimulus guttatus*.

Introduction

It has been widely demonstrated that pollinator-mediated interactions between co-flowering plants can have both direct (e.g. improper pollen transfer; Morales and Traveset 2008; Ashman and Arceo-Gómez 2013) and indirect effects (e.g. via pollinator visitation; Mitchell et al. 2009; Losapio and Schöb

2020) on plant reproductive success. These interactions can in turn have important ecological and evolutionary consequences and influence floral evolution and co-flowering community assembly (Galen 1999; Kay and Sargent 2009; Mitchell et al. 2009; Ashman and Arceo-Gómez 2013; Carvalheiro et al. 2014). Among

Received: 22 June 2021; Editorial decision: 4 October 2021; Accepted: 19 October 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

these, two main types of indirect plant–plant interactions via changes in pollinator preference have been described, i.e. pollinator competition and facilitation (Campbell 1985; Brown *et al.* 2002; Feldman *et al.* 2004; Ghazoul 2006; Mitchell *et al.* 2009). Competition occurs when the presence of one species decreases pollinator visitation to another (e.g. Campbell 1985; Brown *et al.* 2002; Mitchell *et al.* 2009), negatively affecting the pollination success of one or both species; whereas in facilitation pollinator visitation increases for at least one plant species when two or more species flower simultaneously (e.g. Feldman *et al.* 2004; Ghazoul 2006; Carvalheiro *et al.* 2014) leading to an increase in pollination success. Thus, pollinator-mediated interactions between two plants species can result in six different scenarios, i.e. positive outcomes for both partners (+: +), positive for one and negative for the other (+: -), positive and neutral (+: 0), negative and neutral (-: 0) and negative (-: -) or neutral (0: 0) for both partners. Pollinator-mediated interactions however have historically been studied by focusing on only one of the interacting partners (e.g. Campbell 1985; Morales and Traveset 2008; Flanagan *et al.* 2009; Muchhala *et al.* 2009; Arceo-Gómez and Ashman 2011; but see Caruso and Alfaro 2000; Ghazoul 2006; Muchhala *et al.* 2010; Suárez-Mariño *et al.* 2019; Streher *et al.* 2020; Bergamo *et al.* 2020b; Ha *et al.* 2021). Although central in advancing our understanding of the drivers and consequences of pollinator-mediated interactions, studies that focus on only one interacting partner may underestimate the real complexity of these interactions in nature. Specifically, a unidirectional focus overlooks the potential for bidirectional effects between individual species pairs (e.g. Bascompte and Jordano 2007; Vila *et al.* 2009), which can further influence floral evolution and community assembly. In this sense, plant species pairs can experience either similar (both impair or facilitate each other) or asymmetrical effects (e.g. one experiences competition while the other one is facilitated; Caruso and Alfaro 2000; Ghazoul 2006; Muchhala *et al.* 2010; Suárez-Mariño *et al.* 2019). It is also possible that one species will be affected (positively or negatively) while its interacting partner will not (neutral effect). However, the occurrence and directionality of these two-sided interactions have surprisingly received considerably less attention compared to the study of unidirectional effects. Studies that consider bidirectional effects between co-flowering species are necessary if we aim to advance our understanding of the processes that mediate plant species coexistence in diverse co-flowering communities.

The strength and direction of pollinator-mediated interactions can vary spatially due to changes in the composition of the surrounding plant and pollinator community, potentially leading to geographic mosaics in the outcomes of species interactions (e.g. Thompson 1999, 2005; Moeller 2005). For example, changes in pollinators' availability or co-flowering community composition can influence patterns of pollinator-mediated facilitation and competition (Anderson and Johnson 2007; Lázaro *et al.* 2009; Martín-Rodríguez *et al.* 2011). However, very few studies have evaluated geographic variation in the outcomes of pollinator-mediated interactions between co-flowering species (Olesen and Jordano 2002; Burkle *et al.* 2016). It is thus imperative that we fully evaluate the extent of spatial variation in the outcome of pollinator-mediated interactions and how this may lead to spatial mosaics of floral evolution and community assembly (Thompson *et al.* 1999, 2005). Finally, the strength and direction of pollinator-mediated plant–plant interactions have often been determined by evaluating effects on patterns of pollinator visitation alone (e.g. Waser *et al.* 1983; Feldman *et al.* 2004; Mitchell *et al.* 2009). However, studies have

shown that pollinator visitation may not accurately reflect pollination success (King *et al.* 2013; Ballantyne *et al.* 2015; Stavert *et al.* 2016), mainly because pollinators vary in their ability to successfully transfer and deposit conspecific pollen (Benevides *et al.* 2013; King *et al.* 2013; Stavert *et al.* 2016). Pollinators may also vary in the quality of pollen they transport (conspecific vs. heterospecific pollen; Morales and Traveset 2008; Ashman *et al.* 2020). Heterospecific pollen transfer has been shown to impose strong negative effects on males (Muchhala *et al.* 2010) and female reproductive success (Ashman and Arceo-Gómez 2013), potentially strengthening the effects of pollinator competition or offsetting the effects of facilitation via increased pollinator visitation. Thus, heterospecific pollen transfer may have the potential to affect the directionality of pollinator-mediated interactions among co-flowering plants. For instance, the presence of one plant species may lead to increased visitation to another species (i.e. reproductive facilitation via increased visitation) but also lead to an increase in heterospecific pollen (i.e. reproductive competition via pollen interference) and the overall outcome of the interaction may be negative (e.g. Morales and Traveset 2008; Thomson *et al.* 2019). Thus, studies that evaluate the strength and direction of pollinator-mediated interactions by integrating multiple aspects of the pollination process including conspecific and heterospecific pollen deposition as well as pollination success (e.g. pollen tube growth) are key in order to fully evaluate the consequences of these interactions in nature.

In this study, we examine geographical variation in the bidirectional effects of pollinator-mediated interactions between two co-flowering species, *Mimulus guttatus* (Phrymaceae) and *Delphinium uliginosum* (Ranunculaceae). We evaluated these effects across multiple stages of the pollination process (i.e. conspecific and heterospecific pollen deposition, pollen tube growth). These two species differ widely in their floral phenotypes (Fig. 1). Furthermore, *M. guttatus* is typically more abundant and considered a pollinator generalist while *D. uliginosum* tends to be primarily bumblebee-pollinated, which also visits *M. guttatus* flowers (Koski *et al.* 2015). Thus, it is possible to expect that *M. guttatus* may help facilitate pollination success in the less abundant *D. uliginosum* (Wei *et al.* 2021). However, this may occur at the detriment of *M. guttatus* pollination, either via a decrease in conspecific pollen or via an increase in heterospecific pollen deposition (Wei *et al.* 2021). Specifically, we ask the following questions: (i) do patterns of conspecific and heterospecific pollen deposition, and pollen tube growth differ between *M. guttatus* and *D. uliginosum* when they grow alone compared to when they occur together? (ii) is the outcome of species pair interactions the same or asymmetrical? and (iii) does the outcome of these interactions vary across three different sites?

Materials and Methods

Focal species

We use *M. guttatus* as a focal species because it is a key species mediating indirect pollinator interactions in the serpentine seep co-flowering communities in northern California (Koski *et al.* 2015). Due to its long flowering period and high floral abundance *M. guttatus* has the potential to engage in positive and negative pollinator-mediated interactions with a wide array of co-flowering species across the season (Koski *et al.* 2015). *Mimulus guttatus* (MIGU) produces yellow zygomorphic flowers (Fig. 1) that are predominantly pollinated by bumblebees but also attract a variety of other pollinators including solitary bees,



Figure 1. Studied plant species. (A) *Mimulus guttatus* and (B) *Delphinium uliginosum*.

beetles, butterflies and flies (Arceo-Gómez and Ashman 2014a; Koski et al. 2015). We specifically evaluated how *M. guttatus* interacted with *D. uliginosum* via pollinators. We chose *D. uliginosum* due to its high abundance at the seeps and its high pollinator use overlap with *M. guttatus* (Koski et al. 2015) despite strong differences in flower morphology (i.e. size, shape and colour; Fig. 1). *Delphinium uliginosum* (DEUL) is a perennial herb that produces blue-purple flowers that are primarily pollinated by bumblebees at the study sites, but plants are also visited by butterflies and other bees (Koontz and Warnock 2012; Koski et al. 2015; G. Arceo-Gómez, unpubl. data; Fig. 1).

Study sites

This study was conducted at three locations (i.e. serpentine seeps; here identified as BS, RHA, TP9) at the McLaughlin Natural Reserve in Lower Lake California, USA (38°52'16.2"N, 122°25'09.5"W). The seeps are separated by an average of 2.01 ± 0.46 km (mean $\pm$ SD) and are surrounded by vast swathes of grassland, small shrubs and chaparral species. Seep communities are typically composed of a mix of annual and perennial animal-pollinated species. Species richness at the seeps can vary from only a few to more than 30 plant species (Arceo-Gómez and Ashman 2014a; Koski et al. 2015). Serpentine seeps are ecologically similar in that they share similar environmental conditions and unique soil chemistry and have thus been considered as replicates of similar plant-pollinator communities within the same geographic area (Freestone and Harrison 2006; Arceo-Gómez and Ashman 2014a; Arceo-Gómez et al. 2016, 2018). Our sites were selected based on the presence of both focal species and were composed of a core of 39–62 species. Each site was visited twice per week between 16 and 31 May in 2018 (five total visits per site), which falls within the flowering peak of the studied species (Arceo-Gómez et al. 2018). At each site, we delineated 1×2 m plots ensuring plots contained one of three species composition treatments: (i) *M. guttatus* alone (MIGU only), (ii) a mix of *M. guttatus* and *D. uliginosum* (DEUL-MIGU) and (iii) *D. uliginosum* alone (DEUL

only). We established 5–7 plots (separated by at least 5 m) for each species composition treatment per seep for a total of 17–19 plots per treatment across all three seeps (BS: 5 DEUL only, 6 MIGU only, 6 DEUL-MIGU; RHA: 7 DEUL only, 6 MIGU only, 6 DEUL-MIGU; TP9: 6 DEUL only, 6 MIGU only, 6 DEUL-MIGU). Although plots sometimes contained individuals of other plant species in the community ~90 % of flowers (average 138.4 flowers per plot) within the plot belonged to the focal species studied.

Pollen transfer dynamics and post-pollination success

To evaluate differences in pollen transfer dynamics and pollen tube growth between plants in different plot treatments we collected three styles per species per plot during each visit to a site, (each style from a different individual). We collected styles from flowers that were at the end of their lifetime (wilted flowers) and thus were no longer visited. In total we collected 172–195 *M. guttatus* styles (87–90 single and 85–102 mixed plots), and 89–166 styles of *D. uliginosum* (48–96 single and 41–71 mixed plots) across all three seeps. Styles were stored in 70 % ethanol until processing and softened and stained with aniline blue using standard methods in the laboratory (Dafni 1992; Arceo-Gómez et al. 2016). We collected data for a total of 950 flowers (539 of *M. guttatus* and 411 of *D. uliginosum*).

Pollen loads on the stigma were visualized using a compound microscope at 400 $\times$ magnification. We recorded the total number of conspecific pollen (pollen belonging to the recipient species) and heterospecific pollen (pollen belonging to a different species than the recipient). We did not limit our heterospecific pollen counts to only our focal plant species, but that from any other co-flowering plant species (focal and non-focal). To aid in the identification of pollen grains as either *M. guttatus* or *D. uliginosum*, we previously created a pollen reference library by collecting pollen from mature anthers from both species in the field. A total of 178 276 pollen grains were counted across the two species and the three seeps.

We also counted the number of pollen tubes that reached the base of the style under a fluorescent microscope. The number of pollen tubes has been considered a good proxy of pollination success (e.g. Alonso *et al.* 2012; Arceo-Gómez and Ashman 2014b). We performed this for a subset of randomly selected slides from each treatment group. A total of 50 *M. guttatus* (25 from mixed, 25 from single plots), and 50 *D. uliginosum* (25 from mixed, 25 from single plots) slides were counted. *Delphinium uliginosum* flowers have 1–3 styles and thus conspecific pollen loads, heterospecific pollen loads and pollen tubes were counted on all styles and added.

Data analysis

To evaluate the strength and direction of pollinator-mediated interactions between the two co-flowering species (i.e. *M. guttatus* and *D. uliginosum*), we ran three separate analyses. We conducted generalized linear mixed models to evaluate differences in conspecific pollen loads, heterospecific pollen loads and the proportion of pollen tubes that reached the base of the style (i.e. number of pollen tubes/total conspecific pollen load). Evaluating differences in conspecific and heterospecific pollen loads separately allows evaluating the relative importance of the two main mechanisms by which plants interact with each other via pollinators, i.e. via pollinator visitation (hence altering conspecific pollen receipt; Mitchell *et al.* 2009) and via heterospecific pollen interference on the stigma (Ashman and Arceo-Gómez 2013). Further, the evaluation of the proportion of pollen tubes reflects the effect of pollinator-mediated interactions on overall female fitness. For all analyses we used plot species composition treatment (i.e. alone vs. mixed), site and plant species as fixed effects. Plot ID was set as a random factor. Because we were also interested in evaluating whether responses between alone versus mixed treatments vary by site, species or both, we also included the site * treatment, treatment * species and the three-way interaction as fixed effects. We used negative binomial distribution with log link function for conspecific and heterospecific pollen loads and a binomial distribution with logit link function for the proportion of pollen tubes. When a three-way interaction was significant, we used *a priori* contrasts (see Arceo-Gómez and Ashman 2014a) to specifically evaluate differences between single and mixed plot treatments for each species at each one of the three seeps (e.g. Molofsky and Fisher 1993; Rosenthal *et al.* 2000; Strauss and Murch 2004; Arceo-Gómez and Ashman 2014a). All flowers within a plot (flower density) were counted and used as a covariate in the analyses. Model fit (overdispersion) was evaluated using the residual deviance against the degrees of freedom of the model; this resulted in values close to one, which indicates that our models were not overdispersed. All analyses were conducted using the *lme4* (Bates *et al.* 2019), *MASS* (Ripley *et al.* 2020) and *emmeans* packages (Lenth *et al.* 2020) in R (R version 3.6.1; R Development Core Team 2017).

Results

Conspecific pollen deposition

Average conspecific pollen load was significantly higher in *M. guttatus* (mean $\pm$ SD: 262.8 $\pm$ 9.1 pollen grains) compared to *D. uliginosum* (mean $\pm$ SD: 73.8 $\pm$ 3.5 pollen grains; $\chi^2 = 255.70$, $df = 1$, $P < 0.001$). This however is not surprising since the former produces far more ovules (~400 seeds; Arceo-Gómez and Ashman 2014a) compared to the latter (typically less than 50; G. Arceo-Gómez, pers. obs.). Conspecific pollen load size also

differed significantly among sites ($\chi^2 = 19.71$, $df = 2$, $P < 0.001$), but not between species composition treatments (single vs. mixed; $\chi^2 = 0.18$, $df = 1$, $P = 0.67$). However, we found a significant site * treatment ($\chi^2 = 13.1$, $df = 2$, $P = 0.001$), treatment * species ($\chi^2 = 18.36$, $df = 1$, $P < 0.001$) and a marginally significant site * treatment * species interaction ($\chi^2 = 5.34$, $df = 2$, $P < 0.07$; Fig. 2). *A priori* contrasts showed that *D. uliginosum* received significantly more conspecific pollen when growing with *M. guttatus* (MIGU-DEUL plot) compared to when growing alone (DEUL-only plot), although this difference was only significant at one site (i.e. RHA; $P < 0.001$ for both; Fig. 2). On the other hand, *M. guttatus* tended to receive larger conspecific pollen loads when growing alone (single MIGU plots) although these differences were not significant ($P > 0.05$ for all; Fig. 2). Floral density had a positive significant effect on conspecific pollen receipt ($\chi^2 = 5.81$, $df = 1$, $P < 0.05$).

Heterospecific pollen deposition

Average heterospecific pollen load size differed significantly between *M. guttatus* (mean $\pm$ SD: 7.41 $\pm$ 0.72 heterospecific pollen grains) and *D. uliginosum* (mean $\pm$ SD: 5.50 $\pm$ 0.58; $\chi^2 = 12.07$, $df = 1$, $P < 0.001$), among sites ($\chi^2 = 40.88$, $df = 2$, $P < 0.001$), and was marginally significant between plot species composition treatments (single vs. mixed; $\chi^2 = 3.20$, $df = 1$, $P = 0.07$). We also found a significant site * treatment interaction ($\chi^2 = 9.44$, $df = 2$, $P < 0.01$), but the treatment * species interaction was not significant ($\chi^2 = 0.23$, $df = 1$, $P = 0.63$). However, the site * treatment * species interaction was significant ($\chi^2 = 14.94$, $df = 2$, $P < 0.001$; Fig. 3). *A priori* contrasts showed that *M. guttatus* receives significantly larger amounts of heterospecific pollen when growing in mixed (MIGU-DEUL) compared to MIGU-only plots, but only at one site (i.e. RHA, $P < 0.001$; Fig. 3). Single versus mixed plot differences at the remaining two sites were not significant ($P > 0.05$; Fig. 3). We were unable to detect differences in heterospecific pollen load size between *D. uliginosum* plants growing in single versus mixed plots across all sites ($P > 0.05$ for all; Fig. 3). Floral density did not affect heterospecific pollen load size ($\chi^2 = 1.43$, $df = 1$, $P = 0.23$).

Pollen tube growth

The proportion of pollen tubes (pollen tubes/conspecific pollen load) did not differ significantly between *M. guttatus* (mean $\pm$ SD: 0.40 $\pm$ 0.04) and *D. uliginosum* (mean $\pm$ SD: 0.26 $\pm$ 0.24; $\chi^2 = 1.05$, $df = 1$, $P = 0.31$) or between plot species composition treatments (alone vs. mixed; $\chi^2 = 0.76$, $df = 1$, $P = 0.38$). Although we identified differences among sites ($\chi^2 = 7.71$, $df = 2$, $P < 0.05$), we did not find significant effects of the site * treatment ($\chi^2 = 3.20$, $df = 2$, $P = 0.20$), site * species ($\chi^2 = 1.27$, $df = 2$, $P = 0.53$), treatment * species ($\chi^2 = 0.67$, $df = 1$, $P = 0.41$) or site * treatment * species interactions ($\chi^2 = 0.70$, $df = 2$, $P = 0.70$) on the proportion of pollen tubes (Fig. 4). Floral density did not have an effect on the proportion of pollen tubes ($\chi^2 = 0.01$, $df = 1$, $P = 0.92$).

Discussion

Evaluating the processes that mediate plant species coexistence is central for developing a predictive understanding of the mechanisms that govern the assembly of plant communities and how these may change in response to human disturbances (Sargent and Ackerly 2008; Elo *et al.* 2016). In this sense, pollinator competition and facilitation have been commonly identified as the most important single-acting forces in the organization of co-flowering communities at a local scale (after dispersal and




Figure 2. Number (mean $\pm$ SE) of conspecific pollen grains found on styles by *D. uliginosum* and *M. guttatus* in single or mixed plots at three different sites (BS, RHA, TP9). Significant post hoc comparisons among species * site * treatment interactions are shown. ***P < 0.001.




Figure 3. Number (mean $\pm$ SE) of heterospecific pollen grains found on styles of *D. uliginosum* and *M. guttatus* in single or mixed plots at three different sites (BS, RHA, TP9). Significant post hoc comparisons among species * site * treatment interactions are shown. ***P < 0.001.




Figure 4. Proportion of pollen tubes/conspecific pollen found on styles of *D. uliginosum* and *M. guttatus* in single or mixed plots. Post hoc comparisons between species * site * treatment interactions were not significant.

environmental filters; Levine 1999; Choler et al. 2001; Callaway et al. 2002; Brooker 2006; Brooker et al. 2008; Mitchell et al. 2009). Our results, however, provide evidence suggesting a more complex mosaic of pollinator-mediated interactions and that community assembly processes could result from both types of interaction (i.e. competition and facilitation) acting together rather than singly. Specifically, we show evidence of bidirectional interactions between plant species with asymmetrical outcomes that are highly dependent on the site where they occur. Experiments that evaluate the prevalence of these interactions in the field, and that expand across a wide geographical context, are necessary to more fully understand the mechanisms that shape plant communities in nature.

The distribution of plant functional traits within a community has been increasingly used to infer the processes that govern plant community assembly (Westoby and Wright 2006; Kraft and Ackerly 2014; Xu et al. 2018; Navarro-Cano et al. 2021), including those mediated by pollinator-mediated interactions (e.g. Sargent and Ackerly 2008; McEwen and Vamosi 2010; de Jager et al. 2011; Wolowski et al. 2017; Bergamo et al. 2018). However, underlying this approach is the implicit assumption that either competition or facilitation is singly responsible for structuring co-flowering communities. Here we show evidence that both competition and facilitation could act simultaneously. Specifically, *M. guttatus* tended to receive less conspecific pollen (although not significantly; Fig. 2) and more heterospecific pollen (e.g. at RHA; Fig. 3) when co-flowering with *D. uliginosum*. These results are consistent with both forms of pollinator-mediated competition, i.e. via changes in pollinator visitation and via heterospecific pollen interference. On the contrary, *D. uliginosum* received significantly more conspecific pollen when

co-flowering with *M. guttatus* (at RHA; Fig. 2) thus indicating that it may benefit from the presence of *M. guttatus*, at least at some sites (see below). Interestingly, plant–pollinator network studies have long proposed the existence of strong asymmetries in the strength and potential effects of species interactions (Jordano 1987; Vázquez and Aizén 2004; Bascompte et al. 2006). Therefore, it is somewhat surprising that these asymmetries, and their role in community assembly, have not been more often considered when evaluating community-wide consequences of pollinator-mediated interactions. For instance, recent studies that have evaluated the role of phenotypic traits in determining the outcome of community-wide species interactions have also suggested that both, competition and facilitation, likely play a role in the assembly of plant communities (Tur et al. 2016; Bergamo et al. 2018; Navarro-Cano et al. 2021; Wei et al. 2021). Although the importance of the interplay between competition and facilitation in the assembly of co-flowering communities was proposed more than 35 years ago (Rathcke 1983; Mitchell et al. 2009), only few studies have experimentally evaluated the bidirectionality of pollinator-mediated interactions between co-flowering species (i.e. Randall and Hilu 1990; Rahmé et al. 2009; Sieber et al. 2011; Briscoe Runquist 2012; Briscoe Runquist and Stanton 2013; Wassink and Caruso 2013). We thus emphasize the need for experimental studies that evaluate the directionality of pollinator-mediated interactions in the field in order to validate predictions resulting from trait-based assembly and network studies. Such studies would also help to develop a more predictive understanding of the role of competition and facilitation in structuring plant communities.

In our study, the asymmetrical outcome in patterns of pollen receipt observed between *M. guttatus* and *D. uliginosum* may be the

result of differences in several floral characteristics. For instance, *D. uliginosum* flowers have large amounts of nectar while *M. guttatus* flowers offer little to no nectar at our study sites (Robertson et al. 1994; G. Arceo-Gómez, pers. obs.). Hence, bumblebees (primary floral visitors) may be highly attracted to mixed-species plots but preferentially visit the resource rich *D. uliginosum* flowers, to the detriment of *M. guttatus* flowers, thus generating positive effects for the former and negative effects for the latter. Furthermore, *M. guttatus* flowers are more generalized and also visited by other insects such as beetles, flies and butterflies (Koski et al. 2015; Wei et al. 2021), which may be more abundant in mixed-species plots but are less efficient and deposit higher amounts of heterospecific pollen loads. On the contrary, *D. uliginosum* is more specialized and primarily visited by bumblebees that feed on nectar-rich resources, which may result in higher visitor efficiency and fidelity. In fact, bumblebee visitation was more than three times higher for *D. uliginosum* flowers (0.007 visits per flower per min) than for *M. guttatus* (0.002 visits per flower per min; G. Arceo-Gómez, unpubl. data). *Mimulus guttatus* in turn receives a higher visitation from other insects in the community (0.017 visits per flower per min) compared to *D. uliginosum* (0.005 visits per flower per min; G. Arceo-Gómez, unpubl. data). Moreover, *M. guttatus* is a dominant species (70 flowers per m²) in the studied seep communities, particularly compared to *D. uliginosum* (32 flowers per m²; G. Arceo-Gómez, unpubl. data; also see Koski et al. 2015). Abundant species have been proposed to be more prone to competitive interactions while less abundant and more specialized species are often facilitated (Ghazoul 2005, 2006; Bizecki Robson 2013; Zografou et al. 2020; Wei et al. 2021). Overall, these results highlight the dynamic nature of pollinator-mediated interactions (e.g. Rathcke 1983; Feinsinger 1987; Tur et al. 2016), and suggest that their effects can be determined by plant species pollination niches (de Jager et al. 2011; Phillips et al. 2020; Wei et al. 2021). However, the exact mechanisms mediating asymmetrical interactions were not evaluated here and studies that evaluate the factors mediating these asymmetries across plant species pairs are needed in order to fully understand the processes that shape natural plant communities.

Similar to other ecological interactions, the outcomes of pollinator-mediated interactions can vary along species' distribution range, with potential implications on floral diversification (Thompson 1999, 2005; Gómez et al. 2009b; Thompson et al. 2017; Friberg et al. 2019) and community assembly (Briscoe-Runquist et al. 2016; Tur et al. 2016). In spite of this, studies that evaluate plant-plant interactions via pollinators have been typically conducted at a single location (e.g. Ye et al. 2014; Ha and Ivey 2017; Bergamo et al. 2020a, b; Ha et al. 2021), thus neglecting the importance of geographic differences in the outcome of these interactions (but see Sieber et al. 2011). For instance, our results show that *M. guttatus* receives less heterospecific pollen when growing alone, but these effects were not significant at all sites. Similarly, *D. uliginosum* conspecific pollen loads were significantly higher when growing in mixed plots, but only at one site. Interestingly, both of these significant differences were observed at the same site (RHA; Figs 2 and 3), which also had the largest floral density compared to the two other sites (BS: 136.9, RHA: 161.1, TP9: 115.5 flowers per m²). Thus, it is possible that the outcome of pollinator-mediated interactions in these communities is density-dependent (also see Wei et al. 2021). Overall, our results thus suggest that the occurrence of competitive or facilitative interactions, even between the same plant species pair, is highly context-dependent. These differential responses might be explained by differences in plant richness among sites (BS: 62, RHA: 47, TP9: 39 plant species; Arceo-Gómez et al. 2018), differences in pollinator assemblages (Koski et al.

2015) and/or due to changes in specific pollinator preferences among the study sites (Ellis et al. 2021). Geographic variation in the outcome of these interactions can also result from changes in the number and identity of the pollinator assemblage (Gómez et al. 2009a) and plant community composition, as it is also exemplified in studies of diffuse selection (reviewed in Strauss et al. 2005). For instance, a study at the same study sites showed that *M. guttatus* pollen limitation (e.g. pollinator competition) depends strongly on the species richness of the surrounding co-flowering community (Arceo-Gómez and Ashman 2014b). Thus, the outcome of a specific plant-plant interaction at a single location may not reflect how the same plant species interact along their entire distribution range. The differential outcome of these interactions across spatial scales could have important implications for community assembly as they would favour coexistence and co-flowering in some sites while discouraging it in others (e.g. Briscoe-Runquist and Stanton 2013). Spatial variation in the outcome of these interactions can also contribute to generating geographical mosaics of selection (Thompson 1999, 2005), as the relationship between floral traits and fitness would vary spatially with varying intensity of competition and facilitation. For instance, Siebert et al. (2011) reported no evidence for either competition or facilitation at small spatial scales but they did detect facilitation at large scales. To our knowledge, this study is one of few (see also Sieber et al. 2011) to show geographical variation in the outcome of bidirectional pollinator-mediated interactions. Thus, studies that evaluate facilitative and competitive pollinator-mediated interactions across large spatial scales are needed in order to understand their contribution to floral evolution and plant community assembly.

Finally, studies of pollinator-mediated interactions have almost exclusively relied on estimates of flower visitation to infer positive versus negative interactions among plant species (e.g. Ghazoul 2006; Sieber et al. 2011; Ye et al. 2014; van der Kooi et al. 2016) despite that flower visitation alone can be a poor proxy of overall pollination success (e.g. King et al. 2013). In fact, recent studies have shown that flower visitation alone is likely insufficient to capture the full complexity of the pollination and fertilization process that more directly relates to plant reproductive success (Ashman et al. 2020). For example, although heterospecific pollen loads on styles are rather low compared to conspecific pollen loads, which has also been reported for most studied plants (Arceo-Gómez et al. 2019), and differences in heterospecific pollen deposition are not high among the experimental plots, the small loads and differences of heterospecific pollen do not prevent their potential negative effects on plant reproduction. Thus, estimates of conspecific pollen (i.e. pollen limitation) and heterospecific pollen (i.e. pollen interference) receipt, along with estimates of plant fitness (i.e. pollen tube growth, seed production), are necessary in order to more accurately describe the outcomes and underlying mechanisms driving plant-plant interactions via pollinators (e.g. Tur et al. 2016; Ashman et al. 2020). For instance, while we observed a significant increase in *D. uliginosum* conspecific pollen receipt when co-flowering with *M. guttatus* (at one site), this increase did not affect ovule fertilization success (i.e. pollen tube growth). This could be because *D. uliginosum* is not pollen-limited at the study sites and therefore all plants have received enough conspecific pollen to maximize pollen tube production. Similarly, an increase in *M. guttatus* heterospecific pollen receipt in mixed plots compared to single plots (at one site) did not translate to a decrease in *M. guttatus* reproductive success; perhaps due to high *M. guttatus* tolerance to interference by *D. uliginosum* pollen (perhaps due to their relatively large phylogenetic distance; Ashman and Arceo-Gómez 2013; Streher et al. 2020). What is

evident, however, is that in order to accurately describe the outcomes and identify the underlying mechanisms mediating pollinator-mediated interactions, it is necessary to evaluate their effects at multiple stages of the pollination process, i.e. from pollen deposition to ovule fertilization success.

Sources of Funding

This study was funded by a Research Development Committee grant from the East Tennessee State University and a grant from the National Science Foundation-Division Of Environmental Biology (NSF-DEB 1931163) to G.A.-G.

Conflict of Interest

The authors declare that there is no conflict of interest.

Contributions by the Authors

G.A.-G. designed the study. A.S. and G.A.-G. performed the experiment. A.S. and G.A.-G. collected the data. A.S., C.M. and G.A.-G. analysed the data. C.M. and G.A.-G. wrote the manuscript.

Acknowledgements

The authors thank R. Herrmann and J. Daniels for help with data collection in the field and in the lab. C.M. thanks the Max Planck Partner Group for providing him financial support at PUCP.

Data Availability

All data used for analysis in this publication can be found in the following Dryad repository (doi:10.5061/dryad.nzs7h44sm).

Literature Cited

- Alonso C, Herrera CM, Ashman T-L. 2012. A piece of the puzzle: a method for comparing pollination quality and quantity across multiple species and reproductive events. *The New Phytologist* **193**:532–542.
- Anderson B, Johnson SD. 2007. The geographical mosaic of coevolution in a plant–pollinator mutualism. *Evolution* **62**:220–225.
- Arceo-Gómez G, Abdala-Roberts L, Jankowiak A, Kohler C, Meindl GA, Navarro-Fernández CM, Parra-Tabla V, Ashman T-L, Alonso C. 2016. Patterns of among- and within-species variation in heterospecific pollen receipt: the importance of ecological generalization. *American Journal of Botany* **103**:396–407.
- Arceo-Gómez G, Ashman T-L. 2011. Heterospecific pollen deposition: does diversity alter the consequences? *The New Phytologist* **192**:738–746.
- Arceo-Gómez G, Ashman T-L. 2014a. Coflowering community context influences female fitness and alters the adaptive value of flower longevity in *Mimulus guttatus*. *The American Naturalist* **183**:E50–E63.
- Arceo-Gómez G, Ashman T-L. 2014b. Patterns of pollen quantity and quality limitation of pre-zygotic reproduction in *Mimulus guttatus* vary with co-flowering community context. *Oikos* **123**:1261–1269.
- Arceo-Gómez G, Kaczorowski RL, Ashman T-L. 2018. A network approach to understanding patterns of coflowering in diverse communities. *International Journal of Plant Sciences* **179**:569–582.
- Arceo-Gómez G, Schroeder A, Albor C, Ashman T-L, Knight TM, Bennett JM, Suarez B, Parra-Tabla V. 2019. Global geographic patterns of heterospecific pollen receipt help uncover potential ecological and evolutionary impacts across plant communities worldwide. *Scientific Reports* **9**:8086.
- Ashman T-L, Alonso C, Parra-Tabla V, Arceo-Gómez G. 2020. Pollen on stigmas as proxies of pollinator competition and facilitation: complexities, caveats and future directions. *Annals of Botany* **125**:1003–1012.
- Ashman T-L, Arceo-Gómez G. 2013. Toward a predictive understanding of the fitness costs of heterospecific pollen receipt and its importance in co-flowering communities. *American Journal of Botany* **100**:1061–1070.
- Ballantyne G, Baldock KCR, Willmer PG. 2015. Constructing more informative plant–pollinator networks: visitation and pollen deposition networks in a heathland plant community. *Proceedings of the Royal Society B: Biological Sciences* **282**:20151130.
- Bascompte J, Jordano P. 2007. Plant–animal mutualistic networks: the architecture of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**:567–593.
- Bascompte J, Jordano P, Olesen JM. 2006. Asymmetric coevolutionary networks facilitate biodiversity maintenance. *Science* **312**:431–433.
- Bates D, Maechler M, Bolker B, Walker S, Bojesen Christensen RH, Singmann H, Dai B, Scheipl F, Grothendieck G, Green P, Fox J. 2019. lme4: linear mixed-effects models using ‘eigen’ and s4. R package version 1.1-21. <https://cran.r-project.org/package=lme4/> (10 February 2021).
- Benevides CR, Evans DM, Gaglianone MC. 2013. Comparing the structure and robustness of Passifloraceae—floral visitor and true pollinator networks in a lowland Atlantic forest. *Sociobiology* **60**:295–305.
- Bergamo PJ, Streher NS, Wolowski M, Sazima M. 2020b. Pollinator-mediated facilitation is associated with floral abundance, trait similarity and enhanced community-level fitness. *Journal of Ecology* **108**:1334–1346.
- Bergamo PJ, Susin Streher N, Traveset A, Wolowski M, Sazima M. 2020a. Pollination outcomes reveal negative density-dependence coupled with interspecific facilitation among plants. *Ecology Letters* **23**:129–139.
- Bergamo PJ, Wolowski M, Maruyama PK, Vizentin-Bugoni J, Sazima M. 2018. Trait patterns across space and time suggest an interplay of facilitation and competition acting on Neotropical hummingbird-pollinated plant communities. *Oikos* **127**:1690–1700.
- Bizecki Robson D. 2013. An assessment of the potential for pollination facilitation of a rare plant by common plants: *Symphyotrichum sericeum* (Asteraceae) as a case study. *Botany* **91**:34–42.
- Briscoe Runquist RD. 2012. Pollinator-mediated competition between two congeners, *Limnanthes douglasii* subsp. *rosea* and *L. alba* (Limnanthaceae). *American Journal of Botany* **99**:1125–1132.
- Brooker RW. 2006. Plant–plant interactions and environmental change. *The New Phytologist* **171**:271–284.
- Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G, Liancourt P, Tielbörger K, Travis JMJ, Anthelme F, Armas C, Coll L, Corcket E, Delzon S, Forey E, Kikvidze Z, Olofsson J, Pugnaire F, Quiroz CL, Saccone P, Schiffrers K, Seifan M, Touzard B, Michalet R. 2008. Facilitation in plant communities: the past, the present, and the future. *Journal of Ecology* **96**:18–34.
- Brown BJ, Mitchell RJ, Graham SA. 2002. Competition for pollination between an invasive species (purple loosestrife) and a native congener. *Ecology* **83**:2328–2336.
- Briscoe Runquist R, Grossenbacher D, Porter S, Kay K, Smith J. 2016. Pollinator-mediated assemblage processes in California wildflowers. *Journal of Evolutionary Biology* **29**:1045–1058.
- Burkle LA, Myers JA, Belote RT. 2016. The beta-diversity of species interactions: untangling the drivers of geographic variation in plant–pollinator diversity and function across scales. *American Journal of Botany* **103**:118–128.
- Callaway RM, Brooker RW, Choler P, Kikvidze Z, Lortie CJ, Michalet R, Paolini L, Pugnaire FI, Newingham B, Aschehoug ET, Armas C, Kikvidze D, Cook BJ. 2002. Positive interactions among alpine plants increase with stress. *Nature* **417**:844–848.
- Campbell D. 1985. Pollen and gene dispersal: the influences of competition for pollination. *Evolution* **39**:418–431.
- Caruso CM, Alfaro M. 2000. Interspecific pollen transfer as a mechanism of competition: effect of *Castilleja linariaefolia* pollen on seed set of *Ipomopsis aggregata*. *Canadian Journal of Botany* **78**:600–606.
- Carvalho LG, Biesmeijer JC, Benadi G, Fründ J, Stang M, Bartomeus I, Kaiser-Bunbury CN, Baude M, Gomes SI, Merckx V, Baldock KC, Bennett AT, Boada R, Bommarco R, Cartar R, Chacoff N, Dänhardt J, Dicks LV, Dormann CF, Ekroos J, Henson KS, Holzschuh A, Junker RR, Lopezaraiza-Mikel M, Memmott J, Montero-Castaño A, Nelson IL, Petanidou T, Power EF, Rundlöf M, Smith HG, Stout JC, Temitope K, Tscharntke T, Tscheulin T, Vilà M, Kunin WE. 2014. The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. *Ecology Letters* **17**:1389–1399.

- Choler P, Michalet R, Callaway RM. 2001. Facilitation and competition on gradients in alpine plant communities. *Ecology* 82:3295–3308.
- Dafni A. 1992. *Pollination ecology: a practical approach*. Oxford: Oxford University Press.
- de Jager ML, Dreyer LL, Ellis AG. 2011. Do pollinators influence the assembly of flower colours within plant communities? *Oecologia* 166:543–553.
- Ellis AG, Anderson B, Kemp JE. 2021. Geographic mosaics of fly pollinators with divergent color preferences drive landscape-scale structuring of flower color in daisy communities. *Frontiers in Plant Science* 12:617761.
- Elo M, Kareksela S, Haapalehto T, Vuori H, Aapala K, Kotiaho JS. 2016. The mechanistic basis of changes in community assembly in relation to anthropogenic disturbance and productivity. *Ecosphere* 7:e01310.
- Feinsinger P. 1987. Effects of plant species on each other's pollination: is community structure influenced? *Trends in Ecology & Evolution* 2:123–126.
- Feldman TS, Morris WF, Wilson WF. 2004. When can two plant species facilitate each other's pollination? *Oikos* 105:197–207.
- Flanagan RJ, Mitchell RJ, Knutowski D, Karron JD. 2009. Interspecific pollinator movements reduce pollen deposition and seed production in *Mimulus ringens* (Phrymaceae). *American Journal of Botany* 96:809–815.
- Freestone AL, Harrison S. 2006. Regional enrichment of local assemblages is robust to variation in local productivity, abiotic gradients, and heterogeneity. *Ecology Letters* 9:95–102.
- Friberg M, Schwind C, Guimarães PR Jr, Raguso RA, Thompson JN. 2019. Extreme diversification of floral volatiles within and among species of *Lithophragma* (Saxifragaceae). *Proceedings of the National Academy of Sciences of the United States of America* 116:4406–4415.
- Galen C. 1999. Why do flowers vary? The functional ecology of variation in flower size and form within natural plant populations. *Bioscience* 49:631–640.
- Ghazoul J. 2005. Pollen and seed dispersal among dispersed plants. *Biological Reviews of the Cambridge Philosophical Society* 80:413–443.
- Ghazoul J. 2006. Floral diversity and the facilitation of pollination. *Journal of Ecology* 94:295–304.
- Gómez JM, Abdelaziz M, Camacho JP, Muñoz-Pajares AJ, Perfectti F. 2009b. Local adaptation and maladaptation to pollinators in a generalist geographic mosaic. *Ecology Letters* 12:672–682.
- Gómez JM, Perfectti F, Bosch J, Camacho JPM. 2009a. A geographic selection mosaic in a generalized plant–pollinator–herbivore system. *Ecological Monographs* 79:245–263.
- Ha MK, Ivey CT. 2017. Pollinator-mediated interactions in experimental arrays vary with neighbor identity. *American Journal of Botany* 104:252–260.
- Ha MK, Schneider SA, Adler LS. 2021. Facilitative pollinator sharing decreases with floral similarity in multiple systems. *Oecologia* 195:273–286.
- Jordano P. 1987. Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence asymmetries, and coevolution. *The American Naturalist* 129:657–677.
- Kay KM, Sargent SD. 2009. The role of animal pollination in plant speciation: integrating ecology, geography, and genetics. *Annual Review of Ecology, Evolution, and Systematics* 40:637–656.
- King C, Ballantyne G, Willmer PG. 2013. Why flower visitation is a poor proxy for pollination: measuring single-visit pollen deposition, with implications for pollination networks and conservation. *Methods in Ecology and Evolution* 4:811–818.
- Koontz JA, Warnock MJ. 2012. *Delphinium uliginosum*. In: Jepson Flora Project, eds. *Jepson eFlora*. <https://ucjeps.berkeley.edu/eflora/> (20 October 2020).
- Koski MH, Meindl GA, Arceo-Gómez G, Wolowski M, LeCroy KA, Ashman T-L. 2015. Plant-flower visitor networks in a serpentine metacommunity: assessing traits associated with keystone plant species. *Arthropod-Plant Interactions* 9:9–21.
- Kraft NJB, Ackerly DD. 2014. The assembly of plant communities. In: Monson RK, ed. *Ecology and the environment*. New York: Springer, 67–88.
- Lázaro A, Lundgren R, Totland Ø. 2009. Co-flowering neighbors influence the diversity and identity of pollinator groups visiting plant species. *Oikos* 118:691–702.
- Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2020. emmeans: estimated marginal means, aka least-squares means. R package version 1.4.8. <http://cran.r-project.org/package=emmeans/> (28 February 2021).
- Levine JM. 1999. Indirect facilitation: evidence and predictions from a riparian community. *Ecology* 80:1762–1769.
- Losapio G, Schöb C. 2020. Pollination interactions reveal direct costs and indirect benefits of plant–plant facilitation for ecosystem engineers. *Journal of Plant Ecology* 13:107–113.
- Martén-Rodríguez S, Kress WJ, Temeles EJ, Meléndez-Ackerman E. 2011. Plant–pollinator interactions and floral convergence in two species of *Heliconia* from the Caribbean Islands. *Oecologia* 167:1075–1083.
- McEwen JR, Vamosi JC. 2010. Floral colour versus phylogeny in structuring subalpine flowering communities. *Proceedings of the Royal Society B: Biological Sciences* 277:2957–2965.
- Mitchell RJ, Flanagan RJ, Brown BJ, Waser NM, Karron JD. 2009. New frontiers in competition for pollination. *Annals of Botany* 103:1403–1413.
- Moeller DA. 2005. Pollinator community structure and sources of spatial variation in plant–pollinator interactions in *Clarkia xantiana* ssp. *xantiana*. *Oecologia* 142:28–37.
- Molofsky J, Fisher BL. 1993. Habitat and predation effects on seedling survival and growth in shade-tolerant tropical trees. *Ecology* 74:261–265.
- Morales CL, Traveset A. 2008. Interspecific pollen transfer: magnitude, prevalence and consequences for plant fitness. *Critical Reviews in Plant Sciences* 27:221–238.
- Muchhala N, Brown Z, Armbruster WS, Potts MD. 2010. Competition drives specialization in pollination systems through costs to male fitness. *The American Naturalist* 176:732–743.
- Muchhala N, Caiza A, Vizuete JC, Thomson JD. 2009. A generalized pollination system in the tropics: bats, birds and *Aphelandra acanthus*. *Annals of Botany* 103:1481–1487.
- Navarro-Cano JA, Goberna M, Valiente-Banuet A, Verdú M. 2021. Phenotypic structure of plant facilitation networks. *Ecology Letters* 24:509–519.
- Olesen JM, Jordano P. 2002. Geographic patterns in plant–pollinator mutualistic networks. *Ecology* 83:2416–2424.
- Phillips RD, Peakall R, van der Niet T, Johnson SD. 2020. Niche perspectives on plant–pollinator interactions. *Trends in Plant Science* 25:779–793.
- Rahmé J, Widmer A, Karrenberg S. 2009. Pollen competition as an asymmetric reproductive barrier between two closely related *Silene* species. *Journal of Evolutionary Biology* 22:1937–1943.
- Randall JL, Hilu KW. 1990. Interference through improper pollen transfer in mixed stands of *Impatiens capensis* and *I. pallida* (Balsaminaceae). *American Journal of Botany* 77:939–944.
- Rathcke B. 1983. Competition and facilitation among plants for pollination. In: Real L, ed. *Pollination biology*. Orlando, FL: Academic Press, 305–329.
- R Development Core Team. 2017. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/> (15 March 2020).
- Ripley B, Venables B, Bates DM, Hornik K, Gebhardt A, Firth D. 2020. MASS: support functions and datasets for Venables and Ripley's MASS. R package version 7.3-51.6. <http://cran.r-project.org/package=MASS/> (20 February 2021).
- Robertson AW, Diaz A, Macnair MR. 1994. The quantitative genetics of floral characters in *Mimulus guttatus*. *Heredity* 72:300–311.
- Rosenthal R, Rosnow RL, Rubin DB. 2000. *Contrasts and effect sizes in behavioral research*. Cambridge: Cambridge University Press.
- Runquist RB, Stanton ML. 2013. Asymmetric and frequency-dependent pollinator-mediated interactions may influence competitive displacement in two vernal pool plants. *Ecology Letters* 16:183–190.
- Sargent RD, Ackerly DD. 2008. Plant–pollinator interactions and the assembly of plant communities. *Trends in Ecology & Evolution* 23:123–130.
- Sieber Y, Holderegger R, Waser NM, Thomas VFD, Braun S, Erhardt A, Reyer H-U, Wirth LR. 2011. Do alpine plants facilitate each other's pollination? Experiments at a small spatial scale. *Acta Oecologica* 37:369–374.
- Stavert JR, Liñán-Cembrano G, Beggs JR, Howlett BG, Pattemore DE, Bartomeus I. 2016. Hairiness: the missing link between pollinators and pollination. *PeerJ* 4:e2779.
- Strauss SY, Murch P. 2004. Towards an understanding of the mechanisms of tolerance: compensating for herbivore damage by enhancing a mutualism. *Ecological Entomology* 29:234–239.
- Strauss SY, Sahl H, Conner JK. 2005. Toward a more trait-centered approach to diffuse (co)evolution. *The New Phytologist* 165:81–89.
- Streher NS, Bergamo PJ, Ashman T-L, Wolowski M, Sazima M. 2020. Effect of heterospecific pollen deposition on pollen tube growth depends on the phylogenetic relatedness between donor and recipient. *AoB Plants* 12:plaa016.

- Suárez-Mariño A, Arceo-Gómez G, Sosenski P, Parra-Tabla V. 2019. Patterns and effects of heterospecific pollen transfer between an invasive and two native plant species: the importance of pollen arrival time to the stigma. *American Journal of Botany* **106**:1308–1315.
- Thompson JN. 1999. Specific hypotheses on the geographic mosaic of coevolution. *The American Naturalist* **153**:S1–S14.
- Thompson JN. 2005. *The geographic mosaic of coevolution*. Chicago, IL: The University of Chicago Press.
- Thompson JN, Schwind C, Friberg M. 2017. Diversification of trait combinations in coevolving plant and insect lineages. *The American Naturalist* **190**:171–184.
- Thomson JD, Fung HF, Ogilvie JE. 2019. Effects of spatial patterning of co-flowering plant species on pollination quantity and purity. *Annals of Botany* **123**:303–310.
- Tur C, Sáez A, Traveset A, Aizen MA. 2016. Evaluating the effects of pollinator-mediated interactions using pollen transfer networks: evidence of widespread facilitation in south Andean plant communities. *Ecology Letters* **19**:576–586.
- van der Kooij CJ, Pen I, Staal M, Stavenga DG, Elzenga JTM. 2016. Competition for pollinators and intra-communal spectral dissimilarity of flowers. *Plant Biology* **18**:56–62.
- Vázquez DP, Aizén MA. 2004. Asymmetric specialization: a pervasive feature of plant–pollinator interactions. *Ecology* **85**:1251–1257.
- Vilà M, Bartomeus I, Dietzsch AC, Petanidou T, Steffan-Dewenter I, Stout JC, Tscheulin T. 2009. Invasive plant integration into native plant–pollinator networks across Europe. *Proceedings of the Royal Society B: Biological Sciences* **276**:3887–3893.
- Waser NM, Jones CE, Little RJ. 1983. Competition for pollination and floral character differences among sympatric plant species: a review of evidence. In: Jones CE, Little RJ, eds. *Handbook of experimental pollination biology*. New York: Van Nostrand Reinhold, 277–293.
- Wassink E, Caruso CM. 2013. Effect of coflowering *Mimulus ringens* on phenotypic selection on floral traits of gynodioecious *Lobelia siphilitica*. *Botany* **91**:745–751.
- Wei N, Kaczorowski RL, Arceo-Gómez G, O'Neill EM, Hayes RA, Ashman T-L. 2021. Pollinators contribute to the maintenance of flowering plant diversity. *Nature* **597**:688–692.
- Westoby M, Wright IJ. 2006. Land-plant ecology on the basis of functional traits. *Trends in Ecology & Evolution* **21**:261–268.
- Wolowski M, Carvalheiro LG, Freitas L. 2017. Influence of plant–pollinator interactions on the assembly of plant and hummingbird communities. *Journal of Ecology* **105**:332–344.
- Xu J, Chai Y, Wang M, Dang H, Guo Y, Chen Y, Zhang C, Li T, Zhang L, Yue M. 2018. Shifts in plant community assembly processes across growth forms along a habitat severity gradient: a test of the plant functional trait approach. *Frontiers in Plant Science* **9**:180.
- Ye Z-M, Dai W-K, Jin X-F, Gituru RW, Wang Q-F, Yang C-F. 2014. Competition and facilitation among plants for pollination: can pollinator abundance shift the plant–plant interactions? *Plant Ecology* **215**:3–13.
- Zografou K, Swartz MT, Tilden VP, McKinney EN, Eckenrode JA, Sewall BJ. 2020. Stable generalist species anchor a dynamic pollination network. *Ecosphere* **11**:e03225.