

Metalloradical Activation of In Situ-Generated α -Alkynyldiazomethanes for Asymmetric Radical Cyclopropanation of Alkenes

Jing Ke, Wan-Chen Cindy Lee, Xiaoxu Wang, Yong Wang, Xin Wen, and X. Peter Zhang*

Cite This: *J. Am. Chem. Soc.* 2022, 144, 2368–2378

Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: α -Alkynyldiazomethanes, generated in situ from the corresponding sulfonyl hydrazones in the presence of a base, can serve as effective metalloradicalophiles in Co(II)-based metalloradical catalysis (MRC) for asymmetric cyclopropanation of alkenes. With D_2 -symmetric chiral amidoporphyrin 2,6-DiMeO-QingPhyrin as the optimal supporting ligand, the Co(II)-based metalloradical system can efficiently activate different α -alkynyldiazomethanes at room temperature for highly asymmetric cyclopropanation of a broad range of alkenes. This catalytic radical process provides a general synthetic tool for stereoselective construction of alkynyl cyclopropanes in high yields with high both diastereoselectivity and enantioselectivity. Combined computational and experimental studies offer several lines of evidence in support of the underlying stepwise radical mechanism for the Co(II)-catalyzed olefin cyclopropanation involving a unique α -metalloradical intermediate that is associated with two resonance forms of α -Co(III)-propargyl radical and γ -Co(III)-allenyl radical. The resulting enantioenriched alkynyl cyclopropanes, as showcased with several stereospecific transformations, may serve as valuable chiral building blocks for stereoselective organic synthesis.




INTRODUCTION


Radical reactions have been increasingly exploited as complementary synthetic methods to ionic reactions in modern organic synthesis as they enjoy intrinsic attributes such as high reactivity and tolerance of functional groups.¹ Among significant challenges facing this endeavor is the control of reactivity as well as selectivities in the reactions of free organic radicals, especially enantioselectivity.² Rising to these challenges, metalloradical catalysis (MRC) has emerged as a conceptually new approach to the development of stereoselective radical reactions through catalytic generation of metal-supported organic radicals as key catalytic intermediates.^{3–5} As stable 15e-metalloradicals with well-defined low-spin d^7 -configuration, Co(II) complexes of D_2 -symmetric chiral amidoporphyrins [Co(D_2 -Por*)] have been recognized as a family of open-shell catalysts that are effective for asymmetric cyclopropanation of alkenes with diazo compounds.⁶ During the Co(II)-catalyzed cyclopropanation, the initially generated α -metalloalkyl radical intermediates from metalloradical activation of diazo compounds, which are centrally situated inside the pocketlike environment of the chiral porphyrin ligands,⁷ can be precisely governed to perform a sequence of homolytic reactions such as radical addition and radical substitution with olefin substrates, leading to productive formation of cyclopropanes with effective control of diastereoselectivity and enantioselectivity. Except for a few recent examples of using α -aryldiazomethanes,⁸ Co(II)-based radical cyclopropanation has so far been mostly involved with the use

of acceptor- and acceptor/acceptor-substituted diazo compounds as metalloradicalophiles.⁶ In all the previous cases, the key α -Co(III)-supported C-centered radical intermediates are stabilized by $C(sp^2)$ -based carbonyl or aryl substituents through potential H-bonding interactions with the amide units of the catalyst, facilitating the reactivity and stereoselectivity of the catalytic radical process. It was unclear if Co(II)-based metalloradical system could also be applicable to other types of diazo compounds with substituents beyond $C(sp^2)$ -based carbonyl and aryl groups. Specifically, we were intrigued to learn whether α -alkynyldiazomethanes, a common type of diazo compounds containing $C(sp)$ -based alkynyl substituents, could be employed as potential metalloradicalophiles by Co(II)-based metalloradical system for radical olefin cyclopropanation (Scheme 1). As α -alkynyldiazomethanes **1'** are typically generated in situ from corresponding hydrazones **1** under basic conditions,⁹ it would be crucially important to match the rates between the diazomethane generation and the ensuing metalloradical activation in order to preclude their thermal decomposition to form pyrazole and azine side products. Apart from the concern with the effectiveness for

Received: December 14, 2021

Published: January 31, 2022




Scheme 1. Proposed Pathway for Radical Cyclopropanation of Alkenes with α -Alkynyldiazomethanes via Co(II)-MRC

metalloradical activation of α -alkynyldiazomethanes **1'**, the additional issue of regioselectivity could arise from two potential reactivity modes of the resulting α -metalloalkyl radical intermediates that are associated with the two resonance forms of α -Co(III)-propargyl radicals **I** and γ -Co(III)-allenyl radicals **I'**.¹⁰ Furthermore, in the absence of the aforementioned H-bonding interactions, what elements could be utilized during the subsequent radical addition of the

initially generated α -Co(III)-propargyl radicals **I** to the olefin substrates to control the enantioselectivity for the first C–C bond formation? What also under question is the control of chemoselectivity as well as the diastereoselectivity during the final step of radical substitution of γ -Co(III)-alkyl radicals **II** for the second C–C bond formation. Considering the presence of the C $\equiv$ C triple bond, could γ -Co(III)-alkyl radicals **II** undergo the desired 3-*exo-tet* radical cyclization chemoselectively over the potentially competitive 4-*exo-dig* and 5-*endo-dig* cyclization, leading to diastereoselective construction of cyclopropanes? We hoped to address these and related issues through judicious development of Co(II)-based metalloradical catalyst by fine-tuning the environments of D_2 -symmetric chiral amidoporphyrin ligand. If implemented successfully, then it would lead to the development of a new catalytic radical process for asymmetric olefin cyclopropanation with in situ-generated α -alkynyldiazomethanes to construct chiral alkynyl cyclopropanes. In addition to serving as useful intermediates for stereoselective organic synthesis, chiral alkynyl cyclopropanes occur as common substructures in many important natural products and drug molecules (see Figure S1).¹¹

Metal-catalyzed asymmetric cyclopropanation of alkenes with α -alkynyldiazomethanes offers a potentially general approach for stereoselective construction of valuable chiral alkynyl cyclopropanes.^{12–14} While donor/acceptor-substituted diazo compounds alkynyldiazoacetates were successfully

Scheme 2. Ligand Effect on Co(II)-Based Catalytic System for Olefin Cyclopropanation with Alkynyldiazomethane^a

^aCarried out with **1a** (0.10 mmol) and **2a** (0.20 mmol) in the presence of KH (0.40 mmol) by [Co(Por)] (2 mol %) in ethyl acetate (0.6 mL) at 22 °C for 24 h; Tris = 2,4,6-triisopropylbenzenesulfonyl; isolated yields; diastereomeric excess (de) determined by ¹H NMR; enantiomeric excess (ee) of *trans*-isomer determined by chiral HPLC.

Table 1. Scope of Catalytic Asymmetric Cyclopropanation of Alkenes with α -Alkynyldiazomethanes by [Co(P6)]^a

Entry	Alkynyl Diazomethane (1)	Alkene (2)	Yield (%)	dr	ee (%)
1	(-)-3b	styrene	90	79:21	97
2	(-)-3c	styrene	90	96:4	99
3	(-)-3d	styrene	90	85:15	98
4	(-)-3e	styrene	94	90:10	99
5	(-)-3f	styrene	99	94:6	95
6	(-)-3g	styrene	94	90:10	90
7	(-)-3h	styrene	90	93:7	98
8	(-)-3i	styrene	95	80:20	90
9	(-)-3j	styrene	41	91:9	85
10	(-)-3k	styrene	71	96:4	87
11	(-)-3l	styrene	94	91:9	89
12	(-)-3m	styrene	89	90:10	94
13	(-)-3n	styrene	85	81:19	95
14	(-)-3o	styrene	88	78:22	81
15	(-)-3p	styrene	96	85:15	95
16	(-)-3q	styrene	91	90:10	94
17	(-)-3r	styrene	88	88:12	94
18	(-)-3s	styrene	93	84:16	91
19	(-)-3t	styrene	96	92:8	99
20	(-)-3u	styrene	97	96:4	92
21	(-)-3v	styrene	92	90:10	92
22	(-)-3w	styrene	90	88:12	93
23	(-)-3x	styrene	92	98:2	89
24	(-)-3y	styrene	89	96:4	88
25	(-)-3z	styrene	93	96:4	94
26	(-)-3aa ^b	styrene	91	97:3	91
27	(-)-3ab	styrene	79	85:15	93
28	(-)-3ac	styrene	93	97:3	93
29	(-)-3ad	styrene	93	91:9	95
30	(-)-3ae ^c	styrene	32	86:14	81
31	(-)-3af ^c	styrene	40	89:11	85

R = Et, (-)-3ae^c; R = Ph, (-)-3af^c

^aCarried out with **1** (0.10 mmol) and **2** (0.20 mmol) in the presence of KH (0.40 mmol) by [Co(P6)] (2 mol %) in ethyl acetate (0.6 mL) at 22 °C for 24 h; Tris = 2,4,6-triisopropylbenzenesulfonyl; isolated yields; diastereomeric ratio (dr) determined by ¹H NMR analysis of reaction mixture; enantiomeric excess (ee) of *trans*-isomer determined by chiral HPLC. ^bAbsolute configuration determined by X-ray crystallography. ^cCarried out with **2** (1.00 mmol) at 80 °C.

employed by Davies and co-workers as carbene precursors for Rh₂-catalyzed asymmetric cyclopropanation,¹⁵ donor-substituted diazo compounds α -alkynyldiazomethanes, in contrast, have been scarcely explored.¹⁶ A single example of cyclopropanation of *trans*-stilbene with the use of α -alkynyldiazomethane could be found in a recent report by Bi and co-workers on Ag-based catalytic system involving in situ generation of diazo compounds from *N*-nosylhydrazones.^{16c} This underdevelopment is mainly attributed to the inherent propensity of α -alkynyldiazomethanes toward the formation of the stable aromatic pyrazoles.¹⁷ To circumvent the problem, Echavarren and co-workers recently reported an innovative two-step approach for synthesis of alkynyl cyclopropanes from alkenes based on Rh₂-catalyzed decarbenation of 7-alkynyl cycloheptatrienes as alkynylcarbene precursors.¹⁸ To the best of our knowledge, there has been no previous report on catalytic system for asymmetric olefin cyclopropanation with α -alkynyldiazomethanes for stereoselective synthesis of chiral alkynyl cyclopropanes. As a new application of Co(II)-based metalloradical catalysis (MRC), we herein report the development of the first asymmetric catalytic system for olefin cyclopropanation that can effectively utilize α -alkynyldiazomethanes through in situ generation from the corresponding sulfonyl hydrazones in the presence of base. Through identification of an optimal D₂-symmetric chiral amidoporphyrin (D₂-Por*) as the supporting ligand, the Co(II)-based

catalytic system is capable of efficiently activating different alkynyldiazomethanes at room temperature to cyclopropanate a wide range of alkenes with varied electronic and steric properties, delivering alkynyl cyclopropanes in high yields with excellent control of both diastereoselectivity and enantioselectivity. We show the importance of catalyst development through fine-tuning of the ligand environment in achieving high reactivity as well as stereoselectivities. We present combined computational and experimental studies that shed light on the underlying stepwise radical mechanism of the Co(II)-catalyzed cyclopropanation. To demonstrate the synthetic applications of the new catalytic system, we showcase several examples of stereoselective transformations of the resulting enantioenriched alkynyl cyclopropanes.

RESULTS AND DISCUSSION




Catalyst Development. At the outset of the project, α -(phenylethynyl)diazomethane (**1a'**), which was in situ generated from corresponding trisylhydrazone **1a** in the presence of KH, was designated as the representative α -alkynyldiazomethane for asymmetric radical cyclopropanation of styrene (**2a**) using Co(II)-based metalloradical catalysts [Co(Por)] (Scheme 2; see Table S1). It was found that simple achiral metalloradical catalyst [Co(P1)] (P1 = tetraphenylporphyrin) could activate the diazo compound to cyclopropanate styrene, giving desired alkynyl cyclopropane **3a** in low yield

(36%) with inferior control of diastereoselectivity (14% de). With the employment of the Co(II) complex of D_{2h} -symmetric achiral amidoporphyrin [Co(P2)] (P2 = 3,5-Di^tBu-IbuPhyrin) as the catalyst,¹⁹ the catalytic cyclopropanation reaction was significantly enhanced, forming cyclopropane 3a in a higher yield (84%) with improved diastereoselectivity (56% de) in favor of the *trans*-isomer. To evaluate the feasibility of asymmetric induction during the catalytic process, Co(II) complexes of D_2 -symmetric chiral amidoporphyrins were employed as the catalysts. When first-generation chiral metalloradical catalyst [Co(P3)] (P3 = 3,5-Di^tBu-ChenPhyrin)^{6a} was used, the catalytic reaction produced cyclopropane 3a in an almost quantitative yield (98%) with the same level of diastereoselectivity (58% de) while exhibiting significant enantioselectivity (53% ee). To improve the stereoselectivities of the catalytic process, we then applied metalloradical catalyst [Co(P4)] (P4 = 3,5-Di^tBu-ZhuPhyrin),²⁰ which is devised to achieve conformational rigidity through unique intramolecular H-bonding interactions in (S)-(-)-2-tetrahydrofurancarboxamide units. As expected, the catalytic reaction by [Co(P4)] brought about improved enantioselectivity (59% ee) but reduced diastereoselectivity (52% de) and significantly decreased yield (34%) as a result of the more sterically encumbered ligand environment. These results prompted us to explore second-generation chiral metalloradical catalysts bearing cyclopropanecarboxamides with two contiguous stereogenic centers in the hope of further enhancing the asymmetric induction of the catalytic process. Gratifyingly, when [Co(P5)] (P5 = 3,5-Di^tBu-QingPhyrin) was used as the catalyst for the reaction,^{6g} cyclopropane 3a was produced in high yield (80%) with excellent enantioselectivity (94% ee) albeit with low diastereoselectivity (20% de). Inspired by this positive outcome, we then employed the analogous catalyst [Co(P6)] (P6 = 2,6-DiMeO-QingPhyrin) bearing the same chiral amide units but with different achiral *meso*-aryl groups to further enhance the rigidity of the chiral environment. To our delight, the catalytic reaction by [Co(P6)] indeed significantly improved both diastereoselectivity (74% de) and yield (90%) while maintaining the excellent enantioselectivity (96% ee). The remarkable difference in performance among the metalloradical catalysts indicates that subtle alteration in ligand environment can give rise to significant improvement in catalytic reactivity as well as stereoselectivities, manifesting the effectiveness of catalyst development in controlling the radical process.

Substrate Scope. Under the optimized conditions, the scope and versatility of [Co(P6)]-catalyzed asymmetric cyclopropanation were evaluated using different α -alkynyldiazomethanes through in situ generation from the corresponding sulfonyl hydrazones 1 under the basic condition with various olefins (Table 1). Similar to α -alkynyldiazomethane 1a', α -(arylethynyl)diazomethanes bearing substituents with varied electronic and steric properties on the phenyl group could be effectively employed as metalloradicalophiles for the Co(II)-based catalytic system as shown for asymmetric cyclopropanation of styrene (2a) as the representative olefin substrate. For example, α -(arylethynyl)diazomethanes bearing alkyl substituents at different aryl positions such as *p*-Me and *o*-iPr could be efficiently activated by [Co(P6)] at room temperature to cyclopropanate styrene, producing desired alkynyl cyclopropanes 3b and 3c in high yields with good diastereoselectivities and excellent enantioselectivities (entries 1 and 2). Furthermore, [Co(P6)] could activate α -

(arylethynyl)diazomethanes containing an electron-donating -OMe group at all the three different positions of the aryl group for the cyclopropanation reaction, allowing stereoselective construction of corresponding alkynyl cyclopropanes 3d–3f in high yields with high diastereoselectivities and excellent enantioselectivities (entries 3–5). It is interesting to note that the observed diastereoselectivity increases when the position of MeO-substituent in cyclopropanes 3d–3f moves from *para*- to the *meta*- to the *ortho*-position. Likewise, α -(arylethynyl)diazomethanes bearing electron-withdrawing groups could also be suitable metalloradicalophiles as exemplified by asymmetric synthesis of alkynyl cyclopropanes 3g and 3h in high yields with excellent control of stereoselectivities (entries 6 and 7). Moreover, naphthalene-containing α -alkynyldiazomethanes could also be effectively activated by [Co(P6)] under the same conditions for asymmetric cyclopropanation reaction, leading to high-yielding formation of corresponding alkynyl cyclopropane 3i with both high diastereoselectivity and enantioselectivity (entry 8). In addition to α -(arylethynyl)diazomethanes, the Co(II)-based metalloradical system was shown to be applicable to α -alkynyldiazomethanes containing certain nonaryl substituents as demonstrated by catalytic asymmetric cyclopropanation of styrene with alkyl substituted diazo derivative, generating desired cyclopropane 3j with high stereoselectivities albeit in moderate yield (entry 9). Moreover, a triisopropylsilyl (TIPS)-substituted diazo derivative can also serve as suitable substrate, furnishing the corresponding cyclopropane 3k in good yield with high diastereoselectivity and enantioselectivity (entry 10).

In addition to being capable of using different α -alkynyldiazomethanes, the Co(II)-based system for asymmetric cyclopropanation was found to be suitable to a wide range of alkenes under the optimized conditions (Table 1). Like styrene, its derivatives bearing various substituents such as -*tert*-Bu, -OMe, -CN, and -NO₂ groups at different positions could be reliably cyclopropanated by [Co(P6)] with α -alkynyldiazomethane 1a', generating corresponding alkynyl cyclopropanes 3l–3o in similarly high yields with effective control of diastereoselectivities and enantioselectivities (entries 11–14). Additionally, halogenated aromatic olefins could also serve as suitable substrates in [Co(P6)]-catalyzed asymmetric cyclopropanation as exemplified by highly stereoselective generation of alkynyl cyclopropanes 3p–3s with halogen atoms at various positions (entries 15–18). Furthermore, extended aromatic olefins like 2-vinylnaphthalene could be also effectively applied to the catalytic system, affording alkynyl cyclopropane 3t in excellent yield with high diastereoselectivity and outstanding enantioselectivity (entry 19). Moreover, the Co(II)-based cyclopropanation was shown to be compatible with heteroaryl alkenes such as those containing thiophene, pyridine, and benzothiophene, affording corresponding heteroaryl cyclopropanes 3u–3w in high yields with high stereoselectivities (entries 20–22). To further highlight the unique feature of Co(II)-based metalloradical catalysis, even electron-deficient olefins such as acrylamide and ethyl acrylate, which are challenging substrates for catalytic cyclopropanation systems involving electrophilic metallocarbene intermediates, could be effectively cyclopropanated by [Co(P6)] to form desired products 3x and 3y in high yields with high stereoselectivities (entries 23 and 24). In addition to monosubstituted olefins, 1,1-disubstituted olefins like α -methylstyrene and α -bromostyrene could serve as suitable substrates as well, allowing for highly stereoselective

Scheme 3. Mechanistic Studies on Co(II)-Catalyzed Olefin Cyclopropanation with α -AlkynyldiazomethanesA. DFT Calculations on Catalytic Pathways and Associated Energetics of Co(II)-Catalyzed Olefin Cyclopropanation^aB. Trapping of α -Co(III)-Propargyl Radical and γ -Co(III)-Allenyl Radical Intermediates by Spin Trap PBN^bC. Trapping of α -Co(III)-Propargyl Radical Intermediates by TEMPO and Characterizations of α,α -Bis-TEMPO-Trapped Products^c







^aApplied bp86/LANL2DZ for geometry optimization and B3LYP/def2-tzvp for calculations of single-point energies (kcal/mol) along with Grimme's dispersion correction and SMD (ethyl acetate) solvation model. ^bCarried out with **1f** (0.10 mmol), [Co(P1)] (2 mol %) and PBN (0.12 mmol) in the presence of Et₃N (0.20 mmol) at 60 °C in benzene (1.0 mL) for 10 min; Ar = 2-methoxyphenyl. The simulation of the EPR spectrum was carried out by iteration of the isotropic *g*-values and line widths using the EPR simulation program SpinFit Xenon. ^cCarried out with **1** (0.10 mmol) and TEMPO (0.60 mmol) in the presence of KH (0.40 mmol) by [Co(P1)] (2 mol %) in ethyl acetate (0.6 mL) at 22 °C for 24 h; Tris = 2,4,6-triisopropylbenzene sulfonyl; isolated yield; structure determined by X-ray crystallography.

construction of trisubstituted cyclopropanes **3z** and **3aa** with excellent control of the newly generated quaternary carbon stereogenic centers (entries 25 and 26). The absolute configuration of the major enantiomer of alkynyl cyclopropane **3aa** was established as (*S,S*) by X-ray crystallography. When conjugated dienes and enynes were used as the substrates, [Co(P6)] could regio- and chemoselectively cyclopropanate the terminal double bonds, leading to exclusive formation of cyclopropanes **3ab–3ad** in high yields with high stereoselectivities without affecting the internal alkene and alkyne units (entries 27–29). It is interesting to note the C₂-symmetric structure of bisalkynyl cyclopropane **3ad**. Notably, [Co(P6)]-based catalytic system proved to be even effective for asymmetric cyclopropanation of unactivated alkyl-substituted olefins, affording alkynylcyclopropanes **3ae** and **3af** in

moderate yields with high stereoselectivities (entries 30 and 31).

Mechanistic Studies. Combined computational and experimental studies were carried out to shed light on the underlying stepwise radical mechanism of Co(II)-based catalytic system for asymmetric cyclopropanation (Scheme 3). First, density functional theory (DFT) calculations were conducted to examine the details of the catalytic pathway and associated energetics for asymmetric cyclopropanation of styrene (**2a**) with (phenylethynyl)diazomethane (**1a'**) by metalloradical catalyst [Co(P6)] (A) (Scheme 3A; see the Supporting Information for details). The DFT calculations reveal the initial formation of intermediate **B** resulting from the binding of diazomethane **1a'** by catalyst [Co(P6)] through a network of noncovalent attractive interactions, including H-

Scheme 4. Synthetic Applications of Resulting Chiral Alkynyl Cyclopropanes from Co(II)-Catalyzed Olefin Cyclopropanation

A. Stereoselective Synthesis of (Z)-Vinyl Cyclopropane by Hydrogenation^aC. Stereoselective Synthesis of Trisubstituted Cyclopropane by Benzylation^bB. Stereoselective Synthesis of Trisubstituted Cyclopropane by Methylation^bD. Stereoselective Synthesis of Trisubstituted Cyclopropane by Allylation^bE. Stereoselective Synthesis of 1,2,3-Triazole-Substituted Cyclopropane via Click Reaction^cF. Stereoselective Synthesis of 2-Benzylidenetetrahydrothiophene by [1+3+1] Cycloaddition^d

^aCarried out with **3f** (0.10 mmol) and quinoline (2.0 equiv) by Lindlar catalyst (1.2 equiv) in mixed solvent (2.0 mL; hexane:ethyl acetate = 1:1) under H₂ atmosphere at 22 °C for 12 h. ^bCarried out with **3f** (0.10 mmol) and *n*-BuLi (1.6 equiv) in THF (2.0 mL) under N₂ atmosphere at −78 °C for 1 h, followed by addition of electrophile (1.6 equiv) and then stirred at 22 °C for 24 h. ^cCarried out with **3f** (0.10 mmol), NaN₃ (1.5 equiv), and PhI(OAc)₂ (1.0 equiv) in MeCN (2.0 mL) under N₂ atmosphere at 22 °C for 12 h. ^dCarried out with **3a** (0.10 mmol) and Na₂S·9H₂O (6.0 equiv) in DMA (0.5 mL) at 150 °C for 12 h.

bonding and π -stacking interactions, as illustrated in the DFT-optimized structure (Scheme 3A). The binding event, which is slightly exergonic by 1.2 kcal/mol, positions the α -carbon atom of **1a'** in close proximity to the Co(II)-metalloradical center of [Co(P6)] (C...Co: $\sim$ 2.792 Å) for the subsequent activation. Bound diazomethane **1a'** is then further activated by catalyst [Co(P6)] to generate α -Co(III)-propargyl radical intermediate **C** with the release of dinitrogen as byproduct. The metalloradical activation, which is exergonic by 18.2 kcal/mol, is found to be the rate-determining step associated with a relatively high but accessible activation barrier ($\Delta G^{\ddagger}_{TS1}$ = 16.3 kcal/mol). As displayed by the spin plot of intermediate **C** (Scheme 3A), the spin density mainly distributes on α - and γ -carbon atoms in similar amounts (α -C: 0.59; γ -C: 0.41), which can be represented as two resonance forms of α -Co(III)-propargyl radical **C** and γ -Co(III)-allenyl radical **C'**. To rationalize the observed regioselectivity of the catalytic reaction, we calculated the energetics associated with subsequent radical addition to alkene **2a** by both α -Co(III)-propargyl radical form **C** and γ -Co(III)-allenyl radical form **C'**, leading to the formation of γ -Co(III)-benzyl radical intermediate **D** and ϵ -Co(III)-benzyl radical intermediate **D'**, respectively. DFT calculations indicate radical addition of propargyl radical **C** is more favorable than allenyl radical **C'** both kinetically ($\Delta G^{\ddagger}_{TS2}$ = 11.4 kcal/mol; $\Delta G^{\ddagger}_{TS2'}$ = 18.7 kcal/mol) and thermodynamically (ΔG°_{D} = −11.1 kcal/mol; $\Delta G^{\circ}_{D'}$ = −7.2 kcal/mol). According to the DFT calculations, γ -Co(III)-alkyl radical **D** then undergoes radical substitution to produce alkynyl cyclopropane **3a** while regenerating metalloradical catalyst [Co(P6)]. This final step of 3-*exo-tet* radical cyclization, which is exergonic by 15.2 kcal/mol, is found to be

an almost barrierless process. The overall low activation barrier is in accordance with the experimental observation that the catalytic reaction could proceed effectively even at room temperature.

To provide direct evidence for the existence of the key α -Co(III)-propargyl radical intermediate, efforts were made to trap the Co-supported organic radicals for experimental detection and characterization. First, the spin trapping reagent *N*-tert-butyl- α -phenylnitrone (PBN) was added to the reaction mixture of alkynyldiazomethane **1f** with metalloradical catalyst [Co(P1)] in the absence of olefin substrate and was then monitored by X-band electron paramagnetic resonance (EPR) spectroscopy at room temperature (Scheme 3B; see the Supporting Information for details). The observed isotropic EPR spectrum exhibits strong signals with the characteristic splitting pattern at *g*-value of $\sim$ 2.00, which was taken as the evidence for formation of radical III_{[Co(P1)]/If} resulting from PBN trapping of the initially generated Co(III)-propargyl radical intermediate I_{[Co(P1)]/If}.²¹ In accordance with the spin density distribution from DFT calculations (Scheme 3A), the observed broad spectrum (in black) could be near perfectly simulated (in red) as four well-defined triplet of doublet signals (Scheme 3B) by involving four isomeric PBN-trapped radical species that are originated from the two resonance forms of α -Co(III)-propargyl radical intermediate I_{[Co(P1)]/If} and γ -Co(III)-allenyl radical intermediate I'_{[Co(P1)]/If} on the basis of the hyperfine coupling by ¹⁴N (*I* = 1) and ¹H (*I* = 1/2): 81% of O-centered radicals III_{[Co(P1)]/If} from I_{[Co(P1)]/If} as two diastereomers (59% of major diastereomer: *g* = 2.00627, *A*_(N) = 41.9 MHz, *A*_(H) = 9.5 MHz; 22% of minor diastereomer: *g* = 2.00652, *A*_(N) = 38.7 MHz, *A*_(H) = 8.5 MHz) and 19% of O-

centered radicals $\text{III}'_{[\text{Co}(\text{P1})]/1\text{f}}$ from $\text{I}'_{[\text{Co}(\text{P1})]/1\text{f}}$ as two diastereomers (12% of major isomer: $g = 2.00622$, $A_{(\text{N})} = 40.5$ MHz, $A_{(\text{H})} = 6.2$ MHz; 7% of minor isomer: $g = 2.00636$, $A_{(\text{N})} = 38.9$ MHz, $A_{(\text{H})} = 6.0$ MHz).

Besides the spectroscopic observation of the key radical intermediate by EPR, significant efforts were devoted to directly trap α -Co(III)-propargyl radical intermediate by stable TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy) radical for structural characterization (Scheme 3C). When the metal-radical activation of alkynyl diazomethane **1f'** by $[\text{Co}(\text{P1})]$ was conducted in the presence of TEMPO without alkene substrates, we were able to isolate product **4f** in 40% yield, which was shown to contain two geminal TEMPO units at the propargylic position. Formation of bis-TEMPO-trapped product **4f** evidently implies the initial generation of α -Co(III)-propargyl radical $\text{I}_{[\text{Co}(\text{P1})]/1\text{f}}$ which was trapped by TEMPO through radical recombination to generate intermediate $\text{IV}_{[\text{Co}(\text{P1})]/1\text{f}}$. Subsequent radical substitution reaction of $\text{IV}_{[\text{Co}(\text{P1})]/1\text{f}}$ with a second molecule of TEMPO was likely responsible for the final formation of bis-TEMPO-trapped product **4f** upon the homolytic cleavage of the weak Co(III)–C bond. Analogous TEMPO-trapping experiment was carried out with alkynyl diazomethane **1g**, resulting in the isolation of bis-TEMPO-trapped product **4g** in 46% yield, the structure of which was further confirmed by the X-ray crystallography (Scheme 3C). In both reactions, α,α -bis-TEMPO-trapped products were exclusively generated from α -Co(III)-propargyl radical intermediate without the formation of α,γ -bis-TEMPO-trapped products from its γ -Co(III)-allenyl radical resonance form. The observed preference of reactivity at α -propargyl position over γ -allenyl position is in good accordance with the regioselectivity of Co(II)-based catalytic system for asymmetric cyclopropanation, which is also supported by the DFT computational studies.

Synthetic Applications. Considering that the resulting enantioenriched alkynyl cyclopropanes contain both versatile $\text{C}\equiv\text{C}$ triple bond as well as relatively acidic tertiary C–H bonds on the cyclopropane ring, they may serve as useful intermediates for stereoselective organic synthesis. To this end, we carried out several stereoselective transformations using enantioenriched alkynyl cyclopropanes **3f** (95% ee) and **3a** (96% ee) as the model compounds to demonstrate the synthetic utility of this methodology (Scheme 4).²² For example, it was shown that enantioenriched alkynyl cyclopropane **3f** could undergo effective *cis*-hydrogenation with dihydrogen in the presence of Lindlar catalyst, generating vinyl cyclopropane **5f** with (*Z*)-configuration exclusively in almost quantitative yield (99% yield) without any erosion of the original diastereoselectivity (100% ds) and enantioselectivity (100% es) (Scheme 4A). Furthermore, by taking advantage of the difference in acidity between the two tertiary C–H bonds on the cyclopropane ring, we demonstrated that the C–H bond neighboring at the alkynyl substituent in cyclopropane **3f** could be preferentially deprotonated with *n*-BuLi over the C–H bond neighboring at the aryl substituent, leading to site-selective generation of cyclopropyl lithium intermediate **3fLi** (Scheme 4B).^{22a} By adding different alkyl halides as the electrophiles, the in situ-generated cyclopropyl lithium intermediate **3fLi** could undergo subsequent nucleophilic substitution reactions, allowing for stereoselective synthesis of trisubstituted cyclopropanes with three different types of groups while creating an all-carbon quaternary stereogenic center without the formation of the potential allenyl

products. For instance, the in situ reaction of **3fLi** with methyl iodide furnished corresponding 1-methyl-1-alkynyl-2-phenyl cyclopropane **6f** in moderate yield (66%) with notable loss of stereopurities (50% ds and 85% es) presumably due to the small size of the electrophile (Scheme 4B). When the larger benzyl bromide was added as the electrophile, the reaction could deliver corresponding 1-benzyl-1-alkynyl-2-phenyl cyclopropane **7f** in higher yield (72%) with complete retention of the diastereopurity (100% ds) and without significant loss of the original enantiopurity (98% es) (Scheme 4C). The reaction also worked similarly well with allylic bromide as the electrophilic partner, affording desired 1-allyl-1-alkynyl-2-phenyl cyclopropane **8f** in high yield (92%) while maintaining the majority of the stereopurities (93% ds and 88% es) (Scheme 4D). Moreover, oxidative click reaction of cyclopropane **3f** with sodium azide could be successfully achieved in the presence of $\text{PhI}(\text{OAc})_2$, producing 1,2,3-triazole-substituted cyclopropane **9f** in high yield (81%) without any erosion of the original diastereoselectivity (100% ds) and enantioselectivity (100% es) (Scheme 4E).^{22b} Presumably, the in situ-generated azidyl radical first underwent radical addition with the $\text{C}\equiv\text{C}$ bond in cyclopropane **3f** to generate vinyl radical intermediate **A**. Ensuing *5-endo-dig* radical cyclization of intermediate **A** caused the formation of 5-membered aminyl radical intermediate **B**, which gave rise to final product **9f** after hydrogen atom abstraction from the solvent.^{22b} In addition to these demonstrated transformations for stereoselective synthesis of new cyclopropane derivatives, the three-membered cyclopropane ring in the resulting enantioenriched alkynyl cyclopropanes could be expanded to form 5-membered cyclic structure as exemplified by the reaction of enantioenriched alkynyl cyclopropane **3a** with $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ in DMA at high temperature, affording 2-benzylidenetetrahydrothiophene **4a** in high yield (73%) with good configuration control of the trisubstituted olefin (*Z:E* = 85:15) but with only moderate enantioselectivity (36% ee) (Scheme 4F). According to the proposed mechanism,^{22c} the cyclopropane ring in **3a** was first opened by in situ-generated trisulfur radical anion $\text{S}_3^{\bullet-}$ via radical substitution to generate propargylic radical intermediate **C**, which was protonated by H_2O after one-electron reduction and followed by homolysis of the S–S bond to form thiyl radical intermediate **D** while releasing disulfur radical anion $\text{S}_2^{\bullet-}$. The following *5-exo-dig* radical cyclization of intermediate **D** led to the formation of vinyl radical intermediate **E**, which went through another sequence of reduction and protonation to give final product **4a**.^{22c}

CONCLUSIONS

In summary, we have developed the first asymmetric catalytic system that can use in situ-generated α -alkynyl diazomethanes for direct cyclopropanation of alkenes via Co(II)-based metalloradical catalysis (MRC). On the basis of a remarkable ligand effect on the Co(II)-based catalytic system, D_2 -symmetric chiral amidoporphyrin 2,6-DiMeO-QingPhyrin has been identified as the optimal supporting ligand that offers suitable steric, electronic, and chiral environments surrounding the Co(II)-metalloradical center for engaging a network of noncovalent attractive interactions to facilitate the cyclopropanation process. The Co(II)-based metalloradical system is capable of activating different alkynyl diazomethanes under mild conditions for highly asymmetric cyclopropanation of diverse alkenes with varied electronic and steric properties,

affording chiral alkynyl cyclopropanes in high yields with high both diastereoselectivity and enantioselectivity. The combined computational and experimental studies have shed light on the underlying stepwise radical mechanism of the Co(II)-based cyclopropanation system involving a unique α -metalloradical intermediate that is associated with two resonance forms of α -Co(III)-propargyl radical and γ -Co(III)-allenyl radical. In addition to rationalizing the unique profile of reactivity and selectivity, the established mechanism offers a convincing explanation of the regioselectivity toward the formation of alkynyl cyclopropanes via α -Co(III)-propargylic radical form without any complication from potential reaction via γ -Co(III)-allenyl radical form. The resulting enantioenriched alkynyl cyclopropanes, as showcased in several stereospecific transformations, may serve as useful intermediates for stereospecific organic synthesis. Considering the ubiquity of chiral alkynyl cyclopropanes, we believe this Co(II)-catalyzed asymmetric radical cyclopropanation process will find useful applications in organic synthesis.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c13154>.

Experimental details and analytical data for all new compounds (PDF)

Accession Codes

CCDC 2128604 and 2128605 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

X. Peter Zhang – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0001-7574-8409; Email: peter.zhang@bc.edu

Authors

Jing Ke – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0001-8006-6541

Wan-Chen Cindy Lee – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0002-2433-7404

Xiaoxu Wang – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0003-0444-6950

Yong Wang – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0003-0532-3530

Xin Wen – Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States; orcid.org/0000-0002-2589-5117

Complete contact information is available at:
<https://pubs.acs.org/doi/10.1021/jacs.1c13154>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for financial support from the NIH (R01-GM102554) and in part by NSF (CHE-1900375).

■ REFERENCES

- (1) For selected books on radical reactions, see (a) Zard, S. Z. *Radical Reactions in Organic Synthesis*; Oxford University Press: Oxford, UK, 2003. (b) Chatgililoglu, C.; Studer, A. *Encyclopedia of Radicals in Chemistry, Biology, and Materials*. John Wiley & Sons: Chichester, U.K., 2012. For selected reviews on radical reactions, see (c) Zard, S. Z. Recent Progress in the Generation and Use of Nitrogen-Centred Radicals. *Chem. Soc. Rev.* **2008**, 37, 1603–1618. (d) Quiclet-Sire, B.; Zard, S. Z. Fun with Radicals: Some New Perspectives for Organic Synthesis. *Pure Appl. Chem.* **2010**, 83, 519–551. (e) Narayanan, J. M.; Stephenson, C. R. Visible Light Photoredox Catalysis: Applications in Organic Synthesis. *Chem. Soc. Rev.* **2011**, 40, 102–113. (f) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, 113, 5322–5363. (g) Studer, A.; Curran, D. P. Catalysis of Radical Reactions: A Radical Chemistry Perspective. *Angew. Chem., Int. Ed.* **2016**, 55, 58–102.
- (2) For selected reviews on stereoselective radical reactions, see (a) Bar, G.; Parsons, A. F. Stereoselective Radical Reactions. *Chem. Soc. Rev.* **2003**, 32, 251–263. (b) Sibi, M. P.; Manyem, S.; Zimmerman, J. Enantioselective Radical Processes. *Chem. Rev.* **2003**, 103, 3263–3296. (c) Brimiouille, R.; Lenhart, D.; Maturi, M. M.; Bach, T. Enantioselective Catalysis of Photochemical Reactions. *Angew. Chem., Int. Ed.* **2015**, 54, 3872–3890. For selected examples of approaches to controlling radical reactivity and stereoselectivity, see (d) Du, J.; Skubi, K. L.; Schultz, D. M.; Yoon, T. P. A Dual-Catalysis Approach to Enantioselective [2 + 2] Photocycloadditions Using Visible Light. *Science* **2014**, 344, 392–396. (e) Huo, H.; Shen, X.; Wang, C.; Zhang, L.; Röse, P.; Chen, L.-A.; Harms, K.; Marsch, M.; Hilt, G.; Meggers, E. Asymmetric Photoredox Transition-Metal Catalysis Activated by Visible Light. *Nature* **2014**, 515, 100–103. (f) Kainz, Q. M.; Matier, C. D.; Bartoszewicz, A.; Zultanski, S. L.; Peters, J. C.; Fu, G. C. Asymmetric Copper-Catalyzed C–N Cross-Couplings Induced by Visible Light. *Science* **2016**, 351, 681–684. (g) Zhang, W.; Wang, F.; McCann, S. D.; Wang, D.; Chen, P.; Stahl, S. S.; Liu, G. Enantioselective Cyanation of Benzylic C–H Bonds via Copper-Catalyzed Radical Relay. *Science* **2016**, 353, 1014–1018. (h) Kern, N.; Plesniak, M. P.; McDouall, J. J.; Procter, D. J. Enantioselective Cyclizations and Cyclization Cascades of Samarium Ketyl Radicals. *Nat. Chem.* **2017**, 9, 1198–1204. (i) Morrill, C.; Jensen, C.; Just-Baringo, X.; Grogan, G.; Turner, N. J.; Procter, D. J. Biocatalytic Conversion of Cyclic Ketones Bearing α -Quaternary Stereocenters into Lactones in an Enantioselective Radical Approach to Medium-Sized Carbocycles. *Angew. Chem.* **2018**, 130, 3754–3758. (j) Procter, R. S.; Davis, H. J.; Phipps, R. J. Catalytic Enantioselective Minisci-Type Addition to Heteroarenes. *Science* **2018**, 360, 419–422. (k) Biegasiewicz, K. F.; Cooper, S. J.; Gao, X.; Oblinsky, D. G.; Kim, J. H.; Garfinkle, S. E.; Joyce, L. A.; Sandoval, B. A.; Scholes, G. D.; Hyster, T. K. Photoexcitation of Flavoenzymes Enables a Stereoselective Radical Cyclization. *Science* **2019**, 364, 1166–1169. (l) Huang, H.-M.; McDouall, J. J.; Procter, D. J. SmI₂-Catalysed Cyclization Cascades by Radical Relay. *Nat. Catal.* **2019**, 2, 211–218. (m) Cheng, Y.-F.; Liu, J.-R.; Gu, Q.-S.; Yu, Z.-L.; Wang, J.; Li, Z.-L.; Bian, J.-Q.; Wen, H.-T.; Wang, X.-J.; Hong, X.; Liu, X.-Y. Catalytic Enantioselective Desymmetrizing Functionalization of Alkyl Radicals via Cu (i)/CPA Cooperative Catalysis. *Nat. Catal.* **2020**, 3, 401–410. (n) Gu, Q.-S.; Li, Z.-L.; Liu, X.-Y. Copper (i)-Catalyzed Asymmetric Reactions Involving Radicals. *Acc. Chem. Res.* **2020**, 53, 170–181. (o) Li, Z.-L.; Fang, G.-C.; Gu, Q.-S.; Liu, X.-Y. Recent Advances in Copper-Catalysed Radical-Involved Asymmetric 1,2-Difunctionaliza-

tion of Alkenes. *Chem. Soc. Rev.* **2020**, *49*, 32–48. (p) Nakafuku, K. M.; Zhang, Z.; Wappes, E. A.; Stateman, L. M.; Chen, A. D.; Nagib, D. A. Enantioselective Radical C–H Amination for the Synthesis of β -Amino Alcohols. *Nat. Chem.* **2020**, *12*, 697–704. (q) Wang, Y.; Carder, H. M.; Wendlandt, A. E. Synthesis of Rare Sugar Isomers through Site-Selective Epimerization. *Nature* **2020**, *578*, 403–408. (r) Yang, C.-J.; Zhang, C.; Gu, Q.-S.; Fang, J.-H.; Su, X.-L.; Ye, L.; Sun, Y.; Tian, Y.; Li, Z.-L.; Liu, X.-Y. Cu-Catalyzed Intramolecular Radical Enantioconvergent Tertiary B-C(sp³)–H Amination of Racemic Ketones. *Nat. Catal.* **2020**, *3*, 539–546. (s) Zhang, C.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Catalytic Enantioselective C(sp³)–H Functionalization Involving Radical Intermediates. *Nat. Commun.* **2021**, *12*, 475. (t) Zhou, H.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Ligand-Enabled Copper (I)-Catalyzed Asymmetric Radical C(sp³)–C Cross-Coupling Reactions. *ACS Catal.* **2021**, *11*, 7978–7986.

(3) For selected reviews and highlights on Co(II)-based MRC, see (a) Lu, H.; Zhang, X. P. Catalytic C–H Functionalization by Metalloporphyrins: Recent Developments and Future Directions. *Chem. Soc. Rev.* **2011**, *40*, 1899–1909. (b) Pellissier, H.; Clavier, H. Enantioselective Cobalt-Catalyzed Transformations. *Chem. Rev.* **2014**, *114*, 2775–2823. (c) Demarteau, J.; Debuigne, A.; Detrembleur, C. Organocobalt Complexes as Sources of Carbon-Centered Radicals for Organic and Polymer Chemistries. *Chem. Rev.* **2019**, *119*, 6906–6955. (d) Huang, H.-M.; Garduño-Castro, M. H.; Morrill, C.; Procter, D. J. Catalytic Cascade Reactions by Radical Relay. *Chem. Soc. Rev.* **2019**, *48*, 4626–4638. (e) Singh, R.; Mukherjee, A. Metalloporphyrin Catalyzed C–H Amination. *ACS Catal.* **2019**, *9*, 3604–3617. (f) Roy, S.; Das, S. K.; Khatua, H.; Das, S.; Chattopadhyay, B. Road Map for the Construction of High-Valued N-Heterocycles via Denitrogenative Annulation. *Acc. Chem. Res.* **2021**, *54*, 4395–4409.

(4) For selected examples of Ti(III)-based radical processes, see (a) Nugent, W. A.; RajanBabu, T. Transition-Metal-Centered Radicals in Organic Synthesis. Titanium (III)-Induced Cyclization of Epoxy Olefins. *J. Am. Chem. Soc.* **1988**, *110*, 8561–8562. (b) RajanBabu, T.; Nugent, W. A. Selective Generation of Free Radicals from Epoxides Using a Transition-Metal Radical. A Powerful New Tool for Organic Synthesis. *J. Am. Chem. Soc.* **1994**, *116*, 986–997. (c) Gansäuer, A.; Hildebrandt, S.; Michelmann, A.; Dahmen, T.; von Laufenberg, D.; Kube, C.; Fianu, G. D.; Flowers, R. A. Cationic Titanocene (III) Complexes for Catalysis in Single-Electron Steps. *Angew. Chem., Int. Ed.* **2015**, *54*, 7003–7006. (d) Gansäuer, A.; Hildebrandt, S.; Vogelsang, E.; Flowers, R. II Tuning the Redox Properties of the Titanocene (III)/(IV)-Couple for Atom-Economical Catalysis in Single Electron Steps. *Dalton Trans.* **2016**, *45*, 448–452. (e) Hao, W.; Wu, X.; Sun, J. Z.; Siu, J. C.; MacMillan, S. N.; Lin, S. Radical Redox-Relay Catalysis: Formal [3 + 2] Cycloaddition of N-Acylaziridines and Alkenes. *J. Am. Chem. Soc.* **2017**, *139*, 12141–12144. (f) Yao, C.; Dahmen, T.; Gansäuer, A.; Norton, J. Anti-Markovnikov Alcohols via Epoxide Hydrogenation through Cooperative Catalysis. *Science* **2019**, *364*, 764–767. (g) Ye, K.-Y.; McCallum, T.; Lin, S. Bimetallic Radical Redox-Relay Catalysis for the Isomerization of Epoxides to Allylic Alcohols. *J. Am. Chem. Soc.* **2019**, *141*, 9548–9554.

(5) For selected examples of metalloradical-mediated radical processes, see (a) Wayland, B. B.; Poszmik, G.; Mukerjee, S. L.; Fryd, M. Living Radical Polymerization of Acrylates by Organocobalt Porphyrin Complexes. *J. Am. Chem. Soc.* **1994**, *116*, 7943–7944. (b) Zhang, X.-X.; Wayland, B. B. Rhodium (II) Porphyrin Bimetallo-radical Complexes: Preparation and Enhanced Reactivity with CH₄ and H₂. *J. Am. Chem. Soc.* **1994**, *116*, 7897–7898. (c) Chan, K. S.; Li, X. Z.; Dzik, W. I.; de Bruin, B. Carbon-Carbon Bond Activation of 2,2,6,6-Tetramethyl-piperidine-1-oxyl by a Rh^{II} Metalloradical: A Combined Experimental and Theoretical Study. *J. Am. Chem. Soc.* **2008**, *130*, 2051–2061. (d) Chan, Y. W.; Chan, K. S. Metalloradical-Catalyzed Aliphatic Carbon-Carbon Activation of Cyclooctane. *J. Am. Chem. Soc.* **2010**, *132*, 6920–6922. (e) Li, G.; Han, A.; Pulling, M. E.; Estes, D. P.; Norton, J. R. Evidence for Formation of a Co–H Bond from (H₂O)₂Co(dmgBF₂)₂ under H₂: Application to Radical Cyclizations. *J. Am. Chem. Soc.* **2012**, *134*,

14662–14665. (f) Kuo, J. L.; Hartung, J.; Han, A.; Norton, J. R. Direct Generation of Oxygen-Stabilized Radicals by H• Transfer from Transition Metal Hydrides. *J. Am. Chem. Soc.* **2015**, *137*, 1036–1039. (g) Roy, S.; Das, S. K.; Chattopadhyay, B. Cobalt(II)-Based Metalloradical Activation of 2-(Diazomethyl) Pyridines for Radical Transannulation and Cyclopropanation. *Angew. Chem., Int. Ed.* **2018**, *57*, 2238–2243. (h) Roy, S.; Khatua, H.; Das, S. K.; Chattopadhyay, B. Iron (II)-Based Metalloradical Activation: Switch from Traditional Click Chemistry to Denitrogenative Annulation. *Angew. Chem., Int. Ed.* **2019**, *58*, 11439–11443. (i) Das, S. K.; Roy, S.; Khatua, H.; Chattopadhyay, B. Iron-Catalyzed Amination of Strong Aliphatic C(sp³)–H Bonds. *J. Am. Chem. Soc.* **2020**, *142*, 16211–16217. (j) Zhang, Z.; Gevorgyan, V. Co-Catalyzed Transannulation of Pyridotriazoles with Isothiocyanates and Xanthate Esters. *Org. Lett.* **2020**, *22*, 8500–8504. (k) Roy, S.; Das, S. K.; Khatua, H.; Das, S.; Singh, K. N.; Chattopadhyay, B. Iron-Catalyzed Radical Activation Mechanism for Denitrogenative Rearrangement over C(sp³)–H Amination. *Angew. Chem., Int. Ed.* **2021**, *60*, 8772–8780.

(6) (a) Chen, Y.; Fields, K. B.; Zhang, X. P. Bromoporphyrins as Versatile Synthons for Modular Construction of Chiral Porphyrins: Cobalt-Catalyzed Highly Enantioselective and Diastereoselective Cyclopropanation. *J. Am. Chem. Soc.* **2004**, *126*, 14718–14719. (b) Caselli, A.; Gallo, E.; Ragaini, F.; Ricatto, F.; Abbiati, G.; Cenini, S. Chiral Porphyrin Complexes of Cobalt(II) and Ruthenium(II) in Catalytic Cyclopropanation and Amination Reactions. *Inorg. Chim. Acta* **2006**, *359*, 2924–2932. (c) Chen, Y.; Ruppel, J. V.; Zhang, X. P. Cobalt-Catalyzed Asymmetric Cyclopropanation of Electron-Deficient Olefins. *J. Am. Chem. Soc.* **2007**, *129*, 12074–12075. (d) Fantauzzi, S.; Gallo, E.; Rose, E.; Raoul, N.; Caselli, A.; Issa, S.; Ragaini, F.; Cenini, S. Asymmetric Cyclopropanation of Olefins Catalyzed by Chiral Cobalt (II)-Binaphthyl Porphyrins. *Organometallics* **2008**, *27*, 6143–6151. (e) Zhu, S.; Perman, J. A.; Zhang, X. P. Acceptor/Acceptor-Substituted Diazo Reagents for Carbene Transfers: Cobalt-Catalyzed Asymmetric Z-Cyclopropanation of Alkenes with α -Nitrodiazoacetates. *Angew. Chem., Int. Ed.* **2008**, *47*, 8460–8463. (f) Zhu, S.; Xu, X.; Perman, J. A.; Zhang, X. P. A General and Efficient Cobalt(II)-Based Catalytic System for Highly Stereoselective Cyclopropanation of Alkenes with α -Cyanodiazoacetates. *J. Am. Chem. Soc.* **2010**, *132*, 12796–12799. (g) Xu, X.; Lu, H.; Ruppel, J. V.; Cui, X.; Lopez de Mesa, S.; Wojtas, L.; Zhang, X. P. Highly Asymmetric Intramolecular Cyclopropanation of Acceptor-Substituted Diazoacetates by Co(II)-Based Metalloradical Catalysis: Iterative Approach for Development of New-Generation Catalysts. *J. Am. Chem. Soc.* **2011**, *133*, 15292–15295. (h) Paul, N. D.; Mandal, S.; Otte, M.; Cui, X.; Zhang, X. P.; de Bruin, B. Metalloradical Approach to 2H-Chromenes. *J. Am. Chem. Soc.* **2014**, *136*, 1090–1096. (i) Reddy, A. R.; Hao, F.; Wu, K.; Zhou, C. Y.; Che, C. M. Cobalt(II) Porphyrin-Catalyzed Intramolecular Cyclopropanation of N-Alkyl Indoles/Pyrroles with Alkylcarbene: Efficient Synthesis of Polycyclic N-Heterocycles. *Angew. Chem., Int. Ed.* **2016**, *55*, 1810–1815. (j) Chirila, A.; Gopal Das, B.; Paul, N. D.; de Bruin, B. Diastereoselective Radical-Type Cyclopropanation of Electron-Deficient Alkenes Mediated by the Highly Active Cobalt(II) Tetramethyltetraaza[14]Annulene Catalyst. *ChemCatChem* **2017**, *9*, 1413. (k) Wang, J.-Y.; Xie, J.-J.; Lee, W.-C. C.; Wang, D.-S.; Zhang, X. P., Radical Differentiation of Two Esters in Unsymmetric Diazomalonates for Highly Asymmetric Olefin Cyclopropanation. *Chem. Catal.* **2021**, DOI: 10.1016/j.cheecat.2021.11.018.

(7) (a) Dzik, W. I.; Xu, X.; Zhang, X. P.; Reek, J. N.; de Bruin, B. ‘Carbene Radicals’ in Co^{II}(Por)-Catalyzed Olefin Cyclopropanation. *J. Am. Chem. Soc.* **2010**, *132*, 10891–10902. (b) Belof, J. L.; Cioce, C. R.; Xu, X.; Zhang, X. P.; Space, B.; Woodcock, H. L. Characterization of Tunable Radical Metal–Carbenes: Key Intermediates in Catalytic Cyclopropanation. *Organometallics* **2011**, *30*, 2739–2746. (c) Lu, H.; Dzik, W. I.; Xu, X.; Wojtas, L.; de Bruin, B.; Zhang, X. P. Experimental Evidence for Cobalt (III)-Carbene Radicals: Key Intermediates in Cobalt (II)-Based Metalloradical Cyclopropanation. *J. Am. Chem. Soc.* **2011**, *133*, 8518–8521.

- (8) (a) Wang, Y.; Wen, X.; Cui, X.; Wojtas, L.; Zhang, X. P. Asymmetric Radical Cyclopropanation of Alkenes with in Situ-Generated Donor-Substituted Diazo Reagents via Co(II)-Based Metalloradical Catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 1049–1052. (b) Lee, W.-C. C.; Wang, D.-S.; Zhang, C.; Xie, J.; Li, B.; Zhang, X. P. Asymmetric Radical Cyclopropanation of Dehydroaminocarboxylates: Stereoselective Synthesis of Cyclopropyl α -Amino Acids. *Chem.* **2021**, *7*, 1588–1601. (c) Wang, X.; Ke, J.; Zhu, Y.; Deb, A.; Xu, Y.; Zhang, X. P. Asymmetric Radical Process for General Synthesis of Chiral Heteroaryl Cyclopropanes. *J. Am. Chem. Soc.* **2021**, *143*, 11121–11129.
- (9) (a) Noro, M.; Masuda, T.; Ichimura, A. S.; Koga, N.; Iwamura, H. Carbene-Carbene Interconversion between 1-and 3-Phenyl-2-Propynylidenes. *J. Am. Chem. Soc.* **1994**, *116*, 6179–6190. (b) Bowling, N. P.; Halter, R. J.; Hodges, J. A.; Seburg, R. A.; Thomas, P. S.; Simmons, C. S.; Stanton, J. F.; McMahon, R. J. Reactive Carbon-Chain Molecules: Synthesis of 1-Diazo-2,4-Pentadiyne and Spectroscopic Characterization of Triplet Pentadiynylidene ($\text{H}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{H}$). *J. Am. Chem. Soc.* **2006**, *128*, 3291–3302. (c) Wommack, A. J.; Moebius, D. C.; Travis, A. L.; Kingsbury, J. S. Diverse Alkanones by Catalytic Carbon Insertion into the Formyl C–H Bond. Concise Access to the Natural Precursor of Achyrofurin. *Org. Lett.* **2009**, *11*, 3202–3205. (d) Thomas, P. S.; Bowling, N. P.; Burmann, N. J.; McMahon, R. J. Dialkynyl Carbene Derivatives: Generation and Characterization of Triplet *tert*-Butylpentadiynylidene ($\text{tBu}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{H}$) and Dimethylpentadiynylidene ($\text{Me}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{Me}$). *J. Org. Chem.* **2010**, *75*, 6372–6381. (e) Kumar, R.; Turcaud, S.; Micouin, L. The Reaction of Dimethylalkynylaluminum Reagents with Trimethylsilyldiazomethane: Original Reactivity Leading to New α -Silylated Alkynyl Hydrazones. *Org. Lett.* **2014**, *16*, 6192–6195.
- (10) (a) Collin, J.; Lossing, F. Ionization and Dissociation of Allene, Propyne, 1-Butyne, and 1,2-and 1,3-Butadienes by Electron Impact; the C_3H_3^+ Ion. *J. Am. Chem. Soc.* **1957**, *79*, 5848–5853. (b) Fantazier, R. M.; Poutsma, M. L. Reduction of Propargylic Chlorides with Tri-*n*-Butyltin Hydride. The Ambident Behavior of Propargylic Radicals. *J. Am. Chem. Soc.* **1968**, *90*, 5490–5498. (c) Kochi, J. K.; Krusic, P. J. Electron Spin Resonance of Free Radicals from Acetylenes and Allenes. *J. Am. Chem. Soc.* **1970**, *92*, 4110–4114. (d) Kasai, P. H. Electron Spin Resonance Studies of Vinyl, Propargyl, and Butatrienyl Radicals Isolated in Argon Matrices. *J. Am. Chem. Soc.* **1972**, *94*, 5950–5956. (e) Kuhnle, W.; Gey, E.; Ondruschka, B. Theoretical-Studies of Characteristics of Unsaturated Hydrocarbon Radicals of the Allyl and Propargyl Types. *J. Phys. Chem.* **1987**, *268*, 805–814. (f) Herges, R.; Mebel, A. Propargylene. *J. Am. Chem. Soc.* **1994**, *116*, 8229–8237. (g) Adam, W.; Ortega-Schulte, C. M. Spin Delocalization by Triple-Bonded Functionalities in Propargyl and Heteropropargyl Radicals, Assessed from the EPR-Spectral *D* Parameter of 1,3-Cyclopentadienyl Triplet Diradicals. *J. Org. Chem.* **2003**, *68*, 1007–1011. (h) Steinbauer, M.; Lang, M.; Fischer, I.; de Miranda, B. K. C.; Romanzin, C.; Alcaraz, C. The Photoionisation of Propargylene and Diazopropyne. *Phys. Chem. Chem. Phys.* **2011**, *13*, 17956–17959. (i) Maury, J.; Jammi, S.; Vibert, F.; Marque, S. R. A.; Siri, D.; Feray, L.; Bertrand, M. EPR Investigation of Zinc/Iodine Exchange between Propargyl Iodides and Diethylzinc: Detection of Propargyl Radical by Spin Trapping. *J. Org. Chem.* **2012**, *77*, 9081–9086. (j) Giegerich, J.; Petersen, J.; Mitrić, R.; Fischer, I. Photodissociation Dynamics of Propargylene, Hcch. *Phys. Chem. Chem. Phys.* **2014**, *16*, 6294–6302. (k) Osborn, D. L.; Vogelhuber, K. M.; Wren, S. W.; Miller, E. M.; Lu, Y.-J.; Case, A. S.; Sheps, L.; McMahon, R. J.; Stanton, J. F.; Harding, L. B.; Ruscic, B.; Lineberger, W. C. Electronic States of the Quasilinear Molecule Propargylene (HCCCH) from Negative Ion Photoelectron Spectroscopy. *J. Am. Chem. Soc.* **2014**, *136*, 10361–10372. (l) Qiu, S.; Zhang, Y.; Huang, X.; Bao, L.; Hong, Y.; Zeng, Z.; Wu, J. 9-Ethynylfluorenyl Radicals: Regioselective Dimerization and Post Ring-Cyclization Reactions. *Org. Lett.* **2016**, *18*, 6018–6021. (m) Hansmann, M. M.; Melaimi, M.; Bertrand, G. Crystalline Monomeric Allenyl/Propargyl Radical. *J. Am. Chem. Soc.* **2017**, *139*, 15620–15623. (n) Reusch, E.; Kaiser, D.; Schleier, D.; Buschmann, R.; Krueger, A.; Hermann, T.; Engels, B.; Fischer, I.; Hemberger, P. Pentadiynylidene and Its Methyl-Substituted Derivatives: Threshold Photoelectron Spectroscopy of $\text{R}_1\text{-C}_5\text{-R}_2$ Triplet Carbon Chains. *J. Phys. Chem. A* **2019**, *123*, 2008–2017. (o) Dong, X.-Y.; Zhan, T.-Y.; Jiang, S.-P.; Liu, X.-D.; Ye, L.; Li, Z.-L.; Gu, Q.-S.; Liu, X.-Y. Copper-Catalyzed Asymmetric Coupling of Allenyl Radicals with Terminal Alkynes to Access Tetrasubstituted Allenes. *Angew. Chem., Int. Ed.* **2021**, *60*, 2160–2164.
- (11) (a) Zampella, A.; D'Auria, M. V.; Minale, L.; Debitus, C.; Roussakis, C. Callipeltoside A: A Cytotoxic Aminodeoxy Sugar-Containing Macrolide of a New Type from the Marine Lithistida Sponge *Callipelta* sp. *J. Am. Chem. Soc.* **1996**, *118*, 11085–11088. (b) Zampella, A.; D'Auria, M. V.; Minale, L.; Debitus, C. Callipeltosides B and C, Two Novel Cytotoxic Glycoside Macrolides from a Marine Lithistida Sponge *Callipelta* Sp. *Tetrahedron* **1997**, *53*, 3243–3248. (c) Adkins, J. C.; Noble, S. Efavirenz. *Drugs* **1998**, *56*, 1055–1064. (d) Evans, D. A.; Hu, E.; Burch, J. D.; Jaeschke, G. Enantioselective Total Synthesis of Callipeltoside A. *J. Am. Chem. Soc.* **2002**, *124*, 5654–5655. (e) Trost, B. M.; Gunzner, J. L.; Dirat, O.; Rhee, Y. H. Callipeltoside A: Total Synthesis, Assignment of the Absolute and Relative Configuration, and Evaluation of Synthetic Analogues. *J. Am. Chem. Soc.* **2002**, *124*, 10396–10415. (f) Carpenter, J.; Northrup, A. B.; Chung, d.; Wiener, J. J. M.; Kim, S.-G.; MacMillan, D. W. C. Total Synthesis and Structural Revision of Callipeltoside C. *Angew. Chem., Int. Ed.* **2008**, *47*, 3568–3572. (g) Raddatz, R.; Tao, M.; Hudkins, R. L. Histamine H^3 Antagonists for Treatment of Cognitive Deficits in CNS Diseases. *Curr. Top. Med. Chem.* **2010**, *10*, 153–169. (h) Rakhmanina, N. Y.; van den Anker, J. N. Efavirenz in the Therapy of HIV Infection. *Expert Opinion on Drug Metabolism & Toxicology* **2010**, *6*, 95–103. (i) La Ferla, B.; Airolidi, C.; Zona, C.; Orsato, A.; Cardona, F.; Merlo, S.; Sironi, E.; D'Orazio, G.; Nicotra, F. Natural Glycoconjugates with Antitumor Activity. *Nat. Prod. Rep.* **2011**, *28*, 630–648. (j) Frost, J. R.; Pearson, C. M.; Snaddon, T. N.; Booth, R. A.; Turner, R. M.; Gold, J.; Shaw, D. M.; Gaunt, M. J.; Ley, S. V. Callipeltosides A, B and C: Total Syntheses and Structural Confirmation. *Chem. - Eur. J.* **2015**, *21*, 13261–13277.
- (12) For general reviews on metal-catalyzed olefin cyclopropanation with diazo compounds, see (a) Doyle, M. P. Catalytic Methods for Metal Carbene Transformations. *Chem. Rev.* **1986**, *86*, 919–939. (b) Doyle, M. P.; Forbes, D. C. Recent Advances in Asymmetric Catalytic Metal Carbene Transformations. *Chem. Rev.* **1998**, *98*, 911–936. (c) Intrieri, D.; Carminati, D. M.; Gallo, E. The Ligand Influence in Stereoselective Carbene Transfer Reactions Promoted by Chiral Metal Porphyrin Catalysts. *Dalton Trans.* **2016**, *45*, 15746–15761. (d) Mato, M.; Franchino, A.; García-Morales, C.; Echavarren, A. M. Gold-Catalyzed Synthesis of Small Rings. *Chem. Rev.* **2021**, *121*, 8613–8684.
- (13) For asymmetric synthesis of alkynylcyclopropanes from cyclopropanation of conjugated enynes, see ref 8 and (a) Saha, B.; Uchida, T.; Katsuki, T. Highly Enantioselective Intramolecular Cyclopropanation of Alkenyl Diazo Ketones Using Ru(Salen) as Catalyst. *Chem. Lett.* **2002**, *31*, 846–847. (b) Du, H.; Long, J.; Shi, Y. Catalytic Asymmetric Simmons–Smith Cyclopropanation of Silyl Enol Ethers. Efficient Synthesis of Optically Active Cyclopropanol Derivatives. *Org. Lett.* **2006**, *8*, 2827–2829. (c) Suematsu, H.; Kanchiku, S.; Uchida, T.; Katsuki, T. Construction of Aryliridium–Salen Complexes: Enantio- and *Cis*-Selective Cyclopropanation of Conjugated and Nonconjugated Olefins. *J. Am. Chem. Soc.* **2008**, *130*, 10327–10337. (d) Singha, S.; Buchsteiner, M.; Bistoni, G.; Goddard, R.; Fürstner, A. A New Ligand Design Based on London Dispersion Empowers Chiral Bismuth–Rhodium Paddlewheel Catalysts. *J. Am. Chem. Soc.* **2021**, *143*, 5666–5673.
- (14) For asymmetric synthesis of alkynylcyclopropanes from direct functionalization of cyclopropanes, see (a) Krämer, K.; Leong, P.; Lautens, M. Enantioselective Palladium-Catalyzed Carbozincation of Cyclopropanes. *Org. Lett.* **2011**, *13*, 819–821. (b) Teng, H.-L.; Ma, Y.; Zhan, G.; Nishiura, M.; Hou, Z. Asymmetric C(sp)³–H Addition of Terminal Alkynes to Cyclopropanes by a Chiral Gadolinium Catalyst. *ACS Catal.* **2018**, *8*, 4705–4709. (c) Dian, L.; Marek, I. Pd-

Catalyzed Enantioselective Hydroalkynylation of Cyclopropenes. *ACS Catal.* **2020**, *10*, 1289–1293.

(15) Davies, H. M. L.; Boebel, T. A. Asymmetric Synthesis of 1-Alkynylcyclopropane-1-Carboxylates. *Tetrahedron Lett.* **2000**, *41*, 8189–8192.

(16) (a) Skell, P. S.; Klebe, J. Structure and Properties of Propargylene C_3H_2 . *J. Am. Chem. Soc.* **1960**, *82*, 247–248. (b) Padwa, A.; Gareau, Y.; Xu, S. L. A Study of the Rearrangement Chemistry of Alkynyl Carbenes. *Tetrahedron Lett.* **1991**, *32*, 983–986. (c) Liu, Z.; Zhang, X.; Zanon, G.; Bi, X. Silver-Catalyzed Cyclopropanation of Alkenes Using *N*-Nosylhydrazones as Diazo Surrogates. *Org. Lett.* **2017**, *19*, 6646–6649.

(17) (a) Regitz, M.; Maas, G. *Diazo Compounds: Properties and Synthesis*; Academic Press: Orlando, FL, 1986; pp 407–426. (b) Aggarwal, V. K.; Alonso, E.; Bae, I.; Hynd, G.; Lydon, K. M.; Palmer, M. J.; Patel, M.; Porcelloni, M.; Richardson, J.; Stenson, R. A.; Studley, J. R.; Vasse, J.-L.; Winn, C. L. A New Protocol for the in Situ Generation of Aromatic, Heteroaromatic, and Unsaturated Diazo Compounds and Its Application in Catalytic and Asymmetric Epoxidation of Carbonyl Compounds. Extensive Studies to Map out Scope and Limitations, and Rationalization of Diastereo- and Enantioselectivities. *J. Am. Chem. Soc.* **2003**, *125*, 10926–10940. (c) Fulton, J. R.; Aggarwal, V. K.; de Vicente, J. The Use of Tosylhydrazone Salts as a Safe Alternative for Handling Diazo Compounds and Their Applications in Organic Synthesis. *Eur. J. Org. Chem.* **2005**, *2005*, 1479–1492. (d) Seburg, R. A.; Hodges, J. A.; McMahon, R. J. Propynal Equivalents and Diazopropyne: Synthesis of All Mono- ^{13}C Isotopomers. *Helv. Chim. Acta* **2009**, *92*, 1626–1643. (e) Vuluga, D.; Legros, J.; Crousse, B.; Bonnet-Delpon, D. Synthesis of Pyrazoles Through Catalyst-Free Cycloaddition of Diazo Compounds to Alkynes. *Green Chem.* **2009**, *11*, 156–159. (f) Allouche, E. M. D.; Charette, A. B. Cyclopropanation Reactions of Semi-Stabilized and Non-Stabilized Diazo Compounds. *Synthesis* **2019**, *51*, 3947–3963.

(18) Mato, M.; Montesinos-Magraner, M.; Sugranyes, A. R.; Echavarren, A. M. Rh(II)-Catalyzed Alkynylcyclopropanation of Alkenes by Decarbenation of Alkynylcycloheptatrienes. *J. Am. Chem. Soc.* **2021**, *143*, 10760–10769.

(19) Ruppel, J. V.; Jones, J. E.; Huff, C. A.; Kamble, R. M.; Chen, Y.; Zhang, X. P. A Highly Effective Cobalt Catalyst for Olefin Aziridination with Azides: Hydrogen Bonding Guided Catalyst Design. *Org. Lett.* **2008**, *10*, 1995–1998.

(20) Zhu, S.; Ruppel, J. V.; Lu, H.; Wojtas, L.; Zhang, X. P. Cobalt-Catalyzed Asymmetric Cyclopropanation with Diazosulfones: Rigidification and Polarization of Ligand Chiral Environment via Hydrogen Bonding and Cyclization. *J. Am. Chem. Soc.* **2008**, *130*, 5042–5043.

(21) (a) Wang, Y.; Wen, X.; Cui, X.; Zhang, X. P. Enantioselective Radical Cyclization for Construction of 5-Membered Ring Structures by Metalloradical C–H Alkylation. *J. Am. Chem. Soc.* **2018**, *140*, 4792–4796. (b) Wen, X.; Wang, Y.; Zhang, X. P. Enantioselective Radical Process for Synthesis of Chiral Indolines by Metalloradical Alkylation of Diverse $C(sp^3)$ –H Bonds. *Chem. Sci.* **2018**, *9*, 5082–5086.

(22) (a) Ma, S.; He, Q. The Cyclopropyl Effect on the Regioselectivity of Coupling Reactions Involving the Lithiation of 1-Cyclopropyl-2-Arylacetylenes. *Tetrahedron* **2006**, *62*, 2769–2778. (b) Hu, L.; Mück-Lichtenfeld, C.; Wang, T.; He, G.; Gao, M.; Zhao, J. Reaction between Azidyl Radicals and Alkynes: A Straightforward Approach to *NH*-1,2,3-Triazoles. *Chem. - Eur. J.* **2016**, *22*, 911–915. (c) Li, J.-H.; Huang, Q.; Wang, S.-Y.; Ji, S.-J. Trisulfur Radical Anion ($S_3^{\bullet-}$) Involved $[1 + 2 + 2]$ and $[1 + 3 + 1]$ Cycloaddition with Aromatic Alkynes: Synthesis of Tetraphenylthiophene and 2-Benzylidenetetrahydrothiophene Derivatives. *Org. Lett.* **2018**, *20*, 4704–4708.

Recommended by ACS

Controlling Enantioselectivity and Diastereoselectivity in Radical Cascade Cyclization for Construction of Bicyclic Structures

Congzhe Zhang, X. Peter Zhang, *et al.*

JULY 14, 2021

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Catalytic Enantioselective Synthesis of γ -Lactams with β -Quaternary Centers via Merging of C–C Activation and Sulfonyl Radical Migration

Xuan Yu, Guangbin Dong, *et al.*

MAY 17, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

From Serendipity to Rational Design: Heteroleptic Dirhodium Amidate Complexes for Diastereodivergent Asymmetric Cyclopropanation

Fabio Pasquale Caló, Alois Fürstner, *et al.*

APRIL 14, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Carboxylic Acid Directed γ -Lactonization of Unactivated Primary C–H Bonds Catalyzed by Mn Complexes: Application to Stereoselective Natural Product Diversifica...

Arnau Call, Miquel Costas, *et al.*

OCTOBER 13, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >