

EQUIVARIANT GEOMETRY OF ODD-DIMENSIONAL COMPLETE INTERSECTIONS OF TWO QUADRICS

BRENDAN HASSETT AND YURI TSCHINKEL

To Herb Clemens, with admiration

ABSTRACT. Fix a finite group G . We seek to classify varieties with G -action equivariantly birational to a representation of G on affine or projective space. Our focus is odd-dimensional smooth complete intersections of two quadrics, relating the equivariant rationality problem with analogous Diophantine questions over nonclosed fields. We explore how invariants – both classical cohomological invariants and recent symbol constructions – control rationality in some cases.

1. INTRODUCTION

Let $X \subset \mathbb{P}^{2g+1}$ be a smooth complete intersection of two quadrics over an algebraically closed field k of characteristic zero. We are particularly interested in these varieties because they have a rich birational structure, which can be completely understood in small dimensions. They also have beautiful connections to hyperelliptic curves and are key examples in the theory of intermediate Jacobians.

In this paper, we study these varieties from the perspective of their equivariant geometry, for regular generically free actions of finite groups. The main problem is to distinguish such actions up to equivariant birational equivalence, and in particular, to determine which of these are *linearizable*, i.e., equivariantly birational to a linear action on \mathbb{P}^{2g-1} . This shares many similarities with the study of these varieties over nonclosed fields, but has important special features.

To address this problem, we examine various canonical constructions:

- automorphism groups and their induced actions on geometric invariants;
- the structure of varieties of linear subspaces on X and associated pencils of quadric hypersurfaces;
- intermediate Jacobians and their principal homogeneous spaces.

We elaborate on constructions of Reid [Rei72], Desale-Ramanan [DR77], Donagi [Don80], and Bhargava-Gross-Wang [BGW17, Wan18] from a functorial/moduli perspective applicable to equivariant geometry. We also present a new connection with hyperkähler geometry (see Section 5), extending Kummer-type constructions to higher dimensions; connections between Fano and hyperkähler geometry are in the focus of many recent studies, including [FMOS21].

Recent work [BW20, HT21b, HT21a, BW19, KP21] addresses rationality questions for geometrically-rational threefolds over nonclosed fields. Our principal theorem (Theorem 24) demonstrates how these results translate into equivariant contexts: for smooth complete intersections of two quadrics in \mathbb{P}^5 , rationality is governed by the existence of lines.

In Section 2, we present fundamental notions of G -equivariant rationality and related cohomological invariants. We summarize key geometric structures arising from odd-dimensional complete intersections of two quadrics in Section 3. Equivariant constructions and results over nonclosed fields are developed in parallel. The resulting principal homogeneous spaces and their embeddings are explored in Section 4. Section 5 breaks from the main narrative to make a connection with hyperkähler manifolds in small dimensions. The rest of the paper focuses on rationality problems. Two key constructions are presented in Section 6. Section 7 relates existence of fixed points to the analogous questions on rational points over function fields. We close with detailed analysis of the three-dimensional case in Section 8, highlighting both generic behavior and the special properties of examples with large automorphism groups. This brings into sharp relief the similarities and differences between equivariant geometry and geometry over nonclosed fields.

Acknowledgments: The first author was partially supported by Simons Foundation Award 546235 and NSF grant 1701659, the second author by NSF grant 2000099. We thank Aaron Landesman and Daniel Litt for pointing out a gap in a previous version of this paper, as well as the referees for their extraordinary care reviewing our manuscript.

2. ACTIONS AND INVARIANTS

In this section, the base field k is algebraically closed of characteristic zero.

Let G be a finite group and X a smooth projective (connected) variety with a regular G -action; we call such varieties G -varieties. We say that G -varieties X, Y are G -birational if there exists a G -equivariant birational map $X \dashrightarrow Y$; stable G -birationality means G -birationality of $X \times \mathbb{P}^n$ and $Y \times \mathbb{P}^m$, with trivial G -actions on the second factors. Of particular interest are cases when $Y = \mathbb{P}^n$ is projective space, with *linear* G -action, which we describe below.

2.1. Linear actions. An action of a finite group G on \mathbb{P}^n is given by a representation $G \rightarrow \mathrm{PGL}(V)$ where $V = \mathbb{A}^{n+1}$. A linear action on \mathbb{P}^n may have at least two different meanings:

- *strictly linear action*: the projectivization of a linear representation $G \rightarrow \mathrm{GL}(W \oplus 1)$ where $W = \mathbb{A}^n$ and 1 is the trivial representation;
- *linear action*: the projectivization of a linear representation $G \rightarrow \mathrm{GL}(V)$ where $V = \mathbb{A}^{n+1}$;

We record a few obvious facts:

- strictly linear actions admit fixed points;
- if L is a one-dimensional representation of G then the representations V and $V \otimes L$ give rise to the same projective actions;

- using the exact sequence

$$1 \rightarrow \mu_{n+1} \rightarrow \mathrm{SL}_{n+1} \rightarrow \mathrm{PGL}_{n+1} \rightarrow 1$$

any projective action lifts to a linear action for a central extension

$$1 \rightarrow \mu_{n+1} \rightarrow \tilde{G} \rightarrow G \rightarrow 1$$

and the exact sequence

$$1 \rightarrow \mathbb{G}_m \rightarrow \mathrm{GL}_{n+1} \rightarrow \mathrm{PGL}_{n+1} \rightarrow 1$$

lifts any projective action to a representation on a central extension

$$1 \rightarrow \mathbb{G}_m \rightarrow \tilde{G} \rightarrow G \rightarrow 1;$$

- given a projective representation $\rho : G \rightarrow \mathrm{PGL}_{n+1}$, the resulting cohomology class

$$(2.1) \quad \alpha(\rho) \in H^2(G, k^\times), \quad (n+1)\alpha(\rho) = 0,$$

measures the obstruction to lifting ρ to a linear representation of G .

One last observation: Suppose we are given a projective action of G with a fixed point $p \in \mathbb{P}^n$. We obtain a lift

$$\tilde{G} \rightarrow \mathrm{SL}(V)$$

and the preimage of p gives a one-dimensional subspace $L \subset V$. The tensor product $V \otimes L^{-1}$ is also a representation of \tilde{G} on which μ_{n+1} acts trivially, thus descends to a representation of G . Thus we find:

Proposition 1. *A projective representation with fixed point is strictly linear and the class α vanishes.*

If there is an equivariant embedding

$$X \hookrightarrow \mathbb{P}^N$$

such that \mathbb{P}^N admits no fixed points then the same holds true for X .

2.2. Notions of rationality. We say that the G -action on X is *projective* if X admits a G -equivariant birational map to $\mathbb{P}^n, n = \dim(X)$; it is *strictly linear* or *linear* if the G -action on \mathbb{P}^n has the same properties. (We think of these as equivariant analogs of rationality over nonclosed fields.) When G is abelian, the existence of a fixed point is a birational invariant of smooth projective G -varieties [RY00, Appendix]. Thus for abelian actions, the existence of a fixed point is a necessary condition for strict linearity.

The classification of rational G -varieties has been essentially settled in dimension two [DI09], but is largely open in higher dimensions. The *birational* classification of finite group actions on projective space remains a challenging and interesting problem [KT21].

Two strictly linear and generically-free actions of G on projective space need not be G -birational but we shall see that they are necessarily stably G -birational. The key ingredient is:

Proposition 2. *Suppose X and Y are smooth varieties with generically-free G -actions. If there exist G -equivariant vector bundles $E \rightarrow X$ and $F \rightarrow Y$ such that E and F are equivariantly birational then X and Y are equivariantly stably birational.*

This is a corollary of the ‘No-name Lemma’ [CGR06, §4.3]: E is G -equivariantly birational to $\mathbb{A}^{\text{rank}(E)} \times X$ over X , where the affine factor has trivial G -action. One can find examples of X and Y that are not G -birational using the invariants of [RY02].

As a corollary, the following notions of G -equivariant stable birational equivalence coincide:

- $X \times \mathbb{A}^m$ and $Y \times \mathbb{A}^n$ are G -equivariantly birational, where the affine spaces have trivial G -actions;
- $X \times V$ and $Y \times W$ are G -equivariantly birational, where V and W are linear representations of G ;
- $X \times \mathbb{P}^m$ and $Y \times \mathbb{P}^n$ are G -equivariantly birational, where the actions on the projective spaces are strictly linear or admit a fixed point.

The last statement follows from Proposition 1.

2.3. Picard and Brauer groups. We continue to assume that G is a finite group acting regularly on a smooth projective variety X . We refer the reader to [Bri18], [KKV89] and [KKLV89] for background on line bundles and group actions.

A G -linearized line bundle $L \rightarrow X$ consists of a line bundle L over X and an action of G on L , compatible with the action on X , such that the induced action on the fibers is linear. It follows that $\Gamma(X, L^{\otimes N})$, $N \in \mathbb{Z}$, is naturally a representation of G . Conversely, if $X \hookrightarrow \mathbb{P}^n$ is G -equivariant with G acting linearly on \mathbb{P}^n then $L = \mathcal{O}_{\mathbb{P}^n}(-1)|X$ – the restriction of the universal line subbundle on \mathbb{P}^n to X – has a natural linearization. The *equivariant Picard group* $\text{Pic}_G(X)$ parametrizes G -linearized line bundles on X , up to equivariant isomorphism. We have an exact sequence cf. [FW71, Th. 1]

$$(2.2) \quad 0 \rightarrow G^\vee \rightarrow \text{Pic}_G(X) \rightarrow \text{Pic}(X)^G \rightarrow H^2(G, k^\times)$$

where $G^\vee = \text{Hom}(G, k^\times)$. Given a G -equivariant $X \hookrightarrow \mathbb{P}^n$ with G acting projectively on \mathbb{P}^n , the final coboundary morphism applied to $\mathcal{O}_{\mathbb{P}^n}(-1)|X$ vanishes precisely when the class $\alpha = 0$ (see (2.1)). In particular, for $L \in \text{Pic}(X)^G$ some power $L^{\otimes N}$, $N \neq 0$, admits a linearization because $H^2(G, k^\times)$ is a torsion group.

Let $\text{Br}_G(X)$ denote the equivariant Brauer group of X , following Fröhlich and Wall [FW71, FW00]. It parametrizes Azumaya algebras $\mathcal{A} \rightarrow X$ with G -action, compatible with the action on X and linear over the fibers. (Azumaya algebras over commutative rings are direct generalizations of central simple algebras over fields.) We mod out by those of the form $\text{End}(\mathcal{E})$, where \mathcal{E} is a G -equivariant locally-free sheaf on X . The group $\text{Br}_G(X)$ may be computed with a Hochschild-Serre type spectral sequence cf. [FW00, §4], with graded pieces

$$\begin{aligned} & \text{coker}(\text{Pic}(X)^G \rightarrow H^2(G, k^\times)) \\ & \text{ker}(H^1(G, \text{Pic}(X)) \rightarrow H^3(G, k^\times)) \\ & \text{ker} \left(\text{ker}(\text{Br}(X)^G \rightarrow H^2(G, \text{Pic}(X))) \rightarrow \text{coker}(H^1(G, \text{Pic}(X)) \rightarrow H^3(G, k^\times)) \right). \end{aligned}$$

Part of this is summarized in the extension to (2.2) [FW71, Th. 1]:

$$0 \rightarrow G^\vee \rightarrow \text{Pic}_G(X) \rightarrow \text{Pic}(X)^G \rightarrow H^2(G, k^\times) \rightarrow \ker(\text{Br}_G(X) \rightarrow \text{Br}(X)) \rightarrow H^1(G, \text{Pic}(X)) \rightarrow H^3(G, k^\times).$$

Proposition 3. *Let V be a finite-dimensional linear representation of G over k . Then the homomorphism*

$$(2.3) \quad H^2(G, k^\times) = \text{Br}_G(\text{point}) \rightarrow \text{Br}_G(\mathbb{P}(V)),$$

induced by the structure morphism, is an isomorphism.

Suppose that $\rho : G \rightarrow \text{PGL}(V)$ is a projective representation. Then the kernel of (2.3) contains $\alpha(\rho)$.

Corollary 4. *Let X be a smooth projective G -variety and $E \rightarrow X$ a G -equivariant vector bundle. Then the induced homomorphism*

$$\text{Br}_G(X) \rightarrow \text{Br}_G(\mathbb{P}(E))$$

is an isomorphism.

Proof of Proposition 3 and Corollary 4. The first statement of the proposition follows from applying the Hochschild-Serre formalism to $\mathbb{P}(V)$. We are applying $\text{Pic}(\mathbb{P}(V)) = \mathbb{Z}$ (with trivial G -action), $\text{Br}(\mathbb{P}(V)) = 0$, and exact sequence (2.2). For the second, note that projection

$$\mathbb{P}(V) \times \mathbb{P}(V) \rightarrow \mathbb{P}(V)$$

admits the diagonal section so Proposition 1 gives the vanishing of $\alpha(\rho)$ on pullback to $\mathbb{P}(V)$. The corollary follows by computing the étale Leray spectral sequence for $\mathbb{P}(E) \rightarrow X$, using the vanishing underlying Proposition 3. \square

Proposition 5. *Br_G is an equivariant stable birational invariant of smooth projective G -varieties.*

The stable birational invariance of the Brauer group is well-known [CTS21, §5.2].

Proof. We first prove birational invariance. By weak factorization, it suffices to prove the assertion for a blow up

$$\text{Bl}_Z(X) \rightarrow X,$$

where $Z \subset X$ is smooth and irreducible as a G -variety, i.e., G permutes the connected components transitively. In this situation, we observe that

- The exceptional divisor E is G -invariant and irreducible, thus the bottom graded piece of Br_G is unchanged.
- The connected components of E generate a permutation module for G , with trivial H^1 , thus the middle graded piece is unchanged.
- We have $\text{Br}(\text{Bl}_Z(X)) = \text{Br}(X)$ thus the top graded piece is unchanged.

For the stable case, we need to verify that

$$\text{Br}_G(X \times \mathbb{P}^n) = \text{Br}_G(X)$$

provided the G -action on \mathbb{P}^n is linear. We compute this using the Leray spectral sequence associated with the projection $X \times \mathbb{P}^n \rightarrow X$. The vanishing used in the proof of Proposition 3 gives the desired equality. \square

Corollary 6. *Let X be a smooth projective G -variety. Assume there is a G -equivariant embedding*

$$X \hookrightarrow \mathbb{P}^n$$

where the action of G on \mathbb{P}^n is not linear. Then X is not equivariantly birational to projective space with a linear G -action.

Indeed, we can factor

$$\mathrm{Br}_G(\mathrm{point}) \rightarrow \mathrm{Br}_G(\mathbb{P}^n) \rightarrow \mathrm{Br}_G(X)$$

and it suffices to exhibit nonzero elements in the kernel of the first homomorphism. However, the class $[\mathcal{O}_{\mathbb{P}^n}(1)] \in \mathrm{Pic}^G(\mathbb{P}^n)$ maps to a nontrivial $\alpha \in H^2(G, k^\times)$. The spectral sequence above shows that α is in the kernel of $\mathrm{Br}_G(\mathrm{point}) \rightarrow \mathrm{Br}_G(\mathbb{P}^n)$.

Remark 7. The paper [BCDP18, §6] introduces similar ideas via the *Amitsur subgroup*, defined as the image of $\mathrm{Pic}(X)^G \rightarrow H^2(G, k^\times)$.

Much of this extends to a nonclosed field k , except for the interpretation of $\mathrm{Br}_G(\mathrm{Spec}(k))$ as the group cohomology for G . Moreover, we would have to keep track of the Galois actions on μ_n and \bar{G} .

3. ALGEBRAIC GEOMETRY OF PENCILS OF QUADRICS

Here we assume that the ground field is algebraically closed of characteristic zero and all objects are equivariant for the action of a finite group G . The discussion below is valid, with minor changes, when the objects are defined over a field of characteristic zero.

We start with a projective representation $\rho : G \rightarrow \mathrm{PGL}_{2g+2}$ corresponding to a G -action on \mathbb{P}^{2g+1} . Let ρ^* be the dual representation, $\mathrm{Sym}^2(\rho^*)$ the symmetric square, and $\wedge^2(\mathrm{Sym}^2(\rho^*))$ its second exterior power; note that tensor powers of projective representations are well-defined as projective representations.

Consider a smooth complete intersection of two quadrics $X \subset \mathbb{P}^{2g+1}$. We may write

$$X = \{Q_1 = Q_2 = 0\},$$

where Q_1 and Q_2 are basis elements for the distinguished two-dimensional subrepresentation of $\mathrm{Sym}^2(\rho^*)$ generating the ideal of X . (The elements Q_1 and Q_2 need not be invariant under the action of G .) Now $\wedge^2(\mathrm{Sym}^2(\rho))$ has a fixed point – the pencil – so Proposition 1 gives

$$\alpha(\wedge^2(\mathrm{Sym}^2(\rho))) = 4\alpha(\rho) = 0.$$

We write

$$\mathcal{Q} = \mathrm{Bl}_X(\mathbb{P}^{2g+1}) = \{t_1Q_1 + t_2Q_2 = 0\} \subset \mathbb{P}^{2g+1} \times \mathbb{P}^1$$

for the corresponding pencil and

$$q : \mathcal{Q} \rightarrow \mathbb{P}^1$$

for the projection onto the second factor, a fibration in quadric hypersurfaces.

We recall fundamental results from [Rei72] and [Don80]. The singular members of the pencil \mathcal{Q} are given by the degeneracy locus

$$B = \{\det(t_1Q_1 + t_2Q_2) = 0\} \subset \mathbb{P}^1,$$

which consists of $2g + 2$ distinct points b_1, \dots, b_{2g+2} , each corresponding to a nodal fiber \mathcal{Q}_{b_i} . The action of G induces a permutation of the b_i .

Let $F_g(q) \rightarrow \mathbb{P}^1$ denote the relative variety of maximal isotropic subspaces of $q : \mathcal{Q} \rightarrow \mathbb{P}^1$, with Stein factorization

$$F_g(q) \rightarrow C \rightarrow \mathbb{P}^1,$$

where the first arrow is smooth. The double cover $C \rightarrow \mathbb{P}^1$ encodes the two connected components of the variety of maximal isotropic subspaces. The pullback of the point class on \mathbb{P}^1 to C is written g_2^1 , an element of $(\text{Pic}^2(C))^G$. We have natural bijections between

- the branch points b_1, \dots, b_{2g+2} of $C \rightarrow \mathbb{P}^1$;
- the nodes of members of the pencil \mathcal{Q}_t .

The points b_i – regarded as ramification points on C – generate a subgroup of $\text{Pic}(C)$ presented as follows:

- $2b_i = 2b_j = g_2^1$ for all i, j ;
- $b_1 + \dots + b_{2g+2} = (g+1)g_2^1$.

The elements $b_i - b_j$ generate $J(C)[2]$, the two-torsion of the Jacobian of C .

The relative variety of maximal isotropic subspaces $F_g(q)$ is isomorphic to the variety parametrizing $(g-1)$ -dimensional quadric hypersurfaces contained in X , which is stratified by rank

$$K_0(X) \subset K_1(X) \cdots \subset K_g(X) = F_g(q).$$

Consider the variety $F_{g-1}(X) \subset \text{Gr}(g, 2g+2)$ parametrizing $(g-1)$ -dimensional linear subspaces contained in X . We have:

- $F_{g-1}(X)$ is a principal homogeneous space over the Jacobian $J(C)$.
- There is a correspondence

$$\begin{array}{ccc} \widetilde{K_1(X)} & \rightarrow & K_1(X) \\ \downarrow & & \\ F_{g-1}(X) \times F_{g-1}(X) & & \end{array}$$

where the horizontal arrow is the double cover reflecting the support of elements of $K_1(X)$.

- The correspondence induces a morphism

$$(3.1) \quad \text{Sym}^2(F_{g-1}(X)) \rightarrow \text{Pic}^1(C)$$

taking $K_1(X)$ to $C \subset \text{Pic}^1(C)$, with fibers Kummer varieties singular along the 2^{2g} points of $K_0(X)$ over each point of C . Work of X. Wang [Wan18] establishes the formula

$$(3.2) \quad 2[F_{g-1}(X)] = [\text{Pic}^1(C)],$$

as principal homogenous spaces over the Jacobian $J(C)$ of C , i.e., as elements of the Weil-Châtelet group of $J(C)$.

- $K_0(X)$ may be interpreted as a $J(C)[2]$ -principal homogeneous space over C . We have $2^{2g}(2g+2)$ distinguished points of $K_0(X)$ corresponding to the elements lying over the Weierstrass points of C .
- The automorphisms of X act faithfully on $K_0(X)$ but automorphisms lifting the hyperelliptic involution of C fix the $2^{2g}(2g+2)$ distinguished points.

We summarize the implications of the discussion above for the automorphisms:

Proposition 8. *We have an extension*

$$1 \rightarrow J(C)[2] \rightarrow \text{Aut}(X) \rightarrow \text{Aut}(X, C) \rightarrow 1,$$

where $\text{Aut}(X, C)$ is the image of $\text{Aut}(X) \rightarrow \text{Aut}(C)$. Moreover,

- $\text{Aut}(X, C)$ contains the hyperelliptic involution ι in its center,
- $\text{Aut}(X, C)/\langle \iota \rangle \subset \text{Aut}(\mathbb{P}^1)$ acts via permutation on the $2g + 2$ branch points of the cover $C \rightarrow \mathbb{P}^1$, and
- the induced action of $\text{Aut}(X, C)$ on $J(C)[2]$ is induced by this permutation action.

Remark 9. Suppose we diagonalize the forms

$$Q_1 = \sum_{i=1}^{2g+2} x_i^2, \quad Q_2 = \sum_{i=1}^{2g+2} \lambda_i x_i^2.$$

The combinations $Q_2 - \lambda_i Q_1$ correspond to the b_i . The 2-elementary extension

$$(3.3) \quad 1 \rightarrow J(C)[2] \rightarrow H \rightarrow \langle \iota \rangle \rightarrow 1$$

acts on \mathbb{P}^{2g+1} via diagonal $(2g+2) \times (2g+2)$ matrices with ± 1 as entries. The image in the quotient $\langle \iota \rangle$ encodes the determinant of the matrix.

4. PRINCIPAL HOMOGENEOUS SPACES FOR THE JACOBIAN

We maintain the assumptions of Section 3.

4.1. Abstracting the principal homogeneous space. Let $C \rightarrow \mathbb{P}^1$ be a hyperelliptic curve of genus g ; in the equivariant context, we assume that the G -action on C descends to a linear action on \mathbb{P}^1 . The hyperelliptic involution ι induces an involution

$$(4.1) \quad \begin{aligned} \iota : \text{Pic}^1(C) &\rightarrow \text{Pic}^1(C) \\ D &\mapsto g_2^1 - D. \end{aligned}$$

Now suppose that F is a square root of $\text{Pic}^1(C)$, meaning a $J(C)$ principal homogeneous space satisfying Wang's relation (3.2). The automorphisms of F include translations by $J(C)[2]$ and transformations

$$x \mapsto \iota(x) + \tau, \quad \tau \in \text{Pic}^1(C), 2\tau = g_2^1,$$

encoded by the extension (3.3).

Given C , does there exist a smooth complete intersection of two quadrics $X \subset \mathbb{P}^{2g+1}$ whose associated pencil yields C ? A necessary condition is the existence of a $J(C)$ -principal homogeneous space F satisfying Wang's relation (3.2). However, this is not the only way such a variety may arise. Consider a Brauer-Severi variety $P \subset \mathbb{P}^{\binom{2g+3}{2}-1}$, realized geometrically as a 2-Veronese reembedding of \mathbb{P}^{2g+1} , and a pencil of hyperplane sections of P . The base locus X is geometrically a complete intersection of two quadrics in \mathbb{P}^{2g+1} and gives rise to auxiliary varieties $F_{g-1}(X)$ and C as above. However X is generally not embeddable in \mathbb{P}^{2g+1} ; in equivariant terms, a G -action

$$G \times X \rightarrow X$$

may not linearize to \mathbb{P}^{2g+1} . This is often the only obstruction:

Proposition 10. *Let $C \rightarrow \mathbb{P}^1$ be a hyperelliptic curve of genus g . In the equivariant context, we assume the group acts with a fixed point outside the branch locus.*

A square root $F \rightarrow \text{Pic}^1(C)$, as $J(C)$ principal homogeneous spaces, exists if and only if there exists a codimension-two linear section

$$X \subset P \subset \mathbb{P}^{\binom{2g+3}{2}-1},$$

where P is a form of \mathbb{P}^{2g+1} realized as a Veronese variety. Thus there exists

$$X \subset \mathbb{P}^{2g+1},$$

with the group acting linearly in \mathbb{P}^{2g+1} in the equivariant context, if and only if we may choose P such that $[P] = 0 \in H^2(\mathbb{G}_m)$.

The cohomology group is $H^2(G, k^\times)$ in the G -equivariant context and $\text{Br}(k)$ over a nonclosed field k .

Proof. We follow [BGW17]; the case of nonclosed fields is a corollary of their results. Given $p \in \mathbb{P}^1$ (k -rational or G -fixed) that is not a branch point of $C \rightarrow \mathbb{P}^1$, write $p', p'' \in C$ for the points over p . We have an exact sequence for the generalized Jacobian of C with respect to $\{p', p''\}$

$$(4.2) \quad 0 \rightarrow T \rightarrow J_m(C) \rightarrow J(C) \rightarrow 0$$

where the first term

$$T = (\mathbf{R}_{\{p', p''\}/\{p\}} \mathbb{G}_m) / \mathbb{G}_m.$$

Taking two-torsion gives

$$(4.3) \quad 0 \rightarrow \mu_2 \rightarrow J_m(C)[2] \rightarrow J(C)[2] \rightarrow 0.$$

Suppose that L is an étale algebra of degree $2g+2$ over k associated with the branch points of $C \rightarrow \mathbb{P}^1$. We have [BGW17, Prop. 22] identifications

$$(4.4) \quad J_m(C)[2] \Leftrightarrow (\mathbf{R}_{L/k} \mu_2)_{N=1}$$

$$(4.5) \quad J(C)[2] \Leftrightarrow (\mathbf{R}_{L/k} \mu_2)_{N=1} / \mu_2$$

where $N : \mathbf{R}_{L/k} \rightarrow \mu_2$ is the norm map from the restriction of scalars. These act linearly and projectively on \mathbb{P}^{2g+1} respectively. The existence of F is controlled by

$$0 \rightarrow J(C)[2] \rightarrow J(C)[4] \xrightarrow{\times 2} J(C)[2] \rightarrow 0;$$

given $[\text{Pic}^1(C)] \in H^1(J(C)[2])$, the obstruction to a square root $[F] \in H^1(J(C)[4])$ sits in

$$H^2(J(C)[2]) = H^2((\mathbf{R}_{L/k} \mu_2)_{N=1} / \mu_2),$$

i.e., the Steenrod square of $[\text{Pic}^1(C)]$. The vanishing of this class means \mathbb{P}^{2g+1} descends to a Brauer-Severi variety P . Moreover, the obstruction to producing $X \subset \mathbb{P}^{2g+1}$ is controlled by [BGW17, Th. 24]

$$0 \rightarrow J_m(C)[2] \rightarrow J_m(C)[4] \xrightarrow{\times 2} J_m(C)[2] \rightarrow 0;$$

once F exists, this is controlled via (4.3) by a class

$$\alpha \in \text{coker}(H^1(J(C)[2]) \rightarrow H^2(\mu_2)),$$

where $\alpha \equiv [P]$ by the identifications. □

Remark 11. What does this argument yield – in the equivariant context – when there is no fixed point? Assume first that the cocycle in $H^2(G, J(C)[2])$ vanishes. Write $U \subset \mathbb{P}^1$ for the complement of the branch points, with the induced G -action. The generalized Jacobian $J_m(C)$ may still be defined over U using (4.2) with

$$T = (\mathbf{R}_{C \times_{\mathbb{P}^1} U}/U \mathbb{G}_m)/\mathbb{G}_m.$$

We obtain

$$\begin{array}{ccc} \mathcal{X} & \longrightarrow & \mathcal{P} \\ & \searrow & \downarrow \\ & & U \end{array}$$

where the vertical arrow is a Brauer-Severi fibration of relative dimension $2g + 1$. If the cocycle in $H^2(G, J_m(C)[2])$ vanishes then the vertical arrow is a linear \mathbb{P}^{2g+1} fibration.

4.2. Projective geometry. For the moment, we ignore the group action or assume the base field is algebraically closed. Recalling the imbedding $F_{g-1}(X) \subset \mathrm{Gr}(g, 2g+2)$, we have

$$\mathcal{O}_{\mathrm{Gr}(g, 2g+2)}(1)|F_{g-1}(X) = \mathcal{O}_{J(C)}(4\Theta),$$

where Θ is the class of a theta divisor. Note however that the corresponding embedding is *not* linearly normal as

$$4^g = \dim \Gamma(\mathcal{O}_{J(C)}(4\Theta)) > \dim \Gamma(\mathcal{O}_{\mathrm{Gr}(g, 2g+2)}(1)) = \binom{2g+2}{g}, \quad g > 1.$$

For small g , we have

g	4^g	$\binom{2g+2}{g}$
1	4	4
2	16	15
3	64	56
4	256	210

We explain the reason for this discrepancy. Suppose that (J, Θ) is a principally polarized abelian variety and L is a line bundle on J representing Θ . For each $n \in \mathbb{N}$, the Heisenberg extension associated with $n\Theta$

$$1 \rightarrow \mathbb{G}_m \rightarrow \mathcal{G}(L^n) \rightarrow J[n] \rightarrow 1$$

acts on the space of global sections $\Gamma(L^n)$. Recall that the extension data is given by the commutator

$$J[n] \times J[n] \rightarrow \mu_n \subset \mathbb{G}_m$$

associated with the polarization form. Suppose that $n = 4$ and realize $J[2] \subset J[4]$ in the standard way; the commutator pairing for 4Θ is isotropic on $J[2]$, i.e., we may regard

$$\mu_2 \times J[2] \subset \mathcal{G}(L^4)$$

as an abelian subgroup. Thus it is reasonable to diagonalize the theta functions for this group. Indeed, we have

Proposition 12. [BL04, Ex. 6.10.1] *Let $\vartheta \in \Gamma(J, L)$ denote a generator and $\tau_x^* \vartheta$ its translate under $x \in J$. Then the elements*

$$\{2^* \vartheta_x : x \in J[2]\}$$

form a basis for $\Gamma(J, L^4)$, naturally indexed by the 2-torsion elements of J . (Here $2 : J \rightarrow J$ is multiplication by two.)

Assume that either k is nonclosed or that all the varieties and constructions are G -equivariant. For our application, we use the squaring map

$$F_{g-1}(X) \rightarrow \text{Pic}^1(C)$$

introduced in (3.1); thus Proposition 12 applies.

Consider the canonical theta divisor $\vartheta = \text{Sym}^{g-1}(C) \subset \text{Pic}^{g-1}(C)$. We analyze the translates of ϑ by elements in

$$\langle b_1, \dots, b_{2g+2} \rangle \subset \text{Pic}(C)$$

contained in $\text{Pic}^1(C)$. These have the structure of a principal homogeneous space for $J(C)[2]$. The Galois action on $\langle b_1, \dots, b_{2g+2} \rangle$ factors through the permutation representation on the branch points.

Suppose first that g is even; here the principal homogeneous space is trivial with distinguished divisor

$$\vartheta - \frac{g-2}{2} g_2^1,$$

i.e., the canonical theta divisor translated by the g_2^1 . Recall that $J(C)[2]$ corresponds to even partitions

$$S \sqcup S^c = \{b_1, \dots, b_{2g+2}\}, \quad |S| = 2j.$$

These are counted via the combinatorial identity for even g

$$4^g = \binom{2g+2}{g} + \sum_{j=0}^{g/2-1} \binom{2g+2}{2j}.$$

The sections of

$$\Gamma(F_{g-1}(X), \mathcal{O}_{F_{g-1}(X)}(1))$$

correspond to translates associated with sums of g branch points b_1, \dots, b_{2g+2} .

Remark 13. For even g , any square root F of $\text{Pic}^1(C)$ admits a distinguished polarization of type 4Θ . However, given a projective representation $\rho : G \rightarrow \text{PGL}(V)$ note that

$$\alpha(\wedge^g \rho) = g\alpha(\rho).$$

Thus two-torsion $\alpha(V) \in H^2(\mathbb{G}_m)$ vanishes on passage from $X \subset \mathbb{P}(V)$ to $F_{g-1}(X) \subset \mathbb{P}(\wedge^g V)$.

Now take g odd. The odd-degree divisors in $\langle b_1, \dots, b_{2g+2} \rangle$ – a principal homogeneous space of $J(C)[2]$ – correspond to odd partitions

$$S \sqcup S^c = \{b_1, \dots, b_{2g+2}\}, \quad |S| = 2j-1.$$

These index translates of ϑ pulling back to our desired polarization on $F_1(X)$. We have the combinatorial identity for odd g

$$4^g = \binom{2g+2}{g} + \sum_{j=1}^{(g-1)/2} \binom{2g+2}{2j-1}.$$

Again, the sections correspond to translates associated with sums of g branch points.

Remark 14. For odd g , any square root F of $\text{Pic}^1(C)$ has a (Galois or G) invariant divisor class: the pull back of the divisors

$$\vartheta - \sum_{j \in J} b_j - \frac{g-2j-1}{2} g_2^1 \in \text{Pic}^1(C), \quad |J| = 2j-1,$$

to F . However, there may be an obstruction to the existence of a line bundle (defined over k or linearized for G) on X realizing this class. Nonzero two-torsion $\alpha(V) \in H^2(\mathbb{G}_m)$ remains nonzero on passage from $X \subset \mathbb{P}(V)$ to $F_{g-1}(X) \subset \mathbb{P}(\wedge^g V)$.

5. LAGRANGIAN INTERPRETATION IN DIMENSION THREE

In Section 4.1 we discussed how to recover a smooth complete intersection of two quadrics $X \subset \mathbb{P}^{2g+1}$ from the associated hyperelliptic curve C , principal homogeneous space $J(C) \times F \rightarrow F$, and additional cohomological data. We present a geometric framework for these reconstruction results when $g = 2$.

The relationship between hyperkähler manifolds Y and Fano varieties arising as Lagrangian submanifolds is rich and intricate. Lagrangian $\mathbb{P}^n \subset Y$ can be characterized via intersection properties of the Hodge lattice of Y [BHT15, HT13]. Further subtle constructions have been studied in [FMOS21, §1.1]. For example, cubic fourfolds arise as Lagrangian submanifolds of hyperkähler varieties of dimension eight [LLSvS17].

We saw in Section 3 that Kummer varieties arise naturally in the study of $X \subset \mathbb{P}^{2g+1}$. For $g = 2$, Kummer *surfaces* take center stage but generalized Kummer *sixfolds* are most relevant for recovering X . We realize X naturally as a Lagrangian submanifold of a Kummer sixfold naturally arising from the variety of lines $F_1(X)$. Recovering X from $F = F_1(X)$ boils down to understanding certain Lagrangian subvarieties in this Kummer sixfold.

5.1. The basic construction.

Assume that the ground field is algebraically closed.

Let $X \subset \mathbb{P}^5$ denote a smooth complete intersection of two quadrics, $F_1(X)$ its variety of lines, and $\text{Alb}(F_1(X))$ the associated principally polarized abelian surface. The variety of conics on X equals the variety $F_2(q)$ in Section 3. Thus it fibers over a genus two curve C , parametrizing connected components of the varieties of maximal isotropic subspaces in the quadric hypersurfaces cutting out X . We may interpret $\text{Alb}(F_1(X)) \simeq J(C)$.

Consider the Hilbert scheme $F_1(X)^{[4]}$ and the natural map

$$\begin{aligned} F_1(X)^{[4]} &\rightarrow J(C) \\ (\ell_1, \ell_2, \ell_3, \ell_4) &\mapsto \ell_1 + \ell_2 + \ell_3 + \ell_4 - h^2, \end{aligned}$$

where h is the hyperplane class. Here 0 is identified with cycles of lines obtained as codimension-two linear sections of X . The preimage of 0 is a subvariety

$$\text{Kum}(X) \subset F_1(X)^{[4]},$$

a twist of the generalized Kummer sixfold $K_{J(C)}(3)$ associated with $J(C)$. Regarding $F_1(X)$ as a twist of $J(C)$ by a cocycle for $J(C)[4]$, applying this cocycle to $K_{J(C)}(3)$ yields $\text{Kum}(X)$.

Consider the incidence variety

$$Z = \{[x, \ell] : x \in \ell\} \subset X \times F_1(X)$$

and the projection $\pi : Z \rightarrow X$. This is generically finite of degree four.

Proposition 15. *The projection π is flat over X .*

Proof. Indeed, since Z and X are both smooth and projective it suffices to show that π is equidimensional. But if a point $x \in X$ were contained in a positive dimensional family of lines then either

- X admits a ruled hyperplane section through x – a cone over an elliptic quartic curve – which would force X to be singular at x ;
- X admits a ruled surface of degree < 4 through x , violating the Lefschetz hyperplane theorem.

Since X is assumed to be smooth, we conclude the flatness of π . \square

As a corollary, we obtain

Proposition 16. *There is an injective morphism*

$$\begin{aligned} j : X &\hookrightarrow \text{Kum}(X) \\ x &\mapsto \pi^{-1}(x) \end{aligned}$$

realizing X as a Lagrangian subvariety of $\text{Kum}(X)$.

5.2. Numerical invariants. Assume that X is general in the sense that $\text{NS}(J(C))$ is of rank one, generated by $[\Theta]$. Then the Néron-Severi group of $K_{J(C)}(3)$ has rank two and is generated by [Bea83, p. 769]

- θ – subschemes with support along a theta divisor;
- e – where the nonreduced subschemes have class $2e$.

Proposition 17. *The restriction homomorphism*

$$j^* : \text{NS}(\text{Kum}(X)) \rightarrow \text{NS}(X) \simeq \mathbb{Z}h$$

is given by

$$j^*(\theta) = 5h, \quad j^*(e) = 4h.$$

Proof. For the purpose of this computation, we may ignore G -actions or work over an algebraically closed field. The class θ on $J(C) = F_1(X)$ may be realized as the locus $W_\ell \subset F_1(X)$ of lines incident to a fixed line $\ell \subset X$, which sweeps out a divisor on X . This divisor is the exceptional divisor of the projection

$$\pi_\ell : X \dashrightarrow \mathbb{P}^3;$$

the center of the inverse map is a quintic space curve C of genus two so the exceptional locus has degree 20. This yields the first equation.

The locus of nonreduced subschemes on $\text{Kum}(X)$ restricts to the branch locus of $\pi : Z \rightarrow X$. Restricting to a line $\ell \subset X$, we see that

$$\pi^{-1}(\ell) = \ell \sqcup W_\ell,$$

where the latter component has genus two and is realized as a degree-three cover of ℓ . Such a cover has eight branch points, so the branch locus has class $8h$ and we get the second equation. \square

5.3. Reversing the construction. For the moment, let A be an abelian surface over k , not necessarily principally polarized.

- The group $A[4]$ acts on A via translation.
- The semi-direct product $A[4] \rtimes \mu_2$, where μ_2 acts on $A[4]$ via ± 1 , acts on A as well.
- Consider the addition map

$$\alpha : A^{[4]} \rightarrow A$$

and the induced action of G on $A^{[4]}$. Note that $A[4]$ acts on the fibers and the action of μ_2 commutes with addition. Thus G acts on $K_A(3)$ as well.

Note that G admits a distinguished normal 2-elementary subgroup

$$H = A[2] \times \mu_2,$$

as -1 acts trivially on two-torsion elements. Now assume $A = J(C)$ and consider the various Lagrangian threefolds in $K_{J(C)}(3)$. The subgroup $H \simeq (\mathbb{Z}/2\mathbb{Z})^5$ stabilizes each, acting via automorphisms. The full group G gives an orbit of 16 components, with transitive action of

$$G/H \simeq A[4]/A[2] \simeq A[2].$$

Remark 18. Assume that the base field k is arbitrary, of characteristic zero. Let C be a smooth projective curve of genus two over k and F a principal homogeneous space for $J(C)$ such that $2[F] = [\text{Pic}^1(C)]$. Let $K_F(3) \subset F^{[4]}$ denote the generalized Kummer variety lying over the divisor $g_2^1 \in \text{Pic}^2(C)$. We may realize F as the image of a cocycle for $J(C)[4]$; the 16 conjugate Lagrangian threefolds naturally form a principal homogeneous space for $J(C)[4]/J(C)[2] \simeq J(C)[2]$. However, even when this is trivial there is no guarantee that the corresponding X can be defined over k – the obstruction to descent is discussed in Proposition 10.

6. RATIONALITY CONSTRUCTIONS

We continue to assume that $X \subset \mathbb{P}^{2g+1}$ is a smooth complete intersection of two quadrics.

6.1. Simple rational parametrizations. Fix a line $\ell \subset X$ and consider the projection

$$\pi_\ell : X \dashrightarrow \mathbb{P}^{2g-1},$$

a birational map, resolved by blowing up ℓ . The inverse map

$$\mathbb{P}^{2g-1} \rightarrow \mathbb{P}^{2g+1}$$

is obtained as follows: Consider a matrix

$$\begin{pmatrix} l_{00} & l_{01} \\ l_{10} & l_{11} \\ q_0 & q_1 \end{pmatrix},$$

where the l_{ij} are linear and q_i are quadratic. The 2×2 minors generate an ideal I_Z , where Z is the base locus of π_ℓ^{-1} . Cubic forms in I_Z yield the linear series inducing this map. The kernel of the matrix gives a rational map

$$\phi : Z \dashrightarrow \mathbb{P}^1,$$

which is regular for $g = 2, 3$. The generic fiber of ϕ is a quadric hypersurface in \mathbb{P}^{2g-3} ; we may interpret this as ℓ^\perp/ℓ , understood as a subquotient of the generic fiber of the pencil $q : \mathcal{Q} \rightarrow \mathbb{P}^1$ (from Section 3), i.e., the generic fibers of ϕ and q are equivalent in the Witt ring of $k(\mathbb{P}^1)$.

Much more can be said when $g = 2$; we refer the reader to the corresponding case in Section 6.2.

For the $g = 3$ case, the fibration

$$\phi : Z \rightarrow \mathbb{P}^1$$

is a quadric surface fibration. We will denote by C the discriminant curve of this fibration. Specifically, the relative variety of lines factors

$$F_1(\phi) \xrightarrow{\cong} C \rightarrow \mathbb{P}^1,$$

where ϖ is a smooth conic fibration. The following conditions are equivalent:

- ϕ admits a section;
- ϖ admits a section;
- $q : \mathcal{Q} \rightarrow \mathbb{P}^1$ admits an isotropic plane containing $\ell \times \mathbb{P}^1$.

These are elementary properties of quadratic forms. The Amer-Brumer Theorem [Lee, §2] gives a fourth equivalent condition:

- there exists a plane $\mathbb{P}^2 \subset X$.

The fibration ϖ admits a section if and only if we can express

$$F_1(\phi) = \mathbb{P}(\mathcal{E})$$

for some rank-two vector bundle $\mathcal{E} \rightarrow C$. Desale-Ramanan [DR77] use such constructions to analyze rank-two vector bundles of odd degree on hyperelliptic curves, relating automorphisms of X to natural tensor and duality operations on the vector bundles.

Remark 19. Over nonclosed fields, e.g. $k = \mathbb{C}(s)$, and for $g \geq 2$ (resp. $g \geq 3$), it is possible for $X \subset \mathbb{P}^{2g+1}$ (resp. $F_1(X) \subset \mathrm{Gr}(2, 2g+2)$) to admit a rational point even when C admits no divisors of odd degree.

Consider a hyperelliptic curve $C_0 \rightarrow \mathbb{P}^1$ represented as a double cover branched over $g+1$ orbits for an involution on \mathbb{P}^1 . For example, if the involution is

$$[1, t] \mapsto [1, -t]$$

we could take the branch points as the roots $\pm\lambda_1, \dots, \pm\lambda_{g+1}$ of $\prod_{i=1}^{g+1}(t^2 - a_i)$ with the a_i distinct and nonzero. Write

$$X_0 = \left\{ \sum_{i=1}^{2g+2} x_i^2 = \sum_{i=1}^{g+1} \lambda_i (x_{2i-1}^2 - x_{2i}^2) = 0 \right\},$$

with involution given by

$$(x_1, x_2, x_3, x_4, \dots, x_{2g+1}, x_{2g+2}) \mapsto (x_2, x_1, x_4, x_3, \dots, x_{2g+2}, x_{2g+1});$$

the associated hyperelliptic curve C_0 has the desired branch locus.

Choose a quadratic extension L/k and let X and C denote the associated quadratic twists of X_0 and C_0 . By construction, C admits no cycles of odd degree. However, X is geometrically rationally connected. The same holds for $F_1(X)$ provided $g \geq 3$ – indeed, it has ample anticanonical class and thus is geometrically rationally connected. Thus both varieties have k -rational points by the Graber-Harris-Starr Theorem.

Note that [BGW17, Th. 29] implies that $F_{g-1}(X)$ admits no k -rational points.

6.2. Stable rationality constructions. Fix a $(g-1)$ -dimensional subspace $L \subset X$ and look at the projection from L :

$$\pi : X \dashrightarrow \mathbb{P}^{g+1}$$

with generic fiber a projective space \mathbb{P}^{g-2} . We analyze the structure of this bundle.

Suppose that $L = \{x_0 = \dots = x_{g+1}\}$ so that the induced map on linear spaces factors through

$$\mathbb{P}(\mathcal{O}_{\mathbb{P}^{g+1}}(-1) \oplus \mathcal{O}_{\mathbb{P}^{g+1}}^g)$$

so that y_0, \dots, y_{g-1} are trivializing sections of the $\mathcal{O}_{\mathbb{P}^{g+1}}$ factors and the linear series to \mathbb{P}^{2g+1} is given by

$$y_0, \dots, y_{g-1}, zx_0, \dots, zx_{g+1}.$$

The proper transform of X has equations

$$A(x_i; y_j) + zQ(x_j) = B(x_i; y_j) + zR(x_j) = 0,$$

where A and B are bilinear and Q and R are quadratic. Eliminate z to get the relation

$$z = -A/Q = -B/R,$$

which gives

$$AR - BQ = F_0 y_0 + \dots + F_{g-1} y_{g-1} = 0,$$

which is cubic in x_i and linear in y_j ; the F_j are cubic forms in x_0, \dots, x_{g+1} . This is the formula for the \mathbb{P}^{g-2} bundle in the product

$$\mathbb{P}_{x_i}^{g+1} \times \mathbb{P}_{y_j}^{g-1}.$$

Look at the locus C where the morphism

$$\mathrm{Bl}_L(X) \rightarrow \mathbb{P}^{g+1}$$

fails to be flat. The F_j are linear combinations of Q and R with linear coefficients. The locus C is the residual intersection to the locus

$$Z = \{Q = R = 0\}$$

in the intersection of cubics

$$\{F_0 = \cdots = F_{g-1} = 0\}.$$

Proposition 20. *The excess intersection contribution of Z to the intersection of cubics is $3^g - (2g + 1)$ whence C has degree $2g + 1$. The curve C is hyperelliptic, embedded via a generic $(2g + 1)$ -degree polarization D .*

Proof. This is a computation with Fulton's excess intersection formula, encoded by the exact sequence

$$0 \rightarrow N_{Z/\mathbb{P}^{g+1}} \simeq \mathcal{O}_Z(2)^2 \rightarrow \mathcal{O}_Z(3)^g \rightarrow Q \rightarrow 0,$$

where the equivalence of Z equals $c_{g-2}(Q)$. Note that

$$(1 + 3ht)^g / (1 + 2ht)^2 = 1 + (3g - 4)ht + \cdots + \frac{3^g - 2g - 1}{4}h^{g-2}t^{g-2}$$

using the identity

$$\sum_{j+k=0}^{g-2} \binom{g}{k} 3^k (j+1)(-2)^j = \frac{3^g - 2g - 1}{4}.$$

□

Fixing (C, D) , what are the constructions that arise? We analyze the induced morphism

$$\begin{aligned} \gamma : F_{(g-1)}(X) &\rightarrow \text{Pic}^{2g-1}(C) \\ L &\mapsto D. \end{aligned}$$

following [Don80] and [BGW17]. First, translation by two-torsion in $J(C)[2]$ corresponds to an automorphism of X acting trivially on cohomology. Thus we have

$$\gamma([L] + \tau) = \gamma([L]), \quad \tau \in J(C)[2].$$

Automorphisms with determinant -1 may be presented in the form

$$p \mapsto -p + \beta, \quad \beta \in \text{Pic}^1(C), 2\beta = g_2^1.$$

These act by -1 on middle cohomology whence

$$\gamma(-[L] + \beta) = (2g + 1)g_2^1 - D.$$

Moreover, Wang's formula $2[F_1(X)] = [\text{Pic}^1(C)]$ [Wan18] implies these are the only possible relations intertwining γ for a generic curve C .

To summarize our discussion:

Proposition 21. *For each hyperelliptic curve C of genus g and unordered pair of divisors*

$$D, D' \in \text{Pic}^{2g+1}(C), \quad D + D' = (2g + 1)g_2^1,$$

we obtain a group of stable birational equivalences of \mathbb{P}^{g+1} parametrized by the group

$$(\mathbb{Z}/2\mathbb{Z})^{2g+1} \simeq \langle b_1, \dots, b_{2g+2} \rangle / \langle g_2^1 \rangle \subset \text{Pic}(C) / \langle g_2^1 \rangle.$$

Remark 22. When $g = 2$, we obtain birational equivalences of \mathbb{P}^3 . This is the subgroup of the Cremona group on \mathbb{P}^3 associated with an orbit of $F_1(X)$ under the diagonalizable automorphisms of X . If $\ell \subset X$ is a line and $h \in \text{Aut}(X)$ is a diagonalizable automorphism then the associated birational map is

$$\mathbb{P}^3 \xleftarrow{\pi_\ell} X \xrightarrow{h} X \xrightarrow{\pi_\ell} \mathbb{P}^3.$$

7. REDUCTION TO NONCLOSED FIELDS

We are interested in translating rationality criteria for geometrically rational threefolds over nonclosed fields to the equivariant context. For example, the existence of points and subvarieties of prescribed type sometimes suffices to characterize rationality over nonclosed fields. We hope that the corresponding criteria for G -equivariant rationality are valid for varieties with G -action, where G is a finite group. We focus on situations where principal homogeneous spaces over abelian varieties control rationality.

7.1. Representations and monodromy groups. We recall results of [GLLM15] on representations of monodromy groups for curves with group action.

Let Σ denote a smooth projective complex curve of genus $g \geq 2$ and fundamental group T . Consider a finite group G and a surjective homomorphism $p : T \twoheadrightarrow G$ with kernel R . This is associated with a connected covering $\tilde{\Sigma} \rightarrow \Sigma$ with homology [GLLM15, Prop. 1.1]:

$$(7.1) \quad H_1(\tilde{\Sigma}, \mathbb{Q}) \simeq \mathbb{Q} \otimes_{\mathbb{Z}} R/[R, R] \simeq \mathbb{Q}^2 \oplus \mathbb{Q}[G]^{2g-2}.$$

This is an equivariant refinement of the Hurwitz formula due to Chevalley-Weil.

We decompose $\mathbb{Q}[G]$ using the theory of semisimple algebras [GLLM15, §3.2]:

$$\mathbb{Q}[G] \simeq \mathbb{Q} \times \prod_{i=1}^{\ell} A_i,$$

where each $A_i \simeq \text{Mat}_{m_i}(D_i)$, matrices over a division algebra. Moreover, let L_i denote the center of D_i , a number field. The index i for the product encodes types of nontrivial representations of G that are irreducible over \mathbb{Q} . The formula (7.1) therefore yields

$$(7.2) \quad H_1(\tilde{\Sigma}, \mathbb{Q}) \simeq \mathbb{Q}^{2g} \oplus \bigoplus_{i=1}^{\ell} A_i^{2g-2}.$$

Each of the summands A_i^{2g-2} comes with a natural skew-Hermitian structure with respect to an explicit subfield $K_i \subset L_i$ of index ≤ 2 [GLLM15, §3]. For each index i , there is a distinguished algebraic group $\mathcal{G}_{G,i}$, defined over K_i , parametrizing the automorphisms of A_i^{2g-2} preserving this skew-Hermitian structure. (We abuse notation, using the same notation for this group and its restriction of scalars to \mathbb{Q} .) The complex groups that may arise are listed in [GLLM15, Thm. 1.7]:

$$(7.3) \quad \text{Sp}_{(2g-2)n}(\mathbb{C}), \text{O}_{(2g-2)n}(\mathbb{C}), \text{GL}_{(2g-2)n}(\mathbb{C})$$

for some $n \in \mathbb{N}$.

Let $\mathfrak{D}_i \subset A_i$ denote the order arising as the image of $\mathbb{Z}[G]$, $\mathcal{G}_{G,i}(\mathfrak{D}_i)$ the resulting arithmetic group. Note that this arithmetic group depends only on the structure of G and its action on the symplectic form. Write $\Gamma_{G,p}$ for the mapping class group of G -coverings $\tilde{\Sigma} \rightarrow \Sigma$ [GLLM15, p. 1494]. For each $i = 1, \dots, \ell$, it admits a representation

$$\rho_{G,p,i} : \Gamma_{G,p} \rightarrow \mathcal{G}_{G,i} \subset \text{Aut}_{A_i}(A_i^{2g+2}).$$

We use $\mathcal{G}_{G,i}^1$ to denote the elements of $\mathcal{G}_{G,i}$ of reduced norm one over L_i ; $\mathcal{G}_{G,i}^1(\mathfrak{D}_i)$ has finite index in $\mathcal{G}_{G,i}(\mathfrak{D}_i)$ [GLLM15, Prop. 3.9] as the reduced norm takes values in roots of unity.

To summarize, we obtain natural representations of a finite-index subgroup of the mapping class group $\Gamma_{G,p}$ into the arithmetic groups $\mathcal{G}_{G,i}(\mathfrak{D}_i)$ [GLLM15, p. 1528]. Fortunately, there are sufficient conditions guaranteeing that the image of $\rho_{G,p,i}$ contains a finite-index subgroup of our arithmetic group. Assume further that $g \geq 3$ and p factors

$$p : T \xrightarrow{\varphi} F_g \xrightarrow{p'} G,$$

where

- F_g is a free group on g generators;
- φ is surjective;
- the kernel of p' contains one of the free generators of F_g .

Whenever G can be generated by $g - 1$ elements we can find p satisfying these conditions. Under these assumptions, the image of $\rho_{G,p,i}$ contains a finite-index subgroup of $\mathcal{G}_{G,i}^1(\mathfrak{D}_i)$ [GLLM15, Thm. 1.6].

Observe that $\mathcal{G}_{G,i}^1(\mathfrak{D}_i) \subset \mathcal{G}_{G,i}^1$ is Zariski dense by the Borel density theorem [Mor15, 4.5.6, 5.1.11]. Borel's Theorem requires that associated real Lie group has no compact factors; indeed, the assumption $g \geq 3$ guarantees the factors have \mathbb{Q} -rank at least two [GLLM15, p. 1529]. (Information about the real forms arising from these groups may be found in [GLLM15, § 4].)

The fact that the monodromy is large has implications for the structure of $H_1(\tilde{\Sigma}, \mathbb{Q})$. The decomposition (7.2) cannot be refined; any summand of $H_1(\tilde{\Sigma}, \mathbb{Q})$ stable under the action of $\Gamma_{G,p}$ is a direct sum of the \mathbb{Q}^{2g} (coming from $H_1(\Sigma, \mathbb{Q})$) and the A_i^{2g-2} . Using the classification (7.3), we find that for a very general covering $\tilde{\Sigma} \rightarrow \Sigma$ associated with $p : T \rightarrow G$, the Jacobian $J(\tilde{\Sigma})$ admits no factor of dimension less than $g - 1$.

We summarize this, following [GLLM15, Thm. 1.8]:

Fix a finite group G and an integer $g \geq 3$ such that G can be generated by $g - 1$ elements. There exists a family of pairs

$$(\Sigma, \tilde{\Sigma} \rightarrow \Sigma),$$

where Σ is a smooth complex projective curve of genus g and $\tilde{\Sigma} \rightarrow \Sigma$ is a connected G -covering, with the following property: For a very general $\tilde{\Sigma}$, the Jacobian $J(\tilde{\Sigma})$ admits no factors over \mathbb{Q} of dimension less than $g - 1$.

7.2. Statement and proof of results. Let G be a finite group acting generically freely from the right on P , an abelian variety. We do not assume that G has a fixed point on P . We write $\text{Alb}(P)$ for the Albanese of P , the G -equivariant abelian variety parametrizing the group of translations on P . In other words, P is a G -equivariant principal homogeneous space for $\text{Alb}(P)$.

Fix a base curve B , smooth and projective of genus $g \geq 3$ over k . Let $f : \tilde{B} \rightarrow B$ be a connected G covering space, with G acting from the left. Consider the projection

$$P \times \tilde{B} \rightarrow \tilde{B},$$

where the left-hand-side has induced left G -action

$$\gamma \cdot (p, \tilde{b}) = (p\gamma^{-1}, \gamma\tilde{b}),$$

with quotient $P \times_G \tilde{B}$. Consider the induced morphism

$$\pi_f : P \times_G \tilde{B} \rightarrow B$$

whose fibers, away from the branch locus of f , are geometrically isomorphic to P .

Proposition 23. *Assume that $g > \dim(P) + 1$, $g \geq 3$, and B is of general moduli.*

Suppose that for every f as above, π_f has a section. Then the action of G on P has a fixed point.

Proof. The existence of a section for π_f is equivalent to a G -equivariant morphism

$$\phi_f : \tilde{B} \rightarrow P$$

which factors

$$\tilde{B} \hookrightarrow \text{Pic}^1(\tilde{B}) \rightarrow P.$$

Our assumption – that $\dim(P) < g - 1$ – allows us to apply the results of Section 7.1 to deduce $J(\tilde{B})$ has no factors of dimension $\dim(P)$. It follows that there is no nontrivial G -equivariant homomorphism

$$J(\tilde{B}) \rightarrow \text{Alb}(P).$$

This forces ϕ_f to be constant, which forces the triviality of $\text{Pic}^1(\tilde{B})$. \square

8. APPLICATIONS IN DIMENSION THREE

In this section, we present parametrizations arising from the existence of G -invariant linear subspaces

$$L \subset X \subset \mathbb{P}^{2g+1}.$$

The geometry here is both simpler and richer than the geometry over nonclosed fields. There are more possible Galois actions than actions via automorphisms; not every subgroup of \mathfrak{S}_{2g+2} arises as the automorphisms of a configuration of $2g + 2$ points. On the other hand, if one can find a linear subspace

$$L \simeq \mathbb{P}^r \subset X$$

defined over k , one automatically has subspaces $\mathbb{P}^s \subset X$ for all $s \leq r$. This is not the case in the equivariant context, as the underlying representation may be irreducible.

Throughout this section, k is algebraically closed of characteristic zero.

8.1. Review of surface case. We review the classification of G -actions on smooth intersections of two quadrics in dimension 2. The general strategy for attacking this question uses the G -equivariant minimal model program; the most systematic description may be found in [DI09].

Let $X \subset \mathbb{P}^4$ be a smooth quartic del Pezzo surface with a generically free action of a finite group G . There are 16 lines on X , which are permuted by G . If there exists a G -equivariant collection of disjoint lines, it may be blown down to obtain a del Pezzo surface of larger degree. Del Pezzo surfaces of degree 7, 8, or 9 are equivariantly birational to \mathbb{P}^2 or $\mathbb{P}^1 \times \mathbb{P}^1$. In degrees 5 and 6 there are actions not

birational to actions on homogeneous spaces as above; see Section 8 of [DI09] for more details.

For our purposes – to illustrate the aspects common to all dimensions – we focus on examples with generic automorphism group. Any quartic del Pezzo surface may be written in diagonal form

$$x_0^2 + x_1^2 + x_2^2 + x_3^2 + x_4^2 = a_0 x_0^2 + a_1 x_1^2 + a_2 x_2^2 + a_3 x_3^2 + a_4 x_4^2 = 0$$

which admits a diagonal action of $H = \mu_2^5/\mu_2$. Generically, these are the only automorphisms. Hence we focus on subgroups of

$$H \simeq (\mathbb{Z}/2\mathbb{Z})^4$$

acting via sign changes on coordinates of diagonal quadrics defining X . Consider involutions $\iota \in H$; we present one representative for each conjugacy class:

(1,1,1,1,-1): Here X is a double cover of the quadric surface in \mathbb{P}^3

$$(a_4 - a_0)x_0^2 + (a_4 - a_1)x_1^2 + (a_4 - a_2)x_2^2 + (a_4 - a_3)x_3^2$$

branched over the elliptic curve E cut out by

$$x_0^2 + x_1^2 + x_2^2 + x_3^2 = 0.$$

This surface has no fixed lines and has nontrivial cohomology $H^1(G, \text{Pic}(X))$ (from the curve E , cf. [BP13]) and thus is not G -rational or even stably rational. It is clearly minimal as the orbits of lines consist of eight pairs meeting at points. (These are rulings of the quadric surface tangent to E .) Any G -action on a quartic del Pezzo surface containing this involution is G -irrational.

(1,1,1,-1,-1): Here we have four fixed points given by $\{x_3 = x_4 = 0\}$ and eight orbits of disjoint lines. Thus X is birational to a sextic del Pezzo surface, admitting three conic bundle structures. One of these must be fixed under the involution, thus X is equivariantly birational to $\mathbb{P}^1 \times \mathbb{P}^1$ with linear action.

8.2. Reduction to the case of nonclosed fields. We turn to the case of dimension three. As a corollary of results in Section 7, we have:

Theorem 24. *Let $X \subset \mathbb{P}^5$ be a smooth complete intersection of two quadrics with generically free action of a finite group G . Then X is G -equivariantly birational to \mathbb{P}^3 , with G acting projectively, if and only if there is a G -invariant line on X .*

Proof. The action of G on X admits a canonical linearization on $\omega_X^{-1} = \mathcal{O}_X(2)$. It follows that G acts projectively on $\Gamma(\mathcal{O}_X(1))$ and thus on \mathbb{P}^5 . A central extension

$$1 \rightarrow \mu_2 \rightarrow \tilde{G} \rightarrow G \rightarrow 1$$

acts linearly on \mathbb{P}^5 .

An invariant line corresponds to a two-dimensional \tilde{G} -invariant subspace. A complementary \tilde{G} -stable subspace induces a projection

$$\pi_\ell : X \xrightarrow{\sim} \mathbb{P}^3,$$

where \mathbb{P}^3 admits a linear action of \tilde{G} and a projective action of G .

We turn to the converse statement: Suppose there is a G -equivariant birational map

$$X \dashrightarrow \mathbb{P}^3$$

with G acting projectively.

Let B be a curve of large genus satisfying the conditions of Proposition 23. Construct an isotrivial family $\mathcal{X} \rightarrow B$, with generic fiber isomorphic to X , split over a G -covering $\tilde{B} \rightarrow B$. (The G -action on X induces one on $X \times \tilde{B}$; we take quotients to obtain our isotrivial family.) It is birational over B to a fibration $\mathcal{P} \rightarrow B$ that is generically a \mathbb{P}^3 -bundle. The Tsen-Lang Theorem implies it is birational over B to $\mathbb{P}^3 \times B$. The variety of lines $F_1(\mathcal{X}/B)$ is a principal homogeneous space over an abelian surface, over a dense open subset of B . The main result of [HT21a] implies that $F_1(\mathcal{X}/B) \rightarrow B$ admits a rational section. Proposition 23 implies that the associated action of G on $F_1(X)$ necessarily admits a fixed point, which yields a line $\ell \subset X$ invariant under the G -action. \square

Remark 25. Let X have a G -action as above; we do not assume the existence of a fixed point or invariant line. Nevertheless, we can construct the isotrivial fibration

$$\mathcal{X} \rightarrow B$$

which *always* admits a section by the Tsen-Lang (or Graber-Harris-Starr) Theorem. Thus Proposition 23, stated for abelian varieties, is not valid for other classes of varieties such as rationally-connected varieties.

8.3. Relations with the Burnside formalism. The Burnside group formalism is presented in [KT20]. Let G be a finite group acting generically freely on a smooth projective variety Y . For each nontrivial subgroup $H \subset G$, we record data

- the locus $Z \subset Y$ fixed under H ;
- the action of H on the normal bundle of each component of Z .

The character for the action on the normal bundle is called a *symbol*. Consider how this data changes as we blow up Y along a G -stable smooth subvariety. To obtain a G -birational invariant, we impose equivalences on the possible symbols that may arise, the *blowup relations*.

The simplest approach is to discard information about the components of the fixed locus Z , recording only the representation on the normal bundle. (Here we only retain the dimensions of the components.). See [HKT21, Section 5] for refinements via Grothendieck classes of varieties, and Section 6 of that paper for applications to cubic fourfolds.

Proposition 26. *Let G be a finite group acting generically freely on $X \subset \mathbb{P}^5$, a smooth complete intersection of two quadrics. Suppose there exist*

- an element $g \neq 1$ fixing a hyperplane section $S \subset X$;
- a subgroup $\langle g \rangle \subset H \subset G$ acting on S .

Assume that S is not H -birational to either

- (1) $\mathbb{P}(V)$, a projective representation of H ; or
- (2) $\mathbb{P}(E)$ where E is an H -equivariant rank-two vector bundle over a curve.

Then X is not G -birational to \mathbb{P}^3 with a G -action.

The surface case is addressed in [KT21, Prop. 3.9].

Proof. Suppose we have a G -birational map

$$\rho : \mathbb{P}^3 \dashrightarrow X$$

with G acting projectively on the source. By weak factorization, we may assume it is a composition of blowups and blowdowns along smooth G -stable centers.

Consider the center $Z \rightarrow \mathbb{P}^3$ blown up to obtain S ; it is irreducible, fixed by g , and has an action of H . Since $S \dashrightarrow Z$ is birational, we conclude that S is birational to a divisor in \mathbb{P}^3 with nontrivial stabilizer. The locus in \mathbb{P}^3 with nontrivial stabilizers is a union of linear subspaces, and we are in the first case. If Z is a point then S is birational to the projectivization of the tangent space at that point; again, we are in the first case. Finally, suppose Z is a curve; then S is birational to the projectivization of the normal bundle to Z , putting us in the second case. \square

We turn to other representative examples. We follow Avilov [Avi16], who enumerated actions of finite groups on three-dimensional complete intersections of two quadrics known to be equivariantly birational to projective space, a quadric hypersurface, or a Mori fiber space.

Example 27. Suppose that G admits a subgroup $H \simeq C_2 \times C_2$ acting on X via the diagonal matrices

$$\text{diag}(1, 1, 1, 1, \pm 1, \pm 1).$$

Take

$$g = \text{diag}(1, 1, 1, 1, 1, -1)$$

which fixes a del Pezzo surface S . The residual C_2 action on S was considered in Section 8.1. Since there is nontrivial cohomology, i.e. $H^1(C_2, \text{Pic}(S)) \neq 0$, the two possibilities from Proposition 26 are precluded. It follows that X is not equivariantly rational for any group containing H .

We emphasize that Theorem 24 gives a *stronger* conclusion: If X is G -birational to \mathbb{P}^3 then G acts on $F_1(X)$ with fixed points. In particular, every element of G fixes a point of $F_1(X)$ so G has no elements conjugate to $\pm \text{diag}(1, 1, 1, 1, -1, -1)$. Indeed, these correspond to translates by two-torsion, which act freely. For instance, if $G = \langle (1, 1, 1, 1, -1, -1) \rangle$ the fixed locus on X is an elliptic curve. There are no Burnside invariants available as the relevant symbol groups for $G = C_2$ are zero [HKT21, Section 3.1], [KPT21, Section 12].

Remark 28. It would be interesting to have a general theory – in the context of G -equivariant birational geometry – encompassing both Burnside/symbol-type invariants and obstructions arising from Chow-theoretic principal homogeneous spaces for intermediate Jacobians.

8.4. Rational complete intersections need not have points. Theorem 24 shows that equivariant rationality of $X \subset \mathbb{P}^5$ is governed by the existence of invariant lines on X . When G is cyclic, the existence of a G -invariant line guarantees a point on that line fixed by G . For noncyclic groups, an invariant line need *not* have a fixed point. There are examples of G -rational $X \subset \mathbb{P}^5$ with no fixed points.

Let C denote the complex curve

$$y^2 = x^6 + 1.$$

Its automorphism group contains G , a central extension

$$1 \rightarrow \mu_2 = \langle \iota \rangle \rightarrow G \rightarrow D_{12} \rightarrow 1$$

of the dihedral group D_{12} of order twelve. We write

$$G = \langle \sigma, \tau, \iota : \sigma^6 = \tau^2 = \iota^2 = 1, \tau\sigma\tau^{-1}\sigma = \iota \rangle$$

with action

$$\sigma \cdot (x, y) = (\zeta x, y), \quad \tau \cdot (x, y) = (y/x^3, 1/x), \quad \zeta = e^{2\pi i/6},$$

where ι is the hyperelliptic involution. The induced action on the global sections $\Gamma(\omega_C)$ is

$$\sigma \mapsto \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^2 \end{pmatrix}, \quad \tau \mapsto \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \quad \iota \mapsto \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Actually, $\text{Aut}(C) = G$; see [LMF21, Genus two curve 2916.b.11664.1] for more information.

Consider the quotient of C under the unique cyclic subgroup of order three $\langle \sigma^2 \rangle$, with invariants and equation

$$y, z = x^3 - y^2 = z^2 + 1.$$

Let $\rho : C \rightarrow R$ denote the corresponding degree-three morphism. The induced action on R is generically free via the dihedral group of order eight.

The double cover and ρ give a morphism

$$C \hookrightarrow \mathbb{P}^1 \times R.$$

While R is isomorphic to \mathbb{P}^1 as a variety, the action of G on R is not linear. (The subgroup $\langle \tau, \sigma \rangle$ acts on R as a Klein four-group, with no fixed points; such actions are not linearizable.) However, we do have a central extension

$$1 \rightarrow \mu_2 \rightarrow \tilde{G} \rightarrow G \rightarrow 1$$

and \tilde{G} -representations $V = \Gamma(\omega_C)$ and W such that

$$C \hookrightarrow \mathbb{P}(V) \times \mathbb{P}(W) \hookrightarrow \mathbb{P}(V \otimes W) \simeq \mathbb{P}^3,$$

realizing our curve as a $(2, 3)$ divisor. The linear series of cubic forms vanishing on C gives a birational map

$$\mathbb{P}^3 \xrightarrow{\sim} X \subset \mathbb{P}^5,$$

where X is a smooth complete intersection of two quadrics. This blows up C and blows down $\mathbb{P}(V) \times \mathbb{P}(W)$ via projection to the first factor. The map $X \xrightarrow{\sim} \mathbb{P}^3$ is projection from a G -stable line $\ell \simeq \mathbb{P}(V)$.

We claim that X has no G -fixed points. We know that $\mathbb{P}(V)$ has no fixed points so it suffices to check that $\mathbb{P}^3 = \mathbb{P}(V \otimes W)$ has no fixed points. Looking at σ acting on V , any fixed points would necessarily lie on

$$[1, 0] \times W \text{ or } [0, 1] \times W.$$

However, we have already seen that the dihedral group of order eight acts on R without fixed points.

This analysis also shows that the G -action on X does not linearize to the ambient \mathbb{P}^5 . The map $\text{Br}_G(\text{point}) \rightarrow \text{Br}_G(X)$ is not an isomorphism; its kernel contains $0 \neq$

$\alpha(G, W) \in H^2(G, \mu_2)$ (see Proposition 3). This illustrates the general obstruction analysis in Section 4.1.

This answers a question of Avilov [Avi16, Rem. 2], who asked where X – Case 2(ii) of his Theorem 1 – fits in the equivariant birational classification.

8.5. Another special example. We return to another example highlighted by Avilov – Case 2(iv) of [Avi16, Th. 1] – where G fits into an exact sequence

$$1 \rightarrow C_2^5 \rightarrow G \rightarrow \mathfrak{S}_4 \rightarrow 1.$$

The associated binary sextic form is

$$T_0 T_1 (T_0^4 - T_1^4).$$

The hyperelliptic curve

$$C = \{U^2 = T_0 T_1 (T_0^4 - T_1^4)\}$$

has automorphism group

$$\text{Aut}(C) = \langle \sigma, \tau \rangle,$$

where

$$\sigma(T_0, T_1, U) = (e^{3\pi i/4} T_0, e^{\pi i/4} T_1, U) \quad \tau(T_0, T_1, U) = \left(\frac{1}{\sqrt{2}}(T_1 - T_0), \frac{1}{\sqrt{2}}(T_0 + T_1), U \right).$$

The resulting group has relations

$$\sigma^4 = \iota, \tau^2 = \iota^2 = 1, \iota\tau = \tau\iota, (\sigma\tau)^3 = \iota,$$

where ι is the hyperelliptic involution. It sits in a central extension

$$1 \rightarrow \mu_2 = \langle \iota \rangle \rightarrow \text{Aut}(C) \rightarrow \mathfrak{S}_4 \rightarrow 1.$$

This curve appears as [LMF21, Genus two curve 4096.b.65536.1].

We analyze the induced action on the branch points, using coordinate T_0/T_1 :

$$b_1 = \{0\}, b_2 = \{\infty\}, b_3 = \{1\}, b_4 = \{i\}, b_5 = \{-1\}, b_6 = \{-i\}.$$

The generators act via permutations

$$\sigma \mapsto (3456), \quad \tau \mapsto (13)(25)(46)$$

and the hyperelliptic involution acts trivially. Consider the induced action on

$$\text{Pic}^1(C) \supset C$$

as in (4.1). Note that ι fixes only the solutions to $L^2 = g_2^1$, the 16 points (cf. Section 3)

$$b_1, \dots, b_6, b_1 + b_2 + b_3 - g_2^1 = b_4 + b_5 + b_6 - g_2^1, \dots, b_1 + b_5 + b_6 - g_2^1 = b_2 + b_3 + b_4 - g_2^1.$$

None of these is simultaneously fixed by σ and τ , so $\text{Pic}^1(C)$ admits no fixed point for $\text{Aut}(C)$.

As for the complete intersection, we may take equations

$$X = \{Q_1 = Q_2 = 0\} \subset \mathbb{P}^5,$$

where

$$Q_1 = x_0^2 + x_1^2 + ix_2^2 - x_3^2 - ix_4^2, \quad Q_2 = x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_5^2.$$

Proposition 29. *Suppose that G acts on X so that the induced homomorphism*

$$G \rightarrow \text{Aut}(C)$$

is surjective. Then X is not G -equivariantly birational to \mathbb{P}^3 .

This partly answers a question in [Avi16, Rem. 2]; Avilov asked whether these are equivariantly birational to \mathbb{P}^3 .

Proof. By Theorem 24, if X is birational to \mathbb{P}^3 then $F_1(X)$ admits a fixed point. But then $\text{Pic}^1(C)$ would admit one as well via the squaring map $F_1(X) \rightarrow \text{Pic}^1(C)$. This would contradict the computation above.

The argument works under the weaker hypothesis that the image contains the 2-Sylow subgroup

$$\langle \sigma, \tau\sigma^2\tau \rangle \subset \text{Aut}(C)$$

as $\tau\sigma^2\tau \mapsto (12)(46)$. \square

We may have rationality when smaller groups act. We restrict attention to the automorphism of order eight acting on coordinates by

$$\gamma = \begin{pmatrix} \alpha & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad \alpha = e^{3\pi i/4}.$$

This is chosen so that $\gamma \cdot Q_2 = Q_2$ and

$$\gamma \cdot Q_1 = \alpha^2 x_0^2 + x_2^2 + ix_3^2 - x_4^2 - ix_1^2 = -iQ_1.$$

Thus we may interpret γ as a lift of σ^{-1} .

What are the fixed points? The action on the underlying space via the contragredient representation is:

$$\begin{pmatrix} \alpha^7 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

with eigenvectors

$$\begin{aligned} 1 &: [0, 0, 0, 0, 0, 1] \notin X \\ \alpha &: p_1 := [0, 1, \alpha^7, \alpha^6, \alpha^5, 0] \in X \\ \alpha^3 &: [0, 1, \alpha^5, \alpha^2, \alpha^7, 0] \notin X \\ \alpha^5 &: p_2 := [0, 1, \alpha^3, \alpha^6, \alpha, 0] \in X \\ \alpha^7 &: [1, 0, 0, 0, 0, 0], [0, 1, \alpha, \alpha^2, \alpha^3, 0] \notin X. \end{aligned}$$

The last eigenspace is isotropic for $\{Q_2 = 0\}$ and thus meets X in two points $p_3, p_4 := [\pm 2i, 1, \alpha, \alpha^2, \alpha^3, 0]$. The span of p_1 and p_2 is also isotropic for $\{Q_2 = 0\}$.

Proposition 30. *The lines $\ell(p_2, p_3)$ and $\ell(p_2, p_4)$ are invariant under the action.*

Note however that

$$\gamma^4 = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

acts on the variety of lines on X with 16 fixed points, i.e., the lines on the quartic del Pezzo surface $X \cap \{x_5 = 0\}$.

This example is instructive in that the group actions are not associated with Weyl groups:

- $W(D_6)$ is realized as the signed permutation matrices with an even number of (-1) signs, a semidirect product

$$W(D_6) \simeq \mathfrak{S}_6 \ltimes C_2^5,$$

where C_2^5 is the diagonal matrices;

- the semidirect product

$$\mathfrak{S}_6 \ltimes C_2^6 / \text{Diagonal},$$

interpreted as a Weyl group for the projective orthogonal group.

These are not the same; the Weyl group has a nontrivial central element

$$(-1, \dots, -1)$$

whereas the latter group has *no* nontrivial central elements.

However, *neither* of these actions coincide with our situation! Both lack an element γ of order eight, sitting over a four-cycle of \mathfrak{S}_6 , with $\gamma^4 \in H'$ (the 2-elementary group of diagonal automorphisms) having nontrivial determinant. The automorphism group $\text{Aut}(X)$ is NOT a semidirect product of $\text{Aut}(D \subset P^1)$ and H' , which admits NO element γ mapping to

$$\begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix}$$

and whose fourth power is diagonal with entries $(-1, -1, -1, -1, -1, 1)$. Indeed, the candidate in the Weyl group

$$\tilde{\gamma} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

induces

$$\tilde{\gamma}^4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

which has the wrong diagonal entries.

The rationality of these specific examples may be deduced in terms of the group actions:

Proposition 31. *Let γ be an automorphism of order eight on X such that γ^4 fixes a smooth hyperplane section $S \subset X$ and the induced action on S is by an element acting on the degeneracy locus (of both X and S) as a four-cycle. Then S admits a line invariant under γ .*

Proof. The fixed locus of H' corresponds to a del Pezzo S of degree 4 with action associated with the upper 5×5 matrix. Its lines correspond to weights in the D_5 root system:

$$-e_i - e_j - e_k - e_l - e_m, e_i + e_j - e_k - e_l - e_m, e_i + e_j + e_k + e_l - e_m.$$

With respect to the standard basis of the Picard group $\{L, E_1, E_2, E_3, E_4, E_5\}$ projected into $\text{Pic}(S)/\mathbb{Z}K_S$ we have

$$L - E_i \mapsto e_i, \quad 2L - E_j - E_k - E_l - E_m \mapsto -e_i.$$

The induced automorphism has order four. The only possible elements of $W(D_5)$ of order four fix a line. For example, the signed permutations

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}$$

are all conjugate in $W(D_5)$ so it suffices to consider the first. This acts on $\text{Pic}(S)$ by

$$\begin{aligned} L &\mapsto 2L - E_1 - E_3 - E_4 \\ E_1 &\mapsto L - E_3 - E_4 \\ E_2 &\mapsto L - E_1 - E_4 \\ E_3 &\mapsto L - E_1 - E_3 \\ E_4 &\mapsto E_2 \\ E_5 &\mapsto E_5 \end{aligned}$$

which leaves the lines E_5 and $2L - E_1 - E_2 - E_3 - E_4 - E_5$ invariant. \square

The assumption that the induced permutation of the degeneracy locus is a four-cycle is essential. The element

$$\gamma_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

acts via

$$\begin{aligned} L &\mapsto 2L - E_1 - E_4 - E_5 \\ E_1 &\mapsto L - E_4 - E_5 \\ E_2 &\mapsto L - E_1 - E_5 \\ E_3 &\mapsto L - E_1 - E_4 \\ E_4 &\mapsto E_3 \\ E_5 &\mapsto E_2 \end{aligned}$$

with orbits

$$\begin{aligned} E_1 &\mapsto L - E_4 - E_5 \mapsto 2L - E_1 - E_2 - E_3 - E_4 - E_5 \mapsto L - E_2 - E_3 \\ E_2 &\mapsto L - E_1 - E_5 \mapsto L - E_1 - E_2 \mapsto E_5 \\ E_3 &\mapsto L - E_1 - E_4 \mapsto L - E_1 - E_3 \mapsto E_4 \\ L - E_2 - E_4 &\mapsto L - E_3 - E_4 \mapsto L - E_3 - E_5 \mapsto L - E_2 - E_5 \end{aligned}$$

However, γ_1 maps to a product of two transposition in \mathfrak{S}_6 .

REFERENCES

- [Avi16] A. Avilov. Automorphisms of threefolds that can be represented as an intersection of two quadrics. *Mat. Sb.*, 207(3):3–18, 2016.
- [BCDP18] Jérémie Blanc, Ivan Cheltsov, Alexander Duncan, and Yuri Prokhorov. Finite quasisimple groups acting on rationally connected threefolds, 2018. [arXiv:1809.09226](https://arxiv.org/abs/1809.09226).
- [Bea83] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18(4):755–782 (1984), 1983.
- [BGW17] Manjul Bhargava, Benedict H. Gross, and Xiaoheng Wang. A positive proportion of locally soluble hyperelliptic curves over \mathbb{Q} have no point over any odd degree extension. *J. Amer. Math. Soc.*, 30(2):451–493, 2017. With an appendix by Tim Dokchitser and Vladimir Dokchitser.
- [BHT15] Arend Bayer, Brendan Hassett, and Yuri Tschinkel. Mori cones of holomorphic symplectic varieties of K3 type. *Ann. Sci. Éc. Norm. Supér. (4)*, 48(4):941–950, 2015.
- [BL04] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 302 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2004.
- [BP13] Fedor Bogomolov and Yuri Prokhorov. On stable conjugacy of finite subgroups of the plane Cremona group, I. *Cent. Eur. J. Math.*, 11(12):2099–2105, 2013.
- [Bri18] Michel Brion. Linearization of algebraic group actions. In *Handbook of group actions. Vol. IV*, volume 41 of *Adv. Lect. Math. (ALM)*, pages 291–340. Int. Press, Somerville, MA, 2018.
- [BW19] Olivier Benoist and Olivier Wittenberg. Intermediate Jacobians and rationality over arbitrary fields. *Annales scientifiques de l’École normale supérieure*, to appear, 2019.
- [BW20] Olivier Benoist and Olivier Wittenberg. The Clemens-Griffiths method over non-closed fields. *Algebr. Geom.*, 7(6):696–721, 2020.

- [CGR06] V. Chernousov, P. Gille, and Z. Reichstein. Resolving G -torsors by abelian base extensions. *J. Algebra*, 296(2):561–581, 2006.
- [CTS21] Jean-Louis Colliot-Thélène and Alexei Skorobogatov. *The Brauer–Grothendieck Group*, volume 71 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer International Publishing, 2021.
- [DI09] I. V. Dolgachev and V. A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 443–548. Birkhäuser Boston, Boston, MA, 2009.
- [Don80] Ron Donagi. Group law on the intersection of two quadrics. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 7(2):217–239, 1980.
- [DR77] U. V. Desale and S. Ramanan. Classification of vector bundles of rank 2 on hyperelliptic curves. *Invent. Math.*, 38(2):161–185, 1976/77.
- [FMOS21] Laure Flapan, Emanuele Macrì, Kieran G. O’Grady, and Giulia Saccà. The geometry of antisymplectic involutions, I. *Mathematische Zeitschrift*, to appear, 2021.
- [FW71] A. Fröhlich and C. T. C. Wall. Equivariant Brauer groups in algebraic number theory. In *Colloque de Théorie des Nombres (Univ. de Bordeaux, Bordeaux, 1969)*, pages 91–96. Bull. Soc. Math. France, Mém No. 25, 1971.
- [FW00] A. Fröhlich and C. T. C. Wall. Equivariant Brauer groups. In *Quadratic forms and their applications (Dublin, 1999)*, volume 272 of *Contemp. Math.*, pages 57–71. Amer. Math. Soc., Providence, RI, 2000.
- [GLLM15] Fritz Grunewald, Michael Larsen, Alexander Lubotzky, and Justin Malestein. Arithmetic quotients of the mapping class group. *Geom. Funct. Anal.*, 25(5):1493–1542, 2015.
- [HKT21] Brendan Hassett, Andrew Kresch, and Yuri Tschinkel. Symbols and equivariant birational geometry in small dimensions. In *Rationality of Varieties*, volume 342 of *Progress in Mathematics*, pages 201–236. Birkhäuser, Cham., 2021.
- [HT13] Brendan Hassett and Yuri Tschinkel. Hodge theory and Lagrangian planes on generalized Kummer fourfolds. *Mosc. Math. J.*, 13(1):33–56, 189, 2013.
- [HT21a] Brendan Hassett and Yuri Tschinkel. Cycle class maps and birational invariants. *Communications on Pure and Applied Mathematics*, 74(12):2675–2698, 2021.
- [HT21b] Brendan Hassett and Yuri Tschinkel. Rationality of complete intersections of two quadrics over nonclosed fields. *Enseign. Math.*, 67(1-2):1–44, 2021. With an appendix by Jean-Louis Colliot-Thélène.
- [KKLV89] Friedrich Knop, Hanspeter Kraft, Domingo Luna, and Thierry Vust. Local properties of algebraic group actions. In *Algebraische Transformationsgruppen und Invariantentheorie*, volume 13 of *DMV Sem.*, pages 63–75. Birkhäuser, Basel, 1989.
- [KKV89] Friedrich Knop, Hanspeter Kraft, and Thierry Vust. The Picard group of a G -variety. In *Algebraische Transformationsgruppen und Invariantentheorie*, volume 13 of *DMV Sem.*, pages 77–87. Birkhäuser, Basel, 1989.
- [KP21] Alexander Kuznetsov and Yuri Prokhorov. Rationality of Fano threefolds over nonclosed fields. In *Rationality of Varieties*, volume 342 of *Progress in Mathematics*, pages 249–290. Birkhäuser, Cham., 2021.
- [KPT21] Maxim Kontsevich, Vasily Pestun, and Yuri Tschinkel. Equivariant birational geometry and modular symbols. *Journal of the European Mathematical Society*, online first, 2021.
- [KT20] Andrew Kresch and Yuri Tschinkel. Equivariant birational types and Burnside volume. *Annali della Scuola Normale Superiore di Pisa*, to appear, 2020.
- [KT21] Andrew Kresch and Yuri Tschinkel. Equivariant Burnside groups and representation theory, 2021. [arXiv:2108.00518](https://arxiv.org/abs/2108.00518).
- [Lee] D. Leep. The Amer–Brumer theorem over arbitrary fields. <http://www.ms.uky.edu/~leep/Preprints.html>.
- [LLSvS17] Christian Lehn, Manfred Lehn, Christoph Sorger, and Duco van Straten. Twisted cubics on cubic fourfolds. *J. Reine Angew. Math.*, 731:87–128, 2017.
- [LMF21] The LMFDB Collaboration. The L-functions and modular forms database. <http://www.lmfdb.org>, 2021. [Online; accessed 23 July 2021].
- [Mor15] Dave Witte Morris. *Introduction to arithmetic groups*. Deductive Press, 2015.

- [Rei72] Miles Reid. *The complete intersection of two or more quadrics*. PhD thesis, Trinity College, Cambridge, 1972. available at <http://homepages.warwick.ac.uk/~masda/3folds/qu.pdf>.
- [RY00] Zinovy Reichstein and Boris Youssin. Essential dimensions of algebraic groups and a resolution theorem for G -varieties. *Canad. J. Math.*, 52(5):1018–1056, 2000. With an appendix by János Kollár and Endre Szabó.
- [RY02] Z. Reichstein and B. Youssin. A birational invariant for algebraic group actions. *Pacific J. Math.*, 204(1):223–246, 2002.
- [Wan18] Xiaoheng Wang. Maximal linear spaces contained in the base loci of pencils of quadrics. *Algebr. Geom.*, 5(3):359–397, 2018.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, BOX 1917 151 THAYER STREET PROVIDENCE, RI 02912, USA

Email address: `brendan_hassett@brown.edu`

COURANT INSTITUTE, NEW YORK UNIVERSITY, NEW YORK, NY 10012, USA

SIMONS FOUNDATION, 160 FIFTH AVENUE, NEW YORK, NY 10010, USA

Email address: `tschinkel@cims.nyu.edu`