

## Colloquium: Machine learning in nuclear physics

Amber Boehlein<sup>1</sup>, Markus Diefenthaler<sup>1</sup>, Nobuo Sato,  
Malachi Schram<sup>1</sup>, and Veronique Ziegler

*Thomas Jefferson National Accelerator Facility,  
12000 Jefferson Avenue, Newport News, Virginia 23606, USA*

Cristiano Fanelli<sup>1</sup>

*Laboratory for Nuclear Science and Institute for Artificial Intelligence and Fundamental  
Interactions, Massachusetts Institute of Technology,  
Cambridge, Massachusetts 02139, USA*

Morten Hjorth-Jensen<sup>1</sup>

*Facility for Rare Isotope Beams and Department of Physics and Astronomy,  
Michigan State University, East Lansing, Michigan 48824, USA  
and Department of Physics and Center for Computing in Science Education,  
University of Oslo, N-0316 Oslo, Norway*

Tanja Horn<sup>1</sup>

*Department of Physics, The Catholic University of America, Washington, D.C. 20064, USA  
and Thomas Jefferson National Accelerator Facility, 12000 Jefferson Avenue,  
Newport News, Virginia 23606, USA*

Michelle P. Kuchera

*Department of Physics and Department of Mathematics and Computer Science,  
Davidson College, Davidson, North Carolina 28035, USA*

Dean Lee<sup>1</sup>, Witold Nazarewicz<sup>1</sup>, and Peter Ostroumov

*Facility for Rare Isotope Beams and Department of Physics and Astronomy,  
Michigan State University, East Lansing, Michigan 48824, USA*

Kostas Orginos

*Department of Physics, William & Mary, Williamsburg 23185, Virginia, USA  
and Thomas Jefferson National Accelerator Facility,  
12000 Jefferson Avenue, Newport News, Virginia 23606, USA*

Alan Poon<sup>1</sup> and Xin-Nian Wang

*Nuclear Science Division, Lawrence Berkeley National Laboratory,  
1 Cyclotron Road, Berkeley, California 94720, USA*

Alexander Scheinker<sup>1</sup>

*Accelerator Operations and Technology Division Applied Electrodynamics Group,  
Los Alamos National Laboratory, Los Alamos, New Mexico 87544, USA*

Michael S. Smith<sup>1</sup>

*Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 37831-6354, USA*

Long-Gang Pang<sup>1</sup>

*Key Laboratory of Quark and Lepton Physics, Institute of Particle Physics,  
Central China Normal University, Wuhan 430079, China*



(published 8 September 2022)

Advances in machine learning methods provide tools that have broad applicability in scientific research. These techniques are being applied across the diversity of nuclear physics research topics, leading to advances that will facilitate scientific discoveries and societal applications. This Colloquium provides a snapshot of nuclear physics research, which has been transformed by machine learning techniques.

DOI: [10.1103/RevModPhys.94.031003](https://doi.org/10.1103/RevModPhys.94.031003)

## CONTENTS

I. Introduction	2
II. Machine Learning for Nuclear Physics in Broad Strokes	5
III. Nuclear Theory	5
A. Low-energy nuclear theory	5
1. Early applications of machine learning	5
2. Predicting missing data with ML	5
3. Properties of heavy nuclei and nuclear density functional theory	5
4. Nuclear properties with ML	5
5. Nuclear shell-model applications	5
6. Effective field theory and $A$ -body systems	5
7. Nuclear reactions	5
8. Neutron star properties and nuclear matter equation of state	5
B. Medium-energy nuclear theory	6
1. Bayesian inference	6
2. Simultaneous extraction paradigm	6
3. LQCD and experimental global analysis	6
C. Lattice QCD	7
1. The sign problem at nonzero density	7
2. Ensemble generation	7
3. Correlation function estimators	7
4. Miscellany	7
D. High-energy nuclear theory	8
1. Bayesian inference	8
2. Inversion problems with ML	8
3. Other applications of ML methods	8
IV. Experimental Methods	9
A. Streaming detector readout	9
B. Reconstruction and analysis	9
1. Charged particle tracking	9
2. Calorimetry	9
3. Particle identification	9
4. Event and signal classification	9
5. Event reconstruction	9
6. Spectroscopy	9
C. Experimental design	10
1. Design for detector systems	10
2. Interface with theory	10
V. Accelerator Science and Operations	10
A. ML-based surrogate models for accelerator models	10
B. Anomaly detection and classification	10
C. Control optimization	10
D. Adaptive ML for nonstationary systems	10
VI. Nuclear Data	11
A. Overhauling the nuclear data pipeline	11
B. Improving compilations and evaluations	11
C. Building emulators and surrogate models	11
VII. Summary and Perspectives	12
Acknowledgments	12
References	12

## I. INTRODUCTION

This Colloquium represents an up-to-date summary of work in the application of machine learning (ML) in nuclear science, covering topics in nuclear theory, experimental methods, accelerator technology, and nuclear data. An overview of the use of artificial intelligence (AI) techniques in nuclear physics that aimed at identifying commonalities and needs was provided by [Bedaque \*et al.\* \(2021\)](#).

Nuclear physics is a well-established field, with more than a century of fundamental discoveries covering a large span of degrees of freedom, energy scales, and length scales ranging from our basic understanding of fundamental constituents of matter to the structure of stars and the synthesis of the elements in the cosmos; see Fig. 1. Experiments produce data volumes that range in complexity and heterogeneity, thereby posing enormous challenges to their design, their execution, and the statistical data analysis.

Theoretical modeling of nuclear properties is, in most physical cases of interest, limited by the large amount of degrees of freedom in quantum-mechanical calculations. The analysis of experimental data and the theoretical modeling of nuclear systems aim, as is the case in all fields of physics, at uncovering the basic laws of motion in order to make predictions and estimations, as well as at finding correlations and causations for strongly interacting matter. The broad aims of nuclear physics as a field correspond to a highly distributed scientific enterprise. Experimental efforts utilize many laboratories worldwide, each with unique operation, data acquisition, and analysis methods. Similarly, the scales of focus spanned in theoretical nuclear physics lead to broad needs for algorithmic methods and uncertainty quantification. These efforts, utilizing arrays of data types across size and energy scales, create an ideal environment for applications of ML methods.

## II. MACHINE LEARNING FOR NUCLEAR PHYSICS IN BROAD STROKES

Statistics, data science, and ML form important fields of research in modern science. They describe how to learn and make predictions from data and enable the extraction of key information about physical processes and the underlying scientific laws based on large datasets. As such, recent advances in ML capabilities are being applied to advance scientific discoveries in the physical sciences ([Carleo \*et al.\*, 2019](#); [Deiana \*et al.\*, 2021](#)).

Ideally, ML represents the science of building models to perform a task without the instructions being explicitly programmed. This approach introduces in practice a




FIG. 1. Schematic relationships between the topics discussed in this Colloquium. The diagram emphasizes the close connections between theory, computations (both computational science and data science, as well as many elements from computer science), and experiments.

hierarchy of mathematical operations that enable the computer to learn complicated concepts by building them out of simpler ones. In terms of a graphical representation, this can be visualized as a deep network of training and learning operations, often referred to simply as deep learning (Goodfellow, Bengio, and Courville, 2016).

There are many ML approaches; they are often split into two main categories, supervised and unsupervised. In supervised learning, training data are labeled and one lets a specific ML algorithm learn and deduce patterns in the datasets.

This allows one to make predictions about future events and/or data not included in the training set. On the other hand, unsupervised learning is a method for finding patterns and relationship in datasets without any prior knowledge of the system. Many researchers also operate using a third category called reinforcement learning. This is a paradigm of learning inspired by behavioral psychology, where actions are learned to maximize reward. One may encounter reinforcement learning accompanied by supervised deep learning methods such as deep artificial neural network (ANN). Furthermore, what is often referred to as semi-supervised learning entails the development of algorithms that aim at learning from a dataset that includes both labeled and unlabeled data.

Another way to categorize ML tasks is to consider the desired output of a system. Some of the most common tasks are as follows (Bishop, 2006; Hastie, Tibshirani, and Friedman, 2009; Murphy, 2012; Goodfellow, Bengio, and Courville, 2016; Mehta *et al.*, 2019; Cranmer, Seljak, and Terao, 2022).

**Classification.**—Outputs are divided into two or more classes. The goal is to produce a model that assigns inputs into one of these classes. An example is to

identify digits based on pictures of handwritten numbers.

**Regression.**—Finding a functional relationship between an input dataset and a reference dataset. The goal is to construct a function that maps input data into continuous output values.

**Clustering.**—Data are divided into groups with certain common traits without knowledge of the different groups beforehand. This ML task falls under the category of unsupervised learning.

**Generation.**—Building a model to generate data that are akin to a training dataset in both examples and distributions of examples. Most generative models are types of unsupervised learning.

In Table I we list many of the methods encountered in this Colloquium with their respective abbreviations.

The methods that we cover here have three central elements in common, irrespective of whether we deal with supervised, unsupervised, or semisupervised learning. The first element is some dataset (which can be subdivided into training, validation, and test data), while the second element is a model, which is normally a function of some parameters to be determined by the chosen optimization process. The model reflects our prior knowledge of the system (or a lack thereof). As an example, if we know that our data show a behavior similar to what would be predicted by a polynomial, fitting the data to a polynomial of some degree would determine our model. The last element is a so-called cost (or loss, error, penalty, or risk) function that allows us to present an estimate on how good our model is in reproducing the data that it is supposed to train. This is the function that is optimized in order to obtain the best prediction for the data under study. The simplest cost function in a regression analysis (fitting a continuous function to the data) is the so-called mean squared

TABLE I. Table of ML methods discussed in this Colloquium with an indication of the main type of learning (S, supervised; U, unsupervised; semi-S, semisupervised).

Acronym	Method	Description	Learning type
AE, VAE	Autoencoders, Variational autoencoders	ANN capable of learning efficient representations of the input data without any supervision	U
ANN	Artificial neural network	Models for learning defined by connected units (or nodes) and hidden layers with well-defined inputs and outputs	S
BED	Bayesian experimental design	Bayesian inference for experimental design	S
BM	Boltzmann machine	Generative ANN that can learn a probability distribution from sets of changing inputs	U
BMA, BMM	Bayesian model averaging, Bayesian model mixing	Bayesian inference applied to model selection or the combined estimation, or performed over a mixture model	S
BNN	Bayesian neural network	ANN where the parameters of the network are represented by probabilities learned by Bayesian inference	S
BO	Bayesian optimization	Optimization of functions without an <i>a priori</i> knowledge of functional forms.	S and semi-S
CNN	Convolutional neural network	ANN where convolution is used to reduce dimensionalities	S
EMB	Ensemble methods and boosting	Methods based on collections of decision trees as simple learners	S
GAN	Generative adversarial network	System of two ANNs where a generative network generates outputs while a discriminative network evaluates them	U
GP	Gaussian process	Collection of random variables that have a joint Gaussian distribution used in Bayesian inference	Semi-S
KNN	<i>k</i> -nearest neighbors	Nonparametric method where inputs consist of the <i>k</i> closest training examples in a dataset	S
KR	Kernel regression	Extension of linear regression methods to include nonlinear function kernels	S
LR	Logistic regression	Convex optimization method based on maximum likelihood estimate for classification problems	S
LSTM	Long short-term memory	RNN capable of learning long-term dependencies	S
PCA	Principal component analysis	Dimensionality reduction technique based on retaining the largest eigenvalues of the covariance matrix	U
REG	Linear regression	Linear algebra methods used for modeling continuous functions in terms of their explanatory variables	S
RL	Reinforcement learning	Learning achieved by trial and error of desired and undesired events	Neither S nor U
RNN	Recurrent neural network	ANN where connections between nodes allow for temporal dynamic behavior	S
SVM	Support vector machine	Convex optimization techniques with efficient ways to distinguish features in datasets	S

error function, while for a binary classification problem the so-called cross entropy is widely used; see [Bishop \(2006\)](#), [Hastie, Tibshirani, and Friedman \(2009\)](#), [Murphy \(2012\)](#), and [Goodfellow, Bengio, and Courville \(2016\)](#) for more details. We henceforth refer to this element as the assessment of a given method.

Traditionally, the field of ML has mainly focused on predictions and correlations. In ML and prediction-based tasks, we are often interested in developing algorithms that are capable of learning patterns from existing data in an automated fashion, and then using these learned patterns to make predictions or assessments of new data. In some cases, our primary concern is the quality of the predictions or assessments, with perhaps less focus on the underlying patterns (and probability distributions) that were learned in order to make these predictions. However, in many nuclear physics studies, we are equally interested in being able to estimate errors and find causations. In this Colloquium, we

emphasize the role of predictions and correlations as well as error estimation and causations in statistical learning and ML. For general references on these topics and discussions of frequentist and Bayesian methodologies, see [Bishop \(2006\)](#), [Hastie, Tibshirani, and Friedman \(2009\)](#), [Murphy \(2012\)](#), [Gelman \*et al.\* \(2014\)](#), [Goodfellow, Bengio, and Courville \(2016\)](#), and [Myren and Lawrence \(2021\)](#).

Since the aim of this Colloquium is to give an overview of present usage and research of ML in nuclear physics, we utilize material from several sources on the topic, such as the previously mentioned textbooks and recent reviews ([Carleo \*et al.\*, 2019](#); [Mehta \*et al.\*, 2019](#)). We also mention the theory of Bayesian experimental design (BED) ([Chaloner and Verdinelli, 1995](#); [Liepe \*et al.\*, 2013](#)), a theory that is tailored for making optimal decisions under uncertainty.

During the last few years there has been a surge of interest in applying different ML and Bayesian methods in nuclear physics. In particular, a Bayesian approach has gained large

traction since an estimation of errors plays a major role in theoretical studies, such as reliable determinations of parameters entering models for nuclear forces and density functionals. Similarly, in quantum-mechanical few- and many-body studies, a number of research groups have implemented ML-based techniques in order to handle complicated correlations and exploding numbers of degrees of freedom. These studies cover a large set of approaches, ranging from applications of deep learning methods, such as ANNs and restricted BMs for solving the many-particle Schrödinger equation, to ANNs in many-body methods, with the aim of learning many-body correlations. Similarly, there have been several attempts to use ML approaches to extract information about correlations in field theory and lattice quantum chromodynamics (LQCD), ranging from attempts to circumvent the fermion sign problem to learning fermion propagators. For recent ML applications to nuclear theory, see Sec. III.

In experimental nuclear physics, with increasing degrees of freedom and complexity, one faces many of the same challenges as in nuclear theory. As discussed in Sec. IV, ML approaches offer a number of optimization strategies to handle this surge in dimensionality. Many current nuclear physics experiments produce large amounts of data in excess of terabytes, requiring the use of fast algorithms for tractable data collection and analysis. Machine learning methods such as anomaly detection allow for an exploration of data for unforeseen phenomena. Additionally, labeled data may not be available, due to either an inability to label data or a lack of knowledge of the types and behaviors of the reactions taking place. The latter are normally needed in order to generate simulated data that one can use in the training process. Machine learning techniques play also a considerable role in accelerator science and operations and nuclear data science; see Secs. V and VI, respectively.

After these general remarks about ML in nuclear physics, we move on to a description of ongoing and planned research where many of these approaches are applied to multidimensional problems, large datasets, detection and prognostication, design optimization, and real-time operational control.

### III. NUCLEAR THEORY

The aim of this section is to give the reader an overview of recent progress and future research directions in ML approaches and methods applied to nuclear theory. During the last few years, ML methods have been applied to essentially all length and energy scales of interest for nuclear theory, spanning from theories for the strong force to the equation of state for neutron stars. We start our discussion with low-energy nuclear theory, moving up to medium energies, and to high-energy nuclear theory and lattice quantum chromodynamics.

#### A. Low-energy nuclear theory

##### 1. Early applications of machine learning

In a pioneering paper, the St. Louis–Urbana Collaboration (Gazula, Clark, and Bohr, 1992) successfully carried out

computer experiments based on ANNs to study global nuclear properties across the nuclear landscape, including dripline locations, atomic masses, separation energies, and location of shell-stabilized superheavy nuclei. Their work recognized the potential of using ML techniques to describe the variety of nuclear behavior: “The field of nuclear physics, with a wealth of data reflecting both the fundamental principles of quantum mechanics and the behavior of strong, electromagnetic, and weak interactions on the Fermi scale of distances, offers especially fertile ground for testing and exploiting the new concept of adaptive phenomenological analysis based on neural networks.” They concluded that, “Significant predictive ability is demonstrated, opening the prospect that neural networks may provide a valuable new tool for computing nuclear properties and, more broadly, for phenomenological description of complex many-body systems.” Encouraged by the ability of the ANNs to capture the patterns and irregularities of nuclear observables, they extended their investigations to systematics of nuclear spins and parities (Gernoth *et al.*, 1993). The SVM study of nuclear masses, beta-decay lifetimes, and spins and parities of nuclear ground states was reported by Clark and Li (2006) and Costiris *et al.* (2008), and the application of ANNs to beta decays was carried out by Costiris *et al.* (2009).

#### 2. Predicting missing data with ML

Oftentimes it is necessary to be able to accurately calculate observables that have not been measured to supplement the existing databases. To provide quantified interpolations and extrapolations of nuclear data, nuclear models augmented by modern ML techniques have been used. Examples include studies of nuclear masses with EMB (Carnini and Pastore, 2020), KR (Wu and Zhao, 2020; Wu, Guo, and Zhao, 2021), and ANNs (Yüksel, Soydaner, and Bahtiyar, 2021; Lovell *et al.*, 2022); calculations of the nuclear charge radii using ANNs (Wu *et al.*, 2020); estimation of  $\alpha$ -decay rates using EMB and ANNs (Saxena, Sharma, and Saxena, 2021); estimation of fission yields using mixture-density ANNs (Lovell *et al.*, 2019; Lovell, Mohan, and Talou, 2020), BNNs (Wang *et al.*, 2019; Qiao *et al.*, 2021; Wang and Pei, 2021; Wang, Pei *et al.*, 2021), and KNNs (Tong, He, and Yan, 2021); estimation of the total fusion cross sections using ANNs (Akkoyun, 2020); predictions for the isotopic cross sections in proton-induced spallation reactions using BNNs (Ma, Peng *et al.*, 2020); and estimation of gamma strength functions using BO (Heim *et al.*, 2020).

#### 3. Properties of heavy nuclei and nuclear density functional theory

Kohn-Sham density functional theory (DFT) (Kohn and Sham, 1965) is the basic computational approach to multi-electron systems and there is a rich literature on ML applications in the field of the electron DFT (Hautier *et al.*, 2010; Carleo *et al.*, 2019; Ryczko, Strubbe, and Tamblyn, 2019; Schleder *et al.*, 2019; Moreno, Carleo, and Georges, 2020; Nagai, Akashi, and Sugino, 2020). Nuclear DFT, rooted in the self-consistent mean-field approach (Bender, Heenen, and Reinhard, 2003; Schunck, 2019; Yang and Piekarewicz, 2020), is the basic computational framework for the global




FIG. 2. Bayesian calibration of energy density functionals. Univariate and bivariate marginal estimates of the posterior distribution for the 12-dimensional parameter vector of the UNEDF1 EDF. The upper triangle corresponds to the original UNEDF1 dataset; the lower triangle shows the posterior distribution found when the more recent mass measurements are included. From McDonnell *et al.*, 2015.

modeling of all nuclei, including complex exotic nuclei far from stability. An effective interaction in DFT is given by the energy density functional (EDF), whose parameters are adjusted to experimental data. Over the past decade, better and more refined EDFs have been developed with increasingly complex and computationally expensive computer models, often involving BO and ANN ML (Goriely and Capote, 2014; Higdon *et al.*, 2015; McDonnell *et al.*, 2015; Navarro Pérez *et al.*, 2018; Kejzlar *et al.*, 2020; Scamps *et al.*, 2020; Schunck *et al.*, 2020; Bollapragada *et al.*, 2021; Zhang, Feng, and Chen, 2021). Figure 2 shows the posterior distributions for the parameters of the UNEDF1 EDF obtained by McDonnell *et al.* (2015) by means of the Bayesian model calibration. These distributions have been used to provide uncertainty quantification (UQ) on UNEDF1 model predictions, in particular, the  $r$ -process abundance pattern (Sprouse *et al.*, 2020), the nuclear matter equation of state (Du, Steiner, and Holt, 2019), and neutron star properties (Al-Mamun *et al.*, 2021).

Since global DFT computations are expensive, a promising avenue for ML applications is the emulation of DFT results (Akkoyun *et al.*, 2013; Lasseri *et al.*, 2020; Schunck, Quinlan, and Bernstein, 2020; Nandi *et al.*, 2021). Figure 3 shows the results of ANN calculations by Lasseri *et al.* (2020). A committee of ANNs trained on a set of 210 nuclei is capable of predicting the ground-state and excited-state energies of more than 1800 atomic nuclei with significantly less computational cost than full DFT calculations.

#### 4. Nuclear properties with ML

One can improve the predictive power of a given nuclear model by comparing its predictions to existing data. Here a powerful strategy is to estimate model residuals, i.e., deviations between experimental and calculated observables, by developing an emulator using a training set of observables taken from experiment or other theory. An emulator




FIG. 3. DFT emulator with ANNs. (a) Database nuclei (gray region) as a function of  $N$  and  $Z$ . Nuclei included in the 10% training dataset obtained by active learning are marked as dots. (b) Root mean square deviation between the total energy for the testing dataset calculated in DFT ( $E_{HFB}$ ) and with the committee of ANNs. Adapted from Lasseri *et al.*, 2020.

can be constructed by employing Bayesian approaches, such as BNNs and GPs. Global surveys of nuclear observables employing such a strategy were published by Utama, Piekarewicz, and Prosper (2016), Utama and Piekarewicz (2017, 2018), Niu and Liang (2018), Niu, Fang, and Niu (2019), Rodríguez *et al.* (2019b), Ma, Su *et al.* (2020), Pastore *et al.* (2020), and Gao *et al.* (2021) (extrapolations of nuclear masses with NNs); by Neufcourt *et al.* (2018) and Shelley and Pastore (2021) (extrapolations of nuclear masses with GPs); by Utama, Chen, and Piekarewicz (2016) and Wu *et al.* (2020) (studies of nuclear radii); by Niu *et al.* (2019) and Wu *et al.* (2021) (studies of beta-decay rates); and by Rodríguez *et al.* (2019a) (studies of alpha-decay rates). See also Liu *et al.* (2021) and Pastore and Carnini (2021) for more discussions on ANN extrapolations.

By considering several global models and the most recent data, one can apply the powerful techniques of Bayesian model averaging (BMA) and Bayesian model mixing (BMM) to assess model-related uncertainties in the multimodel context (Phillips *et al.*, 2021). Examples of recent model-mixed predictions using BMA include analysis of the neutron dripline in the Ca region (Neufcourt *et al.*, 2019), studies of proton dripline and proton radioactivity (Neufcourt *et al.*, 2020a), quantification of the particle stability of nuclei (Neufcourt *et al.*, 2020b), combination of models calibrated in different domains (Kejzlar *et al.*, 2020), and assessment of the puzzling mass of  $^{80}\text{Zr}$  (Hamaker *et al.*, 2021). Figure 4 shows the posterior probability of existence for all nuclei in the nuclear landscape based on predictions of 11 global mass models, the most recent data on nuclear existence and masses, and three model-averaging strategies to compute the BMA weights.

#### 5. Nuclear shell-model applications

Machine learning methods have been used to provide UQ of shell-model (configuration interaction) calculations based on




FIG. 4. Bayesian extrapolation and model averaging. The quantified landscape of nuclear existence obtained in the BMA calculations using 11 global mass models and three model-averaging strategies is shown. For every nucleus with  $Z, N \geq 8$  and  $Z \leq 119$ , the probability that the nucleus is bound with respect to proton and neutron decay is marked. The domains of nuclei that have been experimentally observed and whose separation energies have been measured (and used for training) are indicated. Adapted from Neufcourt *et al.*, 2020b.

phenomenological two-body matrix elements. Yoshida *et al.* (2018) used Bayesian ML to compute marginal estimates of the posterior distribution for the shell-model Hamiltonian in the  $0p$  space and uncertainty estimates on observables. A similar analysis, but for effective Hamiltonians in the  $1s0d$  shell, was carried out by Fox, Johnson, and Perez (2020) and Magilligan and Brown (2020) (using PCA) and Akkoyun, Laouet, and Benrachi (2021) (using ANNs). Eigenvector continuation (EC) was used by Yoshida and Shimizu (2022) to construct an emulator of the shell-model calculations for a valence space, parameter optimization, and UQ.

## 6. Effective field theory and A-body systems

Bayesian ML, BO, and UQ in *ab initio* nuclear theory were reviewed by Ekström *et al.* (2019) and Ekström (2020). There have been several studies of nucleon-nucleon scattering using Bayesian ML to estimate chiral effective field theory (EFT) truncation errors and low-energy coupling constants (Melendez, Wesolowski, and Furnstahl, 2017; Melendez *et al.*, 2019; Wesolowski *et al.*, 2019; Svensson, Ekström, and Forssén, 2022), and there have also been applications to  $np \rightarrow dy$  (Acharya and Bacca, 2021). Furnstahl *et al.* (2020) used eigenvector continuation (Frame *et al.*, 2018) as an emulator for scattering. The method was improved upon by Drischler *et al.* (2021), who made use of different boundary conditions. The application of EC as a fast emulator for UQ for few- and many-body systems was first explored by König *et al.* (2020) and adapted for coupled cluster calculations and a global sensitivity analysis of  $^{16}\text{O}$  (Ekström and Hagen, 2019) and, with Bayesian history matching (Vernon, Goldstein, and Bower, 2014), for global properties of  $^{208}\text{Pb}$  (Hu *et al.*, 2021). Eigenvector continuation emulators have been used to put rigorous constraints on low-energy constants for the three-nucleon forces (Wesolowski *et al.*, 2021) and to make predictions for the binding of  $A = 6$  nuclei (Djärv *et al.*, 2022). For other applications of eigenvector continuation, see Eklind (2021), Melendez, Drischler *et al.* (2021), Sarkar and Lee (2021), and Zhang and Furnstahl (2021). Connell, Billig, and Phillips (2021) addressed whether BMA improves the

extrapolation of polynomials, which are used as proxies for fixed-order EFT calculations.

Using a scattering amplitude as input data (respecting unitarity, Hermiticity, and analyticity), Sombillo *et al.* (2020, 2021a, 2021b) classified the nature of the poles near threshold with multilayer ANNs. In particular, Sombillo *et al.* (2020) applied the ANNs to predicting the correct nature of the poles in the nucleon-nucleon scattering data from a partial wave dataset. This is an example of a classification problem where the aim is to classify whether the poles represent bound, virtual, or resonant states. Kapschak and Meißner (2021) proposed an iterative ANN perturbation theory to study  $s$ -wave scattering lengths for shallow potentials.

Navarro Pérez *et al.* (2015) propagated the statistical errors in nucleon-nucleon scattering to calculations of light nuclei. The radiative capture rates  $^7\text{Be} + p \rightarrow ^8\text{B} + \gamma$  (Zhang, Nollett, and Phillips, 2015) and  $^3\text{He} + ^4\text{He} \rightarrow ^7\text{Be} + \gamma$  (Zhang, Nollett, and Phillips, 2020) were estimated with BO. Bayesian ML has been applied to neutron- $\alpha$  scattering in chiral EFT (Kravaris *et al.*, 2020).

In other few-body calculations, ANNs were used to determine the deuteron wave function using variational optimization (Keeble and Rios, 2020). Three-body Efimov bound states were studied using ANNs (Saito, 2018), and CNNs were used to classify states of a three-body system (Huber *et al.*, 2021). Variational Monte Carlo calculations optimized with ANNs have been performed for light nuclei with up to six nucleons (Adams *et al.*, 2021; Gnech *et al.*, 2022); see Fig. 5. The latter results are interesting since the representation of the Jastrow factor in terms of ANNs has the potential to introduce additional correlations. The universal approximation theorem (Cybenko, 1989; Hornik, 1991) states that ANNs can represent a wide variety of nonlinear functions when given appropriate weights. Overall, using ANNs to extract complicated correlations in many-body calculations seems to be a promising approach, as recently shown by Pescia *et al.* (2021). They used ANNs for the simulation of strongly interacting systems in the presence of spatial periodicity. This has potential applications for studies of infinite nuclear matter, where periodic boundary conditions are often




FIG. 5. Many-body variational calculations with ANNs. Point-nucleon densities of  ${}^4\text{He}$  for the leading-order pionless-EFT Hamiltonian. The solid points and the shaded area represent the ANN and Green's function Monte Carlo results, respectively. From Adams *et al.*, 2021.

imposed to extract the equation of state of dense fermionic matter, as discussed later.

In larger  $A$ -body systems, the ANN extrapolation of nuclear structure observables have been used for the no-core shell model (Negoita *et al.*, 2019), coupled cluster theory (Jiang, Hagen, and Papenbrock, 2019), and configuration interaction calculations (Yoshida, 2020). Artificial neural networks were used to learn important configurations for symmetry-adapted no-core shell-model calculations (Molchanov *et al.*, 2021). Similarly, ANNs have also been used to invert Laplace transforms required to compute nuclear response functions from Euclidean time Monte Carlo data (Raghavan *et al.*, 2021). Restricted BMs were used to represent many-body contact interactions using auxiliary fields (Rrapaj and Roggero, 2021). Ismail and Gezerlis (2021) used ML techniques to study finite-size effects and extrapolate the unitary gas to the thermodynamic limit at zero range.

Another topic that has great potential for applications to studies of infinite matter and the equation of state (EOS) for dense nuclear matter is the application of ML to many-body methods like coupled cluster theory, Green's function theories, in-medium similarity renormalization group methods; see Hjorth-Jensen, Lombardo, and van Kolck (2017). Common to these methods is that the underlying approximations can be systematically expanded upon by including more complicated correlations. Various approaches, like coupled cluster theory (Hjorth-Jensen, Lombardo, and van Kolck, 2017), sum to infinite order selected many-body contributions such as so-called one-particle-one-hole and two-particle-two-hole correlations. Including three-particle-three-hole correlations is computationally much more involved. Here ML-based methods can be extremely useful, particularly for studies of nuclei from calcium and beyond and infinite matter. Recent atomic and molecular physics studies (Margraf and Reuter, 2018; Wilkins *et al.*, 2019; Townsend and Vogiatzis, 2020; Agarwal *et al.*, 2021) can easily be extended to finite nuclei and infinite matter.

## 7. Nuclear reactions

Low-energy nuclear reaction models have a critical reliance on a wide variety of parameters including nuclear masses,




FIG. 6. Bayesian analysis of the mass-radius relation for neutron stars. The Bayesian uncertainty reflects constraints on an equation of state.  $S$ , symmetry energy;  $M_{\text{max}}$ , masses of three most massive neutron stars;  $\Lambda$ , tidal deformability of GW170817;  $(M, R)$ , high-precision mass-radius measurement;  $I$ , moment of inertia;  $E$ , baryonic rest mass of a star. From Miller, Chirenti, and Lamb, 2020.

nuclear level densities, transmission coefficients, and optical model parameters. While nuclear masses were discussed earlier, there have been some ML-based studies seeking improved predictions of other parameters across the nuclear chart; see Sec. VI. There are also numerous studies using BO for UQ on reaction model parameters; some recent examples include  $R$ -matrix analyses of cross sections (Odell, Brune, and Phillips, 2021), optical model parameters (Lovell and Nunes, 2018; King *et al.*, 2019; Yang *et al.*, 2020; Lovell *et al.*, 2021), and sensitivity analyses (Catacora-Rios *et al.*, 2019, 2020). In the future, it is anticipated that ML will help identify those measurements that most effectively constrain theoretical models, optimize model parameters simultaneously across multiple reaction channels for many isotopes, and provide guidance to theory through global systematic studies that can be efficiently executed with surrogate models.

## 8. Neutron star properties and nuclear matter equation of state

Studies of dense nuclear matter and its pertinent EOS, with its strong implications for studies of neutron stars and studies of supernovae, is a field that has seen considerable progress during the last two decades. The increased wealth of data related to cold dense matter, from laboratory experiments and theoretical simulations to neutron star observations such as the gravitational-wave events GW170817 and GW190814, provide a framework for constraining theoretically the EOS of dense matter.

As an example, we present in Fig. 6 the results of a Bayesian analysis by Miller, Chirenti, and Lamb (2020) based on a posterior probability distribution. The figure displays the mass-radius constraints that correspond to an EOS obtained from Bayesian inference. Here a symmetry energy of  $S = 32 \pm 2$  MeV was adopted, together with the masses of the three most massive neutron stars, the tidal deformability of GW170817, hypothetical masses and radii to a precision of 5%, and similarly a hypothetical measurement of the moment of inertia of a  $1.338M_{\odot}$  star to 10% precision. Miller, Chirenti,

and Lamb showed that the resulting EOS is sensitive to the symmetry energy below the saturation density.

In several studies, Bayesian inference, ML, and/or other statistical approaches have been applied to constrain the EOS and other properties pertaining to infinite matter studies. Drischler, Furnstahl *et al.* (2020) and Drischler, Melendez *et al.* (2020) carried out UQ of many-body calculations of the EOS. In particular, Bayesian ML with GPs was employed to propagate theoretical uncertainties using many-body perturbation theory to fourth order with two- and three-body interactions from chiral effective field theory. Gaussian processes were also used by Essick *et al.* (2021) to constrain the symmetry energy and its slope. Deep learning inference for the EOS was recently studied by Fujimoto, Fukushima, and Murase (2021) with the use of observational data on masses and radii of known neutron stars. Several other groups have used ML methods and/or Bayesian inference to study the nuclear EOS with nuclear data and theoretical calculations (Margueron, Casali, and Gulminelli, 2018a, 2018b; Xu, Xie, and Li, 2020; Newton and Crocombe, 2021), x-ray observations of neutron stars (Näyttälä *et al.*, 2017), gravitational-wave data from neutron star mergers (Lim and Holt, 2019; Capano *et al.*, 2020; Dietrich *et al.*, 2020; Güven *et al.*, 2020; Kunert *et al.*, 2022), or both x-ray and gravitational-wave data (Raaijmakers *et al.*, 2020; Xie and Li, 2020; Al-Mamun *et al.*, 2021; Ayriyan *et al.*, 2021; Han *et al.*, 2021). The properties of dense matter have also been probed using heavy-ion collisions (Morfouace *et al.*, 2019; Xie and Li, 2021) as well as studies at nonzero temperature (Wang *et al.*, 2020). At intermediate energies, CNNs were recently applied by Zhang *et al.* (2021) to determine the impact parameters of heavy-ion collisions at low to intermediate incident energies (up to 100 MeV/nucleon).

## B. Medium-energy nuclear theory

Nuclear femtography is an emerging field in nuclear physics that aims to map out quantum correlation functions (QCFs) that characterize the internal three-dimensional structure of nucleons and nuclei, as well as hadronization in high-energy reactions, in terms of the quark and gluon (collectively called partons) degrees of freedom of quantum chromodynamics (QCD). In contrast to any system found in nature, the partons of QCD are not detectable experimentally due to confinement, which prevents direct access to the QCFs. Nonetheless, certain classes of observables can be factorized in terms of convolutions between short-distance physics calculable in perturbative QCD and long-distance physics that is nonperturbative and encoded in the formulation of QCFs. However, to extract the latter, one faces an inverse problem inherited from the mathematical relation between the experimental observables and the QCFs stemming from the inability to obtain closed form solutions for the QCFs. Therefore, the only practical approach is to parametrize the QCFs and calibrate them via BO or another method.

## 1. Bayesian inference

The traditional approach to implementing Bayesian inference involves the use of theory-inspired parametrizations for

the QCFs, tuned via maximum likelihood estimators (Bishop, 2006; Hastie, Tibshirani, and Friedman, 2009; Murphy, 2012), along with an error analysis based on the Hessian matrix optimization (Bishop, 2006; Murphy, 2012; Pumplin *et al.*, 2002). This approach was developed in the context of QCD global analysis of parton distribution functions (PDFs), which is a type of QCF describing the longitudinal momentum fractions of partons inside nucleons. This approach has been adopted by various groups around the world (Harland-Lang *et al.*, 2015; Accardi *et al.*, 2016; Alekhin, Blümlein, and Moch, 2018; Hou *et al.*, 2021) and has found relatively good success in describing a large bank of high-energy data that are sensitive only to the one-dimensional degrees of freedom in the nucleon.

In recent years, Monte Carlo-based methodologies for Bayesian inference (Hastie, Tibshirani, and Friedman, 2009; Murphy, 2012; Mehta *et al.*, 2019) have gained traction, providing a more reliable uncertainty quantification for QCFs in the Bayesian framework. Ball *et al.* (2010) demonstrated the feasibility of carrying out QCD global analysis by sampling the Bayesian posterior distribution using the data resampling technique. They also introduced ANN parametrizations to extend the flexibility of the QCF modeling and explore the degree of parametrization bias. For the case of PDFs, one found that, in regions where the data maximally constrain the PDFs, ANNs and traditional parametrizations give qualitatively similar results, with the differences becoming increasingly evident in extrapolated regions.

## 2. Simultaneous extraction paradigm

The Monte Carlo approach for Bayesian inference has also recently been applied to a simultaneous extraction of a variety of QCFs, including spin-dependent PDFs (Ethier, Sato, and Melnitchouk, 2017), transverse momentum distributions (Cammarota *et al.*, 2020), and fragmentation functions (Ethier, Sato, and Melnitchouk, 2017; Sato *et al.*, 2020; Moffat *et al.*, 2021), thereby establishing a new paradigm in nuclear femtography. The simultaneous approach is crucial for solving the inverse problem for the QCFs, especially for those quantities that rely on each other. An example of such a situation is in semi-inclusive deep-inelastic scattering, where the experimental observables are sensitive to QCFs describing the internal structure of hadrons as well as QCFs that describe the hadronization process. In principle, each of these types of QCFs can be extracted independently from processes that are solely sensitive to each type of QCFs. However, when included within a global context, they require simultaneous analysis in order to take into account the correlations induced by uncertainties regarding the various interdependent QCFs themselves.

Ethier, Sato, and Melnitchouk (2017) showed that the strange polarization in the nucleon is mostly unconstrained if one takes fully into account the uncertainties on the spin-dependent PDFs and fragmentation functions. Sato *et al.* (2020) and Moffat *et al.* (2021) found a suppression of the unpolarized strange quark PDF relative to lighter quark PDFs to be preferred by the simultaneous extraction of spin-averaged PDFs and fragmentation functions when they included the standard deep-inelastic scattering and

semi-inclusive deep-inelastic scattering electromagnetic observables, semi-inclusive annihilation in  $e^+e^-$  collisions, and lepton-pair production in  $pp$  reactions. [Cammarota \*et al.\* \(2020\)](#) performed the global analysis for all existing single-spin asymmetries, providing the empirical demonstration that the flavor-dependent nucleon tensor charges agree within uncertainties with those computed directly in LQCD.

### 3. LQCD and experimental global analysis

Another new direction that has been recently explored is the inclusion of LQCD data as Bayesian priors that can help overcome the difficulties in deconvoluting QCFs from experimental data. An advantage of this is that the lattice data can in principle have access into domains of the QCFs that are inaccessible experimentally or difficult to determine, particularly processes and kinematics. [Lin \*et al.\* \(2018\)](#) used the isovector tensor charge from LQCD as a Bayesian prior to extract the nucleon transversity distribution. Similarly, [Bringewatt \*et al.\* \(2021\)](#) carried out exploratory studies to include off-the-light-cone matrix elements in a QCD global analysis of spin-averaged and spin-dependent PDFs, thereby demonstrating some level of success and/or tension depending on the type of observables and the associated QCFs. For other studies involving LQCD, see [Karpie \*et al.\* \(2019\)](#) and [Del Debbio \*et al.\* \(2021\)](#) and Sec. III.C. The combination of LQCD results and experimental data in the framework of Bayesian inference provides new avenues for addressing the challenges imposed by the inverse problem in QCFs and facilitates reliable comparisons between data and theory.

## C. Lattice QCD

Lattice field theory is the theoretical framework for understanding the properties of strongly interacting matter. The fundamental theory of strong interactions is QCD, a quantum field theory that requires nonperturbative computations to address low-energy hadronic physics. Lattice QCD provides both a rigorous definition of QCD and a powerful tool for numerical computations. The basic computational task in LQCD is a Monte Carlo evaluation of multidimensional integrals that results in the lattice regularization of QCD.

Sophisticated and powerful QCD algorithms have been developed to take advantage of modern-day supercomputers, leading to many important results aiding experimental efforts to understand the nature of strongly interacting matter. However, despite impressive achievements, computations close to the continuum limit that are required to reduce systematic errors are still out of reach with today's resources. Machine learning methods offer a new avenue for improving current computations as well as allowing for studies of previously impossible questions. Typical LQCD calculations proceed in two stages. An ensemble of gauge field configurations is first generated. Subsequently, certain correlation functions of the fundamental fields are computed by averaging over the ensemble of gauge configurations. Finally, analyses of these correlation functions lead to the desired physical observables. Machine learning techniques are now applied to all these stages of LQCD computations, thus promising to enhance and extend the current state of the art.

## 1. The sign problem at nonzero density

Systems at nonzero density (as in nuclear or neutron matter) or Minkowski time dynamics (parton distribution functions and transport coefficients) cannot be studied with standard Monte Carlo methods due to the fermion sign problem. Recently, it has been shown that, by evaluating the relevant path integral over a field manifold deformed into complex space, the sign problem can be alleviated or even eliminated ([Cristoforetti, Di Renzo, and Scorzato, 2012](#); [Cristoforetti \*et al.\*, 2014](#); [Di Renzo and Eruzzi, 2015](#); [Fujii, Kamata, and Kikukawa, 2015](#); [Fukushima and Tanizaki, 2015](#); [Ulybyshev, Winterowd, and Zafeiropoulos, 2020](#)). To date the choice of manifolds has been guided by either impossibly expensive calculations or human insight into particular models. Machine learning methods have begun to be applied in both supervised and unsupervised learning modes ([Alexandru \*et al.\*, 2017](#); [Alexandru, Bedaque, Lamm, and Lawrence, 2018](#); [Alexandru, Bedaque, Lamm, Lawrence, and Warrington, 2018](#); [Bursa and Kroyter, 2018](#); [Ohnishi, Mori, and Kashiwa, 2019](#); [Wynen \*et al.\*, 2021](#)) to discover the integration manifolds that alleviate the sign problem and, in certain cases, allow for calculations that previously were not possible. Refinement of these methods opens a new avenue for understanding QCD at nonzero density, as well as understanding the real-time dynamics of strongly interacting matter. At nonzero temperature the problem of reconstructing spectral functions from Euclidean correlators arises. Recently both VAEs ([Chen \*et al.\*, 2021](#)) and GPs ([Horak \*et al.\*, 2021](#)) have been used to solve the associated inverse problem.

## 2. Ensemble generation

Generating gauge configuration ensembles close to the continuum limit has also proved to be a daunting computational task ([Schaefer, Sommer, and Virotta, 2011](#)). Present-day computations with lattice spacing below 0.05 fm are severely limited due to a critical slowing down, i.e., large autocorrelation times in the Markov chain used to generate the gauge configuration ensemble. The idea of trivializing maps ([Lüscher, 2010](#)) was introduced several years ago as a possible solution to this problem. It relied on an analytically determined map that can be used to change variables in the path integral, resulting in a trivial integration weight. [Lüscher, 2010](#) showed that such maps exist and can be constructed as solutions to a gradient flow equation whose generator can be set up perturbatively in the flow time. However, practical constructions of such maps have not been carried out even for simple models ([Engel and Schaefer, 2011](#)). A successful application of this program was done in the context of stochastic perturbation theory ([Lüscher, 2015](#)), which is used to perform high-order perturbative computations in QCD. Machine learning methods provide a new approach to discovering these maps. In fact, the idea of remapping variables with complex probability distributions to trivially distributed variables is something that is commonly done in ML. In that sense, adaptation of these methods in the context of Markov chain Monte Carlo (MCMC) approaches to lattice field theory is an interesting idea with enormous potential for solving the critical slowing down problem in LQCD.

Applications of ML methods to lattice field theory calculations have already emerged. The scalar  $\phi^4$  theory in two dimensions is one of the first models studied, where GANs were used to reduce autocorrelation times (Pawlowski and Urban, 2020). Albergo, Kanwar, and Shanahan (2019) and Medvidovic *et al.* (2020) showed that normalizing flows constructed via ANNs can indeed be trained to effectively sample the configuration space of two-dimensional scalar  $\phi^4$  theory up to lattice sizes of  $14^2$ . In addition, Hackett *et al.* (2021) used flow-based methods to sample from multimodal distributions. Nicoli *et al.* (2021) studied thermodynamic properties of scalar  $\phi^4$  theory in two dimensions using normalizing flows to generate field configurations, thus confirming the efficacy of this approach. These works demonstrated that critical slowing down may be eliminated in MCMC simulations for scalar  $\phi^4$  theory. However, further studies and improvements of the approach (Del Debbio, Marsh Rossney, and Wilson, 2021) demonstrated that the training cost of the flow-based models scales badly as the critical point is approached and the correlation length and lattice size grow, indicating that further refinements of the approach are needed for the elimination of critical slowing down. The flow-based methods for Monte Carlo sampling have already been extended to gauge theories (Favoni *et al.*, 2020; Kanwar *et al.*, 2020; Luo *et al.*, 2020, 2021; Boyd, Kanwar *et al.*, 2021; Tomiya and Nagai, 2021), fermionic theories (Albergo *et al.*, 2021), and theories with a sign problem (Lawrence and Yamauchi, 2021). Note that an entirely different approach in using ML methods for optimizing MCMC simulations is the so-called L2HMC approach [see Levy, Hoffman, and Sohl-Dickstein (2018)], which was recently applied to the U(1) gauge theory with fermions by Foreman, Jin, and Osborn (2021). In this approach, ML methods are used to construct a new flow-based map that replaces the Hamiltonian evolution in Hamiltonian Monte Carlo (Bishop, 2006; Murphy, 2012; Li, Dong *et al.*, 2020). This is a promising idea that could have an impact in realistic MCMC calculations in lattice field theory.

### 3. Correlation function estimators

Calculations of observables in LQCD require the computation of quark propagators in the background of a large number of gauge configurations. Quark propagators are computed by solving a linear system of equations with a large and sparse coefficient matrix. Machine learning methods are used (Pederiva and Shindler, 2020) to obtain low-precision solutions in order to construct approximate observables from which full precision results are obtained via the so-called all-mode-averaging technique (Shintani *et al.*, 2015), resulting in an enormous increase in computational efficiency. Along similar lines, boosted decision trees were used as approximate estimators of two- and three-point correlation functions used in calculations of nucleon charges and of the phase acquired by the neutron mass with a small parity violation (Yoon, Bhattacharya, and Gupta, 2019). The same methodology was also employed for correlators used in parton distribution function calculations (Zhang *et al.*, 2020). In addition, GPs were employed in predicting the long-distance behavior of matrix elements in PDF computations in LQCD (Alexandrou

*et al.*, 2020). An interesting new idea was introduced by Detmold *et al.* (2021), who used ML methods to find deformations of the fields in the path integral such that the variance of observables was reduced.

### 4. Miscellany

Finally, there is a large body of work in which ML methods are used to understand properties of lattice field theories. These works include the prediction of lattice action parameters from field configurations (Shanahan, Trewartha, and Detmold, 2018; Blücher *et al.*, 2020), discoveries of the holographic geometry in AdS/CFT descriptions of field theories (Hashimoto *et al.*, 2018a, 2018b; You, Yang, and Qi, 2018; Hashimoto, 2019; Hashimoto, Hu, and You, 2021), and an understanding of the nature of phase transitions (Wetzel and Scherzer, 2017; Bachtis, Aarts, and Lucini, 2020; Blücher *et al.*, 2020; Chernodub *et al.*, 2020; Boyd, Chernodub *et al.*, 2021). A discussion of ML applications to LQCD can also be found in a recent Snowmass report (Blum *et al.*, 2021).

### D. High-energy nuclear theory

At extremely high temperature or density, quarks and gluons are deconfined from nucleons to form a new state of nuclear matter, the quark gluon plasma (QGP). In the early Universe, this state of matter existed for about a few microseconds after the big bang and might also be at the core of some neutron stars or created during violent neutron star mergers. The field of high-energy nuclear physics aims to search for its formation, the phase transition between normal nuclear matter and QGP, locate the critical point(s), and determine the deconfinement temperature in high-energy heavy-ion collisions (HICs). Many properties of QGP, such as shear and bulk viscosity, conductivity, and jet transport coefficient, can be extracted from experimental data using soft and hard probes.

### 1. Bayesian inference

Since the lifetime of QGP in a HIC is short (about  $10^{-23}$  s), experiments detect the relics of QGP: hadrons, photons, leptons, and their momentum distributions and correlations. The properties of QGP can only be extracted indirectly through theoretical models. In practice, each observable in experimental data is entangled to many model parameters. This entanglement hinders the determination of a specific model parameter. Even worse, sometimes different combinations of model parameters produce degenerate outputs when the data are projected into lower dimensions.

Two promising ML tools emerge to solve this seemingly ill-defined inverse problem. The first tool is BO, which tends to fit all model parameters at the same time using all available data. During this global fitting, if several groups of model parameters generate the same output, then the probability of the real model to be one of them decreases. On the contrary, if only one group of model parameters describes the data and a change of parameters leads to large model-data differences, then the posterior distribution peaks at the optimal parameter value with small width, thereby indicating a small model uncertainty.

Bayesian optimization is widely used in high-energy nuclear physics to constrain the QCD equation of state (Pratt *et al.*, 2015), the QGP transport coefficients (Bernhard *et al.*, 2016; Bernhard, Moreland, and Bass, 2019; Paquet, 2021), the fluctuation and correlation (Yousefnia *et al.*, 2021), the heavy quark diffusion coefficient (Xu *et al.*, 2018), the jet transport coefficient (Cao *et al.*, 2021), and the jet energy loss distributions (He, Pang, and Wang, 2019). In many applications, other ML tools are used to assist the Bayesian analysis. For example, the relativistic fluid dynamics simulations of HICs are time consuming, which prevents a fast MCMC random walk in the parameter space. Emulators based on the theory of GPs are employed to approximate model outputs using efficient interpolations with much fewer design points in the parameter space. Since data obtained this way are redundant and correlated, PCA is used to compress data to lower dimensions.

The observables used in Bayesian inference are high-level features designed with personal experience. However, feature engineering is known to be incomplete, insufficient, and sometimes misleading, which may cause important correlations hidden in high-dimensional data to be lost. This is unavoidable due to the high-dimensional character of the experimental data. It is difficult to recognize from exotic low-level features the nonlinearly correlated patterns that are unique and robust for determining a specific model parameter.

## 2. Inversion problems with ML

Deep neural networks are promising ML tools for tackling the difficult inverse problem in HIC. If low-dimensional projections of model outputs are degenerate in the seemingly ill-defined inverse problem, one would expect differences to still exist in high dimensions or in nonlinear correlations between different dimensions. The universal approximation theorem (Cybenko, 1989; Hornik, 1991) ensures that ANNs have enough representation capability to map low-level features to some given model parameters in supervised learning. If information gets totally lost in the dynamical evolution of HICs because of entropy production, the network can never succeed in building this map. If the information of one specific model parameter survives the dynamical evolution and exists in the final output of HICs, the network has a better chance to build this map. In this sense, if the network provides high accuracy predictions in a supervised learning scenario, it indicates that the signal encodes in the complex final state output and the network helps to decode this information. On the other hand, if the network fails, this indicates that either the information on the physical signal gets totally lost or the used network does not have enough representation power. For the latter, it is still possible with a deeper, wider, or more suitable network.

The type of nuclear phase transition is an important input to relativistic hydrodynamic simulations of HIC. At high energies, LQCD predicts that the transition between QGP and hadron resonance gas is a smooth crossover. At intermediate beam energies, it is conjectured to be a first-order phase transition. Different phase transitions lead to different pressure gradients around the transition temperature that drives the QGP expansion (it also depends on the shear and bulk




FIG. 7. Dynamical edge-convolution neural network used for event classification and particle tagging in heavy-ion collisions. From Huang *et al.*, 2021.

viscosity). In this inverse problem, one can determine the phase transition type during the dynamical evolution using final state output. Supervised CNNs (Pang *et al.*, 2018; Du *et al.*, 2020), point cloud networks (Steinheimer *et al.*, 2019), and unsupervised AEs (Wang *et al.*, 2020) are trained to identify the QCD phase transition types using the final state hadrons. It is verified that signals of the phase transition survive the dynamical evolution, and deep learning succeeds in decoding this information from the final hadron distribution. To avoid overfitting to given model parameters, different parameter combinations are used to form a diverse training dataset. Prediction difference analysis is used to interpret which region in the momentum space is most important for the networks to make their decisions.

Data produced in high-energy HIC are lists of particles with their four-momenta and quantum numbers. Early studies used histograms to convert this information into images that were required by two-dimensional CNNs. It was later found that the point cloud network is suitable for this data structure (Steinheimer *et al.*, 2019). To enhance the representation power, a dynamical edge-convolution network followed by a point cloud net is used to identify self-similarity and critical fluctuations in HICs (Huang *et al.*, 2021). Figure 7 shows the architecture of the dynamical edge-convolution network. The input to the network is a list of hadrons. The output has two branches: one for event classification and the other for particle tagging. The KNN algorithm finds the  $k$ -nearest neighbors of each particle in both momentum space and feature space. Repeating the KNN and edge-convolution blocks twice helps one to find long-range multiparticle correlations that are the key to searching for critical fluctuations.

The impact parameter is the transverse distance between colliding nuclei, whose precise determination helps many downstream tasks. It is not possible to measure impact parameters directly through experiment. Several different

ML tools are used to determine the impact parameters of HICs, including shallow neural networks (Bass *et al.*, 1996), ANNs (Omana Kuttan *et al.*, 2020, 2021), SVMs (De Sanctis *et al.*, 2009), and EMB (Li, Wang *et al.*, 2020, 2021; Mallick *et al.*, 2021). In this inverse problem, data from Monte Carlo simulations are used to map the final state output to the impact parameter. The traditional method uses a single observable (the particle multiplicity) to determine the impact parameter. Machine learning methods using high-dimensional data result in much smaller uncertainties.

Another inverse problem associated with initial states is linked with the given nuclear structure, which in turn affects in many ways the final state outputs of the HICs. In a prototype inverse problem, a 34-layer residual neural network is used to predict the deformation parameters of the involved nuclei (Pang, Zhou, and Wang, 2019). Using the simulation data, the network succeeds in extracting the magnitude of the nuclear deformation but fails to extract the sign. The failure denotes a degeneracy raised in the dynamical process of high-energy collisions. In another inverse problem, Bayesian CNNs are employed to identify the  $3 - \alpha$  and  $4 - \alpha$  structures in the colliding light nucleus, from the final output of simulated heavy-ion collisions (He *et al.*, 2021). The overall classification accuracy reaches 95% for  $^{12}\text{C}/^{16}\text{O} + ^{197}\text{Au}$  collisions.

### 3. Other applications of ML methods

High-energy jets lose energy when they traverse through hot deconfined nuclear matter. In the inverse problem, CNNs are employed to predict the energy loss ratio from final state hadrons inside a jet cone (Du, Pablos, and Tywoniuk, 2021a, 2021b), which allows one to study the jets based on the initial energy.

The chiral magnetic effect was expected to arise from possible parity violation of strong interactions. However, all previously proposed observables suffer from large background contamination in heavy-ion collisions. Deep CNNs are used to identify the charge separation associated with the chiral magnetic effect (Zhao *et al.*, 2021). The network is robust to diverse conditions including different collision energies, centralities, and elliptic flow backgrounds.

The interaction between bottom and antibottom quarks in QGP is modeled as a heavy quark potential, whose variational function form is represented by deep neural networks (Shi *et al.*, 2022). The inputs to the networks are the temperature of the QGP and the quark antiquark separation  $r$ . The output is the heavy quark potential. Solving the pertinent differential equations numerically with this potential gives mass spectra whose difference from LQCD calculations defines the cost function. Optimizing this cost function gives one the parameters of the model.

Finally, Mroczeck *et al.* (2022) used active learning (Cohen, 2018) to reduce sampling requirements for training classifiers in searches for acceptable EOS parameters.

## IV. EXPERIMENTAL METHODS

The aim of this section is to give the reader an overview of recent progress and future research directions in ML approaches and methods applied to nuclear physics

experiments. During the last few years, ML methods have been applied to the full chain of experimentation, including the design of experiments, the acquisition of data, the processing chain of converting detector information into observables, and physics analysis.

### A. Streaming detector readout

In triggerless or streaming readout data acquisition systems, detector data are read out in continuous parallel streams that are encoded with information about when and where the data were taken. This simplifies the readout as no custom trigger hardware and firmware is needed and is beneficial for experiments that are limited by event-pileup or overlapping signals from different events. Streaming readout also gives one an opportunity to streamline workflows for on-line and off-line data processing and allows one to take advantage of ML approaches.

The LHCb experiment at CERN has pioneered the idea of seamless data processing from the readout to the analysis, using software stages at early stages of the event selection. Part of this system is a custom boosted decision tree algorithm for the reconstruction of decay products of  $b$  hadrons (Gligorov and Williams, 2013; Likhomanenko *et al.*, 2015). Progress in novel boosted decision tree algorithms allow improvements in the efficiency of decay classifications by up to 80% for high-rate events.

The CLAS12 experiment at Jefferson Lab tested a prototype streaming readout system successfully under beam conditions. The test was limited to the measurement of inclusive electroproduction of neutral pions in a lead tungstate calorimeter and a hodoscope. An unsupervised hierarchical cluster algorithm was utilized in real time with real data taken in streaming readout mode to combine the time, position, and energy information at the hit level, and associate each hit with a cluster membership and an outlier score (Ameli *et al.*, 2021). The implementation (McInnes, Healy, and Astels, 2017) allows one to successfully reject noise hits and to identify clusters for diverse topologies and large hit multiplicities.

### B. Reconstruction and analysis

#### 1. Charged particle tracking

Deep learning approaches using neural networks have the advantage of a high level of flexibility and robustness with a minimum of assumptions about the data. This offers an effective solution to the challenges of charged particle tracking at high luminosity. At high luminosity, tracking suffers from track candidates that share hits (combinatorials). This results in hits wrongly identified as “on track” and produces ghost tracks. In high-luminosity environments, the largest fraction of CPU time in tracking in a traditional analysis is spent on setting up various Kalman filters (Grewal and Andrews, 2014) at each measurement site. These are affine operations involving CPU and memory intensive matrix-matrix multiplications and matrix-vector multiplications. Therefore, the improvement in track seeding resulting from using ANNs and deep learning methods yields substantially faster track reconstruction speed. In addition, the selection of the correct seed results in improved tracking efficiency (Guest, Cranmer,

and Whiteson, 2018; Gavalian *et al.*, 2020). Furthermore, noise rejection algorithms have an impact in selecting the right combinations of hits in seeding (Komiske *et al.*, 2017).

One of the most common deep learning algorithms employed for tracking pattern recognition are CNNs. The CNN features are generally representations of the detector geometry. Machine learning algorithms used for background rejection involve topological properties of tracks to isolate signal from background (Kramárik, 2020). Outlier detection methods are used to remove noise uncorrelated hits and to classify tracks including pileup (Ayyad *et al.*, 2018).

Large liquid scintillator detectors can hold a large target mass for neutrino detection in solar and reactor neutrino experiments. Machine learning can benefit pure liquid scintillator detectors in event position and energy reconstruction (Qian *et al.*, 2021). Similarly, neutron detectors such as NeuLAND track both charged and neutral particles. Simple neural network architectures have been shown to efficiently assist in tracking (Mayer *et al.*, 2021).

## 2. Calorimetry

The GlueX experiment at Jefferson Lab used ANNs to reduce the background in the GlueX forward calorimeter for the detection of photons produced in the decays of hadrons (Barsotti and Shepherd, 2020). Energy deposition characteristics in the calorimeter such as shapes, size, and distribution were employed to discriminate between the signal and background, where the background originates mostly from hadronic interactions that can be difficult to distinguish from low-energy photon interactions. The training was done on data using  $\omega$ -meson decays. The ANN-based algorithm proved to be a powerful tool to reconstruct neutral particles with high efficiency and to provide substantial background rejection capability.

## 3. Particle identification

Particle identification (PID) is done with dedicated detectors capable of identifying certain particle types. For example, Cherenkov detectors are largely used in modern nuclear experiments for identifying charged particles like pions, kaons, and protons corresponding to a wide range in momentum. Cherenkov detectors are typically endowed with single photon detectors and the particle type can be recognized by classifying the corresponding detected hit pattern (Fanelli, 2020). DeepRICH (Fanelli and Pomponi, 2020) is a recently developed custom architecture that combines VAEs, CNNs, and ANNs. The reconstruction performance is close to that of other established methods like FastDIRC (Hardin and Williams, 2016) with fast reconstruction times due to its implementation on graphics processing units that allow for parallel processing of batches of particles during the inference phase.

Derkach *et al.* (2020) used GANs to simulate the Cherenkov detector response. This architecture predicts the multidimensional distribution of the likelihood for particle identification produced by FastDIRC bypassing low-level details. GANs were used by Maevskiy *et al.* (2020) for a fast and accurate simulation of a Cherenkov detector; these

have been trained using real data samples collected by the LHCb Collaboration.

The utilization of jets produced by the hadronization of a quark or gluon in heavy-ion experiments or future electron ion colliders for nuclear physics such as the Electron-Ion Collider (EIC) can be functional for a variety of fundamental topics (Page, Chu, and Aschenauer, 2020). Machine learning has been applied to design experimental observables that are sensitive to jet quenching and parton splitting (Lai *et al.*, 2020, 2021). Chien (2019) showed that a deep CNN trained on jet images allows the study of jet quenching utilizing quark and gluon jet substructures. Moreover, CNNs were used to discriminate between quark and gluon jets by Komiske, Metodiev, and Schwartz (2017). Different deep architectures (CNN, dense ANN, and RNN) were studied by Apolinário *et al.* (2021) for the classification of quenched jets and, in particular, to discriminate between mediumlike and vacuumlike jets. Identification of jets as originating from light-flavor or heavy-flavor quarks is an important aspect in inferring the nature of the particles produced in high-energy collisions.

Much progress has been made in recent years on heavy-flavor tagging, with custom architectures like DeepJet (Bols *et al.*, 2020) and JetVLAD (Bielčíková *et al.*, 2021), and on strange jet tagging (Erdmann, 2020; Nakai, Shih, and Thomas, 2020). Hadronic jets feature multiple tracks and extended energy deposition in both electromagnetic and hadronic calorimeters and can represent a primary source of backgrounds for electrons. Recent studies based on CNNs showed the advantage of using low-level calorimeter data represented as images in identifying electrons (Collado *et al.*, 2021). Electromagnetic showers have been classified using computer vision techniques that take advantage of lower level detector information (De Oliveira, Nachman, and Paganini, 2020). Finally, BNNs have been used for pion, kaon, and proton identification, with tests done on data generated for the BES II experiment (Ye, Jian, and Kai-En, 2008), combining multiple features from different detectors like the drift chamber, time of flight, and shower counter.

## 4. Event and signal classification

In collision-based experiments, events are often categorized by event type for analysis. Although selection algorithms differ across experiments, the selection is typically computationally expensive in traditional analyses. In high-luminosity experiments, real-time triggers are deployed to decide which events to store for analysis. For such algorithms, fast inference speed is required. In low-energy, low-luminosity collisions, such as those at rare isotope facilities, event selection post trigger is a computational bottleneck for large data detectors such as time projection chambers (Adamson *et al.*, 2016; Bradt, 2017).

A common task in scintillator detectors in low-energy experiments is to discriminate between the neutron and  $\gamma$  signals. Neural network analyses of pulse shapes have been shown to effectively discriminate between these signals (Doucet *et al.*, 2020).

In postexperiment analyses, deep ANNs and CNNs (Kuchera *et al.*, 2019; Gavalian *et al.*, 2020; Solli *et al.*, 2021) were used to classify events. Seeding networks with

pretrained architectures designed for image classification allow for fast training when events can be structured similar to images. Additionally, hierarchical clustering (Dalitz *et al.*, 2019) is used for track finding in time projection chambers (TPCs).

In low-energy neutrino experiments, ML techniques are typically used to differentiate different physics signal types or signals from backgrounds. One of the earliest uses of ANN to discern neutral-current from charged-current solar neutrino interactions in a heavy-water-Cherenkov detector was performed by the Sudbury Neutrino Observatory experiment (Brice, 1996).

Direct kinematic measurements of tritium  $\beta$  decays have historically been the most sensitive to investigate the neutrino mass scale (Formaggio, de Gouv  a, and Robertson, 2021). The next-generation tritium  $\beta$ -decay experiment Project 8 plans to measure the  $\beta$ -electron energy spectrum via cyclotron radiation emission spectroscopy (CRES) (Monreal and Formaggio, 2009). The geometry of the detector and its electromagnetic field configuration, as well as the dynamics of the  $\beta$  electrons, would give rise to different variations of the CRES signals. The Project 8 Collaboration is developing machine learning techniques to analyze and improve reconstructions of these CRES events (Ashtari Esfahani *et al.*, 2020).

To answer the question of whether neutrinos are their own antiparticles, i.e., Majorana fermions, there are recently completed, operating, and planned experiments to search for neutrinoless double-beta ( $0\nu\beta\beta$ ) decays in  $^{76}\text{Ge}$  (Alvis *et al.*, 2019; Agostini *et al.*, 2020; Abgrall *et al.*, 2021),  $^{100}\text{Mo}$  (Armstrong *et al.*, 2019),  $^{130}\text{Te}$  (Artusa *et al.*, 2015; Anderson *et al.*, 2021; Adams *et al.*, 2022),  $^{136}\text{Xe}$  (Gomez-Cadenas *et al.*, 2014; Gando *et al.*, 2016; Al Kharusi *et al.*, 2018; Anton *et al.*, 2019), and other isotopes. The observation of this lepton-number-violating decay mode, in which two electrons but no neutrinos are emitted, is evidence that neutrinos are Majorana fermions. As the current limit of this decay mode is on the order of  $10^{26}$  yr, experiments (whether they are large-scale liquid scintillator detectors, semiconductor ionization detectors, cryogenic bolometers, or liquid cryogenic or high-pressure gaseous TPCs) are deploying machine learning techniques to distinguish and remove backgrounds in the signal search region. The “single-site” signature of a  $0\nu\beta\beta$  decay signal is the simultaneous appearance from the same origin of two electrons whose energies add up to the decay’s  $Q$  value. Backgrounds tend to have “multisite” characteristics, as with external gamma rays that Compton scatter at multiple locations in the detector’s active volume. For comprehensive reviews of  $0\nu\beta\beta$  decays, see Dolinski, Poon, and Rodejohann (2019) and Agostini *et al.* (2022).

The use of high-purity germanium (HPGe) detectors enriched in  $^{76}\text{Ge}$  in  $0\nu\beta\beta$ -decay searches has a long history (Avignone and Elliott, 2019). The GERDA experiment pioneered using ANNs to identify single-site events in semicoaxial high-purity germanium detectors (Agostini *et al.*, 2013, 2019). The use of ANNs to differentiate single-site and multisite events in HPGe detectors was also developed by Caldwell *et al.* (2015), Holl *et al.* (2019), and Jany *et al.* (2021).

The NEXT experiment exploits the topological difference between a  $0\nu\beta\beta$ -decay event signal and backgrounds in its  $^{136}\text{Xe}$  high-pressure gas TPC using deep neural networks (Renner *et al.*, 2017), and specifically CNNs (Kekic *et al.*, 2021), for the identification of an event topology similar to that of a signal event. The ML tools developed by NEXT have also been adapted in the conceptual design of a  $^{82}\text{SeF}_6$  TPC for a  $0\nu\beta\beta$ -decay study (Nygren *et al.*, 2018).

In  $0\nu\beta\beta$ -decay experiments, an ample amount of the target isotope can be loaded in the scintillator, but the detector energy resolution is typically worse there than in other types of detectors. The KamLAND-ZEN experiment developed CNNs and RNNs to identify  $^{10}\text{C}$  from cosmic-ray spallation in the liquid scintillator loaded with  $^{136}\text{Xe}$  (Hayashida, 2019; Li *et al.*, 2019). Efforts in developing liquid scintillators that allow the separation of the Cherenkov light and the scintillation will facilitate ML to perform signal-background differentiation in  $0\nu\beta\beta$ -decay experiments or other multipurpose detectors (Fraker, 2018; Grusko, 2018; Askins *et al.*, 2020). Large water-Cherenkov detectors could also benefit from gadolinium loading to enhance their neutron detection capability for supernova detection and searches of the diffuse supernova neutrino background. The use of ML for background discrimination in such a detector has been studied (Maksimovi  , Nieslony, and Wurm, 2021).

## 5. Event reconstruction

For the precise knowledge of the kinematic variables of the deep-inelastic scattering process, various reconstruction methods are combined in collider experiments. Each method is using partial information from the scattered lepton and/or the hadronic final state of deep-inelastic scattering and has its own limitations. Recently it was shown in the H1 and ZEUS collider experiments as well as in simulations of a possible EIC detector that deep learning techniques to reconstruct the kinematic variables can serve as a rigorous method to combine and outperform existing reconstruction methods (Diefenthaler *et al.*, 2021; Arratia *et al.*, 2022).

## 6. Spectroscopy

Gamma-ray spectra may be used for isotope identification and fundamental nuclear structure studies. Timing resolution in high-purity germanium detectors has been optimized by fully connected CNN architectures (Gladen *et al.*, 2020). Deep, fully connected neural network architectures have been shown to successfully identify isotopes (Abdel-Aal and Al-Haddad, 1997; Medhat, 2012; Kamuda, Stinnett, and Sullivan, 2017; Jhung *et al.*, 2020) and fit peaks (Abdel-Aal, 2002) in gamma spectra. Machine learning has also been shown to estimate activity levels in spectra from gamma-emitting samples (Vigneron *et al.*, 1996; Abdel-Aal and Al-Haddad, 1997; Kamuda and Sullivan, 2019). Convolutional neural networks have demonstrated robustness to spectra with unidentified background channels and calibration drifts in the detectors (Kamuda, Zhao, and Huff, 2020). In addition, charged particle detection is routinely used for spectroscopy. For example, Bailey *et al.* (2021) used ML to analyze signals from double-sided silicon strip detectors to determine  $\alpha$  clustering.




FIG. 8. Left panels: Geant4 model of the dRICH detector. The full 3D (left panel) downstream and (right panel) upstream views of the dRICH detector. The BO strategy involves tuning eight main design parameters characterizing the geometry and the optical properties. Right panel:  $\pi/K$  separation power. The  $\pi/K$  separation as a number of  $\sigma$  is given as a function of the charged particle momentum. The improvement in the separation power with the approach is compared to the legacy baseline design. The curves are drawn with 68% confidence interval bands that are barely visible in the log plot. From [Cisbani \*et al.\*, 2020](#).

## C. Experimental design

### 1. Design for detector systems

Physics and detector simulations are critical for both the initial design and the optimization of complex subdetector systems in nuclear physics experiments. These systems are usually characterized by multiple parameters capable of tuning the mechanics, the geometry, and the optics of each component.

Traditionally the global design is studied and characterized after the subsystem prototypes are ready. In the subdetector design phase, constraints from the baseline full detector are taken into account. A well-known phenomenon observed in optimization problems with high-dimensional spaces is the so-called curse of dimensionality ([Houle \*et al.\*, 2010](#)), which corresponds to a combinatorial explosion of possible values to search. Indeed, the detector design optimization can be a large combinatorial problem characterized by accurate and computationally expensive simulations performed in Geant4 ([Agostinelli \*et al.\*, 2003](#)). In this context, ML offers different optimization strategies spanning from reinforcement learning ([Sutton and Barto, 2018](#)) to evolutionary algorithms ([Deb, 2001](#)). These new approaches can potentially “revolutionize the way experimental nuclear physics is currently done” ([Stevens \*et al.\*, 2020](#)). Among these approaches, BO ([Jones, Schonlau, and Welch, 1998](#); [Snoek, Larochelle, and Adams, 2012](#)) has gained popularity in detector design because it offers a derivative-free, principled approach to global optimization of noisy and computationally expensive black-box functions. An automated, highly parallelized, and self-consistent procedure has been developed and tested for a dual-radiator ring imaging Cherenkov (dRICH) design ([Cisbani \*et al.\*, 2020](#)), which has been considered as a case study. Eight main design parameters have been considered to improve the PID performance of the dRICH detector. Examples of design parameters are the refractive index and thickness of the aerogel radiator, the focusing mirror radius, its longitudinal and radial positions, and the three-dimensional

shifts of the photon sensor tiles with respect to the mirror center on a spherical surface.

Gaussian processes have been used for regression, and a surrogate model has been reconstructed as shown in Fig. 8. These studies not only resulted in a statistically significant improvement in the PID performance compared to an existing baseline design, they also shed light on the relevance of different features of the detector for the overall performance.

The future EIC is looking at systematically exploiting AI-based optimization strategies during the design and research and development (R&D) phase of the EIC detector ([Khalek \*et al.\*, 2021](#)).

A pipeline based on machine learning for the LHCb electromagnetic calorimeter R&D optimization has been proposed ([Boldyrev \*et al.\*, 2020, 2021](#)) to evaluate its operational characteristics and determine an optimal configuration with a gain in computational time. Different cell sizes have been considered to characterize the calorimeter granularity. Signal energy deposits in the calorimeter have been studied for different background conditions by changing the number of primary vertices. Energy and spatial reconstruction were based on gradient-boosted decision trees ([Chen and Guestrin, 2016](#)), with the regressor trained to minimize the difference between reconstructed and generated observables. Machine learning methods inside this pipeline are used to fine-tune the parameters of both simulations and reconstructions. The application of tools based on deep neural networks and modern automatic differentiation techniques to implement a full modeling of an experimental design (in order to achieve an end-to-end optimization) was described by [Baydin \*et al.\* \(2021\)](#).

### 2. Interface with theory

Bayesian experimental design provides a general framework to maximize the success of an experiment based on the best information available on the existing data, experimental conditions (including the amount of beam time available, the experimental setup, and the budgetary constraints), and




FIG. 9. BED of a proton Compton scattering experiment. Shown is the expected utility of proton differential cross section measurements defined as the gain in Shannon information based on the experiment. Chiral EFT has been used to predict the functional form of the scattering amplitude. Left panel measurements that do not include EFT truncation estimates. Right panel measurements that include the EFT uncertainty. The interior box marks the experimentally accessible regime. The vertical line marks the cusp at the pion-production threshold. The white circles show the optimal five-point design kinematics. Adapted from [Melendez, Furnstahl \*et al.\*, 2021](#).

theoretical models used to process and interpret the data. The goal of BED is to maximize the expected utility of the outcome. Formally this is done by introducing the utility function that is designed by taking into account all costs and benefits ([Phillips \*et al.\*, 2021](#)). Recent applications of BED were detailed by [Giuliani and Piekarewicz \(2021\)](#) and [Melendez, Furnstahl \*et al.\* \(2021\)](#). Figure 9 shows the utility function for a proton Compton scattering experiment interpreted using chiral EFT. As discussed by [Melendez, Furnstahl \*et al.\* \(2021\)](#), the effect of including EFT truncation errors is significant: it shifts the region of optimal utility to lower energies. [Giuliani and Piekarewicz \(2021\)](#) used the transfer function formalism in BED of experiments, aiming at the extraction of neutron densities.

## V. ACCELERATOR SCIENCE AND OPERATIONS

Research in modern ML techniques for accelerators is relatively new; however, it is an active cross-cutting effort among data scientists, computer scientists, control experts, and accelerator physicists. Machine learning applications are poised to play an important role in accelerator facilities by providing data-driven digital models (twins) for anomaly detection and prognostication, design optimization tools, and real-time operational control (tuning). There have been previous efforts to document opportunities ([Edelen \*et al.\*, 2018](#)) and advanced control ([Scheinker, Emma \*et al.\*, 2020](#)) in ML for accelerators. The following is an overview of some of these activities and new advancements since those works.

### A. ML-based surrogate models for accelerator models

Particle accelerators are complex nonlinear systems that require sophisticated simulation software to capture this dynamic. Owing to their complexity and evolving conditions, it is natural to explore modern data science techniques to

provide surrogate models and/or a fully realized digital twin ([Grieves and Vickers, 2017](#); [Fuller \*et al.\*, 2020](#)). By developing digital models, researchers are able to conduct exploratory research without impacting the physical system. The use of ML to develop these digital models provides the ability to capture nonlinear complex dynamics using techniques such as GP regression ([Williams and Rasmussen, 1996](#)), quantile regression models ([Koenker and Bassett, 1978](#)), and RNNs ([Rumelhart, Hinton, and Williams, 1986](#)). These techniques can also be used to predict the future condition of the accelerator system and study anomalies, and to forecast component fatigue and failures.

For simulation studies, surrogate models are the most popular examples of ML methods used in the accelerator community to map between various accelerator parameters and beam properties at speeds that are orders of magnitude faster than computationally expensive physics models and for optimization studies ([Emma \*et al.\*, 2018](#); [Li \*et al.\*, 2018](#); [Edelen \*et al.\*, 2020](#); [Scheinker, Gessner \*et al.\*, 2020](#); [Hanuka \*et al.\*, 2021](#); [Krančević \*et al.\*, 2021](#); [Zhu \*et al.\*, 2021](#)). Additionally, purely data-driven approaches have been explored to model accelerator components, such as the Fermi National Accelerator Laboratory booster system ([John \*et al.\*, 2021](#); [Kafkes and Schram, 2021](#)).

### B. Anomaly detection and classification

There is a growing amount of research using ML to improve accelerator operations by detecting anomalies and classifying them. These studies explore several techniques that identify and elucidate the source of the anomalies and ultimately prevent them altogether in the future. Although it can be relatively easy to avoid these failures when they are actually happening, it would be far better to avoid them all together. Forecasting these failures can be difficult to predict accurately with an acceptable false positive rate. Predictive diagnostics are important for improving operational efficiency; however, they can also serve as actionable precursors to control systems. There is a growing body of research in this domain; as such, we present a select few recent results.

Machine learning models were used to classify cavity faults in the Continuous Electron Beam Accelerator Facility (CEBAF) ([Solopova \*et al.\*, 2019](#); [Tennant \*et al.\*, 2020](#)). Traditionally, accelerator operators are able to analyze time series data, identify which cavities faulted, and then classify the fault type; however, manually labeling the data is laborious and time consuming. To accelerate the process, [Tennant \*et al.\* \(2020\)](#) developed ML models to identify the faulty cavity and classification fault type. The results showed that the cavity identification and fault classification models have accuracies of 84.9% and 78.2%, respectively.

[Resic, Seviour, and Blokland \(2020\)](#) used several machine learning techniques to predict machine failures at the Spallation Neutron Source facility of Oak Ridge National Laboratory using beam current measurements before the faults actually occurred. They evaluated these pulses using a common set of ML-based classification techniques and showed that they could identify accelerator failures prior to actually failing with almost 80% accuracy. The results were further improved by tuning the classifier parameters and using

pulse properties for refining datasets, leading to nearly 92% accuracy in the classification of bad pulses.

Li, Zacharias *et al.* (2021) examined the potential to predict interlocks (reflecting beam shutoff) using multiple measurements along the accelerator. Recurrence-plot-based CNNs (RPCNNs) were used to convert the measurements into time recurrence plots that are used as inputs to CNNs for classification. Comparisons with EMB methods like random forests indicate that the performance of the random forests and RPCNNs are comparable; however, the RPCNNs are more successful at identifying anomalies that evolve over time.

A novel uncertainty-aware Siamese model to predict upcoming faults (Blokland *et al.*, 2021) that combines the use of uncertainty quantification and a deep Siamese architecture was developed to predict the similarity between beam pulses and provide an uncertainty that includes out of domain errors. The results show that this model outperforms previous results by 4 times in operational regions of interest. Additionally, the model performed better than previous results even for anomalies it had not encountered before. The inclusion of uncertainty quantification is an important step toward having robust and safe machine learning models for operational facilities.

### C. Control optimization

Like the efforts in the areas of anomaly detection and classification, the field of accelerator optimization has recently seen advancements through the application of techniques such as BO, genetic algorithms, particle swarm, and reinforcement learning. By utilizing online data from existing fast diagnostics such as beam position monitors, beam loss monitors, and radio-frequency cavity signals, an effort has been made to develop ML-based controllers. Recent demonstrations include BO, RL, and GPs for accelerator tuning (McIntire *et al.*, 2016; Hao *et al.*, 2019; Li, Rainer, and Cheng, 2019; Duris *et al.*, 2020; Shalloo *et al.*, 2020; Miskovich *et al.*, 2021; Roussel and Edelen, 2021; Wang, Bagri *et al.*, 2021) and polynomial chaos expansion-based surrogate models for uncertainty quantification (Adelmann, 2019). Particle swarm techniques have been evaluated to optimize the tuning of aperiodic ion transport lines and for advanced particle separators (Amthor *et al.*, 2018). The use of multiobjective genetic algorithm-(MOGA-) based optimization was used to minimize competing objectives steering operations of CEBAF linear particle accelerators (linacs) (Terzić *et al.*, 2014) and showed that the dynamic heat load can be reduced by over 20% while keeping the same trip rate. Additional studies using MOGA for accelerator optimization were conducted by Li *et al.* (2018) and Neveu *et al.* (2019). Reinforcement learning tools have also been developed to optimize various elements of the accelerator system (Gao *et al.*, 2019; Bruchon *et al.*, 2020; Hirlaender and Bruchon, 2020; Kain *et al.*, 2020; O’Shea, Bruchon, and Gaio, 2020; John *et al.*, 2021). Additionally, there is new research on transferring the RL policy models to a field-programmable gate array to provide a low latency control response time (John *et al.*, 2021). The accuracy of the previously described ML methods for accelerators quickly degrades for systems that change with time, for which

previously collected training data are no longer accurate. A major open problem faced by the ML community is the challenge of developing ML tools for complex systems where the underlying dynamics is evolving (Shimodaira, 2000; Sugiyama and Kawanabe, 2012; Kurle *et al.*, 2020; Dramsch, Lüthje, and Christensen, 2021). For systems that change slowly with time and for which gathering large amounts of new data are feasible without interrupting operations, it is possible to utilize transfer learning techniques (Goodfellow, Bengio, and Courville, 2016).

The most common transfer learning technique is to update or partially retrain the model using new data (Calandra *et al.*, 2012; Koeswiady *et al.*, 2018; Kurle *et al.*, 2020). Another transfer learning approach is domain transform, in which smaller neural networks are developed using experimental data and used as the input layer of trained ANNs (Zeiler *et al.*, 2010). These transfer learning techniques can be applied to GP-based algorithms in which the prior and parameter correlations are first estimated using simulation studies and then fine-tuned with experimental data. Such transfer learning techniques have been demonstrated on a wide range of systems, including cross-modal implementations (Castrejón *et al.*, 2016) and for electron backscatter diffraction (Shen *et al.*, 2019).

### D. Adaptive ML for nonstationary systems

For accelerators, new data can be acquired in many cases in real time and used to quickly update or retrain specific layers of neural networks. However, in other cases repetitive retraining is not always an option. For example, if beam or accelerator characteristics change significantly, diagnostics such as quadrupole magnet scan-based emittance measurements or wire-scan beam profile measurements can be a time-consuming procedure that can interrupt operations. For quickly time-varying systems adaptive feedback techniques exist that are model independent and automatically compensate for nonmodeled disturbances and changes. Novel adaptive feedback algorithms have been developed to tune large groups of parameters based on noisy scalar measurements with analytic proofs of convergence and analytically known guarantees for parameter update rates, making them especially well suited for particle accelerator problems (Scheinker, 2013; Scheinker and Krstić, 2017). Adaptive methods have been applied in real time to changing accelerators to maximize the free electron laser output power (Scheinker *et al.*, 2019) for real-time on-line multiobjective optimization (Scheinker, Hirlaender *et al.*, 2020), for non-invasive diagnostics (Scheinker and Gessner, 2015), and for on-line RL to learn optimal feedback control policies directly from data (Scheinker and Scheinker, 2021). A limitation of local adaptive methods is getting stuck in local minima in high-dimensional parameter spaces.

Adaptive ML and continuous learning is an area of active research combining the robustness of model-independent algorithms with the global properties of ML tools such as CNNs. The latter have been combined with adaptive feedback for fast automatic longitudinal phase space control of time-varying electron beams (Scheinker *et al.*, 2018). Such adaptive ML tools have the potential to enable truly autonomous

accelerator controls and diagnostics that automatically respond to nonmodeled changes and disturbances in real time and thereby keep the accelerator performance (beam energy and energy spread, beam loss, phase space quality, etc.) at a global optimal.

## VI. NUCLEAR DATA

The “nuclear data pipeline” represents the interconnected steps wherein data are compiled, evaluated, processed, and validated for end-user applications (Romano *et al.*, 2020). Evaluations, the most labor-intensive step in the pipeline, provide a recommended “best” data value (Capote, Smith, and Trkov, 2010) by combining new measurements, previous measurements, and nuclear model predictions. In some cases, it can take several years for data to pass fully through the pipeline, limited by manpower and, in some cases, outdated methodology. Machine learning has the potential to significantly improve each step of the pipeline (compilation, evaluation, processing, validation, and dissemination), which may notably reduce the time lag from data measurement to incorporation into standardized databases [like the Evaluated Nuclear Data File (ENDF) (Brown *et al.*, 2018)] that are critical for basic and applied research. Machine learning can also facilitate the creation of surrogate models or emulators that may improve the extraction of physics information from data measurements, as well as improving the predictability of evaluation models.

### A. Overhauling the nuclear data pipeline

Creating a new ENDF release is a time-consuming and complex process whereby the latest experimental reaction measurements, theoretical reaction calculations, theory-generated reference parameter sets, and integral and benchmark experiments are combined and adjusted in an iterative manner for optimum consistency. An effort is under way (Schnabel *et al.*, 2020, 2021) to build software “containers” to hold all of these components, as well as the current evaluations of individual reactions. By appropriately nesting and interlinking these containers to mimic portions of the nuclear data pipeline (e.g., cross section measurements and models are linked together to produce an evaluation of one reaction, which is linked to other reactions to produce an evaluated library), they can be treated as interlinked nodes in a Bayesian network (BN). GPs can then be used to automatically and self-consistently update the BN components (including the output of a new ENDF reaction data library) when any of the components are updated. The test cases to date have utilized a sparse GP construction integrated into the BN framework to enable modeling of energy-dependent cross sections, physics model deficiencies, and energy-dependent systematic experimental errors. This would be a completely new approach to update the ENDF, and one that could be quickly adapted to incorporate new benchmark experiment types, new theory codes, and new techniques to evaluate individual reactions. When completed, this approach could be modified for use in other fields that closely interlink experiment, theory, and benchmarks.

### B. Improving compilations and evaluations

One utilization of ML to improve evaluations is through a robust identification of outlying data points and problematic datasets. A recent example (Neudecker *et al.*, 2020) was the identification of a problematic  $^{19}\text{F}$  neutron inelastic cross section in the ENDF database via the use of EMB-like random forest regression combined with a Shapley additive explanations (SHAP) feature importance metric (Lundberg and Lee, 2017) of predicted effective neutron multiplication factor  $k_{\text{eff}}$  of critical assemblies. The ML approach identified this problematic cross section, which was missed with traditional validation methods; this technique may be applicable for validating other data libraries against particular features of an end-user application. Another recent study (Whewell, Grosskopf, and Neudecker, 2020) focused on utilizing a variety of ML approaches, including SVMs, LR, and EMB, to find relationships between outlying measurements and underlying details of the experiments used in an evaluation (backgrounds, sample backing and impurities, detector type, incident beam type, etc.). Such an approach, which provides a robust method to reject outlying measurements and guide future experiments, was used in this specific case to improve the evaluation of the  $^{239}\text{Pu}(n, f)$  reaction. Random forests were also used (Neudecker *et al.*, 2021) to combine differential data (differential cross sections), application-specific integral data ( $k_{\text{eff}}$  and neutron-leakage spectra from pulsed-sphere experiments), and nuclear theory to isolate likely root causes of disagreements between integral data and predictions using differential data, to select preferred differential datasets that give better agreement with integral measurements, and to specify which differential or theory developments can best reproduce integral results.

Machine learning can be used to provide quantified interpolations and extrapolations of nuclear data; see the discussion on ML for data mining in Sec. III.A. A good example is offered by the study of giant dipole resonances (GDRs) (Bai *et al.*, 2021). ANNs were first used to classify nuclei into groups with one or two GDR peaks, then two multitask learning neural nets (Zhang and Yang, 2021) were used to learn both GDR energies and widths. The multitask learning approach enables multiple related tasks to be learned simultaneously in a way that optimally utilizes the data for each task. As a result, the accuracy of predictions for measured nuclei was doubled compared to other approaches, and then the unmeasured properties of GDRs of nuclei near the  $\beta$ -stability line were predicted. Another study (Nobre *et al.*, 2021) used ten different ML classifiers with a SHAP metric to automate and correct the assignment of spins to neutron resonances, a critical input to BO used in evaluations.

As discussed in Sec. III.A, UQ in evaluations have also been improved by learning discrepancies from existing theoretical models. Moreover, ML can be used to design experiments that address nuclear data gaps for particular applications [such as criticality experiments (Michaud *et al.*, 2019)]. The new data obtained can be essential for validation efforts in the pipeline, which can provide critical adjustments of evaluations.

Machine learning is also being used to help extract knowledge from published documents. Here CNNs, combined with

edge detection techniques, are beginning to automate the extraction of data (tables, plots, and numbers) from publications (Soto, 2019). Natural language processing (NLP), widely used to process text, is a ML-enhanced textual analytics approach that is now a central ingredient in efforts to revamp numerous standardized databases; this includes extracting keywords from documents as needed for the Nuclear Science References bibliographic database (Pritychenko *et al.*, 2011). The next steps in NLP utilization in nuclear data may focus on the extraction of meaning (i.e., semantics) from documents. In this approach, data and concepts could be characterized as “high value,” and certain theoretical and experimental investigations could be recommended based on latent knowledge in the literature (Tshitoyan *et al.*, 2019).

### C. Building emulators and surrogate models

Phenomenological models form the foundation of many evaluation approaches because they are computationally inexpensive and often require few input parameters. As a result, the power to extrapolate evaluations to unmeasured nuclei, to perform self-consistency checks across the nuclear chart, and to accurately capture the correlations between different nuclear properties is limited. Machine learning has the potential to dramatically improve this by creating emulators or surrogate models that require similar computational resources as phenomenological models but capture the physics of full models. A recent example from reactor physics (Radaideh and Kozlowski, 2020) may provide a blueprint for similar advances in nuclear data. Here deep learning ANNs were used to perform UQ, sensitivity analyses, and uncertainty propagation for nuclear reactors by replacing high-fidelity reactor simulations with surrogate models.

## VII. SUMMARY AND PERSPECTIVES

In recent years, ML techniques have gained considerable traction in scientific discovery. In particular, applications and techniques for so-called fast ML, i.e., high-performance ML methods applied to real-time experimental data processing, hold great promise for enhancing scientific discoveries in many different disciplines (Deiana *et al.*, 2021). These developments cover a broad mix of rapidly evolving fields, from the development of ML techniques to computer and hardware architectures. For a field like nuclear physics, which covers a large range of energy and length scales, spanning from the smallest constituents of matter to the physics of dense astronomical objects such as neutron stars, AI and ML techniques offer possibilities for new discoveries and deeper insights.

This Colloquium has summarized present and planned applications of ML techniques in experimental and theoretical nuclear physics research. The vast range of scales is also indicated by new and planned nuclear physics facilities worldwide, where opportunities to incorporate ML methods are expected to play an important role in the justification and design of experiments, and during operations. In our overview, we have presented several recent experimental developments, including detector control, experimental design simulations,

and accelerator operations. Furthermore, ML techniques play a central role in theoretical nuclear physics and a growing role in the evaluation of nuclear data. Nuclear theory, in particular, has seen an explosion in the application of ML methods in the last few years. Note, however, that this Colloquium presents just a snapshot of ML in nuclear physics as of today. New ML approaches are continually being introduced at a pace that is difficult to keep up with. We thus expect that over the next decades ML will play a significant role in leveraging technologies that are at the frontier of computational science and data science.

Owing to the broad range of scales, nuclear physics is a field where the dimensions of the problems studied quickly exceed the capabilities of traditional computational methods. Machine learning techniques offer promising paths to dimension reductions. Traditionally, many of the standard ML methods have focused on making predictions and finding correlations in the datasets. However, as presented in this Colloquium, to quantify errors and find causations also requires the possibility of being able to determine models for probability distributions. In both experiment and theory there are clear indications that statistical learning methods offer new perspectives for future research directions. Research in statistical learning techniques for both supervised and unsupervised learning, combined with fast ML methods and similar developments, has the potential to change the landscape of nuclear physics.

## ACKNOWLEDGMENTS

The work of A. B, M. D, N. S, M. S, and V. Z. is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No. DE-AC05-06OR23177, under which Jefferson Science Associates, LLC, operates the Thomas Jefferson National Accelerator Facility. N. S. is also supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, in the Early Career Program. The work of C. F. is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Grant No. DE-SC0019999. The work of M. H. J. is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Grant No. DE-SC0021152 and U.S. National Science Foundation (NSF) Grants No. PHY-1404159 and No. PHY-2013047. T. H. is supported by NSF Grant No. PHY-2012430. The work of M. P. K. is supported by the U.S. National Science Foundation under Grant No. PHY-2012865. The work of D. L. is partially supported by the U.S. Department of Energy (Grants No. DE-SC0013365 and No. DE-SC0021152) and the NUCLEI SciDAC-4 Collaboration (Grant No. DE-SC0018083). The work of W. N. is partially supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Awards No. DE-SC0013365 and No. DE-SC0018083 (NUCLEI SciDAC-4 Collaboration), and by the National Science Foundation CSSI program under Grant No. 2004601 (BAND Collaboration). K. O. is supported by Jefferson Science Associates, LLC, under U.S. DOE Contract No. DE-AC05-06OR23177, by U.S. DOE Grant No. DE-FG02-04ER41302, and by Center for Nuclear Femtography Grants No. C2-2020-FEMT-006

and No. C2019-FEMT-002-05. The work of A. P. and X.-N. W. is supported by Lawrence Berkeley National Laboratory under U.S. Department of Energy Federal Prime Agreement No. DE-AC02-05CH11231. The work of A. S. is supported by the U.S. Department of Energy (Grant No. 89233218CNA000001) and by the Los Alamos National Laboratory LDRD Program (Project No. 20210595DR). The work of M. S. S. is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Grant No. DE-AC05-00OR22725. The work of L.-G. P. is supported by the NSFC under Grants No. 11861131009 and No. 12075098.

## REFERENCES

Abdel-Aal, R., 2002, “Comparison of algorithmic and machine learning approaches for the automatic fitting of Gaussian peaks,” *Neural Comput. Appl.* **11**, 17–29.

Abdel-Aal, R., and M. Al-Haddad, 1997, “Determination of radio-isotopes in gamma-ray spectroscopy using abductive machine learning,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **391**, 275–288.

Abgrall, N., *et al.* (LEGEND Collaboration), 2021, “The large enriched germanium experiment for neutrinoless  $\beta\beta$  decay: LEGEND-1000 preconceptual design report,” [arXiv:2107.11462](https://arxiv.org/abs/2107.11462).

Accardi, A., L. T. Brady, W. Melnitchouk, J. F. Owens, and N. Sato, 2016, “Constraints on large- $x$  parton distributions from new weak boson production and deep-inelastic scattering data,” *Phys. Rev. D* **93**, 114017.

Acharya, B., and S. Bacca, 2021, “Gaussian process error modeling for chiral effective-field-theory calculations of  $np \leftrightarrow d\gamma$  at low energies,” [arXiv:2109.13972](https://arxiv.org/abs/2109.13972).

Adams, C., G. Carleo, A. Lovato, and N. Rocco, 2021, “Variational Monte Carlo Calculations of  $A \leq 4$  Nuclei with an Artificial Neural-Network Correlator Ansatz,” *Phys. Rev. Lett.* **127**, 022502.

Adams, D. Q., *et al.*, 2022, “Search for Majorana neutrinos exploiting millikelvin cryogenics with CUORE,” *Nature (London)* **604**, 53–58.

Adamson, P., *et al.* (NO $\nu$ A Collaboration), 2016, “First Measurement of Electron Neutrino Appearance in NO $\nu$ A,” *Phys. Rev. Lett.* **116**, 151806.

Adelmann, A., 2019, “On nonintrusive uncertainty quantification and surrogate model construction in particle accelerator modeling,” *SIAM-ASA J. Uncertain.* **7**, 383–416.

Agarawal, V., S. Roy, A. Chakraborty, and R. Maitra, 2021, “Accelerating coupled cluster calculations with nonlinear dynamics and supervised machine learning,” *J. Chem. Phys.* **154**, 044110.

Agostinelli, S., *et al.*, 2003, “Geant4—A simulation toolkit,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250–303.

Agostini, M., G. Benato, J. A. Detwiler, J. Menéndez, and F. Vissani, 2022, “Toward the discovery of matter creation with neutrinoless double-beta decay,” [arXiv:2202.01787](https://arxiv.org/abs/2202.01787).

Agostini, M., *et al.*, 2013, “Pulse shape discrimination for GERDA phase I data,” *Eur. Phys. J. C* **73**, 2583.

Agostini, M., *et al.*, 2019, “Probing Majorana neutrinos with double- $\beta$  decay,” *Science* **365**, 1445–1448.

Agostini, M., *et al.* (GERDA Collaboration), 2020, “Final Results of GERDA on the Search for Neutrinoless Double- $\beta$  Decay,” *Phys. Rev. Lett.* **125**, 252502.

Akkoyun, S., 2020, “Estimation of fusion reaction cross-sections by artificial neural networks,” *Nucl. Instrum. Methods Phys. Res., Sect. B* **462**, 51–54.

Akkoyun, S., T. Bayram, S. O. Kara, and N. Yildiz, 2013, “Consistent empirical physical formulas for potential energy curves of  $^{38-66}\text{Ti}$  isotopes by using neural networks,” *Phys. Part. Nucl. Lett.* **10**, 528–534.

Akkoyun, S., N. Laouet, and F. Benrachi, 2021, “Improvement studies of an effective interaction for  $N = Z$   $sd$ -shell nuclei by neural networks,” *Bulg. J. Phys.* **48**, 245–256, [https://www.bjp-bg.com/papers/bjp2021\\_3\\_245-256.pdf](https://www.bjp-bg.com/papers/bjp2021_3_245-256.pdf).

Albergo, M. S., G. Kanwar, S. Racanière, D. J. Rezende, J. M. Urban, D. Boyd, K. Cranmer, D. C. Hackett, and P. E. Shanahan, 2021, “Flow-based sampling for fermionic lattice field theories,” *Phys. Rev. D* **104**, 114507.

Albergo, M. S., G. Kanwar, and P. E. Shanahan, 2019, “Flow-based generative models for Markov chain Monte Carlo in lattice field theory,” *Phys. Rev. D* **100**, 034515.

Alekhin, S., J. Blümlein, and S. Moch, 2018, “NLO PDFs from the ABMP16 fit,” *Eur. Phys. J. C* **78**, 477.

Alexandrou, C., G. Iannelli, K. Jansen, and F. Manigrasso, 2020, “Parton distribution functions from lattice QCD using Bayes-Gauss-Fourier transforms,” *Phys. Rev. D* **102**, 094508.

Alexandru, A., P. F. Bedaque, H. Lamm, and S. Lawrence, 2017, “Deep learning beyond Lefschetz thimbles,” *Phys. Rev. D* **96**, 094505.

Alexandru, A., P. F. Bedaque, H. Lamm, and S. Lawrence, 2018, “Finite-density Monte Carlo calculations on sign-optimized manifolds,” *Phys. Rev. D* **97**, 094510.

Alexandru, A., P. F. Bedaque, H. Lamm, S. Lawrence, and N. C. Warrington, 2018, “Fermions at Finite Density in  $2 + 1$  Dimensions with Sign-Optimized Manifolds,” *Phys. Rev. Lett.* **121**, 191602.

Al Kharusi, S., *et al.* (nEXO Collaboration), 2018, “nEXO preconceptual design report,” [arXiv:1805.11142](https://arxiv.org/abs/1805.11142).

Al-Mamun, M., A. W. Steiner, J. Näätälä, J. Lange, R. O’Shaughnessy, I. Tews, S. Gandolfi, C. Heinke, and S. Han, 2021, “Combining Electromagnetic and Gravitational-Wave Constraints on Neutron-Star Masses and Radii,” *Phys. Rev. Lett.* **126**, 061101.

Alvis, S. I., *et al.* (MAJORANA Collaboration), 2019, “A search for neutrinoless double-beta decay in  $^{76}\text{Ge}$  with 26 kg-yr of exposure from the MAJORANA Demonstrator,” *Phys. Rev. C* **100**, 025501.

Ameli, F., *et al.*, 2021, “Streaming readout of the CLAS12 forward tagger using TriDAS and JANA2,” [arXiv:2104.11388v1](https://arxiv.org/abs/2104.11388v1).

Amthor, A., Z. Schillaci, D. Morrissey, M. Portillo, S. Schwarz, M. Steiner, and C. Sumithrarachchi, 2018, “Experimental test of an online ion-optics optimizer,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **895**, 90–99.

Anderson, M. R., *et al.* (SNO+ Collaboration), 2021, “Development, characterisation, and deployment of the SNO+ liquid scintillator,” *J. Instrum.* **16**, P05009.

Anton, G., *et al.* (EXO-200 Collaboration), 2019, “Search for Neutrinoless Double- $\beta$  Decay with the Complete EXO-200 Data-set,” *Phys. Rev. Lett.* **123**, 161802.

Apolinário, L., N. Castro, M. C. Romão, J. Milhano, R. Pedro, and F. Peres, 2021, “Deep learning for the classification of quenched jets,” [arXiv:2106.08869v1](https://arxiv.org/abs/2106.08869v1).

Armstrong, W. R., *et al.* (CUPID Collaboration), 2019, “CUPID pre-CDR,” [arXiv:1907.09376](https://arxiv.org/abs/1907.09376).

Arratia, M., D. Britzger, O. Long, and B. Nachman, 2022, “Reconstructing the kinematics of deep inelastic scattering with deep learning,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **1025**, 166164.

Artusa, D. R., *et al.* (CUORE Collaboration), 2015, “Searching for neutrinoless double-beta decay of  $^{130}\text{Te}$  with CUORE,” *Adv. High Energy Phys.* **879871**.

Ashtari Esfahani, A., *et al.*, 2020, “Cyclotron radiation emission spectroscopy signal classification with machine learning in Project 8,” *New J. Phys.* **22**, 033004.

Askins, M., *et al.* (THEIA Collaboration), 2020, “THEIA: An advanced optical neutrino detector,” *Eur. Phys. J. C* **80**, 416.

Avignone, F. T., and S. R. Elliott, 2019, “The search for double beta decay with germanium detectors: Past, present, and future,” *Front. Phys.* **7**, 6.

Ayriyan, A., D. Blaschke, A. G. Grunfeld, D. Alvarez-Castillo, H. Grigorian, and V. Abgaryan, 2021, “Bayesian analysis of multi-messenger M-R data with interpolated hybrid EoS,” *Eur. Phys. J. A* **57**, 318.

Ayyad, Y., W. Mittig, D. Bazin, S. Beceiro-Novo, and M. Cortesi, 2018, “Novel particle tracking algorithm based on the random sample consensus model for the Active Target Time Projection Chamber (AT-TPC),” *Nucl. Instrum. Methods Phys. Res., Sect. A* **880**, 166–173.

Bachtis, D., G. Aarts, and B. Lucini, 2020, “Mapping distinct phase transitions to a neural network,” *Phys. Rev. E* **102**, 053306.

Bai, J., Z. Niu, B. Sun, and Y. Niu, 2021, “The description of giant dipole resonance key parameters with multitask neural networks,” *Phys. Lett. B* **815**, 136147.

Bailey, S., *et al.*, 2021, “The identification of  $\alpha$ -clustered doorway states in  $^{44,48,52}\text{Ti}$  using machine learning,” *Eur. Phys. J. A* **57**, 108.

Ball, R. D., L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, and M. Ubiali, 2010, “A first unbiased global NLO determination of parton distributions and their uncertainties,” *Nucl. Phys. B* **838**, 136–206.

Barsotti, R., and M. R. Shepherd, 2020, “Using machine learning to separate hadronic and electromagnetic interactions in the GlueX forward calorimeter,” *J. Instrum.* **15**, P05021.

Bass, S. A., A. Bischoff, J. A. Maruhn, H. Stoecker, and W. Greiner, 1996, “Neural networks for impact parameter determination,” *Phys. Rev. C* **53**, 2358–2363.

Baydin, A. G., *et al.*, 2021, “Toward machine learning optimization of experimental design,” *Nucl. Phys. News* **31**, 25–28.

Bedaque, P., *et al.*, 2021, “A.I. for nuclear physics,” *Eur. Phys. J. A* **57**, 100.

Bender, M., P.-H. Heenen, and P.-G. Reinhard, 2003, “Self-consistent mean-field models for nuclear structure,” *Rev. Mod. Phys.* **75**, 121–180.

Bernhard, J. E., J. S. Moreland, and S. A. Bass, 2019, “Bayesian estimation of the specific shear and bulk viscosity of quark-gluon plasma,” *Nat. Phys.* **15**, 1113–1117.

Bernhard, J. E., J. S. Moreland, S. A. Bass, J. Liu, and U. Heinz, 2016, “Applying Bayesian parameter estimation to relativistic heavy-ion collisions: Simultaneous characterization of the initial state and quark-gluon plasma medium,” *Phys. Rev. C* **94**, 024907.

Bielčíková, J., R. K. Elayavalli, G. Ponimatkin, J. H. Putschke, and J. Sivic, 2021, “Identifying heavy-flavor jets using vectors of locally aggregated descriptors,” *J. Instrum.* **16**, P03017.

Bishop, C. M., 2006, *Pattern Recognition and Machine Learning* (Springer-Verlag, Berlin).

Blokland, W., M. Schram, K. Rajput, P. Ramuhalli, C. Peters, Y. Yucesan, A. Zhukov, and T. Jeske, 2021, “Uncertainty aware anomaly detection to predict errant beam pulses in the SNS accelerator,” *arXiv:2110.12006*.

Blücher, S., L. Kades, J. M. Pawłowski, N. Strothoff, and J. M. Urban, 2020, “Towards novel insights in lattice field theory with explainable machine learning,” *Phys. Rev. D* **101**, 094507.

Blum, T., P. Boyle, T. Izubuchi, L. Jin, C. Jung, C. Lehner, M. Lin, Lin, and A. Tomiya, 2021, “Machine learning and lattice QCD,” [https://www.snowmass21.org/docs/files/summaries/CompF/SNOWMASS21-CompF3\\_CompF2-TF5\\_TF0\\_Akio\\_Tomiya-131.pdf](https://www.snowmass21.org/docs/files/summaries/CompF/SNOWMASS21-CompF3_CompF2-TF5_TF0_Akio_Tomiya-131.pdf).

Boldyrev, A., D. Derkach, F. Ratnikov, and A. Shevelev, 2020, “ML-assisted versatile approach to calorimeter R&D,” *J. Instrum.* **15**, C09030.

Boldyrev, A., D. Derkach, F. Ratnikov, and A. Shevelev, 2021, “Machine learning in calorimeter optimization,” *J. Phys. Conf. Ser.* **1740**, 012047.

Bollapragada, R., M. Menickelly, W. Nazarewicz, J. O’Neal, P.-G. Reinhard, and S. M. Wild, 2021, “Optimization and supervised machine learning methods for fitting numerical physics models without derivatives,” *J. Phys. G* **48**, 024001.

Bols, E., J. Kieseler, M. Verzetti, M. Stoye, and A. Stakia, 2020, “Jet flavour classification using deepjet,” *J. Instrum.* **15**, P12012.

Boyda, D., G. Kanwar, S. Racanière, D. J. Rezende, M. S. Albergo, K. Cranmer, D. C. Hackett, and P. E. Shanahan, 2021, “Sampling using  $SU(N)$  gauge equivariant flows,” *Phys. Rev. D* **103**, 074504.

Boyda, D. L., M. N. Chernodub, N. V. Gerasimeniuk, V. A. Goy, S. D. Liubimov, and A. V. Molochkov, 2021, “Finding the deconfinement temperature in lattice Yang-Mills theories from outside the scaling window with machine learning,” *Phys. Rev. D* **103**, 014509.

Bradt, J. W., 2017, “Measurement of isobaric analogue resonances of  $^{47}\text{Ar}$  with the active target time projection chamber,” Ph.D. thesis (Michigan State University).

Brice, S. J., 1996, “Monte Carlo and analysis techniques for the Sudbury Neutrino Observatory,” Ph.D. thesis (University of Oxford).

Bringewatt, J., N. Sato, W. Melnitchouk, J.-W. Qiu, F. Steffens, and M. Constantinou, 2021, “Confronting lattice parton distributions with global QCD analysis,” *Phys. Rev. D* **103**, 016003.

Brown, D., *et al.*, 2018, “ENDF/B-VIII.0: The 8th major release of the nuclear reaction data library with CIELO-project cross sections, new standards and thermal scattering data,” *Nucl. Data Sheets* **148**, 1–142.

Bruchon, N., G. Fenu, G. Gaio, M. Lonza, F. H. O’Shea, F. A. Pellegrino, and E. Salvato, 2020, “Basic reinforcement learning techniques to control the intensity of a seeded free-electron laser,” *Electronics* **9**, 781.

Bursa, F., and M. Kroyter, 2018, “A simple approach towards the sign problem using path optimisation,” *J. High Energy Phys.* **12**, 054.

Calandra, R., T. Raiko, M. P. Deisenroth, and F. M. Pouzols, 2012, “Learning deep belief networks from non-stationary streams,” in *Proceedings of the 22nd International Conference on Artificial Neural Networks, Lausanne, 2012*, edited by A. E. P. Villa, W. Duch, P. Érdi, F. Masulli, and G. Palm (Springer, New York), pp. 379–386.

Caldwell, A., F. Cossavella, B. Majorovits, D. Palioselitis, and O. Volynets, 2015, “Signal recognition efficiencies of artificial neural-network pulse-shape discrimination in HPGe  $0\nu\beta\beta$ -decay searches,” *Eur. Phys. J. C* **75**, 350.

Cammarota, J., L. Gamberg, Z.-B. Kang, J. A. Miller, D. Pitonyak, A. Prokudin, T. C. Rogers, and N. Sato (Jefferson Lab Angular Momentum Collaboration), 2020, “Origin of single transverse-spin asymmetries in high-energy collisions,” *Phys. Rev. D* **102**, 054002.

Cao, S., *et al.* (JETSCAPE Collaboration), 2021, “Determining the jet transport coefficient  $\hat{q}$  from inclusive hadron suppression measurements using Bayesian parameter estimation,” *Phys. Rev. C* **104**, 024905.

Capano, C. D., I. Tews, S. M. Brown, B. Margalit, S. De, S. Kumar, D. A. Brown, B. Krishnan, and S. Reddy, 2020, “Stringent

constraints on neutron-star radii from multimessenger observations and nuclear theory," *Nat. Astron.* **4**, 625–632.

Capote, R., D. L. Smith, and A. Trkov, 2010, "Nuclear data evaluation methodology including estimates of covariances," *EPJ Web Conf.* **8**, 04001.

Carleo, G., I. Cirac, K. Cranmer, L. Daudet, M. Schuld, N. Tishby, L. Vogt-Maranto, and L. Zdeborová, 2019, "Machine learning and the physical sciences," *Rev. Mod. Phys.* **91**, 045002.

Carnini, M., and A. Pastore, 2020, "Trees and forests in nuclear physics," *J. Phys. G* **47**, 082001.

Castrejón, L., Y. Aytar, C. Vondrick, H. Pirsiavash, and A. Torralba, 2016, "Learning aligned cross-modal representations from weakly aligned data," in *Proceedings of the IEEE Computer Vision and Pattern Recognition Conference, Las Vegas, 2016*, pp. 2940–2949, <https://ieeexplore.ieee.org/document/7780690>.

Catacora-Rios, M., G. B. King, A. E. Lovell, and F. M. Nunes, 2019, "Exploring experimental conditions to reduce uncertainties in the optical potential," *Phys. Rev. C* **100**, 064615.

Catacora-Rios, M., G. B. King, A. E. Lovell, and F. M. Nunes, 2020, "Which observable offers better constraints on the optical potential?," [arXiv:2012.06653](https://arxiv.org/abs/2012.06653).

Chaloner, K., and I. Verdinelli, 1995, "Bayesian experimental design: A review," *Stat. Sci.* **10**, 273–304.

Chen, S. Y., H. T. Ding, F. Y. Liu, G. Papp, and C. B. Yang, 2021, "Machine learning spectral functions in lattice QCD," [arXiv: 2110.13521](https://arxiv.org/abs/2110.13521).

Chen, T., and C. Guestrin, 2016, "XGBoost: A scalable tree boosting system," in *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD '16), San Francisco, 2016* (Association for Computing Machinery, New York), pp. 785–794, <https://www.kdd.org/kdd2016/papers/files/rfp0697-chenAemb.pdf>.

Chernodub, M. N., H. Erbin, V. A. Goy, and A. V. Molochkov, 2020, "Topological defects and confinement with machine learning: The case of monopoles in compact electrodynamics," *Phys. Rev. D* **102**, 054501.

Chien, Y.-T., 2019, "Probing heavy ion collisions using quark and gluon jet substructure with machine learning," *Nucl. Phys. A* **982**, 619–622.

Cisbani, E., *et al.*, 2020, "AI-optimized detector design for the future electron-ion collider: The dual-radiator RICH case," *J. Instrum.* **15**, P05009.

Clark, J. W., and H. Li, 2006, "Application of support vector machines to global prediction of nuclear properties," *Int. J. Mod. Phys. B* **20**, 5015–5029.

Cohen, O., 2018, "Active learning tutorial," <https://towardsdatascience.com/active-learning-tutorial-57c3398e34d>.

Collado, J., J. N. Howard, T. Faucett, T. Tong, P. Baldi, and D. Whiteson, 2021, "Learning to identify electrons," *Phys. Rev. D* **103**, 116028.

Connell, M. A., I. Billig, and D. R. Phillips, 2021, "Does Bayesian model averaging improve polynomial extrapolations? Two toy problems as tests," *J. Phys. G* **48**, 104001.

Costiris, N., E. Mavrommatis, K. Gernoth, and J. Clark, 2009, "Decoding beta-decay systematics: A global statistical model for  $\beta^-$  half-lives," *Phys. Rev. C* **80**, 044332.

Costiris, N. J., E. Mavrommatis, K. A. Gernoth, J. W. Clark, and H. Li, 2008, "Statistical global modeling of beta-decay halflives systematics using multilayer feedforward neural networks and support vector machines," in *Proceedings of the 17th Symposium of the Hellenic Nuclear Physics Society on Nuclear Physics, Ioannina, Greece, 2008*, [10.12681/hnps.2602](https://arxiv.org/abs/10.12681/hnps.2602).

Cranmer, K., U. Seljak, and K. Terao, 2022, "Machine learning," *Prog. Theor. Exp. Phys.* 083C01 [https://pdg.lbl.gov/2021/reviews/contents\\_sports.html](https://pdg.lbl.gov/2021/reviews/contents_sports.html).

Cristoforetti, M., F. Di Renzo, G. Eruzzi, A. Mukherjee, C. Schmidt, L. Scorzato, and C. Torrero, 2014, "An efficient method to compute the residual phase on a Lefschetz thimble," *Phys. Rev. D* **89**, 114505.

Cristoforetti, M., F. Di Renzo, and L. Scorzato (AuroraScience Collaboration), 2012, "New approach to the sign problem in quantum field theories: High density QCD on a Lefschetz thimble," *Phys. Rev. D* **86**, 074506.

Cybenko, G., 1989, "Approximation by superpositions of a sigmoidal function," *Math. Control Signals Syst.* **2**, 303.

Dalitz, C., Y. Ayyad, J. Wilberg, L. Aymans, D. Bazin, and W. Mittig, 2019, "Automatic trajectory recognition in active target time projection chambers data by means of hierarchical clustering," *Comput. Phys. Commun.* **235**, 159–168.

Deb, K., 2001, *Multi-objective Optimization Using Evolutionary Algorithms* (John Wiley & Sons, New York), <https://www.wiley.com/en-us/Multi+Objective+Optimization+using+Evolutionary+Algorithms-p-9780471873396>.

Deiana, A. M., *et al.*, 2021, "Applications and techniques for fast machine learning in science," [arXiv:2110.13041](https://arxiv.org/abs/2110.13041).

Del Debbio, L., T. Giani, J. Karpie, K. Orginos, A. Radyushkin, and S. Zafeiropoulos, 2021, "Neural-network analysis of parton distribution functions from Ioffe-time pseudodistributions," *J. High Energy Phys.* **02**, 138.

Del Debbio, L., J. Marsh Rossney, and M. Wilson, 2021, "Efficient modeling of trivializing maps for lattice  $\phi^4$  theory using normalizing flows: A first look at scalability," *Phys. Rev. D* **104**, 094507.

De Oliveira, L., B. Nachman, and M. Paganini, 2020, "Electromagnetic showers beyond shower shapes," *Nucl. Instrum. Methods Phys. Res., Sect. A* **951**, 162879.

Derkach, D., N. Kazeev, F. Ratnikov, A. Ustyuzhanin, and A. Volokhova, 2020, "Cherenkov detectors fast simulation using neural networks," *Nucl. Instrum. Methods Phys. Res., Sect. A* **952**, 161804.

De Sanctis, J., M. Masotti, M. Bruno, M. D'Agostino, E. Geraci, G. Vannini, and A. Bonasera, 2009, "Classification of the impact parameter in nucleus-nucleus collisions by a support vector machine method," *J. Phys. G* **36**, 015101.

Detmold, W., G. Kanwar, H. Lamm, M. L. Wagman, and N. C. Warrington, 2021, "Path integral contour deformations for observables in  $SU(N)$  gauge theory," *Phys. Rev. D* **103**, 094517.

Diefenthaler, M., A. Farhat, A. Verbytskyi, and Y. Xu, 2021, "Deeply learning deep inelastic scattering kinematics," [arXiv:2108.11638](https://arxiv.org/abs/2108.11638).

Dietrich, T., M. W. Coughlin, P. T. H. Pang, M. Bulla, J. Heinzel, L. Issa, I. Tews, and S. Antier, 2020, "Multimessenger constraints on the neutron-star equation of state and the Hubble constant," *Science* **370**, 1450–1453.

Di Renzo, F., and G. Eruzzi, 2015, "Thimble regularization at work: From toy models to chiral random matrix theories," *Phys. Rev. D* **92**, 085030.

Djärv, T., A. Ekström, C. Forssén, and H. T. Johansson, 2022, "Bayesian predictions for  $A = 6$  nuclei using eigenvector continuation emulators," *Phys. Rev. C* **105**, 014005.

Dolinski, M. J., A. W. P. Poon, and W. Rodejohann, 2019, "Neutrinoless double-beta decay: Status and prospects," *Annu. Rev. Nucl. Part. Sci.* **69**, 219–251.

Doucet, E., T. Brown, P. Chowdhury, C. Lister, C. Morse, P. Bender, and A. Rogers, 2020, "Machine learning  $n/\gamma$  discrimination in CLYC scintillators," *Nucl. Instrum. Methods Phys. Res., Sect. A* **954**, 161201.

Dramsch, J. S., M. Lüthje, and A. N. Christensen, 2021, “Complex-valued neural networks for machine learning on non-stationary physical data,” *Comput. Geosci.* **146**, 104643.

Drischler, C., R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, 2020, “How Well Do We Know the Neutron-Matter Equation of State at the Densities inside Neutron Stars? A Bayesian Approach with Correlated Uncertainties,” *Phys. Rev. Lett.* **125**, 202702.

Drischler, C., J. A. Melendez, R. J. Furnstahl, and D. R. Phillips, 2020, “Quantifying uncertainties and correlations in the nuclear-matter equation of state,” *Phys. Rev. C* **102**, 054315.

Drischler, C., M. Quinonez, P. Giuliani, A. Lovell, and F. Nunes, 2021, “Toward emulating nuclear reactions using eigenvector continuation,” *Phys. Lett. B* **823**, 136777.

Du, X., A. W. Steiner, and J. W. Holt, 2019, “Hot and dense homogeneous nucleonic matter constrained by observations, experiment, and theory,” *Phys. Rev. C* **99**, 025803.

Du, Y.-L., D. Pablos, and K. Tywoniuk, 2021a, “Jet tomography in heavy ion collisions with deep learning,” [arXiv:2106.11271](https://arxiv.org/abs/2106.11271).

Du, Y.-L., D. Pablos, and K. Tywoniuk, 2021b, “Deep learning jet modifications in heavy-ion collisions,” *J. High Energy Phys.* **03**, 206.

Du, Y.-L., K. Zhou, J. Steinheimer, L.-G. Pang, A. Motornenko, H.-S. Zong, X.-N. Wang, and H. Stöcker, 2020, “Identifying the nature of the QCD transition in relativistic collision of heavy nuclei with deep learning,” *Eur. Phys. J. C* **80**, 516.

Duris, J., *et al.*, 2020, “Bayesian Optimization of a Free-Electron Laser,” *Phys. Rev. Lett.* **124**, 124801.

Edelen, A., N. Neveu, M. Frey, Y. Huber, C. Mayes, and A. Adelmann, 2020, “Machine learning for orders of magnitude speedup in multiobjective optimization of particle accelerator systems,” *Phys. Rev. Accel. Beams* **23**, 044601.

Edelen, A., *et al.*, 2018, “Opportunities in machine learning for particle accelerators,” [arXiv:1811.03172](https://arxiv.org/abs/1811.03172).

Eklind, C., 2021, “Eigenvector continuation and strongly interacting cold atom systems,” master’s thesis (Chalmers University of Technology), <https://hdl.handle.net/20.500.12380/302455>.

Ekström, A., 2020, “Analyzing the nuclear interaction: Challenges and new ideas,” *Front. Phys.* **8**, 29.

Ekström, A., C. Forssén, C. Dimitrakakis, D. Dubhashi, H. T. Johansson, A. S. Muhammad, H. Salomonsson, and A. Schliep, 2019, “Bayesian optimization in *ab initio* nuclear physics,” *J. Phys. G* **46**, 095101.

Ekström, A., and G. Hagen, 2019, “Global Sensitivity Analysis of Bulk Properties of an Atomic Nucleus,” *Phys. Rev. Lett.* **123**, 252501.

Emma, C., A. Edelen, M. Hogan, B. O’Shea, G. White, and V. Yakimenko, 2018, “Machine learning-based longitudinal phase space prediction of particle accelerators,” *Phys. Rev. Accel. Beams* **21**, 112802.

Engel, G. P., and S. Schaefer, 2011, “Testing trivializing maps in the hybrid Monte Carlo algorithm,” *Comput. Phys. Commun.* **182**, 2107–2114.

Erdmann, J., 2020, “A tagger for strange jets based on tracking information using long short-term memory,” *J. Instrum.* **15**, P01021.

Essick, R., I. Tews, P. Landry, and A. Schwenk, 2021, “Astrophysical Constraints on the Symmetry Energy and the Neutron Skin of  $^{208}\text{Pb}$  with Minimal Modeling Assumptions,” *Phys. Rev. Lett.* **127**, 192701.

Ethier, J. J., N. Sato, and W. Melnitchouk, 2017, “First Simultaneous Extraction of Spin-Dependent Parton Distributions and Fragmentation Functions from a Global QCD Analysis,” *Phys. Rev. Lett.* **119**, 132001.

Fanelli, C., 2020, “Machine learning for imaging Cherenkov detectors,” *J. Instrum.* **15**, C02012.

Fanelli, C., and J. Pomponi, 2020, “DeepRICH: Learning deeply Cherenkov detectors,” *Mach. Learn. Sci. Technol.* **1**, 015010.

Favoni, M., A. Ipp, D. I. Müller, and D. Schuh, 2020, “Lattice gauge equivariant convolutional neural networks,” [arXiv:2012.12901](https://arxiv.org/abs/2012.12901).

Foreman, S., X.-Y. Jin, and J. C. Osborn, 2021, “Deep learning Hamiltonian Monte Carlo,” in *Proceedings of the 9th International Conference on Learning Representations, 2021* [[arXiv:2105.03418](https://arxiv.org/abs/2105.03418)].

Formaggio, J. A., A. L. C. de Gouvêa, and R. G. H. Robertson, 2021, “Direct measurements of neutrino mass,” *Phys. Rep.* **914**, 1–54.

Fox, J. M. R., C. W. Johnson, and R. N. Perez, 2020, “Uncertainty quantification of an empirical shell-model interaction using principal component analysis,” *Phys. Rev. C* **101**, 054308.

Fraker, S., 2018, “Deep learning for liquid-scintillator-based double-beta decay searches,” in *Proceedings of the XXVIII International Conference on Neutrino Physics and Astrophysics, Heidelberg, 2018*, <https://ui.adsabs.harvard.edu/abs/2018npa.confE.311F/abstract>.

Frame, D., R. He, I. Ipsen, D. Lee, D. Lee, and E. Rrapaj, 2018, “Eigenvector Continuation with Subspace Learning,” *Phys. Rev. Lett.* **121**, 032501.

Fujii, H., S. Kamata, and Y. Kikukawa, 2015, “Monte Carlo study of Lefschetz thimble structure in one-dimensional Thirring model at finite density,” *J. High Energy Phys.* **12**, 125; 09 (2016) 172.

Fujimoto, Y., K. Fukushima, and K. Murase, 2021, “Extensive studies of the neutron star equation of state from the deep learning inference with the observational data augmentation,” *J. High Energy Phys.* **03**, 273.

Fukushima, K., and Y. Tanizaki, 2015, “Hamilton dynamics for Lefschetz-thimble integration akin to the complex Langevin method,” *Prog. Theor. Exp. Phys.* **111A01**.

Fuller, A., Z. Fan, C. Day, and C. Barlow, 2020, “Digital twin: Enabling technologies, challenges and open research,” *IEEE Access* **8**, 108952–108971.

Furnstahl, R., A. Garcia, P. Millican, and X. Zhang, 2020, “Efficient emulators for scattering using eigenvector continuation,” *Phys. Lett. B* **809**, 135719.

Gando, A., *et al.* (KamLAND-Zen Collaboration), 2016, “Search for Majorana Neutrinos near the Inverted Mass Hierarchy Region with KamLAND-Zen,” *Phys. Rev. Lett.* **117**, 082503.

Gao, Y., J. Chen, T. Robertazzi, and K. A. Brown, 2019, “Reinforcement learning based schemes to manage client activities in large distributed control systems,” *Phys. Rev. Accel. Beams* **22**, 014601.

Gao, Z.-P., Y.-J. Wang, H.-L. Lü, Q.-F. Li, C.-W. Shen, and L. Liu, 2021, “Machine learning the nuclear mass,” *Nucl. Sci. Technol.* **32**, 109.

Gavalian, G., P. Thomadakis, A. Angelopoulos, V. Ziegler, and N. Chrisochoides, 2020, “Using artificial intelligence for particle track identification in CLAS12 detector,” [arXiv:2008.12860](https://arxiv.org/abs/2008.12860).

Gazula, S., J. Clark, and H. Bohr, 1992, “Learning and prediction of nuclear stability by neural networks,” *Nucl. Phys.* **A540**, 1–26.

Gelman, A., J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari, and D. B. Rubin, 2014, *Bayesian Data Analysis* (Chapman and Hall, London).

Gernoth, K., J. Clark, J. Prater, and H. Bohr, 1993, “Neural network models of nuclear systematics,” *Phys. Lett. B* **300**, 1–7.

Giuliani, P. G., and J. Piekarewicz, 2021, “From noise to information: The transfer function formalism for uncertainty quantification in reconstructing the nuclear density,” *Phys. Rev. C* **104**, 024301.

Gladen, R., V. Chirayath, A. Fairchild, M. Manry, A. Koymen, and A. Weiss, 2020, “Efficient machine learning approach for optimizing

the timing resolution of a high purity germanium detector," *Nucl. Instrum. Methods Phys. Res., Sect. A* **981**, 164505.

Gligorov, V. V., and M. Williams, 2013, "Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree," *J. Instrum.* **8**, P02013.

Gnech, A., C. Adams, N. Brawand, G. Carleo, A. Lovato, and N. Rocco, 2022, "Nuclei with up to  $A = 6$  nucleons with artificial neural network wave functions," *Few-Body Syst.* **63**, 7.

Gomez-Cadenas, J. J., *et al.* (NEXT Collaboration), 2014, "Present status and future perspectives of the NEXT experiment," *Adv. High Energy Phys.* 907067.

Goodfellow, I., Y. Bengio, and A. Courville, 2016, *Deep Learning* (MIT Press, Cambridge, MA).

Goriely, S., and R. Capote, 2014, "Uncertainties of mass extrapolations in Hartree-Fock-Bogoliubov mass models," *Phys. Rev. C* **89**, 054318.

Grewal, M. S., and A. P. Andrews, 2014, *Kalman Filtering: Theory and Practice with MATLAB* (Wiley-IEEE Press, Hoboken, NJ).

Grieves, M., and J. Vickers, 2017, "Digital twin: Mitigating unpredictable, undesirable emergent behavior in complex systems," in *Transdisciplinary Perspectives on Complex Systems: New Findings and Approaches*, edited by F.-J. Kahlen, S. Flumerfelt, and A. Alves (Springer International Publishing, Cham, Switzerland), pp. 85–113.

Gruszko, J., 2018, "NuDot: Double-beta decay with direction reconstruction in liquid scintillator," in *Proceedings of the XXVIII International Conference on Neutrino Physics and Astrophysics, Heidelberg, 2018*, <https://ui.adsabs.harvard.edu/abs/2018npa.confE.481G/abstract>.

Guest, D., K. Cranmer, and D. Whiteson, 2018, "Deep learning and its application to LHC physics," *Annu. Rev. Nucl. Part. Sci.* **68**, 161–181.

Güven, H., K. Bozkurt, E. Khan, and J. Margueron, 2020, "Multi-messenger and multiphysics Bayesian inference for the GW170817 binary neutron star merger," *Phys. Rev. C* **102**, 015805.

Hackett, D. C., C.-C. Hsieh, M. S. Albergo, D. Boyd, J.-W. Chen, K.-F. Chen, K. Cranmer, G. Kanwar, and P. E. Shanahan, 2021, "Flow-based sampling for multimodal distributions in lattice field theory," [arXiv:2107.00734](https://arxiv.org/abs/2107.00734).

Hamaker, A., *et al.*, 2021, "Precision mass measurement of light-weight self-conjugate nucleus  $^{80}\text{Zr}$ ," *Nat. Phys.* **17**, 1408–1412.

Han, M.-Z., J.-L. Jiang, S.-P. Tang, and Y.-Z. Fan, 2021, "Bayesian nonparametric inference of the neutron star equation of state via a neural network," *Astrophys. J.* **919**, 11.

Hanuka, A., C. Emma, T. Maxwell, A. S. Fisher, B. Jacobson, M. J. Hogan, and Z. Huang, 2021, "Accurate and confident prediction of electron beam longitudinal properties using spectral virtual diagnostics," *Sci. Rep.* **11**, 2945.

Hao, Y., Y. Li, M. Balcewicz, L. Neufcourt, and W. Cheng, 2019, "Reconstruction of storage ring's linear optics with Bayesian inference," [arXiv:1902.11157](https://arxiv.org/abs/1902.11157).

Hardin, J., and M. Williams, 2016, "FastDIRC: A fast Monte Carlo and reconstruction algorithm for DIRC detectors," *J. Instrum.* **11**, P10007.

Harland-Lang, L. A., A. D. Martin, P. Motylinski, and R. S. Thorne, 2015, "Parton distributions in the LHC era: MMHT 2014 PDFs," *Eur. Phys. J. C* **75**, 204.

Hashimoto, K., 2019, "AdS/CFT correspondence as a deep Boltzmann machine," *Phys. Rev. D* **99**, 106017.

Hashimoto, K., H.-Y. Hu, and Y.-Z. You, 2021, "Neural ordinary differential equation and holographic quantum chromodynamics," *Mach. Learn. Sci. Technol.* **2**, 035011.

Hashimoto, K., S. Sugishita, A. Tanaka, and A. Tomiya, 2018a, "Deep learning and holographic QCD," *Phys. Rev. D* **98**, 106014.

Hashimoto, K., S. Sugishita, A. Tanaka, and A. Tomiya, 2018b, "Deep learning and the AdS/CFT correspondence," *Phys. Rev. D* **98**, 046019.

Hastie, T., R. Tibshirani, and J. Friedman, 2009, *The Elements of Statistical Learning: Data Mining, Inference and Prediction* (Springer-Verlag, Berlin).

Hautier, G., C. C. Fischer, A. Jain, T. Mueller, and G. Ceder, 2010, "Finding nature's missing ternary oxide compounds using machine learning and density functional theory," *Chem. Mater.* **22**, 3762–3767.

Hayashida, S., 2019, "Application of the RNN in the fundamental physics with KamLAND experiment," in *Proceedings of the 8th Edition of the AI & Big Data Congress, Barcelona, 2019*, <https://www.imedpub.com/proceedings/application-of-the-rnn-in-the-fundamental-physics-with-kamland-experiment-7205.html>.

He, J., W.-B. He, Y.-G. Ma, and S. Zhang, 2021, "Machine-learning-based identification for initial clustering structure in relativistic heavy-ion collisions," [arXiv:2109.06277](https://arxiv.org/abs/2109.06277).

He, Y., L.-G. Pang, and X.-N. Wang, 2019, "Bayesian Extraction of Jet Energy Loss Distributions in Heavy-Ion Collisions," *Phys. Rev. Lett.* **122**, 252302.

Heim, F., P. Scholz, J. Mayer, M. Müller, and A. Zilges, 2020, "Constraining nuclear properties in  $^{94}\text{Mo}$  via a  $^{93}\text{Nb}(p, \gamma)^{94}\text{Mo}$  total cross section measurement," *Phys. Rev. C* **101**, 035807.

Higdon, D., J. D. McDonnell, N. Schunck, J. Sarich, and S. M. Wild, 2015, "A Bayesian approach for parameter estimation and prediction using a computationally intensive model," *J. Phys. G* **42**, 034009.

Hirlaender, S., and N. Bruchon, 2020, "Model-free and Bayesian ensembling model-based deep reinforcement learning for particle accelerator control demonstrated on the FERMI FEL," [arXiv:2012.09737](https://arxiv.org/abs/2012.09737).

Hjorth-Jensen, M., M. P. Lombardo, and U. van Kolck, 2017, Eds., *An Advanced Course in Computational Nuclear Physics: Bridging the Scales from Quarks to Neutron Stars*, Lecture Notes in Physics Vol. 936 (Springer Verlag, Berlin).

Holl, P., L. Hauertmann, B. Majorovits, O. Schulz, M. Schuster, and A. J. Zsigmond, 2019, "Deep learning based pulse shape discrimination for germanium detectors," *Eur. Phys. J. C* **79**, 450.

Horak, J., J. M. Pawłowski, J. Rodríguez-Quintero, J. Turnwald, J. M. Urban, N. Wink, and S. Zafeiropoulos, 2021, "Reconstructing QCD spectral functions with Gaussian processes," [arXiv:2107.13464](https://arxiv.org/abs/2107.13464).

Hornik, K., 1991, "Approximation capabilities of multilayer feedforward networks," *Neural Netw.* **4**, 251–257.

Hou, T.-J., *et al.*, 2021, "New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC," *Phys. Rev. D* **103**, 014013.

Houle, M. E., H.-P. Kriegel, P. Kröger, E. Schubert, and A. Zimek, 2010, "Can shared-neighbor distances defeat the curse of dimensionality?," in *Proceedings of the 22nd International Conference on Scientific and Statistical Database Management (SSDBM 2010), Heidelberg, 2010*, edited by M. Gertz and B. Ludäscher (Springer, New York), pp. 482–500.

Hu, B., *et al.*, 2021, "Ab initio predictions link the neutron skin of  $^{208}\text{Pb}$  to nuclear forces," [arXiv:2112.01125](https://arxiv.org/abs/2112.01125).

Huang, Y.-G., L.-G. Pang, X. Luo, and X.-N. Wang, 2021, "Probing criticality with deep learning in relativistic heavy-ion collisions," [arXiv:2107.11828](https://arxiv.org/abs/2107.11828).

Huber, D., O. V. Marchukov, H.-W. Hammer, and A. G. Volosniev, 2021, "Morphology of three-body quantum states from machine learning," *New J. Phys.* **23**, 065009.

Ismail, N., and A. Gezerlis, 2021, "Machine-learning approach to finite-size effects in systems with strongly interacting fermions," *Phys. Rev. C* **104**, 055802.

Jany, A., *et al.*, 2021, "Fabrication, characterization and analysis of a prototype high purity germanium detector for  $^{76}\text{Ge}$ -based neutrinoless double beta decay experiments," *Eur. Phys. J. C* **81**, 38.

Jhung, S., S. Hur, G. Cho, and I. Kwon, 2020, "A neural network approach for identification of gamma-ray spectrum obtained from silicon photomultipliers," *Nucl. Instrum. Methods Phys. Res., Sect. A* **954**, 161704.

Jiang, W. G., G. Hagen, and T. Papenbrock, 2019, "Extrapolation of nuclear structure observables with artificial neural networks," *Phys. Rev. C* **100**, 054326.

John, J. S., *et al.*, 2021, "Real-time artificial intelligence for accelerator control: A study at the Fermilab booster," [arXiv:2011.07371](https://arxiv.org/abs/2011.07371).

Jones, D. R., M. Schonlau, and W. J. Welch, 1998, "Efficient global optimization of expensive black-box functions," *J. Global Optim.* **13**, 455–492.

Kafkes, D., and M. Schram, 2021, "Developing robust digital twins and reinforcement learning for accelerator control systems at the Fermilab Booster," in *Proceedings of the 12th International Particle Accelerator Conference, Campinas, Brazil, 2021*, L. Liu, J. Byrd, R. Neuenschwander, R. Picoreti, and V. R. W. Schaa (JACoW, Geneva), <https://jacow.org/ipac2021/papers/tupab327.pdf>.

Kain, V., S. Hirlander, B. Goddard, F. M. Velotti, G. Z. Della Porta, N. Bruchon, and G. Valentino, 2020, "Sample-efficient reinforcement learning for CERN accelerator control," *Phys. Rev. Accel. Beams* **23**, 124801.

Kamuda, M., J. Stinnett, and C. J. Sullivan, 2017, "Automated isotope identification algorithm using artificial neural networks," *IEEE Trans. Nucl. Sci.* **64**, 1858–1864.

Kamuda, M., and C. J. Sullivan, 2019, "An automated isotope identification and quantification algorithm for isotope mixtures in low-resolution gamma-ray spectra," *Radiat. Phys. Chem.* **155**, 281–286.

Kamuda, M., J. Zhao, and K. Huff, 2020, "A comparison of machine learning methods for automated gamma-ray spectroscopy," *Nucl. Instrum. Methods Phys. Res., Sect. A* **954**, 161385.

Kanwar, G., M. S. Albergo, D. Boyd, K. Cranmer, D. C. Hackett, S. Racanière, D. J. Rezende, and P. E. Shanahan, 2020, "Equivariant Flow-Based Sampling for Lattice Gauge Theory," *Phys. Rev. Lett.* **125**, 121601.

Karpie, J., K. Orginos, A. Rothkopf, and S. Zafeiropoulos, 2019, "Reconstructing parton distribution functions from Ioffe time data: From Bayesian methods to neural networks," *J. High Energy Phys.* **04**, 057.

Kaspschak, B., and U.-G. Meißner, 2021, "Neural network perturbation theory and its application to the Born series," *Phys. Rev. Research* **3**, 023223.

Keeble, J., and A. Rios, 2020, "Machine learning the deuteron," *Phys. Lett. B* **809**, 135743.

Kejzlar, V., L. Neufcourt, W. Nazarewicz, and P.-G. Reinhard, 2020, "Statistical aspects of nuclear mass models," *J. Phys. G* **47**, 094001.

Kekic, M., *et al.* (NEXT Collaboration), 2021, "Demonstration of background rejection using deep convolutional neural networks in the NEXT experiment," *J. High Energy Phys.* **01**, 189.

Khalek, R. A., *et al.*, 2021, "Science requirements and detector concepts for the Electron-Ion Collider: EIC Yellow Report," [arXiv:2103.05419v3](https://arxiv.org/abs/2103.05419v3).

King, G. B., A. E. Lovell, L. Neufcourt, and F. M. Nunes, 2019, "Direct Comparison between Bayesian and Frequentist Uncertainty Quantification for Nuclear Reactions," *Phys. Rev. Lett.* **122**, 232502.

Koenker, R., and G. Bassett, 1978, "Regression quantiles," *Econometrica* **46**, 33.

Kościedziady, A., S. Bedawi, C. Ou, and F. Karray, 2018, "Non-stationary traffic flow prediction using deep learning," in *Proceedings of the 88th Vehicular Technology Conference, Chicago, 2018* (IEEE, New York), pp. 1–5.

Kohn, W., and L. J. Sham, 1965, "Self-consistent equations including exchange and correlation effects," *Phys. Rev.* **140**, A1133–A1138.

Komiske, P. T., E. M. Metodiev, B. Nachman, and M. D. Schwartz, 2017, "Pileup mitigation with machine learning (PUMML)," *J. High Energy Phys.* **12**, 051.

Komiske, P. T., E. M. Metodiev, and M. D. Schwartz, 2017, "Deep learning in color: Towards automated quark/gluon jet discrimination," *J. High Energy Phys.* **01**, 110.

König, S., A. Ekström, K. Hebeler, D. Lee, and A. Schwenk, 2020, "Eigenvector continuation as an efficient and accurate emulator for uncertainty quantification," *Phys. Lett. B* **810**, 135814.

Kramárik, L. (STAR Collaboration), 2020, "Reconstruction of  $D^0$  meson in  $d + \text{Au}$  collisions at  $\sqrt{s_{NN}} = 200$  GeV by the STAR experiment," *Phys. Scr.* **95**, 074010.

Kranjčević, M., B. Riemann, A. Adelmann, and A. Streun, 2021, "Multiobjective optimization of the dynamic aperture using surrogate models based on artificial neural networks," *Phys. Rev. Accel. Beams* **24**, 014601.

Kravaris, K., K. R. Quinlan, S. Quaglioni, K. A. Wendt, and P. Navrátil, 2020, "Quantifying uncertainties in neutron- $\alpha$  scattering with chiral nucleon-nucleon and three-nucleon forces," *Phys. Rev. C* **102**, 024616.

Kuchera, M. P., R. Ramanujan, J. Z. Taylor, R. R. Strauss, D. Bazin, J. Bradt, and R. Chen, 2019, "Machine learning methods for track classification in the AT-TPC," *Nucl. Instrum. Methods Phys. Res., Sect. A* **940**, 156–167.

Kunert, N., P. T. H. Pang, I. Tews, M. W. Coughlin, and T. Dietrich, 2022, "Quantifying modeling uncertainties when combining multiple gravitational-wave detections from binary neutron star sources," *Phys. Rev. D* **105**, L061301.

Kurle, R., B. Cseke, A. Klushyn, P. van der Smagt, and S. Günnemann, 2020, "Continual learning with Bayesian neural networks for non-stationary data," in *Proceedings of the International Conference on Learning Representations (ICLR 2020)*, 2020, <https://openreview.net/pdf?id=SJlsFpVtDB>.

Kuttan, M. O., J. Steinheimer, K. Zhou, A. Redelbach, and H. Stoecker, 2021, "Deep learning based impact parameter determination for the CBM experiment," *Particles* **4**, 47–52.

Lai, Y. S., J. Mulligan, M. Płoskoń, and F. Ringer, 2021, "The information content of jet quenching and machine learning assisted observable design," [arXiv:2111.14589](https://arxiv.org/abs/2111.14589).

Lai, Y. S., D. Neill, M. Płoskoń, and F. Ringer, 2020, "Explainable machine learning of the underlying physics of high-energy particle collisions," [arXiv:2012.06582](https://arxiv.org/abs/2012.06582).

Lasseri, R., D. Regnier, J. Ebran, and A. Penon, 2020, "Taming Nuclear Complexity with a Committee of Multilayer Neural Networks," *Phys. Rev. Lett.* **124**, 162502.

Lawrence, S., and Y. Yamauchi, 2021, "Normalizing flows and the real-time sign problem," *Phys. Rev. D* **103**, 114509.

Levy, D., M. D. Hoffman, and J. Sohl-Dickstein, 2018, "Generalizing Hamiltonian Monte Carlo with neural networks," in *Proceedings of the International Conference on Learning Representations, Vancouver, British Columbia, Canada, 2018*, <https://openreview.net/pdf?id=B1n8LexRZ>.

Li, A., A. Elagin, S. Fraker, C. Grant, and L. Winslow, 2019, “Suppression of cosmic muon spallation backgrounds in liquid scintillator detectors using convolutional neural networks,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **947**, 162604.

Li, F., Y. Wang, Z. Gao, P. Li, H. Lv, Q. Li, C. Y. Tsang, and M. B. Tsang, 2021, “Application of machine learning in the determination of impact parameter in the  $^{132}\text{Sn} + ^{124}\text{Sn}$  system,” [arXiv:2105.08912](https://arxiv.org/abs/2105.08912).

Li, F., Y. Wang, H. Lü, P. Li, Q. Li, and F. Liu, 2020, “Application of artificial intelligence in the determination of impact parameter in heavy-ion collisions at intermediate energies,” *J. Phys. G* **47**, 115104.

Li, S., M. Zacharias, J. Snuverink, J. Coello de Portugal, F. Perez-Cruz, D. Reggiani, and A. Adelmann, 2021, “A novel approach for classification and forecasting of time series in particle accelerators,” *Information* **12**, 121.

Li, S.-H., C.-X. Dong, L. Zhang, and L. Wang, 2020, “Neural Canonical Transformation with Symplectic Flows,” *Phys. Rev. X* **10**, 021020.

Li, Y., W. Cheng, L. H. Yu, and R. Rainer, 2018, “Genetic algorithm enhanced by machine learning in dynamic aperture optimization,” *Phys. Rev. Accel. Beams* **21**, 054601.

Li, Y., R. Rainer, and W. Cheng, 2019, “Bayesian approach for linear optics correction,” *Phys. Rev. Accel. Beams* **22**, 012804.

Liepe, J., S. Filippi, M. Komorowski, and M. P. H. Stumpf, 2013, “Maximizing the information content of experiments in systems biology,” *PLoS Comput. Biol.* **9**, e1002888.

Likhomanenko, T., P. Ilten, E. Khairullin, A. Rogozhnikov, A. Ustyuzhanin, and M. Williams, 2015, “LHCb topological trigger reoptimization,” *J. Phys. Conf. Ser.* **664**, 082025.

Lim, Y., and J. W. Holt, 2019, “Bayesian modeling of the nuclear equation of state for neutron star tidal deformabilities and GW170817,” *Eur. Phys. J. A* **55**, 209.

Lin, H.-W., W. Melnitchouk, A. Prokudin, N. Sato, and H. Shows, 2018, “First Monte Carlo Global Analysis of Nucleon Transversity with Lattice QCD Constraints,” *Phys. Rev. Lett.* **120**, 152502.

Liu, Y., C. Su, J. Liu, P. Danielewicz, C. Xu, and Z. Ren, 2021, “Improved naive Bayesian probability classifier in predictions of nuclear mass,” *Phys. Rev. C* **104**, 014315.

Lovell, A., A. Mohan, P. Talou, and M. Chertkov, 2019, “Constraining fission yields using machine learning,” *EPJ Web Conf.* **211**, 04006.

Lovell, A., F. M. Nunes, M. Catacora-Rios, and G. King, 2021, “Recent advances in the quantification of uncertainties in reaction theory,” *J. Phys. G* **48**, 014001.

Lovell, A. E., A. T. Mohan, T. M. Sprouse, and M. R. Mumpower, 2022, “Nuclear masses learned from a probabilistic neural network,” [arXiv:2201.00676](https://arxiv.org/abs/2201.00676).

Lovell, A. E., A. T. Mohan, and P. Talou, 2020, “Quantifying uncertainties on fission fragment mass yields with mixture density networks,” *J. Phys. G* **47**, 114001.

Lovell, A. E., and F. M. Nunes, 2018, “Constraining transfer cross sections using Bayes’ theorem,” *Phys. Rev. C* **97**, 064612.

Lundberg, S., and S. Lee, 2017, “A unified approach to interpreting model predictions,” in *Proceedings of the 31st International Conference on Neural Information Processing Systems (NIPS’17), Long Beach, CA, 2017*, edited by U. von Luxburg, I. Guyon, S. Bengio, H. Wallach, and R. Fergus (Curran Associates, Red Hook, NY), pp. 4768–4777, <https://papers.nips.cc/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-Abstract.html>.

Luo, D., G. Carleo, B. K. Clark, and J. Stokes, 2020, “Gauge equivariant neural networks for quantum lattice gauge theories,” [arXiv:2012.05232](https://arxiv.org/abs/2012.05232).

Luo, D., Z. Chen, K. Hu, Z. Zhao, V. M. Hur, and B. K. Clark, 2021, “Gauge invariant autoregressive neural networks for quantum lattice models,” [arXiv:2101.07243](https://arxiv.org/abs/2101.07243).

Lüscher, M., 2010, “Trivializing maps, the Wilson flow and the HMC algorithm,” *Commun. Math. Phys.* **293**, 899–919.

Lüscher, M., 2015, “Instantaneous stochastic perturbation theory,” *J. High Energy Phys.* **04**, 142.

Ma, C., D. Peng, H. Wei, Z. Niu, Y. Wang, and R. Wada, 2020, “Isotopic cross-sections in proton induced spallation reactions based on the Bayesian neural network method,” *Chin. Phys. C* **44**, 014104.

Ma, Y., C. Su, J. Liu, Z. Ren, C. Xu, and Y. Gao, 2020, “Predictions of nuclear charge radii and physical interpretations based on the naive Bayesian probability classifier,” *Phys. Rev. C* **101**, 014304.

Maevskiy, A., D. Derkach, N. Kazeev, A. Ustyuzhanin, and M. Artemev, L. Anderlini (LCHb Collaboration), 2020, “Fast data-driven simulation of cherenkov detectors using generative adversarial networks,” *J. Phys. Conf. Ser.* **1525**, 012097.

Magilligan, A., and B. A. Brown, 2020, “New isospin-breaking ‘USD’ Hamiltonians for the  $sd$  shell,” *Phys. Rev. C* **101**, 064312.

Maksimović, D., M. Nieslony, and M. Wurm, 2021, “CNNs for enhanced background discrimination in DSNB searches in large-scale water-Gd detectors,” [arXiv:2104.13426](https://arxiv.org/abs/2104.13426).

Mallick, N., S. Tripathy, A. N. Mishra, S. Deb, and R. Sahoo, 2021, “Estimation of impact parameter and transverse spherocity in heavy-ion collisions at the LHC energies using machine learning,” [arXiv:2103.01736](https://arxiv.org/abs/2103.01736).

Margraf, J. T., and K. Reuter, 2018, “Making the coupled cluster correlation energy machine learnable,” *J. Phys. Chem. A* **122**, 6343.

Margueron, J., H. Casali, and F. Gulminelli, 2018a, “Equation of state for dense nucleonic matter from metamodeling. I. Foundational aspects,” *Phys. Rev. C* **97**, 025805.

Margueron, J., H. Casali, and F. Gulminelli, 2018b, “Equation of state for dense nucleonic matter from metamodeling. II. Predictions for neutron star properties,” *Phys. Rev. C* **97**, 025806.

Mayer, J., K. Boretzky, C. Douma, E. Hoemann, and A. Zilges, 2021, “Classical and machine learning methods for event reconstruction in NeuLAND,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **1013**, 165666.

McDonnell, J. D., N. Schunck, D. Higdon, J. Sarich, S. M. Wild, and W. Nazarewicz, 2015, “Uncertainty Quantification for Nuclear Density Functional Theory and Information Content of New Measurements,” *Phys. Rev. Lett.* **114**, 122501.

McInnes, L., J. Healy, and S. Astels, 2017, “HDBSCAN: Hierarchical density based clustering,” *J. Open Source Software* **2**, 205.

McIntire, M., *et al.*, 2016, “Bayesian optimization of FEL performance at LCLS,” in *Proceedings of the 7th International Particle Accelerator Conference, Busan, Korea, 2016*, edited by K. S. Kim, I. S. Ko, K. R. Kim, and V. R. W. Schaa (JACoW, Geneva), <https://accelconf.web.cern.ch/ipac2016/doi/JACoW-IPAC2016-WEPOW055.html>.

Medhat, M., 2012, “Artificial intelligence methods applied for quantitative analysis of natural radioactive sources,” *Ann. Nucl. Energy* **45**, 73–79.

Medvidovic, M., J. Carrasquilla, L. E. Hayward, and B. Kulchytskyy, 2020, “Generative models for sampling of lattice field theories,” in *Proceedings of the 34th Conference on Neural Information Processing Systems, Vancouver, British Columbia, Canada, 2020* [[arXiv:2012.01442](https://arxiv.org/abs/2012.01442)].

Mehta, P., M. Bukov, C. Wang, A. G. Day, C. Richardson, C. K. Fisher, and D. J. Schwab, 2019, “A high-bias, low-variance

introduction to machine learning for physicists," *Phys. Rep.* **810**, 1–124.

Melendez, J., C. Drischler, A. Garcia, R. Furnstahl, and X. Zhang, 2021, "Fast & accurate emulation of two-body scattering observables without wave functions," *Phys. Lett. B* **821**, 136608.

Melendez, J. A., R. J. Furnstahl, H. W. Grießhammer, J. A. McGovern, D. R. Phillips, and M. T. Pratola, 2021, "Designing optimal experiments: An application to proton Compton scattering," *Eur. Phys. J. A* **57**, 81.

Melendez, J. A., R. J. Furnstahl, D. R. Phillips, M. T. Pratola, and S. Wesolowski, 2019, "Quantifying correlated truncation errors in effective field theory," *Phys. Rev. C* **100**, 044001.

Melendez, J. A., S. Wesolowski, and R. J. Furnstahl, 2017, "Bayesian truncation errors in chiral effective field theory: Nucleon-nucleon observables," *Phys. Rev. C* **96**, 024003.

Michaud, I., N. Kleedtke, J. Hutchinson, T. Smith, R. Little, T. Grove, and M. Rising, 2019, "Designing critical experiments using Gaussian process optimization," *Trans. Am. Nucl. Soc.* **121**, 1035–1038.

Miller, M. C., C. Chirenti, and F. K. Lamb, 2020, "Constraining the equation of state of high-density cold matter using nuclear and astronomical measurements," *Astrophys. J.* **888**, 12.

Miskovich, S. A., *et al.*, 2021, "Online Bayesian optimization for beam alignment in the SECAR recoil mass separator," in *Proceedings of the 35th Conference on Neural Information Processing Systems (NeurIPS 2021)*, Sydney, 2021, [10.48550/arXiv.2112.01557](https://arxiv.org/abs/2112.01557).

Moffat, E., W. Melnitchouk, T. Rogers, and N. Sato, 2021, "Simultaneous Monte Carlo analysis of parton densities and fragmentation functions," [arXiv:2101.04664](https://arxiv.org/abs/2101.04664).

Molchanov, O. M., K. D. Launey, A. Mercenne, G. H. Sargsyan, T. Dytrych, and J. P. Draayer, 2021, "Machine learning approach to pattern recognition in nuclear dynamics from the *ab initio* symmetry-adapted no-core shell model," [arXiv:2107.01498](https://arxiv.org/abs/2107.01498).

Monreal, B., and J. A. Formaggio, 2009, "Relativistic cyclotron radiation detection of tritium decay electrons as a new technique for measuring the neutrino mass," *Phys. Rev. D* **80**, 051301.

Moreno, J. R., G. Carleo, and A. Georges, 2020, "Deep Learning the Hohenberg-Kohn Maps of Density Functional Theory," *Phys. Rev. Lett.* **125**, 076402.

Morfouace, P., *et al.*, 2019, "Constraining the symmetry energy with heavy-ion collisions and Bayesian analyses," *Phys. Lett. B* **799**, 135045.

Mroczek, D., M. Hjorth-Jensen, J. Noronha-Hostler, P. Parotto, C. Ratti, and R. Vilalta, 2022, "Mapping out the thermodynamic stability of a QCD equation of state with a critical point using active learning," [arXiv:2203.13876](https://arxiv.org/abs/2203.13876).

Murphy, K. P., 2012, *Machine Learning: A Probabilistic Perspective* (MIT Press, Cambridge, MA).

Myren, S., and E. Lawrence, 2021, "A comparison of Gaussian processes and neural networks for computer model emulation and calibration," *Stat. Anal. Data Min.* **14**, 606–623.

Nagai, R., R. Akashi, and O. Sugino, 2020, "Completing density functional theory by machine learning hidden messages from molecules," *npj Comput. Mater.* **6**, 43.

Nakai, Y., D. Shih, and S. Thomas, 2020, "Strange jet tagging," [arXiv:2003.09517v1](https://arxiv.org/abs/2003.09517v1).

Nandi, A., C. Qu, P. L. Houston, R. Conte, and J. M. Bowman, 2021, " $\Delta$ -machine learning for potential energy surfaces: A PIP approach to bring a DFT-based PES to CCSD(T) level of theory," *J. Chem. Phys.* **154**, 051102.

Nätilä, J., M. C. Miller, A. W. Steiner, J. J. E. Kajava, V. F. Suleimanov, and J. Poutanen, 2017, "Neutron star mass and radius measurements from atmospheric model fits to x-ray burst cooling tail spectra," *Astron. Astrophys.* **608**, A31.

Navarro Pérez, R., J. E. Amaro, E. Ruiz Arriola, P. Maris, and J. P. Vary, 2015, "Statistical error propagation in *ab initio* no-core full configuration calculations of light nuclei," *Phys. Rev. C* **92**, 064003.

Navarro Pérez, R., N. Schunck, A. Dyhdalo, R. J. Furnstahl, and S. K. Bogner, 2018, "Microscopically based energy density functionals for nuclei using the density matrix expansion. II. Full optimization and validation," *Phys. Rev. C* **97**, 054304.

Negoita, G. A., *et al.*, 2019, "Deep learning: Extrapolation tool for *ab initio* nuclear theory," *Phys. Rev. C* **99**, 054308.

Neudecker, D., M. Grosskopf, M. Herman, W. Haeck, P. Grechanuk, S. Vander Wiel, M. Rising, A. Kahler, N. Sly, and P. Talou, 2020, "Enhancing nuclear data validation analysis by using machine learning," *Nucl. Data Sheets* **167**, 36–60.

Neudecker, D., *et al.*, 2021, "Informing nuclear physics via machine learning methods with differential and integral experiments," *Phys. Rev. C* **104**, 034611.

Neufcourt, L., Y. Cao, S. Giuliani, W. Nazarewicz, E. Olsen, and O. B. Tarasov, 2020a, "Beyond the proton drip line: Bayesian analysis of proton-emitting nuclei," *Phys. Rev. C* **101**, 014319.

Neufcourt, L., Y. Cao, S. A. Giuliani, W. Nazarewicz, E. Olsen, and O. B. Tarasov, 2020b, "Quantified limits of the nuclear landscape," *Phys. Rev. C* **101**, 044307.

Neufcourt, L., Y. Cao, W. Nazarewicz, E. Olsen, and F. Viens, 2019, "Neutron Drip Line in the Ca Region from Bayesian Model Averaging," *Phys. Rev. Lett.* **122**, 062502.

Neufcourt, L., Y. Cao, W. Nazarewicz, and F. Viens, 2018, "Bayesian approach to model-based extrapolation of nuclear observables," *Phys. Rev. C* **98**, 034318.

Neveu, N., L. Spentzouris, A. Adelmann, Y. Ineichen, A. Kolano, C. Metzger-Kraus, C. Bekas, A. Curioni, and P. Arbenz, 2019, "Parallel general purpose multiobjective optimization framework with application to electron beam dynamics," *Phys. Rev. Accel. Beams* **22**, 054602.

Newton, W. G., and G. Crocombe, 2021, "Nuclear symmetry energy from neutron skins and pure neutron matter in a Bayesian framework," *Phys. Rev. C* **103**, 064323.

Nicolí, K. A., C. J. Anders, L. Funcke, T. Hartung, K. Jansen, P. Kessel, S. Nakajima, and P. Stornati, 2021, "Estimation of Thermodynamic Observables in Lattice Field Theories with Deep Generative Models," *Phys. Rev. Lett.* **126**, 032001.

Niu, Z., and H. Liang, 2018, "Nuclear mass predictions based on Bayesian neural network approach with pairing and shell effects," *Phys. Lett. B* **778**, 48–53.

Niu, Z. M., J. Y. Fang, and Y. F. Niu, 2019, "Comparative study of radial basis function and Bayesian neural network approaches in nuclear mass predictions," *Phys. Rev. C* **100**, 054311.

Niu, Z. M., H. Z. Liang, B. H. Sun, W. H. Long, and Y. F. Niu, 2019, "Predictions of nuclear  $\beta$ -decay half-lives with machine learning and their impact on  $r$ -process nucleosynthesis," *Phys. Rev. C* **99**, 064307.

Nobre, G., D. Brown, S. Scoville, M. Fucci, S. Ruiz, R. Crawford, A. Coles, and M. Vorabbi, 2021, "Expansion of machine-learning method for classifying neutron resonances," Brookhaven National Laboratory Report No. BNL-222202-2021-INRE.

Nygren, D. R., B. J. P. Jones, N. López-March, Y. Mei, F. Psihas, and R. J., 2018, "Neutrinoless double beta decay with  $^{82}\text{SeF}_6$  and direct ion imaging," *J. Instrum.* **13**, P03015–P03015.

Odell, D., C. Brune, and D. Phillips, 2021, "How Bayesian methods can improve  $R$ -matrix analyses of data: The example of the  $d\ell$  reaction," [arXiv:2105.06541](https://arxiv.org/abs/2105.06541).

Ohnishi, A., Y. Mori, and K. Kashiwa, 2019, "Path optimization for the sign problem in field theories using neural network," *J. Phys. Soc. Jpn. Conf. Proc.* **26**, 024011.

Omana Kuttan, M., J. Steinheimer, K. Zhou, A. Redelbach, and H. Stoecker, 2020, "A fast centrality-meter for heavy-ion collisions at the CBM experiment," *Phys. Lett. B* **811**, 135872.

O'Shea, F. H., N. Bruchon, and G. Gaio, 2020, "Policy gradient methods for free-electron laser and terahertz source optimization and stabilization at the FERMI free-electron laser at Elettra," *Phys. Rev. Accel. Beams* **23**, 122802.

Page, B. S., X. Chu, and E. C. Aschenauer, 2020, "Experimental aspects of jet physics at a future EIC," *Phys. Rev. D* **101**, 072003.

Pang, L.-G., K. Zhou, N. Su, H. Petersen, H. Stöcker, and X.-N. Wang, 2018, "An equation-of-state-meter of quantum chromodynamics transition from deep learning," *Nat. Commun.* **9**, 210.

Pang, L.-G., K. Zhou, and X.-N. Wang, 2019, "Interpretable deep learning for nuclear deformation in heavy ion collisions," [arXiv: 1906.06429](https://arxiv.org/abs/1906.06429).

Paquet, J.-F. (JETSCAPE Collaboration), 2021, "Revisiting Bayesian constraints on the transport coefficients of QCD," *Nucl. Phys. A* **1005**, 121749.

Pastore, A., and M. Carnini, 2021, "Extrapolating from neural network models: A cautionary tale," *J. Phys. G* **48**, 084001.

Pastore, A., D. Neill, H. Powell, K. Medler, and C. Barton, 2020, "Impact of statistical uncertainties on the composition of the outer crust of a neutron star," *Phys. Rev. C* **101**, 035804.

Pawlowski, J. M., and J. M. Urban, 2020, "Reducing autocorrelation times in lattice simulations with generative adversarial networks," *Mach. Learn. Sci. Technol.* **1**, 045011.

Pederiva, G., and A. Shindler, 2020, "Machine learning for hadronic correlators from lattice QCD" (unpublished).

Pescia, G., J. Han, A. Lovato, J. Lu, and G. Carleo, 2021, "Neural-network quantum states for periodic systems in continuous space," [arXiv:2112.11957](https://arxiv.org/abs/2112.11957).

Phillips, D. R., *et al.*, 2021, "Get on the BAND wagon: A Bayesian framework for quantifying model uncertainties in nuclear dynamics," *J. Phys. G* **48**, 072001.

Pratt, S., E. Sangaline, P. Sorensen, and H. Wang, 2015, "Constraining the Equation of State of Super-Hadronic Matter from Heavy-Ion Collisions," *Phys. Rev. Lett.* **114**, 202301.

Pritychenko, B., E. Běták, M. A. Kellett, B. Singh, and J. Totans, 2011, "The Nuclear Science References (NSR) database and Web retrieval system," *Nucl. Instrum. Methods Phys. Res., Sect. A* **640**, 213.

Pumplin, J., D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky, and W. K. Tung, 2002, "New generation of parton distributions with uncertainties from global QCD analysis," *J. High Energy Phys.* **07**, 012.

Qian, Z., *et al.*, 2021, "Vertex and energy reconstruction in JUNO with machine learning methods," *Nucl. Instrum. Methods Phys. Res., Sect. A* **1010**, 165527.

Qiao, C. Y., J. C. Pei, Z. A. Wang, Y. Qiang, Y. J. Chen, N. C. Shu, and Z. G. Ge, 2021, "Bayesian evaluation of charge yields of fission fragments of  $^{239}\text{U}$ ," *Phys. Rev. C* **103**, 034621.

Raaijmakers, G., *et al.*, 2020, "Constraining the dense matter equation of state with joint analysis of NICER and LIGO/Virgo measurements," *Astrophys. J. Lett.* **893**, L21.

Radaideh, M. I., and T. Kozlowski, 2020, "Analyzing nuclear reactor simulation data and uncertainty with the group method of data handling," *Nucl. Eng. Technol.* **52**, 287–295.

Raghavan, K., P. Balaprakash, A. Lovato, N. Rocco, and S. M. Wild, 2021, "Machine-learning-based inversion of nuclear responses," *Phys. Rev. C* **103**, 035502.

Renner, J., *et al.*, 2017, "Background rejection in NEXT using deep neural networks," *J. Instrum.* **12**, T01004.

Rescic, M., R. Seviour, and W. Blokland, 2020, "Predicting particle accelerator failures using binary classifiers," *Nucl. Instrum. Methods Phys. Res., Sect. A* **955**, 163240.

Rodríguez, U., C. Z. Vargas, M. Gonçalves, S. B. Duarte, and F. Guzmán, 2019a, "Alpha half-lives calculation of superheavy nuclei with  $Q\alpha$ -value predictions based on the Bayesian neural network approach," *J. Phys. G* **46**, 115109.

Rodríguez, U. B., C. Z. Vargas, M. Gonçalves, S. B. Duarte, and F. Guzmán, 2019b, "Bayesian neural network improvements to nuclear mass formulae and predictions in the superheavy elements region," *Europhys. Lett.* **127**, 42001.

Romano, C., *et al.*, 2020, "Proceedings of the workshop for applied nuclear data: WANDA2020," Oak Ridge National Laboratory Technical Report No. ORNL/TM-2020/1617, <https://www.osti.gov/biblio/1649010-proceedings-workshop-applied-nuclear-data-wanda>.

Roussel, R., and A. Edelen, 2021, "Proximal biasing for Bayesian optimization and characterization of physical systems," in *Proceedings of the Workshop on Machine Learning and the Physical Sciences, 2021*, [https://ml4physicalsciences.github.io/2021/files/NeurIPS\\_ML4PS\\_2021\\_82.pdf](https://ml4physicalsciences.github.io/2021/files/NeurIPS_ML4PS_2021_82.pdf).

Rrapaj, E., and A. Roggero, 2021, "Exact representations of many-body interactions with restricted-Boltzmann-machine neural networks," *Phys. Rev. E* **103**, 013302.

Rumelhart, D. E., G. E. Hinton, and R. J. Williams, 1986, "Learning representations by back-propagating errors," *Nature (London)* **323**, 533–536.

Ryczko, K., D. A. Strubbe, and I. Tamblyn, 2019, "Deep learning and density-functional theory," *Phys. Rev. A* **100**, 022512.

Saito, H., 2018, "Method to solve quantum few-body problems with artificial neural networks," *J. Phys. Soc. Jpn.* **87**, 074002.

Sarkar, A., and D. Lee, 2021, "Self-learning emulators and eigenvector continuation," [arXiv:2107.13449](https://arxiv.org/abs/2107.13449).

Sato, N., C. Andres, J. J. Ethier, and W. Melnitchouk [Jefferson Lab Angular Momentum (JAM) Collaboration], 2020, "Strange quark suppression from a simultaneous Monte Carlo analysis of parton distributions and fragmentation functions," *Phys. Rev. D* **101**, 074020.

Saxena, G., P. K. Sharma, and P. Saxena, 2021, "Modified empirical formulas and machine learning for  $\alpha$ -decay systematics," *J. Phys. G* **48**, 055103.

Scamps, G., S. Goriely, E. Olsen, M. Bender, and W. Ryssens, 2020, "Grid Skyrme functional obtained using a committee of multilayer neural networks," [arXiv:2011.07904](https://arxiv.org/abs/2011.07904).

Schaefer, S., R. Sommer, and F. Virotta (ALPHA Collaboration), 2011, "Critical slowing down and error analysis in lattice QCD simulations," *Nucl. Phys.* **B845**, 93–119.

Scheinker, A., 2013, "Model independent beam tuning," in *Proceedings of the International Particle Accelerator Conference, Shanghai, 2013*, edited by Z. Dai, C. Petit-Jean-Genaz, V. R. W. Schaa, and C. Zhang (JACoW, Geneva), <https://accelconf.web.cern.ch/IPAC2013/papers/tupwa068.pdf>.

Scheinker, A., D. Bohler, S. Tomin, R. Kammering, I. Zagorodnov, H. Schlarb, M. Scholz, B. Beutner, and W. Decking, 2019, "Model-independent tuning for maximizing free electron laser pulse energy," *Phys. Rev. Accel. Beams* **22**, 082802.

Scheinker, A., A. Edelen, D. Bohler, C. Emma, and A. Lutman, 2018, "Demonstration of Model-Independent Control of the

Longitudinal Phase Space of Electron Beams in the Linac-Coherent Light Source with Femtosecond Resolution,” *Phys. Rev. Lett.* **121**, 044801.

Scheinker, A., C. Emma, A. L. Edelen, and S. Gessner, 2020, “Advanced Control Methods for Particle Accelerators (ACM4PA) 2019 Workshop Report,” [arXiv:2001.05461](https://arxiv.org/abs/2001.05461).

Scheinker, A., and S. Gessner, 2015, “Adaptive method for electron bunch profile prediction,” *Phys. Rev. ST Accel. Beams* **18**, 102801.

Scheinker, A., S. Gessner, C. Emma, and A. L. Edelen, 2020, “Adaptive model tuning studies for non-invasive diagnostics and feedback control of plasma wakefield acceleration at FACET-II,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **967**, 163902.

Scheinker, A., S. Hirlaender, F. M. Velotti, S. Gessner, G. Z. Della Porta, V. Kain, B. Goddard, and R. Ramjiawan, 2020, “Online multi-objective particle accelerator optimization of the AWAKE electron beam line for simultaneous emittance and orbit control,” *AIP Adv.* **10**, 055320.

Scheinker, A., and M. Krstić, 2017, *Model-Free Stabilization by Extremum Seeking* (Springer, New York).

Scheinker, A., and D. Scheinker, 2021, “Extremum seeking for optimal control problems with unknown time-varying systems and unknown objective functions,” *Int. J. Adapt. Control Signal Process.* **35**, 1143–1161.

Schleider, G. R., A. C. M. Padilha, C. M. Acosta, M. Costa, and A. Fazzio, 2019, “From DFT to machine learning: Recent approaches to materials science—A review,” *J. Phys. Mater.* **2**, 032001.

Schnabel, G., R. Capote, A. Koning, and D. Brown, 2021, “Nuclear data evaluation with Bayesian networks,” [arXiv:2110.10322](https://arxiv.org/abs/2110.10322).

Schnabel, G., *et al.*, 2020, “Conception and software implementation of a nuclear data evaluation pipeline,” [arXiv:2009.00521v1](https://arxiv.org/abs/2009.00521v1).

Schunck, N., 2019, Ed., *Energy Density Functional Methods for Atomic Nuclei* (IOP Publishing, Bristol, England), pp. 2053–2563.

Schunck, N., J. O’Neal, M. Grosskopf, E. Lawrence, and S. M. Wild, 2020, “Calibration of energy density functionals with deformed nuclei,” *J. Phys. G* **47**, 074001.

Schunck, N., K. R. Quinlan, and J. Bernstein, 2020, “A Bayesian analysis of nuclear deformation properties with Skyrme energy functionals,” *J. Phys. G* **47**, 104002.

Shaloo, R. J., *et al.*, 2020, “Automation and control of laser wake-field accelerators using Bayesian optimization,” *Nat. Commun.* **11**, 6355.

Shanahan, P. E., D. Trewartha, and W. Detmold, 2018, “Machine learning action parameters in lattice quantum chromodynamics,” *Phys. Rev. D* **97**, 094506.

Shelley, M., and A. Pastore, 2021, “A new mass model for nuclear astrophysics: Crossing 200 keV accuracy,” *Universe* **7**, 131.

Shen, Y. F., R. Pokharel, T. J. Nizolek, A. Kumar, and T. Lookman, 2019, “Convolutional neural network-based method for real-time orientation indexing of measured electron backscatter diffraction patterns,” *Acta Mater.* **170**, 118–131.

Shi, S., K. Zhou, J. Zhao, S. Mukherjee, and P. Zhuang, 2022, “Heavy quark potential in the quark-gluon plasma: Deep neural network meets lattice quantum chromodynamics,” *Phys. Rev. D* **105**, 014017.

Shimodaira, H., 2000, “Improving predictive inference under covariate shift by weighting the log-likelihood function,” *J. Stat. Plann. Infer.* **90**, 227–244.

Shintani, E., R. Arthur, T. Blum, T. Izubuchi, C. Jung, and C. Lehner, 2015, “Covariant approximation averaging,” *Phys. Rev. D* **91**, 114511.

Snoek, J., H. Larochelle, and R. P. Adams, 2012, “Practical Bayesian optimization of machine learning algorithms,” in *Proceedings of the 26th Annual Conference on Neural Information Processing Systems, Lake Tahoe, NV*, 2012, edited by C. J. C. Burges, L. Bottou, F. C. N. Pereira, P. L. Bartlett, and K. Q. Weinberger, pp. 2951–2959, <https://papers.nips.cc/paper/2012/hash/05311655a15b75fab86956663e1819cd-Abstract.html>.

Solli, R., D. Bazin, M. Hjorth-Jensen, M. P. Kuchera, and R. R. Strauss, 2021, “Unsupervised learning for identifying events in active target experiments,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **1010**, 165461.

Solopova, A., A. Carpenter, T. Powers, Y. Roblin, C. Tennant, L. Vidyaratne, and K. Iftekharuddin, 2019, “SRF cavity fault classification using machine learning at CEBAF,” in *Proceedings of the 10th International Particle Accelerator Conference (IPAC ’19), Melbourne, 2019*, edited by M. Boland, H. Tanaka, D. Button, R. Dowd, V. R. W. Schaa, and E. Tan (JACoW, Geneva), pp. 1167–1170, <https://accelconf.web.cern.ch/ipac2019/papers/tuxplm2.pdf>.

Sombillo, D. L. B., Y. Ikeda, T. Sato, and A. Hosaka, 2020, “Classifying the pole of an amplitude using a deep neural network,” *Phys. Rev. D* **102**, 016024.

Sombillo, D. L. B., Y. Ikeda, T. Sato, and A. Hosaka, 2021a, “Classifying near-threshold enhancement using deep neural network,” *Few-Body Syst.* **62**, 52.

Sombillo, D. L. B., Y. Ikeda, T. Sato, and A. Hosaka, 2021b, “Unveiling the pole structure of  $S$ -matrix using deep learning,” [arXiv:2104.14182](https://arxiv.org/abs/2104.14182).

Soto, C., 2019, “Visual detection with context for document layout analysis,” in *Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing and the 9th International Joint Conference on Natural Language Processing (EMNLP-IJCNLP), Hong Kong, 2019*, edited by S. Padó and R. Huang (Association for Computational Linguistics, Hong Kong), p. 3464.

Sprouse, T. M., R. Navarro Perez, R. Surman, M. R. Mumpower, G. C. McLaughlin, and N. Schunck, 2020, “Propagation of statistical uncertainties of Skyrme mass models to simulations of  $r$ -process nucleosynthesis,” *Phys. Rev. C* **101**, 055803.

Steinheimer, J., L. Pang, K. Zhou, V. Koch, J. Randrup, and H. Stoecker, 2019, “A machine learning study to identify spinodal clumping in high energy nuclear collisions,” *J. High Energy Phys.* **12**, 122.

Stevens, R., V. Taylor, J. Nichols, A. B. Maccabe, K. Yelick, and D. Brown, 2020, “AI for science,” Argonne National Laboratory, Technical Report No. ANL-20/17 158802, <https://www.osti.gov/biblio/1604756>.

Sugiyama, M., and M. Kawanabe, 2012, *Machine Learning in Non-stationary Environments: Introduction to Covariate Shift Adaptation* (MIT Press, Cambridge, MA).

Sutton, R. S., and A. G. Barto, 2018, *Reinforcement Learning: An Introduction* (MIT Press, Cambridge, MA).

Svensson, I., A. Ekström, and C. Forssén, 2022, “Bayesian parameter estimation in chiral effective field theory using the Hamiltonian Monte Carlo method,” *Phys. Rev. C* **105**, 014004.

Tennant, C., A. Carpenter, T. Powers, A. Shabalina Solopova, L. Vidyaratne, and K. Iftekharuddin, 2020, “Superconducting radio-frequency cavity fault classification using machine learning at Jefferson Laboratory,” *Phys. Rev. Accel. Beams* **23**, 114601.

Terzić, B., A. S. Hofler, C. J. Reeves, S. A. Khan, G. A. Krafft, J. Benesch, A. Freyberger, and D. Ranjan, 2014, “Simultaneous optimization of the cavity heat load and trip rates in linacs using a genetic algorithm,” *Phys. Rev. ST Accel. Beams* **17**, 101003.

Tomiya, A., and Y. Nagai, 2021, “Gauge covariant neural network for 4 dimensional non-Abelian gauge theory,” [arXiv:2103.11965](https://arxiv.org/abs/2103.11965).

Tong, L., R. He, and S. Yan, 2021, “Prediction of neutron-induced fission product yields by a straightforward  $k$ -nearest-neighbor algorithm,” *Phys. Rev. C* **104**, 064617.

Townsend, J., and K. D. Vogiatzis, 2020, “Transferable MP2-based machine learning for accurate coupled-cluster energies,” *J. Chem. Theory Comput.* **16**, 7453–7461.

Tshitoyan, V., J. Dagdelen, L. Weston, A. Dunn, Z. Rong, O. Kononova, K. A. Persson, G. Ceder, and A. Jain, 2019, “Unsupervised word embeddings capture latent knowledge from materials science literature,” *Nature (London)* **571**, 95–98.

Ulybyshev, M., C. Winterowd, and S. Zafeiropoulos, 2020, “Lefschetz thimbles decomposition for the Hubbard model on the hexagonal lattice,” *Phys. Rev. D* **101**, 014508.

Utama, R., W. C. Chen, and J. Piekarewicz, 2016, “Nuclear charge radii: Density functional theory meets Bayesian neural networks,” *J. Phys. G* **43**, 114002.

Utama, R., and J. Piekarewicz, 2017, “Refining mass formulas for astrophysical applications: A Bayesian neural network approach,” *Phys. Rev. C* **96**, 044308.

Utama, R., and J. Piekarewicz, 2018, “Validating neural-network refinements of nuclear mass models,” *Phys. Rev. C* **97**, 014306.

Utama, R., J. Piekarewicz, and H. B. Prosper, 2016, “Nuclear mass predictions for the crustal composition of neutron stars: A Bayesian neural network approach,” *Phys. Rev. C* **93**, 014311.

Vernon, I., M. Goldstein, and R. Bower, 2014, “Galaxy formation: Bayesian history matching for the observable Universe,” *Stat. Sci.* **29**, 81–90.

Vigneron, V., J. Morel, M. Lépy, and J. Martinez, 1996, “Statistical modelling of neural networks in  $\gamma$ -spectrometry,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **369**, 642–647.

Wang, D. Y., H. Bagri, C. Macdonald, S. Kiy, P. Jung, O. Shelbaya, T. Planche, W. Fedorko, R. Baartman, and O. Kester, 2021a, “Accelerator tuning with deep reinforcement learning,” in *Proceedings of the 35th Annual Conference on Neural Information Processing Systems, 2021*, [https://ml4physicalsciences.github.io/2021/files/NeurIPS\\_ML4PS\\_2021\\_125.pdf](https://ml4physicalsciences.github.io/2021/files/NeurIPS_ML4PS_2021_125.pdf).

Wang, R., Y.-G. Ma, R. Wada, L.-W. Chen, W.-B. He, H.-L. Liu, and K.-J. Sun, 2020, “Nuclear liquid-gas phase transition with machine learning,” *Phys. Rev. Research* **2**, 043202.

Wang, Z. A., J. C. Pei, Y. J. Chen, C. Y. Qiao, F. R. Xu, Z. G. Ge, and N. C. Shu, 2021, “Bayesian data fusion of imperfect fission yields for augmented evaluations,” [arXiv:2111.14102](https://arxiv.org/abs/2111.14102).

Wang, Z.-A., and J. Pei, 2021, “Optimizing multilayer Bayesian neural networks for evaluation of fission yields,” *Phys. Rev. C* **104**, 064608.

Wang, Z.-A., J. Pei, Y. Liu, and Y. Qiang, 2019, “Bayesian Evaluation of Incomplete Fission Yields,” *Phys. Rev. Lett.* **123**, 122501.

Wesolowski, S., R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, 2019, “Exploring Bayesian parameter estimation for chiral effective field theory using nucleon-nucleon phase shifts,” *J. Phys. G* **46**, 045102.

Wesolowski, S., I. Svensson, A. Ekström, C. Forssén, R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, 2021, “Rigorous constraints on three-nucleon forces in chiral effective field theory from fast and accurate calculations of few-body observables,” *Phys. Rev. C* **104**, 064001.

Wetzel, S. J., and M. Scherzer, 2017, “Machine learning of explicit order parameters: From the Ising model to SU(2) lattice gauge theory,” *Phys. Rev. B* **96**, 184410.

Whewell, B., M. Grosskopf, and D. Neudecker, 2020, “Evaluating  $^{239}\text{Pu}(\text{n}, \text{f})$  cross sections via machine learning using experimental data, covariances, and measurement features,” *Nucl. Instrum. Methods Phys. Res., Sect. A* **978**, 164305.

Wilkins, D. M., A. Grisafi, Y. Yang, K. U. Lao, R. A. DiStasio, and M. Ceriotti, 2019, “Accurate molecular polarizabilities with coupled cluster theory and machine learning,” *Proc. Natl. Acad. Sci. U.S.A.* **116**, 3401–3406.

Williams, C., and C. Rasmussen, 1996, “Gaussian processes for regression,” in *Advances in Neural Information Processing Systems*, Vol. 8, edited by D. Touretzky, M. C. Mozer, and M. Hasselmo (MIT Press, Cambridge, MA).

Wu, D., C. L. Bai, H. Sagawa, S. Nishimura, and H. Q. Zhang, 2021, “ $\beta$ -delayed one-neutron emission probabilities within a neural network model,” *Phys. Rev. C* **104**, 054303.

Wu, D., C. L. Bai, H. Sagawa, and H. Q. Zhang, 2020, “Calculation of nuclear charge radii with a trained feed-forward neural network,” *Phys. Rev. C* **102**, 054323.

Wu, X. H., L. H. Guo, and P. W. Zhao, 2021, “Nuclear masses in extended kernel ridge regression with odd-even effects,” *Phys. Lett. B* **819**, 136387.

Wu, X. H., and P. W. Zhao, 2020, “Predicting nuclear masses with the kernel ridge regression,” *Phys. Rev. C* **101**, 051301.

Wynen, J.-L., E. Berkowitz, S. Krieg, T. Luu, and J. Ostmeyer, 2021, “Machine learning to alleviate Hubbard-model sign problems,” *Phys. Rev. B* **103**, 125153.

Xie, W.-J., and B.-A. Li, 2020, “Bayesian inference of the symmetry energy of superdense neutron-rich matter from future radius measurements of massive neutron stars,” *Astron. J.* **899**, 4.

Xie, W.-J., and B.-A. Li, 2021, “Bayesian inference of the incompressibility, skewness and kurtosis of nuclear matter from empirical pressures in relativistic heavy-ion collisions,” *J. Phys. G* **48**, 025110.

Xu, J., W.-J. Xie, and B.-A. Li, 2020, “Bayesian inference of nuclear symmetry energy from measured and imagined neutron skin thickness in  $^{116,118,120,122,124,130,132}\text{Sn}$ ,  $^{208}\text{Pb}$ , and  $^{48}\text{Ca}$ ,” *Phys. Rev. C* **102**, 044316.

Xu, Y., J. E. Bernhard, S. A. Bass, M. Nahrgang, and S. Cao, 2018, “Data-driven analysis for the temperature and momentum dependence of the heavy-quark diffusion coefficient in relativistic heavy-ion collisions,” *Phys. Rev. C* **97**, 014907.

Yang, J., and J. Piekarewicz, 2020, “Covariant density functional theory in nuclear physics and astrophysics,” *Annu. Rev. Nucl. Part. Sci.* **70**, 21–41.

Yang, L., *et al.*, 2020, “Bayesian analysis on interactions of exotic nuclear systems,” *Phys. Lett. B* **807**, 135540.

Ye, X., H. Jian, and Z. Kai-En, 2008, “Applying Bayesian neural networks to identify pion, kaon and proton in BESII,” *Chin. Phys. C* **32**, 201–204.

Yoon, B., T. Bhattacharya, and R. Gupta, 2019, “Machine learning estimators for lattice QCD observables,” *Phys. Rev. D* **100**, 014504.

Yoshida, S., 2020, “Nonparametric Bayesian approach to extrapolation problems in configuration interaction methods,” *Phys. Rev. C* **102**, 024305.

Yoshida, S., and N. Shimizu, 2022, “Constructing approximate shell-model wavefunctions by eigenvector continuation,” *Prog. Theor. Exp. Phys.* **053D02**.

Yoshida, S., N. Shimizu, T. Togashi, and T. Otsuka, 2018, “Uncertainty quantification in the nuclear shell model,” *Phys. Rev. C* **98**, 061301.

You, Y.-Z., Z. Yang, and X.-L. Qi, 2018, “Machine learning spatial geometry from entanglement features,” *Phys. Rev. B* **97**, 045153.

Yousefnia, K. V., A. Kotibhaskar, R. Bhalerao, and J.-Y. Ollitrault, 2021, “Bayesian approach to long-range correlations and multiplicity fluctuations in nucleus-nucleus collisions,” [arXiv:2108.03471](https://arxiv.org/abs/2108.03471).

Yüksel, E., D. Soydaner, and H. Bahtiyar, 2021, “Nuclear binding energy predictions using neural networks: Application of the multilayer perceptron,” *Int. J. Mod. Phys. E* **30**, 2150017.

Zeiler, M. D., D. Krishnan, G. W. Taylor, and R. Fergus, 2010, “Deconvolutional networks,” in *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition, San Francisco, 2010* (IEEE, New York), pp. 2528–2535, <https://ieeexplore.ieee.org/document/5539957>.

Zhang, R., Z. Fan, R. Li, H.-W. Lin, and B. Yoon, 2020, “Machine-learning prediction for quasiparton distribution function matrix elements,” *Phys. Rev. D* **101**, 034516.

Zhang, X., and R. J. Furnstahl, 2021, “Fast emulation of quantum three-body scattering,” [arXiv:2110.04269](https://arxiv.org/abs/2110.04269).

Zhang, X., K. M. Nollett, and D. R. Phillips, 2015, “Halo effective field theory constrains the solar  ${}^7\text{Be} + p \rightarrow {}^8\text{B} + \gamma$  rate,” *Phys. Lett. B* **751**, 535–540.

Zhang, X., K. M. Nollett, and D. R. Phillips, 2020, “ $S$ -factor and scattering-parameter extractions from  ${}^3\text{He} + {}^4\text{He} \rightarrow {}^7\text{Be} + \gamma$ ,” *J. Phys. G* **47**, 054002.

Zhang, X., *et al.*, 2021, “Determining impact parameters of heavy-ion collisions at low-intermediate incident energies using deep learning with convolutional neural network,” [arXiv:2111.06597](https://arxiv.org/abs/2111.06597).

Zhang, Y., and Q. Yang, 2021, “A survey on multi-task learning,” [arXiv:1707.08114](https://arxiv.org/abs/1707.08114).

Zhang, Z., X.-B. Feng, and L.-W. Chen, 2021, “Bayesian inference on isospin splitting of nucleon effective mass from giant resonances in  ${}^{208}\text{Pb}$ ,” *Chin. Phys. C* **45**, 064104.

Zhao, Y.-S., L. Wang, K. Zhou, and X.-G. Huang, 2021, “Detecting chiral magnetic effect via deep learning,” [arXiv:2105.13761](https://arxiv.org/abs/2105.13761).

Zhu, J., Y. Chen, F. Brinker, W. Decking, S. Tomin, and H. Schlarb, 2021, “Deep learning-based autoencoder for data-driven modeling of an RF photoinjector,” [arXiv:2101.10437](https://arxiv.org/abs/2101.10437).