



Microbial metabolites in the marine carbon cycle

Mary Ann Moran¹✉, Elizabeth B. Kujawinski²✉, William F. Schroer¹, Shady A. Amin³, Nicholas R. Bates^{4,5}, Erin M. Bertrand⁶, Rogier Braakman⁷, C. Titus Brown⁸, Markus W. Covert⁹, Scott C. Doney¹⁰, Sonya T. Dyhrman^{11,12}, Arthur S. Edison¹³, A. Murat Eren^{14,15}, Naomi M. Levine¹⁶, Liang Li¹⁷, Avena C. Ross¹⁸, Mak A. Saito¹⁹, Alyson E. Santoro¹⁹, Daniel Segrè²⁰, Ashley Shade²¹, Matthew B. Sullivan²² and Assaf Vardi²³

One-quarter of photosynthesis-derived carbon on Earth rapidly cycles through a set of short-lived seawater metabolites that are generated from the activities of marine phytoplankton, bacteria, grazers and viruses. Here we discuss the sources of microbial metabolites in the surface ocean, their roles in ecology and biogeochemistry, and approaches that can be used to analyse them from chemistry, biology, modelling and data science. Although microbial-derived metabolites account for only a minor fraction of the total reservoir of marine dissolved organic carbon, their flux and fate underpins the central role of the ocean in sustaining life on Earth.

Organic molecules dissolved in seawater are cycled through the sunlit ocean by the billion marine microbes living in each litre of surface water¹ (Fig. 1). This process involves almost half of the annual net primary production (NPP) of the ocean; therefore, even minor changes in its output lead to important effects on a global scale. Much remains unknown about the chemicals passing through the labile dissolved organic carbon (DOC) pool, largely because they are highly bioavailable and they are almost simultaneously produced and consumed by diverse marine microbes. Specifically, the surface ocean labile DOC pool has a turnover of about three days², with steady-state concentrations in the range of nanomolar to picomolar^{3,4} for compounds abundant enough to be measured with existing methods and below this range for many others.

This low abundance of microbial metabolites in seawater presents a conundrum, as compounds supporting one of the most important fluxes in the carbon cycle of Earth are among the most challenging to study. In this Review, we summarize our current understanding of the sources and fate of marine microbial metabolites in the surface ocean and describe current and emerging analytical approaches for learning about their roles.

Marine sources of labile DOC

Most labile DOC of marine origin is produced by one of three sources: actively photosynthesizing phytoplankton, senescing and dead phytoplankton, or heterotrophic organisms.

Actively photosynthesizing marine phytoplankton generate exo-metabolites (metabolites released into surrounding seawater) that form a pool of carbon named ‘extracellular release’⁵ or ‘dissolved primary production’⁶ (Fig. 2). These released compounds are resources that are now lost from phytoplankton cells, and different hypotheses for the mechanisms underlying their release have been proposed⁷. For example, the million-fold concentration differential between the inside and outside of phytoplankton cell membranes⁸ could drive a molecular diffusion process by which healthy cells continually leak a fraction of their internal metabolites into seawater⁹. Leakage would be a passive process, and low-molecular-weight (<600 Da) and hydrophobic compounds (due to cell membrane composition) are more likely to leak out^{7,9,10}. In active mechanisms of extracellular release, labile DOC would be excreted rather than lost from living phytoplankton, which enables control over the rates and composition of the released metabolites. Physiological stresses imposed by photosynthesis are likely to drive active carbon export, and several mechanisms related to stress have been proposed. For example, when carbon dioxide (CO₂) is limiting, the principal carbon-fixation enzyme binds oxygen instead, which results in the production and release of toxic photorespiration products¹¹ such as glycolate¹². Conversely, when nutrients are limiting but inorganic carbon and irradiance are not, carbon fixation can outpace macromolecule synthesis¹³ and cause extracellular release of the excess fixed carbon in the form of polysaccharides and organic acids, which are referred to as ‘photosynthetic

¹Department of Marine Sciences, University of Georgia, Athens, GA, USA. ²Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, USA. ³Division of Science, New York University Abu Dhabi, Abu Dhabi, United Arab Emirates. ⁴Bermuda Institute of Ocean Sciences, St George's, Bermuda. ⁵School of Ocean and Earth Sciences, University of Southampton, Southampton, UK. ⁶Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada. ⁷Departments of Earth, Atmospheric and Planetary Sciences, and Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA. ⁸Population Health and Reproduction, School of Veterinary Medicine, University of California, Davis, Davis, CA, USA. ⁹Department of Bioengineering, Stanford University, Stanford, CA, USA. ¹⁰Department of Environmental Sciences, University of Virginia, Charlottesville, VA, USA. ¹¹Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA. ¹²Department of Earth and Environmental Science, Columbia University, Palisades, NY, USA. ¹³Departments of Biochemistry and Genetics, Complex Carbohydrate Research Center, University of Georgia, Athens, GA, USA. ¹⁴Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA, USA. ¹⁵Helmholtz-Institute for Functional Marine Biodiversity (HIFMB), University of Oldenburg, Oldenburg, Germany. ¹⁶Marine and Environmental Biology, Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA. ¹⁷Department of Chemistry, University of Alberta, Edmonton, Alberta, Canada. ¹⁸Department of Chemistry, Queen's University, Kingston, Ontario, Canada. ¹⁹Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, CA, USA. ²⁰Department of Biology and Bioinformatics Program, Boston University, Boston, MA, USA. ²¹Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing, MI, USA. ²²Departments of Microbiology and Civil, Environmental, and Geodetic Engineering, and Center of Microbiome Science, The Ohio State University, Columbus, OH, USA. ²³Department of Plant and Environmental Sciences, The Weizmann Institute of Science, Rehovot, Israel. ✉e-mail: mmoran@uga.edu; ekujawinski@whoi.edu




Fig. 1 | Microbial metabolites drive the carbon cycle in the surface ocean.

Metabolites in the labile DOC are synthesized by phytoplankton and other microbes and released into seawater through exudation, leakage, sloppy feeding and lysis. Within hours to days, they are consumed primarily by heterotrophic bacteria for growth and energy. The respiration of labile DOC adds CO_2 back to the surface ocean, which influences air-sea exchange. A small fraction of labile DOC is converted to non-labile forms and participates in long-term carbon storage in the deep ocean (microbial carbon pump). Fixed carbon that is not processed as labile DOC includes living microbes and metazoa, particulate detritus and refractory DOC, with a fraction transported to the deep ocean via the biological carbon pump (BCP).

overflow^{13–16}. Nutrient stress has also been proposed to drive organic carbon excretion as a by-product of cells increasing their ATP/ADP ratio (that is, reaching a higher intracellular energy state) to mitigate the free-energy costs of transport of low concentration nutrients¹⁷. These nutrient-limitation-driven mechanisms should favour the excretion of carbon-rich compounds¹⁷. At points in the diel cycle and regions in the photic zone where light shock can occur, extracellular release from phytoplankton may occur in response to a redox imbalance^{18,19}. Finally, phytoplankton actively release molecules that trigger behavioural or physiological changes in neighbouring microbes. These metabolites often have distinctive chemical structures and can function as defence compounds²⁰, pheromones²¹ or toxins^{22,23}. Given the different mechanisms and drivers of release, the chemical composition of phytoplankton exometabolites should differ from the composition of endometabolites (those present in the cell). This has indeed been observed for amino acids, carbohydrates and carboxylic acids^{24–26}. Regardless of the release mechanism, the tight coordination in daily activity patterns between phytoplankton and heterotrophic bacteria that track the diel irradiance cycle^{18,27,28} establishes a major role for extracellular products of photosynthetic activity in carbon transfer in the surface ocean (Fig. 3).

Phytoplankton senescence and death liberate endometabolites that form a second large source of labile molecules^{29,30} (Fig. 2). For example, sloppy feeding (escape of organic matter during grazing) or egestion (release of organic matter remaining undigested in guts and food vacuoles of predators)^{31–33} is carried out by protist and zooplankton predators and results in the release of dissolved organic




Fig. 2 | Sources of the labile microbial metabolites in the surface ocean.

Green arrows indicate substrate metabolites derived from primary production, whereas brown arrows indicate those from secondary production. The contributions of the three main sources can only be hypothesized at this time⁶⁰ (see the main text) given large reported ranges in the per cent of primary production released in dissolved form^{5,228}, biases from measurement artefacts (for example, cell disruption during filtration⁵ or differential radiolabelling of microbial intracellular pools¹⁹) and influences of environmental factors such as photosynthetic rate^{19,228}, irradiance levels, nutrient limitations, bloom stage and temperature³². Dotted arrows indicate facilitator and ecological signal metabolites that can indirectly contribute to carbon flux through influences on microbial activity.

matter from phytoplankton prey, which liberates 10–30% of prey carbon. Viral lysis of phytoplankton cells releases various metabolites^{34,35}, the composition of which depends on host reprogramming during infection^{36,37}. Recent evidence suggests that microparasitic fungi also change the intracellular chemistry of their host, similarly modifying the composition of released metabolites³⁸. Senescence or ‘autocatalytic cell death’ triggered by nutrient limitation or other stresses is also a source of metabolites^{10,29} and may be particularly important in declining blooms^{39,40}.

Finally, heterotrophs, including bacteria, archaea and protists, are the third main source of metabolic by-products generated by excretion or viral lysis^{32,41–43} (Fig. 2; heterotrophic activities of mixotrophic protists are counted here⁴⁴). Viral infection can modify bacterial endometabolomes⁴⁵ through reprogramming of host metabolism^{45,46}, which is an important component of labile DOC given estimates that one in three ocean bacteria are infected at any time³⁴. Organic molecules originating from heterotrophic protists and zooplankton can be rich in organic nitrogen⁴⁷ and are readily scavenged by bacteria³³. Heterotrophic marine bacteria and archaea release thousands of different molecules^{48–51} that might serve as bacterial substrates⁵². Liberation of labile compounds also occurs during bacterial solubilization of polymeric components of particulate detritus⁵³, such as proteins and polysaccharides, whose degradation products diffuse from the particle surface^{54–57}.

Fraction of carbon routed through microbial metabolites

Each year, heterotrophic marine bacteria are predicted to process ~20 PgC from the labile DOC pool (Fig. 1), as estimated by both geochemical methods based on DOC reactivity² and ecological approaches based on bacterial carbon demand⁵⁸ (that is, bacterial secondary production plus respiration). This routing of recently fixed carbon to bacteria through labile DOC is one of the largest and most rapid fluxes of organic carbon in the biosphere. Two early estimates of marine carbon processed through the microbe-metabolite network were 10–50% (ref. ⁵⁹) and 40% (ref. ⁵⁸) of ocean NPP, and a recent compilation of bacterial carbon demand values is consistent




Fig. 3 | Diel synchrony in microbial synthesis, release and utilization of metabolites. **a**, Periodicity in expression of a diatom gene (*SDH*) in the DHPS biosynthetic pathway (top) is correlated with DHPS concentrations in phytoplankton metabolomes (bottom) in surface waters of the North Pacific Subtropical Gyre. Expression data are transcripts per litre, normalized to the mean. Error bars represent the standard deviation ($n=3$). Data are from ref. ¹⁸. **b**, Periodicity in relative gene expression by *Trichodesmium* (filled symbols) is correlated with expression by its associated microbiome (open symbols) for nitrogen fixation and metabolism genes (top) and carbon fixation and respiration genes (bottom) in the North Atlantic. Data are from ref. ¹⁷³. **c**, Gene expression patterns for key metabolic processes by dominant members of a coastal microbial community in California, averaged over six diel light cycles. Top: photosynthetically active radiation. Bottom: time of peak relative expression for individual genes. Coloured symbols represent genes with statistically significant diel patterns. Gene expression by the dominant primary producer (*Ostreococcus*, green symbols) is shown separately for (top to bottom) photosynthesis, carbon fixation and translation. Gene expression by five dominant heterotrophic bacterial groups is shown for translation. Grey shading represents night. Data are from ref. ²⁷.

with these (52% of NPP⁶⁰). Partitioning of inputs among the three major sources estimated from published values for labile DOC release by microbial and zooplankton activities suggested that ~40% of labile DOC originates directly from phytoplankton as extracellular release from photosynthesizing cells, ~40% is released from phytoplankton death processes (senescence, sloppy feeding, viral and fungal lysis) and ~20% is released from heterotroph excretion and death processes⁶⁰. The magnitude of this last source is constrained by its derivation from secondary production, and therefore from organic carbon pools diminished by respiratory losses⁶¹.

Molecular composition of labile DOC

The molecules that make up the labile DOC pool of the ocean have been highly challenging to identify. Early studies proposed that amino acids, carbohydrates, osmolytes and small carboxylic acids (in particular the photorespiration product glycolate) are the main substrates for surface ocean bacteria^{1,14,62,63}. These compound classes were suggested by chemical analyses of plankton cells under the assumption that internal metabolite pools are proxies for released pools. Metabolites in intracellular pools are easier to measure than those in external pools because they typically have concentrations that are several orders of magnitude higher than in the surrounding seawater⁶³ (Table 1; micromolar to millimolar^{14,64–67} internal versus picomolar to nanomolar^{3,4,67} external). Furthermore, endometabolite samples can be concentrated by capturing cells on filters. However, paired analyses of endometabolites and exometabolites in phytoplankton cultures reveal that internal pools do not closely correspond to external pools⁶⁷, although viral lysis and sloppy feeding may be exceptions. This selective release of phytoplankton metabolites is not surprising but limits the accuracy of using endometabolomes to predict labile DOC composition. Nonetheless, endometabolite analyses of marine plankton have yielded candidates for the labile DOC inventory (Table 1). Examples include quaternary amines (for example, choline, dimethylglycine and trimethylamine-*N*-oxide), organic sulfur compounds (for example, gonyol, cysteate, dimethylsulfonioacetate and sulfolactate) and amino acid derivatives (for example, homarine, trigonelline and ornithine)^{18,65,67–76} (Table 1).

Labile DOC components can be coarsely categorized on the basis of their physiological and ecological roles in the microbe–metabolite network of the ocean. Here we define three broad categories that capture several of these roles: substrates, facilitators and ecological signals (Box 1).

Substrate metabolites are defined as the compounds that are actively assimilated by marine bacteria for carbon and energy, and transferred among microbes in quantities that sustain growth, reproduction, and the cycles of carbon and other elements. Amino acids^{77,78}, polyamines^{79,80}, carbohydrates^{81–83}, sulfonate and sulfonium compounds^{18,69}, carboxylic acids^{72,75,76,84}, and nucleosides^{70,72,75} all function as bacterial substrates in surface seawater (Table 1). Molecules in this category contribute most directly to flux by serving as the conduits or ‘currencies’ that move carbon among marine microbes.

Facilitator metabolites are defined as molecules that enable biochemical reactions and can be reused and exchanged among microbes (Box 1). Those that have been identified in seawater include the soluble B vitamins (B_1 , B_7 and B_{12})^{85–87}, iron siderophores^{88,89} and other biosynthesized metal-binding molecules^{90,91}, all of which have steady-state concentrations at the limit of bacterial uptake kinetics (<1 pM to tens of pM^{92–95}), although concentrations can fluctuate in time and space. Both substrate and facilitator molecules are also considered ‘public goods’ if they are energetically expensive to synthesize and released by only a subset of the microbial community⁹⁶; public goods can set the stage for metabolic dependencies within the microbial network of the ocean⁹⁷.

Ecological signal metabolites or ‘infochemicals’ are defined as compounds that orchestrate specialized microbial interactions under specific conditions^{98,99}. Sourced and shared by microbes, they serve as the medium for interactions by altering community physiological or behavioural features through chemicals passed among members (Box 1). Microbially produced hormones, such as indole acetic acid¹⁰⁰, and quorum-sensing molecules, such as acyl homoserine lactones¹⁰¹, are among the ecological signal compounds thus far discovered in ocean communities or cultivated marine microorganisms. Molecules that inhibit growth or cause mortality^{42,75,102} are also categorized as ecological signal compounds in this Review because of their role in modifying physiology (albeit negatively) of

other microbes. Both facilitator and ecological signal metabolites can indirectly contribute to labile DOC cycling by affecting the rates and routes of carbon flux without constituting a substantial mass fraction themselves.

Challenges to understanding roles of labile DOC

Although expertise in marine metabolites is growing, we cannot yet unravel the roles of labile DOC molecules in carbon flux and fate in the surface ocean¹⁰³. Roadblocks to progress can be boiled down to one (long) sentence: hidden among the hundreds of thousands of mostly unknown organic molecules embedded in a million-fold higher salt concentration are the currencies of a very large carbon flux through a very small carbon reservoir. Here, we pull apart this sentence and explain in more detail the specific challenges it captures.

Hidden in the highly complex marine DOC pool.... Each of the tremendous number of distinct microbial organisms that occupy surface seawater^{104,105} can release tens to thousands of different molecules^{48,49,51,67,106–109}. Genome-scale metabolic models based on flux balance analysis agree that microbial cells simultaneously maintain many hundreds of different endometabolites¹¹⁰ that can potentially be lost or exported as labile DOC. Some of these are predicted to be ‘costless’ metabolites, such as by-products of anabolic and catabolic pathways released without any fitness costs to the microbe¹¹¹. Untargeted mass spectrometry (MS) and nuclear magnetic resonance (NMR) analyses have indeed uncovered hundreds of thousands of distinct organic features in marine DOC^{112,113}, of which only 1–5% can be identified^{114–116}. Manually intensive, low-throughput identification pipelines are currently the primary approach used to convert these unidentified compounds into chemical insight. Of course, the biological reactivities of all the unknown compounds are also unknown; therefore, biogeochemically important molecules cannot be singled out of this multitude. In short, we cannot identify which of these compounds are important substrates, facilitator metabolites or ecological signal metabolites, or are otherwise uninvolved in the labile DOC cycle.

...embedded in a salty matrix.... Salt is a frustrating problem for the analysis of marine DOC. In MS analysis, non-volatile salts interfere with ion formation and degrade spectral quality¹¹⁷. In NMR analysis, salts reduce probe sensitivity¹¹⁸. Accordingly, the development of methods to remove organic compounds from their salty matrix has been an essential endeavour in chemical oceanography. Strategies that have been explored include water and salt removal via tangential flow filtration (TFF)¹¹⁹, reverse osmosis/electrodialysis¹²⁰ and metabolite capture on solid-phase extraction resins^{121,122}. Yet only 10–40% of marine DOC is captured with these methods, with a bias towards moderate- (>300 Da) and high- (>1,000 Da) molecular-weight compounds. In particular, small and polar metabolites, which are characteristics of many labile biomolecules, are lost during TFF or reverse osmosis/electrodialysis and are not well retained on solid-phase extraction resins such as the styrene-divinylbenzene polymer PPL¹²². Other saline fluids that are more easily analysed for metabolite content have higher concentrations of molecules in a lower concentration of salt, for example, millimolar metabolite concentrations (up to million-fold higher than seawater) in 1–10 ppt salt solutions (up to 30-fold lower than seawater) characteristic of blood and urine. For now, the state-of-the-art techniques used to isolate organic molecules from seawater remain strongly biased against the low-molecular-weight and polar exometabolites¹²², which form the majority of components of labile DOC.

...is a small metabolite reservoir with a very large flux. Labile DOC release is largely balanced over time by its consumption by heterotrophic bacteria, which maintain individual compounds in

the picomolar-to-low-nanomolar concentration range^{18,93,123–125} or lower. Specific uptake affinities for DOC components in oligotrophic marine bacteria are among the highest reported, for example, 150-fold higher than for well-studied model bacteria⁹³. Oligotrophic bacteria (those that can survive on low organic matter concentrations) are so efficient at depleting the labile DOC pool that calculations suggest that they must simultaneously scavenge for at least 34 different substrates to grow at one generation per day⁹². Copiotrophic marine bacteria (those that require higher organic matter concentrations) have lower substrate affinities but higher maximum uptake rates that allow them to rapidly draw down local substrate spikes on the order of 100 nM to 1 μ M (ref. ¹²⁶). It may be possible for a single bacterial taxon to switch between these life history strategies^{127,128}, but it is more probable that genetic makeup (for example, genome size, content and regulation) locks a heterotrophic bacterium into either a high- or low-affinity strategy^{129–131}. Regardless of how affinity is apportioned among cells, the communities of marine bacteria in aggregate are capable of both high affinity and fast uptake to maintain low concentrations of labile DOC components in seawater, and are considerably better at detecting DOC than our state-of-the-art chemical methods. Although targeted chemical methods have the sensitivity to directly measure some classes of metabolites in seawater, for example, amino acids¹³², sugars¹³³ and organic sulfur molecules⁶⁹, untargeted methods that aim to maximize the number and novelty of detected molecules are challenged by the low natural concentrations⁴.

The difficulty of characterizing the ‘small reservoir’ of labile DOC within the total DOC pool is intertwined with its ‘large flux’. Cycling rates of metabolites are not predictable by their concentration in seawater and may even be inversely related¹³⁴. For example, dimethylsulfoniopropionate (DMSP) has been estimated to support up to 10% of total bacterial carbon demand in the marine photic zone¹³⁵ yet has a steady-state concentration in the surface ocean of just ~3 nM (ref. ¹³⁶; Box 2). The most common method for estimating flux of recently fixed carbon into bacterial cells is to measure exudates released from ¹⁴C-labelled phytoplankton cells in the absence of bacterial uptake³², the so-called dissolved primary production. These estimates of carbon available for bacterial scavenging span a wide range (from 4% to 47% of NPP¹⁵) and provide no information on the specific compounds involved. Another approach captures the combined flux of labile DOC from all principal sources by determining the bacterial carbon demand as a percentage of ecosystem NPP. Results from this methodology suggest that ~40–60% of NPP is accessible to heterotrophs across a variety of marine environments, but with considerable uncertainties^{58,60}. Moreover, these methods cannot directly observe carbon flux on the time scales (seconds to minutes) and space scales (micrometres) of many microbial processes^{137–139}.

Analytical methods for marine microbial metabolites

Chemical methodological advances. Although there are now more than 18,500 metabolites catalogued in the Kyoto Encyclopedia of Genes and Genomes (KEGG) compound database¹⁴⁰, an untold number, including many with central roles in biogeochemical cycling, are still missing. A strategic approach is needed to focus on those currencies most relevant in driving the carbon cycle of the ocean. Molecular-level characterization of marine DOC has been a goal for decades in chemical oceanography and organic geochemistry, but has been thwarted by the salt and concentration challenges detailed above as well as by instrument limitations. Metabolite identification has emerged as a substantial bottleneck because the complex mixtures contain numerous isobaric and isomeric compounds that are poorly represented in current reference databases that are biased towards human metabolites and human-associated metabolites. Advances in both MS and NMR have now begun to crack this DOC black box (Fig. 4). The advent of electrospray

Table 1 | Marine microbial metabolites

Metabolite	Exometabolite in seawater	Exometabolite in culture	Endometabolite in marine plankton	Endometabolite in culture	Refs.
1-Methylhistidine		×			75
2'-Deoxyguanosine		×			4
2,3-Dihydroxybenzoate	×				108
3-Dehydroshikimate		×			225
3-Hydroxybutyrate				×	76
3-Mercaptopropionate	×	×			67,108
3-Phosphoglycerate		×			75
4-Acetamidobutanoate		×			75
4-Amino-5-aminomethyl-2-methylpyrimidine					3
4-Amino-5-hydroxymethyl-2-methylpyrimidine	×	×			4
4-Aminobenzoate	×	×			3,108
4-Aminobutanoate		×			75
4-Hydroxybenzaldehyde		×			75
4-Hydroxybenzoate	×	×	×		3,67,108,116
4-Hydroxyphenylacetate		×		×	75,76
4-Hydroxyphenylglycine		×			75
5'-Methylthioadenosine		×	×	×	3,116,225
5'-Uridine monophosphate	×	×			4,108
6-Phosphogluconate	×				108
(6R)-5,6,7,8-Tetrahydrobiopterin		×			4
7-Dehydrocholesterol			×		64
α-Aminoadipate		×			75
α-Ketoglutarate	×				3
α-Ribazole		×			225
β-1,3-Glucan				×	76
Acetate				×	76
Acetyltaurine	×				3
Aconitate			×		64
Adenine	×	×			3,4,64
Adenosine	×	×	×	×	3,4,64,67,70,75,108,116
Adenosine monophosphate			×		4,64,67,70,116
Aminobutyrate			×		64
Alanine	×	×	×	×	4,64,65,76
Arachidonate			×		64
Arginine	×	×	×	×	3,4,64,65,67,70,76,116
Asparagine	×	×	×	×	4,64,76
Aspartate	×	×	×	×	64,65,76
Azelaiate		×			75
Biotin	×		×		108,116
C ₁₆ -hydroxy-glycerophosphocholine		×			75
Caffeine	×		×		3,64,116
Carnosine		×			75
Cyclic guanosine monophosphate			×		64
Chitobiose	×	×	×	×	3,4,64,67,70
Chitotriose	×	×			3,4,70,108
Choline	×		×	×	3,64,65,76,108
Ciliatine	×	×			3,4

Continued

Table 1 | Marine microbial metabolites

Metabolite	Exometabolite in seawater	Exometabolite in culture	Endometabolite in marine plankton	Endometabolite in culture	Refs.
Citrate	×		×		3,64
Citrulline	×	×		×	3,67,75
Creatine			×		64
Cyanocobalamin	×	×			67,108
Cystathionine			×		64
Cysteinolate			×		18
Cysteate	×	×	×	×	4,18,64
Cysteine	×	×			4
Cytidine	×		×		4,64
Cytosine			×		64
Desthiobiotin	×				3,108
Dihydroxyacetone phosphate	×				3
DHPS	×	×	×	×	3,4,18,64,65,69,76
Dimethylglycine			×	×	64,76
Dimethylsulfonioacetate				×	71
DMSP	×	×	×	×	64,65,67,69,73,76,108,116,225,226
Dimethylsulfoxonium propionate	×			×	227
Ectoine	×	×			3,4
Ethanolamine				×	76
Folate	×		×		108,116
Fosfomycin	×				3
Gamma-aminobutyrate	×	×			4
Gluconate			×		64
Glucose				×	76
Glucose-6-phosphate	×			×	3,67,108
Glucosamine-6-phosphate	×	×			4,108
Glucosylglycerol			×	×	64,65
Glutamate	×	×	×	×	3,4,64,65,67,76
Glutamine	×	×	×	×	4,64,65,76,225
Glutathione	×		×	×	3,64,67
Glycerol-3-phosphate	×		×	×	3,64,76
Glycerophosphocholine				×	76
Glycine	×	×		×	4,76
Glycine betaine			×	×	64,65,67,76,116
Glyphosate	×				108
Gonyol			×	×	64,71
Guanine	×		×		3,64,116
Guanosine	×	×	×	×	3,64,70,76,108
Guanosine monophosphate			×		64
Histidine	×	×	×		4,64
Homarine			×	×	64,76
Homoserine	×	×			4
Homoserine betaine		×			4
Hydroxocobalamin			×		64
Indole-3-acetate	×	×	×		64,67,108,116
Indole-3-acetamide			×		64

Continued

Table 1 | Marine microbial metabolites

Metabolite	Exometabolite in seawater	Exometabolite in culture	Endometabolite in marine plankton	Endometabolite in culture	Refs.
Inosine 5'-monophosphate	×				108
Inosine	×	×	×		3,67,70,108,116
Isethionate	×	×	×	×	18,64,65
Isoleucine	×	×	×	×	64,65,70,76,108,116
Kynurenine	×	×	×		3,4,64,108
Leucine	×	×	×	×	3,4,64,70,75,76,108,116
Lysine	×	×	×	×	4,64,65,76
Malate	×	×			3,4
Methionine			×	×	64,116,225
Methylglutarate		×			75
Methyl indole-3-carboxylate			×		64
Muramate	×	×			4
<i>N</i> -(3-oxotetradecanoyl)-L-homoserine lactone		×		×	116
<i>N</i> -acetyl-galactosamine	×	×		×	67,75
<i>N</i> -acetyl-glutamate	×	×			67,108
<i>N</i> -acetylmuramate	×	×			3,4,108
<i>N</i> -acetyltaurine		×			67
<i>N</i> -tetradecanoylaspartate		×			75
Niacin			×		64
Nicotinamide adenine dinucleotide-hydrogen	×				3,108
Nicotinamide adenine dinucleotide-phosphate	×				108
Norvaline		×			75
Oleate		×			75
Ornithine	×	×		×	4,65
Pantothenate	×	×	×		3,4,64,108,116
Phenylacetate		×			75
Phenylalanine	×	×	×		3,4,64,67,108,116
Phosphoglycerate			×		64
Phosphorylcholine				×	76
Proline	×	×	×	×	3,4,64,65,67,70,76,116
Propionate		×		×	225
Putrescine	×	×			4
Pyridoxine	×		×		3,64,116
Pyridoxal			×		64
Pyridoxal phosphate			×		64
Quinolinecarboxylate		×			75
Riboflavin	×	×	×	×	3,64,108,116,225
Ribose 5-phosphate			×		64
Rosmarinate		×			75
<i>S</i> -5'-adenosyl-L-homocysteine	×	×	×		3,4,64
<i>S</i> -adenosyl methionine			×		64
Sarcosine	×	×	×		4,64
Serine		×	×		4,64
Shikimate		×			225
Spermidine	×	×		×	4,67
Sphingamine		×			75
Stearate		×			75

Continued

Table 1 | Marine microbial metabolites

Metabolite	Exometabolite in seawater	Exometabolite in culture	Endometabolite in marine plankton	Endometabolite in culture	Refs.
Suberate		×			75
Succinate	×	×			3,67
Sucrose			×		64
Sulfolactate			×	×	18,64
Syringate	×				3
Taurine	×		×	×	4,18,64,65
Taurocholate	×	×			3,4,67,108
Thiamin				×	67
Thiamin monophosphate				×	225
Threonine	×	×	×		3,64,75
Thymidine	×	×	×		3,64,67,70
Thymine			×		64
Trehalose			×		64
Trigonelline			×		64
Trimethylamine- <i>N</i> -oxide				×	76
Tryptamine	×				3,108
Tryptophan	×	×	×		3,64,67,108,116
Tyrosine	×		×		3,64
Uracil				×	67
Uridine	×	×	×	×	3,64,76
Uridine 5-monophosphate	×				108
Uridine diphosphate-glucosamine			×		64
Uridine diphosphate-glucose			×		64
Valine	×	×	×	×	3,4,64,70,76
Vanillate			×		64
Xanthine	×		×		3,108,116
Xanthosine	×		×		3,4,64,108

Shown are exometabolites identified in seawater and culture medium, and endometabolites identified in cultured cells and plankton communities.

ionization¹⁴¹ enabled the transfer of polar molecules directly into mass spectrometers. Mass resolution and accuracy have dramatically improved with Fourier-transform-based analysers such as Fourier transform ion cyclotron resonance cells¹⁴² and Orbitrap detectors¹⁴³, which now routinely enable mass accuracy to levels below parts per million for thousands of molecules within complex mixtures. Chromatographic columns such as hydrophilic interaction chromatography and new mixed-mode resins can deliver specific fractions of the labile DOC pool for characterization¹⁴⁴. Moreover, picomoles of individual compounds are sufficient to trigger fragmentation spectra for identification. Although typically less sensitive than MS for molecular detection, NMR provides structural information that better enables identification. Higher-field NMR magnets¹⁴⁵ and advanced small-diameter probes provide better analyte detection with lower salt sensitivity¹⁴⁶, and nanomoles of individual molecules are currently required for structural identification. Two-dimensional NMR approaches¹⁴⁷ have been developed to obtain complete covalent geometry of molecules, sometimes with stereochemistry¹⁴⁸. Further advances in MS and NMR technology offer promise for improved DOC characterization in the near future, including better data deconvolution strategies¹⁴⁹ and improved methods for integrating data for unknown compound identification¹⁵⁰. The compounds that have been successfully identi-

fied in marine metabolomes (Table 1) represent only a small fraction of total metabolite diversity.

Although MS and NMR offer the best potential for identifying biologically labile molecules in marine DOC, the salt problem has yet to be solved. Derivatization protocols show promise for this challenge, in particular those that target functional groups common in biologically produced compounds (for example, alcohol and amine groups⁴ and carbonyl moieties¹⁵¹). Recent application of derivatization methods are enabling detection in seawater at nanomolar (NMR and gas chromatography/MS)¹⁵² to picomolar (liquid chromatography/MS)⁴ concentrations. Identification of polar metabolites by direct injection of seawater into mass spectrometers has also been demonstrated recently for marine culture media, with limits of detection averaging 600 nM over a range from 10 nM to 3.2 µM (ref. ¹⁵³).

Chemical methods are now better able to measure bacterial uptake of labile metabolites. Isotopically labelled compounds can be tracked into individual cells using nano-secondary ion MS^{154,155} and chemical-tagging tools^{156–159}. As sampling volume requirements decrease and instrument sensitivities increase, single-cell measurements of internal and external metabolites will draw tighter associations between microbes and molecules^{160,161}. High-resolution magic-angle spinning NMR probes provide access to real-time




Box 1 | Ecological classes of microbial exometabolites

Metabolites are small molecules that are direct products of metabolism. Chemically, they hail from a wide variety of structural classes and span a range of solubilities, molecular weights and functional groups²²⁹. Metabolites can also be classified on the basis of their ecological role in microbial communities; in the marine microbiome they typically have one of three main roles, which are outlined below.

Substrate metabolites sustain biomass production and element cycling in microbial communities. In the surface ocean, molecules in this category include carboxylic acids^{230,231}, glycerols and fatty acids^{230,232}, nitrogen-containing compounds (such as polyamines)^{79,169,230}, C₁-compounds^{169,230,232}, carbohydrates (such as glucose)^{7,233}, and sulfonates and sulfonium compounds (such as DHPS)^{18,69,234}. Substrate metabolites are likely to be synthesized in core biochemical pathways during microbial growth and be conserved across diverse taxonomic groups.

Facilitator metabolites enable or enhance chemical reactions and include molecules such as vitamins (such as B₇) and siderophores (such as catecholate siderophores)^{87,235}. Enzymes are considered to be macromolecules rather than metabolites and are not included here. Facilitator metabolites are also likely to be synthesized in core biochemical pathways.

Ecological signal metabolites alter the phenotype of neighbouring microbes and are typically secondary metabolites produced to support non-growth activities. In the marine microbiome, ecological signal metabolites include chemical cues or infochemicals (such as homoserine lactones involved in quorum sensing), microbial pheromones, and antimicrobials and algicides (such as tropodithetic acid)^{236–240}. Marine bacteria can devote considerable genomic resources to the synthesis of ecological signal metabolites^{241,242}.


Substrates sustain biomass production and element cycling**Facilitators enable chemical reactions****Signals alter microbial phenotypes**

metabolism *in vivo*, even in the presence of high salt concentrations¹⁶². Flux measurements at the required temporal and spatial scales are on the horizon, with advances in tracking isotope incorporation through intracellular metabolic pathways^{163,164}, cellular uptake of individual metabolites¹⁶² and fluid-flow devices coupled to high-resolution imaging¹³⁷. These multiple approaches are chipping away at the barriers to progress.

Biological methodological advances. Labile metabolites are defined as those susceptible to microbial transformation, which makes evidence of biological processing an effective operational definition for labile DOC. Biology-based screening approaches fall

Box 2 | Hunting marine metabolites

Determining the role of an ocean metabolite is a complex and multidisciplinary process, as revealed by the decades of research dedicated to learning the biogeochemistry of just one metabolite: DMSP. Early research established DMSP as a major phytoplankton osmolyte²²⁶, an important substrate for marine bacteria²⁴³ and the precursor of dimethylsulfide (DMS), which is the dominant volatile in the ocean–atmosphere sulfur flux^{244,245}. Yet the biochemical mechanisms of DMSP synthesis and degradation remained unknown until genomic data enabled gene discovery beginning in 2006^{246–249}. Since then, the configuration of the microbial–DMSP network, initially considered a simple flux from phytoplankton to bacteria via the labile DOC pool (light arrows), has been revealed as a highly complex web of synthesis and utilization (dark arrows). New network edges have been discovered, such as the findings that bacteria also synthesize DMSP²⁴⁶ and that phytoplankton also assimilate it from the environment²⁵⁰. New nodes have been discovered, such as chemical relatives of DMSP that affect its fate^{71,227} and roles for viruses in release and transformation^{251,252}. New functions for DMSP have been discovered, such as ecological signals for bacterial chemotaxis²⁵³ and pathogenesis²⁵⁴. DMSP may be the highest-flux single metabolite of the surface ocean carbon cycle^{73,135}, yet biological and chemical studies over decades were required to unravel the intricacies of its dynamics. Factors that regulate the fate of DMSP, including what controls its transformation to climate-active DMS, are yet to be resolved^{163,255,256}. DMSOP, dimethylsulfoxonium propionate.



into two categories: those that use biological signals alone to generate hypotheses regarding important compounds and those that couple biological signals with chemical analysis. In the former category, gene, transcript and protein inventories suggest which molecules are produced and consumed. These biosensor strategies do not require disruption of steady-state interactions and instead survey intact natural communities (using metagenomics, metatranscriptomics and metaproteomics) or model organism systems (using genomics, transcriptomics and proteomics). This approach has offered insight into metabolite flux in phytoplankton blooms^{165,166}, oligotrophic seawater^{167,168}, model communities^{66,74} and ocean-wide surveys^{169,170}. It has been used to construct co-occurrence networks of microbes and metabolites^{171,172}, and has provided insight into the choreographed daily cycles of metabolite-driven phytoplankton–bacteria interactions in the surface ocean^{18,27,167,173} (Fig. 3). The approach can be expanded to regional, global and full-ocean-depth scales to illuminate large-scale patterns in labile DOC transformations. Biological screening methods that rely solely on microbial




Fig. 4 | The marine DOC spectrum. Chemical analysis strategies for marine DOC are targeted to molecules of different size and hydrophobicity, and typically begin with extraction and concentration from seawater. Solid-phase extraction (SPE) includes common resin types such as C18, C8, hydrophilic-lipophilic balance and priority pollutant. ESI-LC/MS, electrospray ionization-liquid chromatography/MS; FTMS, Fourier transformMS (includes Orbitrap mass spectrometers and Fourier transform ion cyclotron resonance mass spectrometers); GC/MS, gas chromatography/MS.

response, however, are ultimately constrained by inadequate and slowly advancing gene annotation, an area for which investments in technology and new approaches are sorely needed.

The second category of biological screening is coupled to chemical analysis. This typically requires steady-state disruption, altering either the accumulation or utilization of DOC by modifications to the microbial community (for example, removing phytoplankton or bacteria^{68,75,100} or adding viruses⁴⁵) or by manipulation of environmental parameters (for example, irradiance¹⁷⁴), followed by chemical analysis of the altered pools. Drawdown studies introduce bacteria into a DOC pool and rely on chemical analysis to identify features that are depleted⁷⁰, such as characterization of bacterial substrate use by different species⁷⁰. Mutant screening studies introduce bacteria with disrupted genes into a DOC pool and rely on chemical analysis to identify features that are no longer depleted, for example, identification of the phytoplankton exometabolite dihydroxypropanesulfonate (DHPS) from its accumulation in a transporter mutant assay⁶⁹. Enzymatic activity assays use selective digestion by high-specificity bacterial enzymes to quantify labile compounds within complex mixtures, for example, hydrolases that degrade laminarin into diagnostic sugar units that are readily measured¹⁷⁵. Other biological screening strategies that are currently used primarily in the field of biochemistry have promise for adoption in ecological studies. Vesicular transport assays embed transporters in synthetic membrane vesicles in an inside-out orientation, trapping target metabolites in the vesicle for chemical analysis¹⁷⁶. Metabolite–protein binding assays detect enzymes that bind to known metabolites, for instance, identification of substrates of *Escherichia coli* enzymes that previously lacked functional annotation¹⁷⁷. Finally, activity-based protein profiling uses chemical probes with reporter tags that mimic metabolites and form covalent bonds with microbial enzymes, for instance, to identify novel catabolic enzymes that degrade cellulose¹⁷⁸. The coupling of biological screening with chemical tools holds promise for pinpointing the hidden chemical currencies of the surface ocean and the genes and enzymes that transform them.

Modelling microbial metabolism. Modelling approaches are being used to extract carbon-cycle-relevant insights from observations of the microbe–metabolite network of the ocean. The challenge for models is to bridge a spatial scale spanning 13 orders of magnitude, from cell metabolism at the scale of 10^{-6} m to ocean flux at the scale of 10^7 m. Component models that focus on specific portions of this spatial scale already exist or are being developed. At the microbial end, earlier models of DOC release by phytoplankton sought

a mechanistic understanding based on parameters such as phytoplankton size, nutrient status and photosynthetic output^{179–181}. More recent models use genomic data to address the biochemical basis for metabolite production and consumption. Metabolic models based on flux balance analysis¹⁸² typically optimize for generation of new biomass, but could optimize for other physiological or ecological traits¹⁸³ such as metabolite release, abiotic stress tolerance or carbon use efficiency¹⁸⁴. Phylometabolic modelling integrates comparative genomics with insights from biochemistry and ecology to reconstruct metabolic innovations that affect metabolite production and consumption, such as complementary organic matter exchange between phytoplankton and bacteria¹⁷. Multicell metabolic models for single colonies or multispecies communities¹⁸⁵ uncover rules governing metabolite exchange and provide parameters for models working at regional to global scales^{186,187}.

In the transition from cellular scales to regional and global scales, hard-fought details of metabolite chemistry and biology must be simplified but not trivialized. The need to identify the optimal balance between excessive detail and oversimplification has emerged as a crucial barrier to incorporating microbial processes into global models¹⁸⁸, along with high computational costs, limited conceptual foundation, and lack of data to formulate and evaluate the more complex models. Nonetheless, global and regional DOC dynamics can be captured through nutrient–phytoplankton–zooplankton–detritus (NPZD) modules, which are widely used to compare historical and future climates^{189,190}. These models approximate the details of microbe–metabolite networks with bulk functions (such as Michaelis–Menten equations for substrate uptake) and use simplified rules to track substrates and energy through core metabolic pathways¹⁸⁹. Conceptual convergence between models at different scales can potentially be leveraged, for example, by linking cell growth output from steady-state flux balance analysis models with resource allocation rules applied to NPZD models.

The recently developed ‘emergent’ models, in which microbial community structure and function emerge from a wider set of possibilities, combine microbial genomic or physiological data with a dynamic physical–chemical ocean model to observe biogeochemical outcomes. Outputs from emergent models have revealed, for example, a predictable assembly of communities based on functional repertoire rather than taxonomic affiliation¹⁹¹ and matched distributions and abundance of model microbes with their real-world counterparts across ocean light and temperature gradients¹⁹². An inherent struggle with these models is parameterizing the bioenergetic cost of a gene or gene function^{191,193,194}. More work is needed to overcome these challenges, but it is clear that successful modelling

of labile DOC flux at regional and global scales will close one of the largest knowledge gaps in the global carbon cycle.

Computational data science approaches. Software solutions that respond to the needs for interdisciplinarity and integration in microbiome science are emerging^{195–198}, including those that build microbe–metabolite networks with machine-learning tools¹⁹⁹. Other strategies address metabolite-related gene annotation by merging information on ‘genes without a metabolite’ with information on ‘metabolites without a gene’. For example, the metabolite annotation and gene integration (MAGI)²⁰⁰ method identifies patterns of metabolite–gene associations by scoring the consensus in occurrence between the two datasets. The crowd-sourced Global Natural Product Social Molecular Networking (GNPS)²⁰¹ database improves metabolite identification and annotation by enabling comparisons of fragmentation spectra. Finally, the Paired Omics Data Platform links matched genomic and metabolomic data deposited in public repositories²⁰². As annotations improve, *in silico* reconstructions can serve as knowledge repositories that facilitate data integration of reactions and pathways and enable predictions of microbial biosynthetic capabilities as environmental conditions, genetic perturbations and fitness functions^{111,203,204} vary. These informatic-centric approaches offer starting points for improved inventories of the microbial metabolites and identification of genes with important roles in surface ocean carbon flux.

Despite both progress and interest, the ability to co-investigate chemical compounds and their genetic determinants across space and time remains a bottleneck to characterizing microbe–metabolite networks. Data sharing is imperative among marine chemists and microbiologists to enable discoverability, integration and interoperability of data across software tools. Moreover, data exchange through software is necessary to leverage a growing public database that merges marine bacterial, archaeal and eukaryotic genomes^{205,206}, metagenomes^{207–209}, metatranscriptomes²¹⁰ and metaproteomes²¹¹, and annotation resources for chemical oceanography^{201,212}. Integrative strategies will lead to characterization of the genes that link microbial activity to the production and consumption of key metabolites.

Microbial currencies in a warming ocean

The well-recognized downward export of particulate organic carbon from the surface ocean to deep ocean waters and sediments (the ‘biological carbon pump’) is globally important because it isolates carbon from the atmospheric pool for hundreds to thousands of years^{213,214}. The microbial remineralization of labile carbon to its inorganic form is globally important because it diverts carbon from the biological carbon pump, reducing net community production and influencing air–sea CO₂ fluxes^{215,216}. The quantitative importance of labile DOC remineralization becomes clear by considering that it constitutes >0.1% of the total DOC pool yet accounts for 86% of total DOC turnover². This implies that rapidly cycled microbial metabolites are among the most important individual conduits of ocean carbon.

Recent studies have provided a framework for predicting how phytoplankton-derived components of labile DOC might be altered under future climate scenarios, despite often complex and species-specific responses. Increasing temperature has been shown to enhance total carbon release by phytoplankton¹⁵. Evidence suggests that increased ribosome efficiency under higher temperatures will decrease phosphorus demand^{217–219} and will lower the N:P stoichiometry of labile DOC as rRNA synthesis needs are lessened. Mismatches between phytoplankton photosynthetic flux (weakly affected by temperature) and metabolism (strongly affected by temperature) can have diverse impacts on cellular resource allocation and metabolism²¹⁹. Increased CO₂ concentrations can decrease photorespiration rates and alter the release of photorespiration products such as glycolate^{220,221}. Finally, a warming climate is predicted

to favour smaller phytoplankton cells, which are better able to compete for nutrients in a stratified ocean^{222,223}, shifting labile DOC chemistry towards metabolite profiles characteristic of cyanobacteria and green algae^{18,224}. As these predictions are being formulated, the microbial functions underlain by the production and cycling of the substrates, facilitators and ecological signals exchanged via the labile DOC pool are being recognized.

Conclusion

Details of the metabolic currencies that transfer carbon between microbes to sustain the surface ocean carbon cycle have until recently been largely invisible to scientists. Importantly, climate–carbon feedback loops mediated by the labile DOC reservoir depend on this microbe–metabolite network. The resilience of the ocean to planetary change will rely on responses of the network to temperature increases, ocean acidification and other linked environmental changes. With advances in the chemical, biomolecular and data sciences, more of these nearly invisible molecules and their roles in the ocean carbon economy are being recognized. The future promises rapid scientific advances in understanding the chemical currencies of the surface ocean carbon cycle, at a time when increased knowledge is needed to safeguard against the effects of global change.

Received: 6 December 2019; Accepted: 23 February 2022;
Published online: 1 April 2022

References

1. Azam, F. et al. The ecological role of water-column microbes. *Mar. Ecol. Prog. Ser.* **10**, 257–263 (1983).
2. Hansell, D. A. Recalcitrant dissolved organic carbon fractions. *Ann. Rev. Mar. Sci.* **5**, 421–445 (2013).
3. Weber, L. et al. Extracellular reef metabolites across the protected Jardines de la Reina, Cuba reef system. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2020.582161> (2020).
4. Widner, B., Kido Soule, M. C., Ferrer-González, F. X., Moran, M. A. & Kujawinski, E. B. Quantification of amine- and alcohol-containing metabolites in saline samples using pre-extraction benzoyl chloride derivatization and ultrahigh performance liquid chromatography tandem mass spectrometry (UHPLC MS/MS). *Anal. Chem.* **93**, 4809–4817 (2021).
5. Fogg, G. E. The ecological significance of extracellular products of phytoplankton photosynthesis. *Bot. Mar.* **26**, 3–14 (1983).
6. Morán, X. A. G., Ducklow, H. W. & Erickson, M. Carbon fluxes through estuarine bacteria reflect coupling with phytoplankton. *Mar. Ecol. Prog. Ser.* **489**, 75–85 (2013).
7. Mühlenbruch, M., Grossart, H. P., Eigemann, F. & Voss, M. Phytoplankton-derived polysaccharides in the marine environment and their interactions with heterotrophic bacteria. *Environ. Microbiol.* **20**, 2671–2685 (2018).
8. Flynn, K. J., Clark, D. R. & Xue, Y. Modeling the release of dissolved organic matter by phytoplankton 1. *J. Phycol.* **44**, 1171–1187 (2008).
9. Björriksen, P. K. Phytoplankton exudation of organic matter: why do healthy cells do it? *Limnol. Oceanogr.* **33**, 151–154 (1988).
10. Myklestad, S. M. Release of extracellular products by phytoplankton with special emphasis on polysaccharides. *Sci. Total Environ.* **165**, 155–164 (1995).
11. Falkowski, P. & Raven, J. *Aquatic Photosynthesis* 2nd edn (Princeton Univ. Press, 2007).
12. Lau, W. W. & Armbrust, E. V. Detection of glycolate oxidase gene *glcD* diversity among cultured and environmental marine bacteria. *Environ. Microbiol.* **8**, 1688–1702 (2006).
13. Wood, A. M. & Van Valen, L. M. Paradox lost? On the release of energy-rich compounds by photoplankton. *Mar. Microb. Food Webs* **4**, 103–116 (1990).
14. Hellebust, J. Excretion of organic compounds by cultured and natural populations of marine phytoplankton. *Estuaries* **10**, 192–206 (1967).
15. Thornton, D. C. Dissolved organic matter (DOM) release by phytoplankton in the contemporary and future ocean. *Eur. J. Phycol.* **49**, 20–46 (2014).
16. Wangersky, P. in *Microbial Ecology* Vol. IV (ed. Kinne, O) 115–220 (Wiley, 1978).
17. Braakman, R., Follows, M. J. & Chisholm, S. W. Metabolic evolution and the self-organization of ecosystems. *Proc. Natl Acad. Sci. USA* **114**, E3091–E3100 (2017).
18. Durham, B. P. et al. Sulfonate-based networks between eukaryotic phytoplankton and heterotrophic bacteria in the surface ocean. *Nat. Microbiol.* **4**, 1706–1715 (2019).

19. Mague, T., Friberg, E., Hughes, D. & Morris, I. Extracellular release of carbon by marine phytoplankton; a physiological approach. *Limnol. Oceanogr.* **24**, 262–279 (1980).
20. Vardi, A. et al. A stress surveillance system based on calcium and nitric oxide in marine diatoms. *PLoS Biol.* **4**, e60 (2006).
21. Gillard, J. et al. Metabolomics enables the structure elucidation of a diatom sex pheromone. *Angew. Chem. Int. Ed.* **52**, 854–857 (2013).
22. Brunson, J. K. et al. Biosynthesis of the neurotoxin domoic acid in a bloom-forming diatom. *Science* **361**, 1356–1358 (2018).
23. Legrand, C., Rengefors, K., Fistarol, G. O. & Graneli, E. Allelopathy in phytoplankton-biochemical, ecological and evolutionary aspects. *Phycologia* **42**, 406–419 (2003).
24. Granum, E., Kirkvold, S. & Mykkestad, S. M. Cellular and extracellular production of carbohydrates and amino acids by the marine diatom *Skeletonema costatum*: diel variations and effects of N depletion. *Mar. Ecol. Prog. Ser.* **242**, 83–94 (2002).
25. Mague, T., Friberg, E., Hughes, D. & Morris, I. Extracellular release of carbon by marine phytoplankton; a physiological approach 1. *Limnol. Oceanogr.* **25**, 262–279 (1980).
26. Puskaric, S. & Mortain-Bertrand, A. Physiology of diatom *Skeletonema costatum* (Grev.) Cleve photosynthetic extracellular release: evidence for a novel coupling between marine bacteria and phytoplankton. *J. Plankton Res.* **25**, 1227–1235 (2003).
27. Aylward, F. O. et al. Microbial community transcriptional networks are conserved in three domains at ocean basin scales. *Proc. Natl Acad. Sci. USA* **112**, 5443–5448 (2015).
28. Frischkorn, K. R., Haley, S. T. & Dyhrman, S. T. Coordinated gene expression between *Trichodesmium* and its microbiome over day–night cycles in the North Pacific Subtropical Gyre. *ISME J.* **12**, 997–1007 (2018).
29. Bidle, K. D. The molecular ecophysiology of programmed cell death in marine phytoplankton. *Ann. Rev. Mar. Sci.* **7**, 341–375 (2015).
30. Vardi, A. et al. Host–virus dynamics and subcellular controls of cell fate in a natural coccolithophore population. *Proc. Natl Acad. Sci. USA* **109**, 19327–19332 (2012).
31. Møller, E. F. & Nielsen, T. G. Production of bacterial substrate by marine copepods: effect of phytoplankton biomass and cell size. *J. Plankton Res.* **23**, 527–536 (2001).
32. Nagata, T. in *Microbial Ecology of the Oceans* (ed. Kirchman, D. L.) 121–152 (Wiley-Liss, 2000).
33. Strom, S. L., Benner, R., Ziegler, S. & Dagg, M. J. Planktonic grazers are a potentially important source of marine dissolved organic carbon. *Limnol. Oceanogr.* **42**, 1364–1374 (1997).
34. Suttle, C. A. Marine viruses—major players in the global ecosystem. *Nat. Rev. Microbiol.* **5**, 801–812 (2007).
35. Zhao, Z. et al. Microbial transformation of virus-induced dissolved organic matter from picocyanobacteria: coupling of bacterial diversity and DOM chemodiversity. *ISME J.* **13**, 2551–2565 (2019).
36. Hurwitz, B. L. & U'Ren, J. M. Viral metabolic reprogramming in marine ecosystems. *Curr. Opin. Microbiol.* **31**, 161–168 (2016).
37. Kuhlisch, C. et al. Viral infection of algal blooms leaves a unique metabolic footprint on the dissolved organic matter in the ocean. *Sci. Adv.* **7**, eabf4680 (2021).
38. Klawonn, I. et al. Characterizing the ‘fungal shunt’: parasitic fungi on diatoms affect carbon flow and bacterial communities in aquatic microbial food webs. *Proc. Natl Acad. Sci. USA* **118**, e2102225118 (2021).
39. Agustí, S., Satta, M. P., Mura, M. P. & Benavent, E. Dissolved esterase activity as a tracer of phytoplankton lysis: evidence of high phytoplankton lysis rates in the northwestern Mediterranean. *Limnol. Oceanogr.* **43**, 1836–1849 (1998).
40. Brussaard, C. et al. Effects of grazing, sedimentation and phytoplankton cell lysis on the structure of a coastal pelagic food web. *Mar. Ecol. Prog. Ser.* **123**, 259–271 (1995).
41. Enke, T. N. et al. Modular Assembly of Polysaccharide-Degrading Marine Microbial Communities. *Curr. Biol.* **29**, 1528–1535.e6 (2019).
42. Selander, E. et al. Predator lipids induce paralytic shellfish toxins in bloom-forming algae. *Proc. Natl Acad. Sci. USA* **112**, 6395–6400 (2015).
43. Steinberg, D. K. & Landry, M. R. Zooplankton and the ocean carbon cycle. *Ann. Rev. Mar. Sci.* **9**, 413–444 (2017).
44. Stoecker, D. K., Hansen, P. J., Caron, D. A. & Mitra, A. Mixotrophy in the marine plankton. *Ann. Rev. Mar. Sci.* **9**, 311–335 (2017).
45. Ankrah, N. Y. et al. Phage infection of an environmentally relevant marine bacterium alters host metabolism and lysate composition. *ISME J.* **8**, 1089–1100 (2014).
46. Howard-Varona, C. et al. Phage-specific metabolic reprogramming of virocells. *ISME J.* **14**, 881–895 (2020).
47. Saba, G. K., Steinberg, D. K. & Bronk, D. A. The relative importance of sloppy feeding, excretion, and fecal pellet leaching in the release of dissolved carbon and nitrogen by *Acartia tonsa* copepods. *J. Exp. Mar. Biol. Ecol.* **404**, 47–56 (2011).
48. Bayer, B. et al. Ammonia-oxidizing archaea release a suite of organic compounds potentially fueling prokaryotic heterotrophy in the ocean. *Environ. Microbiol.* **21**, 4062–4075 (2019).
49. Kujawinski, E. B. et al. Identification of possible source markers in marine dissolved organic matter using ultrahigh resolution mass spectrometry. *Geochim. Cosmochim. Acta* **73**, 4384–4399 (2009).
50. Schulz, S. & Dickschat, J. S. Bacterial volatiles: the smell of small organisms. *Nat. Prod. Rep.* **24**, 814–842 (2007).
51. Wienhausen, G., Noriega-Ortega, B. E., Niggemann, J., Dittmar, T. & Simon, M. The exometabolome of two model strains of the *Roseobacter* group: a marketplace of microbial metabolites. *Front. Microbiol.* **8**, 1985 (2017).
52. Ortega-Retuerta, E. et al. Dissolved organic matter released by two marine heterotrophic bacterial strains and its bioavailability for natural prokaryotic communities. *Environ. Microbiol.* <https://doi.org/10.1111/1462-2920.15306> (2020).
53. Kharbush, J. J. et al. Particulate organic carbon deconstructed: molecular and chemical composition of particulate organic carbon in the ocean. *Front. Mar. Sci.* **7**, 518 (2020).
54. Enke, T. N., Leventhal, G. E., Metzger, M., Saavedra, J. T. & Cordero, O. X. Microscale ecology regulates particulate organic matter turnover in model marine microbial communities. *Nat. Commun.* **9**, 2743 (2018).
55. Reintjes, G., Arnosti, C., Fuchs, B. & Amann, R. Selfish, sharing and scavenging bacteria in the Atlantic Ocean: a biogeographical study of bacterial substrate utilisation. *ISME J.* **13**, 1119–1132 (2019).
56. Smith, D. C., Simon, M., Alldredge, A. L. & Azam, F. Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature* **359**, 139–142 (1992).
57. Thor, P., Dam, H. G. & Rogers, D. R. Fate of organic carbon released from decomposing copepod fecal pellets in relation to bacterial production and ectoenzymatic activity. *Aquat. Microb. Ecol.* **33**, 279–288 (2003).
58. Cole, J. J., Findlay, S. & Pace, M. L. Bacterial production in fresh and saltwater ecosystems: a cross-system overview. *Mar. Ecol. Prog. Ser.* **43**, 1–10 (1988).
59. Azam, F. Microbial control of oceanic carbon flux: the plot thickens. *Science* **280**, 694–696 (1998).
60. Moran, M. A. et al. The ocean’s labile DOC supply chain. *Limnol. Oceanogr.* <https://doi.org/10.1002/lno.12053> (2022).
61. Lindeman, R. L. The trophic-dynamic aspect of ecology. *Ecology* **23**, 399–417 (1942).
62. Hollibaugh, J., Carruthers, A., Fuhrman, J. & Azam, F. Cycling of organic nitrogen in marine plankton communities studied in enclosed water columns. *Mar. Biol.* **59**, 15–21 (1980).
63. Keil, R. G. & Kirchman, D. L. Utilization of dissolved protein and amino acids in the northern Sargasso Sea. *Aquat. Microb. Ecol.* **18**, 293–300 (1999).
64. Boysen, A. K. et al. Particulate metabolites and transcripts reflect diel oscillations of microbial activity in the surface ocean. *mSystems* **6**, e00896-20 (2021).
65. Dawson, H. M. et al. Potential of temperature-and salinity-driven shifts in diatom compatible solute concentrations to impact biogeochemical cycling within sea ice. *Elementa* <https://doi.org/10.1525/elementa.421> (2020).
66. Durham, B. P. Deciphering metabolic currencies that support marine microbial networks. *mSystems* **6**, e0076321 (2021).
67. Fiore, C. L., Longnecker, K., Kido Soule, M. C. & Kujawinski, E. B. Release of ecologically relevant metabolites by the cyanobacterium *Synechococcus elongatus* CCMP 1631. *Environ. Microbiol.* **17**, 3949–3963 (2015).
68. Durham, B. P. et al. Recognition cascade and metabolite transfer in a marine bacteria-phytoplankton model system. *Environ. Microbiol.* **19**, 3500–3513 (2017).
69. Durham, B. P. et al. Cryptic carbon and sulfur cycling between surface ocean plankton. *Proc. Natl Acad. Sci. USA* **112**, 453–457 (2015).
70. Ferrer-González, F. X. et al. Resource partitioning of phytoplankton metabolites that support bacterial heterotrophy. *ISME J.* **15**, 762–773 (2020).
71. Gebser, B., Thume, K., Steinke, M. & Pohnert, G. Phytoplankton-derived zwitterionic gonyol and dimethylsulfonioacetate interfere with microbial dimethylsulfoniopropionate sulfur cycling. *MicrobiologyOpen* **9**, e1014 (2020).
72. Heal, K. R. et al. Marine community metabolomes carry fingerprints of phytoplankton community composition. *mSystems* **6**, e01334-20 (2020).
73. Kiene, R. P., Linn, L. J. & Bruton, J. A. New and important roles for DMSP in marine microbial communities. *J. Sea Res.* **43**, 209–224 (2000).
74. Landa, M., Burns, A. S., Roth, S. J. & Moran, M. A. Bacterial transcriptome remodeling during sequential co-culture with a marine dinoflagellate and diatom. *ISME J.* **11**, 2677 (2017).
75. Shibl, A. A. et al. Diatom modulation of select bacteria through use of two unique secondary metabolites. *Proc. Natl Acad. Sci. USA* **117**, 27445–27455 (2020).
76. Uchimiya, M., Schroer, W., Olofsson, M., Edison, A. S. & Moran, M. A. Diel investments in metabolite production and consumption in a model microbial system. *ISME J.* <https://doi.org/10.1038/s41396-021-01172-w> (2021).

77. Mopper, K. & Lindroth, P. Diel and depth variations in dissolved free amino acids and ammonium in the Baltic Sea determined by shipboard HPLC analysis 1. *Limnol. Oceanogr.* **27**, 336–347 (1982).
78. Pomeroy, L. R., Macko, S., Ostrom, P. & Dunphy, J. The microbial food web in Arctic seawater: concentration of dissolved free amino acids and bacterial abundance and activity in the Arctic Ocean and in Resolute Passage. *Mar. Ecol. Prog. Ser.* **61**, 31–40 (1990).
79. Liu, Q., Lu, X. X., Tolar, B. B., Mou, X. Z. & Hollibaugh, J. T. Concentrations, turnover rates and fluxes of polyamines in coastal waters of the South Atlantic Bight. *Biogeochemistry* **123**, 117–133 (2015).
80. Nishibori, N., Yuasa, A., Sakai, M., Fujihara, S. & Nishio, S. Free polyamine concentrations in coastal seawater during phytoplankton bloom. *Fish. Sci.* **67**, 79–83 (2001).
81. Benner, R. & Kaiser, K. Abundance of amino sugars and peptidoglycan in marine particulate and dissolved organic matter. *Limnol. Oceanogr.* **48**, 118–128 (2003).
82. Borch, N. H. & Kirchmann, D. L. Concentration and composition of dissolved combined neutral sugars (polysaccharides) in seawater determined by HPLC-PAD. *Mar. Chem.* **57**, 85–95 (1997).
83. Ittekkot, V., Brockmann, U., Michaelis, W. & Degens, E. T. Dissolved free and combined carbohydrates during a phytoplankton bloom in the northern North Sea. *Mar. Ecol. Prog. Ser.* **4**, 299–305 (1981).
84. Lau, W. W., Keil, R. G. & Armbrust, E. V. Succession and diel transcriptional response of the glycolate-utilizing component of the bacterial community during a spring phytoplankton bloom. *Appl. Environ. Microbiol.* **73**, 2440–2450 (2007).
85. Bertrand, E. M. & Allen, A. E. Influence of vitamin B auxotrophy on nitrogen metabolism in eukaryotic phytoplankton. *Front. Microbiol.* **3**, 375 (2012).
86. Gómez-Consarnau, L. et al. Mosaic patterns of B-vitamin synthesis and utilization in a natural marine microbial community. *Environ. Microbiol.* **20**, 2809–2823 (2018).
87. Sanudo-Wilhelmy, S. A. et al. Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc. Natl Acad. Sci. USA* **109**, 14041–14045 (2012).
88. Amin, S. A. et al. Photolysis of iron-siderophore chelates promotes bacterial–algal mutualism. *Proc. Natl Acad. Sci. USA* **106**, 17071–17076 (2009).
89. D’Onofrio, A. et al. Siderophores from neighboring organisms promote the growth of uncultured bacteria. *Chem. Biol.* **17**, 254–264 (2010).
90. Kenney, G. E. et al. The biosynthesis of methanobactin. *Science* **359**, 1411–1416 (2018).
91. Lhospice, S. et al. *Pseudomonas aeruginosa* zinc uptake in chelating environment is primarily mediated by the metallophore pseudopaline. *Sci. Rep.* **7**, 17132 (2017).
92. Button, D., Robertson, B., Gustafson, E. & Zhao, X. Experimental and theoretical bases of specific affinity, a cytoarchitecture-based formulation of nutrient collection proposed to supercede the Michaelis–Menten paradigm of microbial kinetics. *Appl. Environ. Microbiol.* **70**, 5511–5521 (2004).
93. Button, D., Robertson, B. R., Lepp, P. W. & Schmidt, T. M. A small, dilute-cytoplasm, high-affinity, novel bacterium isolated by extinction culture and having kinetic constants compatible with growth at ambient concentrations of dissolved nutrients in seawater. *Appl. Environ. Microbiol.* **64**, 4467–4476 (1998).
94. Heal, K. R. et al. Determination of four forms of vitamin B12 and other B vitamins in seawater by liquid chromatography/tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **28**, 2398–2404 (2014).
95. Mawji, E. et al. Hydroxamate siderophores: occurrence and importance in the Atlantic Ocean. *Environ. Sci. Technol.* **42**, 8675–8680 (2008).
96. Morris, J. J., Lenski, R. E. & Zinser, E. R. The black queen hypothesis: evolution of dependencies through adaptive gene loss. *mBio* <https://doi.org/10.1128/mBio.00036-12> (2012).
97. Kazamia, E., Helliwell, K. E., Purton, S. & Smith, A. G. How mutualisms arise in phytoplankton communities: building eco-evolutionary principles for aquatic microbes. *Ecol. Lett.* **19**, 810–822 (2016).
98. Pohnert, G., Steinke, M. & Tollrian, R. Chemical cues, defence metabolites and the shaping of pelagic interspecific interactions. *Trends Ecol. Evol.* **22**, 198–204 (2007).
99. Vos, M. et al. Infochemicals structure marine, terrestrial and freshwater food webs: implications for ecological informatics. *Ecol. Inf.* **1**, 23–32 (2006).
100. Amin, S. et al. Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* **522**, 98–101 (2015).
101. Hmelo, L. & Van Mooy, B. A. Kinetic constraints on acylated homoserine lactone-based quorum sensing in marine environments. *Aquat. Microb. Ecol.* **54**, 127–133 (2009).
102. Schwartz, E. R., Poulin, R. X., Mojib, N. & Kubanek, J. Chemical ecology of marine plankton. *Nat. Prod. Rep.* **33**, 843–860 (2016).
103. Nelson, C. E. & Wear, E. K. Microbial diversity and the lability of dissolved organic carbon. *Proc. Natl Acad. Sci. USA* **111**, 7166–7167 (2014).
104. Amaral-Zettler, L. et al. in *Life in the World’s Oceans: Diversity, Distribution and Abundance* (ed. McIntyre, A.) 223–245 (Wiley, 2010).
105. De Vargas, C. et al. Eukaryotic plankton diversity in the sunlit ocean. *Science* **348**, 1261605 (2015).
106. Hertkorn, N. et al. Natural organic matter and the event horizon of mass spectrometry. *Anal. Chem.* **80**, 8908–8919 (2008).
107. Lechtenfeld, O. J., Hertkorn, N., Shen, Y., Witt, M. & Benner, R. Marine sequestration of carbon in bacterial metabolites. *Nat. Commun.* **6**, 6711 (2015).
108. Longnecker, K., Sievert, S. M., Sylva, S. P., Seewald, J. S. & Kujawinski, E. B. Dissolved organic carbon compounds in deep-sea hydrothermal vent fluids from the East Pacific Rise at 9° 50’ N. *Org. Geochem.* **125**, 41–49 (2018).
109. Romano, S. et al. Exo-metabolome of *Pseudovibrio* sp. FO-BEG1 analyzed by ultra-high resolution mass spectrometry and the effect of phosphate limitation. *PLoS ONE* **9**, e96038 (2014).
110. Förster, J., Famili, I., Fu, P., Palsson, B. Ø. & Nielsen, J. Genome-scale reconstruction of the *Saccharomyces cerevisiae* metabolic network. *Genome Res.* **13**, 244–253 (2003).
111. Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of interspecies interactions in microbial ecosystems. *Nat. Commun.* **10**, 103 (2019).
112. Hertkorn, N. et al. Characterization of a major refractory component of marine dissolved organic matter. *Geochim. Cosmochim. Acta* **70**, 2990–3010 (2006).
113. Steen, A. D. et al. Analytical and computational advances, opportunities, and challenges in marine organic biogeochemistry in an era of ‘omics’. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2020.00718> (2020).
114. Becker, J. W. et al. Closely related phytoplankton species produce similar suites of dissolved organic matter. *Front. Microbiol.* **5**, 111 (2014).
115. Bouslimani, A., Sanchez, L. M., Garg, N. & Dorrestein, P. C. Mass spectrometry of natural products: current, emerging and future technologies. *Nat. Prod. Rep.* **31**, 718–729 (2014).
116. Johnson, W. M. et al. Metabolite composition of sinking particles differs from surface suspended particles across a latitudinal transect in the South Atlantic. *Limnol. Oceanogr.* **65**, 111–127 (2020).
117. Metwally, H., McAllister, R. G. & Konermann, L. Exploring the mechanism of salt-induced signal suppression in protein electrospray mass spectrometry using experiments and molecular dynamics simulations. *Anal. Chem.* **87**, 2434–2442 (2015).
118. Kovacs, H., Moskau, D. & Spraul, M. Cryogenically cooled probes—a leap in NMR technology. *Prog. Nucl. Magn. Reson. Spectrosc.* **2**, 131–155 (2005).
119. Aluwihare, L. I., Repeta, D. J. & Chen, R. F. A major biopolymeric component to dissolved organic carbon in surface sea water. *Nature* **387**, 166–169 (1997).
120. Koprivnjak, J.-F. et al. Chemical and spectroscopic characterization of marine dissolved organic matter isolated using coupled reverse osmosis–electrodialysis. *Geochim. Cosmochim. Acta* **73**, 4215–4231 (2009).
121. Dittmar, T., Koch, B., Hertkorn, N. & Kattner, G. A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater. *Limnol. Oceanogr. Meth.* **6**, 230–235 (2008).
122. Johnson, W. M., Soule, M. C. K. & Kujawinski, E. B. Extraction efficiency and quantification of dissolved metabolites in targeted marine metabolomics. *Limnol. Oceanogr. Meth.* **15**, 417–428 (2017).
123. Lee, C. & Bada, J. L. Amino acids in equatorial Pacific Ocean water. *Earth Planet Sci. Lett.* **26**, 61–68 (1975).
124. Yamashita, Y. & Tanoue, E. Chemical characteristics of amino acid-containing dissolved organic matter in seawater. *Org. Geochem.* **35**, 679–692 (2004).
125. Zubkov, M. V., Tarran, G. A., Mary, I. & Fuchs, B. M. Differential microbial uptake of dissolved amino acids and amino sugars in surface waters of the Atlantic Ocean. *J. Plankton Res.* **30**, 211–220 (2008).
126. Unanue, M. et al. Ecto-enzymatic activity and uptake of monomers in marine bacterioplankton described by a biphasic kinetic model. *Microb. Ecol.* **37**, 36–48 (1999).
127. Nissen, H., Nissen, P. & Azam, F. Multiphasic uptake of D-glucose by an oligotrophic marine bacterium. *Mar. Ecol. Prog. Ser.* **16**, 155–160 (1984).
128. Pedler, B. E., Aluwihare, L. I. & Azam, F. Single bacterial strain capable of significant contribution to carbon cycling in the surface ocean. *Proc. Natl Acad. Sci. USA* **111**, 7202–7207 (2014).
129. Giovannoni, S. J., Thrash, J. C. & Temperton, B. Implications of streamlining theory for microbial ecology. *ISME J.* **8**, 1553–1565 (2014).
130. Norris, N., Levine, N. M., Fernandez, V. I. & Stocker, R. Mechanistic model of nutrient uptake explains dichotomy between marine oligotrophic and copiotrophic bacteria. *PLoS Comp. Biol.* **17**, e1009023 (2021).
131. Stocker, R., Seymour, J. R., Samadani, A., Hunt, D. E. & Polz, M. F. Rapid chemotactic response enables marine bacteria to exploit ephemeral microscale nutrient patches. *Proc. Natl Acad. Sci. USA* **105**, 4209–4214 (2008).

132. Keil, R. G. & Kirchman, D. L. Dissolved combined amino acids—chemical form and utilization by marine bacteria. *Limnol. Oceanogr.* **38**, 1256–1270 (1993).
133. Mopper, K. et al. Determination of sugars in unconcentrated seawater and other natural waters by liquid chromatography and pulsed amperometric detection. *Environ. Sci. Technol.* **26**, 133–138 (1992).
134. Zakem, E. J., Cael, B. & Levine, N. M. A unified theory for organic matter accumulation. *Proc. Natl Acad. Sci. USA* **118**, e2016896118 (2021).
135. Moran, M. A., Reisch, C. R., Kiene, R. P. & Whitman, W. B. Genomic insights into bacterial DMSP transformations. *Ann. Rev. Mar. Sci.* **4**, 523–542 (2012).
136. Kiene, R. P. & Slezak, D. Low dissolved DMSP concentrations in seawater revealed by small-volume gravity filtration and dialysis sampling. *Limnol. Oceanogr. Meth.* **4**, 80–95 (2006).
137. Smriga, S., Fernandez, V. I., Mitchell, J. G. & Stocker, R. Chemotaxis toward phytoplankton drives organic matter partitioning among marine bacteria. *Proc. Natl Acad. Sci. USA* **113**, 1576–1581 (2016).
138. Azam, F. & Malfatti, F. Microbial structuring of marine ecosystems. *Nat. Rev. Microbiol.* **5**, 782–791 (2007).
139. Seymour, J. R., Amin, S. A., Raina, J.-B. & Stocker, R. Zooming in on the phycosphere: the ecological interface for phytoplankton–bacteria relationships. *Nat. Microbiol.* **2**, 17065 (2017).
140. Kanehisa, M. & Goto, S. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27–30 (2000).
141. Fenn, J. B., Mann, M., Meng, C. K., Wong, S. F. & Whitehouse, C. M. Electrospray ionization for mass spectrometry of large biomolecules. *Science* **246**, 64–71 (1989).
142. Marshall, A. G., Hendrickson, C. L. & Jackson, G. S. Fourier transform ion cyclotron resonance mass spectrometry: a primer. *Mass Spectrom. Rev.* **17**, 1–35 (1998).
143. Zubarev, R. A. & Makarov, A. Orbitrap mass spectrometry. *Anal. Chem.* **85**, 5288–5296 (1998).
144. Longnecker, K. & Kujawinski, E. B. Mining mass spectrometry data: using new computational tools to find novel organic compounds in complex environmental mixtures. *Org. Geochem.* **110**, 92–99 (2017).
145. Ardenjaer-Larsen, J. H. et al. Facing and overcoming sensitivity challenges in biomolecular NMR spectroscopy. *Angew. Chem. Int. Ed.* **54**, 9162–9185 (2015).
146. Ramaswamy, V. et al. Microsample cryogenic probes: technology and applications. *eMagRes* <https://doi.org/10.1002/9780470034590.emrstm1315> (2007).
147. Edison, A. S. et al. NMR: unique strengths that enhance modern metabolomics research. *Anal. Chem.* **93**, 478–499 (2020).
148. Anaraki, M. T. et al. NMR assignment of the in vivo *Daphnia magna* metabolome. *Analyst* **145**, 5787–5800 (2020).
149. Qi, Y. et al. Absorption-mode: the next generation of Fourier transform mass spectra. *Anal. Chem.* **84**, 2923–2929 (2012).
150. Liaghati Mobarhan, Y., Soong, R., Lane, D. & Simpson, A. J. In vivo comprehensive multiphase NMR. *Mag. Res. Chem.* **58**, 427–444 (2020).
151. Zhao, S., Li, H., Han, W., Chan, W. & Li, L. Metabolomic coverage of chemical-group-submetabolome analysis: group classification and four-channel chemical isotope labeling LC–MS. *Anal. Chem.* **91**, 12108–12115 (2019).
152. Sogin, E. M., Puskas, E., Dubilier, N. & Liebeke, M. Marine metabolomics: measurement of metabolites in seawater by gas chromatography mass spectrometry. *mSystems* **4**, e00638–19 (2019).
153. Pontrelli, S. & Sauer, U. Salt-tolerant metabolomics for exometabolomic measurements of marine bacterial isolates. *Anal. Chem.* **93**, 7164–7171 (2021).
154. Mayali, X., Weber, P. K., Mabery, S. & Pett-Ridge, J. Phylogenetic patterns in the microbial response to resource availability: amino acid incorporation in San Francisco Bay. *PLoS ONE* **9**, e95842 (2014).
155. Thomas, F. et al. Isotopic tracing reveals single-cell assimilation of a macroalgal polysaccharide by a few marine Flavobacteria and Gammaproteobacteria. *ISME J.* **15**, 3062–3075 (2021).
156. Hatzepichler, R. et al. In situ visualization of newly synthesized proteins in environmental microbes using amino acid tagging and click chemistry. *Environ. Microbiol.* **16**, 2568–2590 (2014).
157. Martinez-Garcia, M. et al. Capturing single cell genomes of active polysaccharide degraders: an unexpected contribution of Verrucomicrobia. *PLoS ONE* **7**, e35314 (2012).
158. Raina, J.-B. et al. Subcellular tracking reveals the location of dimethylsulfoniopropionate in microalgae and visualises its uptake by marine bacteria. *eLife* **6**, e23008 (2017).
159. Vasdekis, A. E. & Stephanopoulos, G. Review of methods to probe single cell metabolism and bioenergetics. *Metab. Eng.* **27**, 115–135 (2015).
160. Baumeister, T., Vallet, M., Kaftan, F., Svatoš, A. & Pohnert, G. Live single-cell metabolomics with matrix-free laser/desorption ionization mass spectrometry to address microalgal physiology. *Front. Plant Sci.* **10**, 172 (2019).
161. Schleyer, G. et al. In plaque-mass spectrometry imaging of a bloom-forming alga during viral infection reveals a metabolic shift towards odd-chain fatty acid lipids. *Nat. Microbiol.* **4**, 527–538 (2019).
162. Judge, M. T. et al. Continuous in vivo metabolism by NMR. *Front. Mol. Biosci.* **6**, 26 (2019).
163. Gao, C. et al. Single-cell bacterial transcription measurements reveal the importance of dimethylsulfoniopropionate (DMSP) hotspots in ocean sulfur cycling. *Nat. Commun.* **11**, 1942 (2020).
164. Szul, M. J., Dearth, S. P., Campagna, S. R. & Zinser, E. R. Carbon fate and flux in *Prochlorococcus* under nitrogen limitation. *mSystems* **4**, e00254–18 (2019).
165. Teeling, H. et al. Substrate-controlled succession of marine bacterioplankton populations induced by a phytoplankton bloom. *Science* **336**, 608–611 (2012).
166. Nowinski, B. & Moran, M. A. Niche dimensions of a marine bacterium are identified using invasion studies in coastal seawater. *Nat. Microbiol.* **6**, 524–532 (2021).
167. Poretsky, R. S. et al. Comparative day/night metatranscriptomic analysis of microbial communities in the North Pacific Subtropical Gyre. *Environ. Microbiol.* **11**, 1358–1375 (2009).
168. Sowell, S. M. et al. Transport functions dominate the SAR11 metaproteome at low-nutrient extremes in the Sargasso Sea. *ISME J.* **3**, 93–105 (2009).
169. Lidbury, I., Murrell, J. C. & Chen, Y. Trimethylamine N-oxide metabolism by abundant marine heterotrophic bacteria. *Proc. Natl Acad. Sci. USA* **111**, 2710–2715 (2014).
170. Tang, K., Jiao, N., Liu, K., Zhang, Y. & Li, S. Distribution and functions of TonB-dependent transporters in marine bacteria and environments: implications for dissolved organic matter utilization. *PLoS ONE* **7**, e41204 (2012).
171. Lima-Mendez, G. et al. Determinants of community structure in the global plankton interactome. *Science* **348**, 1262073 (2015).
172. Reverter, M., Rohde, S., Parchemin, C., Tapissier-Bontemps, N. & Schupp, P. J. Metabolomics and marine biotechnology: coupling metabolite profiling and organism biology for the discovery of new compounds. *Front. Mar. Sci.* **7**, 1062 (2020).
173. Frischkorn, K. R., Rouco, M., Van Mooy, B. A. & Dyhrman, S. T. Epibionts dominate metabolic functional potential of *Trichodesmium* colonies from the oligotrophic ocean. *ISME J.* **11**, 2090–2101 (2017).
174. Vorobev, A. et al. Identifying labile DOM components in a coastal ocean through depleted bacterial transcripts and chemical signals. *Environ. Microbiol.* **20**, 3012–3030 (2018).
175. Becker, S. et al. Laminarin is a major molecule in the marine carbon cycle. *Proc. Natl Acad. Sci. USA* **117**, 6599–6607 (2020).
176. Glavinas, H. et al. Utilization of membrane vesicle preparations to study drug–ABC transporter interactions. *Expert Opin. Drug Metab. Toxicol.* **4**, 721–732 (2008).
177. Piazza, I. et al. A map of protein–metabolite interactions reveals principles of chemical communication. *Cell* **172**, 358–372.e23 (2018).
178. Chauvigné-Hines, L. M. et al. Suite of activity-based probes for cellulose-degrading enzymes. *J. Am. Chem. Soc.* **134**, 20521–20532 (2012).
179. Baklouti, M., Diaz, F., Pinazo, C., Faure, V. & Quéguiner, B. Investigation of mechanistic formulations depicting phytoplankton dynamics for models of marine pelagic ecosystems and description of a new model. *Prog. Oceanogr.* **71**, 1–33 (2006).
180. Kovač, Ž., Platt, T., Sathyendranath, S. & Lomas, M. Extraction of photosynthesis parameters from time series measurements of in situ production: Bermuda Atlantic Time-Series Study. *Remote Sens.* **10**, 915 (2018).
181. Mongin, M., Nelson, D. M., Pondaven, P., Brzezinski, M. A. & Tréguer, P. Simulation of upper-ocean biogeochemistry with a flexible-composition phytoplankton model: C, N and Si cycling in the western Sargasso Sea. *Deep Sea Res. I* **50**, 1445–1480 (2003).
182. Karr, J. R. et al. A whole-cell computational model predicts phenotype from genotype. *Cell* **150**, 389–401 (2012).
183. Malik, A. A. et al. Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. *ISME J.* **14**, 1–9 (2020).
184. Ofaim, S., Sulheim, S., Almaas, E., Sher, D. & Segrè, D. Dynamic allocation of carbon storage and nutrient-dependent exudation in a revised genome-scale model of *Prochlorococcus*. *Front. Genet.* **12**, 91 (2021).
185. Colarusso, A. V., Goodchild-Michelman, I., Rayle, M. & Zomorodi, A. R. Computational modeling of metabolism in microbial communities on a genome-scale. *Curr. Opin. Syst. Biol.* <https://doi.org/10.1016/j.coisb.2021.04.001> (2021).
186. Harvey, E. L. et al. A bacterial quorum-sensing precursor induces mortality in the marine cocolithophore, *Emiliania huxleyi*. *Front. Microbiol.* **7**, 59 (2016).
187. Skalnik, C. J. et al. Whole-colony modeling of *Escherichia coli*. Preprint at *bioRxiv* <https://doi.org/10.1101/2021.04.27.441666> (2021).

188. Reid, A. *Incorporating Microbial Processes into Climate Models* (American Academy of Microbiology, 2012).
189. Nicholson, D. P., Stanley, R. H. & Doney, S. C. A phytoplankton model for the allocation of gross photosynthetic energy including the trade-offs of diazotrophy. *J. Geophys. Res. Biogeosci.* **123**, 1796–1816 (2018).
190. Bonan, G. B. & Doney, S. C. Climate, ecosystems, and planetary futures: the challenge to predict life in Earth system models. *Science* **359**, eaam8328 (2018).
191. Coles, V. et al. Ocean biogeochemistry modeled with emergent trait-based genomics. *Science* **358**, 1149–1154 (2017).
192. Follows, M. J., Dutkiewicz, S., Grant, S. & Chisholm, S. W. Emergent biogeography of microbial communities in a model ocean. *Science* **315**, 1843–1846 (2007).
193. Lane, N. & Martin, W. The energetics of genome complexity. *Nature* **467**, 929–934 (2010).
194. Lynch, M. & Marinov, G. K. The bioenergetic costs of a gene. *Proc. Natl Acad. Sci. USA* **112**, 15690–15695 (2015).
195. Bolduc, B., Youens-Clark, K., Roux, S., Hurwitz, B. L. & Sullivan, M. B. iVirus: facilitating new insights in viral ecology with software and community data sets imbedded in a cyberinfrastructure. *ISME J.* **11**, 7–14 (2017).
196. Eren, A. M. et al. Community-led, integrated, reproducible multi-omics with anvio. *Nat. Microbiol.* **6**, 3–6 (2021).
197. Harcombe, W. R. et al. Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. *Cell Rep.* **7**, 1104–1115 (2014).
198. Saito, M. A. et al. Development of an Ocean Protein Portal for interactive discovery and education. *J. Proteome Res.* **20**, 326–336 (2020).
199. Durán, C. et al. Nonlinear machine learning pattern recognition and bacteria-metabolite multilayer network analysis of perturbed gastric microbiome. *Nat. Commun.* **12**, 1926 (2021).
200. Erbilgin, O. et al. MAGI: a method for metabolite annotation and gene integration. *ACS Chem. Biol.* **14**, 704–714 (2019).
201. Wang, M. et al. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* **34**, 828–837 (2016).
202. Schorn, M. A. et al. A community resource for paired genomic and metabolomic data mining. *Nat. Chem. Biol.* **17**, 363–368 (2021).
203. Goldford, J. E. et al. Emergent simplicity in microbial community assembly. *Science* **361**, 469–474 (2018).
204. Zomorodi, A. R. & Segrè, D. Genome-driven evolutionary game theory helps understand the rise of metabolic interdependencies in microbial communities. *Nat. Commun.* **8**, 1563 (2017).
205. Pachiadaki, M. G. et al. Charting the complexity of the marine microbiome through single-cell genomics. *Cell* **179**, 1623–1635.e11 (2019).
206. Paoli, L. et al. Uncharted biosynthetic potential of the ocean microbiome. Preprint at *bioRxiv* <https://doi.org/10.1101/2021.03.24.436479> (2021).
207. Delmont, T. O. et al. Functional repertoire convergence of distantly related eukaryotic plankton lineages revealed by genome-resolved metagenomics. Preprint at *bioRxiv* <https://doi.org/10.1101/2020.10.15.341214> (2021).
208. Sunagawa, S. et al. Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).
209. Acinas, S. G. et al. Deep ocean metagenomes provide insight into the metabolic architecture of bathypelagic microbial communities. *Commun. Biol.* **4**, 604 (2021).
210. Salazar, G. et al. Gene expression changes and community turnover differentially shape the global ocean metatranscriptome. *Cell* **179**, 1068–1083.e21 (2019).
211. Saito, M. A. et al. Progress and challenges in ocean metaproteomics and proposed best practices for data sharing. *J. Proteome Res.* **18**, 1461–1476 (2019).
212. Chandler, C. et al. Effective management of ocean biogeochemistry and ecological data: the BCO-DMO story. In *EGU General Assembly Conference Abstracts* Vol. 14, 1258 (2012).
213. Ducklow, H. W., Steinberg, D. K. & Buesseler, K. O. Upper ocean carbon export and the biological pump. *Oceanography* **14**, 50–58 (2001).
214. Siegel, D. A., DeVries, T., Doney, S. C. & Bell, T. Assessing the sequestration time scales of some ocean-based carbon dioxide reduction strategies. *Environ. Res. Lett.* **16**, 104003 (2021).
215. Del Giorgio, P. A. & Duarte, C. M. Respiration in the open ocean. *Nature* **420**, 379–384 (2002).
216. Rivkin, R. B. & Legendre, L. Biogenic carbon cycling in the upper ocean: effects of microbial respiration. *Science* **291**, 2398–2400 (2001).
217. Toseland, A. et al. The impact of temperature on marine phytoplankton resource allocation and metabolism. *Nat. Clim. Change* **3**, 979–984 (2013).
218. Brembu, T., Mühlroth, A., Alipanah, L. & Bones, A. M. The effects of phosphorus limitation on carbon metabolism in diatoms. *Phil. Trans. R. Soc. B* **372**, 20160406 (2017).
219. Liang, Y., Koester, J. A., Liefer, J. D., Irwin, A. J. & Finkel, Z. V. Molecular mechanisms of temperature acclimation and adaptation in marine diatoms. *ISME J.* **13**, 2415–2425 (2019).
220. Hennon, G. M., Hernández Limón, M. D., Haley, S. T., Juhl, A. R. & Dyhrman, S. T. Diverse CO₂-induced responses in physiology and gene expression among eukaryotic phytoplankton. *Front. Microbiol.* **8**, 2547 (2017).
221. Shi, D. et al. Interactive effects of light, nitrogen source, and carbon dioxide on energy metabolism in the diatom *Thalassiosira pseudonana*. *Limnol. Oceanogr.* **60**, 1805–1822 (2015).
222. Taylor, A. G. et al. Sharp gradients in phytoplankton community structure across a frontal zone in the California Current ecosystem. *J. Plankton Res.* **34**, 778–789 (2012).
223. Marinov, I., Doney, S. & Lima, I. Response of ocean phytoplankton community structure to climate change over the 21st century: partitioning the effects of nutrients, temperature and light. *Biogeosciences* **7**, 3941–3959 (2010).
224. Becker, K. W. et al. Daily changes in phytoplankton lipidomes reveal mechanisms of energy storage in the open ocean. *Nat. Commun.* **9**, 5179 (2018).
225. Johnson, W. M., Soule, M. C. K. & Kujawski, E. B. Evidence for quorum sensing and differential metabolite production by a marine bacterium in response to DMSP. *ISME J.* **10**, 2304–2316 (2016).
226. Matrai, P. & Keller, M. Total organic sulfur and dimethylsulfoniopropionate in marine phytoplankton: intracellular variations. *Mar. Biol.* **119**, 61–68 (1994).
227. Thume, K. et al. The metabolite dimethylsulfoxonium propionate extends the marine organosulfur cycle. *Nature* **563**, 412–415 (2018).
228. Baines, S. B. & Pace, M. L. The production of dissolved organic matter by phytoplankton and its importance to bacteria: patterns across marine and freshwater systems. *Limnol. Oceanogr.* **36**, 1078–1090 (1991).
229. Peironcely, J. E., Reijmers, T., Coulier, L., Bender, A. & Hankemeier, T. Understanding and classifying metabolite space and metabolite-likeness. *PLoS ONE* **6**, e28966 (2011).
230. Gifford, S. M., Sharma, S., Booth, M. & Moran, M. A. Expression patterns reveal niche diversification in a marine microbial assemblage. *ISME J.* **7**, 281–298 (2013).
231. Poretsky, R. S., Sun, S., Mou, X. & Moran, M. A. Transporter genes expressed by coastal bacterioplankton in response to dissolved organic carbon. *Environ. Microbiol.* **12**, 616–627 (2010).
232. McCarren, J. et al. Microbial community transcriptomes reveal microbes and metabolic pathways associated with dissolved organic matter turnover in the sea. *Proc. Natl Acad. Sci. USA* **107**, 16420–16427 (2010).
233. Ziervogel, K. & Arnosti, C. Polysaccharide hydrolysis in aggregates and free enzyme activity in aggregate-free seawater from the north-eastern Gulf of Mexico. *Environ. Microbiol.* **10**, 289–299 (2008).
234. Landa, M. et al. Sulfur metabolites that facilitate oceanic phytoplankton–bacteria carbon flux. *ISME J.* **13**, 2536–2550 (2019).
235. Amin, S. A., Parker, M. S. & Armbrust, E. V. Interactions between diatoms and bacteria. *Microbiol. Mol. Biol. Rev.* **76**, 667–684 (2012).
236. Cude, W. N. et al. The production of the antimicrobial secondary metabolite indigoidine contributes to competitive surface colonization in the marine *Roseobacter Phaeobacter* sp. strain Y4I. *Appl. Environ. Microbiol.* **78**, 4771–4780 (2012).
237. Edwards, B. R., Bidle, K. D. & Van Mooy, B. A. Dose-dependent regulation of microbial activity on sinking particles by polyunsaturated aldehydes: implications for the carbon cycle. *Proc. Natl Acad. Sci. USA* **112**, 5909–5914 (2015).
238. Gram, L., Grossart, H.-P., Schlingloff, A. & Kjørboe, T. Possible quorum sensing in marine snow bacteria: production of acylated homoserine lactones by *Roseobacter* strains isolated from marine snow. *Appl. Environ. Microbiol.* **68**, 4111–4116 (2002).
239. Seyedsayamdost, M. R., Case, R. J., Kolter, R. & Clardy, J. The Jekyll-and-Hyde chemistry of *Phaeobacter gallaeciensis*. *Nat. Chem.* **3**, 331–335 (2011).
240. Teasdale, M. E., Liu, J., Wallace, J., Akhlaghi, F. & Rowley, D. C. Secondary metabolites produced by the marine bacterium *Halobacillus salinus* that inhibit quorum sensing-controlled phenotypes in gram-negative bacteria. *Appl. Environ. Microbiol.* **75**, 567–572 (2009).
241. Ross, A. C., Gulland, L. E., Dorrestein, P. C. & Moore, B. S. Targeted capture and heterologous expression of the *Pseudoalteromonas* alterochromide gene cluster in *Escherichia coli* represents a promising natural product exploratory platform. *ACS Synth. Biol.* **4**, 414–420 (2014).
242. Ziemert, N. et al. Diversity and evolution of secondary metabolism in the marine actinomycete genus *Salinispora*. *Proc. Natl Acad. Sci. USA* **111**, E1130–E1139 (2014).
243. Kiene, R. P. & Linn, L. J. Distribution and turnover of dissolved DMSP and its relationship with bacterial production and dimethylsulfide in the Gulf of Mexico. *Limnol. Oceanogr.* **45**, 849–861 (2000).

244. Andreae, M. O. & Crutzen, P. J. Atmospheric aerosols: biogeochemical sources and role in atmospheric chemistry. *Science* **276**, 1052–1058 (1997).
245. Charlson, R. J., Lovelock, J. E., Andreae, M. O. & Warren, S. G. Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Nature* **326**, 655–661 (1987).
246. Curson, A. R. et al. Dimethylsulfoniopropionate biosynthesis in marine bacteria and identification of the key gene in this process. *Nat. Microbiol.* **2**, 17009 (2017).
247. Howard, E. C. et al. Bacterial taxa that limit sulfur flux from the ocean. *Science* **314**, 649–652 (2006).
248. Todd, J., Curson, A., Dupont, C., Nicholson, P. & Johnston, A. The *dddP* gene, encoding a novel enzyme that converts dimethylsulfoniopropionate into dimethyl sulfide, is widespread in ocean metagenomes and marine bacteria and also occurs in some *Ascomycete* fungi. *Environ. Microbiol.* **11**, 1376–1385 (2009).
249. Todd, J. D. et al. Structural and regulatory genes required to make the gas dimethyl sulfide in bacteria. *Science* **315**, 666–669 (2007).
250. Vila-Costa, M. et al. Dimethylsulfoniopropionate uptake by marine phytoplankton. *Science* **314**, 652–654 (2006).
251. Evans, C. et al. The relative significance of viral lysis and microzooplankton grazing as pathways of dimethylsulfoniopropionate (DMSP) cleavage: an *Emiliana huxleyi* culture study. *Limnol. Oceanogr.* **52**, 1036–1045 (2007).
252. Bratbak, G. et al. Viral activity in relation to *Emiliana huxleyi* blooms: a mechanism of DMSP release? *Mar. Ecol. Prog. Ser.* **128**, 133–142 (1995).
253. Seymour, J. R., Simó, R., Ahmed, T. & Stocker, R. Chemoattraction to dimethylsulfoniopropionate throughout the marine microbial food web. *Science* **329**, 342–345 (2010).
254. Barak-Gavish, N. et al. Bacterial virulence against an oceanic bloom-forming phytoplankton is mediated by algal DMSP. *Sci. Adv.* **4**, eaau5716 (2018).
255. McParland, E. L. & Levine, N. M. The role of differential DMSP production and community composition in predicting variability of global surface DMSP concentrations. *Limnol. Oceanogr.* **64**, 757–773 (2019).
256. Varaljay, V. A. et al. Single-taxon field measurements of bacterial gene regulation controlling DMSP fate. *ISME J.* **9**, 1677–1686 (2015).

Acknowledgements

This Review arose from discussions supported by the CIFAR and developed further with support from the Moore Foundation (5503 to M.A.M., E.B.K. and A.S.E.; 9335 to S.A.A.), the Simons Foundation (542391 to M.A.M.; 253417 to E.B.K.; 504183 to E.M.B.; 509034SCFY20 to R.B.; 721225 to S.T.D.; 687269 to A.M.E.; 542389 to N.M.L.; 345889 to A.E.S.; and 735079 to A.V.), the US NSF (OCE-1756105 to N.R.B.; Major Research Equipment DBI-1946970 to A.S.E.; OCE-1924554 to M.A.S.; CAREER DEB-1749544 to A.S.; OCE-BSF 1635070 to D.S.; and OCE-1829831 to M.B.S.), the Paul M. Angell Family Foundation (to S.T.D.), and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2019-06159 to L.L.; RGPIN-2015-06078 to A.C.R.). This is NSF Center for Chemical Currencies of a Microbial Planet (C-CoMP) publication number 001.

Author contributions

All authors contributed ideas. M.A.M., E.B.K. and W.F.S. wrote the manuscript with substantial input from all authors.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence should be addressed to Mary Ann Moran or Elizabeth B. Kujawinski.

Peer review information *Nature Microbiology* thanks Pieter Dorrestein, Manuel Liebeke and Bethanie Edwards for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Springer Nature Limited 2022