

Self-duality protected multicriticality in deconfined quantum phase transitions

Da-Chuan Lu ,¹ Cenke Xu,² and Yi-Zhuang You¹

¹*Department of Physics, University of California, San Diego, California 92093, USA*

²*Department of Physics, University of California, Santa Barbara, California 93106, USA*



(Received 24 May 2021; revised 18 October 2021; accepted 18 November 2021; published 29 November 2021)

Duality places an important constraint on the renormalization group flows and the phase diagrams. For self-dual theories, the self-duality can be promoted as a symmetry, this leads to the multicriticalities. This work investigates a description of the deconfined quantum criticality, the $N_f = 2$ QED_3 , as an example of self-dual theories, and its multicritical behavior under perturbative deformations. The multicriticality is described by the theory with Gross-Neveu couplings and falls in a different universality class than the standard deconfined quantum criticality. We systematically calculate the scaling dimensions of various operators in the 3D quantum electrodynamics with the Chern-Simons term and Gross-Neveu couplings by the large- N renormalization group analysis. Specifically, we find certain Drelativistic four-fermion interactions which correspond to the dimer-dimer interactions in the lattice model will drive the deconfined quantum criticality to the first-order transition, this result is consistent with previous numerical studies.

DOI: [10.1103/PhysRevB.104.205142](https://doi.org/10.1103/PhysRevB.104.205142)

I. INTRODUCTION

Duality plays an important role in relating different phases of matter. One famous example is the Kramers-Wannier duality [1] in (1+1)D transverse field Ising model $H = \sum_i -JZ_iZ_{i+1} - hX_i$, which exchanges J and h and maps the ferromagnetic (Ising symmetry breaking) phase to the paramagnetic (Ising symmetric) phase and vice versa. More generally, two theories are dual to each other when they have different ultraviolet (UV) descriptions but flow to the same infrared (IR) theory. A well-known example in (2+1)D is the particle-vortex duality, which states that the XY model is dual to the Abelian Higgs model [2–4]. Recent developments further extend this understanding and discover many theories and their dual partners, altogether they form a web of duality [5].

If the theory remains the same under a duality, the duality will be called a self-duality. For example, the Kramers-Wannier duality is a self-duality for the (1+1)D Ising model at the critical point. Recent studies [6–9] further propose to interpret the self-duality as a categorical symmetry, making connections to the fusion category of anyon excitations in the corresponding bulk topological order in one higher dimension. When the self-duality is imposed as a symmetry, the system is enforced to stay on the phase boundary between the two duality-related phases, leading to the self-duality protected criticality and multicriticality [10–15]. For example, as illustrated in Fig. 1(a), in the presence of the Kramers-Wannier duality (enforcing $J = h$), a generic Ising chain (with all additional duality-allowed terms like $-K(X_iX_{i+1} + Z_{i-1}Z_{i+1})$) can either preserve the self-duality and remain gapless along the Ising critical line ($K < K_c$), or spontaneously break the self-duality and becomes gapped along the first-order transition line ($K > K_c$). The continuous and first-order

Ising transitions are separated by a multicritical point ($K = K_c$), i.e., the tricritical Ising point [16,17]. The multicritical point can be circumvented if the self-duality is explicitly broken (e.g., by $J \neq h$). In this sense, the multicriticality is protected by self-duality.

Similar continuous to first-order transition also happens in higher dimensions between the duality-related quantum phases. Here we will explore the (2+1)D example of self-duality-protected multicriticality. In particular, we will consider the topological transition between the bosonic symmetry protected topological (SPT) phase and the trivial phase, as illustrated in Fig. 1(b), where the two phases across the transition are related by the self-duality [5,18–21] of the quantum electrodynamics in (2+1)D (QED_3) with fermionic matters at flavor number $N_f = 2$. This theory also describes the deconfined quantum critical point (DQCP) [22–24] between the XY antiferromagnet (AFM) and the valence bond solid (VBS) in square-lattice quantum magnets with the easy-plane spin anisotropy, as shown in Fig. 1(c). In this case, the two phases are related by a \mathbb{Z}_2 subgroup of the emergent $\text{O}(4)$ symmetry that maps the two-component XY-AFM order parameter to the two-component VBS order parameter. Imposing these emergent symmetries (including the self-duality) essentially promotes the tuning parameters to the fluctuating scalar fields and prohibits the explicit mass terms. This leads to a unified field theory that describes the continuous-to-first-order transition in these systems. Such multicritical point lies in the universality class described by the QED_3 -Gross-Neveu [2,5,25] (QED_3 -GN) theory. More generally, the Chern-Simons (CS) term for the gauge fields can be included to describe the multicritical point of the exotic quantum phase transitions.

To further investigate the stability of this (2+1)D self-duality-protected multicriticality, we extend the field theory




FIG. 1. Quantum phases related by the duality or emergent symmetry: (a) ferromagnetic (ordered) and the paramagnetic (disordered) phases across the Ising transition are related by the Kramers-Wannier duality, (b) bosonic symmetry protected topological (SPT) and trivial phases are related by the fermionic particle-vortex duality, (c) XY antiferromagnetic (AFM) and valence bond solid (VBS) phases are related by an emergent \mathbb{Z}_2 symmetry. In all phase diagrams, the vertical axis is the relevant perturbation that drives the transition between the duality/symmetry-related phases, and the horizontal axis is always taken to be the square of the transition-driving perturbation. In phase diagrams (b) and (c), the existence of continuous transitions between the adjacent phases is assumed, which corresponds to the $N_f = 2$ QED₃ field theory without four-fermion interactions.

to the large fermion flavor number (large N_f) limit, and use the $1/N_f$ expansion [25–31] to analyze the renormalization group (RG) flow of the fermion quartic operators, including the mass-mass $(\bar{\psi} M \psi)^2$ and current-current $(\bar{\psi} \gamma^\mu M \psi)^2$ interactions, at the QED₃-GN fixed point. Our analysis indicates that the DQCP and the multicriticality can be driven to the first-order transition by current-current interactions. Such current-current interactions can be realized in the lattice spin model as a staggered dimer-dimer interaction (or stagger- Q) as proposed and observed in the recent quantum Monte Carlo (QMC) studies [32,33]. Unlike the conventional dimer-dimer interaction that couples the dimers along the vertical or horizontal directions on the square lattice, the stagger- Q interaction couples the dimers along the diagonal direction. The QMC results indicate that such a stagger- Q interaction may be responsible for driving the DQCP between continuous and first-order transitions (see Sec. IV B for more concrete discussion of the QMC results and our theoretical explanation).

The RG analysis can be further generalized to the QED₃-GN theory with additional Chern-Simons (CS) terms for the gauge field. Although there is a lack of known examples of self-dual theory with a non-zero-level CS term, a similar multicritical point separating the continuous and first-order transition still exists and can be analyzed. The result can be applied to the direct transition between bosonic fraction quantum Hall (FQH) and superfluid (SF) phases in interacting boson systems [34,35].

II. SELF-DUALITY OF $N_f = 2$ QED₃

The fermionic particle-vortex duality [36,37] dualizes a free Dirac fermion theory to $N_f = 1$ QED₃ theory with CS terms and the fermion operator is mapped to the fermion operator combined with gauge fluxes. Since CS terms break parity symmetry, the orientation reversed version of the fermionic particle-vortex duality is obtained by changing the sign of the CS terms. By combining the fermion particle-vortex duality

and its orientation reversed version, one can obtain a duality between two $N_f = 2$ QED₃ theories [18,19,21] described by the following Lagrangians:

$$\begin{aligned} & i\bar{\psi}_1 \not{D}_{a+X} \psi_1 + i\bar{\psi}_2 \not{D}_{a-X} \psi_2 + \frac{1}{4\pi} (a + Y) d(a + Y) \\ & + \frac{2}{4\pi} (X dX - Y dY) \end{aligned} \quad (1)$$

$$\iff i\bar{\chi}_1 \not{D}_{\tilde{a}+Y} \chi_1 + i\bar{\chi}_2 \not{D}_{\tilde{a}-Y} \chi_2 + \frac{1}{4\pi} (\tilde{a} + X) d(\tilde{a} + X), \quad (2)$$

where ψ_i, χ_i are fermion fields, $\not{D}_a \equiv \gamma^\mu (\partial_\mu - ia_\mu)$ is the Dirac operator coupled to the U(1) gauge field a . $ada \equiv \epsilon_{\mu\nu\rho} a_\mu \partial_\nu a_\rho$ is understood as the exterior product $a \wedge da$, and the same applies for other CS terms. We adopt the convention as the lower case letters a, \tilde{a} represent the dynamical U(1) gauge fields which will be integrated over in the path integral, and the upper case letters X, Y represent the background gauge fields which are used to keep track of the U(1)_X and U(1)_Y global symmetries.

The two theories (at least) have the common UV symmetry U(1)_X × U(1)_Y. For the U(1) gauge theories in (2+1)D, they automatically have an emergent global U(1)_M magnetic symmetry due to the Bianchi identity $\epsilon^{\mu\nu\lambda} \partial_\mu F_{\nu\lambda} = 0$ where $F_{\nu\lambda}$ is the gauge field strength. The charged operator of this U(1)_M symmetry is the magnetic monopole operator which creates the gauge flux and its coupling with the background gauge field are $\frac{1}{2\pi} adY, \frac{1}{2\pi} \tilde{a}dX$ in the both hand sides respectively. The symmetry charges of the operators are,

	U(1) _a	U(1) _X	U(1) _Y
\mathcal{M}_a	1	0	1
ψ_1	1	1	0
ψ_2	1	-1	0

	U(1) _a	U(1) _X	U(1) _Y
$\leftrightarrow \mathcal{M}_{\tilde{a}}$	1	1	0
χ_1	1	0	1
χ_2	1	0	-1

and the gauge invariant operators are built from these operators.

Renaming the fermion fields $\psi \leftrightarrow \chi$ will exchange $X \leftrightarrow Y$ and add a background term $\frac{1}{4\pi} (X dX - Y dY)$ to the Lagrangian, the left-hand side (LHS) Eq. (1) and the right-hand side (RHS) of Eq. (2) of the duality will be swapped, therefore, establishes the self-duality.

This self-duality can also be understood as exchanging the “electric charge” and the “magnetic charge.” On the LHS of the duality, the fermion field ψ_i is charged under the U(1)_X flavor symmetry, and the magnetic monopole operator \mathcal{M}_a which creates 2π flux for a is charged under the magnetic U(1)_Y due to the mixed CS term $\frac{1}{2\pi} adY$ (note that \mathcal{M}_a is the bare magnetic monopole operator which is not gauge invariant due to the CS term $\frac{1}{4\pi} ada$, the gauge-invariant operators are the combination of the \mathcal{M}_a and fermion creation operators). However, on the RHS, the fermion field χ_i is charged under U(1)_Y and the magnetic monopole operator $\mathcal{M}_{\tilde{a}}$ is charged under U(1)_X. This suggests that the fermion creation operators (resp. monopole operators) on the LHS become monopole operators (resp. fermion creation operators) on the RHS. More details of the self-duality are presented in Appendix B.

Here is a side-note on the conventions to regularize the fermion path integral. One convention is that integrating out a single Dirac fermion in (2+1)D will contribute a (-1) -level CS term for the negative fermion mass and a 0-level CS term for the positive fermion mass. Physically, fermions are doubled when putting on the lattice, one Dirac fermion is accompanied by a massive fermionic partner, otherwise, the single Dirac fermion will have parity anomaly in (2+1)D [38]. This convention assumes that the massive fermionic partner is *not* integrated out beforehand and it is more explicit on the quantization of the level of Chern-Simons term, this is easier to analyze the symmetry charges of the operators since the magnetic monopole operator has charge k if there is a level- k CS term. We will use this convention in discussing the dualities of quantum field theories, such as the self-duality of $N_f = 2$ QED₃.

Another convention is that integrating out the fermion will contribute a $\frac{\text{sgn}(m)}{2}$ -level CS term, this assumes that the massive fermionic partner has been integrated out beforehand and this is relevant to the analysis of the scaling dimensions of the critical theory since the massive fermionic partner does not involve in the transition. Using the later convention, half level CS term will involve in the massless theory, and now the Chern-Simons level is effectively $-\frac{N_f}{2} + k$ where N_f is the number of fermion flavors. We will adopt this convention in the discussion of renormalization group analysis on the critical behavior of the theory.

Schematically, the fermion theory with the level- k CS term using the first convention is related to that using the second convention by

$$\underbrace{i \sum_{i=1}^{N_f} \bar{\psi}_i \not{D}_a \psi_i + \frac{k}{4\pi} ad a}_{\text{the 1st convention}} \cong \underbrace{i \sum_{i=1}^{N_f} \bar{\psi}_i \not{D}_a \psi_i + \frac{k - N_f/2}{4\pi} ad a}_{\text{the 2nd convention}}. \quad (4)$$

The duality presented in Eqs. (1) and (2) will be equivalent to the self-dual theory presented in Ref. [18] by converting to the second convention of the fermion path integral regularization. However, both conventions have the same gauge-invariant operators and they yield the same response theories in the gapped phases.

A. Phase diagram

The $N_f = 2$ QED₃ has two relevant fermion mass deformations, the *singlet mass* $m \bar{\psi} \mathbb{1} \psi \equiv m(\bar{\psi}_1 \psi_1 + \bar{\psi}_2 \psi_2)$ and the *triplet mass* $m' \bar{\psi} \sigma^3 \psi \equiv m'(\bar{\psi}_1 \psi_1 - \bar{\psi}_2 \psi_2)$, where σ^i is the i th Pauli matrix. Under these mass deformations, one can integrate out the fermions and obtain the following effective theories for the background gauge fields Eq. (1) [18,20],

$$\begin{aligned} \frac{2}{4\pi} (X dX - Y dY) & \quad m > 0, m' = 0 \\ 0 & \quad m < 0, m' = 0, \end{aligned} \quad (5)$$

$$\begin{aligned} \frac{1}{2\pi} ad(Y + X) + \frac{1}{4e^2} f^2 + \dots & \quad m' > 0 m = 0 \\ \frac{1}{2\pi} ad(Y - X) + \frac{1}{4e^2} f^2 + \dots & \quad m' < 0 m = 0. \end{aligned} \quad (6)$$




FIG. 2. The phase diagram of $N_f = 2$ QED₃ theory. The singlet mass m drives the SPT transition between two symmetric phases, and the triplet mass m' drives AFM-VBS transition between two symmetry broken phases.

where e is the electron charge. The \dots represents the gapped degrees of freedom that are not important at low energy since the low-energy physics is dominated by the Maxwell term $\frac{1}{4e^2} f^2$ and the first term which describes the gapless Goldstone boson associated to the broken symmetry $U(1)_{Y+X}$ or $U(1)_{Y-X}$.

When the singlet mass m is nonzero, the two response theories in Eq. (5) differ by a $U(1)_{X,2} \times U(1)_{Y,-2}$ CS term, where the number indicates the level of the CS term, i.e., $\frac{2}{4\pi} (X dX - Y dY)$, which corresponds to the topological response of a bosonic SPT state with $U(1)_X \times U(1)_Y$ symmetry.¹ Therefore the $m > 0$ and $m < 0$ phases should be ascribed to the topological and trivial SPT phases, respectively.² When the triplet mass term m' is nonzero, the effective theories in Eq. (6) describe the Goldstone modes in the spontaneous symmetry breaking (SSB) phases with broken symmetries associated to $Y + X$ and $Y - X$, respectively (two different combinations of the generators of $U(1)_X, U(1)_Y$). In the context of square-lattice easy-plane quantum magnets [39,40], we might interpret $U(1)_{Y+X}$ as the in-plane spin rotation symmetry and $U(1)_{Y-X}$ as the lattice rotation symmetry (ignoring the discrete nature of the actual C_4 rotation), then the $m' > 0$ and $m' < 0$ phases could be identified as the XY-AFM and the VBS phases respectively. Figure 2(a) shows the phase diagram summarizing the above interpretations. Under the duality transformation, the singlet mass is odd ($m \rightarrow -m$) while the triplet mass is even ($m' \rightarrow m'$), which effectively swap the SPT and trivial phases but leaving the AFM and VBS phases unchanged (see Fig. 2). To restore the original phase diagram after the duality transformation, one should exchange $U(1)_X \leftrightarrow U(1)_Y$ and add a background $U(1)_{X,2} \times U(1)_{Y,-2}$ CS term to the Lagrangian.

B. Self-duality as a symmetry

As pointed out in Refs. [19,21], the explicit UV symmetry $U(1)_X \times U(1)_Y$ in Eq. (1) and (2) can be enhanced to

¹Since the gauge-invariant operators in UV are all bosonic (no single-fermion operators), the resulting gapped phases can possibly connect to the bosonic theory.

²Which phase is topological/trivial is only a matter of convention, as the notion of SPT phases is only relative.

the emergent symmetry $\frac{\text{SU}(2)_X \times \text{SU}(2)_Y}{\mathbb{Z}_2} \cong \text{SO}(4)$ in the IR. Together with the self-duality \mathbb{Z}_2^D which exchanges $\text{SU}(2)_X \leftrightarrow \text{SU}(2)_Y$ and attaches a $\text{SU}(2)_{X,1} \times \text{SU}(2)_{Y,-1}$ CS term (which falls back to the $\text{U}(1)_{X,2} \times \text{U}(1)_{Y,-2}$ CS term in the UV), the IR symmetry becomes $\text{SO}(4) \rtimes \mathbb{Z}_2^D \cong \text{O}(4)$. However, as the IR theory is shifted by the $\text{SU}(2)_{X,1} \times \text{SU}(2)_{Y,-1}$ background response under the self-duality transformation, the \mathbb{Z}_2^D and the $\text{SO}(4)$ have the mixed 't Hooft anomaly, thus they cannot be simultaneously coupled to the background gauge fields and promoted to the dynamical ones. Nonetheless, it can be viewed as the boundary of a (3+1)D SPT with the full $\text{O}(4)$ symmetry. With appropriate counterterm in the bulk, the whole system can also have time-reversal symmetry \mathbb{Z}_2^T , altogether gives $\text{O}(4) \times \mathbb{Z}_2^T$ as suggested in Ref. [41].

Note that the singlet mass m is invariant under $\text{SO}(4)$ but is odd under \mathbb{Z}_2^D , while the triplet mass m' explicitly breaks $\text{SO}(4)$ (as it is in the (3, 3) representation [20,41] of $\text{SU}(2)_X \times \text{SU}(2)_Y$) but is even under \mathbb{Z}_2^D . Hence, if both the emergent $\text{SO}(4)$ and the self-duality \mathbb{Z}_2^D symmetries are imposed, no fermion bilinear mass could be included in the Lagrangian.

C. Self-duality protected multicriticality

Although the mass term cannot be added to the Lagrangian, squares of the mass term still can, which may take the form of four-fermion interactions $(\bar{\psi} M^a \psi)^2$, where M^a 's are mass matrices acting on the flavor indices. Adding these mass-squared deformations to the QED theory Eq. (1) could potentially drive the theory to new fixed points [28]. The fate of the self-duality \mathbb{Z}_2^D and the $\text{SO}(4)$ symmetry depends on the RG flow of such mass-squared deformations. If both symmetries are preserved, the theory will remain critical (as no mass deformation is allowed), which describes the continuous transition between AFM and VBS phases (as well as the transition between SPT and trivial phases), which is also known as the $\text{O}(4)$ DQCP. When the self-duality \mathbb{Z}_2^D symmetry is spontaneously broken, the SPT transition becomes first-order. When the emergent $\text{SO}(4)$ symmetry [more specifically the \mathbb{Z}_2 subgroup that swaps $\text{U}(1)_{Y+X}$ and $\text{U}(1)_{Y-X}$] is spontaneously broken, the AFM-VBS transition becomes first-order. These first-order transitions are separated from the continuous transition by the multicritical points/lines. We will analyze the RG flow of the generic four-fermion interactions at these multicritical points, aiming to understand how certain kinds of interactions can drive the DQCP from a continuous transition to a first-order transition.

The multicritical point happens when Dirac fermion masses change the sign. To analyze the scaling dimensions of the operators at the multicritical point, we do not need to include the massive fermionic parton which is served to cancel the subtlety in the fermion path integral regularization. We rewrite Eq. (1) as

$$i\bar{\psi}_1 \not{\partial}_{a+X} \psi_1 + i\bar{\psi}_2 \not{\partial}_{a-X} \psi_2 + \frac{1}{2\pi} adY + \frac{1}{4\pi} (XdX - YdY). \quad (7)$$

The CS terms look different from Eq. (1), because we integrate out the massive fermionic partners beforehand and it corresponds to the second convention as discussed in the last




FIG. 3. Mean-field phase diagram of Eq. (8).

three paragraphs of Sec. II, following from Ref. [18]. Note that the changing of convention will not change the gauge invariant operators as well as the different gapped phases. The background gauge fields X and Y will not affect the dynamics and can be set to zero. Adding the mass-squared deformations amounts to promoting the mass terms m and m' to the dynamic scalar fields ϕ_1 and ϕ_2 , that are coupled to the fermions via Yukawa-type couplings $\phi_a \bar{\psi} M^a \psi$, this can also be seen by using the Hubbard-Stratonovich transformation. Together with their own boson mass terms $r_a \phi_a^2$, the action reads as

$$\begin{aligned} & \sum_{i=1}^2 i\bar{\psi}_i \not{\partial}_a \psi_i + \phi_1 \bar{\psi} \not{1} \psi + \phi_2 \bar{\psi} \sigma^3 \psi \\ & + \sum_{a=1}^2 \frac{1}{2g^2} \phi_a (r_a - \partial^2) \phi_a + \frac{\lambda}{4} (\phi_a \phi_a)^2. \end{aligned} \quad (8)$$

For each scalar field ϕ_a , the boson mass r_a has a corresponding critical value $r_{a,c}$. When $r_a \gg r_{a,c}$, the boson is gapped and $\langle \phi_a \rangle = 0$. When $r_a \ll r_{a,c}$, the boson is condensed, such that $\langle \phi_a \rangle \neq 0$ and the symmetry is spontaneously broken. This will dynamically generate the corresponding fermion mass terms. We may loosely set $r_{a,c} = 0$ and assume the bosons are critical when $r_a = 0$ in the following discussion.

The qualitative phase diagram of Eq. (8) is shown in Fig. 3, which can be considered as the extension of the Fig. 2's origin, since no fermion mass terms m, m' are added in the Eq. (8). In the phase diagram, when $r_1, r_2 \gg 0$ (the blue region), both bosons are gapped, leaving Eq. (8) to be the $N_f = 2$ QED₃ theory at low energy. As discussed previously, this theory has an emergent $\text{O}(4)$ symmetry and describes the continuous DQCP transition between the AFM and VBS phases [i.e., between the $\text{U}(1)_{Y+X}$ and $\text{U}(1)_{Y-X}$ SSB phases] when tuning the triplet fermion mass m' externally. If r_1 is at its critical value and $r_2 \gg 0$ (along the red line), the critical theory becomes $N_f = 2$ QED₃-Gross-Neveu model, which describes the continuous DQCP with emergent $\text{SO}(5)$ symmetry as proposed in Ref. [41]. If instead, r_2 is at its critical value

and $r_1 \gg 0$ (across the blue line), the theory describes the multicriticality between the O(4) DQCP and the first-order AFM-VBS transition. If both r_1 and r_2 are critical (the purple point), the theory describes the multicriticality between the SO(5) DQCP and the first-order AFM-VBS transition.

To see that the ϕ_2 condensed phase (the orange region) corresponds to the first-order AFM-VBS transition, we consider driving the AFM-VBS transition by an external triplet mass m' . The actual mass term seen by the fermion will be $(m' + \langle \phi_2 \rangle) \bar{\psi} \sigma^3 \psi$, meaning that the driving parameter m' needs to overcome the expectation value $\langle \phi_2 \rangle$ in order to change the sign of the triplet mass effectively and switch the system from one phase to another. Therefore $\langle \bar{\psi} \sigma^3 \psi \rangle$ will exhibit the hysteresis behavior as m' is tuned back and forth, which manifests the first-order transition. Without the external driving ($m' = 0$), the ground state will be degenerated between AFM and VBS phases.

On the other hand, the ϕ_1 condensed phase (the green region) is a symmetric gapped phase whose ground state is degenerated between topological and trivial SPT phases, which may as well be interpreted as the first-order SPT transition if the singlet mass m is tuned externally. The ϕ_1 condensed phase and the ϕ_2 condensed phase do not coexist, because they compete with each other to gap out the fermion, and the ground state is determined by the condensate that has a larger vacuum expectation value $|\langle \phi_a \rangle|$. When the competition reaches a balance at $|\langle \phi_1 \rangle| = |\langle \phi_2 \rangle|$ (along the gray dashed line), it triggers a direct transition between the symmetric and the SSB phases (either the in-plane magnetic order or the VBS order), which is of the 3D XY universality.

The multicriticality between the continuous and first-order transitions cannot be circumvented in the presence of the anomalous O(4) symmetry. However, it is possible that the protecting symmetry may be broken spontaneously under other potentially relevant perturbations, such that the O(4) DQCP is not stable in general. In the following, we will explore this possibility by analyzing the effect of generic four-fermion interactions in the QED-GN theory using the large- N renormalization group (RG) approach.

III. LARGE- N RENORMALIZATION GROUP ANALYSIS

A. QED-Gross-Neveu-Chern-Simons theory

We extend Eq. (8) to N_f flavors of Dirac fermions $\psi = (\psi_1, \dots, \psi_{N_f})^\top$ coupled to the dynamical U(1) gauge field, together with Yukawa-type couplings to N_b flavors of scalar bosons ϕ_a ($a = 1, \dots, N_b$). The bosons will have their kinetic terms and can be tuned critical by the r_a parameters. We also add the level- k CS term for the dynamical U(1) gauge field (to be general) and consider the QED₃-Gross-Neveu-Chern-Simons (QED-GN-CS) theory as follows:

$$\begin{aligned} \mathcal{L} = & \bar{\psi} (\mathbb{1}_{N_f} \otimes \gamma^\mu) (\partial_\mu - ia_\mu) \psi + \phi_a \bar{\psi} (M^a \otimes \mathbb{1}_2) \psi \\ & + \frac{1}{2g^2} \phi_a (r_a - \partial^2) \phi_a + \frac{\lambda}{4} (\phi_a \phi_a)^2 \\ & + \frac{ik}{4\pi} \epsilon^{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda + \frac{1}{4e^2} f_{\mu\nu} f^{\mu\nu}. \end{aligned} \quad (9)$$

Here, matrices $\mathbb{1}_{N_f}$, M^a act on the flavor space, while matrices $\mathbb{1}_2$, γ^μ act on the spinor space. We take the γ matrices to be

$(\sigma^3, \sigma^1, \sigma^2)$. M^a s are vertices of Yukawa couplings associated with fermion bilinear masses, which are assumed to be orthogonal to each other such that $\text{tr}(M^a M^b) = \mathbb{M} \delta_{ab}$. The last term is the Maxwell term, with the gauge curvature defined as $f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu$.

The multicritical points/lines in the phase diagram Fig. 3 correspond to tuning one or more scalar bosons to critical. We assume that all scalar fields in the effective theory (9) correspond to the critical bosons (other gapped bosons will be dropped from the effective theory automatically). The theory is tuned to the QED-GN-CS fixed point. The boson mass term $(r_a - r_{a,c}) \phi_a^2$ is a relevant perturbation that drives the system away from the multicriticality. It also is possible that some types of fermion interactions may flow to the boson mass term ϕ_a^2 , as it is equivalent to the mass-mass interaction $(\bar{\psi} (M^a \otimes \mathbb{1}_2) \psi)^2$ under the Hubbard-Stratonovich transform. Such fermion interactions will appear relevant at the QED-GN-CS fixed point and can drive the system away from multicriticality as well.

B. Renormalization of four-fermion interactions

To explore this possibility, we carry out a systematic study of the scaling dimension of four-fermion interactions at the QED-GN-CS fixed point (see Appendix A for technical details). We will follow the large- N_f expansion approach recently developed for the QED₃-GN model in Ref. [31], where the scaling dimensions of fermion and boson bilinear operators were analyzed. Here, we will carry over the analysis to four-fermion operators, which has not been presented yet. To be more general, we also include a CS term, such that our result could potentially be applied to other DQCP such as the superfluid to bosonic fractional quantum Hall transition (described by the QED-GN-CS fixed point at level $k = 1$ [34]).

In particular, our scheme to extend Eq. (8) to large N_f corresponds to generalizing the fermion flavor symmetry group from $\text{SU}(2) \rightarrow \text{SU}(2N)$, such that the fermion flavor number scales as $N_f = 2N$ with $N \rightarrow \infty$. The Yukawa vertices are generalized to

$$\{M^a\} = \{\mathbb{1}_2, \sigma^3\} \rightarrow \{M_N^a\} = \{\mathbb{1}_2, \sigma^3\} \otimes \mathbb{1}_N, \quad (10)$$

where $\{M^a\}$ denotes the set formed by M^a s, similar for $\{V^\alpha\}$. The perturbative interactions are

$$\mathcal{L}_{\text{int}} = u_{\alpha,m} (\bar{\psi} V^\alpha \otimes \mathbb{1}_2 \psi)^2 + u_{\alpha,\mu} (\bar{\psi} V^\alpha \otimes \gamma^\mu \psi)^2, \quad (11)$$

where $V^\alpha = \sigma^\alpha \otimes \mathbb{1}_N$ ($\alpha = 0, 1, 2$, and 3). $u_{\alpha,m}$, $u_{\alpha,\mu}$ represent the coupling coefficient of the mass-mass interactions and the current-current interactions respectively, which can be combined to a vector $u_{\alpha,i} = (u_{\alpha,m}, u_{\alpha,0}, u_{\alpha,1}, u_{\alpha,2})^\top$ in each α channel. The RG equations for $u_{\alpha,i}$ takes the following general form:

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2 N_f} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (12)$$

where the repeated indices are summed over and \mathbf{M} is a matrix with entries given by the $\mathcal{O}(1/N_f)$ corrections, the detailed calculations are presented in Appendix A. One can further diagonalize \mathbf{M} to find the eigenchannels. We take $N_f \rightarrow 2$ to restore the case of Eq. (8). The large- N_f analysis is not

well controlled for small N_f , as subleading corrections may not be sufficiently small. However, in our case, we assume the $N_f = 2$ QED₃ has the IR conformal fixed point which is suggested by the QMC simulation [39] and then perform the analysis on the perturbative four-fermion interactions. It turns out that our large- N_f RG results are consistent with the latest QMC simulation [32,33,42].

The first quadrant, O(4) DQCP. Without the contribution from the critical bosons, there is no relevant channel for $\alpha = 0$. However, for $\alpha = 1, 2$, and 3, it has one relevant channel,

$$\frac{du_{\alpha,i}}{d\ell} = 2.24u_{\alpha,i}, \quad \text{with } u_{\alpha,i} = (3, 1, 1, 1)^T, \quad (13)$$

and the spatiotemporal anisotropic channels are irrelevant. Therefore the mass-mass interaction can be generated from the current-current interaction under the RG flow, which could potentially drive the O(4) DQCP to a first-order transition (if the generated mass-squared interaction is strong enough to overcome the bare r_2 term).

With large- N_f , $u_{\alpha,i}$ are independent parameters. However, for $N_f = 2$ (i.e., $N = 1$), the Fierz identity demands the uniform combination $\sum_{\alpha=1, 2, \text{ and } 3} u_{\alpha,i}$ “fuses” into the $\alpha = 0$ channel, which is irrelevant. Additionally, the explicit $U(1)_X \times U(1)_Y$ symmetry guarantees $u_{1,i} = u_{2,i}$, hence for $N_f = 2$, there is only one independent channel of the relevant four-fermion interaction with $\alpha = 3$.

The positive- r_2 axis, SO(5) DQCP. In this case, the scalar boson associated to the singlet mass is critical, $\{M^a\} = \mathbb{1}_2$. There is still no relevant channel for $\alpha = 0$. For $\alpha = 1, 2$, and 3, it has the same relevant channel as the previous case,

$$\frac{du_{\alpha,i}}{d\ell} = 1.70u_{\alpha,i}, \quad \text{with } u_{\alpha,i} = (3, 1, 1, 1)^T. \quad (14)$$

Hence, the stagger-Q term still overlaps with the relevant channel at SO(5) DQCP fixed point. Similarly, as discussed in the last paragraph, for $N_f = 2$, there is only one independent channel of the relevant four-fermion interaction with $\alpha = 3$.

The positive- r_1 axis and the origin. Both cases are more involved. The positive- r_1 axis describes the transition between the O(4) DQCP and first-order transition, and the origin is a multicritical point where three critical lines joins. Both ϕ_1 and ϕ_2 scalar fields are critical at the origin, such that the Yukawa vertices are $\{M^a\} = \{\mathbb{1}_2, \sigma^3\}$. The eigenchannels will have mixture of V^0, V^3 or V^1, V^2 , because M^a will mix V^0 with V^3 as well as V^1 with V^2 . Considering $\{V^a\} = \{V^0, V^3\}$, there is one relevant channel with $u_{03} \equiv (u_{0,i}; u_{3,i}) = (-0.03, -0.071, -0.071, -0.071; 0.82, 0.32, 0.32, 0.32)^T$, and the RG equation reads

$$\begin{aligned} \frac{du_{03}}{d\ell} &= 1.89u_{03} \quad (\text{positive-}r_1 \text{ axis}), \\ \frac{du_{03}}{d\ell} &= 1.35u_{03} \quad (\text{origin}). \end{aligned} \quad (15)$$

The detailed calculation is presented in Appendix A. With one more critical boson at the origin compared to the positive- r_1 axis, the RG eigenvalue of the relevant interaction is smaller at the SO(5) multicritical point compared to the O(4) multicritical line.




FIG. 4. Extended phase diagram in the presence of relevant interaction u . The $u = 0$ plane corresponds to the phase diagram in Fig. 3.

IV. IMPLICATIONS OF RG ANALYSIS

A. Consequence of the relevant interactions

The RG analysis suggests that the SO(5) and O(4) DQCP may not be stable against the perturbation of certain Lorentz symmetry breaking four-fermion interactions in the field theory. The interaction is relevant and flows to the following form:

$$\mathcal{L}_{\text{int}} = u(3(\bar{\psi}\sigma^3\psi)^2 + (\bar{\psi}\sigma^3\gamma^\mu\psi)^2). \quad (16)$$

Depending on the sign of the coefficient u , the interaction may drive different instabilities of the QED theory. By analyzing all possible Wick decomposition of the interaction term, we found the leading eigen decompositions with both positive and negative interaction strength is $\mathcal{L}_{\text{int}} = u(\bar{\psi}\sigma^3\psi)^2 + \dots - u(\bar{\psi}\psi)^2$. Therefore, if $u < 0$, the interaction favors the condensation of the triplet mass term $\bar{\psi}\sigma^3\psi$, or equivalently the scalar field ϕ_2 that couples to it. In this case, the emergent SO(4) symmetry is spontaneously broken, and the AFM-VBS transition becomes first-order. On the other hand, if $u > 0$, the interaction favors the condensation of the singlet mass term $\bar{\psi}\psi$, or equivalently the corresponding scalar field ϕ_1 , which spontaneously breaks the self-duality and results in the symmetric gapped state. Figure 4 shows the extension of the phase diagram in the presence of four-fermion interaction.

The next leading eigen decompositions of the interaction are the singlet pairing channels $-\frac{2}{3}u|\psi^\dagger\sigma^2\gamma^0\gamma^x\psi|^2$ and $-\frac{2}{3}u|\psi^\dagger\sigma^2\gamma^0\gamma^y\psi|^2$ with slightly less interaction strength. When $u > 0$, the system may condense the Cooper pairs $\psi^\dagger\sigma^2\gamma^0\gamma^x\psi$, breaking the Lorentz symmetry. Since this term commutes with some of the kinetic terms in the Hamiltonian, it will split the Dirac points in the momentum space but will not gap out the fermions. It will also Higgs the U(1) gauge group down to \mathbb{Z}_2 . Therefore it opens the possibility for the gapless \mathbb{Z}_2 spin liquid phase instead of the symmetric gapped phase away from the multicritical point, which provides a candidate scenario for the phase diagram observed in the recent QMC study Ref. [42] where the first-order transition and the gapless \mathbb{Z}_2 spin liquid phase are separated by the multicritical point. Another scenario of the gapless \mathbb{Z}_2 spin liquid phase




FIG. 5. Illustration of the (standard) Q term (in blue) and the stagger- Q term (Q_s , in red) on the square lattice. Both are dimer-dimer interactions, but along different directions.

near the DQCP is recently proposed in Ref. [43]. The Lorentz symmetry is also broken by the Higgs field. However, the fermion flavors are doubled in that proposal compared to ours, thus it describes a different gapless \mathbb{Z}_2 spin liquid phase (see Appendix C for details). For example, the entanglement entropy contributed from the massless degrees of freedom will be different, which could be distinguished in future numerical studies.

B. Role of the stagger- Q perturbation

Recent QMC studies revealed the possibility of tuning the DQCP between continuous and first-order transitions [32,33]. In particular, the stagger- Q term (denoted by Q_s , or the so-called Z deformation) was proposed in Ref. [32] as a modification of the J - Q model,

$$H = H_{JQ} + H_{Q_s},$$

$$H_{JQ} = -J \sum_i P_i^x - Q \sum_i P_i^x P_{i+\hat{x}}^x + (x \leftrightarrow y), \quad (17)$$

$$H_{Q_s} = -Q_s \sum_i P_i^x P_{i+\hat{x}+\hat{y}}^x + (x \leftrightarrow y),$$

where $P_i^x = 1/4 - \mathbf{S}_i \cdot \mathbf{S}_{i+\hat{x}}$ and $P_i^y = 1/4 - \mathbf{S}_i \cdot \mathbf{S}_{i+\hat{y}}$ are the dimer operators on the x and y bonds respectively. The stagger- Q term Q_s favors a staggered VBS pattern, and hence the name. The illustration of the Q term and the stagger- Q term is shown in Fig. 5. Another version of the stagger- Q term that involves three dimers interacting along the diagonal direction is studied in Ref. [33]. The three-dimer stagger- Q term has the same symmetry as the two-dimer stagger- Q term, and shares the similar physical effect (both favors the same staggered VBS order). The QMC phase diagram in Ref. [33] explicitly shows that the stagger- Q term can drive the DQCP to a first-order transition. We will connect this observation to our field-theory analysis.

In the momentum space, the stagger- Q term should correspond to the dimer-dimer interaction near momentum (π, π) , which can be argued as follows. Let $P_{\mathbf{q}}^{x,y} = \sum_i P_i^{x,y} e^{-i\mathbf{q} \cdot \mathbf{r}_i}$ be the dimer operator of momentum \mathbf{q} . A large Q_s term favors the dimer to order in the staggered pattern (along the diagonal direction), which corresponds to the condensation of the dimer order parameter at momentum $\mathbf{q} = (\pi, \pi)$, i.e., $\langle P_{(\pi,\pi)}^x \rangle \neq 0$ or $\langle P_{(\pi,\pi)}^y \rangle \neq 0$. Therefore the effect of the stagger- Q interaction H_{Q_s} can be expressed as

$$H_{Q_s} \sim -Q_s ((P_{(\pi,\pi)}^x)^2 + (P_{(\pi,\pi)}^y)^2), \quad (18)$$

because a large Q_s in Eq. (18) also promotes the ordering of $P_{(\pi,\pi)}^{x,y}$, matching the effect of H_{Q_s} in the real space Eq. (17).

At low energy, the dimer fluctuation near momentum (π, π) should correspond to the spatial component of the Noether current associated with the emergent $U(1)_{Y-X}$ symmetry that rotates the VBS order parameters:

$$P_{(\pi,\pi)}^x \sim j_{VBS}^y, P_{(\pi,\pi)}^y \sim j_{VBS}^x. \quad (19)$$

This mapping was derived in Ref. [44] from the fermionic parton construction. A simple symmetry argument is as follows. We first notice that $P_{(0,0)}^x$ and $P_{(0,0)}^y$ are the VBS order parameters favored by the standard Q term in the J - Q model. They can be combined into a complex order parameter $\Psi_{VBS} = P_{(0,0)}^x + iP_{(0,0)}^y$. The $U(1)_{Y-X}$ rotation corresponds to $\Psi_{VBS} \rightarrow e^{i\theta} \Psi_{VBS}$, therefore the associated current operator should be

$$\begin{aligned} j_{VBS}^x &= i\Psi_{VBS}^\dagger \partial_x \Psi_{VBS} + \text{H.c.} \\ &= P_{(0,\pi)}^y \partial_x P_{(\pi,0)}^x - P_{(\pi,0)}^x \partial_x P_{(0,\pi)}^y, \\ j_{VBS}^y &= i\Psi_{VBS}^\dagger \partial_y \Psi_{VBS} + \text{H.c.} \\ &= P_{(0,\pi)}^y \partial_y P_{(\pi,0)}^x - P_{(\pi,0)}^x \partial_y P_{(0,\pi)}^y. \end{aligned} \quad (20)$$

Thus both j_{VBS}^x and j_{VBS}^y carry the total momentum (π, π) (as a summation of $(\pi, 0)$ and $(0, \pi)$). Under the (site-centered) reflection about the y axis, i.e., $(x, y) \rightarrow (-x, y)$, we have $(P^x, P^y) \rightarrow (-P^x, P^y)$, $(\partial_x, \partial_y) \rightarrow (-\partial_x, \partial_y)$, thus $(j_{VBS}^x, j_{VBS}^y) \rightarrow (j_{VBS}^x, -j_{VBS}^y)$ transforms as a pseudovector. Similarly, under the reflection $(x, y) \rightarrow (x, -y)$, we have $(j_{VBS}^x, j_{VBS}^y) \rightarrow (-j_{VBS}^x, j_{VBS}^y)$. Furthermore, $j_{VBS}^{x,y}$ does not transform under spin rotation symmetry. All these symmetry properties are precisely matched by Eq. (19), which speaks for its validity.

Using the operator correspondence in Eq. (19), Eq. (18) can be casted into

$$H_{Q_s} \sim -Q_s ((j_{VBS}^y)^2 + (j_{VBS}^x)^2), \quad (21)$$

which identifies the stagger- Q term to the current-current interaction in the spatial channel. We can make further connection to the field theory. Since the $U(1)_{Y-X}$ symmetry is generated by $\psi_1^\dagger \psi_1 - \psi_2^\dagger \psi_2$ in the $N_f = 2$ QED₃ theory, the corresponding Noether current should be $j_{VBS}^\mu = \bar{\psi} \sigma^3 \gamma^\mu \psi$, therefore, the current-current interaction in Eq. (21) further translates to the four-fermion interaction in Eq. (11) with $u_{3,i} \propto Q_s(0, 0, 1, 1)^T$. According to the RG analysis above, the current-current interaction will generate the mass-mass interaction and flow towards the combined interaction in Eq. (16).

Since the u term in Eq. (16) corresponds to the stagger- Q term in the lattice model, the original J - Q model may be very close to $u = 0$, i.e., the QED-GN fixed point in the field theory, though u should never be precisely zero. But the stagger- Q term in the lattice model will turn on a non-negligible u term in the field theory which is relevant at the QED-GN fixed point, therefore render the transition first order, as was observed numerically. In fact, according to Eq. (14), our calculation of the scaling dimension of the relevant four fermion term is $1.3 = 3 - 1.7$ at the SO(5) DQCP, which is close to the observed scaling dimension of the stagger- Q deformation of the J - Q model ($\Delta_Z \sim 1.4$ in Ref. [32]).




FIG. 6. Schematic phase diagram of the easy-plane J - Q model (22).

The above field theory understanding also applies to the easy-plane J - Q model [39,40],

$$H = H_{JQ} + H_\Delta, \\ H_\Delta = -J\Delta \sum_i S_i^z S_{i+\hat{x}}^z + (x \leftrightarrow y), \quad (22)$$

where the parameter Δ tunes the easy-plane anisotropy. $\Delta = 0$ is the $SU(2)$ isotropic limit, and $\Delta = 1$ is the $U(1) \rtimes \mathbb{Z}_2$ easy-plane limit.

Tuning Δ away from 0 breaks the spin $SU(2)$ symmetry and the u term should in principle also exist for the easy-plane J - Q model, but because it is more relevant compared to that in the $SU(2)$ symmetric case [according to Eqs. (13) and (14)], the easy-plane J - Q model may be a first-order transition more obviously than the isotropic limit. Based on the phase diagram Fig. 4, the system will either enter an intermediate symmetric gapped phase or exhibit a first-order AFM-VBS transition, in the presence of spin anisotropy. Given the physical meaning of the anisotropy term Δ , we can identify the symmetry gapped phase to the easy-axis anisotropy ($\Delta < 0$) and the first-order transition to the easy-plane anisotropy ($\Delta > 0$). A schematic phase diagram is presented in Fig. 6 for the lattice model Eq. (22). The symmetric gapped phase may as well be interpreted as the Ising ordered phase of $\langle S^z \rangle \neq 0$, since the condensation of ϕ_1 field corresponds to the ordering of $\langle S^z \rangle$. The scenario that the AFM-VBS transition becomes first-order as the easy-plane anisotropy is turned on is consistent with the recent QMC study Ref. [45].

V. SUMMARY

In this work, we studied the $N_f = 2$ QED_3 with self-duality. The $N_f = 2$ QED_3 has $SO(4)$ symmetry in the IR, if imposing the self-duality symmetry, it can be enhanced to $O(4)$. The singlet mass is invariant under $SO(4)$ but self-duality odd and the triplet mass is transformed by $SO(4)$ but self-duality even. Requiring the $O(4)$ symmetry, the theory cannot have explicit mass terms, which enables us to treat the mass terms as fluctuation scalar fields and to investigate the continuous-to-first-order transition driven by the mass fluctuations. The multicritical points (lines) separating the continuous and first-order transitions can be described by the QED -GN theory.

We further analyzed the stability of the theory under the four-fermion interactions. In particular, we focus on the spa-

tial current-current interaction of fermions in the field theory, which corresponds to a class of dimer-dimer interaction (the stagger- Q term) in the lattice spin model [32,33]. This operator has been shown to drive the continuous DQCP to a first-order transition in recent numerical works. Our analysis indicates that such dimer interaction can be relevant at the $O(4)$ DQCP and adjacent multicritical lines, which generally destabilize the continuous DQCP to first-order transitions (or intermediate gapped phases). Our finding provides a theoretical understanding of the numerically observed first-order transition driven by the dimer-dimer interaction. Our analysis also suggests a possibility to have \mathbb{Z}_2 spin liquid in this model [42].

We provide systematically large- N renormalization group calculation of the general $N_f = 2$ QED_3 with Gross-Neveu term in Appendix A. Thanks to viewing the Feynman diagrams as string diagrams of symmetry group representations [46], the complicated diagram at $\mathcal{O}(1/N_f)$ can be expressed by a few group parameters. Scaling dimensions of generic fermion/boson bilinear terms and four-fermion perturbations are presented. We expect these general results will find broader applications in other exotic quantum critical systems.

ACKNOWLEDGMENTS

We acknowledge the helpful discussion with John McGreevy and Anders Sandvik. D.-C.L. and Y.-Z.Y. are supported by a startup fund at UCSD. C.X. is supported by NSF Grant No. DMR-1920434, and the Simons Foundation.

APPENDIX A: LARGE- N RENORMALIZATION GROUP

The theory considered in the main text is the QED_3 with level- k Chern-Simons term and Yukawa coupling between the fermion bilinear terms and the scalar fields,

$$\mathcal{L} = \bar{\psi} (\mathbb{1}_{N_f} \otimes \gamma^\mu) (\partial_\mu - i a_\mu) \psi + \phi_a \bar{\psi} (M^a \otimes \mathbb{1}_2) \psi \\ + \frac{1}{2g^2} \phi_a (r_a - \partial^2) \phi_a + \frac{\lambda}{4} (\phi_a \phi_a)^2 \\ + \frac{ik}{4\pi} \epsilon^{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda + \frac{1}{4e^2} f_{\mu\nu} f^{\mu\nu}, \quad (A1)$$

where $\psi, \bar{\psi}$ represents N_f flavors of two-component Dirac fermion fields, $\mathbb{1}_N, M^a$ act on the N_f -dimensional flavor space while $\mathbb{1}_2, \gamma^\mu, \Gamma^{(m),I}$ act on the two-dimensional spinor space. $\Gamma^{(m),\mu_1, \dots, \mu_m}$ is defined as $\gamma^{[\mu_1} \dots \gamma^{\mu_m]}$ (antisymmetrize the indices) and any product of γ matrices can be reduced to this form. Since the spacetime dimension is 3, $\Gamma^{(i)}$ and $\Gamma^{(3-i)}$ are related by the Levi-Civita tensor. ϕ_a with $a = 1, \dots, N_b$ represent the scalar fields which are coupled to the fermion bilinears via a Yukawa type interaction. The last term in the Lagrangian is the Chern-Simons term with level k .

The bare propagators and vertices can be read off from the Lagrangian (A1),

$$\begin{aligned} \text{---} \rightarrow &= -i \frac{p_\mu (\mathbb{1}_{N_f} \otimes \gamma^\mu)}{p^2}, \quad a \text{ ---} b = D_{ab}^{(0)}(q) = \frac{g^2}{q^2} \delta_{a,b} \\ \mu \text{ } \text{---} \text{---} \nu &= \Pi_{\mu\nu}^{(0)}(q) = \frac{e^2}{q^2} \left(\frac{q^2 \delta_{\mu\nu} - q_\mu q_\nu + \frac{k}{2\pi} e^2 \epsilon_{\mu\nu\rho} q^\rho}{q^2 + (\frac{k}{2\pi})^2 e^4} + \xi \frac{q_\mu q_\nu}{q^2} \right), \end{aligned} \quad (\text{A2})$$

where ξ is the gauge parameter. The vertices are

$$\mu \text{ } \text{---} \text{---} \nu = i \mathbb{1}_{N_f} \otimes \gamma^\mu, \quad a \text{ ---} b = M^a \otimes \mathbb{1}_2. \quad (\text{A3})$$

The bare gauge and critical boson propagator will receive corrections, in the large- N limit, the corrections are dominated by fermion loops, for the gauge propagator,

$$\begin{aligned} \mu \text{ } \text{---} \text{---} \nu &= (-1)[i \mathbb{1}_{N_f} \otimes \gamma^\mu] \left[-i \frac{k_\rho (\mathbb{1}_{N_f} \otimes \gamma^\rho)}{k^2} \right] [i \mathbb{1}_{N_f} \otimes \gamma^\nu] \left[-i \frac{(k+q)_\sigma (\mathbb{1}_{N_f} \otimes \gamma^\sigma)}{(k+q)^2} \right] \\ &= (-1) \text{tr} [\mathbb{1}_{N_f} \otimes \gamma^\mu \gamma^\rho \gamma^\nu \gamma^\sigma] \int \frac{d^3 k}{(2\pi)^3} \frac{k_\rho (k+q)_\sigma}{k^2 (k+q)^2} \\ &= -\frac{N_f |q|}{16} \left(\delta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right). \end{aligned} \quad (\text{A4})$$

where N_f comes from trace over the identity matrix $\mathbb{1}_{N_f}$. Similar for the critical boson propagator,

$$\begin{aligned} a \text{ ---} b &= (-1)[M^a \otimes \mathbb{1}_2] \left[-i \frac{k_\rho (\mathbb{1}_{N_f} \otimes \gamma^\rho)}{k^2} \right] [M^b \otimes \mathbb{1}_2] \left[-i \frac{(k+q)_\sigma (\mathbb{1}_{N_f} \otimes \gamma^\sigma)}{(k+q)^2} \right] \\ &= \text{tr} [M^a M^b \otimes \gamma^\rho \gamma^\sigma] \int \frac{d^3 k}{(2\pi)^3} \frac{k_\rho (k+q)_\sigma}{k^2 (k+q)^2} \\ &= -\text{tr} [M^a M^b] \frac{|q|}{8} \equiv -\mathbb{M} \frac{|q|}{8} \delta_{ab}, \end{aligned} \quad (\text{A5})$$

where in the last step we define $\text{tr}[M^a M^b] = \mathbb{M} \delta_{ab}$, this is true when M^a is irreducible representation. The corrected propagator can be found by using Dyson's equation,

$$\Pi(q) = \{[\Pi^{(0)}(q)]^{-1} - \Sigma^{(0)}(q)\}^{-1}.$$

Note that in the large- N limit, this model flows to an interacting conformal field theory in the infrared limit, where the momentum scale q is much smaller than the coupling constants e and g , therefore the leading order of the dressed gauge and critical boson propagators are

$$\begin{aligned} \mu \text{ } \text{---} \text{---} \nu &= \mu \text{ } \text{---} \text{---} \nu + \mu \text{ } \text{---} \text{---} \nu + \mu \text{ } \text{---} \text{---} \nu + \mu \text{ } \text{---} \text{---} \nu \\ \Pi_{\mu\nu}(q) &\simeq \frac{\mathbb{A}}{N_f |q|} \left(\delta_{\mu\nu} - \xi \frac{q_\mu q_\nu}{q^2} \right) + \frac{\mathbb{B}}{N_f} \frac{\epsilon^{\mu\nu\rho} q_\rho}{q^2} + \mathcal{O}(|q|/e^2) \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} a \text{ ---} b &= a \text{ ---} b + a \text{ ---} b + a \text{ ---} b \\ D_{ab}(q) &\simeq \frac{8}{\mathbb{M} |q|} \delta_{ab} + \mathcal{O}(|q|/g^2) \equiv D(q) \delta_{ab}, \end{aligned} \quad (\text{A7})$$

where $\mathbb{A} = (16^{-1} + 16\kappa^2)^{-1}$, $\mathbb{B} = ((256\kappa)^{-1} + \kappa)^{-1}$, and $\kappa = k/(2\pi N_f)$, a simple check is when $k = 0$, $\mathbb{A} = 16$, $\mathbb{B} = 0$ match the coefficients in the large- N analysis of QED₃ theory. Note that κ is not inverse proportional to the 't Hooft coupling and can be any real number, the large- N limit is to take N_f , k to ∞ while keeping κ fixed. We also keep the gauge parameter ξ in the calculation and check that the final result does not depend on ξ .

1. Basic diagrams for $1/N$ corrections: self-energy

We extract the logarithmic divergences from the diagrams and using k and Λ to denote the external momentum and UV cutoff, respectively, the self-energy corrections are

$$\begin{aligned}
& \text{Diagram: } \text{A wavy line with a dot at the left end and an arrow pointing right.} \\
& = \int \frac{d^3 q}{(2\pi)^3} [\Pi_{\mu\nu}(q)] [\text{i} \mathbb{1}_{N_f} \otimes \gamma^\mu] \left[-\text{i} \frac{(k+q)_\sigma (\mathbb{1}_{N_f} \otimes \gamma^\sigma)}{(k+q)^2} \right] [\text{i} \mathbb{1}_{N_f} \otimes \gamma^\nu] \\
& = (\text{i} \mathbb{1}_{N_f} \otimes \gamma^\mu) k_\mu \frac{\mathbb{A}(1-3\xi)}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.}
\end{aligned} \tag{A8}$$

$$\begin{aligned}
& \text{Diagram: } \text{A horizontal line with a square vertex on the left, a dashed arc above it, and a square vertex on the right.} \\
& = \int \frac{d^3 q}{(2\pi)^3} [D(q) \delta_{ab}] [M^a \otimes \mathbb{1}_2] \left[-i \frac{(k+q)_\sigma (\mathbb{1}_{N_f} \otimes \gamma^\sigma)}{(k+q)^2} \right] [M^b \otimes \mathbb{1}_2] \\
& = (i \mathbb{1}_{N_f} \otimes \gamma^\mu) k_\mu \frac{8C_M}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.}
\end{aligned} \tag{A9}$$

where we define $M^a M^a = C_M \mathbb{1}_{N_f}$ in analogy of the Casimir.

2. Basic diagrams for $1/N$ corrections: vertex corrections

The four-fermion interactions in general can be added to the Lagrangian perturbatively, and assuming the small four-fermion perturbations will not drive the system to other fixed points. The general form for such interactions is

$$K_{(\alpha, (m_1), I), (\beta, (m_2), J)} \bar{\psi}(V^\alpha \otimes \Gamma^{(m_1), I}) \psi \bar{\psi}(V^\beta \otimes \Gamma^{(m_2), J}) \psi.$$

For simplicity and physical relevance, we will consider a subset of the four-fermion interactions with the form

$$\mathcal{L} \supset \mathcal{L}_{\text{int}} = u_{\alpha,(m),I} (\bar{\psi} (V^\alpha \otimes \Gamma^{(m),I}) \psi)^2. \quad (\text{A10})$$

We introduce the diagrams for the interaction vertices as

$$\begin{array}{c} \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \\ \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \\ \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \end{array} = u_{\alpha,(m),I}(V^\alpha \otimes \Gamma^{(m),I}), \quad \begin{array}{c} \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \\ \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \\ \text{Diagram: } \text{Two horizontal arrows pointing right, with a circle in the middle.} \\ \vdots \end{array} = K_{(\alpha,(m_1),I),(\beta,(m_2),J)} \begin{bmatrix} (V^\alpha \otimes \Gamma^{(m_1),I}) \\ \otimes \\ (V^\beta \otimes \Gamma^{(m_2),J}) \end{bmatrix}. \quad (\text{A11})$$


The vertex corrections are

$$\text{Diagram} = u_{\alpha,(m),I} (V^\alpha \otimes \Gamma^{(m),I}) \frac{\mathbb{A}(-3 - 2C_{\gamma,(m),I} + 3\xi)}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} \quad (\text{A12})$$

$$= u_{\alpha, (m), I} (V^\alpha \otimes \Gamma^{(m), I}) \frac{8C_{\gamma, (m), I} C_{M, \alpha}}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A13})$$

where we define $C_{M,\alpha}$, $C_{Y,(m),I}$ as $M^a V^\alpha M^a \equiv C_{M,\alpha} V^\alpha$ and $\gamma^\mu \Gamma^{(m),I} \gamma^\mu \equiv C_{Y,(m),I} \Gamma^{(m),I}$, repeated indices a, μ mean summation.

Note that there are also two-loop diagrams for the $1/N$ corrections, we begin with the calculation of the mass bubbles,



$$\begin{aligned} \Pi_{L,R}^{\text{Anticlockwise}} &= -i u_{\alpha,(m),I} \text{tr} [(V_L)(\mathbb{1}_{N_f} \otimes \gamma^a)(V_R)(V^\alpha \otimes (\gamma^b \Gamma^{(m),I} \gamma^c))] \\ &\quad \frac{1}{128q^3} (q_a q_b q_c - q^2 q_c \delta_{a,b} - q^2 q_b \delta_{a,c} + q^2 q_a \delta_{b,c}), \end{aligned} \quad (\text{A14})$$

$$\begin{aligned} \Pi_{L,R}^{\text{Clockwise}} &= i u_{\alpha,(m),I} \text{tr} [(V_L)(V^\alpha \otimes (\gamma^b \Gamma^{(m),I} \gamma^c))(V_R)(\mathbb{1}_{N_f} \otimes \gamma^a)] \\ &\quad \frac{1}{128q^3} (q_a q_b q_c - q^2 q_c \delta_{a,b} - q^2 q_b \delta_{a,c} + q^2 q_a \delta_{b,c}), \end{aligned} \quad (\text{A15})$$

where V_L, V_R stands for the vertex insertion, they could be gauge-gauge, boson-boson, or gauge-boson. The formula is complicated in general. For the three-dimensional theory, the γ matrices are simply the Pauli matrices and m in $\Gamma^{(m),I}$ is up to 3. Besides, $\Gamma^{(3),\{i_1,i_2,i_3\}} = i \epsilon^{i_1,i_2,i_3} \Gamma^{(0)}$, $\Gamma^{(2),\{i_1,i_2\}} = i \epsilon^{i_1,i_2} \Gamma^{(1),I}$.

a. The gauge-gauge insertion. Only $\Gamma^{(0)}, \Gamma^{(3)}$ will have nonzero contribution, as their relation is discussed previously, we can calculate $\Gamma^{(0)}$ and derive the result for $\Gamma^{(3)}$. The mass bubble result for $\Gamma^{(0)}$ is

$$\begin{aligned} \text{Diagram with shaded triangle} &= u_{\alpha,(0)} \text{tr}(V) \frac{q_l \epsilon^{lij}}{4q}. \end{aligned} \quad (\text{A16})$$

The two-loop diagrams give similar results for $\Gamma^{(0)}, \Gamma^{(3)}$,

$$\begin{aligned} \text{Diagram with shaded triangle} &= \begin{cases} u_{\alpha,(0)} (\mathbb{1}_{N_f} \otimes \Gamma^{(0)}) \frac{\text{tr}(V)}{N_f} \frac{(\mathbb{A}^2 - \mathbb{B}^2)}{4\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} & \text{for } m = 0 \\ u_{\alpha,(3),\{i_1,i_2,i_3\}} (\mathbb{1}_{N_f} \otimes \Gamma^{(3),\{i_1,i_2,i_3\}}) \frac{\text{tr}(V)}{N_f} \frac{(\mathbb{A}^2 - \mathbb{B}^2)}{4\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} & \text{for } m = 3. \end{cases} \end{aligned} \quad (\text{A17})$$

b. The boson-boson insertion. The nonzero contributions will occur only if $\text{tr}(M^i M^j V^\alpha) = -\text{tr}(M^i V^\alpha M^j)$, this requires nontrivial choices of the M^a, V^α . If so, the two-loop contributions are

$$\begin{aligned} \text{Diagram with shaded triangle} &= \begin{cases} -i u_{\alpha,(1),i_1} \text{tr}(M^i V^\alpha M^j) \frac{q_{i_1}}{4q} & \text{for } m = 1 \\ -i u_{\alpha,(2),\{i_1,i_2\}} \text{tr}(M^i V^\alpha M^j) \frac{i q_l \epsilon^{l,i_1,i_2}}{2q} & \text{for } m = 2. \end{cases} \end{aligned} \quad (\text{A18})$$

$$\begin{aligned} \text{Diagram with shaded triangle} &= \begin{cases} u_{\alpha,(1),i_1} (M^i M^j \otimes \Gamma^{(1),i_1}) \frac{-8 \text{tr}(M^i V^\alpha M^j)}{3\pi^2 \mathbb{M}^2} \ln(k/\Lambda) + \text{reg.} & \text{for } m = 1 \\ u_{\alpha,(2),\{i_1,i_2\}} (M^i M^j \otimes \Gamma^{(2),\{i_1,i_2\}}) \frac{-8 \text{tr}(M^i V^\alpha M^j)}{3\pi^2 \mathbb{M}^2} \ln(k/\Lambda) + \text{reg.} & \text{for } m = 2. \end{cases} \end{aligned} \quad (\text{A19})$$

For example, the boson 2-loop will contribute when $M^a = \{\mathbb{1}_2, \sigma^1, \sigma^2\}$ and $V^\alpha = \{\mathbb{1}_2, \sigma^1, \sigma^2, \sigma^3\}$ and it will only contribute to the current-current interaction.

c. Mixed gauge-boson insertion. The mixed gauge-boson insertion will vanish for all the choices of V^α and $\Gamma^{(m),I}$, part of the reason is because $\text{tr}(M^i V^\alpha) = \text{tr}(V^\alpha M^i)$ and it will never have a minus sign.


3. Basic diagrams for $1/N$ corrections: ladder corrections

The four-fermion interaction vertices as depicted in Eq. (A11) will receive $\mathcal{O}(1/N)$ correction from gauge and boson propagators as well,

$$\begin{aligned} &\text{Diagram with two horizontal lines and a wavy line} + \text{Diagram with two horizontal lines and a wavy line} \\ &= K_{(\alpha,(m_1),I),(\beta,(m_2),J)} \begin{bmatrix} V^\alpha \\ V^\beta \end{bmatrix} \otimes \left(\begin{bmatrix} \Gamma^{(m_1),I} \gamma^\mu \\ \Gamma^{(m_2),J} \gamma^\mu \end{bmatrix} + \begin{bmatrix} \Gamma^{(m_1),I} \gamma^\mu \\ \gamma^\mu \Gamma^{(m_2),J} \end{bmatrix} \right) \frac{-2\mathbb{A}}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} \end{aligned} \quad (\text{A20})$$

$$\Rightarrow u_{(\alpha,(m),I)} \begin{bmatrix} V^\alpha \\ V^\alpha \end{bmatrix} \otimes \left(\begin{bmatrix} \Gamma^{(m),I} \gamma^\mu \\ \Gamma^{(m),I} \gamma^\mu \end{bmatrix} + \begin{bmatrix} \Gamma^{(m),I} \gamma^\mu \\ \gamma^\mu \Gamma^{(m),I} \end{bmatrix} \right) \frac{-2\mathbb{A}}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} \quad (\text{A21})$$

where the last equation is the correction for the simplified four-fermion interaction as in Eq. (A10).



$$= K_{(\alpha, (m_1), I), (\beta, (m_2), J)} \left(\begin{bmatrix} V^\alpha M^a \\ \otimes \\ V^\beta M^a \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m_1), I} \gamma^\mu \\ \otimes \\ \Gamma^{(m_2), J} \gamma^\mu \end{bmatrix} - \begin{bmatrix} V^\alpha M^a \\ \otimes \\ M^a V^\beta \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m_1), I} \gamma^\mu \\ \otimes \\ \gamma^\mu \Gamma^{(m_2), J} \end{bmatrix} \right) \frac{-8}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A22})$$

$$= u_{(\alpha, (m), I)} \left(\begin{bmatrix} V^\alpha M^a \\ \otimes \\ V^\alpha M^a \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \Gamma^{(m), I} \gamma^\mu \end{bmatrix} - \begin{bmatrix} V^\alpha M^a \\ \otimes \\ M^a V^\alpha \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \gamma^\mu \Gamma^{(m), I} \end{bmatrix} \right) \frac{-8}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A23})$$

the repeated indices μ, a should be summed over.

4. Examples

Above general calculations will be concrete with certain assumptions.

(1) Since $\Gamma^{(i)}$ and $\Gamma^{(3-i)}$ are related by Levi-Civita symbol in three-dimension, we only need to consider $\Gamma^{(0)} = \mathbb{1}_2$ and $\Gamma^{(1), \mu} = \gamma^\mu$.

(2) For the physical relevance, we consider the four-fermion interactions in the form of $u_{\alpha, m}(\bar{\psi}(V^\alpha \otimes \mathbb{1}_2)\psi)^2$ and $u_{\alpha, \mu}(\bar{\psi}(V^\alpha \otimes \gamma^\mu)\psi)^2$ with $\mu = 0, 1$, and 2 .

(3) We further assume V^α, M^a are represented by Pauli matrices. This kind of interaction vertices arise when doing fermionic parton construction of the spin models, i.e., the spin operators correspond to the fermion bilinears with Pauli matrices inserted in the middle.

(4) We also view V^α 's as the basis of certain vector space and form a set $\{V^\alpha\}$ as well as M^a 's form a set $\{M^a\}$, such that $V^\alpha M^a \in \{V^\alpha\}$. For example, for $\{M^a\} = \{\mathbb{1}_2\}$ or $\{M^a\} = \{\mathbb{1}_2, \sigma^3\}$, $\{V^\alpha\}$ can be $\{V^\alpha\} = \{\mathbb{1}_2, \sigma^3\}$ or $\{V^\alpha\} = \{\mathbb{1}_2, \sigma^1, \sigma^2, \sigma^3\}$.

Since V^α, M^a can be represented by Pauli matrices as assumed, we can exploit the underlying algebraic structure of Pauli matrices. We further define the structure constants when multiplying the V^α, M^a as

$$V^\alpha M^b = \sum_\beta f_\beta^{\alpha b} V^\beta, \quad M^b V^\alpha = \sum_\beta f_\beta^{b \alpha} V^\beta, \quad (\text{A24})$$

$$u_{(\alpha, (m), I)} \left[\begin{bmatrix} V^\alpha \\ \otimes \\ V^\alpha \end{bmatrix} \otimes \left(\begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \Gamma^{(m), I} \gamma^\mu \end{bmatrix} + \begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \gamma^\mu \Gamma^{(m), I} \end{bmatrix} \right) \right] \frac{-2\mathbb{A}}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} \quad (\text{A28})$$

$$= u_{(\alpha, i)} \left[\begin{bmatrix} V^\alpha \\ \otimes \\ V^\alpha \end{bmatrix} \otimes \begin{bmatrix} \Gamma^j \\ \otimes \\ \Gamma^j \end{bmatrix} \sum_\mu (f_j^{i\mu} f_j^{i\mu} + f_j^{i\mu} f_j^{\mu i}) \right] \frac{-2\mathbb{A}}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.} \quad (\text{A29})$$

$$= u_{(\alpha, i)} \left[\begin{bmatrix} V^\alpha \\ \otimes \\ V^\alpha \end{bmatrix} \otimes \begin{bmatrix} \Gamma^j \\ \otimes \\ \Gamma^j \end{bmatrix} (\mathbb{1}_{\dim\{V^\alpha\}} \otimes (\mathbf{F}_j^i + \tilde{\mathbf{F}}_j^i)) \right] \frac{-2\mathbb{A}}{6\pi^2 N_f} \ln(k/\Lambda) + \text{reg.}, \quad (\text{A30})$$

where $\mathbf{F}_j^i \equiv \sum_{\mu=(3,1,2)} f_j^{i\mu} f_j^{i\mu}$, $\tilde{\mathbf{F}}_j^i \equiv \sum_{\mu=(3,1,2)} f_j^{i\mu} f_j^{\mu i}$, $i, j = 0, 3, 1, 2$ and Eq. (A22) can be simplified as

$$u_{(\alpha, (m), I)} \left(\begin{bmatrix} V^\alpha M^a \\ \otimes \\ V^\alpha M^a \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \Gamma^{(m), I} \gamma^\mu \end{bmatrix} - \begin{bmatrix} V^\alpha M^a \\ \otimes \\ M^a V^\alpha \end{bmatrix} \otimes \begin{bmatrix} \Gamma^{(m), I} \gamma^\mu \\ \otimes \\ \gamma^\mu \Gamma^{(m), I} \end{bmatrix} \right) \frac{-8}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A31})$$

$$= u_{(\alpha,i)} \begin{bmatrix} V^\beta \\ \otimes \\ V^\beta \end{bmatrix} \otimes \begin{bmatrix} \Gamma^j \\ \otimes \\ \Gamma^j \end{bmatrix} \left(\left(\sum_a f_\beta^{\alpha a} f_\beta^{\alpha a} \right) \left(\sum_\mu f_j^{i\mu} f_j^{i\mu} \right) - \left(\sum_a f_\beta^{\alpha a} f_\beta^{\alpha a} \right) \left(\sum_\mu f_j^{i\mu} f_j^{\mu i} \right) \right) \frac{-8}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A32})$$


$$= u_{(\alpha,i)} \begin{bmatrix} V^\beta \\ \otimes \\ V^\beta \end{bmatrix} \otimes \begin{bmatrix} \Gamma^j \\ \otimes \\ \Gamma^j \end{bmatrix} (\mathbf{F}_\beta^\alpha(\{M^a\}) \otimes \mathbf{F}_j^i - \tilde{\mathbf{F}}_\beta^\alpha(\{M^a\}) \otimes \tilde{\mathbf{F}}_j^i) \frac{-8}{6\pi^2 \mathbb{M}} \ln(k/\Lambda) + \text{reg.} \quad (\text{A33})$$

Both of the ladder contributions will depend on the structure constants with specific forms, and we define $\mathbf{F}_\beta^\alpha(\{M^a\}) \equiv \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a}$, $\tilde{\mathbf{F}}_\beta^\alpha(\{M^a\}) \equiv \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a}$, $\alpha, \beta \in \{V^\alpha\}$ similar to the above definition for the Γ matrices.

The self-energy corrections and the vertex corrections will be diagonal matrices acting on the vector $u_{\alpha,i}$. The self-energy corrections are the same for every $u_{\alpha,i}$, while the vertex corrections depend on the α, i . As listed in Table I, the coefficient $C_{\gamma,(0)} = 3, C_{\gamma,(1),\mu} = -1$ are distinct for mass-mass and current-current. The structure constants in Eq. (A25) also have these distinctions, this suggests the RG equations are in block forms.

5. Renormalization group equation for four-fermion interactions

The $1/N$ corrections for the four-fermion interaction vertices are



As discussed previously, for generic boson-fermion vertices, the ladder correction diagram of one interaction vertex will contribute to another interaction vertex, therefore, one need to include all the possible interaction vertices as the basis. For example, if $\{M^a\} = \{\mathbb{1}_2, \sigma^3\}$ and $\{V^\alpha\} = \{\mathbb{1}_2, \sigma^1\}$, then σ^2, σ^3 also need to be included in $\{V^\alpha\}$.

We will analyze the example in the main text Sec. III B in detail. Due to the reason provided in the previous paragraph, we choose the interaction vertex to be $\{V^\alpha\} = \{\mathbb{1}_2, \sigma^1, \sigma^2, \sigma^3\} \otimes \mathbb{1}_N$ and $\{M^a\} = \{\mathbb{1}_2, \sigma^3\} \otimes \mathbb{1}_N, \{\mathbb{1}_2, \sigma^3\} \otimes \mathbb{1}_N$. Combining with the Γ

TABLE I. The definition for the coefficients that are universal for chosen fermion-boson vertex and interaction matrix.

Notation	Definition
κ	$k/(2\pi N_f)$
\mathbb{A}	$\mathbb{A} = (16^{-1} + 16\kappa^2)^{-1}$
\mathbb{B}	$\mathbb{B} = ((256\kappa)^{-1} + \kappa)^{-1}$
\mathbb{M}	$\text{tr}[M^a M^b] = \mathbb{M} \delta_{ab}$
C_M	$M^a M^a = C_M \mathbb{1}_{N_f}$
$C_{M,\alpha}$	$M^a V^\alpha M^a \equiv C_{M,\alpha} V^\alpha$
$C_{\gamma,(m),I}$	$\gamma^\mu \Gamma^{(m),I} \gamma^\mu \equiv C_{\gamma,(m),I} \Gamma^{(m),I}$ $C_{\gamma,(0),I} = 3, C_{\gamma,(1),\mu} = -1$
f_c^{ab}	$\sigma^a \sigma^b = f_c^{ab} \sigma^c, \frac{1}{2} \text{tr}(\sigma^a \sigma^b \sigma^c)$
$\mathbf{F}_\beta^\alpha(\{M^a\}), \tilde{\mathbf{F}}_\beta^\alpha(\{M^a\})$	$\mathbf{F}_\beta^\alpha(\{M^a\}) = \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a},$ $\tilde{\mathbf{F}}_\beta^\alpha(\{M^a\}) = \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a}, \alpha, \beta \in \{V^\alpha\}$

matrices, the basis of the interaction vertices $u_{\alpha,i} = (u_{\alpha,m}, u_{\alpha,0}, u_{\alpha,1}, u_{\alpha,2})$ is $4 \times 4 = 16$ dimensional. The RG equation is organized as

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (\text{A35})$$

where α, β are the indices of the flavors, and i, j are the indices of the Γ matrices, $i = 0, 1, 2$, and 3 corresponds to $\{\mathbb{1}_2, \gamma^0, \gamma^1, \gamma^2\} = \{\mathbb{1}_2, \sigma^3, \sigma^1, \sigma^2\}$ in 3D.

The matrix $\mathbf{M}_{(\alpha,i),(\beta,j)}$ contains several parts:

$$\mathbf{M}_{(\alpha,i),(\beta,j)} = \mathbf{M}_{(\alpha,i),(\beta,j)}^{sv} + \mathbf{M}_{(\alpha,i),(\beta,j)}^{gL} + \mathbf{M}_{(\alpha,i),(\beta,j)}^{bL}. \quad (\text{A36})$$

The self-energy and vertex corrections are in the diagonal,

$$\mathbf{M}_{(\alpha,0),(\beta,0)}^{sv} = \left(\frac{16\mathbb{A}}{6\pi^2 N_f} + \frac{-2\text{tr}(V^\alpha)(\mathbb{A}^2 - \mathbb{B}^2)}{N_f} \frac{1}{4\pi^2 N_f} \right. \\ \left. + \frac{-16(C_M + 3C_{M,\alpha})}{6\pi^2 \mathbb{M}} \right) \mathbb{1}_{\alpha,\beta}, \quad (\text{A37})$$

$$\mathbf{M}_{(\alpha,i),(\beta,i)}^{sv} = \left(0 + \frac{-16(C_M - C_{M,\alpha})}{6\pi^2 \mathbb{M}} \right) \mathbb{1}_{\alpha,\beta}, \\ \text{with } i = 1, 2, \text{ and } 3. \quad (\text{A38})$$

The ladder correction from the gauge vertex contributes the off-diagonal part,

$$\mathbf{M}_{(\alpha,0),(\beta,i)}^{gL} = \mathbf{M}_{(\alpha,i),(\beta,0)}^{gL} = \frac{8\mathbb{A}}{6\pi^2 N_f} \mathbb{1}_{\alpha,\beta}, \\ \text{with } i = 1, 2, \text{ and } 3. \quad (\text{A39})$$

The ladder corrections from the boson vertex are complicated, in the (i, j) space, there are two parts,

$$\mathbf{M}_{(\alpha,0),(\beta,i)}^{bL} = \mathbf{M}_{(\alpha,i),(\beta,0)}^{bL} = \frac{16}{6\pi^2 \mathbb{M}} (\mathbf{F}_\beta^\alpha(\{M^a\}) - \tilde{\mathbf{F}}_\beta^\alpha(\{M^a\})),$$

with $i = 1, 2$, and 3 ; (A40)

$$\mathbf{M}_{(\alpha,i),(\beta,j)}^{bL} = \frac{16}{6\pi^2 \mathbb{M}} (-\mathbf{F}_\beta^\alpha(\{M^a\}) - \tilde{\mathbf{F}}_\beta^\alpha(\{M^a\})),$$

with $i, j = 1, 2$, and 3 , $i \neq j$; (A41)

where $\mathbf{F}_\beta^\alpha(\{M^a\}) = \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a}$, $\tilde{\mathbf{F}}_\beta^\alpha(\{M^a\}) = \sum_{a \in \{M^a\}} f_\beta^{\alpha a} f_\beta^{\alpha a}$, $\alpha, \beta \in \{V^\alpha\}$ is defined previously. This can be simplified if we take subset of $\{V^\alpha\}$ with proper $\{M^a\}$, and restrict the indices α and β in the subset.

The first quadrant, continuous O(4) DQCP. There is no critical boson in the system, $\{M^a\} = \{\}$. There is no mixture in the flavor space of the eigenchannel. For $V^a = \mathbb{1}_{2N}$,

$$\frac{du_{0,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(0,i),(0,j)} \right) u_{0,j},$$

$$\mathbf{M} = \frac{1}{256\kappa^2 + 1} \begin{pmatrix} \frac{4(512\kappa^2 - 1)}{256\kappa^2 + 1} & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \quad (A42)$$

In our case, $2N = 2$, $\kappa = 0$, the RG equation becomes

$$\frac{du_{0,i}}{d\ell} = \begin{pmatrix} -1 - \frac{128}{3\pi^2} & \frac{32}{3\pi^2} & \frac{32}{3\pi^2} & \frac{32}{3\pi^2} \\ \frac{32}{3\pi^2} & -1 & 0 & 0 \\ \frac{32}{3\pi^2} & 0 & -1 & 0 \\ \frac{32}{3\pi^2} & 0 & 0 & -1 \end{pmatrix} u_{0,j} \quad (A43)$$

and the eigenvalues of this matrix are all negative, meaning the perturbation is irrelevant among all the channels.

For $V^\alpha = \sigma^\alpha \otimes \mathbb{1}_N$, $\alpha = 1, 2$, and 3 , the RG equations are the same for different α s

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\alpha,j)} \right) u_{\alpha,j},$$

$$\mathbf{M} = \frac{1}{256\kappa^2 + 1} \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \quad (A44)$$

In our case, there is one relevant channel, and plug that into Eq. (A44), we get

$$u_{\alpha,i} = g_\alpha(3, 1, 1, 1)^T, \quad \frac{dg_\alpha}{d\ell} = 2.24g_\alpha. \quad (A45)$$

Follow the same procedure, we will present the RG equations and the relevant channel results for other cases.

The r_2 axis, continuous SO(5) DQCP. The boson corresponding to the singlet mass is critical, $\{M^a\} = \{\mathbb{1}_{2N}\}$. There is also no mixture in the flavor space. For $V^\alpha = \mathbb{1}_{2N}$,

$$\frac{du_{0,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(0,i),(0,j)} \right) u_{0,j},$$

$$\mathbf{M} = \begin{pmatrix} \frac{4(512\kappa^2 - 1)}{(256\kappa^2 + 1)^2} - \frac{1}{2} & \frac{1}{256\kappa^2 + 1} & \frac{1}{256\kappa^2 + 1} & \frac{1}{256\kappa^2 + 1} \\ \frac{1}{256\kappa^2 + 1} & 0 & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{256\kappa^2 + 1} & -\frac{1}{4} & 0 & -\frac{1}{4} \\ \frac{1}{256\kappa^2 + 1} & -\frac{1}{4} & -\frac{1}{4} & 0 \end{pmatrix}. \quad (A46)$$

There is no relevant channel in this case.

Again, for $V^\alpha = \sigma^\alpha \otimes \mathbb{1}_N$, $\alpha = 1, 2$, and 3 , the RG equations are the same for different α s

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\alpha,j)} \right) u_{\alpha,j},$$

$$\mathbf{M} = \begin{pmatrix} \frac{2}{256\kappa^2 + 1} - \frac{1}{2} & \frac{1}{256\kappa^2 + 1} & \frac{1}{256\kappa^2 + 1} & \frac{1}{256\kappa^2 + 1} \\ \frac{1}{256\kappa^2 + 1} & 0 & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{256\kappa^2 + 1} & -\frac{1}{4} & 0 & -\frac{1}{4} \\ \frac{1}{256\kappa^2 + 1} & -\frac{1}{4} & -\frac{1}{4} & 0 \end{pmatrix} \quad (A47)$$

and the relevant channel is the same as the case of the first quadrant, but with a smaller eigenvalue,

$$u_{\alpha,i} = g_\alpha(3, 1, 1, 1)^T, \quad \frac{dg_\alpha}{d\ell} = 1.70g_\alpha. \quad (A48)$$

The r_1 axis, transition between the O(4) DQCP and first-order transition. The boson corresponding to the triplet mass is critical, $\{M^a\} = \{\sigma^3 \otimes \mathbb{1}_N\}$. There are mixture between V^0, V^3 and also between V^1, V^2 , we will present the RG equation for V^0, V^3 and V^1, V^2 separately. For $\{V^0, V^3\}$,

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (A49)$$

$$\mathbf{M} = \begin{pmatrix} \left(\frac{4(512\kappa^2 - 1)}{(256\kappa^2 + 1)^2} - \frac{1}{2} \right) & 0 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 \\ 0 & \frac{2}{256\kappa^2 + 1} - \frac{1}{2} & 0_2 & -\frac{1}{4}\sigma^1 & -\frac{1}{4}\sigma^1 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & 0_2 & 0_2 & -\frac{1}{4}\sigma^1 & -\frac{1}{4}\sigma^1 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & -\frac{1}{4}\sigma^1 & 0_2 & 0_2 & -\frac{1}{4}\sigma^1 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & -\frac{1}{4}\sigma^1 & -\frac{1}{4}\sigma^1 & -\frac{1}{4}\sigma^1 & 0_2 \end{pmatrix}, \quad (A50)$$

where the 2×2 matrices act on the $\{V^0, V^3\}$ space, 0_2 is 2×2 matrix with all entries being 0. The only relevant channel is

$$u_{\alpha,i} = g_{(0,3)}((-0.03, 0.82), (-0.071, 0.32), (-0.071, 0.32), (-0.071, 0.32))^T, \quad \frac{dg_{(0,3)}}{d\ell} = 1.89g_{(0,3)}. \quad (\text{A51})$$

For $\{V^1, V^2\}$,

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (\text{A52})$$

$$\mathbf{M} = \begin{pmatrix} \left(\frac{2}{256\kappa^2+1} + \frac{1}{4}\right)\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & -\frac{1}{4}\mathbb{1}_2 & 0_2 & 0_2 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & 0_2 & -\frac{1}{4}\mathbb{1}_2 & 0_2 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & 0_2 & 0_2 & -\frac{1}{4}\mathbb{1}_2 \end{pmatrix}. \quad (\text{A53})$$

There are two relevant channels,

$$u_{\alpha,i} = g_{(1,2)}^{(1)}((-3, 3), (-1, 1), (-1, 1), (-1, 1))^T, \quad \frac{dg_{(1,2)}^{(1)}}{d\ell} = 2.78g_{(1,2)}^{(1)}, \quad (\text{A54})$$

$$u_{\alpha,i} = g_{(1,2)}^{(2)}((4.1, 4.1), (1, 1), (1, 1), (1, 1))^T, \quad \frac{dg_{(1,2)}^{(2)}}{d\ell} = 2.02g_{(1,2)}^{(2)}. \quad (\text{A55})$$

The first channel is antisymmetric combination of V^1, V^2 and the second is symmetric combination.

The origin, multicritical point. Both bosons are critical, the boson-fermion vertices are $\{M^a\} = \{\mathbb{1}_{2N}, \sigma^3 \otimes \mathbb{1}_N\}$. Again, there will be mixture between V^0, V^3 and also between V^1, V^2 . For $\{V^0, V^3\}$,

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (\text{A56})$$

$$\mathbf{M} = \begin{pmatrix} \left(\frac{4(512\kappa^2-1)}{(256\kappa^2+1)^2} - 1\right) & 0 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 \\ 0 & \frac{2}{256\kappa^2+1} - 1 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & 0_2 & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & 0_2 & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & 0_2 & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & -\frac{1}{4}(\mathbb{1}_2 + \sigma^1) & 0_2 & 0_2 \end{pmatrix}. \quad (\text{A57})$$

The relevant channel is the same as previous case with a smaller eigenvalue,

$$u_{\alpha,i} = g_{(0,3)}((-0.03, 0.82), (-0.071, 0.32), (-0.071, 0.32), (-0.071, 0.32))^T, \quad \frac{dg_{(0,3)}}{d\ell} = 1.35g_{(0,3)}. \quad (\text{A58})$$

For $\{V^1, V^2\}$,

$$\frac{du_{\alpha,i}}{d\ell} = \left(-1 + \frac{64}{3\pi^2(2N)} \mathbf{M}_{(\alpha,i),(\beta,j)} \right) u_{\beta,j}, \quad (\text{A59})$$

$$\mathbf{M} = \begin{pmatrix} \left(\frac{2}{256\kappa^2+1} - \frac{1}{4}\right)\mathbb{1}_2 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 \\ (256\kappa^2 + 1)^{-1}\mathbb{1}_2 - \frac{1}{4}\sigma^1 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 & -\frac{1}{4}\mathbb{1}_2 \end{pmatrix}. \quad (\text{A60})$$

There are two relevant channels,


$$u_{\alpha,i} = g_{(1,2)}^{(1)}((-3, 3), (-1, 1), (-1, 1), (-1, 1))^T, \quad \frac{dg_{(1,2)}^{(1)}}{d\ell} = 2.24g_{(1,2)}^{(1)}, \quad (\text{A61})$$

$$u_{\alpha,i} = g_{(1,2)}^{(2)}((4.1, 4.1), (1, 1), (1, 1), (1, 1))^T, \quad \frac{dg_{(1,2)}^{(2)}}{d\ell} = 1.49g_{(1,2)}^{(2)}. \quad (\text{A62})$$

The first relevant channel is the antisymmetric combination of V^1, V^2 , it is interesting that this relevant channel has the same scaling dimension as the relevant channel V^3 with $(3, 1, 1, 1)^T$ in the $\text{O}(4)$ DQCP (A45).

6. Mass scaling

Combining the diagrams in previous sections allows us to calculate the scaling dimension for the fermion mass term, which corresponds to the vertex $\bar{\psi}V^\alpha \otimes \Gamma^{(0)}\psi \equiv \bar{\psi}V^\alpha \otimes \mathbb{1}_2\psi$. As discussed in the main text, we use the $N_f = 2$ QED₃ description of DQCP, and consider its large- N generalization. The vertex of singlet mass is thus $V^\alpha = \mathbb{1}_{2N}$ and for the triplet mass is $V^\alpha = \sigma^3 \otimes \mathbb{1}_N$. The diagram equation for the corrections of the mass scaling dimension is



(A63)

For M^a being full rank, $\mathbb{M} = 2N$, C_M equals to the number of critical boson N_b . For singlet mass term, $C_{M,\alpha} = N_b$, but for the triplet mass, $C_{M,\alpha}$ depends on the choices of M^a , the coefficient is calculated explicitly by $C_{M,\alpha} = \text{tr}[(\sum_a M^a V^\alpha M^a) V^\alpha] / \text{tr}[V^\alpha V^\alpha]$. For boson associated to singlet mass, the result is simple, $M^a = \mathbb{1}$, with $a = 1, \dots, N_b$, $C_{M,\alpha} = N_b$.

For the mass scaling, $m = 0$, $C_{\gamma,(0),\{\}} = 3$, and there is no two-loop correction by critical boson. Collecting the logarithmic divergent part, we get

$$\Delta_{\bar{\psi}\mathbb{1}_{2N}\psi} = 2 - \frac{128(512\kappa^2 - 1)}{3\pi^2(2N)(1 + 256\kappa^2)^2} + \frac{16N_b}{3\pi^2(2N)}, \quad (\text{A64})$$

$$\Delta_{\bar{\psi}(\sigma^3 \otimes \mathbb{1}_N)\psi} = 2 - \frac{64}{3\pi^2(2N)(1 + 256\kappa^2)} + \frac{4(3C_{M,\alpha} + N_b)}{3\pi^2(2N)}, \quad (\text{A65})$$

where the last term in each equation comes from the critical boson contribution. This general result agrees with previous work with certain parameters.

7. Boson mass scaling

We can also calculate the scaling dimension of the boson operator ϕ_a^2 . Following Ref. [31], we define the scalar two-point function as $G_{ab}^\phi \equiv \langle \phi_a(p) \phi_b(-p) \rangle$, and its $\mathcal{O}(1/N)$ 1PI scalar self-energy contribution is represented by $\Sigma_{ab}^{\phi(1)}(p)$. From the Dyson's equation, the two-point function to $\mathcal{O}(1/N)$ is

$$\begin{aligned} G_{ab}^\phi &= D_{ab}(p) + D_{ac}(p) \Sigma_{cd}^{\phi(1)}(p) G_{db}^\phi(p) \\ &\simeq D_{ab}(p) + D_{ac}(p) \Sigma_{cd}^{\phi(1)}(p) D_{db}(p), \end{aligned} \quad (\text{A66})$$

where the self-energy is obtained by summing over the basic diagrams for fermion mass scaling but with nontrivial choices of M^a 's. Because of the coupling $\phi_a \bar{\psi} M^a \psi$, the self-energy corrections depend on M^a 's and can therefore change the scaling dimensions of the corresponding bosons. The self-energy has the following generic form:


$$\Sigma_{ab}^{\phi(1)}(p) = \delta_{ab} \frac{c_a |p|}{\pi^2 N} \ln \left(\frac{\Lambda^2}{p^2} \right). \quad (\text{A67})$$

For example, for $M^a = \mathbb{1}$, $c = \frac{2}{3} - \frac{16(512\kappa^2 - 1)}{3(256\kappa^2 + 1)^2} \xrightarrow{\kappa \rightarrow 0} 6$, and for M^a is traceless, $c = \frac{2}{3} - \frac{8}{3(256\kappa^2 + 1)} \xrightarrow{\kappa \rightarrow 0} -2$. The self-energy will contribute to the scaling dimension of the ϕ^2 in the following diagram:



(A68)

where the shaded bubble is the self-energy correction $\Sigma_{ab}^{\phi(1)}(p)$. There is one more diagram at $\mathcal{O}(1/N)$ will contribute to the scaling of ϕ^2 :




(A69)

the fermion “box” is the summation of fermions running clockwise and anticlockwise. The scaling dimension of ϕ^2 is combining Eqs. (A68) and (A69), this gives

$$\Delta_{\phi_a^2} = 2 - \frac{16c_a}{\pi^2 N} + \frac{8}{\pi^2 N}. \quad (\text{A70})$$

Note that the hourglass diagram [the first diagram in Eq. (A71)] will not contribute to the anomalous dimension, a simple argument is that similar diagram with one internal boson line appears in the self-energy correction (second and third diagram in Eq. (A71)) and it contributes to the anomalous dimension, while the hourglass diagram has two internal boson line, the power in the denominator is larger by 1, hence, it will not contribute to the anomalous dimension.



(A71)

With $\mathcal{O}(1/N)$ correction, the scaling dimension of the boson operator ϕ_a^2 are listed in the Table II. These scaling dimensions are not trustworthy for small fermion flavors N , but they show a trend for the scaling dimensions when having different boson-fermion vertices in large N .

TABLE II. The scaling dimensions of ϕ_a^2 with several choices of the boson-fermion vertices, these choices correspond to the axis and origin of the phase diagram Fig. 3 in the main text.

$\{M^a\}$	$\Delta_{\phi_a^2}$
$\{\mathbb{1}_2\}$	$2 - \frac{8}{3\pi^2 N} + \frac{256(512\kappa^2 - 1)}{3\pi^2(256\kappa^2 + 1)^2 N} \xrightarrow{\kappa \rightarrow 0} 2 - \frac{88}{\pi^2 N}$
$\{\sigma^3\}$	$2 - \frac{8}{3\pi^2 N} + \frac{128}{3\pi^2(256\kappa^2 + 1)N} \xrightarrow{\kappa \rightarrow 0} 2 + \frac{40}{\pi^2 N}$
$\{\mathbb{1}_2, \sigma^3\}$	$\{2 - \frac{40}{3\pi^2 N} + \frac{256(512\kappa^2 - 1)}{3\pi^2(256\kappa^2 + 1)^2 N}, 2 - \frac{40}{3\pi^2 N} + \frac{128}{3\pi^2(256\kappa^2 + 1)N}\} \xrightarrow{\kappa \rightarrow 0} \{2 - \frac{296}{3\pi^2 N}, 2 + \frac{88}{3\pi^2 N}\}$

APPENDIX B: DETAILS OF $N_f = 2$ QED₃ AND SELF-DUALITY

The single flavor fermion coupled to the U(1) gauge field is dual to free fermion theory and this is dubbed as fermion/fermion duality [5,19],

$$i\bar{\Psi}\not{D}_{A_1}\Psi \iff i\bar{\chi}\not{D}_{a_1}\chi - \frac{2}{4\pi}b_1db_1 + \frac{1}{2\pi}a_1db_1 + \frac{1}{2\pi}A_1db_1 - \frac{1}{4\pi}A_1dA_1 - 2CS_g, \quad (B1)$$

$$i\bar{\Psi}\not{D}_{A_2}\Psi \iff i\bar{\chi}\not{D}_{a_2}\chi + \frac{1}{4\pi}a_2da_2 + \frac{2}{4\pi}b_2db_2 - \frac{1}{2\pi}a_2db_2 - \frac{1}{2\pi}A_2db_2 + 2CS_g, \quad (B2)$$

where CS_g denotes the gravitational Chern-Simons term which will vanish in the flat space-times. The second line is the orientation reversed (time-reversal) version of the first one. We can then product them together on each side with the substitution $A_1 \rightarrow A, A_2 \rightarrow A - 2X$. Next, adding the counterterms $\frac{1}{2\pi}Ad(Y - X) + \frac{1}{4\pi}(XdX - YdY) + \frac{1}{4\pi}AdA + 2CS_g$ to both sides and gauging A , after integrating out most of the gauge fields, we get,

$$i\bar{\Psi}_1\not{D}_a\Psi_1 + i\bar{\Psi}_2\not{D}_{a-2X}\Psi_2 + \frac{1}{4\pi}ada + \frac{1}{2\pi}ad(Y - X) + \frac{1}{4\pi}(XdX - YdY) + 2CS_g \iff \quad (B3)$$

$$i\bar{\chi}_1\not{D}_{\tilde{a}-2Y}\chi_1 + i\bar{\chi}_2\not{D}_{\tilde{a}}\chi_2 + \frac{1}{4\pi}\tilde{a}d\tilde{a} + \frac{1}{2\pi}\tilde{a}d(X - Y) + \frac{1}{4\pi}(YdY - XdX) + 2CS_g, \quad (B4)$$

the self-duality exchanges $X \leftrightarrow Y$ and $\chi_i \leftrightarrow \Psi_i$. After relabeling the dynamical gauge fields a and \tilde{a} , it gives back Eqs. (1) and (2).

The self-duality exchanges the monopole symmetry and the Cartan subgroup of the flavor symmetry. It is also the duality between strong and weak couplings, this can be seen from the duality transformations amongst the derivation and their corresponding transformations in the (3+1)D bulk. Considering the (2+1)D U(1) gauge matter theories live at the boundary of (3+1)D U(1) gauge theory with the coupling

constant τ ,

$$I(A) = \frac{1}{8\pi} \int_X d^4x \sqrt{g} \left(\frac{2\pi}{e^2} F_{mn} F^{mn} + \frac{i\theta}{4\pi} \epsilon_{mnpq} F^{mn} F^{pq} \right) \quad (B5)$$

$$= \frac{i}{8\pi} \int_X d^4x \sqrt{g} (\bar{\tau} F_{mn}^+ F^{+mn} - \tau F_{mn}^- F^{-mn}),$$

$$\tau = \frac{\theta}{2\pi} + \frac{2\pi i}{e^2}, \quad (B6)$$

where g is the metric for the space-time, the theory and the transformation properties are well-defined also in the curved space-time. $F = dA$ and F is decomposed into self-dual and anti-self-dual pieces, $F_{mn}^\pm = \frac{1}{2}(F_{mn} \pm (\star F)_{mn})$ with $(\star F)_{mn} = \frac{1}{2}\epsilon_{mnpq} F^{pq}$, also $(\star F)_{mn}(\star F)^{mn} = F_{mn} F^{mn}$. The S transformation and T transformation act as

$$S : \quad \tau \rightarrow \tau' = -\frac{1}{\tau}, \quad \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

$$\int_{\partial X} J \cdot A \rightarrow \int_{\partial X} J \cdot a - \frac{1}{2\pi} adA'; \quad (B7)$$

$$-S : \quad \tau \rightarrow \tau' = -\frac{1}{\tau}, \quad \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

$$\int_{\partial X} J \cdot A \rightarrow \int_{\partial X} J \cdot a + \frac{1}{2\pi} adA'; \quad (B8)$$

$$T[k] : \quad \tau \rightarrow \tau' = \tau + k, \quad \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix},$$

$$\int_{\partial X} J \cdot A \rightarrow \int_{\partial X} J \cdot A - \frac{k}{4\pi} AdA. \quad (B9)$$

The $SL(2, \mathbb{Z})$ matrix acts on the coupling constant τ as

$$\tau \rightarrow \tau' = \frac{a\tau + b}{c\tau + d}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}). \quad (B10)$$

The fermion/fermion duality in the derivation of the $N_f = 2$ QED₃ self-duality is essential in connecting the left-hand-side and the right-hand-side (the other procedures, adding the counterterms and gauging the background gauge fields are the same for both hand sides). Using the above notation, the fermion/fermion duality and its orientation reversed version are

$$T[1] \circ (-S) \circ T[2] \circ (-S), \quad \tau \rightarrow \frac{1}{2} - \frac{1}{2(2\tau - 1)}, \quad (B11)$$

$$S \circ T[-2] \circ S \circ T[-1], \quad \tau \rightarrow \frac{1}{2} - \frac{1}{2(2\tau - 1)}. \quad (B12)$$

Take the coupling of the bulk theory $\tau = \frac{1}{2} + \frac{2\pi i}{e^2}$, under the duality $\tau \rightarrow \frac{1}{2} - \frac{1}{2(2\tau - 1)} = \frac{1}{2} + \frac{e^2 i}{8\pi}$. If $e \rightarrow 0$, which is the weak coupling limit, the dual theory has the strong coupling with $\tau \rightarrow \frac{1}{2} + 0i$. This suggests the fermion/fermion duality is a strong-weak duality, and similar calculation can be done for the $N_f = 2$ QED₃, which involves the $U(1) \times U(1)$ gauge theory in the bulk.

APPENDIX C: CONNECTION TO THE GAPLESS \mathbb{Z}_2 SPIN LIQUID IN REE [43]

1. Matrix form of fermion operators

The two flavor Nambu spinor can be written in matrix form,

$$\chi_i = \begin{pmatrix} f_{i\uparrow} & -f_{i\downarrow}^\dagger \\ f_{i\downarrow} & f_{i\uparrow}^\dagger \end{pmatrix}. \quad (C1)$$

The $SU(2)$ gauge symmetry and physical spin symmetry act as

$$SU(2)_g : \chi_i \rightarrow \chi_i U_{g,i}^\dagger, \quad (C2)$$

$$SU(2)_s : \chi_i \rightarrow U_{s,i} \chi_i. \quad (C3)$$

In majorana basis, one has

$$\chi_i = \frac{1}{\sqrt{2}}(\chi_0 + i\chi_a \sigma^a). \quad (C4)$$

Note that there is a discrepancy in the conventional notation and this, but it is merely relabeling,

$$\begin{aligned} \begin{pmatrix} f_{i\uparrow}^\dagger \\ f_{i\downarrow}^\dagger \\ f_{i\downarrow} \\ f_{i\uparrow} \end{pmatrix} &= \begin{pmatrix} 1 & i & 0 & 0 \\ 1 & -i & 0 & 0 \\ 0 & 0 & 1 & i \\ 0 & 0 & 1 & -i \end{pmatrix} \begin{pmatrix} \chi_{1,1} \\ \chi_{1,2} \\ \chi_{2,1} \\ \chi_{2,2} \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 & i \\ 1 & 0 & 0 & -i \\ 0 & i & -1 & 0 \\ 0 & -i & -1 & 0 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \\ \chi_2 \\ \chi_3 \end{pmatrix} \end{aligned} \quad (C5)$$

the relabeling is

$$\begin{pmatrix} \chi_0 \\ \chi_1 \\ \chi_2 \\ \chi_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} \chi_{1,1} \\ \chi_{1,2} \\ \chi_{2,1} \\ \chi_{2,2} \end{pmatrix}. \quad (C6)$$

$$\chi_a \rightarrow \chi_{a_1, a_2}.$$

2. Hamiltonian and Higgs fields

Define the 4×2 matrix operator,

$$X_{\alpha, v; \beta} = \frac{1}{\sqrt{2}}(\chi_{0,v} \mathbb{1}_{\alpha\beta} + i\chi_{a,v} \sigma_{\alpha\beta}^a), \quad (C7)$$

$$X_v = \frac{1}{\sqrt{2}}(\chi_{0,v} \sigma^0 + i\chi_{a,v} \sigma^a) \quad (C8)$$

with $\chi_{a,v}$, $a = 0 \sim 3$, $v = 1, 2$. The γ matrices act on the spinor index m in $\chi_{m,a,v}$, and it is left implicit. The mean-field Lagrangian is

$$\begin{aligned} \mathcal{L} &= i\text{Tr}\{\bar{X} \gamma^\mu \partial_\mu X\} \\ &= i\text{Tr}\{(\chi_{0,v}^T \gamma^0 \sigma^0 - i\chi_{a,v}^T \gamma^0 \sigma^a) \gamma^\mu \partial_\mu (\chi_{0,v} \sigma^0 + i\chi_{b,v} \sigma^b)\} \\ &= \sum_{a,v} i\chi_{a,v}^T \gamma^0 \gamma^\mu \partial_\mu \chi_{a,v}, \end{aligned}$$

where $\gamma^\mu = \{\sigma^2, \sigma^3, \sigma^1\}$ and $\bar{X} = X^\dagger \gamma^0$.

Let's now proceed to translate the Lagrangian for the Higgs fields in Ref. [43], the matrix μ^i acts on the v indices, one of

the \mathbb{Z}_2 Higgs field is

$$\begin{aligned} &\Phi_1^a \text{Tr}\{\sigma^a \bar{X} \mu^z \gamma^x X\} \\ &= \Phi_1^c \text{Tr}\left\{\sigma^c (\chi_{0,v}^T \gamma^0 \sigma^0 - i\chi_{a,v}^T \gamma^0 \sigma^a) \mu_{v,w}^z \gamma^x (\chi_{0,w} \sigma^0 \right. \\ &\quad \left. + i\chi_{b,w} \sigma^b)\right\} \\ &= \Phi_1^c i(\chi_{0,v}^T \gamma^0 \gamma^x \mu_{v,w}^z \chi_{c,w} - \chi_{c,v}^T \gamma^0 \gamma^x \mu_{v,w}^z \chi_{0,w}), \end{aligned} \quad (C9)$$

one can also get the matrices that act on the index a ,

$$c = 1, \delta_{i,0} \delta_{j,1} - \delta_{i,1} \delta_{j,0} = -i(\sigma^{02} + \sigma^{32}), \quad (C10)$$

$$c = 2, \delta_{i,0} \delta_{j,1} - \delta_{i,1} \delta_{j,0} = -i(\sigma^{20} + \sigma^{23}), \quad (C11)$$

$$c = 3, \delta_{i,0} \delta_{j,1} - \delta_{i,1} \delta_{j,0} = -i(\sigma^{12} + \sigma^{21}). \quad (C12)$$

To compare with our model, we need to change the basis following Eq. (C6),

$$c = 1, -iM^1 = -i(\sigma^{12} + \sigma^{21}), \quad (C13)$$

$$c = 2, -iM^2 = -i(\sigma^{20} + \sigma^{23}), \quad (C14)$$

$$c = 3, -iM^3 = -i(\sigma^{02} + \sigma^{32}), \quad (C15)$$

and using the basis χ_{m,v,a_1,a_2} , therefore, the Higgs field becomes

$$\Phi_1^c \chi^T [(\gamma^0 \gamma^x) \otimes \mu^z \otimes M^c] \chi. \quad (C16)$$

The other \mathbb{Z}_2 Higgs field is

$$\Phi_2^c \chi^T [(\gamma^0 \gamma^y) \otimes \mu^x \otimes M^c] \chi \quad (C17)$$

and the $U(1)$ Higgs field is

$$\Phi_3^c \chi^T [\gamma^0 (\gamma^y k_x + \gamma^x k_y) \otimes \mu^y \otimes M^c] \chi. \quad (C18)$$

3. The Higgs configuration

Reference [43] proposes the staggered flux state is obtained when $\langle \Phi_3 \rangle \propto (0, 0, \delta\phi)$ and the $Z2A_{ZZ13}$ state follows from $\langle \Phi_1 \rangle \propto (\gamma_1 - \gamma_2, \gamma_1 + \gamma_2, 0)$ and $\langle \Phi_2 \rangle \propto (-\gamma_1 - \gamma_2, \gamma_1 - \gamma_2, 0)$. Recall that, $\gamma^\mu = \{\sigma^2, \sigma^3, \sigma^1\}$ and

$$c = 1, -iM^1 = -i(\sigma^{12} + \sigma^{21}), \quad (C19)$$

$$c = 2, -iM^2 = -i(\sigma^{20} + \sigma^{23}), \quad (C20)$$

$$c = 3, -iM^3 = -i(\sigma^{02} + \sigma^{32}). \quad (C21)$$

When condensing the Higgs fields, it corresponds to generate the mass for the combination of the fermion bilinears,

$$\Phi_1^{1,2} : \sigma^{1312} + \sigma^{1321}, \sigma^{1320} + \sigma^{1323}, \quad (C22)$$

$$\Phi_2^{1,2} : \sigma^{3112} + \sigma^{3121}, \sigma^{3120} + \sigma^{3123}, \quad (C23)$$

$$\Phi_3^3 : \sigma^{1202} k_y + \sigma^{1232} k_y, \sigma^{3202} k_x + \sigma^{3232} k_x, \quad (C24)$$

and the kinetic terms are

$$\sigma^{1000} k_x, \sigma^{3000} k_y. \quad (C25)$$

The only Pauli matrix commutes with the above matrices is σ^{0230} , which is also the symmetry generator.

a. Our model. In our model, the kinetic terms are

$$\sigma^{100}k_x, \sigma^{300}k_y \quad (C26)$$

and the pairing terms are

$$\sigma^{323}, \sigma^{321}, \sigma^{123}, \sigma^{121} \quad (C27)$$

and the Pauli matrices that commute with the above are

$$\sigma^{012}, \sigma^{020}, \sigma^{032}. \quad (C28)$$

4. Basis rotation

We can match both theories by examining their symmetry generators. The only matrix σ^{0230} that commutes with other matrices in Ref. [43] can be rotated to

$$\sigma^{0012}, \text{ by } e^{i\frac{\pi}{4}\sigma^{0222}}; \quad (C29)$$

$$\sigma^{0020}, \text{ by } e^{i\frac{\pi}{4}\sigma^{0210}}; \quad (C30)$$

$$\sigma^{0332}, \text{ by } e^{i\frac{\pi}{4}\sigma^{0102}}; \quad (C31)$$

where the rotation is generated by $\sigma^I \rightarrow e^{-i\frac{\pi}{4}\sigma^I} \sigma^I e^{i\frac{\pi}{4}\sigma^I}$. The \mathbb{Z}_2 Higgs fields in Eq. (C22) will be rotated to

$$\begin{aligned} \Phi_1^{1,2} &: \sigma^{1312} + \sigma^{1321}, \sigma^{1102} + \sigma^{1323}; \\ \Phi_2^{1,2} &: \sigma^{3112} + \sigma^{3121}, -\sigma^{3302} + \sigma^{3123} \text{ by } e^{i\frac{\pi}{4}\sigma^{0222}}; \end{aligned} \quad (C32)$$

$$\Phi_1^{1,2} : \sigma^{1102} + \sigma^{1321}, \sigma^{1320} + \sigma^{1323};$$

$$\Phi_2^{1,2} : -\sigma^{3302} + \sigma^{3121}, \sigma^{3120} + \sigma^{3123} \text{ by } e^{i\frac{\pi}{4}\sigma^{0210}}; \quad (C33)$$

$$\Phi_1^{1,2} : -\sigma^{1210} + \sigma^{1321}, -\sigma^{1222} + \sigma^{1323};$$

$$\Phi_2^{1,2} : \sigma^{3112} - \sigma^{3023}, \sigma^{3120} + \sigma^{3021} \text{ by } e^{i\frac{\pi}{4}\sigma^{0102}}. \quad (C34)$$

If one takes the second index as labeling the original theory and the dual theory in our model, some terms of the \mathbb{Z}_2 Higgs fields in Ref. [43] correspond to the pairing in the form of $\sigma^{121}, \sigma^{123}, \sigma^{321}, \sigma^{323}$ that appear in both the original theory and the dual theory according to the Eq. (C34). For example,

$$\begin{aligned} \cdots + \chi^T \sigma^{1321} \chi &= \cdots + \chi_1^T \sigma^{121} \chi_1 - \chi_2^T \sigma^{121} \chi_2 \\ &\sim \cdots + \psi^\dagger \sigma^2 \gamma^0 \gamma^x \psi - \tilde{\psi}^\dagger \sigma^2 \gamma^0 \gamma^x \tilde{\psi}, \end{aligned} \quad (C35)$$

$$\begin{aligned} \cdots + \chi^T \sigma^{3021} \chi &= \cdots + \chi_1^T \sigma^{321} \chi_1 + \chi_2^T \sigma^{321} \chi_2 \\ &\sim \cdots + \psi^\dagger \sigma^2 \gamma^0 \gamma^y \psi + \tilde{\psi}^\dagger \sigma^2 \gamma^0 \gamma^y \tilde{\psi}, \end{aligned} \quad (C36)$$

where ψ is the original fermion and $\tilde{\psi}$ is the dual fermion, they are corresponding to the pairing fermion bilinears that appear in the \mathbb{Z}_2 Higgs fields (C34). However, the dual fermion pairings are not explicit in the self-dual $N_f = 2$ QED₃ theory and the linear combinations with another fermion bilinears are crucial to obtain the \mathbb{Z}_2 Higgs fields in Ref. [43], for example, $\Phi_1^1 \chi^T (-\sigma^{1210} + \sigma^{1321}) \chi$ in the first line of Eq. (C34).

-
- [1] H. A. Kramers and G. H. Wannier, *Phys. Rev.* **60**, 252 (1941).
[2] A. Karch and D. Tong, *Phys. Rev. X* **6**, 031043 (2016).
[3] M. E. Peskin, *Ann. Phys.* **113**, 122 (1978).
[4] C. Dasgupta and B. I. Halperin, *Phys. Rev. Lett.* **47**, 1556 (1981).
[5] N. Seiberg, T. Senthil, C. Wang, and E. Witten, *Ann. Phys.* **374**, 395 (2016).
[6] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, *J. High Energy Phys.* **01** (2019) 026.
[7] W. Ji and X.-G. Wen, *Phys. Rev. Research* **2**, 033417 (2020).
[8] R. Thorngren and Y. Wang, [arXiv:1912.02817](https://arxiv.org/abs/1912.02817) [hep-th].
[9] T. Lichtenman, R. Thorngren, N. H. Lindner, A. Stern, and E. Berg, *Phys. Rev. B* **104**, 075141 (2021).
[10] D. Aasen, R. S. K. Mong, and P. Fendley, *J. Phys. A: Math. Gen.* **49**, 354001 (2016).
[11] M. Buican and A. Gromov, *Commun. Math. Phys.* **356**, 1017 (2017).
[12] M. Bal, D. J. Williamson, R. Vanhove, N. Bultinck, J. Haegeman, and F. Verstraete, *Phys. Rev. Lett.* **121**, 177203 (2018).
[13] W. Ji and X.-G. Wen, *Phys. Rev. Research* **1**, 033054 (2019).
[14] L. Kong and H. Zheng, *J. High Energy Phys.* **02** (2020) 150.
[15] W.-Q. Chen, C.-M. Jian, L. Kong, Y.-Z. You, and H. Zheng, *Phys. Rev. B* **102**, 045139 (2020).
[16] M. Blume, V. J. Emery, and R. B. Griffiths, *Phys. Rev. A* **4**, 1071 (1971).
[17] B. Nienhuis, A. N. Berker, E. K. Riedel, and M. Schick, *Phys. Rev. Lett.* **43**, 737 (1979).
[18] C. Xu and Y.-Z. You, *Phys. Rev. B* **92**, 220416(R) (2015).
[19] P.-S. Hsin and N. Seiberg, *J. High Energy Phys.* **09** (2016) 095.
[20] M. Cheng and C. Xu, *Phys. Rev. B* **94**, 214415 (2016).
[21] F. Benini, P.-S. Hsin, and N. Seiberg, *J. High Energy Phys.* **04** (2017) 135.
[22] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, *Science* **303**, 1490 (2004).
[23] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, *Phys. Rev. B* **70**, 144407 (2004).
[24] M. Levin and T. Senthil, *Phys. Rev. B* **70**, 220403(R) (2004).
[25] J. A. Gracey, *J. Phys. A: Math. Gen.* **25**, L109 (1992).
[26] C. Xu and S. Sachdev, *Phys. Rev. Lett.* **100**, 137201 (2008).
[27] C. Xu, *Phys. Rev. B* **78**, 054432 (2008).
[28] C.-M. Jian, A. Rasmussen, Y.-Z. You, and C. Xu, [arXiv:1708.03050](https://arxiv.org/abs/1708.03050) [cond-mat.str-el].
[29] Y.-Z. You, Y.-C. He, A. Vishwanath, and C. Xu, *Phys. Rev. B* **97**, 125112 (2018).
[30] T. Alanne and S. Blasi, *Phys. Rev. D* **98**, 116004 (2018).
[31] R. Boyack, A. Rayyan, and J. Maciejko, *Phys. Rev. B* **99**, 195135 (2019).
[32] B. Zhao, J. Takahashi, and A. W. Sandvik, *Phys. Rev. Lett.* **125**, 257204 (2020).
[33] B. Zhao, J. Takahashi, and A. W. Sandvik, *Chin. Phys. B* **29**, 057506 (2020).

- [34] M. Barkeshli and J. McGreevy, *Phys. Rev. B* **89**, 235116 (2014).
- [35] M. Barkeshli, N. Y. Yao, and C. R. Laumann, *Phys. Rev. Lett.* **115**, 026802 (2015).
- [36] D. T. Son, *Phys. Rev. X* **5**, 031027 (2015).
- [37] M. A. Metlitski and A. Vishwanath, *Phys. Rev. B* **93**, 245151 (2016).
- [38] E. Witten, *Phys. Rev. B* **94**, 195150 (2016).
- [39] Y. Q. Qin, Y.-Y. He, Y.-Z. You, Z.-Y. Lu, A. Sen, A. W. Sandvik, C. Xu, and Z. Y. Meng, *Phys. Rev. X* **7**, 031052 (2017).
- [40] N. Ma, G.-Y. Sun, Y.-Z. You, C. Xu, A. Vishwanath, A. W. Sandvik, and Z. Y. Meng, *Phys. Rev. B* **98**, 174421 (2018).
- [41] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, *Phys. Rev. X* **7**, 031051 (2017).
- [42] J. Yang, A. W. Sandvik, and L. Wang, [arXiv:2104.08887](https://arxiv.org/abs/2104.08887).
- [43] H. Shackleton, A. Thomson, and S. Sachdev, *Phys. Rev. B* **104**, 045110 (2021).
- [44] W. Wang, D.-C. Lu, X. Y. Xu, Y.-Z. You, and Z. Y. Meng, *Phys. Rev. B* **100**, 085123 (2019).
- [45] N. Desai and R. K. Kaul, *Phys. Rev. B* **102**, 195135 (2020).
- [46] P. Cvitanović, *Group Theory: Birdtracks, Lie's, and Exceptional Groups* (Princeton University Press, Princeton, NJ, 2008).