

Deconfined quantum critical point with nonlocality

Yichen Xu¹, Xiao-Chuan Wu¹, and Cenke Xu¹*Department of Physics, University of California, Santa Barbara, California 93106, USA*

(Received 6 July 2022; accepted 11 October 2022; published 18 October 2022)

The deconfined quantum critical point (DQCP) between the Néel and valence bond solid order was originally proposed in quantum spin systems with a local Hamiltonian. In the last few years analogs of DQCPs with nonlocal interactions have been explored, which can lead to rich possibilities. The nonlocal interactions can either arise from an instantaneous long-range interaction in the Hamiltonian, or from gapless modes that reside in one higher spatial dimension. Here, we consider another mechanism of generating nonlocal interactions by coupling the DQCP to the “hot spots” of a Fermi surface. We demonstrate that at least within a substantial energy window, the physics of the DQCP is controlled by a new fixed point with a dynamical exponent $z > 1$.

DOI: [10.1103/PhysRevB.106.155131](https://doi.org/10.1103/PhysRevB.106.155131)

I. INTRODUCTION

A deconfined quantum critical point (DQCP) occurs between two phases that spontaneously break two different symmetries that do not contain each other as a subgroup. The original DQCP was proposed as a direct unfine-tuned continuous quantum phase transition between the collinear Néel and the valence bond solid (VBS) orders on the square lattice [1,2]. Various analogs of the original DQCP were studied, for example, the transition between the superfluid and various density waves of a quantum boson system can be described in a similar framework as that of the DQCP with an easy-plane anisotropy [3,4]; later the DQCP was also generalized to lattices where the spin order and VBS order both have a different structure from the original DQCP [5]. In all these examples the two ordered phases separated by the DQCP break very different 0-form symmetries; but nowadays one can generalize the notion of DQCP to situations that involve higher-form symmetries. For example, a direct transition between a magnetic order and a topological order can be viewed as a DQCP between a phase with spontaneous breaking of a 0-form symmetry and another phase with spontaneous breaking of a(n) (emergent) 1-form symmetry [6–15]. In the past two decades, a lot of progress has been made towards understanding various aspects of the DQCP, including its connection to mixed ‘t Hooft anomaly and higher-dimensional symmetry protected topological phases [16], as well as a duality web that connects different Lagrangian descriptions of the DQCP [17–24], etc.

Despite all the theoretical progress, the nature of the original DQCP proposed on a square lattice has always remained controversial. Very encouraging evidence of DQCP was found in the numerics on a two-dimensional (2D) lattice quantum spin model dubbed the “ J - Q ” model [25,26], as well as loop models in the 3D Euclidean space [27,28], but numerical simulations have also observed unusual scaling behaviors [29,30] and other complexities [31]. Recently, the DQCP has also been challenged by the “conformal bootstrap” method of analyzing conformal field theories (CFTs): The critical exponents

obtained from numerical simulations seem incompatible with the bounds given by a conformal bootstrap [32,33]. Though these should not exclude the possibility that the DQCP still exists in other lattice models with critical exponents that are consistent with the conformal bootstrap bounds, a consensus on the nature of the DQCP awaits further efforts.

In this paper, rather than trying to address the infrared nature of the original DQCP, we explore a possible continuous quantum phase transition close to the originally proposed DQCP, starting with the transition between the easy-plane Néel order and the VBS order. In particular, we will discuss the effect of nonlocality on DQCP. Nonlocality of a system can directly arise from a long-range instantaneous interaction in the Hamiltonian [34,35], or from coupling to gapless modes in one higher dimension, when the system is realized at the boundary of a bulk [36–47]. It was shown that, by coupling to the bulk quantum critical modes, the transition between the Néel and VBS order could be driven to a new fixed point [41,42]. Nonlocality arising from holography was also explored in Ref. [48].

Here, we explore nonlocality arising from a more realistic mechanism. Nonlocality in space-time usually translates to nonanalyticity in the momentum-frequency space. It is well known that, based on the Hertz-Millis theory [49,50], by coupling an order parameter ϕ to a Fermi surface, the dynamics of the order parameter acquires a singular contribution in the momentum-frequency space. In particular, when the order parameter carries a finite momentum that connects two “hot spots” of the Fermi surface, after formally integrating out the fermions, the order parameter acquires a singular term $\sim \sum_{\omega, q} |\omega| |\phi_{\omega, q}|^2$. Within the framework of the Hertz-Millis theory, this singular term renders the original $\sum_{\omega, q} \omega^2 |\phi_{\omega, q}|^2$ term in the Lagrangian irrelevant, and leads to a $z = 2$ Landau-Ginzburg theory of the order parameter ϕ . But the effect of the coupling to the hot spots will be more complex in the case of DQCP, as the physical order parameter ϕ is now a composite operator of the deconfined degrees of freedom at the DQCP. We note that novel physics arising from

coupling to a background Fermi surface in a one-dimensional setup was explored recently [51].

II. EASY-PLANE DQCP COUPLED WITH HOT SPOTS

Let us first inspect the easy-plane DQCP between the in-plane Néel order and a VBS order on a square lattice. The order parameters involved in this transition include a two-component in-plane Néel order (N_x, N_y) at momentum (π, π) , and a two-component VBS order parameter (V_x, V_y) at momentum $(\pi, 0)$ and $(0, \pi)$, respectively. Since all these order parameters carry a finite momentum, in principle they would acquire a singular term in the form sketched above when the easy-plane DQCP occurs with a background Fermi surface, assuming their momenta connect hot spots of the Fermi surface. The Lagrangian that describes the easy-plane DQCP is an easy-plane $\mathbb{CP}^1$ model, and it is known that this theory enjoys a self-duality [17], i.e., the in-plane Néel order parameter (N_x, N_y) is a bilinear of the $\mathbb{CP}^1$ field $(N_x, N_y) \sim (z^\dagger \sigma^x z, z^\dagger \sigma^y z)$, and the VBS order parameter along the x and y direction is a bilinear of the dual $\mathbb{CP}^1$ field (the vortex of z_1 and z_2 , respectively): $(V_x, V_y) \sim (v^\dagger \sigma^x v, v^\dagger \sigma^y v)$. Then after we integrate out the background Fermi surface according to the Hertz-Millis theory, the action that describes the transition becomes

$$\begin{aligned} \mathcal{S} = & \int d^2x d\tau \sum_{\alpha=1,2} |(\partial - iA)z_\alpha|^2 + r|z_\alpha|^2 + u|z_\alpha|^4 \\ & + \sum_{\omega, \mathbf{q}} \sum_{i=x,y} g|\omega| |(z^\dagger \sigma^i z)_{\omega, \mathbf{q}}|^2, \end{aligned} \quad (1)$$

and the dual action reads

$$\begin{aligned} \mathcal{S}_d = & \int d^2x d\tau \sum_{\alpha=1,2} |(\partial - i\tilde{A})v_\alpha|^2 + \tilde{r}|v_\alpha|^2 + \tilde{u}|v_\alpha|^4 \\ & + \sum_{\omega, \mathbf{q}} \sum_{i=x,y} \tilde{g}|\omega| |(v^\dagger \sigma^i v)_{\omega, \mathbf{q}}|^2, \end{aligned} \quad (2)$$

where $\tilde{r} = -r$. The actions above will be the starting point of our study; higher-order singular terms beyond the Hertz-Millis theory that also arise from integrating out the background fermions will be briefly discussed later. In this paper we will show that, although the bare values of g and $\tilde{g}$ can differ, they may actually flow to a fixed point where $g_* = \tilde{g}_*$. Hence this fixed point not only corresponds to a direct in-plane Néel-to-VBS transition, but our calculation suggests that this new fixed point may still have the self-duality as the originally proposed easy-plane DQCP [17], but we do not make a statement about the presence of the enlarged emergent $O(4)$ symmetry that can be perceived through the low-energy effective nonlinear sigma model of the easy-plane DQCP [52], as well as the duality web [18–24].

In order to study the theory Eq. (1) in a controllable fashion, we follow the standard procedure (see, for example, Refs. [53,54]) by introducing the Hubbard-Stratonovich auxiliary fields λ_α and Φ^i to decompose the two quartic terms of z_α ,




FIG. 1. One-loop Feynman diagrams that contribute to $\beta(g)$. Here, the solid, dashed, dotted, and wavy lines represent the correlators of $z_{a,\alpha}$, λ_α , Φ^i , and gauge field A_μ , respectively.

and consider the following large- N generalization of Eq. (1) at the critical point $r = 0$:

$$\begin{aligned} \mathcal{S} = & \int d^2x d\tau \sum_{a=1}^N \sum_{\alpha=1,2} |(\partial - iA)z_{a,\alpha}|^2 + i\lambda_\alpha |z_{a,\alpha}|^2 \\ & + i \sum_{i=x,y} \Phi^i (z_a^\dagger \sigma^i z_a). \end{aligned} \quad (3)$$

With large N , the correlators of the Hubbard-Stratonovich fields, and the gauge field read

$$\begin{aligned} \langle \lambda_\alpha(\vec{q}) \lambda_{\alpha'}(-\vec{q}) \rangle &= \frac{8}{N} |\vec{q}| \delta_{\alpha,\alpha'}, \\ \langle A_\mu(\vec{q}) A_\nu(-\vec{q}) \rangle &= \frac{16}{2N} \left(\frac{\delta_{\mu\nu} - q_\mu q_\nu / q^2}{|q|} \right), \\ \langle \Phi^i(\vec{q}) \Phi^{i'}(-\vec{q}) \rangle &= g |v| \delta_{i,i'}, \end{aligned} \quad (4)$$

where $\vec{q} = (v, \mathbf{q})$. We assume that g is at the order of $1/N$.

We proceed by calculating the renormalization group (RG) flow of g using the momentum-shell RG by integrating out the modes with momentum within $\Lambda/b < |k| < \Lambda$ (the calculations are repeated with the dimensional regularization as well); the most relevant Feynman diagrams are listed in Fig. 1. Figures 1(a)–1(d) are the standard contributions to

the leading-order $1/N$ expansion of the $\mathbb{CP}^{N-1}$ model [54]. The key of the calculation is as follows: In the large- N limit, the parameter g is exactly marginal, as $z_a^\dagger \sigma^i z_a$ has a scaling dimension $[z_a^\dagger \sigma^i z_a] = 1$ for $i = x, y$, while Φ^i has a scaling dimension $[\Phi^i] = 2$. With finite N , the scaling dimension of $z_a^\dagger \sigma^i z_a$ receives a negative correction at the order of $1/N$: $[z_a^\dagger \sigma^i z_a] = 1 - 28/(3\pi^2 N)$, which makes g weakly relevant with large but finite N . Hence the beta function of g should take the form

$$\frac{dg}{d \ln b} = \beta(g) = \frac{56}{3\pi^2 N} g - C g^2. \quad (5)$$

When C is positive and order unity, g will flow to a fixed point at the order of $g_* \sim 1/N$.

Figures 1(e) and 1(f) potentially contribute to the coefficient C in the beta function above. Figure 1(e) has a vanishing contribution at the easy-plane DQCP under consideration right now, due to the matrix identity $\sum_{j=x,y} \sigma^j \sigma^i \sigma^j = 0$ for $i = x, y$. Figure 1(f) can be interpreted as the self-energy correction to $z_{a,\alpha}$,

$$\begin{aligned} \Sigma(\omega, \mathbf{k}) &= \sum_{j=x,y} g \sigma^j \sigma^j \int_{\Lambda/b}^{\Lambda} \frac{d^3 q}{(2\pi)^3} \frac{|v|}{(q+k)^2} \\ &= g \frac{\omega^2}{2\pi^2} \ln b \times \sigma^0 + \dots \end{aligned} \quad (6)$$

Here, $\vec{q} = (v, \mathbf{q})$, $\vec{k} = (\omega, \mathbf{k})$. The ellipses in the equation represent terms which do not contribute at order $\ln b$. This self-energy correction will modify the Gaussian part of the action of $z_{a,\alpha}$ to

$$\mathcal{L} = z_{a,\alpha}^* (|\partial_\tau|^{2[1-g/(4\pi^2)]} - \partial_x^2) z_{a,\alpha} + \dots \quad (7)$$

This result implies that after coupling to the background Fermi surface, the space-time scaling of the original easy-plane DQCP is modified, which should now be

$$\tau \rightarrow b^{-z} \tau, \quad \mathbf{x} \rightarrow b^{-1} \mathbf{x}, \quad (8)$$

where $z = 1 + \frac{g}{4\pi^2} + O(g^2)$ is the dynamical exponent. Here, we remind the readers that, in the original Hertz-Millis theory, when an order parameter is coupled to hot spots of a Fermi surface, the Gaussian part of the Landau-Ginzburg theory of the order parameter has a dynamical exponent $z = 2$. Here, although the order parameter is a composite operator of $z_{a,\alpha}$, the dynamical scaling exponent z is still modified due to its coupling to the Fermi surface.

The wave-function renormalization in Fig. 1(f) is in fact equivalent to the modification of the space-time scaling, plus a correction to the scaling dimension of $z_{a,\alpha}$: $\Delta[z_{a,\alpha}] = g/(8\pi^2)$. Eventually the beta function of g reads

$$\beta(g) \equiv \frac{dg}{d \ln b} = \left(\frac{56}{3\pi^2 N} - \frac{g}{2\pi^2} \right) g, \quad (9)$$

where the first term arises from Figs. 1(a)–1(d), while the second term is the additional wave-function renormalization from Fig. 1(f) as described above. Indeed, for $g > 0$, the theory flows to a new fixed point $g_* = \frac{112}{3N} + O(\frac{1}{N^2})$. Several two-loop diagrams such as the Aslamazov-Larkin diagrams appear to be also at the $1/N$ order, but careful evaluation shows that these diagrams either do not contribute to the beta

function as they do not lead to a logarithmic divergence, or their contributions cancel out with each other [54].

The same calculation applies to $\tilde{g}$ in Eq. (2). Hence although the bare values of g and $\tilde{g}$ in Eqs. (1) and (2) can be different, the RG equations above suggest that they would flow to a fixed point where $g_* = \tilde{g}_*$. Hence our calculation suggests that at this fixed point the self-duality of the original easy-plane DQCP still holds. Another more technical note is that the VBS order parameter $V_x \sim v^\dagger \sigma^x v$ is also the monopole operator of gauge field A_μ in Eq. (1), hence the $\tilde{g}$ term in Eq. (2) also corresponds to a correction to the action of the gauge field A_μ . But since we expect $\tilde{g}$ to flow to a fixed point at order $1/N$, at the self-consistent level we can ignore this singular correction and use the gauge field propagator in the large- N limit for our calculation.

At this new RG fixed point, we obtain scaling dimensions for the following operators:

$$\begin{aligned} [\lambda^+] &= 2 - \frac{80}{3\pi^2 N} - \frac{3g_*}{4\pi^2} = 2 - \frac{164}{3\pi^2 N} + O\left(\frac{1}{N^2}\right), \\ [\lambda^-] &= 2 + \frac{16}{3\pi^2 N} + \frac{g_*}{4\pi^2} = 2 + \frac{44}{\pi^2 N} + O\left(\frac{1}{N^2}\right), \\ [z_a^\dagger \sigma^{x,y} z_a] &= 1. \end{aligned} \quad (10)$$

Here, we have defined the operators $\lambda^\pm = (\lambda_1 \pm \lambda_2)/2$. Some two-loop diagrams such as the ones considered in Ref. [55] contribute to the evaluation of $[\lambda^+]$. The critical exponent ν is inferred from the scaling dimension of λ^+ :

$$\nu^{-1} = 2 + z - [\lambda^+] = 1 + \frac{64}{\pi^2 N} + O\left(\frac{1}{N^2}\right). \quad (11)$$

These standard $1/N$ expansions may not be extremely reliable at the physically relevant case with small N , but the scaling dimensions of $z_a^\dagger \sigma^{x,y} z_a$ should be exactly 1 at $g = g_*$. This is due to the fact that the system remains scaling invariant at the fixed point with nonzero g_* , and the singular frequency dependence $|\omega|$ in Eq. (1) cannot be renormalized, then to keep the system scaling invariant the in-plane Néel order parameter (N_x, N_y) must have a scaling dimension 1 in the Euclidean space-time.

We note that based on the Hertz-Millis theory there is another singular interaction $|\omega| |(z^\dagger \sigma^z z)_{\omega, \mathbf{q}}|^2$ that would also be generated by coupling the z component of the Néel order to the background Fermi surface, but this term is irrelevant with the large- N generalization of the easy-plane DQCP, as the scaling dimension of $z^\dagger \sigma^z z$ is 2 with large N .

We would also like to comment on the validity of the Hertz-Millis theory. It was noted in Ref. [56] that, when we couple an order parameter ϕ to a Fermi surface, besides generating singular terms at the quadratic order of ϕ , similar higher-order terms $\sim \phi^n$ with space-time singularity are also generated after integrating out the fermions. It was shown in Ref. [56] that direct power counting suggests these higher-order terms are marginal at the $z = 2$ Gaussian fixed point of the Hertz-Millis theory, hence it is no longer justified to ignore these terms. In fact, in our case, once we identify ϕ as $z^\dagger \sigma^i z$, the higher-order terms pointed out in Ref. [56] are still marginal at the new fixed point, since the scaling dimension $[z^\dagger \sigma^i z]$ is precisely 1. But this does not mean that the physics at the new fixed point

we derived is not observable. Let us return to the original theory with a bosonic field ϕ coupled with N_F copies of Fermi surfaces,

$$\mathcal{L}_{\text{BF}} = \sum_{l=1}^{N_F} f_{l1}^\dagger (\partial_\tau - i\mathbf{v}_1 \cdot \nabla) f_{l1} + f_{l2}^\dagger (\partial_\tau - i\mathbf{v}_2 \cdot \nabla) f_{l2} + u\phi \left[\sum_l [f_{l1}^\dagger T f_{l2} + (1 \leftrightarrow 2)] \right], \quad (12)$$

where 1,2 label two points of the Fermi surface connected by the momentum of ϕ , and T is a flavor matrix. The parameter g in Eq. (1) is about $g \sim N_F u^2$, and since $g \sim 1/N$, we need $u \sim \sqrt{1/(NN_F)}$. If we fix N , the higher-order singular terms considered in Ref. [56] will be at the order of $1/N_F^{n/2-1}$. Hence with large N_F , although the ultimate fate of these higher-order singular terms in the infrared limit is unclear, there could be a large energy window where the physics is controlled by the fixed point g_* derived above.

We can also compute the self-energy of the fermions at the hot spot to the leading nontrivial order of u :

$$\Sigma_F(\omega, \mathbf{k}) \sim 2u^2 \sigma^0 \int \frac{d\nu d^2\mathbf{q}}{(2\pi)^3} \frac{1}{i(\omega - \nu) - \mathbf{v}_2 \cdot (\mathbf{k} - \mathbf{q})} \times \frac{1}{(\nu^2 + \mathbf{q}^2)^{1-\eta/2}}. \quad (13)$$

We have taken the correlator of the bosonic field $N_{x,y}(\vec{q}) \sim z^\dagger \sigma^{x,y} z$ to be $1/(\nu^2 + \mathbf{q}^2)^{1-\eta/2}$, where η is the anomalous dimension of the in-plane Néel order parameter $N_{x,y}$ at the purely bosonic easy-plane DQCP. Carrying out the integral, we obtain

$$\Sigma_F(\omega, 0) \sim -iu^2 \text{sgn}(\omega) |\omega|^\eta \sigma^0. \quad (14)$$

Generally we expect the fermions at the hot spots to have non-Fermi-liquid-like self-energy for a considerable energy window.

III. SU(2)-INVARIANT DQCP COUPLED WITH HOT SPOTS

Here, we briefly discuss the SU(2)-invariant DQCP coupled to a background Fermi surface, which can be studied in the same way as the easy-plane case. Here, we only need one Hubbard-Stratonovich field λ^+ to decompose the quartic term, and we obtain the following large- N theory at the critical point:

$$\mathcal{S} = \int d^2x d\tau \sum_{a=1}^N \sum_{\alpha=1,2} |(\partial - iA)z_{a,\alpha}|^2 + i\lambda^+ |z_{a,\alpha}|^2 + i \sum_{i=x,y,z} \Phi^i(z_a^\dagger \sigma^i z_a). \quad (15)$$

Now the Fermi surfaces are coupled to all three components of the Néel order parameter $\vec{N} = z^\dagger \vec{\sigma} z$. The new dynamic exponent is now $z = 1 + \frac{3g}{8\pi^2} + O(g^2)$, and the beta function of g is

$$\beta(g) \equiv \frac{dg}{d \ln b} = \frac{16}{\pi^2 N} g - \frac{g^2}{4\pi^2}. \quad (16)$$

If $g > 0$, the theory flows to a new RG fixed point at $g_* = \frac{64}{N}$. At the new RG fixed point, we have the following scaling dimensions,

$$[\lambda^+] = 2 - \frac{24}{\pi^2 N} - \frac{9g_*}{8\pi^2} = 2 - \frac{96}{\pi^2 N} + O\left(\frac{1}{N^2}\right),$$

$$[z_a^\dagger \sigma^{x,y,z} z_a] = 1, \quad (17)$$

and the critical exponent

$$\nu^{-1} = 2 + z - [\lambda^+] = 1 + \frac{120}{\pi^2 N} + O\left(\frac{1}{N^2}\right). \quad (18)$$

Again, the Néel order parameter has a scaling dimension $[\vec{N}] = 1$ exactly at the new fixed point. Though it is not so convenient to directly compute the RG flow of the singular term of the VBS order parameter due to the lack of a dual Lagrangian for the SU(2)-invariant DQCP, we expect the scaling dimension of the VBS order parameter should also be 1 at the new fixed point. To elaborate, the VBS order parameter V_x, V_y should still acquire a singular term $V_x |\partial_\tau| V_x + (x \rightarrow y)$ from coupling to the Fermi surface. This term is singular (nonanalytic) in the frequency space, and also long range in the temporal direction. The form of the singular term $|\partial_\tau|$ cannot be renormalized under RG. And if this term flows to a fixed point, this term $V_x |\partial_\tau| V_x + (x \rightarrow y)$ remains scaling invariant at the fixed point, hence the VBS order parameter V^x should have precisely scaling dimension 1.

Reference [2] pointed out that at the DQCP there are two length scales, one for the correlation length ξ of the order parameter N_x , and the other is the thickness of the VBS domain wall (ξ_{VBS}). In our system these two length scales still exist, and the relation between these two length scales should be similar to what was pointed out in Ref. [2]: $\xi_{\text{VBS}} \sim \xi f(\lambda \xi^{d+z-\Delta})$, where λ is the strength of the fourfold monopole operator of the gauge field that is the minimal topological defect allowed by symmetry on a square lattice, and Δ is the scaling dimension of the fourfold monopole, which we assume is greater than $d + z$ (which means that the fourfold monopole is irrelevant at the DQCP). z is the dynamical exponent computed in this work. Following the discussions in Ref. [2], one can further infer that $\xi_{\text{VBS}} \sim \xi^{(\Delta-z)/2}$ close to the DQCP. Since we expect Δ to be greater than $2 + z$, ξ_{VBS} diverges faster than ξ near the DQCP.

The quantum critical modes at the new fixed point with $z > 1$ will make a contribution $C \sim T^{2/z}$ to the specific heat, which is different from ordinary $(2+1)d$ QCP with $z = 1$. But the background Fermi surface would contribute a specific heat linear with temperature T , hence we expect that the contribution from the quantum critical modes will be overshadowed by the background Fermi surface.

IV. DISCUSSION

In this paper we discussed the fate of the DQCP when it occurs with a background Fermi surface. We demonstrated that with a large number of copies of Fermi surfaces, there is a substantial energy window where the easy-plane DQCP is controlled by a self-dual fixed point with a dynamical exponent $z > 1$. We did not pursue a full renormalization group analysis of the boson-fermion coupled theory, but an analysis

such as the ones discussed in Refs. [57,58] when the order parameter ϕ is a composite operator of deconfined degrees of freedom is very much worth studying in the future.

Many insights of the DQCP, including the emergent symmetry, 't Hooft anomaly, as well as a possible phase diagram and RG flow, can be gained from the nonlinear sigma model (NLSM) approach that unifies all the order parameters in one action [52,59–64]. The very key term in the NLSM is a topological term. The Néel and VBS order parameters can also be treated on an equal footing in the U(1) and SU(2) spin-liquid language [22]. In the future it is also worth exploring the consequence of coupling the DQCP to a Fermi surface using these different formalisms.

Besides the DQCP, our study is also meaningful to the interaction-driven metal-insulator transition (MIT) where the insulator phase has a certain density wave order. The basic formalism of the theory describing this MIT involves introducing bosonic partons that carry the electric charge,

and fermionic partons that carry the spin. This MIT is interpreted as a superfluid-to-density wave transition of the charged bosonic parton sector [65,66] (the “superfluid” phase of the bosonic sector of the phase diagram corresponds to the metallic phase [67]), which is also described by a $\mathbb{CP}^{N-1}$ model in which the bosonic matter fields are vortices of the charged bosonic parton. There are multiple components of the vortex fields whose condensate corresponds to the degenerate density wave patterns of the insulator phase. When the density wave order parameter couples to the hot spots of the Fermi surface of the fermionic spinon sector, the same singular terms such as the one considered in our current paper will arise. Our study indicates that the physics at this MIT could be controlled by a new fixed point with dynamical exponent $z > 1$.

ACKNOWLEDGMENT

This work is supported by NSF under Grant No. DMR-1920434, and the Simons Investigator program.

-
- [1] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, *Science* **303**, 1490 (2004).
 - [2] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, *Phys. Rev. B* **70**, 144407 (2004).
 - [3] L. Balents, L. Bartosch, A. Burkov, S. Sachdev, and K. Sengupta, *Prog. Theor. Phys. Suppl.* **160**, 314 (2005).
 - [4] A. A. Burkov and L. Balents, *Phys. Rev. B* **72**, 134502 (2005).
 - [5] C.-M. Jian, A. Thomson, A. Rasmussen, Z. Bi, and C. Xu, *Phys. Rev. B* **97**, 195115 (2018).
 - [6] Z. Nussinov and G. Ortiz, *Ann. Phys.* **324**, 977 (2009).
 - [7] O. Aharony, N. Seiberg, and Y. Tachikawa, *J. High Energy Phys.* **08** (2013) 115.
 - [8] S. Gukov and A. Kapustin, *arXiv:1307.4793*.
 - [9] A. Kapustin and R. Thorngren, *arXiv:1308.2926*.
 - [10] A. Kapustin and R. Thorngren, *arXiv:1309.4721*.
 - [11] A. Kapustin and N. Seiberg, *J. High Energy Phys.* **04** (2014) 001.
 - [12] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, *J. High Energy Phys.* **02** (2015) 172.
 - [13] P.-S. Hsin, H. T. Lam, and N. Seiberg, *SciPost Phys.* **6**, 039 (2019).
 - [14] N. Seiberg, *SciPost Phys.* **8**, 050 (2020).
 - [15] J. McGreevy, *arXiv:2204.03045*.
 - [16] A. Vishwanath and T. Senthil, *Phys. Rev. X* **3**, 011016 (2013).
 - [17] O. I. Motrunich and A. Vishwanath, *Phys. Rev. B* **70**, 075104 (2004).
 - [18] C. Xu and Y.-Z. You, *Phys. Rev. B* **92**, 220416(R) (2015).
 - [19] D. F. Mross, J. Alicea, and O. I. Motrunich, *Phys. Rev. Lett.* **117**, 016802 (2016).
 - [20] A. C. Potter, C. Wang, M. A. Metlitski, and A. Vishwanath, *Phys. Rev. B* **96**, 235114 (2017).
 - [21] P.-S. Hsin and N. Seiberg, *J. High Energy Phys.* **09** (2016) 095.
 - [22] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, *Phys. Rev. X* **7**, 031051 (2017).
 - [23] T. Senthil, D. T. Son, C. Wang, and C. Xu, *Phys. Rep.* **827**, 1 (2019).
 - [24] Y. Q. Qin, Y.-Y. He, Y.-Z. You, Z.-Y. Lu, A. Sen, A. W. Sandvik, C. Xu, and Z. Y. Meng, *Phys. Rev. X* **7**, 031052 (2017).
 - [25] A. W. Sandvik, *Phys. Rev. Lett.* **98**, 227202 (2007).
 - [26] R. G. Melko and R. K. Kaul, *Phys. Rev. Lett.* **100**, 017203 (2008).
 - [27] A. Nahum, P. Serna, J. T. Chalker, M. Ortuño, and A. M. Somoza, *Phys. Rev. Lett.* **115**, 267203 (2015).
 - [28] A. Nahum, J. T. Chalker, P. Serna, M. Ortuño, and A. M. Somoza, *Phys. Rev. X* **5**, 041048 (2015).
 - [29] A. W. Sandvik, *Phys. Rev. Lett.* **104**, 177201 (2010).
 - [30] H. Shao, W. Guo, and A. W. Sandvik, *Science* **352**, 213 (2016).
 - [31] J. Zhao, Y.-C. Wang, Z. Yan, M. Cheng, and Z. Y. Meng, *Phys. Rev. Lett.* **128**, 010601 (2022).
 - [32] F. Kos, D. Poland, D. Simmons-Duffin, and A. Vichi, *J. High Energy Phys.* **11** (2015) 106.
 - [33] Y. Nakayama and T. Ohtsuki, *Phys. Rev. Lett.* **117**, 131601 (2016).
 - [34] A. W. Sandvik, *Phys. Rev. Lett.* **104**, 137204 (2010).
 - [35] S. Yang, D.-X. Yao, and A. W. Sandvik, *arXiv:2001.02821*.
 - [36] T. Grover and A. Vishwanath, *arXiv:1206.1332*.
 - [37] L. Zhang and F. Wang, *Phys. Rev. Lett.* **118**, 087201 (2017).
 - [38] C. Ding, L. Zhang, and W. Guo, *Phys. Rev. Lett.* **120**, 235701 (2018).
 - [39] L. Weber, F. Parisen Toldin, and S. Wessel, *Phys. Rev. B* **98**, 140403(R) (2018).
 - [40] L. Weber and S. Wessel, *Phys. Rev. B* **100**, 054437 (2019).
 - [41] Y. Xu, X.-C. Wu, C.-M. Jian, and C. Xu, *Phys. Rev. B* **101**, 184419 (2020).
 - [42] C.-M. Jian, Y. Xu, X.-C. Wu, and C. Xu, *SciPost Phys.* **10**, 033 (2021).
 - [43] M. A. Metlitski, *SciPost Phys.* **12**, 131 (2022).
 - [44] S. Liu, H. Shapourian, A. Vishwanath, and M. A. Metlitski, *Phys. Rev. B* **104**, 104201 (2021).
 - [45] F. Parisen Toldin, *Phys. Rev. Lett.* **126**, 135701 (2021).
 - [46] F. Parisen Toldin and M. A. Metlitski, *Phys. Rev. Lett.* **128**, 215701 (2022).
 - [47] R. Ma, L. Zou, and C. Wang, *SciPost Phys.* **12**, 196 (2022).

- [48] G. La Nave and P. W. Phillips, *Phys. Rev. D* **94**, 126018 (2016).
- [49] J. A. Hertz, *Phys. Rev. B* **14**, 1165 (1976).
- [50] A. J. Millis, *Phys. Rev. B* **48**, 7183 (1993).
- [51] B. Danu, M. Vojta, T. Grover, and F. F. Assaad, [arXiv:2204.00029](#).
- [52] T. Senthil and M. P. A. Fisher, *Phys. Rev. B* **74**, 064405 (2006).
- [53] R. K. Kaul and S. Sachdev, *Phys. Rev. B* **77**, 155105 (2008).
- [54] S. Benvenuti and H. Khachatryan, *J. High Energy Phys.* **05** (2019) 214.
- [55] X.-G. Wen and Y.-S. Wu, *Phys. Rev. Lett.* **70**, 1501 (1993).
- [56] A. Abanov and A. Chubukov, *Phys. Rev. Lett.* **93**, 255702 (2004).
- [57] M. A. Metlitski and S. Sachdev, *Phys. Rev. B* **82**, 075128 (2010).
- [58] A. Schliefl, P. Lunts, and S.-S. Lee, *Phys. Rev. X* **7**, 021010 (2017).
- [59] C. Xu and A. W. W. Ludwig, *Phys. Rev. Lett.* **110**, 200405 (2013).
- [60] Z. Bi, A. Rasmussen, Y. BenTov, and C. Xu, [arXiv:1605.05336](#).
- [61] R. Ma and C. Wang, *Phys. Rev. B* **102**, 020407(R) (2020).
- [62] A. Nahum, *Phys. Rev. B* **102**, 201116(R) (2020).
- [63] Z. Wang, M. P. Zaletel, R. S. K. Mong, and F. F. Assaad, *Phys. Rev. Lett.* **126**, 045701 (2021).
- [64] L. Zou, Y.-C. He, and C. Wang, *Phys. Rev. X* **11**, 031043 (2021).
- [65] Y. Xu, X.-C. Wu, M. Ye, Z.-X. Luo, C.-M. Jian, and C. Xu, *Phys. Rev. X* **12**, 021067 (2022).
- [66] S. Musser, T. Senthil, and D. Chowdhury, [arXiv:2111.09894](#).
- [67] T. Senthil, *Phys. Rev. B* **78**, 045109 (2008).