

From Centaurs to comets: 40 Years

*Nuno Peixinho^a, Audrey Thirouin^b, Stephen C. Tegler^c,
Romina P. Di Sisto^{d,e}, Audrey Delsanti^f,
Aurélie Guilbert-Lepoutre^g, James G. Bauer^h*

^aCITEUC—Center for Earth and Space Research of the University of Coimbra, Geophysical and Astronomical Observatory of the University of Coimbra, Coimbra, Portugal ^bLowell Observatory, Flagstaff, AZ, United States ^cDepartment of Physics and Astronomy, Northern Arizona University, Flagstaff, AZ, United States ^dInstituto de Astrofísica de La Plata, CCT—CONICET—Universidad Nacional de La Plata, La Plata, Argentina ^eFacultad de Ciencias Astronómicas y Geofísicas, UNLP, La Plata, Argentina ^fAix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, Marseille, France ^gLGL-TPE, UMR 5276, CNRS, Claude Bernard Lyon 1 University, ENS Lyon, Villeurbanne Cedex, France ^hDepartment of Astronomy, University of Maryland, College Park, MD, United States

14.1 Introduction

Centaurs are a group of solar system minor bodies loosely classified as those orbiting around the Sun between the orbits of Jupiter and Neptune that are not trapped in any mean-motion resonance with a planet. They have been discovered 15 years before the Trans-Neptunian objects (TNOs), or Kuiper belt objects, except for Pluto/Charon. First announced as “slow-moving object Kowal,” on November 4, 1977, and then with the provisional designation 1977 UB, on November 14, when it was shown that a very high eccentricity orbit was not viable for it, (2060) Chiron was the first Centaur to be discovered ([Kowal and Gehrels, 1977](#)). Chiron was also precovered in photographic plates from 1941 and even from 1895 at the Boyden Observatory, South Africa ([Davies, 2001](#)).

[Fernández \(1980\)](#) suggested that Chiron might be a comet diffusing inwards, through planetary perturbations—from a cometary belt beyond Neptune—the by-then undetected Trans-Neptunian belt. But the orbit of Chiron was not its only unusual characteristic. [Tholen et al. \(1988\)](#) found that its brightness increased more than expected and, later on,

Meech and Belton (1990) directly detected a low-surface-brightness coma for Chiron. Then, it was hypothesized that Chiron was potentially an ice-rich object, a comet nucleus 180 km across, darkened by C-class carbonaceous soil that has been scattered inward from the Oort cloud (Hartmann et al., 1990). A second idea was that Chiron is most likely a bright member of what is known today as the Trans-Neptunian belt (Duncan et al., 1988). Finally, Hahn and Bailey (1990) concluded that Chiron-type orbits are short-lived and chaotic, with an ejection half-life of $t_{1/2} \sim 1$ Myr, therefore Chiron's current orbit should be a transition between a TNO and a short-period comet.

In 1992, 7 months before the discovery of the first TNO, a second Centaur was announced: 1992 AD, now known as (5145) Pholus (Scotti et al., 1992). No sign of coma was identified, but Pholus became the reddest known object in the solar system (Hainaut et al., 1992).

Given their close link, Centaurs have been mostly studied together with TNOs. Previous to this book, the most extensive review of this field was the book *The Solar System Beyond Neptune* (Barucci et al., 2008), preceded by works as the *Trans-Neptunian Objects and Comets* (Jewitt et al., 2008) and the pioneering *The first decadal review of the Edgeworth-Kuiper belt* (Davies and Barrera, 2004). Other reviews, such as Nesvorný (2018), Ferrari (2018), Luu and Jewitt (2002), and Jewitt (1999), are also of highly recommended reading. In this chapter, we will propose a brief overview of the current state of the art regarding composition, rotational and physical properties, and activity regarding the Centaur population as well as their links to the Trans-Neptunian and comet populations.

14.2 Centaurs as progeny of TNOs

Levison and Duncan (1997) and Duncan and Levison (1997) found that a belt of TNOs on rather circular and low inclined orbits—presently known as: classical TNOs—would create a dynamically excited population through scattering interactions with Neptune, hence called the scattered disk objects (SDOs), and that population should be the main source of Centaurs. Integrating different models for that scattered population for 4.5 Gyr provides injection rates of $\sim(1-5) \times 10^{-10}$ SDOs/year and an estimated number of Centaurs larger than ~ 1 km of the order of 10^6-10^8 (Di Sisto and Brunini, 2007; Volk and Malhotra, 2008). Plutinos, the TNOs in the 3:2 mean-motion resonance with Neptune, should also be a source of Centaurs (Morbidelli, 1997; Yu and Tremaine, 1999), contributing ~ 3 times less than SDOs (Di Sisto and Brunini, 2007), and the 2:1 resonant TNOs might also contribute (Tiscareno and Malhotra, 2009; de Elía et al., 2008). Although very secondary as source, every $\sim 60-200$ years a kilometer-sized Neptune Trojan enters the Centaurs region (Horner and Lykawka, 2010). Jupiter Trojans were also found to be a minor source of Centaurs (Di Sisto et al., 2019).

The mean lifetime of Centaurs is strongly dependent on their definition, because lifetime correlates with perihelion distance and semimajor axis and with the initial inclination of the progenitors, and can be as short as few Myr up to tens of Myr (Tiscareno and Malhotra, 2003; Di Sisto and Brunini, 2007; Volk and Malhotra, 2008; Bailey and Malhotra, 2009).

Interesting is the very low probability for a high inclination Centaur, or even a retrograde one, originating from the Trans-Neptunian region (Volk and Malhotra, 2013) and the suggestion that a heterogeneous group of up to 200 Centaurs with diameters $D \gtrsim 150$ km, inclinations

$i > 70$ degrees, and perihelia $q > 15$ AU should exist, originating mainly in the Oort cloud (Brasser et al., 2012; de la Fuente Marcos and de la Fuente Marcos, 2014).

Another very interesting aspect of Centaurs is the existence of binaries. From the 81 known Trans-Neptunian binaries, (42355) Typhon/Echidna and (65489) Ceto/Phorcys cross the orbit of Neptune. Brunini (2014) concludes that a small population of contact binary Centaurs may exist and Araujo et al. (2018) find that Typhon/Echidna could survive close encounters with the planets and even reach the terrestrial planets region as a binary.

14.3 Centaurs as progenitors of Jupiter-family comets

On early days, the expression short-period comets ($P < 200$ years) was frequently used and the Trans-Neptunian belt was thought to be their main source. Centaurs were merely a transient phase. The usage of short-period comets was replaced by the more concrete Jupiter-family comets (JFCs) and Halley-type comets (HTCs)—Chiron and Encke type are also used—using not the orbital period as a criterion but the Tisserand parameter relative to Jupiter, T_J (Tisserand, 1896; Kresák, 1972; Levison, 1996). JFCs are those with $2 \lesssim T_J \lesssim 3$.

It is interesting to notice that the studies regarding the source of JFCs have been made, mostly, regarding the direct link between them and TNOs. Centaurs are taken as a short-lived transient phase not decoupled from this JFC-TNO link. Right after the discovery of Pluto, Leonard (1930) speculated about the existence of Trans-Neptunian and Trans-Plutonian objects but without linking them explicitly to comets. The existence of a Trans-Neptunian region as a source of JFCs was first suggested by Edgeworth (1943, 1949) and Kuiper (1951), being properly addressed much later by Fernández (1980) and Duncan et al. (1988).

Relatively little attention has been given to the dynamics of the Centaur population itself. Tiscareno and Malhotra (2003) find that one-third of the Centaurs will actually be injected into the JFC population, while two-thirds will be ejected from the solar system or will enter the Oort cloud. About 20% of them have lifetimes as Centaurs shorter than 1 Myr, while another 20% have lifetimes exceeding 100 Myr. Horner et al. (2004b) found even that about one-fifth of the Centaurs that became JFCs will become Earth-crossing objects. Some Centaurs may even become temporarily captured as Trojans of the giant planets for up to 100 kyr (Horner and Evans, 2006). Most interesting is the recent finding that active Centaurs tend to evolve to JFCs, while the inactive ones tend to evolve to HTCs (Fernández et al., 2018).

14.4 Centaurs by themselves

14.4.1 Classifying Centaurs

There is no strict definition of Centaurs, which poses some evident problems when comparing different works. The most frequent classification forces them to have both perihelion and semimajor axis between the orbits of Jupiter and Neptune ($q > 5.2$ AU, $a < 30.1$ AU) (e.g., Gladman et al., 2008). With the previous definition, the most strict, as of February 1,

2019¹ there are 292 Centaurs, of which 26 are also classified as comets. Also widely used is the Deep Ecliptic Survey (DES) definition, in which the perihelion needs to be between the orbits of Jupiter and Neptune but without any constraints regarding the semimajor axis (Elliot et al., 2005). With this definition, there are 360 Centaurs listed² (see Fig. 14.1).




FIG. 14.1 Semimajor axis and eccentricity plot of known Centaurs. *Large gray dots* indicate known active Centaurs. *Black dots* indicate Centaurs with both perihelion and semimajor axis between those of Jupiter and Neptune, that is, the more strict definition. *White dots* indicate objects classified as Centaurs by DES alone. The *curves* where objects have perihelion equal to the semimajor axis of Jupiter, q_J , of Saturn, q_S , and of Neptune, q_N , are drawn. Several objects are out of scale.

14.4.2 Surface properties

14.4.2.1 Colors

When photometry is acquired almost simultaneously in various bands, or the eventual rotational brightness variations are negligible or averaged down, the magnitude difference between those bands defines a color. Centaurs' colors are usually measured using the Johnson UBV_RJHK photometric system—from $\approx 3600 \text{ \AA}$ to $\approx 2.2 \text{ \mu m}$ —or the SDSS *ugriz* system (e.g., Schwamb et al., 2018). In the most used color, Centaurs range from a neutral/gray solar-like $B-R = 1.0 \text{ mag}$ to an extraordinarily red $B-R = 2.0 \text{ mag}$. Since complex organic molecules are known to efficiently absorb shorter wavelengths of optical light, the redder the object the richer the object might be in complex organic molecules, although other compounds are capable of doing so. Surface colors can also be used to obtain the low-resolution reflectance spectrum, at a relatively low observation time cost and constraint, at least, the spectral continuum of the studied object (see Delsanti et al., 2004; Dalle Ore et al., 2015). Two large surveys post their

¹ See https://ssd.jpl.nasa.gov/sbdb_query.cgi.

² See <http://www.boulder.swri.edu/~buie/kbo/desclass.html>.

measurements of colors for these objects online the Tegler, Romanishin, and Consolmagno's survey (TRC),³ and the Minor Bodies of the Outer Solar System Survey (MBOSS).⁴

Early on, in the observations of Centaurs and TNOs, a controversy arose as to whether these objects exhibited a unimodal color distribution (Luu and Jewitt, 1996; Jewitt and Luu, 2001; Hainaut and Delsanti, 2002) or a bimodal color distribution (Tegler and Romanishin, 1998, 2003; Tegler et al., 2003). Models attempting to explain the two possible behaviors were very different. Luu and Jewitt (1996) put forth a model of steady radiation reddening and stochastic impacts to explain the unimodal colors of Centaurs and TNOs, but it has been shown that such simple model could not explain the color diversity as it was observed (Jewitt and Luu, 2001; Thébault and Doressoundiram, 2003; Thébault, 2003). Posterior modeling and laboratory works argue that it is, in fact, possible to reproduce the whole range of colors observed with an appropriate combination of initial albedo, meteoritic bombardment, and space weathering (Kaňuchová et al., 2012). On the other hand, Tegler et al. (2003) argue in favor of a thermally driven composition gradient in the primordial Trans-Neptunian belt that would result in bimodal colors. Brown et al. (2011) and Wong and Brown (2016) argue in favor of H₂S, rather than simple organic molecules, as the main reddening agent, where objects located beyond the H₂S sublimation line, hence not depleted of it, would suffer radiation reddening contrarily to the depleted ones that would result in gray surfaces.

We now see that Centaurs and different dynamical classes of TNOs, as well as different sizes, exhibit different colors (e.g., Tegler et al., 2008, 2016; Peixinho et al., 2012; Fraser and Brown, 2012; Wong and Brown, 2017). Consensus is growing toward a scenario where migration of outer planets disrupted the primordial disk of icy planetesimals, objects were scattered onto dynamically hot orbits and then onto Centaur orbits to present us with the bimodal color we see today. In contrast, dynamically cold TNOs were far enough away from the migrating outer planets that they were not scattered onto dynamically hot TNO orbits and so represent a third compositional class. Nonetheless, recent observations call the bimodality of Centaurs and, therefore, models to explain the bimodality into question. The 99.5% confidence level for the bimodality found by Peixinho et al. (2003) and Tegler et al. (2008) drops to 81% with today's larger sample (Tegler et al., 2016, using TRC survey data). Even reanalyzing TRC data for the 50 objects that follow Gladman et al.'s more strict definition of a Centaur, we get a confidence level of 91.6%, that is, no different conclusion. Looking at Fig. 14.2 we can see how the red population has a larger spreading today. It is interesting that Brown et al. (2011) predicted that the red population should have a broader color distribution than the gray one, given the several reddening agents with sublimation lines in the 20–35 AU region of the primordial Trans-Neptunian belt—although in their most recent paper, they claim only one species, H₂S, is serving as the reddening agent. It is, evidently, premature to rule out irradiation of simple organic molecules into complex organic molecules in favor of irradiation H₂S without further evidence, and without studying what would happen when ices are not in their pure form but in a certainly more realistic mixture with refractory material and how would that affect the sublimation lines.

³ See <http://www.physics.nau.edu/~teglar/research/survey.htm>.

⁴ See <http://www.eso.org/~ohainaut/MBOSS>.

Further B–R observations of Centaurs are essential. Confirmation of bimodal Centaur colors will bolster models like that of [Wong and Brown \(2016\)](#). On the other hand, refuting bimodal colors for Centaurs will require a rethinking of the important processes acting on Centaurs. The fact that red Centaurs have a smaller orbital inclination angle distribution than the neutral ones ([Tegler et al., 2016](#)), suggesting a distinct origin, is also puzzling when dynamical models say that Centaurs do not preserve their inclinations ([Volk and Malhotra, 2013](#)). Moreover, the recent discovery that Jupiter Trojans and Neptune Trojans possess similar colors, when they were presumably populated by very distinct regions of the primordial planetesimal disk, evidences that we are still missing something crucial ([Jewitt, 2018](#)).




FIG. 14.2 Centaur B–R histograms as they were in 2003 and as they are in 2018 from TRC data. In the larger 2018 sample, the red population became broader and the gap between the gray and red populations became a dip. Note that we are using the more strict definition of Centaurs.

14.4.2.2 Spectra

Although surface colors provide a good proxy to the general surface properties of our objects, the dedicated investigation tool for a comprehensive surface composition study is, of course, reflectance spectroscopy. The technical limit to this method is set by the collecting power of the current large telescopes and the sensitivity of the instruments. With an 8–10 m class telescope, visible spectroscopy with a signal-to-noise ratio (SNR) above ~ 20 is possible only for objects with $V < 20\text{--}22$ mag. Near-infrared (NIR) spectroscopy can be performed at $\text{SNR} \gtrsim 10$ for objects with $J < 18\text{--}20$ mag. Therefore, only a subset of the currently known Centaurs can be explored with this technique.

Visible reflectance spectra ($\lambda \sim 400\text{--}700$ nm) generally show a linear increasing continuum. Most of the ices expected to be present in these regions of the solar system have absorption bands in the NIR. Signatures of water ice, H_2O , are present at 1.5, 1.65, and 2.0 μm ; of methane ice, CH_4 , at 1.7 and 2.3 μm ; of methanol ice, CH_3OH , at 2.27 μm ; and of ammonia ice, NH_3 at 2.0 and 2.25 μm . Unfortunately, observations in the region $\lambda > 3 \mu\text{m}$, where ices of interest have very deep signatures (including also CO and CO_2) are not feasible. However, some predictions on the surface composition have been tempted by [Dalle Ore et al. \(2015\)](#).

(5145) Pholus masterfully opened the era of the spectroscopic exploration of the outer solar system belt and what we know today about other objects is just a briefer story. Since Pholus,

about two dozens of Centaurs were studied with spectrographs from the largest facilities available from the ground. Its first spectrum showed to be linear between 400 nm and 1 μ m, being the steepest red slope detected so far on a solar system body (Fink et al., 1992; Binzel, 1992; Hoffmann et al., 1993). Such slope was incompatible with the conventional silicate or meteoritic analogs that prevailed in asteroids studies at that time, but rather compatible with a mixture of tholins (Khare et al., 1989), suggesting the presence of organic material processed by irradiation. Binzel (1992) suggested that this body has a primitive composition and is at either the beginning or the end state of its thermal evolution. Although on an unstable orbit, Pholus has been undisturbed for at least 10^5 years, that is, long enough to hold, possibly, a thin surface crust of organics due to space weathering. If Pholus just entered the Centaur region, it should start to experience heating and surface processing. In alternative, it might have experienced past cometary activity, maybe on a different orbit in that region, and rather be at its thermal end state with an insulating mantle of organic tholins. Using Hapke (1993) theory, laboratory data, and new observations, Cruikshank et al. (1998) conclude that the red color of Pholus is caused by the irradiation processing of simple molecules (CH_4 , H_2O , CO , NH_3 , N_2 , and CH_3OH) and describe the full spectrum of Pholus with the presence of silicate olivine, carbon, a complex refractory solid (organic molecules), and a mixture of H_2O and CH_3OH (or an equivalent molecule), highlighting that this inventory includes all the basic components of comets. A nice display of Pholus NIR spectrum can be found in Barucci et al. (2011).

Spectroscopically, Centaurs seem to exhibit the same variety of features as their TNO progenitors, except for methane, seen only among the large TNOs. Water ice has been detected on (2060) Chiron, (8405) Asbolus, (10199) Chariklo, (31824) Elatus, (32532) Thereus, (52872) Okyrhoe, (55576) Amycus, and 2007 UM₁₂₆, and methanol has been detected on (5145) Pholus, (54598) Bienor, (83982) Crantor, and 2008 FC₇₆ (see Table 14.1).

14.4.3 Rotations, shapes, and sizes

14.4.3.1 Light curves

The rotational brightness variation, commonly named *light curve*, is the periodic variation of an object brightness as a function of time due to its rotation. The light curve can be produced by various mechanisms: (i) albedo variations on the body surface, (ii) elongated shape, and/or (iii) contact/close binary (e.g., Sheppard and Jewitt, 2002; Lacerda, 2005). Therefore, light curves carry a lot of information about the rotational and physical properties of small bodies.

The first two basic parameters derived from a light curve are the rotational period of the object (P), and the peak-to-peak light curve amplitude (Δm). Assuming a triaxial object in hydrostatic equilibrium, one can estimate the lower limit to its density using the figures of equilibrium from Chandrasekhar (1987). Assuming the light curve is dominated by the shape of the object, one can estimate its elongation assuming a certain viewing angle. A light curve can also be caused by a spherical object with albedo variation(s) on its surface and so one can extract constraints. Finally, a light curve with a U- and V-shape at the maximum and minimum of brightness and $\Delta m > 0.9$ mag is likely due to a contact/close binary, and thus we can infer the binarity of the object and derive the physical characteristics of both system's components (Leone et al., 1984; Lacerda et al., 2014b; Thirouin and Sheppard, 2018).

Only $\sim 5\%$ of the known Centaurs have been observed for complete/partial light curves (Table 14.1) showing, nonetheless, a large variety of amplitudes and rotational periods. From Maxwellian fits the mean rotation period of Centaurs is 8.1 hours, being slower than the mean period of TNOs, that is, 8.45 hours (Thirouin et al., 2016; Thirouin and Sheppard, 2019). The mean amplitude is ~ 0.15 mag. Because of Centaurs relatively short orbital periods, compared to the TNOs, the viewing angle between the rotation axis and the line of sight will change significantly over the course of a couple of years, and thus the light curve amplitude will vary, as seen for Bienor and Pholus (Tegler et al., 2005a; Fernández-Valenzuela et al., 2017). In such cases, it is possible to derive the object's pole orientation. So far, no strong correlations between P and Δm and orbital parameters have been detected among Centaurs (Duffard et al., 2009). But the correlation between light curve amplitude and absolute magnitude (smaller objects are more elongated than the larger ones) noticed among TNOs is not present in the Centaur population which infer for a different evolution/collisional history (Duffard et al., 2009; Thirouin, 2013; Benecchi and Sheppard, 2013).

14.4.3.2 Size distribution

The radius of an object (in kilometer) is given by: $p_R \times R^2 = 2.24 \times 10^{16} \times 10^{0.4 \times (R_\odot - H_R)}$, where $R_\odot$ is the R-filter magnitude of the Sun and p_R and H_R are, respectively, the geometric albedo and the absolute magnitude of the object in the R-filter (Russell, 1916).

Several techniques can be used to derive or constrain the albedo and absolute magnitude and by extension the size of an object. Stellar occultations can be used to derive shape, size, albedo, detection of rings and satellites, and constraints about atmosphere, such as for Chiron and Chariklo (Braga-Ribas et al., 2014; Ruprecht et al., 2015; Ortiz et al., 2015). With data from the Spitzer Space Telescope, the Herschel Space Observatory, the Wide-field Infrared Survey Explorer mission, and the Atacama Large Millimeter Array, it has been possible to perform thermal modeling in order to estimate the radiometric size and albedo (Stansberry et al., 2008; Lellouch et al., 2013, 2017; Bauer et al., 2013; Lacerda et al., 2014a; Duffard et al., 2014). Generally, Centaurs have a low albedo, of a few percent, and only a handful have an albedo up to $\sim 20\%$ (Duffard et al., 2014; Lacerda et al., 2014a; Lellouch et al., 2017). Also, it was noticed that the mean albedo of the red Centaurs is $\sim 8\% - 12\%$, whereas for the gray ones is lower $\sim 5\% - 6\%$ (Bauer et al., 2013; Duffard et al., 2014).

Several surveys have been dedicated to the search of small bodies (e.g., Petit et al., 2008; Trujillo, 2008; Adams et al., 2014; Weryk et al., 2019; Sheppard et al., 2016; Bannister et al., 2018). However, these surveys are mostly focused on the search of TNOs or Near-Earth Asteroids (NEOs) and the discovery of Centaurs is mainly serendipitous. Size distributions have been estimated, but they are not debiased (Bauer et al., 2013), or used either the maximum likelihood technique (Jedicke and Herron, 1997) or the Monte Carlo simulations (Sheppard et al., 2000)—these two, however, find a similar distribution with $\alpha \sim 0.6$. Adams et al. (2014) proposed a debiased size distribution, using DES data, and inferred that the best fit for Centaurs with absolute magnitudes $7.5 < H < 11$ mag. is a power law with $\alpha = 0.42 \pm 0.02$ suggesting a knee in the size distribution around $H = 7.2$ mag. Similarly, using the OSSOS simulator, Lawler et al. (2018) confirmed that a single slope for the size distribution is not able to match the observations, and thus a break is needed at smaller sizes. Such a break is also required to explain the size distribution of the TNOs (Bernstein et al., 2004; Shankman et al., 2013).

TABLE 14.1 B–R colors of “strict” Centaurs from TRC survey and their albedos, light curves studies, and detected ices from other works.

Object	B–R	Albedo	Single peak P (hours)	Double peak P (hours)	Δm (mag)	Ices?	References
29P Schwassmann-Wachmann 1	–	$4.61^{+5.22}_{-1.90}$	–	–	–	–	S08
(2060) 1977 UB Chiron	–	$7.57^{+1.03}_{-0.87}$	–	5.9180 ± 0.0001	0.088 ± 0.003	Water	B89, O15, G16, SS08
(5145) 1992 AD Pholus	1.97 ± 0.11	$15.5^{+7.6}_{-4.9}$	–	9.98	$0.15/0.60$	Methanol	B92, H92, F01, T05, D14
	–	–	–	–	0.15	–	RT99
(7066) 1993 HA ₂ Nessus	–	$8.6^{+7.5}_{-3.4}$	–	–	0.5	–	RT99
	–	–	–	–	<0.2	–	D98, D14
	–	–	5.68 ± 0.19	–	0.76 ± 0.08	–	K06
1994 TA	1.92 ± 0.10	–	–	–	–	–	TR00
(10370) 1995 DW ₂ Hylonome	1.16 ± 0.09	$5.1^{+3.0}_{-1.7}$	–	–	–	–	TR98, RT99, D14
(8405) 1995 GO Asbolus	1.22 ± 0.05	$5.6^{+1.9}_{-1.5}$	–	8.9351 ± 0.003	0.55	Water	D98, K00, D14
	–	–	–	–	0.34	–	RT99
	–	–	–	–	0.14 ± 0.10	–	R07
(10199) 1997 CU ₂₆ Chariklo	1.25 ± 0.05	$5.63^{+0.76}_{-0.65}$	–	–	<0.1	Water	D98, TR98, RT99, S08
	–	$5.59^{+1.69}_{-1.17}$	–	7.004 ± 0.036	0.11	–	F14, G16
(49036) 1998 QM ₁₀₇ Pelion	1.25 ± 0.04	–	–	–	–	–	TR00
(52872) 1998 SG ₃₅ Okyrhoe	1.21 ± 0.02	$5.6^{+1.2}_{-1.0}$	–	16.6	0.2	Water	B03, D14, TR03
	–	–	4.68 or 6.08	–	0.07 ± 0.01	–	T10
(52975) 1998 TF ₃₅ Cyllarus	1.72 ± 0.12	$13.9^{+15.7}_{-6.4}$	–	–	–	–	D14, TRC03
(29981) 1999 TD ₁₀	–	$4.40^{+1.42}_{-0.96}$	–	–	–	–	S08
(31824) 1999 UG ₅ Elatus	1.75 ± 0.05	$4.9^{+2.8}_{-1.6}$	13.25	–	0.24	Water	D14, TR03, G01
		–	13.41 ± 0.04	–	0.102 ± 0.005	–	B02
(60558) 2000 EC ₉₈ Echeclus	1.39 ± 0.04	$5.2^{+0.70}_{-0.71}$	13.401	26.802	0.24 ± 0.06	–	R05, D14

Continued

TABLE 14.1 B–R colors of “strict” Centaurs from TRC survey and their albedos, light curves studies, and detected ices from other works—cont’d

Object	B–R	Albedo	Single peak P (hours)	Double peak P (hours)	Δm (mag)	Ices?	References
2000 FZ ₅₃	1.17 ± 0.05	–	–	–	–	–	TRC03
2000 GM ₁₃₇	–	$4.3^{+2.6}_{-1.6}$	–	–	–	–	D14
(54598) 2000 QC ₂₄₃ Bienor	1.15 ± 0.08	$4.3^{+1.6}_{-1.2}$	4.57 ± 0.02	9.14 ± 0.04	0.08 to 0.75	Methanol	O03, F17, D14
		–	–	9.17	0.34 ± 0.08	–	R07
		–	$(4.723 \text{ or } 4.594) \pm 0.001$	–	0.38 ± 0.02	–	K06
(63252) 2001 BL ₄₁	1.21 ± 0.03	$4.3^{+2.8}_{-1.4}$	–	–	–	–	D09, TRC03
(88269) 2001 KF ₇₇	1.81 ± 0.04	–	–	–	–	–	TRC03
(32532) 2001 PT ₁₃ Thereus	1.18 ± 0.01	$8.3^{+1.6}_{-1.3}$	4.1546 ± 0.0005	8.3091 ± 0.0001	0.16 ± 0.02	Water	O03, D14
	–	–	–	8.34	0.16	–	FD03
	–	–	–	8.34	0.34 ± 0.08	–	R07
	–	–	–	8.4	0.15	–	F01
(119315) 2001 SQ ₇₃	1.13 ± 0.02	$4.8^{+3.0}_{-1.8}$	–	–	–	–	D14, TRC03
(148975) 2001 XA ₂₅₅	–	–	–	–	~ 0.13	–	T13
	–	–	>7	>14	~ 0.2	–	H18
2001 XZ ₂₅₅	1.91 ± 0.07	–	–	–	–	–	TRC03
(42355) 2002 CR ₄₆ Typhon	–	$5.09^{+1.24}_{-0.80}$	9.67	–	0.07 ± 0.01	–	T10
(55567) 2002 GB ₁₀ Amycus	1.82 ± 0.03	$8.3^{+1.6}_{-1.5}$	9.76	–	0.16 ± 0.01	Water	TRC03, T10
(83982) 2002 GO ₉ Crantor	1.86 ± 0.02	$11.8^{+7.09}_{-?}$	$(6.97 \text{ or } 9.67) \pm 0.03$	–	0.14 ± 0.04	Methanol	O03, TR03, S08
	–	–	–	–	0.34	–	RT99
(95626) 2002 GZ ₃₂	1.03 ± 0.04	–	–	5.80 ± 0.03	0.15 ± 0.03	–	D08
(250112) 2002 KY ₁₄	1.75 ± 0.02	–	–	–	–	–	TRC16
(73480) 2002 PN ₃₄	–	$4.25^{+0.83}_{-0.65}$	–	–	–	–	S08
2002 PQ ₁₅₂	1.85 ± 0.05	–	–	–	–	–	TRC16

(523597) 2002 QX ₄₇	1.08 ± 0.04	–	–	–	–	–	TRC16
(119976) 2002 VR ₁₃₀	–	9.3 ^{+6.6} _{–3.6}	–	–	–	–	D14
(120061) 2003 CO ₁	–	4.9 ^{+0.5} _{–0.6}	3.53/4.13/4.99 ± 0.01/6.30	–	0.10 ± 0.05	–	O06, D14
	–	–	4.51	–	0.06 ± 0.01	–	T10
(65489) 2003 FX ₁₂₈ Ceto	–	7.67 ^{+1.38} _{–1.10}	–	4.43 ± 0.03	0.13 ± 0.02	–	D08, S08
(136204) 2003 WL ₇	1.23 ± 0.04	5.3 ± 1.0	8.24	–	0.04 ± 0.01	–	T10, TRC16, D14
2004 QQ ₂₆	–	4.4 ^{+3.9} _{–1.4}	–	–	–	–	D14
(447178) 2005 RO ₄₃	1.24 ± 0.03	5.6 ^{+3.6} _{–2.1}	–	–	–	–	D14, TRC16
(145486) 2005 UJ ₄₃₈	1.64 ± 0.04	25.6 ^{+9.7} _{–7.6}	8.32	–	0.11 ± 0.01	–	T10, TRC16, D14
(248835) 2006 SX ₃₆₈	1.22 ± 0.02	5.2 ^{+0.7} _{–0.6}	–	–	–	–	TRC16, D14
(309139) 2006 XQ ₅₁	1.15 ± 0.03	–	–	–	–	–	TRC16
(341275) 2007 RG ₂₈₃	1.26 ± 0.03	–	–	–	–	–	TRC16
2007 RH ₂₈₃	1.15 ± 0.03	–	–	–	–	–	TRC16
2007 TK ₄₂₂	1.22 ± 0.04	–	–	–	–	–	TRC16
(25012) 2007 UL ₁₂₆	1.75 ± 0.02	5.7 ^{+1.1} _{–0.7}	3.56 or 4.2	7.12 or 8.4	0.11 ± 0.01	–	T10
2007 UM ₁₂₆	1.13 ± 0.03	–	–	–	–	Water	TRC16
2007 VH ₃₀₅	1.18 ± 0.02	–	–	–	–	–	TRC16
(281371) 2008 FC ₇₆	1.60 ± 0.03	6.7 ^{+1.7} _{–1.1}	–	–	~0.1	Methanol	T13, TRC16, D14
	–	–	5.93 ± 0.05	11.86 ± 0.05	0.04 ± 0.01	–	H18
(315898) 2008 QD ₄	1.20 ± 0.02	–	–	–	~0.09	–	T13, TRC16
	–	–	>7	>14	~0.15	–	H18
(309737) 2008 SJ ₂₃₈	1.60 ± 0.02	–	–	–	–	–	TRC16

Continued

TABLE 14.1 B–R colors of “strict” Centaurs from TRC survey and their albedos, light curves studies, and detected ices from other works—cont’d

Object	B–R	Albedo	Single peak P (hours)	Double peak P (hours)	Δm (mag)	Ices?	References
(309741) 2008 UZ ₆	1.52 ± 0.04	–	–	–	–	–	TRC16
(342842) 2008 YB ₃	1.23 ± 0.02	–	–	–	~0.18	–	T13, TRC16
(346889) 2009 QV ₃₈ Rhiphonos	1.37 ± 0.02	–	–	–	–	–	TRC16
(349933) 2009 YF ₇	1.18 ± 0.03	–	–	–	–	–	TRC16
2010 BK ₁₁₈	–	–	–	–	~0.15	–	T13
(382004) 2010 RM ₆₄	1.56 ± 0.02	–	–	–	–	–	TRC16
2010 TH	1.18 ± 0.03	–	–	–	–	–	TRC16
2010 TY ₅₃	–	–	–	–	<0.14	–	BS13
(471339) 2011 ON ₄₅	1.81 ± 0.04	–	–	–	–	–	TRC16
(449097) 2012 UT ₆₈	1.68 ± 0.03	–	–	–	–	–	TRC16
(463368) 2012 VU ₈₅	1.70 ± 0.07	–	–	–	–	–	TRC16
(523676) 2013 UL ₁₀	1.17 ± 0.03	–	–	–	–	–	TRC16
(459865) 2013 XZ ₈	1.17 ± 0.03	–	–	–	–	–	TRC16
(459971) 2014 ON ₆	1.55 ± 0.03	–	–	–	–	–	TRC16

Notes: Tegler, Romanishin, and Consolmagno’s (TRC) survey: <http://www.physics.nau.edu/~teglar/research/survey.htm>.
B89: Bus et al. (1989); B92: Buie and Bus (1992); B02: Bauer et al. (2002); B03: Bauer et al. (2003); BS13: Benecchi and Sheppard (2013); D98: Davies et al. (1998); D08: Dotto et al. (2008); D09: Duffard et al. (2009); D14: Duffard et al. (2014); F01: Farnham (2001); F14: Fornasier et al. (2014); F17: Fernández et al. (2017); FD03: Farnham and Davies (2003); G01: Gutiérrez et al. (2001); G16: Galiazzo et al. (2016); H92: Hoffmann et al. (1992); H18: Hromakina et al. (2018); K00: Kern et al. (2000); K06: Kern (2006); O03: Ortiz et al. (2003); O06: Ortiz et al. (2006); O15: Ortiz et al. (2015); R05: Rousselot et al. (2005); R07: Rabinowitz et al. (2007); RT99: Romanishin and Tegler (1999); S08: Stansberry et al. (2008); SJ02: Sheppard and Jewitt (2002); T05: Tegler et al. (2005b); T10: Thirouin et al. (2010); T13: Thirouin (2013); TR00: Tegler and Romanishin (2000); TR03: Tegler and Romanishin (2003); TRC03: Tegler et al. (2003); TRC16: Tegler et al. (2016).

14.4.4 Moons and rings

Binary and multiple systems have been found across many minor body populations. The Centaurs are no exception with two known binary systems: Ceto/Phorcys—formerly, (65489) 2003 FX₁₂₈—and Typhon/Echidna—formerly, (42355) 2002 CR₄₆. In the case of Ceto/Phorcys, the components have a radii of 87 and 66 km and are separated by about 1800 km and the system has a density of $\sim 1.4 \text{ g cm}^{-3}$ (Grundy et al., 2007). Typhon/Echidna has a similar semimajor axis of ~ 1600 km and the radii of the components are 76 and 42 km, with a density of only 0.44 g cm^{-3} (Grundy et al., 2008). Despite finding many binary systems among TNOs, only two were found among Centaurs and there is no evidence of contact binary Centaurs so far. But, it is important to keep in mind that the Centaurs have not been the topic of any large-scale survey for light curve studies.

Recently, to everyone's surprise, thanks to stellar occultations, a ring system was discovered around the largest Centaur: Chariklo (Braga-Ribas et al., 2014). Then, Ortiz et al. (2015), Ruprecht et al. (2015), and Fernández-Valenzuela et al. (2017) evaluated the possibility of ring systems around Chiron and Bienor. The formation and stability of ring systems around objects in the size scale of ~ 100 km is still a mystery. Several hypotheses have been considered to explain the formation of these rings: debris disks from a rotational fission, debris disks due to the disruption of a satellite because of impact, and/or tidal forces among other options (Hyodo et al., 2016; Pan and Wu, 2016). Finally, their stability is also an open question (Sicardy et al., 2018).

14.4.5 Cometary activity

As already mentioned, Chiron, the first object identified as a Centaur, was found to exhibit magnitude variations on short timescales. That the first Centaur discovered was an active cometary body was no coincidence. As we saw in Section 14.3, the dynamical evidence for a link between Centaurs and JFCs is compelling.

From the ~ 290 – 360 Centaurs known, depending on the classification, 29 have shown evidence of activity, that is, $\sim 8\%$ – 9% . In addition to the dynamical evidence, the Centaur population has the largest fraction of objects with confirmed activity relative to other populations, besides those of actual comet populations, evidently. Neither main-belt comets—a.k.a. active asteroids—nor NEOs approach the fraction of active bodies embedded within their dynamically defined populations that the Centaurs do. Many of the unique attributes of the Centaur population may be linked with their cometary activity and behavior, and may suggest that activity among the Centaurs is more common than the fraction observed at a given epoch. Several attempts have been made to link the bimodal color distribution with current and past activity of Centaurs as we saw in Section 14.4.2.1. A significant color-based correlation with Centaurs' albedos has been established (Stansberry et al., 2008; Tegler et al., 2008; Bauer et al., 2013; Lacerda et al., 2014a). The darker population matching the neutrally colored populations marks a convergence similar to comet surfaces. In fact, all small body populations nearer the Sun are composed entirely of dark and neutrally colored objects. A plausible explanation for the lack of red and bright surfaces among Centaurs and JFCs is the destruction of these very same red surfaces when entering the inner solar system over a lifetime comparable to that of Centaurs (Jewitt, 2002), which may even take less than 10 years when Centaurs get $q < \sim 10$ au

(Jewitt, 2015), but TNOs show that something more complex is going on. As if doubts existed still, the match between Centaur and JFC size distribution slopes (Fernández et al., 2013; Bauer et al., 2013) also evidences a probably similar origin or evolution.

In fact, technically, Chiron was not the first Centaur discovered, rather, that distinction falls to a more active body, 29P/Schwassmann-Wachmann 1, discovered in 1927. The pattern of behavior of active Centaurs varies with each Centaur. While Chiron's activity varies over large timescales (Duffard et al., 2002, and references therein) Echeclus' outbursts are shorter lived, but more periodic (Choi et al., 2006; Jaeger et al., 2011; Seccull et al., 2019), while 29P's activity is constant, though punctuated with outbursts. Jewitt (2009, 2015) identified that active Centaurs tend to have lower perihelion (see Fig. 14.1), though notable exceptions, like Chiron, exist. In fact, Fernández et al. (2018) found that active and inactive Centaurs have experienced different dynamical evolutions. They suggest that inactive Centaurs deactivated because of a larger permanence far from the Sun, while a recent drastic drop in perihelion distance could be responsible for renewing the activity in the active ones. The brightness variations of the other 26 are lower but morphological features are still identifiable (Jewitt, 2009). Brightness variations and extended morphology are not the sole indicators of possible activity. Larger bodies may exhibit temporary ring structures and dust grain atmospheres as we have seen over Chariklo and Chiron. No outburst has yet been detected on Chariklo but, as mentioned in Section 14.4.4, such kind of events may both be the origin and eventual demise of the rings. The structure surrounding Chiron could also be a manifestation of, or evolve from, a ballistic dust atmosphere (Meech and Belton, 1990).

The activity in Centaurs tends to be recurrent but, unlike standard cometary behavior, Echeclus, 29P, and Chiron all exhibit outbursts and coma decoupled from the Centaur's perihelion. The mechanism that triggers the sublimation is not clear. It may be that localized regions of surface ice are directly exposed through a combination of seasonal and diurnal variations on such bodies, rather than gases being released from the subsurface, or that volatiles may be driven from water ice via the amorphous-crystalline transition as the thermal wave reaches localized subsurface water ice reservoirs (see Prialnik, 1992; Jewitt, 2009, and Section 14.4.6). Dynamical models show that excursions of Centaurs over the inner solar system and subsequent returns to Centaur orbits are common (e.g., Horner et al., 2004a). Deriving any notion of how volatiles, or hypervolatiles, are distributed on Centaurs must accommodate the perseverance of a fraction of the ices throughout such terrestrial-planet-region excursions, possibly in subsurface reservoirs that may later be exposed.

14.4.6 Interiors

The internal structure of Centaurs is, in theory, the result of processes that occurred at three different stages of their history (see Prialnik et al., 2008; Coradini et al., 2008; Sarid and Prialnik, 2009; Guilbert-Lepoutre et al., 2015, for reviews). First, the internal structure may reflect the formation of Centaurs themselves, a process which is not fully constrained yet. Some Centaurs could be primordial aggregates of planetesimals relatively unaltered by collisions (Belton, 2014), while some others could be the outcomes of disruptive collision of larger bodies. Indeed, Stern (1996) and Davis and Farinella (1997) suggested that JFCs—therefore, Centaurs too—could be fragments from parent bodies with initial sizes up to ~100 km. The “outbursty”

activity of Echeclus could be interpreted as being due to internal heterogeneities, possibly reflecting its rubble-pile nature (either primordial or collisional, see [Rousselot et al., 2016](#)). In fact, [Rosenberg and Prialnik \(2010\)](#) showed that internal heterogeneities could potentially be detected through the activity pattern of a comet, where erratic behaviors would be induced by blocks of varying thermophysical properties randomly distributed inside the nucleus.

The second stage of processing is related to radiogenic heating, especially their early heating. The effectiveness of ^{26}Al in heating the interior of an icy object strongly depends on the time delay between its formation and the formation of calcium-aluminum-rich inclusions (CAIs) (see [Prialnik and Podolak, 1995](#); [Prialnik et al., 2008](#)), which remains unknown. [Prialnik et al. \(1987\)](#) found that early radiogenic heating by ^{26}Al could lead to the complete crystallization of amorphous ice inside comets and Centaurs, and even melting of large nuclei. Therefore, the early thermal evolution of Centaurs could have led to phase transitions and chemical differentiation occurring mostly in the core, while an outer layer might remain pristine. We should stress that the internal structure inherited from the first two stages remain speculative: the effects of accretion could be inferred from size, shape, or within the activity pattern, but most observations cannot provide us with this information. Early heating should have mostly affected the core of icy objects, which we cannot access from remote observations of their surface. In addition, the third stage, dominated by cometary activity, could erase many of the effects of the two previous stages.

Whatever stratigraphy is produced after these early stages of formation and evolution, the internal structure of Centaurs should be further modified by cometary activity. Water ice is thermodynamically stable in the giant planet region, that is, it does not sublime, hence it cannot drive the activity of Centaurs. [Jewitt \(2009\)](#) suggested that crystalline-amorphous water ice transition, releasing trapped volatiles, could be the source of Centaurs activity. [Guilbert-Lepoutre \(2012\)](#) studied this scenario concluding that: (i) crystallization at the surface of Centaurs can be triggered at heliocentric distances as large as 16 AU; (ii) cometary activity due to crystallization could be triggered up to $\sim 10\text{--}12$ AU, which is broadly consistent with the observations of active Centaurs; and (iii) as the crystallization front propagates from the surface toward the interior of Centaurs, cometary activity should stop in a relatively short period of time—outgassing is typically sustained for hundreds to thousands of years. The simple fact that active Centaurs do exist shows that a chemical differentiation, due to thermophysical processing, is happening in the giant planet region.

14.5 Discussion

If Centaurs were discovered later than TNOs, they probably would not have been named as a class and would be just seen as low perihelion SDOs. Yet, they do not cease to surprise and puzzle us revealing themselves to be as rich as TNOs alone.

Identifying the important large-scale processes in the outer solar system is of the utmost importance to understand how the entire solar system has formed and how it has evolved. Centaurs are an essential class of minor bodies to help us do so. No longer simply because they are closer, and therefore brighter, than TNOs, allowing for more detailed observational studies, but also because they exhibit peculiar and distinct behaviors that seem to be at the

core of the deep physical and chemical changes that affect icy bodies when entering the inner solar system.

The condition of transition phase between TNOs and JFCs attributed to Centaurs seems unquestionable. However, if dynamical models predict that orbital inclinations are not preserved among Centaurs, how come red Centaurs have a smaller orbital inclination angle distribution than the neutral ones? Are dynamical models that reliable? If Centaurs are scattered inwards from the Trans-Neptunian region through interaction with Neptune, how can they possibly have a different collisional history than TNOs? Indeed puzzling, but only with more data we can make it indisputable. There are no red surfaces among JFCs, therefore, red Centaurs must change their colors when becoming JFCs. Is it not a remarkable coincidence that neutral Centaurs, that should have formed beyond Jupiter in the primordial disk, do not change their colors when becoming JFCs? Too remarkable to leave us comfortable. Models for scenarios of distinct compositions over the primordial planetesimal disk of icy bodies seem increasingly more sound and refined, and yet the evidence for the two distinct color groups of Centaurs has faded. Is it not a paradox? Bearing in mind the principle of explosion, that is, from a false premise anything follows, models, however, elegant they may be, need to be evidence based. It is through observational data they can be demonstrated or refuted. Paradoxically, about 20 years ago we had a seemingly elegant model capable of explaining a continuous color distribution and colors started to appear noncontinuous, whereas today colors might be continuous and we have an apparently elegant model capable of explaining their discontinuity. Confirming or refuting is imperative. Consensus is not a demonstration.

Acknowledgments

CITEUC is funded by National Funds through FCT—Foundation for Science and Technology (project: UID/Multi/00611/2019) and FEDER—European Regional Development Fund through COMPETE 2020—Operational Program Competitiveness and Internationalization (project: POCI-01-0145-FEDER-006922). Audrey Thirouin is partly funded by Lowell Observatory funds and by the National Science Foundation, grant number AST-1734484.

References

- Adams, E.R., Gulbis, A.A.S., Elliot, J.L., Benecchi, S.D., Buie, M.W., Trilling, D.E., Wasserman, L.H., 2014. De-biased populations of Kuiper belt objects from the Deep Ecliptic Survey. *Astron. J.* 148, 55. <https://doi.org/10.1088/0004-6256/148/3/55>.
- Araujo, R.A.N., Galiazzo, M.A., Winter, O.C., Sfair, R., 2018. The journey of Typhon-Echidna as a binary system through the planetary region. *Mon. Not. R. Astron. Soc.* 476, 5323–5331. <https://doi.org/10.1093/mnras/sty583>.
- Bailey, B.L., Malhotra, R., 2009. Two dynamical classes of Centaurs. *Icarus* 203, 155–163. <https://doi.org/10.1016/j.icarus.2009.03.044>.
- Bannister, M.T., Gladman, B.J., Kavelaars, J.J., Petit, J.-M., Volk, K., Chen, Y.-T., Alexandersen, M., Gwyn, S.D.J., Schwamb, M.E., Ashton, E., Benecchi, S.D., Cabral, N., Dawson, R.I., Delsanti, A., Fraser, W.C., Granvik, M., Greenstreet, S., Guilbert-Lepoutre, A., Ip, W.-H., Jakubik, M., Jones, R.L., Kaib, N.A., Lacerda, P., Van Laerhoven, C., Lawler, S., Lehner, M.J., Lin, H.W., Lykawka, P.S., Marsset, M., Murray-Clay, R., Pike, R.E., Rousselot, P., Shankman, C., Thirouin, A., Vernazza, P., Wang, S.-Y., 2018. OSSOS. VII. 800+ Trans-Neptunian objects—the complete data release. *Astrophys. J. Suppl. Ser.* 236, 18. <https://doi.org/10.3847/1538-4365/aab77a>.
- Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R., 2008. *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ.

- Barucci, M.A., Alvarez-Candal, A., Merlin, F., Belskaya, I.N., de Bergh, C., Perna, D., DeMeo, F., Fornasier, S., 2011. New insights on ices in Centaur and Transneptunian populations. *Icarus* 214, 297–307. <https://doi.org/10.1016/j.icarus.2011.04.019>.
- Bauer, J.M., Meech, K.J., Fernández, Y.R., Farnham, T.L., Roush, T.L., 2002. Observations of the Centaur 1999 UG₅: evidence of a unique outer solar system surface. *Publ. Astron. Soc. Pac.* 114, 1309–1321. <https://doi.org/10.1086/344586>.
- Bauer, J.M., Meech, K.J., Fernández, Y.R., Pittichova, J., Hainaut, O.R., Boehnhardt, H., Delsanti, A.C., 2003. Physical survey of 24 Centaurs with visible photometry. *Icarus* 166, 195–211. <https://doi.org/10.1016/j.icarus.2003.07.004>.
- Bauer, J.M., Grav, T., Blauvelt, E., Mainzer, A.K., Masiero, J.R., Stevenson, R., Kramer, E., Fernández, Y.R., Lisse, C.M., Cutri, R.M., Weissman, P.R., Dailey, J.W., Masci, F.J., Walker, R., Waszczak, A., Nugent, C.R., Meech, K.J., Lucas, A., Pearman, G., Wilkins, A., Watkins, J., Kulkarni, S., Wright, E.L., WISE Team, PTF Team, 2013. Centaurs and scattered disk objects in the thermal infrared: analysis of WISE/NEOWISE observations. *Astrophys. J.* 773, 22. <https://doi.org/10.1088/0004-637X/773/1/22>.
- Belton, M.J.S., 2014. The size-distribution of scattered disk TNOs from that of JFCs between 0.2 and 15 km effective radius. *Icarus* 231, 168–182. <https://doi.org/10.1016/j.icarus.2013.12.001>.
- Benecchi, S.D., Sheppard, S.S., 2013. Light curves of 32 large transneptunian objects. *Astron. J.* 145, 124. <https://doi.org/10.1088/0004-6256/145/5/124>.
- Bernstein, G.M., Trilling, D.E., Allen, R.L., Brown, M.E., Holman, M., Malhotra, R., 2004. The size distribution of Trans-Neptunian bodies. *Astron. J.* 128, 1364–1390. <https://doi.org/10.1086/422919>.
- Binzel, R.P., 1992. The optical spectrum of 5145 Pholus. *Icarus* 99, 238–240. [https://doi.org/10.1016/0019-1035\(92\)90185-A](https://doi.org/10.1016/0019-1035(92)90185-A).
- Braga-Ribas, F., Sicardy, B., Ortiz, J.L., Snodgrass, C., Roques, F., Vieira-Martins, R., Camargo, J.I.B., Assafin, M., Duffard, R., Jehin, E., Pollock, J., Leiva, R., Emilio, M., Machado, D.I., Colazo, C., Lellouch, E., Skottfelt, J., Gillon, M., Ligier, N., Maquet, L., Benedetti-Rossi, G., Gomes, A.R., Kervella, P., Monteiro, H., Sfair, R., El Moutamid, M., Tancredi, G., Spagnotto, J., Mauray, A., Morales, N., Gil-Hutton, R., Roland, S., Ceretta, A., Gu, S.-H., Wang, X.-B., Harpsøe, K., Rabus, M., Manfroid, J., Opitom, C., Vanzi, L., Mehret, L., Lorenzini, L., Schneider, E.M., Melia, R., Lecacheux, J., Colas, F., Vachier, F., Widemann, T., Almenares, L., Sandness, R.G., Char, F., Perez, V., Lemos, P., Martinez, N., Jørgensen, U.G., Dominik, M., Roig, F., Reichart, D.E., Lacluyze, A.P., Haislip, J.B., Ivarsen, K.M., Moore, J.P., Frank, N.R., Lambas, D.G., 2014. A ring system detected around the Centaur (10199) Chariklo. *Nature* 508, 72–75. <https://doi.org/10.1038/nature13155>.
- Brasser, R., Schwamb, M.E., Lykawka, P.S., Gomes, R.S., 2012. An Oort cloud origin for the high-inclination, high-perihelion Centaurs. *Mon. Not. R. Astron. Soc.* 420, 3396–3402. <https://doi.org/10.1111/j.1365-2966.2011.20264.x>.
- Brown, M.E., Schaller, E.L., Fraser, W.C., 2011. A hypothesis for the color diversity of the Kuiper belt. *Astrophys. J.* 739, L60. <https://doi.org/10.1088/2041-8205/739/2/L60>.
- Brunini, A., 2014. On the dynamical evolution and end states of binary Centaurs. *Mon. Not. R. Astron. Soc.* 437, 2297–2302. <https://doi.org/10.1093/mnras/stt2045>.
- Buie, M.W., Bus, S.J., 1992. Physical observations of (5145) Pholus. *Icarus* 100, 288–294. [https://doi.org/10.1016/0019-1035\(92\)90101-C](https://doi.org/10.1016/0019-1035(92)90101-C).
- Bus, S.J., Bowell, E., Harris, A.W., Hewitt, A.V., 1989. 2060 Chiron: CCD and electronographic photometry. *Icarus* 77, 223–238. [https://doi.org/10.1016/0019-1035\(89\)90087-0](https://doi.org/10.1016/0019-1035(89)90087-0).
- Chandrasekhar, S., 1987. *Ellipsoidal Figures of Equilibrium*. Dover, New York, NY.
- Choi, Y.-J., Weissman, P., Chesley, S., Bauer, J., Stansberry, J., Tegler, S., Romanishin, W., Consolmagno, G., 2006. Comet 174P/Echeclus. *Cent. Bureau Electron. Telegr.* 563, 1.
- Coradini, A., Capria, M.T., de Sanctis, M.C., McKinnon, W.B., 2008. The structure of Kuiper belt bodies: link with comets. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 243–256.
- Cruikshank, D.P., Roush, T.L., Bartholomew, M.J., Geballe, T.R., Pendleton, Y.J., White, S.M., Bell, J.F., Davies, J.K., Owen, T.C., de Bergh, C., Tholen, D.J., Bernstein, M.P., Brown, R.H., Tryka, K.A., Dalle Ore, C.M., 1998. The composition of centaur 5145 Pholus. *Icarus* 135, 389–407. <https://doi.org/10.1006/icar.1998.5997>.
- Dalle Ore, C.M., Barucci, M.A., Emery, J.P., Cruikshank, D.P., de Bergh, C., Roush, T.L., Perna, D., Merlin, F., Dalle Ore, L.V., 2015. The composition of “ultra-red” TNOs and centaurs. *Icarus* 252, 311–326. <https://doi.org/10.1016/j.icarus.2015.01.014>.
- Davies, J. (Ed.), 2001. *Beyond Pluto*. Cambridge University Press, Cambridge, p. 246.

- Davies, J.K., Barrera, L.H. (Eds.), 2004. The First Decadal Review of the Edgeworth-Kuiper Belt. Kluwer Academic Publishers, Dordrecht (Reprinted from: *Earth Moon Planets* 92 (1–4), 2003).
- Davies, J.K., McBride, N., Ellison, S.L., Green, S.F., Ballantyne, D.R., 1998. Visible and infrared photometry of six centaurs. *Icarus* 134, 213–227. <https://doi.org/10.1006/icar.1998.5931>.
- Davis, D.R., Farinella, P., 1997. Collisional evolution of Edgeworth-Kuiper belt objects. *Icarus* 125, 50–60. <https://doi.org/10.1006/icar.1996.5595>.
- de Elía, G.C., Brunini, A., di Sisto, R.P., 2008. Collisional and dynamical evolution of Plutinos. *Astron. Astrophys.* 490, 835–842. <https://doi.org/10.1051/0004-6361:200809865>.
- de la Fuente Marcos, C., de la Fuente Marcos, R., 2014. Large retrograde centaurs: visitors from the Oort cloud? *Astrophys. Space Sci.* 352, 409–419. <https://doi.org/10.1007/s10509-014-1993-9>.
- Delsanti, A., Hainaut, O., Jourdeuil, E., Meech, K.J., Boehnhardt, H., Barrera, L., 2004. Simultaneous visible-near IR photometric study of Kuiper belt object surfaces with the ESO/Very Large Telescopes. *Astron. Astrophys.* 417, 1145–1158. <https://doi.org/10.1051/0004-6361:20034182>.
- Di Sisto, R.P., Brunini, A., 2007. The origin and distribution of the Centaur population. *Icarus* 190, 224–235. <https://doi.org/10.1016/j.icarus.2007.02.012>.
- Di Sisto, R.P., Ramos, X.S., Gallardo, T., 2019. The dynamical evolution of escaped Jupiter Trojan asteroids, link to other minor body populations. *Icarus* 319, 828–839. <https://doi.org/10.1016/j.icarus.2018.10.029>.
- Dotto, E., Perna, D., Barucci, M.A., Rossi, A., de Bergh, C., Doressoundiram, A., Fornasier, S., 2008. Rotational properties of Centaurs and Trans-Neptunian objects, lightcurves and densities. *Astron. Astrophys.* 490, 829–833. <https://doi.org/10.1051/0004-6361:200809615>.
- Duffard, R., Lazzaro, D., Pinto, S., Carvano, J., Angeli, C., Alvarez-Candal, A., Fernández, S., 2002. New activity of Chiron: results from 5 years of photometric monitoring. *Icarus* 160, 44–51. <https://doi.org/10.1006/icar.2002.6938>.
- Duffard, R., Ortiz, J.L., Thirouin, A., Santos-Sanz, P., Morales, N., 2009. Transneptunian objects and Centaurs from light curves. *Astron. Astrophys.* 505, 1283–1295. <https://doi.org/10.1051/0004-6361/200912601>.
- Duffard, R., Pinilla-Alonso, N., Santos-Sanz, P., Vilenius, E., Ortiz, J.L., Mueller, T., Fornasier, S., Lellouch, E., Mommert, M., Pal, A., Kiss, C., Mueller, M., Stansberry, J., Delsanti, A., Peixinho, N., Trilling, D., 2014. “TNOs are Cool”: a survey of the Trans-Neptunian region. XI. A Herschel-PACS view of 16 Centaurs. *Astron. Astrophys.* 564, A92. <https://doi.org/10.1051/0004-6361/201322377>.
- Duncan, M.J., Levison, H.F., 1997. A scattered comet disk and the origin of Jupiter Family Comets. *Science* 276, 1670–1672. <https://doi.org/10.1126/science.276.5319.1670>.
- Duncan, M., Quinn, T., Tremaine, S., 1988. The origin of short-period comets. *Astrophys. J.* 328, L69. <https://doi.org/10.1086/185162>.
- Edgeworth, K.E., 1943. The evolution of our planetary system. *J. Br. Astron. Assoc.* 53, 181–188.
- Edgeworth, K.E., 1949. The origin and evolution of the solar system. *Mon. Not. R. Astron. Soc.* 109, 600–609. <https://doi.org/10.1093/mnras/109.5.600>.
- Elliot, J.L., Kern, S.D., Clancy, K.B., Gulbis, A.A.S., Millis, R.L., Buie, M.W., Wasserman, L.H., Chiang, E.I., Jordan, A.B., Trilling, D.E., Meech, K.J., 2005. The deep ecliptic survey: a search for Kuiper belt objects and Centaurs. II. Dynamical classification, the Kuiper belt plane, and the core population. *Astron. J.* 129, 1117–1162. <https://doi.org/10.1086/427395>.
- Farnham, T.L., 2001. The rotation axis of the Centaur 5145 Pholus. *Icarus* 152, 238–245. <https://doi.org/10.1006/icar.2001.6656>.
- Farnham, T.L., Davies, J.K., 2003. The rotational and physical properties of the Centaur (32532) 2001 PT₁₃. *Icarus* 164, 418–427. [https://doi.org/10.1016/S0019-1035\(03\)00141-6](https://doi.org/10.1016/S0019-1035(03)00141-6).
- Fernández, J.A., 1980. On the existence of a comet belt beyond Neptune. *Mon. Not. R. Astron. Soc.* 192, 481–491. <https://doi.org/10.1093/mnras/192.3.481>.
- Fernández, Y.R., Kelley, M.S., Lamy, P.L., Toth, I., Groussin, O., Lisse, C.M., A’Hearn, M.F., Bauer, J.M., Campins, H., Fitzsimmons, A., Licandro, J., Lowry, S.C., Meech, K.J., Pittichová, J., Reach, W.T., Snodgrass, C., Weaver, H.A., 2013. Thermal properties, sizes, and size distribution of Jupiter-family cometary nuclei. *Icarus* 226, 1138–1170. <https://doi.org/10.1016/j.icarus.2013.07.021>.
- Fernández, J.A., Licandro, J., Moreno, F., Sosa, A., Cabrera-Lavers, A., de León, J., Birtwhistle, P., 2017. Physical and dynamical properties of the anomalous comet 249P/LINEAR. *Icarus* 295, 34–45. <https://doi.org/10.1016/j.icarus.2017.04.004>.

- Fernández, J.A., Helal, M., Gallardo, T., 2018. Dynamical evolution and end states of active and inactive Centaurs. *Planet. Space Sci.* 158, 6–15. <https://doi.org/10.1016/j.pss.2018.05.013>.
- Fernández-Valenzuela, E., Ortiz, J.L., Duffard, R., Morales, N., Santos-Sanz, P., 2017. Physical properties of centaur (54598) Bienor from photometry. *Mon. Not. R. Astron. Soc.* 466, 4147–4158. <https://doi.org/10.1093/mnras/stw3264>.
- Ferrari, C., 2018. Thermal properties of icy surfaces in the outer solar system. *Space Sci. Rev.* 214, 111. <https://doi.org/10.1007/s11214-018-0546-x>.
- Fink, U., Hoffmann, M., Grundy, W., Hicks, M., Sears, W., 1992. The steep red spectrum of 1992 AD: an asteroid covered with organic material? *Icarus* 97, 145–149. [https://doi.org/10.1016/0019-1035\(92\)90064-E](https://doi.org/10.1016/0019-1035(92)90064-E).
- Fornasier, S., Lazzaro, D., Alvarez-Candal, A., Snodgrass, C., Tozzi, G.P., Carvano, J.M., Jimenez-Teja, Y., Silva, J.S., Bramich, D.M., 2014. The Centaur 10199 Chariklo: investigation into rotational period, absolute magnitude, and cometary activity. *Astron. Astrophys.* 568, L11. <https://doi.org/10.1051/0004-6361/201424439>.
- Fraser, W.C., Brown, M.E., 2012. The Hubble wide field camera 3 test of surfaces in the outer solar system: the compositional classes of the Kuiper belt. *Astrophys. J.* 749, 33. <https://doi.org/10.1088/0004-637X/749/1/33>.
- Galiazzo, M., de la Fuente Marcos, C., de la Fuente Marcos, R., Carraro, G., Maris, M., Montalto, M., 2016. Photometry of Centaurs and Trans-Neptunian objects: 2060 Chiron (1977 UB), 10199 Chariklo (1997 CU₂₆), 38628 Huya (2000 EB₁₇₃), 28978 Ixion (2001 KX₇₆), and 90482 Orcus (2004 DW). *Astrophys. Space Sci.* 361, 212. <https://doi.org/10.1007/s10509-016-2801-5>.
- Gladman, B., Marsden, B.G., Vanlaerhoven, C., 2008. Nomenclature in the outer solar system. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 43–57.
- Grundy, W.M., Stansberry, J.A., Noll, K.S., Stephens, D.C., Trilling, D.E., Kern, S.D., Spencer, J.R., Cruikshank, D.P., Levison, H.F., 2007. The orbit, mass, size, albedo, and density of (65489) Ceto/Phorcys: a tidally-evolved binary Centaur. *Icarus* 191, 286–297. <https://doi.org/10.1016/j.icarus.2007.04.004>.
- Grundy, W.M., Noll, K.S., Virtanen, J., Muinonen, K., Kern, S.D., Stephens, D.C., Stansberry, J.A., Levison, H.F., Spencer, J.R., 2008. (42355) Typhon Echidna: scheduling observations for binary orbit determination. *Icarus* 197, 260–268. <https://doi.org/10.1016/j.icarus.2008.04.004>.
- Guilbert-Lepoutre, A., 2012. Survival of amorphous water ice on centaurs. *Astron. J.* 144, 97. <https://doi.org/10.1088/0004-6256/144/4/97>.
- Guilbert-Lepoutre, A., Besse, S., Mousis, O., Ali-Dib, M., Höfner, S., Koschny, D., Hager, P., 2015. On the evolution of comets. *Space Sci. Rev.* 197, 271–296. <https://doi.org/10.1007/s11214-015-0148-9>.
- Gutiérrez, P.J., Ortiz, J.L., Alexandrino, E., Roos-Serote, M., Doressoundiram, A., 2001. Short term variability of Centaur 1999 UG5. *Astron. Astrophys.* 371, L1–L4. <https://doi.org/10.1051/0004-6361:20010418>.
- Hahn, G., Bailey, M.E., 1990. Rapid dynamical evolution of giant Comet Chiron. *Nature* 348, 132–136. <https://doi.org/10.1038/348132a0>.
- Hainaut, O.R., Delsanti, A.C., 2002. Colors of minor bodies in the outer solar system. A statistical analysis. *Astron. Astrophys.* 389, 641–664. <https://doi.org/10.1051/0004-6361:20020431>.
- Hainaut, O., Smette, A., Zeilinger, W., 1992. 1992 AD. *Int. Astron. Union Circ.* 5450.
- Hapke, B., 1993. Theory of reflectance and emittance spectroscopy. In: *Topics in Remote Sensing*. Cambridge University Press, Cambridge.
- Hartmann, W.K., Tholen, D.J., Meech, K.J., Cruikshank, D.P., 1990. 2060 Chiron: colorimetry and cometary behavior. *Icarus* 83, 1–15. [https://doi.org/10.1016/0019-1035\(90\)90002-Q](https://doi.org/10.1016/0019-1035(90)90002-Q).
- Hoffmann, M., Fink, U., Grundy, W.M., Hicks, M., 1992. Photometric and spectroscopic observations of 5145 1992 AD, In: Brahic, A., Gerard, J.-C., Surdej, J. (Eds.), *Liege International Astrophysical Colloquia*, vol. 30, p. 203.
- Hoffmann, M., Fink, U., Grundy, W.M., Hicks, M., 1993. Photometric and spectroscopic observations of 5145 Pholus. *J. Geophys. Res.* 98, 7403–7407. <https://doi.org/10.1029/93JE00120>.
- Horner, J., Evans, N.W., 2006. The capture of Centaurs as Trojans. *Mon. Not. R. Astron. Soc.* 367, L20–L23. <https://doi.org/10.1111/j.1745-3933.2006.00131.x>.
- Horner, J., Lykawka, P.S., 2010. The Neptune Trojans—a new source for the Centaurs? *Mon. Not. R. Astron. Soc.* 402, 13–20. <https://doi.org/10.1111/j.1365-2966.2009.15702.x>.
- Horner, J., Evans, N.W., Bailey, M.E., 2004a. Simulations of the population of Centaurs—I. The bulk statistics. *Mon. Not. R. Astron. Soc.* 354, 798–810. <https://doi.org/10.1111/j.1365-2966.2004.08240.x>.
- Horner, J., Evans, N.W., Bailey, M.E., 2004b. Simulations of the population of Centaurs—II. Individual objects. *Mon. Not. R. Astron. Soc.* 355, 321–329. <https://doi.org/10.1111/j.1365-2966.2004.08342.x>.

- Hromakina, T., Perna, D., Belskaya, I., Dotto, E., Rossi, A., Bisi, F., 2018. Photometric observations of nine Transneptunian objects and Centaurs. *Mon. Not. R. Astron. Soc.* 474, 2536–2543. <https://doi.org/10.1093/mnras/stx2904>.
- Hyodo, R., Charnoz, S., Genda, H., Ohtsuki, K., 2016. Formation of Centaurs' rings through their partial tidal disruption during planetary encounters. *Astrophys. J.* 828, L8. <https://doi.org/10.3847/2041-8205/828/1/L8>.
- Jaeger, M., Prosseri, E., Vollmann, W., Sato, H., Sostero, G., Guido, E., 2011. Comet 174P/Echeclus = (60558) Echeclus. *Int. Astron. Union Circ.* 9213, 2.
- Jedicke, R., Herron, J.D., 1997. Observational constraints on the Centaur population. *Icarus* 127, 494–507. <https://doi.org/10.1006/icar.1997.5696>.
- Jewitt, D., 1999. Kuiper belt objects. *Annu. Rev. Earth Planet. Sci.* 27, 287–312. <https://doi.org/10.1146/annurev.earth.27.1.287>.
- Jewitt, D.C., 2002. From Kuiper belt object to cometary nucleus: the missing ultrared matter. *Astron. J.* 123, 1039–1049. <https://doi.org/10.1086/338692>.
- Jewitt, D., 2009. The active Centaurs. *Astron. J.* 137, 4296–4312. <https://doi.org/10.1088/0004-6256/137/5/4296>.
- Jewitt, D., 2015. Color systematics of Comets and related bodies. *Astron. J.* 150, Article ID: 201, 18 pp. <https://doi.org/10.1088/0004-6256/150/6/201>.
- Jewitt, D., 2018. The Trojan color conundrum. *Astron. J.* 155, 56. <https://doi.org/10.3847/1538-3881/aaa1a4>.
- Jewitt, D.C., Luu, J.X., 2001. Colors and spectra of Kuiper belt objects. *Astron. J.* 122, 2099–2114. <https://doi.org/10.1086/323304>.
- Jewitt, D., Morbidelli, A., Rauer, H., 2008. Trans-Neptunian objects and comets. In: *Saas-Fee Advanced Course 35. Swiss Society for Astrophysics and Astronomy Series*. ISBN: 978-3-540-71957-1.
- Kaňuchová, Z., Brunetto, R., Melita, M., Strazzulla, G., 2012. Space weathering and the color indexes of minor bodies in the outer solar system. *Icarus* 221, 12–19. <https://doi.org/10.1016/j.icarus.2012.06.043>.
- Kern, S.D., 2006. A Study of Binary Kuiper Belt Objects (Ph.D. thesis). Massachusetts Institute of Technology.
- Kern, S.D., McCarthy, D.W., Buie, M.W., Brown, R.H., Campins, H., Rieke, M., 2000. Compositional variation on the surface of Centaur 8405 Asbolus. *Astrophys. J.* 542, L155–L159. <https://doi.org/10.1086/312932>.
- Khare, B.N., Thompson, W.R., Murray, B.G.J.P.T., Chyba, C.F., Sagan, C., Arakawa, E.T., 1989. Solid organic residues produced by irradiation of hydrocarbon-containing H₂O and H₂O/NH₃ ices: infrared spectroscopy and astronomical implications. *Icarus* 79, 350–361. [https://doi.org/10.1016/0019-1035\(89\)90082-1](https://doi.org/10.1016/0019-1035(89)90082-1).
- Kowal, C.T., Gehrels, T., 1977. Slow-moving object Kowal. *Int. Astron. Union Circ.* 3129, 1.
- Kresák, L., 1972. On the dividing line between cometary and asteroidal orbits, In: Chebotarev, G.A., Kazimirchak-Polonskaia, E.I., Marsden, B.G. (Eds.), *The Motion, Evolution of Orbits, and Origin of Comets*, IAU Symposium, vol. 45, p. 503.
- Kuiper, G.P., 1951. On the origin of the solar system. *Proc. Natl. Acad. Sci. USA* 37, 1–14. <https://doi.org/10.1073/pnas.37.1.1>.
- Lacerda, P., 2005. The Shapes and Spins of Kuiper Belt Objects (Ph.D. thesis). Leiden Observatory, Leiden University, Leiden, The Netherlands.
- Lacerda, P., Fornasier, S., Lellouch, E., Kiss, C., Vilenius, E., Santos-Sanz, P., Rengel, M., Müller, T., Stansberry, J., Duffard, R., Delsanti, A., Guilbert-Lepoutre, A., 2014a. The Albedo-color diversity of transneptunian objects. *Astrophys. J. Lett.* 793, L2. <https://doi.org/10.1088/2041-8205/793/1/L2>.
- Lacerda, P., McNeill, A., Peixinho, N., 2014b. The unusual Kuiper belt object 2003 SQ₃₁₇. *Mon. Not. R. Astron. Soc.* 437, 3824–3831. <https://doi.org/10.1093/mnras/stt2180>.
- Lawler, S.M., Shankman, C., Kavelaars, J.J., Alexandersen, M., Bannister, M.T., Chen, Y.-T., Gladman, B., Fraser, W.C., Gwyn, S., Kaib, N., Petit, J.-M., Volk, K., 2018. OSSOS. VIII. The transition between two size distribution slopes in the scattering disk. *Astron. J.* 155, 197. <https://doi.org/10.3847/1538-3881/aab8ff>.
- Lellouch, E., Santos-Sanz, P., Lacerda, P., Mommert, M., Duffard, R., Ortiz, J.L., Müller, T.G., Fornasier, S., Stansberry, J., Kiss, C., Vilenius, E., Mueller, M., Peixinho, N., Moreno, R., Groussin, O., Delsanti, A., Harris, A.W., 2013. “TNOs are Cool”: a survey of the Trans-Neptunian region. IX. Thermal properties of Kuiper belt objects and Centaurs from combined Herschel and Spitzer observations. *Astron. Astrophys.* 557, A60. <https://doi.org/10.1051/0004-6361/201322047>.
- Lellouch, E., Moreno, R., Müller, T., Fornasier, S., Santos-Sanz, P., Moullet, A., Gurwell, M., Stansberry, J., Leiva, R., Sicardy, B., Butler, B., Boissier, J., 2017. The thermal emission of Centaurs and Trans-Neptunian objects at millimeter wavelengths from ALMA observations. *Astron. Astrophys.* 608, A45. <https://doi.org/10.1051/0004-6361/201731676>.
- Leonard, F.C., 1930. The new planet Pluto. *Leaflet Astron. Soc. Pac.* 1, 121.

- Leone, G., Paolicchi, P., Farinella, P., Zappala, V., 1984. Equilibrium models of binary asteroids. *Astron. Astrophys.* 140, 265–272.
- Levison, H.F., 1996. Comet taxonomy. In: Rettig, T., Hahn, J.M. (Eds.), *Completing the Inventory of the Solar System*, ASP Conf. Ser., vol. 107. Astronomical Society of the Pacific, pp. 173–191.
- Levison, H.F., Duncan, M.J., 1997. From the Kuiper belt to Jupiter-family comets: the spatial distribution of ecliptic comets. *Icarus* 127, 13–32. <https://doi.org/10.1006/icar.1996.5637>.
- Luu, J., Jewitt, D., 1996. Color diversity among the Centaurs and Kuiper belt objects. *Astron. J.* 112, 2310. <https://doi.org/10.1086/118184>.
- Luu, J.X., Jewitt, D.C., 2002. Kuiper belt objects: relics from the accretion disk of the Sun. *Ann. Rev. Astron. Astrophys.* 40, 63–101. <https://doi.org/10.1146/annurev.astro.40.060401.093818>.
- Meech, K.J., Belton, M.J.S., 1990. The atmosphere of 2060 Chiron. *Astron. J.* 100, 1323. <https://doi.org/10.1086/115600>.
- Morbidelli, A., 1997. Chaotic diffusion and the origin of comets from the 2/3 resonance in the Kuiper belt. *Icarus* 127, 1–12. <https://doi.org/10.1006/icar.1997.5681>.
- Nesvorný, D., 2018. Dynamical evolution of the early solar system. *Ann. Rev. Astron. Astrophys.* 56, 137–174. <https://doi.org/10.1146/annurev-astro-081817-052028>.
- Ortiz, J.L., Gutiérrez, P.J., Casanova, V., Sota, A., 2003. A study of short term rotational variability in TNOs and Centaurs from Sierra Nevada observatory. *Astron. Astrophys.* 407, 1149–1155. <https://doi.org/10.1051/0004-6361:20030972>.
- Ortiz, J.L., Gutiérrez, P.J., Santos-Sanz, P., Casanova, V., Sota, A., 2006. Short-term rotational variability of eight KBOs from Sierra Nevada Observatory. *Astron. Astrophys.* 447, 1131–1144. <https://doi.org/10.1051/0004-6361:20053572>.
- Ortiz, J.L., Duffard, R., Pinilla-Alonso, N., Alvarez-Candal, A., Santos-Sanz, P., Morales, N., Fernández-Valenzuela, E., Licandro, J., Campo Bagatin, A., Thirouin, A., 2015. Possible ring material around centaur (2060) Chiron. *Astron. Astrophys.* 576, A18. <https://doi.org/10.1051/0004-6361/201424461>.
- Pan, M., Wu, Y., 2016. On the mass and origin of Chariklo's rings. *Astrophys. J.* 821, 18. <https://doi.org/10.3847/0004-637X/821/1/18>.
- Peixinho, N., Doressoundiram, A., Delsanti, A., Boehnhardt, H., Barucci, M.A., Belskaya, I., 2003. Reopening the TNOs color controversy: centaurs bimodality and TNOs unimodality. *Astron. Astrophys.* 410, L29–L32. <https://doi.org/10.1051/0004-6361:20031420>.
- Peixinho, N., Delsanti, A., Guilbert-Lepoutre, A., Gafeira, R., Lacerda, P., 2012. The bimodal colors of Centaurs and small Kuiper belt objects. *Astron. Astrophys.* 546, A86. <https://doi.org/10.1051/0004-6361/201219057>.
- Petit, J.-M., Kavelaars, J.J., Gladman, B., Lored, T., 2008. Size distribution of multikilometer transneptunian objects. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 71–87.
- Prialnik, D., 1992. Crystallization, sublimation, and gas release in the interior of a porous comet nucleus. *Astrophys. J.* 388, 196. <https://doi.org/10.1086/171143>.
- Prialnik, D., Podolak, M., 1995. Radioactive heating of porous comet nuclei. *Icarus* 117, 420–430. <https://doi.org/10.1006/icar.1995.1166>.
- Prialnik, D., Bar-Nun, A., Podolak, M., 1987. Radiogenic heating of comets by ^{26}Al and implications for their time of formation. *Astrophys. J.* 319, 993. <https://doi.org/10.1086/165516>.
- Prialnik, D., Sarid, G., Rosenberg, E.D., Merk, R., 2008. Thermal and chemical evolution of comet nuclei and Kuiper belt objects. *Space Sci. Rev.* 138, 147–164. <https://doi.org/10.1007/s11214-007-9301-4>.
- Rabinowitz, D.L., Schaefer, B.E., Tourtellotte, S.W., 2007. The diverse solar phase curves of distant icy bodies. I. Photometric observations of 18 Trans-Neptunian objects, 7 Centaurs, and Nereid. *Astron. J.* 133, 26–43. <https://doi.org/10.1086/508931>.
- Romanishin, W., Tegler, S.C., 1999. Rotation rates of Kuiper-belt objects from their light curves. *Nature* 398, 129–132. <https://doi.org/10.1038/18168>.
- Rosenberg, E.D., Prialnik, D., 2010. The effect of internal inhomogeneity on the activity of comet nuclei—application to Comet 67P/Churyumov-Gerasimenko. *Icarus* 209, 753–765. <https://doi.org/10.1016/j.icarus.2010.03.037>.
- Rousselot, P., Petit, J.-M., Poulet, F., Sergeev, A., 2005. Photometric study of Centaur (60558) 2000 EC 98 and Trans-Neptunian object (55637) 2002 UX 25 at different phase angles. *Icarus* 176, 478–491. <https://doi.org/10.1016/j.icarus.2005.03.001>.

- Rousselot, P., Korsun, P.P., Kulyk, I., Guilbert-Lepoutre, A., Petit, J.-M., 2016. A long-term follow up of 174P/Echeclus. *Mon. Not. R. Astron. Soc.* 462, S432–S442. <https://doi.org/10.1093/mnras/stw3054>.
- Ruprecht, J.D., Bosh, A.S., Person, M.J., Bianco, F.B., Fulton, B.J., Gulbis, A.A.S., Bus, S.J., Zangari, A.M., 2015. 29 November 2011 stellar occultation by 2060 Chiron: symmetric jet-like features. *Icarus* 252, 271–276. <https://doi.org/10.1016/j.icarus.2015.01.015>.
- Russell, H.N., 1916. On the Albedo of the planets and their satellites. *Astrophys. J.* 43, 173–196. <https://doi.org/10.1086/142244>.
- Sarid, G., Prialnik, D., 2009. From KBOs to Centaurs: the thermal connection. *Meteorit. Planet. Sci.* 44, 1905–1916. <https://doi.org/10.1111/j.1945-5100.2009.tb02000.x>.
- Schwamb, M.E., Fraser, W.C., Bannister, M.T., Marsset, M., Pike, R.E., Kavelaars, J.J., Benecchi, S.D., Lehner, M.J., Wang, S.-Y., Thirouin, A., Delsanti, A., Peixinho, N., Volk, K., Alexandersen, M., Chen, Y.-T., Gladman, B., Gwyn, S.D.J., Petit, J.-M., 2018. Col-OSSOS: the colours of the outer solar system origins survey. *Astrophys. J. Suppl. Ser.* 243, Article ID: 12, 24 pp. <https://doi.org/10.3847/1538-4365/ab2194>.
- Scotti, J.V., Rabinowitz, D.L., Shoemaker, C.S., Shoemaker, E.M., Levy, D.H., King, T.M., Helin, E.F., Alu, J., Lawrence, K., McNaught, R.H., Frederick, L., Tholen, D., Mueller, B.E.A., 1992. 1992 AD. *Int. Astron. Union Circ.* 5434.
- Seccull, T., Fraser, W.C., Puzia, T.H., Fitzsimmons, A., Cupani, G., 2019. 174P/Echeclus and its blue coma observed post-outburst. *Astron. J.* 157, 88. <https://doi.org/10.3847/1538-3881/aafbe4>.
- Shankman, C., Gladman, B.J., Kaib, N., Kavelaars, J.J., Petit, J.M., 2013. A possible divot in the size distribution of the Kuiper belt's scattering objects. *Astrophys. J.* 764, L2. <https://doi.org/10.1088/2041-8205/764/1/L2>.
- Sheppard, S.S., Jewitt, D.C., 2002. Time-resolved photometry of Kuiper belt objects: rotations, shapes, and phase functions. *Astron. J.* 124, 1757–1775. <https://doi.org/10.1086/341954>.
- Sheppard, S.S., Jewitt, D.C., Trujillo, C.A., Brown, M.J.I., Ashley, M.C.B., 2000. A wide-field CCD survey for Centaurs and Kuiper belt objects. *Astron. J.* 120, 2687–2694. <https://doi.org/10.1086/316805>.
- Sheppard, S.S., Trujillo, C., Tholen, D.J., 2016. Beyond the Kuiper belt edge: new high Perihelion Trans-Neptunian objects with moderate semimajor axes and eccentricities. *Astrophys. J.* 825, L13. <https://doi.org/10.3847/2041-8205/825/1/L13>.
- Sicardy, B., Leiva, R., Renner, S., Roques, F., El Moutamid, M., Santos-Sanz, P., Desmars, J., 2018. Ring dynamics around non-axisymmetric bodies with application to Chariklo and Haumea. *Nat. Astron.* <https://doi.org/10.1038/s41550-018-0616-8>.
- Stansberry, J., Grundy, W., Brown, M., Cruikshank, D., Spencer, J., Trilling, D., Margot, J.-L., 2008. Physical properties of Kuiper belt and centaur objects: constraints from the Spitzer Space Telescope. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 161–179.
- Stern, S.A., 1996. Signatures of collisions in the Kuiper disk. *Astron. Astrophys.* 310, 999–1010.
- Tegler, S.C., Romanishin, W., 1998. Two distinct populations of Kuiper-belt objects. *Nature* 392, 49–51. <https://doi.org/10.1038/32108>.
- Tegler, S.C., Romanishin, W., 2000. Extremely red Kuiper-belt objects in near-circular orbits beyond 40 AU. *Nature* 407, 979–981. <https://doi.org/10.1038/35039572>.
- Tegler, S.C., Romanishin, W., 2003. Resolution of the Kuiper belt object color controversy: two distinct color populations. *Icarus* 161, 181–191. [https://doi.org/10.1016/S0019-1035\(02\)00021-0](https://doi.org/10.1016/S0019-1035(02)00021-0).
- Tegler, S.C., Romanishin, W., Consolmagno, G.J., 2003. Color patterns in the Kuiper belt: a possible primordial origin. *Astrophys. J.* 599, L49–L52. <https://doi.org/10.1086/381076>.
- Tegler, S.C., Romanishin, W., Consolmagno, G.J., Rall, J., Worhatch, R., Nelson, M., Weidenschilling, S., 2005a. The period of rotation, shape, density, and homogeneous surface color of the Centaur 5145 Pholus. *Icarus* 175, 390–396. <https://doi.org/10.1016/j.icarus.2004.12.011>.
- Tegler, S.C., Romanishin, W., Consolmagno, G.J., Rall, J., Worhatch, R., Nelson, M., Weidenschilling, S., 2005b. The period of rotation, shape, density, and homogeneous surface color of the Centaur 5145 Pholus. *Icarus* 175, 390–396. <https://doi.org/10.1016/j.icarus.2004.12.011>.
- Tegler, S.C., Bauer, J.M., Romanishin, W., Peixinho, N., 2008. Colors of Centaurs. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 105–114.
- Tegler, S.C., Romanishin, W., Consolmagno, G.J., J., S., 2016. Two color populations of Kuiper belt and Centaur objects and the smaller orbital inclinations of red Centaur objects. *Astron. J.* 152, 210. <https://doi.org/10.3847/0004-6256/152/6/210>.

- Thébault, P., 2003. A numerical check of the collisional resurfacing scenario. *Earth Moon Planets* 92, 233–241. <https://doi.org/10.1023/B:MOON.0000031941.77871.c5>.
- Thébault, P., Doressoundiram, A., 2003. Colors and collision rates within the Kuiper belt: problems with the collisional resurfacing scenario. *Icarus* 162, 27–37. [https://doi.org/10.1016/S0019-1035\(03\)00004-6](https://doi.org/10.1016/S0019-1035(03)00004-6).
- Thirouin, A., 2013. Study of trans-neptunian objects using photometric techniques and numerical simulations (Ph.D. thesis). University of Granada.
- Thirouin, A., Sheppard, S.S., 2018. The Plutino population: an abundance of contact binaries. *Astron. J.* 155, 248. <https://doi.org/10.3847/1538-3881/aac0ff>.
- Thirouin, A., Sheppard, S.S., 2019. Lightcurves and rotational properties of the pristine cold classical Kuiper belt objects. *Astron. J.* 157, Article ID: 228, 12 pp. <https://doi.org/10.3847/1538-3881/ab18a9>.
- Thirouin, A., Ortiz, J.L., Duffard, R., Santos-Sanz, P., Aceituno, F.J., Morales, N., 2010. Short-term variability of a sample of 29 Trans-Neptunian objects and Centaurs. *Astron. Astrophys.* 522, A93. <https://doi.org/10.1051/0004-6361/200912340>.
- Thirouin, A., Sheppard, S.S., Noll, K.S., Moskovitz, N.A., Ortiz, J.L., Doressoundiram, A., 2016. Rotational properties of the Haumea family members and candidates: short-term variability. *Astron. J.* 151, 148. <https://doi.org/10.3847/0004-6256/151/6/148>.
- Tholen, D.J., Hartmann, W.K., Cruikshank, D.P., Lilly, S., Bowell, E., Hewitt, A., 1988. (2060) Chiron. *Int. Astron. Union Circ.* 4554, 2.
- Tiscareno, M.S., Malhotra, R., 2003. The dynamics of known Centaurs. *Astron. J.* 126, 3122–3131. <https://doi.org/10.1086/379554>.
- Tiscareno, M.S., Malhotra, R., 2009. Chaotic diffusion of resonant Kuiper belt objects. *Astron. J.* 138, 827–837. <https://doi.org/10.1088/0004-6256/138/3/827>.
- Tisserand, F., 1896. *Traité de mécanique céleste*, t. IV. In-4. Gauthier-Villars et fils.
- Trujillo, C.A., 2008. Future surveys of the Kuiper belt. In: Barucci, M.A., Boehnhardt, H., Cruikshank, D.P., Morbidelli, A., Dotson, R. (Eds.), *The Solar System Beyond Neptune*. University of Arizona Press, Tucson, AZ, pp. 573–585.
- Volk, K., Malhotra, R., 2008. The scattered disk as the source of the Jupiter family comets. *Astrophys. J.* 687, 714–725. <https://doi.org/10.1086/591839>.
- Volk, K., Malhotra, R., 2013. Do Centaurs preserve their source inclinations? *Icarus* 224, 66–73. <https://doi.org/10.1016/j.icarus.2013.02.016>.
- Weryk, R.J., Schunova-Lilly, E., Jedicke, R., Fraser, W.C., Volk, K., Wainscoat, R.J., 2019. The Population of Trans-Neptunian Objects Detected by Pan-STARRS1 (in preparation).
- Wong, I., Brown, M.E., 2016. A hypothesis for the color bimodality of Jupiter Trojans. *Astron. J.* 152, 90. <https://doi.org/10.3847/0004-6256/152/4/90>.
- Wong, I., Brown, M.E., 2017. The bimodal color distribution of small Kuiper belt objects. *Astron. J.* 153, 145. <https://doi.org/10.3847/1538-3881/aa60c3>.
- Yu, Q., Tremaine, S., 1999. The dynamics of Plutinos. *Astron. J.* 118, 1873–1881. <https://doi.org/10.1086/301045>.