

ANNALS OF K-THEORY

vol. 4 no. 4 2019

**A Dolbeault–Hilbert complex
for a variety with isolated singular points**

John Lott



A JOURNAL OF THE K-THEORY FOUNDATION

A Dolbeault–Hilbert complex for a variety with isolated singular points

John Lott

Given a compact Hermitian complex space with isolated singular points, we construct a Dolbeault-type Hilbert complex whose cohomology is isomorphic to the cohomology of the structure sheaf. We show that the corresponding K-homology class coincides with the one constructed by Baum, Fulton and MacPherson.

1. Introduction

The program of doing index theory, or more generally elliptic theory, on singular varieties goes back at least to [Singer 1971, §4]. This program takes various directions — for example, the relation between L^2 -cohomology and intersection homology. In this paper we consider a somewhat different direction, which is related to the arithmetic genus. This is motivated by work of Baum, Fulton and MacPherson [Baum et al. 1975; 1979].

Let X be a projective complex algebraic variety and let $\mathcal{S}$ be a coherent sheaf on X . In [Baum et al. 1979], the authors associated to $\mathcal{S}$ an element $[\mathcal{S}]_{\text{BFM}} \in K_0(X)$ of the topological K-homology of X . This class enters into their Riemann–Roch theorem for singular varieties. In particular, under the map $p : X \rightarrow \text{pt}$, the image $p_*[\mathcal{S}]_{\text{BFM}} \in K_0(\text{pt}) \cong \mathbb{Z}$ is expressed in terms of sheaf cohomology by $\sum_i (-1)^i \dim(H^i(X; \mathcal{S}))$.

In view of the isomorphism between topological K-homology and analytic K-homology [Baum and Douglas 1982; Baum et al. 2007], the class $[\mathcal{S}]_{\text{BFM}}$ can be represented by an “abstract elliptic operator” in the sense of [Atiyah 1970]. This raised the question of how to find an explicit cycle in analytic K-homology, even if X is singular, that represents $[\mathcal{S}]_{\text{BFM}}$. The most basic case is when $\mathcal{S}$ is the structure sheaf $\mathcal{O}_X$. If X is smooth then the operator representing $[\mathcal{O}_X]_{\text{BFM}}$ is $\bar{\partial} + \bar{\partial}^*$. Hence we are looking for the right analog of this operator when X may be singular.

A second related question is to find a Hilbert complex, in the sense of [Brüning and Lesch 1992], whose cohomology is isomorphic to $H^*(X; \mathcal{O}_X)$. We want the

Research partially supported by NSF grant DMS-1810700.

MSC2010: 19K33, 19L10, 32W05, 58J10.

Keywords: Dolbeault, singular, variety, Riemann–Roch.

complex to be intrinsic to X . Also, if X is smooth then we want to recover the $\bar{\partial}$ -complex on $(0, \star)$ -forms.

In this paper, we answer these questions when X has isolated singular points. To see the nature of the problem, suppose that X is a complex curve, whose normalization has genus g . In this case, the Riemann–Roch theorem says

$$\dim(H^0(X; \mathcal{O}_X)) - \dim(H^1(X; \mathcal{O}_X)) = 1 - g - \sum_{x \in X_{\text{sing}}} \delta_x, \quad (1.1)$$

where δ_x is a certain positive integer attached to the singular point x [Hartshorne 1977, p. 298]. To find the appropriate Hilbert complex, it is natural to start with the Dolbeault complex $\Omega_c^{0,0}(X_{\text{reg}}) \xrightarrow{\bar{\partial}} \Omega_c^{0,1}(X_{\text{reg}})$ of smooth compactly supported forms on X_{reg} and look for a closed operator extension, where X_{reg} is endowed with the induced Riemannian metric from its projective embedding. For the minimal closure $\bar{\partial}_s$, one finds $\text{Index}(\bar{\partial}_s) = 1 - g$. Taking a different closure can only make the index go up [Brüning et al. 1990], whereas in view of (1.1) we want the index to go down. (Considering complete Riemannian metrics on X_{reg} does not help.) However, on the level of indices, we can get the right answer by enhancing the codomain by $\bigoplus_{x \in X_{\text{sing}}} \mathbb{C}^{\delta_x}$.

Now let X be a compact Hermitian complex space of pure dimension n . For technical reasons, we assume that the singular set X_{sing} consists of isolated singularities. (In the bulk of the paper we allow coupling to a holomorphic vector bundle, but in this introduction we only discuss the case when the vector bundle is trivial.) Let $\bar{\partial}_s$ be the minimal closed extension of the $\bar{\partial}$ -operator on $X_{\text{reg}} = X - X_{\text{sing}}$. Its domain $\text{Dom}(\bar{\partial}_s^{0,\star})$ can be localized to a complex of sheaves $\underline{\text{Dom}}(\bar{\partial}_s^{0,\star})$. Let $\underline{H}^{0,\star}(\bar{\partial}_s)$ denote the cohomology, a sum of skyscraper sheaves on X if $\star > 0$. We write $\mathcal{O}_s$ for $\underline{H}^{0,0}(\bar{\partial}_s)$, which is the sheaf of germs of weakly holomorphic functions on X , the latter being in the sense of [Whitney 1972, Section 4.3]. Then $\mathcal{O}_s/\mathcal{O}_X$ is also a sum of skyscraper sheaves on X . Its vector space of global sections will be written as $(\mathcal{O}_s/\mathcal{O}_X)(X)$. Both $\underline{H}^{0,\star}(\bar{\partial}_s)$ and $\mathcal{O}_s/\mathcal{O}_X$ can be computed using a resolution of X [Ruppenthal 2018, Corollary 1.2].

Define vector spaces T^* by

$$\begin{aligned} T^0 &= \text{Dom}(\bar{\partial}_s^{0,0}), \\ T^1 &= \text{Dom}(\bar{\partial}_s^{0,1}) \oplus (\mathcal{O}_s/\mathcal{O}_X)(X), \\ T^\star &= \text{Dom}(\bar{\partial}_s^{0,\star}) \oplus (\underline{H}^{0,\star-1}(\bar{\partial}_s))(X), \quad \text{if } 2 \leq \star \leq n. \end{aligned} \quad (1.2)$$

To define a differential on T^* , let $\Delta_s^{0,\star}$ be the Laplacian associated to $\bar{\partial}_s$. Let $P_{\text{Ker}(\Delta_s^{0,\star})}$ be orthogonal projection onto the kernel of $\Delta_s^{0,\star}$. As elements of $\text{Ker}(\Delta_s^{0,\star})$ are $\bar{\partial}_s$ -closed, for each $x \in X_{\text{sing}}$ there is a well-defined map $\text{Ker}(\Delta_s^{0,\star}) \rightarrow (\underline{H}^{0,\star}(\bar{\partial}_s))_x$ to the stalk of $\underline{H}^{0,\star}(\bar{\partial}_s)$ at x . For $\star > 0$, putting these together for all $x \in X_{\text{sing}}$, and

precomposing with $P_{\text{Ker}(\Delta_s^{0,\star})}$, gives a linear map $\gamma : \text{Dom}(\bar{\partial}_s^{0,\star}) \rightarrow (\underline{H}^{0,\star}(\bar{\partial}_s))(X)$. For $\star = 0$, we similarly define $\gamma : \text{Dom}(\bar{\partial}_s^{0,0}) \rightarrow (\mathcal{O}_s/\mathcal{O}_X)(X)$. Define a differential $d : T^* \rightarrow T^{*+1}$ by

$$\begin{aligned} d(\omega) &= (\bar{\partial}_s \omega, \gamma(\omega)), & \text{if } \star = 0, \\ d(\omega, a) &= (\bar{\partial}_s \omega, \gamma(\omega)), & \text{if } \star > 0. \end{aligned} \tag{1.3}$$

Theorem 1.4. *The cohomology of (T, d) is isomorphic to $H^*(X; \mathcal{O}_X)$.*

Theorem 1.4 can be seen as an extension of [Ruppenthal 2018, Corollary 1.3], which implies the result when X is normal and has rational singularities. To prove Theorem 1.4, we construct a certain resolution of $\mathcal{O}_X$ by fine sheaves. The cohomology of the complex $(\tilde{T}, \tilde{d})$ of global sections is then isomorphic to $H^*(X; \mathcal{O}_X)$. The complex $(\tilde{T}, \tilde{d})$ is not quite the same as (T, d) but we show that they are cochain-equivalent, from which the theorem follows.

The spectral triple $(C(X), T, d + d^*)$ defines an element $[\mathcal{O}_X]_{\text{an}} \in K_0(X)$ of the analytic K-homology of X .

Theorem 1.5. *If X is a projective algebraic variety with isolated singularities then $[\mathcal{O}_X]_{\text{an}} = [\mathcal{O}_X]_{\text{BFM}}$ in $K_0(X)$.*

There has been some interesting earlier work on the questions addressed in this paper. Ancona and Gaveau [1994] gave a resolution of the structure sheaf of a normal complex space X , assuming that the singular locus is smooth, in terms of differential forms on a resolution of X . The construction depended on the choice of resolution. Fox and Haskell [2000] discussed using a perturbed Dolbeault operator on an ambient manifold to represent the K-homology class of the structure sheaf. Andersson and Samuelsson [2012] gave a resolution of the structure sheaf by certain currents on X that are smooth on X_{reg} . After this paper was written, Bei and Piazza [2019] posted a preprint which also has a proof of Proposition 5.1.

The structure of the paper is the following. In Section 2, given a holomorphic vector bundle V on X , we recall the definition of the minimal closure $\bar{\partial}_{V,s}$ and show that $\bar{\partial}_{V,s} + \bar{\partial}_{V,s}^*$ gives an element of the analytic K-homology group $K_0(X)$, in the unbounded formalism for the Kasparov KK-group $\text{KK}(C(X); \mathbb{C})$. In Section 3 we construct a resolution of the sheaf $\underline{V}$ by fine sheaves. Their global sections give a Hilbert complex. In Section 4 we deform this to the complex (T_V, d_V) . Section 5 has the proof of Theorem 1.5. More detailed descriptions appear at the beginning of the sections.

2. Minimal closure and compact resolvent

In this section we consider a holomorphic vector bundle V on a compact complex space X with isolated singularities. We define the minimal closure $\bar{\partial}_{V,s}$. We show

that the spectral triple $(C(X), \bar{\partial}_{V,s} + \bar{\partial}_{V,s}^*, \Omega_{L^2}^{0,*}(X_{\text{reg}}; V))$ gives a well-defined element of the analytic K-homology group $K_0(X)$, in the unbounded formalism. The main issue is to show that $\bar{\partial}_{V,s} + \bar{\partial}_{V,s}^*$ has a compact resolvent. When V is trivial, this was shown in [Øvrelid and Ruppenthal 2014].

Let X be a reduced compact complex space of pure dimension n . For each $x \in X$, there is a neighborhood U of x with an embedding of U into some domain $U' \subset \mathbb{C}^N$, as the zero set of a finite number of holomorphic functions on U' .

Let $\mathcal{O}_X$ be the analytic structure sheaf of X . Let X_{sing} be the set of singular points of X and put $X_{\text{reg}} = X - X_{\text{sing}}$.

We equip X with a Hermitian metric g on X_{reg} which satisfies the property that for each $x \in X$, there are U and U' as above, along with a smooth Hermitian metric G on U' , so that $g|_{X_{\text{reg}} \cap U} = G|_{X_{\text{reg}} \cap U}$.

Let V be a finite dimensional holomorphic vector bundle on X or, equivalently, a locally free sheaf $\underline{V}$ of $\mathcal{O}_X$ -modules. For each $x \in X$, there are U and U' as above so that $V|_U$ is the restriction of a trivial holomorphic bundle $U' \times \mathbb{C}^N$ on U' . Let h be a Hermitian inner product on $V|_{X_{\text{reg}}}$ which satisfies the property that for each $x \in X$, there are such U and U' so that $h|_{X_{\text{reg}} \cap U}$ is the restriction of a smooth Hermitian metric on $U' \times \mathbb{C}^N$.

Let $\bar{\partial}_{V,s}$ be the minimal closed extension of the $\bar{\partial}_V$ -operator on X_{reg} . That is, the domain of $\bar{\partial}_{V,s}$ is the set of $\omega \in \Omega_{L^2}^{0,*}(X_{\text{reg}}; V)$ so that there are a sequence of compactly supported smooth forms $\omega_i \in \Omega^{0,*}(X_{\text{reg}}; V)$ on X_{reg} and some $\eta \in \Omega_{L^2}^{0,*+1}(X_{\text{reg}}; V)$ such that

$$\begin{aligned} \lim_{i \rightarrow \infty} \omega_i &= \omega \quad \text{in } \Omega_{L^2}^{0,*}(X_{\text{reg}}; V), \\ \lim_{i \rightarrow \infty} \bar{\partial}_{V,s} \omega_i &= \eta \quad \text{in } \Omega_{L^2}^{0,*+1}(X_{\text{reg}}; V). \end{aligned}$$

We then put $\bar{\partial}_{V,s} \omega = \eta$, which is uniquely defined.

Hereafter we assume that X_{sing} is finite.

Proposition 2.1. *The spectral triple $(C(X), \bar{\partial}_{V,s} + \bar{\partial}_{V,s}^*, \Omega_{L^2}^{0,*}(X_{\text{reg}}; V))$ gives a well-defined element of the analytic K-homology group $K_0(X)$.*

Proof. Put $D_V = \bar{\partial}_{V,s} + \bar{\partial}_{V,s}^*$, with dense domain $\text{Dom}(\bar{\partial}_{V,s}) \cap \text{Dom}(\bar{\partial}_{V,s}^*)$. Put $D = \bar{\partial}_s + \bar{\partial}_s^*$, the case when V is the trivial complex line bundle. Put

$$\mathcal{A} = \{f \in C(X) : f(\text{Dom}(D_V)) \subset \text{Dom}(D_V) \text{ and } [D_V, f] \text{ is bounded}\}. \quad (2.2)$$

Using the local trivializations of V , it follows that

$$\mathcal{A} = \{f \in C(X) : f(\text{Dom}(D)) \subset \text{Dom}(D) \text{ and } [D, f] \text{ is bounded}\}. \quad (2.3)$$

To satisfy the definitions of unbounded analytic K-homology [Baaĳ and Julg 1983; Forsyth et al. 2014; Kaad 2019], we first need to show that $\mathcal{A}$ is dense in $C(X)$.

Given $F \in C(X)$ and $\epsilon > 0$, we can construct $f \in C(X)$ such that

- for each $x_j \in X_{\text{sing}}$, there is a neighborhood $U_j \subset X$ of x_j on which f is constant, with $f(x_j) = F(x_j)$;
- f is smooth on X_{reg} ;
- $\sup_{x \in X} |f(x) - F(x)| < \epsilon$.

Then $f(\text{Dom}(D)) \subset \text{Dom}(D)$ and $\|[D, f]\| \leq \text{const} \cdot \|\nabla_h f\|_\infty < \infty$. It follows that $\mathcal{A}$ is dense in $C(X)$.

To prove the proposition, it now suffices to prove the following lemma.

Lemma 2.4. *The operator $(D_V + i)^{-1}$ is compact.*

Proof. If V is trivial then the lemma is true [Øvrelid and Ruppenthal 2014]. We use a parametrix construction to prove it for general V .

Let us first prove the lemma for a special inner product h' on V . We write $X_{\text{sing}} = \{x_j\}_{j=1}^r$. For each j , let U_j be a neighborhood of x_j on which V is trivialized as above, with $\overline{U_j} \cap \overline{U_k} = \emptyset$ for $j \neq k$. Choose open sets with smooth boundary $x_j \in Z_j \subset Y_j \subset W_j \subset U_j$, with $\overline{Z_j} \subset Y_j$, $\overline{Y_j} \subset W_j$ and $\overline{W_j} \subset U_j$. Let $\phi_j \in C(X)$ be identically 1 on Y_j , with support in W_j , and smooth on $U_j - Y_j$. Let $\eta_j \in C(X)$ be identically 1 on W_j , with support in U_j , and smooth on $U_j - Y_j$, so that η_j is 1 on the support of ϕ_j .

Define an inner product h' on V by first taking it to be a trivial inner product on each U_j , in terms of our given trivializations, and then extending it smoothly to the rest of X_{reg} . Let V_j be the extension of the trivialization $U_j \times \mathbb{C}^N$ to a product bundle on $X \times \mathbb{C}^N$ on X , as a smooth vector bundle with trivial inner product. Let $D_{V_j} = D \otimes I_N$ be the corresponding operator. As $(D + i)^{-1}$ is compact [Øvrelid and Ruppenthal 2014], the same is true for D_{V_j} . Let D_{APS} be the operator $\bar{\partial}_V + \bar{\partial}_V^*$ on $X - \bigcup_j Z_j$, with Atiyah–Patodi–Singer boundary conditions [Atiyah et al. 1973]. (The paper [Atiyah et al. 1973] assumes a product structure near the boundary, but this is not necessary.) Then $(D_{\text{APS}} + i)^{-1}$ is compact. Put $\phi_0 = 1 - \sum_j \phi_j$, with support in $X - \bigcup_j \overline{Z_j}$. Pick $\eta_0 \in C(X)$ with support in $X - \bigcup_j \overline{Z_j}$, and smooth on X_{reg} , such that η_0 is one on the support of ϕ_0 .

For $\omega \in \Omega_{L^2}^{0,*}(X_{\text{reg}}; V)$, put

$$Q\omega = \eta_0(D_{\text{APS}} + i)^{-1}(\phi_0\omega) + \sum_j \eta_j(D_{V_j} + i)^{-1}(\phi_j\omega). \quad (2.5)$$

Then Q is compact and

$$\begin{aligned} (D_V + i)Q\omega &= \omega + [D, \eta_0](D_{\text{APS}} + i)^{-1}(\phi_0\omega) + \sum_j [D, \eta_j](D_{V_j} + i)^{-1}(\phi_j\omega), \end{aligned} \quad (2.6)$$

so

$$(D_V + i)^{-1} = Q - (D_V + i)^{-1} \left([D, \eta_0](D_{\text{APS}} + i)^{-1} \phi_0 + \sum_j [D, \eta_j](D_{V_j} + i)^{-1} \phi_j \right). \quad (2.7)$$

As $[D, \eta_0]$, $[D, \eta_j]$ and $(D_V + i)^{-1}$ are bounded, it follows that $(D_V + i)^{-1}$ is compact.

As $(D_V + i)^{-1}$ (for the inner product h') is compact, the spectral theorem for compact operators and the functional calculus imply that $(I + D_V^2)^{-1}$ is compact. Writing $\Delta_{V,s} = D_V^2$, there is then a Hodge decomposition

$$\Omega_{L^2}^{0,*}(X_{\text{reg}}; V) = \text{Ker}(\Delta_{V,s}^{0,*}) \oplus \text{Im}(\bar{\partial}_{V,s}) \oplus \text{Im}(\bar{\partial}_{V,s}^*), \quad (2.8)$$

where the right-hand side is a sum of orthogonal closed subspaces. In particular,

- (1) $\text{Im}(\bar{\partial}_{V,s})$ is closed,
- (2) $\text{Ker}(\bar{\partial}_{V,s}) / \text{Im}(\bar{\partial}_{V,s})$ is finite dimensional and
- (3) the map $\bar{\partial}_{V,s} : \Omega_{L^2}^{0,*}(X_{\text{reg}}; V) / \text{Ker}(\bar{\partial}_{V,s}) \rightarrow \text{Im}(\bar{\partial}_{V,s})$ is invertible and the inverse is compact, i.e., sends bounded sets to precompact sets.

(The inverse map $\text{Im}(\bar{\partial}_{V,s}) \rightarrow \Omega_{L^2}^{0,*}(X_{\text{reg}}; V) / \text{Ker}(\bar{\partial}_{V,s}) \cong \text{Im}(\bar{\partial}_{V,s}^*)$ is DG , where G is the Green's operator for $\Delta_{V,s}$.) As the L^2 -inner products on $\Omega_{L^2}^{0,*}(X_{\text{reg}}; V)$ coming from h' and h are relatively bounded, the above three properties also hold for h . It follows that there is a Hodge decomposition relative to the inner product h , and $(I + D_V^2)^{-1}$ is compact. Hence $(D_V + i)^{-1}$ is compact.

This completes the proof of the lemma, and hence the proposition. $\square$

3. Resolution

In this section we construct a certain resolution of the sheaf of holomorphic sections of a holomorphic vector bundle V on X . To begin, we define a sheaf $\underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,*})$ on X , following [Ruppenthal 2018, Section 2.1].

Given an open set $U \subset X$ and a compact subset $K \subset U$, we write U_{reg} for $U \cap X_{\text{reg}}$ and K_{reg} for $K \cap X_{\text{reg}}$.

Let V be a finite dimensional holomorphic vector bundle on X equipped with a Hermitian metric, in the sense of Section 2. There is a sheaf $\underline{\Omega}_{V, L_{\text{loc}}^2}^{0,*}$ on X whose sections over an open set $U \subset X$ are the locally square integrable V -valued forms of degree $(0, \star)$ on U_{reg} , i.e., they are square integrable on K_{reg} for any compact set $K \subset U$. Convergence means L^2 -convergence on each such K_{reg} . By definition, the sections of $\underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,*})$ over U are the elements $\omega \in \Omega_{L_{\text{loc}}^2}^{0,*}(U_{\text{reg}}; V)$ so that there are

- a sequence $f_i \in \Omega_{C_c^\infty}^{0,*}(U_{\text{reg}}; V)$ and
- some $\eta \in \Omega_{L_{\text{loc}}^2}^{0,*+1}(U_{\text{reg}}; V)$

such that for any compact $K \subset U$, we have

- $\lim_{i \rightarrow \infty} f_i = \omega$ in $\Omega_{L^2}^{0,\star}(K_{\text{reg}}; V)$ and
- $\lim_{i \rightarrow \infty} \bar{\partial}_V f_i = \eta$ in $\Omega_{L^2}^{0,\star+1}(K_{\text{reg}}; V)$.

Then we put $\bar{\partial}_V \omega = \eta$.

This gives a complex of fine sheaves

$$\dots \xrightarrow{\bar{\partial}_V} \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star-1}) \xrightarrow{\bar{\partial}_V} \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star}) \xrightarrow{\bar{\partial}_V} \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star+1}) \xrightarrow{\bar{\partial}_V} \dots \quad (3.1)$$

The cohomology of the complex is the sheaf $\underline{H}^{0,\star}(\bar{\partial}_{V,s})$. For $\star > 0$, it is a direct sum of skyscraper sheaves, with support in X_{sing} . We write $\underline{V}_s$ for $\underline{H}^{0,0}(\bar{\partial}_{V,s})$, i.e., the kernel of $\bar{\partial}_V$ acting on $\underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,0})$. Then $\underline{V}_s/\underline{V}$ is also a direct sum of skyscraper sheaves with support in X_{sing} .

Although we do not need it here, there is a description of these skyscraper sheaves in terms of a resolution of X . Suppose that $\pi : M \rightarrow X$ is a resolution. From [Ruppenthal 2018, Corollary 1.2], if $x \in X$ then we can identify the stalk $(\underline{H}^{0,q}(\bar{\partial}_{V,s}))_x$ with $V_x \otimes (R^q \pi_* \mathcal{O}_M)_x$. In particular, we can identify $\underline{V}_s$ with $\underline{V} \otimes_{\mathcal{O}_X} \pi_* \mathcal{O}_M$ or, more intrinsically, with the sheaf of weakly holomorphic sections of V , i.e., bounded holomorphic sections of $V|_{X_{\text{reg}}}$.

There is a quotient morphism of sheaves:

$$q : \underline{\text{Ker}}(\bar{\partial}_{V,s}^{0,\star}) \rightarrow \underline{H}^{0,\star}(\bar{\partial}_{V,s}).$$

As $\underline{H}^{0,\star}(\bar{\partial}_{V,s})$ is an injective sheaf for $\star > 0$, we can extend q to a morphism $\alpha : \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star}) \rightarrow \underline{H}^{0,\star}(\bar{\partial}_{V,s})$. More specifically, if x is a singular point then the stalk $(\underline{H}^{0,\star}(\bar{\partial}_{V,s}))_x$ is a finite dimensional complex vector space, so we are extending the quotient map $q_x : (\underline{\text{Ker}}(\bar{\partial}_{V,s}^{0,\star}))_x \rightarrow (\underline{H}^{0,\star}(\bar{\partial}_{V,s}))_x$ from the germs of $\bar{\partial}_V$ -closed V -valued forms at x , to the germs of forms in the domain of $\bar{\partial}_{V,s}$.

Considering $\underline{H}^{0,\star}(\bar{\partial}_{V,s})$ to be a complex of sheaves with zero differential, α is a morphism of complexes that is an isomorphism on cohomology in degree $\star > 0$ by construction. Let $\underline{\text{cone}}(\alpha_V)$ be the mapping cone of α_V , with $\underline{\text{cone}}^{0,\star}(\alpha_V) = \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star}) \oplus \underline{H}^{0,\star-1}(\bar{\partial}_{V,s})$ and differential $d_{\text{cone}}(\omega, h) = (\bar{\partial}_V \omega, \alpha_V(\omega))$. It has vanishing cohomology in degree $\star > 1$. Define a complex of sheaves $\underline{\mathcal{C}}_V^{0,\star}$ by

$$\underline{\mathcal{C}}_V^{0,\star} = \begin{cases} \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,0}), & \star = 0, \\ \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,1}), & \star = 1, \\ \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star}) \oplus \underline{H}^{0,\star-1}(\bar{\partial}_{V,s}), & \star > 1, \end{cases} \quad (3.2)$$

where the differential in degree $\star = 0$ is $\bar{\partial}_V$, the differential in degree $\star = 1$ is $(\bar{\partial}_V, \alpha_V)$, and the differential in degrees $\star > 1$ is d_{cone} . Then $\underline{\mathcal{C}}_V$ is a resolution of $\underline{V}_s$ by fine sheaves.

There is a short exact sequence of sheaves

$$0 \rightarrow \underline{V} \rightarrow \underline{V}_s \rightarrow \underline{V}_s/\underline{V} \rightarrow 0. \quad (3.3)$$

We can think of $\underline{V}_s/\underline{V}$ as a resolution of itself, when concentrated in degree zero. Together with the resolution of $\underline{V}_s$ from (3.2), we can construct a resolution of $\underline{V}$ as follows. As $\underline{V}_s/\underline{V}$ is a finite sum of skyscraper sheaves, we can extend the quotient map $\underline{V}_s \rightarrow \underline{V}_s/\underline{V}$ to a morphism $\beta_V : \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,0}) \rightarrow \underline{V}_s/\underline{V}$. Define a complex of sheaves $\tilde{\mathcal{C}}_V$ by

$$\tilde{\mathcal{C}}_V^{0,\star} = \begin{cases} \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,0}), & \star = 0, \\ \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,1}) \oplus \underline{V}_s/\underline{V}, & \star = 1, \\ \underline{\text{Dom}}(\bar{\partial}_{V,s}^{0,\star}) \oplus \underline{H}^{0,\star-1}(\bar{\partial}_{V,s}), & \star > 1, \end{cases} \quad (3.4)$$

where the differential in degree $\star = 0$ is $(\bar{\partial}_V, \beta_V)$, the differential in degree $\star = 1$ sends (ω, v) to $(\bar{\partial}_V \omega, \alpha_V(\omega))$, and the differential in degrees $\star > 1$ is d_{cone} . Then $\tilde{\mathcal{C}}_V$ is a resolution of $\underline{V}$ by fine sheaves; see [Iversen 1986, proof of Proposition I.6.10].

Taking global sections of $\tilde{\mathcal{C}}_V^{0,\star}$ gives a cochain complex $(\tilde{T}_V, \tilde{d}_V)$:

$$\begin{aligned} 0 \rightarrow \text{Dom}(\bar{\partial}_{V,s}^{0,0}) &\rightarrow \text{Dom}(\bar{\partial}_{V,s}^{0,1}) \oplus (\underline{V}_s/\underline{V})(X) \\ &\rightarrow \text{Dom}(\bar{\partial}_{V,s}^{0,2}) \oplus (\underline{H}^{0,1}(\bar{\partial}_{V,s}))(X) \rightarrow \cdots \\ &\rightarrow \text{Dom}(\bar{\partial}_{V,s}^{0,n}) \oplus (\underline{H}^{0,n-1}(\bar{\partial}_{V,s}))(X) \rightarrow 0. \end{aligned} \quad (3.5)$$

For the last term, we use the fact that in terms of a resolution $\pi : M \rightarrow X$, we have $(\underline{H}^{0,n}(\bar{\partial}_{V,s}))_x = V_x \otimes (R^n \pi_* \mathcal{O}_M)_x = 0$.

Proposition 3.6. *The cohomology of $(\tilde{T}_V, \tilde{d}_V)$ is isomorphic to $H^*(X; \underline{V})$.*

Proof. This holds because $\tilde{\mathcal{C}}_V$ is a resolution of $\underline{V}$ by fine sheaves. $\square$

Put arbitrary inner products on the finite dimensional vector spaces $(\underline{V}_s/\underline{V})(X)$ and $(\underline{H}^{0,\star}(\bar{\partial}_{V,s}))(X)$.

4. Hilbert complex

The differential $\tilde{d}_V$ in the Hilbert complex $(\tilde{T}_V, \tilde{d}_V)$ of the previous section involved somewhat arbitrary choices of α_V and β_V . In this section we replace $(\tilde{T}_V, \tilde{d}_V)$ by a more canonical Hilbert complex (T_V, d_V) .

For brevity of notation, we put

$$A_V^* = \begin{cases} (\underline{V}_s/\underline{V})(X), & \star = 0, \\ (\underline{H}^{0,\star}(\bar{\partial}_{V,s}))(X), & \star > 0. \end{cases} \quad (4.1)$$

Then the complex $\tilde{T}_V$ has entries $\tilde{T}_V^{0,\star} = \text{Dom}(\bar{\partial}_{V,s}^{0,\star}) \oplus A_V^{\star-1}$. Combining α_V and β_V , we have constructed a linear map $\gamma_V : \text{Dom}(\bar{\partial}_{V,s}^{0,\star}) \rightarrow A_V^\star$ so that the differential of $\tilde{T}_V$ is given by

$$\tilde{d}_V(\omega, a) = (\partial_V \omega, \gamma_V(\omega)). \quad (4.2)$$

Note that $\gamma_V \circ \bar{\partial}_{V,s} = 0$.

Let $P_{\text{Ker}(\Delta_{V,s}^{0,\star})}$ be orthogonal projection onto $\text{Ker}(\Delta_{V,s}^{0,\star}) \subset \Omega_{L^2}^{0,\star}(X_{\text{reg}}; V)$. Define a new differential d_V on $\tilde{T}_V$ by

$$d_V(\omega, a) = (\partial_V \omega, \gamma_V(P_{\text{Ker}(\Delta_{V,s}^{0,\star})} \omega)). \quad (4.3)$$

Call the resulting cochain complex (T_V, d_V) .

As in (2.8), there is a Hodge decomposition

$$\text{Dom}(\bar{\partial}_{V,s}^{0,\star}) = \text{Ker}(\Delta_{V,s}^{0,\star}) \oplus \text{Im}(\bar{\partial}_{V,s}) \oplus \text{Im}(\bar{\partial}_{V,s}^*). \quad (4.4)$$

Here the terms on the right-hand side of (4.4) are the intersections of $\text{Dom}(\bar{\partial}_{V,s}^{0,\star})$ with the corresponding terms in (2.8). In particular, $\text{Ker}(\Delta_{V,s}^{0,\star})$ and $\text{Im}(\bar{\partial}_{V,s})$ are the same, while the elements of $\text{Im}(\bar{\partial}_{V,s}^*)$ now lie in an H^1 -space. Put

$$\mathcal{I}_V = \bar{\partial}_{V,s}|_{\text{Im}(\bar{\partial}_{V,s}^*)} : \text{Im}(\bar{\partial}_{V,s}^*) \rightarrow \text{Im}(\bar{\partial}_{V,s}), \quad (4.5)$$

an isomorphism.

Define a linear map $m_V : \text{Dom}(\bar{\partial}_{V,s}^{0,\star}) \oplus A_V^{\star-1} \rightarrow \text{Dom}(\bar{\partial}_{V,s}^{0,\star}) \oplus A_V^{\star-1}$ by saying that if

$$(h, \omega_1, \omega_2, a) \in \text{Ker}(\Delta_{V,s}^{0,\star}) \oplus \text{Im}(\bar{\partial}_{V,s}) \oplus \text{Im}(\bar{\partial}_{V,s}^*) \oplus A_V^{\star-1}, \quad (4.6)$$

then

$$m_V(h, \omega_1, \omega_2, a) = (h, \omega_1, \omega_2, a + \gamma_V(\mathcal{I}_V^{-1}(\omega_1))). \quad (4.7)$$

Its inverse is given by

$$m_V^{-1}(h, \omega_1, \omega_2, a) = (h, \omega_1, \omega_2, a - \gamma_V(\mathcal{I}_V^{-1}(\omega_1))). \quad (4.8)$$

Proposition 4.9. *The linear maps m_V and m_V^{-1} are chain maps between (T_V, d_V) and $(T_V, \tilde{d}_V)$, i.e., $m_V \circ d_V = \tilde{d}_V \circ m_V$ and $m_V^{-1} \circ \tilde{d}_V = d_V \circ m_V^{-1}$.*

Proof. We check that $m_V \circ d_V = \tilde{d}_V \circ m_V$; the proof that $m_V^{-1} \circ \tilde{d}_V = d_V \circ m_V^{-1}$ is similar. Given $(h, \omega_1, \omega_2, a)$ as in (4.6), we have

$$\begin{aligned} d_V(h, \omega_1, \omega_2, a) &= (0, \bar{\partial}_V \omega_2, 0, \gamma_V(h)), \\ m_V(d_V(h, \omega_1, \omega_2, a)) &= (0, \bar{\partial}_V \omega_2, 0, \gamma_V(h) + \gamma_V(\omega_2)), \\ m_V(h, \omega_1, \omega_2, a) &= (h, \omega_1, \omega_2, a + \gamma_V(\mathcal{I}_V^{-1}(\omega_1))), \\ \tilde{d}_V(m_V(h, \omega_1, \omega_2, a)) &= (0, \bar{\partial}_V \omega_2, 0, \gamma_V(h) + \gamma_V(\omega_2)). \end{aligned} \quad (4.10)$$

This proves the proposition. □

Theorem 4.11. *The cohomology of (T_V, d_V) is isomorphic to $H^*(X; \underline{V})$.*

Proof. This follows from Propositions 3.6 and 4.9. $\square$

We can now reprove a result from [Fulton 1998, Example 18.3.3 on p. 362].

Proposition 4.12. *In terms of a resolution $\pi : M \rightarrow X$, we have*

$$\sum_{i=0}^n (-1)^i \dim(H^i(X; \mathcal{O}_X)) = \int_M \text{Td}(TM) - \dim((\pi_* \mathcal{O}_M / \mathcal{O}_X)(X)) + \sum_{i=1}^n (-1)^{i-1} \dim((R^i \pi_* \mathcal{O}_M)(X)). \quad (4.13)$$

Proof. Let (T_1, d_1) denote the complex (T_V, d_V) when the vector bundle V is the trivial bundle. From Theorem 4.11, the left-hand side of (4.13) is the index of $d_1 + d_1^*$. We can deform the chain complex (T_1, d_1) to make the differential equal to $\bar{\partial}_s \oplus 0$ without changing the index. The new index is

$$\sum_{i=0}^n (-1)^i \dim(H^i(\bar{\partial}_s)) - \dim((\mathcal{O}_s / \mathcal{O}_X)(X)) + \sum_{i=1}^{n-1} (-1)^{i-1} \dim((\underline{H}^{0,i}(\bar{\partial}_s))(X)). \quad (4.14)$$

From [Pardon and Stern 1991], we have $H^i(\bar{\partial}_s) \cong H^{0,i}(M)$, so

$$\sum_{i=0}^n (-1)^i \dim(H^i(\bar{\partial}_s)) = \sum_{i=0}^n (-1)^i \dim(H^{0,i}(M)) = \int_M \text{Td}(TM). \quad (4.15)$$

From [Ruppenthal 2018, Corollary 1.2], $\mathcal{O}_s \cong \pi_* \mathcal{O}_M$ and $\underline{H}^{0,i}(\bar{\partial}_s) \cong R^i \pi_* \mathcal{O}_M$. The proposition follows. $\square$

Remark 4.16. We can write $\int_M \text{Td}(TM) = \int_X \pi_* \text{Td}(TM)$, where we are integrating a top-degree form on X_{reg} . It is not so clear what the relevant theory of characteristic classes on X should be, for which this would be an example. We have in mind a Chern–Weil theory on X_{reg} with control on how the forms behave near X_{sing} . We note that there is a rational homology class $\pi_*(PD[\text{Td}(TM)])$ on X , where $PD[\text{Td}(TM)] \in H_{\text{even}}(M; \mathbb{Q})$ is the Poincaré dual of $[\text{Td}(TM)] \in H^{\text{even}}(M; \mathbb{Q})$, and if X is connected then $\int_M \text{Td}(TM)$ can be identified with the degree-zero component of $\pi_*(PD[\text{Td}(TM)])$.

5. K-homology class

In this section we prove Theorem 1.5. We first show that if $\pi : M \rightarrow X$ is a resolution of singularities, with a simple normal crossing divisor, then the K-homology class $[\bar{\partial}_s + \bar{\partial}_s^*] \in K_0(X)$ from Proposition 2.1, with V trivial, equals the pushforward $\pi_*[\bar{\partial}_M + \bar{\partial}_M^*]$. We then prove Theorem 1.5.

Proposition 5.1. *Let $\pi : M \rightarrow X$ be a resolution of singularities, with $\pi^{-1}(X_{\text{sing}})$ being a simple normal crossing divisor. Then $[\bar{\partial}_s + \bar{\partial}_s^*] = \pi_*[\bar{\partial}_M + \bar{\partial}_M^*]$.*

Proof. The method of proof comes from [Haskell 1987]. Consider the following part of the K-homology exact sequence for the pair (X, X_{sing}) :

$$K_0(X_{\text{sing}}) \xrightarrow{\alpha} K_0(X) \xrightarrow{\beta} K_0(X, X_{\text{sing}}). \quad (5.2)$$

Lemma 5.3. *We have $\beta([\bar{\partial}_s + \bar{\partial}_s^*]) = \beta(\pi_*[\bar{\partial}_M + \bar{\partial}_M^*])$ in $K_0(X, X_{\text{sing}})$.*

Proof. Put $D = \pi^{-1}(X_{\text{sing}}) \subset M$. Since it has simple normal crossings, there is a small regular neighborhood of D whose closure C' is homotopy equivalent to D . We can also assume that $C = \pi(C')$ is homotopy equivalent to X_{sing} [Milnor 1968, Theorem 2.10]. As $[\bar{\partial}_M + \bar{\partial}_M^*]$ is independent of the choice of Hermitian metric on M , we can choose a Hermitian metric on M so that π restricts to an isometry from $M - C'$ to $X - C$.

Consider the commutative diagram

$$\begin{array}{ccccccc} K_0(M) & \longrightarrow & K_0(M, D) & \xrightarrow{\cong} & K_0(M, C') & \xrightarrow{\cong} & \text{KK}(C_0(M - C'); \mathbb{C}) \\ \pi_* \downarrow & & \downarrow & & \downarrow & & \downarrow \\ K_0(X) & \xrightarrow{\beta} & K_0(X, X_{\text{sing}}) & \xrightarrow{\cong} & K_0(X, C) & \xrightarrow{\cong} & \text{KK}(C_0(X - C); \mathbb{C}) \end{array} \quad (5.4)$$

Starting with $[\bar{\partial}_M + \bar{\partial}_M^*] \in K_0(M)$ and going along the top row, its image in $\text{KK}(C_0(M - C'); \mathbb{C})$ is the restriction of the analytic K-homology class, i.e., one only acts by functions that vanish on C' . The right vertical arrow of the diagram is an isomorphism coming from the bijection between $M - C'$ and $X - C$. By the commutativity of the diagram, we now know what $\beta(\pi_*[\bar{\partial}_M + \bar{\partial}_M^*])$ is as an element of $\text{KK}(C_0(X - C); \mathbb{C})$. However, this is isomorphic to the restriction of $[\bar{\partial}_s + \bar{\partial}_s^*] \in K_0(X)$ to an element of $\text{KK}(C_0(X - C); \mathbb{C})$ (since π gives an isometry between $M - C'$ and $X - C$). The latter restriction is the same as $\beta([\bar{\partial}_s + \bar{\partial}_s^*])$. This proves the lemma. $\square$

Returning to the proof of Proposition 5.1, we know now that $[\bar{\partial}_s + \bar{\partial}_s^*] - \pi_*[\bar{\partial}_M + \bar{\partial}_M^*]$ lies in the kernel of β , and so lies in the image of α . For the purpose of the proof, we can assume that X is connected. Let $a : \text{pt} \rightarrow X$ be an arbitrary fixed embedding and let $a_* : K_0(\text{pt}) \rightarrow K_0(X)$ be the induced homomorphism. The connectedness of X implies that $\text{Im}(\alpha) = \text{Im}(a_*)$. Let $b : X \rightarrow \text{pt}$ be the unique point map. Consider $\text{pt} \xrightarrow{a} X \xrightarrow{b} \text{pt}$ and the induced homomorphisms $K_0(\text{pt}) \xrightarrow{a_*} K_0(X) \xrightarrow{b_*} K_0(\text{pt})$. Then the map b_* restricts to an isomorphism between $\text{Im}(a_*)$ and $K_0(\text{pt})$. Hence, to prove the proposition, it suffices to show that $b_*[\bar{\partial}_s + \bar{\partial}_s^*] = b_*(\pi_*[\bar{\partial}_M + \bar{\partial}_M^*])$ in $K_0(\text{pt}) \cong \mathbb{Z}$.

Now $b_*[\bar{\partial}_s + \bar{\partial}_s^*]$ is the index of $\bar{\partial}_s + \bar{\partial}_s^*$, i.e., $\sum_{i=0}^n (-1)^i \dim(H^i(\bar{\partial}_s))$, while $b_*(\pi_*[\bar{\partial}_M + \bar{\partial}_M^*])$ is the index of $\bar{\partial}_M + \bar{\partial}_M^*$, i.e., $\sum_{i=0}^n (-1)^i \dim(H^i(\bar{\partial}_M))$. From [Pardon and Stern 1991], these are equal term-by-term. This proves the proposition. $\square$

Proof of Theorem 1.5. Suppose that X is a connected projective algebraic variety. In terms of the resolution $\pi : M \rightarrow X$, it was pointed out in [Baum et al. 1975, p. 104] that there is an identity in $K_0(X)$:

$$[\mathcal{O}_X]_{\text{BFM}} - \pi_*[\mathcal{O}_M]_{\text{BFM}} = \sum_j n_j [\mathcal{O}_{V_j}]_{\text{BFM}}. \quad (5.5)$$

Here the n_j are certain integers and the V_j are irreducible subvarieties of the singular locus of X . In our case of isolated singularities, the V_j are just the points x_j in X_{sing} . As $[\mathcal{O}_M]_{\text{BFM}} = [\bar{\partial}_M + \bar{\partial}_M^*]$, Proposition 5.1 implies that

$$[\mathcal{O}_X]_{\text{BFM}} = [\bar{\partial}_s + \bar{\partial}_s^*] + \sum_j n_j [\mathcal{O}_{V_j}]_{\text{BFM}}. \quad (5.6)$$

Let (T_1, d_1) denote the complex (T_V, d_V) when the vector bundle V is the trivial bundle. Let $[\mathcal{O}_X]_{\text{an}} \in K_0(X)$ be the K-homology class coming from the operator $d_1 + d_1^*$. We can deform the chain complex (T_1, d_1) to make the differential equal to $\bar{\partial}_s \oplus 0$ without changing the K-homology class arising from the complex. Then (5.6) implies that $[\mathcal{O}_X]_{\text{an}}$ and $[\mathcal{O}_X]_{\text{BFM}}$ have the same image in $K_0(X, X_{\text{sing}})$; cf. the proof of Lemma 5.3. Let $b : X \rightarrow \text{pt}$ be the unique point map. As in the proof of Proposition 5.1, to conclude that $[\mathcal{O}_X]_{\text{an}} = [\mathcal{O}_X]_{\text{BFM}}$ in $K_0(X)$, it now suffices to show that $b_*[\mathcal{O}_X]_{\text{an}} = b_*[\mathcal{O}_X]_{\text{BFM}}$ in $K_0(\text{pt}) \cong \mathbb{Z}$. Now $b_*[\mathcal{O}_X]_{\text{an}}$ is the index of $d_1 + d_1^*$ which, from Theorem 4.11, equals the arithmetic genus $\sum_{i=0}^n (-1)^i \dim(H^i(X; \mathcal{O}_X))$. On the other hand, from [Baum et al. 1979, Section 3], we also have $b_*[\mathcal{O}_X]_{\text{BFM}} = \sum_{i=0}^n (-1)^i \dim(H^i(X; \mathcal{O}_X))$. This proves the theorem. $\square$

Remark 5.7. We mention some of the issues involved in extending the present paper to nonisolated singularities. First, it seems to be open whether $\bar{\partial}_s + \bar{\partial}_s^*$ has compact resolvent, so the unbounded KK-formalism may not be applicable. However, it is known that the unreduced cohomology of the $\bar{\partial}_s$ -complex is finite dimensional, being isomorphic to the cohomology of a resolution [Pardon and Stern 1991]. Hence the $\bar{\partial}_s$ -complex is Fredholm and one could use the bounded KK-description of K-homology, although it would be more cumbersome.

We expect that Proposition 5.1 still holds if X has nonisolated singularities. It is known that taking resolutions $\pi : M \rightarrow X$, the pushforward $\pi_*[\bar{\partial}_M + \bar{\partial}_M^*] \in K_0(X)$ is independent of the choice of resolution [Hilsum 2018].

One could ask for an extension of Theorem 4.11 to the case of nonisolated singularities. As an indication, one would expect that taking products of complex spaces would lead to tensor products of the cochain complexes. In particular, suppose that Z is a smooth Hermitian manifold and X has isolated singular points. Then the cochain complex for $Z \times X$ would have contributions from differential forms along the singular locus.

In a related vein, in principle one can apply (5.5) inductively to get an expression for $[\mathcal{O}_X]_{\text{BFM}}$.

Acknowledgments

I thank Paul Baum and Peter Haskell for discussions. I especially thank Peter for pointing out the relevance of [Haskell 1987]. I also thank the referee for helpful comments.

References

- [Ancona and Gaveau 1994] V. Ancona and B. Gaveau, “Differential forms and resolution on certain analytic spaces, III: Spectral resolution”, *Ann. Mat. Pura Appl.* (4) **166** (1994), 175–202. MR Zbl
- [Andersson and Samuelsson 2012] M. Andersson and H. Samuelsson, “A Dolbeault–Grothendieck lemma on complex spaces via Koppelman formulas”, *Invent. Math.* **190**:2 (2012), 261–297. MR Zbl
- [Atiyah 1970] M. F. Atiyah, “Global theory of elliptic operators”, pp. 21–30 in *Proceedings of the International Conference on Functional Analysis and Related Topics* (Tokyo, 1969), University of Tokyo Press, 1970. MR Zbl
- [Atiyah et al. 1973] M. F. Atiyah, V. K. Patodi, and I. M. Singer, “Spectral asymmetry and Riemannian geometry”, *Bull. London Math. Soc.* **5** (1973), 229–234. MR Zbl
- [Baaj and Julg 1983] S. Baaj and P. Julg, “Théorie bivariante de Kasparov et opérateurs non bornés dans les C^* -modules hilbertiens”, *C. R. Acad. Sci. Paris Sér. I Math.* **296**:21 (1983), 875–878. MR Zbl
- [Baum and Douglas 1982] P. Baum and R. G. Douglas, “ K homology and index theory”, pp. 117–173 in *Operator algebras and applications, I* (Kingston, ON, 1980), edited by R. V. Kadison, Proc. Sympos. Pure Math. **38**, Amer. Math. Soc., Providence, RI, 1982. MR Zbl
- [Baum et al. 1975] P. Baum, W. Fulton, and R. MacPherson, “Riemann–Roch for singular varieties”, *Inst. Hautes Études Sci. Publ. Math.* **45** (1975), 101–145. MR Zbl
- [Baum et al. 1979] P. Baum, W. Fulton, and R. MacPherson, “Riemann–Roch and topological K theory for singular varieties”, *Acta Math.* **143**:3–4 (1979), 155–192. MR Zbl
- [Baum et al. 2007] P. Baum, N. Higson, and T. Schick, “On the equivalence of geometric and analytic K -homology”, *Pure Appl. Math. Q.* **3**:1 (2007), 1–24. MR Zbl
- [Bei and Piazza 2019] F. Bei and P. Piazza, “On analytic Todd classes of singular varieties”, preprint, 2019. arXiv
- [Brüning and Lesch 1992] J. Brüning and M. Lesch, “Hilbert complexes”, *J. Funct. Anal.* **108**:1 (1992), 88–132. MR Zbl
- [Brüning et al. 1990] J. Brüning, N. Peyerimhoff, and H. Schröder, “The $\bar{\partial}$ -operator on algebraic curves”, *Comm. Math. Phys.* **129**:3 (1990), 525–534. MR Zbl
- [Forsyth et al. 2014] I. Forsyth, B. Mesland, and A. Rennie, “Dense domains, symmetric operators and spectral triples”, *New York J. Math.* **20** (2014), 1001–1020. MR Zbl
- [Fox and Haskell 2000] J. Fox and P. Haskell, “Perturbed Dolbeault operators and the homology Todd class”, *Proc. Amer. Math. Soc.* **128**:12 (2000), 3715–3721. MR Zbl
- [Fulton 1998] W. Fulton, *Intersection theory*, 2nd ed., *Ergebnisse der Mathematik (3)* **2**, Springer, 1998. MR Zbl

- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Math. **52**, Springer, 1977. MR Zbl
- [Haskell 1987] P. Haskell, “Index theory of geometric Fredholm operators on varieties with isolated singularities”, *K-Theory* **1**:5 (1987), 457–466. MR Zbl
- [Hilsum 2018] M. Hilsum, “Une preuve analytique de la conjecture de J. Rosenberg”, preprint, 2018, available at <https://hal.archives-ouvertes.fr/hal-01841905>.
- [Iversen 1986] B. Iversen, *Cohomology of sheaves*, Springer, 1986. MR Zbl
- [Kaad 2019] J. Kaad, “On the unbounded picture of KK -theory”, preprint, 2019. arXiv
- [Milnor 1968] J. Milnor, *Singular points of complex hypersurfaces*, Ann. of Math. Studies **61**, Princeton University Press, 1968. MR Zbl
- [Øvrelid and Ruppenthal 2014] N. Øvrelid and J. Ruppenthal, “ L^2 -properties of the $\bar{\partial}$ and the $\bar{\partial}$ -Neumann operator on spaces with isolated singularities”, *Math. Ann.* **359**:3–4 (2014), 803–838. MR Zbl
- [Pardon and Stern 1991] W. L. Pardon and M. A. Stern, “ L^2 - $\bar{\partial}$ -cohomology of complex projective varieties”, *J. Amer. Math. Soc.* **4**:3 (1991), 603–621. MR Zbl
- [Ruppenthal 2018] J. Ruppenthal, “ L^2 -Serre duality on singular complex spaces and rational singularities”, *Int. Math. Res. Not.* **2018**:23 (2018), 7198–7240. MR Zbl
- [Singer 1971] I. M. Singer, “Future extensions of index theory and elliptic operators”, pp. 171–185 in *Prospects in mathematics* (Princeton, NJ, 1970), Ann. of Math. Studies **70**, Princeton University Press, 1971. MR Zbl
- [Whitney 1972] H. Whitney, *Complex analytic varieties*, Addison-Wesley, Reading, MA, 1972. MR Zbl

Received 23 Apr 2019. Revised 20 Jun 2019. Accepted 11 Jul 2019.

JOHN LOTT: lott@berkeley.edu

Department of Mathematics, University of California, Berkeley, CA, United States

ANNALS OF K-THEORY

msp.org/akt

EDITORIAL BOARD

Joseph Ayoub	ETH Zürich, Switzerland joseph.ayoub@math.uzh.ch
Paul Balmer	University of California, Los Angeles, USA balmer@math.ucla.edu
Guillermo Cortiñas	Universidad de Buenos Aires and CONICET, Argentina gcorti@dm.uba.ar
Hélène Esnault	Freie Universität Berlin, Germany liveesnault@math.fu-berlin.de
Eric Friedlander	University of Southern California, USA ericmf@usc.edu
Max Karoubi	Institut de Mathématiques de Jussieu – Paris Rive Gauche, France max.karoubi@imj-prg.fr
Moritz Kerz	Universität Regensburg, Germany moritz.kerz@mathematik.uni-regensburg.de
Huaxin Lin	University of Oregon, USA livehlin@uoregon.edu
Alexander Merkurjev	University of California, Los Angeles, USA merkurev@math.ucla.edu
Birgit Richter	Universität Hamburg, Germany birgit.richter@uni-hamburg.de
Jonathan Rosenberg	(Managing Editor) University of Maryland, USA jmr@math.umd.edu
Marco Schlichting	University of Warwick, UK schlichting@warwick.ac.uk
Charles Weibel	(Managing Editor) Rutgers University, USA weibel@math.rutgers.edu
Guoliang Yu	Texas A&M University, USA guoliangyu@math.tamu.edu

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

Annals of K-Theory is a journal of the K-Theory Foundation (ktheoryfoundation.org). The K-Theory Foundation acknowledges the precious support of Foundation Compositio Mathematica, whose help has been instrumental in the launch of the Annals of K-Theory.

See inside back cover or msp.org/akt for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$550/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Annals of K-Theory (ISSN 2379-1681 electronic, 2379-1683 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

AKT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY



mathematical sciences publishers
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANNALS OF K-THEORY

2019

vol. 4

no. 4

On the intersection motive of certain Shimura varieties: the case of Siegel threefolds	525
JÖRG WILDESHAUS	
A Baum–Connes conjecture for singular foliations	561
IAKOVOS ANDROULIDAKIS and GEORGES SKANDALIS	
Witt groups of abelian categories and perverse sheaves	621
JÖRG SCHÜRMANN and JON WOOLF	
A generalized Vaserstein symbol	671
TARIQ SYED	
A Dolbeault–Hilbert complex for a variety with isolated singular points	707
JOHN LOTT	