

JGR Space Physics

RESEARCH ARTICLE

10.1029/2021JA029248

Key Points:

- Coordinated observations of atmospheric waves from Global-scale Observations of the Limb and Disk and ionospheric total electron content perturbations are presented
- Perturbations consistent with a large-scale wave are seen moving horizontally at the same rate in both data sets
- The amplitude and phase relationship between atmospheric and ionospheric parameters are not always clear

Correspondence to:

S. L. England,
englands@vt.edu

Citation:

England, S. L., Greer, K. R., Zhang, S.-R., Evans, S., Solomon, S. C., Eastes, R. W., et al. (2021). First comparison of traveling atmospheric disturbances observed in the middle thermosphere by Global-scale Observations of the Limb and Disk to traveling ionospheric disturbances seen in ground-based total electron content observations. *Journal of Geophysical Research: Space Physics*, 126, e2021JA029248. <https://doi.org/10.1029/2021JA029248>

Received 12 FEB 2021
Accepted 10 MAY 2021

First Comparison of Traveling Atmospheric Disturbances Observed in the Middle Thermosphere by Global-Scale Observations of the Limb and Disk to Traveling Ionospheric Disturbances Seen in Ground-Based Total Electron Content Observations

Scott L. England¹ , Katelynn R. Greer² , Shun-Rong Zhang³ , Scott Evans⁴, Stanley C. Solomon⁵ , Richard W. Eastes² , William E. McClintock² , and Alan G. Burns⁵ 

¹Aerospace and Ocean Engineering, Virginia Polytechnic Institute and State University, Blacksburg, VA, USA,

²Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, CO, USA, ³MIT Haystack Observatory, Westford, MA, USA, ⁴Computational Physics, Inc., Springfield, VA, USA, ⁵High Altitude Observatory, National Center for Atmospheric Research, Boulder, CO, USA

Abstract Traveling ionospheric disturbances (TIDs) and their neutral counterparts known as traveling atmospheric disturbances (TADs) are believed to play a role in communicating inputs to other locations in the fluid. While these two phenomena are believed to be connected, they may not have a one-to-one correspondence as the geomagnetic field influences the TID but has no direct impact on the TAD. The relative amplitudes of the perturbations seen in the ionosphere and atmosphere have been observed but rarely together. This study reports results from a 3-day campaign to observe TIDs and TADs simultaneously over a broad latitudinal region over the eastern United States using a combination of Global-scale Observations of the Limb and Disk (GOLD) and a distributed network of ground-based Global Navigation Satellite System (GNSS) receivers. These results demonstrate that GOLD and the ground-based total electron content (TEC) observations can see the atmospheric and ionospheric portions of a large-scale traveling disturbance. The phase difference in the perturbations to the GOLD airglow brightness, O/N₂ and thermospheric disk temperature are consistent with an atmospheric gravity wave moving through this region. The ionospheric signatures move at the same rate as those in the atmosphere, but their amplitudes do not have a simple correspondence to the amplitude of the signal seen in the atmosphere. This campaign demonstrates a proof-of-concept that this combination of observations is able to provide information on TIDs and TADs, including quantifying their impact on the temperature and chemical composition of the upper atmosphere.

Plain Language Summary Waves are a ubiquitous feature of the Earth's atmosphere. Just as with waves on the surface of water, these act to transport energy and momentum from the region in which they are produced to other parts of the fluid. A specific type of these waves in the upper atmosphere, often with spatial scales of hundreds to thousands of km, are known as traveling atmospheric disturbances (TADs). A signature that is often associated with these TADs is seen in the charged particle environment around the Earth, that are referred to as traveling ionospheric disturbances (TIDs). While it is widely accepted that TADs and TIDs are related, the exact relationship between their amplitudes and other quantities is not well established. This paper describes a campaign during which observations of the atmosphere (from NASA's GOLD mission) and charged particles (from ground-based observations) are combined in order to identify the relationship between the TAD and TID parameters. The GOLD observations of the perturbations to the upper atmospheric temperature, composition and its emissions are all consistent with what is known about TADs. At the same time, the ground-based observations of the charged particles reveal a TID that moves at the same speed as the TAD.

1. Introduction

Atmospheric gravity waves are believed to play some role in redistributing energy and momentum within the Earth's atmosphere and communicating impacts in one region to physically distant locations. In the

thermosphere, gravity waves are believed to be a key component of the wave spectrum that is present, and can significantly impact this region through dissipating momentum, generating mixing, local heating, and flux of heat and minor constituents (e.g., Yiğit & Medvedev, 2015). Many of the gravity waves seen in the thermosphere are believed to have propagated up from the lower atmosphere (as both primary and secondary waves, e.g., Vadas & Fritts, 2006; Vadas & Liu, 2009; Yiğit et al., 2014), while others are believed to have been generated at thermospheric altitudes through processes such as Joule heating in the auroral regions (e.g., Richmond, 1978) and the solar terminator (e.g., Galushko et al., 1998). During geomagnetic storms, particularly large amplitude gravity waves may be generated by impulsive heating of the atmosphere at high-latitudes (e.g., Shiokawa et al., 2007).

Atmospheric gravity waves are often described alongside their ionospheric counterparts, where the former is referred to as traveling atmospheric disturbance (TAD) and the latter as a traveling ionospheric disturbance (TID). TIDs are typically grouped into two categories according to their horizontal wavelengths/spatial sizes. Large-scale TIDs are typically classified as having wavelengths over 1,000 km. Large-scale TIDs are often believed to be accompanied by TADs generated in the auroral regions (e.g., Bruinsma et al., 2006). Medium-scale TIDs are typically classified as having horizontal wavelengths in the 100–1,000 km range. These medium-scale TIDs are believed to often be accompanied by TADs generated in the lower regions of the atmosphere (e.g., Azeem et al., 2017), while others are not (especially at midlatitude during the night).

Surveys of both atmospheric and ionospheric observations have revealed that TIDs/TADs are present under almost all conditions, at some amplitude. At the midlatitudes that will be considered in this study, results such as Livneh et al. (2009) have revealed TID-like fluctuations in the ionosphere that are omnipresent, with periods in the 30–60-min range. Tsugawa et al. (2007) presented an analysis of ground-based total electron content (TEC) variations, showing abundant waves with horizontal wavelengths in the 300–1,000 km range over North America, consistent with medium-scale TIDs. Recently, Tsuchiya et al. (2020) used 10 years of ground-based observations from a variety of sites and found a variation in the horizontal wavenumber spectrum observed with location. Mandal and Pallamraju (2020) also identified that the vertical propagation of gravity waves in the thermosphere varies with season at low-latitudes, which may impact the ability of some waves originating the lower atmosphere to form TAD and TID perturbations.

Direct observations of TADs are relatively rare, compared to the broad survey ionospheric data such as ground-based TEC that have permitted extensive studies of TIDs. In-situ atmospheric observations of TADs are most commonly available in the upper thermosphere where sustained spaceflight is possible (e.g., Bruinsma & Forbes, 2008; Garcia et al., 2016; Park et al., 2014). A large fraction of the observations at altitudes below these orbital altitudes come from observations of TIDs, from which the existence and properties of a TAD are inferred (e.g., Djuth et al., 2010; Galushko et al., 1998). Negrea et al. (2016) used an extensive ionospheric data set and established that the TID characteristics seen were consistent with the theoretical properties of acoustic gravity waves. Concurrent ion and neutral observations of waves have been made in situ (e.g., Earle et al., 2008; Paulino et al., 2018) which have demonstrated that the TIDs are often a good proxy for the TAD. Given that TID observations are readily available and that they are often used to infer the TAD, further study of the relationship between these two using simultaneous observations is of importance and will be the focus of this study.

NASA's Global-scale Observations of the Limb and Disk (GOLD) mission of opportunity provides measurements of the middle thermosphere over the daylit disk of the Earth visible from geostationary orbit over the Amazon river delta (Eastes et al., 2017). It has been demonstrated that by using a dedicated campaign mode, GOLD is able to observe TADs as they move through this region (England et al., 2020). This provides thermospheric measurements over a latitudinally broad region which, when combined with distributed ground-based ionospheric observations, offers an opportunity to examine the relationship between TIDs and TADs over low to midlatitudes. Furthermore, the ability for both GOLD and distributed ground-based receivers to measure this region continuously offers the opportunity to remove any spatial-temporal ambiguity that can come into play with low earth orbit spacecraft data. England et al. (2020) presented results of an extremely limited (6 h) campaign that took place during the commissioning phase of the GOLD instrument. The selection of the viewing location on the disk meant that virtually no ground-based data were available during this campaign. Here, we report results of a follow-on campaign that built upon lessons learned and focused on establishing how well we can compare the TADs observed by GOLD to TIDs seen in




Figure 1. Location of the GOLD Channel B observations during the campaign. A purple plus symbol marks the location of each of the 376 data bins along the length of the image. The light blue dots show the locations of the GPS dTEC data that are focused on in this study. dTEC, differential total electron content.

ground-based TEC observations. Key changes to the campaign setup will be described in more detail in the next section, but include: (a) extending the duration from 1 to 3 days; (b) providing overlap with ground sites; (c) identification of perturbations to both the N_2 airglow and the temperature of the middle thermosphere, which provide substantially more data with which to characterize the waves present. While previous studies have compared spatial features in the GOLD nighttime data (which are primarily ionospheric) with ground-based TEC observations (e.g., Cai et al., 2020), this study will focus on spatial features seen in the GOLD daytime data (which are primarily thermospheric) with ground-based TEC.

2. Experimental Setup and Data Used

This section will discuss the observational campaign performed by GOLD, and its associated data; the geophysical conditions present during the campaign; and the ground-based TEC data used to provide large-scale context and daytime ionospheric parameters.

2.1. Experimental Setup

GOLD is a 2-channel far-ultraviolet (FUV) imaging spectrometer that observes the Earth from onboard the SES-14 geostationary communications satellite, which is located at -47.5° longitude (Eastes et al., 2017, 2019; McClintock et al., 2017). During regular operations, when GOLD's field-of-view (FOV) is illuminated by the sun, the instrument scans the disk of the Earth, building up an image of the Earth's full disk with a 30-min cadence. GOLD makes observations between ~ 134 and 167 nm, which allows identification of the prominent dayglow features of O at 135.6 nm and LBH bands of N_2 . While the ratio of the brightness of these two features conveys information related to the composition of the thermosphere (column O/ N_2 ratio), the shape of the N_2 LBH emission bands can be used to determine the temperature of the thermosphere. During October 17–19, 2019, Channel B of the GOLD instrument was used to perform a special mode campaign, designed to identify the impacts of atmospheric waves on the middle thermosphere and their relation to similar fluctuations in the ionosphere. This basic experimental setup follows that described in England et al. (2020), but with several important changes. The prior campaign demonstrated that GOLD is able to detect wave-like signatures in the airglow when it changes from regular operations in which it scans the Earth's disk, and instead stares at one location (one near vertical line as shown in Figure 1) for several hours. This staring has the effect of increasing the signal by a factor of over 100 times compared to the scanning mode while sacrificing the instrument's ability to continue regular data collection. To both meet this need for a non-scanning mode, and not interrupt the regular GOLD data collection, a single channel of the instrument (Channel B) was used for this campaign. As regular operations were not interrupted, this enabled a longer campaign, spanning 3 consecutive days (October 17–19, 2019), with campaign data taken with Channel B from 12:30 to 20:00 h UTC. As these special campaign operations require a departure from the regular operating sequence for GOLD Channel B, a limit of 3 days was selected to balance both the impact on instrument commanding and the desire to determine how repeatable any effects seen may be from 1 day to the next. In the initial campaign described by England et al. (2020), the availability of both GOLD channels allowed for the identification of the wave propagation direction, velocity, and wavelength. With only a single channel, this is not possible. To both meet this need and provide information on any ionospheric perturbations that were not available during the initial campaign, concurrent data from ground-based observatories were considered when designing this campaign. By positioning the Channel B slit off-nadir, centered on the United States East Coast region, many sites observing TEC using GNSS radio signals are available and fill both needs described above. Figure 1 shows the location of the observations from Channel B that will be the focus of this study. The Level 1b data from this campaign are available from <https://gold.cs.ucf.edu/data/campaigns/>, and other Level 2 data are available from <https://gold.cs.ucf.edu/data>.

2.2. GOLD Airglow Brightness

To determine the fluctuations in the airglow brightness, this study uses the Level 1b (L1b) data from the GOLD instrument, which are described in the Data Products Users Guide (see <https://gold.cs.ucf.edu>). The L1b files are available for this campaign at 2-min cadence and provide the spectra in instrument counts versus wavelength and latitude. The L1b processing includes geometric corrections for the detector and optics, filtering of the counts based on the detector pulse heights, a correction based on the detector deadtime (maximum count rate) and data are binned on a regular wavelength scale. England et al. (2020) also used the Level 1b data, which provide the observed signal with the corrections described above at the highest available temporal and spatial cadence (spatial resolution of 48-km zonal by 17-km meridional at nadir and 2-min temporal cadence). From that prior study, it is evident that the perturbations to the airglow are relatively small in amplitude (below 1%) and are able to be seen in the L1b data that we will use here.

2.3. GOLD Disk Temperatures

GOLD makes measurements of the thermospheric temperature by observing the shape of the N₂ LBH bands in the ~137–160-nm wavelength range (see Eastes et al., 2017). This technique was first demonstrated by Aksnes et al. (2006) using limb observations from the High-resolution Ionospheric and Thermospheric Spectrograph (HITS) on the Advanced Research and Global Observation Satellite (ARGOS). The LBH emission provides information on the neutral temperatures in the middle thermosphere, with the maximum weighting to an altitude around 160 km (i.e., near the altitude of peak emission). Here, we consider these disk temperature data, which are Level 2 products. For the disk scanning mode (e.g., Figure 1), these are available at a 30-min cadence, but for Channel B these are available at a 10-min cadence, with a resolution corresponding to 40 spatial pixels along the slit for which temperature can be derived (emission angle constraints prevent determining temperatures very close to the limb as seen in Figure 1b). At the higher count rate per spatial-temporal bin that is obtained during the special mode campaign, the average random uncertainty per bin is 9.4 K (discussed further in Section 3.3).

2.4. GOLD Column O/N₂ Ratios

GOLD makes measurements of the thermospheric column O/N₂ ratio by observing the ratio of the brightness of the 135.6 nm O doublet to the brightness of the N₂ LBH band in the ~137–160-nm wavelength range (see Eastes et al., 2017). Use of disk observations of these emissions to determine the thermospheric column O/N₂ ratio was first demonstrated by Strickland et al. (1995) and has been widely used with data such as from TIMED-GUVI (e.g., Zhang et al., 2004). The O/N₂ ratio is indicative of the ratio of the column density of these two species from infinity down to an altitude corresponding to an N₂ column density of 10¹⁷ cm⁻², which is typically around 140 km. Here, we use the GOLD Level 2 O/N₂ data. These are available at the same resolution and cadence as the disk temperatures described above.

2.5. Differential TEC

Ground-based estimates of TEC from GNSS radio signals are routinely produced using algorithms developed at MIT Haystack Observatory (Rideout & Coster, 2006; Vierinen et al., 2016). Previous studies have used these data to identify traveling ionospheric disturbances, which have revealed that this is most easily detected in differential TEC (dTEC) values, rather than the absolute TEC values (e.g., Zhang et al., 2019). Following the methodology in Zhang et al. (2019), we adopt a cut-off elevation angle of 30°, below which ground-satellite ray paths are excluded. Differential TEC values are computed by first removing the background TEC using a low-pass Savitzky-Golay filter (Savitzky & Golay, 1964). The Savitzky-Golay filter uses a 60-min running window of data to eliminate artifacts related to the motion of the GNSS satellites and identify the slowly varying background TEC. The detrended TEC is then found by subtracting this filtered data from the original. This detrending analysis is applied to each of the TEC data segments of individual line-of-sight measurements between a GNSS satellite and a receiver. Examples of these data are shown in Figure 2 as functions of time and latitude at three different longitudes around the GOLD observations, plus a region that follows the location of the GOLD slit to within ±5° of longitude, corresponding to the symbols shown in Figure 1.




Figure 2. Global Navigation Satellite System differential total electron content (dTEC) for October 18, 2019 as functions of latitude and universal time. The range of longitudes included is indicated in panels (a–c). Panel (d) shows the dTEC following the locations of the GOLD slit, as described in the text and shown in Figure 1. The overlaid black lines in panels (b and d) mark locations of local maxima used to find the approximate phase velocity of the features seen, as described in the text.

2.6. Geophysical Conditions

The F10.7 index (available from NASA's OMNIWeb at <https://omniweb.gsfc.nasa.gov>) is between 64 and 66 throughout the 3 days of the special mode campaign, representing extremely low and steady solar EUV flux values. Of more note is the impact that any heating of the thermosphere associated with the auroral electrojet may have, as this is believed to be a major source of TADs (e.g., Shiokawa et al., 2007). Figure 3 shows the 1-min SME auroral electrojet index from SuperMAG (Gjerloev, 2012; Newell & Gjerloev, 2011) for October 17–19, 2019. The overall SME index is moderate throughout, but it is worth noting that there is a slightly elevated electrojet on the third day, both before and during the hours of the daytime thermospheric observations.

3. Method and Results

This section describes the analysis of each data set, starting with the differential TEC which is used to guide the analysis of the GOLD airglow and temperature data. The middle day of the campaign (October 18, 2019) will be shown throughout to demonstrate the methodology, and results from all 3 days, combining all three data sets are shown in Figure 6.




Figure 3. One-minute SME Auroral Electrojet Index for the 3 days of the campaign during October 2019. The time-periods corresponding to the GOLD observations are highlighted with the yellow-shaded regions.

3.1. Differential TEC

Given the elevation angle constraint described in Section 2.4, the dTEC data are confined to the land and coastal regions. As seen in Figure 1, the GOLD observations cross over the land and ocean. The greatest overlap in the available data between the TEC and GOLD observations occurs over the east coast of the United States, corresponding to panel d of Figure 2. This region has good data coverage from around 35° to 50° north. Further north, there is very little dTEC data available, and further south the GOLD FOV lies over the ocean where few TEC observations are available (see Figure 1). Hence, we will focus primarily on the variations seen in this latitude band at this time.

As noted in previous studies, the dTEC observations often reveal wave-like oscillations associated with TIDs. Many of these are evident in Figure 2, with different propagation directions and periods. Using the data from the three adjacent longitude regions (panels a–c), it is possible to characterize the propagation and scale sizes of some of these oscillations. Focusing on the latitude region from 35° to 50° north, clear features are seen near 18 and 22 h UT moving in a southward direction in all three panels (highlighted in panel b), with a third feature around 14 h UT visible in panels a and b. As these features are clear in all three panels and appear to be the strongest observed, we will focus on these. Using the data shown in all three panels, it is possible to estimate the phase speed of the TID seen. This is done in four steps. (1) Points close to the positive phases of the wave are identified by eye, and these are used as initial inputs. (2) Data for 1 h around each location are then chosen. (3) A cubic-spline interpolation to these values is used to provide data at regular spacing and the local maxima is identified. (4) The results of step 3 are compared to the location of the maximum value in the data from step 2 to ensure the results of step 3 are not numerical artifacts. A least-squares linear fit to these is used to determine an estimate of the latitudinal phase velocity of this TID at $\sim 7^\circ/\text{h}$ (shown with the solid black lines in panel b).

Leveraging the other available dTEC data in panels a and c, it is possible to identify the horizontal wavelength and phase velocity of the observed TID. This is done by using the observations from the three longitudes shown, and assuming that the wave has plane wavefronts. The phase difference between the three longitudes provides information on the azimuth of the wave phase velocity. In this case, we see no appreciable difference, indicating the wave propagation direction is nearly due south. With this, and estimating the dTEC observations to be coming from around 300-km altitude and above, $7^\circ/\text{h}$ corresponds to a phase speed of ~ 210 m/s. The wavelength estimated in this way is within the range of large-scale TIDs reported previously (see Section 1), although it is worth noting that the phase velocity identified is slow compared to many larger-scale TIDs reported. The wave speed and heading are indicative of a wave source in the auroral region, but a lower-atmospheric source or secondary gravity wave remain other plausible possibilities.

Clearly other wave features are seen in Figure 2. A full analysis of all of these is beyond the scope of this study. Instead, we will focus on this clearest wave feature and investigate if a corresponding signature in the atmosphere is observed simultaneously by GOLD. This will be done in-part by using the propagation rate determined from the dTEC observations and comparing this to that seen in the GOLD observations. As the GOLD slit is not perfectly aligned with the geographic meridional direction, it is important to select




Figure 4. Example spectrum made from summing 100 Level 1b exposures from the GOLD data from October 18, 2019. The light blue symbols highlight the region used to identify the atomic oxygen feature and the dark blue highlight the corresponding background area. The yellow symbols highlight the region used to identify the nitrogen features and the dark purple highlight the corresponding background area.

manner to the LBH bands; Zhang et al., 2018) are identified. These are highlighted in Figure 4 by the light blue and yellow symbols. The brightness in each of these (measured in Level 1b counts) is summed up at each spatial pixel and each point in time. To determine the background level, wavelength ranges adjacent to these are used. These are highlighted in Figure 4 by the dark blue and purple symbols. The brightness in each of these is summed up and the value subtracted from the in-band emissions found above (after normalizing for the number of spectral bins). Step 2: Identify and remove the strong local-time variation associated with solar zenith angle (note that the emission angle is constant for each pixel during this campaign). This is done by fitting a third-order polynomial to the time series of data for the two emissions identified above at each spatial pixel. The brightness variations associated with this low-order fit are then removed. It is worth noting that this third-order polynomial performs the same basic function as the removal of the Savitzky-Golay-filtered TEC data in that both act to remove the diurnally varying background signal (in TEC and in airglow brightness). Step 3: Identify and remove any change in brightness that occurs at all points along the disk simultaneously that are associated with small changes in solar EUV flux during the day. The mean value of the brightness across the pixels on the image is found and removed from the detrended data. The output of steps 2 and 3 is a residual signal with a mean value of zero. Following England et al. (2020), we then reduce the extremely noisy 2-min L1b residual data with a 10-min rolling median filter on the time series at each spatial pixel. This removes outlying high and low data points, while preserving a much higher temporal cadence than the oscillations seen in the dTEC values in Figures 2 and 5.

Figure 5 panels a and b show the detrended L1b values for the oxygen and nitrogen emissions, respectively. The large variations in brightness seen at the very northernmost extent of the GOLD data, above $\sim 60^\circ$ latitude could be associated with variations in the aurora that lies close to the edge of the FOV, and therefore we will focus on the region equatorward of this. Similar to the dTEC perturbations seen in Figure 2, fluctuations in the brightness in both oxygen and nitrogen emissions are seen, which appear to move southwards with time. The amplitude of these fluctuations is small compared to the mean brightness. In the case of the oxygen airglow, the mean brightness on this day is 643 counts, whereas the fluctuations are ~ 10 counts, or $\sim 0.4\%$ of the mean. This small amplitude is consistent with that reported by England et al. (2020). It is worth noting that the latitudes for which dTEC observations are available lie well away from the equatorial anomaly, which avoids any impact this may have on either these or the 135.6-nm airglow.

Given the low amplitude of the features observed, it is worthwhile determining the detection limit of the GOLD instrument in this mode. Assuming that the 135.6-nm signal is dominated by Poisson noise, 643 counts would give a per pixel signal-to-noise ratio of 25.4. Thus, a signal of level 3.9% would have a

data that follow the GOLD field of view as best as possible. These are shown in panel d of Figure 2, and correspond to within $\pm 5^\circ$ of longitude of the GOLD field of view. Using these data, the rate of propagation of the features seen parallel to the GOLD field of view is determined using the same process described above. The resultant best-fit phase speed of $7.3^\circ/\text{h}$ parallel to the GOLD field of view is found, and corresponding lines shown in panel d of Figure 2. This rate (and corresponding rates for the two other days of the campaigns) will be used when examining the GOLD observations. It is worth noting that this process will tend to identify the features shown in panel d of Figure 2, and minimize any other wave features present.

3.2. GOLD Airglow Radiances

GOLD is able to see perturbations to the FUV airglow caused by atmospheric waves in the middle thermosphere. To identify these perturbations, we follow the method of England et al. (2020), but here extend it to include the N_2 LBH bands, in addition to the OI 135.6 nm doublet. This is done in three steps. Step 1: Identify these two emission features and subtract any background emission. Wavelength bins within the GOLD spectrum that correspond to the 135.6 nm doublet and the nitrogen band (including the atomic nitrogen line at 149.3 nm that responds in a similar




Figure 5. Perturbations to the airglow, composition, and temperature observed on October 18, 2019. Panel (a) shows the perturbation in the oxygen airglow. Panel (b) shows the perturbation in the nitrogen airglow. Panel (c) shows the perturbation in the oxygen minus twice the nitrogen airglow to highlight the difference. Panel (d) shows the oxygen airglow minus twice the nitrogen airglow, where the universal times at each latitude have been offset to correspond to the time at 47° latitude, using the latitudinal phase propagation rates derived from the dTEC data, as described in the text. Panels (e and f) are as (c and d), but for the column O/N₂ perturbations derived from the oxygen and nitrogen airglow brightnesses. Panels (f and h) are as (c and d), but for the perturbations in the disk temperature derived from the shape of the LBH band emission. dTEC, differential total electron content.

signal-to-noise ratio (SNR) of 1. To identify a signal as small as 0.4% with an SNR of 1, such a feature would need to be present in ~100 pixels. The data shown in Figure 5 span 350 pixels in space and the wave-like features appear to fill the entire observable latitude range, allowing for unambiguous determination of these features, despite the low-amplitude signal. This will be especially important when considering the data in Figure 6, where these data have been reduced to a single line-plot.




Figure 6. Panels (a–d) show the mean perturbation in the 135.6-nm airglow compared to the LBH airglow, column O/N_2 , disk temperatures, and dTEC perturbations, respectively. All data are averaged over 20–50° latitude, and are shifted in UT to a common value at 47° north as in Figure 5. Panels (a–d) correspond to the results for October 17, 2019. Panels (e–h) show the same as (a–d), but for October 18, 2019. Panels (i–l) show the same as a through d, but for October 19, 2019. In each panel, the red line corresponds to the oxygen airglow. The airglow values are in counts. The disk temperatures are in K and are multiplied by 0.2 to be visible on this scale. The column O/N_2 are dimensionless ratios and are multiplied by 200 to be visible on this scale. The dTEC values are in TECu and are multiplied by 20 to be visible on this scale. dTEC, differential total electron content.

Comparing the variations in the oxygen airglow with those in the nitrogen, these fluctuations appear to be out-of-phase with one another such that an increase in the brightness of the oxygen airglow is approximately correspondent with a decrease in the brightness of the nitrogen airglow. This is consistent with the effect of vertical advection, which tends to increase (decrease) oxygen relative to nitrogen when such advection is downwards (upwards). To investigate this in a more quantitative manner, Figure 5 panel c shows the difference in the oxygen airglow and the nitrogen airglow. As the amplitude of the fluctuation in the nitrogen appears lower everywhere below the aurora, we multiply it by two and then subtract it from the oxygen perturbation. This signature of the oxygen-nitrogen difference appears almost exactly in phase with the oxygen airglow, confirming the out-of-phase relationship of the emissions from the two species. This general out-of-phase relationship is seen on all 3 days of the campaign (see Figure 6 and Section 3.5), and while not perfect, suggests that the perturbation to the airglow layer has a vertical extent that is large compared to the width of the layer itself.

Visually comparing to Figure 2 it is not immediately clear if these are the very same fluctuations as seen in dTEC or not. To investigate this further, we can establish if the perturbations are moving at the same rate as those shown in Figure 2. This is done by shifting the GOLD data at each latitude back to a reference at 47°, using the phase propagation rate found from the dTEC data. The choice of 47° latitude is arbitrary, and simply falls within the range of good overlap between GOLD and the dTEC values—the use of any other latitude would simply shift the epoch for the horizontal axis. If the perturbations seen by GOLD are moving at the same rate as those seen in the dTEC data, the result should show perturbations that appear vertical in this latitude-modified universal time coordinate system. This is shown for the oxygen-nitrogen differences in Figure 5d. While the perturbations are not perfectly vertical in this figure, it is clear that the airglow perturbations are moving in latitude at the same rate as those seen in dTEC, supporting the hypothesis that the two data sets are seeing related features in the atmosphere and ionosphere, respectively. The correspondence between each data set in this shifted coordinate system is explored in more detail in Section 3.5.

3.3. GOLD Disk Temperatures

Having established that GOLD observes fluctuations in the brightness of the airglow that appear to move with those seen in the TEC data, we now consider the GOLD observations of the temperature of the middle thermosphere. As described in Section 2.3, these are available over a slightly reduced region relative to the airglow brightness data, but still cover all of the latitudes seen in the TEC data. As with the airglow brightness the temperature exhibits a strong local time dependence, and so the same detrending procedure as described in steps 2 and 3 in Section 3.2 is applied to the temperature data. The resulting detrended temperatures are shown in Figure 5 panel g. It is clear that similar perturbations are seen in the temperatures as the airglow brightness, but with some shift in the phase of the signature (approximately one-quarter wavelength). For the disk temperature data included in Figure 5, the average random uncertainty per spatial-temporal bin is 9.4 K. The temperature perturbation seen appears to be around twice this value, and also is present across many bins at once, yielding a relatively unambiguous determination of this moving feature. Following the analysis described above, Figure 5h shows the temperature data that have been shifted using the rate from the dTEC data. This again supports the hypothesis that the perturbation seen by GOLD is related to that seen in the dTEC data.

3.4. GOLD Column O/N₂

Given the variations seen in the oxygen and nitrogen airglow features described in Section 3.2, it is expected that measurable changes in the column O/N₂ will be present during this campaign. Using the Level 2 GOLD data provides an opportunity to determine the amplitude of this variation and compare this to the signatures identified in the temperature and ionospheric observations. As with the airglow brightness the O/N₂ exhibits a strong local time dependence, and so the same detrending procedure as described in steps 2 and 3 in Section 3.2 is applied to the O/N₂ data. The resulting detrended column O/N₂ are shown in Figure 5 panel e. These follow a similar pattern to those in panels c and d that show the differences in the brightnesses of the oxygen and nitrogen airglow, as may be expected.

When interpreting such small changes in column O/N₂ derived from far-UV observations, it is important to acknowledge that the 135.6 nm observations used to determine O/N₂ contain a small contribution from O⁺ radiative recombination, in addition to the photoelectron impact on O. Using the dTEC observations, we can estimate the impact this may have, relative to the O/N₂ perturbations reported here. First, we select a location corresponding to approximately the middle of where the perturbations are seen (40° north, 77° east, and 15 h UT). At this location, we use an ionospheric O⁺ density and electron temperature profile from IRI-2016 (Bilitza, 2018) and the radiative recombination airglow coefficients from Meléndez-Alvira et al. (1999) to estimate the total emission from O⁺ radiative recombination in this region to be ~19 R, compared to the mean GOLD observed value of 1,000 R. From the magnitude of the perturbation seen in TEC, we can estimate that this O⁺ radiation recombination may vary by 1.7 R, or 0.17% of the mean brightness. Comparing this to the observed variation in 135.6 (Section 3.2), which is around 0.4% of the mean, we see that this radiative recombination may account for a little less than half of the total signal. This is consistent with a comparison of the signatures seen in the dTEC and 135.6 nm observations, which appear similar but not identical as the 135.6 nm signature is a mixture of the O and O⁺ perturbations.

3.5. Correspondence Between Observed Parameters

For a more quantitative comparison of the three data sets, we make use of the UT shift shown in Figure 5 to align all of the major perturbations such that they occur at all latitudes at approximately the same shifted UT. This then allows us to average the signal over a range of latitudes, which compensates for the relatively low magnitude of the perturbation in the airglow brightness data. In effect, this emphasizes only the features moving at the rate of those shown in Figure 2, but as described in Section 3.2, it increases the signal-to-noise ratio for those signatures. A latitude range of 20–50° is chosen as the perturbations are clear throughout this region, and all data sets have good coverage throughout this latitude range. Figure 6 shows the results of this for all 3 days of the campaign.

Examining Figure 6 panels a, e, and i, a clear antiphase relationship is seen between the oxygen and nitrogen airglow (both of which come from layers that peak in the 140–180-km altitude region). This is further

verified by examining Figure 6 panels b, f, and j, which show that the perturbations in the retrieved column O/N_2 look extremely similar to the variations in the oxygen airglow brightness. This signature is a clear sign of vertical advection, which in combination with the different altitude distributions of oxygen and nitrogen in the middle thermosphere, acts to increase the column abundance of one of these relative to the other. The oscillatory nature of the signatures would also be consistent with this advection being caused by an atmospheric wave moving through this region.

Examining Figure 6 panels c, g, and k, we see the relationship between the oxygen airglow and temperature perturbations. While not perfect, a clear relationship between the two is apparent. Rather than a simple in phase or out-of-phase relationship, the positive and negative perturbations in the temperature appear to lag those seen in the oxygen airglow brightness by approximately one-quarter of the wave period. This phase relationship is consistent with an atmospheric gravity wave propagating southwards with time, where we may expect to see the increase in atomic oxygen that is associated with downward advection one-quarter wave period before the increased temperatures (e.g., Andrews et al., 1987).

Examining Figure 6 panels d, h, and l, the relationship between the perturbations in the oxygen airglow brightness and dTEC is less clear (it is worth noting that the TEC layer height is more variable than the airglow, but typically corresponds to the 200–350-km altitude region at these latitudes). Nonetheless, some features appear in common, especially on the first 2 days of the campaign. On these 2 days, the increases in dTEC and oxygen airglow brightness appear to be roughly in phase, although the amplitudes of the perturbations do not appear to have a simple correspondence such that sometimes a large increase in airglow is seen at the same time as a small increase in dTEC, and vice versa. It is also worth noting that the rate of motion of the signatures seen in dTEC, airglow brightness, column O/N_2 , and disk temperatures are all approximately the same (the rate of motion in dTEC is used to detrend all data sets into the shifted UT frame used here).

The following section will explore the consistency between the magnitude of the neutral atmosphere feature seen with our understanding of the oxygen airglow. For this, we note from Figure 6 the typical magnitudes seen in these parameters. From Figure 6 panels a, e, and i, we see a typical amplitude of the oxygen airglow brightness of 2.5 counts, compared to a mean value of 640 counts, which corresponds to $\sim 0.4\%$. From panels c, g, and k, we see a typical disk temperature amplitude of 13 K (compared to a mean temperature of 710 K). The correspondence of these will be examined in the next section.

For a more quantitative assessment of the correspondence between the various parameters shown in Figure 6, we examine the relationship between sets of these parameters. This is done in the following steps: (1) interpolate all quantities onto a common 10-min temporal cadence (to match the lowest cadence) using a cubic-spline interpolation; (2) shift the temperature data by 1 h, corresponding to the apparent one-quarter wave period shift seen on October 18 in Figure 6; (3) plot corresponding pairs of data in the scatterplots shown in Figure 7; (4) as all perturbations are extremely small we can expect an approximately linear relationship between different quantities, which is found with a least-squares fit to the data in Figure 7 (shown by the pink solid lines); (5) compute the coefficient of determination (R^2) value for each set of parameters shown.

Examining panel a of Figure 7, we see the near antiphase relationship between the oxygen and nitrogen airglow brightness perturbations that is expected from the earlier results. The best-fit linear trend shows a reasonable, but certainly not perfect agreement with the data, with a coefficient of determination of 0.51. Given this, it is worth considering how closely we may expect the fit to be to the data. From Section 3.2, we can tell that a signal with the apparent amplitude of that observed would have a SNR per 10-min bin of ~ 3 for the oxygen airglow and ~ 2 for the nitrogen airglow. Also, it is worth noting that these estimates assume the signal is present across the entire latitude range selected (which appears to be the case based on Figure 5), and that the features moving at the rate determined from the dTEC perturbations characterize the only airglow perturbations present. This latter condition is likely not the case, based on Figure 2. With these two factors taken into consideration an R^2 value of no more than about 0.51 may be expected.

Examining panel b of Figure 7, we see the expected good anticorrelation between the column O/N_2 ratio and the nitrogen airglow brightness. The higher R^2 value of 0.67 suggests that any impact of O^+ radiative




Figure 7. Correlations between the perturbations in the oxygen and nitrogen airglow brightness, disk temperatures, column O/N_2 , and dTEC for October 18, 2019 data shown in Figure 6. Panel (a) shows the nitrogen airglow brightness versus the oxygen airglow brightness. Panel (b) shows the O/N_2 ratio versus the nitrogen airglow brightness. Panel (c) shows the nitrogen airglow brightness versus the disk temperature. Panel (d) shows the dTEC versus the oxygen airglow brightness. All data sets have been interpolated onto a 10-min cadence, and the temperature data were shifted by 1 h, corresponding to approximately one-quarter wave period. The pink lines show the least-squares linear fits to each pair of data. Coefficients of determination for each of these linear fits are shown. dTEC, differential total electron content.

recombination in the determination of column O/N_2 perturbations does not have a significant impact on the pattern of O/N_2 perturbations that are found.

Examining panel c of Figure 7, we see some anticorrelation between the nitrogen airglow and the time-shifted disk temperature perturbations. While the R^2 value of 0.44 reflects a somewhat poor fit, no correspondence between these parameters is seen with the 1-h time shift in the temperature data (R^2 value of 4×10^{-5} in this case). This confirms that the apparent one-quarter period shift expected by theory and seen in Figure 6 is a reasonable fit to the data.

Finally, examining panel d of Figure 7, we see some positive correlation between the oxygen airglow and the dTEC perturbations. The low R^2 value of 0.27 reflects what was noted in regard to Figure 6, that the perturbations in the ionosphere do not simply look like those in the neutral atmosphere, although they are apparently moving southwards at the same rate. In principle, it is possible to use the relative phase of the signals seen at the airglow layer (near 150-km altitude), the TEC layer (near 300-km altitude), and the estimate of the horizontal wavelength and period from Section 3.1 to estimate the vertical wavelength of the feature seen. Around in panel d of Figure 7, 17 h universal time the phase shift appears around 20 min. Using this the vertical wavelength is estimated at around 2,000 km, or in essence there is less than one vertical wavelength in the column of atmosphere that exists above the UV airglow layer.

3.6. GLOW Model Results

Following England et al. (2020), it is possible to use the magnitude of the oxygen airglow brightness perturbation to estimate the amplitude of the underlying atmospheric perturbation. With the addition of the disk temperature data included in this study, we can also establish how consistent the estimated atmospheric perturbation is with the variation in the brightness of the airglow seen. Following Greer et al. (2018), we use the thermosphere-ionosphere-electrodynamics GCM (TIEGCM v2.0; Maute, 2017) model to provide




Figure 8. Results of the GLOW airglow simulations of the O 135.6 nm doublet at 45° north and 80° west for the geophysical conditions on October 18, 2019. The amplitude of the simulated airglow perturbation is shown as a function of the change in the neutral atmosphere, expressed in terms of the temperature. Simulations correspond to an average over 12–20 UT. The solid lines show the linear least-squares fit to these data.

information on the background atmosphere, onto which sinusoidal perturbations are applied that modify the atmospheric density and temperature in the middle thermosphere, as described in Section 2 of Greer et al. (2018). We then use the Global Airglow model (GLOW; Solomon, 2017) to simulate the 135.6 nm O airglow GOLD would observe at 45° north and 80° west, for the given geophysical and solar conditions corresponding to October 18, 2019. Comparisons between GOLD and TIEGCM-GLOW simulations have been presented in detail by Greer et al. (2020), and while the absolute brightness seen by GOLD is different to that simulated, the relative changes spatially and temporally match well. By varying the amplitude of the induced perturbation, we can find the amplitude of a wave in the middle thermosphere that would produce a perturbation in the airglow of the same amplitude as observed by GOLD. Figure 8 shows the results of these simulations. At these small atmospheric perturbation levels, the response of the airglow is linear, as shown in Figure 8. The ~0.4% amplitude of the oxygen airglow perturbation noted in the previous section is consistent with an atmospheric temperature perturbation of around 22.4 K. To compare this with the observed disk temperatures, it is necessary to acknowledge that the amplitude of any realistic gravity wave perturbation will vary with altitude, and the 22.4 K value reported here refers to the perturbation at 150-km altitude, close to the peak of where the disk far-UV airglow originates. This is illustrated in more detail in Figure 9, which reproduces the atmospheric perturbations used and the airglow perturbation from GLOW that results. The variation in the wave amplitude and phase with altitude matches that used in Greer et al. (2018). The amplitude of the temperature perturbation varies from around 12 K near 130 km to almost 36 K near 175 km. Similarly, the corresponding neutral density perturbation varies across this region, and has a value of 2.15% at 150-km altitude.

Examining Figure 9 in more detail, we see the phase offset between the oxygen airglow brightness and temperature perturbations, that match what is seen in the GOLD observations. The resultant oxygen and nitrogen airglow brightness perturbations reflect the overall antiphase relationship seen by GOLD. The disk temperature perturbation observed by GOLD was 13 K, which is similar to the simulated variation near 130 km, and is generally less than that in the best-fit simulation for October 18, shown in Figure 9. However, it is worth noting that the GOLD observation represents an altitude-integral of the temperature perturbation, weighted by the LBH airglow emission profile that peaks near 150 km, but extends over several tens of km above and below this altitude. Given this, and the tilted phase-fronts of any realistic gravity wave, the temperature perturbation seen by GOLD is expected to be less than the simulated value at the peak of the layer near 150 km. In principle, it is possible to use knowledge of an estimate of the vertical wavelength (such as that estimated above) to provide a more quantitative estimate of the relationship between the temperature perturbation observed by GOLD and that simulated at a range of altitudes. However, it must be recognized that the vertical wavelength will vary with background temperature (visible in the simulation in Figure 9), which is known to increase substantially between the lower region of the airglow profile (130–150 km) and the F-region peak (300–350 km). Further, at altitudes where molecular diffusion becomes significant, perturbations become near constant with altitude. Without knowledge of the altitude profile of temperature, and thus the change in the vertical wavelength of an atmospheric gravity wave throughout this region it is not possible to provide a more quantitative relationship between the observed temperature perturbation and that simulated at any one altitude, other than to note they are generally consistent in this case. Taking all of these factors into consideration, we see that a gravity wave similar to that shown in Figure 9 is consistent with all of the GOLD observed features.




Figure 9. Impacts of the best-match gravity wave-like perturbation on the oxygen and nitrogen airglow brightness. Panel (a) shows the perturbation in temperature and panel (b) shows the perturbation in neutral density. Panel (c) shows the resultant impact of the wave-like perturbation from panels (a) and (b) on the oxygen and nitrogen airglow simulated using the GLOW model at 45° latitude and -80° longitude for the geophysical conditions on October 18. Note the variation in the amplitude of the atmospheric perturbation with altitude.

4. Discussion and Conclusions

Results from a 3-day campaign with the GOLD instrument to observe atmospheric waves in the atmosphere during October 2019 are presented. In the operating mode used for this campaign, GOLD obtained data during daytime by using one channel in a fixed viewing direction, maximizing the signal and providing information along a line that is approximately aligned the meridional direction. To provide overlap with ground-based data, while being near nadir viewing geometry, GOLD Channel B was positioned over the eastern coast of the US. There, ground-based GPS TEC data are then available to help identify wave perturbations in ionospheric parameters, providing supporting information for the GOLD observations.

Perturbations are seen in the differential TEC data that are indicative of TIDs. As indicated in prior studies, wave-like TIDs are ubiquitous in GNSS dTEC data sets, and the 3 days of this campaign are no exception. Given how clearly such features are seen at midlatitudes in the dTEC we use these data to determine the phase propagation of the waves on each of the 3 days. On October 18, presented in detail, the waves appear to move $\sim 7^\circ/\text{h}$ and are headed almost due south ($7.3^\circ/\text{h}$ along the GOLD field-of-view), with a horizontal wavelength of $\sim 1,500$ km and a phase speed of 210 m s^{-1} . These properties are indicative of a slow-moving large-scale TID, consistent with the relatively low levels of geomagnetic activity present on this day. While determining the wave source is beyond the data sets at hand, the phase propagation direction is consistent with a source near the auroral region, but a lower-atmospheric source or secondary gravity wave source at intermediate altitude remain other possibilities.

GOLD sees similar wave-like perturbations in the airglow brightness associated with oxygen and nitrogen species, column O/N_2 ratio derived from those brightnesses, and disk temperatures derived from the shape of the nitrogen LBH band. Using the phase propagation rates determined for the dTEC signature, we

demonstrate that these are moving at approximately the same rate, but the dTEC and GOLD oxygen signatures are not quite in phase with one another.

The phase and amplitude relationships between each quantity are examined by taking an average of the values over 20–50° latitude, after all have been adjusted using the phase propagation rates determined from dTEC. The oxygen and nitrogen airglow brightness perturbations are in antiphase with one another, producing a clear variation in the column O/N₂ ratio that is consistent with a response to vertical advection. The temperature perturbations appear shifted relative to the oxygen airglow by around one-quarter wave period, with the temperature perturbations lagging or appearing slightly northwards of those in the LBH, consistent with an atmospheric gravity wave propagating southwards (and thus consistent with the original picture from the dTEC). The correspondence in the phase between the airglow and dTEC is only partial, but on two out of the 3 days the perturbations in dTEC appear to be mostly in phase with those in the oxygen airglow. Less correspondence in the amplitudes of the signals seen in dTEC and the airglow are seen (as measured by the changes in the R^2 values reported in Section 3.5), suggesting a complex relationship between the amplitude of these signals, even though both appear to be related in the measures described above. While the source of this complex relationship is unclear from these observations alone, one possibility is that the airglow signatures observed by GOLD and the TEC observations from the ground both represent altitude-averaged quantities with only limited overlap in altitude. The far-UV airglow features GOLD observes are emitted over a range of altitudes and are strongest in the 130–180-km region, representative of the middle thermosphere. The TEC values are also an altitude integrated quantity and are most highly weighted to F-region altitudes, typically in the 250–350-km region, which overlaps with the upper thermosphere. Neither GOLD nor the ground-based TEC provide precise constraints on these altitude ranges, but it is reasonable to expect that any changes in the wave properties (such as change in amplitude) with altitude could account for a difference in the phase or amplitude seen in these two data sets. Additionally, the analysis presented here focuses on only the strongest wave perturbations seen in the dTEC data and identifies features moving at the same latitudinal rate as these. Any other waves or propagating features present could be categorized as noise in this analysis and contribute to an apparent lack of agreement. A final consideration is that perturbations in the F-region electron density may tend to be aligned with the geomagnetic field, whereas the neutral perturbations are not. The interplay of this, and the measure of TEC, which is a vertical column will likely vary between the higher latitudes where the field is aligned nearly vertically and lower latitudes where it is not. This may provide another source of discrepancy between the dTEC perturbations and those observed in the neutral atmosphere with GOLD.

Using the GLOW airglow model, we demonstrated the consistency of the signatures observed by GOLD with a gravity wave perturbation to the atmosphere. The model, with a background atmosphere that is perturbed by a gravity wave-like structure is able to reproduce fluctuations in the airglow brightness that have the same kind of phase relationship as that observed. The best-fit amplitude of the atmospheric perturbation is 22.4 K temperature and 2.15% density variation at 150 km. Given the altitude range of the airglow emission, the tilt in any phase of the perturbation (corresponding to an unknown vertical wavelength), and the growth in the amplitude of a perturbation with height, the observed disk temperature perturbation is expected to be smaller than that simulated at the 150-km altitude, which is the case. The phase between the density, temperature, and airglow perturbations are all consistent with the simulated gravity wave-like feature.

This campaign yields a number of important findings. First is that the GOLD disk temperature data show the signatures of the TADs very clearly—arguably better than the airglow brightness data used in the prior study by England et al. (2020). This is of particular interest as observations of the temperature in the middle thermosphere are extremely sparse, and thus estimates of the amplitude of wave perturbations of these temperatures are similarly rare. Second is that the phase relationships between the observed perturbations in the oxygen and nitrogen airglow, the column O/N₂ and the disk temperatures are all consistent with the perturbations coming from an atmospheric gravity wave or TAD. Third is that the TAD and TID seen in the dTEC data propagate at the same rate but have a complex relationship between their amplitudes that is not easy to determine from just a sample of 3 days. With these findings in mind, as GOLD moves into its extended mission (beginning November 2020), campaigns such as that described here will be performed more frequently. Lessons learned from this campaign and the methods described here will allow a much larger

data set of TAD and TID pairs to be identified, under different conditions and for different seasons, with the hope that the relationship between these two can be characterized.

Data Availability Statement

The GOLD data are available from the GOLD Science Data Center (<http://gold.cs.ucf.edu/>) and NASA's Space Physics Data Facility (<https://spdf.gsfc.nasa.gov>). The F10.7 data are available from NASA's OMNI-Web (<https://omniweb.gsfc.nasa.gov>). The SME Auroral Electrojet Index data are available from the SuperMAG at <http://supermag.jhuapl.edu> and we gratefully acknowledge the SuperMAG collaborators.

Acknowledgments

This research was supported by NASA contract 80GSFC18C0061. GPS TEC data products and access through the Madrigal distributed data system are provided to the community (<http://www.openmadrigal.org>) by the Massachusetts Institute of Technology (MIT) under support from the U.S. National Science Foundation (NSF) Grant AGS-AGS-195273. S. R. Zhang also acknowledges AFOSR MURI Grant FA9559-16-1-0364.

References

Aksnes, A., Eastes, R., Budzien, S., & Dymond, K. (2006). Neutral temperatures in the lower thermosphere from N₂ Lyman-Birge-Hopfield (LBH) band profiles. *Geophysical Research Letters*, *33*, L15103. <https://doi.org/10.1029/2006GL026255>

Andrews, D. G., Holton, J. R., & Leovy, C. B. (1987). *Middle atmosphere dynamics, international geophysics series*. Academic Press.

Azeem, I., Vadas, S. L., Crowley, G., & Makela, J. J. (2017). Traveling ionospheric disturbances over the United States induced by gravity waves from the 2011 Tohoku tsunami and comparison with gravity wave dissipative theory. *Journal of Geophysical Research: Space Physics*, *122*, 3430–3447. <https://doi.org/10.1002/2016JA023659>

Bilitza, D. (2018). IRI the international standard for the ionosphere. *Advances in Radio Science*, *16*, 1–11. <https://doi.org/10.5194/ars1612018>

Bruinsma, S., Forbes, J. M., Nerem, R. S., & Zhang, X. (2006). Thermosphere density response to the 20–21 November 2003 solar and geomagnetic storm from CHAMP and GRACE accelerometer data. *Journal of Geophysical Research*, *111*, A06303. <https://doi.org/10.1029/2005JA011284>

Bruinsma, S. L., & Forbes, J. M. (2008). Medium- to large-scale density variability as observed by CHAMP. *Space Weather*, *6*, S08002. <https://doi.org/10.1029/2008SW000411>

Cai, X., Burns, A. G., Wang, W., Coster, A., Qian, L., Liu, J., et al. (2020). Comparison of GOLD nighttime measurements with total electron content: Preliminary results. *Journal of Geophysical Research: Space Physics*, *125*, e2019JA027767. <https://doi.org/10.1029/2019JA027767>

Djuth, F. T., Zhang, L. D., Livneh, D. J., Seker, I., Smith, S. M., Sulzer, M. P., et al. (2010). Arecibo's thermospheric gravity waves and the case for an ocean source. *Journal of Geophysical Research*, *115*, A08305. <https://doi.org/10.1029/2009JA014799>

Earle, G. D., Musumba, A. M., & Vadas, S. L. (2008). Satellite-based measurements of gravity wave-induced midlatitude plasma density perturbations. *Journal of Geophysical Research*, *113*, A03303. <https://doi.org/10.1029/2007JA012766>

Eastes, R. W., McClintock, W. E., Burns, A. G., Anderson, D. N., Andersson, L., Codrescu, M., et al. (2017). The global-scale observations of the limb and disk (GOLD) mission. *Space Science Reviews*, *212*(1–2), 383–408. <https://doi.org/10.1007/s11214-017-0392-2>

Eastes, R. W., Solomon, S. C., Daniell, R. E., Anderson, D. N., Burns, A. G., England, S. L., et al. (2019). Global-scale observations of the equatorial ionization anomaly. *Geophysical Research Letters*, *46*, 9318–9326. <https://doi.org/10.1029/2019GL084199>

England, S. L., Greer, K. R., Solomon, S. C., Eastes, R. W., McClintock, W. E., & Burns, A. G. (2020). Observation of thermospheric gravity waves in the Southern Hemisphere with GOLD. *Journal of Geophysical Research: Space Physics*, *125*, e2019JA027405. <https://doi.org/10.1029/2019JA027405>

Galushko, V. G., Paznukhov, V. V., Yampolski, Y. M., & Foster, J. C. (1998). Incoherent scatter radar observations of AGW/TID events generated by the moving solar terminator. *Annales Geophysicae*, *16*, 821–827. <https://doi.org/10.1007/s00585-998-0821-3>

Garcia, R. F., Bruinsma, S., Massarweh, L., & Doornbos, E. (2016). Medium-scale gravity wave activity in the thermosphere inferred from GOCE data. *Journal of Geophysical Research: Space Physics*, *121*, 8089–8102. <https://doi.org/10.1002/2016JA022797>

Gjerloev, J. W. (2012). The SuperMAG data processing technique. *Journal of Geophysical Research*, *117*, A09213. <https://doi.org/10.1029/2012JA017683>

Greer, K. R., Eastes, R., Solomon, S., McClintock, W., Rusch, D., & Burns, A. (2020). Variations of lower thermospheric FUV emissions based on GOLD observations and GLOW modeling. *Journal of Geophysical Research: Space Sciences*, *125*, e2020JA027810. <https://doi.org/10.1029/2020JA027810>

Greer, K. R., England, S. L., Becker, E., Rusch, D., & Eastes, R. (2018). Modeled gravity wave-like perturbations in the brightness of far ultraviolet emissions for the GOLD mission. *Journal of Geophysical Research: Space Physics*, *123*, 5821–5830. <https://doi.org/10.1029/2018JA025501>

Livneh, D. J., Seker, I., Djuth, F. T., & Mathews, J. D. (2009). Omnipresent vertically coherent fluctuations in the ionosphere with a possible worldwide-midlatitude extent. *Journal of Geophysical Research*, *114*, A06303. <https://doi.org/10.1029/2008JA013999>

Mandal, S., & Pallamraju, D. (2020). Thermospheric gravity wave characteristics in the daytime over low-latitude during geomagnetically quiet and disturbed conditions. *Journal of Atmospheric and Solar-Terrestrial Physics*, *211*, 105470. <https://doi.org/10.1016/j.jastp.2020.105470>

McClintock, W. E., Solomon, S. C., Eastes, R., Anderson, D. N., Andersson, L., Burns, A. G., et al. (2017). *Global-scale Observations of the Limb and Disk (GOLD): Science implementation*. American Geophysical Union Fall Meeting, SA31A-2561.

Meléndez-Alvira, D. J., Meier, R. R., Picone, J. M., Feldman, P. D., & McLaughlin, B. M. (1999). Analysis of the oxygen nightglow measured by the Hopkins Ultraviolet Telescope: Implications for ionospheric partial radiative recombination rate coefficients. *Journal of Geophysical Research*, *104*(A7), 14901–14913. <https://doi.org/10.1029/1999JA900136>

Negrea, C., Zabolotn, N., Bullett, T., Fuller-Rowell, T., Fang, T.-W., & Codrescu, M. (2016). Characteristics of acoustic gravity waves obtained from Dynasonde data. *Journal of Geophysical Research: Space Physics*, *121*, 3665–3680. <https://doi.org/10.1002/2016JA022495>

Newell, P. T., & Gjerloev, J. W. (2011). Evaluation of SuperMAG auroral electrojet indices as indicators of substorms and auroral power. *Journal of Geophysical Research*, *116*, A12211. <https://doi.org/10.1029/2011JA016779>

Park, J., Lühr, H., Lee, C., Kim, Y. H., Jee, G., & Kim, J.-H. (2014). A climatology of medium-scale gravity wave activity in the midlatitude/low-latitude daytime upper thermosphere as observed by CHAMP. *Journal of Geophysical Research: Space Physics*, *119*, 2187–2196. <https://doi.org/10.1002/2013JA019705>

- Paulino, I., Moraes, J. F., Maranhão, G. L., Wrasse, C. M., Buriti, R. A., Medeiros, A. F., et al. (2018). Intrinsic parameters of periodic waves observed in the OI6300 airglow layer over the Brazilian equatorial region. *Annales de Geophysique*, *36*, 265–273. <https://doi.org/10.5194/angeo362652018>
- Richmond, A. D. (1978). Gravity wave generation, propagation, and dissipation in the thermosphere. *Journal of Geophysical Research*, *83*(A9), 4131–4145. <https://doi.org/10.1029/JA083iA09p04131>
- Rideout, W., & Coster, A. (2006). Automated GPS processing for global total electron content data. *GPS Solutions*, *10*, 219–228. <https://doi.org/10.1007/s1029100600295>
- Savitzky, A., & Golay, M. J. E. (1964). Smoothing and differentiation of data by simplified least squares procedures. *Analytical Chemistry*, *36*, 1627–1639. <https://doi.org/10.1021/ac60214a047>
- Shiokawa, K., Lu, G., Otsuka, Y., Ogawa, T., Yamamoto, M., Nishitani, N., & Sato, N. (2007). Ground observation and AMIE-TIEG-CM modeling of a storm-time traveling ionospheric disturbance. *Journal of Geophysical Research*, *112*, A05308. <https://doi.org/10.1029/2006JA011772>
- Solomon, S. C. (2017). Global modeling of thermospheric airglow in the far ultraviolet. *Journal of Geophysical Research: Space Physics*, *122*, 7834–7848. <https://doi.org/10.1002/2017JA024314>
- Strickland, D. J., Evans, J. S., & Paxton, L. J. (1995). Satellite remote sensing of thermospheric O/N₂ and solar EUV, 1, Theory. *Journal of Geophysical Research*, *100*, 12217. <https://doi.org/10.1029/95JA00574>
- Tsuchiya, S., Shiokawa, K., Otsuka, Y., Nakamura, T., Yamamoto, M., Connors, M., et al. (2020). Wavenumber spectra of atmospheric gravity waves and medium-scale traveling ionospheric disturbances based on more than 10-year airglow images in Japan, Russia, and Canada. *Journal of Geophysical Research: Space Physics*, *125*, e2019JA026807. <https://doi.org/10.1029/2019JA026807>
- Tsugawa, T., Otsuka, Y., Coster, A. J., & Saito, A. (2007). Medium-scale traveling ionospheric disturbances detected with dense and wide TEC maps over North America. *Geophysical Research Letters*, *34*, L22101. <https://doi.org/10.1029/2007GL031663>
- Vadas, S. L., & Fritts, D. C. (2006). Influence of solar variability on gravity wave structure and dissipation in the thermosphere from tropospheric convection. *Journal of Geophysical Research*, *111*, A10S12. <https://doi.org/10.1029/2005JA011510>
- Vadas, S. L., & Liu, H.-L. (2009). The generation of large-scale gravity waves and neutral winds in the thermosphere from the dissipation of convectively-generated gravity waves. *Journal of Geophysical Research*, *114*, A10310. <https://doi.org/10.1029/2009JA014108>
- Vierinen, J., Coster, A. J., Rideout, W. C., Erickson, P. J., & Norberg, J. (2016). Statistical framework for estimating GNSS bias. *Atmospheric Measurement Techniques*, *9*, 1303–1312. <https://doi.org/10.5194/amt-9-1303-2016>
- Yiğit, E., & Medvedev, A. S. (2015). Internal wave coupling processes in Earth's atmosphere. *Advances in Space Research*, *55*(4), 983–1003. <https://doi.org/10.1016/j.asr.2014.11.020>
- Yiğit, E., Medvedev, A. S., England, S. L., & Immel, T. J. (2014). Simulated variability of the high-latitude thermosphere induced by small-scale gravity waves during a sudden stratospheric warming. *Journal of Geophysical Research: Space Physics*, *119*, 357–365. <https://doi.org/10.1002/2013JA019283>
- Zhang, S.-R., Coster, A. J., Erickson, P. J., Goncharenko, L. P., Rideout, W., & Vierinen, J. (2019). Traveling ionospheric disturbances and ionospheric perturbations associated with solar flares in September 2017. *Journal of Geophysical Research: Space Physics*, *124*, 5894–5917. <https://doi.org/10.1029/2019JA026585>
- Zhang, Y., Paxton, L. J., Morrison, D., & Schaefer, B. (2018). Storm-time variations of atomic nitrogen 149.3 nm emission. *Journal of Atmospheric and Solar-Terrestrial Physics*, *169*, 78–82. <https://doi.org/10.1016/j.jastp.2018.01.023>
- Zhang, Y., Paxton, L. J., Morrison, D., Wolven, B., Kil, H., Meng, C.-I., et al. (2004). O/N₂ changes during 1–4 October 2002 storms: IMAGE SI-13 and TIMED/GUVI observations. *Journal of Geophysical Research*, *109*, A10308. <https://doi.org/10.1029/2004JA010441>