

Cite this: DOI: 10.1039/d2ob00162d

Received 26th January 2022,  
Accepted 4th April 2022

DOI: 10.1039/d2ob00162d  
[rsc.li/obc](http://rsc.li/obc)

## Applications of catalysis in hydroboration of imines, nitriles, and carbodiimides

Adineh Rezaei Bazkiae,<sup>a</sup> Michael Findlater  <sup>\*b</sup> and Anne E. V. Gorden  <sup>\*a</sup>

The catalytic hydroboration of imines, nitriles, and carbodiimides is a powerful method of preparing amines which are key synthetic intermediates in the synthesis of many value-added products. Imine hydroboration has perennially featured in notable reports while nitrile and carbodiimide hydroboration have gained attention recently. Initial developments in catalytic hydroboration of imines and nitriles employed precious metals and typically required harsh reaction conditions. More recent advances have shifted toward the use of base metal and main group element catalysis and milder reaction conditions. In this survey, we review metal and nonmetal catalyzed hydroboration of these unsaturated organic molecules and group them into three distinct categories: precious metals, base metals, and main group catalysts. The TON and TOF of imine hydroboration catalysts are reported and summarized with a brief overview of recent advances in the field. Mechanistic and kinetic studies of some of these protocols are also presented.

## 1. Introduction

First reported by H. C. Brown in 1956, hydroboration is the addition of a boron–hydrogen bond across an unsaturated moiety.<sup>1</sup> In these reactions, the boron containing group acts as

a functional handle to form a new carbon–carbon bond or to introduce functional groups such as alcohols, amides, amines, or halogens.<sup>2</sup> Usually, the reaction proceeds without the addition of a catalyst and the borane reagent, such as diborane ( $B_2H_6$ ) or a borane adduct ( $BH_3\cdot THF$ ) and, reacts rapidly at room temperature to afford an anti-Markovnikov organoborane. Nonetheless, when using pinacolborane ( $HBpin$ ) and catecholborane ( $HBcat$ ), the boron is bonded to a heteroatom which results in increased electron density at the boron center.<sup>3</sup> As a consequence, the Lewis acidity of the boron

<sup>a</sup>Department of Chemistry and Biochemistry, Texas Tech University, Lubbock, Texas 79409, USA. E-mail: anne.gorden@ttu.edu

<sup>b</sup>Department of Chemistry and Biochemistry, University of California Merced, Merced, California 95343, USA. E-mail: michael.findlater@ucmerced.edu



Adineh Rezaei Bazkiae

she was with Bayer as industrial chemist intern. In 2021, she joined the group of Prof. Anne Gorden to continue her Ph.D. program. Her research interests mainly include organometallic chemistry and catalysis.



Michael Findlater

2021, he moved to the University of California, Merced, and was promoted to full professor. His research interests include organometallic chemistry, catalysis using earth-abundant elements, electrosynthesis, and the functionalization of  $CO_2$ .

center decreases, thus requiring increased temperatures or the addition of some catalyst or co-reagent for hydroboration to proceed efficiently.<sup>4,5</sup>

The first catalytic hydroboration process was disclosed in a report by Kono and Ito in 1975. They reported that Wilkinson's catalyst [ $\text{Rh}(\text{PPh}_3)_3\text{Cl}$ ] can undergo oxidative addition with catecholborane (HBcat).<sup>6</sup> A decade later, Manning and Nöth demonstrated that Wilkinson's catalyst along with HBcat catalyzed hydroboration of a C=C bond selectively in the presence of a keto group.<sup>7</sup>

To date, catalysts ranging from main group elements<sup>8–13</sup> and transition metals<sup>14–20</sup> have been extensively developed for the hydroboration of unsaturated organic molecules.<sup>21–26</sup> These continue to be of interest in chemical syntheses of complex molecules as well as the production of commodity and agrochemicals. Amongst unsaturated species, nitrogen containing compounds such as imines, nitriles, carbodiimides *etc.* have garnered considerable attention in the application of hydroboration processes. This is due to primary and secondary amines being extensively present in nature. They are important compounds in chemistry and biology with versatile application in the synthesis of agrochemicals,<sup>27</sup> drug molecules<sup>28,29</sup> and polymer materials.<sup>30</sup> Furthermore, borylated amine products which form in the hydroboration of nitrogenous compounds can act as an excellent synthetic surrogates. Traditional protocols for such transformations, for example, in the production of amine products employ metal hydrides<sup>31–33</sup> and pressurized hydrogen gas as hydrogen source<sup>34–42</sup> are well-exploited. Nevertheless, these methods can suffer from poor selectivity and limited functional group tolerance, the deleterious formation of inorganic byproducts, and present potential safety concerns. Hydroboration offers an efficient pathway into the preparation of amines through: (1) chemoselective reduction and, (2) with modest reaction conditions and energy requirements.<sup>43</sup>



Anne E. V. Gorden

ate professor in 2011. She moved to Texas Tech University in 2020. Research in the Gorden group has combined interests in organic syntheses, supramolecular chemistry, *f*-elements, and catalysis for sustainable chemistry.

The milestone discovery of imine hydroboration, which was first introduced by the Westcott group,<sup>44</sup> led to the development of a plethora of transition metal and main group element complexes as the catalyst ranging from precious,<sup>45–48</sup> to rare earth<sup>49,50</sup> and even base metal complexes.<sup>51,52</sup> A  $\beta$ -diketiminato magnesium complex was reported in 2013 as a main group element pre-catalyst in the hydroboration of imines.<sup>53</sup> Following such pioneering works, other metal and non-metal main group complexes were developed in related hydroboration transformations.<sup>53–65</sup> Catalyst free imine hydroboration was also recently introduced by the Rit group.<sup>66</sup>

Nitrile hydroboration, operating through the formation of an imine intermediate,<sup>49</sup> was only introduced less than a decade ago by Nikonov and co-workers who employed an imido-hydrido complex of Mo(IV).<sup>67</sup> Significant progress has been made in the development of nitrile hydroboration using main group elements<sup>31,63,68–72</sup> and transition metals (d- and f-blocks) in recent years.<sup>47,73–76</sup> One report in 2019 exploited a thorium amide complex in nitrile hydroboration by Eisen and co-workers featuring a broad substrate scope and high TOF (500  $\text{h}^{-1}$ ). This was an unusual example of actinide catalysis and takes advantage of the low cost of thorium and the stability of the Th(IV) oxidation state.<sup>49</sup> There are also reports involving the further transformation of *N*-boryl amines in the coupling reactions to form *N*-arylation or *N*-acylation products.<sup>74,76,77</sup>

In contrast to the advances in both nitrile and imine hydroboration, analogous heteroallene ( $\text{E} = \text{C} = \text{E}'$ :  $\text{E}' = \text{RN}$ ,  $\text{RN}$ ;  $\text{RN}$ ,  $\text{O}$ ) hydroboration chemistry, such as carbodiimides, are not well characterized.<sup>78,79</sup> In 1994, the single and double hydroboration of carbodiimide was demonstrated in the presence of 9-borabicyclo[3.3.1]-nonane (9-BBN) without the need for a catalyst. Amidinate and bis(boryl)aminal products were obtained, albeit under forcing thermal conditions (160 °C).<sup>80</sup> Almost three decades later, there remains only a handful of reports of carbodiimide hydroboration.<sup>62,63,81,82</sup>

Numerous catalysts have been reported to effect the hydroboration of imines, nitriles, and carbodiimides. Notably, the current state-of-the-art still relies mainly upon precious metal elements. However, the limited availability, high and volatile cost and toxicity of precious metal catalysts has shifted recent focus towards the development of inexpensive and environmentally benign first-row transition metal complexes as alternatives.<sup>83</sup> This review outlines the advances and challenges in this field as well as new avenues to efficiently catalyze hydroboration of imines, nitriles and carbodiimides.

## 2. Hydroboration of imines

### 2.1. Precious metal catalyzed hydroboration of imines

Transition metals such as Pt, Pd, Ru, Rh, and Ir (known as precious metals) have been the most active and catalytically effective metal complexes in transformations of unsaturated organic molecules for decades.<sup>84,85</sup> This is in large part owing

to their capacity to accommodate  $2e^-$  redox processes.<sup>86</sup> As such, they have found extensive application in both academia and industry in transformations such as hydroboration,<sup>87</sup> hydrosilylation,<sup>88,89</sup> amination<sup>90,91</sup> amongst others.

The first transition metal-catalyzed imine hydroboration was reported by Baker and Westcott in 1995.<sup>44</sup> A wide variety of metal complexes of Cu, Ag, and Au were studied.  $[\text{AuCl}(\text{L})]_n$  complexes (1 mol%) were found to be the most catalytically efficient and afforded *N*-borylamine products in the presence of HBcat (Fig. 1). The hydroboration reaction was monitored by  $^1\text{H}$  NMR over several half-lives and the results were normalized to give an estimate of the relative rates.  $[\text{AuCl}(\mu\text{-DPPF})]_n$  exhibited the highest relative rate in hydroboration of  $\text{PhCH}_2\text{N}=\text{CHPh}$  in  $\text{THF-d}_8$ .<sup>44</sup>

Subsequent work from the same research group in 1998 reported that 2 mol%  $\text{RhCl}(\text{PPh}_3)_3$  could be employed at room temperature (25 °C) in the hydroboration of  $\text{PhN}=\text{C}(\text{CH}_3)\text{Ph}$  in the presence of  $\text{B}_2\text{cat}'_2$  ( $\text{cat}' = 4\text{-But-1,2-O}_2\text{C}_6\text{H}_3$ ).<sup>92</sup> This reaction produced equal amounts of *N*-borylenamine (Scheme 1a) and *N*-borylamine (Scheme 1b) products as determined by NMR spectroscopy. The reaction was proposed to follow the pathway shown below (Scheme 1): (1) oxidative addition of the diboron species to the metal center; (2) coordination of the imine with subsequent regioselective insertion into the M–B bond; (3)  $\beta$ -H elimination to generate the borylated amine.<sup>92</sup> Baker and Westcott also introduced the first application of Rh-catalyzed (Wilkinson's catalyst) hydroboration of an imine using HBpin in 2001.<sup>44,45</sup>

Subsequently in 2009, a boron-substituted analogue of the Shvo hydrogenation catalyst was introduced by Clark and co-workers in the hydroboration of aldehydes, ketones, and





Fig. 1 Au(I) complexes used in hydroboration of imines.<sup>44</sup>



Scheme 1 Rh catalyzed hydroboration of ketimine.<sup>92</sup>

imines (Fig. 2b). This is an example of a ligand–metal bifunctional catalyst which forms the boron-substituted catalyst *in situ* upon the addition of HBpin to the ruthenium dimer complex (Fig. 3). Clark's report only elaborates on a small number of imine substrates which undergo hydroboration (Table 1). Hydroboration occurs under 2–4 mol% catalyst loading at 70 °C in 17–120 h. The presence of electron-withdrawing groups within the substrate was found to increase the reaction rate.<sup>48</sup>

A more comprehensive report applying Ru for metal catalyzed imine hydroboration emerged in 2016 from the Gunanathan group (Fig. 2d), in which a commercially available  $[\text{Ru}(\text{p-cymene})\text{Cl}_2]_2$  precatalyst at low catalyst loadings (0.1 mol%) was employed. This readily available complex cata-




Fig. 2 Precious metal catalysts in imine hydroboration.




Fig. 3 Ruthenium catalytic active species.<sup>48</sup>

Table 1 Catalytic hydroboration of imines<sup>48</sup>


Entry	Substrate	Time (h)	Yield <sup>a,b</sup> (%)
1 <sup>b</sup>		120	82
2		22	86
3		17	74

<sup>a</sup> Isolated yield of amines. <sup>b</sup> 4 mol% Ru catalyst is used.

lyzed imine hydroboration *via* a proposed monohydrido-bridged intermediate  $[(\eta^6-p\text{-cymene})\text{RuCl}_2(\mu\text{-H}\cdot\mu\text{-Cl})]$  to form boronate amines. This reaction can be efficiently carried out at 60 °C in as little as 15 h with a much wider substrate scope compared to earlier reports employing Ru catalysts (10 examples). The formed boronate amines are conveniently hydrolyzed using silica gel in methanol for 6 h at 50 °C. The resulting secondary amines were isolated using column chromatography providing good yields (83–92%, Scheme 2).<sup>47</sup>

In 2017, a series of commercially available Re complexes were applied in several catalytic reactions by the Westcott group. The high-oxidation state complex of  $[(\text{PPh}_3)_2\text{Re}(\text{O})_2\text{I}]$  was found to be an efficient precatalyst in imine hydroboration; however, the scope of substrate studied was limited (4 examples). *N*-Benzylidenebenzylamine was found to be reduced smoothly using 5 mol% Re catalyst, 1.1 eq. HBpin at 80 °C within 2 h. However, aldimines derived from aromatic primary amines were found to be quite sluggish in this reaction (Table 2). Surprisingly, (*E*)-*N*-benzylideneprop-2-yn-1-amine proceeded in a chemoselective fashion which resulted in hydroboration exclusively at the C=N, while the alkyne group remained intact. It is proposed that the mechanism of this reaction is similar to that outlined by Fernandes and co-workers, through 1,2-addition across Re=O pi bond to generate a reactive rhenium hydride species along with a [Re]-OBpin group (Scheme 3).<sup>93</sup> This is the only Re-catalyzed imine hydroboration reported to date.<sup>94</sup>


Remarkably, a photocatalytic inverse hydroboration of imine was introduced in 2018 by Zhu and coworkers to produce  $\alpha$ -amino organoborons.<sup>46</sup> These  $\alpha$ -amino organoborons are potentially quite versatile building blocks in the syn-



**Scheme 2** Hydroboration of imine catalyzed by  $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$  and the conversion of amine boronates to secondary amine.<sup>47</sup>

**Table 2** Rhenium-catalyzed hydroboration of aldimines<sup>94</sup>


Entry	Substrate	Product	Yield
1			100
2			25
3			100
4			100




**Scheme 3** Addition of HBpin to  $[(\text{PPh}_3)_2\text{Re}(\text{O})_2\text{I}]$ .<sup>93</sup>

thesis of protease inhibitors and compounds for medicinal chemistry. In this work, they utilized N-heterocyclic (NHC) carbene boranes in the catalytic system along with the  $^*\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$  complex (Fig. 2e) in the presence of phenylmethanethiol and NaH as the inorganic base under the irradiation of 33 W CFL at room temperature (Scheme 4). The photoexcited  $^*\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$  complex readily accepts one electron from thiol ( $\text{C}_6\text{H}_5\text{CH}_2\text{SH}$ ) to form thiyl radical which can then reactively generate NHC-boryl radicals. This method is different from conventional strategies using toxic radical initiators to form active radical species.<sup>46</sup> Two possible pathways were proposed for inverse imine hydroboration which is shown in Scheme 5. Intermediate a (Scheme 5) was trapped by 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO).

Path (a) proposed a single electron transfer (SET) reduction of the imine with strongly reductive  $\text{Ir}^{\text{II}}$  followed by radical-radical C–B cross-coupling of the NHC-boryl radical with the  $\alpha$ -amino-benzyl radical. Path (b) suggested that NHC-boryl radical addition to the imine occurs first, with subsequent SET reduction of the nitrogen-centered radical. Both experimental methods and density functional theory (DFT) studies were employed to study the mechanism and revealed that a key factor in the reaction is the thiol organocatalyst; it facilitates the formation of boryl radical and suppresses the ionic



**Scheme 4** Selective catalytic inverse hydroboration of imine with N-heterocyclic carbene (NHC) boranes.<sup>46</sup>



**Scheme 5** Mechanistic hypothesis of iridium catalyzed inverse imine hydroboration.<sup>46</sup>

reduction pathway.<sup>46</sup> Both radical intermediates (Scheme 5a & b) were detected in the reaction process. However, the radical chain pathway (path a) is considered a minor contributor as the quantum yield of the model reaction was determined as 0.35.<sup>46</sup>

## 2.2. Base metal catalyzed hydroboration of imines

Although catalysis is a fundamental pillar of green chemistry, the use of sustainable and less toxic reagents and catalysts represents an additional improvement to this strategy.<sup>95</sup> The scarcity and toxicity of precious metals has increased the demand for inexpensive and more environmentally benign surrogates. Base metal catalysts, like first row transition metal complexes, are generally less expensive and have much lower toxicity.<sup>3</sup>


The first report of a base metal catalyzed imine hydroboration was disclosed in 2016. The Gordon group reported Ni(bpy)(cod) (Fig. 4a) to be an effective catalyst in imine hydroboration.<sup>51</sup> The metal center in this example is supported by a redox-active bipyridine ligand and this complex has been deployed in a range of both catalytic and stoichiometric transformations: ring-opening of aziridines,<sup>96</sup>  $\alpha$ -amino acid *N*-carboxy anhydrides,<sup>97</sup> coupling of CO with 1,2-diphenylacetylene<sup>98</sup> and, the dimerization of 1,4-substituted allenes<sup>99</sup> amongst others. It was proposed that the addition of pinacol-

borane as an external reductant in these reactions helped to facilitate the release of the reduced substrate from the nickel center and regenerate the catalytically relevant [Ni(bpy)] moiety. To illustrate the redox non-innocent nature of the ligand, a single-crystal X-ray diffraction experiment revealed a molecular structure consistent with a Ni<sup>I</sup>(bpy)<sup>+</sup> fragment with a nickel center in a pseudo-tetrahedral environment. This catalyst can reduce aryl-substituted imines in excellent yields (up to 96%) at 25 °C. The reaction conditions are mild with low catalyst loading (0.3 mol%) and the reaction is complete within 2.5 h; however, the scope of substrate examined was limited to only three examples (Table 3).<sup>51</sup>

A heterogeneous cobalt(II) coordination polymer precatalyst (Fig. 4b) was developed by the Zhang group in 2018.<sup>52</sup> This precatalyst is effective in imine hydroboration in the absence of any air- or moisture-sensitive organometallic reagents. Several aromatic imines with fluoro-, nitro- and ester-groups were studied as substrates. In all cases, the corresponding amine products could be obtained in good to excellent yield (up to 96%). The reaction conditions are presented as described in Scheme 6 with 0.1 mol% cobalt coordination polymer as catalyst along with 2 mol% KO<sup>t</sup>Bu in imine hydroboration at 70 °C in THF. This Co<sup>II</sup> coordination polymer catalytic system has the benefit of being air stable, highly efficient, and reusable.<sup>52</sup>

In 2020, a cationic [(iminophosphine)nickel(allyl)]<sup>+</sup> complex (Fig. 4c) was reported to effect catalytic imine hydroboration by Schmidt and coworkers. With loadings as low as 5 mol%, smooth conversion of *N*-allylimine substrates to their corresponding amine products was observed. Yields as high as 92% were obtained in the presence of 1.2 eq. HBpin at room temperature. Slightly sluggish reaction rates were observed in substrates with electron withdrawing groups.<sup>100</sup>

In that same year, the Gade group reported an iron alkyl precatalyst (Fig. 4d) in the enantioselective hydroboration of *N*-alkyl imines. A chiral bis(oxazolinylmethylidene)isoindoline pincer ligand was deployed in the asymmetric reduction of acyclic *N*-alkyl imines. Low catalyst loading of 0.5–3 mol% is used in the presence of HBpin at –40 °C and a wide range of substrates studied afforded excellent yields (31–99%) with up to >99% ee.<sup>101</sup>




**Fig. 4** Base metal catalysts in imine hydroboration.

**Table 3** Catalytic hydroboration of imines with Ni(bpy)(cod)<sup>51</sup>

Entry	Substrate	Catalyst (mol %)	Time (h)	Yield <sup>a</sup> (%)
1		0.3	2.5	83
2		0.3	2.5	89
3		0.3	2.5	96

<sup>a</sup> General condition: C<sub>6</sub>D<sub>6</sub> as solvent, room temperature. <sup>1</sup>H NMR yields using *tert*-butylbenzene as an internal standard.



**Scheme 6** Cobalt(II) coordination polymer catalyzed hydroboration of imine.<sup>52</sup> Reaction conditions: Imines (1.0 mmol), HBpin (1.1 mmol), cobalt CP (0.1 mol%), KOtBu (2 mol%), and THF (1 mL), 70 °C, 16 h, N<sub>2</sub>. Yields of isolated products of amines with GC conversion of imines are shown in parentheses.

At the beginning of 2021, the Findlater group reported the use of <sup>d</sup>P<sub>2</sub>BIANFe(Tol) as precatalyst in the rapid hydroboration of aldimines and ketimines to afford the corresponding secondary amines after hydrolysis through silica and hexanes. Low catalyst loadings (1 mol%) are employed in the presence of HBpin and NaO<sup>t</sup>Bu at 70 °C. In general, yields ranged from 46–99%, and most notably, the reaction was also effective in substrates featuring steric encumbrance with good to moderate yields observed (43–63%). Products can be readily isolated as ammonium salts. The synthetic utility of the reaction was demonstrated using gram scale reactions in which 90% isolated yield was obtained in a gram-scale reaction of *N*-benzylideneaniline.<sup>102</sup> The mechanism of the reaction was probed using kinetic, stoichiometric, and temperature-dependent rate experiments which suggest the formation of metal hydride species in the reaction mechanism. Subsequently, formation of an iron(boryl)(amine) was proposed to occur upon insertion of the C=N bond into the iron-hydride species.<sup>102</sup>

The first example of a zinc-catalyzed imine hydroboration was reported in 2021 by the Xu group.<sup>103</sup> In this example, a zinc dihydride complex stabilized by N-heterocyclic carbene ligands was shown to be highly efficient in catalytic hydroboration of nitriles and imines under low catalyst loading (0.5 mol%) and neat (solvent-free) conditions at room temperature. A wide scope of aldimines underwent rapid hydroboration. Most of the substrates reduced within 10–40 min with 82–92% isolated yields. However, aliphatic and sterically hindered substrates required longer reaction time (6–16 h) with 84–92% isolated yields.<sup>103</sup>

### 2.3. Rare earth metal catalyzed hydroboration of imines

Despite the widespread application of transition metal complexes as catalysts in imine hydroboration chemistry, rare earth metals (the f-element lanthanides and actinides) have not been as widely studied (Table 4). A boratabenzene metal


**Table 4** Transition metal catalyzed hydroboration of imines; comparison of reaction condition

Catalyst	Catalyst mol %	Solvent	Temp(°C)/ time (h)	TON	TOF (h <sup>-1</sup> )	Ref.
Ru 2016	0.1	Toluene	60/15	990	66	47
Ru 2009	2	Toluene	70/17–120	41	<1	48
Ir	1	CH <sub>3</sub> CN	RT/24	73	3	46
Re	5	C <sub>6</sub> D <sub>6</sub>	80/2	20	10	94
Ni(bpy)(cod)	0.3	C <sub>6</sub> D <sub>6</sub>	RT/2.5	200	80	51
Co	0.1	THF	70/16	990	62	52
Th	1	C <sub>6</sub> D <sub>6</sub>	80/12–24	100	8.3	49
Y	50	Toluene	75/48	1.87	<1	104
Er	2	Toluene	110/6	47	<8	50
Fe	1	Toluene	70/0.5	99	198	102
(Findlater)	0.5–3	Toluene	–40–RT/ 14–56	66	4.7	101
Fe (Gade)	—	—	RT/15 min	100	400	103
Zn	0.5	—	—	—	—	—


complex capable of reducing unsaturated organic molecules was first reported in 2011 (Fig. 5a).

The Xia group reported 1-*H*-boratabenzene complex of Y (Yttrium) in the hydroboration of alkenes, imines, and carbodiiimides as a metal ion dependent hydroboration. Boratabenzene is isoelectronic with the cyclopentadienide anion (Cp<sup>\*</sup>), and the catalytic behavior of boratabenzene metal complexes are greatly influenced by the boron substituent, due to the higher reactivity of the B–H bond as compared to the C–H bond on the ubiquitous Cp ligand (Fig. 6). The yttrium complex [C<sub>5</sub>H<sub>5</sub>BH]<sub>2</sub>YCl was prepared *via* the metathesis reaction between [C<sub>5</sub>H<sub>5</sub>BH]<sub>2</sub>Li and anhydrous YCl<sub>3</sub> in toluene. The hydroboration of benzylidene-*n*-propylamine, using [C<sub>5</sub>H<sub>5</sub>BH]<sub>2</sub>YCl, resulted in the amino-substituted boratabenzene complex [C<sub>5</sub>H<sub>5</sub>BN(<sup>t</sup>Pr)CH<sub>2</sub>Ph]<sub>2</sub>YCl in excellent yield (94%).<sup>104</sup>


In 2018, the Wang group reported a series of amidate-functionalized N-heterocyclic carbene (NHC) rare-earth metal (Er, Y, Gd, Dy) amido complexes in the hydroboration of imines and nitriles.<sup>50</sup> These complexes were synthesized *via* reaction



**Fig. 5** Rare earth metal catalysts in imine hydroboration.




**Fig. 6** 6 $\pi$ -Electron cyclopentadienyl.<sup>104</sup> aromatic anions: boratabenzene and

Scheme 7 Synthesis of rare earth metal complexes.<sup>50</sup>

between ligand precursor, ( $L = 1-(C_6H_5C=ONCH_2CH_2)-3-(CH_3)_3C_6H_2(N(CH_2)_2NC)$ ),  $RECl_3$  ( $RE = Er, Y, Dy, Gd$ ) and  $KN(SiMe_3)_2$  (KHMDS) in a one-pot fashion as depicted in Scheme 7. All complexes were crystallized to afford pure products from tetrahydrofuran (THF)/*n*-hexane and characterized by IR spectroscopy and elemental analysis. The complexes of Er and Gd turned out to be the most effective catalysts with high product yields of 96% and 95% respectively, in the hydroboration of benzylideneaniline (2 mol% catalyst loading).<sup>50</sup> The optimized reaction conditions were established to be 2 mol% catalyst, 1.5 eq. of HBpin, 0.5 ml toluene at a reaction temperature of 110 °C for 6 h. Various imine substrates were subjected to these hydroboration conditions employing the Er complex as the catalyst under study. The reaction afforded the desired secondary amines in excellent yields (50–99%) after work-up with column chromatography (methanol/silica gel). Several different functional groups including halogens, ethers and hydroxyl groups exhibited little effect on the progress of the reaction and were well tolerated. However, the presence of the strongly electron-withdrawing  $-NO_2$  group resulted in a decreased yield of 50%, and thus, it was proposed that the strongly electron-withdrawing nature of this group helped to disfavor the coordination of the imine to the metal center. Additionally, the chelation of the oxygen atoms of the nitro group to the metal center of the catalyst is also a possibility that would reduce the catalyst efficiency.<sup>50</sup> A plausible reaction mechanism was proposed as depicted in Scheme 8. The proposed mechanism postulated the formation of a rare-earth metal hydride species *via*  $\sigma$ -bond metathesis as a key step. Subsequently, imine  $C=N$  insertion into the  $Er-H$  bond is followed by the reaction with another HBpin molecule to release the product and regenerate the active metal hydride species.<sup>50</sup>

Subsequently, the Eisen group developed an N-heterocyclic iminato thorium complex (Fig. 5c) capable of catalyzing the hydroboration of aldimines in 2019. The Th(IV) precatalyst was synthesized by the protonolysis of a seven-membered N-heterocyclic iminato ligand and a metallacyclic thorium amide complex  $[(Me_3Si)_2N]_2Th[\kappa_2-(N,C)-CH_2Si(CH_3)_2N(SiMe_3)]$ . Hydroboration of *N*,*1*-diphenylmethanimine yielded 95% hydroborated amine product in the presence of 1 mol% Th catalyst upon heating to 80 °C after 24 h. A wide range of aldimines bearing electron-donating, withdrawing, neutral and aliphatic groups were studied with excellent yields (80–100%) to form hydroborated secondary amines.<sup>49</sup>

Scheme 8 Proposed reaction mechanism for imine hydroboration with amidate-functionalized N-heterocyclic carbene (NHC) Er complex.<sup>50</sup>


However, this catalytic system was incapable of the hydroboration of ketimines.<sup>49</sup> Stoichiometric reactivity studies, kinetic and temperature dependence measurements were performed in order to gain insight into the mechanism of the reaction. An analysis of the reaction kinetics revealed a first-order dependence on the catalyst concentration, HBpin, and no dependence on imine concentration (eqn (1)).

$$\frac{\delta p}{\delta t} = k_{obs} \times [Th]^1 [HBpin]^1 [PhCNPh]^0 \quad (1)$$

The activation parameters were calculated from Eyring and Arrhenius plots and  $\Delta S^\ddagger$ ,  $\Delta H^\ddagger$ , and  $E_a$  values of  $-36.46$  (0.8) e. u.,  $13.42$  (1.06) kcal mol $^{-1}$ , and  $14.05$  (0.77) kcal mol $^{-1}$  respectively were obtained.<sup>49</sup> A highly organized four-centered transition state with concerted bond-cleavage and bond-formation can be inferred from the  $\Delta S^\ddagger$  and  $\Delta H^\ddagger$  values. One plausible hydroboration mechanism was proposed in which the reaction initiates *via*  $\sigma$ -bond metathesis between the  $Th-N(SiMe_3)_2$  and HBpin. Subsequently, imine  $C=N$  insertion followed by the concerted addition of HBpin *via*  $Th-N/HBpin$   $\sigma$ -bond metathesis to afford amine product and regenerate the catalytically active species (Scheme 9). Unfortunately, no hydride signal ( $Th-H$ ) or deuteride were detected by NMR spectroscopy which would have indicated a rapid exchange equilibrium between the hydride at the metal and at the HBpin.<sup>49</sup>

#### 2.4. Main group element catalyzed hydroboration of imines


Catalysis employing main group elements has been widely studied in the context of hydroboration (Table 5). In 2012, the Crudden group introduced a Lewis acid-base adduct of DABCO·B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and Ph<sub>3</sub>C<sup>+</sup>/DABCO (DABCO = 1,4-diazabicyclo [2.2.2]octane) in hydroboration of aldimines and ketimines. The reaction between DABCO·B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> or Ph<sub>3</sub>C<sup>+</sup>/DABCO with HBpin result in three-coordinate boronium ions *in situ* which can catalyze imine substrates (Scheme 10a). Relatively broad substrate scope of aldimines and ketimines (13 examples) were studied with isolated yields of 70–96%. The reaction mecha-



**Scheme 9** Proposed mechanism of Th-catalyzed imine hydroboration.<sup>49</sup>

nism proposed to begin with the transfer of a boreonium ion from catalyst to imine. The resulting boron-activated iminium ion is then reduced by HBpin which is then assisted by the Lewis base DABCO according to the low temperature NMR studies (Scheme 10b).<sup>105</sup>

Later in 2013, the Hill group investigated the catalytic potential of alkaline earth metal complexes of  $\beta$ -diketiminato magnesium alkyl  $[\text{LMgBu}]$  ( $\text{L} = \text{CH}[\text{CMe}(\text{NDipp})_2]$  Dipp = 2,6-diisopropylphenyl).<sup>53</sup> This Mg complex was previously reported to be an efficient catalyst in hydroboration of aldehydes,<sup>106</sup> ketones<sup>106</sup> and pyridines.<sup>107</sup> This complex was also demonstrated to be highly effective precatalyst in imine hydroboration.<sup>53</sup> The reduction of imines proceeded smoothly in the presence of HBpin under mild reaction conditions of 5–10 mol% catalyst loading at 25–70 °C.<sup>53</sup> Imines with primary *N*-alkyl substituents, such as benzyl, *n*-butyl, 2-methoxyethyl underwent facile hydroboration under mild reaction conditions. However, the reaction was sluggish for the reduction of substrates bearing secondary (cyclohexyl) or tertiary (*tert*-butyl) alkyl substituents (1–3 days). The slow rate of reaction is compounded by competing catalyst decomposition over the reaction time period.<sup>53</sup> Interestingly, hydroboration of the pyridine-substituted substrate 3-(PhCH=N)C<sub>5</sub>H<sub>4</sub>N underwent



**Scheme 10** (a) Generation of boreonium ion *in situ*. (b) Mechanism of boreonium ion-catalyzed imine reduction.<sup>105</sup>


exclusive hydroboration at the exocyclic imine residue at 50 °C. Selected hydroboration examples of alkyl- and aryl-substituted imines are presented in Scheme 11.<sup>53</sup> Both electron-donating and electron-withdrawing substituents in the *ortho*-position were found to decrease the reaction rate. Ketimines derived from both aryl and alkyl ketones were also underwent successful reduction under increased temperature or catalyst loading.

Careful analysis of stoichiometric studies revealed the formation of (<sup>7</sup>Bu)Bpin and  $[\{\text{LMgH}\}_2]$  as identified by <sup>1</sup>H NMR spectroscopy.<sup>53</sup> This is in equilibrium with a magnesium borohydride complex of the  $[\text{H}({}^7\text{Bu})\text{Bpin}]^-$  ion (Scheme 12). Subsequent reaction with one equivalent of PhCH=NPh instantaneously afforded a bright-orange solution which faded over a period of three hours at room temperature. The <sup>1</sup>H NMR spectrum of the resulting colorless solution showed complete disappearance of the aldimine and MgH singlets and the formation of an amido complex. Addition of another equivalent of HBpin to this solution at room temperature led to the formation of the hydroborated product PhCH<sub>2</sub>N(Bpin)Ph.<sup>53</sup>


Kinetic studies reveal the reaction to be second order in [imine] and zero order in [HBpin]. In parallel room-tempera-

**Table 5** Main group element catalyzed hydroboration of imines; comparison of reaction conditions

Catalyst	Catalyst mol %	Solvent	Tem(°C)/time (h)	TON	TOF (h <sup>-1</sup> )	Ref.
Boreonium ion	5	PhCF <sub>3</sub>	RT/45 min–24 h	19.2	19.2	105
Mg	5–10	C <sub>6</sub> D <sub>6</sub>	25–70/0.2–96	19.8	99	53
Si	1	CH <sub>3</sub> CN	65/48–72	44	0.88	108
DAP (2017)	2	THF	25/16	47	3	55
TAP	0.1	THF	RT/16	937	58.5	56
2,4,6-BArF <sub>9</sub>	2	CH <sub>2</sub> Cl <sub>2</sub>	RT-60/4–24	47.5	12	57
BArF <sub>3</sub>	1	C <sub>6</sub> D <sub>6</sub>	RT/18	99	5.5	58
Al-Li	5	C <sub>6</sub> D <sub>6</sub>	RT/0.5–2	14	29	54
<sup>m</sup> NHOs	2	THF	70/12	48.5	4	116
LiBr	3	THF	RT/1	32	32	117
NaH	3	THF	60/24	33	1.37	118
PhSSPh	10	CH <sub>3</sub> CN	CFL (12 W)/14 h	9.9	0.71	120
<i>n</i> -BuLi	6	THF	RT/1 h	16.5	16.5	114



**Scheme 11** Scope of the magnesium-catalyzed hydroboration for a variety of alkyl- and aryl-substituted imines.<sup>53</sup> <sup>a</sup> NMR yield calculated by integration *versus* an internal standard of tetrakis(tri-methylsilyl)silane  
<sup>b</sup> 100% pyridine hydroboration. A complex mixture of 1,2- and 1,4-dihydropyridine and partial imine hydroboration products formed.




**Scheme 12** Stoichiometric step-by-step hydroboration of  $PhCH=NPh$  by complex  $[Mg]$ .<sup>53</sup>

ture experiments using the same catalyst/imine stock solution and 10 equivalents of HBpin, the reaction rate is decreased considerably due to the strong inhibition exhibited by HBpin *via* coordination to the magnesium center. According to stoichiometric imine insertion and HBpin  $\sigma$ -bond metathesis, the rate law of the reaction is reported as below:<sup>53</sup>

$$\text{rate} = -k \frac{[\text{cat}][\text{imine}]^2}{k_{\text{inhib}}[\text{HBpin}]_0} = -k' \frac{[\text{cat}][\text{imine}]^2}{[\text{HBpin}]_0}$$


Main group catalyzed imine hydroboration was extensively studied in 2017 by the Sen group. The catalytic activity of a



**Fig. 7** Main group element catalysis in imine hydroboration.

benzamidinato silane complex,  $[PhC(N^tBu)_2SiHCl_2]$  (Fig. 7c) was investigated. This silicon(IV) based catalyst can be easily synthesized *via* a single step in high yield.<sup>108</sup> Electron-donating, -withdrawing and electronically neutral aldimine substrates underwent efficient hydroboration under 1 mol% catalyst loading in the presence of HBpin at 65 °C. Corresponding secondary amine products were obtained after treating with silica/methanol at 65 °C. The mechanism is proposed to occur *via*  $\sigma$ -bond metathesis at the silicon center, followed by the formation of a four-membered transition state and addition of an Si-H bond to the C=N bond (Scheme 13). The mechanism was supported by DFT calculations. The energy barrier was calculated to be 40.8 kcal mol<sup>-1</sup> for the bond making and breaking between the Si-N and B-H bond. This high energy barrier explains why the reaction requires elevated temperatures to proceed (65 °C).<sup>109</sup>


A later study in 2017 reported by Speed and coworkers employed a chiral diazaphospholene catalyst in asymmetric imine hydroboration. Synthesis of this chiral catalyst is relatively straightforward; a three-step process from the commercially available 1-(1-naphthyl)ethylamine through reductive cyclization with  $PBr_3$  in the presence of cyclohexene to give the bromide diazaphospholene. No chromatographic purification is required (Scheme 14).<sup>55</sup> Several diazaphospholene (DAP) catalysts with various chiral groups were then explored in the hydroboration of imine substrates as the precursor to therapeutic entrasagiline. A diazaphospholene catalyst bearing larger and more rigid 1-naphthyl aromatic side chains demon-




**Scheme 13** The catalytic cycle and reaction mechanism of imine hydroboration by a silicon-amidinate catalyst, calculated at the PBE/TZVP level of theory with DFT.  $\Delta G$  and  $\Delta G^\#$  represent the Gibbs free energy of reaction and the Gibbs free energy of activation respectively. All values are in  $\text{kcal mol}^{-1}$ .<sup>59</sup>

strated the highest enantioselectivity for the hydroboration of alkyl imines \_enantiomer ratios of up to 88:12\_ in the presence of pinacolborane.<sup>55</sup> Twenty examples of imines bearing various functional groups were investigated under this asymmetric catalytic system (2 mol% catalyst). The enantiomeric ratios are among the best reports in alkyl imine hydroboration with pinacolborane (the enantiomeric ratios were determined by HPLC analysis on a chiral stationary phase).<sup>55</sup>

Concurrently, Johnson and Speed reported 1,2,4,5-triazaphospholenes (TAPs) halide catalysts (Fig. 7e) in imine hydroboration. A series of triazaphospholene halides and alkoxide were synthesized through the reaction between bulky amidrazones and phosphorus tribromide in the presence of triethylamine which cleanly provided triazaphospholene bromide. Triazaphospholene hydrides obtained subsequently under treatment with sodium benzyloxide in toluene.<sup>56</sup>




**Scheme 14** Synthesis of chiral diazaphospholene catalyst.<sup>55</sup>



**Fig. 8** Synthesized triazaphospholene halides and alkoxide for imine hydroboration.<sup>56</sup>

Triazaphospholenes are analogous to N-heterocyclic carbenes and yet there are few reports of them used in catalysis. Among the synthesized catalysts (Fig. 8), pyrrolo triazaphospholene halides appeared to be the most effective due to the ease of isolation of the corresponding amidrazone in imine hydroboration.<sup>56</sup> Nineteen imine substrates underwent hydroboration including those which contained cyclopropyl and alkyne groups. Heterocycles such as pyridine and furan were well tolerated under the reaction conditions.<sup>56</sup> Two pharmaceutical substrates (fendiline and rasagiline) were also demonstrated to undergo reduction using this protocol indicating its potential for applicability toward commercially important compounds.<sup>56</sup>

The mechanism of this reaction was probed through stoichiometric reactions and DFT calculations which indicated that phosphorus serves as a Lewis acid and there is a possible interaction between triazaphospholene catalyst and imine (Scheme 15).<sup>56</sup> DFT calculations with the LC- $\omega$ PBE functional and the exchange-hole dipole moment (XDM) dispersion model indicated a possible interaction of HBpin with the nitrogen bearing a *tert*-butyl group. Subsequent hydride transfer through a six-membered transition state leading to complex **III** (Scheme 15) would be exothermic, with an activation barrier of 23.0 kcal mol<sup>-1</sup> above complex **II**. Finally, the triazaphospholene cation is regenerated through B-N bond formation and dissociation of the borylated amine (modest barrier of 11.7 kcal mol<sup>-1</sup>) (Scheme 15).<sup>56</sup>




**Scheme 15** Proposed catalytic cycle and enthalpies calculated by DFT in triazaphospholenes catalyzed imine hydroboration. Energies given in  $\text{kcal mol}^{-1}$  relative to starting materials.<sup>56</sup>

Following the work of Johnson and Ferguson, the Melen group reported an example of a boron-based catalyst in imine reduction: tris(2,4,6-trifluorophenyl)-borane ( $2,4,6\text{-BAr}^F_3$ ). This catalyst is capable of reducing aldimines *via* hydroboration.<sup>57</sup> A range of electron donating, withdrawing, electronically neutral, and aliphatic aldimines were reduced under low catalyst loading and mild reaction conditions. For example, *N*-benzylideneaniline gave great yield of 95% (by *in situ*  $^1\text{H}$  NMR spectroscopy) to borylated amine at room temperature within 4 h. This Lewis acidic boron-based catalyst prevents any unwanted side-reactions with unsaturated frameworks in contrast to other Lewis acid boranes.<sup>110</sup>


Similarly, another boron Lewis acid, tris[3,5-bis(trifluoromethyl)phenyl]borane ( $\text{BAr}^F_3$ ), catalyzed imine hydroboration with pinacolborane (Scheme 16). The reaction enables effective hydroboration of both aldimines and ketimines within 18 h at room temperature. This catalyst does not require an external Lewis base and gives moderate to excellent yield of 30–99%.<sup>58</sup> In comparison with other Lewis acidic boranes,  $\text{BAr}^F_3$  was found to be much more reactive. The difference in catalytic activity can be attributed to decreased steric effects since both  $\text{BAr}^F_3$  and  $\text{B}(\text{C}_6\text{F}_5)_3$  have similar Lewis acidity (Fig. 9).<sup>111</sup>

A broad scope of ketimine substrates with a range of substituents underwent efficient hydroboration to afford amine products after aqueous workup (77–99% yield). Stoichiometric reactions between *N*,1-diphenylethan-1-imine and  $\text{BAr}^F_3$  formed the expected Lewis pair  $\text{PhCCH}_3=\text{NPh}(\text{BAr}^F_3)$  in  $\text{CD}_2\text{Cl}_2$  instantly. However, no reaction was observed between equimolar amounts of *N*,1-diphenylethan-1-imine and  $\text{HBpin}$ , as monitored by  $^1\text{H}$  and  $^{11}\text{B}$  NMR spectroscopy. Based on these findings and previous literature reports a proposed mechanism is presented in Scheme 17, which exclude the potential formation of borenium ion as it was not observed in the NMR spectra.<sup>58</sup>


In 2018, the catalytic activity of aluminum complexes were studied in transformations such as C–C cross coupling,<sup>112</sup>



**Scheme 16** Imine hydroboration catalyzed by Lewis acid tris[3,5-bis(trifluoromethyl)phenyl]borane ( $\text{BAr}^F_3$ ).<sup>58</sup>




**Fig. 9** Boron Lewis acid catalysts.<sup>111</sup>



**Scheme 17** Postulated catalytic cycle for  $\text{BAr}^F_3$  catalyzed imine hydroboration.<sup>58</sup>

catalytic hydroelementation such as alkyne and carbonyl hydroboration<sup>113</sup> etc. Following up on earlier reports, Mulvey and McLellan studied the catalytic activity of bimetallic lithium aluminates and neutral aluminum counterparts in hydroboration reactions. Hydroboration of *N*-benzylideneethylamine under 10 mol% of Al complexes at room temperature revealed the bimetallic complexes to be more efficient than monometallic analogues. This was the first report of the use of Al complexes in catalytic imine hydroboration. In addition, benzophenone imine was reduced in higher yield (up to 80%) in comparison to an aldimine substrate.<sup>54</sup> Stoichiometric experiments were carried out to gain insight into the reaction mechanism.

Results indicate that bimetallic lithium aluminate complexes *h* and *i* (Fig. 7) proceeds through deprotonation. Proposed mechanism under catalyst *i* is depicted in Scheme 18




**Scheme 18** Proposed mechanism for hydroboration of benzophenone imine catalyzed by *i*.<sup>54</sup>

which is started with deprotonation of benzophenone imine, followed by hydroboration and subsequently protonolysis to generate product and catalytically active species. The higher activity of bimetallic complexes is attributed to the key role of bi-metallic (Li-Al) cooperativity in these complexes which increases the polarization of reaction intermediates. This study revealed the importance of anionic ate complexes among main group catalysts with higher conversion in shorter timescale.<sup>54</sup>

Later on in 2019, the Xue group reported *n*-butyllithium (*n*-BuLi) as catalyst in imines hydroboration. A wide scope of substrates (18 examples) studied under 6 mol% *n*-BuLi at room temperature within 1–24 h. Isolated yields of 76–91% obtained. However, most of the substrates were among aldimines.<sup>114</sup> In the same year, the Speed group reported diazaphosphonium triflate in asymmetric catalysis of cyclic imines *via* hydroboration. 0.2–1 mol% catalyst loading is used and enantiomeric ratios of up to 97:3 obtained to form aryl/heteroaryl pyrrolidines and piperidines (22 examples- at 35 °C). Enantiomeric ratios determined by HPLC on a chiral stationary phase with 71–94% isolated yields. This is the first example of an asymmetric reaction catalyzed by a phosphonium cation.<sup>115</sup>

In 2020, a mesoionic N-heterocyclic olefin (<sup>m</sup>NHO) compound was developed and reported to be highly effective in imine hydroboration. Although, only two imine substrates were studied with this catalytic system.<sup>116</sup> In the same year, the lithium bromide catalyzed imine hydroboration was also disclosed. *N*-Benzylideneaniline was used as the model substrate with which to optimize the reaction conditions. This catalytic process was effective in a broad scope of substrate, both aldimines and ketimines bearing electron donating, withdrawing, electronically neutral substituents, and heteroatoms were reduced under mild low catalyst load (3 mol%) at room temperature. Chemoselective experiments with various functional groups (*i.e.* esters, amides, nitriles, or alkenes) revealed exclusive reduction of the imine moiety to form amine products. A plausible reaction mechanism was proposed using supporting DFT calculations (Scheme 19).<sup>117</sup>



Scheme 19 Plausible mechanism of imine hydroboration using LiBr.<sup>117</sup>


In 2020, that same group reported a readily available NaH catalytic system for aldimine hydroboration. Only 3 mol% NaH was required for this transformation to form amine products under minimal solvent. The reaction is carried out in THF at 60 °C. This reaction time was considerably longer than the prior study. This protocol is chemoselective in the presence of other reducible groups, including esters, amides, nitriles, alkyl halides, and epoxide. The mechanism of the reaction also was elucidated by means of DFT calculations.<sup>118</sup>

A recent study described how protic additives (*i.e.*, alcohols or water) can promote alkyl imine reduction with pinacolborane.<sup>119</sup> However, aniline derived imines were not found to be effectively reduced upon addition of a stoichiometric amount of alcohol or water. This can be attributed to the reduced basicity of the aniline derived imines. Furthermore, amine impurities can also promote imine reduction in some cases resulting in a slow autocatalytic reaction.<sup>119</sup>

In 2021, an inverse imine hydroboration was reported by the Kawamoto group using NHC-boranes (1,3-dimethylimidazole borohydride). This hydroboration is free of any photoredox catalyst, and it is prompted by visible light irradiation of an acetonitrile solution of substrate in the presence of diphenyl disulfide (PhSSPh, 10 mol%). A relatively broad substrate scope (12 examples) was studied with high isolated yields of 69–95%. The mechanism proposed proceeds through a radical chain reaction *via* a nucleophilic boryl radical (Scheme 20).<sup>120</sup>

In the same year, borane-dimethyl sulfide (Me<sub>2</sub>S-BH<sub>3</sub>) catalyzed hydroboration of imines was reported by the Vidović group. They proposed that Me<sub>2</sub>S-BH<sub>3</sub> acts as a hydride donor in the reduction of various imines with good to moderate isolated yield of 44–88% (14 examples).<sup>121</sup>

Noticeably, a catalyst free and solvent free imine hydroboration was report in 2019. This methodology is capable and compatible using 1.1–1.3 eq. HBpin. A wide scope of aldimines ranging from electronically diverse and sterically hindered examples underwent reduction at ambient temperature, although the reaction time is longer as compared to catalyzed protocols (6–24 h). Moreover, this catalytic system was unsuc-



Scheme 20 Chain mechanism for the inverse hydroboration of imines.<sup>120</sup>

cessful in ketimine reduction under the same reaction conditions (at room temperature). Ketimine hydroboration was observed at a higher temperature of 60 °C, but this was much less effective than previously reported catalyzed reactions. This protocol offers a chemoselective conversion of imine over alkene, alkyne, ketone, nitro, and nitrile.<sup>66</sup>

### 3. Hydroboration of nitriles

#### 3.1. Transition metal catalyzed hydroboration of nitriles

The reduction of nitriles to primary amines has been carried out through hydroelementation reactions such as hydroboration and hydrosilylation. Like imines, this method resolves the problems associated with high-pressure setups, eliminates the generation of inorganic wastes, and possesses improved selectivity. Nitrile hydrosilylation forms both the monosilylated imine and disilylamine products; however, hydroboration produces the diborane amine selectively.<sup>47</sup> Nitriles have a strong C≡N bond dissociation energy (750.0 kJ mol<sup>-1</sup>) which makes them hard to reduce under the typical reaction conditions.<sup>82</sup> However, the presence of a catalyst can facilitate the process.

The first catalytic nitrile hydroboration was reported by the Nikonov group in 2012. An imido-hydrido complex of Mo(IV) was developed and deployed in the hydroboration of acetonitrile and benzonitrile using catecholborane. Reaction progress was monitored employing <sup>1</sup>H NMR which indicated complete conversion was achieved in the presence of 5 mol% catalyst within 12 h at 22 °C.<sup>67</sup>

To date, more than a dozen transition metal catalysts have been reported to be effective in nitrile hydroboration. In 2015, a series of proton-switchable bifunctional ruthenium complexes was introduced by the Szymczak group.<sup>75</sup> These metal complexes contain pincer ligands. At 5 mol% loading, the Ru complex (example Fig. 10b) could convert nitrile substrates to borylated amines in the presence of 2 eq. HBpin and C<sub>6</sub>D<sub>6</sub> as solvent. A wide range of *p*-substituted aryl nitriles can thus be hydroborated in moderate to excellent yields. Functional groups such as methyl esters, trifluoromethyl, 2-pyridyl, benzyl ethers and 2-furyl were all well tolerated. In addition, stoichiometric reactions with H<sub>2</sub> and HBpin revealed B–H bond activation and is facilitated through Ru<sup>II</sup>–O bond.<sup>75</sup> This provides Ru–H species available for reaction with the neighboring atoms. The reaction is also revealed to be first order with respect to catalyst and zero order in both HBpin and benzonitrile. This proposed reaction mechanism is presented in Scheme 21 depicting the critical role of the bifunctional ligand in the reaction: (1) B–H bond heterolysis, (2) Bpin binding to oxygens of the ligand and cooperative interaction with substrate, and (3) a proton-switchable assist at the metal center.<sup>75</sup>

Another example of Ru catalyzed nitrile hydroboration was reported by the Gunanathan group in 2016 (Fig. 10c). Using 1 mol% of a commercially available [RuCl<sub>2</sub>(*p*-cymene)]<sub>2</sub> catalyst, diboryl amines could be prepared under solvent-free conditions at 60 °C. The reaction proceeds *via* the *in situ* for-






Fig. 10 Transition metal catalysis in nitrile hydroboration.



Scheme 21 Proposed mechanism in Ru catalyzed nitrile hydroboration.<sup>75</sup>

mation of imines.<sup>47</sup> Interestingly, this catalyst is capable of the chemoselective hydroboration of nitriles over esters. Moreover, stoichiometric reaction between the Ru complex and HBpin produces a singlet resonance in the metal-hydride region ( $\delta_{\text{Ru-H}} = -10.18$  ppm) which confirms the immediate formation of a mono hydrido-bridged dinuclear complex of form:  $[(\eta^6\text{-}p\text{-cymene})\text{-RuCl}] (\mu\text{-H}\text{-}\mu\text{-Cl})$  in the reaction mixture (Scheme 22).<sup>47</sup>




**Scheme 22** Mono hydrido-bridged dinuclear complex of  $\{[(\eta^6\text{-}p\text{-cymene})\text{-RuCl}_2](\mu\text{-H}\text{-}\mu\text{-Cl})\}$  in nitrile hydroboration.<sup>47</sup>


An iron-indium cooperative catalyst was introduced by Nakazawa:  $[\text{Fe}(\text{CH}_3\text{CN})_6][\text{cis-Fe}(\text{CO})_4(\text{InCl}_3)_2]$ . The double hydroboration of organonitriles could be accomplished with 5–10 mol% of this iron–indium catalyst using several different reducing agents HBpin, HBCat and 9-BBN at 80 °C within 24 h. Both aliphatic and aromatic substrates underwent hydroboration with good to moderate isolated yields of 47–87% (Scheme 23). A plausible reaction mechanism is presented in Scheme 24 and involves the dissociation of one  $\text{InCl}_3$  of the catalyst to afford a monoindium iron complex. Subsequent ligand exchange occurs between CO and the organonitrile substrate.<sup>122</sup>

Several base metal catalysts were reported in 2017, among them examples of Co<sup>123</sup> and Ni<sup>124</sup> complexes. The Fout group reported a Co<sup>I</sup> system which was effective in catalytic nitrile double hydroboration (Fig. 10e).<sup>123</sup> The scope of substrate explored in this report was limited (5 examples). The highest yield reported was 85% in the hydroboration of 3,5-difluorobenzenonitrile where the product was isolated as the corresponding ammonium salt.<sup>123</sup>


Later in the same year, an interesting tetradentate  $\alpha$ -diimine cobalt hydride catalyst was reported by the Trovitch group.<sup>73</sup> This catalyst was prepared from the addition of  $\text{NaEt}_3\text{BH}$  to the precursor dihalide complex,  $(^{\text{Ph}_2\text{PPr}}\text{DI})\text{CoCl}_2$  (Scheme 25). X-ray diffraction experiments and DFT calculations revealed that the monohydride,  $(^{\text{Ph}_2\text{PPr}}\text{DI})\text{CoH}$  complex



**Scheme 23** Double hydroboration of organonitriles in the presence of  $[\text{Fe}(\text{CH}_3\text{CN})_6][\text{cis-Fe}(\text{CO})_4(\text{InCl}_3)_2]$ .<sup>122</sup>



**Scheme 24** Proposed catalytic cycle for double hydroboration of organonitriles in the presence of  $[\text{Fe}(\text{CH}_3\text{CN})_6][\text{cis-Fe}(\text{CO})_4(\text{InCl}_3)_2]$  catalyst.<sup>122</sup>



**Scheme 25** Monohydride cobalt catalyst in nitrile hydroboration.<sup>73</sup>

(Scheme 25) contains a radical monoanion  $\alpha$ -diimine ligand and a Co<sup>II</sup> center with a distorted square pyramidal geometry.<sup>73</sup> Addition of 1 mol% of the Co hydride catalyst to nitrile substrates in the presence of HBpin afforded the corresponding diboryl amines in good to moderate isolated yields of 30–88% within 24 h.<sup>73</sup>

The Shimada group employed commercially available nickel salts of bis(acetylacetonato)nickel(II) and found them to be highly effective in nitrile hydroboration.<sup>124</sup> Initially, the catalytic activity of different metal salts in hydroboration of benzonitrile with HBCat at room temperature was explored (Table 6).<sup>124</sup> Bis(2,2,6,6-tetramethyl-3,5-heptanedionato)nickel (II) (Table 6-entry 3) was revealed to be the most effective catalyst amongst other metal salts. Nitriles with various functional groups underwent reduction with excellent yield up to 95% isolated yield under 0.5–5 mol% catalyst loading with HBCat (benzene as solvent/at room temperature). The proposed reaction mechanism is presented in Scheme 26 which is through oxidative addition of Ni(0) species to HBCat, leading to the formation of boryl hydride intermediate.<sup>124</sup>


In 2018, an N-heterocyclic carbene-supported complex of Er was disclosed by the Wang group. This rare-earth metal catalyst (2 mol%) was found to exhibit excellent catalytic activity (89–99% isolated yield) in nitrile hydroboration in toluene.<sup>50</sup>

The Tobita group reported a Ru complex to be an effective catalyst in nitrile hydroboration. This 16-electron ruthenium complex bears a bis(silyl)xanthene chelate ligand xantsil; Ru

**Table 6** Hydroboration of PhCN with HBcat catalysed by different metal salts<sup>124</sup>

Entry	Catalyst	Yield <sup>a,b</sup>
1	$\text{Ni}(\text{acac})_2$	>99(49) <sup>c</sup>
2	$\text{M} = \text{Ni}$ , $\text{R} = \text{CF}_3$	>99(19) <sup>c</sup>
3	$\text{M} = \text{Ni}$ , $\text{R} = \text{tBu}$	>99(78) <sup>c</sup>
4	$\text{NiCl}_2$	ND
5	$\text{Ni}(\text{cod})_2$	87
6	$\text{Co}(\text{acac})_2$	37
7	$\text{Cu}(\text{acac})_2$	ND
8	$\text{Mn}(\text{acac})_2$	ND
9	$\text{Pd}(\text{acac})_2$	ND


<sup>a</sup> Catalyst (0.01 mmol), PhCN (0.20 mmol), and HBcat (0.44 mmol) in 5 mL  $\text{C}_6\text{H}_6$  at room temperature for 12 h. <sup>b</sup>  $^1\text{H}$  NMR yield based on an internal standard ( $\text{PhSiMe}_3$ ). <sup>c</sup> The values in parenthesis represent the yields at a 1 mol% catalyst loading.



**Scheme 26** Possible mechanism of hydroboration of nitrile catalyzed by bis(2,2,6,6-tetramethyl-3,5-heptanedionato)nickel(II).<sup>124</sup>

$[\kappa_3(\text{Si},\text{O},\text{Si})\text{-xantsil}](\text{CO})(\text{PCyp}_3)$  ( $\text{Cyp} = \text{cyclopentyl}$ ).<sup>74</sup> Interestingly, this complex functions as a catalyst in both double and single hydroboration of nitriles. HBpin and 9-BBN (9-borabicyclo[3.3.1]nonane) were employed to form bis(boryl) amines and *N*-borylimines, respectively in mostly excellent yield up to 99%. Beyond the hydroboration pathway, this study also describes deborylative C–N coupling reactions of borylated amines utilizing  $\text{Pd}(\text{dba})_2$  ( $\text{dba} = \text{dibenzylideneacetone}$ ) as catalyst (Scheme 27).<sup>74,125</sup>

A complex of the actinide thorium was recently introduced by Eisen and coworkers as a nitrile hydroboration catalyst (Fig. 10k).<sup>49</sup> This metallacyclic thorium amide complex,  $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{Th}[\kappa_2\text{-}(\text{N},\text{C})\text{-CH}_2\text{Si}(\text{CH}_3)_2\text{N}(\text{SiMe}_3)]$ , is highly effective in nitrile hydroboration with very good TOF of




**Scheme 27** One-pot synthesis of *N,N*-diarylamines or *N*-arylimines from nitriles by double or single hydroboration and C–N coupling.<sup>74</sup> Conditions for synthesis of bis(boryl)amines: nitrile (0.29 mmol), HBpin (0.63 mmol), Ru catalyst (7  $\mu\text{mol}$ , 2 mol%), 40–60  $^\circ\text{C}$ , THF; bromoarenes (0.61 mmol),  $\text{KO}^\ddagger\text{Bu}$  (0.61 mmol),  $\text{Pd}(\text{dba})_2$  (6.1  $\mu\text{mol}$ , 2 mol%), CyJohnPhos (14  $\mu\text{mol}$ , 5 mol%), 60  $^\circ\text{C}$ , 12 h, THF. Conditions for synthesis of *N*-borylimines: nitrile (0.099–0.29 mmol), 9-BBN (0.12–0.32 mmol for the monomer), Cat. (0.7–1.4  $\mu\text{mol}$ , 0.5–0.7 mol%), 1 h, rt, THF; bromoarenes (0.11–0.32 mmol),  $\text{KO}^\ddagger\text{Bu}$  (0.12–0.32 mmol),  $\text{Pd}(\text{dba})_2$  (2.1–6  $\mu\text{mol}$ , 2 mol%), CyJohnPhos (4–14  $\mu\text{mol}$ , 4–8 mol%), 60  $^\circ\text{C}$ , 12 h, THF. Isolated yields based on nitriles.


up to 500  $\text{h}^{-1}$ . A wide scope of substrates bearing different functional groups were studied under low catalyst loading (0.1 mol%) with excellent yields (75–100%).<sup>49</sup>

The mechanism of this reaction was studied using kinetic, stoichiometric, and temperature dependence experiments. Taken together, these results were interpreted to generate a proposed mechanism in which a facile activation of the pre-catalyst occurs *via* insertion of the HBpin into the thorium metallacycle to afford the active thorium hydride complex as depicted in Scheme 28. The activation parameters for the hydroboration of  $\text{PhCH}_2\text{CN}$  were determined from Eyring and Arrhenius plots. ( $\Delta H^\ddagger = 6.99$  (0.44) kcal  $\text{mol}^{-1}$   $\Delta S^\ddagger = 47.25$  (1.23) e.u.,  $E_a = 7.66$  (0.44) kcal  $\text{mol}^{-1}$ ).<sup>49</sup>

In the same year, another report of cobalt catalyzed nitrile hydroboration was disclosed by the Trovitch group.<sup>126</sup> This low-spin Co(II) catalyst (1 mol%) is capable of the hydroboration of nitrile substrates efficiently with TOF of up to 380  $\text{h}^{-1}$ , and 14 nitrile examples were reduced under this method with mostly excellent yields obtained at ambient temperate within 2 h. DFT calculations revealed catalyst activation to occur through a Co–N bond hydroboration, implying that through



**Scheme 28** Proposed mechanism for Th catalysed hydroboration of nitriles.<sup>49</sup>




**Scheme 29** Co(III) hydride species in nitrile hydroboration.<sup>126</sup>


the course of the catalytic cycle all steps involve high-spin Co ( $\mu$ )-species (Scheme 29).<sup>126</sup>

Recently, the Panda group reported examples of Ti catalysts in nitrile hydroboration. These were the first reports of titanium acting as catalyst in the synthesis of organo-boron compounds (Schemes 30–32).<sup>77,127</sup> The titanium amido bond in these catalysts are labile allowing the facile formation of a metal hydride (Schemes 30 & 31).<sup>127</sup> The metal hydride species can effectively catalyze the hydroboration of unsaturated C–X bonds. Hydroboration of nitriles proceeds in excellent yield in neat conditions under 10 mol% catalyst loading at 65 °C. A wide scope of substrates bearing electron donating, electron withdrawing, neutral, and heteroatoms are reduced in 80–99% yield within 2–10 h using both HBpin and HBcat. Hydrolysis of diboryl amine products was performed under treatment with 0.05 M aqueous HCl for 4 h at room temperature to afford the corresponding amines.<sup>127</sup>

A cobalt catalyzed nitrile hydroboration reaction was reported by the Khalimon group in 2020. This group utilized the commercially available  $\text{Co}(\text{acac})_2$  (acac = acetylacetone) in conjunction with dpephos (bis[(2-diphenylphosphino)phenyl]ether) ligand in nitrile hydroboration at ambient temperature. *N,N*-Diborylamines contain labile B–N bonds and can be uti-




**Scheme 30** Selected examples of Ti catalyzed double hydroboration of nitriles in the presence of HBpin.<sup>127</sup> Reaction conditions: Catalyst (10 mol%), nitrile substrates (1 eq.), HBpin (2.2 eq.). The reaction mixture was heated to 65 °C. Isolated yields.




**Scheme 31** Selected examples of Ti catalyzed double hydroboration of nitriles in the presence of HBcat.<sup>127</sup> Reaction conditions: Catalyst (10 mol%), nitrile substrates (1 eq.), HBcat (2.2 eq.). The reaction mixture was heated at 65 °C. Yields were calculated based on <sup>1</sup>H NMR (400 MHz) integration of characteristic product peak using 50 mol% of 1,3,5-trimethoxybenzene as an internal standard.

lized in the synthesis of a variety of N-containing organic molecules.<sup>128</sup> To this end, a one-pot transformation of nitriles to aldimine was achieved *via* hydroboration and subsequent *in situ* reaction of diborylamine with aldehydes to give aldimines. The aldimine products were purified by flash column chromatography using silica gel neutralized with  $\text{NET}_3$ .<sup>128</sup>

Khalimon and co-workers proposed that the reaction mechanism proceeds through the formation of a (dpephos)Co<sup>I</sup>-H species.<sup>128</sup> This would involve the migratory insertion of a bound nitrile into the Co-H bond and the subsequent reaction with HBpin to generate the *N*-borylimine RCH = N(Bpin). This species subsequently undergoes insertion into the Co-H



**Scheme 32** Selected examples of Ti catalyzed double hydroboration of nitriles in the presence of H-Bpin.<sup>127</sup> Reaction conditions: Catalyst (3 mol%), nitriles (1 eq.), H-Bpin (2.2 eq.), neat condition at 60 °C; yield was calculated by  $^1\text{H}$  NMR (400 MHz) with respect to HMB (15 mol%) as the internal standard.



**Scheme 33** Proposed mechanism for  $\text{Co}(\text{acac})_2$  catalyzed hydroboration of nitriles.<sup>128</sup>

bond to yield *N*-borylamide and finally the *N,N*-diborylamine (Scheme 33).<sup>128</sup>

Very recently in 2021, Wang and Guo reported a mixture of cobalt(II) iodide and potassium *tert*-butoxide to be effective in double hydroboration of nitriles.<sup>129</sup> The alkoxide-pinacolborane combination activates cobalt(II) iodide to generate a metastable heterotopic cobalt catalyst *in situ* employing 0.5–2 mol%  $\text{CoI}_2$  and 10 mol%  $\text{KO}^t\text{Bu}$ . A wide range of aliphatic and aromatic nitriles were studied and isolated yields (as ammonium salts) as high as 99% could be achieved; catalytic reaction mixtures were stirred under  $\text{N}_2$  at 30 °C for 4–24 h. Substrates with reductive groups of formyl and acetyl were further investigated. Both the cyano and carbonyl groups were reduced under standard conditions with a moderate yield of 74% and 64% of the doubly reduced product, respectively.<sup>129</sup>

Another base metal catalyst deployed in nitrile diborylation was reported by the Findlater group using  $^{\text{dpp}}\text{BIANFe}(\text{Tol})$  pre-catalyst (1 mol%) under solvent-free and additive-free conditions (70 °C or RT). In this investigation, a small substrate scope was studied. The reaction was found to tolerate bromo- and chloro-substituents and aliphatic (cyclic) substrates. All substrates studied afforded diborylated amines in good to excellent yields of 84–99%. Products can be conveniently isolated as ammonium salts.<sup>102</sup>

Subsequently, manganese catalyzed hydroboration of nitriles introduced in 2021.<sup>130,131</sup> The Wangelin group

reported the manganese bis(hexamethyldisilazide) as pre-catalyst in nitrile dihydroboration under mild reaction condition (5 mol%  $\text{Mn}(\text{hmds})_2$ , 20 °C within 20 h). A broad scope of aromatic and aliphatic nitriles substrates studied to form primary amines under this protocol with high yields (26 examples, 36–99%). Study of the reaction mechanism revealed the efficient catalytic activity resides in the ability of the metal center to act as Lewis acid. This is corroborated by isolation of an adduct upon the reaction between  $\text{Mn}(\text{hmds})_2$  and PhCN in hexanes.<sup>131</sup>

In 2022, the Tamm group reported N-heterocyclic carbene-phosphinidene complex of Ru ( $[(\eta^6-p\text{-cymene})\{\text{IMes}\}\text{P}]\text{RuCl}$ ) in nitrile hydroboration. 2.1 eq. H-Bpin is used under 3 mol% catalyst loading and neat condition at 60 °C for 11 h. Isolated yields of borylated amines are reported (82–99%\_15 examples). Preliminary mechanistic, kinetic studies and stoichiometric reactions revealed the initial formation of monohydride complex of  $[(\eta^6-p\text{-cymene})\{\text{IMes}\}\text{P}]\text{RuH}$  as the catalytically active species.<sup>132</sup>


### 3.2. Main group element catalyzed hydroboration of nitriles

The first example of a main group catalyst in nitrile hydroboration was introduced by Hill and co-workers in 2016 with the report of a single  $\beta$ -diketiminato *n*-butylmagnesium pre-catalyst. In this work, 15 nitrile examples underwent hydroboration using 10 mol% catalyst at 60 °C, over 0.5–30 h with good to excellent isolated yields (43–96%). Dihydroboration reaction was facile for 1-cyanoethyl substrate with isolated yield of 70% within 0.5 h. A sluggish reaction was observed in the case of diphenylacetonitrile substrate (43% after 30 h) which was attributed to steric encumbrance at the substrate.<sup>71</sup>



The Panda group reported an aluminum alkyl complex pre-catalyst in 2018 capable of nitrile hydroboration to afford diborylamines under solvent-free conditions and mild temperatures (60 °C).<sup>133</sup> This aluminum complex was prepared by the treatment of  $[2\text{-F-C}_6\text{H}_4\text{NHP}(\text{Se})\text{Ph}_2]$  and  $\text{AlMe}_3$  in toluene (Scheme 34).<sup>133</sup>

In this work, an extensive range of substrates (up to 30 examples) were studied under this protocol using both H-Bpin and H-BCat as the borylating agent (Schemes 35 & 36).<sup>133</sup> The end products were isolated using aqueous HCl (0.05 M) at room temperature for two hours, which gave a fine white powder of substituted benzyl ammonium chloride (Scheme 37). Moreover, the reaction mechanism is proposed through the formation of an aluminum hydride as the active species.<sup>133</sup>


In the same year, an unsymmetrical magnesium(I)  $\beta$ -diketiminate complex was synthesized by the Ma group. This




**Scheme 34** Synthesis of aluminum(III) complex.<sup>133</sup>



**Scheme 35** Selected examples of aluminum-catalyzed hydroboration of nitriles with HBpin.<sup>133</sup> Reaction conditions: Al catalyst (1 mol%), nitrile (1 eq.), HBpin (2.2 eq.), yield was calculated via <sup>1</sup>H NMR (400 MHz) integration of characteristic product signal present in the reaction.



**Scheme 36** Selected examples of aluminum-catalyzed hydroboration of nitriles with HBcat.<sup>133</sup> Reaction conditions: Al catalyst (1 mol%), nitriles (1 eq.), HBcat (2.2 eq.), yield was calculated based on <sup>1</sup>H NMR.



**Scheme 37** Hydrolysis of diborylamines to primary amines.<sup>133</sup>


alkaline earth-based catalyst was found to be effective with various nitrile substrates for hydroboration with a range of functional groups. Surprisingly, this complex has good solubility even in hexanes which makes it a highly efficient catalyst in almost all organic solvents.<sup>134</sup>

A year later, in 2019, closely related Al catalysts were independently reported by Ma and Yang<sup>31</sup> and Roesky<sup>82</sup> almost simultaneously. These groups employed (<sup>DiPP</sup>NacNac)Al(Bu)<sub>2</sub> and (EtPhNacNac)AlH<sub>2</sub> as efficient catalysts in nitrile hydroboration under solvent free conditions.<sup>31,82</sup> These reactions can be used to convert both aliphatic and aromatic nitriles to the corresponding *N,N*-diborylamines with 2 eq. HBpin and 3–5 mol% catalyst loading with good to excellent yields. (<sup>DiPP</sup>NacNac)Al(Bu)<sub>2</sub> is comparatively more effective as the yields are typically greater than 90% for all substrates studied. The reaction was equally effective for both aliphatic and aromatic substrates. (<sup>DiPP</sup>NacNac)Al(Bu)<sub>2</sub> catalyzed hydroboration was shown to be less sensitive to the electronic and steric properties of nitrile substrates which afforded higher yields and milder reaction conditions (60 °C, 0.25–4 h as compared with 80 °C, 6–24 h). Both methods are believed to proceed through a key nitrile insertion step into the Al–H bond of the catalytically active species.<sup>31,82</sup>


A well-defined aluminum dihydride complex bearing conjugated bis-guanidinate ligand was reported as catalytically relevant in nitrile dihydroboration by the Nembenna group in 2020 (Fig. 11). This conjugated bis-guanidine (CBG) supported aluminum dihydride complex was synthesized *via* treatment of CBG free ligand, LH, with 1 eq. of Alane, H<sub>3</sub>Al·NMe<sub>2</sub>Et in toluene at 80 °C (Scheme 38).<sup>135</sup>

In this case, catalytic studies were initiated with the model substrate benzonitrile which was used to optimize reaction conditions. For the model system, a 97% yield was obtained under 3 mol% catalyst loading, 2 eq. HBpin at 60 °C with neat conditions.

A wide range of organic nitriles were investigated using this catalytic approach; a number of diverse substituents such as




**Fig. 11** Reported main group element catalysts in nitrile hydroboration.




**Scheme 38** Synthesis of conjugated bis-guanidinate supported aluminum dihydride complex.<sup>135</sup>

OMe, Cl, and F are all well tolerated.<sup>135</sup> Substrates bearing the reducible carbonyl functional group were incompatible with this system. These reactions resulted in 83% and 52% diborylated products using 4-formylbenzonitrile and 4-acetylbenzonitrile as substrates respectively. Moreover, gram-scale hydroboration of benzonitrile (1 g) was also successful (Scheme 39). The mechanism of this reaction is proposed to proceed through the formation of an intermediate imine complex followed by association of HBpin to the Al center and addition of H-B across the Al-N bond *via* a four-membered transition state (Scheme 40). In addition, the stoichiometric reaction between the catalyst and trimethylacetonitrile in  $\text{CDCl}_3$  at 70 °C provided supporting evidence for the postulated aluminum imine intermediate.<sup>135</sup>

Subsequently, the von Wangelin group reported lithium hexamethyldisilazide ( $\text{LiN}(\text{SiMe}_3)_2$ ) or Li(hmds) as a catalyst in nitrile hydroboration. Here, the optimized reaction conditions were found to be 10 mol% catalyst loading at 20–70 °C to




**Scheme 39** Large-scale hydroboration of benzonitrile with HBpin.<sup>135</sup>



**Scheme 40** Proposed mechanism for hydroboration of nitrile using Al catalyst.<sup>135</sup>

obtain yields range of 64–99% for both aromatic and aliphatic substrates within 20 h.<sup>136</sup> This readily available compound is suitably Lewis acidic to activate the nitrile substrates while also Lewis basic enough to convert the borane to an-ate complex.<sup>136</sup> Various alkali metal alkoxides and amide bases with bulky groups were studied in benzonitrile hydroboration, and  $(\text{LiN}(\text{SiMe}_3)_2)_2$  gave excellent yields of 70–96% of the desired benzylamine products at room temperature.<sup>136</sup> Further optimization of the reaction using various borylating reagents such as  $\text{NH}_3\text{BH}_3$ ,  $\text{NMe}_3\text{BH}_3$ , HBpin and HBcat identified HBpin to be the most suitable. A broad substrate scope was investigated and the conditions were found to tolerate iodo-, bromo-, and thiomethyl-substituents even in the presence of heteroarenes such as thiophene, furan, and pyridine.<sup>136</sup> However, higher temperatures were required to facilitate reactions of the electron-rich 4-*N,N*-dimethylaminobenzonitrile and the bulky *ortho*-tolunitrile. Additionally, the reaction is highly chemoselective for nitriles over styrenic substrates. Kinetic studies revealed a first order rate with respect to both lithium catalyst and nitrile. The proposed mechanism is based upon stoichiometric and kinetic analysis and invokes the formation of Lewis acid–base adduct, the amidinate  $[\text{RC}(\text{NSiMe}_3)_2\text{Li}\cdot\text{NCR}]$  (Scheme 41).<sup>136</sup>

A highly efficient solvent-free double hydroboration of nitriles was reported by the Findlater group in 2020. Commercially available sodium triethylborohydride was employed as the catalyst and diboryl amine products were easily obtained in up to 99% yield using 5 mol% catalyst. An array of substrates (22 examples) was reported to undergo facile hydroboration. Substrates with aliphatic and aromatic groups, and heteroatoms underwent hydroboration with excellent isolated yields of 74–99%. High TOF of 2400  $\text{h}^{-1}$  is



**Scheme 41** Proposed reaction mechanism of Li catalyzed hydroboration of nitriles in the presence of HBpin.<sup>136</sup>


obtained in hydroboration of 4-(trifluoromethyl)benzonitrile. However, there is no reaction for substrates with nitro and pyridine groups which could be the result of inductive effects from nitrogen atoms in the backbone of the substrate which disfavor the formation of B–N bond. The mechanism of the reaction was probed through the independent preparation of the proposed boryl-imine intermediate, and the subsequent addition of a second equivalent of HBPin.<sup>69</sup>

In 2021, Ma and Yang reported *n*-BuLi (5 mol%) as an efficient catalyst in nitrile dihydroboration with HBPin under solvent-free condition. Broad substrate scope (15 examples) with good functional group compatibility were studied with NMR yield of 78–99% at 60 °C within 3–12 h. Large scale synthesis went successful with 89% yield of benzonitrile. In addition, high chemoselectivity of imine over nitrile reduction was observed under this procedure.<sup>137</sup>

Very recently in 2022, Mandal group reported a Bicyclic (alkyl)(amino)carbene (BICAAC) catalyst in metal free and solvent free hydroboration of nitriles. A wide scope of substrates including aromatic, heteroaromatic, and aliphatic nitriles (40 examples) were studied using 5 mol% catalyst with 30–92% isolated yield as ammonium salt. Gram-scale synthesis of benzonitrile was also successful with 70% isolated yield. Mechanism of the reaction studied using both experiments and DFT calculations which suggest the B–H addition to the carbene center, as hydride source.<sup>138</sup>

## 4. Main group and transition metal catalyzed hydroboration of carbodiimides

The first example of carbodiimide hydroboration was reported in 2011 by the Xia group. In this work, 1-*H*-boratabenzene rare-earth metal complexes of Yttrium (Y) or Lutetium (Lu) efficiently catalyzed hydroboration of carbodiimide substrates. The presence of a B–H bond makes this complex special as the B–H bond is much more reactive than the analogous C–H bond. Hydroboration of *N,N'*-diisopropylcarbodiimide with the Yttrium complex of  $[C_5H_5BH]_2YCl$  yields boratabenzene yttrium chloride  $[C_5H_5BN-(^iPr)CHN(^iPr)][C_5H_5BH]YCl$  in 90% at room temperature (Scheme 42). Another 1-*H*-boratabenzene complex was synthesized using lutetium,  $[C_5H_5BH]_2LuCl$ . Catalytic hydroboration of *N,N'*-diisopropyl-carbodiimide was



**Scheme 42** Hydroboration of *N,N'*-diisopropylcarbodiimide with 1-*H*-boratabenzene rare-earth metal complexes.<sup>104</sup>

also performed successfully in 87% yield at room temperature. However this report was limited to a single substrate.<sup>104</sup>

In 2016, more extensive research on carbodiimide hydroboration was reported by the Hill group. Alkyl- and aryl-substituted carbodiimides were studied using a  $\beta$ -diketiminato magnesium alkyl complex,  $[CH\{C(Me)NDipp\}_2Mg^+Bu]$ . Substrates with reduced steric hindrance provided more efficient conversion to the desired products.



Prolonged reaction times and elevated temperatures were required when using substrates with bulky groups (Table 7). The reaction is proposed to proceed through B–H/Mg–C metathesis to initiate the catalytic reaction. This is then followed by the formation of hydridomagnesium species (Scheme 43).<sup>139</sup>

In 2017, the Hill group modified their own  $\beta$ -diketiminato magnesium hydride  $[HC\{Me\}CNDipp]MgH$  catalyst *via* hydride abstraction with  $HB(C_6F_5)_3$ .<sup>140</sup> The resulting magnesium compound  $[HC\{Me\}CNDipp]_2Mg(HB(C_6F_5)_3)$  has been applied in the catalytic hydroboration of *N,N'*-diisopropyl

**Table 7** Magnesium-catalyzed hydroboration of commercially available carbodiimide<sup>139</sup>

Entry	R, R'	t (h)	T (°C)	Conv. <sup>a</sup> (%)	Yield <sup>b</sup> (%)
1	<sup>i</sup> Pr	15	60	90	89
2	Cy	15.5	60	90	69
3	<sup>t</sup> Bu	60	80	88	87
4	2,6- <i>i</i> Pr <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	24	80	80	77
5	<i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	15	80	81	63
6	<sup>t</sup> Bu, Et	20	60	75	87
7	(CH <sub>2</sub> ) <sub>3</sub> NMe <sub>2</sub> , Et	5.5	60	90	87

<sup>a</sup> Determined by <sup>1</sup>H NMR spectroscopy in C<sub>6</sub>D<sub>6</sub>. <sup>b</sup> Preparative-scale experiments performed in toluene.




**Scheme 43** Proposed mechanism in Mg catalyzed carbodiimide hydroboration.<sup>139</sup>

carbodiimide with HBpin. The product, bis(*N*-boryl)aminal  $\text{H}_2\text{C}(\text{N}\{\text{Bpin}\}^{\text{i}}\text{Pr})_2$  was obtained in >90% conversion at room temperature in 30 h or within 6 h at 60 °C.<sup>140</sup>

Virtually simultaneously, the Parkin group described the synthesis of a terminal magnesium hydride compound with tris[(1-isopropylbenzimidazol-2-yl)-dimethylsilyl]methyl ligand, [Tism<sup>PriBenz</sup>]MgH.<sup>141</sup> This complex features a carbatriene motif provided by the tris[(1-isopropylbenzimidazol-2-yl)-dimethylsilyl]methyl ligand (Fig. 12c). Significantly, [Tism<sup>PriBenz</sup>]MgH is an effective catalyst at room temperature. Hydroboration of *N,N*'-dicyclohexycarbodiimide and *N,N*'-diisopropylcarbodiimide is carried out under 10 mol% of this catalyst in a J. Young NMR tube. 1.5 eq. HBpin is used in  $\text{C}_6\text{D}_6$  as solvent in the presence of mesitylene as internal standard. The reaction is monitored by  $^1\text{H}$  NMR. Substrate consumption was complete in 1 hour and TOFs of 6.5 and 8.5  $\text{h}^{-1}$  were obtained respectively.<sup>141</sup>

In the same year, a magnesium hydridotriphenylborate  $[\text{Mg}(\text{thf})_6][\text{HBPh}_3]_2$  complex was synthesized as a solvent-separated ion pair (SSIP) as colorless crystals (Scheme 44). In this report, one example of carbodiimide hydroboration was disclosed employing this catalyst.<sup>142</sup> Dihydroboration of *N,N*-diisopropyl carbodiimide was achieved under 1 mol% catalyst loading at 60 °C within 12 h (Scheme 45).<sup>142</sup>

The first Al catalyst successfully deployed in carbodiimide hydroboration was reported in 2019.<sup>82</sup> Here, a  $\beta$ -diketiminato bis-*n*-butylaluminum of  $\text{LAl}(n\text{-Bu})_2$  ( $\text{L} = (\text{ArNCMe})_2\text{CH}$ , Ar =




**Scheme 45** Catalytic dihydroboration of *N,N*'-diisopropyl carbodiimide.<sup>142</sup>


2,6- $^{\text{i}}\text{Pr}_2\text{C}_6\text{H}_4$ ) was found to be effective in single hydroboration of carbodiimides. Hydroboration of *N,N*-diisopropylcarbodiimide in the presence of 3 mol% Al catalyst under neat conditions at 60 °C resulted in 99% yield. Furthermore, bis(*tert*-butyl)carbodiimide, bis(cyclohexyl)carbodiimide, and bis(2,6-diisopropylphenyl)-carbodiimide were reduced to the corresponding singly hydroborated *N*-borylformamidine [RNCHN(BPin)R] in excellent yields (Scheme 46).<sup>82</sup>

Roesky, Ma and Yang synthesized organoaluminum complexes (Scheme 47) which allow access to monohydroboration of commercially available carbodiimides in the presence of 1.1 eq.


HBpin, 4 mol% catalyst under neat condition at 80 °C.<sup>81</sup> Various aluminum hydride and aluminum alkyls were synthesized and investigated as potential catalysts in these transformations. Hydroborated products using four different aluminum complexes (4 mol%) under neat condition are summarized in Tables 8 and 9.<sup>81</sup> Surprisingly, the formation of mono-hydroborated C-borylformamidine product was also observed (Scheme 48) with the extension of heating time. This product has not been observed under previous reports. The reaction was proposed to proceed through an aluminum formamidate compound, which is depicted in Scheme 49.<sup>81</sup>



**Fig. 12** Highlighted reported catalysts in carbodiimide hydroboration.



**Scheme 44** Synthesis of  $[\text{Mg}(\text{thf})_6][\text{HBPh}_3]_2$  catalyst via  $\beta$ -SiH abstraction.<sup>142</sup>



**Scheme 47** Organoaluminium complexes in carbodiimide hydroboration (Ar = 2,6-Et<sub>2</sub>C<sub>6</sub>H<sub>3</sub>, Dipp).<sup>81</sup>

**Table 8** Optimization for the monohydroboration of different carbodiimides using aluminum complex as the catalyst<sup>81</sup>


Entry <sup>a</sup>	RNCNR	t (h)	T (°C)	Yield <sup>b</sup> (%)
1	iPr	12	70	79
2			80	96
3	Cy	12	70	81
4			80	97
5	tBu	60	70	34
6			80	69
7	2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	4	70	96
8		2	80	99

<sup>a</sup> Reaction conditions: 0.04 mmol catalyst, 1.1 mmol HBpin, 1 mmol carbodiimide. <sup>b</sup> <sup>1</sup>H NMR analysis.

**Table 9** Catalytic monohydroboration of different carbodiimides using aluminum complexes as a precatalyst<sup>81</sup>

Entry <sup>a</sup>	Catalyst	RNCNR	t (h)	Yield <sup>b</sup> (%)
1		iPr	24	99
2		Cy	24	95
3		tBu	40	96
4		2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	48	95
5		iPr	24	98
6		Cy	24	99
7		tBu	60	31
8		2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	36	44
9		iPr	24	93
10		Cy	24	91
11		tBu	40	73
12		2,6-iPr <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	48	77

<sup>a</sup> Reaction conditions: 0.04 mmol catalyst, 1.1 mmol HBpin, 1 mmol carbodiimide, at 80 °C. <sup>b</sup> By <sup>1</sup>H NMR analysis.




**Scheme 49** Suggested mechanism for the hydroboration of carbodiimides catalyzed by aluminum hydride and aluminum methyl complexes.<sup>81</sup>

In 2019, 9-borabicyclo[3.3.1]nonane dimer was reported as a metal-free catalyst in carbodiimide hydroboration with pinacolborane.<sup>143</sup> (H-BBN)<sub>2</sub> is usually applied as a hydroboration agent and is rarely used as a catalyst. However, this report detailed that in some cases it can be a highly active catalyst (more so than some reported metal-based catalysts).<sup>143</sup> A broad substrate scope was studied using this catalyst; including aliphatic, aromatic, sterically encumbered and asymmetric substrates. Excellent conversion of 99% was obtained in most of the substrates under low catalyst loading of 0.5–2 mol% at 25–60 °C.

The highest activity (>99% conversion) was observed in hydroboration of *N,N'*-dicyclohexycarbodiimide using 0.5 mol% catalyst in 2 h at 25 °C. The lowest conversion of 23% was observed for the substrate with SiMe<sub>3</sub> group (SiMe<sub>3</sub>N=C=NSiMe<sub>3</sub>). The reaction times varied widely 0.2–90 h (Table 10).

Stoichiometric reactions and kinetic analysis were performed to better understand the mechanism of the reaction. The boron amidinates CH(NiPr)<sub>2</sub>BBN and CH(NDipp)<sub>2</sub>BBN were prepared through the direct reaction of the corresponding carbodiimide and (H-BBN)<sub>2</sub>. Stoichiometric reactions with equimolar amounts of HBpin and boron-amidinate were carried out at room temperature as depicted in Scheme 50. Formamidinato-H-BBN adduct was not observed in the presence of CH(NDipp)<sub>2</sub>BBN which might be less energetically favored due to bulky Dipp groups.<sup>143</sup>

The experimental studies were supported by DFT calculations and resulted in the proposed mechanism depicted in Scheme 51 which involves the formation of an adduct between carbodiimide and H-BBN (**a** in Scheme 51). Followed by the corresponding formamidinate complex (**b** in Scheme 51). The formamidinate compounds can form adducts with HBpin and rearrange to the energetically more stable heterocyclic intermediate (**c** in Scheme 51). The hydride transfer to BBN leads to the formation of adduct **d**. Finally, release of the H-BBN molecule, and rotation about the C–N single bond, affords the final product.<sup>143</sup>




**Scheme 48** Hydroboration of carbodiimide in the presence of different Al complexes.<sup>81</sup>


Table 10 Monohydroboration of carbodiimides using 9-BBN<sup>143</sup>

Entry <sup>a</sup>	R, R'	Catalyst loading	t (h)	T (°C)	Conv. <sup>b</sup> (%)	RN=C=NR' + H-Bpin
						Catalyst (H-BBN) <sub>2</sub>
1	iPr	2.5	0.2	25	>99	
2	iPr	0.5	2	25	>99	
3	Cy	2.5	0.2	25	>99	
4	Cy	0.5	2	25	>99	
5	tBu	2.5	3.5	25	>99	
6	tBu	0.5	2.5	60	99	
7	p-Tol	2.5	1	60	>99	
8	p-Tol	0.5	2.5	60	>99	
9	Dipp	2.5	2.5	25	>99	
10	Dipp	0.5	2.5	60	>99	
11	SiMe <sub>3</sub>	2.5	3.5	60	23	
12	Et, tBu	2.5	20	60	>99	
13	Et, tBu	0.5	90	60	93	
14	Et, (CH <sub>2</sub> ) <sub>3</sub> NMe <sub>2</sub> <sup>c</sup>	2.5	15	60	>99	
15	Et, (CH <sub>2</sub> ) <sub>3</sub> NMe <sub>2</sub> <sup>c</sup>	0.5	20	60	>99	

<sup>a</sup> Reaction condition: (RNCR', 0.20 mmol), HBpin (0.20 mmol) (H-BBN)<sub>2</sub> as a catalyst in benzene. <sup>b</sup> Determined by <sup>1</sup>H NMR spectroscopy. <sup>c</sup> Isomer mixture 43 : 57.




Scheme 51 Proposed catalytic cycle for the monohydroboration of carbodiimides catalyzed by (H-BBN)<sub>2</sub>.<sup>143</sup>



Scheme 50 Stoichiometric reactions between (a) CH(NiPr)<sub>2</sub>BBN and HBpin, (b) CH(NDipp)<sub>2</sub>BBN and HBpin.<sup>143</sup>

In 2020, an unprecedented organometallic hafnium hydride complex was introduced by the Eisen group. This Lewis acid catalyst is effective in carbodiimide hydroboration. This complex was prepared *via* the proteolysis of hafnium tetra-*tert*-butyl and N-heterocyclic imidazolin-2-imine ligands (Scheme 52).

The electronic properties of these ancillary ligands decrease the oxophilicity of the metal center along with their steric demand which prevents possibility of the formation of bridging Hf–O–Hf macrocyclic complexes. Highly selective product formation (monoborylation) was achieved even with excess amounts of HBpin. Catalytic reactions proceed with singly reduced amidinate product formed within one hour for aliphatic carbodiimides. However, longer reaction times were required for aromatic carbodiimides. The hydroboration reaction was inhibited by the presence of electron donating substi-




Scheme 52 Synthesis of mono(imidazoline-2-iminato) hafnium(IV) complexes.<sup>144</sup>


tuents at the backbone of aromatic carbodiimides (Scheme 53). Based upon an examination of stoichiometric reactions it is proposed that the first step of the catalytic process is  $\sigma$ -bond metathesis between the hafnium complex and HBpin to form a Hf–H species. Subsequently, carbodiimide insertion takes place into the Hf–H bond which produces the hafnium amidinate complex (Scheme 54).<sup>144</sup>

Later in 2021, the Von Wangelin group reported a Mn(II) pre-catalyst supported by hmds ligands (Mn(hmds)<sub>2</sub>) in the hydroboration of a wide array of functional groups including carbodiimides.<sup>131</sup> Mn(hmds)<sub>2</sub> (5 mol%) is used in the presence of 1.1 eq. HBpin at 20–80 °C for the selective monoreduction of carbodiimides within 20 h (78–95%).

Bulky carbodiimides with tBu and Dipp groups were cleanly converted to products, 78% and 83% yield respectively but required elevated reaction temperature. It was noteworthy, that no double reduction products were observed even in the presence of excess amounts of HBpin. Regioselective hydroboration was observed with unsymmetrical carbodiimide (*N*-*tert*-butyl-*N*'-ethylcarbodiimide) which is exclusively converted to the ethylaminoborionate isomer in 92% yield.<sup>131</sup>



**Scheme 53** Monohydroboration of carbodiimides using Hf complexes.<sup>144</sup> Reaction conditions: 0.155 mmol of carbodiimide, 0.155 mmol of HBpin, and 0.0015 mmol of catalyst in 600  $\mu$ L of  $C_6D_6$  at 70 °C. The reactions are monitored by  $^1H$  NMR.



**Scheme 54** Plausible mechanism for the catalytic hydroboration of carbodiimide.<sup>144</sup>

In the same year of 2021, Ma and Yang reported *n*-BuLi as catalyst in monohydroboration of carbodiimides under neat condition at 80 °C. Both aliphatic and aromatic carbodiimides (3 examples) underwent reduction with yield of 97–99% within 24–48 h.<sup>137</sup>

The Sen group reported lithium complexes, 2,6-di-*tert*-butyl phenolate lithium and 1,1'-dilithioferrocene (Fig. 12k) in carbodiimide hydroboration in 2021. 4 examples studied using 4 mol% catalyst loading, neat condition at 80 °C with  $^1H$  NMR yield of 30–99%.<sup>145</sup>

## 5. Conclusions and outlook

In summary, we have reviewed the recent developments in the catalytic hydroboration of imines, nitriles, and carbodiimides.

Both steric and electronic features of these molecules tend to make their interactions with catalysts less favorable compared to aldehydes and ketones. In particular, the steric constraints make hydroboration challenging in nitrile substrates. HBpin, HBcat and 9-BBN have been deployed as borylating agent in most catalytic applications; within this subset of boranes reports employing HBpin are dominant, which is attributed to its Lewis acidity and relative stability which reduces the difficulty in handling other boranes which exhibit air- and moisture-instability. These hydroboration transformations underwent mainly through: oxidative addition of B–H bond,<sup>6,92,124</sup> strong base-induced nucleophilic attack,<sup>105</sup> radical pathway<sup>46,120</sup> or sigma-bond metathesis.<sup>144</sup>

Imine substrates were studied with a wide array of catalysts ranging from main group elements to transition metals. The highest reported TOF of 400  $h^{-1}$  was achieved employing zinc dihydrides catalyst with N-heterocyclic carbene ligands in the aldimine hydroboration. In contrast, amongst reported main group element catalysts, the  $\beta$ -diketiminato magnesium alkyl [ $LMg^nBu$ ] ( $L = CH[CMe(NDipp)]_2$ ) complex afforded the highest TOF (99  $h^{-1}$ ). Most transition metal systems share common mechanistic details and, at some stage of the reaction, the formation of a metal hydride species is implied. These pathways are typically supported by either experimental or computational studies. It is important to note that the underlying mechanisms of paramagnetic catalysts require further study. Despite impressive advances in the field, there are still challenges which need to be overcome: (1) there is an ongoing need to extend and develop current base metal catalysts. Clearly, ligand design principles related to non-innocence are of interest in emulating the efficacy of precious metal catalysis. (2) The use of ketimine substrates is underdeveloped compared to aldimines, often ketimines are sluggish or simply do not react under mild reaction conditions. (3) Currently, there remains only a handful of examples of chemoselective hydroboration in the presence of a large number of functional groups such as carbonyl and alkene have remained elusive. (4) Asymmetric hydroboration of imines need to be further studied.

Nitrile hydroboration has steadily gained attention in the last decade. There are many reports utilizing both transition metals and main group elements as catalysts.<sup>67,71,122,134</sup> Ru and Co catalysts are more studied among precious metals and base metal elements, respectively.<sup>47,73,75</sup> Metallacyclic thorium amide complex  $[(Me_3Si)_2N]_2Th[\kappa^2=(N,C)-CH_2Si(CH_3)_2N(SiMe_3)]$  gained one of the highest TOFs (500  $h^{-1}$ ) among transition metal catalysts.<sup>49</sup> Main group element catalysts in nitrile hydroboration are mainly dominated by Al complexes.<sup>31,82,133,135</sup> However, sodium triethylborohydride ( $NaHBET_3$ ) gained one the highest TOFs (2400  $h^{-1}$ ) among main group catalysts.<sup>69</sup> Along with the promising developments in nitrile hydroboration there remain several important challenges as follows: (1) there are few examples of rare earth metal catalysts. (2) Selectively accessing singly hydroborylated products is difficult, as most reports focus on double hydroboration of nitriles. (3) Chemoselectivity in the presence of

reducible functional groups, such as  $\text{NO}_2$ , carbonyl, alkene, ester *etc.* are limited and most reports exhibit limited tolerance (halogens, electron donating groups, and heteroatoms *etc.*). Future work is expected to focus more on designing catalysts based upon rare earth metals and employ the catalysis in both mono and dihydroboration with improvement in chemoselectivity.

Carbodiimide reduction is not very well studied, and Mg catalysts mostly dominate the field. 9-BBN which is usually reported as hydroboration reagent has been shown to act as a catalyst in carbodiimide monohydroboration. However, mono (imidazolin-2-iminato) Hafnium complex revealed to have the highest TOF ( $102 \text{ h}^{-1}$ ) among the other catalysts in carbodiimide hydroboration.<sup>144</sup> There are still challenges associated with hydroboration of these nitrogenous compounds: (1) there are only handful reports of catalysts exist which are mainly based on main group elements. (2) Substrate scope is very limited. (3) Dihydroboration of carbodiimide has not been investigated. Therefore, there are still opportunities to develop catalysts based on transition metals to achieve both mono and dihydroboration along with chemoselective studies.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was funded through a grant from the National Science Foundation, award number OAC-1934725 (AG) and a grant from the Welch Foundation, award number 23B571 (MF/ARB).

## Notes and references

- 1 H. C. Brown and B. S. Rao, *J. Am. Chem. Soc.*, 1956, **78**, 5694–5695.
- 2 J. V. Obligacion and P. J. Chirik, *Nat. Rev. Chem.*, 2018, **2**, 15–34.
- 3 A. Y. Khalimon, P. M. Farha and G. I. Nikonov, *Dalton Trans.*, 2015, **44**, 18945–18956.
- 4 I. Beletskaya and A. Pelter, *Tetrahedron*, 1997, **53**, 4957–5026.
- 5 C. E. Tucker, J. Davidson and P. Knochel, *J. Org. Chem.*, 1992, **57**, 3482–3485.
- 6 H. Kono, K. Ito and Y. Nagai, *Chem. Lett.*, 1975, **4**, 1095–1096.
- 7 D. Männig and H. Nöth, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 878–879.
- 8 V. K. Jakhar, M. K. Barman and S. Nembenna, *Org. Lett.*, 2016, **18**, 4710–4713.
- 9 J. Schneider, C. P. Sindlinger, S. M. Freitag, H. Schubert and L. Wesemann, *Angew. Chem.*, 2017, **129**, 339–343.
- 10 A. Bismuto, S. P. Thomas and M. J. Cowley, *Angew. Chem., Int. Ed.*, 2016, **55**, 15356–15359.
- 11 N. Villegas-Escobar, H. F. Schaefer III and A. Toro-Labbé, *J. Phys. Chem. A*, 2020, **124**, 1121–1133.
- 12 M. K. Sharma, M. Ansari, P. Mahawar, G. Rajaraman and S. Nagendran, *Dalton Trans.*, 2019, **48**, 664–672.
- 13 M. K. Barman, A. Baishya and S. Nembenna, *Dalton Trans.*, 2017, **46**, 4152–4156.
- 14 A. Singh, S. Shafiei-Haghghi, C. R. Smith, D. K. Unruh and M. Findlater, *Asian J. Org. Chem.*, 2020, **9**, 416–420.
- 15 S. R. Tamang and M. Findlater, *J. Org. Chem.*, 2017, **82**, 12857–12862.
- 16 S. R. Tamang, D. Bedi, S. Shafiei-Haghghi, C. R. Smith, C. Crawford and M. Findlater, *Org. Lett.*, 2018, **20**, 6695–6700.
- 17 S. R. Tamang and M. Findlater, *Dalton Trans.*, 2018, **47**, 8199–8203.
- 18 S. R. Tamang, A. Singh, D. K. Unruh and M. Findlater, *ACS Catal.*, 2018, **8**, 6186–6191.
- 19 R. J. Newland, J. M. Lynam and S. M. Mansell, *ChemComm*, 2018, **54**, 5482–5485.
- 20 A. Baishya, S. Baruah and K. Geetharani, *Dalton Trans.*, 2018, **47**, 9231–9236.
- 21 L. Wenbo and L. Zhan, *Chin. J. Org. Chem.*, 2020, **40**, 3596–3604.
- 22 J. Chen, J. Guo and Z. Lu, *Chin. J. Chem.*, 2018, **36**, 1075–1109.
- 23 W. Fan, L. Li and G. Zhang, *J. Org. Chem.*, 2019, **84**, 5987–5996.
- 24 J. Guo, Z. Cheng, J. Chen, X. Chen and Z. Lu, *Acc. Chem. Res.*, 2021, **54**, 6452–6455.
- 25 S. R. Tamang and M. Findlater, *Molecules*, 2019, **24**, 3194–3218.
- 26 K. Saha and S. Ghosh, *Dalton Trans.*, 2022, **51**, 2631–2640.
- 27 R. Mesnage, C. Benbrook and M. N. Antoniou, *Food Chem. Toxicol.*, 2019, **128**, 137–145.
- 28 M. A. Blaskovich, *J. Med. Chem.*, 2016, **59**, 10807–10836.
- 29 R. Jastrząb, L. Łomozik and B. Tylkowski, *Phys. Sci. Rev.*, 2016, **1**, 0003–0026.
- 30 A. Yang, C. Ching, M. Easler, D. E. Helbling and W. R. Dichtel, *ACS Mater. Lett.*, 2020, **2**, 1240–1245.
- 31 W. Liu, Y. Ding, D. Jin, Q. Shen, B. Yan, X. Ma and Z. Yang, *Green Chem.*, 2019, **21**, 3812–3815.
- 32 C. Gunanathan, M. Hölscher and W. Leitner, *Eur. J. Inorg. Chem.*, 2011, **2011**, 3381–3386.
- 33 M. S. Newman and T. Fukunaga, *J. Am. Chem. Soc.*, 1960, **82**, 693–696.
- 34 Y. Liu and H. Du, *J. Am. Chem. Soc.*, 2013, **135**, 6810–6813.
- 35 H. Li, A. Al-Dakhil, D. Lupp, S. S. Gholap, Z. Lai, L.-C. Liang and K.-W. Huang, *Org. Lett.*, 2018, **20**, 6430–6435.
- 36 S. Chakraborty and D. Milstein, *ACS Catal.*, 2017, **7**, 3968–3972.

37 G. Lu, P. Zhang, D. Sun, L. Wang, K. Zhou, Z.-X. Wang and G.-C. Guo, *Chem. Sci.*, 2014, **5**, 1082–1090.

38 S. Chakraborty and H. Berke, *ACS Catal.*, 2014, **4**, 2191–2194.

39 S. Werkmeister, K. Junge and M. Beller, *Org. Process Res. Dev.*, 2014, **18**, 289–302.

40 J. Long, B. Yin, Y. Li and L. Zhang, *AIChE J.*, 2014, **60**, 3565–3576.

41 J. Long, K. Shen and Y. Li, *ACS Catal.*, 2017, **7**, 275–284.

42 H. Bauer, M. Alonso, C. Färber, H. Elsen, J. Pahl, A. Causero, G. Ballmann, F. De Proft and S. Harder, *Nat. Catal.*, 2018, **1**, 40–47.

43 S. R. Tamang, A. Singh, D. Bedi, A. R. Bazkiae, A. A. Warner, K. Glogau, C. McDonald, D. K. Unruh and M. Findlater, *Nat. Catal.*, 2020, **3**, 154–162.

44 R. T. Baker, J. C. Calabrese and S. A. Westcott, *J. Organomet. Chem.*, 1995, **498**, 109–117.

45 C. M. Vogels, P. E. O'Connor, T. E. Phillips, K. J. Watson, M. P. Shaver, P. G. Hayes and S. A. Westcott, *Can. J. Chem.*, 2001, **79**, 1898–1905.

46 N. Zhou, X. A. Yuan, Y. Zhao, J. Xie and C. Zhu, *Angew. Chem.*, 2018, **130**, 4054–4058.

47 A. Kaithal, B. Chatterjee and C. Gunanathan, *J. Org. Chem.*, 2016, **81**, 11153–11161.

48 L. Koren-Selfridge, H. N. Londino, J. K. Vellucci, B. J. Simmons, C. P. Casey and T. B. Clark, *Organometallics*, 2009, **28**, 2085–2090.

49 S. Saha and M. S. Eisen, *ACS Catal.*, 2019, **9**, 5947–5956.

50 Z. Huang, S. Wang, X. Zhu, Q. Yuan, Y. Wei, S. Zhou and X. Mu, *Inorg. Chem.*, 2018, **57**, 15069–15078.

51 A. E. King, S. C. E. Stieber, N. J. Henson, S. A. Kozimor, B. L. Scott, N. C. Smythe, A. D. Sutton and J. C. Gordon, *Eur. J. Inorg. Chem.*, 2016, **2016**, 1635–1640.

52 J. Wu, H. Zeng, J. Cheng, S. Zheng, J. A. Golen, D. R. Manke and G. Zhang, *J. Org. Chem.*, 2018, **83**, 9442–9448.

53 M. Arrowsmith, M. S. Hill and G. Kociok-Köhn, *Chem. – Eur. J.*, 2013, **19**, 2776–2783.

54 V. A. Pollard, M. Á. Fuentes, A. R. Kennedy, R. McLellan and R. E. Mulvey, *Angew. Chem., Int. Ed.*, 2018, **57**, 10651–10655.

55 M. R. Adams, C. H. Tien, R. McDonald and A. W. Speed, *Angew. Chem.*, 2017, **129**, 16887–16890.

56 C.-H. Tien, M. R. Adams, M. J. Ferguson, E. R. Johnson and A. W. Speed, *Org. Lett.*, 2017, **19**, 5565–5568.

57 J. R. Lawson, L. C. Wilkins and R. L. Melen, *Chemistry*, 2017, **23**, 10997–11000.

58 Q. Yin, Y. Soltani, R. L. Melen and M. Oestreich, *Organometallics*, 2017, **36**, 2381–2384.

59 X. Zhu and H. Du, *Org. Biomol. Chem.*, 2015, **13**, 1013–1016.

60 C. Bakewell, *Dalton Trans.*, 2020, **49**, 11354–11360.

61 M. R. Adams, C. H. Tien, B. S. Huchenski, M. J. Ferguson and A. W. Speed, *Angew. Chem., Int. Ed.*, 2017, **56**, 6268–6271.

62 T. K. Panda, I. Banerjee and S. Sagar, *Appl. Organomet. Chem.*, 2020, **34**, 5765–5775.

63 X. He, B. Yan, C. Ni, Y. Zhao, Z. Yang and X. Ma, *Asian J. Org. Chem.*, 2021, **10**, 196–201.

64 K. Nakaya, S. Takahashi, A. Ishii, K. Boonpalit, P. Surawatanawong and N. Nakata, *Dalton Trans.*, 2021, **50**, 14810–14819.

65 C. R. Aversa-Fleener, D. K. Chang and A. L. Liberman-Martin, *Organometallics*, 2021, **40**, 4050–4054.

66 V. K. Pandey, S. N. R. Donthireddy and A. Rit, *Chem. – Asian J.*, 2019, **14**, 3255–3258.

67 A. Y. Khalimon, P. Farha, L. G. Kuzmina and G. I. Nikonov, *ChemComm*, 2012, **48**, 455–457.

68 C. J. Major, K. L. Bamford, Z.-W. Qu and D. W. Stephan, *ChemComm*, 2019, **55**, 5155–5158.

69 D. Bedi, A. Brar and M. Findlater, *Green Chem.*, 2020, **22**, 1125–1128.

70 A. B. Prasad, J. B. Kanth and M. Periasamy, *Tetrahedron*, 1992, **48**, 4623–4628.

71 C. Weetman, M. D. Anker, M. Arrowsmith, M. S. Hill, G. Kociok-Köhn, D. J. Liptrot and M. F. Mahon, *Chem. Sci.*, 2016, **7**, 628–641.

72 A. Harinath, J. Bhattacharjee and T. K. Panda, *Adv. Synth. Catal.*, 2019, **361**, 850–857.

73 H. Ben-Daat, C. L. Rock, M. Flores, T. L. Groy, A. C. Bowman and R. J. Trovitch, *ChemComm*, 2017, **53**, 7333–7336.

74 T. Kitano, T. Komuro and H. Tobita, *Organometallics*, 2019, **38**, 1417–1420.

75 J. B. Geri and N. K. Szymczak, *J. Am. Chem. Soc.*, 2015, **137**, 12808–12814.

76 D. Hayrapetyan and A. Y. Khalimon, *Chem. – Asian J.*, 2020, **15**, 2575–2587.

77 J. Bhattacharjee, A. Harinath, K. Bano and T. K. Panda, *ACS Omega*, 2020, **5**, 1595–1606.

78 R. K. Sahoo, N. Sarkar and S. Nembenna, *Angew. Chem.*, 2021, **133**, 12098–12107.

79 Y. Yang, M. D. Anker, J. Fang, M. F. Mahon, L. Maron, C. Weetman and M. S. Hill, *Chem. Sci.*, 2017, **8**, 3529–3537.

80 R. Boese, R. Köster and M. Yalpani, *Z. Naturforsch., B: J. Chem. Sci.*, 1994, **49**, 1453–1458.

81 Q. Shen, X. Ma, W. Li, W. Liu, Y. Ding, Z. Yang and H. W. Roesky, *Chem. – Eur. J.*, 2019, **25**, 11918–11923.

82 Y. Ding, X. Ma, Y. Liu, W. Liu, Z. Yang and H. W. Roesky, *Organometallics*, 2019, **38**, 3092–3097.

83 M. D. Greenhalgh, A. S. Jones and S. P. Thomas, *ChemCatChem*, 2015, **7**, 190–222.

84 E. A. Redina, K. V. Vikanova, G. I. Kapustin, I. V. Mishin, O. P. Tkachenko and L. M. Kustov, *Eur. J. Org. Chem.*, 2019, 4159–4170.

85 A. C. F. Caires, A. E. Mauro, A. C. Moro, A. d. O. Legendre and S. R. Ananias, *Quim. Nova*, 2006, **29**, 750–754.

86 V. Lyaskovskyy and B. de Bruin, *ACS Catal.*, 2012, **2**, 270–279.

87 C. C. Chong and R. Kinjo, *ACS Catal.*, 2015, **5**, 3238–3259.

88 S. Diez-Gonzalez and S. P. Nolan, *Org. Prep. Proced. Int.*, 2007, **39**, 523–559.

89 K. Riener, M. P. Högerl, P. Gigler and F. E. Kühn, *ACS Catal.*, 2012, **2**, 613–621.

90 J. Jiao, K. Murakami and K. Itami, *ACS Catal.*, 2016, **6**, 610–633.

91 Y. Park, Y. Kim and S. Chang, *Chem. Rev.*, 2017, **117**, 9247–9301.

92 T. M. Cameron, R. T. Baker and S. A. Westcott, *ChemComm*, 1998, 2395–2396.

93 A. C. Fernandes, J. A. Fernandes, F. A. A. Paz and C. C. Romão, *Dalton Trans.*, 2008, 6686–6688.

94 R. Arévalo, C. M. Vogels, G. A. MacNeil, L. Riera, J. Pérez and S. A. Westcott, *Dalton Trans.*, 2017, **46**, 7750–7757.

95 P. T. Anastas and M. M. Kirchhoff, *Acc. Chem. Res.*, 2002, **35**, 686–694.

96 B. L. Lin, C. R. Clough and G. L. Hillhouse, *J. Am. Chem. Soc.*, 2002, **124**, 2890–2891.

97 T. J. Deming, *J. Am. Chem. Soc.*, 1998, **120**, 4240–4241.

98 H. Hoberg and A. Herrera, *Angew. Chem.*, 1980, **92**, 951–952.

99 L. Stehling and G. Wilke, *Angew. Chem.*, 1985, **97**, 505–506.

100 I. Hossain and J. Schmidt, *Eur. J. Inorg. Chem.*, 2020, **2020**, 1877–1884.

101 C. K. Blasius, N. F. Heinrich, V. Vasilenko and L. H. Gade, *Angew. Chem., Int. Ed.*, 2020, **59**, 15974–15977.

102 A. R. Bazkiae, M. Wiseman and M. Findlater, *RSC Adv.*, 2021, **11**, 15284–15289.

103 X. M. Wang and X. Xu, *RSC Adv.*, 2021, **11**, 1128–1133.

104 Y. Yuan, X. Wang, Y. Li, L. Fan, X. Xu, Y. Chen, G. Li and W. Xia, *Organometallics*, 2011, **30**, 4330–4341.

105 P. Eisenberger, A. M. Bailey and C. M. Crudden, *J. Am. Chem. Soc.*, 2012, **134**, 17384–17387.

106 M. Arrowsmith, T. J. Hadlington, M. S. Hill and G. Kociok-Köhn, *ChemComm*, 2012, **48**, 4567–4569.

107 M. Arrowsmith, M. S. Hill, T. Hadlington, G. Kociok-Köhn and C. Weetman, *Organometallics*, 2011, **30**, 5556–5559.

108 S. S. Sen, H. W. Roesky, D. Stern, J. Henn and D. Stalke, *J. Am. Chem. Soc.*, 2010, **132**, 1123–1126.

109 M. K. Bisai, S. Pahar, T. Das, K. Vanka and S. S. Sen, *Dalton Trans.*, 2017, **46**, 2420–2424.

110 J. R. Lawson, L. C. Wilkins and R. L. Melen, *Chemistry*, 2017, **23**, 10997.

111 T. J. Herrington, A. J. Thom, A. J. White and A. E. Ashley, *Dalton Trans.*, 2012, **41**, 9019–9022.

112 H. Minami, T. Saito, C. Wang and M. Uchiyama, *Angew. Chem.*, 2015, **127**, 4748–4751.

113 Z. Yang, M. Zhong, X. Ma, S. De, C. Anusha, P. Parameswaran and H. W. Roesky, *Angew. Chem.*, 2015, **127**, 10363–10367.

114 D. Yan, X. Wu, J. Xiao, Z. Zhu, X. Xu, X. Bao, Y. Yao, Q. Shen and M. Xue, *Org. Chem. Front.*, 2019, **6**, 648–653.

115 T. Lundrigan, E. N. Welsh, T. Hynes, C.-H. Tien, M. R. Adams, K. R. Roy, K. N. Robertson and A. W. Speed, *J. Am. Chem. Soc.*, 2019, **141**, 14083–14088.

116 Z. Zhang, S. Huang, L. Huang, X. Xu, H. Zhao and X. Yan, *J. Org. Chem.*, 2020, **85**, 12036–12043.

117 H. Kim, H. T. Kim, J. H. Lee, H. Hwang and D. K. An, *RSC Adv.*, 2020, **10**, 34421–34427.

118 H. Kim, J. H. Lee, H. Hwang and D. K. An, *New J. Chem.*, 2020, **44**, 11330–11335.

119 B. S. Huchenski and A. W. Speed, *Org. Biomol. Chem.*, 2019, **17**, 1999–2004.

120 T. Kawamoto, T. Morioka, K. Noguchi, D. P. Curran and A. Kamimura, *Org. Lett.*, 2021, **23**, 1825–1828.

121 M. M. Kamal, Z. Liu, S. Zhai and D. Vidović, *Molecules*, 2021, **26**, 5443–5452.

122 M. Ito, M. Itazaki and H. Nakazawa, *Inorg. Chem.*, 2017, **56**, 13709–13714.

123 A. D. Ibrahim, S. W. Entsminger and A. R. Fout, *ACS Catal.*, 2017, **7**, 3730–3734.

124 G. Nakamura, Y. Nakajima, K. Matsumoto, V. Srinivas and S. Shimada, *Catal. Sci. Technol.*, 2017, **7**, 3196–3199.

125 N. Sarkar, M. Mahato and S. Nembenna, *Eur. J. Inorg. Chem.*, 2020, **2020**, 2295–2301.

126 C. Ghosh, S. Kim, M. R. Mena, J.-H. Kim, R. Pal, C. L. Rock, T. L. Groy, M.-H. Baik and R. J. Trovitch, *J. Am. Chem. Soc.*, 2019, **141**, 15327–15337.

127 I. Banerjee, S. Anga, K. Bano and T. K. Panda, *J. Organomet. Chem.*, 2019, **902**, 120958–120967.

128 K. A. Gudun, A. Slamova, D. Hayrapetyan and A. Y. Khalimon, *Chem. – Eur. J.*, 2020, **26**, 4963–4968.

129 C. Li, S. Song, Y. Li, C. Xu, Q. Luo, Y. Guo and X. Wang, *Nat. Commun.*, 2021, **12**, 3813–3821.

130 R. Thenarukandiyil, V. Satheesh, L. J. Shimon and G. de Ruiter, *Chem. – Asian J.*, 2021, **16**, 999–1006.

131 P. Ghosh and A. Jacobi von Wangenheim, *Angew. Chem., Int. Ed.*, 2021, **60**, 16035–16043.

132 J. Bhattacharjee, D. Bockfeld and M. Tamm, *J. Org. Chem.*, 2022, **87**, 1098–1109.

133 T. K. Panda, A. Harinath and J. Bhattacharjee, *Adv. Synth. Catal.*, 2018, **47**, 12613–12622.

134 J. Li, M. Luo, X. Sheng, H. Hua, W. Yao, S. A. Pullarkat, L. Xu and M. Ma, *Org. Chem. Front.*, 2018, **5**, 3538–3547.

135 N. Sarkar, S. Bera and S. Nembenna, *J. Org. Chem.*, 2020, **85**, 4999–5009.

136 P. Ghosh and A. J. von Wangenheim, *Org. Chem. Front.*, 2020, **7**, 960–966.

137 B. Yan, X. He, C. Ni, Z. Yang and X. Ma, *ChemCatChem*, 2021, **13**, 851–854.

138 S. K. Mandal, N. Gautam, R. Logdi, P. Sreejyothi, N. Rajendran and A. Tiwari, *ChemComm*, 2022, 1104–1108.

139 C. Weetman, M. S. Hill and M. F. Mahon, *Chem. – Eur. J.*, 2016, **22**, 7158–7162.

140 M. D. Anker, M. Arrowsmith, R. L. Arrowsmith, M. S. Hill and M. F. Mahon, *Inorg. Chem.*, 2017, **56**, 5976–5983.

141 M. Rauch, S. Ruccolo and G. Parkin, *J. Am. Chem. Soc.*, 2017, **139**, 13264–13267.

142 D. Mukherjee, S. Shirase, T. P. Spaniol, K. Mashima and J. Okuda, *ChemComm*, 2016, **52**, 13155–13158.

143 A. Ramos, A. Antiñolo, F. Carrillo-Hermosilla, R. Fernández-Galán and D. García-Vivó, *ChemComm*, 2019, **55**, 3073–3076.

144 M. Khononov, N. Fridman, M. Tamm and M. S. Eisen, *Eur. J. Org. Chem.*, 2020, 3153–3160.

145 M. K. Bisai, K. Gour, T. Das, K. Vanka and S. S. Sen, *J. Organomet. Chem.*, 2021, **949**, 121924–121930.